

Hospital Readmission Analysis Dashboard

Total Encounters	Unique Patients	30-Day Readmission Rate
101766	71518	11.16


KEY INSIGHTS

- Patients with 3+ emergency visits showed highest readmission rates
- Short hospital stays were associated with increased readmission risk
- Diabetes patients had slightly higher readmission percentages
- Age groups 30–80 demonstrated elevated readmission trends
- Medium medication count patients represented largest readmission volume


RECOMMENDATIONS

- Improve follow-up monitoring for high emergency visit patients
- Strengthen discharge planning for short-stay patients
- Prioritize preventive care for diabetes patients
- Focus care management on high-risk age groups
- Monitor medication adherence to reduce readmission risk


Readmission Breakdown




30-Day Readmission Rate by Age Group (%)




30-Day Readmission Rate by Length of Stay (%)



Readmission Rate by Emergency Visits (%)




Medication Impact on Readmission



Readmission Distribution by Diabetes Status



30-Day Readmission Rate by A1C Result



Highest Risk Patient Segments

Age	Stay Group	Emergency V Patients	Readmitted	Readmission
[30-40]	Short Stay (1-3+ Emergency)	101	34	33.66
[70-80]	Short Stay (1-3+ Emergency)	103	31	30.1
[40-50]	Short Stay (1-3+ Emergency)	183	46	25.14
[40-50]	Medium Stay (3+ Emergency)	132	30	22.73

Patients with 3+ emergency visits and short hospital stays showed the highest readmission risk, especially among ages 30–80.