

A Machine Learning Approach to Urban Growth Prediction of Juba, South Sudan (2024-2033)

NYOMBE BENSON JALLE 17-CIT-104

Supervised By: Dr. Manzu Gerald Kenyi, Ph.D

**Department of Information Technology
School of Computer Science and Information Technology,
University of Juba**



**Graduation Thesis submitted in partial fulfillment of the requirements leading to the award
of Bachelor of Science (Hons) in Information Technology**

March 2025

Abstract.

Urban growth in rapidly developing cities, such as Juba in South Sudan, poses significant challenges for sustainable development and effective urban planning. This study aims to predict the future patterns of urban growth in Juba up to the year 2040 by analyzing historical land use and land cover (LULC) changes from 1990 to 2023 using advanced machine learning and geospatial techniques. The methodology integrates satellite imagery, Geographic Information systems (GIS), and predictive modeling approaches, including the random forest model, to forecast urban expansion. This model allows classification and the simulation of spatial dynamics and the estimation of transition probabilities between different land use types, providing a robust framework for understanding urban growth trajectories.

The study will employ a Random Forest classifier to accurately classify LULC into distinct categories, followed by rigorous validation using metrics such as overall accuracy, kappa coefficient, and ROC-AUC. Scenario analysis further explores the impact of various factors, such as population growth and infrastructure development, on future urban expansion. The findings are expected to offer critical insights for urban planners and policymakers in Juba, enabling them to make informed decisions that balance developmental needs with environmental sustainability. This research contributes to the growing body of knowledge on urban growth prediction in rapidly expanding cities, with practical implications for managing urban sprawl and ensuring sustainable development in Juba and similar contexts.

Map showing the region of interest.




Table of contents.

1. Chapter one: Introduction.	4
1.1 Background	4
1.2 Problem Statement	6
1.3 Objectives of the Study	Error! Bookmark not defined.
1.3.1 Primary Objective	10
1.3.2 Specific Objectives	10
1.4 Significance of Study	10
1.5 Research Questions	11
1.6 Limitations and scope of the study.	12
Chapter 2: Literature review.	12
2.1 Introduction	13
2.2 Land Use and Land Cover (LULC) Dynamics in Juba	13
2.3 Methodologies for Urban Growth Prediction	13
2.3.1 Remote Sensing and GIS	Error! Bookmark not defined.
2.3.2 Cellular Automata (CA) Models	17
2.3.3 Markov Chain Models	18
2.3.4 Artificial Neural Networks (ANN)	16
2.3.5 Logistic Regression	18
2.4 Contribution of the Current Study	20
Chapter 3: Materials and Methods	20
3.1 Overview	22
3.2 Data Collection	22
3.3 Data Preprocessing	Error! Bookmark not defined.
3.4 Land Use and Land Cover Classification	Error! Bookmark not defined.
3.5 Urban Growth Modeling	Error! Bookmark not defined.
3.6 Validation and Accuracy Assessment	32
3.7 Scenario Analysis	Error! Bookmark not defined.
3.8 Software and Tools	Error! Bookmark not defined.

Chapter one: Introduction.

Background

Urbanization is a defining characteristic of modern development, particularly in rapidly expanding cities within developing nations, like Juba in South Sudan. The process of urban growth introduces both opportunities and challenges, shaping the trajectory of economic and social progress ([Justin, 2017; Justin and and De Vries, 2019](#)).

In regions such as South Sudan, the pace of urbanization is accelerated by a combination of population growth, migration trends, and economic transitions. While this expansion fosters economic opportunities and infrastructural advancements, it also places considerable strain on essential resources, including housing, transportation networks, public utilities, and environmental sustainability ([Zeng et al., 2016; Jiang and Lv, 2022](#)).

In many developing countries, the challenges associated with urban expansion are compounded by weak governance structures, fragile institutional frameworks, and a lack of comprehensive urban planning mechanisms. These factors contribute to the unregulated sprawl of cities, making it increasingly difficult to implement effective development strategies ([Zeng et al., 2016; Justin, 2017; Wild, Jok and Patel, 2018; Jiang and Lv, 2022](#)).

Addressing these complexities requires a multi-faceted approach that integrates sustainability principles while simultaneously tackling pressing concerns such as affordable housing, efficient public transportation, equitable resource distribution, and social inclusion. Aligning urban development strategies with global frameworks, particularly the United Nations Sustainable Development Goals (SDGs), is crucial in fostering long-term sustainability. In this context, SDG 11 (sustainable cities and communities), SDG 6 (clean water and sanitation), and SDG 13 (climate action) serve as guiding principles for ensuring balanced urban expansion ([Zeng et al., 2016; Magbool et al., 2020](#)).

Historical Context of South Sudan's Urban Growth

The urbanization trajectory of South Sudan is deeply intertwined with its socio-political history. Decades of conflict and political instability have shaped the nation's development, culminating in its secession from Sudan. On July 9, 2011, South Sudan officially became an independent nation following a referendum, making it the world's newest sovereign state and Africa's 54th country.

This historic event marked the beginning of a new chapter for the nation, one defined by both aspirations for progress and the daunting task of post-conflict reconstruction ([UN-HABITAT *et al.*, 2023](#)). As a newly established country emerging from prolonged civil wars, South Sudan has faced significant hurdles in building its economy, infrastructure, and governance structures. The post-independence period has been characterized by efforts to stabilize the nation, foster economic growth, and rebuild critical institutions. However, these ambitions have been met with persistent challenges, including ongoing political tensions, economic instability, and weak governance capacity ([UN-HABITAT *et al.*, 2023](#)).

Juba: South Sudan's Epicenter of Rapid Urbanization

Juba, the capital and largest city of South Sudan, has been at the forefront of the country's urban expansion, experiencing substantial demographic and infrastructural transformations ([Justin, 2017; Justin and De Vries, 2019; Salvatore *et al.*, 2023](#)). The city's growth has been driven by multiple interrelated factors, including rural-to-urban migration, economic activities, and large-scale reconstruction efforts following years of civil conflict. After South Sudan gained independence, Juba became a major destination for displaced populations, returning refugees, and migrants seeking employment, security, and access to essential services ([UN-HABITAT *et al.*, 2023](#)). The influx of people into Juba has led to a surge in housing demand, transforming the city's spatial landscape. Previously undeveloped areas and agricultural lands are being rapidly converted into residential, commercial, and industrial zones to accommodate the growing population ([UN-HABITAT *et al.*, 2023](#)). However, this expansion has largely been unplanned, resulting in sprawling informal settlements that lack adequate infrastructure and basic services. The unregulated nature of urban expansion has created significant challenges for city planners, including inefficient land use, encroachments on environmentally sensitive areas, and the proliferation of slums ([Justin and De Vries, 2019; UN-HABITAT *et al.*, 2023](#)).

Infrastructure Struggles and Urban Challenges

As Juba continues to expand, its infrastructure development has struggled to keep pace with the growing demands of its residents. The city faces critical deficits in essential urban services, including roads, housing, sanitation systems, and public transportation. Many neighborhoods experience chronic congestion due to poorly maintained road networks and inadequate transportation options. The lack of a reliable waste management system has exacerbated environmental pollution, while deficiencies in water supply and electricity distribution remain

major concerns for both residents and businesses ([UN-HABITAT et al., 2023](#)).

Furthermore, Juba's evolving urban environment reflects stark contrasts between modern development and persistent infrastructural deficits. High-rise buildings, commercial hubs, and new business centers are emerging alongside informal settlements, highlighting the disparities in urban planning and resource distribution. The rapid transformation of Juba's cityscape underscores both the opportunities and challenges associated with urban growth. On one hand, the city's expansion presents prospects for economic development, increased investments, and improved public services. On the other hand, the lack of coordinated planning and sustainable urban policies threatens long-term resilience and livability ([UN-HABITAT et al., 2023](#)).

Therefore, Juba's rapid urbanization calls for urgent need for comprehensive and sustainable urban planning. As the city continues to grow, it is imperative to implement policies that balance development with environmental conservation, social equity, and infrastructural efficiency. Addressing these challenges requires a collaborative approach involving government agencies, urban planners, and international development organizations. By integrating modern urban planning techniques, investing in resilient infrastructure, and aligning growth strategies with global sustainability frameworks, Juba can navigate its urbanization trajectory in a manner that promotes long-term prosperity and stability

1.2 Problem Statement

Urban growth, as conceptualized by the United Nations Human Settlements Program (UN-Habitat), refers to the progressive expansion of towns and cities in terms of both spatial extent and population size ([UN-HABITAT, 2006](#)). This dynamic phenomenon unfolds through two primary mechanisms: inward growth, which is characterized by an increase in population density and intensified land use within pre-existing urban boundaries, and outward growth, which involves the physical expansion of urban areas into surrounding rural or undeveloped regions. These growth patterns are particularly pronounced in rapidly urbanizing regions, where economic, social, and demographic factors collectively contribute to the transformation of urban landscapes ([Sexton et al., 2013](#)). However, urban growth is a multifaceted process that extends beyond simple expansion, significantly influencing both human settlements and the environment at large ([Munthali and Murayama, 2011](#); [Khawaldah, Farhan and Alzboun, 2020](#)).

Urban growth and land use changes

As urban areas expand, they inevitably encroach upon previously undeveloped lands, leading to substantial alterations in Land Use and Land Cover (LULC). This transformation, broadly referred to as Land Use and Land Cover Change (LULCC), denotes modifications to the Earth's surface brought about by both human activities and natural processes over time and space ([Rodríguez-Cuenca, Alonso and Tamés-Noriega, 2015](#)). The driving forces behind LULCC include the growing demand for residential, commercial, and infrastructural development, which often results in the conversion of agricultural land and natural ecosystems into urbanized spaces.

Research indicates that urban areas across the globe are expanding at an unprecedented pace, with projections forecasting further and more extensive growth in the coming decades ([Ishtiaque, Shrestha and Chhetri, 2017](#)). Given the rapid nature of urbanization, the necessity of monitoring and predicting LULCC through change detection techniques has become increasingly crucial. Change detection methodologies facilitate the systematic tracking, analysis, and management of evolving spatial distributions of land resources, enabling informed decision-making for sustainable urban planning ([Turner and Ruscher, 1988](#)). Failure to adequately monitor these transformations can lead to haphazard urban expansion, inefficient infrastructure deployment, and heightened susceptibility to environmental hazards such as flooding, pollution, and resource depletion.

Rapid urban expansion in Juba

Juba, the capital city of South Sudan, serves as a prime example of a rapidly urbanizing metropolis experiencing both densification within its core and sprawling expansion into peripheral areas. The surge in population within the city's established boundaries has exerted immense pressure on public infrastructure and essential services, leading to a multitude of urban challenges, including overcrowding, traffic congestion, inadequate sanitation, and public health concerns. Simultaneously, the outward expansion of Juba has resulted in the transformation of surrounding agricultural land, forests, and wetlands into built-up areas, thereby significantly disrupting local ecosystems and biodiversity ([Gudo, Deng and Qureshi, 2022a](#)).

One of the most pressing challenges in managing Juba's urban growth is the lack of comprehensive and up-to-date spatial data on LULCC ([Gudo, Deng and Qureshi, 2022b](#); [UN-HABITAT et al., 2023](#)). The city's rapid and largely unregulated expansion, coupled with insufficient monitoring systems, presents substantial difficulties for urban planners and policymakers attempting to anticipate and address the implications of land use changes on infrastructure development and environmental sustainability. Consequently, large swathes of agricultural and natural lands are

being converted into urban spaces, leading to altered hydrological patterns, exacerbated environmental degradation, and increased pollution levels([Gudo, Deng and Qureshi, 2022b](#)).

Challenges in urban monitoring and sustainable development

Traditional methods of monitoring urban growth, such as manual land surveys and historical satellite imagery analysis, often fall short in capturing the complexities associated with rapid urbanization, particularly in cities characterized by informal settlements and unregulated construction. The limitations inherent in these conventional approaches underscore the growing necessity for more sophisticated change detection methodologies and geospatial analytical techniques to effectively map and predict urban expansion patterns ([Ishtiaque, Shrestha and Chhetri, 2017](#); [Khawaldah, Farhan and Alzboun, 2020](#)) .

Emerging technologies, such as machine learning, remote sensing, and Geographic Information Systems (GIS), have proven to be invaluable tools for urban planners in their efforts to assess real-time land cover changes, detect informal settlements, and identify infrastructural deficits. The integration of these advanced analytical techniques can significantly enhance the efficiency of land-use planning, risk assessment, and environmental management, ensuring that urban expansion is conducted in a manner that is both sustainable and adaptive to future challenges.

Environmental and social-economic challenges

The unregulated and unchecked expansion of urban areas presents a host of human and ecological challenges. On the human side, the continuous influx of people into urban environments necessitates the provision of adequate housing, efficient transportation networks, and robust social services, all of which must evolve in tandem with growing demand. Failure to adequately address these needs can precipitate housing shortages, inefficient public transportation systems, and widening socio-economic disparities.

From an environmental perspective, rapid urbanization results in the depletion of natural resources, habitat loss, deforestation, increased surface temperatures, and alterations in local climate patterns. Additionally, the large-scale conversion of natural landscapes into impervious urban surfaces has far-reaching consequences for hydrological systems, disrupting water cycles, exacerbating flood risks, and contributing to groundwater depletion. These ecological disruptions pose serious long-term sustainability challenges, emphasizing the urgency of adopting urban planning strategies that strike a balance between developmental needs and environmental conservation ([Munthali and Murayama, 2011](#); [Sexton et al., 2013](#); [Munthali et al., 2019](#); [Khawaldah, Farhan and Alzboun,](#)

[2020](#)).

Summary

Like other rapidly evolving urban cities in developing countries, Juba's ongoing urban expansion can be seen in a two-faced reality: on one hand, it signifies a profound opportunity for economic development, while on the other, it poses significant challenges for sustainable resource management. Rapid urbanization, if unchecked, can lead to issues such as infrastructure strain, environmental degradation, and the displacement of vulnerable populations. Therefore, the effective prediction of urban growth is essential for informed decision-making and sustainable development. To mitigate the adverse consequences related to rapid urbanization, it is crucial to implement integrated urban planning strategies that prioritize the development of resilient infrastructure, efficient land-use policies, and climate-conscious urban designs. Leveraging cutting-edge geospatial technologies, remote sensing data, and predictive analytics can play a pivotal role in this effort.

In this study, we propose the prediction of Juba's urban growth from 2024 to 2033 based on Random Forest, a powerful and widely used Machine Learning model for spatial prediction tasks (mainly characterized into classification and regression tasks). By analyzing Landsat 5 and 7 data spanning from 1994 to 2023, this approach allows us to predict land-use changes, and urban sprawl with high accuracy. Random Forest excels in handling large, complex datasets and capturing non-linear relationships between urban growth drivers, such as population density, land cover changes, and socioeconomic factors.

When trained on historical land-use patterns, and by recognizing these patterns, the Random Forest model can predict future urban growth trends. These predictions are vital for urban planners and policymakers to better monitor, forecast, and guide Juba's expansion in a manner that fosters economic growth while minimizing environmental and social challenges.

Achieving sustainable urban development in Juba will necessitate collaborative efforts between governmental agencies, academic researchers such as in this study, and local communities, ensuring that urban expansion aligns with broader sustainability objectives and contributes to the overall well-being of its inhabitants. Only through proactive and data-driven urban management approaches, such as this Random Forest-based prediction model, can Juba effectively navigate the complexities of its rapid growth trajectory. This study will contribute to creating resilient, sustainable, and economically viable urban environments, ensuring that Juba can emerge as a

model for future cities in the region.

1.3 Objectives of the Study

1.3.1 Primary objective

In this study, we seek to develop a comprehensive understanding of both historical and future urban growth patterns in Juba, South Sudan, by analyzing past trends from 1994 to 2023 and predicting future expansion from 2024 up to the year 2033. This includes utilizing advanced predictive models, specifically Machine Learning's Random Forest and advanced geospatial pre-processing techniques such as cloud masking and ground control points (GCPs) to collect, extract, and manipulate data to inform urban planning and development strategies.

1.3.2 Specific objectives

1. To collect historical time series data from Landsat 5 and 7 spanning 1994 to 2023.
2. To develop a Random Forest machine learning model that identifies historical urban growth trends in Juba from 1994 to 2023, focusing on changes in land use and land cover.
3. To train the model to generate visible historical urban growth patterns merged for the years 1994-2003, 2004-2013 and 2014-2023.
4. To evaluate the performance of the trained model (Random Forest) using a confusion matrix as a metric.
5. To predict future urban growth patterns in Juba for the period 2024-2033 using the trained model.

1.4 Significance of the study

Urbanization in Juba, South Sudan, is rapidly transforming the city and the landscape of its neighborhood, presenting both opportunities for economic development and challenges related to infrastructure, environmental sustainability, and resource management. Understanding past trends and forecasting future urban expansion will be vital for informed decision-making and sustainable urban planning.

This study is significant as it leverages Machine Learning's Random Forest and advanced open-source geospatial pre-processing techniques using both Google Earth Engine (GEE) and Quantum Geographic Information System (GIS), including cloud masking and ground control points (GCPs), to analyze urban growth patterns. By utilizing Landsat 5 and 7 data from 1994 to 2023, this research provides an Informatics-informed data-driven foundation for predicting Juba's urban expansion from 2024 to 2033. The findings will enable urban planners, policymakers, and researchers to:

1. **Enhance Urban Planning:** By identifying historical land-use trends and forecasting future urban growth, national, state, and city planners and policy makers can implement data-driven policies that promote sustainable development and efficient land management.
2. **Support Infrastructure Development:** Accurate predictions of Juba's urban expansion can inform the strategic placement of roads, utilities, housing, disaster preparedness, and public services, ensuring that infrastructure development aligns with population growth.
3. **Mitigate Environmental and Social Risks:** Understanding patterns of urbanization allows stakeholders to anticipate and address potential challenges, such as deforestation, loss of green spaces, flood planning, and informal settlements, which can impact climate resilience and socio-economic stability.
4. **Advance Geospatial and Machine Learning Applications:** This study contributes to the growing field of geospatial machine learning by demonstrating the effectiveness of Random Forest in modeling and predicting urban growth, setting a benchmark for future studies in the region.
5. **Provide a Decision-Support Tool:** The predictive model developed in this study can serve as a decision-support system, enabling government agencies, urban developers, and policymakers to make proactive, evidence-based choices for Juba's future growth.

By combining historical analysis with predictive modeling, this research will not only deepen the understanding of Juba's urban transformation but also provide practical insights for managing its expansion in a sustainable and resilient manner.

1.5 Research Questions

To achieve the research objectives, the study shall address two (2) key questions. These are:

1. What are the historical trends and patterns of urban growth in Juba from 1994 to the

present (2023) based on Landsat 5 and 7 data?

2. How well does the trained Random Forest model perform in detecting historical urban growth trends in Juba from 1994-2003?
3. What are the anticipated (predicted) future patterns of urban growth in Juba from 2024-2033?
4. How can predictive urban growth modeling support decision-making for sustainable land management and infrastructure planning in Juba?

Chapter 2: Literature review.

2.1 Introduction

Urban growth prediction is an important component of urban planning and sustainable development, particularly in rapidly urbanizing regions such as Juba, the capital of Central Equatoria State that doubles as the Capital city of South Sudan. Juba has experienced significant changes in land use and land cover (LULC) over the past few decades as noted by ([Gudo, Deng and Qureshi, 2022a](#); [UN-HABITAT et al., 2023](#)), driven by factors such as population growth, civil unrest, and economic development. This literature review explores the methodologies and findings of recent studies on urban growth prediction in general and Juba in particular.

2.4 Methodologies for Urban Growth Prediction

Urban growth prediction methodologies employ a variety of models and technologies to simulate future urban expansion patterns. These approaches aim to generate accurate forecasts that support urban planning, infrastructure development, and policy-making, ensuring sustainable urban growth. By analyzing historical trends and incorporating spatial and statistical data, these methodologies provide insights into the drivers of urbanization and help stakeholders make informed decisions.

Machine Learning Models (Random Forest)

Machine learning models have become indispensable tools in urban growth prediction, offering the capability to analyze complex patterns in large spatial datasets. Among these, Random Forest (RF) has gained significant traction for land use classification due to its ability to leverage historical remote sensing data and spatial attributes to identify urban expansion trends ([Balzter et al., 2015](#); [Ouma et al., 2024](#); [Sulistiyono et al., 2024](#)). Similarly, Support Vector Machines (SVM) have proven effective in distinguishing land cover types and detecting urbanization dynamics by identifying decision boundaries in high-dimensional feature spaces ([Bouaziz, Eisold and Guermazi,](#)

[2017; Trabelsi et al., 2023](#)). Meanwhile, Artificial Neural Networks (ANNs), with their capacity to model nonlinear relationships, have been increasingly employed to enhance predictive accuracy in urban growth forecasting ([Feng et al., 2011](#)).

Ensemble learning techniques, particularly Random Forest, have been instrumental in improving predictive performance and robustness in urban expansion modeling. Introduced by Breiman (2001), Random Forest is a powerful ensemble method that constructs multiple decision trees during training, each trained on a random subset of the dataset ([Breiman, 2001](#)). This bagging (bootstrap aggregating) approach reduces model variance and enhances generalizability, making RF especially effective for noisy or high-dimensional datasets. Unlike traditional decision trees, which are prone to overfitting, RF leverages randomness at multiple stages—both in bootstrap sampling and feature selection—to generate de-correlated trees. At each node, only a subset of features is considered for splitting, allowing the model to uncover complex interactions between spatial and socio-economic factors influencing urbanization ([Balzter et al., 2015; Sulistiyono et al., 2024](#)).

A key advantage of RF is its built-in mechanism for estimating model accuracy through Out-of-Bag (OOB) error. In each iteration, approximately one-third of the data is left out of the bootstrap sample, allowing for internal cross-validation without the need for a separate test set ([Sulistiyono et al., 2024](#)). This feature makes RF a robust and efficient choice for LULCC modeling, particularly in scenarios where data availability is limited ([Breiman, 2001](#)). Given its ability to capture spatial heterogeneity and account for uncertainties, Random Forest remains a preferred machine learning model for urban growth prediction, aiding policymakers and urban planners in making data-driven decisions for sustainable development.

Cellular Automata (CA)

Cellular Automata (CA) models are widely used in Land Use and Land Cover Change (LULCC) studies due to their ability to simulate spatial dynamics through local interactions. CA is a mathematical modeling approach that represents landscapes as grids of interacting cells, where each cell's state evolves based on predefined transition rules and the states of its neighboring cells. This method effectively captures the self-organizing behavior of urban systems, producing self-similar patterns and highlighting both spatial and quantitative changes over time ([Khawaldah, Farhan and Alzboun, 2020; Das and Angadi, 2022](#)).

In the context of urban expansion, CA models simulate land use changes by integrating various influencing factors such as population density, infrastructure development, road networks, and exclusion zones. These models are particularly effective in capturing the nonlinear nature of urban growth, where small-scale local interactions contribute to large-scale spatial transformations ([Khawaldah, Farhan and Alzboun, 2020; Das and Angadi, 2022](#)).

Variants of CA, such as Logistic Regression–Cellular Automata (LR-CA) and Artificial Neural Network–Cellular Automata (ANN-CA), enhance prediction accuracy by incorporating statistical and deep learning techniques ([Feng et al., 2011](#)) (Feng et al., 2011; Justin and De Vries, 2019). By leveraging historical land cover changes and spatial constraints, these hybrid models refine urban growth simulations and provide more accurate projections for future development.

CA-based models have been successfully applied to predict urban growth patterns in various cities, demonstrating their ability to replicate real-world spatial dynamics. Their adaptability and capacity to integrate diverse data sources make them valuable tools in urban planning and LULCC studies.

Markov Chain Models

Markov Chain models are widely utilized for predicting land-use and land-cover change (LULCC) by estimating the probability of transition from one land-use type to another over a given time period. These models analyze historical land-use data to determine the likelihood of changes occurring in the future, making them particularly effective for assessing long-term urban growth patterns ([Al-Hameedi et al., 2021](#)). By computing transition matrices, Markov Chain models can identify areas with a high probability of urban expansion and forecast potential land-use distributions.

One of the key strengths of Markov Chain models is their ability to provide quantitative insights into urbanization trends, allowing planners to anticipate spatial transformations and make informed policy decisions ([Das and Angadi, 2022](#)). However, because Markov models rely solely on historical transition probabilities, they do not inherently account for spatial constraints, environmental factors, or socioeconomic influences. To address this limitation, they are often integrated with Cellular Automata (CA) models, which introduce spatial dynamics by simulating interactions between neighboring land parcels ([Somvanshi et al., 2020; Al-Hameedi et al., 2021; Das and Angadi, 2022](#)). This hybrid approach enhances the predictive accuracy of urban growth models by incorporating both probabilistic and spatial elements.

Markov Chain models have been successfully applied in various urban settings to forecast future expansion scenarios and assess land-use sustainability. Studies have demonstrated their effectiveness in identifying high-risk zones for urban sprawl, enabling proactive planning interventions to mitigate adverse environmental impacts ([Somvanshi et al., 2020](#); [Al-Hameedi et al., 2021](#); [Das and Angadi, 2022](#)). By simulating different urban growth scenarios, these models contribute to strategic land-use planning, infrastructure development, and environmental conservation efforts, ensuring a more sustainable and resilient approach to urban expansion.

2.4.5 Artificial Neural Networks (ANN)

Artificial Neural Networks (ANNs) are advanced computational models inspired by the structure and function of the human brain ([Somvanshi et al., 2020](#); [Al-Hameedi et al., 2021](#)). They are designed to recognize patterns, process complex data, and make predictions based on input variables. In the context of urban growth prediction, ANNs are used to model intricate relationships between various factors, such as socio-economic conditions, land-use policies, environmental factors, and infrastructure development. Their ability to process large datasets and capture non-linear interactions makes them particularly effective in understanding urban dynamics ([Somvanshi et al., 2020](#); [Al-Hameedi et al., 2021](#)).

Studies employing ANNs have shown remarkable success in accurately forecasting urban growth patterns. By leveraging Geographic Information System (GIS) data, ANNs can incorporate spatial and temporal elements into their predictions, enhancing the model's reliability. These models are particularly useful in identifying areas that are likely to experience urban expansion, as well as understanding the key drivers of this growth, such as population density, economic development, and policy decisions ([Somvanshi et al., 2020](#); [Al-Hameedi et al., 2021](#)).

The application of ANNs in urban growth modeling has provided valuable insights into future land-use trends, helping urban planners and policymakers make informed decisions regarding infrastructure development and land-use management. ANNs have proven instrumental in identifying potential urban sprawl areas, understanding the complex factors influencing growth, and assessing the sustainability of urban expansion ([Somvanshi et al., 2020](#); [Al-Hameedi et al.,](#)

[2021](#)). Through their predictive capabilities, ANNs enable the formulation of more effective strategies for managing urbanization, ensuring that cities can grow in a way that is both sustainable and responsive to the needs of their populations.

Google Earth Engine and other Models

Geospatial and remote sensing techniques play a crucial role in urban growth prediction by providing accurate land cover data and spatial analysis tools. Satellite imagery from sources such as Landsat and Sentinel enables the detection of land use changes over time through spectral analysis. Google Earth Engine (GEE) facilitates large-scale spatial analysis and geospatial data processing, while Geographic Information Systems (GIS) provide visualization and spatial integration capabilities for urban development assessments ([Bruno Soares, Daly and Dessai, 2018](#); [Nyasulu et al., 2022](#)).

Hybrid and ensemble models combine different methodologies to improve predictive accuracy. By integrating machine learning classification techniques with CA-based spatial simulation, these models refine urban growth predictions ([Cifuentes et al., 2020](#); [Nyasulu et al., 2022](#)). Techniques such as stacking and weighted averaging utilize multiple predictive models, enhancing overall forecasting reliability and robustness.

Statistical and econometric models also contribute to urban growth prediction. Markov Chain Analysis is frequently used to predict land use transitions by analyzing historical changes and estimating probabilistic shifts in land cover. Geographically Weighted Regression (GWR) accounts for spatial variations in urban growth determinants, providing more localized and detailed predictions ([Zhao et al., 2009](#); [Micheal, 2025](#)).

Scenario-based approaches offer valuable insights into potential future urban expansion under different policy frameworks. Urban growth scenarios such as business-as-usual, compact city development, and urban sprawl simulations help planners evaluate the impacts of different urbanization strategies. Additionally, agent-based models (ABM) simulate individual or group decision-making processes, illustrating how various factors influence urban expansion patterns ([Dai et al., 2020](#); [Gobin and Van Herzele, 2023](#)).

Land Use and Land Cover (LULC) Dynamics in Juba

Several studies have examined land use and land cover (LULC) changes in Juba, consistently highlighting the rapid expansion of built-up areas at the expense of agricultural land and natural ecosystems. One study by Gudo et al. focused on the spatiotemporal dynamics of LULC changes in Jubek State, which encompasses Juba. Using Landsat satellite imagery from 2000 to 2017, the study analyzed transformations across six major land cover categories: built-up areas, waterbodies, agricultural land, forest land, barren land, and grassland ([Gudo, Deng and Qureshi, 2022b](#)).

The findings revealed a substantial increase in built-up areas, particularly between 2009 and 2017, driven by factors such as population growth, rural-to-urban migration, and economic expansion. Urbanization in Juba followed a linear trajectory alongside population growth, with patterns influenced by security conditions, the availability of natural resources, and broader socio-economic and political factors. The prolonged civil unrest in South Sudan further exacerbated these trends, as large numbers of people migrated to Juba in search of safety and economic opportunities. This influx heightened demand for housing, infrastructure, and public services, accelerating the city's spatial expansion.

In addition to documenting urban growth, the study also identified a critical challenge in predicting future expansion: the lack of consistent and reliable data. Political instability and conflict in the region have disrupted systematic data collection efforts, limiting the availability of high-quality geospatial information ([Gudo, Deng and Qureshi, 2022b](#)).

These constraints have made it difficult to develop accurate predictive models for urban growth, underscoring the need for improved data acquisition strategies and methodological approaches to enhance urban planning and decision-making in Juba.

2.3 Land use and Land cover change dynamics in Africa

A study conducted by Faisal Koko, Auwalu et al., investigated the patterns of land use and land cover (LULC) changes in Lagos, Nigeria, employing maximum likelihood classification and post-classification change detection techniques ([Faisal Koko et al., 2021](#)). These methods were used to

analyze the city's urban expansion and the dynamics of land cover transformation over several decades.

The study identified three distinct phases of urban growth in Lagos and revealed a significant increase in the city's built-up area. In 1990, the built-up area covered approximately 496 km², which expanded to around 860 km² by the year 2000. The urban footprint continued to grow, reaching 1,113 km² in 2010 and surpassing 1,256 km² by 2021. This rapid expansion was accompanied by a marked decline in other land cover types, with vegetation cover decreasing by approximately 398 km², water bodies shrinking by around 246 km², and bare soil reducing by 115 km² between 1990 and 2020 ([Faisal Koko et al., 2021](#)).

The study highlighted the environmental consequences of these shifts, particularly the heightened risk of urban flooding. The loss of vegetation and water bodies, combined with the increase in impervious surfaces, has disrupted natural drainage systems, exacerbating both natural and human-induced flooding events. Given these findings, the researchers emphasized the need for sustainable urban planning and proactive flood mitigation strategies to address the growing challenges posed by Lagos' rapid urbanization.

Another study by Ouma et al., explored the application of a Random Forest Regression (RFR) model for predicting land use and land cover change (LULCC) in the Gaborone Dam catchment, Botswana, for the period 2019–2030 ([Ouma et al., 2024](#)). The study compared the performance of RFR with two other modeling approaches: logistic regression–cellular automata (LR-CA) and artificial neural network–cellular automata (ANN-CA). The researchers proposed RFR as the preferred model due to its ability to capture complex interactions between land use intensity and the nonlinear relationships between various change-driving factors.

To forecast LULCC, the study incorporated a range of influencing variables categorized into physiographic factors, such as elevation, slope, and aspect, and proximity-neighborhood factors, which included distances to water bodies, roads, and urban areas. The historical LULC patterns from 1986 to 2019 were simulated at five-year intervals to assess the performance of each model. The results demonstrated that RFR significantly outperformed ANN-CA and LR-CA, achieving an accuracy of 84.9%, compared to 62.1% and 60.7%, respectively ([Ouma et al., 2024](#)).

Using the RFR model, the study projected notable changes in land cover between 2019 and 2030. The analysis indicated a decline in vegetation cover by 8.9%, while bare soil increased by the same margin. Built-up areas expanded by 2.49%, whereas cropland shrank by 2.8%. Water bodies exhibited minimal change; however, the study found a strong correlation between urban expansion and diminishing dam water capacity. This suggests that increasing population density and urbanization may exert further pressure on water resources.

The study's methodology underscores the potential of machine learning models like RFR in understanding the land–water nexus within catchment areas. These insights can be instrumental in formulating effective policies for sustainable catchment management, urban planning, and water resource conservation in regions experiencing rapid environmental and demographic changes.

Novelty of the Study

This study introduces a novel approach to predicting urban growth in Juba, South Sudan, by integrating advanced machine learning techniques with geospatial technologies. It aims to address the unique challenges of rapid urbanization in a developing, post-conflict urban center. While urban growth prediction models such as Random Forest (RF), Cellular Automata (CA), and Markov Chains have been widely used globally, their application in Juba—a city undergoing rapid, often unplanned expansion due to population growth, civil unrest, and economic pressures—offers a distinct contribution to the field.

The novelty of this research lies in its context-specific application in Juba, a city where consistent geospatial data is scarce due to prolonged political instability and conflict. By leveraging historical Landsat 5 and 7 data from 1994 to 2023, this study pioneers a data-driven approach in a region where urban growth analyses have been limited. This work fills a critical gap in urban planning knowledge for South Sudan, offering a unique perspective on urban growth in a resource-constrained setting.

This research also integrates Random Forest with advanced geospatial pre-processing techniques, such as cloud masking and ground control points (GCPs), using Google Earth Engine (GEE) and Quantum GIS. These methods ensure robust data quality despite challenges such as cloud cover and limited ground-truthing in Juba, setting a new standard for predictive modeling in data-scarce environments. Additionally, the study uses a 29-year historical dataset (1994–2023) to train the

Random Forest model, segmented into decadal patterns, and provides a decade-long future prediction (2024–2033), offering urban planners a detailed temporal framework to anticipate and manage growth.

Another key aspect of this study is its focus on sustainability and resilience. Beyond prediction, the study aims to inform integrated urban planning strategies by forecasting land-use changes and urban sprawl with high accuracy. This directly addresses Juba's dual challenges of economic opportunity and sustainability issues, such as infrastructure strain, environmental degradation, and displacement. The study provides actionable insights that align with global sustainability goals while being locally relevant.

Finally, the development of a Random Forest-based predictive model as a decision-support system is particularly innovative for Juba. Unlike studies that focus solely on model accuracy, this research evaluates performance using a confusion matrix and translates predictions into practical tools for policymakers. This approach enhances infrastructure development, disaster preparedness, and environmental risk mitigation, offering a proactive approach to urban planning in a rapidly transforming city.

Chapter 3: Materials and Methods

3.1 Overview

This study utilizes a machine learning framework to predict urban growth and land cover dynamics in Juba, South Sudan, for the period 2024 to 2033. By leveraging historical satellite imagery from 1994 to 2023, the study applies advanced geospatial analysis techniques to assess past trends and forecast future urban expansion. The methodology integrates satellite data processing, machine learning classification, and predictive modeling to generate insights into land-use changes.

Using Google Earth Engine (GEE) as the primary processing platform, the study employs the Random Forest (RF) algorithm to classify land cover into four categories: Urban, Bare Land, Water, and Vegetation. Historical classifications are generated for three distinct time periods (1994–2003, 2004–2013, and 2014–2023), providing a basis for long-term trend analysis. These classifications inform predictive modeling, which projects future land cover changes based on observed patterns. The study also incorporates spatial analysis and visualization tools, such as Quantum GIS (QGIS), to interpret and present the results. By employing statistical evaluations and validation techniques, the methodology ensures accuracy and reliability in predicting urban expansion. The approach not only enhances understanding of Juba's urbanization dynamics but also serves as a decision-support tool for sustainable urban planning in South Sudan.




Fig. 1 A flow chart of the methodology.

Data Acquisition and Preprocessing Overview

This study relies on satellite imagery and ground truth points to develop a robust dataset for urban growth prediction in Juba, South Sudan. For 1994-2003, Landsat 5 Collection 2, Tier 1, Level 2 surface reflectance (SR_B1 to SR_B7) and thermal data (ST_B6) are used, while Landsat 7 data of the same tier and level cover 2004-2013 and 2014-2023. Ground Control Points (GCPs) are provided as feature-collections with a landcover property from 0-3 for each class, which define training signatures. The region of interest (ROI) is specified as a geometry asset. Preprocessing ensures data consistency: optical bands are scaled to reflectance using the equation

$$R_{scaled} = \frac{(DN * Multiplicative Factor) + Additive Offset}{\sin(\theta)}$$

Where:

R_{scaled} is the scaled reflectance value, DN is the raw digital number from the sensor, the *Multiplicative Factor* is 0.0000275 (specific to Landsat Collection 2 Level 2 data), the *Additive Offset* is -0.2, and θ is the solar elevation angle in radians, accounting for solar illumination variations across image acquisition dates.

The Thermal band is converted to kelvin with:

$$T_{kelvin} = (DN * Thermal\ Multiplicative\ Factor) + Thermal\ Additive\ Offset$$

Where:

T_{kelvin} is the temperature in kelvin, the *Thermal Multiplicative Factor* is 0.00341802, and the *Thermal Additive Offset* is 149.0, converting raw thermal data to a physically meaningful scale.

Clouds and shadows are masked using the QA_PIXEL band, a critical component of Landsat Collection 2 Level 2 data. The QA_PIXEL band is a 16-bit unsigned integer layer that encodes quality information for each pixel in a bit-packed format, where each bit or combination of bits flags specific conditions such as clouds, shadows, water, or snow. For Landsat 5 and 7, this band is generated during atmospheric correction and quality assessment processes, providing a standardized way to identify and filter out contaminated pixels. Each bit position corresponds to a specific condition, and its value (0 or 1) indicates whether that condition is absent or present. In this study, the focus is on masking clouds and cloud shadows, which obscure surface features and could skew classification results. The bit assignments for Landsat Collection 2 Level 2 data are as follows (relevant bits highlighted):

- Bit 0: Fill (1 if data is invalid, e.g., outside the image extent)
- Bit 1: Dilated Cloud (1 if near a cloud, expanding the cloud boundary)
- Bit 2: Cirrus (1 if cirrus clouds are detected, less relevant for Landsat 5/7)
- Bit 3: Cloud (1 if a cloud is present)
- Bit 4: Cloud Shadow (1 if a cloud shadow is present)
- Bits 5-15: Additional flags (e.g., snow, water, confidence levels)

In the masking process, we are only concerned with Bit 3 and Bit 4, which represents clouds and cloud shadows. The process in Google Earth Engine involves extracting the cloud shadow information from the QA_PIXEL band using bitwise operation. Since the QA_PIXEL band

encodes multiple quality indicators within a 16-bit integer we use bitwise shifting and logical operations to isolate and remove cloud-affected pixels through:

$$Mask = \left(\frac{QA_{PIXEL}}{2^3} \bmod 2 = 0 \right) \wedge \left(\frac{QA_{PIXEL}}{2^4} \bmod 2 = 0 \right)$$

Where:

- QA_{PIXEL} is the quality assurance band for each pixel
- 2^3 represents the bit position for cloud (Bit 3)
- 2^4 represents the bit position for cloud shadow (Bit 4)
- The modulo operation, $\bmod 2$ checks whether the bit is set as (1) or not (0)
- $\frac{QA_{PIXEL}}{2^3} \bmod 2 = 0$ ensures the cloud (bit 3) is not set
- $\frac{QA_{PIXEL}}{2^4} \bmod 2 = 0$ ensures the cloud shadow (Bit 4) is not set
- The pixels where both conditions are true (bit is 0) are retained
- $\wedge$ is the logical AND operator

Alternatively, we represent the above equation as:

$$Mask = ((QA_{PIXEL} \gg 3) \bmod 2 = 0) \wedge ((QA_{PIXEL} \gg 4) \bmod 2 = 0) \wedge$$

Where:

- $\gg$ is the bitwise right shift operator, shifting the relevant bit to the least significant position before applying the modulo operation.

Median composites are then created by computing the median value of unmasked (clear) pixels across the time range for each period, clipped to the ROI to reduce temporal noise. These median composites represent the periods 1994-2003, 2003-2013, and 2014-2023, which are fed for Random Forest training.

Random Forest Classification

Historical classifications utilize a Random Forest (RF) classifier, an ensemble method that enhances accuracy by combining multiple decision trees.

For a dataset with (n) features (spectral bands SR_B1 to SR_B7 and ST_B6, where) and (m) training samples (GCPs), RF constructs (T) trees, each trained on a bootstrapped subset of the data and a random subset of features at each node. The final classification for a pixel with

feature vector $x = [x_1, x_2, \dots, x_n]$ is determined by majority voting. The classification is represented by:

$$h(x) = \text{mode}\{h_1(x), h_2(x), \dots, h_T(x)\}$$

Where:

- $x = [x_1, x_2, \dots, x_n]$ is the feature vector for a pixel, consisting of $n = 8$ spectral bands: SR_B1 to SR_B7 and ST_B6.
- $h_t(x)$ is the class (0 for Urban, 1 for Bare, 2 for water, or 3 for vegetation) predicted by the t -th decision tree in the ensemble.
- $T = 50$ is the number of decision trees used in the Random Forest, chosen to balance computational efficiency and robustness.
- *mode* represents the majority voting function, selecting the most frequently predicted class among all trees.

Historical Land Cover Classification

For 1994-2003, the Landsat 5 median composite is classified using the RF model trained on 2003 GCPs (urban2003, bare2003, water2003, vegetation2003), assigning pixels to Urban (0), Bare (1), Water (2), or Vegetation (3) via the majority vote from 50 trees. Validation employs a confusion matrix, with overall accuracy calculated as:

$$\text{Overall Accuracy} = \frac{\sum_{i=0}^3 \text{True Positives}_i}{\text{Total Number of Pixels in Validation Set}}$$

Where:

- True Positives_i is the number of pixels correctly classified as class (i) in a withheld validation subset (for instance 20% of GCPs),
- The denominator, *Total Number of Pixels in Validation Set* is the total number of validation pixels, providing a measure of classification reliability.

For 2004-2013 and 2014-2023, the Landsat 7 composite is classified using urban2013 and 2003 GCPs for non-urban classes, adapting to recent changes while retraining the RF model to reflect urban expansion.

Area Computation

Post-classification, areas are quantified in square kilometers. For each period, the area A_c of class (c) is calculated as:

$$A_c = Pixel\ Count_c * (Pixel\ Resolution^2 / 1,000,000)$$

Where:

- $Pixel\ Count_c$ is the number of pixels classified as class (c),
- $Pixel\ Resolution = 30m$ (Landsat's spatial resolution),
- $Pixel\ Resolution^2 = 30*30=900m^2$ and divided by 1,000,000 converts square meters to square kilometer, yielding 0.0009 km² per pixel.

Thus, in order to ensure consistency with the ROI's total area while establishing a baseline for trend analysis, the equation simplifies to:

$$A_c = Pixel\ Count_c * 0.0009km^2$$

Prediction of Land Cover (2024-2033)

Future land cover for 2024-2033 is predicted by extrapolating spectral and area trends based on the full historical dataset from 1994 to 2023, incorporating data points from 2003 (1994-2003 composite), 2013 (2004-2013 composite), and 2023 (2014-2023 composite). To simplify, the annual spectral change rate per band is computed across the 20-year span from 2003 to 2023:

$$\Delta S_{band} = \frac{S_{2023,band} - S_{2003,band}}{20}$$

Where:

- $S_{year,band}$ is the median value of the specified band (e.g., SR_B1) in the composite for the respective period, and
- 20 is the number of years from 2003 to 2023, providing a straightforward annual change rate.

A simulated 2033 composite is generated by applying this rate forward from 2023:

$$S_{2033} = S_{2023,band} + (\Delta S_{band} * 10)$$

Where:

- $\Delta S_{band} * 10$ is the total change over the 10 years from 2023 to 2033, assuming a linear trend.

This composite is classified using the RF model trained on 2014-2023 data, reflecting the most recent spectral characteristics and training points.

For area prediction, a simplified linear growth rate is calculated based on the total change from 2003 to 2033 as:

$$r_c = \frac{A_{c,2023} - A_{c,2003}}{20}$$

Where:

- $A_{c,2003}$ and $A_{c,2023}$ are areas of class (c) from the 1994-2003 and 2014-2023 classifications, and
- 20 is the number of years, giving the annual area change in km².

The predicted area for class (c) in year (t) (2024-2033) is then computed as:

$$A_{c,t} = A_{c,2023} + (r_c * (t - 2023))$$

Where:

- $A_{c,2023}$ is the area in 2023,
- r_c is the annual area change, and
- $t - 2023$ is the number of years from 2023 to (t) (for instance 1 for 2024, 10 for 2033), providing a linear year-by-year projection to 2033.

Evaluation Metric

The performance of the Random Forest classifier is evaluated using a confusion matrix, a widely used metric in classification tasks to assess accuracy and error rates. In this project, the confusion matrix is generated within GEE to validate the classifications for each period (1994-2003, 2004-2013, 2014-2023) against ground truth data from GCPs. It provides a tabular summary of predicted versus actual class labels, enabling the calculation of overall accuracy and per-class performance.

Definition: For the four land cover classes (Urban [0], Bare [1], Water [2], Vegetation [3]), the confusion matrix is a 4x4 table where rows represent actual classes (from validation GCPs) and columns represent predicted classes (from the RF classifier). Each cell C_{ij} contains the number of pixels where the actual class is (i) and the predicted class is (j). Diagonal elements (C_{ii}) are correct predictions (true positives), while the off-diagonal elements are misclassifications. This is represented by:

$$Overall Accuracy = \frac{\sum_i^3 C_{ii}}{\sum_{i=0}^3 \sum_{j=0}^3 C_{ij}}$$

Where:

- C_{ii} is the number of true positives for class (i) – diagonal elements, and
- $\sum_{i=0}^3 \sum_{j=0}^3 C_{ij}$ is the total number of pixels in the validation set (sum of all matrix elements).

Producer's Accuracy (PA)

PA measures the proportion of actual pixels in a class correctly classified, reflecting the classifier's ability to identify all instances of that class (omission error inverse).

$$PA_i = \frac{C_{ii}}{\sum_{j=0}^3 C_{ij}}$$

Where:

- C_{ii} is the number of true positives for class (i), and
- $\sum_{j=0}^3 C_{ij}$ is the total number of pixels predicted as class (i) (row total)

User's Accuracy (UA)

UA measures the proportion of predicted pixels in a class that are correct, reflecting the reliability of the classifier's predictions for that class (commission error inverse), represented as:

$$UA_i = \frac{C_{ii}}{\sum_{j=0}^3 C_{ij}}$$

Where:

- C_{ii} is the number of true positives for class (i), and
- $\sum_{j=0}^3 C_{ij}$ is the total number of pixels predicted as class (i) (column total)

Visualizations and Further Processing

Initial visualizations such as classified maps for 1994-2003, 2004-2013, 2014-2023, and 2024-2033, plus time series charts (2003-2033), bar and pie charts, and confusion matrices are generated in GEE. For further processing and enhanced mapping, data is exported from GEE to Google Drive. Classified images are saved as GeoTIFFs (e.g., Collection_1994_2003.tif), preserving the 30 m resolution and class values (0-3). GCPs are exported as Shapefiles (e.g., GCPs_2003.shp) for

reference, and area statistics and chart data are saved as CSVs. Further processing to generate visualization map files are done using QGIS.

Software and Applications

This study employs a combination of software tools and programming languages to process satellite imagery, classify land cover, predict urban growth, and visualize results for Juba, South Sudan. The primary platforms used are Google Earth Engine (GEE) and QGIS, with JavaScript serving as the programming language within GEE. These tools facilitate data acquisition, preprocessing, classification, and visualization, ensuring a comprehensive workflow for urban growth prediction.

Google Earth Engine (GEE)

GEE, as a cloud-based geospatial processing platform, is designed for large-scale environmental data analysis. It provides access to extensive satellite imagery archives, including Landsat 5 and Landsat 7 data, used in this study. The platform supports data acquisition, preprocessing, classification using the Random Forest algorithm (selected for this study), and visualization. It also offers preprocessing capabilities such as cloud and shadow masking, band scaling, and composite creation. The built-in Random Forest classifier is used for land cover classification and prediction, and results are visualized through maps, time series charts, and accuracy assessments. The GEE Code Editor, accessible via <https://code.earthengine.google.com>, serves as the development environment where the project script was written and managed, with integration to Google Drive for exporting processed outputs.

JavaScript in GEE

JavaScript is the primary scripting language for implementing the methodology in GEE. It enables the definition of functions, data handling, and algorithm implementation. Custom functions are written for image preprocessing, cloud masking, classification, and area computation. The language facilitates the loading of Landsat imagery, filtering by date and region of interest, and managing training data as feature collections. JavaScript commands allow seamless export of classified images and tabular data to Google Drive, while the GEE Earth Engine API supports efficient geospatial processing.

Random Forest in GEE


The Random Forest (RF) classifier is integrated into GEE as part of its supervised classification tools under the `ee.Classifier` namespace. It is implemented using the SMILE (Statistical Machine Intelligence and Learning Engine) library and optimized for large-scale geospatial data analysis. The RF classifier is instantiated using JavaScript (`ee.Classifier.smileRandomForest(50)`), where 50 decision trees balance accuracy and efficiency. Training data, consisting of labeled ground control points (GCPs), is used to train the classifier, which is then applied to median composites for historical periods and the predicted 2033 dataset. Validation is performed using confusion matrices to assess classification accuracy, and results are exported for further analysis.

QGIS

QGIS is an open-source Geographic Information System (GIS) software used for advanced visualization and spatial analysis of the study's outputs. It enables the import of GeoTIFFs and Shapefiles while maintaining spatial resolution and metadata. The platform provides tools for symbology customization, comparative time series analysis, and cartographic enhancements such as legends and scale bars. QGIS also supports supplementary processing, including CSV integration for area statistics and spatial data manipulation. Its robust visualization capabilities enhance interpretability, making it a valuable complement to GEE.


Integration and Workflow

The integration of GEE and QGIS ensures an efficient workflow for urban growth prediction. GEE handles cloud-based data processing, classification, and initial visualization, while QGIS refines and presents the results offline. JavaScript scripts orchestrate the RF classification process in GEE, exporting outputs to Google Drive for import into QGIS. This workflow leverages GEE's computational scalability and QGIS's visualization capabilities, providing a robust pipeline from raw satellite data to actionable urban planning insights.



Elevation

Height (m)



834

443

Contour Lines

0 3.5 7 10.5 14 km



QGIS ©Map Created By: Nyombe Benson | Supervised By: Dr. Manzu Gerald

REFERENCES

1. Al-Hameedi, W.M.M. et al. (2021) 'Remote Sensing-Based Urban Sprawl Modeling Using Multilayer Perceptron Neural Network Markov Chain in Baghdad, Iraq', *Remote Sensing*, 13(20), p. 4034. Available at: <https://doi.org/10.3390/rs13204034>.
2. Balzter, H. et al. (2015) 'Mapping CORINE Land Cover from Sentinel-1A SAR and SRTM Digital Elevation Model Data using Random Forests', *Remote Sensing*, 7(11), pp. 14876–14898. Available at: <https://doi.org/10.3390/rs71114876>.
3. Bouaziz, M., Eisold, S. and Guermazi, E. (2017) 'Semiautomatic approach for land cover classification: a remote sensing study for arid climate in southeastern Tunisia', *Euro-Mediterranean Journal for Environmental Integration*, 2(1), p. 24. Available at: <https://doi.org/10.1007/s41207-017-0036-7>.
4. Breiman, L. (2001) 'Random Forests', *Machine Learning*, 45(1), pp. 5–32. Available at: <https://doi.org/10.1023/A:1010933404324>.
5. Bruno Soares, M., Daly, M. and Dessai, S. (2018) 'Assessing the value of seasonal climate forecasts for decision-making', *WIREs Climate Change*, 9(4), p. e523. Available at: <https://doi.org/10.1002/wcc.523>.
6. Cifuentes, J. et al. (2020) 'Air Temperature Forecasting Using Machine Learning Techniques: A Review', *Energies*, 13(16), p. 4215. Available at: <https://doi.org/10.3390/en13164215>.
7. Dai, E. et al. (2020) 'Agent-based model of land system: Theory, application and modelling framework', *Journal of Geographical Sciences*, 30(10), pp. 1555–1570. Available at: <https://doi.org/10.1007/s11442-020-1799-3>.
8. Das, S. and Angadi, D.P. (2022) 'Land use land cover change detection and monitoring of urban growth using remote sensing and GIS techniques: a micro-level study', *GeoJournal*, 87(3), pp. 2101–2123. Available at: <https://doi.org/10.1007/s10708-020-10359-1>.
9. Faisal Koko, A. et al. (2021) 'Analyzing urban growth and land cover change scenario in Lagos, Nigeria using multi-temporal remote sensing data and GIS to mitigate flooding', *Geomatics, Natural Hazards and Risk*, 12(1), pp. 631–652. Available at: <https://doi.org/10.1080/19475705.2021.1887940>.
10. Feng, Y. et al. (2011) 'Modeling dynamic urban growth using cellular automata and particle swarm optimization rules', *Landscape and Urban Planning*, 102(3), pp. 188–196. Available at: <https://doi.org/10.1016/j.landurbplan.2011.04.004>.
11. Gobin, A. and Van Herzele, A. (2023) 'A Data-Driven Farm Typology as a Basis for Agricultural Land Use Decisions', *Land*, 12(11), p. 2032. Available at: <https://doi.org/10.3390/land12112032>.
12. Gudo, A.J.A., Deng, J. and Qureshi, A.S. (2022a) 'Analysis of Spatiotemporal Dynamics of Land Use/Cover Changes in Jubei State, South Sudan', *Sustainability*, 14(17), p. 10753. Available at: <https://doi.org/10.3390/su141710753>.
13. Gudo, A.J.A., Deng, J. and Qureshi, A.S. (2022b) 'Analysis of Spatiotemporal Dynamics of Land Use/Cover Changes in Jubei State, South Sudan', *Sustainability*, 14(17), p. 10753. Available at: <https://doi.org/10.3390/su141710753>.
14. Ishtiaque, A., Shrestha, M. and Chhetri, N. (2017) 'Rapid Urban Growth in the Kathmandu Valley, Nepal: Monitoring Land Use Land Cover Dynamics of a Himalayan City with Landsat Imageries', *Environments*, 4(4), p. 72. Available at: <https://doi.org/10.3390/environments4040072>.
15. Jiang, F. and Lv, Y. (2022) 'Changes in the spatial distribution of population in urban agglomerations: A spatial-temporal perspective'. In Review. Available at: <https://doi.org/10.21203/rs.3.rs-1244609/v1>.
16. Justin, L. (2017) South Sudan Governance: A Call for Federalism to Address : Marginalization and Prevent Armed Conflict Recurrence - 21_Justin Laku.pdf. Available at: https://sonar.ch/documents/324250/preview/21_Justin%20Laku.pdf (Accessed: 30 March 2025).
17. Justin, P.H. and De Vries, L. (2019) 'Governing Unclear Lines: Local Boundaries as a (Re)source of Conflict in South Sudan', *Journal of Borderlands Studies*, 34(1), pp. 31–46. Available at: <https://doi.org/10.1080/08865655.2017.1294497>.
18. Khawaldah, H., Farhan, I. and Alzboun, N. (2020) 'Simulation and prediction of land use and land cover change using GIS, remote sensing and CA-Markov model', *Global Journal of Environmental Science and Management*, 6, pp. 215–232. Available at: <https://doi.org/10.22034/gjesm.2020.02.07>.
19. Maqbool, A. et al. (2020) 'Disaster Mitigation in Urban Pakistan Using Agent Based Modeling with GIS', *ISPRS International*

Journal of Geo-Information, 9(4), p. 203. Available at: <https://doi.org/10.3390/ijgi9040203>.

20. Micheal, D. (2025) Schools closed after 12 students fainted daily over heat exhaustion: Official - Eye Radio, Eye Radio. Available at: <https://www.eyeradio.org/schools-closed-after-12-students-fainted-daily-over-heat-exhaustion-official/> (Accessed: 29 March 2025).

21. Munthali, K.G. and Murayama, Y. (2011) 'Land use/cover change detection and analysis for Dzalanyama forest reserve, Lilongwe, Malawi', *Procedia - Social and Behavioral Sciences*, 21, pp. 203–211. Available at: <https://doi.org/10.1016/j.sbspro.2011.07.035>.

22. Munthali, M.G. et al. (2019) 'Local Perception of Drivers of Land-Use and Land-Cover Change Dynamics across Dedza District, Central Malawi Region', *Sustainability*, 11(3), p. 832. Available at: <https://doi.org/10.3390/su11030832>.

23. Nyasulu, C. et al. (2022) 'Towards Resilient Agriculture to Hostile Climate Change in the Sahel Region: A Case Study of Machine Learning-Based Weather Prediction in Senegal', *Agriculture*, 12(9), p. 1473. Available at: <https://doi.org/10.3390/agriculture12091473>.

24. Ouma, Y.O. et al. (2024) 'Land-Use Change Prediction in Dam Catchment Using Logistic Regression-CA, ANN-CA and Random Forest Regression and Implications for Sustainable Land–Water Nexus', *Sustainability*, 16(4), p. 1699. Available at: <https://doi.org/10.3390/su16041699>.

25. Rodríguez-Cuenca, B., Alonso, M.C. and Tamés-Noriega, A. (2015) 'LULC database updating from VHR images and LIDAR data using evidence theory', in: 2015 Conference of the International Fuzzy Systems Association and the European Society for Fuzzy Logic and Technology (IFSA-EUSFLAT-15), Atlantis Press, pp. 987–993. Available at: <https://doi.org/10.2991/ifsaeusflat-15.2015.139>.

26. Sexton, J.O. et al. (2013) 'Urban growth of the Washington, D.C.–Baltimore, MD metropolitan region from 1984 to 2010 by annual, Landsat-based estimates of impervious cover', *Remote Sensing of Environment*, 129, pp. 42–53. Available at: <https://doi.org/10.1016/j.rse.2012.10.025>.

27. Somvanshi, S.S. et al. (2020) 'Monitoring spatial LULC changes and its growth prediction based on statistical models and earth observation datasets of Gautam Budh Nagar, Uttar Pradesh, India', *Environment, Development and Sustainability*, 22(2), pp. 1073–1091. Available at: <https://doi.org/10.1007/s10668-018-0234-8>.

28. Sulistiyono, N. et al. (2024) 'Identification of mangrove forests using Sentinel-1 Synthetic Aperture Radar (SAR) satellite imagery in Medan Belawan District, Medan City', *IOP Conference Series: Earth and Environmental Science*, 1352(1), p. 012051. Available at: <https://doi.org/10.1088/1755-1315/1352/1/012051>.

29. Trabelsi, N. et al. (2023) 'Classification and Spatio-Temporal Change Detection of Land Use/Land Cover Using Remote Sensing and Geographic Information System in the Manouba Region, NE Tunisia', *Journal of Geographic Information System*, 15(06), pp. 652–668. Available at: <https://doi.org/10.4236/jgis.2023.156033>.

30. Turner, M.G. and Ruscher, C.L. (1988) 'Changes in landscape patterns in Georgia, USA', *Landscape Ecology*, 1(4), pp. 241–251. Available at: <https://doi.org/10.1007/BF00157696>.

31. UN-HABITAT (2006) Meeting Development Goals in Small Urban Centres - Water and Sanitation in the World's Cities 2006 | UN-Habitat. Available at: <https://unhabitat.org/meeting-development-goals-in-small-urban-centres-water-and-sanitation-in-the-worlds-cities-2006> (Accessed: 30 March 2025).

32. UN-HABITAT et al. (2023) Juba Strategic Plan | UN-Habitat. Available at: <https://unhabitat.org/juba-strategic-plan> (Accessed: 30 March 2025).

33. Wild, H., Jok, J.M. and Patel, R. (2018) 'The Militarization of Cattle Raiding in South Sudan: How a Traditional Practice Became a Tool for Political Violence', *Journal of International Humanitarian Action*, 3(1). Available at: <https://doi.org/10.1186/s41018-018-0030-y>.

34. Zeng, Y. et al. (2016) 'MULTI-AGENT BASED SIMULATION OF OPTIMAL URBAN LAND USE ALLOCATION IN THE MIDDLE REACHES OF THE YANGTZE RIVER, CHINA', *The International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, XLI-B8, pp. 1089–1092. Available at: <https://doi.org/10.5194/isprs-archives-XLI-B8-1089-2016>.

35. Zhao, S. et al. (2009) 'A spatial resolution threshold of land cover in estimating regional terrestrial carbon sequestration'. *Biogeochemistry: Modelling, Terrestrial*. Available at: <https://doi.org/10.5194/bgd-6-7983-2009>.

