

Infomap Bioregions >AMPHIBIANS About





#### Rioregion 1

42161 records of 719 unique species in 167 cells

| Most common species      |       | Most indicative species  |       |
|--------------------------|-------|--------------------------|-------|
| Name                     | Count | Name                     | Score |
| Trachycephalus venulosus | 600   | Lithobates palmipes      | 3.29  |
| Hypsiboas punctatus      | 598   | Scinax ruber             | 2.92  |
| Rhinella marina          | 591   | Rhinella margaritifera   | 2.63  |
| Scinax ruber             | 581   | Adenomera hylaedactyla   | 2.45  |
| Hypsiboas geographicus   | 577   | Hypsiboas geographicus   | 2.28  |
| Rhinella margaritifera   | 572   | Hypsiboas punctatus      | 2.18  |
| Adenomera hylaedactyla   | 562   | Elachistocleis ovalis    | 2.05  |
| Lithobates palmipes      | 559   | Dendropsophus minutus    | 2.01  |
| Dendropsophus minutus    | 552   | Trachycephalus venulosus | 2.01  |
| Elachistocleis ovalis    | 547   | Rhinella marina          | 1.94  |

| Bioregion 2 |
|-------------|
|-------------|

27267 records of 553 unique species in 333 cells

| Most common species        |       | Most indicative species    |       |
|----------------------------|-------|----------------------------|-------|
| Name                       | Count | Name                       | Score |
| Kassina senegalensis       | 970   | Hildebrandtia ornata       | 2.08  |
| Phrynobatrachus natalensis | 824   | Hemisus marmoratus         | 2.02  |
| Tomopterna cryptotis       | 820   | Ptychadena oxyrhynchus     | 2.02  |
| Hemisus marmoratus         | 793   | Kassina senegalensis       | 1.99  |
| Ptychadena oxyrhynchus     | 650   | Phrynobatrachus natalensis | 1.97  |

http://localhost:3003/ Page 1 of 29

| Ptychadena mascareniensis  | 625 | Amietophrynus maculatus    | 1.91 |
|----------------------------|-----|----------------------------|------|
| Amietophrynus maculatus    | 608 | Tomopterna cryptotis       | 1.88 |
| Hoplobatrachus occipitalis | 605 | Hoplobatrachus occipitalis | 1.82 |
| Hildebrandtia ornata       | 562 | Amietophrynus regularis    | 1.77 |
| Amietophrynus regularis    | 522 | Ptychadena mascareniensis  | 1.76 |

### Bioregion 3

13083 records of 103 unique species in 313 cells

| Most common species   |       | Most indicative species |       |
|-----------------------|-------|-------------------------|-------|
| Name                  | Count | Name                    | Score |
| Rana arvalis          | 1547  | Triturus cristatus      | 1.32  |
| Rana temporaria       | 1383  | Rana temporaria         | 1.31  |
| Bufo bufo             | 1362  | Pelobates fuscus        | 1.31  |
| Lissotriton vulgaris  | 1015  | Pelophylax lessonae     | 1.30  |
| Pelophylax ridibundus | 906   | Pelophylax ridibundus   | 1.29  |
| Triturus cristatus    | 633   | Hyla arborea            | 1.29  |
| Pelobates fuscus      | 605   | Bufotes viridis         | 1.28  |
| Hyla arborea          | 493   | Lissotriton vulgaris    | 1.28  |
| Bufotes viridis       | 452   | Bufo bufo               | 1.16  |
| Pelophylax lessonae   | 423   | Rana arvalis            | 1.14  |

### Bioregion 4

10452 records of 178 unique species in 160 cells

| Most common species       |       | Most indicative species   |       |
|---------------------------|-------|---------------------------|-------|
| Name                      | Count | Name                      | Score |
| Notophthalmus viridescens | 742   | Notophthalmus viridescens | 2.74  |
| Anaxyrus americanus       | 521   | Ambystoma maculatum       | 2.73  |
| Pseudacris crucifer       | 450   | Lithobates palustris      | 2.72  |
| Lithobates clamitans      | 417   | Pseudacris crucifer       | 2.68  |
| Lithobates sylvaticus     | 393   | Lithobates clamitans      | 2.63  |
| Lithobates catesbeianus   | 372   | Hyla chrysoscelis         | 2.41  |
| Hyla chrysoscelis         | 346   | Anaxyrus americanus       | 2.34  |
| Ambystoma maculatum       | 334   | Lithobates pipiens        | 1.23  |
| Lithobates pipiens        | 323   | Lithobates catesbeianus   | 1.19  |
| Lithobates palustris      | 271   | Lithobates sylvaticus     | 0.827 |

### Bioregion 5

9224 records of 468 unique species in 120 cells

| Most common species |       | Most indicative species |       |  |
|---------------------|-------|-------------------------|-------|--|
| Name                | Count | Name                    | Score |  |

http://localhost:3003/ Page 2 of 29

| Fejervarya limnocharis     | 574 | Microhyla pulchra          | 3.51 |
|----------------------------|-----|----------------------------|------|
| Microhyla fissipes         | 491 | Microhyla butleri          | 3.27 |
| Duttaphrynus melanostictus | 331 | Microhyla fissipes         | 3.27 |
| Microhyla heymonsi         | 261 | Hoplobatrachus rugulosus   | 3.24 |
| Hoplobatrachus rugulosus   | 252 | Microhyla heymonsi         | 2.95 |
| Microhyla butleri          | 214 | Kaloula pulchra            | 2.83 |
| Kaloula pulchra            | 187 | Occidozyga lima            | 2.74 |
| Occidozyga lima            | 183 | Fejervarya limnocharis     | 1.65 |
| Microhyla pulchra          | 176 | Duttaphrynus melanostictus | 1.62 |
| Polypedates leucomystax    | 176 | Polypedates leucomystax    | 1.37 |

| ior |  |  |  |
|-----|--|--|--|
|     |  |  |  |
|     |  |  |  |

11131 records of 282 unique species in 69 cells

| Most common species    |       | Most indicative species |       |
|------------------------|-------|-------------------------|-------|
| Name                   | Count | Name                    | Score |
| Leptodactylus fuscus   | 271   | Scinax fuscomarginatus  | 4.61  |
| Physalaemus cuvieri    | 271   | Rhinella schneideri     | 3.63  |
| Rhinella schneideri    | 271   | Physalaemus cuvieri     | 3.31  |
| Leptodactylus latrans  | 270   | Dendropsophus nanus     | 2.91  |
| Rhinella granulosa     | 269   | Scinax x-signatus       | 2.28  |
| Dendropsophus nanus    | 268   | Rhinella granulosa      | 2.19  |
| Scinax fuscomarginatus | 266   | Elachistocleis ovalis   | 2.18  |
| Dendropsophus minutus  | 266   | Leptodactylus latrans   | 2.17  |
| Scinax x-signatus      | 264   | Dendropsophus minutus   | 2.15  |
| Elachistocleis ovalis  | 263   | Leptodactylus fuscus    | 2.07  |

### Bioregion 7

7375 records of 259 unique species in 56 cells

| Most common species          |       | Most indicative species      |       |
|------------------------------|-------|------------------------------|-------|
| Name                         | Count | Name                         | Score |
| Afrixalus quadrivittatus     | 208   | Hymenochirus boettgeri       | 9.43  |
| Hylarana albolabris          | 199   | Hyperolius ocellatus         | 9.40  |
| Hyperolius cinnamomeoventris | 190   | Hyperolius tuberculatus      | 9.36  |
| Xenopus epitropicalis        | 178   | Xenopus epitropicalis        | 9.31  |
| Hyperolius ocellatus         | 172   | Xenopus fraseri              | 8.56  |
| Leptopelis notatus           | 162   | Leptopelis notatus           | 7.73  |
| Hymenochirus boettgeri       | 159   | Chiromantis rufescens        | 7.68  |
| Chiromantis rufescens        | 157   | Hylarana albolabris          | 6.49  |
| Hyperolius tuberculatus      | 155   | Hyperolius cinnamomeoventris | 5.63  |
| Xenopus fraseri              | 154   | Afrixalus quadrivittatus     | 4.82  |

http://localhost:3003/ Page 3 of 29

## Bioregion 8

5562 records of 185 unique species in 113 cells

| Most common species        |       | Most indicative species    |       |
|----------------------------|-------|----------------------------|-------|
| Name                       | Count | Name                       | Score |
| Litoria caerulea           | 343   | Litoria novaehollandiae    | 5.93  |
| Litoria rubella            | 312   | Litoria latopalmata        | 5.93  |
| Limnodynastes tasmaniensis | 232   | Limnodynastes fletcheri    | 5.93  |
| Platyplectrum ornatum      | 200   | Litoria inermis            | 5.93  |
| Litoria inermis            | 158   | Platyplectrum ornatum      | 5.87  |
| Litoria novaehollandiae    | 149   | Limnodynastes tasmaniensis | 5.86  |
| Litoria latopalmata        | 148   | Litoria longipes           | 5.83  |
| Limnodynastes fletcheri    | 127   | Litoria rothii             | 5.79  |
| Litoria rothii             | 122   | Litoria caerulea           | 5.43  |
| Litoria longipes           | 116   | Litoria rubella            | 4.39  |

#### Bioregion 9

5152 records of 190 unique species in 61 cells

| Most common species       |     | Most indicative species   |       |
|---------------------------|-----|---------------------------|-------|
| Name Count                |     | Name                      | Score |
| Leptodactylus mystacinus  | 205 | Hypsiboas pulchellus      | 9.68  |
| Leptodactylus latrans     | 194 | Leptodactylus latinasus   | 9.14  |
| Physalaemus biligonigerus | 165 | Elachistocleis bicolor    | 8.82  |
| Odontophrynus americanus  | 161 | Leptodactylus gracilis    | 8.56  |
| Leptodactylus gracilis    | 156 | Physalaemus biligonigerus | 8.03  |
| Leptodactylus latinasus   | 150 | Rhinella arenarum         | 7.74  |
| Rhinella schneideri       | 129 | Odontophrynus americanus  | 5.35  |
| Elachistocleis bicolor    | 127 | Leptodactylus mystacinus  | 4.47  |
| Rhinella arenarum         | 124 | Rhinella schneideri       | 2.28  |
| Hypsiboas pulchellus      | 119 | Leptodactylus latrans     | 2.06  |

### Bioregion 10

4899 records of 282 unique species in 94 cells

| Most common species        |     | Most indicative species  |       |  |
|----------------------------|-----|--------------------------|-------|--|
| Name Count                 |     | Name                     | Score |  |
| Euphlyctis cyanophlyctis   | 398 | Zakerana syhadrensis     | 5.06  |  |
| Duttaphrynus stomaticus    | 397 | Duttaphrynus stomaticus  | 5.06  |  |
| Fejervarya limnocharis     | 354 | Hoplobatrachus tigerinus | 4.94  |  |
| Microhyla ornata           | 327 | Euphlyctis cyanophlyctis | 4.94  |  |
| Hoplobatrachus tigerinus   | 317 | Polypedates maculatus    | 4.92  |  |
| Duttaphrynus melanostictus | 272 | Hoplobatrachus crassus   | 4.80  |  |

http://localhost:3003/ Page 4 of 29

| Sphaerotheca breviceps | 270 | Sphaerotheca breviceps     | 4.79 |
|------------------------|-----|----------------------------|------|
| Polypedates maculatus  | 227 | Microhyla ornata           | 4.62 |
| Zakerana syhadrensis   | 201 | Duttaphrynus melanostictus | 1.92 |
| Hoplobatrachus crassus | 185 | Fejervarya limnocharis     | 1.47 |

#### **Bioregion 11**

3508 records of 299 unique species in 61 cells

| Most common species        |     | Most indicative species    |       |  |
|----------------------------|-----|----------------------------|-------|--|
| Name Count                 |     | Name                       | Score |  |
| Polypedates leucomystax    | 182 | Polypedates macrotis       | 10.7  |  |
| Fejervarya limnocharis     | 165 | Odorrana hosii             | 10.7  |  |
| Fejervarya cancrivora      | 134 | Hylarana nicobariensis     | 10.1  |  |
| Duttaphrynus melanostictus | 100 | Kaloula baleata            | 9.66  |  |
| Limnonectes kuhlii         | 96  | Phrynoidis asper           | 9.51  |  |
| Polypedates macrotis       | 95  | Fejervarya cancrivora      | 6.40  |  |
| Odorrana hosii             | 90  | Limnonectes kuhlii         | 5.68  |  |
| Kaloula baleata            | 89  | Polypedates leucomystax    | 4.46  |  |
| Hylarana nicobariensis     | 78  | Duttaphrynus melanostictus | 1.54  |  |
| Phrynoidis asper           | 74  | Fejervarya limnocharis     | 1.50  |  |

#### Bioregion 12

3349 records of 442 unique species in 21 cells

| Most common species      |    | Most indicative species  |       |
|--------------------------|----|--------------------------|-------|
| Name Count               |    | Name                     | Score |
| Physalaemus cuvieri      | 63 | Hypsiboas semilineatus   | 28.5  |
| Leptodactylus latrans    | 53 | Hypsiboas crepitans      | 4.97  |
| Scinax x-signatus        | 52 | Physalaemus cuvieri      | 3.31  |
| Siphonops annulatus      | 50 | Leptodactylus mystacinus | 3.27  |
| Dendropsophus minutus    | 47 | Rhinella schneideri      | 2.59  |
| Hypsiboas crepitans      | 46 | Siphonops annulatus      | 1.96  |
| Leptodactylus mystacinus | 46 | Scinax x-signatus        | 1.93  |
| Elachistocleis ovalis    | 46 | Leptodactylus latrans    | 1.83  |
| Rhinella schneideri      | 45 | Elachistocleis ovalis    | 1.64  |
| Hypsiboas semilineatus   | 45 | Dendropsophus minutus    | 1.63  |

# Bioregion 13

3470 records of 102 unique species in 83 cells

| Most common species |       | Most indicative species |       |  |
|---------------------|-------|-------------------------|-------|--|
| Name                | Count | Name                    | Score |  |
| Ambystoma tigrinum  | 305   | Anaxyrus punctatus      | 6.11  |  |

http://localhost:3003/ Page 5 of 29

| Anaxyrus woodhousii     | 256 | Spea multiplicata       | 5.81 |
|-------------------------|-----|-------------------------|------|
| Anaxyrus cognatus       | 254 | Scaphiopus couchii      | 5.77 |
| Spea bombifrons         | 215 | Anaxyrus cognatus       | 5.67 |
| Anaxyrus punctatus      | 208 | Spea bombifrons         | 5.29 |
| Lithobates catesbeianus | 183 | Anaxyrus woodhousii     | 4.91 |
| Scaphiopus couchii      | 167 | Ambystoma tigrinum      | 3.79 |
| Spea multiplicata       | 162 | Lithobates pipiens      | 1.48 |
| Lithobates pipiens      | 160 | Pseudacris maculata     | 1.48 |
| Pseudacris maculata     | 151 | Lithobates catesbeianus | 1.42 |

#### Bioregion 14

3237 records of 18 unique species in 114 cells

| Most common species        |       | Most indicative species    |        |
|----------------------------|-------|----------------------------|--------|
| Name                       | Count | Name                       | Score  |
| Salamandrella keyserlingii | 1505  | Rana dybowskii             | 1.10   |
| Rana amurensis             | 918   | Rana amurensis             | 1.08   |
| Rana arvalis               | 230   | Salamandrella keyserlingii | 1.00   |
| Rana dybowskii             | 176   | Rana pirica                | 0.901  |
| Bufo bufo                  | 124   | Strauchbufo raddei         | 0.264  |
| Strauchbufo raddei         | 108   | Hyla japonica              | 0.201  |
| Hyla japonica              | 60    | Rana arvalis               | 0.175  |
| Bufo gargarizans           | 52    | Bufo gargarizans           | 0.161  |
| Rana pirica                | 18    | Bufo bufo                  | 0.109  |
| Lissotriton vulgaris       | 10    | Lissotriton vulgaris       | 0.0129 |

### Bioregion 15

2453 records of 52 unique species in 70 cells

| Most common                | species | Most indicative s          | Most indicative species |  |
|----------------------------|---------|----------------------------|-------------------------|--|
| Name                       | Count   | Name                       | Score                   |  |
| Strauchbufo raddei         | 439     | Bombina orientalis         | 4.27                    |  |
| Pelophylax nigromaculatus  | 277     | Rana chensinensis          | 4.08                    |  |
| Rana chensinensis          | 271     | Kaloula borealis           | 3.71                    |  |
| Bufo gargarizans           | 266     | Strauchbufo raddei         | 3.67                    |  |
| Hyla japonica              | 222     | Hyla immaculata            | 2.93                    |  |
| Salamandrella keyserlingii | 121     | Bufo gargarizans           | 2.82                    |  |
| Rana amurensis             | 118     | Pelophylax nigromaculatus  | 2.80                    |  |
| Kaloula borealis           | 112     | Hyla japonica              | 2.55                    |  |
| Bombina orientalis         | 93      | Rana amurensis             | 0.476                   |  |
| Hyla immaculata            | 55      | Salamandrella keyserlingii | 0.276                   |  |

### Bioregion 16

http://localhost:3003/ Page 6 of 29

| 1 | E71 | rocord | o of | 122 | uniaua | snecies | in | 20 colle | _ |
|---|-----|--------|------|-----|--------|---------|----|----------|---|
| ı | 5/4 | record | s ot | 432 | unique | species | ın | 20 Cella | S |

| Most common species       |       | Most indicative species   |       |
|---------------------------|-------|---------------------------|-------|
| Name                      | Count | Name                      | Score |
| Rhinella marina           | 30    | Craugastor megacephalus   | 67.8  |
| Craugastor fitzingeri     | 30    | Pristimantis ridens       | 67.8  |
| Leptodactylus savagei     | 25    | Craugastor fitzingeri     | 67.8  |
| Rhaebo haematiticus       | 25    | Leptodactylus savagei     | 62.8  |
| Agalychnis callidryas     | 22    | Rhaebo haematiticus       | 58.5  |
| Craugastor megacephalus   | 22    | Espadarana prosoblepon    | 54.8  |
| Espadarana prosoblepon    | 21    | Smilisca phaeota          | 48.5  |
| Smilisca phaeota          | 20    | Agalychnis callidryas     | 32.4  |
| Pristimantis ridens       | 20    | Leptodactylus melanonotus | 17.6  |
| Leptodactylus melanonotus | 20    | Rhinella marina           | 1.97  |

#### Bioregion 17

1607 records of 306 unique species in 32 cells

| Most common species   |       | Most indicative species |       |
|-----------------------|-------|-------------------------|-------|
| Name                  | Count | Name                    | Score |
| Hylophorbus rufescens | 112   | Hylarana garritor       | 18.2  |
| Litoria infrafrenata  | 104   | Sphenophryne cornuta    | 18.2  |
| Hylarana papua        | 96    | Hylarana papua          | 17.8  |
| Hylarana garritor     | 81    | Callulops robustus      | 17.8  |
| Callulops robustus    | 44    | Hylophorbus rufescens   | 17.7  |
| Platymantis papuensis | 44    | Litoria pygmaea         | 17.7  |
| Litoria thesaurensis  | 41    | Platymantis papuensis   | 14.0  |
| Sphenophryne cornuta  | 39    | Litoria infrafrenata    | 13.4  |
| Hylarana daemeli      | 39    | Hylarana daemeli        | 12.4  |
| Litoria pygmaea       | 37    | Litoria thesaurensis    | 8.66  |

### Bioregion 18

1714 records of 20 unique species in 64 cells

| Most common species   |       | Most indicative species |       |
|-----------------------|-------|-------------------------|-------|
| Name                  | Count | Name                    | Score |
| Lithobates sylvaticus | 766   | Anaxyrus hemiophrys     | 2.20  |
| Pseudacris maculata   | 342   | Lithobates sylvaticus   | 1.56  |
| Anaxyrus hemiophrys   | 160   | Pseudacris maculata     | 1.33  |
| Lithobates pipiens    | 140   | Lithobates pipiens      | 0.517 |
| Anaxyrus americanus   | 71    | Spea bombifrons         | 0.343 |
| Ambystoma tigrinum    | 66    | Ambystoma tigrinum      | 0.327 |
| Spea bombifrons       | 35    | Anaxyrus americanus     | 0.309 |

http://localhost:3003/ Page 7 of 29

| Anaxyrus boreas   | 30 | Anaxyrus boreas   | 0.234 |
|-------------------|----|-------------------|-------|
| Anaxyrus cognatus | 24 | Anaxyrus cognatus | 0.213 |
| Hyla chrysoscelis | 22 | Hyla chrysoscelis | 0.149 |

## Bioregion 19

2111 records of 82 unique species in 51 cells

| Most common species     |       | Most indicative species |       |
|-------------------------|-------|-------------------------|-------|
| Name                    | Count | Name                    | Score |
| Ambystoma macrodactylum | 312   | Aneides vagrans         | 6.52  |
| Anaxyrus boreas         | 262   | Ambystoma macrodactylum | 6.42  |
| Pseudacris regilla      | 176   | Pseudacris regilla      | 6.24  |
| Rana luteiventris       | 172   | Rana luteiventris       | 6.06  |
| Lithobates catesbeianus | 124   | Spea intermontana       | 5.16  |
| Lithobates sylvaticus   | 110   | Anaxyrus boreas         | 5.01  |
| Lithobates pipiens      | 92    | Lithobates catesbeianus | 0.943 |
| Spea intermontana       | 87    | Lithobates pipiens      | 0.835 |
| Aneides vagrans         | 82    | Ambystoma tigrinum      | 0.656 |
| Ambystoma tigrinum      | 54    | Lithobates sylvaticus   | 0.550 |

### Bioregion 20

1058 records of 33 unique species in 17 cells

| Most common species     |       | Most indicative species |       |
|-------------------------|-------|-------------------------|-------|
| Name                    | Count | Name                    | Score |
| Nannophryne variegata   | 660   | Eupsophus calcaratus    | 3.08  |
| Eupsophus calcaratus    | 96    | Eupsophus emiliopugini  | 3.08  |
| Batrachyla leptopus     | 66    | Batrachyla antartandica | 3.08  |
| Batrachyla antartandica | 57    | Hylorina sylvatica      | 3.08  |
| Rhinoderma darwinii     | 45    | Rhinoderma darwinii     | 3.08  |
| Batrachyla taeniata     | 25    | Nannophryne variegata   | 3.07  |
| Eupsophus emiliopugini  | 24    | Batrachyla leptopus     | 3.04  |
| Pleurodema thaul        | 17    | Batrachyla taeniata     | 2.66  |
| Hylorina sylvatica      | 15    | Calyptocephalella gayi  | 2.06  |
| Calyptocephalella gayi  | 10    | Pleurodema thaul        | 1.69  |

### Bioregion 21

1022 records of 238 unique species in 20 cells

| Most common species       |       | Most indicative species |       |
|---------------------------|-------|-------------------------|-------|
| Name                      | Count | Name                    | Score |
| Ptychadena mascareniensis | 51    | Mantidactylus femoralis | 39.9  |
| Laliostoma labrosum       | 26    | Mantidactylus ulcerosus | 39.9  |
|                           |       |                         |       |

http://localhost:3003/ Page 8 of 29

| Mantidactylus remoralis          | 20 | Agiyptodactyius madagascariensis | 39.9 |
|----------------------------------|----|----------------------------------|------|
| Boophis doulioti                 | 20 | Boophis doulioti                 | 39.9 |
| Mantidactylus curtus             | 19 | Boophis brachychir               | 39.9 |
| Mantidactylus ulcerosus          | 17 | Scaphiophryne calcarata          | 39.9 |
| Aglyptodactylus madagascariensis | 16 | Laliostoma labrosum              | 39.9 |
| Guibemantis liber                | 15 | Guibemantis liber                | 39.9 |
| Boophis brachychir               | 14 | Mantidactylus curtus             | 39.9 |
| Scaphiophryne calcarata          | 14 | Ptychadena mascareniensis        | 2.74 |

#### **Bioregion 22**

1027 records of 101 unique species in 12 cells

| Most common species        |       | Most indicative species    |       |
|----------------------------|-------|----------------------------|-------|
| Name                       | Count | Name                       | Score |
| Hyperolius concolor        | 30    | Leptopelis spiritusnoctis  | 45.8  |
| Amietophrynus regularis    | 30    | Afrixalus dorsalis         | 36.7  |
| Ptychadena bibroni         | 29    | Hyperolius concolor        | 22.4  |
| Hoplobatrachus occipitalis | 29    | Phrynobatrachus calcaratus | 13.3  |
| Phrynobatrachus latifrons  | 28    | Xenopus tropicalis         | 13.0  |
| Xenopus tropicalis         | 27    | Phrynobatrachus latifrons  | 8.83  |
| Leptopelis spiritusnoctis  | 27    | Chiromantis rufescens      | 8.48  |
| Afrixalus dorsalis         | 26    | Ptychadena bibroni         | 6.58  |
| Phrynobatrachus calcaratus | 26    | Amietophrynus regularis    | 3.29  |
| Chiromantis rufescens      | 25    | Hoplobatrachus occipitalis | 2.82  |

### **Bioregion 23**

626 records of 27 unique species in 11 cells

| Most common species       |       | Most indicative species   |       |
|---------------------------|-------|---------------------------|-------|
| Name                      | Count | Name                      | Score |
| Platymantis weberi        | 215   | Ceratobatrachus guentheri | 9.47  |
| Platymantis solomonis     | 144   | Batrachylodes trossulus   | 9.47  |
| Ceratobatrachus guentheri | 43    | Batrachylodes wolfi       | 9.47  |
| Litoria thesaurensis      | 38    | Platymantis guppyi        | 9.47  |
| Batrachylodes trossulus   | 27    | Platymantis weberi        | 9.42  |
| Discodeles guppyi         | 22    | Platymantis solomonis     | 9.34  |
| Discodeles bufoniformis   | 21    | Discodeles bufoniformis   | 9.03  |
| Batrachylodes wolfi       | 13    | Batrachylodes vertebralis | 8.79  |
| Platymantis guppyi        | 13    | Discodeles guppyi         | 8.01  |
| Batrachylodes vertebralis | 13    | Litoria thesaurensis      | 4.18  |

## Bioregion 24

1196 records of 232 unique species in 19 cells

http://localhost:3003/ Page 9 of 29

| Most common species         |       | Most indicative species     |       |
|-----------------------------|-------|-----------------------------|-------|
| Name                        | Count | Name                        | Score |
| Rhinella marina             | 47    | Incilius valliceps          | 31.9  |
| Leptodactylus fragilis      | 45    | Scinax staufferi            | 28.4  |
| Smilisca baudinii           | 44    | Leptodactylus melanonotus   | 23.6  |
| Leptodactylus melanonotus   | 42    | Hypopachus variolosus       | 21.9  |
| Incilius valliceps          | 42    | Smilisca baudinii           | 19.6  |
| Scinax staufferi            | 40    | Leptodactylus fragilis      | 16.5  |
| Trachycephalus venulosus    | 40    | Lithobates berlandieri      | 12.2  |
| Lithobates berlandieri      | 37    | Dendropsophus microcephalus | 2.95  |
| Hypopachus variolosus       | 37    | Rhinella marina             | 1.97  |
| Dendropsophus microcephalus | 31    | Trachycephalus venulosus    | 1.71  |

### Bioregion 25

850 records of 35 unique species in 17 cells

704 records of 83 unique species in 15 cells

Limnonectes leytensis

Platymantis corrugatus

Philautus surdus

| Most common species     |       | Most indicative species |       |
|-------------------------|-------|-------------------------|-------|
| Name                    | Count | Name                    | Score |
| Hyla japonica           | 64    | Hynobius naevius        | 31.8  |
| Hynobius naevius        | 60    | Rhacophorus schlegelii  | 31.2  |
| Glandirana rugosa       | 60    | Rana ornativentris      | 31.1  |
| Lithobates catesbeianus | 57    | Hynobius yatsui         | 31.1  |
| Rhacophorus schlegelii  | 57    | Cynops pyrrhogaster     | 30.6  |
| Cynops pyrrhogaster     | 51    | Rana tagoi              | 30.0  |
| Rana tagoi              | 51    | Rana japonica           | 29.4  |
| Rana japonica           | 48    | Glandirana rugosa       | 28.1  |
| Rana ornativentris      | 47    | Hyla japonica           | 5.04  |
| Hynobius yatsui         | 45    | Lithobates catesbeianus | 2.11  |

| Most common species     |       | Most indicative species |       |
|-------------------------|-------|-------------------------|-------|
| Name                    | Count | Name                    | Score |
| Fejervarya cancrivora   | 51    | Limnonectes leytensis   | 39.9  |
| Rhinella marina         | 47    | Philautus surdus        | 39.9  |
| Polypedates leucomystax | 46    | Platymantis corrugatus  | 39.9  |
| Fejervarya vittigera    | 39    | Kaloula conjuncta       | 37.7  |
| Kaloula conjuncta       | 34    | Fejervarya vittigera    | 33.1  |
| Kaloula picta           | 29    | Kaloula picta           | 33.1  |

http://localhost:3003/ Page 10 of 29

Occidozyga laevis

Fejervarya cancrivora

Polypedates leucomystax

18.1

8.70

4.03

25

22

20

| Occidozyga laevis | 15 | Rhinella marina | 1.82 |
|-------------------|----|-----------------|------|
| , 6               |    |                 |      |

| Bioreg                                       | ion 27                  |  |
|--|-------------------------|--|
| 808 records of 25 unique species in 18 cells |                         |  |
| Most common species                          | Most indicative species |  |
|  |                         |  |

| Most common species       |       | Most indicative species   |       |
|---------------------------|-------|---------------------------|-------|
| Name                      | Count | Name                      | Score |
| Notaden nichollsi         | 129   | Uperoleia russelli        | 15.8  |
| Litoria maini             | 126   | Platyplectrum spenceri    | 14.2  |
| Platyplectrum spenceri    | 124   | Litoria maini             | 14.1  |
| Neobatrachus sutor        | 117   | Neobatrachus sutor        | 14.1  |
| Litoria rubella           | 93    | Notaden nichollsi         | 12.6  |
| Litoria platycephala      | 51    | Neobatrachus wilsmorei    | 12.4  |
| Neobatrachus wilsmorei    | 37    | Pseudophryne occidentalis | 8.28  |
| Pseudophryne occidentalis | 32    | Litoria platycephala      | 5.33  |
| Uperoleia russelli        | 23    | Litoria rubella           | 3.48  |
| Litoria caerulea          | 22    | Litoria caerulea          | 0.925 |

### Bioregion 28

538 records of 127 unique species in 13 cells

| Most common species       |       | Most indicative species   |       |
|---------------------------|-------|---------------------------|-------|
| Name                      | Count | Name                      | Score |
| Hyla arenicolor           | 27    | Eleutherodactylus nitidus | 67.8  |
| Craugastor augusti        | 26    | Plectrohyla bistincta     | 53.4  |
| Hyla eximia               | 21    | Hyla eximia               | 52.8  |
| Spea multiplicata         | 19    | Tlalocohyla smithii       | 52.4  |
| Ambystoma velasci         | 18    | Incilius occidentalis     | 39.1  |
| Eleutherodactylus nitidus | 18    | Craugastor augusti        | 22.5  |
| Plectrohyla bistincta     | 17    | Ambystoma velasci         | 21.9  |
| Tlalocohyla smithii       | 16    | Hyla arenicolor           | 18.7  |
| Hypopachus variolosus     | 16    | Hypopachus variolosus     | 16.5  |
| Incilius occidentalis     | 14    | Spea multiplicata         | 7.70  |

|  | Bioregion 29 |
|--|--------------|
|--|--------------|

552 records of 31 unique species in 23 cells

| Most common species        |       | Most indicative species    |       |
|----------------------------|-------|----------------------------|-------|
| Name                       | Count | Name                       | Score |
| Bufotes boulengeri         | 149   | Discoglossus scovazzi      | 13.7  |
| Pelophylax saharicus       | 128   | Salamandra algira          | 13.7  |
| Amietophrynus mauritanicus | 74    | Amietophrynus mauritanicus | 13.5  |
| I halo montalionalio       | 40    | Dalambulay ashariaya       | 107   |

http://localhost:3003/ Page 11 of 29

| пута тнегтитопаль     | 40 | геторпутах запансиз | 14.7  |
|-----------------------|----|---------------------|-------|
| Amietophrynus xeros   | 27 | Bufotes boulengeri  | 11.8  |
| Bufo bufo             | 24 | Discoglossus pictus | 10.2  |
| Discoglossus scovazzi | 21 | Hyla meridionalis   | 7.16  |
| Discoglossus pictus   | 18 | Pleurodeles waltl   | 1.95  |
| Salamandra algira     | 8  | Amietophrynus xeros | 0.691 |
| Pleurodeles waltl     | 6  | Bufo bufo           | 0.213 |

#### Bioregion 30

584 records of 38 unique species in 14 cells

| Most common species         |       | Most indicative species     |       |
|-----------------------------|-------|-----------------------------|-------|
| Name                        | Count | Name                        | Score |
| Tomopterna cryptotis        | 90    | Kassina maculifer           | 21.0  |
| Chiromantis kelleri         | 71    | Hildebrandtia macrotympanum | 20.4  |
| Poyntonophrynus lughensis   | 59    | Amietophrynus blanfordii    | 18.7  |
| Hildebrandtia macrotympanum | 56    | Chiromantis kelleri         | 18.2  |
| Amietophrynus xeros         | 54    | Duttaphrynus dodsoni        | 17.7  |
| Duttaphrynus dodsoni        | 54    | Poyntonophrynus lughensis   | 17.3  |
| Ptychadena anchietae        | 41    | Kassina somalica            | 13.5  |
| Kassina somalica            | 31    | Amietophrynus xeros         | 2.29  |
| Amietophrynus blanfordii    | 24    | Tomopterna cryptotis        | 2.22  |
| Kassina maculifer           | 13    | Ptychadena anchietae        | 1.68  |

# Bioregion 31

387 records of 135 unique species in 12 cells

| Most common species    |       | Most indicative species |       |
|------------------------|-------|-------------------------|-------|
| Name                   | Count | Name                    | Score |
| Rhinella spinulosa     | 54    | Gastrotheca marsupiata  | 35.8  |
| Pleurodema marmoratum  | 40    | Telmatobius marmoratus  | 35.2  |
| Pleurodema cinereum    | 22    | Pleurodema cinereum     | 34.5  |
| Gastrotheca marsupiata | 19    | Pleurodema marmoratum   | 34.3  |
| Telmatobius marmoratus | 14    | Nymphargus bejaranoi    | 29.3  |
| Rhinella arenarum      | 11    | Rhinella veraguensis    | 26.9  |
| Rhinella veraguensis   | 10    | Hypsiboas andinus       | 18.8  |
| Hypsiboas andinus      | 8     | Rhinella spinulosa      | 18.7  |
| Nymphargus bejaranoi   | 7     | Rhinella arenarum       | 2.61  |
| Leptodactylus gracilis | 6     | Leptodactylus gracilis  | 1.25  |

## Bioregion 32

184 records of 92 unique species in 2 cells

| Most common species | Most indicative species |
|---------------------|-------------------------|
|---------------------|-------------------------|

http://localhost:3003/ Page 12 of 29

| Name                          | Count | Name                          | Score |
|-------------------------------|-------|-------------------------------|-------|
| Pseudophilautus reticulatus   | 6     | Ramanella obscura             | 339   |
| Pseudophilautus fergusonianus | 6     | Pseudophilautus reticulatus   | 339   |
| Ramanella obscura             | 5     | Pseudophilautus fergusonianus | 339   |
| Pseudophilautus folicola      | 4     | Taruga longinasus             | 339   |
| Euphlyctis hexadactylus       | 4     | Pseudophilautus folicola      | 339   |
| Duttaphrynus melanostictus    | 4     | Pseudophilautus singu         | 339   |
| Taruga longinasus             | 4     | Pseudophilautus femoralis     | 339   |
| Pseudophilautus singu         | 4     | Euphlyctis hexadactylus       | 8.93  |
| Pseudophilautus femoralis     | 4     | Euphlyctis cyanophlyctis      | 3.29  |
| Euphlyctis cyanophlyctis      | 4     | Duttaphrynus melanostictus    | 1.87  |

#### **Bioregion 33**

330 records of 74 unique species in 5 cells

| Most common species      |       | Most indicative species  |       |
|--------------------------|-------|--------------------------|-------|
| Name                     | Count | Name                     | Score |
| Leptodactylus fuscus     | 16    | Relictivomer pearsei     | 84.8  |
| Hypsiboas crepitans      | 15    | Pleurodema brachyops     | 17.5  |
| Leptodactylus bolivianus | 14    | Engystomops pustulosus   | 17.3  |
| Rhinella granulosa       | 14    | Hypsiboas crepitans      | 6.38  |
| Rhinella marina          | 14    | Leptodactylus bolivianus | 3.48  |
| Relictivomer pearsei     | 12    | Leptodactylus fuscus     | 2.07  |
| Trachycephalus venulosus | 12    | Rhinella granulosa       | 1.93  |
| Elachistocleis ovalis    | 11    | Rhinella marina          | 1.72  |
| Pleurodema brachyops     | 11    | Elachistocleis ovalis    | 1.54  |
| Engystomops pustulosus   | 11    | Trachycephalus venulosus | 1.51  |

### Bioregion 34

181 records of 65 unique species in 4 cells

| Most common species           |       | Most indicative species       |       |
|-------------------------------|-------|-------------------------------|-------|
| Name                          | Count | Name                          | Score |
| Hypsiboas heilprini           | 28    | Osteopilus vastus             | 72.7  |
| Osteopilus vastus             | 24    | Eleutherodactylus ruthae      | 72.7  |
| Osteopilus pulchrilineatus    | 22    | Eleutherodactylus abbotti     | 72.7  |
| Eleutherodactylus ruthae      | 7     | Osteopilus dominicensis       | 72.7  |
| Eleutherodactylus abbotti     | 6     | Peltophryne guentheri         | 72.7  |
| Eleutherodactylus inoptatus   | 6     | Eleutherodactylus pictissimus | 72.7  |
| Osteopilus dominicensis       | 6     | Eleutherodactylus audanti     | 72.7  |
| Peltophryne guentheri         | 4     | Osteopilus pulchrilineatus    | 72.7  |
| Eleutherodactylus pictissimus | 4     | Hypsiboas heilprini           | 70.2  |
| Eleutherodactylus audanti     | 3     | Eleutherodactylus inoptatus   | 62.3  |

http://localhost:3003/ Page 13 of 29

#### Bioregion 35

299 records of 32 unique species in 8 cells

| Most common species       |       | Most indicative species   |       |
|---------------------------|-------|---------------------------|-------|
| Name                      | Count | Name                      | Score |
| Limnodynastes dorsalis    | 31    | Litoria adelaidensis      | 65.6  |
| Pseudophryne occidentalis | 29    | Litoria moorei            | 65.6  |
| Neobatrachus pelobatoides | 28    | Neobatrachus albipes      | 65.6  |
| Pseudophryne guentheri    | 24    | Neobatrachus pelobatoides | 63.4  |
| Neobatrachus kunapalari   | 22    | Limnodynastes dorsalis    | 61.7  |
| Heleioporus albopunctatus | 21    | Pseudophryne guentheri    | 60.6  |
| Litoria adelaidensis      | 15    | Heleioporus albopunctatus | 59.9  |
| Neobatrachus sutor        | 14    | Neobatrachus kunapalari   | 57.8  |
| Litoria moorei            | 12    | Pseudophryne occidentalis | 31.2  |
| Neobatrachus albipes      | 11    | Neobatrachus sutor        | 7.02  |

#### Bioregion 36

156 records of 13 unique species in 3 cells

| Most common species   |       | Most indicative species |       |
|-----------------------|-------|-------------------------|-------|
| Name                  | Count | Name                    | Score |
| Hylarana arfaki       | 80    | Litoria congenita       | 22.5  |
| Litoria congenita     | 23    | Litoria aruensis        | 22.3  |
| Litoria bicolor       | 22    | Hylarana arfaki         | 17.9  |
| Litoria infrafrenata  | 8     | Lechriodus melanopyga   | 7.12  |
| Litoria aruensis      | 7     | Litoria bicolor         | 5.38  |
| Lechriodus melanopyga | 7     | Xenorhina oxycephala    | 1.70  |
| Hylophorbus rufescens | 2     | Litoria infrafrenata    | 1.44  |
| Litoria thesaurensis  | 2     | Litoria thesaurensis    | 0.592 |
| Xenorhina oxycephala  | 1     | Callulops robustus      | 0.565 |
| Callulops robustus    | 1     | Hylophorbus rufescens   | 0.442 |

### Bioregion 37

230 records of 32 unique species in 8 cells

| Most common species   |       | Most indicative species |       |
|-----------------------|-------|-------------------------|-------|
| Name                  | Count | Name                    | Score |
| Pelophylax bergeri    | 20    | Pelophylax hispanicus   | 96.4  |
| Bufo bufo             | 19    | Pelophylax bergeri      | 92.5  |
| Pelophylax hispanicus | 18    | Rana italica            | 88.2  |
| Bufotes balearicus    | 17    | Hyla sarda              | 87.2  |
| Unda inkannaadia      | 15    | Discoolsoniasadiis      | 000   |

http://localhost:3003/ Page 14 of 29

30/11/15 21:58 Infomap Bioregions

| пунанценнечна       | 10 | DISCORIOSSOS SALUOS | 03.0 |
|---------------------|----|---------------------|------|
| Rana dalmatina      | 14 | Bufotes balearicus  | 72.1 |
| Triturus carnifex   | 14 | Hyla intermedia     | 66.4 |
| Discoglossus sardus | 14 | Triturus carnifex   | 25.0 |
| Rana italica        | 13 | Rana dalmatina      | 6.25 |
| Hyla sarda          | 12 | Bufo bufo           | 1.26 |

299 records of 166 unique species in 2 cells

| Most common species         |       | Most indicative species     |       |
|-----------------------------|-------|-----------------------------|-------|
| Name                        | Count | Name                        | Score |
| Leptodactylus fuscus        | 8     | Hypodactylus latens         | 254   |
| Pristimantis gaigei         | 5     | Pristimantis gaigei         | 97.8  |
| Pristimantis w-nigrum       | 5     | Craugastor raniformis       | 72.7  |
| Hypsiboas crepitans         | 5     | Leptodactylus colombiensis  | 72.7  |
| Craugastor raniformis       | 4     | Pristimantis w-nigrum       | 66.9  |
| Leptodactylus colombiensis  | 4     | Craugastor longirostris     | 63.6  |
| Dendropsophus microcephalus | 4     | Hypsiboas crepitans         | 4.25  |
| Craugastor longirostris     | 4     | Dendropsophus microcephalus | 2.24  |
| Hypodactylus latens         | 4     | Leptodactylus fuscus        | 2.07  |
| Elachistocleis ovalis       | 4     | Elachistocleis ovalis       | 1.12  |

313 records of 112 unique species in 4 cells

| Most common species               |       | Most indicative species           |       |
|-----------------------------------|-------|-----------------------------------|-------|
| Name                              | Count | Name                              | Score |
| Eleutherodactylus cystignathoides | 15    | Incilius cristatus                | 136   |
| Rhinella marina                   | 11    | Craugastor decoratus              | 136   |
| Ambystoma tigrinum                | 11    | Chiropterotriton chondrostega     | 136   |
| Lithobates berlandieri            | 11    | Eleutherodactylus longipes        | 106   |
| Craugastor augusti                | 9     | Eleutherodactylus cystignathoides | 70.2  |
| Chiropterotriton chondrostega     | 8     | Craugastor augusti                | 14.0  |
| Smilisca baudinii                 | 8     | Lithobates berlandieri            | 11.4  |
| Incilius cristatus                | 7     | Smilisca baudinii                 | 11.2  |
| Eleutherodactylus longipes        | 7     | Ambystoma tigrinum                | 2.78  |
| Craugastor decoratus              | 6     | Rhinella marina                   | 1.44  |

## Bioregion 40

| 244 records of 43 unique species in 8 cells |  |  |  |  |
|---|--|--|--|--|
| Most common species Most indicative species |  |  |  |  |
| Name Count Name Score                       |  |  |  |  |

http://localhost:3003/ Page 15 of 29

| Rana kukunoris          | 48 | Scutiger mammatus       | 42.4  |
|-------------------------|----|-------------------------|-------|
| Scutiger boulengeri     | 43 | Bufo tibetanus          | 38.4  |
| Scutiger mammatus       | 28 | Scutiger glandulatus    | 35.3  |
| Bufo tibetanus          | 19 | Nanorana pleskei        | 33.9  |
| Batrachuperus tibetanus | 16 | Scutiger boulengeri     | 27.6  |
| Nanorana pleskei        | 12 | Batrachuperus tibetanus | 27.1  |
| Scutiger glandulatus    | 10 | Rana kukunoris          | 26.8  |
| Bufo minshanicus        | 7  | Bufo minshanicus        | 15.6  |
| Bufo andrewsi           | 7  | Bufo andrewsi           | 3.41  |
| Strauchbufo raddei      | 6  | Strauchbufo raddei      | 0.459 |

#### Bioregion 41

306 records of 67 unique species in 5 cells

| Most common species        |       | Most indicative species    |       |
|----------------------------|-------|----------------------------|-------|
| Name                       | Count | Name                       | Score |
| Fejervarya limnocharis     | 43    | Buergeria japonica         | 47.3  |
| Buergeria japonica         | 28    | Microhyla okinavensis      | 47.3  |
| Microhyla okinavensis      | 26    | Babina okinavana           | 47.3  |
| Lithobates catesbeianus    | 17    | Rhacophorus aurantiventris | 47.3  |
| Bufo gargarizans           | 9     | Echinotriton andersoni     | 47.3  |
| Babina okinavana           | 9     | Cynops ensicauda           | 47.3  |
| Rhacophorus aurantiventris | 8     | Fejervarya limnocharis     | 1.65  |
| Echinotriton andersoni     | 8     | Bufo gargarizans           | 0.975 |
| Cynops ensicauda           | 8     | Lithobates catesbeianus    | 0.938 |
| Polypedates leucomystax    | 7     | Polypedates leucomystax    | 0.726 |

### Bioregion 42

214 records of 61 unique species in 4 cells

| Most common species            |       | Most indicative species        |       |
|--------------------------------|-------|--------------------------------|-------|
| Name                           | Count | Name                           | Score |
| Osteopilus septentrionalis     | 28    | Eleutherodactylus varleyi      | 72.7  |
| Eleutherodactylus limbatus     | 11    | Eleutherodactylus atkinsi      | 72.7  |
| Peltophryne peltocephala       | 11    | Eleutherodactylus thomasi      | 72.7  |
| Lithobates catesbeianus        | 10    | Peltophryne peltocephala       | 72.7  |
| Eleutherodactylus planirostris | 10    | Eleutherodactylus limbatus     | 72.7  |
| Eleutherodactylus varleyi      | 10    | Peltophryne empusa             | 65.4  |
| Peltophryne empusa             | 9     | Eleutherodactylus pinarensis   | 65.4  |
| Eleutherodactylus pinarensis   | 9     | Osteopilus septentrionalis     | 38.4  |
| Eleutherodactylus atkinsi      | 9     | Eleutherodactylus planirostris | 18.6  |
| Eleutherodactylus thomasi      | 6     | Lithobates catesbeianus        | 0.847 |

http://localhost:3003/ Page 16 of 29

#### Bioregion 43

160 records of 116 unique species in 3 cells

| Most common species      |       | Most indicative species  |       |
|--------------------------|-------|--------------------------|-------|
| Name                     | Count | Name                     | Score |
| Rhinella marina          | 5     | Hyloxalus infraguttatus  | 407   |
| Trachycephalus jordani   | 5     | Engystomops pustulatus   | 407   |
| Leptodactylus labrosus   | 4     | Pristimantis walkeri     | 305   |
| Epipedobates machalilla  | 3     | Epipedobates machalilla  | 305   |
| Engystomops pustulatus   | 3     | Trachycephalus jordani   | 291   |
| Trachycephalus venulosus | 3     | Leptodactylus labrosus   | 233   |
| Pristimantis achatinus   | 3     | Hypsiboas rosenbergi     | 76.3  |
| Hyloxalus infraguttatus  | 3     | Pristimantis achatinus   | 76.3  |
| Hypsiboas rosenbergi     | 3     | Rhinella marina          | 1.97  |
| Pristimantis walkeri     | 3     | Trachycephalus venulosus | 1.20  |

#### **Bioregion 44**

147 records of 88 unique species in 1 cells

| Most common species    |       | Most indicative species |       |
|------------------------|-------|-------------------------|-------|
| Name                   | Count | Name                    | Score |
| Pseudoeurycea mixteca  | 5     | Thorius narisovalis     | 407   |
| Thorius arboreus       | 4     | Pseudoeurycea juarezi   | 407   |
| Pseudoeurycea juarezi  | 4     | Thorius macdougalli     | 407   |
| Thorius macdougalli    | 4     | Thorius arboreus        | 407   |
| Craugastor augusti     | 4     | Plectrohyla crassa      | 407   |
| Thorius narisovalis    | 4     | Thorius boreas          | 407   |
| Plectrohyla crassa     | 3     | Pseudoeurycea conanti   | 407   |
| Exerodonta sumichrasti | 3     | Pseudoeurycea mixteca   | 339   |
| Pseudoeurycea conanti  | 3     | Exerodonta sumichrasti  | 174   |
| Thorius boreas         | 3     | Craugastor augusti      | 18.7  |

### Bioregion 45

153 records of 126 unique species in 1 cells

| Most common species       |       | Most indicative species   |       |
|---------------------------|-------|---------------------------|-------|
| Name                      | Count | Name                      | Score |
| Pristimantis vicarius     | 3     | Pristimantis vicarius     | 678   |
| Pristimantis ockendeni    | 2     | Osornophryne bufoniformis | 678   |
| Pristimantis repens       | 2     | Pristimantis repens       | 678   |
| Pristimantis ortizi       | 2     | Pristimantis ortizi       | 678   |
| Pristimantis myersi       | 2     | Pristimantis myersi       | 678   |
| Osornophryne bufoniformis | 2     | Ameerega bilinguis        | 339   |

http://localhost:3003/ Page 17 of 29

| Leptodactylus leptodactyloides | 2 | Pristimantis w-nigrum          | 71.4 |
|--------------------------------|---|--------------------------------|------|
| Ameerega bilinguis             | 2 | Pristimantis ockendeni         | 20.6 |
| Hypsiboas cinerascens          | 2 | Leptodactylus leptodactyloides | 3.43 |
| Pristimantis w-nigrum          | 2 | Hypsiboas cinerascens          | 2.96 |

#### **Bioregion 46**

168 records of 27 unique species in 6 cells

| Most common species        |       | Most indicative species    |       |
|----------------------------|-------|----------------------------|-------|
| Name                       | Count | Name                       | Score |
| Pleurodema bufoninum       | 71    | Alsodes gargola            | 28.7  |
| Rhinella spinulosa         | 25    | Pleurodema bufoninum       | 25.4  |
| Rhinella arenarum          | 19    | Alsodes australis          | 12.3  |
| Odontophrynus occidentalis | 15    | Rhinella rubropunctata     | 11.5  |
| Pleurodema thaul           | 6     | Rhinella spinulosa         | 6.57  |
| Leptodactylus mystacinus   | 4     | Odontophrynus occidentalis | 6.06  |
| Alsodes australis          | 3     | Pleurodema thaul           | 5.55  |
| Alsodes gargola            | 3     | Rhinella arenarum          | 3.43  |
| Batrachyla taeniata        | 2     | Batrachyla taeniata        | 1.98  |
| Rhinella rubropunctata     | 2     | Leptodactylus mystacinus   | 0.252 |

### Bioregion 47

130 records of 107 unique species in 1 cells

| Most common species       |       | Most indicative species   |       |
|---------------------------|-------|---------------------------|-------|
| Name                      | Count | Name                      | Score |
| Bolitoglossa walkeri      | 3     | Caecilia occidentalis     | 678   |
| Pristimantis viridicans   | 3     | Dendropsophus columbianus | 678   |
| Gastrotheca angustifrons  | 3     | Pristimantis viridicans   | 678   |
| Dendropsophus columbianus | 3     | Colostethus mertensi      | 678   |
| Pristimantis boulengeri   | 3     | Pristimantis buckleyi     | 509   |
| Pristimantis buckleyi     | 3     | Bolitoglossa walkeri      | 509   |
| Caecilia occidentalis     | 3     | Gastrotheca angustifrons  | 407   |
| Espadarana prosoblepon    | 2     | Pristimantis boulengeri   | 407   |
| Colostethus mertensi      | 2     | Centrolene buckleyi       | 226   |
| Centrolene buckleyi       | 2     | Espadarana prosoblepon    | 52.2  |

## Bioregion 48

96 records of 7 unique species in 7 cells

| Most common species      |       | Most indicative species |       |
|--------------------------|-------|-------------------------|-------|
| Name                     | Count | Name                    | Score |
| Duttaphrynus dhufarensis | 36    | Euphlyctis ehrenbergii  | 44.6  |

http://localhost:3003/ Page 18 of 29

| Duttaphrynus arabicus  | 33 | Duttaphrynus dhufarensis | 44.2   |
|------------------------|----|--------------------------|--------|
| Euphlyctis ehrenbergii | 15 | Duttaphrynus arabicus    | 43.4   |
| Bufo tihamicus         | 6  | Bufo tihamicus           | 42.4   |
| Hyla savignyi          | 3  | Duttaphrynus dodsoni     | 1.64   |
| Duttaphrynus dodsoni   | 2  | Hyla savignyi            | 1.39   |
| Pelophylax ridibundus  | 1  | Pelophylax ridibundus    | 0.0613 |

### Bioregion 49

97 records of 81 unique species in 1 cells

| Most common species     |       | Most indicative species |       |
|-------------------------|-------|-------------------------|-------|
| Name                    | Count | Name                    | Score |
| Pristimantis proserpens | 5     | Pristimantis bambu      | 407   |
| Engystomops randi       | 5     | Engystomops randi       | 407   |
| Leptodactylus lineatus  | 2     | Pristimantis proserpens | 291   |
| Gastrotheca weinlandii  | 2     | Ceratophrys stolzmanni  | 204   |
| Pristimantis bambu      | 2     | Gastrotheca weinlandii  | 163   |
| Gastrotheca longipes    | 2     | Gastrotheca longipes    | 116   |
| Ceratophrys stolzmanni  | 2     | Leptodactylus wagneri   | 45.2  |
| Leptodactylus wagneri   | 2     | Pristimantis w-nigrum   | 42.8  |
| Engystomops petersi     | 2     | Engystomops petersi     | 23.9  |
| Pristimantis w-nigrum   | 2     | Leptodactylus lineatus  | 1.84  |

### Bioregion 50

114 records of 22 unique species in 4 cells

| Most common species             |       | Most indicative species         |       |
|---------------------------------|-------|---------------------------------|-------|
| Name                            | Count | Name                            | Score |
| Eleutherodactylus johnstonei    | 30    | Eleutherodactylus martinicensis | 67.8  |
| Rhinella marina                 | 27    | Eleutherodactylus johnstonei    | 42.4  |
| Eleutherodactylus martinicensis | 16    | Eleutherodactylus coqui         | 12.3  |
| Osteopilus septentrionalis      | 6     | Osteopilus septentrionalis      | 7.68  |
| Lithobates catesbeianus         | 6     | Pleurodema brachyops            | 4.24  |
| Pleurodema brachyops            | 5     | Eleutherodactylus planirostris  | 3.48  |
| Scinax ruber                    | 3     | Glandirana rugosa               | 2.99  |
| Glandirana rugosa               | 3     | Rhinella marina                 | 1.77  |
| Eleutherodactylus planirostris  | 2     | Lithobates catesbeianus         | 0.474 |
| Eleutherodactylus coqui         | 2     | Scinax ruber                    | 0.301 |

| Bioregion 51  |       |      |       |  |
|---|-------|------|-------|--|
| 121 records of 75 unique species in 1 cells  Most common species  Most indicative species |       |      |       |  |
| Name  | Count | Name | Score |  |

http://localhost:3003/ Page 19 of 29

| Pristimantis nervicus        | 13 | Pristimantis nervicus        | 157   |
|------------------------------|----|------------------------------|-------|
| Leptodactylus fuscus         | 4  | Dendropsophus mathiassoni    | 52.2  |
| Ctenophryne geayi            | 3  | Physalaemus fischeri         | 12.4  |
| Physalaemus fischeri         | 3  | Scinax rostratus             | 5.94  |
| Scinax rostratus             | 3  | Ctenophryne geayi            | 0.970 |
| Dendropsophus mathiassoni    | 3  | Leptodactylus bolivianus     | 0.919 |
| Leptodactylus mystaceus      | 3  | Leptodactylus mystaceus      | 0.803 |
| Siphonops annulatus          | 3  | Phyllomedusa hypochondrialis | 0.690 |
| Phyllomedusa hypochondrialis | 3  | Leptodactylus fuscus         | 0.637 |
| Leptodactylus bolivianus     | 3  | Siphonops annulatus          | 0.570 |

# Bioregion 52

89 records of 67 unique species in 1 cells

| Most common species        |       | Most indicative species    |       |
|----------------------------|-------|----------------------------|-------|
| Name                       | Count | Name                       | Score |
| Scinax flavidus            | 4     | Scinax flavidus            | 254   |
| Leptodactylus fuscus       | 4     | Strabomantis ingeri        | 254   |
| Elachistocleis ovalis      | 3     | Craugastor raniformis      | 72.7  |
| Craugastor raniformis      | 2     | Leptodactylus colombiensis | 72.7  |
| Leptodactylus colombiensis | 2     | Pristimantis w-nigrum      | 53.6  |
| Bolitoglossa altamazonica  | 2     | Bolitoglossa altamazonica  | 6.24  |
| Siphonops annulatus        | 2     | Hypsiboas crepitans        | 3.40  |
| Pristimantis w-nigrum      | 2     | Leptodactylus fuscus       | 2.07  |
| Strabomantis ingeri        | 2     | Elachistocleis ovalis      | 1.68  |
| Hypsiboas crepitans        | 2     | Siphonops annulatus        | 1.23  |

### Bioregion 53

119 records of 94 unique species in 1 cells

| Most common species           |       | Most indicative species       |       |
|-------------------------------|-------|-------------------------------|-------|
| Name                          | Count | Name                          | Score |
| Siphonops annulatus           | 4     | Hyloxalus idiomelus           | 509   |
| Rhinella poeppigii            | 2     | Osteocephalus leoniae         | 339   |
| Dendropsophus rossalleni      | 2     | Osteocephalus elkejungingerae | 78.3  |
| Osteocephalus elkejungingerae | 2     | Rhinella poeppigii            | 35.1  |
| Pristimantis diadematus       | 2     | Pristimantis diadematus       | 27.5  |
| Pristimantis ockendeni        | 2     | Pristimantis ockendeni        | 15.4  |
| Hyloxalus idiomelus           | 2     | Dendropsophus rossalleni      | 9.33  |
| Osteocephalus leoniae         | 2     | Dendropsophus parviceps       | 2.74  |
| Trachycephalus venulosus      | 2     | Siphonops annulatus           | 2.47  |
| Dendropsophus parviceps       | 2     | Trachycephalus venulosus      | 1.00  |

http://localhost:3003/ Page 20 of 29

#### Bioregion 54

86 records of 28 unique species in 2 cells

| Most common species           |       | Most indicative species       |       |
|-------------------------------|-------|-------------------------------|-------|
| Name                          | Count | Name                          | Score |
| Leptodactylus albilabris      | 9     | Eleutherodactylus cochranae   | 226   |
| Eleutherodactylus antillensis | 9     | Eleutherodactylus wightmanae  | 226   |
| Eleutherodactylus coqui       | 8     | Eleutherodactylus richmondi   | 226   |
| Eleutherodactylus cochranae   | 7     | Eleutherodactylus lentus      | 226   |
| Osteopilus septentrionalis    | 6     | Eleutherodactylus gryllus     | 226   |
| Rhinella marina               | 6     | Leptodactylus albilabris      | 226   |
| Eleutherodactylus wightmanae  | 5     | Eleutherodactylus antillensis | 204   |
| Eleutherodactylus richmondi   | 4     | Eleutherodactylus coqui       | 164   |
| Eleutherodactylus lentus      | 3     | Osteopilus septentrionalis    | 25.6  |
| Eleutherodactylus gryllus     | 3     | Rhinella marina               | 1.31  |

#### Bioregion 55

85 records of 69 unique species in 1 cells

| Most common species         |       | Most indicative species     |       |
|-----------------------------|-------|-----------------------------|-------|
| Name                        | Count | Name                        | Score |
| Scinax flavidus             | 3     | Hyloscirtus jahni           | 678   |
| Leptodactylus fuscus        | 3     | Atelopus oxyrhynchus        | 678   |
| Hyloscirtus jahni           | 3     | Aromobates capurinensis     | 678   |
| Hypsiboas crepitans         | 3     | Flectonotus pygmaeus        | 339   |
| Aromobates capurinensis     | 2     | Scinax flavidus             | 254   |
| Gastrotheca nicefori        | 2     | Gastrothecanicefori         | 104   |
| Atelopus oxyrhynchus        | 2     | Hypsiboas crepitans         | 6.81  |
| Flectonotus pygmaeus        | 2     | Dendropsophus microcephalus | 2.98  |
| Dendropsophus microcephalus | 2     | Leptodactylus fuscus        | 2.07  |
| Elachistocleis ovalis       | 2     | Elachistocleis ovalis       | 1.50  |

## Bioregion 56

67 records of 57 unique species in 1 cells

| Most common species           |       | Most indicative species   |       |
|-------------------------------|-------|---------------------------|-------|
| Name                          | Count | Name                      | Score |
| Pristimantis lymani           | 3     | Atelopus pachydermus      | 678   |
| Pristimantis cryptomelas      | 2     | Pristimantis percnopterus | 678   |
| Gastrotheca monticola         | 2     | Pristimantis lymani       | 509   |
| Atelopus pachydermus          | 2     | Pristimantis cryptomelas  | 452   |
| Pristimantis percnopterus     | 2     | Noblella heyeri           | 339   |
| Osteocephalus elkeiungingerae | 2     | Pristimantis muscosus     | 339   |

http://localhost:3003/ Page 21 of 29

| Engystomops petersi   | 2 | Gastrotheca monticola         | 339  |
|-----------------------|---|-------------------------------|------|
| Siphonops annulatus   | 2 | Osteocephalus elkejungingerae | 104  |
| Noblella heyeri       | 2 | Engystomops petersi           | 39.9 |
| Pristimantis muscosus | 1 | Siphonops annulatus           | 1.65 |

#### Bioregion 57

73 records of 32 unique species in 1 cells

| Most common species         |       | Most indicative species     |       |
|-----------------------------|-------|-----------------------------|-------|
| Name                        | Count | Name                        | Score |
| Microbatrachella capensis   | 6     | Microbatrachella capensis   | 339   |
| Amietia fuscigula           | 5     | Xenopus gilli               | 339   |
| Tomopterna delalandii       | 5     | Amietophrynus pantherinus   | 339   |
| Strongylopus grayii         | 5     | Breviceps montanus          | 254   |
| Xenopus laevis              | 4     | Vandijkophrynus angusticeps | 170   |
| Vandijkophrynus angusticeps | 4     | Tomopterna delalandii       | 84.8  |
| Amietophrynus rangeri       | 4     | Amietia fuscigula           | 73.7  |
| Breviceps montanus          | 3     | Strongylopus grayii         | 27.8  |
| Xenopus gilli               | 3     | Amietophrynus rangeri       | 21.2  |
| Amietophrynus pantherinus   | 3     | Xenopus laevis              | 3.26  |

### Bioregion 58

77 records of 62 unique species in 1 cells

| Most common species       |       | Most indicative species   |       |
|---------------------------|-------|---------------------------|-------|
| Name                      | Count | Name                      | Score |
| Pristimantis rhabdolaemus | 3     | Phrynopus juninensis      | 678   |
| Siphonops annulatus       | 3     | Gastrotheca stictopleura  | 339   |
| Rhinella poeppigii        | 3     | Pristimantis rhabdolaemus | 226   |
| Rhinella spinulosa        | 3     | Ranitomeya sirensis       | 113   |
| Engystomops freibergi     | 3     | Gastrotheca testudinea    | 79.8  |
| Hemiphractus helioi       | 2     | Hemiphractus helioi       | 75.4  |
| Ranitomeya sirensis       | 2     | Rhinella poeppigii        | 70.2  |
| Gastrotheca stictopleura  | 2     | Rhinella spinulosa        | 18.7  |
| Phrynopus juninensis      | 2     | Engystomops freibergi     | 10.3  |
| Gastrotheca testudinea    | 2     | Siphonops annulatus       | 2.47  |

# Bioregion 59

69 records of 12 unique species in 1 cells

| Most common species |       | Most indicative species |       |
|---------------------|-------|-------------------------|-------|
| Name                | Count | Name                    | Score |
|                     |       |                         |       |

http://localhost:3003/ Page 22 of 29

| Ptychadena mascareniensis | 12 | Hypogeophis rostratus     | 170  |
|---------------------------|----|---------------------------|------|
| Hypogeophis rostratus     | 10 | Grandisonia alternans     | 170  |
| Sooglossus sechellensis   | 9  | Grandisonia larvata       | 170  |
| Sechellophryne gardineri  | 7  | Tachycnemis seychellensis | 170  |
| Grandisonia alternans     | 7  | Sechellophryne gardineri  | 170  |
| Grandisonia larvata       | 6  | Grandisonia sechellensis  | 170  |
| Tachycnemis seychellensis | 5  | Grandisonia brevis        | 170  |
| Sooglossus thomasseti     | 4  | Sooglossus thomasseti     | 170  |
| Grandisonia sechellensis  | 3  | Sooglossus sechellensis   | 170  |
| Grandisonia brevis        | 3  | Ptychadena mascareniensis | 2.74 |

| Most indicative species |
|-------------------------|
|                         |

| Most common species      |       | Most indicative species  |       |
|--------------------------|-------|--------------------------|-------|
| Name                     | Count | Name                     | Score |
| Fejervarya limnocharis   | 12    | Hylarana elberti         | 170   |
| Polypedates leucomystax  | 11    | Limnonectes kadarsani    | 170   |
| Hylarana elberti         | 7     | Oreophryne jeffersoniana | 170   |
| Kaloula baleata          | 6     | Fejervarya verruculosa   | 170   |
| Fejervarya cancrivora    | 6     | Litoria everetti         | 170   |
| Limnonectes kadarsani    | 6     | Hylarana florensis       | 170   |
| Oreophryne jeffersoniana | 5     | Kaloula baleata          | 9.88  |
| Fejervarya verruculosa   | 5     | Fejervarya cancrivora    | 4.35  |
| Litoria everetti         | 5     | Polypedates leucomystax  | 4.09  |
| Hylarana florensis       | 4     | Fejervarya limnocharis   | 1.65  |

|            | Bioregion 61                      |
|------------|-----------------------------------|
| 65 records | s of 12 unique species in 3 cells |

| Most common species     |       | Most indicative species |       |
|-------------------------|-------|-------------------------|-------|
| Name                    | Count | Name                    | Score |
| Bufotes oblongus        | 24    | Bufotes turanensis      | 78.0  |
| Bufotes turanensis      | 23    | Bufotes oblongus        | 65.6  |
| Duttaphrynus stomaticus | 4     | Rana pseudodalmatina    | 28.3  |
| Pelophylax ridibundus   | 3     | Bufo eichwaldi          | 21.2  |
| Pelobates syriacus      | 2     | Iranodon persicus       | 21.2  |
| Rana pseudodalmatina    | 2     | Pelobates syriacus      | 2.42  |
| Hyla arborea            | 2     | Triturus karelinii      | 1.77  |
| Triturus karelinii      | 1     | Duttaphrynus stomaticus | 0.846 |
| Bufo eichwaldi          | 1     | Hyla arborea            | 0.337 |
| Iranodon persicus       | 1     | Pelophylax ridibundus   | 0.276 |

http://localhost:3003/ Page 23 of 29

#### Bioregion 62

62 records of 22 unique species in 2 cells

| Most common species        |       | Most indicative species    |       |
|----------------------------|-------|----------------------------|-------|
| Name                       | Count | Name                       | Score |
| Limnonectes modestus       | 12    | Oreophryne moluccensis     | 170   |
| Platymantis papuensis      | 9     | Hylarana moluccana         | 170   |
| Litoria infrafrenata       | 8     | Rhacophorus rhyssocephalus | 170   |
| Hylarana moluccana         | 6     | Callulops dubius           | 170   |
| Limnonectes grunniens      | 5     | Callulops boettgeri        | 170   |
| Oreophryne moluccensis     | 5     | Limnonectes modestus       | 53.6  |
| Fejervarya cancrivora      | 2     | Platymantis papuensis      | 26.8  |
| Rhacophorus rhyssocephalus | 1     | Limnonectes grunniens      | 22.9  |
| Callulops dubius           | 1     | Litoria infrafrenata       | 9.62  |
| Callulops boettgeri        | 1     | Fejervarya cancrivora      | 1.45  |

#### Bioregion 63

84 records of 46 unique species in 2 cells

| Most common species         |       | Most indicative species     |       |
|-----------------------------|-------|-----------------------------|-------|
| Name                        | Count | Name                        | Score |
| Scinax ruber                | 4     | Hyalinobatrachium orientale | 382   |
| Leptodactylus fuscus        | 4     | Engystomops pustulosus      | 18.8  |
| Rhinella marina             | 3     | Hypsiboas crepitans         | 5.10  |
| Lithobates palmipes         | 3     | Scinax ruber                | 3.01  |
| Hypsiboas crepitans         | 3     | Lithobates palmipes         | 2.65  |
| Trachycephalus venulosus    | 3     | Leptodactylus fuscus        | 2.07  |
| Hyalinobatrachium orientale | 3     | Adenomera hylaedactyla      | 1.96  |
| Hypsiboas punctatus         | 3     | Hypsiboas punctatus         | 1.64  |
| Engystomops pustulosus      | 3     | Trachycephalus venulosus    | 1.51  |
| Adenomera hylaedactyla      | 3     | Rhinella marina             | 1.48  |

### Bioregion 64

68 records of 40 unique species in 1 cells

| Most common species       |       | Most indicative species   |       |
|---------------------------|-------|---------------------------|-------|
| Name                      | Count | Name                      | Score |
| Pleurodema brachyops      | 4     | Dendropsophus yaracuyanus | 509   |
| Leptodactylus fuscus      | 3     | Pleurodema brachyops      | 25.4  |
| Hypsiboas crepitans       | 3     | Scinax rostratus          | 19.3  |
| Dendropsophus yaracuyanus | 3     | Engystomops pustulosus    | 12.6  |
| Leptodactylus bolivianus  | 3     | Leptodactylus fragilis    | 8.62  |
| Rhinella marina           | 3     | Hypsiboas crepitans       | 5.10  |

http://localhost:3003/ Page 24 of 29

| Scinax rostratus       | 3 | Leptodactylus bolivianus | 2.99 |
|------------------------|---|--------------------------|------|
| Scinax ruber           | 3 | Scinax ruber             | 2.26 |
| Engystomops pustulosus | 2 | Leptodactylus fuscus     | 1.55 |
| Leptodactylus fragilis | 2 | Rhinella marina          | 1.48 |

#### **Bioregion 65**

51 records of 22 unique species in 2 cells

| Most common species     |       | Most indicative species |       |
|-------------------------|-------|-------------------------|-------|
| Name                    | Count | Name                    | Score |
| Platymantis boulengeri  | 4     | Platymantis gilliardi   | 509   |
| Platymantis gilliardi   | 4     | Platymantis boulengeri  | 509   |
| Discodeles guppyi       | 4     | Platymantis nexipus     | 509   |
| Platymantis schmidti    | 4     | Platymantis akarithymus | 509   |
| Litoria thesaurensis    | 4     | Platymantis schmidti    | 509   |
| Hylarana daemeli        | 4     | Platymantis mimicus     | 509   |
| Platymantis nexipus     | 3     | Discodeles guppyi       | 78.3  |
| Platymantis akarithymus | 3     | Hylarana daemeli        | 35.7  |
| Litoria infrafrenata    | 3     | Litoria thesaurensis    | 23.7  |
| Platymantis mimicus     | 3     | Litoria infrafrenata    | 10.8  |

#### Bioregion 66

59 records of 18 unique species in 1 cells

| Most common species        |       | Most indicative species    |       |
|----------------------------|-------|----------------------------|-------|
| Name                       | Count | Name                       | Score |
| Rhinella spinulosa         | 7     | Alsodes pehuenche          | 291   |
| Odontophrynus occidentalis | 7     | Alsodes nodosus            | 291   |
| Pleurodema thaul           | 5     | Rhinella arunco            | 121   |
| Alsodes pehuenche          | 5     | Calyptocephalella gayi     | 77.5  |
| Rhinella arenarum          | 5     | Pleurodema thaul           | 46.9  |
| Rhinella arunco            | 5     | Odontophrynus occidentalis | 28.7  |
| Calyptocephalella gayi     | 4     | Rhinella spinulosa         | 18.7  |
| Xenopus laevis             | 4     | Pleurodema bufoninum       | 10.9  |
| Pleurodema bufoninum       | 3     | Rhinella arenarum          | 9.14  |
| Alsodes nodosus            | 3     | Xenopus laevis             | 2.80  |

### Bioregion 67

44 records of 32 unique species in 1 cells

| Most common species     |       | Most indicative species |         |
|-------------------------|-------|-------------------------|---------|
| Name                    | Count | Name                    | Score   |
| Strabomantis biporcatus | 2     | Strabomantis biporcatus | 1.02e+3 |

http://localhost:3003/ Page 25 of 29

| Scinax ruber                      | 2 | Hyalinobatrachium guairarepanense | 1.02e+3 |
|-----------------------------------|---|-----------------------------------|---------|
| Pleurodema brachyops              | 2 | Gastrotheca walkeri               | 678     |
| Phyllomedusa trinitatis           | 2 | Phyllomedusa trinitatis           | 339     |
| Hyalinobatrachium guairarepanense | 2 | Scinax rostratus                  | 25.8    |
| Hypsiboas crepitans               | 2 | Pleurodema brachyops              | 25.4    |
| Scinax rostratus                  | 2 | Engystomops pustulosus            | 25.1    |
| Leptodactylus bolivianus          | 2 | Hypsiboas crepitans               | 6.81    |
| Gastrotheca walkeri               | 2 | Leptodactylus bolivianus          | 3.98    |
| Engystomops pustulosus            | 2 | Scinax ruber                      | 3.01    |

#### Bioregion 68

32 records of 29 unique species in 1 cells

| Most common species       |       | Most indicative species   |         |
|---------------------------|-------|---------------------------|---------|
| Name                      | Count | Name                      | Score   |
| Rhinella spinulosa        | 2     | Telmatobius carrillae     | 1.02e+3 |
| Ranitomeya sirensis       | 2     | Pristimantis meridionalis | 1.02e+3 |
| Gastrotheca peruana       | 2     | Phrynopus barthlenae      | 1.02e+3 |
| Telmatobius carrillae     | 1     | Telmatobius hockingi      | 1.02e+3 |
| Rhinella multiverrucosa   | 1     | Rhinella multiverrucosa   | 1.02e+3 |
| Siphonops annulatus       | 1     | Gastrotheca peruana       | 678     |
| Pristimantis meridionalis | 1     | Osteocephalus leoniae     | 339     |
| Phrynopus barthlenae      | 1     | Ranitomeya sirensis       | 170     |
| Telmatobius hockingi      | 1     | Rhinella spinulosa        | 18.7    |
| Osteocephalus leoniae     | 1     | Siphonops annulatus       | 1.23    |

### Bioregion 69

31 records of 25 unique species in 1 cells

| Most common species            |       | Most indicative species        |       |
|--------------------------------|-------|--------------------------------|-------|
| Name                           | Count | Name                           | Score |
| Eleutherodactylus planirostris | 3     | Eleutherodactylus tonyi        | 678   |
| Eleutherodactylus tonyi        | 3     | Eleutherodactylus sisyphodemus | 678   |
| Eleutherodactylus johnstonei   | 2     | Eleutherodactylus andrewsi     | 678   |
| Rhinella marina                | 2     | Eleutherodactylus junori       | 678   |
| Osteopilus wilderi             | 1     | Osteopilus wilderi             | 678   |
| Eleutherodactylus albipes      | 1     | Eleutherodactylus albipes      | 339   |
| Eleutherodactylus andrewsi     | 1     | Eleutherodactylus planirostris | 52.2  |
| Eleutherodactylus sisyphodemus | 1     | Eleutherodactylus johnstonei   | 28.3  |
| Eleutherodactylus junori       | 1     | Osteopilus septentrionalis     | 12.8  |
| Osteopilus septentrionalis     | 1     | Rhinella marina                | 1.31  |

Bioregion 70

http://localhost:3003/ Page 26 of 29

| 25 records of 23 unique species in 1 cell | 25 | records | of 23 un | iaue sn | ecies in | 1 cells |
|---|----|---------|----------|---------|----------|---------|
|---|----|---------|----------|---------|----------|---------|

| Most common species          |       | Most indicative              | Most indicative species |  |
|------------------------------|-------|------------------------------|-------------------------|--|
| Name                         | Count | Name                         | Score                   |  |
| Pristimantis petrobardus     | 2     | Pristimantis petrobardus     | 1.02e+3                 |  |
| Atelopus peruensis           | 2     | Gastrotheca aratia           | 1.02e+3                 |  |
| Gastrotheca aratia           | 1     | Atelopus peruensis           | 1.02e+3                 |  |
| Telmatobius brevipes         | 1     | Pristimantis pinguis         | 1.02e+3                 |  |
| Pristimantis pinguis         | 1     | Pristimantis phalaroinguinis | 1.02e+3                 |  |
| Pristimantis phalaroinguinis | 1     | Centrolene hesperium         | 1.02e+3                 |  |
| Centrolene hesperium         | 1     | Telmatobius brevipes         | 1.02e+3                 |  |
| Leptodactylus labrosus       | 1     | Pristimantis chimu           | 1.02e+3                 |  |
| Pristimantis chimu           | 1     | Phrynopus thompsoni          | 1.02e+3                 |  |
| Phrynopus thompsoni          | 1     | Leptodactylus labrosus       | 145                     |  |

#### Bioregion 71

21 records of 5 unique species in 1 cells

| Most common species    |       | Most indicative species |       |
|------------------------|-------|-------------------------|-------|
| Name                   | Count | Name                    | Score |
| Leiopelma hochstetteri | 8     | Leiopelma archeyi       | 254   |
| Litoria raniformis     | 6     | Leiopelma hochstetteri  | 226   |
| Leiopelma archeyi      | 3     | Litoria aurea           | 28.3  |
| Litoria aurea          | 2     | Litoria raniformis      | 19.3  |
| Litoria ewingii        | 2     | Litoria ewingii         | 8.77  |

### Rioregion 72

20 records of 7 unique species in 1 cells

| Most common species        |       | Most indicative species    |       |
|----------------------------|-------|----------------------------|-------|
| Name                       | Count | Name                       | Score |
| Duttaphrynus melanostictus | 7     | Fejervarya nicobariensis   | 233   |
| Fejervarya nicobariensis   | 4     | Ingerana charlesdarwini    | 145   |
| Hylarana nicobariensis     | 3     | Fejervarya andamanensis    | 145   |
| Limnonectes doriae         | 2     | Limnonectes doriae         | 30.6  |
| Kaloula baleata            | 2     | Hylarana nicobariensis     | 10.1  |
| Ingerana charlesdarwini    | 1     | Kaloula baleata            | 5.64  |
| Fejervarya andamanensis    | 1     | Duttaphrynus melanostictus | 2.81  |

| Bioregion 73                              |       |                         |       |
|---|-------|-------------------------|-------|
| 10 records of 1 unique species in 1 cells |       |                         |       |
| Most common species                       |       | Most indicative species |       |
| Name                                      | Count | Name                    | Score |

http://localhost:3003/ Page 27 of 29

| Rana asiatica | 10 | Rana asiatica | 102 |
|---------------|----|---------------|-----|

| Bioregion 74                              |       |                         |       |
|---|-------|-------------------------|-------|
| 11 records of 1 unique species in 1 cells |       |                         |       |
| Most common species                       |       | Most indicative species |       |
| Name                                      | Count | Name                    | Score |
| Platymantis pelewensis                    | 11    | Platymantis pelewensis  | 157   |

| 8 records of 7 unique species in 1 cells |       |                          |            |  |
|--|-------|--------------------------|------------|--|
| Most common species                      |       | Most indication          | ve species |  |
| Name                                     | Count | Name                     | Score      |  |
| Hyperolius molleri                       | 2     | Hyperolius thomensis     | 1.02e+3    |  |
| Hyperolius thomensis                     | 1     | Phrynobatrachus dispar   | 1.02e+3    |  |
| Leptopelis palmatus                      | 1     | Ptychadena newtoni       | 1.02e+3    |  |
| Phrynobatrachus dispar                   | 1     | Leptopelis palmatus      | 1.02e+3    |  |
| Ptychadena newtoni                       | 1     | Hyperolius molleri       | 1.02e+3    |  |
| Phrynobatrachus leveleve                 | 1     | Phrynobatrachus leveleve | 509        |  |
| Schistometopum thomense                  | 1     | Schistometopum thomense  | 509        |  |

| Bioregion 76                                |       |                       |       |  |
|---|-------|-----------------------|-------|--|
| 7 records of 2 unique species in 1 cells    |       |                       |       |  |
| Most common species Most indicative species |       |                       |       |  |
| Name  | Count | Name                  | Score |  |
| Platymantis vitianus                        | 4     | Platymantis vitianus  | 407   |  |
| Platymantis vitiensis                       | 3     | Platymantis vitiensis | 382   |  |

| 7 records of 7 unique species in 1 cells  Most common species |       | Most indicative species     |         |
|---|-------|-----------------------------|---------|
| Name  | Count | Name                        | Score   |
| Hylophorbus rainerguentheri                                   | 1     | Litoria singadanae          | 2.04e+3 |
| Litoria singadanae  | 1     | Hylophorbus rainerguentheri | 2.04e+3 |
| Litoria eucnemis  | 1     | Litoria dux                 | 2.04e+3 |
| Hylarana papua  | 1     | Mantophryne infulata        | 1.02e+3 |
| Mantophryne infulata  | 1     | Austrochaperina mehelyi     | 678     |
| Litoria dux   | 1     | Litoria eucnemis            | 53.6    |
| Austrochaperina mehelyi                                       | 1     | Hylarana papua              | 20.8    |

http://localhost:3003/ Page 28 of 29

#### Bioregion 78

6 records of 3 unique species in 1 cells

| Most common species        |       | Most indicative species    |       |
|----------------------------|-------|----------------------------|-------|
| Name                       | Count | Name                       | Score |
| Lyciasalamandra helverseni | 3     | Lyciasalamandra helverseni | 678   |
| Hyla arborea               | 2     | Pelophylax cerigensis      | 678   |
| Pelophylax cerigensis      | 1     | Hyla arborea               | 2.70  |

#### Bioregion 79

5 records of 3 unique species in 1 cells

| Most common species  |       | Most indicative species |       |
|----------------------|-------|-------------------------|-------|
| Name                 | Count | Name                    | Score |
| Rhinella atacamensis | 3     | Telmatobius hauthali    | 678   |
| Rhinella spinulosa   | 1     | Rhinella atacamensis    | 509   |
| Telmatobius hauthali | 1     | Rhinella spinulosa      | 6.22  |

#### Bioregion 80

5 records of 5 unique species in 1 cells


| Most common species       |       | Most indicative species   |         |
|---------------------------|-------|---------------------------|---------|
| Name                      | Count | Name                      | Score   |
| Telmatobius halli         | 1     | Telmatobius halli         | 2.04e+3 |
| Telmatobius philippii     | 1     | Telmatobius fronteriensis | 2.04e+3 |
| Telmatobius fronteriensis | 1     | Telmatobius philippii     | 2.04e+3 |
| Rhinella spinulosa        | 1     | Telmatobius dankoi        | 2.04e+3 |
| Telmatobius dankoi        | 1     | Rhinella spinulosa        | 18.7    |


http://localhost:3003/ Page 29 of 29

**Infomap Bioregions** 

**>**mammals

About





### Bioregion 1

 ${\bf 635177}$  records of  ${\bf 393}$  unique species in  ${\bf 268}$  cells

| Most common species       |       | Most indicative species   |       |
|---------------------------|-------|---------------------------|-------|
| Name                      | Count | Name                      | Score |
| Meles meles               | 46299 | Talpa europaea            | 1.24  |
| Sciurus vulgaris          | 41263 | Capreolus capreolus       | 1.24  |
| Erinaceus europaeus       | 31423 | Sciurus vulgaris          | 1.24  |
| Vulpes vulpes             | 29962 | Pipistrellus pipistrellus | 1.23  |
| Lutra lutra               | 28925 | Lutra lutra               | 1.23  |
| Capreolus capreolus       | 27477 | Erinaceus europaeus       | 1.22  |
| Pipistrellus pipistrellus | 25626 | Meles meles               | 1.21  |
| Oryctolagus cuniculus     | 23623 | Sciurus carolinensis      | 1.07  |
| Sciurus carolinensis      | 18580 | Oryctolagus cuniculus     | 0.807 |
| Talpa europaea            | 18174 | Vulpes vulpes             | 0.647 |

http://localhost:3003/ Page 1 of 28

# Bioregion 2

298374 records of 269 unique species in 39 cells

| Most common species    |       | Most indicative species |       |
|------------------------|-------|-------------------------|-------|
| Name                   | Count | Name                    | Score |
| Phascolarctos cinereus | 26029 | Petaurus australis      | 2.20  |
| Vulpes vulpes          | 21199 | Petauroides volans      | 2.19  |
| Tachyglossus aculeatus | 13302 | Wallabia bicolor        | 2.18  |
| Vombatus ursinus       | 12698 | Phascolarctos cinereus  | 2.18  |
| Wallabia bicolor       | 12397 | Vombatus ursinus        | 2.16  |
| Petaurus australis     | 11811 | Macropus giganteus      | 2.11  |
| Petauroides volans     | 11757 | Petaurus breviceps      | 2.10  |
| Trichosurus vulpecula  | 11407 | Trichosurus vulpecula   | 1.88  |
| Petaurus breviceps     | 10373 | Tachyglossus aculeatus  | 1.87  |
| Macropus giganteus     | 9103  | Vulpes vulpes           | 0.814 |

## Bioregion 3

 $277816\,\text{records}$  of  $807\,\text{unique}$  species in  $423\,\text{cells}$ 

| Most common species       |       | Most indicative species   |       |
|---------------------------|-------|---------------------------|-------|
| Name                      | Count | Name                      | Score |
| Peromyscus maniculatus    | 17600 | Dipodomys ordii           | 3.24  |
| Peromyscus leucopus       | 7473  | Sciurus niger             | 3.22  |
| Thomomys bottae           | 4964  | Dipodomys merriami        | 3.21  |
| Reithrodontomys megalotis | 4437  | Thomomys bottae           | 3.21  |
| Dipodomys merriami        | 3935  | Peromyscus maniculatus    | 3.20  |
| Sigmodon hispidus         | 3737  | Reithrodontomys megalotis | 3.15  |
| Dipodomys ordii           | 3725  | Peromyscus leucopus       | 3.15  |
| Canis latrans             | 3629  | Canis latrans             | 3.12  |
| Sciurus niger             | 3587  | Microtus pennsylvanicus   | 2.82  |
| Microtus pennsylvanicus   | 3512  | Sigmodon hispidus         | 2.69  |

### **Bioregion 4**

http://localhost:3003/ Page 2 of 28

/ 0403 records or 14/7 unique species in 202 cens

| Most common species    |       | Most indicative species |       |
|------------------------|-------|-------------------------|-------|
| Name                   | Count | Name                    | Score |
| Glossophaga soricina   | 1584  | Tapirus terrestris      | 35.9  |
| Artibeus lituratus     | 1561  | Mazama americana        | 35.8  |
| Carollia perspicillata | 1552  | Uroderma bilobatum      | 35.8  |
| Sturnira lilium        | 1473  | Carollia perspicillata  | 35.5  |
| Artibeus jamaicensis   | 1465  | Artibeus lituratus      | 35.4  |
| Tapirus terrestris     | 1353  | Carollia brevicauda     | 35.0  |
| Desmodus rotundus      | 1238  | Sturnira lilium         | 34.1  |
| Carollia brevicauda    | 825   | Glossophaga soricina    | 33.7  |
| Mazama americana       | 771   | Desmodus rotundus       | 32.5  |
| Uroderma bilobatum     | 739   | Artibeus jamaicensis    | 30.9  |

# Bioregion 5

44874 records of 148 unique species in 24 cells

| Most common species    |       | Most indicative species |       |
|------------------------|-------|-------------------------|-------|
| Name                   | Count | Name                    | Score |
| Macropus robustus      | 9180  | Petrogale xanthopus     | 6.23  |
| Macropus rufus         | 6321  | Macropus fuliginosus    | 4.53  |
| Petrogale xanthopus    | 5635  | Macropus robustus       | 4.36  |
| Macropus fuliginosus   | 4914  | Capra hircus            | 4.23  |
| Capra hircus           | 4094  | Macropus rufus          | 3.90  |
| Vulpes vulpes          | 1655  | Nyctophilus geoffroyi   | 0.820 |
| Oryctolagus cuniculus  | 1627  | Mus musculus            | 0.372 |
| Mus musculus           | 1201  | Tachyglossus aculeatus  | 0.299 |
| Tachyglossus aculeatus | 749   | Oryctolagus cuniculus   | 0.280 |
| Nyctophilus geoffroyi  | 632   | Vulpes vulpes           | 0.180 |

| <br>IO | ra | $\alpha$ 1 | ı 6 |
|--------|----|------------|-----|
|        |    |            |     |

| 39011 records of 263 unique species in 93 cells |                         |
|---|-------------------------|
| ·   |                         |
| Most common species                             | Most indicative species |
|   |                         |
|   |                         |

http://localhost:3003/ Page 3 of 28

| Name                        | Count | Name                        | Score |
|-----------------------------|-------|-----------------------------|-------|
| Oryctolagus cuniculus       | 4522  | Notomys alexis              | 12.1  |
| Macropus rufus              | 3211  | Pseudomys hermannsburgensis | 11.9  |
| Pseudomys hermannsburgensis | 2928  | Camelus dromedarius         | 11.2  |
| Mus musculus                | 2922  | Sminthopsis macroura        | 8.83  |
| Canis lupus                 | 1905  | Bos taurus                  | 4.07  |
| Camelus dromedarius         | 1525  | Macropus rufus              | 4.02  |
| Macropus robustus           | 1486  | Mus musculus                | 1.84  |
| Notomys alexis              | 1482  | Oryctolagus cuniculus       | 1.58  |
| Sminthopsis macroura        | 1381  | Macropus robustus           | 1.43  |
| Bos taurus                  | 915   | Canis lupus                 | 1.15  |

# Bioregion 7

38226 records of 1102 unique species in 322 cells

| Most common species   |       | Most indicative species |       |
|-----------------------|-------|-------------------------|-------|
| Name                  | Count | Name                    | Score |
| Mastomys natalensis   | 991   | Mus musculoides         | 57.6  |
| Loxodonta africana    | 619   | Praomys jacksoni        | 57.5  |
| Lemniscomys striatus  | 550   | Lophuromys aquilus      | 57.4  |
| Rattus rattus         | 504   | Mastomys natalensis     | 57.4  |
| Mus musculoides       | 444   | Lemniscomys striatus    | 57.4  |
| Praomys jacksoni      | 362   | Loxodonta africana      | 57.3  |
| Arvicanthis niloticus | 353   | Oenomys hypoxanthus     | 57.2  |
| Lophuromys aquilus    | 347   | Hipposideros caffer     | 56.5  |
| Hipposideros caffer   | 300   | Arvicanthis niloticus   | 51.4  |
| Oenomys hypoxanthus   | 278   | Rattus rattus           | 3.71  |

# Bioregion 8

28221 records of 223 unique species in 50 cells

| Most common species |       | Most indicative species |       |
|---------------------|-------|-------------------------|-------|
| Name                | Count | Name                    | Score |
|                     |       |                         |       |

http://localhost:3003/ Page 4 of 28

| Macropus agilis       | 1777 | Macropus agilis       | 31.0 |
|-----------------------|------|-----------------------|------|
| Isoodon macrourus     | 1465 | Dasyurus hallucatus   | 29.7 |
| Melomys burtoni       | 1405 | Pseudomys nanus       | 28.7 |
| Rattus tunneyi        | 1344 | Zyzomys argurus       | 28.4 |
| Zyzomys argurus       | 1261 | Pseudomys delicatulus | 27.8 |
| Dasyurus hallucatus   | 1035 | Rattus tunneyi        | 26.6 |
| Pseudomys delicatulus | 919  | Melomys burtoni       | 26.3 |
| Pseudomys nanus       | 889  | Isoodon macrourus     | 16.4 |
| Macropus robustus     | 736  | Macropus robustus     | 1.80 |
| Canis lupus           | 682  | Canis lupus           | 1.05 |

### Bioregion 9

21656 records of 156 unique species in 9 cells

| Most common species |       | Most indicative species |       |
|---------------------|-------|-------------------------|-------|
| Name                | Count | Name                    | Score |
| Gazella gazella     | 3824  | Gazella gazella         | 14.9  |
| Canis aureus        | 2209  | Hyaena hyaena           | 14.4  |
| Vulpes vulpes       | 1617  | Hystrix indica          | 14.4  |
| Gazella dorcas      | 1263  | Gazella dorcas          | 14.2  |
| Canis lupus         | 1182  | Canis aureus            | 13.9  |
| Hyaena hyaena       | 1164  | Capra ibex              | 13.0  |
| Capra ibex          | 1034  | Sus scrofa              | 0.966 |
| Sus scrofa          | 988   | Canis lupus             | 0.842 |
| Meles meles         | 867   | Vulpes vulpes           | 0.423 |
| Hystrix indica      | 836   | Meles meles             | 0.275 |

# Bioregion 10

13757 records of 907 unique species in 205 cells

| Most common species   |       | Most indicative species |       |
|-----------------------|-------|-------------------------|-------|
| Name                  | Count | Name                    | Score |
| Rattus exulans        | 400   | Eonycteris spelaea      | 143   |
| Cynopterus brachyotis | 379   | Cynopterus brachyotis   | 143   |

http://localhost:3003/ Page 5 of 28

| Rattus rattus             | 362 | Rattus everetti           | 141  |
|---------------------------|-----|---------------------------|------|
| Macroglossus minimus      | 286 | Rattus tanezumi           | 136  |
| Suncus murinus            | 241 | Panthera tigris           | 135  |
| Rattus tanezumi           | 209 | Suncus murinus            | 124  |
| Rousettus amplexicaudatus | 198 | Rattus exulans            | 96.2 |
| Eonycteris spelaea        | 164 | Rousettus amplexicaudatus | 88.8 |
| Panthera tigris           | 162 | Macroglossus minimus      | 78.9 |
| Rattus everetti           | 158 | Rattus rattus             | 6.61 |

# Bioregion 11

16294 records of 163 unique species in 171 cells

| Most common species     |       | Most indicative species |       |
|-------------------------|-------|-------------------------|-------|
| Name                    | Count | Name                    | Score |
| Myodes rutilus          | 2047  | Myodes rutilus          | 24.1  |
| Sorex cinereus          | 1313  | Microtus oeconomus      | 20.6  |
| Microtus oeconomus      | 1222  | Gulo gulo               | 17.2  |
| Canis lupus             | 816   | Lynx canadensis         | 14.9  |
| Sorex monticolus        | 656   | Ursus arctos            | 10.0  |
| Ursus arctos            | 530   | Sorex cinereus          | 9.13  |
| Microtus pennsylvanicus | 502   | Sorex monticolus        | 6.13  |
| Gulo gulo               | 490   | Microtus pennsylvanicus | 3.46  |
| Tamiasciurus hudsonicus | 431   | Tamiasciurus hudsonicus | 3.18  |
| Lynx canadensis         | 407   | Canis lupus             | 1.09  |

# Bioregion 12

10525 records of 124 unique species in 8 cells

| Most common species   |       | Most indicative species |       |
|-----------------------|-------|-------------------------|-------|
| Name                  | Count | Name                    | Score |
| Isoodon obesulus      | 1430  | Isoodon obesulus        | 25.2  |
| Macropus fuliginosus  | 1142  | Macropus fuliginosus    | 6.76  |
| Oryctolagus cuniculus | 719   | Antechinus flavipes     | 3.90  |
| Mus musculus          | 650   | Rattus fuscipes         | 1.69  |

http://localhost:3003/ Page 6 of 28

| Tachyglossus aculeatus   | 583 | Trichosurus vulpecula    | 1.54  |
|--------------------------|-----|--------------------------|-------|
| Trichosurus vulpecula    | 513 | Tachyglossus aculeatus   | 1.49  |
| Vulpes vulpes            | 508 | Mus musculus             | 1.29  |
| Rattus fuscipes          | 390 | Pseudocheirus peregrinus | 1.28  |
| Antechinus flavipes      | 261 | Oryctolagus cuniculus    | 0.795 |
| Pseudocheirus peregrinus | 257 | Vulpes vulpes            | 0.355 |

# Bioregion 13

4913 records of 158 unique species in 48 cells

| Most common species        |       | Most indicative species    |       |
|----------------------------|-------|----------------------------|-------|
| Name                       | Count | Name                       | Score |
| Abrothrix longipilis       | 581   | Geoxus valdivianus         | 98.5  |
| Abrothrix olivaceus        | 454   | Loxodontomys micropus      | 98.0  |
| Oligoryzomys longicaudatus | 442   | Abrothrix longipilis       | 97.9  |
| Puma concolor              | 382   | Abrothrix olivaceus        | 97.3  |
| Loxodontomys micropus      | 175   | Chelemys macronyx          | 96.7  |
| Geoxus valdivianus         | 132   | Pudu puda                  | 96.7  |
| Lama guanicoe              | 130   | Eligmodontia morgani       | 93.2  |
| Eligmodontia morgani       | 105   | Lama guanicoe              | 88.4  |
| Chelemys macronyx          | 104   | Oligoryzomys longicaudatus | 87.6  |
| Pudu puda                  | 102   | Puma concolor              | 20.9  |

# Bioregion 14

5107 records of 325 unique species in 46 cells

| Most common species       |       | Most indicative species |       |
|---------------------------|-------|-------------------------|-------|
| Name                      | Count | Name                    | Score |
| Syconycteris australis    | 260   | Echymipera kalubu       | 220   |
| Rattus exulans            | 151   | Melomys rufescens       | 218   |
| Macroglossus minimus      | 134   | Nyctimene albiventer    | 208   |
| Rousettus amplexicaudatus | 121   | Dobsonia moluccensis    | 206   |
| Dobsonia moluccensis      | 117   | Phalanger orientalis    | 199   |
| Nvctimene albiventer      | 116   | Svconvcteris australis  | 106   |

http://localhost:3003/ Page 7 of 28

| •                    |     | • •                       |      |
|----------------------|-----|---------------------------|------|
| Phalanger orientalis | 103 | Rousettus amplexicaudatus | 83.5 |
| Melomys rufescens    | 102 | Hipposideros diadema      | 66.3 |
| Echymipera kalubu    | 96  | Macroglossus minimus      | 56.9 |
| Hipposideros diadema | 87  | Rattus exulans            | 55.9 |

# Bioregion 15

5605 records of 429 unique species in 51 cells

| Most common species   |       | Most indicative species |       |
|-----------------------|-------|-------------------------|-------|
| Name                  | Count | Name                    | Score |
| Phyllotis xanthopygus | 296   | Akodon albiventer       | 191   |
| Galea musteloides     | 116   | Calomys lepidus         | 185   |
| Graomys griseoflavus  | 110   | Phyllotis osilae        | 180   |
| Tadarida brasiliensis | 95    | Akodon boliviensis      | 169   |
| Phyllotis osilae      | 94    | Galea musteloides       | 140   |
| Calomys musculinus    | 90    | Phyllotis xanthopygus   | 137   |
| Akodon albiventer     | 90    | Calomys musculinus      | 118   |
| Akodon boliviensis    | 88    | Graomys griseoflavus    | 113   |
| Mus musculus          | 86    | Tadarida brasiliensis   | 9.98  |
| Calomys lepidus       | 86    | Mus musculus            | 0.826 |

## Bioregion 16

**5542** records of **126** unique species in **14** cells

| Most common species   |       | Most indicative species |       |
|-----------------------|-------|-------------------------|-------|
| Name                  | Count | Name                    | Score |
| Mus musculus          | 546   | Tarsipes rostratus      | 97.5  |
| Tarsipes rostratus    | 320   | Cercartetus concinnus   | 36.7  |
| Macropus fuliginosus  | 286   | Phascogale tapoatafa    | 16.7  |
| Cercartetus concinnus | 283   | Vespadelus regulus      | 8.07  |
| Rattus fuscipes       | 276   | Antechinus flavipes     | 6.53  |
| Phascogale tapoatafa  | 238   | Macropus fuliginosus    | 4.43  |
| Chalinolobus gouldii  | 193   | Rattus fuscipes         | 3.14  |
| Manadalia rasidia     | 100   | Chalinalahus sauldii    | 2.00  |

http://localhost:3003/ Page 8 of 28

| vespaueius i eguius | 170 | CHAIIIIOIODUS goululi | ∠.77 |
|---------------------|-----|-----------------------|------|
| Rattus rattus       | 170 | Mus musculus          | 2.84 |
| Antechinus flavipes | 167 | Rattus rattus         | 2.27 |

### **Bioregion 17**

3834 records of 85 unique species in 7 cells

| Most common species   |       | Most indicative species |        |
|-----------------------|-------|-------------------------|--------|
| Name                  | Count | Name                    | Score  |
| Tursiops truncatus    | 2890  | Tursiops truncatus      | 14.3   |
| Phocoena phocoena     | 89    | Geomys pinetis          | 5.50   |
| Geomys pinetis        | 55    | Kogia breviceps         | 3.52   |
| Peromyscus gossypinus | 45    | Peromyscus gossypinus   | 0.939  |
| Oryzomys palustris    | 43    | Oryzomys palustris      | 0.748  |
| Sigmodon hispidus     | 42    | Phocoena phocoena       | 0.457  |
| Delphinus delphis     | 31    | Delphinus delphis       | 0.444  |
| Sciurus carolinensis  | 30    | Sigmodon hispidus       | 0.184  |
| Kogia breviceps       | 29    | Sciurus niger           | 0.142  |
| Sciurus niger         | 26    | Sciurus carolinensis    | 0.0277 |

# Bioregion 18

415 records of 50 unique species in 6 cells

| Most common species    |       | Most indicative        | Most indicative species |  |
|------------------------|-------|------------------------|-------------------------|--|
| Name                   | Count | Name                   | Score                   |  |
| Rattus rattus          | 102   | Zalophus wollebaeki    | 561                     |  |
| Mus musculus           | 69    | Arctocephalus forsteri | 45.2                    |  |
| Erinaceus europaeus    | 40    | Homo sapiens           | 29.5                    |  |
| Homo sapiens           | 24    | Rattus rattus          | 7.30                    |  |
| Zalophus wollebaeki    | 20    | Mus musculus           | 1.92                    |  |
| Trichosurus vulpecula  | 17    | Capra hircus           | 0.930                   |  |
| Rattus norvegicus      | 15    | Rattus norvegicus      | 0.737                   |  |
| Arctocephalus forsteri | 14    | Trichosurus vulpecula  | 0.716                   |  |
| <b>.</b>               | 10    | - ·                    | 0.707                   |  |

http://localhost:3003/ Page 9 of 28

| Capra nırcus | 10 | Erinaceus europaeus | 0./0/ |
|--------------|----|---------------------|-------|
| Sus scrofa   | 9  | Sus scrofa          | 0.330 |

### **Bioregion 19**

67 records of 17 unique species in 2 cells

| Most common species   |       | Most indicative species |       |
|-----------------------|-------|-------------------------|-------|
| Name                  | Count | Name                    | Score |
| Lepus europaeus       | 10    | Rupicapra rupicapra     | 46.2  |
| Felis catus           | 9     | Homo sapiens            | 25.1  |
| Oryctolagus cuniculus | 8     | Felis catus             | 6.16  |
| Rupicapra rupicapra   | 7     | Lepus europaeus         | 4.00  |
| Trichosurus vulpecula | 6     | Cervus elaphus          | 3.69  |
| Cervus elaphus        | 6     | Capra hircus            | 2.85  |
| Erinaceus europaeus   | 4     | Trichosurus vulpecula   | 2.58  |
| Mus musculus          | 3     | Oryctolagus cuniculus   | 1.27  |
| Capra hircus          | 3     | Mus musculus            | 0.853 |
| Homo sapiens          | 2     | Erinaceus europaeus     | 0.721 |

# Bioregion 20

3312 records of 202 unique species in 78 cells

| Most common species   |       | Most indicative species |       |
|-----------------------|-------|-------------------------|-------|
| Name                  | Count | Name                    | Score |
| Gerbillus gerbillus   | 224   | Dipodillus campestris   | 242   |
| Jaculus jaculus       | 207   | Gerbillus gerbillus     | 232   |
| Mus musculus          | 134   | Gerbillus pyramidum     | 201   |
| Acomys cahirinus      | 132   | Jaculus jaculus         | 171   |
| Gerbillus pyramidum   | 131   | Meriones libycus        | 162   |
| Meriones libycus      | 115   | Meriones crassus        | 155   |
| Meriones crassus      | 115   | Psammomys obesus        | 147   |
| Vulpes vulpes         | 101   | Acomys cahirinus        | 130   |
| Psammomys obesus      | 90    | Mus musculus            | 1.70  |
| Dipodillus campestris | 87    | Vulpes vulpes           | 0.451 |

http://localhost:3003/ Page 10 of 28

## Bioregion 21

3328 records of 163 unique species in 8 cells

| Most common species      |       | Most indicative species  |       |
|--------------------------|-------|--------------------------|-------|
| Name                     | Count | Name                     | Score |
| Melomys cervinipes       | 126   | Dendrolagus lumholtzi    | 428   |
| Rattus leucopus          | 126   | Rattus leucopus          | 353   |
| Rhinolophus megaphyllus  | 122   | Uromys caudimaculatus    | 272   |
| Melomys burtoni          | 114   | Rattus sordidus          | 220   |
| Rattus fuscipes          | 106   | Melomys cervinipes       | 57.3  |
| Uromys caudimaculatus    | 95    | Melomys burtoni          | 30.1  |
| Isoodon macrourus        | 92    | Rhinolophus megaphyllus  | 26.6  |
| Rattus sordidus          | 81    | Isoodon macrourus        | 14.5  |
| Miniopterus schreibersii | 69    | Miniopterus schreibersii | 7.62  |
| Dendrolagus lumholtzi    | 66    | Rattus fuscipes          | 5.23  |

## Bioregion 22

1295 records of 56 unique species in 7 cells

| Most common species   |       | Most indicative species |        |
|-----------------------|-------|-------------------------|--------|
| Name                  | Count | Name                    | Score  |
| Mirounga leonina      | 1004  | Mirounga leonina        | 51.8   |
| Orcinus orca          | 122   | Orcinus orca            | 27.4   |
| Otaria flavescens     | 32    | Otaria flavescens       | 22.0   |
| Mus musculus          | 11    | Eligmodontia typus      | 4.03   |
| Eubalaena australis   | 10    | Eubalaena australis     | 3.96   |
| Eligmodontia typus    | 7     | Arctocephalus gazella   | 2.83   |
| Hydrurga leptonyx     | 7     | Hydrurga leptonyx       | 2.39   |
| Arctocephalus gazella | 6     | Graomys griseoflavus    | 1.81   |
| Graomys griseoflavus  | 6     | Tursiops truncatus      | 0.0712 |
| Tursiops truncatus    | 5     | Mus musculus            | 0.0311 |

D:----:-- 00

http://localhost:3003/ Page 11 of 28

#### ाणा स्थाणा ४०

2164 records of 257 unique species in 50 cells

| Most common species    |       | Most indicative species |       |
|------------------------|-------|-------------------------|-------|
| Name                   | Count | Name                    | Score |
| Mus musculus           | 170   | Calomyscus baluchi      | 337   |
| Vulpes vulpes          | 105   | Meriones persicus       | 323   |
| Cricetulus migratorius | 85    | Tatera indica           | 225   |
| Meriones persicus      | 68    | Nesokia indica          | 195   |
| Meriones libycus       | 59    | Cricetulus migratorius  | 190   |
| Meriones crassus       | 48    | Gerbillus nanus         | 152   |
| Nesokia indica         | 47    | Meriones libycus        | 109   |
| Calomyscus baluchi     | 44    | Meriones crassus        | 85.1  |
| Gerbillus nanus        | 43    | Mus musculus            | 2.84  |
| Tatera indica          | 42    | Vulpes vulpes           | 0.618 |

### Bioregion 24

1668 records of 82 unique species in 7 cells

| Most common species    |       | Most indicative species |       |
|------------------------|-------|-------------------------|-------|
| Name                   | Count | Name                    | Score |
| Potorous tridactylus   | 109   | Sarcophilus harrisii    | 482   |
| Perameles gunnii       | 99    | Thylogale billardierii  | 398   |
| Dasyurus viverrinus    | 83    | Perameles gunnii        | 377   |
| Trichosurus vulpecula  | 78    | Dasyurus viverrinus     | 283   |
| Sarcophilus harrisii   | 77    | Globicephala melas      | 70.6  |
| Globicephala melas     | 71    | Potorous tridactylus    | 43.3  |
| Macropus rufogriseus   | 70    | Isoodon obesulus        | 13.9  |
| Thylogale billardierii | 69    | Rattus lutreolus        | 13.5  |
| Rattus lutreolus       | 64    | Macropus rufogriseus    | 5.70  |
| Isoodon obesulus       | 60    | Trichosurus vulpecula   | 3.07  |

### **Bioregion 25**

1093 records of 103 unique species in 13 cells

Most common species Most indicative species

http://localhost:3003/ Page 12 of 28

| Name                     | Count | Name                     | Score |
|--------------------------|-------|--------------------------|-------|
| Sus scrofa               | 178   | Microtus montebelli      | 322   |
| Paguma larvata           | 159   | Urotrichus talpoides     | 322   |
| Nyctereutes procyonoides | 142   | Lepus brachyurus         | 297   |
| Apodemus speciosus       | 69    | Apodemus speciosus       | 258   |
| Procyon lotor            | 55    | Paguma larvata           | 254   |
| Urotrichus talpoides     | 43    | Apodemus argenteus       | 250   |
| Lepus brachyurus         | 36    | Nyctereutes procyonoides | 52.0  |
| Apodemus argenteus       | 35    | Cervus nippon            | 5.61  |
| Microtus montebelli      | 23    | Procyon lotor            | 4.34  |
| Cervus nippon            | 22    | Sus scrofa               | 3.74  |

### Bioregion 26

941 records of 136 unique species in 19 cells

| Most common species       |       | Most indicative species   |         |
|---------------------------|-------|---------------------------|---------|
| Name                      | Count | Name                      | Score   |
| Rattus rattus             | 42    | Hemicentetes semispinosus | 1.36e+3 |
| Tadarida pumila           | 28    | Eliurus webbi             | 1.36e+3 |
| Indri indri               | 28    | Microgale talazaci        | 1.36e+3 |
| Microgale talazaci        | 26    | Setifer setosus           | 1.36e+3 |
| Setifer setosus           | 24    | Propithecus diadema       | 1.36e+3 |
| Hemicentetes semispinosus | 23    | Indri indri               | 1.32e+3 |
| Eliurus webbi             | 22    | Hapalemur griseus         | 1.30e+3 |
| Hapalemur griseus         | 21    | Tenrec ecaudatus          | 1.19e+3 |
| Tenrec ecaudatus          | 21    | Tadarida pumila           | 283     |
| Propithecus diadema       | 21    | Rattus rattus             | 7.30    |

# Bioregion 27

1331 records of 73 unique species in 22 cells

| Most common species |       | Most indicative species |       |
|---------------------|-------|-------------------------|-------|
| Name                | Count | Name                    | Score |

http://localhost:3003/ Page 13 of 28

| Artibeus jamaicensis    | 215 | Monophyllus redmani     | 266  |
|-------------------------|-----|-------------------------|------|
| Monophyllus redmani     | 121 | Erophylla sezekorni     | 264  |
| Erophylla sezekorni     | 110 | Pteronotus quadridens   | 259  |
| Macrotus waterhousii    | 106 | Brachyphylla cavernarum | 141  |
| Molossus molossus       | 68  | Macrotus waterhousii    | 130  |
| Pteronotus parnellii    | 64  | Artibeus jamaicensis    | 33.4 |
| Eptesicus fuscus        | 60  | Molossus molossus       | 23.5 |
| Brachyphylla cavernarum | 60  | Pteronotus parnellii    | 22.9 |
| Tadarida brasiliensis   | 54  | Tadarida brasiliensis   | 7.81 |
| Pteronotus quadridens   | 34  | Eptesicus fuscus        | 5.60 |

# Bioregion 28

1205 records of 165 unique species in 38 cells

| Most common species      |       | Most indicative species  |       |
|--------------------------|-------|--------------------------|-------|
| Name                     | Count | Name                     | Score |
| Meriones meridianus      | 62    | Ochotona pallasi         | 923   |
| Apodemus uralensis       | 51    | Cricetulus longicaudatus | 862   |
| Allactaga sibirica       | 47    | Phodopus roborovskii     | 821   |
| Dipus sagitta            | 47    | Allactaga sibirica       | 819   |
| Ochotona pallasi         | 44    | Dipus sagitta            | 804   |
| Cricetulus longicaudatus | 42    | Meriones meridianus      | 795   |
| Meriones unguiculatus    | 42    | Meriones unguiculatus    | 693   |
| Mus musculus             | 38    | Apodemus uralensis       | 645   |
| Cricetulus migratorius   | 36    | Cricetulus migratorius   | 220   |
| Phodopus roborovskii     | 32    | Mus musculus             | 1.74  |

## **Bioregion 29**

1600 records of 148 unique species in 3 cells

| Most common species  |       | Most indicative species |       |
|----------------------|-------|-------------------------|-------|
| Name                 | Count | Name                    | Score |
| Chaetodipus goldmani | 93    | Chaetodipus artus       | 483   |
|                      | 00    | Cl. I II I              | 475   |

http://localhost:3003/ Page 14 of 28

| Cnaetodipus pernix   | 72 | Cnaetodipus goldmani | 400  |
|----------------------|----|----------------------|------|
| Peromyscus eremicus  | 88 | Chaetodipus pernix   | 396  |
| Thomomys bottae      | 66 | Peromyscus merriami  | 201  |
| Liomys pictus        | 60 | Liomys pictus        | 63.7 |
| Chaetodipus artus    | 51 | Peromyscus eremicus  | 21.6 |
| Onychomys torridus   | 45 | Onychomys torridus   | 21.1 |
| Dipodomys merriami   | 40 | Glossophaga soricina | 14.1 |
| Glossophaga soricina | 39 | Thomomys bottae      | 8.07 |
| Peromyscus merriami  | 36 | Dipodomys merriami   | 6.18 |

# Bioregion 30

678 records of 79 unique species in 11 cells

| Most common species        |       | Most indicative species    |        |
|----------------------------|-------|----------------------------|--------|
| Name                       | Count | Name                       | Score  |
| Phocoena phocoena          | 176   | Eubalaena glacialis        | 91.8   |
| Megaptera novaeangliae     | 121   | Balaenoptera physalus      | 89.3   |
| Balaenoptera acutorostrata | 55    | Megaptera novaeangliae     | 86.5   |
| Balaenoptera physalus      | 45    | Balaenoptera acutorostrata | 55.1   |
| Canis latrans              | 27    | Phocoena phocoena          | 14.8   |
| Microtus pennsylvanicus    | 17    | Lynx canadensis            | 4.26   |
| Vulpes vulpes              | 16    | Canis latrans              | 2.32   |
| Eubalaena glacialis        | 11    | Clethrionomys gapperi      | 2.00   |
| Lynx canadensis            | 10    | Microtus pennsylvanicus    | 1.36   |
| Clethrionomys gapperi      | 8     | Vulpes vulpes              | 0.0909 |

# Bioregion 31

1143 records of 160 unique species in 43 cells

| Most common species |       | Most indicative species |       |
|---------------------|-------|-------------------------|-------|
| Name                | Count | Name                    | Score |
| Myodes rutilus      | 100   | Microtus fortis         | 506   |
| Myodes rufocanus    | 91    | Ochotona hyperborea     | 497   |
| Sorex caecutiens    | 66    | Apodemus peninsulae     | 433   |

http://localhost:3003/ Page 15 of 28

| Tamias sibiricus    | 65 | Sorex caecutiens   | 356   |
|---------------------|----|--------------------|-------|
| Microtus oeconomus  | 60 | Tamias sibiricus   | 351   |
| Apodemus peninsulae | 34 | Myodes rufocanus   | 322   |
| Ochotona hyperborea | 33 | Apodemus agrarius  | 31.0  |
| Apodemus agrarius   | 30 | Myodes rutilus     | 24.1  |
| Mus musculus        | 24 | Microtus oeconomus | 20.7  |
| Microtus fortis     | 23 | Mus musculus       | 0.682 |

### Bioregion 32

 $662\,\text{records}$  of  $143\,\text{unique}$  species in  $12\,\text{cells}$ 

| Most common species     |       | Most indicative species |         |
|-------------------------|-------|-------------------------|---------|
| Name                    | Count | Name                    | Score   |
| Niviventer confucianus  | 44    | Apodemus chevrieri      | 1.30e+3 |
| Apodemus draco          | 32    | Apodemus draco          | 1.22e+3 |
| Niviventer fulvescens   | 24    | Anourosorex squamipes   | 1.01e+3 |
| Anourosorex squamipes   | 24    | Dremomys pernyi         | 868     |
| Eothenomys melanogaster | 23    | Eothenomys melanogaster | 831     |
| Crocidura attenuata     | 22    | Niviventer confucianus  | 744     |
| Callosciurus erythraeus | 21    | Crocidura attenuata     | 666     |
| Apodemus chevrieri      | 21    | Niviventer fulvescens   | 300     |
| Micromys minutus        | 17    | Callosciurus erythraeus | 209     |
| Dremomys pernyi         | 16    | Micromys minutus        | 7.05    |

### **Bioregion 33**

538 records of 43 unique species in 9 cells

| Most common species    |       | Most indicative species |       |
|------------------------|-------|-------------------------|-------|
| Name                   | Count | Name                    | Score |
| Erinaceus europaeus    | 102   | Phocarctos hookeri      | 528   |
| Arctocephalus forsteri | 92    | Arctocephalus forsteri  | 297   |
| Trichosurus vulpecula  | 55    | Hydrurga leptonyx       | 43.7  |
| Phocarctos hookeri     | 47    | Homo sapiens            | 16.0  |
| Oryctolagus cuniculus  | 34    | Trichosurus vulpecula   | 2.31  |

http://localhost:3003/ Page 16 of 28

| Mus musculus      | 32 | Erinaceus europaeus   | 1.80  |
|-------------------|----|-----------------------|-------|
| Felis catus       | 26 | Felis catus           | 1.75  |
| Rattus rattus     | 20 | Rattus rattus         | 1.43  |
| Homo sapiens      | 13 | Mus musculus          | 0.892 |
| Hydrurga leptonyx | 13 | Oryctolagus cuniculus | 0.527 |

### Bioregion 34

427 records of 60 unique species in 4 cells

| Most common species   |       | Most indicative species |       |
|-----------------------|-------|-------------------------|-------|
| Name                  | Count | Name                    | Score |
| Callicebus lucifer    | 159   | Callicebus lucifer      | 349   |
| Lagothrix lagotricha  | 66    | Cebuella pygmaea        | 285   |
| Cebus albifrons       | 59    | Lagothrix lagotricha    | 229   |
| Alouatta seniculus    | 38    | Cebus albifrons         | 126   |
| Cebuella pygmaea      | 19    | Callicebus cupreus      | 109   |
| Callicebus cupreus    | 10    | Alouatta seniculus      | 57.7  |
| Uroderma magnirostrum | 3     | Uroderma magnirostrum   | 7.30  |
| Sturnira lilium       | 3     | Noctilio albiventris    | 4.74  |
| Molossus molossus     | 3     | Molossus molossus       | 1.40  |
| Noctilio albiventris  | 3     | Sturnira lilium         | 0.692 |

### Bioregion 35

379 records of 40 unique species in 6 cells

| Most common species    |       | Most indicative species |       |
|------------------------|-------|-------------------------|-------|
| Name                   | Count | Name                    | Score |
| Lasiurus cinereus      | 82    | Monachus schauinslandi  | 524   |
| Herpestes javanicus    | 68    | Herpestes javanicus     | 362   |
| Rattus rattus          | 38    | Stenella longirostris   | 170   |
| Mus musculus           | 38    | Lasiurus cinereus       | 45.7  |
| Rattus exulans         | 28    | Rattus exulans          | 32.9  |
| Megaptera novaeangliae | 20    | Megaptera novaeangliae  | 30.7  |

http://localhost:3003/ Page 17 of 28

| Monachus schauinslandi | 15 | Rattus rattus | 3.38  |
|------------------------|----|---------------|-------|
| Stenella longirostris  | 10 | Mus musculus  | 1.32  |
| Sus scrofa             | 10 | Felis catus   | 0.668 |
| Felis catus            | 8  | Sus scrofa    | 0.456 |

### **Bioregion 36**

 $562\,\text{records}$  of  $125\,\text{unique}$  species in  $14\,\text{cells}$ 

| Most common species        |       | Most indicative species    |         |
|----------------------------|-------|----------------------------|---------|
| Name                       | Count | Name                       | Score   |
| Erinaceus concolor         | 49    | Spermophilus xanthoprymnus | 1.02e+3 |
| Sciurus anomalus           | 45    | Sciurus anomalus           | 835     |
| Lepus europaeus            | 22    | Erinaceus concolor         | 421     |
| Vulpes vulpes              | 22    | Rhinolophus hipposideros   | 2.95    |
| Mus musculus               | 18    | Rattus rattus              | 2.09    |
| Apodemus sylvaticus        | 16    | Martes foina               | 1.81    |
| Spermophilus xanthoprymnus | 14    | Lepus europaeus            | 1.80    |
| Rattus rattus              | 14    | Apodemus sylvaticus        | 1.57    |
| Martes foina               | 12    | Mus musculus               | 1.04    |
| Rhinolophus hipposideros   | 12    | Vulpes vulpes              | 0.449   |

### **Bioregion 37**

265 records of 21 unique species in 3 cells

| Most common species     |       | Most indicativ          | Most indicative species |  |
|-------------------------|-------|-------------------------|-------------------------|--|
| Name                    | Count | Name                    | Score                   |  |
| Tadarida teniotis       | 33    | Plecotus teneriffae     | 1.74e+3                 |  |
| Oryctolagus cuniculus   | 32    | Pipistrellus maderensis | 1.74e+3                 |  |
| Mus musculus            | 27    | Atelerix algirus        | 99.6                    |  |
| Pipistrellus maderensis | 24    | Hypsugo savii           | 72.0                    |  |
| Hypsugo savii           | 22    | Tadarida teniotis       | 54.8                    |  |
| Atelerix algirus        | 21    | Suncus etruscus         | 19.1                    |  |
| Plecotus teneriffae     | 20    | Nyctalus leisleri       | 9.78                    |  |
| Nyctalus leisleri       | 17    | Mus musculus            | 2.33                    |  |
|                         |       |                         |                         |  |

http://localhost:3003/ Page 18 of 28

| Suncus etruscus   | 13 | Rattus norvegicus     | 1.98 |
|-------------------|----|-----------------------|------|
| Rattus norvegicus | 13 | Oryctolagus cuniculus | 1.53 |

# Bioregion 38

182 records of 15 unique species in 3 cells

| Most common species   |       | Most indicative species |         |
|-----------------------|-------|-------------------------|---------|
| Name                  | Count | Name                    | Score   |
| Atlantoxerus getulus  | 44    | Crocidura canariensis   | 1.30e+3 |
| Oryctolagus cuniculus | 34    | Atlantoxerus getulus    | 806     |
| Atelerix algirus      | 27    | Atelerix algirus        | 96.0    |
| Crocidura canariensis | 24    | Pipistrellus kuhlii     | 13.5    |
| Mus musculus          | 18    | Hypsugo savii           | 4.91    |
| Pipistrellus kuhlii   | 13    | Oryctolagus cuniculus   | 1.22    |
| Rattus rattus         | 6     | Mus musculus            | 1.16    |
| Rattus norvegicus     | 6     | Rattus rattus           | 0.995   |
| Felis silvestris      | 3     | Felis silvestris        | 0.743   |
| Hypsugo savii         | 2     | Rattus norvegicus       | 0.684   |

# Bioregion 39

156 records of 45 unique species in 2 cells

| Most common species     |       | Most indicative species |         |
|-------------------------|-------|-------------------------|---------|
| Name                    | Count | Name                    | Score   |
| Apodemus semotus        | 20    | Apodemus semotus        | 2.73e+3 |
| Episoriculus fumidus    | 16    | Episoriculus fumidus    | 2.69e+3 |
| Niviventer culturatus   | 15    | Niviventer culturatus   | 2.68e+3 |
| Microtus kikuchii       | 12    | Anourosorex yamashinai  | 2.45e+3 |
| Anourosorex yamashinai  | 12    | Microtus kikuchii       | 2.29e+3 |
| Tamiops maritimus       | 8     | Petaurista alborufus    | 2.15e+3 |
| Eothenomys melanogaster | 8     | Niviventer coxingi      | 1.43e+3 |
| Petaurista alborufus    | 6     | Tamiops maritimus       | 674     |
| Rattus rattus           | 5     | Eothenomys melanogaster | 636     |
|                         |       |                         |         |

http://localhost:3003/ Page 19 of 28

01/12/15 00:41 Infomap Bioregions

Niviventer coxingi 3 Kattus rattus 1.83

## Bioregion 40

125 records of 6 unique species in 4 cells

| Most common species     |       | Most indicative species |       |
|-------------------------|-------|-------------------------|-------|
| Name                    | Count | Name                    | Score |
| Arctocephalus gazella   | 107   | Arctocephalus gazella   | 473   |
| Mirounga leonina        | 10    | Arctocephalus australis | 21.4  |
| Rangifer tarandus       | 4     | Leptonychotes weddellii | 14.5  |
| Hydrurga leptonyx       | 2     | Hydrurga leptonyx       | 6.41  |
| Arctocephalus australis | 1     | Mirounga leonina        | 4.84  |
| Leptonychotes weddellii | 1     | Rangifer tarandus       | 1.58  |

# Bioregion 41

82 records of 17 unique species in 4 cells

| Most common species |       | Most indicative species |         |
|---------------------|-------|-------------------------|---------|
| Name                | Count | Name                    | Score   |
| Callorhinus ursinus | 22    | Sorex pribilofensis     | 2.34e+3 |
| Vulpes lagopus      | 16    | Callorhinus ursinus     | 485     |
| Eumetopias jubatus  | 15    | Eumetopias jubatus      | 164     |
| Sorex pribilofensis | 9     | Vulpes lagopus          | 85.7    |
| Enhydra lutris      | 4     | Odobenus rosmarus       | 12.9    |
| Mus musculus        | 3     | Enhydra lutris          | 7.25    |
| Phoca vitulina      | 2     | Rangifer tarandus       | 3.84    |
| Rangifer tarandus   | 2     | Phoca vitulina          | 2.47    |
| Phocoena phocoena   | 1     | Phocoena phocoena       | 0.674   |
| Odobenus rosmarus   | 1     | Mus musculus            | 0.388   |

## Bioregion 42

| 159 records of 25 unique species in 5 cells |                         |
|---|-------------------------|
| Most common species                         | Most indicative species |
|   |                         |

http://localhost:3003/ Page 20 of 28

| Name                    | Count | Name                    | Score   |
|-------------------------|-------|-------------------------|---------|
| Pteropus tonganus       | 27    | Pteropus ornatus        | 2.12e+3 |
| Miniopterus australis   | 23    | Notopteris macdonaldi   | 2.12e+3 |
| Pteropus ornatus        | 14    | Pteropus samoensis      | 2.12e+3 |
| Emballonura semicaudata | 12    | Pteropus anetianus      | 1.94e+3 |
| Pteropus anetianus      | 11    | Pteropus tonganus       | 1.74e+3 |
| Hipposideros cervinus   | 10    | Emballonura semicaudata | 1.59e+3 |
| Notopteris macdonaldi   | 10    | Miniopterus macrocneme  | 424     |
| Miniopterus macrocneme  | 8     | Hipposideros cervinus   | 164     |
| Pteropus samoensis      | 6     | Miniopterus australis   | 21.1    |
| Rattus exulans          | 5     | Rattus exulans          | 17.8    |

## Bioregion 43

51 records of 30 unique species in 3 cells

| Most common species     |       | Most indicative species |       |
|-------------------------|-------|-------------------------|-------|
| Name                    | Count | Name                    | Score |
| Eubalaena australis     | 12    | Hyperoodon planifrons   | 530   |
| Mirounga leonina        | 5     | Eubalaena australis     | 398   |
| Graomys griseoflavus    | 3     | Pontoporia blainvillei  | 281   |
| Chaetophractus villosus | 2     | Chaetophractus villosus | 129   |
| Eligmodontia typus      | 2     | Otaria flavescens       | 115   |
| Pontoporia blainvillei  | 2     | Eligmodontia typus      | 96.4  |
| Otaria flavescens       | 2     | Graomys griseoflavus    | 75.7  |
| Leopardus geoffroyi     | 1     | Ctenomys haigi          | 63.6  |
| Ctenomys haigi          | 1     | Leopardus geoffroyi     | 38.8  |
| Hyperoodon planifrons   | 1     | Mirounga leonina        | 21.6  |

## Bioregion 44

101 records of 46 unique species in 5 cells

| Most common species |       | Most indicative species |         |
|---------------------|-------|-------------------------|---------|
| Name                | Count | Name                    | Score   |
| Marmota himalayana  | 12    | Pantholops hodgsonii    | 4.77e+3 |

http://localhost:3003/ Page 21 of 28

| Procapra picticaudata | 8 | Vulpes ferrilata      | 3.98e+3 |
|-----------------------|---|-----------------------|---------|
| Ochotona curzoniae    | 8 | Procapra picticaudata | 3.82e+3 |
| Vulpes vulpes         | 6 | Ochotona curzoniae    | 3.47e+3 |
| Pantholops hodgsonii  | 5 | Marmota himalayana    | 3.18e+3 |
| Lepus oiostolus       | 5 | Equus kiang           | 3.18e+3 |
| Vulpes ferrilata      | 5 | Pseudois nayaur       | 2.39e+3 |
| Pseudois nayaur       | 4 | Lepus oiostolus       | 1.99e+3 |
| Equus kiang           | 4 | Capra hircus          | 1.58    |
| Capra hircus          | 2 | Vulpes vulpes         | 0.500   |

# Bioregion 45

88 records of 34 unique species in 3 cells

| Most common species  |       | Most indicative species |         |
|----------------------|-------|-------------------------|---------|
| Name                 | Count | Name                    | Score   |
| Meriones rex         | 10    | Gazella arabica         | 5.73e+3 |
| Gerbillus famulus    | 7     | Gerbillus famulus       | 5.73e+3 |
| Acomys cahirinus     | 6     | Meriones rex            | 5.21e+3 |
| Rhinopoma hardwickii | 6     | Gerbillus cheesmani     | 747     |
| Mus musculus         | 5     | Rhinopoma hardwickii    | 661     |
| Gazella arabica      | 4     | Acomys cahirinus        | 133     |
| Rattus rattus        | 4     | Rhinolophus clivosus    | 115     |
| Rhinolophus clivosus | 3     | Lepus capensis          | 3.52    |
| Lepus capensis       | 3     | Rattus rattus           | 2.92    |
| Gerbillus cheesmani  | 3     | Mus musculus            | 1.42    |

### Bioregion 46

45 records of 7 unique species in 4 cells

| Most common species     |       | Most indicative species |         |
|-------------------------|-------|-------------------------|---------|
| Name                    | Count | Name                    | Score   |
| Leptonychotes weddellii | 27    | Leptonychotes weddellii | 1.55e+3 |
| Lobodon carcinophaga    | 10    | Lobodon carcinophaga    | 544     |
| Avetacanhalisa saralla  | 2     | Avatacanhalisa caralla  | EQ. (   |

http://localhost:3003/ Page 22 of 28

| Arctocephalus gazella      | S | Arctocephalus gazella      | JZ.0 |
|----------------------------|---|----------------------------|------|
| Mirounga leonina           | 2 | Hydrurga leptonyx          | 12.7 |
| Megaptera novaeangliae     | 1 | Balaenoptera acutorostrata | 6.52 |
| Hydrurga leptonyx          | 1 | Megaptera novaeangliae     | 4.66 |
| Balaenoptera acutorostrata | 1 | Mirounga leonina           | 3.83 |

### **Bioregion 47**

33 records of 24 unique species in 3 cells

| Most common species   |       | Most indicative species |         |
|-----------------------|-------|-------------------------|---------|
| Name                  | Count | Name                    | Score   |
| Spermophilus suslicus | 4     | Mustela eversmanni      | 1.43e+4 |
| Sicista subtilis      | 3     | Sicista subtilis        | 7.16e+3 |
| Apodemus agrarius     | 2     | Spermophilus suslicus   | 3.82e+3 |
| Arvicola amphibius    | 2     | Erinaceus roumanicus    | 622     |
| Mustela eversmanni    | 2     | Erinaceus concolor      | 105     |
| Erinaceus roumanicus  | 2     | Apodemus agrarius       | 51.7    |
| Mustela nivalis       | 1     | Arvicola amphibius      | 5.47    |
| Ondatra zibethicus    | 1     | Ondatra zibethicus      | 3.44    |
| Erinaceus concolor    | 1     | Mustela nivalis         | 1.16    |
| Lepus europaeus       | 1     | Lepus europaeus         | 1.00    |

# Bioregion 48

35 records of 12 unique species in 2 cells

| Most common species     |       | Most indicativ        | Most indicative species |  |
|-------------------------|-------|-----------------------|-------------------------|--|
| Name                    | Count | Name                  | Score                   |  |
| Dicrostonyx hudsonius   | 9     | Dicrostonyx hudsonius | 3.37e+3                 |  |
| Microtus pennsylvanicus | 8     | Phenacomys ungava     | 254                     |  |
| Synaptomys borealis     | 4     | Lepus arcticus        | 141                     |  |
| Clethrionomys gapperi   | 3     | Synaptomys borealis   | 71.3                    |  |
| Phenacomys ungava       | 3     | Pusa hispida          | 50.9                    |  |
| Sorex cinereus          | 2     | Clethrionomys gapperi | 14.7                    |  |
| Pusa hispida            | 1     | Vulpes lagopus        | 13.1                    |  |
|                         |       |                       |                         |  |

http://localhost:3003/ Page 23 of 28

| Lepus arcticus | 1 | Microtus pennsylvanicus | 12.6 |
|----------------|---|-------------------------|------|
| Vulpes lagopus | 1 | Myodes gapperi          | 3.35 |
| Myodes gapperi | 1 | Sorex cinereus          | 3.16 |

# Bioregion 49

22 records of 11 unique species in 1 cells

| Most common species      |       | Most indicative species  |         |
|--------------------------|-------|--------------------------|---------|
| Name                     | Count | Name                     | Score   |
| Pteropus niger           | 6     | Pteropus niger           | 9.54e+3 |
| Mormopterus acetabulosus | 6     | Mormopterus acetabulosus | 9.54e+3 |
| Macaca fascicularis      | 2     | Pteropus subniger        | 9.54e+3 |
| Stenella longirostris    | 1     | Trichechus senegalensis  | 3.18e+3 |
| Tenrec ecaudatus         | 1     | Tenrec ecaudatus         | 398     |
| Suncus murinus           | 1     | Stenella longirostris    | 233     |
| Trichechus senegalensis  | 1     | Macaca fascicularis      | 133     |
| Pteropus subniger        | 1     | Taphozous mauritianus    | 91.8    |
| Taphozous mauritianus    | 1     | Suncus murinus           | 34.2    |
| Rattus norvegicus        | 1     | Rattus norvegicus        | 0.836   |

## Bioregion 50

21 records of 7 unique species in 1 cells

| Most common species  |       | Most indicative species |         |
|----------------------|-------|-------------------------|---------|
| Name                 | Count | Name                    | Score   |
| Microtus abbreviatus | 10    | Microtus abbreviatus    | 5.21e+3 |
| Vulpes lagopus       | 6     | Phoca largha            | 102     |
| Phoca vitulina       | 1     | Vulpes lagopus          | 70.7    |
| Odobenus rosmarus    | 1     | Callorhinus ursinus     | 48.5    |
| Phoca largha         | 1     | Ursus maritimus         | 39.8    |
| Ursus maritimus      | 1     | Odobenus rosmarus       | 28.5    |
| Callorhinus ursinus  | 1     | Phoca vitulina          | 2.71    |

http://localhost:3003/ Page 24 of 28

# Bioregion 51

22 records of 18 unique species in 1 cells

| Most common species     |       | Most indicative species |         |
|-------------------------|-------|-------------------------|---------|
| Name                    | Count | Name                    | Score   |
| Hydropotes inermis      | 2     | Lepus sinensis          | 3.58e+3 |
| Crocidura suaveolens    | 2     | Hydropotes inermis      | 248     |
| Mus musculus            | 2     | Pipistrellus javanicus  | 220     |
| Eptesicus serotinus     | 2     | Callosciurus erythraeus | 219     |
| Lepus sinensis          | 1     | Crocidura suaveolens    | 92.5    |
| Rattus rattus           | 1     | Eptesicus serotinus     | 17.9    |
| Pipistrellus javanicus  | 1     | Nyctalus noctula        | 5.97    |
| Callosciurus erythraeus | 1     | Rattus rattus           | 3.65    |
| Meles meles             | 1     | Mus musculus            | 2.84    |
| Nyctalus noctula        | 1     | Meles meles             | 0.607   |

## Bioregion 52

10 records of 6 unique species in 1 cells

| Most common species    |       | Most indicative species |         |
|------------------------|-------|-------------------------|---------|
| Name                   | Count | Name                    | Score   |
| Dasyurus spartacus     | 4     | Dasyurus spartacus      | 1.43e+4 |
| Sminthopsis archeri    | 2     | Sminthopsis archeri     | 4.77e+3 |
| Tachyglossus aculeatus | 1     | Dorcopsis muelleri      | 1.43e+3 |
| Dorcopsis muelleri     | 1     | Sminthopsis virginiae   | 36.6    |
| Sminthopsis virginiae  | 1     | Melomys burtoni         | 8.31    |
| Melomys burtoni        | 1     | Tachyglossus aculeatus  | 0.916   |

### **Bioregion 53**

10 records of 9 unique species in 1 cells

| Most common species |       | Most indicative species |         |
|---------------------|-------|-------------------------|---------|
| Name                | Count | Name                    | Score   |
| Tupaia nicobarica   | 2     | Tupaia nicobarica       | 2.86e+4 |
| Dteronic melanotic  | 1     | Dteronic melanotic      | 2860+1  |

http://localhost:3003/ Page 25 of 28

| r teropus meianotus | - | r teropus meianotus | <b>2.000</b> i = |
|---------------------|---|---------------------|------------------|
| Hipposideros ater   | 1 | Pongo abelii        | 4.77e+3          |
| Capra hircus        | 1 | Tadarida mops       | 4.77e+3          |
| Macaca fascicularis | 1 | Macaca fascicularis | 199              |
| Tadarida mops       | 1 | Panthera tigris     | 166              |
| Panthera tigris     | 1 | Hipposideros ater   | 119              |
| Pongo abelii        | 1 | Capra hircus        | 4.75             |
| Canis lupus         | 1 | Canis lupus         | 1.36             |

## Bioregion 54

7 records of 7 unique species in 1 cells

| Most common species   |       | Most indicative species |         |
|-----------------------|-------|-------------------------|---------|
| Name                  | Count | Name                    | Score   |
| Nyctalus noctula      | 1     | Myomimus personatus     | 5.73e+4 |
| Myomimus personatus   | 1     | Spermophilus citellus   | 1.36e+3 |
| Mus musculus          | 1     | Nyctalus noctula        | 11.9    |
| Mustela nivalis       | 1     | Felis silvestris        | 10.9    |
| Vulpes vulpes         | 1     | Mustela nivalis         | 4.66    |
| Spermophilus citellus | 1     | Mus musculus            | 2.84    |
| Felis silvestris      | 1     | Vulpes vulpes           | 1.00    |

### **Bioregion 55**

6 records of 6 unique species in 1 cells

| Most common species |       | Most indicative species |         |
|---------------------|-------|-------------------------|---------|
| Name                | Count | Name                    | Score   |
| Pentalagus furnessi | 1     | Pentalagus furnessi     | 5.73e+4 |
| Rattus rattus       | 1     | Tokudaia osimensis      | 5.73e+4 |
| Rattus norvegicus   | 1     | Balaenoptera omurai     | 5.73e+4 |
| Apodemus argenteus  | 1     | Apodemus argenteus      | 1.27e+3 |
| Tokudaia osimensis  | 1     | Rattus rattus           | 7.30    |
| Balaenoptera omurai | 1     | Rattus norvegicus       | 5.01    |

http://localhost:3003/ Page 26 of 28

### Bioregion 56

5 records of 4 unique species in 1 cells

| Most common species      |       | Most indicative species  |         |
|--------------------------|-------|--------------------------|---------|
| Name                     | Count | Name                     | Score   |
| Dicrostonyx unalascensis | 2     | Dicrostonyx unalascensis | 1.91e+4 |
| Eumetopias jubatus       | 1     | Eumetopias jubatus       | 120     |
| Phoca vitulina           | 1     | Phoca vitulina           | 13.6    |
| Vulpes vulpes            | 1     | Vulpes vulpes            | 0.500   |

### Bioregion 57

6 records of 6 unique species in 1 cells

| Most common species    |       | Most indicative species |         |
|------------------------|-------|-------------------------|---------|
| Name                   | Count | Name                    | Score   |
| Tursiops truncatus     | 1     | Nyctalus azoreum        | 5.73e+4 |
| Nyctalus azoreum       | 1     | Physeter macrocephalus  | 249     |
| Delphinus delphis      | 1     | Delphinus delphis       | 41.4    |
| Erinaceus europaeus    | 1     | Tursiops truncatus      | 14.3    |
| Felis silvestris       | 1     | Felis silvestris        | 10.9    |
| Physeter macrocephalus | 1     | Erinaceus europaeus     | 1.80    |

### Bioregion 58

5 records of 4 unique species in 1 cells

| Most common species |       | Most indic         | Most indicative species |  |
|---------------------|-------|--------------------|-------------------------|--|
| Name                | Count | Name               | Score                   |  |
| Barytherium grave   | 2     | Barytherium grave  | 2.86e+4                 |  |
| Jaculus jaculus     | 1     | Gazella leptoceros | 2.39e+3                 |  |
| Gazella dorcas      | 1     | Jaculus jaculus    | 92.3                    |  |
| Gazella leptoceros  | 1     | Gazella dorcas     | 21.5                    |  |

### **Bioregion 59**

E manual of E ....io... on oilo in 4 collo

http://localhost:3003/ Page 27 of 28

o records or o unique species in a cens

| Most common species   |       | Most indicative species |         |
|-----------------------|-------|-------------------------|---------|
| Name                  | Count | Name                    | Score   |
| Clethrionomys rutilus | 1     | Pusa sibirica           | 5.73e+4 |
| Mustela sibirica      | 1     | Sorex daphaenodon       | 2.73e+3 |
| Mustela erminea       | 1     | Mustela sibirica        | 1.40e+3 |
| Pusa sibirica         | 1     | Clethrionomys rutilus   | 318     |
| Sorex daphaenodon     | 1     | Mustela erminea         | 5.29    |

| 5 records of 5 unique species in 1 cells |       |                         |         |  |
|--|-------|-------------------------|---------|--|
| Most common species                      |       | Most indicative species |         |  |
| Name                                     | Count | Name                    | Score   |  |
| Rattus norvegicus                        | 1     | Tokudaia muenninki      | 5.73e+4 |  |
| Suncus murinus                           | 1     | Crocidura dsinezumi     | 5.21e+3 |  |
| Rattus rattus                            | 1     | Suncus murinus          | 205     |  |
| Tokudaia muenninki                       | 1     | Rattus rattus           | 7.30    |  |
| Crocidura dsinezumi                      | 1     | Rattus norvegicus       | 5.01    |  |

http://localhost:3003/ Page 28 of 28