Multiple Probabilities of Direction: A Rationale for Prior-odds Adjustment

Margherita Calderan[1], Filippo Gambarota[2], Enrico Toffalini[1], and Gianmarco Altoè[2]

[1]Department of General Psychology, University of Padova

[2]Department of Developmental Psychology and Socialization, University of Padova

Author Note

Margherita Calderan https://orcid.org/0009-0005-5668-5162

Filippo Gambarota https://orcid.org/0000-0002-6666-1747

Enrico Toffalini https://orcid.org/0000-0002-1404-5133

Gianmarco Altoè https://orcid.org/0000-0003-1254-9528

Correspondence concerning this article should be addressed to Margherita Calderan, Department of General Psychology, University of Padova, Via Venezia 8, Padova 35131, Italy, Email: margherita.calderan@unipd.itFilippo Gambarota, Department of Developmental Psychology and Socialization, University of Padova, Via Venezia 8, Padova 35131, Italy, Email: filippo.gambarota@unipd.itEnrico Toffalini, Department of General Psychology, University of Padova, Via Venezia 8, Padova 35131, Italy, Email: enrico.toffalinia@unipd.itGianmarco Altoè, Department of Developmental Psychology and Socialization, University of Padova, Via Venezia 8, Padova 35131, Italy, Email: gianmarco.altoe@unipd.it

Abstract

In psychological research, Bayesian analyses are often summarized using posterior indices and then interpreted with conventional cutoffs across many parameters. The probability of direction (*pd*) is one such index, defined as the posterior probability that a parameter is greater or smaller than a null value. When *pd* is thresholded and used repeatedly, multiplicity becomes consequential but is rarely handled explicitly. Here, we examine the operating characteristics of *pd* under repeated use by leveraging its correspondence with the frequentist *p*-value under common settings. Simulations show that extreme *pd* can arise by chance and that misleading conclusions become more frequent as the number of tests increases. We further express *pd* in terms of posterior odds and propose prior-odds adjustments that provide multiplicity control while preserving Bayesian coherence. To support practical implementation, we provide an R package that applies the proposed prior-odds corrections.

Keywords: probability of direction, multiple testing, prior-odds

Multiple Probabilities of Direction: A Rationale for Prior-odds Adjustment

Over the past two decades, the proportion of articles reporting Bayesian analyses in psychology has increased considerably (Jevremov & Pajić, 2024; Pfadt et al., 2025; Van De Schoot et al., 2017). Unfortunately, this expansion has been accompanied by practices reminiscent of the "null ritual" in frequentist statistics, where widespread use was associated with automatic application rather than principled reasoning (Cohen, 1994; Gigerenzer, 2004). For instance, researchers sometimes treat Bayes factor thresholds (e.g., BF > 3) as dichotomous decision rules or claim that an effect "exists" because the posterior probability of a positive effect exceeds a conventional benchmark such as 95% (Heck et al., 2023; Hoijtink et al., 2016; Kelter, 2023; Kruschke, 2018; Tendeiro et al., 2024; Tendeiro & Kiers, 2019). While such summaries are convenient, their use as default cutoffs encourages all-or-none interpretations. When these binary judgments are applied repeatedly across many parameters, the issue of multiplicity arises.

In the frequentist framework, multiplicity is typically framed as Type I error (false-positive) problem: if each individual test is conducted at level α, the probability of at least one false positive increases with the number of tests. Classical approaches therefore control either the familywise error rate (the probability of at least one Type I error) or the false discovery rate (the expected proportion of false positives among rejected hypotheses) [**REFs**]. However, multiplicity concerns extend beyond false positives. When researchers examine many parameters and highlight the most extreme estimates, selection effects can produce exaggerated magnitudes (Altoè et al., 2020; Gelman & Tuerlinckx, 2000; Zwet & Cator, 2021). Bayesian shrinkage methods address these distortions by pulling extreme estimates toward more plausible values. Such shrinkage can arise through hierarchical models with partial pooling (Berry & Hochberg, 1999; Gelman et al., 2012, 2013; Kruschke, 2011) and/or through informative priors that reduce overestimation in low signal-to-noise settings (Dawid, 1994; Zwet & Gelman, 2022).

Regularization improves estimation, but it does not by itself calibrate repeated threshold-based decisions across many parameters. If the same cutoff is applied to many parameters (e.g., declaring an effect whenever posterior probability exceeds 0.95), the chance of at

least one spurious finding can remain inflated, even if shrinkage reduces it. Bayesian approaches that directly target this issue typically rely on either simultaneous credible statements (Box & Tiao, 1992; Gelman & Tuerlinckx, 2000; Pruzek, 2016; Schnell et al., 2020) or priors on the hypothesis/model space that scale with the number of tested effects (Jeffreys, 1938; Scott & Berger, 2006; Scott & Berger, 2010; Westfall et al., 1997). Nevertheless, psychological researchers often conflate selection effects that distort estimates with the distinct problem of making many simultaneous decisions. As a result, shrinkage is sometimes treated as sufficient for multiplicity control (Chen et al., 2020; Erčulj & Štrumbelj, 2015; Hogeveen et al., 2022; Limbachia et al., 2021; Mathar et al., 2022).

Other justifications invokes the *subjectivist* interpretation of Bayesian inference, which treats posterior probabilities as rational degrees of personal belief (De Finetti, 2017; Edwards et al., 1963; Lindley, 1972; Rozeboom, 1960). However, this interpretation fits poorly with common psychological practice, where researchers typically apply default software priors (often weakly informative) rather than priors encoding substantive domain knowledge (Smid & Winter, 2020; Van De Schoot et al., 2017). When default priors are combined with sufficient data, Bayesian and frequentist procedures often converge to similar conclusions (Albers et al., 2018; Bernardo & Smith, 1994; Gelman et al., 2013). This convergence makes it both natural and useful to evaluate the frequentist operating characteristics of Bayesian decision rules under multiplicity.

Accordingly, we adopt a *falsificationist* perspective, which treats Bayesian procedures as tools to be calibrated rather than as solely expressions of belief (Box, 1980; Gelman & Hennig, 2017; Gelman & Shalizi, 2013; Mayo, 2018; Rubin, 1984). Here, we focus on the Probability of Direction (*pd*), a Bayesian index that can be mapped directly to the frequentist *p*-value in common settings (Berger & Sellke, 1987; Casella & Berger, 1987; Makowski et al., 2019; Marsman & Wagenmakers, 2017). This correspondence allows us to evaluate *pd*'s behavior under repeated use and to compare Bayesian and frequentist approaches to multiplicity control on common ground.

The paper proceeds in four steps. First, we introduce *pd* and make explicit its connection to the frequentist *p*-value, which provides a reference point for calibration. Second, we

characterize *pd*'s null distribution within a three-sided hypothesis testing framework (Goeman et al., 2010) and examine its multiple-testing behavior through simulations that quantify false-positive rates. Third, we relate *pd* to directional Bayes factors and formalize the role of prior odds. Finally, we present prior-odds adjustments for multiplicity following Jeffreys (Jeffreys, 1938; see also Westfall et al., 1997), discuss implementation under dependence, and provide guidance for reporting both raw and multiplicity-adjusted *pd*.

Probability of Direction and *p*-value

The Probability of Direction (*pd*) for continuous distributions can be defined as the maximum between the posterior probabilities that a parameter is greater or smaller than the null value:

$$pd = \max(\Pr(\theta < \theta_{null} \mid y),\ \Pr(\theta > \theta_{null} \mid y)).$$

As proposed and popularized in applied Bayesian workflows, *pd* offers an intuitive summary of directional evidence by quantifying the proportion of posterior that lies on the dominant side (Makowski et al., 2019); it ranges from 0.50 (no directional preference) to 1 (all on one side).

Figure 1

Posterior distribution with shaded directional posterior area. In this example, the probability of direction (pd) is the greater posterior probability on either side of 0.



Because *pd* is a functional of the posterior, it inherits dependence on the full model specification (likelihood and prior), so its long-run error behavior is not a property of *pd* "in

isolation" but of the overall Bayesian procedure that generates the posterior. A convenient starting point for calibration is the mapping between *pd* and the frequentist *p*-value. Under flat (or sufficiently diffuse symmetric) priors in typical hypothesis-testing scenarios (e.g., testing means), *pd* is simply the posterior on the dominant side of θ_{null}, which corresponds to the complement of the one-sided *p*-value (Casella & Berger, 1987; Marsman & Wagenmakers, 2017):

$$pd = \max(Pr(\theta < \theta_{null} \mid y), Pr(\theta > \theta_{null} \mid y)) = 1 - p,$$

with the two-sided correspondence being $p_{two-sided} = 2p = 2(1 - pd)$.

Probability of Direction as a test of existence and direction

The *pd* can be embedded in a three-sided testing framework (Goeman et al., 2010), where thresholding $pd > c$ yields a claim that an effect exists *and* has a particular sign. The three-sided testing framework partitions the parameter space into three mutually exclusive regions:

$H_- : \theta < 0$ (inferiority),

$H_0 : \theta = 0$ (equivalence),

$H_+ : \theta > 0$ (superiority).

Within this partition, *pd* maps naturally onto the two directional regions because it is the posterior probability of the dominant sign, i.e., the larger of $\Pr(\theta > 0 \mid y)$ and $\Pr(\theta < 0 \mid y)$. Thus, $pd > c$ implies either $\Pr(\theta > 0 \mid y) > c$ or $\Pr(\theta < 0 \mid y) > c$. When $pd > c$ supports $\theta > 0$, the posterior probability assigned to nonpositive values is bounded by

$$\Pr(\theta \leq 0 \mid y) < 1 - c$$

(analogously, when $pd > c$ supports $\theta < 0$, $\Pr(\theta \geq 0 \mid y) < 1 - c$). This decision rule does more than compare H_+ versus H_-, it also excludes 0 by ensuring that posterior mass on the null-or-opposite-sign region remains below $1 - c$. Since $\Pr(\theta \leq 0 \mid y) < 1 - c$, zero must lie below the $(1 - c)$-quantile of the posterior distribution. Equivalently, the $(1 - c)$-quantile is strictly positive. As a result, the central credible interval (bounded by the $(1 - c)$ and c quantiles) excludes 0 (see Figure 2).

Figure 2

Posterior distributions illustrating the relationship between the probability of direction (pd) and credible-interval (CI), using $c = 0.975$ and $\theta_{null} = 0$. ***Left:*** *A narrow posterior whose 95% CI excludes 0.* ***Right:*** *A wider posterior whose 95% CI overlaps 0.*



Pure directional testing conditions on the assumption that $\theta \neq 0$ and evaluates only the sign of the effect by comparing H_+ versus H_-; $H_0 : \theta = 0$ is not among the hypotheses under consideration. In contrast, the three-sided framework explicitly includes H_0 and separates decisions about *existence* (excluding zero) from decisions about direction (positive vs negative). Error types follow directly from this distinction. Under the three-sided framework, a Type I error occurs when the true parameter lies in H_0 but H_0 is rejected in favor of H_+ or H_-.

Simulation setup

Our simulations implement the three-sided Type I error definition by generating data under the global null (all $\theta = 0$) and computing $\Pr(pd > c \mid \theta = 0)$. Operationally, this quantifies how often the procedure falsely excludes zero from the credible region, that is, incorrectly concludes that an effect exists (with a particular sign) when the true parameter is zero. Specifically, we conduct a Monte Carlo simulation in which, for each condition, we generated families of size $m \in \{1, 2, 5, 10, 25\}$, with $n \in \{30, 50, 100\}$ observations per dataset, and repeated the procedure 10,000 times. In every replication, data were generated under the global null: each dataset had true mean 0 and unit variance, $y \sim \mathcal{N}(0, 1)$. Each dataset was analyzed using a

conjugate normal model with known variance $\sigma^2 = 1$ and prior $\theta \sim \mathcal{N}(0, \tau^2)$, where $\tau \in \{0.2, 0.5, 1, 1.5, 2\}$ spans skeptical to flat priors. We drew posterior samples of θ from the resulting normal posterior and computed pd as $pd = \max(\Pr(\theta > 0 \mid y), \Pr(\theta < 0 \mid y))$ using these draws. For the frequentist comparison, we conducted two-sided one-sample z-tests.

Prior influence

Under the null hypothesis, two-sided p-values are uniformly distributed on $[0, 1]$, and because pd can be mapped to the p-value, this implies a corresponding null reference behavior for pd on $[0.5, 1]$ in the continuous case. However, because the pd is a posterior probability about the sign of θ, its null distribution is not determined by the likelihood alone: it can shift depending on how strongly the prior concentrates probability near 0 (i.e., how readily the prior "expects" near-null effects). Showing this prior dependence at $m = 1$ makes the target of later multiplicity analyses explicit: even before any multiplicity is introduced, a fixed cutoff such as $pd > 0.975$ does not have a universal "null meaning" across priors and sample sizes (see Figure 3). Put differently, the first simulation establishes the baseline calibration of "extreme pd" under the chosen model. With this baseline in hand, multiplicity can be introduced as a second, distinct mechanism: even if each individual test has only a small chance of producing an extreme pd under H_0, the probability that at least one of $pd_1, \ldots, pd_m$ is extreme increases with m. To make this separation explicit, the simulation extends from $m = 1$ to $m = 20$ under the global null and summarizes familywise behavior by computing the probability of at least one $pd > 0.975$ across the family of tests (Figure 4).

Figure 3

Null reference distributions for p-values and probability of direction. The ***panel A*** *shows the histogram of two-sided one-sample z-test p-values from 10,000 replications under the global null ($\theta = 0$), with binwidth 0.025. The* ***panel B*** *shows the corresponding empirical distributions of the probability of direction (pd) computed from a conjugate normal posterior, stratified by sample size ($n = 30, 50, 100$; rows) and prior scale ($\theta \sim \mathcal{N}(0, \tau^2)$ with $\tau \in \{0.2, 0.5, 1, 1.5, 2\}$; columns). The vertical dashed line marks the commonly used evidential cutoff (pd = 0.975), which corresponds to a two-sided p-value of 0.05 under the standard conversion $p = 2(1 - pd)$.*



Figure 4

Familywise error rates under the global null for p-values and probability of direction. The figure shows the estimated familywise error rate (fwer), defined as the probability of at least one false positive across m tests, as a function of the number of tests (m), based on 10,000 replications under the global null ($\theta = 0$ for all tests). The pink curve corresponds to declaring at least one discovery when at least one two-sided one-sample z-test p-value < 0.05, whereas the blue curve corresponds to declaring at least one discovery when at least one pd > 0.975 under the conjugate normal Bayesian model. Rows vary the per-test sample size ($n = 30, 50, 100$), and columns vary the prior scale ($\theta \sim \mathcal{N}(0, \tau^2)$ with $\tau \in \{0.2, 0.5, 1, 1.5, 2\}$).



…

aggiungo riassunto risultati simulazione

…

Probability of Direction, posterior odds, and prior-odds adjustment

From posterior probabilities to posterior odds

As previously noted, the probability of direction (pd) is defined as the maximum of the two posterior tail probabilities: $pd = \max(\Pr(\theta > \theta_{null} \mid y), \Pr(\theta < \theta_{null} \mid y))$. To give pd an evidential interpretation, consider the directional hypotheses $H_+ : \theta > 0$ and $H_- : \theta < 0$. If the prior on θ is continuous and symmetric about 0 (i.e., $\pi(\theta) = \pi(-\theta)$), then H_+ and H_- have equal prior probability: $\Pr(H_+) = \Pr(H_-) = 0.5$. In this setting, pd is simply the posterior probability of the favored direction. Without loss of generality, suppose the posterior favors H_+, so that $\Pr(H_+ \mid y) = pd$ and $\Pr(H_- \mid y) = 1 - pd$. By standard Bayesian updating, these posterior odds factorize into a Bayes factor and prior odds:

$$\frac{pd}{1 - pd} = \mathrm{BF}_{+/-} \times \frac{\Pr(H_+)}{\Pr(H_-)},$$

where $\mathrm{BF}_{+/-}$ is the directional Bayes factor (the ratio of integrated likelihoods under H_+ and H_-). Under prior symmetry, the prior odds equal 1, so $\frac{pd}{1-pd} = \mathrm{BF}_{+/-}$ (Marsman & Wagenmakers, 2017). Thus, the quantity $\frac{pd}{1-pd}$ computed from posterior draws can be interpreted directly as the evidence favoring one direction over the other.

Multiplicity adjustment via post-hoc prior-odds

When assessing a single directional claim under a symmetric prior centered at the null, prior odds equal 1, encoding prior indifference between positive and negative effects. However, when m hypotheses are assessed simultaneously and the same default prior odds are used for each, the implied prior distribution over the number of non-null directional effects across the family can become unrealistically optimistic as m increases.

A coherent approach to multiplicity control is to decrease the prior odds assigned to any particular directional claim as a function of m, while leaving the likelihood, and hence the Bayes factor component, unchanged. Because the analysis is symmetric in the sign of θ and pd is defined

as the posterior probability of the dominant direction, the same adjustment can be applied regardless of whether the posterior favors $\theta > 0$ or $\theta < 0$. This yields an adjusted posterior-odds expression of the form

$$\text{Posterior odds}_{\text{adjusted}} = \frac{pd}{1 - pd} \times \text{ adjustment factor}(m).$$

One common formalization introduces a "global null" event Q (all null hypotheses are true) and its complement $\neg Q$ (at least one null is false), and assigns equal prior probability $\Pr(Q) = \Pr(\neg Q) = 0.5$ (Jeffreys, 1938; Westfall et al., 1997). Under independence and a common per-hypothesis probability k that a null is false, this implies

$$(1 - k)^m = 0.5 \quad \Rightarrow \quad k = 1 - 0.5^{1/m}.$$

As m increases, k decreases, encoding increasing prior skepticism about any particular deviation from the null hypothesis (Figure 5).

Figure 5

Per-hypothesis prior probability k s a function of the total number of hypotheses m, with the overall prior probability of at least one effect fixed at $q = 0.5$. The per-hypothesis prior is given by $k = 1 - 0.5^{1/m}$.



Although derived from a Bayesian prior-odds argument, the behavior of the resulting procedure under independence closely resembles frequentists multiplicity corrections (Westfall et

al., 1997). In this simulation, we compare this method with the Šidák adjustment ($p_{adj} = 1 - (1 - p)^m$), which gives exact frequentist family-wise error rate control under independence (Figure 6).

Figure 6

Familywise error rates under the global null for adjusted p-values and probability of direction. The figure shows the estimated familywise error rate (fwer), defined as the probability of at least one false positive across m tests, as a function of the number of tests (m), based on 10,000 replications under the global null ($\theta = 0$ for all tests). The red curve corresponds to declaring at least one discovery when at least one two-sided one-sample z-test p-value < 0.05, whereas the blue curve corresponds to declaring at least one discovery when at least one pd > 0.975 under the conjugate normal Bayesian model. Rows vary the per-test sample size ($n = 30, 50, 100$), and columns vary the prior scale ($\theta \sim \mathcal{N}(0, \tau^2)$ with $\tau \in \{0.2, 0.5, 1, 1.5, 2\}$).



…

aggiungo riassunto risultati simulazione

…

Choosing the global null probability

The framework for multiplicity adjustment via adjusted prior odds naturally allows researchers to incorporate substantive prior information based on background knowledge. One approach is to specify a global prior probability that all effects are null, $q = \Pr(\text{all null})$, and translate that belief into the per-hypothesis prior probability k. Under the assumption of independence, each hypothesis is non-null with probability k, such that $\Pr(\text{all null}) = (1-k)^m$. Domain knowledge guides the choice of q: if one expects that no real effects are present, q may be large (e.g., 0.5), whereas studies targeting well-established effects might justify a smaller q (e.g., 0.1). This leads to the general calibration:

$$(1-k)^m = q \quad \Rightarrow \quad k = 1 - q^{1/m}.$$

However, with small m and low q values, this formula can paradoxically inflate rather than reduce posterior probabilities. This occurs when the per-hypothesis prior probability of the null hypothesis ($\Pr(\text{one null}) = q^{1/m}$) falls below 0.5, effectively favoring the alternative hypothesis a priori. Because we conceive this adjustment as a safeguard against inflated false positives, it is crucial to ensure it never acts in the opposite direction. To this end, the procedure should include a safeguard: if $\Pr(\text{one null}) < 0.5$, the original posterior probabilities should remain unmodified.

```
#' Prior-odds adjustment for Probability of Direction (pd)
#'
#' Adjusts a vector of Probability of Direction (pd) values using a global
#' prior probability that all tested hypotheses are null, q. The adjustment
#' converts q into a per-hypothesis prior probability H0 = q^(1/m),
#' where m is the family size, and then reweights each pd accordingly.
#'
```

```r
prior_adj = function(pd, q = 0.5, m = length(pd)) {

  stopifnot(
    "`pd` must be numeric"        = is.numeric(pd),
    "`pd` must be in [0.5, 1]"    = all(pd >= 0.5 & pd <= 1),
    "`q` must be a single number" = length(q) == 1L && is.finite(q),
    "`q` must be in (0, 1)"       = (q > 0) && (q < 1),
    "`m` must be >= 1"            = length(m) == 1L && is.finite(m) && (m >= 1)
  )

  H0 = q^(1 / m)
  H1 = 1 - H0

  if (H0 >= 0.5) {
    pd_adj = (pd * H1) / (pd * H1 + (1 - pd) * H0)
  } else {
    warning("Pr(H0) < 0.5; returning unadjusted pd")
    pd_adj = pd
  }

  return(pd_adj)
}
```

Once q is fixed, the prior odds for each individual hypothesis become a deterministic function of the family size m: larger families imply smaller k, and therefore higher prior probability toward the null for each test.

To illustrate this behavior, we simulated a mixed data-generating process in which the

global null holds with probability $\Pr(\text{all } \theta = 0) = 0.5$, otherwise $\theta = 0.25$. We repeated the previous simulation procedure with $\tau = 2$, $n \in \{30, 50, 100\}$, and analysis priors $q \in \{0.50, 0.40, 0.30, 0.20, 0.10\}$.

Across all values of m, Šidák-adjusted p-value keep the family-wise error rate (FWER; Figure 7) close to the nominal level, but this comes with a power cost (Figure 8) that becomes more pronounced as the family size increases. In contrast, the adjusted *pd* procedure attains higher power for smaller q (i.e., when the prior places less mass on the global null), but this gain is accompanied by inflated FWER.

Notably, setting $q = 0.4$ yields FWER and power trajectories that closely track the Šidák adjustment across all m, suggesting that, under this calibration, the Bayesian adjustment can effectively emulate classical strong FWER control.

Figure 7

Familywise error rate for adjusted p-value and probability of direction as a function of the number of tests (m), and prior $\Pr(\textit{all } \theta = 0)$. *The red curve corresponds to declaring at least one discovery when at least one two-sided one-sample z-test p-value < 0.05, whereas the blue curve corresponds to declaring at least one discovery when at least one pd > 0.975 under the conjugate normal Bayesian model.*



Figure 8

Power for adjusted p-value and probability of direction as a function of the number of tests (m), and prior $\Pr(all\ \theta = 0)$. *The red curve corresponds to declaring at least one discovery when at least one two-sided one-sample z-test p-value < 0.05, whereas the blue curve corresponds to declaring at least one discovery when at least one pd > 0.975 under the conjugate normal Bayesian model.*



…

Discussion

The proposed multiplicity adjustment can incorporate researcher-specific priors that may draw on evidence from previous studies, theoretical constraints, measurement reliability, or structured expert elicitation [**REFs**]. For transparency, researchers should report the raw *pd* for each parameter alongside at least one multiplicity-aware quantity derived from the prior-odds adjustment. When possible, presenting both the conservative benchmark (i.e., pd when $\Pr(\text{all } \theta = 0) = 0.4$) and an informed adjustment helps clarify how sensitive conclusions are to the assumed multiplicity structure.

RICORDA AGGIUNGERE PRIOR PARAM + PRIOR HPS

Because strict independence-based adjustments can be overly conservative when hypotheses are correlated, a practical option is to replace the nominal number of tests m with an effective number m_{eff} that reflects the dependence structure. This can be done by computing a correlation matrix R capturing correlation among tests (e.g., posterior distributions from the same model) and mapping R to m_{eff} via eigenvalue-based methods (Cheverud, 2001; Del Giudice, 2021; Li & Ji, 2005). Importantly, the resulting m_{eff} will generally depend on how R is defined and estimated, so it should be treated as a sensitivity choice rather than a uniquely determined quantity. More generally, researchers can also specify hypothesis-specific values (or "classes" of hypotheses) with different null prior probabilities, for example, distinguishing target tests from exploratory ones (Guo & Heitjan, 2010).

A key advantage of this procedure is that it can be applied to *pd* computed from posterior distributions obtained under different models. In principle, the most coherent approach is to analyze all data relevant to a study within a single joint model that represents dependence explicitly, and to base inference on the resulting joint posterior distribution [**REFs**]. In practice, however, this is not always feasible, and researchers may therefore report *pd* across several analyses within the same paper.

Software implementation

For implementation, it might be best to integrate the adjustment directly into bayestestR, since it already promotes p_direction. We could reach out to its developer to discuss adding this feature instead of creating a separate package.

References

Albers, C. J., Kiers, H. A., & Ravenzwaaij, D. van. (2018). Credible confidence: A pragmatic view on the frequentist vs Bayesian debate. *Collabra: Psychology*, *4*(1).

Altoè, G., Bertoldo, G., Zandonella Callegher, C., Toffalini, E., Calcagnì, A., Finos, L., & Pastore, M. (2020). Enhancing statistical inference in psychological research via prospective and retrospective design analysis. *Frontiers in Psychology*, *Volume 10 - 2019*. https://doi.org/10.3389/fpsyg.2019.02893

Berger, J. O., & Sellke, T. (1987). Testing a point null hypothesis: The irreconcilability of p values and evidence. *Journal of the American Statistical Association*, *82*(397), 112–122.

Bernardo, J. M., & Smith, A. F. M. (1994). *Bayesian Theory* (1st ed.). Wiley. https://doi.org/10.1002/9780470316870

Berry, D. A., & Hochberg, Y. (1999). Bayesian perspectives on multiple comparisons. *Journal of Statistical Planning and Inference*, *82*(1), 215–227. https://doi.org/10.1016/S0378-3758(99)00044-0

Box, G. E. P. (1980). Sampling and bayes' inference in scientific modelling and robustness. *Journal of the Royal Statistical Society. Series A (General)*, *143*(4), 383–430. http://www.jstor.org/stable/2982063

Box, G. E. P., & Tiao, G. C. (1992). *Bayesian inference in statistical analysis* (1st ed.). Wiley. https://doi.org/10.1002/9781118033197

Casella, G., & Berger, R. L. (1987). Reconciling bayesian and frequentist evidence in the one-sided testing problem. *Journal of the American Statistical Association*, *82*(397), 106–111. https://doi.org/10.1080/01621459.1987.10478396

Chen, G., Taylor, P. A., Cox, R. W., & Pessoa, L. (2020). Fighting or embracing multiplicity in

neuroimaging? Neighborhood leverage versus global calibration. *NeuroImage*, *206*, 116320. https://doi.org/10.1016/j.neuroimage.2019.116320

Cheverud, J. M. (2001). A simple correction for multiple comparisons in interval mapping genome scans. *Heredity*, *87*(1), 52–58. https://doi.org/https://doi.org/10.1046/j.1365-2540.2001.00901.x

Cohen, J. (1994). The earth is round (p < .05). *American Psychologist*, *49*(12), 997–1003. https://doi.org/10.1037/0003-066X.49.12.997

Dawid, A. P. (1994). Selection paradoxes of bayesian inference. *Lecture Notes-Monograph Series*, *24*, 211–220.

De Finetti, B. (2017). *Theory of Probability: A Critical Introductory Treatment* (A. Machí & A. Smith, Eds.; 1st ed.). Wiley. https://doi.org/10.1002/9781119286387

Del Giudice, M. (2021). Effective dimensionality: A tutorial. *Multivariate Behavioral Research*, *56*(3), 527–542.

Edwards, W., Lindman, H., & Savage, L. J. (1963). Bayesian statistical inference for psychological research. *Psychological Review*, *70*(3), 193.

Erčulj, F., & Štrumbelj, E. (2015). Basketball shot types and shot success in different levels of competitive basketball. *PLOS ONE*, *10*(6), 1–14. https://doi.org/10.1371/journal.pone.0128885

Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., & Rubin, D. B. (2013). *Bayesian data analysis* (3rd ed.). Chapman; Hall/CRC. https://doi.org/10.1201/b16018

Gelman, A., & Hennig, C. (2017). Beyond subjective and objective in statistics. *Journal of the Royal Statistical Society Series A: Statistics in Society*, *180*(4), 967–1033. https://doi.org/10.1111/rssa.12276

Gelman, A., Hill, J., & Yajima, M. (2012). Why we (usually) don't have to worry about multiple comparisons. *Journal of Research on Educational Effectiveness*, *5*(2), 189–211. https://doi.org/10.1080/19345747.2011.618213

Gelman, A., & Shalizi, C. R. (2013). Philosophy and the practice of bayesian statistics. *British*

Journal of Mathematical and Statistical Psychology, *66*(1), 8–38. https://doi.org/https://doi.org/10.1111/j.2044-8317.2011.02037.x

Gelman, A., & Tuerlinckx, F. (2000). Type s error rates for classical and bayesian single and multiple comparison procedures. *Computational Statistics*, *15*(3), 373–390. https://doi.org/10.1007/s001800000040

Gigerenzer, G. (2004). Mindless statistics. *The Journal of Socio-Economics*, *33*(5), 587–606. https://doi.org/https://doi.org/10.1016/j.socec.2004.09.033

Goeman, J. J., Solari, A., & Stijnen, T. (2010). Three-sided hypothesis testing: Simultaneous testing of superiority, equivalence and inferiority. *Statistics in Medicine*, *29*(20), 2117–2125. https://doi.org/https://doi.org/10.1002/sim.4002

Guo, M., & Heitjan, D. F. (2010). Multiplicity-calibrated bayesian hypothesis tests. *Biostatistics*, *11*(3), 473–483. https://doi.org/10.1093/biostatistics/kxq012

Heck, D. W., Boehm, U., Böing-Messing, F., Bürkner, P.-C., Derks, K., Dienes, Z., Fu, Q., Gu, X., Karimova, D., Kiers, H. A., et al. (2023). A review of applications of the bayes factor in psychological research. *Psychological Methods*, *28*(3), 558.

Hogeveen, J., Mullins, T. S., Romero, J. D., Eversole, E., Rogge-Obando, K., Mayer, A. R., & Costa, V. D. (2022). The neurocomputational bases of explore-exploit decision-making. *Neuron*, *110*(11), 1869–1879.e5. https://doi.org/10.1016/j.neuron.2022.03.014

Hoijtink, H., Kooten, P. van, & Hulsker, K. (2016). Why bayesian psychologists should change the way they use the bayes factor. *Multivariate Behavioral Research*, *51*(1), 2–10.

Jeffreys, H. (1938). Significance tests when several degrees of freedom arise simultaneously. *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences*, *165*(921), 161–198. https://doi.org/10.1098/rspa.1938.0052

Jevremov, T., & Pajić, D. (2024). Bayesian method in psychology: A bibliometric analysis. *Current Psychology*, *43*(10), 8644–8654.

Kelter, R. (2023). How to choose between different bayesian posterior indices for hypothesis testing in practice. *Multivariate Behavioral Research*, *58*(1), 160–188.

Kruschke, J. K. (2011). Bayesian assessment of null values via parameter estimation and model comparison. *Perspectives on Psychological Science*, *6*(3), 299–312. https://doi.org/10.1177/1745691611406925

Kruschke, J. K. (2018). Rejecting or Accepting Parameter Values in Bayesian Estimation. *Advances in Methods and Practices in Psychological Science*, *1*(2), 270–280. https://doi.org/10.1177/2515245918771304

Li, J., & Ji, L. (2005). Adjusting multiple testing in multilocus analyses using the eigenvalues of a correlation matrix. *Heredity*, *95*(3), 221–227.

Limbachia, C., Morrow, K., Khibovska, A., Meyer, C., Padmala, S., & Pessoa, L. (2021). Controllability over stressor decreases responses in key threat-related brain areas. *Communications Biology*, *4*(1), 42. https://doi.org/10.1038/s42003-020-01537-5

Lindley, D. V. (1972). Bayesian statistics a review. In *Bayesian statistics* (pp. 1–74). Society for Industrial; Applied Mathematics. https://doi.org/10.1137/1.9781611970654.ch1

Makowski, D., Ben-Shachar, M. S., Chen, S. H. A., & Lüdecke, D. (2019). Indices of effect existence and significance in the bayesian framework. *Frontiers in Psychology*, *Volume 10 - 2019*. https://doi.org/10.3389/fpsyg.2019.02767

Marsman, M., & Wagenmakers, E.-J. (2017). Three insights from a bayesian interpretation of the one-sided p value. *Educational and Psychological Measurement*, *77*(3), 529–539. https://doi.org/10.1177/0013164416669201

Mathar, D., Erfanian Abdoust, M., Marrenbach, T., Tuzsus, D., & Peters, J. (2022). The catecholamine precursor tyrosine reduces autonomic arousal and decreases decision thresholds in reinforcement learning and temporal discounting. *PLOS Computational Biology*, *18*(12). https://doi.org/10.1371/journal.pcbi.1010785

Mayo, D. G. (2018). *Statistical inference as severe testing: How to get beyond the statistics wars*. Cambridge University Press.

Pfadt, J. M., Bartoš, F., Godmann, H. R., Waaijers, M., Groot, L., Heo, I., et al. (2025, October 30). *A methodological metamorphosis: The rapid rise of bayesian inference and open science*

practices in psychology. OSF Preprints. https://doi.org/10.31234/osf.io/ck3js_v1

Pruzek, R. M. (2016). *An introduction to bayesian inference and its applications* (pp. 305–334). Routledge.

Rozeboom, W. W. (1960). The fallacy of the null-hypothesis significance test. *Psychological Bulletin*, *57*(5), 416–428. https://doi.org/10.1037/h0042040

Rubin, D. B. (1984). Bayesianly justifiable and relevant frequency calculations for the applied statistician. *The Annals of Statistics*, *12*(4), 1151–1172. https://doi.org/10.1214/aos/1176346785

Schnell, P. M., Fiecas, M., & Carlin, B. P. (2020). **credsubs** : Multiplicity-adjusted subset identification. *Journal of Statistical Software*, *94*(7). https://doi.org/10.18637/jss.v094.i07

Scott, J. G., & Berger, J. O. (2006). An exploration of aspects of bayesian multiple testing. *Journal of Statistical Planning and Inference*, *136*(7), 2144–2162.

Scott, J. G., & Berger, J. O. (2010). Bayes and empirical-Bayes multiplicity adjustment in the variable-selection problem. *The Annals of Statistics*, *38*(5), 2587–2619. https://doi.org/10.1214/10-AOS792

Smid, S. C., & Winter, S. D. (2020). Dangers of the defaults: A tutorial on the impact of default priors when using bayesian SEM with small samples. *Frontiers in Psychology*, *Volume 11 - 2020*. https://doi.org/10.3389/fpsyg.2020.611963

Tendeiro, J. N., & Kiers, H. A. (2019). A review of issues about null hypothesis bayesian testing. *Psychological Methods*, *24*(6), 774.

Tendeiro, J. N., Kiers, H. A., Hoekstra, R., Wong, T. K., & Morey, R. D. (2024). Diagnosing the misuse of the bayes factor in applied research. *Advances in Methods and Practices in Psychological Science*, *7*(1), 25152459231213371.

Van De Schoot, R., Winter, S. D., Ryan, O., Zondervan-Zwijnenburg, M., & Depaoli, S. (2017). A systematic review of bayesian articles in psychology: The last 25 years. *Psychological Methods*, *22*(2), 217.

Westfall, P. H., Johnson, W. O., & Utts, J. M. (1997). A bayesian perspective on the bonferroni

adjustment. *Biometrika*, *84*(2), 419–427.

Zwet, E. van, & Cator, E. A. (2021). The significance filter, the winner's curse and the need to shrink. *Statistica Neerlandica*, *75*(4), 437–452. https://doi.org/10.1111/stan.12241

Zwet, E. van, & Gelman, A. (2022). A proposal for informative default priors scaled by the standard error of estimates. *The American Statistician*, *76*(1), 1–9. https://doi.org/10.1080/00031305.2021.1938225