# From Probability of Direction to Multiple Decisions: A Prior-Odds Adjustment

Margherita Calderan[1], Filippo Gambarota[2], Enrico Toffalini[1], and Gianmarco Altoè[2]

[1]Department of General Psychology, University of Padova

[2]Department of Developmental Psychology and Socialization, University of Padova

## Author Note

Margherita Calderan https://orcid.org/0009-0005-5668-5162

Filippo Gambarota https://orcid.org/0000-0002-6666-1747

Enrico Toffalini https://orcid.org/0000-0002-1404-5133

Gianmarco Altoè https://orcid.org/0000-0003-1254-9528

Correspondence concerning this article should be addressed to Margherita Calderan, Department of General Psychology, University of Padova, Via Venezia 8, Padova 35131, Italy, Email: margherita.calderan@unipd.it

**Abstract**

In psychological research, Bayesian analyses are often summarized using posterior indices and then interpreted with conventional cutoffs across many parameters. The probability of direction (*pd*) is one such index, defined as the posterior probability that a parameter is greater or smaller than a null value. When *pd* is thresholded and used repeatedly, multiplicity becomes consequential but is rarely handled explicitly. This paper examines the operating characteristics of *pd*-based decisions under repeated use, leveraging its correspondence with the frequentist *p*-value in common settings. Simulations show that extreme *pd* values can occur by chance and that misleading conclusions become more likely as the number of tests increases. We then express *pd* in terms of posterior odds and propose a principled prior-odds adjustment that induces multiplicity control while preserving the underlying Bayes-factor component. An accompanying R package implements the proposed prior-odds correction to support routine applied use.

*Keywords:* probability of direction, multiple testing, prior-odds

## From Probability of Direction to Multiple Decisions: A Prior-Odds Adjustment

Over the past two decades, the proportion of articles reporting Bayesian analyses in psychology has increased considerably (Jevremov & Pajić, 2024; Pfadt et al., 2025; Van De Schoot et al., 2017). Unfortunately, this expansion has been accompanied by practices reminiscent of the "null ritual" in frequentist statistics, where widespread use was associated with automatic application rather than principled reasoning (Cohen, 1994; Gigerenzer, 2004). For instance, researchers sometimes treat Bayes factor thresholds (e.g., BF > 3) as dichotomous decision rules or claim that an effect "exists" because the posterior probability of a positive effect exceeds a conventional benchmark such as 95% (Heck et al., 2023; Hoijtink et al., 2016; Kelter, 2023; Kruschke, 2018; Tendeiro et al., 2024; Tendeiro & Kiers, 2019). While such summaries are convenient, their use as default cutoffs encourages all-or-none interpretations. When these binary judgments are applied repeatedly across many parameters, the issue of multiplicity arises.

In the frequentist framework, multiplicity is typically framed as Type I error (false-positive) problem: if each individual test is conducted at level $\alpha$, the probability of at least one false positive increases with the number of tests. Classical approaches therefore control either the familywise error rate (the probability of at least one Type I error) or the false discovery rate (the expected proportion of false positives among rejected hypotheses) (Benjamini & Hochberg, 1995; Westfall & Young, 1993). Multiplicity concerns extend beyond false positives: when researchers examine many parameters and highlight the most extreme estimates, selection effects can produce exaggerated magnitudes (Altoè et al., 2020; Gelman & Tuerlinckx, 2000; Zwet & Cator, 2021). Bayesian shrinkage methods address these distortions by pulling extreme estimates toward more plausible values. Such shrinkage can arise through hierarchical models with partial pooling (Berry & Hochberg, 1999; Gelman et al., 2012, 2013; Kruschke, 2011) and/or through informative priors that reduce overestimation in low signal-to-noise settings (Dawid, 1994; Zwet & Gelman, 2022).

Regularization improves estimation, but it does not by itself calibrate repeated threshold-based decisions across many parameters. If the same cutoff is applied to many

parameters (e.g., declaring an effect whenever posterior probability exceeds 0.95), the chance of at least one spurious finding can remain inflated, even if shrinkage reduces it. Bayesian approaches that directly target this issue typically rely on either simultaneous credible statements (Box & Tiao, 1992; Gelman & Tuerlinckx, 2000; Pruzek, 2016; Schnell et al., 2020) or priors on the hypothesis/model space that scale with the number of tested effects (Jeffreys, 1938; Scott & Berger, 2006; Scott & Berger, 2010; Westfall et al., 1997). Nevertheless, psychological researchers often conflate selection effects that distort estimates with the distinct problem of making many simultaneous decisions. As a result, shrinkage is sometimes treated as sufficient for multiplicity control (Chen et al., 2020; Erčulj & Štrumbelj, 2015; Hogeveen et al., 2022; Limbachia et al., 2021; Mathar et al., 2022).

Other justifications invokes the *subjectivist* interpretation of Bayesian inference, which treats posterior probabilities as rational degrees of personal belief (De Finetti, 2017; Edwards et al., 1963; Lindley, 1972; Rozeboom, 1960). However, this interpretation fits poorly with common psychological practice, where researchers typically apply default software priors (often weakly informative) rather than priors encoding substantive domain knowledge (Smid & Winter, 2020; Van De Schoot et al., 2017). When these priors are combined with sufficient data, Bayesian and frequentist procedures often converge to similar conclusions (Albers et al., 2018; Bernardo & Smith, 1994; Gelman et al., 2013). This convergence makes it both natural and useful to evaluate the frequentist operating characteristics of Bayesian decision rules under multiplicity.

Accordingly, we adopt a *falsificationist* perspective, which treats Bayesian procedures as tools to be calibrated rather than as solely expressions of belief (Box, 1980; Gelman & Hennig, 2017; Gelman & Shalizi, 2013; Mayo, 2018; Rubin, 1984). Here, we focus on the Probability of Direction (*pd*), a Bayesian index that can be mapped directly to the frequentist *p*-value in common settings (Berger & Sellke, 1987; Casella & Berger, 1987; Makowski et al., 2019; Marsman & Wagenmakers, 2017). This correspondence provides a common metric for evaluating *pd*-based decision rules under repeated use and for comparing Bayesian and frequentist approaches to multiplicity control.

The paper proceeds in four steps. First, we introduce *pd* and make explicit its connection to the frequentist *p*-value, which we use as a calibration reference. Second, we derive the null behavior of *pd* and study the multiple-testing properties of *pd*-thresholded decisions in simulations, focusing on family-wise error rate. Third, we express *pd* in terms of posterior odds, relate these odds to directional Bayes factors, and clarify the contribution of prior odds. Finally, building on Jeffreys' global-null formulation (Jeffreys, 1938; see also Westfall et al., 1997), we propose prior-odds adjustments for multiplicity, discuss practical implementation (including departures from independence), and provide guidance for reporting both raw and multiplicity-adjusted *pd* values.

**Probability of direction and *p*-value**

The probability of direction (*pd*) for continuous distributions can be defined as the maximum between the posterior probabilities that a parameter is greater or smaller than the null value (Figure 1): $pd = \max(\Pr(\theta < \theta_{null} \mid y),\ \Pr(\theta > \theta_{null} \mid y))$.

**Figure 1**

*Posterior distribution with directional posterior mass on either side of $\theta_{null}$, here defined as $\theta_{null} = 0$, shown in different colors; the probability of direction (pd) is the larger of the two posterior probabilities.*




As proposed and popularized in applied Bayesian workflows [**REFs**], *pd* offers an intuitive summary of directional evidence by quantifying the proportion of posterior that lies on the

dominant side (Makowski et al., 2019); it ranges from 0.50 (no directional preference) to 1 (all on one side). Because *pd* is a functional of the posterior, it inherits dependence on the full model specification (likelihood and prior), so the long-run error behavior of *pd*-based decisions is not a property of *pd* "in isolation" but of the overall Bayesian procedure that generates the posterior.

A convenient starting point for calibration is the mapping between *pd* and the frequentist $p$-value. Under flat (or sufficiently diffuse symmetric) priors in typical hypothesis-testing scenarios (e.g., testing means), *pd* is simply the posterior on the dominant side of $\theta_{null}$, which corresponds to the complement of the one-sided $p$-value (Casella & Berger, 1987; Marsman & Wagenmakers, 2017), with the two-sided correspondence being $p_{two-sided} = 2p = 2(1 - pd)$.

**Probability of direction as a test of existence and direction**

The *pd* can be embedded in a testing framework, where thresholding $pd > c$ yields a claim that an effect exists and has a particular sign. For example, when $pd > c$ supports $\theta > 0$, the posterior probability assigned to nonpositive values is bounded by

$$\Pr(\theta \leq 0 \mid y) < 1 - c,$$

and, analogously, when $pd > c$ supports $\theta < 0$, it follows that $\Pr(\theta \geq 0 \mid y) < 1 - c$.

This rule therefore does more than compare $H_+$ versus $H_-$: it also controls posterior mass in the null-and-opposite-sign region, ensuring that it remains below $1 - c$. In particular, $\Pr(\theta \leq 0 \mid y) < 1 - c$ implies that 0 lies below the $1 - c$-quantile of the posterior distribution (equivalently, the $1 - c$-quantile is strictly positive). Consequently, the central credible interval bounded by the $1 - c$ and $c$ quantiles excludes 0.

Finally, $c$ can be set so that the interval between the $1 - c$ and $c$ posterior quantiles coincides with the usual equal-tailed $1 - \alpha$ central credible interval, whose endpoints are the $\alpha/2$ and $1 - \alpha/2$ posterior quantiles. Equating endpoints gives $1 - c = \alpha/2$ and $c = 1 - \alpha/2$; therefore $pd > c$ can be read as excluding 0 from a two-sided $100(1 - \alpha)\%$ central interval. With $\alpha = 0.05$, this yields $c = 0.975$ (and $1 - c = 0.025$), mirroring the common two-sided 5% convention (Figure 2).

**Figure 2**

*Posterior distributions illustrating the relationship between the probability of direction (pd) and credible-interval (CI), using $c = 0.975$ and $\theta_{null} = 0$.*




Pure directional testing conditions on the assumption that $\theta \neq 0$ and evaluates only the sign of the effect by comparing $H_+$ versus $H_-$; $H_0 : \theta = 0$ is not among the hypotheses under consideration. In contrast, here we explicitly include $H_0$ and separates decisions about existence and direction (excluding zero, e.g. $H_+$) from decisions about direction (including zero, e.g. $H_{-0}$; Goeman et al. (2010)).

Importantly, although we set $H_0 : \theta = \theta_{null} = 0$ throughout the manuscript for simplicity, $\theta_{null}$ can be defined as the smallest effect size of interest (Lakens et al., 2018). The testing framework thus shifts from evaluating whether an effect differs from zero to whether it exceeds a substantively meaningful threshold.

## Simulation

In our simulations we generate data under the global null (all $\theta = 0$) and we compute $\Pr(pd > c \mid \theta = 0)$. Operationally, this quantifies how often the procedure falsely excludes zero from the credible region (bounded by $1 - c$ and $c$), that is, incorrectly concludes that an effect exists (with a particular sign). We conduct a Monte Carlo simulation in which, for each condition, we generated families of size $m \in \{1, 2, 5, 10, 20\}$, with $n \in \{30, 50, 100\}$ observations per dataset, and repeated the procedure 10,000 times. In every replication, data were generated under

the global null: each dataset had true mean 0 and unit variance, $y \sim \mathcal{N}(0, 1)$. Each dataset was analyzed using a conjugate normal model with known variance $\sigma^2 = 1$ and prior $\theta \sim \mathcal{N}(0, \tau^2)$, where $\tau \in \{0.2, 0.5, 1, 1.5, 2\}$ spans skeptical to flat priors. We drew posterior samples of $\theta$ from the resulting normal posterior and computed *pd* as $pd = \max(\Pr(\theta > 0 \mid y), \Pr(\theta < 0 \mid y))$ using these draws. For the frequentist comparison, we conducted two-sided one-sample $z$-tests.

***Prior influence***

Under the null hypothesis, two-sided $p$-values are uniformly distributed on $[0, 1]$, and because *pd* can be mapped to the $p$-value, this implies a corresponding null reference behavior for *pd* on $[0.5, 1]$ in the continuous case. However, because the *pd* is a posterior probability about the sign of $\theta$, its null distribution is not determined by the likelihood alone: it can shift depending on how strongly the prior concentrates probability near 0 (i.e., how readily the prior "expects" near-null effects). Showing this prior dependence at $m = 1$ makes the target of later multiplicity analyses explicit: even before any multiplicity is introduced, a fixed cutoff such as $pd > 0.975$ does not have a universal "null meaning" across priors and sample sizes.

As shown in Figure 3, *pd* exhibits behavior broadly analogous to a $p$-value when samples are large and the prior places relatively little mass near 0. However, increasing prior skepticism (i.e., concentrating more prior mass around 0) reduces the frequency of extreme *pd* values.

**Figure 3**

*Null reference distributions for p-values and probability of direction. Panel A shows the histogram of two-sided one-sample z-test p-values from 10,000 replications under the global null ($\theta = 0$), with binwidth 0.025. Panel B shows the corresponding empirical distributions of the probability of direction (pd) computed from a conjugate normal posterior, stratified by sample size ($n = 30, 50, 100$; rows) and prior scale ($\theta \sim \mathcal{N}(0, \tau^2)$ with $\tau \in \{0.2, 0.5, 1, 1.5, 2\}$; columns). The vertical dashed line marks the commonly used evidential cutoff (pd = 0.975), which corresponds to a two-sided p-value of 0.05 under the standard conversion $p = 2(1 - pd)$.*



***Behavior under multiplicity***

Building on this baseline, multiplicity can be introduced as a second, distinct mechanism: even when each individual test has only a small probability of producing an extreme $pd$ under $H_0$, the probability that at least one among $pd_1, \ldots, pd_m$ is extreme increases with the number of tests $m$. We quantify this familywise error rate (FWER) behavior by estimating the frequency of $pd > 0.975$ for families ranging from $m = 1$ to $m = 20$.

As expected, $pd$-based decisions behaves broadly analogously to their frequentist counterpart when sample sizes are large and the prior places relatively little mass near 0 (Figure 4). By contrast, as the prior becomes more skeptical (i.e., increasingly concentrated around 0), the FWER decreases.

**Figure 4**

*Familywise error rates under the global null for p-values and probability of direction. The figure shows the estimated familywise error rate (FWER), defined as the probability of at least one false positive across m tests, as a function of the number of tests (m), based on 10,000 replications under the global null ($\theta = 0$ for all tests). The pink curve corresponds to declaring at least one discovery when at least one two-sided one-sample z-test p-value < 0.05, whereas the blue curve corresponds to declaring at least one discovery when at least one pd > 0.975 under the conjugate normal Bayesian model. Rows vary the per-test sample size ($n = 30, 50, 100$), and columns vary the prior scale ($\theta \sim \mathcal{N}(0, \tau^2)$ with $\tau \in \{0.2, 0.5, 1, 1.5, 2\}$).*



The simulation results suggest that Type I error could, in principle, be reduced by adopting a more skeptical within-model prior for each parameter. However, using within-model prior skepticism as a general remedy for multiplicity is typically not the most appropriate response, because it conflates two separate goals: (i) encoding substantive prior knowledge about a specific parameter value and (ii) calibrating a decision rule's error rate under repeated testing. For this reason, the approach developed below targets multiplicity at the level where it arises (the competition among hypotheses), by adjusting the prior odds assigned across competing hypotheses while leaving the within-model prior for any single parameter unchanged.

## Probability of direction, posterior odds, and prior-odds adjustment

### *From posterior probabilities to posterior odds*

To connect $pd$ to posterior odds, consider the directional hypotheses $H_+ : \theta > 0$ and $H_- : \theta < 0$. These hypotheses partition the parameter space based on the sign of $\theta$. If the prior for $\theta$ is continuous and symmetric about 0, then the two directions are equally plausible a priori: $\Pr(H_+) = \Pr(H_-) = 1/2$.

Without loss of generality, assume the posterior favors $H_+$, so that $pd = \Pr(H_+ \mid y) = \Pr(\theta > 0 \mid y)$ (if the posterior instead favors $H_-$, we simply relabel the hypotheses), then the complement is given by $1 - pd = \Pr(\theta \leq 0 \mid y)$. The posterior odds comparing these two hypothesis are therefore:

$$\frac{pd}{1 - pd} = \frac{\Pr(H_+ \mid y)}{\Pr(H_{-0} \mid y)}.$$

For a continuous posterior distribution, the probability of any single point (e.g., $\Pr(\theta = 0 \mid y)$) is 0, which means that one could write $\frac{pd}{1-pd} = \frac{\Pr(H_+|y)}{\Pr(H_-|y)}$. However, we retain the notation $H_{-0}$ in the denominator to explicitly indicate that the hypothesis includes the point null value of zero.

Applying Bayes' rule, these posterior odds can be decomposed into a Bayes factor and prior odds:

$$\frac{\Pr(H_+ \mid y)}{\Pr(H_{-0} \mid y)} = \mathrm{BF}_{+/-0} \times \frac{\Pr(H_+)}{\Pr(H_{-0})},$$

where BF+/−0 quantifies the relative likelihood of the data under $H+$ versus $H_{-0}$. Under prior

symmetry, $\frac{\Pr(H_+)}{\Pr(H_{-0})} \approx 1$, which yields $\frac{pd}{1-pd} = \mathrm{BF}_{+/-0}$ (Casella & Berger, 1987; Marsman & Wagenmakers, 2017). Thus, the odds $\frac{pd}{(1-pd)}$ computed from posterior draws can be interpreted directly as evidence favoring $\theta > 0$ over $\theta \leq 0$ (or $\theta < 0$ over $\theta \geq 0$).

***Multiplicity adjustment via post-hoc prior-odds***

When assessing a single directional claim under a symmetric prior centered at the null, $\frac{\Pr(H_+)}{\Pr(H_{-0})} = 1$ encoding prior indifference. However, when $m$ hypotheses are tested simultaneously, assigning prior probability $\frac{1}{2}$ to each directional claim implies a joint prior that rapidly concentrates probability on the event that at least one effect departs from 0. For instance, with $m = 10$ the implied prior probability that all effects are not in the favored direction (i.e., in the opposite direction with zero included) is $0.5^{10} \approx 0.001$.

In this scenario, a principled way to address multiplicity is to decrease the prior odds assigned to any particular directional claim as a function of $m$, while leaving the the Bayes factor component unchanged. Because the analysis is symmetric in the sign of $\theta$ and $pd$ is defined as the posterior probability of the dominant direction, the same adjustment can be applied regardless of whether the posterior favors $\theta > 0$ or $\theta < 0$.

Concretely, let $\Pr(H_{1_m})$ and $\Pr(H_{0_m})$ denote the multiplicity-adjusted prior probabilities assigned to the favored direction and its complement (i.e., opposite direction with zero included), respectively (with $\Pr(H_{0_m}) + \Pr(H_{1_m}) = 1$). The adjusted posterior odds are then:

$$\frac{pd_{\text{adj}}}{1 - pd_{\text{adj}}} = \frac{pd}{1 - pd} \times \frac{\Pr(H_{1_m})}{\Pr(H_{0_m})},$$

so that, equivalently,

$$pd_{\text{adj}} = \frac{pd \cdot \Pr(H_{1_m})}{pd \cdot \Pr(H_{1_m}) + (1 - pd) \cdot \Pr(H_{0_m})}.$$

Thus, the multiplicity adjustment acts through the prior-odds factor $\Pr(H_{1_m})/\Pr(H_{0_m})$.

***Choosing the prior-odds***

Following the global-null formulation of Jeffreys (1938; see also Westfall et al., 1997), we introduce a global null event $Q$ and its complement $\neg Q$, and assign equal prior probability

$\Pr(Q) = \Pr(\neg Q) = 1/2$. Define, for each $j = 1, \ldots, m$, the "favored-direction" alternative $H_{1,j}$ as the directional hypothesis corresponding to the posterior-dominant sign (i.e., the direction used in $pd_j = \Pr(H_{1,j} \mid y)$), and let $H_{0,j}$ denote its complement (opposite direction with 0 included). We then set $Q = \bigcap_{j=1}^{m} H_{0,j}$ (no parameter lies in its favored direction) and $\neg Q$ as its complement (at least one parameter lies in its favored direction). Under independence and a common per-hypothesis prior probability $k = \Pr(H_{1,j})$ for all $j$, this implies $\Pr(Q) = (1-k)^m$, so $(1-k)^m = 0.5$ and therefore $k = 1 - 0.5^{1/m}$. As $m$ increases, the per-hypothesis prior probability $k = \Pr(H_{1,j})$ decreases, encoding increasing prior skepticism (Figure 5).

**Figure 5**

*Per-hypothesis prior probability k s a function of the total number of hypotheses m, with the overall prior probability of at least one effect fixed at* $q = 0.5$. *The per-hypothesis prior* $\Pr(H_{1,j})$ *is given by* $k = 1 - 0.5^{1/m}$.




In this simulation, we compare this method with the Šidák $p$-value adjustment ($p_{adj} = 1 - (1-p)^m$), which gives exact frequentist family-wise error rate control under independence. While grounded in Bayesian prior-odds logic, the procedure exhibits behavior similar to frequentist multiplicity correction, though it tends to be more conservative even when $n$ and $\tau$ are large (Figure 6).

**Figure 6**

*Familywise error rates under the global null for adjusted p-values and probability of direction. The figure shows the estimated familywise error rate (FWER), defined as the probability of at least one false positive across m tests, as a function of the number of tests (m), based on 10,000 replications under the global null ($\theta = 0$ for all tests). The red curve corresponds to declaring at least one discovery when at least one two-sided one-sample z-test p-value < 0.05, whereas the blue curve corresponds to declaring at least one discovery when at least one pd > 0.975 under the conjugate normal Bayesian model. Rows vary the per-test sample size ($n = 30, 50, 100$), and columns vary the prior scale ($\theta \sim \mathcal{N}(0, \tau^2)$ with $\tau \in \{0.2, 0.5, 1, 1.5, 2\}$).*



Having linked the multiplicity adjustment to the global null event $Q$, the same construction yields a more general calibration directly in terms of $\Pr(Q)$. Specifically, instead of fixing $\Pr(Q) = 0.5$, researchers may encode background knowledge by choosing $q = \Pr(Q)$, the prior probability that all $m$ hypotheses are null (i.e., opposite direction with 0 included). Under a priori independence and a common per-hypothesis prior probability $k = \Pr(H_{1,j})$ for the favored-direction alternative, $\Pr(Q) = (1-k)^m$, so choosing $q$ implies

$$(1-k)^m = q \quad \Rightarrow \quad k = 1 - q^{1/m}.$$

Prior knowledge guides $q$: when few (if any) non-null effects are expected, $q$ should be large; when effects are well established, $q$ can be smaller. For small $m$ and sufficiently small $q$, however, the calibration can become anti-conservative: $\Pr(H_{0,j}) = 1 - k = q^{1/m}$ may fall below $1/2$, which implies $k > 1/2$ and therefore increases rather than decreases the odds for $H_{1,j}$. Because the goal of the adjustment is to guard against inflated false positives, it should never move posterior probabilities in this direction; accordingly, a simple safeguard is to leave posterior probabilities unchanged whenever $\Pr(H_{0,j}) < 1/2$ (equivalently, whenever $q^{1/m} < 1/2$).

Once $q$ is fixed, the prior odds assigned to each individual hypothesis become a deterministic function of the family size $m$: larger families imply smaller $k$, and consequently a higher prior probability on the null for each test. To illustrate this dependence, we consider two data-generating regimes: a global-null scenario in which $\theta_1 = \cdots = \theta_m = 0$, and an alternative scenario in which all but one effect are null ($\theta_1 = \cdots = \theta_{m-1} = 0$, $\theta_m = 0.2$). We then repeat the previous simulation procedure with $\tau = 2$ and $n = 50$, i.e. a setting in which *pd* and *p*-value behavior is broadly comparable, while varying only the number of tests $m \in \{1, 2, 3, 4, 5, 10\}$ and the prior parameter $q \in \{0.50, 0.40, 0.30, 0.20, 0.10\}$.

**Figure 7**

*Familywise error rate (Panel A), power (Panel B), and difference between method (Panel C) for adjusted* p*-values and probability of direction as a function of the number of tests ($m$) and prior* Pr(*all* $\theta = 0$). *The Šidák-adjusted* p*-value curve (pink) declares at least one discovery when any two-sided one-sample* z*-test satisfies* $p < 0.05$*, whereas the adjusted* pd *curve (blue) declares at least one discovery when any* $pd > 0.975$ *under the conjugate normal Bayesian model. Panel C shows the difference (adjusted pd minus Šidák-adjusted p-value) for FWER (under the global null) and power (under the global alternative).*



Across all values of $m$, Šidák-adjusted $p$-values maintain the family-wise error rate (FWER; Figure 7, Panel A) near the nominal level, but this control comes at the expense of diminished power (Figure 7, Panel B). By contrast, the adjusted *pd* procedure exhibits FWER and power trajectories that depend on the prior parameter $q$. For smaller values of $q$ ($q \in \{0.30, 0.20, 0.10\}$), the tradeoff is favorable: the increase in power is larger than the corresponding increase in FWER, implying gains relative to the loss of error control (Figure 7, Panel C). Notably, $q = 0.4$ yields FWER and power curves that closely mirror the Šidák adjustment across the full range of $m$, suggesting that this calibration allows the Bayesian adjustment to approximate classical strong FWER control (Figure 7).

## Application

## Discussion

Although *pd* is a descriptive posterior index, thresholding it into a decision rule and applying that rule repeatedly across many parameters induces a multiplicity problem, as the probability of at least one false directional claim increases with the number of decisions. The multiplicity adjustment proposed here targets this decision-level error while allowing researcher-specific knowledge to enter transparently through the prior-odds specification.

An advantage of this procedure is its applicability to *pd* values computed from posterior distributions obtained under different models. While the most coherent approach analyzes all relevant data within a single joint model that explicitly represents dependence (Gelman et al., 2012, 2013; Kruschke, 2011), this is not always feasible in practice, and researchers often report *pd* across multiple analyses within the same manuscript [**REFs**]. The procedure outlined here is designed for exactly these scenarios and also generalizes straightforwardly to non-zero null values by defining directional hypotheses relative to a smallest effect size of interest (Lakens et al., 2018).

The procedure does assume independence between hypotheses when allocating global null probability across $m$ tests, treating each as providing independent evidence. When hypotheses are correlated through shared parameters, overlapping data structures, or measurement dependencies, independence-based adjustments overestimate the multiplicity penalty, yielding overly

conservative estimates (Pesarin & Salmaso, 2010; Westfall & Young, 1993). A practical solution is to replace the nominal number of tests $m$ with an effective number $m_{\text{eff}}$ that reflects the dependence structure. This can be achieved by computing a correlation matrix $R$ that captures correlation among tests (e.g., posterior distributions from the same model) and mapping $R$ to $m_{\text{eff}}$ via eigenvalue-based methods (Cheverud, 2001; Del Giudice, 2021; Li & Ji, 2005). The resulting $m_{\text{eff}}$ will generally depend on how $R$ is defined and estimated, so it should be treated as a sensitivity parameter rather than a uniquely determined quantity.

Another important caveat is that this framework does not, by itself, deliver evidence for a point null (in the sense of assigning posterior support to $\theta = 0$); instead, it evaluates directional claims defined on inequalities (e.g., $\theta > \theta_{null}$ vs. $\theta \leq \theta_{null}$), where the complement includes the null value by construction. This contrasts with other Bayes factor approaches that represent the point null as a distinct model component by assigning it positive prior mass, thereby allowing direct statements of evidence in favor of the point-null hypothesis (Heck et al., 2023; Ly et al., 2016; Morey & Rouder, 2011).

For applied reporting, a two-tier strategy is recommended: (i) report the unadjusted *pd* for each parameter to preserve transparency, and (ii) report at least one multiplicity-adjusted measure based on the proposed prior-odds correction. When *pd* values are computed from posterior distributions obtained from different models, a conservative default is to report the adjustment induced by $\Pr(Q) = 0.4$, a skeptical but defensible choice that yields corrections analogous to the Šidák adjustment for $p$-values. When substantive knowledge motivates alternative priors, report both the conservative and the informed adjustment to demonstrate robustness to plausible prior assumptions.

Despite its limitations, our method provides a coherent, practical way to recalibrate *pd*-based decisions for multiple comparisons, making reported directional claims easier to interpret and less likely to reflect chance patterns across many parameters.

## References

Albers, C. J., Kiers, H. A., & Ravenzwaaij, D. van. (2018). Credible confidence: A pragmatic view on the frequentist vs Bayesian debate. *Collabra: Psychology*, *4*(1).

Altoè, G., Bertoldo, G., Zandonella Callegher, C., Toffalini, E., Calcagnì, A., Finos, L., & Pastore, M. (2020). Enhancing statistical inference in psychological research via prospective and retrospective design analysis. *Frontiers in Psychology*, *Volume 10 - 2019*. https://doi.org/10.3389/fpsyg.2019.02893

Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodological)*, *57*(1), 289–300. https://doi.org/10.2307/2346101

Berger, J. O., & Sellke, T. (1987). Testing a point null hypothesis: The irreconcilability of p values and evidence. *Journal of the American Statistical Association*, *82*(397), 112–122.

Bernardo, J. M., & Smith, A. F. M. (1994). *Bayesian Theory* (1st ed.). Wiley. https://doi.org/10.1002/9780470316870

Berry, D. A., & Hochberg, Y. (1999). Bayesian perspectives on multiple comparisons. *Journal of Statistical Planning and Inference*, *82*(1), 215–227. https://doi.org/10.1016/S0378-3758(99)00044-0

Box, G. E. P. (1980). Sampling and bayes' inference in scientific modelling and robustness. *Journal of the Royal Statistical Society. Series A (General)*, *143*(4), 383–430. http://www.jstor.org/stable/2982063

Box, G. E. P., & Tiao, G. C. (1992). *Bayesian inference in statistical analysis* (1st ed.). Wiley. https://doi.org/10.1002/9781118033197

Casella, G., & Berger, R. L. (1987). Reconciling bayesian and frequentist evidence in the one-sided testing problem. *Journal of the American Statistical Association*, *82*(397), 106–111. https://doi.org/10.1080/01621459.1987.10478396

Chen, G., Taylor, P. A., Cox, R. W., & Pessoa, L. (2020). Fighting or embracing multiplicity in neuroimaging? Neighborhood leverage versus global calibration. *NeuroImage*, *206*, 116320.

https://doi.org/10.1016/j.neuroimage.2019.116320

Cheverud, J. M. (2001). A simple correction for multiple comparisons in interval mapping genome scans. *Heredity*, *87*(1), 52–58. https://doi.org/https://doi.org/10.1046/j.1365-2540.2001.00901.x

Cohen, J. (1994). The earth is round (p < .05). *American Psychologist*, *49*(12), 997–1003. https://doi.org/10.1037/0003-066X.49.12.997

Dawid, A. P. (1994). Selection paradoxes of bayesian inference. *Lecture Notes-Monograph Series*, *24*, 211–220.

De Finetti, B. (2017). *Theory of Probability: A Critical Introductory Treatment* (A. Machí & A. Smith, Eds.; 1st ed.). Wiley. https://doi.org/10.1002/9781119286387

Del Giudice, M. (2021). Effective dimensionality: A tutorial. *Multivariate Behavioral Research*, *56*(3), 527–542.

Edwards, W., Lindman, H., & Savage, L. J. (1963). Bayesian statistical inference for psychological research. *Psychological Review*, *70*(3), 193.

Erčulj, F., & Štrumbelj, E. (2015). Basketball shot types and shot success in different levels of competitive basketball. *PLOS ONE*, *10*(6), 1–14. https://doi.org/10.1371/journal.pone.0128885

Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., & Rubin, D. B. (2013). *Bayesian data analysis* (3rd ed.). Chapman; Hall/CRC. https://doi.org/10.1201/b16018

Gelman, A., & Hennig, C. (2017). Beyond subjective and objective in statistics. *Journal of the Royal Statistical Society Series A: Statistics in Society*, *180*(4), 967–1033. https://doi.org/10.1111/rssa.12276

Gelman, A., Hill, J., & Yajima, M. (2012). Why we (usually) don't have to worry about multiple comparisons. *Journal of Research on Educational Effectiveness*, *5*(2), 189–211. https://doi.org/10.1080/19345747.2011.618213

Gelman, A., & Shalizi, C. R. (2013). Philosophy and the practice of bayesian statistics. *British Journal of Mathematical and Statistical Psychology*, *66*(1), 8–38.

https://doi.org/https://doi.org/10.1111/j.2044-8317.2011.02037.x

Gelman, A., & Tuerlinckx, F. (2000). Type s error rates for classical and bayesian single and multiple comparison procedures. *Computational Statistics*, *15*(3), 373–390. https://doi.org/10.1007/s001800000040

Gigerenzer, G. (2004). Mindless statistics. *The Journal of Socio-Economics*, *33*(5), 587–606. https://doi.org/https://doi.org/10.1016/j.socec.2004.09.033

Goeman, J. J., Solari, A., & Stijnen, T. (2010). Three-sided hypothesis testing: Simultaneous testing of superiority, equivalence and inferiority. *Statistics in Medicine*, *29*(20), 2117–2125. https://doi.org/https://doi.org/10.1002/sim.4002

Heck, D. W., Boehm, U., Böing-Messing, F., Bürkner, P.-C., Derks, K., Dienes, Z., Fu, Q., Gu, X., Karimova, D., Kiers, H. A. L., Klugkist, I., Kuiper, R. M., Lee, M. D., Leenders, R., Leplaa, H. J., Linde, M., Ly, A., Meijerink-Bosman, M., Moerbeek, M., … Hoijtink, H. (2023). A review of applications of the Bayes factor in psychological research. *Psychological Methods*, *28*(3), 558–579. https://doi.org/10.1037/met0000454

Hogeveen, J., Mullins, T. S., Romero, J. D., Eversole, E., Rogge-Obando, K., Mayer, A. R., & Costa, V. D. (2022). The neurocomputational bases of explore-exploit decision-making. *Neuron*, *110*(11), 1869–1879.e5. https://doi.org/10.1016/j.neuron.2022.03.014

Hoijtink, H., Kooten, P. van, & Hulsker, K. (2016). Why bayesian psychologists should change the way they use the bayes factor. *Multivariate Behavioral Research*, *51*(1), 2–10.

Jeffreys, H. (1938). Significance tests when several degrees of freedom arise simultaneously. *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences*, *165*(921), 161–198. https://doi.org/10.1098/rspa.1938.0052

Jevremov, T., & Pajić, D. (2024). Bayesian method in psychology: A bibliometric analysis. *Current Psychology*, *43*(10), 8644–8654.

Kelter, R. (2023). How to choose between different bayesian posterior indices for hypothesis testing in practice. *Multivariate Behavioral Research*, *58*(1), 160–188.

Kruschke, J. K. (2011). Bayesian assessment of null values via parameter estimation and model

comparison. *Perspectives on Psychological Science*, *6*(3), 299–312. https://doi.org/10.1177/1745691611406925

Kruschke, J. K. (2018). Rejecting or Accepting Parameter Values in Bayesian Estimation. *Advances in Methods and Practices in Psychological Science*, *1*(2), 270–280. https://doi.org/10.1177/2515245918771304

Lakens, D., Scheel, A. M., & Isager, P. M. (2018). Equivalence testing for psychological research: A tutorial. *Advances in Methods and Practices in Psychological Science*, *1*(2), 259–269. https://doi.org/10.1177/2515245918770963

Li, J., & Ji, L. (2005). Adjusting multiple testing in multilocus analyses using the eigenvalues of a correlation matrix. *Heredity*, *95*(3), 221–227.

Limbachia, C., Morrow, K., Khibovska, A., Meyer, C., Padmala, S., & Pessoa, L. (2021). Controllability over stressor decreases responses in key threat-related brain areas. *Communications Biology*, *4*(1), 42. https://doi.org/10.1038/s42003-020-01537-5

Lindley, D. V. (1972). Bayesian statistics a review. In *Bayesian statistics* (pp. 1–74). Society for Industrial; Applied Mathematics. https://doi.org/10.1137/1.9781611970654.ch1

Ly, A., Verhagen, J., & Wagenmakers, E.-J. (2016). Harold jeffreys's default bayes factor hypothesis tests: Explanation, extension, and application in psychology. *Journal of Mathematical Psychology*, *72*, 19–32. https://doi.org/https://doi.org/10.1016/j.jmp.2015.06.004

Makowski, D., Ben-Shachar, M. S., Chen, S. H. A., & Lüdecke, D. (2019). Indices of effect existence and significance in the bayesian framework. *Frontiers in Psychology*, *Volume 10 - 2019*. https://doi.org/10.3389/fpsyg.2019.02767

Marsman, M., & Wagenmakers, E.-J. (2017). Three insights from a bayesian interpretation of the one-sided p value. *Educational and Psychological Measurement*, *77*(3), 529–539. https://doi.org/10.1177/0013164416669201

Mathar, D., Erfanian Abdoust, M., Marrenbach, T., Tuzsus, D., & Peters, J. (2022). The catecholamine precursor tyrosine reduces autonomic arousal and decreases decision

thresholds in reinforcement learning and temporal discounting. *PLOS Computational Biology*, *18*(12). https://doi.org/10.1371/journal.pcbi.1010785

Mayo, D. G. (2018). *Statistical inference as severe testing: How to get beyond the statistics wars*. Cambridge University Press.

Morey, R. D., & Rouder, J. N. (2011). Bayes factor approaches for testing interval null hypotheses. *Psychological Methods*, *16*(4), 406–419. https://doi.org/10.1037/a0024377

Pesarin, F., & Salmaso, L. (2010). The permutation testing approach: A review. *Statistica*, *70*(4), 481–509. https://doi.org/10.6092/issn.1973-2201/3599

Pfadt, J. M., Bartoš, F., Godmann, H. R., Waaijers, M., Groot, L., Heo, I., et al. (2025, October 30). *A methodological metamorphosis: The rapid rise of bayesian inference and open science practices in psychology*. OSF Preprints. https://doi.org/10.31234/osf.io/ck3js_v1

Pruzek, R. M. (2016). *An introduction to bayesian inference and its applications* (pp. 305–334). Routledge.

Rozeboom, W. W. (1960). The fallacy of the null-hypothesis significance test. *Psychological Bulletin*, *57*(5), 416–428. https://doi.org/10.1037/h0042040

Rubin, D. B. (1984). Bayesianly justifiable and relevant frequency calculations for the applied statistician. *The Annals of Statistics*, *12*(4), 1151–1172. https://doi.org/10.1214/aos/1176346785

Schnell, P. M., Fiecas, M., & Carlin, B. P. (2020). **credsubs** : Multiplicity-adjusted subset identification. *Journal of Statistical Software*, *94*(7). https://doi.org/10.18637/jss.v094.i07

Scott, J. G., & Berger, J. O. (2006). An exploration of aspects of bayesian multiple testing. *Journal of Statistical Planning and Inference*, *136*(7), 2144–2162.

Scott, J. G., & Berger, J. O. (2010). Bayes and empirical-Bayes multiplicity adjustment in the variable-selection problem. *The Annals of Statistics*, *38*(5), 2587–2619. https://doi.org/10.1214/10-AOS792

Smid, S. C., & Winter, S. D. (2020). Dangers of the defaults: A tutorial on the impact of default priors when using bayesian SEM with small samples. *Frontiers in Psychology*, *Volume 11 -*

*2020*. https://doi.org/10.3389/fpsyg.2020.611963

Tendeiro, J. N., & Kiers, H. A. (2019). A review of issues about null hypothesis bayesian testing. *Psychological Methods*, *24*(6), 774.

Tendeiro, J. N., Kiers, H. A., Hoekstra, R., Wong, T. K., & Morey, R. D. (2024). Diagnosing the misuse of the bayes factor in applied research. *Advances in Methods and Practices in Psychological Science*, *7*(1), 25152459231213371.

Van De Schoot, R., Winter, S. D., Ryan, O., Zondervan-Zwijnenburg, M., & Depaoli, S. (2017). A systematic review of bayesian articles in psychology: The last 25 years. *Psychological Methods*, *22*(2), 217.

Westfall, P. H., Johnson, W. O., & Utts, J. M. (1997). A bayesian perspective on the bonferroni adjustment. *Biometrika*, *84*(2), 419–427.

Westfall, P. H., & Young, S. S. (1993). *Resampling-based multiple testing: Examples and methods for p-value adjustment*. Wiley.

Zwet, E. van, & Cator, E. A. (2021). The significance filter, the winner's curse and the need to shrink. *Statistica Neerlandica*, *75*(4), 437–452. https://doi.org/10.1111/stan.12241

Zwet, E. van, & Gelman, A. (2022). A proposal for informative default priors scaled by the standard error of estimates. *The American Statistician*, *76*(1), 1–9. https://doi.org/10.1080/00031305.2021.1938225