


Curvature and Torsion in the Continuum Theory of Lattice Defects

Marvin Henke, Dr. Nikodem Szpak
Faculty of Physics, University Duisburg-Essen, 47048 Duisburg

UNIVERSITÄT
DUISBURG
ESSEN

Offen im Denken

Continuum Theory - Graphene



Tight-binding Hamiltonian

$$H = \sum_{\vec{r}} \sum_{j=1}^3 (a_{\vec{r}}^{\dagger} b_{\vec{r}+\vec{d}_j} + b_{\vec{r}+\vec{d}_j}^{\dagger} a_{\vec{r}})$$

Fourier transform

$$i\partial_t \vec{\psi}(\vec{k}) = \mathbf{h}(\vec{k}) \vec{\psi}(\vec{k})$$
$$\mathbf{h}(\vec{k}) = \begin{pmatrix} 0 & f(\vec{k}) \\ f^*(\vec{k}) & 0 \end{pmatrix}, \quad f(\vec{k}) = \sum_{j=1}^3 e^{i\vec{k} \cdot \vec{d}_j}$$


Linear dispersion around Dirac points

$$\vec{k} = \vec{K}^{(s)} + \delta\vec{k} \implies \mathbf{h}(\vec{k}) \approx -\frac{3}{2} \delta\vec{k} \cdot \vec{\sigma}$$

Continuum limit

$$i\partial_t \vec{\phi}(\vec{r}) = i c \vec{\sigma}^i \partial_i \vec{\phi}(\vec{r}), \quad \rho = \vec{\phi}^{\dagger} \vec{\phi}, \quad \vec{j} = -c \vec{\phi}^{\dagger} \vec{\sigma} \vec{\phi}$$
$$\Updownarrow$$
$$i\gamma^{\mu} \partial_{\mu} \phi = 0, \quad \gamma^{\mu} = \sigma^{\mu} \sigma^3, \quad \mu \in \{0, 1, 2\}$$

Burgers vector in 5-7 defects



One 5-7 defect causes a lattice dislocation which can be described by the Burgers vector $\vec{b} = \sqrt{3} \vec{u}_x$


TODO:
In depth visual explanation of torsion and how it is analogous to the burgers vector

Riemannian Geometry

Riemannian Manifold $\mathcal{M}$ equipped with metric $g_{\mu\nu}$ and connection $\Gamma^{\lambda}_{\mu\nu}$.


Metric

Defines distances and angles.



Connection

Connects nearby **tangent spaces**, which allows the definition of a derivative on **vector fields** which take on values in the tangent space.



The unique torsion-free, metric compatible connection is called the Levi-Civita connection:

$$\{\lambda_{\mu\nu}\} = \frac{1}{2} g^{\lambda\rho} (\partial_{\nu} g_{\rho\mu} + \partial_{\mu} g_{\rho\nu} - \partial_{\rho} g_{\mu\nu})$$

Covariant derivative

Takes into account the structure of the manifold and transforms like a tensor.

$$\nabla_{\mu} v^{\nu} = \partial_{\mu} v^{\nu} + \Gamma^{\nu}_{\mu\lambda} v^{\lambda}$$

Autoparallels

A curve whose tangent vector is parallel-transported along itself.

$$u^{\mu} \nabla_{\mu} u^{\alpha} = 0$$


Curvature

The Riemann curvature tensor describes how parallel transport of vectors depends on the path.

$$\Delta v^{\mu} = R_{\alpha\beta}{}^{\mu}{}_{\nu} v^{\nu} dy^{\alpha} dz^{\beta}$$

It can be expressed in terms of the connection:

$$R_{\alpha\beta}{}^{\mu}{}_{\nu} = \partial_{\alpha} \Gamma^{\mu}_{\beta\nu} - \partial_{\beta} \Gamma^{\mu}_{\alpha\nu} + \Gamma^{\mu}_{\alpha\lambda} \Gamma^{\lambda}_{\beta\nu} - \Gamma^{\mu}_{\beta\lambda} \Gamma^{\lambda}_{\alpha\nu}$$



TODO:
Torsion <-> metric mapping
- Riemann curvature
- Autoparallels
-> Problem with parallel transport and tangent space in general

TODO:
Numerical implementation of the tight binding hamiltonian
- details on defect modeling
- unitary timeevolution

TODO:
Numerical results
(tight binding vs. autoparallels)
Plot Scattering angle over defects

TODO:
Outlook
- Dirac equation on curved backgroun with torsion
- Numeric simulations of this for some torsion fields

TODO:
References