

Formalizing the homotopy extension property in LEAN

Mara Masike Silge

Born on 24. April 2003 in Werne

July 10, 2026

Bachelor's Thesis Mathematics

Advisor: Prof. Dr. Floris van Doorn

Second Advisor: Prof. Dr. Markus Hausmann

MATHEMATISCHES INSTITUT

MATHEMATISCH-NATURWISSENSCHAFTLICHE FAKULTÄT DER
RHEINISCHEN FRIEDRICH-WILHELMS-UNIVERSITÄT BONN

Zusammenfassung

Motivation: Whitehead theorem oder so?

Contents

1	Vorbemerkung	4
1.1	Motivation	4
2	Homotopy Extension Property	5
2.1	The Homotopy Extension Property	5
2.2	HEP for relative CW-complexes	8
3	Lean and mathlib	12
3.1	Lean's Dependent Type Theory	12
3.2	Structures	14
3.3	Junk Values	14
3.4	API lemma	14
4	Lean formalisation of the Homotopy Extension Property	15
4.1	CW-complexes in Mathlib	15
4.2	HEP Definition	15
4.3	Retraction	16
5	Methodes and Usage of AI	18
	Literatur	19

List of Symbols

- (I_n) The indexinset for cells of dimension n
- X_n The n -*skeleton* of a CW-complex X

1 Vorbemerkung

In this work, if not specified in any other way, every space has a topology and every map is a continuous map.

1.1 Motivation

The goal of this thesis is to formalize the *homotopy extension property (HEP)* in Lean. I started with the proof of the *retraction criterion* and worked towards the result that every relative CW complex satisfies the HEP. Our motivation for studying this property is that it is needed for two major theorems in algebraic topology. The first is the *cellular approximation theorem*.

Theorem 1.1 (Cellular Approximation Theorem, [MR1867354]). *Every map $f: X \rightarrow Y$ of CW complexes is homotopic to a cellular map. If f is already cellular on a subcomplex $A \subset X$, the homotopy may be taken to be stationary on A .*

Here a map is called *cellular* if it sends the n -skeleton of X to the n -skeleton of Y for all $n \in \mathbb{N}$.

Equipping S^k with the CW structure consisting of one 0-cell and one k -cell, it has no cells in dimensions $0 < n < k$. Hence any map $S^n \rightarrow S^k$ with $n < k$ is homotopic to one landing in the n -skeleton, which is a single point, and is therefore nullhomotopic. This yields the corollary $\pi_n(S^k) = 0$ for all $n < k$.

Secondly, the result that relative CW complexes satisfy the HEP is one of the main steps towards the proof of *Whitehead's theorem*.

Theorem 1.2 (Whitehead's Theorem, [MR1867354]). *If a map $f: X \rightarrow Y$ between connected CW complexes induces isomorphisms $f_*: \pi_n(X) \rightarrow \pi_n(Y)$ for all n , then f is a homotopy equivalence. In case f is the inclusion of a subcomplex $X \hookrightarrow Y$, the conclusion is stronger: X is a deformation retract of Y .*

Whitehead's theorem states that, in order to prove that two connected CW complexes are homotopy equivalent, it suffices to exhibit a map between them that induces isomorphisms on all homotopy groups, i.e. a weak homotopy equivalence. Showing that two spaces are homotopy equivalent is in general a difficult task, and Whitehead's theorem provides a powerful tool for such problems.

2 Homotopy Extension Property

2.1 The Homotopy Extension Property

There are several equivalent ways to define the homotopy extension property, which will be discussed in this first chapter. This first definition is closest to what I formalized in Lean.

Definition 2.1. A pair of topological spaces (X, A) with $A \subseteq X$ has the *homotopy extension property (HEP)* if for every topological space Y , every map $f: X \rightarrow Y$, and every homotopy $H: A \times [0, 1] \rightarrow Y$ satisfying $H(a, 0) = f(a)$ for all $a \in A$, there exists a homotopy $H': X \times [0, 1] \rightarrow Y$ such that

$$\begin{aligned} H'(x, 0) &= f(x) & \text{for all } x \in X, \\ H'(a, t) &= H(a, t) & \text{for all } (a, t) \in A \times [0, 1]. \end{aligned}$$

Remark 2.2. Note, that for any space X , the pair (X, X) and the pair $(X, \emptyset)$ have the *HEP*.

– unsauber, gute Quelle finden

Definition 2.3. A pair of two maps (f, H) , which agree on the space A respectively $A \times \{0\}$, same as in the previous definition, is an *extension problem*.

Remark 2.4. Another equivalent way of phrasing this definition is to say that (X, A) has the *HEP* if and only if every map defined on $X \cup_A (A \times [0, 1])$ extends to a map on $X \times [0, 1]$. Note that this is not, in general, the same as requiring that every map defined on $X \times \{0\} \cup (A \times [0, 1])$ extends to a map on $X \times [0, 1]$. This is because the natural map

$$X \cup_A (A \times [0, 1]) \rightarrow X \times \{0\} \cup (A \times [0, 1])$$

is not necessarily a homeomorphism. In general, it is only a continuous bijection. However, if A is closed, then this map is a homeomorphism and therefore yields another equivalent definition.

The equivalence from theorem 2.4 is proven in the following lemma.

Lemma 2.5. *Let A be a subset of a topological space X .*

(i) (X, A) has the HEP

(ii) *for every topological space Y and every continuous map $g: X \times \{0\} \cup A \times [0, 1] \rightarrow Y$ there exists an extension of g to a map $G: X \times [0, 1] \rightarrow Y$*

Then (i) $\implies$ (ii) and under the additional condition, that A is closed, also the inverse direction (ii) $\implies$ (i) holds.

Proof. (i) $\Rightarrow$ (ii): Let $g : X \times \{0\} \cup A \times [0, 1] \rightarrow Y$. We define maps $f : X \rightarrow Y, x \mapsto g(x, 0)$ and $H : A \times [0, 1] \rightarrow Y, (x, t) \mapsto g(x, t)$. It is easy to check, that f and H agree on $A \times [0, 1]$, so the *HEP* for the pair (X, A) yields an extension of those two maps. We can chose G to be exactly this extended homotopy.

(A closed $\wedge$ (ii)) $\Rightarrow$ (i): Let $f : X \rightarrow Y$ and $H : A \times [0, 1] \rightarrow Y$. We define the map

$$g : X \times \{0\} \cup A \times [0, 1] \rightarrow Y, (x, t) \mapsto \begin{cases} f(x), & \text{if } t = 0, \\ H(x, t), & \text{else} \end{cases}$$

Clearly g is well-defined, since f and H agree on A . To proof continuity of g , we need that $A \subseteq X$ is closed:

$A \subseteq X$ closed $\implies A \times [0, 1] \subset X \times [0, 1]$ closed. Furthermore, $X \times \{0\} \subseteq X \times [0, 1]$ is closed. So g is continuous, because it is continuous on two closed subsets and therefor also on their union.

By assumption there exists an extension $G : X \times [0, 1]$ of g , which we can choose to be the extended homotopy for the definition of the *HEP*. $\square$

Lemma 2.6. *Let A be a subset of a topological space X . Then the following are equivalent:*

- (i) *for every $g : A \rightarrow Y$ there exists an extension $G : X \rightarrow Y$*
- (ii) *A is a retract of X , i.e. $\exists r : X \rightarrow A$, such that r is the identity on $A \subset X$.*

Proof. (i) $\Rightarrow$ (ii): We look at the special case of (i), when $Y = A$ and $g = id_A$. Then the extension G is the desired retraction.

(ii) $\Rightarrow$ (i): Let $g : A \rightarrow Y$. Then we define $G : X \rightarrow Y$ to be the map g precomposed with a retraction $r : X \rightarrow A$. Since r is the identity on A this indeed is an extension of g . $\square$

To proof, that a space pair actually satisfies the *HEP* it is a convenient way to use the *retraction criterion*. This follows from the previous two lemma. Often, the retraction criterion is only phrased for closed subsets $A \subset X$, but I assumed the closeness condition on A , as in theorem 2.5, only for the implication, where it is needed.

Definition 2.7. Let X be a topological space and $A \subset X$ a subset. A retraction is a continuous map $r : X \rightarrow A$, such that precomposition with the inclusion $i : A \hookrightarrow X$ yields the identity on A : $r \circ i : A \rightarrow A = id_A$.

Lemma 2.8 (Retraction Criterion). *Let X be a topological space and $A \subseteq X$ a subspace. Consider the following two statements:*

- (i) *the pair (X, A) has the *HEP**

(ii) the subspace $X \times \{0\} \cup A \times [0, 1] \subseteq X \times [0, 1]$ is a retract of $X \times [0, 1]$

Then (1) $\implies$ (2). If, in addition, A is closed in X , the converse (2) $\implies$ (1) holds as well.

Proof. (i) $\implies$ (ii): If (X, A) has the HEP, then by "(i) $\implies$ (ii)" from theorem 2.5 we get: for every map $g: X \times \{0\} \cup A \times [0, 1] \rightarrow Y$ there exists an extension of g to a map on $X \times [0, 1]$. Now the statement follows directly from theorem 2.5.

(A closed $\wedge$ (ii)) $\implies$ (i): For this direction we first use that $X \cup A \times [0, 1] \subseteq X \times [0, 1]$ is closed and apply the backward direction of theorem 2.6. This yields the same intermediate step as in the previous part of this proof / the forward direction. With the same closedness as before, we apply the backward direction of theorem 2.5 and conclude the proof. $\square$

It follows from the retraction criterion, that the HEP is preserved under taking the product with a space.

Corollary 2.9. *If (X, A) has the HEP, so does $(W \times X, W \times A)$ for any space W .*

Proof. If $r: X \times [0, 1] \rightarrow X \times \{0\} \cup A \times [0, 1] \subseteq X \times [0, 1]$ is a retract, so is

$id_W \times r: W \times X \times [0, 1] \rightarrow W \times X \times \{0\} \cup W \times A \times [0, 1] \subseteq W \times X \times [0, 1]$.

Continuity follows, because id_W is the product of two continuous maps. Furthermore id_W is fixed, where required, since id_W is fixed on all of W . $\square$

-> ToDo: noch in Lean schreiben

Corollary 2.10. *If (X, A) has the HEP and J is a discret set, then $(J \times X, J \times A)$ has the HEP.*

Proof. Let $r: X \times [0, 1] \rightarrow X \times \{0\} \cup A \times [0, 1] \subseteq X \times [0, 1]$ be a retract, then also

$$\begin{aligned} id_J \times r: J \times X \times [0, 1] &\rightarrow J \times (X \times \{0\} \cup A \times [0, 1]) \\ &= J \times X \times \{0\} \cup J \times A \times [0, 1] \end{aligned}$$

is a retract. $\square$

Theorem 2.11. *The HEP is preserved under homeomorphisms, i.e. given two spacepairs (X, A) and (Y, B) . If there exists a homeomorphism from X to Y , that maps A to B , and (X, A) has the HEP, then so does (Y, B) .*

Proof. Let $\phi: X \rightarrow Y$, be a homeomorphism, that maps A to B and let $r_X: X \times [0, 1] \rightarrow X \times \{0\} \cup A \times [0, 1]$ be a retraction, which we obtain from the Retraction Criterion for the pair (X, A) . $pr_X: X \times [0, 1] \rightarrow X$ and

$pr_I: X \times [0, 1] \rightarrow X$ are the projections onto the first respectively second component. We use those maps, to define r_Y .

$$\begin{aligned} r_Y: Y \times [0, 1] &\rightarrow Y \times \{0\} \cup B \times [0, 1] \\ (y, t) &\mapsto (\phi \circ pr_X \circ r_X(\phi^{-1}(y), t), pr_I \circ r_X(\phi^{-1}(y), t)) \end{aligned}$$

It is left to check, that r_Y is a retraction, which yields the *HEP* for the space pair (Y, B) . The map r_Y ...

- is continuous: the two component functions are compositions of continuous maps.
- is fixed on $Y \times \{0\} \cup B \times [0, 1]$: Let (y, t) be a point in this subspace of $Y \times [0, 1]$. Then, $(\phi^{-1}(y), t)$ is in $X \times \{0\} \cup A \times [0, 1]$, because ϕ^{-1} maps Y to X and B to A . So $(\phi^{-1}(y), t)$ lies in exactly the space, which is fixed under r_X . After applying ϕ onto the first component, we observe, that $r_Y(y, t) = (y, t)$.
- has image in $Y \times \{0\} \cup B \times [0, 1]$: Let (y, t) be a point in $Y \times [0, 1]$. $r_X(\phi^{-1}(y), t)$ is in the image of r_X , which is $X \times \{0\} \cup A \times [0, 1]$. It follows from the definition of r_Y , that either the second component of $r_Y(y, t)$ is zero, or the first one lies in B .

□

2.2 HEP for relative CW-complexes

To proof the *HEP* for relative CW-complexes, we will generalize the space pair in three steps and proof the *HEP* for each of these pairs. The general strategy is as follows.

1. Any disc relative its boundary has the *HEP*.
2. The pair of two consecutive skeletons has the *HEP*, i.e (X_{n+1}, X_n) has the *HEP* for all n .
3. Any relative CW-complex has the *HEP*.

Before we start proofing the *HEP* for any space pair, I will give some definitinons, introduce notation for CW-complexes and proof some basic properties for them, which are not yet in mathlib.

I copied the following definition of CW-complexes from the bachelor thesis of Hannah Scholz [Sch24]. Her formalisation of CW complexes, which can be found as the "classical CW-complex" in mathlib, is explained in this thesis. She chose the historical definition of CW-complexes, presented by Whithead in [Whi18].

Definition 2.12. Let X be a Hausdorff space. A *CW-complex* on X consists of a family of indexing sets $(I_n)_{n \in \mathbb{N}}$ and a family of maps $(Q_i^n : D_i^n \rightarrow X)_{n \geq 0, i \in I_n}$ s.t.

1. $Q_i^n|_{D_i^n} : D_i^n \rightarrow Q_i^n(D_i^n)$ is a homeomorphism. We call $e_i^n := Q_i^n(D_i^n)$ an (*open*) n -cell (or a cell of dimension n) and $\bar{e}_i^n := Q_i^n(D_i^n)$ a *closed* n -cell.
2. For all $n, m \in \mathbb{N}$, $i \in I_n$ and $j \in I_m$ where $(n, i) \neq (m, j)$ the cells e_i^n and e_j^m are disjoint.
3. For each $n \in \mathbb{N}$, $i \in I_n$, $Q_i^n(\partial D_i^n)$ is contained in the union of a finite number of closed cells of dimension less than n .
4. $A \subseteq X$ is closed iff $Q_i^n(D_i^n) \cap A$ is closed for all $n \in \mathbb{N}$ and $i \in I_n$.
5. $\bigcup_{n \geq 0} \bigcup_{i \in I_n} Q_i^n(D_i^n) = X$.

We call Q_i^n a *characteristic map* and $\partial e_i^n := Q_i^n(\partial D_i^n)$ the *frontier of the* n -cell for any i and n . Additionally we define $X_n := \bigcup_{m < n+1} \bigcup_{i \in I_m} \bar{e}_i^m$ and call it the n -skeleton of X for $-1 \leq n \leq \infty$

Definition 2.13. CW-complex Hanna – ToDo

– Charakteristic map

The more modern way to think about CW-complexes is the following: – ToDo

The modern definition yields quotientmaps from every D^n to a corresponding cell of dimension n . The weak topology on a CW complex agrees with the quotient-topology induced by those quotientmaps. – ToDo : Könnte man auch als Theorem mit reinnehmen, aber brauche ich im Finalen Beiweis vermutlich nicht – Furthermore, we can define a quotient map onto all of the n -skeleton of a relative CW complex in the following way : is a quotientspace of the disjoint union of the base space together with all D^k in dimension $k \leq n$. We will proof, that the weak topology of the relative CW complex indeed is the same as the topology induced by this quotientmap onto the skeleton.

Lemma 2.14. Let (X, C) be a relative CW-complex with skeleton X_n . There exists a quotient map $C \dot{\cup} \coprod_{n \leq m} \coprod_{i \in J_n} D^n \rightarrow X_m$ for every $m \in \mathbb{N}$.

Proof. We define the map

$$Q : C \dot{\cup} \coprod_{n \leq m} \coprod_{i \in J_n} D^n \rightarrow X_m$$

$$x \mapsto \begin{cases} Q_i^n(x), & \text{if } x \in \text{the } i\text{th copy of } D^n, \\ x, & \text{else } (x \in C) \end{cases}$$

Surjectivity of Q follows directly from the modern definition, and it is also already a theorem in mathlib, which writes the m -skeleton as the union of the base-space C together with all cells in dimension up to dimension m . To prove, Q defines a quotient map, it remains to prove that the topologies agree.

For $A \subset X_m$ closed in the weak topology of a relative CW complex, we get by definition, that $Q_i^n(D^n) \cap C$ is closed for all $n \in \mathbb{N}$ and $i \in J_n$ and $A \cap C$ closed. Together with the characteristic maps and the identity being continuous, this is all we need, for $Q^{-1}(A)$ to be closed as well.

Assume $A \subset X_m$ closed in the topology, which is induced by Q . □

Theorem 2.15. $\forall n \in \mathbb{N}$: $(D^m, \partial D^m)$ has the HEP.

There are two different attempts to prove this theorem. One is quite visual via the retraction criterion and the other one constructs the extended homotopy explicitly. Here, I will give the constructive proof, because that one was easier to formalize in LEAN.

Proof. Given a homotopy $H : \partial D^m \times [0, 1] \rightarrow Y$ and a map $f : D^m \rightarrow Y$, which agree on $\partial D^m \times \{0\}$ identified with ∂D^m . We assemble these to a map $h : D_2^m \rightarrow Y$, where D_2^m denotes a disc with radius 2 in $\mathbb{R}^m$. h is defined in the following way.

$$h(x) = \begin{cases} f(x) & \text{if } \|x\| \leq 1 \\ H(x/\|x\|, \|x\| - 1) & \text{else} \end{cases}$$

Using h we define the desired homotopy $H' : D^m \times [0, 1] \rightarrow Y$ to be

$$H'(x, t) = h((1 + t)x)$$

It is left to check that this actually defines a continuous extension of the homotopy H and that it starts with f .

-> wenn zu kurz, dann hier ausführlicher □

Since the definition of CW-complexes in Mathlib uses n -dimensional cubes instead of closed balls as preimage of a closed cell under the characteristic map, we will state the *HEP* for cubes relative to their boundary.

Corollary 2.16. $\forall n \in \mathbb{N}$: $([-1, 1]^n, \partial[-1, 1]^n)$ has the HEP.

Proof. This statement follows from the observation, that D^m is homeomorphic to $[-1, 1]^n$. One constructs a homeomorphism via the radial projection, which maps the boundary of the sphere to the boundary of the cube. All points in the interior are scaled along the rays. In that way, the cube inherits the *HEP* from the sphere. □

In the next step, we will prove, that for every relative CW complex (X, A) with skeleton X_n and for all $n \in \mathbb{N}$, the pair (X_{n+1}, X_n) has the *HEP*. – The key-idea of the proof is to apply the retraction criterion. To construct the retraction, we apply a retraction for each of the added cells relative their boundary at the same time. On the base space, we use the identity. Later, we will need this result in order to construct a retraction on the entire CW-complex.

Theorem 2.17. *Let (X, C) be a relative CW-complex with skeleton X_n , then $\forall n \in \mathbb{N} : (X_{n+1}, X_n)$ has the *HEP*.*

Before provig this theorem, we need some preparation. The key-idea of the proof is, to apply the retraction criterion, i.e. we want to construct a map on $X_{n+1} \times [0, 1]$.

Proof. We proof the *HEP*, using the retraction criterion.

First, we define the map and then proof afterwards, that it has the required properties, i.e. is surjective and the weak topology agrees with the inuced topology of the quotientmap. For the definition of the map, we use the retraction r_C from the *HEP* for cubes relative boundary, which is defined on $[-1, 1]^n \times [0, 1]$. Even tough r_C depends on the dimension of the cube, for reasons of convenience, we use the same notation for all of those maps. Let J_n denote the indexing set for cells in dimension $n \in \mathbb{N}$ and $Q_i^n : D^n \rightarrow X$ be the charakteristic map of the i th n -cell.

Informally spoken, the map r

$$r : X_{n+1} \times [0, 1] \rightarrow X_{n+1} \times \{0\} \cup X_n \times [0, 1]$$

$$(x, t) \mapsto \begin{cases} r_C(), & \text{if } t = 0, \\ H(x, t), & \text{else} \end{cases}$$

□

3 Lean and mathlib

After taking a closer look at the informal mathematics behind my formalisation in the previous chapter, I now want to give a brief overview of what Lean is. I will explain some of the conventions and concepts used in formalization that are relevant to this project.

Lean is a proof assistant, i.e. a tool that checks and confirms the individual reasoning steps of mathematical proofs written in Lean’s formal language. It verifies that assumptions, axioms, functions, and theorems are only applied where permitted, and that new results are derived using valid logical inferences. In *The Lean Theorem Prover (system description)* the main designers of Lean explain that the correct verification of mathematical proofs in Lean rests upon a “small trusted kernel based on dependent type theory” (leanTP). I will elaborate the term “dependent type theory” later in this section.

Lean is one of several proof assistants, alongside others such as Rocq or Isabelle. However, as Alex Kontorovich elaborates in *The Shape of Math to Come* [Kon25], it is remarkable how quickly Lean is gaining relevance, not only in computer science and mathematical logic, but also in mainstream mathematical research.

Since July 2023, the non-profit *Lean Focused Research Organization (FRO)* has supported and coordinated the ongoing development of the language. Its stated goal is “enhanced scalability, usability, documentation, and proof automation while guiding Lean toward long-term self-sustainability” [FRO]. The same source also provides a brief overview of Lean’s development history. The first version was created by *Leonardo de Moura* in 2013. Since then, the language itself has evolved. The first stable release of Lean 4 appeared in September 2023, and both the community and the digital library *mathlib* have grown considerably. Today, mathlib 4 contains more than 2,000,000 lines of code and has over 770 contributors. Current statistics can be found at [\[1\]](#).

3.1 Lean’s Dependent Type Theory

The digital book *Theorem Proving in Lean 4* teaches how to verify proofs in Lean and explains the underlying logical system. Its second chapter, on *Dependent Type Theory*, serves as the basis for the following description.

The first and probably most important fact is that every expression in Lean has a type. Using the command `#check`, one can display the type of an expression. In the example below, I first define `n` to be the natural number 3 and then check its type, which is $\mathbb{N}$ by the definition above. As stated, every expression in Lean has a type, so the natural numbers themselves can be checked as well. Their type is called `Type`. The classical objects $\mathbb{N}, \mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$, as well as the finite-dimensional Euclidean spaces, all have type `Type`. Continuing the same procedure and checking the type of `Type` yields

`Type` : `Type` 1. This can be repeated indefinitely: in general, `Type` k has type `Type` $(k+1)$ for every natural number k . I want to present one more `Type`, which is called `Prop`. As the name already implies, `Prop` is the type used for propositions, e.g. $2 = 1$. We can combine propositions to build new ones. For example we can connect two propositions with a logical "and" in order to obtain a new expression of type `Prop`. `Prop` itself has type `Type`.

```
def n : ℕ := 3
#check n
      n : ℕ
#check ℕ
      ℕ : Type
#check EuclideanSpace ℝ (Fin 3)
      EuclideanSpace ℝ (Fin 3) : Type
#check Prop
      Prop : Type
#check Type
      Type : Type 1
#check 2 = 1
      2 = 1 : Prop
#check Prop
      Prop : Type
```

Type theory not only allows one to define an element, such as the natural number 3 above, but also to construct new types from existing ones. Functions are one such example. To illustrate this, I start by choosing two arbitrary but fixed types α and β . Writing `Type*` admits types of any universe. Lean then automatically introduces a so-called universe variable, here `u_1`. A function with domain α and codomain β has type $\alpha \rightarrow \beta$. This newly constructed type again has a type of its own, whose universe is the maximum of the universe levels of α and β .

```
variable (α β : Type*) (f : α → β)
#check α
      α : Type u_1
#check f
      f : α → β
#check α → β
      α → β : Type (max u_1 u_2)
```

Functions such as the addition of natural numbers, which one would usually write as a map $\mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$, have in Lean the type $\mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbb{N}$. This means that after applying a single natural number to the addition function, we obtain a function $\mathbb{N} \rightarrow \mathbb{N}$ that adds the first number. Since $\mathbb{N}$ has type `Type`, the reasoning of the previous paragraph shows that the function type $\mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbb{N}$ again has type `Type`. Note the distinction between the two levels: the addition function itself has type $\mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbb{N}$, whereas this function type in turn has type `Type`.

```
#check Nat.add
      Nat.add :  ℕ → ℕ → ℕ
#check ℕ → ℕ → ℕ
      ℕ → ℕ → ℕ :  Type
```

Even though I do not want to illustrate in detail, what makes the type theory of lean a dependent type theory, let me

3.2 Structures

3.3 Junk Values

If you write a division by zero in a formalized theorem, Lean returns the result 0.

```
example : 1 / 0 = 0 := by norm_num
```

This makes division in Lean always well-defined, even though certain inputs have no genuine mathematical meaning. As explained in [MLS25], the use of so-called *junk values* is common practice. The same convention applies, for instance, to the inversion of square matrices, which is always defined and returns the zero matrix whenever the matrix is singular, or to the derivative of a non-differentiable function, which is defined to be zero. Many further examples of this kind exist. At first glance this may seem concerning, since such values have no mathematical justification and could appear to enable incorrect proofs. Indeed, one can state and prove propositions that, in classical mathematics, would look false or ill-defined. This issue is resolved by adding suitable hypotheses such as $n \neq 0$ or `Differentiable k f` to any statement about the object in question. They ensure we only use the result of the theorem in applicable situations, such as, if our number is nonzero or our function is differentiable. Consequently, although the convention may seem alarming initially, relying on junk values is standard practice in Lean formalizations, precisely because it simplifies both definitions and statements. Returning to the first example, it is what allows division to be expressed as a total map $\mathbb{Q} \rightarrow \mathbb{Q} \rightarrow \mathbb{Q}$.

3.4 API lemma

4 Lean formalisation of the Homotopy Extension Property

4.1 CW-complexes in Mathlib

1. NOTIZEN:
2. Zellen im CW complex sind würfel anstatt spheres
3. CW -complexe leben in einem größeren Raum ->

4.2 HEP Definition

Already in the mathematical section, the definition of the *homotopy extension property* was quite long and technical. The same is true for its formalization. Currently I have two versions of the definition in my project, since we decided within the work progress, that a different phrasing might be more applicable. I will later on discuss the pros and cons for each decision. The first version of the definition is the following:

```
agreeOn.{u} {X Y : Type u} (f : X → Y) (H : X × ℝ → Y) (A : Set X)
  : Prop :=
  ∀ (a : A), f a = H (a, 0)
```

```
HomotopyExtension.{u} {X Y : Type u} [TopologicalSpace X] [
  TopologicalSpace Y] (H' : X × ℝ → Y) (f : X → Y) (H : X × ℝ
  → Y) (A : Set X) : Prop :=
  ContinuousOn H' { p : X × ℝ | p.2 ∈ unitInterval } ∧ { p : X ×
  ℝ | p.2 ∈ unitInterval }.MapsTo H' rangeH' ∧ (∀ (x : X), f x =
  H' (x, 0)) ∧ (∀ (a : A) (t : unitInterval), H (a, t) = H' (a, t))
```

```
HEP.{u} (X : Type u) [TopologicalSpace X] (A : Set X) : Prop :=
  ∀ (Y : Type u) [TopologicalSpace Y], ∀ (f : X → Y), ∀ (H : X ×
  ℝ → Y), Continuous f → ContinuousOn H {p : X × ℝ | (p.1 ∈ A)
  ∧ (p.2 ∈ (unitInterval))} → agreeOn f H A → ∃ (H' : X × ℝ →
  Y), HomotopyExtension H' f H A
```

To shorten the definition a little and in order to make it more readable, I first defined `agreeOn`. This tells, that the function `f` and the homotopy `H`, which we want to extend, have the same values on the subset `A` respectively $A \times 0$. The second "predefinition" I made, is called `HomotopyExtension` and combines all the properties, our desired homotopy should have. Namely, continuity and that it agrees with the two functions given in the begin. While the benefit of the first predefinition might be discussed, the second clearly shortens our definition.

If we now take a closer look at the definition of the homotopy extension property, it is pretty similar to the mathematical definition theorem 2.1 of the *HEP*. Still, there is one point, I want to discuss. Defining the homotopies

as a map on $X \times \mathbb{R}$ instead of $X \times \text{unitInterval}$ makes it easier to work with the function. We apply it to any real number in the second coordinated, without the need to provide a prove that this real number lies between zero and one. Still, this does not change anything about the defined object. `ContinuousOn` $H' \{ p : X \times \mathbb{R} \mid p.2 \in \text{unitInterval} \}$ and $\forall (a : A) (t : \text{unitInterval}), H(a, t) = H'(a, t)$ make it irrelevant, what happens outside of the compact interval, but instead puts precise focus on what we need for our definition.

Let us now compare this definition with the later version. `HEP'` :

```
open Set.Notation
def HEP' {Y : Type*} [TopologicalSpace Y] (X A : Set Y) : Prop :=
  HEP X (X ↓ ∩ A)
```

The difference to the first version of the definition is, that in our definition `HEP X A`, A is a subset of the type X . That means that the pair of spaces does not live in an ambient space. Whereelse, in the definition `HEP'`, both sets X and A are sets of the same type Y .

4.3 Retraction

Even though defining a retraction r from B to A via the construction in theorem 2.7 is the convenient approach, I chose a different phrasing for the formalization. Instead of working directly with the classical definition, I introduced a structure that bundles a map with two sets and three defining properties. Concretely, I defined the following structure:

```
structure RetractionOn {X : Type*} [TopologicalSpace X] (r : X → X)
  (B A : Set X) : Prop where
  continuousOn : ContinuousOn r B
  mapsTo : B.MapsTo r A
  fixesOn : ∀ a ∈ A, r a = a
```

The first thing to notice is, that the map r is a map from a Type X into itself and it is not a retraction on all of its domain. As the name `RetractionOn` implies, it just "retracts" a subset B to a subset A of the Type X . The first property `continuousOn` ensures, it is a continuous Map on the relevant subset B . The codomain of the retraction is supposed to be A , so the property `mapsTo` ensures, that r maps all elements of B to an element in A . Furthermore the classical definition states, that precomposition with the inclusion $A \hookrightarrow B$ should give the identity id_A . In our case there is not really an inclusion map, since both - image and preimage of r - are already elements in the same Type X . So in our setting this condition implies, that each element in A needs to be fixed by applying r . This property is called `fixedOn`.

Note, that the classical definition requires A to be a subset of B , since otherwise there would not be an inclusion map $A \hookrightarrow B$. This condition never occurs in the formalized version of the definition. But only if it is satisfied,

the structure actually gives that the map r is a retraction from B to A in the classical sence. You can see, that leaving out this condition actually yields a more general structure. If we additional assumed the subset condition, we would also get, that $r(B) = A$, which is a straight forward observation from the standard definition. But this is not in general true for the structure defined above.

So why did I leave out this condition? First of all it makes the definition itself but also proofs using it shorter. When we work with a retraction, the instance $A \subset B$ usually is true. Therefore it sufficed for all proofs in my thesis. For example, proving the retraction criterion only requires exactly those three properties. When one wants to proof more general theorems about retractions it might be necessary to add $A \subset B$ as an additional assumption. One can say, that the defined structure takes a *junk value* as explained in the section 3.3. In the case that $A \not\subset B$ we still obtain an object of the defined structure, same as the division by zero still returns the number zero.

5 Methodes and Usage of AI

References

- [] *Mathlib Statistics*. URL: https://leanprover-community.github.io/mathlib_stats.html (visited on 07/02/2026).
- [FRO] Lean FRO. *FRO - About us*. URL: <https://lean-lang.org/fro/about/> (visited on 07/02/2026).
- [Kon25] Alex Kontorovich. *The Shape of Math To Come*. 2025. arXiv: 2510.15924 [math.HO]. URL: <https://arxiv.org/abs/2510.15924>.
- [MLS25] Alex Meiburg, Leonardo A. Lessa, and Rodolfo R. Soldati. *A Formalization of the Generalized Quantum Stein’s Lemma in Lean*. 2025. arXiv: 2510.08672 [quant-ph]. URL: <https://arxiv.org/abs/2510.08672>.
- [Sch24] Hannah Scholz. “Formalisation of CW-complexes”. Bachelor’s Thesis Mathematics. Rheinischen Friedrich-Wilhelms-Universität Bonn, 2024. URL: <https://florisvandoorn.com/theses/HannahScholz.pdf>.
- [Whi18] J. H. C. Whitehead. “Combinatorial homotopy. I”. In: *Bulletin (new series) of the American Mathematical Society* 55.3 (2018), pp. 213–245. ISSN: 0273-0979.