

STATISTICAL COMPUTATIONAL METHODS

Seminar Nr. 3

Computer Simulations of Random Variables and Monte Carlo Studies; Inverse Transform Method, Rejection Method, Special Methods

1. Use the inverse transform method to generate a $Geo(p), p \in (0, 1)$, variable.
2. Use a special method to generate a $Poiss(\lambda), \lambda > 0$, variable.
3.
 - a) Use the inverse transform method to generate an $Exp(\lambda), \lambda > 0$, variable.
 - b) Use the inverse transform method to generate an $Gam(\alpha, \lambda), \alpha \in \mathbb{N}, \lambda > 0$, variable.
Hint: A Gamma $Gam(\alpha, \lambda)$ variable is the sum of α independent $Exp(1/\lambda)$ variables.
 - c) Use the rejection method to generate an $Exp(\lambda), \lambda > 0$, variable.

4. Application: Forecasting for new software release

An IT company is testing a new software to be released. Every day, software engineers find a random number of errors and correct them. On each day t , the number of errors found, X_t , has a $Poisson(\lambda_t)$ distribution, where the parameter λ_t is the lowest number of errors found during the previous k days,

$$\lambda_t = \min\{X_{t-1}, X_{t-2}, \dots, X_{t-k}\}.$$

If some errors are still undetected after $tmax$ days (i.e. if not all errors are found in $tmax - 1$ days), the software is withdrawn and goes back to development. Generate a Monte Carlo study to estimate

- a) the time it will take to find all errors;
 - b) the total number of errors found in this new release;
 - c) the probability that the software will be sent back to development.
- (Try $k = 4, [X_{t-1}, X_{t-2}, X_{t-3}, X_{t-4}] = [10, 5, 7, 6], tmax = 10$.)