

# The Probabilistic Method

exercise solutions by

MARCEL K. GOH

14 FEBRUARY 2022

## 1. The Basic Method

**Exercise 1.1.** Prove that if there is a real  $p$ , with  $0 \leq p \leq 1$  such that

$$\binom{n}{k} p^{\binom{k}{2}} + \binom{n}{t} (1-p)^{\binom{t}{2}} < 1,$$

then the Ramsey number  $r(k, t)$  satisfies  $r(k, t) > n$ . Using this, show that

$$r(4, t) \geq \Omega(t^{3/2}/(\ln t)^{3/2}).$$

*Proof.* We follow the proof of Proposition 1.1.1 in the book. We consider a random graph on  $n$  vertices, where each edge is present with probability  $p$ . Let  $K$  be the event that there is a clique of size  $k$  in the graph, and let  $I$  be the event that there is an independent set of size  $t$  in the graph. By the union bound,

$$\mathbf{P}\{K \cup I\} \leq \mathbf{P}\{K\} + \mathbf{P}\{I\} \leq \sum_{|S|=k} p^{\binom{k}{2}} + \sum_{|S|=t} (1-p)^{\binom{t}{2}} = \binom{n}{k} p^{\binom{k}{2}} + \binom{n}{t} (1-p)^{\binom{t}{2}} < 1.$$

This means that  $\mathbf{P}\{\neg K \cap \neg I\} > 0$  and since the sample space is finite, there exists a graph on  $n$  vertices with no clique of size  $k$  and no independent set of size  $t$  and therefore  $r(k, t) > n$ .

Next we show that  $r(4, t) > (et/\ln t)^{3/2}$  for large enough  $t$ . Note that

$$\binom{n}{4} p^6 + \binom{n}{t} (1-p)^{\binom{t}{2}} \leq n^4 p^6 + \frac{e^t n^t}{t^t} (1-p)^{t^2/4},$$

by the inequality

$$\frac{n^k}{k^k} \leq \binom{n}{k} \leq \frac{e^k n^k}{k^k}.$$

Setting  $n = et^{3/2}/(\ln t)^{3/2}$ , we have

$$\begin{aligned} n^4 p^6 + \frac{e^t n^t}{t^t} (1-p)^{t^2/4} &= \left( \frac{tp}{e \ln t} \right)^6 + \frac{e^t t^{3t/2}}{t^t e^{3t/2} (\ln t)^{3t/2}} (1-p)^{t^2/4} \\ &= \left( \frac{tp}{e \ln t} \right)^6 + \frac{t^{t/2}}{e^{t/2} (\ln t)^{3t/2}} (1-p)^{t^2/4} \\ &\leq \left( \frac{tp}{e \ln t} \right)^6 + \left( \frac{t(1-p)^{t/2}}{e (\ln t)^3} \right)^{t/2} \\ &\leq \left( \frac{tp}{e \ln t} \right)^6 + \left( \frac{t}{e^{pt/2+1} (\ln t)^3} \right)^{t/2}, \end{aligned}$$

where in the last line we used the inequality  $1-p \leq e^{-p}$ . Choosing  $p = 2 \ln t/t$ , we simply need  $t$  large enough such that

$$\left( \frac{t}{e^{\ln t+1} (\ln t)^3} \right)^{t/2} = \left( \frac{1}{e (\ln t)^3} \right)^{t/2} < 1 - \frac{2}{e^6},$$

which can be done since the left-hand side goes to 0. ■