

Answers to Selected Exercises in *Principles of Mathematical Analysis**

Solutions by

MARCEL K. GOH

CHAPTER 2. BASIC TOPOLOGY

1. Prove that the empty set is a subset of every set.

Proof. Let S be an arbitrary set. Then every element of \emptyset is an element of S . So $\emptyset \subseteq S$. ■

2. A complex number z is said to be algebraic if there are integers a_0, \dots, a_n , not all zero, such that

$$a_0 z^n + a_1 z^{n-1} + \dots + a_{n-1} z + a_n = 0.$$

Prove that the set of all algebraic numbers is countable. *Hint:* For every positive integer N , there are only finitely many equations with

$$n + |a_0| + |a_1| + \dots + |a_n| = N$$

Proof. Let A denote the set of all algebraic numbers and partition A as follows: For each $z \in A$, calculate the positive integer N that corresponds to its equation and place it in a set E_N . So

$$A = \bigcup_{N \in \mathbf{N}} E_N,$$

where each E_N is finite. Then apply the corollary of Theorem 2.12 to find that A is countable. ■

3. Prove that there exist real numbers which are not algebraic.

Proof. Let A denote the set of all algebraic numbers and suppose, towards a contradiction, that all real numbers are algebraic. Then $\mathbf{R} \subseteq A$. But the set of real numbers is uncountable and we know from Problem 2 that A is countable. The contradiction completes the proof. ■

4. Is the set of all irrational real numbers countable?

The answer is no.

Proof. If $\mathbf{R} \setminus \mathbf{Q}$ is countable, then $\mathbf{R} = \mathbf{Q} \cup (\mathbf{R} \setminus \mathbf{Q})$, is countable, which we know to be false. ■

5. Construct a bounded set of real numbers with exactly three limit points.

$\{\frac{1}{n} : n \in \mathbf{N}\} \cup \{\frac{1}{n} + 2 : n \in \mathbf{N}\} \cup \{\frac{1}{n} + 4 : n \in \mathbf{N}\}$ has limit points 0, 2, and 4. It is bounded above by 6 and below by 0.

* Walter Rudin. 1986. *Principles of Mathematical Analysis*, 3rd ed. McGraw-Hill, Inc., USA.