

MATH 140 Fall 2026

notes by

MARCEL GOH

A note on these notes. This document will periodically be updated with the material that we cover. The timestamps in the left margin indicate when the notes from each day start. Subsections labelled with a * are optional. These notes should *not* be used as a substitute for the textbook or for attending class. (For instance, on the blackboard I will often draw graphs or diagrams, and these for the most part will be omitted from the notes.) The best way to do well in this class is to do as many exercises from Stewart's textbook as you can stomach. This document may contain errors, typographical or mathematical. Please email me if you find any.

I. Foundations	2
1. Rates of change	3
2. Limits	5
3. Rules for evaluating limits	18
4. Continuity	29
5. Derivatives	34
II. Techniques	46
6. Algebraic laws for differentiation	47
7. Derivatives of the exponential and trigonometric functions	52

I. FOUNDATIONS

*Que si cependant la Geometrie a toujours
quelque obscurité essentiel, qu'on ne puisse dissiper,
& ce fera uniquement, à ce que je crois, du côté de l'Infini,
c'est que de ce côté-là la Geometrie tient à la Phisique,
à la nature intime des Corps que nous connoissons peu,
& peut-être aussi à une Metaphisique trop élevée,
dont il ne nous est permis que d'appercevoir quelques rayons.*

— BERNARD LE BOVIER DE FONTENELLE, *Éléments de la Geometrie de l'Infini* (1727)

1. Rates of change

31.VIII This set of notes assumes that the reader is familiar with all the pre-calculus concepts that one learns in the later years of secondary school: functions, trigonometry, exponential functions, logarithms, etc. However, given that the very definition of what differential calculus is requires us to talk about functions and their inputs, it is perhaps prudent to begin with a refresher on real-valued functions with real-valued outputs.

A *function* is a collection of ordered pairs (x, y) with the rule that if (x, y) and (x, z) are both in the collection, then $y = z$. We can view x as the *input* to the function, and y as the *output* of the function; under this view, the rule above is simply that the function never outputs two different values when given the same input. People like to use the letter f (or g or h , but really we can use any letter) to denote a function, and $f(x)$ to denote its value evaluated at a given input x . However, many particular functions are difficult to write without involving the input variable x in the expression, and in these cases we'll write $f(x)$ to mean the function as well.

Here are some examples of functions:

- a) The function $f(x) = 3x$, defined on all real numbers, that outputs triple its input value.
- b) The function that assigns to each real number x its square x^2 .
- c) The function $y = f(x)$ that equals 0 whenever x is a negative real number, and equals x whenever $x \geq 0$.
- d) The function that takes as input any nonzero real number x and outputs its reciprocal $1/x$.
- e) The function $f(x)$, defined on all real numbers, given by

$$f(x) = \begin{cases} 1, & \text{if } x \text{ is rational;} \\ 0, & \text{if } x \text{ is irrational.} \end{cases}$$

Note that in each case above, we were careful to specify what the possible inputs to the function were. Going back to the definition of a function as a collection of ordered pairs (x, y) , the set of all x such that (x, y) is in the function is called the *domain* of f ; this is precisely the set of all valid inputs to the function f . (The set of all y such that (x, y) is in f is called the *range* of f , but this definition won't be as important to us.)

The title of this section is “rates of change,” but determining the rate at which a given function changes is actually what this *entire class* will essentially be about. We want to study how much the output of a function changes when you change its input.

For example, consider a car moving from Montreal to Quebec City. As time goes on, it gets further away from Montreal and closer to Quebec City. So we can model the trip as a function $d(t)$, where for each positive real number t we let $d(t)$ equal the distance of the car from Montreal (in kilometres), exactly t

minutes since it set off. The rate of change of this function at any given point in time t is the car's *velocity* at that point.

How do we calculate velocity? Well, we know that it is measured in units such as “kilometres per hour” or “metres per second,” so it is not too hard to see that velocity is calculated by dividing distance over time.

Let's go back to the car example. Suppose your car is equipped with a very good GPS that can measure your distance from Montreal; however, it does not have a working speedometer. Your friend (who is driving) asks you, “How fast am I driving?” You have taken high school physics, so you know how to solve this problem. You get out a stopwatch and start it exactly when the GPS measures your distance at 135.2 kilometres from Montreal. After one minute on the stopwatch, the GPS reads 137.4, so you report to your friend that she is driving $137.4 - 135.2 = 2.2$ kilometres per minute, or 132 kilometres per hour.

But your friend is not satisfied. “You told me my average speed over the course of a minute, not my speed at the moment I asked you the question. During that minute, I had to speed up to overtake a U-Haul and then slow down because someone cut me off in the fast lane.”

So you try and do better. You get out the stopwatch again, and when you start it the GPS reads 137.7. You wait ten seconds, then observe that the GPS reads 138.1. So you report your friend's new speed at $138.1 - 137.7 = 0.4$ kilometres every ten seconds, or 2.4 kilometres per minute, or 144 kilometres per hour.

Your friend is a little more satisfied, but wonders if you can do better. So you try one last time. You start the stopwatch when the GPS reads 138.5, wait exactly one second, and then check the GPS again. To everyone's surprise and disappointment, it still reads 138.5, because it only measures distance to the tenth part of a kilometre. So the calculation would report that the car has moved 0 kilometres in that one second, that is, it's going 0 kilometres per hour.

Obviously in this example you were hamstrung by the precision of your measuring instruments, but it is still worth pondering the following point. Your friend was asking for her speed at some particular instant. However, during any particular instant she does not move, nor does time move. So any naïve calculation as to her speed would go along the lines of $0/0$, which is an undefined value.

Yet it is undeniable that something's “speed at a particular instant” is a concept that holds meaning in everyday life. If you are in a plane and your friend is in a car below you, then it's certainly not nonsensical to say that you are moving faster than your friend at that moment, even though technically it could also be said that both of you have moved “0 kilometres over 0 hours.”

The way we make this precise is suggested by your technique in the first scenario. To avoid dividing by zero, we let time elapse just a tiny bit, then measure how much the object's position has changed in that minute interval, and perform the division. Assuming we have arbitrarily sophisticated measuring instruments, we can keep taking smaller and smaller intervals, and arrive at something closer and closer to the “instantaneous velocity.” But it is worth

stating that “instantaneous velocity” is a theoretical concept, and not one that can be measured exactly. So when you are driving in a real car, and your speedometer reads some value, say, 127 kilometres per hour, what it’s really doing is measuring your car’s change in distance over a very small period of time, and reporting to you the quotient (not unlike what you were doing in that first scenario!).

Secants and tangents. The average rate of change of a function $f(x)$ between the input x and the input $x + h$ (for some small $h > 0$) is the total amount it changes, namely $f(x + h) - f(x)$, divided by the amount the input changed, namely $(x + h) - x = h$. If you graph the function $f(x)$ on the (x, y) plane, then geometrically what you are doing is drawing a line between the points $(x, f(x))$ and $(x + h, f(x + h))$ and calculating its *slope* (the “rise over run”). Such a line cutting through the function’s graph is called a *secant line*, because it *cuts* through the function, at the very least at those two aforementioned points.

By taking h smaller and smaller (but never zero!) we continue to obtain secant lines, but the distance between the two points $(x, f(x))$ and $(x + h, f(x + h))$ will get smaller and smaller. You can imagine these points getting so close that it’s as if the *line touches the function exactly once*, at the point $(x, f(x))$ and nowhere else. This line, if it exists, will be called the *tangent line*, but we’re not ready to precisely define it just yet.

All we note at this point is that if we admit the concept of the “instantaneous rate of change,” and allow that it can be approximated by taking the the average rate of change over smaller and smaller intervals, then the slope of this tangent line (if it exists!) is exactly the instantaneous rate of change that we seek. So it seems that the slope of the tangent line should be the value of the quotient

$$\frac{f(x + h) - f(x)}{(x + h) - x} = \frac{f(x + h) - f(x)}{h}$$

when $h = 0$, if only we could plug in $h = 0$. But we cannot, since division by zero is undefined.

What we have learned from this preliminary section is that to understand rates of change, we shall have to develop a precise concept of what it means to take a function close to some input at which it is not necessarily defined. This is the notion of a *limit*.

2. Limits

02.IX In this section we establish the notion of a limit, which will be a rigorous way of formalising the idea of a function approaching a value at some point where it may or may not be defined. To understand the definition of a limit, it might be helpful to think about it as a generalisation of the concept of a continuous function f equalling some value y at the input $f(x)$.

Consider the very simple function $f(x) = x^2$. We know $f(3) = 3^2 = 9$. But since the function is continuous, it is also the case that if you want the output

to be close to 9, then all you have to do is get the input close to 3. This is true in a *quantitative* sense!

If you want to get $f(x)$ to be within 0.1 of 9, then you just need to get x within 0.01 of the input value 3. How do we know this? Well $2.99^2 = 8.9401$ and $3.01^2 = 9.0601$, so (since the function x^2 is nondecreasing for positive inputs) for any x in the range $(2.99, 3.01)$, the value of $f(x)$ is in the range $(8.9, 9.1)$.

To sum up the previous paragraph in one sentence, you could say that in order for $f(x)$ to get 0.1-close to 9, it suffices to pick any input x that is 0.01-close to 3. But since $f(3) = 9$ and f is continuous, we can do much better. To guarantee that $f(x)$ be 0.01-close to 9, we just need to ensure that x is within 0.0001 of 3. And in fact, for any small number $\epsilon > 0$, there exists some δ such that whenever $x \in (3 - \delta, 3 + \delta)$, we must have $f(x) \in (9 - \epsilon, 9 + \epsilon)$. (In math, we often use the Greek letters ϵ and δ to denote very small numbers.)

The quintessential non-example. Here is an example to illustrate what we *don't* mean when we are talking about a function getting arbitrarily close to a number when its input gets close to some number. Let

$$f(x) = \sin\left(\frac{\pi}{x}\right);$$

this function is defined everywhere except at $x = 0$. The question is, does it “approach” anything when $x = 0$? This function oscillates between -1 and 1 , but as x approaches 0 it oscillates very wildly. You might be fooled into thinking that as x approaches 0, the function f approaches 0. This is because

$$0 = \sin(\pi) = \sin(2\pi) = \sin(3\pi) = \cdots,$$

so $f(1) = 0$, $f(1/2) = 0$, $f(1/3) = 0$, and so on. But we also have

$$1 = \sin\left(\frac{\pi}{2}\right) = \sin\left(\frac{5\pi}{2}\right) = \sin\left(\frac{9\pi}{2}\right) = \cdots,$$

giving us $f(2) = 1$, $f(2/5) = 1$, $f(2/9) = 1$, etc., so this is a case of x approaching 0 but $f(x)$ *not* getting closer and closer to 0.

So the notion we end up defining for f getting close to something as x approaches something should not accidentally apply to this instance of f above. We are just about ready to state this notion formally.

The formal definition of a limit. First we define what a *neighbourhood* of a number is; it is simply some (open) interval containing that number. For example, the interval $(2.5, 3.5)$ is a neighbourhood of the number 3. So is $(2.9, 3.2)$, but $(10, 100)$ is not, and neither is $(1.5, 3)$, since we are considering open intervals.

Suppose that f is defined on all inputs in some neighbourhood of the number a , except possibly at a itself. (In other words, suppose that there is some neighbourhood of a contained in the domain of f , except possibly the point a .)

We say that the *limit* of the function f at the input a is equal to L , and write

$$\lim_{x \rightarrow a} f(x) = L,$$

if for every $\epsilon > 0$, there exists some $\delta > 0$ such that whenever $x \neq a$ and $x \in (a - \delta, a + \delta)$, we have $f(x) \in (L - \epsilon, L + \epsilon)$.

(If the use of the “element of” symbol $\in$ disturbs you, you can write the latter two expressions as $a - \delta < x < a + \delta$ and $L - \epsilon < f(x) < L + \epsilon$, respectively. Another way to write the definition of a limit is: For every $\epsilon > 0$ there is some $\delta > 0$ such that whenever $0 < |x - a| < \delta$, we have $|f(x) - L| < \epsilon$. Convince yourself that this means the exact same thing mathematically as the definition in terms of intervals that we had above—where is the condition $x \neq a$ encoded in the second definition?)

Because the formal definition of a limit is rather cumbersome, involving parameters ϵ and δ , in many cases we can make do with the following “simplified” or “intuitive” definition of a limit. *The limit of a function f at the input a is L if we can make $f(x)$ as close as we like to L by requiring that x be sufficiently close (but not necessarily equal to) a .*

In the example at the beginning of this section, we were convincing ourselves that

$$\lim_{x \rightarrow 3} x^2 = 9.$$

This should not be a mindblowing fact, since we of course have $3^2 = 9$. But we were careful in defining the concept of a limit in such a way that it now applies to many more situations as well, not just when $f(a)$ equals L .

A slightly harder example. Let us now show an example of a function f and an input a such that $\lim_{x \rightarrow a} f(x)$ exists and equals some number L , even when f is not defined at the input a itself. Suppose we let

$$f(x) = \frac{3x^2 - 2x - 1}{x - 1}.$$

Then we see that $f(x)$ is not defined at the input $x = 1$, since $1 - 1 = 0$ and dividing by 0 is illegal. Nevertheless, if you graph this function f using a graphing calculator or something like Desmos, you will see that the function f appears very simple. It is defined everywhere except at $x = 1$, and besides that little gap, its graph is simply a line; it is the unique line passing through the points $(0, 1)$ and $(2, 7)$. So even though f is not defined when $x = 1$, we might suspect from this picture that $\lim_{x \rightarrow 1} f(x) = 4$. Let us state this as a proposition and prove it rigorously (via the formal definition of a limit).

Proposition 1. *We have*

$$\lim_{x \rightarrow 1} \frac{3x^2 - 2x - 1}{x - 1} = 4.$$

Proof. Let $f(x) = (3x^2 - 2x - 1)/(x - 1)$ again for short. Suppose that $\epsilon > 0$ is given to us; we need to prescribe some $\delta > 0$ such that whenever $0 < |x - 1| < \delta$, we have $0 < |f(x) - 4| < \epsilon$. We shall show that picking $\delta = \epsilon/3$ works.

Here's why. Assume that $0 < |x - 1| < \delta = \epsilon/3$. Multiplying the second inequality by 3 gives us

$$|3x - 3| < \epsilon.$$

We want a term of -4 inside the absolute-value bars, so let's just add and subtract 1 in that expression to make that happen:

$$|3x + 1 - 4| < \epsilon \tag{1}$$

Now we're almost done—we would be done if we had $f(x)$ instead of $3x + 1$. But note that

$$3x^2 - 2x - 1 = (3x + 1)(x - 1),$$

so it's time to use the other assumption we had, which was $|x - 1| > 0$, to see that $(x - 1)/(x - 1) = 1$. In other words, under the assumption $|x - 1| > 0$, we have

$$3x + 1 = (3x + 1) \cdot 1 = \frac{(3x + 1)(x - 1)}{x - 1} = \frac{3x^2 - 2x - 1}{x - 1} = f(x).$$

Substituting this into (1) gives us $0 < |f(x) - 4| < \epsilon$, which is precisely what we wanted to show. ■

Picking $\delta = \epsilon/3$ seemed to require a great deal of algebraic ingenuity here; let's try to convince ourselves that it's not so mysterious a choice. If you draw the graph of f , you'll note that the slope near $x = 1$ is 3 (well it's actually 3 everywhere!). So to guarantee no more than ϵ error in the output, we need to be three times as precise in the input: we can have no more than $\epsilon/3$ error in the input. This leads to the choice $\delta = \epsilon/3$. Note that the proof would also work if we picked δ even smaller, e.g., $\delta = \epsilon/10$. But $\delta = \epsilon/3$ is the largest one can take δ for the proof to work, and it is a common stylistic choice for a mathematician to pick a maximal value such as this.

Before moving on, we note here that you may at some point encounter authors who write “ $f(x) \rightarrow L$ as $x \rightarrow a$.” This is just an alternative way of writing $\lim_{x \rightarrow a} f(x) = L$. We will try to stick with the $\lim_{x \rightarrow a}$ version in these notes, as that is the convention the textbook uses, but there is a strong argument for the $f(x) \rightarrow L$ notation as well, as it makes it clear that $f(x)$ is not required to actually equal L at any stage, it just needs to get arbitrarily close to it.

o4.IX **Formal negation of limit definition.** To fully understand the meaning of $\lim_{x \rightarrow a} f(x)$, it may be enlightening for us to consider the negation of this statement. Concretely, what does it mean for a number L *not* to be the limit of a function f at some input a ? Negating long mathematical statements (such as the definition of limit) properly is not an easy task, and the untrained mathematician is liable to make errors especially concerning quantifiers such as “for all” and “there exists.”

The correct formal negation is the following: *The real number L is not a limit of f at the input a if there exists some $\epsilon > 0$ such that for all $\delta > 0$, there is some x in the domain of f satisfying $0 < |x - a| < \delta$ with the property that $|f(x) - L| \geq \epsilon$.*

This is a fairly convoluted statement, but if you ponder it for a while you will convince yourself that it does mean the opposite of what the limit definition says. When we write $\lim_{x \rightarrow a} f(x) = L$, we mean that you can get the values of $f(x)$ arbitrarily close to L by taking x sufficiently close to a , and the opposite of this statement is there is some fixed positive distance ϵ such that even if you take x arbitrarily close to a , there will always be some badly-behaved choices of x that result in $f(x)$ being too far (at a distance at least ϵ) from $f(x)$.

(Note that the negation of the mathematical statement $\lim_{x \rightarrow a} f(x) = L$ should not be recorded as $\lim_{x \rightarrow a} f(x) \neq L$, since even writing down the left-hand side of this inequality suggests that it exists (is a real number), and as we shall see later in this section, the limit of a function f need not always exist at some given a .)

Here's an example to illustrate how you disprove the claim that the limit of a function f at some a is L .

Proposition 2. *Let*

$$f(x) = \begin{cases} -1, & \text{if } x < 0; \\ 1, & \text{if } x > 0 \end{cases}.$$

Then the limit of f at 0 is not 0.

Proof. We just need to find some $\epsilon > 0$ such that for any choice of $\delta > 0$, there will always be some real number x with $0 < |x - 0| < \delta$ but $|f(x) - 0| \geq \epsilon$. It turns out that choosing $\epsilon = 1/2$ works.

To see this, let $\delta > 0$ be given. We need to pick some x with $|x - 0| < \delta$ but with $|f(x) - 0| \geq 1/2$. This is not so hard—we have lots of choices! Choosing $x = \delta/2$ works, since $f(x) = f(\delta/2) = 1$, and $|1| \geq 1/2$.

Hence the limit of f at 0 is not 0. ■

***A game with the Adversary.** Let f be a function, let L be a number, and let a be a number such that f is defined in some neighbourhood of a . The statement “ $\lim_{x \rightarrow a} f(x) = L$ ” is either true or false (the “law of the excluded middle” tells us that every mathematical statement is either true or false). We can interpret its truthhood or falsity as the result of a game played between ourselves and a supernatural Adversary. Here's the how the game goes.

Game L (*The limit game*). Let f be a function, L a number, and a a number with a neighbourhood contained in the domain of f .

L1. [Adversary's turn.] The Adversary is allowed to pick any $\epsilon > 0$.

L2. [Our turn.] We can then pick any $\delta > 0$.

L3. [Adversary's turn.] The Adversary is then allowed to pick any x in the domain of f that satisfies

$$0 < |x - a| < \delta.$$

(This means that x is at distance less than δ from a , but doesn't equal a .)

L4. [Verdict?] If

$$|f(x) - L| < \epsilon,$$

that is, if the distance between $f(x)$ and L is less than ϵ , then we win; otherwise the Adversary wins. ■

If $\lim_{x \rightarrow a} f(x) = L$ is true, then we can always win; the very definition of limit tells us that no matter what ϵ is chosen in step L1 by the Adversary, in step L2 there is always a choice of δ that will force the Adversary into letting us win in step L3. On the other hand, if $\lim_{x \rightarrow a} f(x) = L$ is false (either because this limit does not exist or because it equals something else), then the Adversary can always win. There is a choice of ϵ in step L1 such that no matter what we pick in step L2, the Adversary can beat us by making a further nefarious choice for x in step L3.

Nonexistent limits. We alluded earlier to the possibility of a function f not having any limits at some point a . We say that the limit of f at a *does not exist* if for all real numbers L , the limit of f at a is not L . In fact, the function f from the previous proposition has this property at $a = 0$. The proof is largely unchanged from the easier case above when we had just set L to 0, requiring only a little more care in the second paragraph.

Proposition 3. For $x \neq 0$, let

$$f(x) = \frac{|x|}{x}.$$

Then the limit of f as x approaches 0 does not exist.

Proof. We can re-express f by writing

$$f(x) = \begin{cases} -1, & \text{if } x < 0; \\ 1, & \text{if } x > 0 \end{cases},$$

which may be clearer to some readers (and also shows this is the same function from Proposition 2). Let L be arbitrary. We need to show that the limit of f at 0 is not L . To do this, we shall pick $\epsilon > 0$ such that for any choice of $\delta > 0$, there is some real number x with $0 < |x - 0| < \delta$ but $|f(x) - L| \geq \epsilon$. Just as in the previous proof for the special case $L = 0$, the choice $\epsilon = 1/2$ works.

Let $\delta > 0$ be given. We need to pick some x with $|x - 0| < \delta$ but with $|f(x) - L| \geq 1/2$. Here it matters whether L is positive or negative (if $L = 0$ then just appeal to the proof of Proposition 2). If L is positive, then choose $x = -\delta/2$. We have $0 < |-\delta/2 - 0| < \delta$ as required, and also

$$|f(-\delta/2) - L| = |-1 - L| > |-1 - 0| \geq 1/2 = \epsilon.$$

On the other hand, if L is negative, we instead choose $x = \delta/2$, so we still have $0 < |\delta/2 - 0| < \delta$ as needed, and this time

$$|f(\delta/2) - L| = |1 - L| > |1 - 0| \geq 1/2 = \epsilon.$$

Either way, we have satisfied the criterion of the limit at f at a not being L .

But the previous paragraph works for all numbers L , so the limit of f at a cannot equal any number! Hence we have proved that it does not exist. ■

In the example just given, we now know that the limit of f at 0 does not exist. However, we often think of a limit intuitively as the function “approaching” some value, and in this sense it’s wrong to say that f doesn’t approach anything at all when the input gets close to 0. When you approach from the right, it is equal to 1. And when you approach from the left, the function equals -1 . This leads us to the concept of a one-sided limit.

One-sided limits. Let f be a function, let L be a number, and let a be a number such that f is defined to the right of a . (More precisely, we assume that f is defined on the interval $(a, a + \eta)$ for some $\eta > 0$.) We say that *limit of $f(x)$ as x approaches a from the right is L* , and write

$$\lim_{x \rightarrow a^+} f(x) = L,$$

if for all $\epsilon > 0$ there is $\delta > 0$ such that any number x in the domain of f with

$$a < x < a + \delta$$

must satisfy

$$|f(x) - L| < \epsilon.$$

We might say in this case that L is the *right-hand limit* of f .

Likewise, let f be a function, L be a number, and suppose that a is such that f is defined to the *left* of a . We say that *limit of $f(x)$ as x approaches a from the left is L* , and write

$$\lim_{x \rightarrow a^-} f(x) = L,$$

if for all $\epsilon > 0$ there is $\delta > 0$ such that any number x in the domain of f with

$$a - \delta < x < a$$

must satisfy

$$|f(x) - L| < \epsilon.$$

If this is true then we say L is the *left-hand limit* of f .

More loosely, L is the right-hand limit of f at a if you can get $f(x)$ to be arbitrarily close to L by requiring that x be sufficiently close to a and *greater than a* , and it’s the left-hand limit of f at a if you can get $f(x)$ as close as you like to L by requiring that x be sufficiently close to a and *smaller than a* .

Both of these definitions are relaxations of the definition of limit; it is “easier” to be a left-hand limit than it is to be a *bona fide* limit). For example, we saw in Proposition 3 that 1 is not a limit of f from that proposition, because

f has no limits. However, once you have digested the definitions of one-sided limits, you should be able to check that 1 is the right-hand limit of f at 0, and -1 is the left-hand limit at 0.

If the actual limit of f at a point a is L , then it is a consequence of the definitions that the right-hand limit and the left hand limit are also L . Conversely, if the one-sided limits both equal the same L , then the limit is actually L . This can be summed up symbolically in the following statement, which we'll call a theorem since it comes in handy now and again.

Theorem 4. *For any function f and numbers a and L , we have*

$$\lim_{x \rightarrow a} f(x) = L$$

if and only if

$$\lim_{x \rightarrow a^+} f(x) = L \quad \text{and} \quad \lim_{x \rightarrow a^-} f(x) = L. \quad \blacksquare$$

A simple application of Theorem 4 is to prove that the function $f(x) = |x|$ goes to 0 as x approaches 0. We'll see this as an example in the next section, but it's really not a difficult proof, so you could try it now. Show that both one-sided limits approach 0, then simply invoke the theorem.

Here's a related theorem (no less useful) that tells us what happens if the one-sided limits of a function do not agree at some given point. It shows that the example exhibited by Proposition 3 is an instance of a very general phenomenon.

Theorem 5. *If*

$$\lim_{x \rightarrow a^+} f(x) = L, \quad \lim_{x \rightarrow a^-} f(x) = M,$$

and $L \neq M$, then $\lim_{x \rightarrow a} f(x)$ does not exist.

Proof. We skip this proof as it's beyond the scope of this course (you would see it in, e.g., MATH 242 or MATH 254), but it is not so different in spirit to the proof of Proposition 3. The motivated student is encouraged to try it as an advanced exercise. (If you try for a while and get stuck, email me or come to office hours with what you have and I could help you fill in the details.) $\blacksquare$

If we were told Theorem 5 from the start, we could have used it to prove Proposition 3 without much ado.

Proof of Proposition 3 using Theorem 5. Recall that we are trying to prove that the limit of the function $f(x) = |x|/x$ does not exist as x approaches 0. Since $|x| = x$ for all $x > 0$, we see that

$$\lim_{x \rightarrow 0^+} \frac{|x|}{x} = \lim_{x \rightarrow 0^+} \frac{x}{x} = \lim_{x \rightarrow 0^+} 1 = 1.$$

(The last equality is easy to see if you just graph the constant function 1. If you are a serious student and insist on a rigorous proof, then please do Exercise 24 of

Section 2.4 in the textbook, which invites you to furnish a proof that $\lim_{x \rightarrow a} c = c$ for all numbers a and c .)

On the other hand, since $|x| = -x$ for all $x < 0$, we have

$$\lim_{x \rightarrow 0^-} \frac{|x|}{x} = \lim_{x \rightarrow 0^-} \frac{-x}{x} = \lim_{x \rightarrow 0^+} -1 = -1.$$

So the left- and right-hand limits do not agree, and Theorem 5 applies in this situation to tell us that the limit does not exist. ■

One-sided limits not agreeing is one way that the limit of a function at a given point can fail to exist. Another way is for the limit to get arbitrarily large, which is the situation we tackle next.

09.IX **Infinite limits and vertical asymptotes.** We say that the *limit of $f(x)$ is infinity as x approaches a from the right*, and write

$$\lim_{x \rightarrow a^+} f(x) = \infty,$$

if for every large (positive) number M , there exists $\delta > 0$ such that every x satisfying

$$a < x < a + \delta$$

has

$$f(x) > M.$$

If, instead, for every large positive number M there is $\delta > 0$ such that every x with

$$a < x < a + \delta$$

satisfies

$$f(x) < -M,$$

then we say that the limit of $f(x)$ is *minus infinity* as x approaches a from the right, and write

$$\lim_{x \rightarrow a^-} f(x) = -\infty.$$

The same definition can be made for left-hand limits by replacing the assumption $a < x < a + \delta$ with the assumption $a - \delta < x < a$ above. This makes for four situations in total, and if any one of these situations occurs, then f is said to have a *vertical asymptote* at the line $x = a$.

Just as in the case of finite limits, when the left- and right-hand limits agree (and are both the same type of infinity), then we simply say that the *limit* is infinity (or minus infinity, accordingly). In other words, we have

$$\lim_{x \rightarrow a} f(x) = \infty$$

if and only if

$$\lim_{x \rightarrow a^-} f(x) = \infty \quad \text{and} \quad \lim_{x \rightarrow a^+} f(x) = \infty,$$

and we have

$$\lim_{x \rightarrow a} f(x) = -\infty$$

if and only if

$$\lim_{x \rightarrow a^-} f(x) = -\infty \quad \text{and} \quad \lim_{x \rightarrow a^+} f(x) = -\infty.$$

(Let p and q be mathematical statements. The phrase “ p if q ” means that q implies p , and the phrase “ p only if q ” means that p implies q —think about it for a second, it makes sense! So the statement “ p if and only if q ” means that p is true exactly when q is true, and vice versa.)

One should not interpret the symbol ∞ above as a number, since there is no such number. When we write $\lim_{x \rightarrow a} f(x) = \infty$, we simply mean that this expression can be made arbitrarily large as by requiring that x be sufficiently close to a . It should also be stressed that the equality symbol in this expression should *not* be interpreted as meaning that the limit of f at a exists, because the terminology of a limit “existing” is reserved for when the limit equals a number (and, once again, ∞ is not a number). As with finite limits, sometimes authors write, e.g., $f(x) \rightarrow -\infty$ as $x \rightarrow a$ instead of $\lim_{x \rightarrow a} f(x) = -\infty$. This notation is also valid, and in some sense preferable as it does not suggest that $f(x)$ ever “equals” some “number” ∞ .

Let’s look at a couple of simple examples of functions with vertical asymptotes.

Proposition 6. *We have*

$$\lim_{x \rightarrow 0^+} \frac{1}{x} = \infty$$

and

$$\lim_{x \rightarrow 0^-} \frac{1}{x} = -\infty.$$

Proof. We prove both statements separately. The first asserts that $1/x$ can be made arbitrarily large by taking $x \rightarrow 0$ from the right. To prove it, let $M > 0$ be an arbitrarily large number (given to us). Pick $\delta = 1/M$, and assume that $0 < x < \delta = 1/M$. Because $x > 0$, we are allowed to write $1/x$, and the fact that $x < 1/M$ implies that $1/x > M$, which is what we needed to prove in the definition of infinite limit.

The second statement is proved extremely similarly, but we include the full proof for completeness. Let M be given, and pick $\delta = 1/M$. Assume that $-\delta < x < 0$; the right inequality here assures us that the expression $1/x$ is well-defined, and the left inequality here can be rewritten as $x > -\delta = -1/M$. Rearranging the equation now yields $1/x < -M$, which is what we needed to show in order to prove that the limit is $-\infty$. ■

To sum up this proof, we can make $1/x$ arbitrarily large to the right of 0 by requiring that $x > 0$ be sufficiently close to 0, and we can make $1/x$ an arbitrarily

large negative number by requiring that $x < 0$ be sufficiently close to (the left of) 0.

Here's another example where the left- and right-hand limits agree on the type of infinity.

Proposition 7. *The limit of*

$$f(x) = \frac{1}{(x-3)^2}$$

is infinity as x approaches 3.

Proof. We can do the left- and right-hand limits separately, but it is not hard to prove this whole thing all at once (i.e., to consider both sides of 3 at the same time).

Let $M > 0$ be given. We want to prescribe $\delta > 0$ such that all $x \neq 3$ with

$$3 - \delta < x < 3 + \delta$$

satisfy

$$\frac{1}{(x-3)^2} > M.$$

(The assumption $3 - \delta < x < 3 + \delta$ is what we mean by considering both sides of 3 at the same time; note that this assumption together with the assumption that $x \neq 3$ can be summed up in the single inequality $0 < |x - 3| < \delta$.) Setting $\delta = 1/\sqrt{M}$ works this time, so our assumption on x will be

$$3 - \frac{1}{\sqrt{M}} < x < 3 + \frac{1}{\sqrt{M}},$$

and that $x \neq 3$ of course, so that $f(x)$ is even defined. The difference $x - 3$ may be positive or negative; we don't know which. But what we do know, from our assumption, is that the distance $|x - 3|$ between x and 3 satisfies

$$|x - 3| < \frac{1}{\sqrt{M}}.$$

Squaring both sides (and using the fact that $|x - 3|^2 = (x - 3)^2$, since the square of a negative number is positive anyway) gives us

$$(x - 3)^2 < \frac{1}{M},$$

and rearranging yields

$$f(x) = \frac{1}{(x-3)^2} > M,$$

which is what we needed to show. (The number M was arbitrary in this proof, so we can keep repeating with larger and larger M to show that $f(x)$ can be made arbitrarily large in the way shown above.) ■

Once again, picking $\delta = 1/\sqrt{M}$ made the algebra work out nicely, but you could have written a valid proof by picking, say, $\delta = 1/M$ instead, but then there'd be some issues with having to assume that $M > 1$. (Morally this is O.K., since we're only worried about what happens when M is very large, but *technically* our definition had $M > 0$.) Don't stress too much about picking the "best" value for δ . Generally, you find δ by first trying out various values of M and seeing what δ is necessary to deal with that M ; after a few tries you should be able to come up with a formula that works generally. The broader thing to take away from reading this proof or trying to do it yourself is that no matter what M is given, you can pick *some* δ that makes the proof work out.

11.IX **Limits at infinity and horizontal asymptotes.** There is one last type of limit that remains to be defined. This is the case in which the values of $f(x)$ can be taken arbitrarily close to a value by making x sufficiently large, either in the positive direction or the negative direction. Formally, we write

$$\lim_{x \rightarrow \infty} f(x) = L$$

and say that the *limit of $f(x)$ is L as x approaches infinity* if for all $\epsilon > 0$ there exists some large $M > 0$ such that every x with $x > M$ has $|f(x) - L| < \epsilon$. Likewise, we say the *limit of $f(x)$ is L as x approaches negative infinity* and write

$$\lim_{x \rightarrow -\infty} f(x) = L$$

if the same condition applies, except the inequality $x > M$ is replaced by $x < -M$. In either of these situations, we say that $y = f(x)$ has a *horizontal asymptote* at the line $y = L$.

Note that the same example from Proposition 6 also exhibits a horizontal asymptote.

Proposition 8. *We have*

$$\lim_{x \rightarrow \infty} 1/x = 0 \quad \text{and} \quad \lim_{x \rightarrow -\infty} 1/x = 0.$$

Proof. Let $\epsilon > 0$ and pick $M = 1/\epsilon$. Then any number x with $x > M$ has

$$\left| \frac{1}{x} - 0 \right| = \frac{1}{x} < \frac{1}{M} = \epsilon,$$

so $\lim_{x \rightarrow \infty} 1/x = 0$.

For the other inequality, the proof is very similar. Let $\epsilon > 0$ and again we choose $M = 1/\epsilon$. Now any (negative) number x with $x < -M$ has

$$\left| \frac{1}{x} - 0 \right| = \left| \frac{1}{x} \right| = \frac{1}{-x} < \frac{1}{M} = \epsilon,$$

so $\lim_{x \rightarrow -\infty} 1/x = 0$. ■

Since ∞ is not a number, it does not make sense to write an expression such as, say, $1/\infty$. On the other hand, you may have heard people say, informally, something like “anything divided by infinity is zero.” This statement can be made into a proper mathematical one by expressing it in terms of limits. It is true that for any fixed number K , the limit $\lim_{x \rightarrow \infty} K/x$ equals 0. (As an exercise, prove this directly by very slightly modifying the proof of Proposition 8 above. Hint: The proposition already gives a proof when $K = 1$. So for the general case, instead of setting $M = 1/\epsilon$, you might try choosing something very similar but involving K .)

Infinite limits at infinity. As with limits as x approaches some finite a , the infinite limit is said not to exist if it does not equal any finite number L . There is one special case of this worth mentioning, which is when the function $f(x)$ can be made arbitrarily large (positive or negative) by requiring that x be sufficiently large (positive or negative). Hence there are four cases of infinite limits at infinity, namely

$$\begin{aligned} \lim_{x \rightarrow \infty} f(x) = \infty, & \quad \lim_{x \rightarrow -\infty} f(x) = \infty, \\ \lim_{x \rightarrow \infty} f(x) = -\infty, & \quad \text{and} \quad \lim_{x \rightarrow -\infty} f(x) = -\infty. \end{aligned}$$

Tedious as it is, let us formally define all four of these statements, for the sake of completeness.

a) We say that $f(x)$ goes to infinity as x goes to infinity and write

$$\lim_{x \rightarrow \infty} f(x) = \infty$$

if for all $M > 0$, there exists $N > 0$ such that for all x with $x > N$, we have $f(x) > M$.

b) We say that $f(x)$ goes to infinity as x goes to negative infinity and write

$$\lim_{x \rightarrow -\infty} f(x) = \infty$$

if for all $M > 0$, there exists $N > 0$ such that for all x with $x < -N$, we have $f(x) > M$.

c) We say that $f(x)$ goes to negative infinity as x goes to infinity and write

$$\lim_{x \rightarrow \infty} f(x) = -\infty$$

if for all $M > 0$, there exists $N > 0$ such that for all x with $x > N$, we have $f(x) < -M$.

d) We say that $f(x)$ goes to negative infinity as x goes to negative infinity and write

$$\lim_{x \rightarrow -\infty} f(x) = -\infty$$

if for all $M > 0$, there exists $N > 0$ such that for all x with $x < -N$, we have $f(x) < -M$.

That was a mouthful. But by this point it is hoped that the reader would be comfortable enough with these formal definitions to be able to formulate, say (a), (c), and (d) after being shown (b).

We finish off this very long section by presenting an example of a function that seems to grow fairly slowly $x \rightarrow \infty$, but nevertheless still goes to infinity as x is taken to be large.

Proposition 9. *We have*

$$\lim_{x \rightarrow \infty} \sqrt[3]{x} = \infty$$

and

$$\lim_{x \rightarrow -\infty} \sqrt[3]{x} = -\infty.$$

Proof. Let $M > 0$ be a large positive number given to us. We want to pick N such that for all $x > N$, we have $\sqrt[3]{x} = x^{1/3} > M$. After a moment's pause we see that choosing $N = M^3$ works, since if we assume that $x > N$, then

$$x^{1/3} > N^{1/3} = (M^3)^{1/3} = M.$$

The negative infinite limit is proved similarly; we choose $N = M^3$ this time as well, so that whenever $x < -N$, we have

$$x^{1/3} < (-N)^{1/3} = (-M^3)^{1/3} = -M.$$

This completes the proof. ■

3. Rules for evaluating limits

In the previous section, we already encountered two rules (namely, Theorems 4 and 5) that can help us find limits of functions or determine that they do not exist. This section will be devoted to establishing a few more useful laws. First we record a list of algebraic laws, which show that the limit behaves well under a number of algebraic operations such as addition, multiplication, etc.

Theorem 10 (*Algebraic limit laws*). *Let c be any fixed number, and suppose that f and g are functions such that*

$$\lim_{x \rightarrow a} f(x) \quad \text{and} \quad \lim_{x \rightarrow a} g(x)$$

both exist. Then

- i) $\lim_{x \rightarrow a} (f(x) + g(x)) = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x);$
- ii) $\lim_{x \rightarrow a} (f(x) - g(x)) = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x);$
- iii) $\lim_{x \rightarrow a} cf(x) = c \cdot \lim_{x \rightarrow a} f(x);$

- iv) $\lim_{x \rightarrow a} (f(x)g(x)) = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x)$;
- v) $\lim_{x \rightarrow a} f(x)/g(x) = \lim_{x \rightarrow a} f(x) / \lim_{x \rightarrow a} g(x)$, provided that $\lim_{x \rightarrow a} g(x) \neq 0$;
- vi) for all positive integers n , $\lim_{x \rightarrow a} f(x)^n = (\lim_{x \rightarrow a} f(x))^n$; and
- vii) for all positive integers n , $\lim_{x \rightarrow a} \sqrt[n]{f(x)} = \sqrt[n]{\lim_{x \rightarrow a} f(x)}$, (provided that $\lim_{x \rightarrow a} f(x) > 0$ in the case that n is even, so that the n th root is well-defined).

These statements are all true even when $x \rightarrow a$ is replaced with $x \rightarrow \infty$, $x \rightarrow -\infty$, $x \rightarrow a^+$, or $x \rightarrow a^-$.

The identities above are known as (i) the sum law, (ii) the difference law, (iii) the constant multiple law, (iv) the product law, (v) the quotient law, (vi) the power law, and (vii) the root law, respectively. They can be expressed in words as well: the limit of a sum is the sum of the limits, the limit of a difference is the difference of the limits, and so on.

14.IX *Proof.* We only prove (i) in these notes. Laws (ii) through (v) are proved in Appendix F of the textbook. Law (vi) is a consequence of repeating law (iv) n times (formally, you would need to use the *principle of mathematical induction*, which is outside the scope of this course), and the proof of law (vii) is assigned to you in the textbook as Exercise 69 of Section 2.5. All of these proofs only deal with the case that $x \rightarrow a$ for some finite number a , as opposed to $x \rightarrow \infty$; it is an instructive exercise to try and adapt the proof below for the $x \rightarrow \infty$ case, to test that you have fully digested this proof and the definition of limits at infinity.

Enough talk. On to the proof of (i). We shall make use of the *triangle inequality*, which says that for any numbers a , b , and c , we have

$$|a - c| \leq |a - b| + |b - c|.$$

In other words, the length of any side in a triangle is always smaller than the sum of the other two sides.

Suppose $\lim_{x \rightarrow a} f(x) = L_1$ and $\lim_{x \rightarrow a} g(x) = L_2$. We want to show that the limit of $f(x) + g(x)$ as $x \rightarrow a$ is $L_1 + L_2$. Let $\epsilon > 0$. We wish to pick a $\delta > 0$ such that for all x with $0 < |x - a| < \delta$, we have

$$|f(x) + g(x) - (L_1 + L_2)| < \epsilon.$$

Well, since $\lim_{x \rightarrow a} f(x) = L_1$, we can make $f(x)$ arbitrarily close to L_1 by prescribing that x be sufficiently close to a . So there is some $\delta_1 > 0$ such that if we assume $0 < |x - a| < \delta_1$, then

$$|f(x) - L_1| < \frac{\epsilon}{2}.$$

(We choose $\epsilon/2$ so that there is some additional room for error when we deal with g .) Likewise, there is some $\delta_2 > 0$ such that if we assume $0 < |x - a| < \delta_2$, then

$$|g(x) - L_2| < \frac{\epsilon}{2}.$$

Now, set $\delta = \min\{\delta_1, \delta_2\}$. This guarantees both $\delta \leq \delta_1$ and $\delta \leq \delta_2$, so that if we assume $0 < |x - a| < \delta$, then the inequalities

$$0 < |x - a| < \delta_1 \quad \text{and} \quad 0 < |x - a| < \delta_2$$

both hold. As a result, by our observations in the previous paragraph, both of the inequalities

$$|f(x) - L_1| < \frac{\epsilon}{2} \quad \text{and} \quad |g(x) - L_2| < \frac{\epsilon}{2}$$

are true as well. Finally we see that

$$\begin{aligned} |f(x) + g(x) - (L_1 + L_2)| &= |f(x) - L_1 + g(x) - L_2| \\ &\leq |f(x) - L_1| + |g(x) - L_2| \\ &\leq \frac{\epsilon}{2} + \frac{\epsilon}{2} \\ &= \epsilon, \end{aligned}$$

where the second line follows from the aforementioned triangle inequality. This completes the proof. ■

These laws are extremely powerful in general, allowing us to express the limit of a complex expression in terms of the limits of simpler expressions. For example, using laws (i), (iii), and (iv), we can deduce that

$$\lim_{x \rightarrow 3} 4x^2 - 7 = 4 \cdot \left(\lim_{x \rightarrow 3} x \right) \left(\lim_{x \rightarrow 3} x \right) - \lim_{x \rightarrow 3} 7. \quad (2)$$

We already encountered an expression of the form $\lim_{x \rightarrow a} c$, where c is a constant number, in an earlier proof. It should be obvious that this limit is c , and that $\lim_{x \rightarrow a} x = a$. Let us state these two facts, along with two more as a theorem, so that we can easily invoke them later on. (We continue the numbering from Theorem 10 in order to remain consistent with the textbook.)

Theorem 11 (*Special limits*). For any positive integer n and any numbers a and c ,

viii) $\lim_{x \rightarrow a} c = c$;

ix) $\lim_{x \rightarrow a} x = a$;

x) $\lim_{x \rightarrow a} x^n = a^n$; and

xi) $\lim_{x \rightarrow a} \sqrt[n]{x} = \sqrt[n]{a}$.

Laws (viii) and (ix) remain true when $x \rightarrow a$ is replaced by $x \rightarrow \infty$ or $x \rightarrow -\infty$.

Proof. As mentioned before, the proof of law (viii) is presented to you as Exercise 24 in Section 2.4; the proof of law (ix) is Exercise 23 in the same section. This is in the case where x approaches some finite a ; you might like to prove the corresponding limits at infinity to test your understanding of those definitions.

Laws (x) and (xi) are easy to prove once we have all the other laws established. To prove law (x), apply the power law with $f(x) = x$ and then law (ix) to obtain

$$\lim_{x \rightarrow a} x^n = \left(\lim_{x \rightarrow a} x \right)^n = a^n.$$

To prove law (xi), apply the root law with $f(x) = x$ and then law (ix) to obtain

$$\lim_{x \rightarrow a} \sqrt[n]{x} = \sqrt[n]{\lim_{x \rightarrow a} x} = \sqrt[n]{a}. \quad \blacksquare$$

With these new laws in hand, we can revisit the polynomial $4x^2 - 7$ from earlier and conclude that

$$\lim_{x \rightarrow 3} (4x^2 - 7) = 4 \cdot 3^2 - 7 = 29.$$

In fact, we don't even need to break up x^2 into $x \cdot x$ like we did in (2), as law (x) takes care of monomials such as these.

When applying the limit laws to determine what happens when $x \rightarrow 3$ to a simple polynomial such as $4x^2 - 7$, it seemed like we could just substitute in $x = 3$ in the expression and get the right answer. This turns out to be true in the general case $x \rightarrow a$ as long as a is in the domain of f .

Theorem 12 (*Direct substitution*). *Let f be a polynomial or a rational function (a quotient of two polynomials). If a is in the domain of f , then*

$$\lim_{x \rightarrow a} f(x) = f(a).$$

Proof. Every rational function f is of the form

$$f(x) = \frac{a_m x^m + a_{m-1} x^{m-1} + \cdots + a_1 x + a_0}{b_n x^n + b_{n-1} x^{n-1} + \cdots + b_1 x + b_0},$$

where m and n are nonnegative integers, and $a_0, a_1, \dots, a_m, b_0, b_1, \dots, b_n$ are real (possibly negative or zero) coefficients. Taking the limit of both sides, the quotient law deals with the fraction, the sum law splits the limit of the sum into the sum of limits, and from there we use law (x) and the constant multiple law to break up these terms further. We then apply laws (viii) and (ix) as to finish off the proof. $\blacksquare$

This theorem applies to functions involving roots as well, thanks to the root law. (It more generally applies to all *continuous* functions—we will discuss the

notion of continuity very soon.) Using the direct substitution rule is fairly easy whenever its hypotheses are satisfied, so we will not dwell on this. There is, however, a more advanced version of this rule that is useful in certain cases, such as when $f(x)$ is a rational function and a is *not* in the domain of f (e.g., because plugging in $x = a$ makes the denominator 0). Then we may have to invoke the following theorem in order to use the direct substitution rule.

Theorem 13. *Let a be a number, and suppose that f and g are two functions that are equal for all x in some interval containing a , except possibly at a itself. Then if $\lim_{x \rightarrow a} f(x)$ and $\lim_{x \rightarrow a} g(x)$ both exist, we must have*

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x).$$

**Proof.* This proof is a bit more advanced than you should fairly be expected to follow in this course, though if you take a second-year mathematics class such as MATH 242 one day, you'd find it a routine exercise by the end of that course. We include it in these notes for completeness.

We know that f and g agree in some interval containing a , except possibly at a itself. To be concrete, this means there exists some number $\eta > 0$ be such that $f(x) = g(x)$ for every x with $0 < |x - a| < \eta$. Suppose $\lim_{x \rightarrow a} f(x) = L$ and $\lim_{x \rightarrow a} g(x) = M$. We want to show that $L = M$; we shall show that $|L - M| < \epsilon$ for all $\epsilon > 0$, which is an equivalent statement (a nonnegative quantity that is less than any positive quantity must be zero).

Let $\epsilon > 0$. We will use an “ $\epsilon/2$ trick” again, just as in the proof of the sum law above. Since $\lim_{x \rightarrow a} f(x) = L$, there is δ_1 such that for all x with $0 < |x - a| < \delta_1$, we have

$$|f(x) - L| < \epsilon/2. \quad (3)$$

Likewise, since $\lim_{x \rightarrow a} g(x) = M$, there is δ_2 such that for all x with $0 < |x - a| < \delta_2$, we have

$$|g(x) - M| < \epsilon/2. \quad (4)$$

Now pick $\delta = \min\{\eta, \delta_1, \delta_2\}$, so that every x with $0 < |x - a| < \delta$ has $f(x) = g(x)$ and satisfies both (3) and (4). We now combine all three of these facts and invoke the triangle inequality to obtain

$$\begin{aligned} |L - M| &= |L - g(x) + g(x) - M| \\ &\leq |L - g(x)| + |g(x) - M| \\ &= |f(x) - L| + |g(x) - M| \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} \\ &= \epsilon. \end{aligned}$$

As we said above, since this proof shows that $|L - M| < \epsilon$ for every $\epsilon > 0$, we must have $|L - M| = 0$; that is, $L = M$. ■

If we'd had these theorems from the start, we wouldn't have had to use so much elbow grease in proving Proposition 1. Here's how we could have proceeded.

Proof of Proposition 1 using Theorems 12 and 13. Let

$$f(x) = \frac{3x^2 - 2x - 1}{x - 1};$$

recall that we are trying to show that $\lim_{x \rightarrow 1} f(x) = 4$. We factor $3x^2 - 2x - 1 = (3x + 1)(x - 1)$ and observe that for any $x \neq 1$,

$$f(x) = \frac{3x^2 - 2x - 1}{x - 1} = 3x + 1.$$

Letting $g(x) = 3x + 1$, on any open interval containing 1 we have $f(x) = g(x)$ everywhere except at $x = 1$, so Theorem 13 applies to inform us that $\lim_{x \rightarrow 1} f(x) = \lim_{x \rightarrow 1} g(x)$. Moreover, $g(x)$ is a polynomial with $x = 1$ in its domain, so

$$\lim_{x \rightarrow 1} g(x) = 3 \cdot 1 + 1 = 4$$

by Theorem 12. ■

Here's another example in which we use the root law.

Proposition 14. *We have*

$$\lim_{x \rightarrow 0} \frac{\sqrt{x^2 + 4} - 2}{x^2} = \frac{1}{4}.$$

Proof. We cannot use the quotient law, since $x^2 \rightarrow 0$ as $x \rightarrow 0$. If we can massage the expression to obtain a factor of x^2 in the numerator, then (in an open interval containing 0 but not at 0 itself), we can cancel the x^2 in the denominator and perhaps make some headway. This strategy is facilitated by the observation that

$$\left(\sqrt{x^2 + 4} - 2\right)\left(\sqrt{x^2 + 4} + 2\right) = \left(\sqrt{x^2 + 4}\right)^2 - 2^2 = x^2.$$

Hence we have

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{\sqrt{x^2 + 4} - 2}{x^2} &= \lim_{x \rightarrow 0} \frac{\sqrt{x^2 + 4} - 2}{x^2} \cdot \frac{\sqrt{x^2 + 4} + 2}{\sqrt{x^2 + 4} + 2} \\ &= \lim_{x \rightarrow 0} \frac{x^2}{x^2(\sqrt{x^2 + 4} + 2)} \\ &= \lim_{x \rightarrow 0} \frac{1}{\sqrt{x^2 + 4} + 2}. \end{aligned}$$

We no longer have the issue of the limit of the denominator being 0, and can happily apply the algebraic limit laws as well as the direct substitution property:

$$\begin{aligned}
 \lim_{x \rightarrow 0} \frac{1}{\sqrt{x^2 + 4} + 2} &= \frac{1}{\lim_{x \rightarrow 0} (\sqrt{x^2 + 4} + 2)} \\
 &= \frac{1}{\lim_{x \rightarrow 0} \sqrt{x^2 + 4} + \lim_{x \rightarrow 0} 2} \\
 &= \frac{1}{\sqrt{\lim_{x \rightarrow 0} (x^2 + 4)} + 2} \\
 &= \frac{1}{\sqrt{4} + 2} \\
 &= \frac{1}{4} \quad \blacksquare
 \end{aligned}$$

Using one-sided limits. Recall Theorem 4, which states that a limit exists and equals some L if and only if both one-sided limits exist and equal L . This can be used to compute $\lim_{x \rightarrow 0} |x|$, even though the algebraic limit laws do not apply to the function $|x|$ directly.

Proposition 15. *We have*

$$\lim_{x \rightarrow 0} |x| = 0.$$

Proof. Towards the invocation of Theorem 4, we calculate the one-sided limits of $|x|$ using the direct substitution rule. When $x > 0$, we have $|x| = x$, so

$$\lim_{x \rightarrow 0^+} |x| = \lim_{x \rightarrow 0^+} x = 0,$$

since for all $x < 0$ we instead have $|x| = -x$, so

$$\lim_{x \rightarrow 0^-} |x| = \lim_{x \rightarrow 0^-} -x = 0.$$

Both one-sided limits agree, so we conclude that the limit is 0. $\blacksquare$

Evaluating limits at infinity. Recall that most of the algebraic limit laws established earlier in this section apply to limits at infinity. Hence we can combine the root and power laws with Proposition 8 to prove the following useful theorem.

Theorem 16. *Let $r > 0$ be a rational number. Then*

$$\lim_{x \rightarrow \infty} \frac{1}{x^r} = 0.$$

If $r > 0$ is such that x^r is defined for negative values of x as well, then

$$\lim_{x \rightarrow -\infty} \frac{1}{x^r} = 0.$$

Proof. Let $r > 0$ be a rational number. This means there are positive integers m and n such that $r = m/n$, so we can write

$$\lim_{x \rightarrow \infty} \frac{1}{x^r} = \lim_{x \rightarrow \infty} \sqrt[n]{\left(\frac{1}{x}\right)^m} = \sqrt[n]{\left(\lim_{x \rightarrow \infty} \frac{1}{x}\right)^m} = 0,$$

where the third equality follows from the root and power laws, and the last follows from Proposition 8.

This works for $x \rightarrow -\infty$ as well, as long as x^r is defined for negative values of x . (If n is even and m is odd, x^m is negative and the n th root of a negative is not defined, so it does not make sense to write $x^r = \sqrt[n]{x^m}$ in this case.) ■

When studying the behaviour of a function $f(x)$ as $x \rightarrow \infty$, we can always multiply the whole expression by x^r/x^r for any r , since we are interested in the case where x is large and don't have to worry about the denominator possibly being 0. This is very useful in practice when applying Theorem 16.

Here's a simple example.

16.IX **Proposition 17.** We have

$$\lim_{x \rightarrow \infty} \frac{5x^2 - 3x + 4}{3x^2 + 10x - 3} = \frac{5}{3}$$

Proof. Since we're only interested in the behaviour when x is very large, we can assume $x > 0$, so that x^2/x^2 is defined. We divide by this function, obtaining

$$\lim_{x \rightarrow \infty} \frac{5x^2 - 3x + 4}{3x^2 + 10x - 3} = \lim_{x \rightarrow \infty} \frac{(5x^2 - 3x + 4)/x^2}{(3x^2 + 10x - 3)/x^2} = \lim_{x \rightarrow \infty} \frac{5 - 3/x + 4/x^2}{3 + 10/x - 3/x^2}.$$

Invoking the quotient and sum laws now, we take the limits of each term in the numerator and denominator individually, and Theorem 16 applies to give us

$$\lim_{x \rightarrow \infty} \frac{5 - 3/x + 4/x^2}{3 + 10/x - 3/x^2} = \lim_{x \rightarrow \infty} \frac{5 - 0 + 0}{3 + 0 - 0} = \frac{5}{3}.$$

(We technically used the constant multiple law here too, e.g., when performing the operation $\lim_{x \rightarrow \infty} 4/x^2 = 4 \cdot \lim_{x \rightarrow \infty} 1/x^2$.) ■

The limit as $x \rightarrow -\infty$ of the function in the above proposition is also $5/3$ (work it out!), but it is not always the case that the limits as $x \rightarrow \infty$ and $x \rightarrow -\infty$ agree.

Proposition 18. Let

$$f(x) = \frac{\sqrt{2x^2 + 1}}{5x - 1}.$$

We have

$$i) \lim_{x \rightarrow \infty} f(x) = \sqrt{2}/5;$$

- ii) $\lim_{x \rightarrow -\infty} f(x) = -\sqrt{2}/5$;
- iii) $\lim_{x \rightarrow (1/5)^+} f(x) = \infty$; and
- iv) $\lim_{x \rightarrow (1/5)^-} f(x) = -\infty$.

Proof. To prove part (i), we perform the same strategy. The largest power of x in the denominator is just x , so we divide top and bottom by x , noting that for $x > 0$ one has $x = \sqrt{x^2}$:

$$\begin{aligned}
 \lim_{x \rightarrow \infty} \frac{(\sqrt{2x^2 + 1})/x}{(5x - 1)/x} &= \lim_{x \rightarrow \infty} \frac{(\sqrt{2x^2 + 1})/\sqrt{x^2}}{5 - 1/x} \\
 &= \lim_{x \rightarrow \infty} \frac{(\sqrt{2 + 1/x^2})}{5 - 1/x} \\
 &= \frac{\sqrt{2 + 0}}{5 - 0} \\
 &= \frac{\sqrt{2}}{5}
 \end{aligned}$$

It looks like we can do the same thing when $x < 0$, and morally we can, except we have to remember that for negative x one instead has $x = -\sqrt{x^2}$. So

$$\begin{aligned}
 \lim_{x \rightarrow -\infty} \frac{(\sqrt{2x^2 + 1})/x}{(5x - 1)/x} &= \lim_{x \rightarrow -\infty} \frac{(\sqrt{2x^2 + 1})/(-\sqrt{x^2})}{5 - 1/x} \\
 &= \lim_{x \rightarrow -\infty} \frac{-(\sqrt{2 + 1/x^2})}{5 - 1/x} \\
 &= \frac{-\sqrt{2 + 0}}{5 - 0} \\
 &= \frac{-\sqrt{2}}{5},
 \end{aligned}$$

settling claim (ii). We might as well take a moment here to practise finding infinite limits. The input $x = 1/5$ is outside the domain of f , since it makes the denominator 0, so we should investigate the behaviour of this function around this point. Note that when $x = 1/5$ exactly, the numerator is a positive number (since squaring a positive number preserves its positivity, and all the other operations in that expression do as well—if you were extra insecure, you’d calculate the numerator fully and see that it equals $\sqrt{27}/5$, a positive quantity). Hence slightly to the left of $x = 1/5$ and slightly to the right, f is also positive. When x just slightly to the right of $1/5$, the denominator is a very small positive number, and we can make this denominator arbitrarily small by taking x sufficiently close to x from the right. So

$$\lim_{x \rightarrow (1/5)^+} f(x) = \infty,$$

since the numerator is positive as well. On the other hand, when x is very close to $1/5$ from the left, the denominator is a very small negative number, and again

this can be taken arbitrarily close to 0 by taking x sufficiently close to x from the left. So

$$\lim_{x \rightarrow (1/5)^-} = -\infty. \quad \blacksquare$$

The examples we've seen of limits at infinity have been finite, so before moving on, let's see an example where the limit at infinity is infinite.

Proposition 19. *We have*

$$\lim_{x \rightarrow \infty} x - x^2 = -\infty.$$

Proof. We might be tempted to try the difference law here, but since the limits are both infinite we get $\infty - \infty$, which is nonsense. A better thing to do is to factor $x - x^2 = x(1 - x)$. When x gets large, x is a large positive number and $(1 - x)$ is a large negative number. Hence the product is a (very) large negative number. We conclude that the limit must be $-\infty$. $\blacksquare$

The squeeze theorem. We end this section by highlighting the following intuitive theorem, which also goes by the names *sandwich theorem* or *two policemen and a drunk theorem*.

Theorem 20 (*Squeeze theorem*). *Let a be a number and suppose that for all $x \neq a$ in an open interval containing a , we have*

$$f(x) \leq g(x) \leq h(x).$$

If

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} h(x) = L,$$

then $\lim_{x \rightarrow a} g(x)$ exists and equals L .

Proof. This proof is given in Appendix F of the textbook, but we include it in the notes as well for completeness.

Let $\epsilon > 0$. Since $\lim_{x \rightarrow a} f(x) = L$, there is $\delta_1 > 0$ such that every x with $0 < |x - a| < \delta_1$ satisfies

$$|f(x) - L| < \epsilon. \quad (5)$$

Likewise, there is $\delta_2 > 0$ such that every x with $0 < |x - a| < \delta_2$ satisfies

$$|h(x) - L| < \epsilon. \quad (6)$$

Let $\eta > 0$ be a number such that for all x with $0 < |x - a| < \eta$, we have $f(x) \leq g(x) \leq h(x)$. Subtracting L from this inequality gives us

$$f(x) - L \leq g(x) - L \leq h(x) - L \quad (7)$$

for all such x . Hence, letting $\delta = \min\{\delta_1, \delta_2, \eta\}$, every x with $0 < |x - a| < \delta$ satisfies all three of (5), (6), and (7). Combining these three inequalities yields

$$-\epsilon < f(x) - L \leq g(x) - L \leq h(x) - L < \epsilon,$$

which implies that

$$|g(x) - L| < \epsilon,$$

as desired. ■

Near the beginning of the course we encountered the pathological function $\sin(\pi/x)$ and convinced ourselves that it does not possess a limit as $x \rightarrow 0$. However, by combining the squeeze theorem with Proposition 15, we can show that a modification of $\sin(\pi/x)$ does in fact approach a limit.

Proposition 21. *We have*

$$\lim_{x \rightarrow 0} x \sin\left(\frac{\pi}{x}\right) = 0.$$

Proof. It looks like this follows from the product law, but we cannot actually apply that law because $\lim_{x \rightarrow 0} \sin(\pi/x)$ does not exist.

The thing to note is that the function $\sin$ is bounded from above by the constant function 1 and bounded from below by the constant function -1 . In other words, for all real numbers t we have

$$-1 \leq \sin(t) \leq 1.$$

So for all $x > 0$ we have

$$-x \leq x \sin\left(\frac{\pi}{x}\right) \leq x$$

and for all $x < 0$ we have

$$x \leq x \sin\left(\frac{\pi}{x}\right) \leq -x.$$

In other words, the inequality

$$-|x| \leq x \sin\left(\frac{\pi}{x}\right) \leq |x|$$

holds for all $x \neq 0$. By Proposition 15, $\lim_{x \rightarrow 0} |x| = 0$, so we also have $\lim_{x \rightarrow 0} -|x| = 0$ by the constant multiple law. Hence by the squeeze theorem (with $f(x) = -|x|$, $g(x) = x \sin(\pi/x)$, and $h(x) = |x|$), we have

$$\lim_{x \rightarrow 0} x \sin\left(\frac{\pi}{x}\right) = 0.$$

which is what we wanted to show. ■

We will see more applications of the squeeze theorem in the future, for example, when we work out the derivative (whatever that is) of the sine function.

4. Continuity

18.IX We have spent the entire course so far discussing the limit of functions $f(x)$ at numbers a . We now understand that this is the value (if any) that the outputs $f(x)$ must approach as x is taken closer and closer to a . This definition of a limit does not concern itself with $f(a)$ equalling this value that the function approaches, or in fact equalling any value at all. But in many cases, $f(a)$ does equal the limit of f at a , and this situation turns out to be pleasant and important enough to warrant significant attention.

We say that a function f is *continuous* at a number a if

- i) $f(a)$ is defined (i.e., a is in the domain of f);
- ii) $\lim_{x \rightarrow a} f(x)$ exists; and
- iii) $\lim_{x \rightarrow a} f(x) = f(a)$.

If f is defined near a (on some interval containing a but not necessarily at a itself) and yet fails to be continuous by violating one or more of these three conditions, then we say f is *discontinuous* at a .

Clearly by this definition there are various ways for a function to be discontinuous, and some of these cases have special names.

- a) If $\lim_{x \rightarrow a} f(x)$ exists but $f(a)$ is either undefined or does not equal this limit, we say that f has a *removable discontinuity* at a , because we could eliminate this discontinuity by (re)defining the value of f at the single point a . For instance, the function $f(x) = 3x^2 - 2x - 1/(x - 1)$ we saw before has a removable discontinuity at $x = 1$.
- b) If (ii) fails to hold because $\lim_{x \rightarrow a} f(x) = \pm\infty$, then we say f has an *infinite discontinuity* at a (regardless of whether $f(a)$ is defined or not). The function $f(x) = 1/(x - 3)^2$ has an infinite discontinuity at $x = 3$.
- c) Lastly, if the one-sided limits of f at a both exist, but (ii) fails because they do not agree, then we say f has a *jump discontinuity* at a —the function “jumps” from having some value directly to the left of a to having some other value directly to the right of a . The function

$$f(x) = \begin{cases} 0, & \text{if } x \leq 0; \\ 1, & \text{if } x > 0 \end{cases} \quad (8)$$

exhibits this behaviour at $x = 0$.

The appearance of one-sided limits in the definition of a jump discontinuity suggests that we should have a special name for the general case when $f(a)$ equals a one-sided limit of f at a but perhaps not the other. We say that a function f is *continuous from the right* at a number a if

$$\lim_{x \rightarrow a^+} f(x),$$

and *continuous from the left* at a if

$$\lim_{x \rightarrow a^-} f(x).$$

For example, the function $f(x)$ appearing in (8) is continuous from the left but not from the right. Of course, a function is continuous at a if and only if it is continuous from the right and from the left at a , by Theorem 4.

Since the definition of continuous is given directly in terms of the definition of limit, we can use the algebraic limit laws in order to prove corresponding algebraic laws for continuity.

Theorem 22. *Suppose that f and g are continuous at a . Then*

- i) $f + g$;
- ii) $f - g$;
- iii) cf , where c is any real number; and
- iv) fg

are all continuous at a . Furthermore, if $g(a) \neq 0$, then

- v) f/g

is also continuous at a .

Proof. We just prove (i), leaving the rest to the reader as they are similarly easy now that we have access to the algebraic limit laws.

Let f and g be continuous at a . This means that

$$\lim_{x \rightarrow a} f(x) = f(a) \quad \text{and} \quad \lim_{x \rightarrow a} g(x) = g(a).$$

By the sum law for limits, we have

$$\begin{aligned} \lim_{x \rightarrow a} (f + g)(x) &= \lim_{x \rightarrow a} (f(x) + g(x)) \\ &= \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x) \\ &= f(a) + g(a) \\ &= (f + g)(a), \end{aligned}$$

which is what we needed to prove. ■

So if we know certain functions are continuous, then combining them algebraically in the five ways listed above preserves the property of being continuous. This begs the question of which basic functions are continuous. We already know about one class of examples, because the direct substitution property we learned in the previous section directly asserts that any rational function is continuous at all points in its domain. (Go back and read this theorem and you'll see for yourself!) In particular, since every polynomial is a rational function and the domain of a polynomial is the entire real line, polynomials are continuous everywhere.

The following theorem presents a more exhaustive catalogue of continuous functions.

Theorem 23. If a function f is a

- i) polynomial (e.g., $3x^5 - 4x + 1$);
- ii) rational function (e.g., $(4x^2 - 3)/(5x - 1)$);
- iii) root function (e.g., $\sqrt{x}$, $\sqrt[3]{x}$);
- iv) trigonometric function (e.g., $\sin x$, $\tan x$);
- v) inverse trigonometric function (e.g., $\arcsin x$, $\arccos x$);
- vi) exponential function (e.g., e^x , 2^x); or
- vii) logarithmic function (e.g., $\log_{10} x$, $\ln x$);

then f is continuous at all points in its domain.

Proof. We shall skip the proofs of (iv) through (vii) since giving them rigorously would take far too much time, but as we noted before, (i) and (ii) are simply a restatement of the direct substitution property. We have also already seen (iii), since limit law (xi) from Theorem 11 asserts precisely that root functions are continuous. ■

For any set S of numbers, we say that f is *continuous on S* if f is continuous at a for all elements a of S . In particular, we say that f is continuous on the interval (c, d) if f is continuous at a for all $c < a < d$. (Similar definitions can be made for intervals of the form $(c, d]$, $[c, d)$, and $[c, d]$ by changing $<$ to $\leq$ accordingly.)

Knowing which types of functions are continuous now lets us determine many limits without straining our brain very much at all.

Proposition 24. We have

$$\lim_{x \rightarrow \pi/2} \frac{\sin x - 4 \cos x}{3x} = \frac{2}{3\pi}.$$

Proof. We expand

$$\lim_{x \rightarrow \pi/2} \frac{\sin x - 4 \cos x}{3x} = \frac{\lim_{x \rightarrow \pi/2} \sin x - 4 \cdot \lim_{x \rightarrow \pi/2} \cos x}{\lim_{x \rightarrow \pi/2} 3x}$$

by using the algebraic limit laws. In the numerator, we apply the fact that the sine and cosine functions are continuous, and in the denominator we use the direct substitution property of the polynomial $3x$. This yields

$$\lim_{x \rightarrow \pi/2} \frac{\sin x - 4 \cos x}{3x} = \frac{\sin(\pi/2) - 4 \cos(\pi/2)}{3\pi/2} = \frac{1 - 4 \cdot 0}{3\pi/2} = \frac{1}{3\pi/2} = \frac{2}{3\pi}. \quad \blacksquare$$

Often we wish to combine functions not just by adding or multiplying them, etc., but instead by *composing* them. For instance, the function $\sin(\pi/x)$ is the composition $f \circ g$ where $f(x) = \sin x$ and $g(x) = \pi/x$. For this operation we have the following theorem.

Theorem 25. Suppose that $g(x)$ is continuous at a and that $f(x)$ is continuous at $g(a)$. Then $f \circ g$ is continuous at a ; in other words,

$$\lim_{x \rightarrow a} (f \circ g)(x) = (f \circ g)(a) = f(g(a)).$$

Proof. This proof, which proceeds directly via the formal definition of the limit, is found in Appendix F of the textbook. ■

This theorem shows that, erratic though its behaviour may be near the vertical line $x = 0$, the function $\sin(\pi/x)$ is continuous at all $a \neq 0$, since it is the composition of the sine function with a rational function, which is continuous on its domain. We can also use the theorem to compute limits that seem difficult to compute otherwise.

Proposition 26. We have

$$\lim_{x \rightarrow 1} \arccos\left(\frac{1}{\sqrt{x+1}}\right) = \frac{\pi}{4}.$$

Proof. First, we observe that $g(x) = 1/\sqrt{x+1}$ is continuous on its domain. To see this, note that by Theorem 25, since $\sqrt{x}$ and $x+1$ are both continuous, their composition $\sqrt{x+1}$ is, and by the quotient law so is $1/\sqrt{x+1}$ at all points except at $a = -1$.

To apply Theorem 25 again with $f(x)$ and $g(x)$, we must check that $\arccos$ is continuous at $g(1) = 1/\sqrt{1+1} = 1/\sqrt{2}$. Well, $\arccos$ is continuous on its domain, which is the interval $[-1, 1]$, and $1/\sqrt{2}$ falls within this interval. So by Theorem 25, we have

$$\lim_{x \rightarrow 1} f(g(x)) = f(g(a)) = \arccos\left(\frac{1}{\sqrt{2}}\right) = \frac{\pi}{4},$$

which is what we wanted to show. ■

21.IX **The intermediate value theorem.** The remainder of this section is devoted to a very important theorem. It expresses the intuitive idea that for a continuous function to get from one value to another, it must pass through all values in between.

Theorem 27 (*Intermediate value theorem*). Let f be a function that is continuous on some closed interval $[a, b]$ and either

$$f(a) < c < f(b) \quad \text{or} \quad f(b) < c < f(a),$$

then there is some x in the open interval (a, b) such that $f(x) = c$.

The proof of this theorem is non-trivial and takes several lectures of a second-year math course to set up properly. In this class we'll simply use it as a black box.

An easy first application is to show that certain polynomials have roots.

Proposition 28. *The polynomial*

$$f(x) = 3x^4 - 3x^3 + 4x^2 - 9x + 4$$

has a root.

Proof. We wish to show that

$$3x^4 - 3x^3 + 4x^2 - 9x + 4 = 0$$

has a solution. There is a quartic formula (akin to the quadratic formula but for polynomials of degree 4), but it is terrifically complicated. One could also try to factorise the polynomial but it is difficult to figure out where to start. But we are just tasked with showing that there *exists* a root; we are not asked to find out what the root is exactly. So we can use the intermediate value theorem.

Note that the constant term of the polynomial is 4, so $f(0) = 4$. This means that at some point, the graph of the function is above the x -axis. Now let's try to find an input that makes f negative. An easy thing to try is $f(1)$, since we just have to add up all the coefficients:

$$f(1) = 3 - 3 + 4 - 9 + 4 = -1$$

Since f is a polynomial, it is continuous everywhere and in particular is continuous on the interval $[0, 1]$. By the intermediate value theorem with $a = 0$ and $b = 1$ then, since

$$f(0) = 4 > 0 > -1 = f(1),$$

there must be some $0 < x < 1$ such that $f(x) = 0$. This completes the proof. ■

Here is a fun “real-world” application.

Theorem 29 (*Temperature theorem*). *Show that at any given point in time, there is some location in the world that has the same temperature as the opposite point on the globe (assume that temperature varies continuously with position on the globe).*

Proof. For any point x on the globe, let $f(x)$ be the temperature at that point. Pick any starting point p and call the opposite point on the globe $-p$. (Strictly speaking, we're now working in 3-dimensional space. Think of the centre of the earth as the origin. Then the negative of any point is just its opposite point on the globe. Don't worry about this point too much. We can still define continuity for functions on 3-dimensional space, and the intermediate value theorem still holds on curves—the equivalent of intervals—in space.)

If $f(p) = f(-p)$, then we're done (the point we sought is just p). If $f(p) \neq f(-p)$, then we have to be a bit more clever and invoke the intermediate value theorem. Let

$$g(x) = f(x) - f(-x)$$

for all points x on the earth. Since f is continuous, by the limit laws g is continuous as well. For the point p that we chose initially, we have

$$g(p) = f(p) - f(-p) \neq 0.$$

We don't know if $g(p)$ is positive or negative. We just know it's not zero. But now what about $g(-p)$? It turns out that

$$g(-p) = f(-p) - f(-(-p)) = f(-p) - f(p) = -g(p),$$

so whatever sign $g(p)$ had, $g(-p)$ has the opposite sign.

We're almost there. Draw any line on the surface of the earth (this is actually a curve, since the earth is round) between p and $-p$. The function g is continuous on this curve, so the intermediate theorem applies. At one endpoint of the curve, $-p$, the function g has one sign, and at the other endpoint g has the opposite sign. So by the intermediate value theorem, there must be some point x on this curve with $g(x) = 0$. But what does this mean for f ? By the original definition of g , we have

$$0 = g(x) = f(x) - f(-x),$$

so $f(x) = f(-x)$, and this point x has the same temperature as its opposite point on the globe. ■

This theorem applies not just to the temperature, but to any continuous function, and we didn't use the whole globe, but just a single circle on it. The general version of Theorem 29 for any function on a sphere in any dimension is called the Borsuk–Ulam theorem; it was first proved in 1933.

5. Derivatives

Equipped now with a more robust idea of what it means for a function to approach a value but not necessarily get there, we revisit the rate-of-change problem introduced in the first lecture, which is the broad motivating topic for this course as a whole.

Recall that we were considering a function f and wanted to know the rate at which it is changing near some input a (for now, assume that $f(a)$ is well defined). We convinced ourselves geometrically that this rate of change could be modelled as the slope of a line “tangent” to the graph of $f(a)$ at the point $(a, f(a))$. We couldn't rigorously define this line at the time, but it seemed that its slope should be the value of the quotient

$$\frac{f(x+h) - f(x)}{h}$$

“when” $h = 0$, even though we knew that h cannot be zero, being in the denominator.

Now that we are comfortable with limits, however, it is abundantly clear to us that we can overcome this impasse by declaring the slope to be

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h},$$

provided this slope is defined. And of course, once we know the slope, we simply remember that this so-called “tangent line” is supposed to touch the function at the point $(a, f(a))$, so we already know a point on the line. Knowing just one point as well as the slope determines the line completely. Hence the *tangent line* of a function f at the point $P = (a, f(a))$ can be defined to be the line through P with slope

$$m = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h},$$

so long as this limit exists.

Here’s an example demonstrating how to find the equation of a tangent line.

Proposition 30. *For the function*

$$f(x) = x^3,$$

the tangent line passing through the point $(-1, f(-1)) = (-1, -1)$ is given by the equation

$$y = 3x + 2,$$

or alternatively

$$3x - y + 2 = 0.$$

Proof. To find the slope of the tangent line at $a = -1$, we compute

$$m = \lim_{h \rightarrow 0} \frac{f(-1+h) - f(-1)}{h} = \lim_{h \rightarrow 0} \frac{(h-1)^3 - (-1)^3}{h}.$$

Expanding $(h-1)^3$ and then simplifying gives us

$$m = \lim_{h \rightarrow 0} \frac{h^3 - 3h^2 + 3h - 1 - (-1)}{h} = \lim_{h \rightarrow 0} \frac{h^3 - 3h^2 + 3h}{h},$$

and since we’re only interested in values of h near 0 but not *at* 0, we can divide by h on the top and bottom and apply the direct substitution property to get

$$m = \lim_{h \rightarrow 0} (h^2 - 2h + 3) = 0 - 0 + 3 = 3.$$

So the slope of the line is 3 and the equation of the line must be of the form

$$y = 3x + b$$

for some number b . We solve for this b by substituting one known point on the line, namely $(-1, -1)$. We have

$$-1 = 3(-1) + b,$$

whence $b = 2$ and we see that the tangent line can be expressed by the equation

$$3x - y + 2 = 0,$$

as claimed. ■

- 23.IX **Secant lines.** Although one of the primary innovations of differential calculus is the ability to derive tangent lines, we take a brief detour to talk about secant lines and what they represent. Suppose $f(x)$ is a function and we graph it on the plane by setting $y = f(x)$. Then y is a quantity that depends on x . If x changes from x_1 to x_2 , then the *change in x* (also called the *increment of x*) is the quantity

$$\Delta x = x_2 - x_1.$$

The change in $y = f(x)$ that corresponds to this change in x is, similarly, denoted

$$\Delta y = f(x_2) - f(x_1).$$

Then the quotient

$$\frac{\Delta y}{\Delta x} = \frac{f(x_2) - f(x_1)}{x_2 - x_1}$$

is called the *average rate of change of y with respect to x* over the interval $[x_1, x_2]$. Geometrically, this is the slope of the *secant line* that passes through the two points $(x_1, f(x_1))$ and $(x_2, f(x_2))$. We already saw above that the slope of the tangent line is the limit of the slopes of secant lines as the distance between the two points goes to 0. The majority of this course will be about the manifold applications of this definition.

But for some problems, the answer lies directly in the simpler calculation associated with the secant line. Take this one for instance. The fuel usage of a car is often measured in litres per hundred kilometres (so a smaller number means better fuel economy). Suppose that between kilometre 20 and kilometre 320 on Highway 40, your car goes from a full tank (40 litres) down to 19 litres of gasoline in that time. If we wish to know the average rate of change of g with respect to d , then we divide

$$\frac{\Delta g}{\Delta d} = \frac{40 - 19}{320 - 20} = \frac{21 \text{ litres}}{300 \text{ kilometres}} = \frac{7 \text{ litres}}{100 \text{ kilometres}}$$

to conclude that the average rate of fuel consumption is 7 L/100km.

If we had a fuller description or a graph of the function of fuel usage with respect to time, then we'd be able to take secant lines cutting the graph at closer

and closer points. In the limit, we'd obtain a tangent line, whose slope indicates the fuel usage at a specific given point, but this idea of *instantaneous* rate of change of fuel with respect to distance has very little real-world meaning, so we omit it.

(Well, that's not true. The instantaneous rate of change of fuel usage with respect to distance does have some real-world meaning, although the interpretation is perhaps more amusing than actually useful. A litre equals one thousandth of a cubic metre, and a kilometre is of course a thousand metres. So we see that the unit L/100 km equals 10^{-8} m^2 or 0.01 mm^2 —it measures area! This might be baffling the first time you see it, but the best way to interpret this, borrowing numbers from the previous example, is to imagine all 21 litres of fuel (which is three-dimensional) stretched out over the 300 km stretch of highway as a very long skinny rectangular prism. The average fuel usage, then, is the average cross-sectional area of this rectangular prism—namely 0.07 square millimetres in the example. If we now imagine the amount of fuel used varying smoothly as the driver manages the throttle, then the thickness of this imaginary long rectangular prism of liquid varies along the drive as well, and the instantaneous rate of change of fuel with respect to distance at any point is exactly the cross-sectional area of the rectangular prism at that point.)

Instantaneous velocity. A far more useful application of the concept of instantaneous rate of change is when calculating distance travelled over time. This is the notion of *speed* or *velocity*, which we are well acquainted with from a young age.

Let $s(t)$ be the position of some object at time t . We deduced earlier that, whatever the instantaneous velocity is at some time a , it can be approximated by taking averages

$$\frac{s(a+h) - s(a)}{h},$$

over smaller and smaller time intervals h . Now that we know what a limit is, we can simply *define* the *instantaneous velocity* at some time $t = a$ of an object with position function $s(t)$ to be

$$v(a) = \lim_{h \rightarrow 0} \frac{s(a+h) - s(a)}{h}. \quad (9)$$

This is best illustrated by an example. Ha Ling Peak in the Rocky Mountains is 2407 metres above sea level, and the town of Canmore below it is at roughly 1400 metres above sea level. (Let's say it's an even 1400 metres just for the sake of the example.) At the top of the peak there is a sign telling you not to throw rocks off the cliff as they could hit someone down below and cause grave injury. Suppose you could throw a rock off Ha Ling Peak with enough horizontal velocity to make it to town. Let's find out what its vertical velocity would be once it reached the ground, ignoring factors such as air drag, etc.

The elevation of the rock after t seconds of free fall is modelled by the function

$$s(t) = 2407 - 4.9t^2,$$

assuming its initial vertical velocity is zero, and it starts falling from 2407 metres above sea level. First, we find out how long it takes for the rock to reach the ground in town, which is at 1400 metres. We need to solve the equation

$$1400 = 2407 - 4.9t^2;$$

this rearranges to

$$t = \sqrt{\frac{1400}{4.9}} \approx 14.3$$

seconds.

So we want to calculate $v(14.3)$, where $v(a)$ is the instantaneous velocity at time $t = a$. First we need to derive a formula for v . Plugging our formula for s into the definition (g) of v , we solve the resulting limit to obtain

$$\begin{aligned} v(a) &= \lim_{h \rightarrow 0} \frac{2407 - 4.9(a+h)^2 - (2407 - 4.9a^2)}{h} \\ &= \lim_{h \rightarrow 0} \frac{-4.9a^2 - 9.8ah - h^2 + 4.9a^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{-9.8ah - h^2}{h} \\ &= \lim_{h \rightarrow 0} (-9.8a - h) \\ &= -9.8a. \end{aligned}$$

The negative sign here simply expresses that the rock will be moving downwards (losing elevation). From here we can simply plug in $a = 14.3$ to conclude that the vertical velocity of the rock right before it hits the ground is

$$v(14.3) = -9.8(14.3) = -140.5$$

metres per second. (Of course in real life this could not happen because the terminal velocity of the rock is likely much lower than this, and you would have to throw the rock at well over 100 metres per second to even reach the town in the first place. Also because you would never do this for fear of landing in Alberta jail.)

The derivative. In the situation above, the actual equation of the tangent line was not totally necessary to solve the corresponding motivating problem. We only needed to know the slope of this line. Hence for a given function $f(x)$ we have a special name for the slope

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} \tag{10}$$

of the tangent line at the input a ; it is called the *derivative* of the function f at a , and denoted $f'(a)$. Sometimes it's useful to write $x = a + h$, so that

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}.$$

This is a completely equivalent alternative definition of the derivative which we'll use from time to time, but more often than not, the definition (10) leads to simpler calculations.

Here's an example.

Proposition 31. *Let*

$$f(x) = x^2 - x + 1.$$

We have $f'(3) = 5$, and more generally for every number a we have

$$f'(a) = 2a - 1.$$

Proof. If we do the derivation for general a , we can then just plug in $a = 3$ to get the answer of $2(3) - 1 = 5$. But let's do the case $a = 3$ first just to get a sense of the calculations, then tackle the general case after.

We have

$$\begin{aligned} f'(3) &= \lim_{h \rightarrow 0} \frac{f(3+h) - f(3)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(3+h)^2 - (3+h) + 1 - (3^2 - 3 + 1)}{h} \\ &= \lim_{h \rightarrow 0} \frac{3^2 + 6h + h^2 - 3 - h + 1 - 3^2 + 3 - 1}{h} \\ &= \lim_{h \rightarrow 0} \frac{5h + h^2}{h} \\ &= \lim_{h \rightarrow 0} (5 + h) \\ &= 5 + 0 \\ &= 5, \end{aligned}$$

and for the general case at a , we simply replace 3 with a everywhere to obtain

$$\begin{aligned} f'(a) &= \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(a+h)^2 - (a+h) + 1 - (a^2 - a + 1)}{h} \\ &= \lim_{h \rightarrow 0} \frac{a^2 + 2ah + h^2 - a - h + 1 - a^2 + a - 1}{h} \\ &= \lim_{h \rightarrow 0} \frac{2ah - h + h^2}{h} \\ &= \lim_{h \rightarrow 0} (2a - 1 + h) \\ &= 2a - 1 \end{aligned}$$

instead. ■

25.IX **The derivative as a function.** The second calculation in the previous proposition completely characterised the values of $f'(a)$ for different values of a . This, along with the notation f' we employed, suggests that f' is itself a function—plug in different inputs a and you get different outputs. We like to indicate inputs of functions by x , so substituting x for a in the definition (10) of the derivative, we obtain

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}, \quad (11)$$

wherever this limit exists. Hence taking the derivative or *differentiating* the function f resulted in another function, f' .

Let us calculate the derivatives of several simple functions, to get a sense of this idea.

Proposition 32. *The derivative of*

$$f(x) = 2x^3 - x$$

is

$$f'(x) = 6x^2 - 1.$$

Proof. We simply compute

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{2(x+h)^3 - (x+h) - (2x^3 - x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{2(x^3 + 3x^2h + 3xh^2 + h^3) - x - h - 2x^3 + x}{h} \\ &= \lim_{h \rightarrow 0} \frac{2x^3 + 6x^2h + 6xh^2 + 2h^3 - h - 2x^3}{h} \\ &= \lim_{h \rightarrow 0} \frac{6x^2h + 6xh^2 + 2h^3 - h}{h} \\ &= \lim_{h \rightarrow 0} (6x^2 + 6xh + 2h^2 - 1) \\ &= 6x^2 - 1. \quad \blacksquare \end{aligned}$$

Recall that the derivative $f'(x)$ is only defined to exist at points x for which the limit (32) exists. This sometimes means that the domain of a derivative is smaller than the domain of the original function.

Proposition 33. *The function*

$$f(x) = \sqrt{x}$$

is defined on $[0, \infty)$, but its derivative, which is given by the formula

$$f'(x) = \frac{1}{2\sqrt{x}},$$

is only defined on $(0, \infty)$.

Proof. It is clear that $\sqrt{x}$ is well defined for all $x \geq 0$. We now calculate

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{\sqrt{x+h} - \sqrt{x}}{h} \\ &= \lim_{h \rightarrow 0} \frac{\sqrt{x+h} - \sqrt{x}}{h} \cdot \frac{\sqrt{x+h} + \sqrt{x}}{\sqrt{x+h} + \sqrt{x}} \\ &= \lim_{h \rightarrow 0} \frac{x+h-x}{h(\sqrt{x+h} + \sqrt{x})} \\ &= \lim_{h \rightarrow 0} \frac{1}{\sqrt{x+h} + \sqrt{x}} \\ &= \frac{1}{2\sqrt{x}}, \end{aligned}$$

as claimed. But we see that this is undefined for $x = 0$ since we cannot have 0 in the denominator, so we conclude that 0 is not in the domain of f' , which proves the final assertion in the proposition. ■

Notes on notation. Suppose we use the notation $y = f(x)$ to show that the variable y depends on the variable x . Then some common alternative ways to denote the derivative $f'(x)$ are

$$f'(x) = \frac{dy}{dx} = \frac{d}{dx}f(x) = y' = \frac{df}{dx} = Df(x) = D_x f(x) = \dot{y},$$

sorted from left to right in rough descending order of how often you'll see these notations used in these notes.

The reason that there are so many different ways of writing down the derivative symbolically is largely historical; the theory of differential calculus was built up slowly over centuries, and the notations mathematicians came up with along the way were well suited for different contexts and purposes. Differential calculus was first invented in the late 1600s by Isaac Newton and Gottfried Wilhelm Leibniz independently (Leibniz's account was published first, but Newton's private notes predated those of Leibniz by several years). Leibniz used the notation dy/dx , while Newton favoured the “flyspeck” notation $\dot{y}$. We will not use the latter notation very much but it appears often in physics literature. The *prime notation* $f'(x)$ is probably the most popular notation in modern times. It was first used by Leonhard Euler but popularised by Joseph-Louis Lagrange, and first appeared in print in 1749.

Returning to the Leibniz notation dy/dx , we stress that this should not be regarded as a ratio; it is simply another way of writing $f'(x)$. It is reminiscent of the expression $\Delta y/\Delta x$ for the rate of change, which is apt because

$$\frac{dy}{dx} = \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x}.$$

The symbol d/dx is called the *differentiation operator*, because you can apply it to a function f to obtain its derivative (df/dx) . (Recall that the operation of taking a derivative is called differentiation.) The term *operator* refers to a special type of function that takes as input a function and outputs another function, which is what d/dx does, in an abstract sense. When we evaluate the function f' at some number a , it is easy to write $f'(a)$ for this value. It is a little more cumbersome to write this number if we're using Leibniz notation; we write instead

$$\left. \frac{dy}{dx} \right|_{x=a} \quad \text{or} \quad \left. \frac{dy}{dx} \right]_{x=a}.$$

Differentiability. A function f is said to be *differentiable at some number a* if $f'(a)$ (which is defined in terms of a limit) exists. It is said to be *differentiable* on a set S if f is differentiable at every number in S . (In practice, the set S is most often taken to be an open interval (a, b) , an interval of the form (a, ∞) or $(-\infty, b)$, or else the whole real line. If it's differentiable on the whole real line, we might say f is *differentiable everywhere*.)

We already saw in Proposition 33 that the function $\sqrt{x}$, though it is defined on $[0, \infty)$, is only differentiable on $(0, \infty)$. We now proceed to consider a couple more examples.

Proposition 34. *The function*

$$f(x) = |x|$$

is not differentiable at $x = 0$. However, it is differentiable at every other real number.

Proof. We wish to calculate

$$f'(0) = \lim_{h \rightarrow 0} \frac{|0 + h| - |0|}{h} = \lim_{h \rightarrow 0} \frac{|h|}{h}.$$

But we have encountered this limit on the right-hand side before! It was the subject of Proposition 3, and our very first example of a limit that does not exist. Thus $f'(0)$ does not exist.

To prove the second assertion, assume that $x \neq 0$ and show that $f'(x)$ does exist. This is a simple exercise and we leave it to the reader [*Hint*: Split up the proof into one case where $x > 0$ and the other case where $x < 0$.] ■

Graphically, the function $|x|$ does not have a derivative at 0 because the slope of the tangent line near 0 jumps drastically from being -1 for negative x to 1 for positive x . Anytime you see such a “kink” in a graph, you’ll know the graph is not differentiable there.

But that is not the only way a function can fail to be differentiable. Consider the function

$$f(x) = \begin{cases} \sqrt{x}, & \text{if } x \geq 0; \\ -\sqrt{-x}, & \text{if } x < 0. \end{cases}$$

This function is continuous everywhere and if you graph it you’ll see that the picture is “smooth” everywhere; there are no kinks. And yet we saw in Proposition 33 that its derivative did not exist at 0 because the limit defining the derivative approaches infinity at $a = 0$. (This was strictly speaking only for the positive half of f , but you can convince yourself that the negative half does the same thing near 0 as well.) In other words, this example is not differentiable at 0 because its *tangent line is vertical* there.

There is an important point here to be made about the relationship between differentiability and continuity. Both $|x|$ and the function f from the previous paragraph are continuous at 0 but not differentiable there. Therefore continuity at a point does not guarantee differentiability at that point. However, the opposite implication *does* hold.

Theorem 35. Suppose f is a function and a is a number such that $f(a)$ is defined. If f is differentiable at a , then it is continuous at a .

Proof. Assume that $f'(a)$ exists; that is, the limit

$$\lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

equals some well defined number. Then we have

$$\begin{aligned} \lim_{x \rightarrow a} f(x) &= \lim_{x \rightarrow a} f(x) - f(a) + f(a) \\ &= \lim_{x \rightarrow a} \left(\frac{f(x) - f(a)}{x - a} \cdot (x - a) \right) + f(a) \\ &= \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} \cdot \lim_{x \rightarrow a} (x - a) + f(a) \\ &= f'(a) \cdot 0 + f(a) \\ &= f(a), \end{aligned}$$

which is exactly the criterion for f to be continuous at a . ■

Recall the function $\sin(\pi/x)$, which does not have a limit as $x \rightarrow 0$. Using the squeeze theorem, we saw that if we modify the function slightly by multiplying x , the resulting function $x \sin(\pi/x)$, while still not *defined* at 0, does go to 0 as $x \rightarrow 0$. So if we artificially adjoin the point $(0, 0)$ to this graph, then the resulting function is continuous. However, it is not differentiable, as we now proceed to show.

Proposition 36. *The function*

$$f(x) = \begin{cases} x \sin(\pi/x), & \text{if } x \neq 0; \\ 0, & \text{if } x = 0 \end{cases}$$

is not differentiable at 0.

Proof. We compute

$$\begin{aligned} f'(0) &= \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(0+h) \sin(\pi/(0+h)) - 0}{h} \\ &= \lim_{h \rightarrow 0} \frac{h \sin(\pi/h)}{h} \\ &= \lim_{h \rightarrow 0} \sin\left(\frac{\pi}{h}\right), \end{aligned}$$

and we have already seen that this limit does not exist. **■**

To illustrate that the matter of differentiability is fairly subtle, consider that if we multiply $\sin(\pi/x)$ by x^2 instead of x (and artificially adjoin $(0,0)$ as before, then the result *is* differentiable at 0.

Proposition 37. *The function*

$$f(x) = \begin{cases} x^2 \sin(\pi/x), & \text{if } x \neq 0; \\ 0, & \text{if } x = 0 \end{cases}$$

is both continuous and differentiable at 0.

Proof. To prove that $f(x)$ is continuous at 0, we need to show that

$$\lim_{x \rightarrow 0} x^2 \sin\left(\frac{\pi}{x}\right) = 0.$$

This follows from the squeeze theorem and the fact that $-x^2 \leq f(x) \leq x^2$ for all numbers x . (Fill in the details here as an exercise.)

To show that f is differentiable at 0, we just need to show that $f'(0)$ exists. So we compute

$$\begin{aligned} f'(0) &= \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(0+h)^2 \sin(\pi/(0+h)) - 0}{h} \\ &= \lim_{h \rightarrow 0} \frac{h^2 \sin(\pi/h)}{h} \\ &= \lim_{h \rightarrow 0} h \sin\left(\frac{\pi}{h}\right), \end{aligned}$$

but this last limit equals 0, by Proposition 21. **■**

Higher derivatives. We said earlier that the operator d/dx takes in a function and spits out another function (which may have a different domain than the original one). It is natural, then, to consider applying the operator again to the resulting function. If the original function is f , then its derivative is denoted by df/dx or f' , and taking the derivative again gives us *second derivative* of f , denoted by

$$\frac{d^2}{dx^2}f(x) \quad \text{or} \quad f''(x).$$

If we write $y = f(x)$, which we often do when considering the graph of the function f , then we can also denote the second derivative by d^2y/dx^2 or y'' .

We can keep repeating the process, obtaining the third derivative, fourth derivative, and so on. We denote the n th derivative of f by

$$\frac{d^n}{dx^n}f(x) \quad \text{or} \quad f^{(n)}(x),$$

because writing the $'$ symbol n times can become tedious (though it's not uncommon to see $f'''(x)$ instead of $f^{(3)}(x)$).

There are many uses for higher derivatives, which we'll see as the course progresses. At this juncture we'll just note one function whose second derivative has a very intuitive interpretation. This is the position function $s(t)$ of a moving object with respect to time. We already convinced ourselves earlier that its derivative $s'(t)$ describes the object's velocity with respect to time, since velocity is the rate of change of position. Taking another derivative tells us how much the velocity changes with respect to time, so $s''(t)$ is the object's *acceleration*.

For instance, the elevation of the rock in our Ha Ling Peak example was described by the function

$$s(t) = 2407 - 4.9t^2,$$

and we calculated its velocity function

$$v(t) = s'(t) = -9.8t.$$

Taking the derivative once more (we'll spare the limit calculations, because soon we'll develop strategies to compute derivatives very quickly) gives us the acceleration

$$a(t) = s''(t) = -9.8,$$

which is just a constant, namely the constant of acceleration due to gravity.

This discussion concludes the section we've devoted to defining the derivative and various other notions pertaining to it. It also concludes the first third of this course, which introduced a number of foundational topics, namely limits, continuity, and differentiation. In the next third of the course, we build up a toolbox of techniques that will allow us to quickly calculate derivatives of a large class of functions that are commonly used in science and engineering.

II. TECHNIQUES

*Sit a quantitas data constans,
erit da æqualis 0,
et dāx æqualis adx:
si sit y æqualis v
(seu ordinata quævis curvæ Y,
æqualis cuivis ordinatæ respondentī V)
erit dy æqualis dv.*

— GOTTFRIED WILHELM LEIBNIZ, *Nova methodus pro maximis et minimis* (1684)

6. Algebraic rules for differentiation

28.IX Early in our study of limits we proved simple algebraic laws like $\lim(f + g) = \lim f + \lim g$. The goal of this section is to repeat this exercise for derivatives, though as we shall see, some of the algebraic rules for differentiation are not as simple as their counterparts in the limit world.

We begin with the easy stuff. A constant function has a completely flat graph; it does not change with respect to its input. Hence its derivative should be 0 everywhere. Let us prove this simple fact.

Theorem 38. *Let c be any number and consider it to be a constant function of x (no matter what number x it is given, the function just outputs the number c). Then*

$$\frac{d}{dx}c = 0.$$

Proof. Let $f(x) = c$ for all x . For any number x ,

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{c - c}{h} = 0. \quad \blacksquare$$

Next, we record the sum, difference, and constant multiple rules for differentiation, which follow directly from the corresponding limit laws.

Theorem 39 (*Sum, difference, and constant multiple rules*). *Suppose that f and g are both differentiable at some number x . Then the function $f + g$ and $f - g$ are also differentiable at x , and we have*

$$(f + g)'(x) = f'(x) + g'(x) \quad \text{as well as} \quad (f - g)'(x) = f'(x) - g'(x).$$

Furthermore, for any constant c , the function cf is differentiable at x , and

$$(cf)'(x) = cf'(x).$$

Proof. For the sum rule, we simply expand

$$\begin{aligned} (f + g)'(x) &= \lim_{h \rightarrow 0} \frac{(f + g)(x + h) - (f + g)(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x + h) + g(x + h) - f(x) - g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h} + \lim_{h \rightarrow 0} \frac{g(x + h) - g(x)}{h} \\ &= f'(x) + g'(x), \end{aligned}$$

where the third equality is a consequence of the sum law for limits. The difference and constant multiple rules are proved similarly. Try them as an exercise. $\blacksquare$

The sum law in particular can be used to prove the power rule for monomials of the form x^n , where n is a positive integer.

Theorem 40 (*Power rule*). Let $n \geq 1$ be an integer and let

$$f(x) = x^n.$$

Then for any number a , we have

$$f'(a) = na^{n-1}.$$

Proof. This proof will make use of the identity

$$x^n - a^n = (x - a)(x^{n-1} + ax^{n-2} + a^2x^{n-3} + \cdots + a^{n-3}x^2 + a^{n-2}x + a^{n-1}).$$

It can be verified by simply multiplying out the right-hand side. Now we expand

$$\begin{aligned} f'(a) &= \lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} \\ &= \lim_{x \rightarrow a} (x^{n-1} + ax^{n-2} + a^2x^{n-3} + \cdots + a^{n-3}x^2 + a^{n-2}x + a^{n-1}). \end{aligned}$$

Note that there are exactly n terms in the limit, and in each term, the power of a and the power of x together add up to $n - 1$. So by applying the sum law for limits to distribute the limit over the sum before taking $x \rightarrow a$ in each term, we obtain

$$f'(a) = na^{n-1},$$

which is what we wanted. ■

Using the theorems we've proved so far in this section, we can very easily recompute the derivative of the function $x^2 - x + 1$ that appeared in Proposition 31. We simply have

$$\frac{d}{dx}(x^2 + x + 1) = \frac{d}{dx}x^2 - \frac{d}{dx}x + \frac{d}{dx}1 = 2x - 1,$$

which we proved with greater difficulty earlier. Other polynomials are dealt with similarly. Let's see one more example.

Suppose we want to find all points on the curve

$$y = -\frac{x^3}{3} + 9x + 5$$

at which the tangent line is parallel to the line $y = 5x$ (if any such points exist). In other words, where does this curve have a tangent line with a slope of 5? We compute the derivative

$$\frac{dy}{dx} = 9 - x^2 = (3 - x)(3 + x).$$

From here it's easy to see that this equals 5 when $x = 2$ or $x = -2$. (Alternatively you could solve for $9 - x^2 = 5$, which is equivalent to $x^2 = 4$, which has the same solutions.) To find the exact points on the curve, we now plug these points into the original function. When $x = 2$, we have

$$y = -\frac{2^3}{3} + 9 \cdot 2 + 5 = 23 - \frac{8}{3} = \frac{61}{3},$$

and when $x = -2$ we have

$$y = -\frac{(-2)^3}{3} + 9 \cdot (-2) + 5 = \frac{8}{3} - 13 = \frac{-31}{3}.$$

We conclude that the two points on the curve $y = -x^3/3 + 9x + 5$ at which the tangent line has slope equal to 5 are $(2, 61/3)$ and $(-2, -31/3)$.

Returning to the formula given for the power rule, you may have noticed something interesting. In Proposition 33, we showed that

$$\frac{d}{dx} x^{1/2} = \frac{d}{dx} \sqrt{x} = \frac{1}{2\sqrt{x}} = \frac{1}{2} x^{-1/2}.$$

One might speculate, then, that the power rule holds even when n is not a positive integer. This is in fact true, but we don't have the tools to prove it yet, so we defer it until later.

The product and quotient rules. The next algebraic rules to consider are multiplication and division. Because $\lim fg = (\lim f)(\lim g)$ and $\lim f/g = (\lim f)/(\lim g)$ wherever these limits (and the division) make sense, you might guess that we have something like $(fg)' = f' \cdot g'$. This turns out to be off the mark.

To derive the correct answer, you need to draw a rectangle. Imagine f and g as quantities depending on some independent variable x . To make your life easy, assume both f and g are positive. Draw a rectangle whose length is f and whose height is g , so the area of this rectangle is exactly the quantity fg we want to study. In particular, we want to know how much its area changes if you perturb x very slightly. Suppose that if you increase x very slightly, the value of f increases very slightly and the value of g increases very slightly as well. Call these increases Δf and Δg , respectively. (In general, f could actually decrease, or g could decrease, or both, but this is just a sketch for the sake of intuition, so ignore these possibilities for now.) What has happened to the area of the rectangle; in other words, what is $\Delta(fg)$? Well, three new rectangles have been added to it whose areas are $f \cdot \Delta g$, $g \cdot \Delta f$, and $(\Delta f)(\Delta g)$. Leibniz reasoned that the area $(\Delta f)(\Delta g)$, because it is “infinitely smaller” than the other terms, can simply be ignored. Hence he claimed that

$$\Delta(fg) = f\Delta g + g\Delta f,$$

whence

$$\frac{\Delta(fg)}{\Delta x} = f \frac{\Delta g}{\Delta x} + g \frac{\Delta f}{\Delta x},$$

and since $d(fg)/dx$ is the limit of $\Delta(fg)/\Delta x$ as $\Delta x \rightarrow 0$, we can simply replace all instances of Δ with d above to obtain

$$\frac{d}{dx}(fg) = f \frac{dg}{dx} + g \frac{df}{dx}.$$

Leibniz was correct, though his “proof” does not hold up to modern standards. (If we are throwing away the quantity $(\Delta f)(\Delta g)$ simply because its area is “infinitely small” as $\Delta x \rightarrow 0$, why not throw away the other thin rectangles too?—these areas also go to 0.) Let us now restate the product rule in alternative notation (using f' and g') and also give a rigorous proof with our modern definition of derivative.

Theorem 41 (*Product rule*). *If f and g are both differentiable at x , then the product fg is also differentiable at x , and*

$$(fg)'(x) = f(x)g'(x) + g(x)f'(x).$$

Proof. There is only one step in this proof that is not entirely obvious. First we expand

$$\begin{aligned} (fg)'(x) &= \lim_{h \rightarrow 0} \frac{(fg)(x+h) - (fg)(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \end{aligned}$$

The nonobvious thing to do is to add $-g(x)f(x+h) + g(x)f(x+h)$ into the numerator. This will help us factor things out and make the formulas for $f'(x)$ and $g'(x)$ both appear. Picking up where we left off, we have

$$\begin{aligned} (fg)'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - g(x)f(x+h) + g(x)f(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h)(g(x+h) - g(x)) + g(x)(f(x+h) - f(x))}{h} \\ &= \left(\lim_{h \rightarrow 0} f(x+h) \right) \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} + g(x) \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \left(\lim_{h \rightarrow 0} f(x+h) \right) g'(x) + g(x) f'(x). \end{aligned} \tag{12}$$

The last thing to deal with is the limit $\lim_{h \rightarrow 0} f(x+h)$. Since f is differentiable at x , it is also continuous at x by Theorem 35. So

$$\lim_{h \rightarrow 0} f(x+h) = f(x),$$

and substituting this into (12) proves the theorem. ■

30.IX Next we set our sights on proving a quotient rule, that is, a formula for $(f/g)'$. Because we can write a quotient f/g as $f \cdot (1/g)$, supposing we can figure out what the derivative of $1/g$ is, we can use the product rule to compute what the quotient rule should be.

The formula for the derivative of the reciprocal of a function is more complicated than you might expect, though if you have taken calculus before in high school and know the derivative of $1/x$ as well as the chain rule, then you should understand why this formula makes sense. If this is your first time taking calculus, don't worry, we'll learn those other things soon, and you can revisit this at a later stage. For now, let's just get on with the theorem and its proof.

Theorem 42. Suppose that g is differentiable at x and that $g(x) \neq 0$, so that $1/g(x)$ is defined. Then the function $1/g$ is differentiable at x as well, and

$$\left(\frac{1}{g}\right)'(x) = \frac{-g'(x)}{g(x)^2}.$$

Proof. Though the formula is a little bit gruesome, the proof is not. Expanding, putting everything over a common denominator, and then rearranging yields

$$\begin{aligned} \left(\frac{1}{g}\right)'(x) &= \lim_{h \rightarrow 0} \frac{1/g(x+h) - 1/g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{g(x) - g(x+h)}{hg(x+h)g(x)} \\ &= -\lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \cdot \lim_{h \rightarrow 0} \frac{1}{g(x+h)g(x)} \\ &= -g'(x) \cdot \lim_{h \rightarrow 0} \frac{1}{g(x+h)g(x)}. \end{aligned} \tag{13}$$

From here we once again use Theorem 35 to see that g is continuous at x , so that

$$\lim_{h \rightarrow 0} \frac{1}{g(x+h)g(x)} = \frac{1}{g(x)^2},$$

and substituting this fact into (13) now yields the claimed identity. ■

Now that we have the reciprocal rule, a general quotient rule for functions f and g can be derived simply by applying the product rule with the functions f and $1/g$.

Theorem 43 (Quotient rule). Suppose that f and g are differentiable at x , and assume that $g(x) \neq 0$, so that f/g is defined at x . Then f/g is differentiable at x , and

$$\left(\frac{f}{g}\right)'(x) = \frac{g(x)f'(x) - f(x)g'(x)}{g(x)^2}.$$

Proof. Since $f/g = f \cdot (1/g)$, we use the product rule to obtain

$$\left(f \cdot \frac{1}{g}\right)'(x) = f(x)\left(\frac{1}{g}\right)'(x) + \left(\frac{1}{g}\right)(x)f'(x).$$

By Theorem 42, we have

$$\left(\frac{1}{g}\right)'(x) = \frac{-g'(x)}{g(x)^2};$$

substituting this into (14) gives us

$$\begin{aligned} \left(f \cdot \frac{1}{g}\right)'(x) &= f(x) \left(\frac{-g'(x)}{g(x)^2}\right) + \frac{f'(x)}{g(x)} \\ &= \frac{f'(x)}{g(x)} - \frac{f(x)g'(x)}{g(x)^2} \\ &= \frac{g(x)f'(x) - f(x)g'(x)}{g(x)^2}, \end{aligned}$$

which is exactly what we wanted. ■

We are now able to find derivatives of any rational function. Here's an example. Suppose we are asked to find the points on the curve

$$y = \frac{x}{x^2 + 1}$$

at which the tangent line to the curve is horizontal, i.e., has slope equal to 0. Since x^2 is always nonnegative, the denominator here is always nonzero, so y is defined for all values of x . By the quotient rule, the derivative of y is

$$y' = \frac{(x^2 + 1) - x(2x)}{(x^2 + 1)^2} = \frac{1 - x^2}{(x^2 + 1)^2}.$$

This equals 0 when $x^2 = 1$, or when x equals either 1 or -1 . When $x = 1$, we have

$$y = \frac{1}{1^2 + 1} = \frac{1}{2},$$

and when $x = -1$, we have

$$y = \frac{-1}{(-1)^2 + 1} = \frac{-1}{2},$$

so we see that the two points on the curve $y = x/(x^2 + 1)$ whose tangent lines are horizontal are $(1, 1/2)$ and $(-1, -1/2)$.

7. Derivatives of the exponential and trigonometric functions

We now have a great number of rules that allow us to find the derivatives of algebraic expressions like $f + g$ and fg , assuming we know the derivatives of f and g . But our inventory of basic functions whose derivative we know is still rather meagre. We only know how to deal with polynomials and rational functions, and adding, subtracting, multiplying and dividing polynomials or rational

functions will only give us more rational functions. So we'll have to go back to basics in order to build up our stockpile of functions.

The first class of functions we look at is exponential functions like 2^x or 10^x . More generally, let b be any number; we shall take it for granted that the quantity b^x varies smoothly as one changes the inputs x . So we can consider the function

$$f(x) = b^x$$

and ask how it changes as x varies. Using the formula $b^{x+y} = b^x b^y$, we compute

$$f'(x) = \lim_{h \rightarrow 0} \frac{b^{x+h} - b^x}{h} = \lim_{h \rightarrow 0} \frac{b^x b^h - b^x}{h} = \lim_{h \rightarrow 0} \frac{b^x (b^h - 1)}{h} = b^x \lim_{h \rightarrow 0} \frac{b^h - 1}{h}.$$

The limit on the right-hand side doesn't depend on x at all, and we see that the original function $f(x) = b^x$ has reappeared on the right-hand side. Plugging in $x = 0$, we obtain the identity

$$\left. \frac{d}{dx} b^x \right|_{x=0} = f'(0) = \lim_{h \rightarrow 0} \frac{b^h - 1}{h},$$

which tells us that

$$\frac{d}{dx} b^x = b^x \cdot \left. \frac{d}{dx} b^x \right|_{x=0}, \quad (14)$$

assuming the derivative exists. This means that the derivative of an exponential function is proportional to the function itself; the slope of the tangent line is some constant (namely the derivative of the function at zero) times the function's height at that point.

In light of (14), it is natural to wonder if there is some base b such that the derivative

$$\lim_{h \rightarrow 0} \frac{b^h - 1}{h}$$

equals 1; if this is true for some b , then the derivative of b^x would be b^x again, which seems like a pleasant thing to have. To see that such a base does exist, note that if you take $b = 2$ and estimate the resulting limit (44) by taking smaller and smaller values of h , the limit seems to be something less than 1 (in fact, it is roughly 0.693). Likewise, if you take $b = 3$ and do the same thing, the limit seems to be larger than 1 (in fact, it is roughly 1.099). So by the intermediate value theorem, there must be some number between 2 and 3, call it e , with

$$\lim_{h \rightarrow 0} \frac{e^h - 1}{h} = 1.$$

By taking better approximations one can compute the value

$$e \approx 2.71828 \dots$$

with better precision, but the numerical value of e doesn't come up in this class very often. A more important observation to record at this point is that this number e satisfies

$$\frac{d}{dx}e^x = e^x.$$

We'll prove more facts about e and its associated exponential function e^x (which is so important that we call it *the* exponential function) as the course progresses.

02.X For now let's see a more complicated example involving exponentials. If nothing else, this will help us practise using the product and quotient rules from earlier.

Proposition 44. *Let*

$$f(x) = \frac{e^{2x} - 1}{2e^x},$$

which is defined for all numbers x . We have

$$f'(x) = \frac{e^{2x} + 1}{2e^x}.$$

Proof. We would like to use the quotient rule, but first we should deal with the fact that there is an expression in the formula for $f(x)$ we don't immediately know the derivative of, namely e^{2x} . We'll learn how to properly deal with function compositions such as these soon, but for now, just note that $e^{2x} = e^x e^x$ and invoke the product rule to obtain

$$\frac{d}{dx}e^{2x} = \frac{d}{dx}(e^x e^x) = e^x e^x + e^x e^x = 2e^{2x}. \quad (15)$$

Now we're in a better position to invoke the quotient rule on f itself. We compute

$$\begin{aligned} f'(x) &= \frac{2e^x \cdot (d/dx)(e^{2x} - 1) - (e^{2x} - 1)(d/dx)(2e^x)}{(2e^x)^2} \\ &= \frac{2e^x \cdot 2e^{2x} - (e^{2x} - 1)2e^x}{4e^{2x}} \\ &= \frac{2e^x(2e^{2x} - e^{2x} + 1)}{4e^{2x}} \\ &= \frac{e^{2x} + 1}{2e^x}, \end{aligned}$$

which is exactly the claim. ■

So the derivative f' of $f(x) = (e^{2x} - 1)/2e^x$ admits an expression that is very similar to the formula for f itself. If you have some spare time and want to practise the quotient rule some more, try finding $f''(x)$. You might notice a pattern that will make it easy to determine $f^{(3)}$ and $f^{(4)}$ as well. We'll revisit these functions at a later date.

***Formally defining the exponential.** This small subsection is intended for the reader who is unhappy with the fact that we have swept the continuity of b^x under the rug. The problem with trying to prove such a thing now is that we don't yet have a proper definition of what the expression b^x means when x is not a rational number. If x is rational—say $x = p/q$ for positive integers p and q —then $b^x = \sqrt[q]{b^p}$, and b^p is simply the product of b with itself p times. But this breaks apart if x is irrational. For example, if $x = \pi$, what does it mean to multiply x by itself “ π times?” The reason we cannot formally resolve this yet is that a precise definition of exponentials requires the concept of the integral, which is only introduced in MATH 141. Fortunately, most of you will take this next semester, but if you have never seen integrals before, the following paragraph will be gibberish to you.

To define b^x for general real values of b and x , we start by defining the (natural) logarithm of any $x > 0$ as the value of the definite integral

$$\ln x = \int_1^x \frac{1}{t} dt.$$

It can be shown that this is continuous, and there is a unique number x that makes this integral x . We call this number e , and if we define $\exp$ as the inverse of $\ln$, which exists because $\ln$ as we have defined it is one-to-one, then we have $\exp(1) = e$. (This is the same e that we defined earlier in the section.) From here, one can prove that

$$\exp(x) = (\exp(1))^x = e^x,$$

and finally we *define*

$$b^x = \exp(\ln(b^x)) = e^{x \ln b}.$$

By this definition, it is clear that this is a continuous function of x , since $\ln$ is and therefore its inverse $\exp$ must be. From here you can rederive all the facts about exponentials and logarithms that you already know.

Trigonometric derivatives. So much for exponentials. The rest of this section is devoted to finding formulas for the derivatives of the six basic trigonometric functions

$$\sin, \quad \cos, \quad \tan, \quad \csc, \quad \sec, \quad \text{and} \quad \cot.$$

Because the latter four functions can all be expressed as quotients involving $\sin$ and $\cos$, we only need to differentiate $\sin$ and $\cos$ from scratch; the rest of the formulas will then follow from the quotient rule.

Differentiating the sine and cosine functions will require us to recall some important facts from trigonometry class. First, note that $\sin$ is odd and $\cos$ is even; in other words,

$$\sin(-\theta) = -\sin \theta \quad \text{and} \quad \cos(-\theta) = \cos \theta.$$

Next, by drawing a unit circle and invoking the Pythagorean theorem, we see that

$$\sin^2 \theta + \cos^2 \theta = 1$$

for all θ . It is a bit harder to prove the *addition formulas*

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \sin \beta \cos \alpha \quad (16)$$

and

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta; \quad (17)$$

if you want to practise your trigonometric skills, these facts are proved in Exercises 89 to 91 of Appendix D of the book. (In those exercises, you actually end up proving

$$\sin(\alpha - \beta) = \sin \alpha \cos \beta - \sin \beta \cos \alpha$$

instead of (16), but from this subtraction formula one can recover (16) by substituting $-\beta$ instead of β , and remembering that the sine function is odd.)

In addition to the aforementioned trigonometric propositions, we shall also need the following two special limits.

Theorem 45. *We have*

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1 \quad (18)$$

and

$$\lim_{\theta \rightarrow 0} \frac{\cos \theta - 1}{\theta} = 0. \quad (19)$$

Proof. We prove these statements in order. First, we calculate the right-hand limit $\lim_{\theta \rightarrow 0^+} \sin \theta / \theta$. We shall do this using the squeeze theorem. (This proof is significantly easier if you have access to a piece of paper on which to draw. Alternatively, refer to Figure 6 in Section 3.3 of the textbook.)

Assume that $0 < \theta < \pi/2$, which we can do since we want to take θ to 0. Draw a line segment AC along the x -axis of length 1, and draw a line segment of length 1 AB above it so that the angle BAC is θ . Enclosing this angle with an arc of a circle of radius 1, we now have a slice of a circle. To calculate the area of this slice, we note that the arc of the circle has length θ , so its length is $\theta/(2\pi)$ proportion of the full circumference of the circle. Thus the area of the slice must be $\theta/(2\pi)$ times the area of a full circle with radius 1, which is π . In other words, the area of the slice is $\theta/2$.

Now draw a line from B down to the x -axis so that it intersects AC perpendicularly. Call this intersection point D . The right triangle ABD lies entirely within the slice of the circle we considered above. The area of this right triangle is easy to calculate, since it is a right triangle with hypotenuse 1 and angle θ . Thus AD has length $\cos \theta$ and BD has length $\sin \theta$, and the triangle ABD has area $(\sin \theta \cos \theta)/2$. Since this triangle lies completely within the circle slice, we have

$$\frac{\sin \theta \cos \theta}{2} \leq \frac{\theta}{2}. \quad (20)$$

Lastly, draw a vertical line extending up from the point C (perpendicular to the x -axis). At the same time, extend the line AB so that it intersects with this new vertical line; call this intersection point E . The triangle ACE completely contains the circle slice from earlier. Moreover, the length of AC is 1 and we have a right angle at C , so the length of CE must be $\tan \theta$. This shows that the area of ACE is $\tan \theta/2$, and hence

$$\frac{\theta}{2} \leq \tan \theta, \quad (21)$$

since this bigger triangle contains the circle slice. Combining the inequalities (20) and (21) and dividing through by 2, we obtain

$$\sin \theta \cos \theta \leq \theta \leq \tan \theta.$$

Now, dividing by $\sin \theta$ yields

$$\cos \theta \leq \frac{\theta}{\sin \theta} \leq \frac{\tan \theta}{\sin \theta} = \frac{1}{\cos \theta},$$

and taking the reciprocal of the whole inequality gives us

$$\frac{1}{\cos \theta} \geq \frac{\sin \theta}{\theta} \geq \cos \theta.$$

By the squeeze theorem, we must have

$$\lim_{\theta \rightarrow 0^+} \frac{1}{\cos \theta} \geq \lim_{\theta \rightarrow 0^+} \frac{\sin \theta}{\theta} \geq \lim_{\theta \rightarrow 0^+} \cos \theta,$$

but both the left- and right-hand sides go to 1, so $\sin \theta/\theta \rightarrow 1$ as well, as $x \rightarrow 0$ from the right.

To see that the limit is also 1 from the left, note that

$$\frac{\sin(-\theta)}{-\theta} = \frac{-\sin(-\theta)}{\theta} = \frac{\sin \theta}{\theta},$$

since $\sin$ is an odd function. In short, the function $\sin \theta/\theta$ is even, so its one-sided limits are equal. This proves (18).

It still remains to prove (19); fortunately, this can be derived from (18) without much further ado. We expand and multiply by the conjugate of $\cos \theta - 1$, namely $\cos \theta + 1$, on the top and bottom to obtain

$$\begin{aligned} \lim_{\theta \rightarrow 0} \frac{\cos \theta - 1}{\theta} &= \lim_{\theta \rightarrow 0} \frac{\cos \theta - 1}{\theta} \cdot \frac{\cos \theta + 1}{\cos \theta + 1} \\ &= \lim_{\theta \rightarrow 0} \frac{\cos^2 \theta - 1}{\theta(\cos \theta + 1)}. \end{aligned}$$

From here, we use the fact that $\sin^2 \theta + \cos^2 \theta = 1$ to write

$$\begin{aligned} \lim_{\theta \rightarrow 0} \frac{\cos^2 \theta - 1}{\theta(\cos \theta + 1)} &= \lim_{\theta \rightarrow 0} \frac{\sin^2 \theta}{\theta(\cos \theta + 1)} \\ &= \lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} \cdot \lim_{\theta \rightarrow 0} \frac{\sin \theta}{\cos \theta + 1} \\ &= 1 \cdot \frac{0}{1 + 1} \\ &= 0, \end{aligned}$$

where the third equality follows from (18). ■

We are now able to find formulas for all six trigonometric functions. We state this as one megatheorem.

Theorem 46 (*Derivatives of trigonometric functions*). *We have*

- i) $\sin' = \cos$;
- ii) $\cos' = -\sin$;
- iii) $\tan' = \sec^2$;
- iv) $\csc' = -\csc \cdot \cot$;
- v) $\sec' = \sec \cdot \tan$; and
- vi) $\cot' = -\csc^2$.

wherever these derivatives exist. (The derivatives $\sin'$ and $\cos'$ exist everywhere; $\tan'$ and $\sec'$ are defined wherever $\cos \neq 0$, that is, everywhere except at inputs of the form $n\pi/2$ for all odd integers n ; and $\csc$ and $\cot$ are defined wherever $\sin \neq 0$, so they are defined everywhere except at integer multiples of π .)

Proof. By the addition formula (16) for sine, we have

$$\begin{aligned} \sin'(x) &= \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin x}{h} \\ &= \lim_{h \rightarrow 0} \frac{\sin x \cos h + \sin h \cos x - \sin x}{h} \\ &= \lim_{h \rightarrow 0} \frac{\sin x(\cos h - 1) + \sin h \cos x}{h} \\ &= \sin x \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} + \cos x \lim_{h \rightarrow 0} \frac{\sin h}{h} \\ &= \sin x \cdot 0 + \cos x \cdot 1 \\ &= \cos x \end{aligned}$$

for all numbers x , where the fifth line follows from the special limits computed

in Theorem 45. Likewise, by the addition formula (17) for cosine,

$$\begin{aligned}
 \cos'(x) &= \lim_{h \rightarrow 0} \frac{\cos(x+h) - \cos x}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\cos x \cos h - \sin x \sin h - \cos x}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\cos x(\cos h - 1) - \sin x \sin h}{h} \\
 &= \cos x \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} - \sin x \lim_{h \rightarrow 0} \frac{\sin h}{h} \\
 &= \cos x \cdot 0 - \sin x \cdot 1 \\
 &= -\sin x
 \end{aligned}$$

for all x , where again we have invoked Theorem 45 in the fifth line. This proves both (i) and (ii).

To prove (iii), we apply the quotient rule to the identity $\tan = \sin / \cos$. Wherever the cosine function is not 0, we have

$$\tan' = \left(\frac{\sin}{\cos} \right)' = \frac{\cos \cdot \sin' - \sin \cdot \cos'}{\cos^2} = \frac{\cos^2 - \sin \cdot (-\sin)}{\cos^2} = \frac{1}{\cos^2} = \sec^2,$$

where we have once again used the Pythagorean identity $\cos^2 + \sin^2 = 1$.

Because the cosecant, secant, and cotangent functions are defined as reciprocals of the sine, cosine, and tangent functions, respectively, we shall use the reciprocal rule $(1/g)' = -g'/g^2$ given by Theorem 42 to prove (iv), (v), and (vi). For the cosecant function, we compute

$$\csc' = \left(\frac{1}{\sin} \right)' = \frac{-\sin'}{\sin^2} = \frac{-\cos}{\sin^2} = -\csc \cdot \cot,$$

which settles (iv). Finally we have

$$\sec' = \left(\frac{1}{\cos} \right)' = \frac{-\cos'}{\cos^2} = \frac{-(-\sin)}{\cos^2} = \sec \cdot \tan$$

and

$$\cot' = \left(\frac{1}{\tan} \right)' = \frac{-\tan'}{\tan^2} = \frac{-\sec^2 \cdot \cos^2}{\sin^2} = -\csc^2,$$

which proves the rest of the theorem. ■

Depending on how much you trust your memory, you may be able to store all these formulas safely in your brain. At minimum, you should commit the derivatives of sine and cosine to memory, along with the quotient and reciprocal rules. Then you can rederive the rest of the trigonometric derivatives in a matter of seconds. Whatever strategem you employ, a good sanity check to keep in mind

is that the “co-” functions $\cos$, $\csc$, and $\cot$ are the ones whose derivatives have a negative sign in their formulas.

We end off this section by observing a curious pattern. The derivative of $\sin$ is $\cos$ and the derivative of $\cos$ is $-\sin$. Hence $\sin''' = -\cos$ and $\sin^{(4)}$ is $\sin$. Repeating this process again, we see that $\sin^{(n)} = \sin$ whenever n is a multiple of 4. This allows us to compute huge higher derivatives of the sine function (and the cosine function) without having to actually take the derivative so many times. For instance, say we want to know the 439th derivative of $\sin$. By long division, we have

$$439 = 436 + 3 = 109 \cdot 4 + 3.$$

The number 436 is a multiple of 4, meaning that $\sin^{(436)}$ is just $\sin$. Taking the derivative three more times, we get $\sin^{(437)} = \cos$, $\sin^{(438)} = -\sin$, so $\sin^{(439)} = -\cos$.