

Certainty assessment							№ of patients		Effect		Certainty	Importance
№ of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	machine learning prediction	real cumulative cases	Relative (95% CI)	Absolute (95% CI)		
Linear regression												
4	randomised trials	serious ^a	not serious ^a	not serious	not serious ^a	none	21	21	-	SMD 0.17 SD higher (0.43 lower to 0.78 higher)	⊕⊕⊕○ Moderate	CRITICAL
Convolutional neural network with Logistic regression												
1	randomised trials	not serious	not serious	not serious	serious ^b	publication bias strongly suspected ^b	4	4	-	SMD 0 SD (1.39 lower to 1.39 higher)	⊕⊕○○ Low	CRITICAL
Long Short-Term Memory (LSTM) recurrent neural networks (RNN)												
7	randomised trials	serious ^c	not serious	not serious	not serious ^{c,d}	none	68	68	-	SMD 0.49 SD higher (0.14 higher to 0.83 higher)	⊕⊕⊕○ Moderate	CRITICAL
Stochastic, PCA and logistic models												
7	randomised trials	not serious	not serious	not serious	serious ^{a,f}	none	40	40	-	SMD 0 SD (0.44 lower to 0.44 higher)	⊕⊕⊕○ Moderate	IMPORTANT
Time series models												
8	randomised trials	not serious	serious ^a	not serious	serious ^a	none	59	59	-	SMD 0.01 SD lower (0.37 lower to 0.35 higher)	⊕⊕○○ Low	CRITICAL
Multilayer perceptron												
2	randomised trials	serious ^h	not serious	not serious	serious ^h	none	20	20	-	SMD 0.47 SD higher (0.16 lower to 1.1 higher)	⊕⊕○○ Low	CRITICAL
Ensemble family												
4	randomised trials	not serious	not serious	not serious	serious ⁱ	none	20	20	-	SMD 0.01 SD lower (0.63 lower to 0.61 higher)	⊕⊕⊕○ Moderate	IMPORTANT
Reinforcement learning												
3	randomised trials	serious ⁱ	not serious	not serious	serious ⁱ	publication bias strongly suspected ⁱ	0	0	-	SMD 0 SD (0 to 0)	⊕○○○ Very low	NOT IMPORTANT
Gate recurrent unit												
2	randomised trials	not serious	not serious	not serious	serious ^k	publication bias strongly suspected ⁱ	0	0	-	0 (0 to 0)	⊕⊕○○ Low	NOT IMPORTANT