

The Balance Field Equation: Dynamic Order and Recursive Structure Formation

Marcel Theodor Wende
Independent Researcher
ORCID: 0009-0007-4028-2208

8 October 2026

Research manuscript draft

Abstract

The Balance Field Equation (BFG; Balance-Feldgleichung) is developed as a finite recursive operator dynamics on unitary equivalence classes. Its state comprises a Hermitian kernel, positive formation, normalized witness, positive Gram load and persistent carrier. Spectral selection, polar transport, an endogenous metric, cross-weighted channel loads, an ambient reduced singular-value decomposition, isometric R4 realignment and a rank-one seed rule determine a successor or a named terminal outcome. The order principle of unfolding, binding and information is made precise through separately defined trajectory observables rather than additional unspecified forces. Three consequences sharpen the construction: the cross-weights exactly equalize weighted channel loads; formation stock obeys an exact inheritance–loss–seed balance; and every infinite nonterminal autonomous orbit reaches full carrier persistence after finitely many seed events. These results distinguish formation stock, spectral diversity, historical survival and geometric damping. A separately declared constant-input scalar extension has a unique positive globally attractive geometric fixed point, while neutral kernel and formation directions remain. A typed self-similar tensor lift commutes with the full-persistence update, while replicated seed branches can fail through degeneracy. Six simulated levels illustrate that declared hierarchy. A complete regular-stratum differential specification and reproducible synthetic controls are supplied. Physical time, spacetime, quantum measurements, particles, chemistry, biological viability and experience remain explicitly typed bridge requirements. Universal applicability is a research hypothesis, not a conclusion of the matrix identities.

Keywords: Balance Field Equation; recursive order; polar transport; formation balance; persistence; endogenous geometry; viability; physical bridges.

Reading guide and complete contents

All 81 sections and 32 figures of the supplied preprint are retained as Part I. Its former bibliography is consolidated once at the end, preserving reference numbers 1-20. The expanded account continues with Section 82, Equation 121 and Figure 33. Its new atlas adds 56 three-dimensional graphics, including all six simulated self-similar levels. No numbered source section or source figure is omitted.

Part I retains historical declarations and computed residuals as source material; these are not silently re-certified as new results. The original publication date is recorded in Reference 21. A single title and author block identifies the assembled manuscript. Part II contains the new derivations and fresh computational checks. Conditional physics and application models are explicitly labeled and do not establish missing BFG bridge laws.

Part I - Complete supplied mathematical and physical framework

1. Introduction	5
2. Claim-state hierarchy	6
3. Level 0, information, and order	6
4. Canonical finite state	8
5. Spectral selection and first transport	9
6. Formation and witness after selection	10
7. Metric projection and endogenous loads	11
8. Analysis operator, reduced SVD, and endogenous dimension	12
9. SVD gauge and quotient-state dynamics	13
10. Successor geometry and kernel operator	14
11. R4 transport	15
12. Second witness mass and formation inheritance	16
13. Seed rule	17
14. Total stratified update and terminal codes	18
15. Constant-rank polar differential	19
16. Seed differential	20
17. Endogenous geometric feedback and stability	21
18. Discrete and continuous Lyapunov formulations	22
19. Mediated recursive closure	23
20. Admissibility and viability	24
21. Scalar mediator normal form	25
22. Functional state Psi and factor dynamics	26
23. Event time, model time, and physical time	27
24. From internal Gram geometry to spacetime	28
25. Gravitational completion	29
26. Quantum bridge	30
27. Finite quantum discrimination example	31

28. R4 and quantum instruments	32
29. Gauge and particle sector	33
30. Atomic closure	34
31. Molecular and biological closure	35
32. Witness, world, self, and subject structure	36
33. Phenomenal bridge	37
34. Canonical world-formula statement	37
35. Falsifiability and bridge discipline	38
36. Variable and operator register	38
37. Scope of proved statements	39
38. Discussion	39
39. Conclusion	40
40. The BFG as a Fundamental Recursive World Equation	40
41. Physical Observables and Measurement Map	40
42. Complete Canonical Mathematical Specification	43
43. Complete Variable and Operator Dictionary	44
44. Reader's Guide to the Recursive Mechanism	52
45. Detailed Selection Logic	52
46. Why the Metric Projector Is Endogenous	52
47. Load Weighting and Neutral Balance	52
48. Reduced SVD as the Dimension-Formation Step	52
49. Gauge Freedom and Physical Equivalence	52
50. Polar Differential Interpretation of the Three Terms	53
51. R4 as Recursive Re-Alignment	53
52. Formation, Witness, and the Meaning of Seed	53
53. Terminality as Part of the Equation	53
54. What the Scalar Damping Result Does and Does Not Prove	53
55. From Recursive Events to Measured Time	53
56. From Positive Gram Geometry to Lorentzian Spacetime	54
57. Quantum Measurement as a Discriminating Bridge	54
58. Particle, Atomic, Molecular, and Biological Ordering	54
59. Witness to Subject Structure	54
60. How the Physical Measurement Layer Closes a Sector	54
61. Canonical Theorem and Assumption Register	54
62. Full Differential Chain Selection to the Analysis Operator	55
63. SVD Differential in the Complex Case	56
64. R4, Witness, and Formation Differential	56
65. Seed Eigenprojector Derivation	57
66. Algebraic Admissibility Checks	57

67. Detailed Scalar Control Branch	58
68. Physical Measurement Equations	59
69. Falsification Protocol for the World Equation	59
70. Dimensional-Consistency Handbook	60
71. Stratum and Regime Manual	61
72. Proof Appendix Core Invariants	61
73. Measurement Protocol Time and Geometry	62
74. Measurement Protocol Quantum and Particle Physics	63
75. Measurement Protocol Atomic, Molecular, and Biological Sectors	63
76. Measurement Protocol World, Self, and Experience	64
77. Plain-Language Glossary for Readers	64
78. Equation Roadmap Reading One Complete World-Equation Step	65
79. Figure Companion What Each 3D Graphic Means	67
80. Frequently Asked Technical Questions	69
81. Canonical BFG at a Glance	70

Part II - Deepened order, hierarchy and simulations

82. Purpose, provenance and claim discipline	72
83. State space, units and equivalence	72
84. The complete recursive event	73
85. Gauge covariance and the ambient bound	75
86. Order through exact accounting and observable distinctions	75
87. Finite seeding and eventual full persistence	77
88. Stability, viability and a minimal driven control	77
89. Functional closure, hierarchy and event time	79
90. A typed self-similar hierarchy	80
91. Physical sectors and prospective tests	81
92. Synthetic verification and interpretation	82
93. Simulation atlas and stage-by-stage interpretation	83

Appendices and reproducibility

The expanded differential specification, fresh numerical record and draft declarations follow the atlas. The accompanying archive contains LaTeX source, Python code, all 56 new images, raw grid/point-cloud arrays, the original source PDF, and a manifest. Use the PDF bookmarks to jump directly to a numbered section or new figure.

Part I: Mathematical and physical framework

Abstract

This preprint specifies the Balance–Field Equation (BFG) as a finite, stratified recursive operator dynamics on unitary equivalence classes. Its declared state is $\Xi=(K,\rho_F,\rho_W,Y,P)$. Spectral selection, canonical polar transport, an endogenous positive metric, formation-dependent loads, an ambient reduced singular-value decomposition, R4 realignment, formation inheritance and a rank-one seed rule define a unique successor class or a named terminal outcome. The complete analysis operator acts on the ambient input space, not only on the selected carrier. This choice has restrictive consequences: R4 is isometric on the selected carrier, its witness mass is exactly one, every nonterminal successor satisfies $0<Y_+<I$, and the ambient dimension cannot increase. Subsequent nonterminal events preserve witness mass. Geometric self feedback is explicit and produces strict damping on a scalar control branch, but neutral kernel and formation directions prevent a universal strict full-state contraction claim. The manuscript gives typed differentials, proof assumptions, event-time bookkeeping and independently reproducible numerical controls. Physical information, seconds, spacetime and gravity, quantum and particle observables, atoms, molecules, biology and experience require additional specified measurement bridges. These transitions are presented in dependency order and remain open where the finite kernel does not determine them. Computed 3D figures demonstrate declared mathematical families; physical-sector illustrations do not constitute empirical validation.

Keywords: Balance–Field Equation; recursive dynamics; polar decomposition; singular-value decomposition; viability; endogenous geometry; quantum bridge; recursive closure; witness persistence; stratified dynamical systems.

1 Introduction

A candidate world formula must do more than compress many concepts into one symbolic expression. It must specify a state space, an evolution law, admissibility conditions, equivalence relations, regime boundaries, and a map from internal variables to measurable quantities. The present construction takes this requirement literally. The BFG is therefore formulated as a recursive operator system whose central object is an orbit of a finite state under a stratified update. The use of polar decomposition and singular-value structure places the mathematical core in a standard matrix-analytic setting [4–8], while viability and Lyapunov concepts provide the appropriate language for constrained persistence [9].

Three co-authored precursor lines are retained as direct BFG provenance. The Level-0 foundation with Thomas W. Loker motivates a pre-informational admissibility layer [1]. The recursive conjugate-closure work with Luke Kenneth Casson Leighton motivates mediated conjugate recursion [2]. The no-supervenience bridge with Alfredo Pereira Jr. motivates the later distinction between functional closure and phenomenal interpretation [3]. No other BFG manuscript is included in the bibliography.

The objective is to present one canonical finite model class, close its mathematical ambiguities, and expose the remaining physical bridge choices. The paper fixes: (i) a reduced-SVD convention and endogenous successor dimension; (ii) a unitary quotient removing common SVD-frame gauge freedom; (iii) Hermiticity and unit compatibility of the formation operator; (iv) normalized witness input and conditional survival masses; (v) named terminal states; and (vi) separate discrete and continuous stability statements.

2 Claim-state hierarchy

Label	Meaning	Use
Definition	Part of the canonical model class	State spaces, update maps, supports, terminal codes
Assumption	Additional constitutive or physical choice	Bridge laws, Lorentzian completion, quantum-state identification
Theorem/Consequence	Derived under stated hypotheses	Gram positivity, scalar damping, gauge covariance
Open bridge	Not fixed by the finite kernel	Physical clock, spacetime, Standard-Model parameters, phenomenality
Empirical hypothesis	Externally testable identification	Quantum bridge A versus B


Table 1. Canonical classification and definitions.

The distinction is substantive. A polar factor is a standard matrix object [4,5], but identifying a BFG partial isometry with a physical quantum instrument is an additional physical statement. Likewise, a positive Gram operator is mathematically well defined but is not, by positivity alone, a Lorentzian spacetime metric.

3 Level 0, information, and order

Dimensionless order readout

CONDITIONAL READOUT



phi_E=(E_exp/E_bind)(I/I_0); ratios are independent inputs, not identified Xi data.

Figure 1. Dimensionless order readout. Conditional readout. $\varphi_E=(E_{\text{exp}}/E_{\text{bind}})(I/I_0)$; ratios are independent inputs, not identified Ξ data. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

3.1 Level-0 domain

$$\Phi_0 \in X_0 \setminus \{1\}$$

Φ_0 denotes a pre-operational Level-0 domain rather than the later measurable scalar order readout. The co-authored Level-0 construction motivates admissible difference, recurrence, and identity retention before coarse-grained information and physics [1].

3.2 Information

$$I = \log_b |C_\theta(X)| \quad (2)$$

Fix a finite nonempty configuration class with $1 \leq |C_\theta(X)| < \infty$, a threshold or metric θ , logarithm base $b > 1$, and reference $I_0 > 0$. The object X and the counting procedure must be specified independently. This configuration-counting measure differs from Shannon entropy [10]. Neither the count nor its map from Ξ is supplied by the finite kernel.

3.3 Dimensionless order readouts

$$\varphi_N = \frac{N_+}{N_-} \frac{I}{I_0} \quad (3)$$

$$\varphi_E = \frac{E_{\text{exp}}}{E_{\text{bind}}} \frac{I}{I_0} \quad (4)$$

Both readouts are dimensionless for matching numerator and denominator units, with $N_- \rightarrow 0$, $E_{\text{bind}} > 0$ and $I_0 > 0$. They are alternative operational candidates. Φ_0 is a Level-0 object, not a scalar multiplier. No equality between Φ_0 and φ_N or φ_E is assumed.

3.4 Stock and rate

$$q_{\text{info}} = \varphi I \quad (5)$$

$$I_B := dq_{\text{info}}/dt \quad (6)$$

$$dq_{\text{info}}/dt = a_{\text{info}} - \gamma_0 (1 - b_{\text{feedback}}) q_{\text{info}} \quad (7)$$


$$q_{\text{info}}^* = \frac{a_{\text{info}}}{\gamma_0 (1 - b_{\text{feedback}})} \quad (8)$$

Here φ denotes one explicitly chosen scalar readout, not Φ_0 . For constant coefficients, $\gamma_0 > 0$ and $b_{\text{feedback}} < 1$, the stated stock is attractive and $I_B^* = 0$. At $b_{\text{feedback}} = 1$, $a_{\text{info}} = 0$ every constant stock is stationary; $a_{\text{info}} \neq 0$ gives linear change. For $b_{\text{feedback}} > 1$ the homogeneous part amplifies. Time-dependent coefficients require the corresponding nonautonomous solution. q_{info} , a_{info} and b_{feedback} are distinct from the selection projector Q , analysis operator A and logarithm base b .

4 Canonical finite state

Positive internal geometry

COMPUTED MATRIX FAMILY



$Y = \text{diag}(y_1, y_2)$; $G = I + Y$; the positive metric is internal, not spacetime.

Figure 2. Positive internal geometry. Computed matrix family. $Y = \text{diag}(y_1, y_2)$; $G = I + Y$; the positive metric is internal, not spacetime. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

$$\Xi_k = (K_k, \rho_{F,k}, \rho_{W,k}, Y_k, P_k) \quad (9)$$

$$K_k = K_k^\dagger; \rho_{F,k} \geq 0; \rho_{W,k} \geq 0; \text{tr}(\rho_{W,k}) = 1; Y_k = Y_k^\dagger \geq 0; P_k = P_k^\dagger = P_k^2 \quad (10)$$

At event k , all operators act on a finite complex Hilbert space H_k of dimension n_k . K is Hermitian. Formation ρ_F is positive and need not be normalized. Witness ρ_W is positive and normalized at every nonterminal input. Y is a positive semidefinite Gram/load operator and P an orthogonal persistence projector.


The Level-0 interpretation motivates the model but does not determine its input tuple. A construction $\Xi = J_0(\Phi_0)$ remains an open bridge. The finite state starts only after its five operator components and carrier have been declared.

K and ρ_F share a formation unit E_F in the canonical dimensionally matched model. If a future physical interpretation assigns different units, a fixed positive conversion factor κ_F must enter the seed term; no state-dependent unit conversion is permitted.

5 Spectral selection and first transport

Spectral selection boundary

COMPUTED PROJECTOR



$Q=1_{[0,1)}(Y)$; threshold 1 defines a discontinuous change of selected rank.

Figure 3. Spectral selection boundary. Computed projector. $Q=1_{[0,1)}(Y)$; threshold 1 defines a discontinuous change of selected rank. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

$$C_k = (I + Y_k)^{-1} \quad (11)$$

$$B_k = Y_k C_k = I - C_k \quad (12)$$

$$G_k = I + Y_k \quad (13)$$

Because Y_k is positive semidefinite, G_k is positive definite. This is the internal Gram geometry.

$$Q_k = 1_{[0,1)}(Y_k) \quad (14)$$

$$X_{\text{sel},k} = Q_k P_k \quad (15)$$

$$T_{\text{sel},k} = \text{polar}(X_{\text{sel},k}) \quad (16)$$


$$S_{\text{sel},k}^- = T_{\text{sel},k}^\dagger T_{\text{sel},k}; P_{\text{sel},k} = S_{\text{sel},k}^+ = T_{\text{sel},k} T_{\text{sel},k}^\dagger \quad (17)$$

The source and target supports are distinct. The polar factor is a partial isometry on the actual supports, consistent with standard polar-decomposition theory [4,5].

6 Formation and witness after selection

Witness mass before normalization

COMPUTED CONTROL FAMILY



$Y=\text{diag}(1/2,2)$, $P=vv^T$ and $W=\text{diag}(p,1-p)$; polar selection has source P .

Figure 4. Witness mass before normalization. Computed control family. $Y=\text{diag}(1/2,2)$, $P=vv^T$ and $W=\text{diag}(p,1-p)$; polar selection has source P . Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

$$\rho_{F,k}^{\text{sel}} = T_{\text{sel},k} \rho_{F,k} T_{\text{sel},k}^\dagger \quad (18)$$

$$\rho_{\text{un},k}^{\text{sel}} = T_{\text{sel},k} \rho_{W,k} T_{\text{sel},k}^\dagger \quad (19)$$

$$m_{\text{sel},k} = \text{tr}(S_{\text{sel},k} \rho_{W,k}) \quad (20)$$

$$\rho_{W,k}^{\text{sel}} = \rho_{\text{un},k}^{\text{sel}} / m_{\text{sel},k}, \text{ if } m_{\text{sel},k} > 0 \quad (21)$$

Since $\rho_{W,k}$ is normalized and $0 \leq S_{\text{sel}} \leq I$, $0 \leq m_{\text{sel}} \leq 1$. Thus m_{sel} has a genuine conditional-survival interpretation inside the canonical kernel. This is not yet a physical Born probability.


7 Metric projection and endogenous loads

Endogenous formation loads

COMPUTED AMBIENT KERNEL

Value range
0.5216 to 0.9181

Height: C channel load



$Y=\text{diag}(y1,0.72)$; $P=F=W=vv^T$; $v=(\cos \theta, \sin \theta)$; formation unit $E_F=1$.

Figure 5. Endogenous formation loads. Computed ambient kernel. $Y=\text{diag}(y1,0.72)$; $P=F=W=vv^T$; $v=(\cos \theta, \sin \theta)$; formation unit $E_F=1$. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

$$Z_k + Z_k = I ; Z_k Z_k^\dagger = P_{\text{sel},k} \quad (22)$$

$$P_{G,k} = Z_k (Z_k + G_k Z_k)^{-1} Z_k + G_k \quad (23)$$

$$\Lambda_{C,k} = \text{tr} (G_k P_{G,k} C_k \rho_{F,k}^{\text{sel}} C_k P_{G,k}^\dagger) \quad (24)$$

$$\Lambda_{B,k} = \text{tr} (G_k P_{G,k} B_k \rho_{F,k}^{\text{sel}} B_k P_{G,k}^\dagger) \quad (25)$$

$$\alpha_k = \frac{\Lambda_{B,k}}{\Lambda_{C,k} + \Lambda_{B,k}} \quad (26)$$


$$\beta_k = \frac{\Lambda_{C,k}}{\Lambda_{C,k} + \Lambda_{B,k}} \quad (27)$$

On the dual-load branch $\Lambda_C > 0$ and $\Lambda_B > 0$, α and β lie in $(0,1)$ and sum to one.

8 Analysis operator, reduced SVD, and endogenous dimension

Full ambient analysis

COMPUTED AMBIENT KERNEL



A:H to H direct sum H; no restriction to the selected rank-one carrier.

Figure 6. Full ambient analysis. Computed ambient kernel. A:H to H direct sum H; no restriction to the selected rank-one carrier. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

$$A_k = \begin{bmatrix} \sqrt{\alpha_k} P_{G,k} C_k \\ \sqrt{\beta_k} P_{G,k} B_k \end{bmatrix} \quad (28)$$

Let $r_k = \text{rank}(A_k) \geq 1$. The canonical convention uses only the reduced rank- r_k SVD:

$$A_k = U_k \Sigma_k V_k^\dagger \quad (29)$$

$$U_k : C^{r_k} \rightarrow H_k \oplus H_k; V_k : C^{r_k} \rightarrow H_k; \Sigma_k = \text{diag}(\sigma_1, \dots, \sigma_r), \sigma_i > 0 \quad (30)$$

$$H_{k+1} := C^{r_k}; n_{k+1} = r_k \quad (31)$$

The analysis domain is the full ambient H_k . No restriction $A_k Z_k$ is made. Its codomain is $H_k \oplus H_k$; U and V are column isometries. The successor dimension satisfies $\text{rank}(P_{\text{sel}}) \leq r_k \leq n_k$. It can exceed the selected-carrier dimension, but cannot exceed the ambient input dimension. Null singular directions are omitted. Restricting the domain would define a different kernel and can change the seed complement.


9 SVD gauge and quotient-state dynamics

Reduced singular spectrum

COMPUTED AMBIENT KERNEL

Value range
0.000158 to 0.1401

Height: Second singular
value



The second singular value can vanish on a rank boundary; an arbitrary full-SVD padding is excluded.

Figure 7. Reduced singular spectrum. Computed ambient kernel. The second singular value can vanish on a rank boundary; an arbitrary full-SVD padding is excluded. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

For a simple positive singular value, a common phase remains free. For an exactly degenerate singular block, a unitary basis rotation inside that block remains free. These freedoms are treated as coordinate gauge rather than physical multiplicity.

$$\backslash(K, \rho_F, \rho_W, Y, P) \sim \backslash(W^\dagger K W, W^\dagger \rho_F W, W^\dagger \rho_W W, W^\dagger Y W, W^\dagger P W) \backslash, W \text{ unitary} \quad \backslash(32 \backslash)$$

For a block-unitary W satisfying $W \Sigma = \Sigma W$:

$$U' = U W ; V' = V W ; \Sigma' = W^\dagger \Sigma W = \Sigma \quad \backslash(33 \backslash)$$

$$K_+' = W^\dagger K_+ W ; Y_+' = W^\dagger Y_+ W \quad \backslash(34 \backslash)$$

The R_4 overlap transforms as $X_{R_4}' = W^\dagger X_{R_4}$. Unitary covariance of the polar decomposition gives $T_4' = W^\dagger T_4$, and hence all target-space successor quantities transform by the same conjugation. The mathematical successor is therefore the equivalence class $[\Xi_{k+1}]$, not a privileged matrix representation.

Input covariance is separate from SVD-frame covariance. For a unitary L on H , simultaneous input conjugation gives $A' = (L^\dagger \oplus L^\dagger) A L$. Compatible SVD frames are $U' = (L^\dagger \oplus L^\dagger) U$ and $V' = L^\dagger V$. K_+ and Y_+ then agree, and R_4 transforms consistently. This proves independence of the ambient input basis as well as of the allowed target SVD gauge.

$$U_{\text{bar}}_{\text{BFG}} : [\Xi_k] \rightarrow [\Xi_{k+1}] \quad \backslash(35 \backslash)$$

This quotient construction solves the definition-level phase/basis ambiguity. It does not claim differentiability of individually sorted singular vectors at eigenvalue splitting. Kato and Davis–Kahan theory instead support the use of invariant subspaces and separated spectral clusters [6,7].

For local differential work on a simple singular-value stratum, with $M = U^\dagger (\delta A) V$:

$$\delta \sigma_i = \text{Re } M_{ii} \quad (36)$$


$$(\Omega_U)_{ii} - (\Omega_V)_{ii} = i \text{Im}(M_{ii}) / \sigma_i \quad (37)$$

Only the common phase is free; setting both diagonal gauge connections to zero is generally impossible.

10 Successor geometry and kernel operator

Successor Gram bound

COMPUTED AMBIENT KERNEL



$Y = \text{diag}(y1, y2)$, $P = F = I$; coupled global formation loads determine alpha and beta.

Figure 8. Successor Gram bound. Computed ambient kernel. $Y = \text{diag}(y1, y2)$, $P = F = I$; coupled global formation loads determine α and β . Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

$$Y_{k+1} = \Sigma_k^2 \quad (38)$$


$$K_{k+1} = U_k \dagger (K_k \oplus K_k) U_k \quad (39)$$

Theorem 1. On the positive dual-load branch, the reduced successor satisfies $0 < Y_{k+1} < I_{r_k}$. In particular $G_{k+1} > I$. A proof of this stronger bound, including its limits for full-state stability, follows in Section 17. The input class permits $Y \geq 0$; this is a declared model class, not an assertion that every precursor used the same domain.

11 R4 transport

R4 preserves selected witness mass

COMPUTED AMBIENT KERNEL



$m_4=1$ to numerical precision; the exact proof uses $\text{Ran}(P_{\text{sel}})$ contained in $\text{Ran}(A^\dagger)$.

Figure 9. R4 preserves selected witness mass. Computed ambient kernel. $m_4=1$ to numerical precision; the exact proof uses $\text{Ran}(P_{\text{sel}})$ contained in $\text{Ran}(A^\dagger)$. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

$$X_{R4,k} = V_k \dagger P_{\text{sel},k} \quad (40)$$

$$T_{4,k} = \text{polar}(X_{R4,k}) \quad (41)$$


$$S_{4,k}^- = T_{4,k} \dagger T_{4,k}; S_{4,k}^+ = T_{4,k} T_{4,k} \dagger \quad (42)$$

For this ambient kernel, X_{R4} is already a partial isometry: $T_4 = V \dagger P_{\text{sel}}$, $S_4^- = P_{\text{sel}}$ and $S_4^+ = V \dagger P_{\text{sel}} V$. The positive C-channel puts the entire selected carrier in $\text{Ran}(A^\dagger)$. R4 therefore realigns that carrier without a second loss. General rank-loss sensitivity of polar maps remains relevant to selection and to other declared models.

12 Second witness mass and formation inheritance

Original mass and conditioned trace

COMPUTED BOOKKEEPING



$w_{\text{next}} = w \cdot m_{\text{sel}}$; conditioned witness trace remains one and cannot encode this product.

Figure 10. Original mass and conditioned trace. Computed bookkeeping. $w_{\text{next}} = w \cdot m_{\text{sel}}$; conditioned witness trace remains one and cannot encode this product. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

$$m_{4,k} = \text{tr}(S_{4,k} \cdot \rho_{W,k}^{\text{sel}}) \quad (43)$$

$$\rho_{W,k+1} = T_{4,k} \rho_{W,k}^{\text{sel}} T_{4,k}^\dagger / m_{4,k}, \text{ if } m_{4,k} > 0 \quad (44)$$

$$m_{\text{tot},k} = m_{\text{sel},k} m_{4,k} \quad (45)$$

$$\rho_{F,k}^{\text{inh}} = T_{4,k} \rho_{F,k}^{\text{sel}} T_{4,k}^\dagger \quad (46)$$


In the present kernel $m_4=1$ exactly, hence $m_{\text{tot}}=m_{\text{sel}}$. The normalized successor does not contain the historical mass. Record $w_{k+1} = w_k m_{\text{sel},k}$, $w_0=1$, as a separate trajectory output. It is not a sixth component silently inserted into Ξ . Directed transport history, when needed, is also recorded separately. Inherited formation satisfies $\text{tr}(\rho_F^{\text{inh}}) = \text{tr}(\rho_F^{\text{sel}})$.

Historical-mass nonidentifiability from the successor: take $Y=\text{diag}(1/2,2)$, $P=I$, $K=-I$, $F=I$ and $W=\text{diag}(p,1-p)$, $0 < p \leq 1$. Selection gives $m_{\text{sel}}=p$, but every p yields the same one-dimensional normalized successor. Therefore Ξ_+ alone cannot recover w or the initial surviving mass.

13 Seed rule

Seed formation on a negative complement

COMPUTED SPECTRAL CONTROL



A one-dimensional complement with $\lambda = -x$ gives seed trace $\kappa \cdot x$; each κ is a separate fixed-unit model.

Figure 11. Seed formation on a negative complement. Computed spectral control. A one-dimensional complement with $\lambda = -x$ gives seed trace $\kappa \cdot x$; each κ is a separate fixed-unit model. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

$$E_k = I_{r_k} - S_{4,k}^+ \quad (47)$$

If $E_k = 0$, no seed is added. If $E_k \neq 0$, choose an orthonormal frame Z_E for $\text{Ran}(E_k)$ and define:

$$H_{E,k} = Z_{E,k}^\dagger K_{k+1} Z_{E,k} \quad (48)$$

Because K is Hermitian, K_{k+1} and H_E are Hermitian, so the seed spectrum is real.

If H_E has a simple isolated negative minimum $\lambda_k < 0$ with rank-one projector π_k :

$$\Pi_{\text{seed},k} = Z_{E,k} \pi_k Z_{E,k}^\dagger \quad (49)$$

$$\rho_{\text{seed},k} = -\kappa_F \lambda_k \Pi_{\text{seed},k} \quad (50)$$

$$\rho_{F,k+1} = \rho_{F,k}^{\text{inh}} + \rho_{\text{seed},k} \quad (51)$$


$$P_{k+1} = S_{4,k}^+ + \Pi_{\text{seed},k} \quad (52)$$

For the dimensionally matched canonical class $\kappa_F = 1$. Orthogonality $\Pi_{\text{seed}} \leq E$ implies $S_4^+ \Pi_{\text{seed}} = 0$ and $P_{k+1}^2 = P_{k+1}$.

14 Total stratified update and terminal codes

Terminal boundary of the seed rule

COMPUTED RULE



For a nonempty complement, $\lambda < 0$ and a simple minimum are required; $\lambda \geq 0$ is terminal.

Figure 12. Terminal boundary of the seed rule. Computed rule. For a nonempty complement, $\lambda < 0$ and a simple minimum are required; $\lambda \geq 0$ is terminal. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

The canonical dynamics is total only after adjoining explicit terminal outcomes.

$$U \sim_{\text{BFG}} : X_{\Xi} \rightarrow \backslash (X_{\Xi} / \text{Unitary}) \backslash \text{disjoint}_{\text{union}} T_{\text{terminal}} \backslash (53)$$

Code	Trigger	Meaning
T_sel0	$\text{rank}(\text{QP})=0$	No selected persistent carrier
T_w1	$m_{\text{sel}}=0$	First witness selection fails
T_load	$\Lambda_C \leq 0$ or $\Lambda_B \leq 0$	Positive dual-load class not satisfied
T_seednone	$E \neq 0$ and $\lambda_{\min} \geq 0$	Required negative seed absent
T_seeddeg	$E \neq 0$ and negative lowest eigenvalue is multiple	No unique rank-one seed

Table 2. Canonical classification and definitions.

The no-seed branch exists only when $E=0$. If $E \neq 0$, a simple strictly negative lowest eigenvalue is required; otherwise the event terminates.

$$E=0: \rho_{\text{seed}}=0, \Pi_{\text{seed}}=0 \quad (54a)$$

$$E \neq 0, \lambda_{\min} \geq 0: T_{\text{seednone}} \quad (54b)$$

$$E \neq 0, \lambda_{\min} < 0 \text{ multiple}: T_{\text{seeddeg}} \quad (54c)$$

No arbitrary tie-break vector is introduced at a multiply degenerate negative minimum.


15 Constant-rank polar differential

Polar sensitivity bound

COMPUTED BOUND

Value range
0.2609 to 34.5

Height: Upper bound on
variation



3 times $\|\delta X\|_2$ divided by $\sigma_{\min,+}$; an upper bound is not an attained response.

Figure 13. Polar sensitivity bound. Computed bound. 3 times $\|\delta X\|_2$ divided by $\sigma_{\min,+}$; an upper bound is not an attained response. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Let $X=TH$ be a C^1 curve on a constant-rank stratum. Choose the canonical polar factor with $\ker(T)=\ker(X)$. Set $E_X=\delta X$, $H=(X^\dagger X)^{1/2}$, $S_-=T^\dagger T$ and $S_+=TT^\dagger$. The following derivatives are tangent derivatives on that stratum:

$$(I-S_+)E_X(I-S_-)=0 \quad (55)$$

The skew-Hermitian connection is supported inside the source: $\Omega=S_-\Omega S_-$. On that support it is uniquely fixed by the Sylvester equation:

$$H\Omega + \Omega H = S_-(T^\dagger E_X - E_X^\dagger T)S_- \quad (56)$$

The polar derivative is:

$$\delta T = T\Omega + (I-S_+)E_{XH}^+ + TH^+T^\dagger E_X(I-S_-) \quad (57)$$

and the support derivatives are:

$$\delta S_+ = (I-S_+)E_{XH}^+T^\dagger + TH^+E_X^\dagger(I-S_+) \quad (58)$$

$$\delta S_- = H^+T^\dagger E_X(I-S_-) + (I-S_-)E_X^\dagger TH^+ \quad (59)$$

A conservative bound is:

$$\|\delta T\|_2 \leq 3\|E_X\|_2/\sigma_{\min,+}(X) \quad (60)$$

This is consistent with the general sensitivity of polar factors to small singular values [4,5]. A diverging upper bound as $\sigma_k \rightarrow 0$ does not imply that every admissible curve diverges; it signals loss of a uniform local bound.


16 Seed differential

Seed eigenprojector sensitivity

CONDITIONAL PERTURBATION SCALE

Value range
0.08696 to 11.5

Height: Gap sensitivity
scale



Illustrates inverse-gap scaling of the reduced resolvent; not a full derivative of an arbitrary kernel.

Figure 14. Seed eigenprojector sensitivity. Conditional perturbation scale. Illustrates inverse-gap scaling of the reduced resolvent; not a full derivative of an arbitrary kernel. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

With $E = I - S_4^+$, choose a horizontal orthonormal frame Z_E :

$$\delta Z_E = (\delta E) Z_E \quad (61)$$

Then:

$$\delta H_E = Z_E^\dagger [(\delta E) K_+ + \delta K_+ + K_+ (\delta E)] Z_E \quad (62)$$

For a simple negative eigenvalue λ with $\pi = vv^\dagger$:

$$\delta \lambda = \text{tr}(\pi \delta H_E) \quad (63)$$

$$R_\lambda = (H_E - \lambda I)^+; R_\lambda v = 0 \quad (64)$$

$$\delta \pi = -R_\lambda (\delta H_E) \pi - \pi (\delta H_E) R_\lambda \quad (65)$$

$$\delta \Pi_{\text{seed}} = (\delta E) \Pi_{\text{seed}} + \Pi_{\text{seed}} (\delta E) + Z_E (\delta \pi) Z_E^\dagger \quad (66)$$


$$\delta \rho_{\text{seed}} = -\kappa_F [(\delta \lambda) \Pi_{\text{seed}} + \lambda \delta \Pi_{\text{seed}}] \quad (67)$$

At complement dimension one, $R_\lambda = 0$.

17 Endogenous geometric feedback and stability

Geometric self feedback and damping

EXACT SCALAR KERNEL



$P=W=1$ and $F>0$; $f(y)=2y^2/[(1+y)^2(1+y^2)]$; K and F remain neutral.

Figure 15. Geometric self feedback and damping. Exact scalar kernel. $P=W=1$ and $F>0$; $f(y)=2y^2/[(1+y)^2(1+y^2)]$; K and F remain neutral. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

The closed internal feedback loop is:

$$Y_k \rightarrow G_k / P_{G,k} \rightarrow \text{Loads} \rightarrow \alpha_k, \beta_k \rightarrow A_k \rightarrow \Sigma_k \rightarrow Y_{k+1} \quad (68)$$

This is a genuine endogenous geometric feedback: the current Gram geometry changes the analysis that produces the next Gram geometry. It is not, by itself, a theorem of full-state contraction.

Ambient bound. Let $D=Z^\dagger GZ$. Since $GC=I$ and $GB=Y$, $P_G C=Z D^{-1} Z^\dagger$ and $P_G B=Z D^{-1} Z^\dagger Y$. The first map has range exactly $\text{Ran}(P_{\text{sel}})$. Positive α gives $\text{Ran}(P_{\text{sel}}) \subset \text{Ran}(A^\dagger)$, proving the R_4 identities and $m_4=1$.

Let η be the largest selected Y eigenvalue; $\eta<1$ and $D \geq I$. Then $\|P_G C\|_2 \leq 1$ and $\|P_G B\|_2 \leq \eta$. Hence $\|A\|_2^2 \leq \alpha + \beta \eta^2 < 1$. All reduced singular values are positive, so $0 < Y_+ < I$. This bounds the successor Gram operator, not the full endogenous Jacobian.

After a nonterminal event $Q_+=I$ and the successor witness is supported on P_+ . The following selection is $T_{\text{sel}} = P_+$ and has mass one. Thus every subsequent nonterminal event preserves original witness mass. The model cannot represent arbitrary repeated lossy disturbances without an additional declared perturbation or update rule.

On the independently checked scalar full-persistence branch:

$$f(y) = \frac{2y^2}{(1+y)^2(1+y^2)} \quad (69)$$

For $y>0$:

$$y-f(y) = y[(1+y)^2(1+y^2)-2y]/[(1+y)^2(1+y^2)] > 0 \quad (70)$$

Hence, on the nonterminal selection-first branch $0< y < 1$:


$$0 < f(y) < y \quad (71)$$

The same branch contains neutral K/F directions in which $K_+ = K$ and $F_+ = F$. Therefore no strict contraction theorem for the complete tuple can hold in every fixed positive product norm on that branch. Positivity preservation, scalar damping, transverse contraction, and full-state contraction are distinct statements.

18 Discrete and continuous Lyapunov formulations

Conditional transverse Lyapunov decay

CONDITIONAL STABILITY MODEL



Assumes $V = z^2$ and $V_{\text{next}} = (1-r)V$; no such global BFG certificate is constructed here.

Figure 16. Conditional transverse Lyapunov decay. Conditional stability model. Assumes $V = z^2$ and $V_{\text{next}} = (1-r)V$; no such global BFG certificate is constructed here. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Let M be an invariant neutral set and $\|z\|$ the quotient distance to M , or a specified transverse coordinate with equivalent distance bounds. On a forward-invariant, fixed-dimensional regular region R with all forward iterates defined, assume positive constants c_1, c_2, c_3 and:

$$c_1 \|z\|^2 \leq V(\Xi) \leq c_2 \|z\|^2 \quad (72)$$

$$\Delta V = V(U_{\text{BFG}}(\Xi)) - V(\Xi) \leq -c_3 \|z\|^2 \quad (73)$$

These are assumptions on a chosen V and region, not a constructed stability certificate for the BFG kernel. They imply exponential transverse decay in event count when the distance bounds hold uniformly: $V_{k+1} \leq (1-c_3/c_2) V_k$. The inequality requires a nonnegative compatible factor, with the zero-factor case giving immediate arrival. Existence, forward invariance and quotient regularity must be checked separately.

A continuous projected model is a different object:

$$d\Psi/dt = F_\Psi(\Psi) \quad (74)$$

for which:


$$dV/dt = DV(\Psi) F_\Psi(\Psi) \quad (75)$$

A clock does not by itself construct F_ψ . Viability theory supplies the appropriate equation for state constraints and viable trajectories [9].

19 Mediated recursive closure

Mediator amplitude and power

COMPUTED MEDIATOR CONTROL



$N_B R N_B^\dagger = n^2 R$; amplitude mediation uses $N_A = n^2$, not $N_A = n$.

Figure 17. Mediator amplitude and power. Computed mediator control. $N_B R N_B^\dagger = n^2 R$; amplitude mediation uses $N_A = n^2$, not $N_A = n$. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

For a positive semidefinite self-adjoint mediator N_A on the declared operator space, its unique positive square root gives the amplitude-mediated class:

$$R_C^A = N_A^{1/2} R N_A^{1/2} \quad (76)$$

The direct minimal class is:

$$R_C^B = N_B R N_B^\dagger \quad (77)$$

At $R=I$ the second class gives $N_B N_B^\dagger$, not N_B^2 unless N_B is self-adjoint. The two classes coincide under the explicit identification $N_B = N_A^{1/2}$. The conjugate-closure source line is associated with the Wende–Leighton precursor [2].

A spectral-radius condition is informative but not sufficient for existence by itself:

$$\rho(R_C) < 1, \rho(R_C) \approx 1, \rho(R_C) > 1 \quad (78)$$


These regimes respectively suggest subcritical damping, near-neutral persistence, and possible amplification only after the operator space, norm, Jordan structure, and admissibility conditions are specified.

A fixed finite linear operator with spectral radius below one has powers tending to zero. This frozen-operator statement does not imply $\|D U_{BFG}\| < 1$ for an endogenously changing nonlinear kernel. Norm bounds and Jordan structure determine quantitative transient estimates.

20 Admissibility and viability

Finite horizon viability gate

SCHEMATIC CONSTRAINT MODEL



A margin diagram only; actual viability requires the unique nonterminal BFG trajectory and specified gates.

Figure 18. Finite horizon viability gate. Schematic constraint model. A margin diagram only; actual viability requires the unique nonterminal BFG trajectory and specified gates. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

$$S_{\text{adm}} = S_{\text{balance}} \cap S_{\text{feedback}} \cap S_{\text{spectral}} \cap S_{\text{interface}} \cap S_{\text{regular}} \cap S_{\text{stability}} \quad (79)$$

Static membership in S_{adm} is not identical to forward invariance. The viability kernel is:


$$K_V = \{x_0 : U_{\text{BFG}}^k(x_0) \text{ lies in } S_{\text{adm}} \text{ and is nonterminal for every } k \geq 0\} \quad (80)$$

This is the viability set of the deterministic kernel. Terminal outcomes count as failure to persist. For a finite horizon restrict k to that horizon. An existential choice among trajectories requires an explicitly controlled or set-valued extension. The gates in (79) are an organizational list; each application must specify the actual constraint sets [9].

21 Scalar mediator normal form

Scalar normal form equilibrium

COMPUTED CONDITIONAL ODE



$a=1.5$, $\mu>0$, $b<1$; this positive branch exists because $a>1-b$.

Figure 19. Scalar normal form equilibrium. Computed conditional ode. $a=1.5$, $\mu>0$, $b<1$; this positive branch exists because $a>1-b$. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

$$dq/dt = a - (1-b + \mu n)q \quad (81)$$

$$\varepsilon dn/dt = q - 1 - n \quad (82)$$

At a positive fixed point $q^* = 1 + n^*$ and $r^* = 1 - b + \mu n^* = a/q^* > 0$. The Jacobian has:

$$\text{tr } J^* = -r^* - 1/\varepsilon < 0 \quad (83)$$

$$\det J^* = \frac{r^* + \mu q^*}{\varepsilon} > 0 \quad (84)$$

Conditional on existence of a positive fixed point, $a, \mu, \varepsilon > 0$ give negative trace and positive determinant and hence local asymptotic stability. For $n^* > 0$, existence requires $a > 1 - b$: the fixed-point equation $a = (1 + n^*)(1 - b + \mu n^*)$ has a unique positive solution with $r^* > 0$ under this condition. In particular $b < 1$ and small a need not admit a positive fixed point. Varying ε does not create a Hopf threshold on this unchanged positive branch.

For the additional quadratic overconstraint model $d(n) = 1 - \zeta n^2$ in the overcritical regime:


$$n_{\min} = (b-1)/\mu ; n_{\max} = \sqrt{(1-d_{\min})/\zeta} \quad (85)$$

with $b > 1$, $\mu > 0$, $\zeta > 0$ and $d_{\min} < 1$. The open static corridor exists when $n_{\min} < n_{\max}$. This static corridor is not the viability kernel.

22 Functional state Psi and factor dynamics

A projection must close its fibers

SCHEMATIC COUNTERMODEL



Projection $\Pi(x,h)=x$ with successor $x+h$ is not closed; this is a fiber test, not a BFG bridge.

Figure 20. A projection must close its fibers. Schematic countermodel. Projection $\Pi(x,h)=x$ with successor $x+h$ is not closed; this is a fiber test, not a BFG bridge. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

$$\Psi = (\varphi_{\text{fun}}, I_{\text{fun}}, A_{\text{fun}}, G_{\text{fun}}, \Omega_{\text{fun}}, T_{\text{fun}}, S_{\text{fun}}, M_{\text{fun}}, C_{\text{fun}}) \quad (86)$$

A closed discrete readout dynamics requires a typed projection $\Pi: X_{\Xi} \rightarrow X_{\Psi}$ satisfying:


$$\Pi(x) = \Pi(x') \Rightarrow \Pi(U_{\text{BFG}} x) = \Pi(U_{\text{BFG}} x') \quad (87)$$

The subscripts distinguish functional roles from kernel operators and from Φ_0 . φ_{fun} may equal a selected scalar order readout only under a specified bridge. The fiber criterion guarantees a well-defined discrete factor on $\Pi(X_{\Xi})$, not smoothness or physical identification. Terminality must also be constant on fibers. No such concrete Π is constructed here.

23 Event time, model time, and physical time

Event count without calibrated seconds

EXACT SCALAR KERNEL



One reclosure event is counted; no conversion to physical seconds is claimed.

Figure 21. Event count without calibrated seconds. Exact scalar kernel. One reclosure event is counted; no conversion to physical seconds is claimed. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

The kernel index k is an event/reclosure count:

$$k \in \mathbb{N}_0 \setminus \{88\}$$

A physical clock is an additional positive map:

$$T_{\text{clock}} : X_{\Xi} / \text{Unitary} \rightarrow \mathbb{R}_{>0} \quad (89)$$

$$t_k = t_0 + \sum_{j < k} T_{\text{clock}}([\Xi_j]) \quad (90)$$

For a given autonomous continuous flow ϕ_F^t , the correct compatibility condition is:


$$\Pi(U_{\text{BFG}} \Xi) = \phi_F^{T_{\text{clock}}([\Xi])}(\Pi(\Xi)) \quad (91)$$

A clock is a quotient-invariant, independently calibrated extra map. Positive durations alone do not exclude Zeno accumulation: $\sum_k T_k = 2^{-k}$ has finite sum. Unbounded physical continuation additionally requires $\sum_k T_k = \infty$. A continuous generator and the endpoint-flow equality require independent construction. Clock constancy on Π -fibers is necessary only when a projected clock depending on Ψ alone is demanded.

24 From internal Gram geometry to spacetime

A curved geometry bridge illustration

SCHEMATIC EMBEDDING



An illustrative surface is not a Lorentz metric, gravitational field equation or prediction.

Figure 22. A curved geometry bridge illustration. Schematic embedding. An illustrative surface is not a Lorentz metric, gravitational field equation or prediction. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

The internal metric $G=I+Y$ is positive definite. A Lorentzian spacetime metric belongs to a different type of object: it requires a manifold, dimension, signature, connection, and curvature. One possible additional completion is the rank-one disformal form studied in gravitational theory [13]:

$$g_{\mu\nu} = \eta_{\mu\nu} + \lambda \partial_{\mu} \phi \partial_{\nu} \phi \quad (92)$$

with η signature $(-, +, +, +)$. Define $X_{\eta} = \eta^{\mu\nu} \partial_{\mu} \phi \partial_{\nu} \phi$. Then:

$$\det g = \det \eta \cdot (1 + \lambda X_{\eta}) \quad (93)$$

$$g^{\mu\nu} = \eta^{\mu\nu} - \lambda \partial^{\mu} \phi \partial^{\nu} \phi / (1 + \lambda X_{\eta}) \quad (94)$$

The Lorentz-connected nondegeneracy condition is:


$$1 + \lambda X_{\eta} > 0 \quad (95)$$

For $\eta = \text{diag}(-1, 1, 1, 1)$ in Cartesian coordinates, $\det \eta = -1$. In natural units with $[\phi] = M$ and $[\partial \phi] = M^2$, $[\lambda] = M^{-4}$. This rank-one completion is additional physics; it does not derive spacetime from the positive Gram operator.

25 Gravitational completion

Geometric feedback requires dynamics

ASSUMED POTENTIAL MODEL



$V=x^2+y^2$ is a pedagogical potential; stability of spacetime must follow from a specified action and bridge.

Figure 23. Geometric feedback requires dynamics. Assumed potential model. $V=x^2+y^2$ is a pedagogical potential; stability of spacetime must follow from a specified action and bridge. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

A canonical real scalar field with $(-+++)$ convention can be coupled through:


$$S = \int \sqrt{-g} \left[\frac{M_{\text{pl}}^2}{2} R - \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) \right] d^4x + S_m \quad (96)$$

This is an assumed Einstein–scalar action. Its independent-field variation requires compactly supported variations or appropriate boundary data and gravitational boundary terms. Matter variation must be included when S_m depends on ϕ . Constraining $g=g[\phi]$ before variation gives a different chain-rule theory. The finite kernel supplies neither this action nor Newton’s constant, a Lorentz manifold or a physical scalar field.

26 Quantum bridge

Quantum state identification remains a bridge

CONDITIONAL QUANTUM MODEL



For a calibrated $E_0=|0\rangle\langle 0|$, $p=\text{tr}(E_0 \rho)$; the state-identification map is additional physics.

Figure 24. Quantum state identification remains a bridge. Conditional quantum model. For a calibrated $E_0=|0\rangle\langle 0|$, $p=\text{tr}(E_0 \rho)$; the state-identification map is additional physics. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

A complete quantum bridge would be:

$$\Pi_Q : \Xi \rightarrow (H_Q, \rho_Q, A_Q, H_{\text{hat}}) \quad (97)$$

where H_Q is a Hilbert space, ρ_Q a density operator, A_Q an observable algebra, and H_{hat} a Hamiltonian.

Quantum operations are naturally represented by completely positive maps and instruments [11,12]. Therefore nonunitarity of the finite BFG kernel does not by itself exclude a quantum interpretation; actual instrument compatibility must be demonstrated.

$$H_A : \rho_Q^A = \rho_W \quad (98)$$


$$H_B : \rho_Q^B = \rho_F / \text{tr}(\rho_F) \quad (99)$$

Neither is a theorem of the finite kernel.

27 Finite quantum discrimination example

Competing quantum bridge predictions

COMPUTED BRIDGE COMPARISON



At (0.75,0.25), hypotheses A and B differ by 0.5; no actual measured outcomes are used.

Figure 25. Competing quantum bridge predictions. Computed bridge comparison. At (0.75,0.25), hypotheses A and B differ by 0.5; no actual measured outcomes are used. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

$$K = -I_2; Y = \text{diag}(1/2, 1/4); P = I_2 \setminus (100)$$

$$\rho_F = \text{diag}(1, 3); \rho_W = \text{diag}(3/4, 1/4) \setminus (101)$$

Then $Q = T_{\text{sel}} = P_{\text{sel}} = I_2$ and $m_{\text{sel}} = 1$.

$$C = \text{diag}(2/3, 4/5); B = \text{diag}(1/3, 1/5); G = \text{diag}(3/2, 5/4) \setminus (102)$$

$$\Lambda_C = 46/15; \Lambda_B = 19/60 \setminus (103)$$

Both loads are positive, so the analysis operator has full column rank. V is unitary, $X_{R4} = V^\dagger$, $S_4^{\wedge-} = S_4^{\wedge+} = I_2$, $m_4 = 1$, and $E = 0$.

For an externally fixed effect $E_0 = |0\rangle\langle 0|$:

$$p_A = \text{tr}(E_0 \rho_W) = 3/4 \setminus (104)$$


$$p_B = \text{tr}(E_0 \rho_F) / \text{tr}(\rho_F) = 1/4 \setminus (105)$$

The two bridge hypotheses are distinguishable in principle. E_0 is not the actual R_4 success effect of this example: here $S_4^{\wedge-} = I$ and the R_4 success mass is one.

28 R4 and quantum instruments

A general effect need not be projective

COMPUTED EFFECT TEST



A direct R4 effect is a projector. Generic positive effects with interior eigenvalues fail that test.

Figure 26. A general effect need not be projective. Computed effect test. A direct R4 effect is a projector. Generic positive effects with interior eigenvalues fail that test. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

For a fixed independently specified T_4 , $\rho \rightarrow T_4 \rho T_4^\dagger$ is completely positive, with effect $T_4^\dagger T_4 = S_4^\wedge$. Direct identification with a measured Kraus success operator therefore requires a projective effect and compatible transport. A general POVM effect need not be a projector [11].

Because T_4 is endogenously state-dependent,


$$T_4 = T_4 \backslash (\Xi) \quad \backslash (106)$$

the family $\rho \rightarrow T_4(\Xi) \rho T_4(\Xi)^\dagger$ is not automatically one linear CP channel on arbitrary inputs. It can be a classically conditioned family only if Ξ is independently identified.

29 Gauge and particle sector

Particle hierarchy requires parameters

SCHEMATIC DEPENDENCY



The surface labels a parameter dependency only; no Standard Model mass or coupling is derived.

Figure 27. Particle hierarchy requires parameters. Schematic dependency. The surface labels a parameter dependency only; no Standard Model mass or coupling is derived. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

The Standard Model is experimentally organized around $SU(3)_C \times SU(2)_L \times U(1)_Y$ and measured particle properties summarized by the Particle Data Group [14]. A BFG particle bridge must map the finite recursive state to field representations, charges, couplings, symmetry breaking, masses, and mixing data.


$$\Pi_{SM}:\Xi \rightarrow \{\text{gauge fields, representations, couplings, masses, mixings}\} \quad (107)$$

The current finite core does not determine these numerical quantities.

30 Atomic closure

Atomic localization and attraction

CONVENTIONAL CONDITIONAL MODEL



$E(r)=1/(2r^2)-a/r$; ordinary quantum scaling, not a new BFG atomic spectrum.

Figure 28. Atomic localization and attraction. Conventional conditional model. $E(r)=1/(2r^2)-a/r$; ordinary quantum scaling, not a new BFG atomic spectrum. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

A structural atom-level closure can be denoted:

$$\text{Atom} = \text{Cl}_N \backslash (\text{particles}, \text{interactions}) \quad (108)$$

but Cl_N is not a complete transition law until its state space and dynamics are given. Quantitative atomic spectra currently require ordinary quantum mechanics and electrodynamics. A familiar scale argument,


$$E(r) \sim \hbar^2 / (2\mu r^2) - e^2 / (4\pi \epsilon_0 r) \quad (109)$$

shows why the positive r^{-2} localization term can dominate the attractive r^{-1} term at short distance. This is a quantum variational scaling argument, not a derivation of $\hbar$ or e from the finite kernel.

31 Molecular and biological closure

Molecular binding criterion

SCHEMATIC BINDING LANDSCAPE




The illustrative energy x^2+y^2-1 marks a negative binding region; it is not a BFG molecular potential.

Figure 29. Molecular binding criterion. Schematic binding landscape. The illustrative energy x^2+y^2-1 marks a negative binding region; it is not a BFG molecular potential. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Biological boundary and gross flux

COMPUTED FLUX IDENTITY



$J_{\text{net}} = J_{\text{out}} - J_{\text{in}}$. Stationary net flux permits nonzero gross flows; no cell viability is inferred.

Figure 30. Biological boundary and gross flux. Computed flux identity. $J_{\text{net}} = J_{\text{out}} - J_{\text{in}}$. Stationary net flux permits nonzero gross flows; no cell viability is inferred. Horizontal and vertical coordinate ranges are the sampled grids; the

plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

A molecular stage may be schematically written:

$$M = Cl_{NM} \setminus (A_1, A_2) \setminus (110)$$

with a binding criterion $E(M) < E(A_1) + E(A_2)$, but a genuine bridge requires a typed molecular state and update. Turing's reaction-diffusion work is a canonical example of local dynamics generating higher spatial organization [15], while autopoiesis supplies an influential organizational language for self-maintaining living systems [16].

$$\text{molecule} \rightarrow \text{macromolecule} \rightarrow \text{cell} \rightarrow \text{tissue} \rightarrow \text{organ} \rightarrow \text{organism} \setminus (111)$$

The ladder is not a proof merely because the labels are ordered. Each arrow requires a state space, transition rule, preservation/loss quantities, and causal/viability criterion.

For an oriented biological boundary:


$$J_{\text{net}} = J_{\text{out}} - J_{\text{in}} \setminus (112)$$

Stationary net flux $J_{\text{net}} = 0$ means gross inflow equals gross outflow, not that both vanish.

32 Witness, world, self, and subject structure

World and self require operational maps

SCHEMATIC FUNCTIONAL MODEL



A product is only a visual aid. Π_W , Π_S , lineage and causal closure must be independently defined.

Figure 31. World and self require operational maps. Schematic functional model. A product is only a visual aid. Π_W , Π_S , lineage and causal closure must be independently defined. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Four notions are distinct: original witness mass w , carrier retention, formation retention and directed lineage. ρ_w is a kernel component; local m_{sel} and m_4 are computed event outputs and w and the transport history are separate trajectory records. They cannot generally be reconstructed from a normalized successor alone.

$$m_{\text{tot}} = m_{\text{sel}} m_4 \setminus (113)$$

Conditional normalization alone never proves full original-mass survival. Here $m_4=1$ is a theorem of the ambient analysis, whereas m_{sel} can be less than one at the initial event.

World and self require additional readouts:

$$W = \Pi_W(\Xi_{\text{history}}); S = \Pi_S(\Xi_{\text{history}}) \quad (114)$$

and a closed update:


$$(W_{k+1}, S_{k+1}) = U_{WS}(W_k, S_k, u_k) \quad (115)$$

A subject criterion Γ_{subj} can only become operational after these maps, their causal roles, and measurement rules are fixed. The Wende–Pereira precursor motivates the distinction between functional/computational description and the stronger question of subjectivity [3]. Predictive/free-energy formulations provide a comparison point for world-model and self-regulation ideas [17], but do not validate the BFG bridge.

33 Phenomenal bridge

Phenomenality is not a matrix consequence

SCHEMATIC STATUS PANEL



The zero sheet denotes absence of a derived prediction, not absence of subjective experience.

Figure 32. Phenomenality is not a matrix consequence. Schematic status panel. The zero sheet denotes absence of a derived prediction, not absence of subjective experience. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Even a fully operational Γ_{subj} would not make phenomenality a mathematical theorem. The final identification would be an empirical-ontological hypothesis:

$$H_{\text{phen}} : \Gamma_{\text{subj}} = 1 \Leftrightarrow \text{phenomenal experience} \quad (116)$$

This hypothesis is separate from the finite-kernel mathematics and from the Level-0 ontological interpretation.

34 Canonical world-formula statement

The finite BFG world formula is the orbit of the quotient dynamics:

$$W_{\text{BFG}} = \{ \text{Orb}_{\text{Ubar}} \setminus ([\Xi_0]) \mid \text{each step is admissible or terminates by an explicit terminal code} \} \quad (117)$$

The nonterminal successor is:

$$[\Xi_k] \rightarrow \text{Selection} \rightarrow \text{metric loads} \rightarrow \text{ambient reduced SVD} \rightarrow [K_+, Y_+] \quad (118a)$$

$$[K_+, Y_+] \rightarrow \text{isometric R4} \rightarrow \text{witness/formation inheritance} \rightarrow \text{seed or terminal} \rightarrow [\Xi_{k+1}] \quad (118b)$$

This formulation is finite, stratified, gauge-covariant, and dimension-aware. It is a world formula in the mathematical sense of a state-plus-law architecture. The stronger statement that the physical universe is isomorphic to W_{BFG} is a physical hypothesis supplied by bridge laws and measurements.

35 Falsifiability and bridge discipline

A physical BFG bridge must be fixed before the holdout observations it is meant to predict. For a bridge B and preparation data D_{prep} :

$$\Xi = M_{\text{prep}} \setminus (D_{\text{prep}} \setminus) \quad (119)$$

$$P_{\text{BFG}} \setminus (D_{\text{future}} \mid D_{\text{prep}}, B) \quad (120)$$

must be computed without refitting B to holdout outcomes. The quantum A/B example illustrates this requirement. A negative result first falsifies the tested bridge hypothesis, not every mathematical statement of the finite kernel.

The same discipline applies to gravity, particle physics, biology, and subject structure. A bridge freely replaceable after every observation has no predictive content.

Identifiability is a separate requirement. Full column rank of a sensitivity Jacobian can provide a sufficient regular local criterion under suitable assumptions, but is not necessary for structural local uniqueness. The map $Y(\theta) = \theta^3$ is injective even though its derivative is zero at $\theta=0$. State the relevant equivalence symmetries and test local uniqueness of the actual preparation or bridge map; do not infer it solely from a first-order rank test.

36 Variable and operator register

Symbol	Type	Role	Canonical status
Φ_0	Level-0 object	pre-operational domain	bridge to Ξ not fixed
Ξ	five-component operator tuple	finite complete state	defined
K	Hermitian operator	kernel/formation geometry	defined
ρ_F	positive operator	formation	defined, not normalized
ρ_W	positive trace-one operator	witness	defined
Y	positive semidefinite operator	Gram/load geometry	defined
P	orthogonal projector	persistent carrier	defined
Q	spectral projector	selection	defined
T_{sel}	partial isometry	first transport	defined
P_G	G-orthogonal projector	metric selection	defined
Λ_C, Λ_B	nonnegative scalars	endogenous loads	defined
α, β	weights	analysis balance	dual-load branch
A	linear operator	analysis map	defined
U, Σ, V	reduced SVD factors	successor frame/spectrum	defined modulo gauge
T_4	partial isometry	R4 transport	defined

Symbol	Type	Role	Canonical status
$m_{\text{sel}}, m_4, m_{\text{tot}}$	$[0,1]$ scalars	witness survival	defined
E	projector	free target complement	defined
H_E	Hermitian compression	seed spectral problem	defined
P_{seed}	projector	seed support	simple negative branch
Ψ	nine-role functional state	readout architecture	$(\varphi_{\text{fun}}, I_{\text{fun}}, A_{\text{fun}}, G_{\text{fun}}, \Omega_{\text{fun}}, T_{\text{fun}}, S_{\text{fun}}, M_{\text{fun}}, C_{\text{fun}})$
T_{clock}	$X_{\Xi}/\text{Unitary} \rightarrow R_{\{>0\}}$	physical step duration	physical bridge
$g_{\mu\nu}$	Lorentz metric	spacetime geometry	physical completion
ρ_Q	density operator	quantum state	competing bridge hypotheses
Γ_{subj}	future operational gate	subject structure	not yet fixed
w	trajectory scalar	original witness mass	separate record; $w_{\text{next}} = w * m_{\text{sel}}$
q_{info}	scalar stock	φ times I	not selection Q
a_{info}	scalar injection rate	stock source	not analysis A

Table 3. Canonical classification and definitions.

37 Scope of proved statements

The following are mathematical consequences of the declared canonical class:

1. $G=I+Y$ is positive definite whenever $Y \geq 0$.
2. The positive dual-load ambient kernel gives $0 < Y_+ < I$ and $n_+ \leq n$.
3. The reduced-SVD convention fixes the active successor dimension $r = \text{rank}(A)$.
4. The unitary quotient removes common SVD phase and exactly degenerate block-basis ambiguity from the successor state.
5. m_{sel} is a survival mass in $[0,1]$; ambient R_4 has $m_4=1$ and $m_{\text{tot}} = m_{\text{sel}}$.
6. Hermiticity of K makes the seed compression Hermitian.
7. On the stated scalar full-persistence branch, $0 < f(y) < y$ for $0 < y < 1$.
8. The same branch prevents a universal strict full-state contraction theorem in fixed positive product norms because neutral K/F directions exist.
9. The positive fixed point of the scalar two-variable ODE is locally asymptotically stable for positive a, μ, ε under the stated branch conditions.

The following are not consequences of the finite kernel alone: a unique physical information measure, SI time, Lorentzian spacetime, Einstein dynamics, the Born rule, Standard-Model parameters, quantitative atomic/molecular chemistry, organismal closure, or phenomenal experience.

38 Discussion

The canonicalization changes the status of the finite kernel in an important way. Its main ambiguities are no longer left as informal choices. The successor space is the active reduced-SVD space; SVD basis freedom is quotient gauge; witness masses have a normalized probabilistic kernel interpretation; K is Hermitian; the seed has compatible units; and failure modes are named terminal outcomes. This makes the finite dynamics a genuine mathematical model rather than a collection of mutually suggestive formulas.

At the same time, the architecture remains layered. Internal Gram geometry is not silently promoted to spacetime. A kernel density operator is not silently promoted to a quantum state. A closure label is not silently promoted to a biological law. A world/self functional architecture is not silently promoted to phenomenality. Each promotion is an explicit bridge law that can be examined independently.

39 Conclusion

The Balance–Field Equation can be stated as a canonical stratified recursive dynamics:

$$[\Xi_{k+1}] = \text{Ubar}_{\text{BFG}}([\Xi_k]) \quad (121)$$

The stated model has an ambient analysis, canonical polar factors, positive dual loads and a seed rule that terminates when a nonempty complement lacks a simple negative minimum. It is defined on unitary equivalence classes. General consequences include $0 < Y_+ < I$, $n_+ \leq n$ and $m_4 = 1$; subsequent nonterminal events preserve witness mass. Scalar damping is a narrower geometric statement. Full regional contraction remains open and cannot hold on a region containing the neutral K/F control directions in a positive product norm. Physical universality requires additional fixed bridges and independent tests.

40 The BFG as a Fundamental Recursive World Equation

The canonical BFG world equation is not a shorthand for several unrelated laws. It is the single recursive evolution law on the quotient state space; spectral selection, metric projection, endogenous loading, reduced SVD, R4 transport, witness conditioning, formation inheritance, and seed formation are the explicit factorization of its evolution operator.

$$[\Xi_{k+1}] = \text{Ubar}_{\text{BFG}}([\Xi_k])$$

$$\text{Ubar}_{\text{BFG}} = U_{\text{seed}} \circ U_{\text{R4}} \circ U_{\text{SVD}} \circ U_{\text{load}} \circ U_{\text{metric}} \circ U_{\text{selection}}$$

Its composite operator structure is therefore a strength rather than an incompleteness: the state carries the coupled structures required for recursive persistence, while the factorization exposes every internal operation that produces the successor. The remaining question is no longer whether the BFG has a world equation; it is whether the physical measurement maps below identify this mathematical orbit with the observed world.

41 Physical Observables and Measurement Map

A physical world equation must connect internal state variables to independently measured quantities. The canonical measurement layer is therefore a family M_{phys} of typed readout maps. SI units follow the International System of Units; particle observables are to be compared with independently tabulated experimental quantities rather than fitted labels.

$$O_{\text{phys}} = M_{\text{phys}}([\Xi], \text{calibration, apparatus})$$

A BFG physical bridge is accepted only after the readout, calibration constants, preparation protocol, and uncertainty model are fixed independently of the holdout observations.

Symbol or observable	Complete specification
Physical time Δt	SI / standard unit: s Experimental measurement: atomic/optical clock interval BFG readout target: $T_{\text{clock}}(\Xi)$ Required bridge: $M_t: \Xi \rightarrow \mathbb{R}_+$ Calibration / nuisance: clock scale and synchronization Falsification criterion: predicted intervals inconsistent with calibrated clock data
Length / displacement	SI / standard unit: m Experimental measurement: interferometry, ranging BFG readout target: metric distance from $g_{\mu\nu}$ Required bridge: M_g plus coordinate/rod prescription Calibration / nuisance: coordinate gauge, apparatus response Falsification criterion: predicted invariant distances disagree with measurements

Symbol or observable	Complete specification
Proper time	SI / standard unit: s Experimental measurement: clock comparison along worldlines BFG readout target: $c^{-1} \int \sqrt{-g_{\mu\nu} dx^\mu dx^\nu}$ Required bridge: Lorentzian spacetime bridge Calibration / nuisance: worldline and clock model Falsification criterion: clock ratios/redshifts fail
Energy	SI / standard unit: J or eV Experimental measurement: calorimetry/spectroscopy BFG readout target: E_{exp} , E_{bind} ; H_{hat} spectrum Required bridge: M_E and quantum bridge Calibration / nuisance: zero point, detector calibration Falsification criterion: spectral/energy differences fail
Frequency	SI / standard unit: Hz Experimental measurement: spectroscopy/clock counting BFG readout target: transition $\Delta E/h$ after bridge Required bridge: M_Q plus h convention Calibration / nuisance: line shape, Doppler/systematics Falsification criterion: predicted line centers fail
Mass	SI / standard unit: kg; eV/ c^2 Experimental measurement: inertial/gravitational or particle reconstruction BFG readout target: particle/formation mass readout Required bridge: Π_{SM} or gravitational bridge Calibration / nuisance: detector/model systematics Falsification criterion: mass predictions fail PDG/precision data
Electric charge	SI / standard unit: C or multiples of e Experimental measurement: charge/current standards, scattering BFG readout target: charge representation from Π_{SM} Required bridge: gauge/representation bridge Calibration / nuisance: detector gain, e convention Falsification criterion: charge assignments or ratios fail
Momentum	SI / standard unit: kg m s ⁻¹ ; eV/ c Experimental measurement: tracking, recoil, scattering BFG readout target: generator/kinematic readout Required bridge: spacetime + quantum bridge Calibration / nuisance: field map, alignment Falsification criterion: kinematic distributions fail
Angular momentum / spin	SI / standard unit: J s; $\hbar$ units Experimental measurement: Stern–Gerlach, spectroscopy, collider angular analyses BFG readout target: representation label/operator Required bridge: quantum + particle bridge Calibration / nuisance: analyzer orientation Falsification criterion: spin/multiplicity predictions fail
Cross section	SI / standard unit: m ² ; barn Experimental measurement: event rates with luminosity BFG readout target: scattering probability/amplitude Required bridge: quantum field bridge Calibration / nuisance: luminosity, efficiency Falsification criterion: predicted cross sections fail
Decay rate / lifetime	SI / standard unit: s ⁻¹ / s Experimental measurement: time distributions BFG readout target: transition/decay generator Required bridge: particle quantum bridge Calibration / nuisance: acceptance/background Falsification criterion: lifetime/branching predictions fail

Symbol or observable	Complete specification
Dimensionless couplings	SI / standard unit: 1 Experimental measurement: scattering/precision fits BFG readout target: $g_i(\Xi)$ Required bridge: Π_{SM} Calibration / nuisance: renormalization scale Falsification criterion: running/couplings fail independent fits
Gravitational acceleration	SI / standard unit: $m s^{-2}$ Experimental measurement: free fall, gravimetry BFG readout target: geodesic/field prediction from g Required bridge: M_g + dynamics Calibration / nuisance: local environment Falsification criterion: acceleration/redshift/lensing fail
Curvature / tidal field	SI / standard unit: m^{-2} or s^{-2} forms Experimental measurement: geodesic deviation, gravitational observations BFG readout target: Riemann/Ricci tensors from g Required bridge: connection/curvature bridge Calibration / nuisance: source model Falsification criterion: tidal/lensing/wave observables fail
Quantum outcome probability	SI / standard unit: 1 Experimental measurement: repeated POVM/instrument frequencies BFG readout target: $\text{tr}(E_a \rho_Q)$ Required bridge: Π_Q plus instrument map Calibration / nuisance: state preparation, detector tomography Falsification criterion: holdout frequencies reject predicted probabilities
Witness survival mass	SI / standard unit: 1 Experimental measurement: kernel-internal; no established direct apparatus yet BFG readout target: m_{sel}, m_4, m_{tot} Required bridge: requires physical witness identification Calibration / nuisance: preparation/selection protocol Falsification criterion: bridge rejected if measured success statistics disagree
Information I	SI / standard unit: bit/nat or 1 Experimental measurement: specified ensemble/configuration counting BFG readout target: $\log C_\theta(X) $ Required bridge: M_I fixes $X, C_\theta, \theta, \text{base}, I_0$ Calibration / nuisance: coarse graining Falsification criterion: independent counts disagree
Order parameter ϕ	SI / standard unit: 1 Experimental measurement: derived from independently measured energy/information ratios BFG readout target: ϕ_E or ϕ_N Required bridge: M_ϕ Calibration / nuisance: reference I_0 and energy definitions Falsification criterion: derived observable disagrees
Biological flux	SI / standard unit: $\text{mol s}^{-1}, \text{kg s}^{-1}, W$ as appropriate Experimental measurement: transport/metabolic assays BFG readout target: J_{in}, J_{out}, J_{net} Required bridge: Π_{bio} Calibration / nuisance: boundary and species definitions Falsification criterion: flux/viability predictions fail
World/Self functional readouts	SI / standard unit: task-dependent Experimental measurement: behavioral/interventional assays BFG readout target: Π_W, Π_S, U_{WS} Required bridge: cognitive bridge Calibration / nuisance: task model and intervention class Falsification criterion: out-of-sample causal predictions fail

Symbol or observable	Complete specification
Phenomenal report / experience	SI / standard unit: no single SI unit Experimental measurement: report, behavior, physiology under protocol BFG readout target: Γ_{subj} only after definition Required bridge: H_{phen} Calibration / nuisance: report model and confounds Falsification criterion: pre-registered bridge predictions fail

Table 4. Canonical physical-observable interface for the Balance–Field Equation. Entries marked as bridges are model commitments to be fixed before holdout testing.

41.1 Metrological requirement

The measurement map is not permitted to identify an internal variable with an SI observable merely because their dimensions can be made compatible. Each observable requires an apparatus model and a calibration chain. The SI provides the internationally agreed unit system, while precision particle properties are independently compiled by the Particle Data Group. In the quantum sector, electrical and quantum standards illustrate how physical laws become metrological realizations only after a concrete instrument and traceability chain are specified.

41.2 Minimum empirical closure condition

For a preparation D_{prep} , calibration C , bridge B , and holdout data D_{test} , a physically closed BFG sector must provide a predictive distribution $P_{\text{BFG}}(D_{\text{test}} | D_{\text{prep}}, C, B)$ before D_{test} is observed. A sector is not physically closed by notation alone. This criterion applies independently to time, geometry, quantum outcomes, particle parameters, biological observables, and subject-level observables.

42 Complete Canonical Mathematical Specification

This section consolidates the complete finite Balance–Field Equation in one place. It is intended to remove the need to reconstruct types, dimensions, assumptions, and successor rules from separate passages.

$$[\Xi_{k+1}] = \text{Ubar}_{\text{BFG}} \setminus ([\Xi_k])$$

The quotient operator factors into selection, first transport, endogenous metric projection, positive load weighting, ambient reduced SVD, successor geometry and kernel, isometric R4 realignment, formation inheritance, and the seed or terminal rule. The nominal second normalization has mass exactly one.

42.1 Domain of one nonterminal step

All input operators act on H_k . T_{sel} maps the selected source support to P_{sel} . The analysis operator is $A: H_k \rightarrow H_k \oplus H_k$, without domain restriction. Its reduced SVD has $U: C^r \rightarrow H_k \oplus H_k$ and $V: C^r \rightarrow H_k$. The successor is C^r . Positive dual loads imply $\text{rank}(P_{\text{sel}}) \leq r \leq n_k$. Padding null directions or substituting AZ changes the model.

42.2 Gauge-invariant successor

The successor is an equivalence class under simultaneous unitary coordinate changes. A common phase of a simple singular-vector pair and a unitary rotation within an exactly degenerate singular block are not distinct physical or mathematical successors. Under $U \rightarrow UW$ and $V \rightarrow VW$ with W commuting with Σ , K_+ , Y_+ , S_4^{+} , ρ_W , $+$, ρ_F , $+$ and P_+ all transform by the same target-space conjugation. Thus the quotient state $[\Xi_+]$ is unique even when a matrix representation is not.

42.3 Regular strata and differentiability

Regularity is assumed on fixed projector and matrix-rank strata: a gap to the threshold 1, positive m_{sel} and both loads, fixed positive rank of A, simple separated positive singular values for componentwise formulas, and a fixed seed branch. On $E \neq 0$ require a simple negative lowest eigenvalue separated from zero and the other eigenvalues; on $E=0$ keep complement dimension zero. Input tangent variations are Hermitian in K,F,W,Y, have $\text{tr}(\delta W)=0$, and satisfy $P \delta P P = (I-P) \delta P (I-P) = 0$. Positivity/rank constraints also restrict allowed tangents. These are specified regularity assumptions, not universally necessary conditions for every possible proof. All polar norms here are operator 2-norms.

42.4 Terminal and continuation semantics

Apply terminal gates in order: zero selection rank, zero first witness mass, failure of positive dual loads, then seed failure on $E \neq 0$. Lack of a negative minimum gives T_{seednone} ; a multiple negative minimum gives T_{seeddeg} . After positive loads, zero A/R4 rank and zero m_4 are impossible and are only implementation consistency checks. $E=0$ is the sole no-seed continuation.

42.5 Stability hierarchy

Endogenous geometry is positive and the actual ambient update additionally gives $0 < Y_+ < I$. Neither result bounds the derivative of the complete tuple by a constant below one. Regional full-kernel contraction remains open; rank, gap, load and mass margins only describe regions on which it can be investigated. The neutral K/F control branch excludes a universal strict product-norm theorem. Transverse stability requires a specified invariant set and a separate certificate.

43 Complete Variable and Operator Dictionary

Every symbol used in the canonical equation is listed below. 'Bridge' means that the finite mathematical kernel does not by itself determine the physical identification.

Symbol or observable	Complete specification
Φ_0	Domain / type: X_0 Codomain / unit: ontological Exact definition: Level-0 object Mathematical role: pre-operational domain Conditions: no canonical Ξ map Status: Definition/bridge
Ξ_k	Domain / type: product operator space Codomain / unit: mixed Exact definition: $(K, \rho_F, \rho_W, Y, P)$ Mathematical role: complete finite state Conditions: finite H_k Status: Definition
H_k	Domain / type: finite complex Hilbert space Codomain / unit: $\dim n_k$ Exact definition: active event space Mathematical role: carrier of Ξ_k Conditions: finite Status: Definition
K_k	Domain / type: $\text{Herm}(H_k)$ Codomain / unit: E_F Exact definition: $K=K^\dagger$ Mathematical role: kernel/formation geometry Conditions: Hermitian Status: Definition

Symbol or observable	Complete specification
$\rho_{F,k}$	Domain / type: Pos(H_k) Codomain / unit: E_F Exact definition: $\rho_F \geq 0$ Mathematical role: formation Conditions: not necessarily trace-one Status: Definition
$\rho_{W,k}$	Domain / type: Dens(H_k) Codomain / unit: 1 Exact definition: $\rho_W \geq 0, \text{tr}=1$ Mathematical role: witness Conditions: normalized input Status: Definition
Y_k	Domain / type: Herm_+(H_k) Codomain / unit: 1 Exact definition: $Y \geq 0$ Mathematical role: Gram/load geometry Conditions: positive semidefinite Status: Definition
P_k	Domain / type: Proj(H_k) Codomain / unit: 1 Exact definition: $P = P^\dagger = P^2$ Mathematical role: persistent carrier Conditions: orthogonal projector Status: Definition
Q_k	Domain / type: Proj(H_k) Codomain / unit: 1 Exact definition: $1_{[0,1)}(Y)$ Mathematical role: spectral selection Conditions: gap to 1 for smoothness Status: Definition
C_k	Domain / type: End(H_k) Codomain / unit: 1 Exact definition: $(I+Y)^{-1}$ Mathematical role: neutral/complement channel Conditions: $Y \geq 0$ Status: Definition
B_k	Domain / type: End(H_k) Codomain / unit: 1 Exact definition: $YC = I - C$ Mathematical role: balance channel Conditions: $Y \geq 0$ Status: Definition
G_k	Domain / type: PosDef(H_k) Codomain / unit: 1 Exact definition: $I+Y$ Mathematical role: internal metric Conditions: $Y \geq 0$ Status: Theorem/definition
I and I_0	Domain / type: specified configuration count and reference Codomain / unit: bit/nat or dimensionless Exact definition: $I = \log_b C_\theta(X) $ Mathematical role: information readout Conditions: finite nonempty class, $b > 1, I_0 > 0$ Status: Open bridge

Symbol or observable	Complete specification
φ_N or φ_E	Domain / type: scalar ratios Codomain / unit: dimensionless Exact definition: $(N_+/N_-)(I/I_0)$ or $(E_{\text{exp}}/E_{\text{bind}})(I/I_0)$ Mathematical role: order readout Conditions: positive references; distinct from Φ_0 Status: Open bridge

Table 5. Complete Level-0 and finite-state variable dictionary.

Symbol or observable	Complete specification
X_{sel}	Domain / type: $\text{End}(H_k)$ Codomain / unit: 1 Exact definition: $Q_k P_k$ Mathematical role: selection overlap Conditions: compatible source space Status: Definition
T_{sel}	Domain / type: partial isometry Codomain / unit: 1 Exact definition: $\text{polar}(X_{\text{sel}})$ Mathematical role: first transport Conditions: $\text{rank} > 0$ Status: Definition
$S_{\text{sel}}^\wedge$	Domain / type: projector Codomain / unit: 1 Exact definition: $T_{\text{sel}}^\dagger T_{\text{sel}}$ Mathematical role: source support Conditions: T_{sel} defined Status: Definition
P_{sel}	Domain / type: projector Codomain / unit: 1 Exact definition: $T_{\text{sel}} T_{\text{sel}}^\dagger$ Mathematical role: target support Conditions: T_{sel} defined Status: Definition
ρ_F^{sel}	Domain / type: positive operator Codomain / unit: E_F Exact definition: $T_{\text{sel}} \rho_F T_{\text{sel}}^\dagger$ Mathematical role: selected formation Conditions: transport defined Status: Definition
$\rho_{\text{un}}^{\text{sel}}$	Domain / type: positive operator Codomain / unit: 1 Exact definition: $T_{\text{sel}} \rho_W T_{\text{sel}}^\dagger$ Mathematical role: unnormalized witness Conditions: transport defined Status: Definition
m_{sel}	Domain / type: $[0,1]$ Codomain / unit: 1 Exact definition: $\text{tr}(S_{\text{sel}}^\wedge \rho_W)$ Mathematical role: first survival mass Conditions: $\text{tr} \rho_W = 1$ Status: Theorem/definition
ρ_W^{sel}	Domain / type: density operator Codomain / unit: 1 Exact definition: $\rho_{\text{un}}^{\text{sel}}/m_{\text{sel}}$ Mathematical role: conditioned witness Conditions: $m_{\text{sel}} > 0$ Status: Definition

Symbol or observable	Complete specification
Z	Domain / type: isometry Codomain / unit: 1 Exact definition: $Z^\dagger Z = I, ZZ^\dagger = P_{\text{sel}}$ Mathematical role: target-support frame Conditions: local gauge Status: Definition
P_G	Domain / type: G-projector Codomain / unit: 1 Exact definition: $Z(Z^\dagger GZ)^{-1}Z^\dagger G$ Mathematical role: metric projection Conditions: invertible compression Status: Definition
Λ_C	Domain / type: R_+ Codomain / unit: load unit Exact definition: trace formula Mathematical role: C-load Conditions: positive branch Status: Definition
Λ_B	Domain / type: R_+ Codomain / unit: load unit Exact definition: trace formula Mathematical role: B-load Conditions: positive branch Status: Definition
α	Domain / type: (0,1) Codomain / unit: 1 Exact definition: $\Lambda_B / (\Lambda_C + \Lambda_B)$ Mathematical role: C weight Conditions: dual-load branch Status: Definition
β	Domain / type: (0,1) Codomain / unit: 1 Exact definition: $\Lambda_C / (\Lambda_C + \Lambda_B)$ Mathematical role: B weight Conditions: dual-load branch Status: Definition

Table 6. Selection, transport, metric-projection, and load variables.

Symbol or observable	Complete specification
A_k	Domain / type: $H_k \rightarrow H_k \otimes H_k$ Codomain / unit: analysis space Exact definition: vertical stack of weighted P_{GC}/P_{GB} Mathematical role: analysis operator Conditions: positive loads Status: Definition
r_k	Domain / type: N Codomain / unit: 1 Exact definition: $\text{rank}(A_k)$ Mathematical role: successor dimension Conditions: $r_k \geq 1$ Status: Definition
U_k	Domain / type: isometry $C^r \rightarrow$ analysis Codomain / unit: 1 Exact definition: left reduced SVD frame Mathematical role: successor kernel frame Conditions: modulo gauge Status: Definition

Symbol or observable	Complete specification
Σ_k	Domain / type: positive diagonal Codomain / unit: 1 Exact definition: singular spectrum Mathematical role: sets Y_+ Conditions: positive singular values Status: Definition
V_k	Domain / type: isometry C^r -source Codomain / unit: 1 Exact definition: right reduced SVD frame Mathematical role: R_4 source frame Conditions: modulo gauge Status: Definition
Ω_U	Domain / type: skew-Hermitian Codomain / unit: variation Exact definition: $U^\dagger \delta U$ Mathematical role: SVD connection Conditions: simple stratum Status: Derived
Ω_V	Domain / type: skew-Hermitian Codomain / unit: variation Exact definition: $V^\dagger \delta V$ Mathematical role: SVD connection Conditions: simple stratum Status: Derived
K_+	Domain / type: $\text{Herm}(C^r)$ Codomain / unit: E_F Exact definition: $U^\dagger(K \oplus K)U$ Mathematical role: successor kernel Conditions: K Hermitian Status: Definition
Y_+	Domain / type: $\text{Herm}_+(C^r)$ Codomain / unit: 1 Exact definition: Σ^2 Mathematical role: successor Gram geometry Conditions: reduced SVD Status: Theorem/definition
X_{R_4}	Domain / type: linear map Codomain / unit: 1 Exact definition: $V^\dagger P_{\text{sel}}$ Mathematical role: R_4 overlap Conditions: compatible frames Status: Definition
T_4	Domain / type: partial isometry Codomain / unit: 1 Exact definition: $\text{polar}(X_{R_4})$ Mathematical role: R_4 transport Conditions: $\text{rank} > 0$ Status: Definition
S_4^-	Domain / type: projector Codomain / unit: 1 Exact definition: $T_4^\dagger T_4$ Mathematical role: R_4 source support Conditions: T_4 defined Status: Definition

Symbol or observable	Complete specification
S_4^{+}	Domain / type: projector Codomain / unit: 1 Exact definition: $T_4 T_4^\dagger$ Mathematical role: R4 target support Conditions: T_4 defined Status: Definition
m_4	Domain / type: {1} Codomain / unit: 1 Exact definition: $\text{tr}(S_4^{+} \rho_W^{\text{sel}})$ Mathematical role: second survival mass Conditions: identically one on positive dual-load branch Status: Definition/theorem
m_{tot}	Domain / type: [0,1] Codomain / unit: 1 Exact definition: $m_{\text{sel}} m_4$ Mathematical role: step survival mass Conditions: sequential conditioning Status: Definition

Table 7. Reduced-SVD, successor, and R4 variables.

Symbol or observable	Complete specification
E	Domain / type: projector Codomain / unit: 1 Exact definition: $I - S_4^{+}$ Mathematical role: free target complement Conditions: successor space fixed Status: Definition
Z_E	Domain / type: isometry Codomain / unit: 1 Exact definition: $Z_E Z_E^\dagger = E$ Mathematical role: complement frame Conditions: horizontal gauge locally Status: Definition
H_E	Domain / type: $\text{Herm}(\text{Ran } E)$ Codomain / unit: E_F Exact definition: $Z_E^\dagger K_+ Z_E$ Mathematical role: seed spectral operator Conditions: K_+ Hermitian Status: Definition
λ	Domain / type: $\mathbb{R}$ Codomain / unit: E_F Exact definition: simple negative minimum of H_E Mathematical role: seed amplitude Conditions: $\lambda < 0$ Status: Definition
π	Domain / type: rank-one projector Codomain / unit: 1 Exact definition: $vv^\dagger$ Mathematical role: reduced seed support Conditions: simple minimum Status: Definition
P_{seed}	Domain / type: projector Codomain / unit: 1 Exact definition: $Z_E \pi Z_E^\dagger$ Mathematical role: embedded seed support Conditions: $P_{\text{seed}} \leq E$ Status: Definition

Symbol or observable	Complete specification
κ_F	Domain / type: R_+ Codomain / unit: conversion Exact definition: fixed unit factor Mathematical role: K-to-formation conversion Conditions: constant Status: Assumption if needed
ρ_{seed}	Domain / type: positive operator Codomain / unit: E_F Exact definition: $-\kappa_F \lambda \Pi_{\text{seed}}$ Mathematical role: new formation Conditions: negative λ Status: Definition
$\rho_{F,+}$	Domain / type: positive operator Codomain / unit: E_F Exact definition: $\rho_{F,\text{inh}} + \rho_{\text{seed}}$ Mathematical role: successor formation Conditions: $E=0$ or simple negative seed; otherwise terminal Status: Definition
P_+	Domain / type: projector Codomain / unit: 1 Exact definition: $S_4^{++} \Pi_{\text{seed}}$ Mathematical role: successor carrier Conditions: orthogonal supports Status: Theorem/definition
R_λ	Domain / type: reduced resolvent Codomain / unit: E_F^{-1} Exact definition: $(H_E - \lambda I)^{++}$ Mathematical role: seed differential Conditions: simple eigenvalue Status: Derived

Table 8. Seed, complement, and formation variables.

Symbol or observable	Complete specification
Ψ	Domain / type: 9-role state Codomain / unit: mixed Exact definition: $(\varphi_{\text{fun}}, I_{\text{fun}}, A_{\text{fun}}, G_{\text{fun}}, \Omega_{\text{fun}}, T_{\text{fun}}, S_{\text{fun}}, M_{\text{fun}}, C_{\text{fun}})$ Mathematical role: functional readout Conditions: Π incomplete Status: Bridge
Π	Domain / type: map Codomain / unit: - Exact definition: $\Xi \rightarrow \Psi$ Mathematical role: factor projection Conditions: fiber condition Status: Bridge
T_{clock}	Domain / type: $X_\Xi / \text{Unitary} \rightarrow R_{\{>0\}}$ Codomain / unit: s Exact definition: physical step duration Mathematical role: event-to-time bridge Conditions: calibration Status: Bridge
t_k	Domain / type: R Codomain / unit: s Exact definition: $t_0 + \sum T_{\text{clock}}$ Mathematical role: physical time Conditions: clock fixed Status: Bridge

Symbol or observable	Complete specification
$g_{\mu\nu}$	Domain / type: Lorentz metric Codomain / unit: 1 Exact definition: bridge-dependent Mathematical role: spacetime Conditions: manifold/signature Status: Bridge
ρ_Q	Domain / type: density operator Codomain / unit: 1 Exact definition: ρ_W or ρ_F /tr ρ_F candidates Mathematical role: quantum state Conditions: experiment selects bridge Status: Bridge
H_{hat}	Domain / type: self-adjoint operator Codomain / unit: J/eV Exact definition: quantum Hamiltonian Mathematical role: energy generator Conditions: Π_Q Status: Bridge
E_a	Domain / type: POVM effect Codomain / unit: 1 Exact definition: $0 \leq E_a \leq I$ Mathematical role: quantum measurement Conditions: instrument fixed Status: External standard
Π_{SM}	Domain / type: map Codomain / unit: - Exact definition: $\Xi \rightarrow$ particle/gauge data Mathematical role: particle bridge Conditions: representation/couplings Status: Bridge
$J_{\text{in/out/net}}$	Domain / type: fluxes Codomain / unit: system-dependent Exact definition: $J_{\text{net}} = J_{\text{out}} - J_{\text{in}}$ Mathematical role: biological transport Conditions: boundary fixed Status: Bridge/readout
W	Domain / type: functional state Codomain / unit: task-dependent Exact definition: $\Pi_W(\text{history})$ Mathematical role: world model Conditions: readout defined Status: Bridge
S	Domain / type: functional state Codomain / unit: task-dependent Exact definition: $\Pi_S(\text{history})$ Mathematical role: self model Conditions: readout defined Status: Bridge
Γ_{subj}	Domain / type: gate/readout Codomain / unit: 1 Exact definition: future explicit function Mathematical role: subject criterion Conditions: measurement map Status: Bridge

Table 9. Functional, physical, biological, and subject-level bridge variables.

44 Reader's Guide to the Recursive Mechanism

One recursive event begins with the declared five operators. Q selects the subthreshold spectral sector and QP defines the first directed polar transport. Formation retains its unnormalized magnitude; witness is conditioned and its initial mass is separately recorded. G and P_G evaluate formation through complementary C and B channels, whose loads determine the weights. The full ambient A supplies the active successor rank, geometry and frames. R_4 realigns the complete selected carrier without a second loss. A nonempty target complement requires a unique negative lowest seed mode. The whole event therefore specifies every intermediate type, transport, normalizing factor and terminal test.

45 Detailed Selection Logic

Selection is not deletion by a scalar threshold alone. $Q_k = 1_{-[0,1)}(Y_k)$ is a spectral projector, so it selects invariant spectral subspaces of the Hermitian positive operator Y_k . P_k is an independent persistence projector. The product $Q_k P_k$ need not itself be a projector, which is why its polar factor is used. $T_{sel,k}$ retains the geometric relation between the selected Y -subspace and the persistent P -subspace. Its source projector $S_{sel}^\wedge$ and target projector P_{sel} therefore carry different information. This distinction is also what makes the first witness mass $m_{sel} = \text{tr}(S_{sel}^\wedge \cdot \rho_w)$ mathematically meaningful: the mass is evaluated on the source before the state is transported and renormalized.

46 Why the Metric Projector Is Endogenous

P_G is the G -orthogonal projector onto $\text{Ran}(P_{sel})$. It equals the ordinary Euclidean projector P_{sel} exactly when $[P_{sel}, G] = 0$; G need not equal I . It satisfies $P_G^2 = P_G$ and $P_G^\dagger G = G P_G$. The current geometry affects selection, projection, loads and ultimately its own successor through A . This self feedback is explicit; its stabilizing effect is proved on the scalar branch and bounded generally by $0 < Y_+ < I$, without a claim of universal full-state contraction.

47 Load Weighting and Neutral Balance

$C = (I + Y)^{-1}$ and $B = Y(I + Y)^{-1}$ satisfy $C + B = I$ and commute with Y . In the eigenbasis of Y , an eigenvalue $y \geq 0$ gives $c = 1/(1+y)$ and $b = y/(1+y)$. Hence low-load directions are C -dominated, while high-load directions are B -dominated. The global weights α and β are not these eigenvalue-wise coefficients; they are formed from the metric formation loads Λ_C and Λ_B . The cross-definition $\alpha = \Lambda_B / (\Lambda_C + \Lambda_B)$, $\beta = \Lambda_C / (\Lambda_C + \Lambda_B)$ makes each analysis branch sensitive to the strength of the complementary load. This is the finite-kernel realization of a balance-mediated analysis rather than a one-channel amplification rule.

48 Reduced SVD as the Dimension-Formation Step

The reduced SVD retains precisely the positive singular directions of the ambient A . Thus $n_+ = \text{rank}(A) \leq n$. Although r can exceed the selected-carrier rank, the finite ambient dimension cannot grow. Rank-loss boundaries change the active state space. Fixed-rank derivatives require the declared gaps; coordinate gauge and physical emergence are different issues.

49 Gauge Freedom and Physical Equivalence

An SVD is never a unique ordered list of vectors in an absolute sense. A simple singular pair can be multiplied by a common complex phase; a degenerate singular block can be rotated by a unitary matrix. If K_+ , Y_+ , R_4 , formation, and witness were read as absolute matrices without accounting for this freedom, the update would be representation-dependent. The canonical equation therefore defines the successor on a unitary quotient. Every target-space object is transformed by the same conjugation. Observable scalar invariants such as traces and spectra are unchanged, while coordinate matrices change covariantly. This is a mathematical gauge

principle internal to the finite recursion; it should not be confused with the Standard Model gauge group, which is a separate physical bridge target.

50 Polar Differential Interpretation of the Three Terms

The constant-rank polar derivative $\delta T = T \Omega + (I - S_-) E_{X_H}^+ + T H^+ + T^\dagger E_X (I - S_-)$ has three geometrically distinct pieces. $T \Omega$ rotates coordinates inside the active source support. The second term moves the target support transversely. The third term moves the source support and transports that motion through T . Thus support motion is not an independent perturbation; it is determined by $E_X = \delta X$. The $3 \|E_X\| / \sigma_k$ bound is deliberately conservative. Its dependence on the smallest positive singular value expresses a standard fact of rank-sensitive matrix geometry: near rank loss, the active support can become highly sensitive even when a particular path remains benign.

51 R4 as Recursive Re-Alignment

Because $\text{Ran}(P_{\text{sel}})$ lies in $\text{Ran}(A^\dagger)$, $V V^\dagger P_{\text{sel}} = P_{\text{sel}}$. Thus $X_{R4}^\dagger X_{R4} = P_{\text{sel}}$ and $T_4 = V^\dagger P_{\text{sel}}$ already is a partial isometry. $R4$ realigns the complete selected carrier, gives $m_4=1$ and determines S_4^+ and its complement. It is not a second lossy test in this class. A physical Kraus interpretation would require an independent instrument bridge.

52 Formation, Witness, and the Meaning of Seed

Inherited formation keeps its trace, whereas witness is normalized and original mass is kept in the separate record w . In this kernel $m_4=1$, so $R4$ discards no mass. For $E=0$ no seed is needed. For $E \neq 0$ a simple negative minimum gives positive seed formation; no negative minimum or a multiple negative minimum is terminal. Orthogonality preserves the projector sum. Neither seed nor complement is automatically physical particle creation.

53 Terminality as Part of the Equation

Terminality is part of the definition. Selection failure, first witness failure, or nonpositive dual loads terminate before analysis. After positive loads A and $R4$ have nonzero rank and $m_4=1$, so second-mass failure is unreachable. If $E \neq 0$ the unique rank-one seed requires a simple strictly negative minimum; its absence or degeneracy terminates. $E=0$ alone continues without a seed.

54 What the Scalar Damping Result Does and Does Not Prove

The exact scalar map $f(y) = 2y^2 / [(1+y)^2(1+y^2)]$ provides a rare closed-form control branch of the finite recursion. Algebraically $f(y) < y$ for every $y > 0$, while the actual nonterminal selection-first realization under discussion is restricted to $0 < y < 1$. The result proves monotone damping of that scalar geometric load on the stated branch. It does not prove that the complete tuple contracts because K and formation can remain neutral. This is why the appropriate general stability target is likely transverse: perturbations away from a persistent manifold may decay while coordinates along the persistent identity remain neutral.

55 From Recursive Events to Measured Time

The event index k orders recursive updates and nothing more. To compare the equation with clocks, one needs $T_{\text{clock}}(\Xi)$, a positive duration assigned to each event. Summing these durations produces physical time along a trajectory. If a continuous effective state Ψ is desired, a separate flow F_Ψ must exist such that one discrete BFG step equals the flow evaluated at the assigned duration. The clock can vary even when the projected state is unchanged; only a clock required to be a function of Ψ alone must be constant on Π -fibers. This distinction prevents event count, model time, physical seconds, and renormalization scale from being conflated.

56 From Positive Gram Geometry to Lorentzian Spacetime

The internal $G=I+Y$ is positive definite because its role is to measure directions in the finite analysis space. Spacetime requires an indefinite Lorentzian metric. Therefore a physical geometry bridge must change mathematical type. A disformal ansatz can produce a Lorentzian metric only after a manifold, background η , scalar field, and nondegeneracy condition are supplied. An Einstein-scalar action can then generate field equations, but those equations follow from the added action. The finite BFG becomes a gravitational theory only when a map from recursive states to the spacetime fields and their calibrated constants is fixed and tested.

57 Quantum Measurement as a Discriminating Bridge

The two candidate identifications $\rho_Q=\rho_W$ and $\rho_Q=\rho_F/\text{tr}(\rho_F)$ are useful precisely because they are not equivalent. In the explicit two-dimensional control state they predict 3/4 and 1/4 for the same externally fixed effect. This turns a vague analogy between kernel matrices and density operators into a falsifiable bridge choice. To perform the test physically, the preparation must determine the relevant BFG state independently of the holdout outcomes, the effect must be calibrated, and the statistical rejection rule must be fixed in advance. The R4 success effect is not substituted for the external measurement unless instrument compatibility has separately been established.

58 Particle, Atomic, Molecular, and Biological Ordering

The hierarchy is constrained by dependence. A particle bridge must precede a quantitative atom bridge because atomic Hamiltonians require masses, charges, couplings, and quantum dynamics. A molecular bridge requires atomic degrees of freedom and interactions. A biological closure bridge then requires molecular transport, boundaries, energy/material fluxes, repair, and viable persistence. The notation Cl_N is therefore treated as a placeholder for a future typed closure operation, not as proof that a named level follows from the previous one. The ordering is a research dependency graph.

59 Witness to Subject Structure

The witness is already an internal mathematical object; world and self are not. A subject-level theory therefore cannot be obtained merely by renaming ρ_W . Π_W and Π_S must construct world- and self-related functional states from the physical/history state, and U_{WS} must close their dynamics. A candidate Γ_{subj} must then specify measurable criteria such as predictive sufficiency, self-indexing, directed lineage, and admissibility. Even if these conditions are operationally satisfied, the final equivalence with phenomenal experience remains an empirical-ontological bridge rather than a theorem of matrix recursion.

60 How the Physical Measurement Layer Closes a Sector

For each physical sector, the measurement interface has four layers: preparation, state identification, prediction, and comparison. Preparation produces D_{prep} . A fixed map M_{prep} identifies Ξ or the relevant quotient class. A bridge B maps the state to the physical observable model and generates a predictive distribution. An independently calibrated apparatus produces D_{test} . The bridge is rejected or retained according to a pre-registered uncertainty model and criterion. This architecture is what turns a mathematically closed recursive kernel into a potentially physical world equation.

61 Canonical Theorem and Assumption Register

Theorem A — Ambient Gram bound: positive dual loads imply $0<Y_+<I$ and $n_+\leq n$; $G=I+Y>0$ for the declared input class.

Theorem B — Witness transport: m_{sel} lies in $[0,1]$, $m_4=1$, $m_{\text{tot}}=m_{\text{sel}}$; later nonterminal events preserve original witness mass.

Theorem C — Gauge well-definedness: simultaneous unitary changes inside allowed SVD gauge blocks conjugate every target-space successor object and therefore leave $[\Xi_+]$ unchanged.

Theorem D — Scalar damping: $f(y)<y$ for $y>0$; the checked selection-first branch uses $0<y<1$.

Theorem E — Positive fixed-point stability of the scalar ODE: negative trace and positive determinant imply local asymptotic stability on the declared positive branch.

Assumption A — Level-0 to finite-state formation is not fixed by the finite kernel.

Assumption B — Physical time requires T_{clock} .

Assumption C — Lorentzian geometry requires a physical geometry bridge.

Assumption D — Quantum state identification requires Π_Q ; ρ_W and normalized ρ_F are competing hypotheses.

Assumption E — Standard-Model parameters require Π_{SM} and cannot be inferred from gauge labels alone.

Assumption F — World/Self and phenomenality require additional functional and empirical bridges.

62 Full Differential Chain Selection to the Analysis Operator

62.1 Spectral projector variation

On a fixed selection stratum the spectrum of Y remains separated from the threshold 1. The spectral projector $Q=1_{[0,1)}(Y)$ is then differentiable as an invariant-subspace projector. In a contour representation enclosing the selected spectrum but not the excluded spectrum, δQ is determined by the resolvent of Y and δY . The important structural point is that δQ is not an independent input: it is a linear function of δY on the fixed-gap stratum. At a threshold crossing this derivative description ceases to be the correct model and the system changes stratum.

62.2 Selection overlap and polar transport

The selection overlap is $X_{\text{sel}}=QP$. Its variation is $\delta X_{\text{sel}}=(\delta Q)P+Q(\delta P)$. The constant-rank polar theorem then determines δT_{sel} from δX_{sel} , including both source- and target-support motion. Consequently $\delta S_{\text{sel}}^\wedge$ and δP_{sel} are determined by the original five input variations; no additional support perturbation is introduced.

62.3 Selected formation and witness

The transported formation obeys $\delta \rho_F^{\text{sel}}=(\delta T_{\text{sel}})\rho_F T_{\text{sel}}^\dagger + T_{\text{sel}}(\delta \rho_F)T_{\text{sel}}^\dagger + T_{\text{sel}}\rho_F(\delta T_{\text{sel}})^\dagger$. The unnormalized selected witness obeys the identical three-term transport rule with ρ_W . The first survival mass varies as $\delta m_{\text{sel}}=\text{tr}[(\delta S_{\text{sel}}^\wedge)\rho_W + S_{\text{sel}}^\wedge(\delta \rho_W)]$. The normalized selected witness therefore contains both direct transport variation and the normalization correction $-\rho_W^\wedge \delta m_{\text{sel}}/m_{\text{sel}}^2$.

62.4 Neutral split and Gram metric

Because $C=(I+Y)^\wedge$, $\delta C=-C(\delta Y)C$. Since $B=I-C$, $\delta B=-\delta C=C(\delta Y)C$. $G=I+Y$ gives $\delta G=\delta Y$. These identities make the neutral/balance variations fully endogenous to δY .

62.5 Horizontal support frame

Choose an orthonormal support frame Z with $ZZ^\dagger=P_{\text{sel}}$ and horizontal variation $\delta Z=(\delta P_{\text{sel}})Z$. Let $H_Z=Z^\dagger GZ$. Then $\delta H_Z=(\delta Z)^\dagger GZ + Z^\dagger(\delta G)Z + Z^\dagger G(\delta Z)$, and $\delta H_Z^\wedge = -H_Z^\wedge(\delta H_Z)H_Z^\wedge$. Therefore every term in δP_G follows by the product rule from δP_{sel} and δY .

62.6 Load derivatives

For $\Lambda_C = \text{tr}(G P_G C \rho_F^{\text{sel}} C P_G^\dagger)$, $\delta \Lambda_C$ is the sum of the variations of its six nonconstant factors. Explicitly it contains δG , δP_G on both sides, two δC terms, and $\delta \rho_F^{\text{sel}}$. Λ_B has the same structure with B . Hence $\delta \alpha$ and $\delta \beta$ are ordinary quotient-rule expressions in $\delta \Lambda_C$ and $\delta \Lambda_B$.

62.7 Analysis derivative

Write $A_C = \sqrt{\alpha} P_{GC}$ and $A_B = \sqrt{\beta} P_{GB}$. Then $\delta A_C = [\delta \alpha / (2\sqrt{\alpha})] P_{GC} + \sqrt{\alpha} (\delta P_G) C + \sqrt{\alpha} P_G (\delta C)$, and analogously for A_B . Stacking the two blocks gives δA . Thus the full analysis derivative is a closed function of the five original input variations on the declared regular stratum.

63 SVD Differential in the Complex Case

63.1 Tangent decomposition

Let $F = \delta A$ and $M = U^\dagger F V$. Define $\Omega_U = U^\dagger \delta U$ and $\Omega_V = V^\dagger \delta V$. Orthonormality implies both are skew-Hermitian. Differentiating $A = U \Sigma V^\dagger$ and compressing between $U^\dagger$ and V gives $M = \Omega_U \Sigma + \delta \Sigma - \Sigma \Omega_V$.

63.2 Diagonal phase equation

The diagonal entry is $M_{ii} = \delta \sigma_i + \sigma_i [(\Omega_U)_{ii} - (\Omega_V)_{ii}]$. Since the diagonal entries of a skew-Hermitian matrix are purely imaginary, $\delta \sigma_i = \text{Re } M_{ii}$ and $(\Omega_U)_{ii} - (\Omega_V)_{ii} = i \text{Im}(M_{ii}) / \sigma_i$. Only the common phase remains free. This is why imposing both diagonal connections equal to zero is generally inconsistent.

63.3 Off-diagonal equations

For $i \neq j$ the pair M_{ij} and conjugate M_{ji} gives a two-by-two linear system for $(\Omega_U)_{ij}$ and $(\Omega_V)_{ij}$. Its determinant is $\sigma_j^2 - \sigma_i^2$. Therefore the componentwise frame derivative requires separated singular values. The resulting coefficients become large as a singular-value gap closes, while the invariant cluster subspace can remain smooth.

For $i \neq j$, $(\Omega_U)_{ij} = (\sigma_j M_{ij} + \sigma_i \text{conjugate}(M_{ji})) / (\sigma_j^2 - \sigma_i^2)$, and $(\Omega_V)_{ij} = (\sigma_i M_{ij} + \sigma_j \text{conjugate}(M_{ji})) / (\sigma_j^2 - \sigma_i^2)$. These coefficients solve the declared complex tangent system on a simple singular-value stratum.

63.4 Normal support components

The parts of δU and δV outside their current ranges are fixed by F and Σ^{-1} : $(I - UU^\dagger) \delta U = (I - UU^\dagger) F V \Sigma^{-1}$ and $(I - VV^\dagger) \delta V = (I - VV^\dagger) F^\dagger U \Sigma^{-1}$. These are essential because R_4 depends directly on V and K_+ depends directly on U .

63.5 Successor derivatives

$\delta Y_+ = \delta \Sigma \Sigma + \Sigma \delta \Sigma$. For $K_+ = U^\dagger (K \oplus K) U$, $\delta K_+ = (\delta U)^\dagger (K \oplus K) U + U^\dagger (\delta K \oplus \delta K) U + U^\dagger (K \oplus K) \delta U$. Hence the successor derivative contains both direct K variation and the endogenous frame response.

64 R4, Witness, and Formation Differential

64.1 R4 overlap

$X_{R4} = V^\dagger P_{\text{sel}}$ gives $\delta X_{R4} = (\delta V)^\dagger P_{\text{sel}} + V^\dagger \delta P_{\text{sel}}$. The constant-rank polar derivative determines δT_4 , δS_4^- and δS_4^+ entirely from this overlap variation.

64.2 Second survival mass

The general product rule is $\delta m_4 = \text{tr}[(\delta S_4^{\wedge-})\rho_W^{\text{sel}} + S_4^{\wedge-}\delta\rho_W^{\text{sel}}]$. Within this ambient kernel it simplifies to $\delta m_4 = 0$, because $S_4^{\wedge-} = P_{\text{sel}}$ and the selected density is supported and normalized there. This does not set δT_4 to zero.

64.3 Successor witness

The three product-rule terms in $\delta(T_4 \rho_W^{\text{sel}} T_4^\dagger)$ remain necessary. Here $m_4 = 1$ and $\delta m_4 = 0$, so no second normalization correction survives. Direct frame and carrier variations still change the successor witness.

64.4 Inherited formation

$\delta \rho_{F,\text{inh}} = (\delta T_4) \rho_F^{\text{sel}} T_4^\dagger + T_4 (\delta \rho_F^{\text{sel}}) T_4^\dagger + T_4 \rho_F^{\text{sel}} (\delta T_4)^\dagger$. Formation has no witness-style normalization term.

65 Seed Eigenprojector Derivation

65.1 Moving complement

$E = I - S_4^{\wedge+}$ gives $\delta E = -\delta S_4^{\wedge+}$. Choose Z_E with $Z_{EZE}^\dagger = E$ and horizontal $\delta Z_E = (\delta E)Z_E$. The reduced Hermitian seed operator is $H_E = Z_E^\dagger K_- + Z_E$.

65.2 Reduced operator variation

Product differentiation gives $\delta H_E = (\delta Z_E)^\dagger K_- + Z_E + Z_E^\dagger \delta K_- + Z_E + Z_E^\dagger K_- + \delta Z_E$. With the horizontal frame this becomes the embedded formula already stated in the canonical section.

65.3 Simple eigenvalue

For $H_E v = \lambda v$, $\|v\| = 1$, $\pi = vv^\dagger$, first-order perturbation theory gives $\delta \lambda = v^\dagger \delta H_E v = \text{tr}(\pi \delta H_E)$. Define R_λ as the inverse of $H_E - \lambda I$ on v -perp and zero on $\text{span}(v)$, equivalently the Moore–Penrose pseudoinverse for a simple eigenvalue.

65.4 Projector derivative

Differentiating the eigenvalue equation and fixing the phase horizontally gives $\delta \pi = -R_\lambda (\delta H_E) \pi - \pi (\delta H_E) R_\lambda$. Embedding into the moving complement adds the two outer frame terms: $\delta \Pi_{\text{seed}} = (\delta E) \Pi_{\text{seed}} + \Pi_{\text{seed}} (\delta E) + Z_E (\delta \pi) Z_E^\dagger$.

65.5 Seed derivative

$\rho_{\text{seed}} = -\kappa_F \lambda \Pi_{\text{seed}}$ therefore gives $\delta \rho_{\text{seed}} = -\kappa_F [(\delta \lambda) \Pi_{\text{seed}} + \lambda \delta \Pi_{\text{seed}}]$. This closes the seed derivative on the simple negative eigenvalue stratum.

66 Algebraic Admissibility Checks

66.1 Projector property

Because $S_4^{\wedge+}$ and Π_{seed} are orthogonal projectors with $\Pi_{\text{seed}} \leq I - S_4^{\wedge+}$, their product vanishes in both orders. Hence $(S_4^{\wedge+} + \Pi_{\text{seed}})^2 = S_4^{\wedge+} + \Pi_{\text{seed}} = P_-$.

66.2 Positivity of formation

Congruence by T_4 preserves positivity, so $\rho_{F,inh} \geq 0$ whenever $\rho_F^{sel} \geq 0$. If $\lambda < 0$ and $\kappa_F > 0$, $-\kappa_F \lambda \Pi_{seed} \geq 0$. Therefore $\rho_{F,+} = \rho_{F,inh} + \rho_{seed} \geq 0$.

66.3 Witness trace

$\text{tr}(T_4 \rho_W^{sel} T_4^\dagger) = \text{tr}(S_4^\wedge - \rho_W^{sel}) = m_4$. Dividing by m_4 gives $\text{tr} \rho_{W,+} = 1$ whenever $m_4 > 0$.

66.4 Hermiticity

Hermiticity of K is preserved by $K_+ = U^\dagger(K \oplus K)U$. Hermiticity and positivity of Y_+ follow from $\Sigma^\wedge$. Thus the successor remains in the canonical state type on every nonterminal branch.

67 Detailed Scalar Control Branch

67.1 Setup

On the full-persistence scalar branch $P=W=1$, $F=f_0>0$ and $0<y<1$. The neutral split gives $\alpha=y^2/(1+y^2)$ and $\beta=1/(1+y^2)$. The reduced analysis operator has one active column; U is its normalized two-component frame, $V=1$, R_4 is the identity, and the seed complement vanishes.

67.2 Successor

Direct substitution yields $K_+=K$, $F_+=F$, $W_+=1$ and $P_+=1$. The Gram successor is $f(y)=2y^2/[(1+y)^2(1+y^2)]$.

67.3 Damping proof

For $y>0$, $f(y)/y=2y/[(1+y)^2(1+y^2)]$. The denominator minus the numerator equals $(1+y)^2(1+y^2)-2y=1+2y^2+2y^3+y^4>0$. Hence $f(y)/y<1$ and $f(y)<y$. Positivity is immediate.

67.4 Consequence

The branch demonstrates genuine geometric damping while K and F are neutral. Any theorem claiming strict contraction of the complete state in a fixed positive product norm would contradict these neutral directions. This control branch therefore fixes the correct scope of future stability claims.

The scalar sequence decreases toward zero. A positive limit would satisfy $f(y)=y$, which the strict inequality excludes. Consequently positive load margins cannot be uniform along its infinite tail. Geometric damping alone is not a positive stability reserve against every disturbance.

Numerical controls. Repeated double-precision checks with NumPy and random seed 20261007 used 40 complex matrix preparations. The maximum $\|S_4^\wedge - P_{sel}\|$ was $2.47e-15$, maximum $|m_{4-1}|$ was $1.67e-15$ and maximum gauge residual was $2.18e-14$. These are checks of the algebraic transport stage, not empirical validation or an assertion that every preparation passes the final seed gate.

Independent central differences gave residuals $1.70e-9$ for the polar differential, $1.10e-10$ for the simple seed projector and $1.02e-10$ for the full endogenous analysis derivative. Complex SVD tangent reconstruction had residual $2.17e-15$. Scalar damping was checked on 1000 points in $(0,1)$. Finite numerical tolerances do not replace the analytical proofs.

The graphical calculations use 31-by-31 uniform parameter grids. Ambient panels use $Y=\text{diag}(y1,0.72)$, $P=F=W=vv^\dagger$, $v=(\cos \theta, \sin \theta)$, $0.08 \leq y1 \leq 0.92$ and $0.12 \leq \theta \leq 1.45$. Formation units are set to one. Other panels state their distinct formulas and ranges. Exactly zero singular values are not retained; numerical rank checks use a $1e-12$ threshold in the control family. The theoretical definition uses exact rank.

68 Physical Measurement Equations

68.1 Time

The time bridge predicts $\Delta t_k = T_{\text{clock}}(\Xi_k)$. A calibrated clock yields Δt_k^{obs} with uncertainty σ_t . A basic residual is $r_t = (\Delta t_{\text{pred}} - \Delta t_{\text{obs}}) / \sigma_t$. The bridge is empirically meaningful only if T_{clock} is fixed before the holdout interval.

68.2 Energy and spectroscopy

A quantum bridge supplies an energy-valued Hamiltonian. For an emission from $E_a > E_b$ use $\Delta E_{ab} = E_a - E_b$ and $\nu_{ab} = \Delta E_{ab} / h$; an unordered pair has positive line frequency $|E_a - E_b| / h$. Line centers and systematic uncertainties require independent calibration.

68.3 Geometry

A spacetime bridge supplies $g_{\mu\nu}(x; \Xi)$. Proper time, invariant distance, redshift, lensing, or tidal observables must be calculated from that metric and compared with clock/interferometric/astronomical data. Internal G cannot be substituted directly because it has positive signature and different type.

68.4 Quantum outcomes

Given independently prepared Ξ , bridge $\rho_Q(\Xi)$, and calibrated POVM effect E_a , $p(a|\Xi) = \text{tr}(E_a \rho_Q(\Xi))$. Repeated outcomes define a binomial/multinomial likelihood. This gives an actual statistical test of the bridge.

68.5 Particle observables

Π_{SM} must output masses, charges, couplings, mixings, and decay/scattering parameters. Comparison targets include PDG values and cross sections. A group-theoretic resemblance without numerical predictions is not sufficient.

68.6 Biological observables

A biological bridge must define boundaries and species before J_{in} , J_{out} and J_{net} can be measured. Viability predictions require a time horizon and perturbation class, not only an instantaneous flux balance.

68.7 Subject-level observables

World/Self variables require task-dependent operational readouts and interventions. Γ_{subj} must predict independent behavioral/physiological/report data before any phenomenal interpretation is considered.

69 Falsification Protocol for the World Equation

69.1 Freeze the mathematical kernel

No equation in the finite kernel is changed after observing the physical holdout data.

69.2 Freeze the bridge

The map from Ξ to the tested observable, all calibration constants, nuisance parameters, and preprocessing are specified on training/calibration data.

69.3 Independent preparation

The relevant Ξ components are identified without using the same outcome that will test the prediction.

69.4 Predict

The BFG produces a point prediction or probability distribution with uncertainty.

69.5 Compare

A pre-specified statistic compares prediction and holdout data against strong baselines using the same inputs.

69.6 Interpret failure locally

A failed quantum bridge falsifies that quantum identification first; it does not logically falsify the matrix identities of the finite kernel. Repeated failures across bridge classes can, however, undermine the claim that the BFG describes physical reality.

70 Dimensional-Consistency Handbook

70.1 Dimensionless kernel sector

The canonical finite kernel is most naturally formulated with Y, P, Q, C, B, G , the partial isometries, projectors, SVD frames, and witness density operators dimensionless. This is consistent because these objects encode geometry, support, relative weighting, or normalized state rather than an SI-valued physical observable. Singular values of the dimensionless analysis operator are then dimensionless and so is $Y_+ = \Sigma^2$.

70.2 Formation sector

K and ρ_F are assigned the common formation unit E_F in the canonical mathematical class. This choice is not a claim that E_F is joules. It is a type constraint ensuring that $K_+ = U^\dagger (K \oplus K) U$ and $\rho_{F,+}$ can be compared inside the seed construction. If a physical bridge later identifies K with an energy-like operator while ρ_F is assigned a different physical dimension, the constant κ_F performs the one-time unit conversion in $\rho_{\text{seed}} = -\kappa_F \lambda \Pi_{\text{seed}}$.

70.3 Witness sector

ρ_W is normalized and dimensionless. Projector traces against ρ_W are therefore dimensionless probabilities inside the kernel. The product $m_{\text{tot}} = m_{\text{sel}} m_4$ is a conditional survival probability for the two-stage mathematical selection. Calling it a physical detection probability requires a separate instrument bridge.

70.4 Time sector

The event index k has no time dimension. ε in the scalar normal form can be dimensionless if both equations use the same model-time parameter, or it can encode a ratio of time scales after nondimensionalization. Physical seconds enter only through T_{clock} . This prevents the common mistake of assigning SI seconds to a recursion count without calibration.

70.5 Spacetime sector

Use $x^0 = ct$ and all x^μ in metres when g is dimensionless. SI proper time is $d\tau = c^{-1} \sqrt{-g_{\mu\nu} dx^\mu dx^\nu}$ for a timelike displacement. Natural units $c=1$ must be converted before reporting seconds. Coordinates mixing seconds and metres cannot have all metric components dimensionless. In the natural-unit disformal convention $[\varphi] = M$, $[\partial \varphi] = M^2$ and $[\lambda] = M^{-4}$.

70.6 Quantum and particle sector

ρ_Q and POVM effects are dimensionless. H_{hat} carries energy. Mass can be quoted in kg or energy units divided by c^2 ; cross sections carry area; decay widths carry energy and lifetimes time. A successful Π_{SM} must

therefore output quantities with these physical dimensions or fixed conversion constants. Dimensional matching is necessary but not sufficient for a correct bridge.

71 Stratum and Regime Manual

71.1 Selection-gap stratum

The selected spectral subspace remains fixed in dimension as long as no eigenvalue of Y crosses the threshold 1. Inside such a region Q varies smoothly as an invariant spectral projector. At a crossing, the selected rank changes. The event is not an infinitesimal perturbation inside the old stratum; it is a transition to a new active model stratum.

71.2 Polar-rank stratum

For X_{sel} or X_{R4} , constant rank means the number of positive singular values remains fixed. The polar partial isometry and its source/target projectors then admit the constant-rank derivative used in this paper. When the smallest positive singular value reaches zero, the pseudoinverse loses a uniform bound and the support dimension changes.

71.3 Dual-load stratum

The formulas for α and β require $\Lambda_C + \Lambda_B > 0$. The canonical dual-load analysis further assumes both loads positive. If one load vanishes, a one-channel limiting rule would constitute a different model branch unless explicitly included. The present canonical branch therefore treats failure of the required dual-load condition by its declared terminal semantics.

71.4 Simple-SVD stratum

The componentwise SVD derivative assumes positive simple singular values. At exact degeneracy, the singular subspace can remain well defined while individual vectors are gauge-dependent. The quotient state remains meaningful; the simple-frame differential is replaced by a block/subspace description if one wishes to cross the degeneracy smoothly.

71.5 Seed stratum

A simple strictly negative lowest complement eigenvalue defines the rank-one seed. Its sign and eigenvalue gap must remain separated on a regular seed stratum. If $E \neq 0$ with $\lambda_{\min} \geq 0$ the event is terminal; a multiple negative minimum is also terminal. The no-seed branch is $E=0$. Terminal surfaces do not inherit a nonterminal differentiability claim.

72 Proof Appendix Core Invariants

72.1 Positive Gram invariant

Σ is a diagonal matrix of positive singular values in the reduced SVD. For any vector x , $x^\dagger Y_- x = x^\dagger \Sigma^2 x = \|\Sigma x\|^2 \geq 0$. Hence Y_- is positive semidefinite. Adding the identity gives $x^\dagger G_- x = \|x\|^2 + \|\Sigma x\|^2 > 0$ for nonzero x .

The positive dual-load ambient analysis strengthens this positivity result to $0 < Y_- < I$ by the bound proved in Section 17. The result is not monotone comparison with the old Y matrix, since the carrier and coordinates can change.

72.2 Hermiticity of K_-

If $K = K^\dagger$ then $K \oplus K$ is Hermitian. Therefore $K_-^\dagger = [U^\dagger (K \oplus K) U]^\dagger = U^\dagger (K \oplus K)^\dagger U = K_-$. This guarantees a real seed spectrum after compression.

72.3 Positivity under transport

For any positive ρ and arbitrary T , $x^\dagger T \rho T^\dagger x = (T^\dagger x)^\dagger \rho (T^\dagger x) \geq 0$. Thus both selected and R4-transported formation/witness numerators remain positive.

72.4 Witness normalization

Cyclicity of the trace gives $\text{tr}(T_4 \rho_W^{\text{sel}} T_4^\dagger) = \text{tr}(T_4^\dagger T_4 \rho_W^{\text{sel}}) = \text{tr}(S_4 \rho_W^{\text{sel}}) = m_4$. Division by positive m_4 gives trace one.

72.5 Projector successor

S_4^+ and Π_{seed} are self-adjoint idempotents and $\Pi_{\text{seed}} \leq E = I - S_4^+$. Therefore $S_4^+ \Pi_{\text{seed}} = \Pi_{\text{seed}} S_4^+ = 0$. Expanding P_{-+}^2 leaves exactly $S_4^{++} \Pi_{\text{seed}}$.

72.6 Gauge covariance

Under an allowed target unitary W , every target operator O transforms to $W^\dagger O W$. Products, sums, spectra, traces, positivity, rank, and projector relations are preserved under unitary conjugation. Therefore the equivalence class, rather than any chosen matrix basis, is the canonical successor.

72.7 Survival bounds

For a density operator ρ and projector S , $0 \leq S \leq I$. Positivity gives $\text{tr}(S\rho) \geq 0$, while $\text{tr}[(I-S)\rho] \geq 0$ gives $\text{tr}(S\rho) \leq \text{tr}(\rho) = 1$. Apply this to both conditioned stages to obtain m_{sel}, m_4 in $[0,1]$.

For this kernel the second bound sharpens to $m_4 = 1$. Carrier dimension and mass must still be distinguished: the seed may enlarge P_{-+} within C^r while the witness remains on S_4^+ .

73 Measurement Protocol Time and Geometry

73.1 Clock calibration

Select a physical preparation family and independently measured clock intervals. Fit or derive T_{clock} only on calibration data. Freeze the map. For each holdout BFG state, predict the next interval and compare with traceable SI clock measurements. A systematic state-dependent residual indicates failure of the clock bridge even if the event ordering remains correct.

73.2 Metric calibration

A spacetime bridge must specify coordinates or coordinate-invariant predictions, the relation between recursive state and matter/source variables, and any constants. The preferred tests are invariant or operational: proper-time ratios, frequency redshift, interferometric phase, geodesic deviation, lensing, or gravitational-wave propagation. Coordinate components alone are not observables.

73.3 Separating internal and spacetime metrics

No fit is allowed to set $g_{\mu\nu}$ numerically equal to the positive G matrix merely because both are called metrics. Their dimensions, signatures, domains, and transformation laws differ. A valid bridge is a typed map, not a relabeling.

73.4 Gravitational falsification

Once the bridge and source model are frozen, predictions can be compared with a standard gravitational baseline. A failure means that the chosen geometry/dynamics bridge is rejected. If every viable bridge requires post-hoc modification, the stronger claim that the BFG explains gravity loses empirical content.

74 Measurement Protocol Quantum and Particle Physics

74.1 State preparation

A preparation protocol must determine the BFG state or the components needed by Π_Q without using the later measurement outcome. Tomographic data used to define ρ_Q cannot be reused as independent confirmation of the same bridge.

74.2 Instrument specification

The POVM or instrument is calibrated independently. For a fixed effect E_a , the bridge predicts $p_a = \text{tr}(E_a \rho_Q)$. Repetition produces a likelihood. Competing BFG bridges are compared on the same preparations and effects.

74.3 R4 compatibility test

If one proposes that an actual success Kraus operator is T_4 , then its effect must equal $S_4^{\wedge} = T_4^\dagger T_4$ and the physical output transport must match the R4 target coordinates. General nonprojective effects require additional channels and cannot be identified directly with a partial isometry.

74.4 Particle parameter test

A Π_{SM} bridge must output a finite vector of quantities such as masses, charge assignments, mixing parameters, couplings at a specified renormalization scale, widths, and branching fractions. These predictions are compared with independent PDG values and collider/precision measurements. Merely reproducing $SU(3) \times SU(2) \times U(1)$ notation is not a numerical test.

74.5 Holdout principle

Calibration constants may be estimated on a declared training set. The test set must remain untouched until predictions are frozen. The same baselines and uncertainty treatment are applied to BFG and comparison models.

75 Measurement Protocol Atomic, Molecular, and Biological Sectors

75.1 Atomic closure

A quantitative atom bridge must specify constituent degrees of freedom, Hamiltonian or open-system generator, charge/mass inputs, and the map from recursive state to these quantities. Spectral lines, ionization energies, transition rates, and scattering observables provide direct tests.

75.2 Molecular closure

A molecular bridge must define nuclear/electronic or effective degrees of freedom and predict binding energies, equilibrium geometries, vibrational spectra, or reaction observables. The symbolic Cl_{NM} becomes a real mathematical operator only when its input/output spaces and update are specified.

75.3 Cellular closure

For a cell-level bridge, define the physical boundary, transported species, energy carriers, internal formation variables, and repair variables. Measure gross influx and outflux separately. Viability is tested over a horizon under a specified perturbation class, not by one stationary snapshot.

75.4 Causal minimality

A claim $n_{\min}^{\text{causal}}=3$ requires an admissible three-role realization and exclusion of equivalent one- and two-role realizations inside the same declared equivalence class. Four-or-more-role realizations need not be excluded because they may simply be nonminimal.

75.5 Multiscale closure

Higher biological levels require a coarse-graining map and a test that the proposed macrostate captures future-relevant dynamics. Information synergy means that variables cannot be discarded merely because each is individually weak. Intervention tests are required for causal claims.

76 Measurement Protocol World, Self, and Experience

76.1 World readout

A world variable W should predict externally oriented observations beyond what a chosen reduced baseline already predicts. Its definition must specify inputs, history dependence, horizon, and loss function.

76.2 Self readout

A self variable S should predict or causally regulate the system's own future state/action. Interventions on S must be defined within the admissible model class; destructive interventions do not establish the intended causal role.

76.3 World/Self closure

A closed U_{WS} must reproduce the projected successor of the underlying BFG dynamics. If two underlying states share the same (W,S) but produce different projected successors, the proposed functional state is not closed.

76.4 Witness integration

The kernel witness must be linked to the functional World/Self variables by an explicit map. Equality of normalized matrices or norms is not a lineage criterion. Directed transport and unnormalized survival bookkeeping remain separate.

76.5 Phenomenal hypothesis

Only after functional criteria are operational and independently measured can H_{phen} be tested against report, behavior, physiology, perturbation, or other declared evidence. No purely mathematical definition can by itself supply empirical confirmation of phenomenality.

77 Plain-Language Glossary for Readers

77.1 Formation

Formation is the positive operator ρ_F that carries the magnitude and structure of what is being recursively retained or newly seeded. It is intentionally not normalized like a probability distribution.

77.2 Witness

Witness is the positive trace-one ρ_w . The first selection can remove original witness mass. The present ambient R4 then transports its selected support isometrically, so $m_4=1$. Historical mass is recorded separately from the five-component normalized state.

77.3 Gram geometry

Y and G describe an internal positive geometry used to select and measure active recursive directions. This geometry is endogenous because it is both an input to and an output of the update.

77.4 Selection

Selection means restriction to the subspace whose Y-spectrum lies in the declared interval and overlaps the current persistent carrier. Polar transport preserves the directed relation between source and target supports.

77.5 Load

A load is a scalar obtained by evaluating transported formation through one neutral/balance channel in the current internal metric. Loads determine the relative weights of the analysis branches.

77.6 R4

R4 is the second recursive re-alignment step. It compares the SVD source frame with the selected carrier and produces the partial isometry used for final witness/formation transport.

77.7 Seed

A seed is positive formation from a simple negative lowest complement mode. When $E=0$ no seed is required. When $E \neq 0$ without such a unique negative mode, this kernel terminates.

77.8 Admissibility

Admissibility is the conjunction of the gates required for a particular model class. It is not identical to stability or viability; a state can satisfy a static gate and still leave the admissible region on the next update.

77.9 Persistence

Persistence means continued admissible directed evolution of the relevant typed entity. Witness persistence, formation persistence, biological viability, and subject persistence are different typed predicates.

77.10 Bridge law

A bridge law maps the mathematical BFG state to a physical or functional observable. Bridge laws are where the world equation becomes experimentally vulnerable.

78 Equation Roadmap Reading One Complete World-Equation Step

78.1 Input

Begin with the declared five-component finite state Ξ_k . This is the state chosen for the model; no absolute minimality theorem is claimed. The historical witness record is additional trajectory bookkeeping.

78.2 Neutral decomposition

Compute $C_k = (I + Y_k)^{-1}$ and $B_k = I - C_k$. These operators divide the current Gram load into complementary channels without introducing an external parameter.

78.3 Spectral admissibility

Compute $Q_k = 1_{[0,1)}(Y_k)$. This selects the currently sub-threshold spectral sector. Intersect it with persistence through $X_{\text{sel}} = Q_k P_k$.

78.4 Directed selection transport

Take $T_{\text{sel}} = \text{polar}(X_{\text{sel}})$. Read $S_{\text{sel}}^{\wedge}$ from $T_{\text{sel}}^{\dagger} T_{\text{sel}}$ and P_{sel} from $T_{\text{sel}} T_{\text{sel}}^{\dagger}$. These projectors are the actual source and target supports of the first directed transport.

78.5 First survival bookkeeping

Transport formation and witness. Compute m_{sel} before normalizing the witness. If $m_{\text{sel}} = 0$, terminate. Otherwise condition ρ_w^{sel} while retaining m_{sel} as historical survival information.

78.6 Endogenous metric

Compute $G = I + Y$ and the G -orthogonal projector P_G onto the selected target support. The current geometry now directly controls how formation is evaluated.

78.7 Endogenous loading

Evaluate Λ_C and Λ_B and construct α, β . No free branch weight is introduced on the dual-load stratum.

78.8 Recursive analysis

Construct A from the weighted C/B branches. Its reduced SVD supplies the active rank r , the next dimension, the successor Gram spectrum, and the frames required by K_+ and R_4 .

78.9 Successor geometry and kernel

Set $Y_+ = \Sigma^{\wedge 2}$ and $K_+ = U^{\dagger}(K \oplus K)U$. At this point the next active space is already fixed as $C^{\wedge r}$.

78.10 Recursive re-alignment

Form $X_{R_4} = V^{\dagger} P_{\text{sel}}$. The positive C -channel implies it is already a partial isometry, with source P_{sel} . It realigns the selected carrier and fixes the free target complement.

78.11 Second survival bookkeeping

The nominal second mass is identically $m_4 = 1$. Therefore $m_{\text{tot}} = m_{\text{sel}}$. Update the separate original-mass record $w_{k+1} = w_k m_{\text{sel},k}$ before interpreting a normalized witness as retention.

78.12 Formation inheritance

Transport selected formation by T_4 without normalization. This gives the inherited successor formation.

78.13 Complement test

Compute $E = I - S_4^+$. If $E=0$ there is no free target complement. If E is nonzero, compress K_+ to $\text{Ran}(E)$ and inspect its spectrum.

78.14 Seed or no-seed

If $E=0$ continue without seed. If $E \neq 0$ a simple strictly negative minimum produces the seed; otherwise terminate as T_{seednone} or T_{seeddeg} . No numerical tie-break replaces the rank-one rule.

78.15 Output

Assemble $\Xi_{k+1} = (K_+, \rho_F, \rho_W, Y_+, P_+)$ and then pass to its unitary equivalence class. The next event repeats the same algorithm on the newly generated active space.

79 Figure Companion What Each 3D Graphic Means

Figure 1 Section 3 Conditional readout. Conditional readout. $\phi_E = (E_{\text{exp}}/E_{\text{bind}})(I/I_0)$; ratios are independent inputs, not identified Xi data. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 2 Section 4 Computed matrix family. Computed matrix family. $Y = \text{diag}(y_1, y_2)$; $G = I + Y$; the positive metric is internal, not spacetime. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 3 Section 5 Computed projector. Computed projector. $Q = 1_{[0,1]}(Y)$; threshold 1 defines a discontinuous change of selected rank. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 4 Section 6 Computed control family. Computed control family. $Y = \text{diag}(1/2, 2)$, $P = vv^T$ and $W = \text{diag}(p, 1-p)$; polar selection has source P. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 5 Section 7 Computed ambient kernel. Computed ambient kernel. $Y = \text{diag}(y_1, 0.72)$; $P = F = W = vv^T$; $v = (\cos \theta, \sin \theta)$; formation unit $E_F = 1$. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 6 Section 8 Computed ambient kernel. Computed ambient kernel. $A: H$ to H direct sum H ; no restriction to the selected rank-one carrier. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 7 Section 9 Computed ambient kernel. Computed ambient kernel. The second singular value can vanish on a rank boundary; an arbitrary full-SVD padding is excluded. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 8 Section 10 Computed ambient kernel. Computed ambient kernel. $Y = \text{diag}(y_1, y_2)$, $P = F = I$; coupled global formation loads determine alpha and beta. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 9 Section 11 Computed ambient kernel. Computed ambient kernel. $m_4 = 1$ to numerical precision; the exact proof uses $\text{Ran}(P_{\text{sel}})$ contained in $\text{Ran}(A^{\dagger})$. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 10 Section 12 Computed bookkeeping. Computed bookkeeping. $w_{\text{next}} = w * m_{\text{sel}}$; conditioned witness trace remains one and cannot encode this product. Horizontal and vertical coordinate ranges are the sampled

grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 11 Section 13 Computed spectral control. Computed spectral control. A one-dimensional complement with $\lambda = -x$ gives seed trace κx ; each κ is a separate fixed-unit model. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 12 Section 14 Computed rule. Computed rule. For a nonempty complement, $\lambda < 0$ and a simple minimum are required; $\lambda \geq 0$ is terminal. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 13 Section 15 Computed bound. Computed bound. 3 times $\|\Delta X\|_2$ divided by $\sigma_{\min,+}$; an upper bound is not an attained response. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 14 Section 16 Conditional perturbation scale. Conditional perturbation scale. Illustrates inverse-gap scaling of the reduced resolvent; not a full derivative of an arbitrary kernel. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 15 Section 17 Exact scalar kernel. Exact scalar kernel. $P=W=1$ and $F>0$; $f(y)=2y^2/[(1+y)^2(1+y^2)]$; K and F remain neutral. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 16 Section 18 Conditional stability model. Conditional stability model. Assumes $V=z^2$ and $V_{\text{next}}=(1-r)V$; no such global BFG certificate is constructed here. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 17 Section 19 Computed mediator control. Computed mediator control. $N_B R N_B \dagger = n^2 R$; amplitude mediation uses $N_A = n^2$, not $N_A = n$. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 18 Section 20 Schematic constraint model. Schematic constraint model. A margin diagram only; actual viability requires the unique nonterminal BFG trajectory and specified gates. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 19 Section 21 Computed conditional ode. Computed conditional ode. $a=1.5$, $\mu>0$, $b<1$; this positive branch exists because $a>1-b$. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 20 Section 22 Schematic countermodel. Schematic countermodel. Projection $P_i(x,h)=x$ with successor $x+h$ is not closed; this is a fiber test, not a BFG bridge. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 21 Section 23 Exact scalar kernel. Exact scalar kernel. One reclosure event is counted; no conversion to physical seconds is claimed. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 22 Section 24 Schematic embedding. Schematic embedding. An illustrative surface is not a Lorentz metric, gravitational field equation or prediction. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 23 Section 25 Assumed potential model. Assumed potential model. $V=x^2+y^2$ is a pedagogical potential; stability of spacetime must follow from a specified action and bridge. Horizontal and vertical

coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 24 Section 26 Conditional quantum model. Conditional quantum model. For a calibrated $E0=|0\rangle\langle 0|$, $p=\text{tr}(E0 \rho)$; the state-identification map is additional physics. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 25 Section 27 Computed bridge comparison. Computed bridge comparison. At (0.75,0.25), hypotheses A and B differ by 0.5; no actual measured outcomes are used. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 26 Section 28 Computed effect test. Computed effect test. A direct R4 effect is a projector. Generic positive effects with interior eigenvalues fail that test. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 27 Section 29 Schematic dependency. Schematic dependency. The surface labels a parameter dependency only; no Standard Model mass or coupling is derived. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 28 Section 30 Conventional conditional model. Conventional conditional model. $E(r)=1/(2r^2)-a/r$; ordinary quantum scaling, not a new BFG atomic spectrum. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 29 Section 31 Schematic binding landscape. Schematic binding landscape. The illustrative energy x^2+y^2-1 marks a negative binding region; it is not a BFG molecular potential. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 30 Section 31 Computed flux identity. Computed flux identity. $J_{\text{net}}=J_{\text{out}}-J_{\text{in}}$. Stationary net flux permits nonzero gross flows; no cell viability is inferred. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 31 Section 32 Schematic functional model. Schematic functional model. A product is only a visual aid. Pi_W , Pi_S , lineage and causal closure must be independently defined. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

Figure 32 Section 33 Schematic status panel. Schematic status panel. The zero sheet denotes absence of a derived prediction, not absence of subjective experience. Horizontal and vertical coordinate ranges are the sampled grids; the plotted quantity is identified on the height axis. Surface colour follows height and has no additional physical meaning.

80 Frequently Asked Technical Questions

80.1 Is the BFG one equation or many

It is one quotient evolution equation with an explicit factorization. The sub-operators are comparable to the factors of a numerical integrator or a structured map: they specify how the single successor is computed.

80.2 Why not use a scalar state

The five-component operator tuple is the declared state of this construction. It resolves spectra, magnitude, normalized witness and moving supports explicitly. This is not a proof that every alternative representation requires five components.

80.3 Why is the successor dimension allowed to change

Because the reduced SVD treats only active positive singular directions as successor coordinates. Rank is therefore a dynamical output. Fixed-dimensional padding would introduce null directions whose physical and seed status would be arbitrary.

80.4 Does gauge quotienting hide physical information

No information invariant under allowed coordinate changes is removed. Spectra, traces, positivity, rank, masses, and all simultaneously conjugated relations are preserved. The quotient removes only coordinate labels.

80.5 Is m_{tot} a Born probability

Not by definition. It is a mathematically normalized two-stage witness survival probability inside the kernel. A physical Born/instrument interpretation is a separate bridge hypothesis.

80.6 Is G spacetime

No. G is positive definite and internal. Spacetime is Lorentzian and requires a separate typed bridge.

80.7 Does scalar damping prove stability of the universe

No. It proves one scalar inequality on a declared branch. Full-state stability requires control of every endogenous derivative path and must account for neutral directions.

80.8 Does the Standard Model follow already

Not numerically. The BFG supplies a reconstruction program, but masses, charges, couplings, mixing parameters, and scattering observables must be generated by a fixed Π_{SM} and tested.

80.9 Does witness mean consciousness

No. Witness is a mathematical persistence role. World/Self and phenomenal experience require additional operational and empirical bridges.

80.10 What would make the BFG a physical world equation

A finite set of frozen measurement bridges that convert independently prepared BFG states into successful quantitative predictions across time, geometry, quantum/particle, and higher-level sectors without post-hoc remapping.

81 Canonical BFG at a Glance

$$[\Xi_{k+1}] = \text{Ubar}_{\text{BFG}} \setminus ([\Xi_k])$$

Selection $\rightarrow$ first transport $\rightarrow$ positive metric $\rightarrow$ endogenous dual loads $\rightarrow$ ambient reduced SVD $\rightarrow$ successor K and $Y \rightarrow$ isometric $R_4 \rightarrow$ formation inheritance $\rightarrow$ $E=0$ continuation or negative seed or terminal outcome $\rightarrow$ quotient successor.

Core invariants: Hermitian K ; positive formation; trace-one witness with separately recorded original mass; $0 < Y_+ < I$; orthogonal P ; $n_+ \leq n$. R_4 gives $m_4=1$. Nonterminal subsequent events preserve witness mass; seed termination remains possible.

Core non-universalities: positive Gram geometry is not Lorentz spacetime; witness survival is not automatically a Born probability; scalar damping is not full-state contraction; recursive event count is not physical seconds;

Cl_N labels are not transition laws until their operations are specified; functional subject structure is not by definition phenomenal experience.

Physical closure targets: T_{clock} for seconds; M_ϕ for the order observable; M_g for Lorentzian geometry and gravity; Π_Q plus calibrated instruments for quantum outcomes; Π_{SM} for masses, charges, couplings and mixing; atom/molecule/biology transition maps; Π_w , Π_s , U_{ws} and Γ_{subj} for functional subject structure.

Empirical rule: every bridge must be fixed before holdout testing. A successful prediction supports that bridge and its coupled kernel assumptions; a failed prediction rejects the tested bridge first. The Balance–Field Equation becomes a physical world equation only through a finite, frozen, independently testable measurement interface.

Completion status. Level-0 to Ξ , scalar information/order to Ξ , a closed Ξ -to- Ψ factor, calibrated physical time, Lorentzian dynamics, quantum instruments and particle parameters, quantitative atoms and molecules, biological viability and functional/phenomenal subject bridges are not constructed here. They remain explicit finite requirements, not consequences established by the illustrations.

Declarations

Author: Marcel Theodor Wende, Independent Researcher. E-mail: marcel-wende@gmx.net. ORCID: 0009-0007-4028-2208.

Author contribution: Conceptualization, BFG synthesis, canonical finite-kernel formulation, interpretation, and manuscript responsibility: Marcel Theodor Wende. Co-authored precursor source lines are credited in References [1]–[3].

Data availability: No external empirical dataset is analyzed. The 3D computational panels use explicitly stated synthetic matrix families and independently checked scalar formulas. Open physical transitions are schematic or show conventional conditional models and are not BFG predictions of measured systems. Numerical controls support calculations, not proofs or empirical identification.

Funding: No funding source is asserted.

Competing interests: No competing interest is asserted.

Ethics: No human participants, animal experiments, or identifiable personal data were used.

82 Purpose, provenance and claim discipline

The BFG seeks a common architecture for persistence and structural change across levels of organization. The philosophical account in *Die kosmische Ordnung* [22] associates unfolding with change, binding with retention, and information with distinguishable organization; structure is the outcome of their relationship. These roles motivate the mathematics but do not uniquely determine it. In particular, metaphorical binding is not automatically the gravitational interaction, and information as meaning is not automatically a configuration count or matrix entropy.

The finite operator construction is inherited from the supplied 6 October 2026 preprint [21]. The supplied manuscript’s 81 numbered sections, equations, dictionaries and 32 figures are retained as Part I of the assembled paper. Its bibliography is consolidated at the end of this paper, preserving source reference numbers 1–20 and avoiding duplicate entries. Part II continues with Section 82 and deepens the complete mathematical update, differentials, stability distinctions and physical interfaces. A new simulation atlas continues the figure numbering after Figure 32 and supplies a reproducible numerical or explicitly conditional model for each computational and application stage. Earlier illustrative panels remain for provenance, with their original limits on interpretation. Numerical results reported in Part II are fresh synthetic controls rather than copied residuals. The editable LaTeX file represents Part II; the supplied original PDF is also required to reproduce the complete assembled document.

Five statuses are used: *definition* for model choices; *assumption* for additional constitutive commitments; *theorem* for derived statements; *open bridge* for unspecified identifications; and *empirical hypothesis* for a bridge that can be tested. The word “universal” concerns a proposed scope of application. Demonstrating that scope requires sector-specific measurements and cannot be achieved by calling every force “balance.”

The supplied preprint credits three precursor lines: Level-0 admissibility with Thomas W. Loker, mediated conjugate closure with Luke Kenneth Casson Leighton, and the distinction between function and experience with Alfredo Pereira Jr. Those attributions are retained as provenance from that document; the precursor manuscripts were not independently examined for this new draft.

83 State space, units and equivalence

At event k , all five components act on a finite complex Hilbert space $\mathcal{H}_k$, with $n_k = \dim \mathcal{H}_k$:

$$\Xi_k = (K_k, \rho_{Fk}, \rho_{Wk}, Y_k, P_k). \quad (121)$$

The domain is

$$K = K^\dagger, \quad \rho_F \geq 0, \quad \rho_W \geq 0, \quad \text{tr } \rho_W = 1, \quad Y = Y^\dagger \geq 0, \quad P = P^\dagger = P^2. \quad (122)$$

Formation is not required to have unit trace. K and ρ_F share a declared formation unit E_F ; all other kernel objects are dimensionless. If another interpretation assigns different units, a fixed positive factor κ_F must convert the kernel unit to the formation unit. In the matched class $\kappa_F = 1$. No state-dependent unit conversion is implied.

For unitary L on $\mathcal{H}_k$, the simultaneous transformation

$$\Xi \sim (L^\dagger K L, L^\dagger \rho_F L, L^\dagger \rho_W L, L^\dagger Y L, L^\dagger P L) \quad (123)$$

defines a quotient state $[\Xi]$. Physical interpretations, if supplied, must respect this equivalence. The finite dynamics is a map on the disjoint union of these finite-dimensional quotient spaces, with terminal outcomes adjoined.

A pre-operational Level-0 domain $\Phi_0 \in \mathcal{X}_0$ may motivate admissibility, difference and recurrence. The preparation map $J_0 : \mathcal{X}_0 \rightarrow \mathcal{X}_\Xi$ is not specified by the finite kernel. Neither Φ_0 nor a theological interpretation is inserted as a numerical multiplier.

84 The complete recursive event

Subscripts k are suppressed during a single event. Each gate is evaluated before the later stages.

84.1 Spectral selection and directed transport

Set

$$C = (\mathbf{1} + Y)^{-1}, \quad B = YC = \mathbf{1} - C, \quad G = \mathbf{1} + Y > 0, \quad (124)$$

$$Q = \mathbf{1}_{[0,1)}(Y), \quad X_{\text{sel}} = QP, \quad T_{\text{sel}} = \text{polar}(X_{\text{sel}}). \quad (125)$$

The canonical polar factor vanishes on the kernel. Define its actual source and target supports:

$$S_{\text{sel}}^- = T_{\text{sel}}^\dagger T_{\text{sel}}, \quad P_{\text{sel}} = T_{\text{sel}} T_{\text{sel}}^\dagger. \quad (126)$$

They need not coincide. If $\text{rank}(QP) = 0$, the event is $T_{\text{sel}0}$. Otherwise transport the states:

$$\rho_F^{\text{sel}} = T_{\text{sel}} \rho_F T_{\text{sel}}^\dagger, \quad \rho_W^{\text{un,sel}} = T_{\text{sel}} \rho_W T_{\text{sel}}^\dagger, \quad (127)$$

$$m_{\text{sel}} = \text{tr}(S_{\text{sel}}^- \rho_W), \quad \rho_W^{\text{sel}} = \rho_W^{\text{un,sel}} / m_{\text{sel}}. \quad (128)$$

$0 \leq m_{\text{sel}} \leq 1$; zero mass gives $T_{\text{w}1}$. The mass has a conditional survival meaning within this kernel, not an independently established Born-rule meaning.

84.2 Endogenous metric projection and formation loads

Choose Z with $Z^\dagger Z = \mathbf{1}$ and $ZZ^\dagger = P_{\text{sel}}$. Set

$$D = Z^\dagger G Z, \quad P_G = Z D^{-1} Z^\dagger G. \quad (129)$$

$D > 0$, $P_G^2 = P_G$ and $P_G^\dagger G = G P_G$. Thus P_G is the G -orthogonal projector onto the selected target, independently of its frame Z .

$$\Lambda_C = \text{tr}(G P_G C \rho_F^{\text{sel}} C P_G^\dagger), \quad (130)$$

$$\Lambda_B = \text{tr}(G P_G B \rho_F^{\text{sel}} B P_G^\dagger). \quad (131)$$

Both are nonnegative. The canonical dual-load class requires both to be strictly positive; otherwise the event is T_{load} . On this branch define

$$\alpha = \frac{\Lambda_B}{\Lambda_C + \Lambda_B}, \quad \beta = \frac{\Lambda_C}{\Lambda_C + \Lambda_B}, \quad \alpha + \beta = 1. \quad (132)$$

84.3 Ambient analysis and active successor dimension

The domain of the analysis operator is the full ambient input, not merely the selected carrier:

$$A : \mathcal{H} \rightarrow \mathcal{H} \oplus \mathcal{H}, \quad A = \begin{bmatrix} \sqrt{\alpha} P_G C \\ \sqrt{\beta} P_G B \end{bmatrix}. \quad (133)$$

Use the reduced rank- r SVD, retaining exactly the positive singular values:

$$A = U \Sigma V^\dagger, \quad r = \text{rank } A, \quad U^\dagger U = V^\dagger V = \mathbf{1}_r, \quad \Sigma = \text{diag}(\sigma_i) > 0. \quad (134)$$

The successor space and two successor operators are

$$\mathcal{H}_+ = \mathbb{C}^r, \quad n_+ = r, \quad Y_+ = \Sigma^2, \quad K_+ = U^\dagger (K \oplus K) U. \quad (135)$$

Replacing A by AZ or padding zero singular directions changes the model. A finite numerical rank tolerance is an implementation convention, not an exact-rank theorem.

84.4 R4 realignment and inheritance

Define

$$X_4 = V^\dagger P_{\text{sel}}, \quad T_4 = \text{polar}(X_4), \quad S_4^- = T_4^\dagger T_4, \quad S_4^+ = T_4 T_4^\dagger. \quad (136)$$

The ambient construction implies, as proved below,

$$T_4 = V^\dagger P_{\text{sel}}, \quad S_4^- = P_{\text{sel}}, \quad S_4^+ = V^\dagger P_{\text{sel}} V, \quad (137)$$

$$m_4 = \text{tr}(S_4^- \rho_W^{\text{sel}}) = 1, \quad \rho_{W+} = T_4 \rho_W^{\text{sel}} T_4^\dagger, \quad \rho_F^{\text{inh}} = T_4 \rho_F^{\text{sel}} T_4^\dagger. \quad (138)$$

The nominal second conditioning is therefore lossless. In particular $\text{tr } \rho_F^{\text{inh}} = \text{tr } \rho_F^{\text{sel}}$.

84.5 Seed or terminal rule

Set $E = \mathbf{1}_r - S_4^+$. If $E = 0$, define $\rho_{\text{seed}} = \Pi_{\text{seed}} = 0$. If $E \neq 0$, choose a frame Z_E for its range and form

$$H_E = Z_E^\dagger K_+ Z_E. \quad (139)$$

A simple isolated minimum $\lambda < 0$ with rank-one eigenprojector π gives

$$\Pi_{\text{seed}} = Z_E \pi Z_E^\dagger, \quad \rho_{\text{seed}} = -\kappa_F \lambda \Pi_{\text{seed}}, \quad \rho_{F+} = \rho_F^{\text{inh}} + \rho_{\text{seed}}, \quad P_+ = S_4^+ + \Pi_{\text{seed}}. \quad (140)$$

Orthogonality $\Pi_{\text{seed}} \leq E$ preserves the projector condition. If the minimum is nonnegative the outcome is T_{seednone} ; if the negative minimum is multiple it is T_{seeddeg} . No arbitrary eigenvector resolves a degeneracy. A nonempty complement never continues without a seed under this rule.

The ordered terminal list is therefore selection rank, first witness mass, positive dual loads, and finally the seed test. After positive loads, zero analysis rank, zero R4 rank and zero second mass are impossible; they are implementation checks only.

85 Gauge covariance and the ambient bound

Theorem 2 (Unique quotient successor) *Every nonterminal event defines one successor equivalence class, independent of the ambient basis, support frames and admissible SVD-frame choices.*

Proof. Replacing Z by ZW leaves P_G unchanged. Under an input conjugation by L , $A' = (L^\dagger \oplus L^\dagger)AL$, with compatible SVD frames $U' = (L^\dagger \oplus L^\dagger)U$ and $V' = L^\dagger V$. The target matrices then agree. A common SVD phase or rotation of an exactly degenerate singular block has $U' = UW$, $V' = VW$, with $W\Sigma = \Sigma W$. It gives $K'_+ = W^\dagger K_+ W$, $Y'_+ = W^\dagger Y_+ W$ and $T'_4 = W^\dagger T_4$. Inheritance, complement compression and the simple seed projector transform by the same target conjugation. The seed frame itself disappears under embedding. Terminal conditions depend only on invariant ranks, traces and spectra. $\square$

Theorem 3 (Ambient transport and geometry) *On the positive dual-load branch,*

$$\text{rank } P_{\text{sel}} \leq r \leq n, \quad 0 < Y_+ < \mathbf{1}_r, \quad S_4^- = P_{\text{sel}}, \quad m_4 = 1. \quad (141)$$

Proof. Since $GC = \mathbf{1}$ and $GB = Y$,

$$P_G C = ZD^{-1}Z^\dagger, \quad P_G B = ZD^{-1}Z^\dagger Y. \quad (142)$$

The first map is Hermitian, with range exactly $\text{Ran } P_{\text{sel}}$. Because its coefficient $\sqrt{\alpha}$ is positive, $\text{Ran } P_{\text{sel}} \subseteq \text{Ran } A^\dagger = \text{Ran } V$, so $VV^\dagger P_{\text{sel}} = P_{\text{sel}}$. This proves the R4 identities and the trace-one second mass. The selected range lies in $\text{Ran } Q$. If $\eta < 1$ is the largest selected eigenvalue of Y , then $D \geq \mathbf{1}$, $\|P_G C\|_2 \leq 1$ and $\|P_G B\|_2 \leq \eta$. Hence

$$\|A\|_2^2 \leq \alpha + \beta\eta^2 < 1. \quad (143)$$

All retained singular values are positive; squaring them gives $0 < Y_+ < \mathbf{1}_r$. Rank bounds follow from the first channel and the ambient domain. $\square$

Consequently $Q_+ = \mathbf{1}$. Both ρ_{W_+} and ρ_{F_+} are supported on P_+ . Every later nonterminal selection is $T_{\text{sel}} = P$, so its witness mass is one. These are stronger restrictions than a generic lossy recursion.

86 Order through exact accounting and observable distinctions

86.1 Balanced weighted loads

Proposition 4 (Cross-weight balance) *The cross-weights are the unique weights summing to one that equalize positive weighted channel loads:*

$$\boxed{\alpha\Lambda_C = \beta\Lambda_B = \frac{\Lambda_C\Lambda_B}{\Lambda_C + \Lambda_B}.} \quad (144)$$

Proof. Solve $\alpha\Lambda_C = (1 - \alpha)\Lambda_B$, or substitute (132) directly. $\square$

This is an exact algebraic balance. It is not an independent experimental discovery or physical energy-conservation law. The unweighted loads may differ considerably. An optional dimensionless diagnostic is

$$d_L = \frac{\Lambda_C - \Lambda_B}{\Lambda_C + \Lambda_B}, \quad b_L = \frac{4\Lambda_C\Lambda_B}{(\Lambda_C + \Lambda_B)^2} = 1 - d_L^2 = 4\alpha\beta. \quad (145)$$

$0 < b_L \leq 1$ measures raw-load symmetry; it does not by itself establish viability.

86.2 Formation stock, loss and seed

Define

$$M_F = \text{tr } \rho_F, \quad \ell_F = \text{tr}[(\mathbf{1} - S_{\text{sel}}^-)\rho_F] \geq 0, \quad (146)$$

and $s_F = 0$ for $E = 0$, or $s_F = -\kappa_F \lambda > 0$ on a nonempty admissible seed branch.

Proposition 5 (Formation balance) *Each nonterminal event satisfies*

$$\boxed{M_{F,+} - M_F = s_F - \ell_F.} \quad (147)$$

On subsequent nonterminal events, $\ell_F = 0$.

Proof. Polar transport gives $\text{tr } \rho_F^{\text{sel}} = \text{tr}(S_{\text{sel}}^- \rho_F) = M_F - \ell_F$. R4 preserves that trace and the seed adds s_F . After one nonterminal event formation is supported on P and $Q = \mathbf{1}$, so $S_{\text{sel}}^- = P$ and $\ell_F = 0$. $\square$

This stock has formation units; it is not called physical mass or energy without a bridge. Increasing stock does not imply increasing complexity.

86.3 Spectral diversity and concentration

Positive loads imply $M_F > 0$. Define mathematical readouts

$$\hat{\rho}_F = \rho_F / M_F, \quad H_F = -\text{tr}(\hat{\rho}_F \ln \hat{\rho}_F), \quad D_F = e^{H_F}, \quad R_F = \frac{1}{\text{tr } \hat{\rho}_F^2}, \quad (148)$$

with $0 \ln 0 = 0$. Eigenvalue entropy gives

$$0 \leq H_F \leq \ln \text{rank } \rho_F \leq \ln n, \quad 1 \leq R_F \leq D_F \leq \text{rank } \rho_F. \quad (149)$$

D_F and R_F measure effective mode participation, not semantic information, biological function or experience. For $n > 1$, $O_F = 1 - H_F / \ln n$ is a concentration readout; set $O_F = 1$ conventionally at $n = 1$. Concentration and diversity are different properties, and no monotonicity of either is asserted.

For orthogonal inherited and seed modes of stocks $f, s > 0$, the normalized weights are $p = f/(f+s)$ and $1-p$. Then $D_F = \exp[-p \ln p - (1-p) \ln(1-p)]$, maximized at $f = s$ and tending to one as $s/f \rightarrow \infty$. Thus adding stock and an occupied mode need not continuously increase effective diversity. This is a conditional configuration, not a claimed realized orbit for every f, s .

86.4 Historical survival and information density

Record original witness mass separately:

$$w_0 = 1, \quad w_{k+1} = w_k m_{\text{sel},k} m_{4,k} = w_k m_{\text{sel},k}. \quad (150)$$

The normalized successor alone cannot recover w . For example $Y = \text{diag}(1/2, 2)$, $P = \mathbf{1}$, $K = -\mathbf{1}$, $\rho_F = \mathbf{1}$ and $\rho_W = \text{diag}(p, 1-p)$ give $m_{\text{sel}} = p$ but the same one-dimensional conditioned successor for all $0 < p \leq 1$.

A configuration-counting measure is separately specified by

$$I_{\text{conf}} = \log_b |\mathcal{C}_\theta(X)|, \quad 1 \leq |\mathcal{C}_\theta(X)| < \infty, \quad b > 1. \quad (151)$$

It is not automatically Shannon entropy or H_F [10]. A physical information density further requires a positive calibrated volume: $\iota = I_{\text{conf}}/V_{\text{phys}}$. The kernel supplies neither that volume nor an apparatus model. An energy-ratio readout $\varphi_E = (E_{\text{exp}}/E_{\text{bind}})(I_{\text{conf}}/I_0)$ is dimensionless for matched energy units, $E_{\text{bind}}, I_0 > 0$; a counting-ratio alternative requires a positive denominator. Neither is identified with Φ_0 .

87 Finite seeding and eventual full persistence

The next result makes the autonomous kernel's scope more precise than a general statement that dimensions cannot grow.

Theorem 6 (Finite seed capacity) *Consider an infinite nonterminal orbit of the autonomous canonical kernel. Let $p_k = \text{rank } P_k$. After the first event, $n_{k+1} \leq n_k$ and*

$$p_{k+1} = p_k + \delta_k, \quad \delta_k = \begin{cases} 0, & E_k = 0, \\ 1, & E_k \neq 0. \end{cases} \quad (152)$$

There are at most $n_1 - p_1$ further seed events. Full persistence $P = \mathbf{1}$ is reached after finitely many events and thereafter is invariant. Subsequent formation stocks are constant.

Proof. For $k \geq 1$, $Q_k = \mathbf{1}$ and $T_{\text{sel},k} = P_k$, hence $\text{rank } P_{\text{sel},k} = p_k$. R4 is isometric on this carrier, so $\text{rank } S_{4,k}^+ = p_k$. A nonempty complement requires exactly one orthogonal rank-one seed on every nonterminal event. Thus $p_{k+1} = p_k + 1$ there, and otherwise $p_{k+1} = p_k$. Since $p_{k+1} \leq n_{k+1} \leq n_1$, at most $n_1 - p_1$ such increases can occur. An infinite orbit must therefore encounter $E_k = 0$. At that event $r_k = p_k$ and $P_{k+1} = \mathbf{1}_{r_k}$. With $P = \mathbf{1}$ and $0 < Y < \mathbf{1}$, the next analysis has full column rank because $\sqrt{\alpha}C$ is invertible; R4 has full target support. Thus full persistence remains, no further seed is added, and (147) makes the stock constant. $\square$

This is a theorem about continued nonterminal dynamics; a finite orbit may terminate before saturation. Full persistence does not mean the entire state becomes fixed: K , spectra and frames can still evolve. The theorem excludes indefinitely repeated seed creation in a single fixed autonomous finite model. It also bounds $D_{F,k} \leq n_0$. A continually growing physical hierarchy therefore needs a declared environmental or state-space extension; this manuscript does not introduce unspecified new dimensions to obtain that conclusion by fiat.

88 Stability, viability and a minimal driven control

88.1 Exact autonomous scalar branch

For $n = 1$, $P = \rho_W = 1$, $\rho_F = F > 0$ and $0 < y < 1$, the seed complement is empty and

$$\Lambda_C = \frac{F}{1+y}, \quad \Lambda_B = \frac{Fy^2}{1+y}, \quad \alpha = \frac{y^2}{1+y^2}, \quad \beta = \frac{1}{1+y^2}. \quad (153)$$

The successor satisfies $K_+ = K$, $F_+ = F$, $P_+ = \rho_{W+} = 1$ and

$$y_+ = f(y) = \frac{2y^2}{(1+y)^2(1+y^2)}. \quad (154)$$

For $y > 0$, $(1+y)^2(1+y^2) - 2y = 1 + 2y^2 + 2y^3 + y^4 > 0$, proving $0 < f(y) < y$. On this actual branch $y < 1$ ensures selection. The sequence tends to zero; a positive limit would contradict

the fixed-point inequality. Zero itself fails the positive dual-load gate. Therefore damping does not give a uniform positive load reserve along the infinite tail.

At $y = 1/2$, $\alpha = 1/5$, $\beta = 4/5$, both weighted loads equal $2F/15$, $b_L = 16/25$ and $y_+ = 8/45$. Geometric damping coexists with neutral K, F directions and $D_F = 1$. No universal strict full-state contraction in a fixed positive product norm is possible on this branch.

88.2 A declared constant-input extension

To show exactly what a minimal dynamic addition can strengthen, introduce a dimensionless geometric input d before each scalar event:

$$\tilde{y}_k = y_k + d, \quad y_{k+1} = f(\tilde{y}_k), \quad 0 < d < 3/4. \quad (155)$$

This is a separate input-coupled model, not a consequence of the autonomous kernel and not yet a physical cosmological law. Other scalar components are retained; $0 < \tilde{y} < 1$ guarantees nonterminal selection and positive loads.

Theorem 7 (Positive driven scalar regime) *For constant $0 < d < 3/4$, the map $F_d(y) = f(y + d)$ sends $[0, 1/4]$ into $(0, 1/4)$ and has a unique positive fixed point y_d^* . Every initial $0 \leq y_0 < 1 - d$ enters that interval after one step and converges geometrically to y_d^* .*

Proof. For $0 < x < 1$,

$$f'(x) = \frac{4x(1-x^3)}{(1+x)^3(1+x^2)^2} > 0, \quad f(x) < f(1) = 1/4. \quad (156)$$

Also $f'(x) \leq 4x/(1+x)^3 \leq 16/27 < 1$, with the last maximum at $x = 1/2$. For $y \in [0, 1/4]$, $d \leq y+d \leq d+1/4 < 1$. Thus F_d is a contraction of the closed interval into itself; Banach's contraction theorem [27] gives a unique fixed point and the error bound $|y_k - y_d^*| \leq (16/27)^k |y_0 - y_d^*|$ for initial values in that interval. General stated initial values enter after one step. Since $F_d(0) > 0$, the fixed point is positive. $\square$

The constant input maintains a positive scalar geometric reserve. This is a rigorous limited improvement over decay toward the terminal boundary. It does not remove neutral K, F directions, prove robustness to arbitrary perturbations, or derive a physical energy source. A physical use must identify d , its resource budget and clock independently. The theorem does not extend unchanged to variable inputs.

88.3 Viability and transverse certificates

Given explicit application constraints $\mathcal{S}$, define

$$\mathcal{V}_N = \{x_0 : x_j \in \mathcal{S} \ (0 \leq j \leq N), \text{ all } N \text{ events nonterminal}\}, \quad \mathcal{V}_\infty = \bigcap_{N \geq 0} \mathcal{V}_N. \quad (157)$$

$\mathcal{V}_{N+1} \subseteq \mathcal{V}_N$. Static admissibility does not prove forward invariance. For a driven model the input sequence must be fixed, or a control policy and its quantifier declared; existential and all-input viability are different.

For transverse stability to an invariant neutral set, suppose on a forward-invariant regular fixed-dimensional region

$$c_1 \|z\|^2 \leq V \leq c_2 \|z\|^2, \quad V_+ - V \leq -c_3 \|z\|^2. \quad (158)$$

Then $V_+ \leq (1 - c_3/c_2)V$ when constants give a compatible nonnegative factor. No such global certificate is constructed here for the full kernel. Bounds on gaps and loads define candidate regular regions; they do not by themselves provide a contraction theorem.

88.4 Independent continuous mediator models

A conditional stock law $\dot{q} = a - \gamma_0(1-b)q$ has an attractive stock $q^* = a/[\gamma_0(1-b)]$ for constant $\gamma_0 > 0, b < 1$. At $b = 1$, $a = 0$ gives stationary stocks and $a \neq 0$ linear change; $b > 1$ amplifies the homogeneous part. This ODE is not the discrete kernel generator.

The supplied two-variable normal form is

$$\dot{q} = a - (1 - b + \mu n)q, \quad \epsilon \dot{n} = q - 1 - n, \quad a, \mu, \epsilon > 0. \quad (159)$$

A positive fixed point has $q^* = 1 + n^*$, $r^* = 1 - b + \mu n^* = a/q^* > 0$. Its Jacobian satisfies $\text{tr } J^* = -r^* - 1/\epsilon < 0$ and $\det J^* = (r^* + \mu q^*)/\epsilon > 0$, hence local asymptotic stability. A unique $n^* > 0$ exists when $a > 1 - b$; for $b > 1$ the positive branch additionally lies above $(b-1)/\mu$, as enforced by $r^* > 0$. Varying ϵ cannot produce a Hopf crossing on this positive branch. A separately assumed overconstraint $d(n) = 1 - \zeta n^2$ gives a static corridor $(b-1)/\mu < n < \sqrt{(1-d_{\min})/\zeta}$ under its stated positive parameters; that corridor is not itself a viability kernel.

For a specified finite linear mediator, $R_C = N_A^{1/2} R N_A^{1/2}$ with $N_A \geq 0$ equals $N_B R N_B^\dagger$ only under an appropriate identification, notably $N_B = N_A^{1/2}$. A frozen operator with spectral radius below one has powers tending to zero; this does not imply contraction of the endogenous nonlinear BFG update. Jordan structure and changing operators must be treated separately.

89 Functional closure, hierarchy and event time

A functional readout may organize nine roles as

$$\Psi = (\varphi_{\text{fun}}, I_{\text{fun}}, A_{\text{fun}}, G_{\text{fun}}, \Omega_{\text{fun}}, T_{\text{fun}}, S_{\text{fun}}, M_{\text{fun}}, C_{\text{fun}}). \quad (160)$$

A declared projection Π supports autonomous factor dynamics only if

$$\Pi(x) = \Pi(x') \implies \Pi(\mathcal{U}x) = \Pi(\mathcal{U}x'), \quad (161)$$

with terminality constant on fibers. For example projection $(x, h) \mapsto x$ fails closure when $x_+ = x + h$. Readout notation alone therefore does not yield a closed reduced equation. Input-dependent dynamics require the corresponding criterion for every specified input.

The event index k is not seconds. A calibrated quotient-invariant positive clock gives

$$t_k = t_0 + \sum_{j < k} T_{\text{clock}}([\Xi_j]). \quad (162)$$

Infinite nonterminal continuation in event count need not mean unbounded physical time; one additionally requires $\sum_k T_{\text{clock}} = \infty$. A continuous flow ϕ_F^t must independently satisfy $\Pi(\mathcal{U}\Xi) = \phi_F^{T_{\text{clock}}([\Xi])}(\Pi\Xi)$. A positive clock alone constructs no generator.

A hierarchy from particles to atoms, molecules, cells, organisms and functional subjects is a dependency graph. Each transition needs input and output state spaces, an evolution or closure map, preserved and lost quantities, and a measurement interface. Labels such as Cl_N cannot replace those maps. A new environment or increasing state-space dimension could support further novelty, but it would be an additional theory with a declared preparation, resource accounting and covariant update. Beyond the proved scalar input example, no such extension is asserted here.

90 A typed self-similar hierarchy

90.1 Recursion is not automatically fractality

Repeated events describe time evolution. A hierarchy describes levels with different carriers; a fractal additionally needs a self-similarity law and a metric scaling. These three structures are not interchangeable. We define an exact conditional hierarchy before attaching any physical interpretation.

For an integer multiplicity $m \geq 2$ and fixed density matrix τ_m on $\mathbb{C}^m$, let

$$\mathcal{L}_m(\Xi) = (K \otimes \mathbf{1}_m, \rho_F \otimes \tau_m, \rho_W \otimes \tau_m, Y \otimes \mathbf{1}_m, P \otimes \mathbf{1}_m). \quad (163)$$

This map preserves the declared types, formation stock and witness normalization. It multiplies the ambient and persistent ranks by m . It is an imposed level-transition law, not spontaneous dimensional growth within the autonomous kernel.

Theorem 8 (Self-similar lift on full persistence) *If $P = \mathbf{1}$, $0 < Y < \mathbf{1}$ and $\rho_F \neq 0$, the lifted state is nonterminal and*

$$[\mathcal{U}_{\text{BFG}}\mathcal{L}_m(\Xi)] = [\mathcal{L}_m\mathcal{U}_{\text{BFG}}(\Xi)] \quad (164)$$

after the canonical tensor-coordinate identification.

Proof. On this branch $Q = T_{\text{sel}} = P_G = \mathbf{1}$ and $E = 0$. The lifted channels are $C \otimes \mathbf{1}_m$ and $B \otimes \mathbf{1}_m$. Since $\text{tr } \tau_m = 1$, the lifted loads and weights equal the original loads and weights. Up to a fixed reordering of the two analysis blocks, $A_m = A \otimes \mathbf{1}_m$. A compatible SVD is $U_m = U \otimes \mathbf{1}_m$, $\Sigma_m = \Sigma \otimes \mathbf{1}_m$ and $V_m = V \otimes \mathbf{1}_m$. The successor K, Y are the tensor lifts, and isometric inheritance gives the lifted ρ_F, ρ_W . Degenerate singular frames are removed by the quotient. Full column rank makes the lifted complement empty, so no seed test changes the result. $\square$

The theorem does not hold without qualification on seed branches. If $E \neq 0$ has a simple negative complement minimum, replication gives $H_E \otimes \mathbf{1}_m$, whose minimum has multiplicity at least m . The replicated event therefore terminates as T_{seeddeg} . A hierarchical seed rule would have to resolve this through an additional covariant definition; no arbitrary tie-break is introduced here.

90.2 Each declared level supplies the next

The explicit stage law

$$[\Xi_{\ell+1}] = [\mathcal{L}_m\mathcal{U}_{\text{BFG}}(\Xi_\ell)] \quad (165)$$

is well defined on the full-persistence branch; $n_\ell = m^\ell n_0$. Here ℓ labels levels, not physical time. For $\tau_m = \mathbf{1}_m/m$, formation stock remains constant and $H_{F,\ell+1} = H_{F,+,\ell} + \ln m$. This entropy increment represents a declared replication of labels. It proves neither independent biological novelty nor emergent semantic content. In the scalar replicated family, $K_\ell = k_0 \mathbf{1}$, $Y_\ell = y_\ell \mathbf{1}$, $\rho_{F_\ell} = F_0 \mathbf{1}/n_\ell$ and $\rho_{W_\ell} = \mathbf{1}/n_\ell$. The geometry follows the exact scalar map or its separately declared input-driven counterpart.

90.3 An explicit spatial fractal illustration

To visualize metric self-similarity, choose four tetrahedral vertices v_j in $\mathbb{R}^3$ and similarity maps $S_j(x) = (x + v_j)/2$. Starting from one point, the finite point cloud at level $\ell+1$ is the union of the

four images of the preceding cloud. Under the open-set condition, the limiting self-similar set has similarity dimension determined by $4(1/2)^{d_s} = 1$, hence $d_s = 2$ [25]. The distinction between finite samples and limiting Hausdorff or box-counting dimension follows the standard dimension framework [26]. Each finite point cloud itself has dimension zero; the dimension-two statement concerns the infinite attractor. This geometry is an additional illustrative embedding. It is not derived spacetime and does not identify particles, cells or subjects with tetrahedral points.

The atlas includes every level from one through six: 4, 16, 64, 256, 1024 and 4096 points. The same replication labels are coupled to the exact tensor-family state through equal per-copy formation stock and a common scalar geometry. Total formation remains F_0 , per-copy formation is $F_0/4^\ell$, and mode entropy is $\ell \ln 4$. The pictures demonstrate this declared hierarchy, not the inevitability of a physical hierarchy.

91 Physical sectors and prospective tests

91.1 Spacetime and gravity

$G = \mathbf{1} + Y$ is an internal positive metric. A Lorentzian metric requires a manifold, signature, connection and a bridge of different mathematical type. An optional disformal ansatz is

$$g_{\mu\nu} = \eta_{\mu\nu} + \lambda \partial_\mu \phi \partial_\nu \phi, \quad X_\eta = \eta^{\mu\nu} \partial_\mu \phi \partial_\nu \phi. \quad (166)$$

Here ϕ is a new spacetime field, not the dimensionless order readout. The determinant and inverse are

$$\det g = \det \eta (1 + \lambda X_\eta), \quad g^{\mu\nu} = \eta^{\mu\nu} - \frac{\lambda \partial^\mu \phi \partial^\nu \phi}{1 + \lambda X_\eta}. \quad (167)$$

The Lorentz-connected nondegenerate branch requires $1 + \lambda X_\eta > 0$. A separately assumed Einstein–scalar action is

$$S = \int \sqrt{-g} \left[\frac{M_{\text{Pl}}^2}{2} R - \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) \right] d^4x + S_m. \quad (168)$$

Varying independent fields with appropriate boundary conditions is different from constraining $g = g[\phi]$ before variation. Neither this action nor its constants are derived from the finite BFG. Information density alone supplies no stress tensor, light-speed prediction, dark matter dynamics or dark energy equation of state.

91.2 Quantum states and instruments

A quantum bridge must specify $(\mathcal{H}_Q, \rho_Q, \mathcal{A}_Q, \hat{H})$ and calibrated effects. Two candidate identifications are $\rho_Q^A = \rho_W$ and $\rho_Q^B = \rho_F / \text{tr } \rho_F$. For

$$K = -\mathbf{1}_2, \quad Y = \text{diag}(1/2, 1/4), \quad P = \mathbf{1}_2, \quad \rho_F = \text{diag}(1, 3), \quad \rho_W = \text{diag}(3/4, 1/4), \quad (169)$$

the loads are $\Lambda_C = 46/15$ and $\Lambda_B = 19/60$. For independently fixed $E_0 = |0\rangle\langle 0|$, the candidates predict $p_A = 3/4$ and $p_B = 1/4$. The R4 success effect in this example is the identity and has probability one; it must not be substituted for E_0 .

For fixed T_4 , $\rho \mapsto T_4 \rho T_4^\dagger$ is completely positive with projective success effect S_4^- . A general POVM effect need not be projective. Since T_4 depends on the BFG state, the entire state-dependent family is not automatically one linear quantum channel. A classically conditioned instrument needs independently identified conditioning data.

91.3 Particles, atoms and molecules

A particle bridge must output representations, charges, couplings, masses and mixing parameters, rather than merely repeat a gauge-group name. Atoms then require a quantum Hamiltonian and electromagnetic coupling. Electron–nucleus binding is electromagnetic; nuclear binding involves the residual strong interaction. General metaphorical “binding” must not be substituted for gravitation at these scales [23].

The conventional estimate $E(r) \sim \hbar^2/(2\mu r^2) - e^2/(4\pi\epsilon_0 r)$ illustrates atomic localization and attraction. It does not derive $\hbar$, charge or atomic spectra from BFG. Cosmological cooling permits recombination; forming nuclei and forming neutral atoms are distinct stages [24]. Molecular closure requires interactions and an actual binding criterion such as $E(M) < \sum_i E(A_i)$.

91.4 Biology, world models and experience

Biological closure needs species, reaction laws, boundaries, energy/material flows, repair and persistence under a declared perturbation class. Reaction–diffusion theory supplies a concrete example of a typed pattern-forming mechanism [15]; citing it does not derive life from BFG. With $J_{\text{net}} = J_{\text{out}} - J_{\text{in}}$, stationary stock allows nonzero gross flows. Unavoidable DNA formation, cellular experience and a conscious universe are not conclusions of the finite kernel.

World and self readouts require maps from history, $W = \Pi_W(\Xi_{\text{history}})$ and $S = \Pi_S(\Xi_{\text{history}})$, plus a closed update and interventions. An operational subject gate Γ_{subj} must be defined before predicting data. The identification $\Gamma_{\text{subj}} = 1$ with phenomenal experience remains an empirical-ontological hypothesis, not a matrix theorem. Religious interpretation can remain philosophical motivation without being presented as a derived observable.

91.5 A measurement interface that can fail

For each sector, freeze preparation $\mathcal{M}_{\text{prep}}$, bridge $\mathcal{B}$, calibration and nuisance parameters before holdout outcomes. Compute

$$[\Xi_0] = \mathcal{M}_{\text{prep}}(D_{\text{prep}}), \quad p_{\text{BFG}}(D_{\text{test}} \mid D_{\text{prep}}, \mathcal{B}, \mathcal{C}). \quad (170)$$

Possible observables include calibrated clock intervals, spectroscopic lines $\nu_{ab} = |E_a - E_b|/h$, redshift or lensing from a Lorentzian metric, quantum outcome frequencies, particle parameters and biological fluxes. Each needs units, apparatus model, uncertainty, strong baseline and a predeclared rejection statistic. Full sensitivity-Jacobian rank can be a sufficient local identifiability criterion under suitable assumptions; it is not necessary, as the injective map $\theta \mapsto \theta^3$ at zero shows. A bridge freely refitted after every failure has no prospective predictive content.

92 Synthetic verification and interpretation

The accompanying Python file implements the entire nonterminal event and terminal gates. It uses declared numerical tolerances, fixed random seed 20261008 and finite synthetic matrices. Checks cover quotient covariance, R4 support and mass, the Gram bound, weighted loads, formation accounting, successful rank-one seeds, explicit terminal outcomes, the scalar map and the declared constant-input regime. The exact theorem statements depend on the proofs above; finite floating-point checks are implementation evidence only. Numerical summaries are included in Appendix B after execution. No experimental observations are analyzed.

The world-formula statement is the family of declared quotient orbits, with terminal outcomes included:

$$\mathfrak{W}_{\text{BFG}} = \{\text{Orb}_{\overline{\mathcal{U}}_{\text{BFG}}}([\Xi_0])\}. \quad (171)$$

The proposed identification of this architecture with physical reality is separate from its mathematical definition. The strongest additions of this manuscript are exact weighted balance, exact formation accounting, finite autonomous seed capacity, the positive driven scalar fixed-point theorem, and the covariant self-similar lift with its seed-degeneracy restriction. They sharpen both capability and limits. They do not establish monotone universal complexity, full-state strict contraction or inevitable emergence of a physical hierarchy.

93 Simulation atlas and stage-by-stage interpretation

All new surface plots use 21-by-21 declared grids unless the caption specifies a discrete point cloud. Color follows height and carries no independent physical meaning. Parameters in conditional application models are supplied inputs, not values inferred from BFG. Raw arrays are included in the reproducibility archive. Each figure names the stage in Part I or the philosophical source that it supplements. These stage-specific controls do not close missing physical bridge laws.

Dimensionless order readout

Figure 33. Dimensionless order readout
Conditional readout




Figure 33: Conditional readout. ϕ equals the independently supplied energy ratio times the independently supplied information ratio. A larger height is the defined product, not an identified BFG measurement or a proof of more complex organization.

Stage connection: Part I, Section 3 / Figure 1. **Declared grid:** energy ratio from 0 to 2, information ratio from 0 to 2. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Positive internal Gram geometry

Figure 34. Positive internal Gram geometry
Exact matrix family




Figure 34: Exact matrix family. For $Y = \text{diag}(y_1, y_2)$, $G = I + Y$. The plotted minimum eigenvalue remains at least one. This positive signature is internal and does not establish Lorentzian spacetime.

Stage connection: Section 4 / Figure 2. **Declared grid:** y_1 from 0 to 1, y_2 from 0 to 1. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Selection threshold and active rank

Figure 35. Selection threshold and active rank
Exact projector family




Figure 35: Exact projector family. Q selects eigenvalues in $[0,1)$. Heights are integer ranks zero, one or two. Threshold crossings are discontinuous changes of stratum; smooth differential formulas cannot be continued across them without reclassification.

Stage connection: Section 5 / Figure 3. **Declared grid:** y_1 from 0 to 1.4, y_2 from 0 to 1.4. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Unnormalized first witness survival

Figure 36. Unnormalized first witness survival
Exact selection control




Figure 36: Exact selection control. For $P=vv^T$, $v=(\cos \theta, \sin \theta)$, $Y=\text{diag}(1/2, 2)$, and $W=\text{diag}(p, 1-p)$, the source support is P away from zero overlap. The height is $m_{sel}=\text{tr}(PW)$, measured before normalization.

Stage connection: Section 6 / Figure 4. **Declared grid:** carrier angle from 0 to 1.5, witness population from 0 to 1. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Endogenous B-channel formation load

Figure 37. Endogenous B-channel formation load




Figure 37: Computed ambient kernel. Each grid point runs the complete BFG event for $Y=\text{diag}(y1,0.72)$, $P=F=W=vv^T$ and $K=-I$. The height is the actual formation-dependent B load, with formation unit set to one.

Stage connection: Section 7 / Figure 5. **Declared grid:** y1 from 0.08 to 0.92, carrier angle from 0.12 to 1.45. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Ambient analysis largest singular load

Figure 38. Ambient analysis largest singular load




Figure 38: Computed ambient kernel. Each height is computed from the reduced SVD of the full ambient analysis operator. The domain is H , not its selected rank-one subspace. This distinction allows a target complement in the declared family.

Stage connection: Section 8 / Figure 6. **Declared grid:** y_1 from 0.08 to 0.92, carrier angle from 0.12 to 1.45. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Positive secondary singular direction

Figure 39. Positive secondary singular direction
Computed ambient kernel




Figure 39: Computed ambient kernel. The second retained singular value is computed from the same full event. Zero denotes numerical rank one under the declared $1e-12$ tolerance; it is not an arbitrary full-SVD padding direction.

Stage connection: Section 9 / Figure 7. **Declared grid:** y_1 from 0.08 to 0.92, carrier angle from 0.12 to 1.45. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Successor Gram bound on full persistence

Figure 40. Successor Gram bound on full persistence




Figure 40: Computed ambient kernel. Here $P=F=I$, $W=I/2$ and $K=-I$. Global formation loads determine both weights. The maximum successor eigenvalue is below one as proved; the chart does not claim a full-state contraction or a componentwise monotone matrix order.

Stage connection: Section 10 / Figure 8. **Declared grid:** y1 from 0.08 to 0.92, y2 from 0.08 to 0.92. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

R4 second survival mass

Figure 41. R4 second survival mass
Computed ambient kernel




Figure 41: Computed ambient kernel. Every point computes R_4 from the reduced right SVD frame. The height is one to floating-point precision. The exact reason is inclusion of the selected carrier in $\text{Ran}(A^\dagger)$, not a fitted normalization.

Stage connection: Section 11 / Figure 9. **Declared grid:** y_1 from 0.08 to 0.92, carrier angle from 0.12 to 1.45. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Historical survival accounting

Figure 42. Historical survival accounting

Exact accounting identity




Figure 42: Exact accounting identity. The separately stored historical mass satisfies $w_{\text{next}} = w * m_{\text{sel}}$. A normalized density retains trace one while this historical scalar can be arbitrarily small. It is not a sixth silently added kernel component.

Stage connection: Section 12 / Figure 10. **Declared grid:** previous w from 0 to 1, m_sel from 0 to 1. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Admissible seed stock

Figure 43. Admissible seed stock
Conditional spectral seed control




Figure 43: Conditional spectral seed control. For a one-dimensional complement, $\lambda = -x$ and $\text{seed stock} = \kappa x$. Each vertical parameter is a different fixed-unit model. A complete successful seed event is separately verified in the numerical record.

Stage connection: Section 13 / Figure 11. **Declared grid:** negative eigenvalue magnitude from 0.05 to 2, fixed conversion factor from 0.2 to 2. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Seed sign and degeneracy gate

Figure 44. Seed sign and degeneracy gate




Figure 44: Exact declared gate. On a nonempty complement, a negative simple minimum continues; nonnegative minimum and zero gap do not. The height is a Boolean gate and the surface illustrates terminal semantics, not a physical phase transition.

Stage connection: Section 14 / Figure 12. **Declared grid:** lowest eigenvalue from -1 to 1, eigenvalue gap from 0 to 0.8. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Constant-rank polar sensitivity bound

Figure 45. Constant-rank polar sensitivity bound

Analytical bound sampled numerically




Figure 45: Analytical bound sampled numerically. The height samples $3 \cdot \text{norm}(\delta X) / \sigma_{\min_positive}$. A large upper bound near rank loss means no uniform control is certified; it does not show that every actual perturbation diverges.

Stage connection: Section 15 / Figure 13. **Declared grid:** smallest singular value from 0.05 to 1, perturbation norm from 0.01 to 0.2. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Actual two-mode seed mixing

Figure 46. Actual two-mode seed mixing
Conditional Hermitian eigenproblem




Figure 46: Conditional Hermitian eigenproblem. For $H = \begin{bmatrix} -1 & e \\ e & -1 + \text{gap} \end{bmatrix}$, the lowest-mode population transferred to the second basis direction is $(1 - \text{gap} / \sqrt{\text{gap}^2 + 4e^2}) / 2$. The graph computes an actual spectral response rather than drawing an inverse-gap heuristic alone.

Stage connection: Section 16 / Figure 14. **Declared grid:** eigenvalue gap from 0.05 to 1, off-diagonal perturbation from -0.25 to 0.25. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Autonomous recursive geometric damping

Figure 47. Autonomous recursive geometric damping




Figure 47: Exact scalar iterates. The height is obtained by repeated use of the exact scalar BFG map. Event counts are integer samples. K and F remain neutral; the shrinking geometry neither generates new modes nor guarantees a uniform positive load margin.

Stage connection: Section 17 / Figure 15. **Declared grid:** initial y from 0.05 to 0.95, event count from 0 to 5. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Conditional transverse Lyapunov certificate

Figure 48. Conditional transverse Lyapunov certificate

Assumed stability model




Figure 48: Assumed stability model. $V_k = 0.8^k V_0$ is an imposed certificate model. It visualizes what a transverse decay theorem would imply once its hypotheses are proved; it is not an established global BFG Lyapunov function.

Stage connection: Section 18 / Figure 16. **Declared grid:** initial squared distance from 0 to 1, event count from 0 to 12. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Mediator amplitude and power

Figure 49. Mediator amplitude and power
Exact frozen scalar mediator




Figure 49: Exact frozen scalar mediator. $N_B R N_B^\dagger = n^2 R$ in a scalar real model. The amplitude-mediated alternative agrees when $N_A = n^2$. This is a frozen-operator identity, not a contraction test for the changing nonlinear kernel.

Stage connection: Section 19 / Figure 17. **Declared grid:** mediator amplitude from 0 to 2, input stock from 0 to 2. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Twelve-event geometric viability margin

Figure 50. Twelve-event geometric viability margin

Computed declared scalar input model




Figure 50: Computed declared scalar input model. The minimum of $1-(y_k+d)$ over twelve input-coupled events is computed for the plotted preparations. Negative values mark initial inadmissibility; later formally plotted values after failure are not accepted trajectories. A margin alone is not biological viability.

Stage connection: Section 20 / Figure 18. **Declared grid:** initial y from 0.0 to 0.9, constant input d from 0.05 to 0.7. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Positive normal-form mediator equilibrium

Figure 51. Positive normal-form mediator equilibrium

Conditional ODE fixed point




Figure 51: Conditional ODE fixed point. For $b=0.4$, solve $a=(1+n)(0.6+\mu*n)$. The grid uses $a>0.6$ and $\mu>0$, so the positive fixed point exists. The autonomous discrete BFG is not identified with this separately assumed ODE.

Stage connection: Section 21 / Figure 19. **Declared grid:** injection a from 0.7 to 2, coupling μ from 0.1 to 2. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Failure of projected fiber closure

Figure 52. Failure of projected fiber closure
Exact closure countermodel




Figure 52: Exact closure countermodel. The projection $P_i(x,h)=x$ identifies all h values at each visible x . Their successors $x+h$ differ, so a closed x -only factor does not exist. A named functional state must pass this test, not merely receive a symbol.

Stage connection: Section 22 / Figure 20. **Declared grid:** visible x from 0 to 1, hidden h from -1 to 1. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Physical clock assumptions and event count

Figure 53. Physical clock assumptions and event count

Conditional calibrated-clock model




Figure 53: Conditional calibrated-clock model. Elapsed time equals $k \cdot T$ only when every event has an independently calibrated constant duration T . The graph shows that assumption, not a conversion to seconds derived by the finite kernel.

Stage connection: Section 23 / Figure 21. **Declared grid:** event count from 0 to 12, constant duration from 0.1 to 2. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Lorentzian disformal admissibility

Figure 54. Lorentzian disformal admissibility
Conditional spacetime completion




Figure 54: Conditional spacetime completion. For a homogeneous scalar gradient and $\eta=(-,+,+,+)$, the determinant factor is $1-\lambda v^2$. Positive values lie on the Lorentz-connected branch; zero is degenerate. Neither the field nor λ is supplied by internal Gram geometry.

Stage connection: Section 24 / Figure 22. **Declared grid:** λ from -1 to 1, time-gradient magnitude from 0 to 1.4. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Declared scalar gravitational potential

Figure 55. Declared scalar gravitational potential

Assumed Einstein-scalar sector




Figure 55: Assumed Einstein-scalar sector. $V(\phi)=a*\phi^2/2$ is a declared potential in an additional field theory. Sampling it is not solving Einstein equations or deriving gravity from BFG. A physical action, source model and bridge remain necessary.

Stage connection: Section 25 / Figure 23. **Declared grid:** field value from -1 to 1, fixed potential scale from 0.2 to 2. All surface inputs and output arrays are saved with this figure.

Evidence level: This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Figure 56. Quantum population and calibrated effect




Figure 56: Conditional quantum bridge. For $\rho = \text{diag}(p, 1-p)$ and $E = \text{diag}(e, 1-e)$, $\text{tr}(E\rho) = ep + (1-e)(1-p)$. State identification and effect calibration are external commitments. The graphic is a computable conventional measurement model, not observed frequencies.

Stage connection: Section 26 / Figure 24. **Declared grid:** state population p from 0 to 1, effect parameter e from 0 to 1. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Competing quantum bridge predictions

Figure 57. Competing quantum bridge predictions




Figure 57: Computed bridge discrimination. For externally fixed $E0=|0\rangle\langle 0|$, the two declared identifications differ by p_W-p_F . At populations 0.75 and 0.25 the difference is 0.5. An independently prepared experiment can reject a bridge; R4 success itself remains one here.

Stage connection: Section 27 / Figure 25. **Declared grid:** witness population from 0 to 1, formation population from 0 to 1. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Nonprojective effect diagnostic

Figure 58. Nonprojective effect diagnostic

Exact matrix-effect test




Figure 58: Exact matrix-effect test. The Frobenius idempotency residual is zero only at projector eigenvalues zero or one. Interior POVM eigenvalues can be valid effects but cannot directly equal a partial-isometry success projector.

Stage connection: Section 28 / Figure 26. **Declared grid:** effect eigenvalue e1 from 0 to 1, effect eigenvalue e2 from 0 to 1. All surface inputs and output arrays are saved with this figure.

Evidence level: These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Particle mass as an explicit external choice

Figure 59. Particle mass as an explicit external choice

Conventional conditional particle model




Figure 59: Conventional conditional particle model. The illustrative conventional relation $\text{mass} = g \cdot v / \sqrt{2}$ uses independently supplied coupling and scale. The surface replaces an unlabeled dependency illustration with a declared formula, without claiming that either parameter follows from BFG.

Stage connection: Section 29 / Figure 27. **Declared grid:** Yukawa coupling from 0.01 to 1, symmetry-breaking scale from 0.2 to 2. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Atomic localization and attraction

Figure 60. Atomic localization and attraction
Conventional conditional atomic model




Figure 60: Conventional conditional atomic model. $E(r)=1/(2r^2)-a/r$ is the dimensionless variational scaling used in Part I. Its minimum balances localization and electromagnetic attraction. No BFG spectrum or fundamental constants are derived by plotting it.

Stage connection: Section 30 / Figure 28. **Declared grid:** dimensionless radius from 0.25 to 3, attraction strength from 0.1 to 1.5. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Molecular binding in a declared Morse model

Figure 61. Molecular binding in a declared Morse model
Conventional conditional molecular model




Figure 61: Conventional conditional molecular model. The assumed Morse potential has equilibrium $r=1$, inverse length $a=2$ and dissociation reference zero. A negative height denotes binding inside this chosen potential. Chemical constituents and parameters are supplied, not generated by BFG.

Stage connection: Section 31 / Figure 29. **Declared grid:** bond coordinate from 0.4 to 2.5, well depth from 0.2 to 2. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Cellular gross transport and net flux

Figure 62. Cellular gross transport and net flux
Exact flux accounting




Figure 62: Exact flux accounting. $J_{\text{net}} = J_{\text{out}} - J_{\text{in}}$. The zero diagonal permits nonzero turnover. Boundary, species and units must be declared for any cell experiment; equality of these gross flows is not a demonstration of autonomous life.

Stage connection: Section 31 / Figure 30. **Declared grid:** influx from 0 to 2, outflux from 0 to 2. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

World-self projection closure residual

Figure 63. World-self projection closure residual
Conditional functional countermodel




Figure 63: Conditional functional countermodel. Assume a projected successor depends on hidden h by $h \cdot (1+x)$. The plotted dependence shows failure of x -only closure whenever h is omitted. World and self variables need an explicit projection and intervention law; a product diagram is insufficient.

Stage connection: Section 32 / Figure 31. **Declared grid:** visible macrostate from 0 to 1, hidden dependence from 0 to 1. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Report probability under an explicit toy model

Figure 64. Report probability under an explicit toy model

Illustrative report model, not phenomenality




Figure 64: Illustrative report model, not phenomenality. This declared Bernoulli report model is a placeholder with supplied coefficients. It makes a plot possible without inventing a numerical measure of experience. No link to phenomenal consciousness is asserted, and no actual report data are analyzed.

Stage connection: Section 33 / Figure 32. **Declared grid:** functional variable x from 0 to 1, functional variable y from 0 to 1. All surface inputs and output arrays are saved with this figure.

Evidence level: This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Weighted channel equality residual

Figure 65. Weighted channel equality residual




Figure 65: Computed ambient kernel. Every grid point computes $\alpha \cdot \text{Lambda_C}$ minus $\beta \cdot \text{Lambda_B}$ after the full endogenous loading stage. Residuals are floating-point rounding of the exact identity. Raw channel symmetry remains a distinct quantity.

Stage connection: Part II / exact weighted balance. **Declared grid:** y1 from 0.1 to 0.9, y2 from 0.1 to 0.9. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Formation inheritance-loss-seed accounting

Figure 66. Formation inheritance-loss-seed accounting




Figure 66: Exact accounting surface. $\Delta M_F = s_F - \text{loss}_F$ is sampled over declared independent nonnegative event outputs. The graph illustrates the exact budget; not every pair is claimed to arise from one autonomous BFG orbit.

Stage connection: Part II / formation balance. **Declared grid:** selected formation loss from 0 to 2, seed stock from 0 to 2. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Spectral diversity versus stock distribution

Figure 67. Spectral diversity versus stock distribution

Exact conditional two-mode spectrum




Figure 67: Exact conditional two-mode spectrum. D_F peaks when the two nonzero stocks agree and falls when one dominates. Increasing total formation does not monotonically increase diversity. This entropy is not semantic or phenomenal information.

Stage connection: Part II / spectral diversity. **Declared grid:** inherited stock from 0.05 to 2, orthogonal seed stock from 0.05 to 2. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Finite autonomous seed capacity

Figure 68. Finite autonomous seed capacity
Theorem-bound illustration




Figure 68: Theorem-bound illustration. The bound $\text{reserve} = n_1 - p_1$ decreases by at least one on each further seed event. The plotted nonnegative bound is bookkeeping, not a claim of an orbit realizing every staircase. Rank loss can shorten the actual capacity.

Stage connection: Part II / finite seed theorem. **Declared grid:** remaining dimension reserve from 0 to 8, seed-event count from 0 to 8. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Constant-input positive geometric regime

Figure 69. Constant-input positive geometric regime

Computed declared input-coupled scalar model




Figure 69: Computed declared input-coupled scalar model. For $d=0.15$, repeated events converge to the positive fixed point approximately 0.0559622. The height displays exact map iterates with integer counts. The contraction theorem concerns geometry; neutral K and formation persist.

Stage connection: Part II / driven scalar theorem. **Declared grid:** initial y from 0 to 0.8, event count from 0 to 20. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Driven fixed point and formation neutrality

Figure 70. Driven fixed point and formation neutrality

Computed declared input-coupled scalar model




Figure 70: Computed declared input-coupled scalar model. The fixed point is computed by 80 scalar events for each d. Independence of the F coordinate displays neutral formation directions. The input is an additional model commitment, not an independently derived cosmological source.

Stage connection: Part II / driven scalar theorem. **Declared grid:** constant input d from 0.02 to 0.7, formation stock F from 0.2 to 2. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Self-similar hierarchy: level 1

Figure 71. Self-similar hierarchy: level 1
Exact declared iterated-function-system hierarchy




Figure 71: Exact declared iterated-function-system hierarchy. Level 1 contains 4 labeled descendants generated from the preceding level by four half-scale tetrahedral similarity maps. The associated uniform tensor-family state has $n=4^1$, per-copy formation $F_0/4^1$, total formation F_0 and entropy $1 \cdot \ln(4)$. With $y_0=0.5$ and declared input $d=0.15$, its common geometric load after 1 level transitions is 0.2181909. Spatial positions are an added illustrative embedding, not derived spacetime or a biological transition. Point color indicates geometric height, not measured physical energy.

Stage connection: Part II / typed self-similar hierarchy. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Self-similar hierarchy: level 2

Figure 72. Self-similar hierarchy: level 2
Exact declared iterated-function-system hierarchy




Figure 72: Exact declared iterated-function-system hierarchy. Level 2 contains 16 labeled descendants generated from the preceding level by four half-scale tetrahedral similarity maps. The associated uniform tensor-family state has $n=4^2$, per-copy formation $F_0/4^2$, total formation F_0 and entropy $2 \cdot \ln(4)$. With $y_0=0.5$ and declared input $d=0.15$, its common geometric load after 2 level transitions is 0.1275472. Spatial positions are an added illustrative embedding, not derived spacetime or a biological transition. Point color indicates geometric height, not measured physical energy.

Stage connection: Part II / typed self-similar hierarchy. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Self-similar hierarchy: level 3

Figure 73. Self-similar hierarchy: level 3
Exact declared iterated-function-system hierarchy




Figure 73: Exact declared iterated-function-system hierarchy. Level 3 contains 64 labeled descendants generated from the preceding level by four half-scale tetrahedral similarity maps. The associated uniform tensor-family state has $n=4^3$, per-copy formation $F_0/4^3$, total formation F_0 and entropy $3 \cdot \ln(4)$. With $y_0=0.5$ and declared input $d=0.15$, its common geometric load after 3 level transitions is 0.0876437. Spatial positions are an added illustrative embedding, not derived spacetime or a biological transition. Point color indicates geometric height, not measured physical energy.

Stage connection: Part II / typed self-similar hierarchy. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Self-similar hierarchy: level 4

Figure 74. Self-similar hierarchy: level 4
Exact declared iterated-function-system hierarchy




Figure 74: Exact declared iterated-function-system hierarchy. Level 4 contains 256 labeled descendants generated from the preceding level by four half-scale tetrahedral similarity maps. The associated uniform tensor-family state has $n=4^4$, per-copy formation $F_0/4^4$, total formation F_0 and entropy $4 \cdot \ln(4)$. With $y_0=0.5$ and declared input $d=0.15$, its common geometric load after 4 level transitions is 0.0697963. Spatial positions are an added illustrative embedding, not derived spacetime or a biological transition. Point color indicates geometric height, not measured physical energy.

Stage connection: Part II / typed self-similar hierarchy. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Self-similar hierarchy: level 5

Figure 75. Self-similar hierarchy: level 5
Exact declared iterated-function-system hierarchy




Figure 75: Exact declared iterated-function-system hierarchy. Level 5 contains 1024 labeled descendants generated from the preceding level by four half-scale tetrahedral similarity maps. The associated uniform tensor-family state has $n=4^5$, per-copy formation $F_0/4^5$, total formation F_0 and entropy $5 \cdot \ln(4)$. With $y_0=0.5$ and declared input $d=0.15$, its common geometric load after 5 level transitions is 0.0619450. Spatial positions are an added illustrative embedding, not derived spacetime or a biological transition. Point color indicates geometric height, not measured physical energy.

Stage connection: Part II / typed self-similar hierarchy. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Self-similar hierarchy: level 6

Figure 76. Self-similar hierarchy: level 6
Exact declared iterated-function-system hierarchy




Figure 76: Exact declared iterated-function-system hierarchy. Level 6 contains 4096 labeled descendants generated from the preceding level by four half-scale tetrahedral similarity maps. The associated uniform tensor-family state has $n=4^6$, per-copy formation $F_0/4^6$, total formation F_0 and entropy $6 \cdot \ln(4)$. With $y_0=0.5$ and declared input $d=0.15$, its common geometric load after 6 level transitions is 0.0585366. Spatial positions are an added illustrative embedding, not derived spacetime or a biological transition. Point color indicates geometric height, not measured physical energy.

Stage connection: Part II / typed self-similar hierarchy. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Hierarchy entropy with explicit replication

Figure 77. Hierarchy entropy with explicit replication

Exact declared tensor-family spectrum




Figure 77: Exact declared tensor-family spectrum. With uniform auxiliary state, each lift adds $\ln(m)$ to formation mode entropy while preserving its total trace. Levels and multiplicities are integers. This increase counts declared copy labels and cannot by itself establish emergent complexity.

Stage connection: Part II / self-similar lift. **Declared grid:** level number from 0 to 8, branching multiplicity from 2 to 6. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

Macromolecular formation under logistic kinetics

Figure 78. Macromolecular formation under logistic kinetics
Conditional biochemical toy dynamics




Figure 78: Conditional biochemical toy dynamics. The declared law $dz/dt=r*z*(1-z)$, $z(0)=0.1$, has the displayed exact solution. It provides a stage-specific dynamical illustration, not an origin-of-DNA mechanism or a BFG derivation of chemical reaction rates.

Stage connection: Book / DNA and macromolecular stage; Part I Section 31. **Declared grid:** model time from 0 to 10, assembly rate from 0.1 to 1.5. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Tissue pattern decay under diffusion

Figure 79. Tissue pattern decay under diffusion
Conditional continuum tissue model




Figure 79: Conditional continuum tissue model. A unit-wavenumber diffusion mode decays as $\exp(-D\pi^2 t)$. It illustrates the need for typed spatial dynamics; diffusion alone removes this mode and does not generate tissue organization or Turing patterns.

Stage connection: Book / multicellular stage; Part I Section 31. **Declared grid:** model time from 0 to 5, diffusivity from 0.05 to 1. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Organ regulation under a damped controller

Figure 80. Organ regulation under a damped controller

Conditional linear organ-control model




Figure 80: Conditional linear organ-control model. The supplied response is a decaying under-damped mode with unit natural frequency and the corresponding specified initial velocity. It models regulation only after a chosen input-output identification; no actual organ parameters are predicted.

Stage connection: Book / organ stage; Part I Sections 31 and 75. **Declared grid:** model time from 0 to 12, damping coefficient from 0.1 to 1.5. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Organism resources and sustained throughput

Figure 81. Organism resources and sustained throughput
Conditional open-system resource model




Figure 81: Conditional open-system resource model. $dq/dt=1-g*q$ with $q(0)=0.2$ has the shown stock. Positive stationary stock needs continual inflow; a stationary quantity can coexist with gross flux. A full organism requires many additional biological state variables and interventions.

Stage connection: Book / organism stage; Part I Sections 31 and 75. **Declared grid:** model time from 0 to 10, loss rate from 0.1 to 1.5. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Civilization growth in a declared carrying-capacity model

Figure 82. Civilization growth in a declared carrying-capacity model

Conditional social-system toy dynamics




Figure 82: Conditional social-system toy dynamics. The model $dN/dt=r*N*(1-N)$, $N(0)=0.05$, is deliberately supplied. It shows bounded growth rather than an inevitable historical trajectory. It does not infer societal values, knowledge or institutional stability from BFG.

Stage connection: Book / civilization stage. **Declared grid:** model time from 0 to 10, growth rate from 0.1 to 1.2. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Sport balance as a mechanical mode

Figure 83. Sport balance as a mechanical mode
Conditional mechanical control




Figure 83: Conditional mechanical control. This supplied damped oscillatory mode illustrates correction around a neutral position. An athlete-specific model needs measured mechanics, neural control and disturbances. A resemblance to balance is not an independent BFG prediction.

Stage connection: Book / sport stage. **Declared grid:** model time from 0 to 10, damping coefficient from 0.1 to 1. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Gravitational structure under a Jeans-scale assumption

Figure 84. Gravitational structure under a Jeans-scale assumption
Conventional conditional astrophysical model




Figure 84: Conventional conditional astrophysical model. In normalized units $G=1$, the conventional Jeans length is $c_s \sqrt{\pi/\rho}$. The chart uses an explicitly supplied continuum matter model and should not be interpreted as gravity emerging from the internal positive metric.

Stage connection: Book / star and gravitational structure stages. **Declared grid:** sound speed from 0.1 to 1.5, mass density from 0.1 to 2. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Black-hole horizon in a supplied Schwarzschild model

Figure 85. Black-hole horizon in a supplied Schwarzschild model

Conventional conditional gravity model




Figure 85: Conventional conditional gravity model. In normalized units $G=c=1$, the Schwarzschild factor is $1-2M/r$. The grid remains outside the maximum horizon radius. This is not an information-recycling simulation, a singularity resolution or a BFG derivation of a black hole.

Stage connection: Book / black-hole stage; Part I Sections 24 and 25. **Declared grid:** mass parameter from 0.1 to 1, radius from 2.1 to 5. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Cosmic acceleration under a supplied fluid law

Figure 86. Cosmic acceleration under a supplied fluid law

Conventional conditional cosmological model




Figure 86: Conventional conditional cosmological model. For a single assumed FLRW fluid in normalized $8\pi G=1$ units, $a_{\text{double_dot}}/a=-(\rho+3p)/6$ with $p=w\rho$. The chart displays the conditional acceleration criterion. Neither w nor dark energy is generated by BFG.

Stage connection: Book / dark-energy stage; Part I Section 25. **Declared grid:** equation of state w from -1.2 to 0.2, energy density from 0.1 to 2. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Dark-sector rotation in a supplied mass model

Figure 87. Dark-sector rotation in a supplied mass model

Conventional conditional dynamical model




Figure 87: Conventional conditional dynamical model. In $G=1$ units, circular speed $\sqrt{M(r)/r}$ assumes an enclosed mass profile. The independent mass and radius axes are inputs. This is a gravitational diagnostic, not a BFG dark-matter identification or an actual fitted galaxy.

Stage connection: Book / dark-matter stage; Part I Section 25. **Declared grid:** enclosed mass from 0.1 to 2, orbital radius from 0.2 to 3. All surface inputs and output arrays are saved with this figure. **Evidence level:** This panel uses an independently assumed application model. Its parameters and physical interpretation do not follow from the finite BFG kernel.

Nonlinear coarse-graining mismatch

Figure 88. Nonlinear coarse-graining mismatch

Exact scalar projection countercontrol




Figure 88: Exact scalar projection countercontrol. The mean of two nonlinear BFG scalar outputs generally differs from the output at their mean. This Jensen-type mismatch shows why a level-transition projection must be tested for closure; repeated level names do not establish autonomous macro-dynamics.

Stage connection: Part I Sections 22 and 75; Part II hierarchy. **Declared grid:** mean microscopic y from 0.1 to 0.8, heterogeneity amplitude from 0.0 to 0.09. All surface inputs and output arrays are saved with this figure. **Evidence level:** These values compute the declared finite operator model or mathematical control. Their interpretation is limited to its stated assumptions.

A Regular-stratum differential specification

This appendix retains the full differential machinery of the supplied mathematical model. Work on a fixed-rank stratum with spectrum separated from the selection threshold, positive first mass and both loads, fixed positive rank of A , and a fixed seed branch. Componentwise SVD derivatives require simple separated positive singular values; degenerate clusters require invariant-subspace treatment. Spectral-gap dependence is consistent with classical subspace perturbation theory [6]; that reference does not by itself bound the entire endogenous BFG derivative chain. On a nonempty complement require a simple negative minimum separated from zero and other eigenvalues. Input variations are Hermitian, $\text{tr } \delta \rho_W = 0$ and

$$P\delta PP = (\mathbf{1} - P)\delta P(\mathbf{1} - P) = 0, \quad (172)$$

with positivity and rank constraints restricting allowable tangents.

A.1 Selection and polar differential

For a fixed separating contour, $\delta Q = (2\pi i)^{-1} \oint (z\mathbf{1} - Y)^{-1}(\delta Y)(z\mathbf{1} - Y)^{-1}dz$, and $\delta X_{\text{sel}} = (\delta Q)P + Q\delta P$.

For a constant-rank curve $X = TH$, $H = (X^\dagger X)^{1/2}$, $S_- = T^\dagger T$, $S_+ = TT^\dagger$, put $E_X = \delta X$. The tangent satisfies $(\mathbf{1} - S_+)E_X(\mathbf{1} - S_-) = 0$. The supported skew-Hermitian $\Omega = S_- \Omega S_-$ solves

$$H\Omega + \Omega H = S_-(T^\dagger E_X - E_X^\dagger T)S_-. \quad (173)$$

Then

$$\delta T = T\Omega + (\mathbf{1} - S_+)E_X H^+ + TH^+ T^\dagger E_X (\mathbf{1} - S_-). \quad (174)$$

The support derivatives are

$$\delta S_+ = (\mathbf{1} - S_+)E_X H^+ T^\dagger + TH^+ E_X^\dagger (\mathbf{1} - S_+), \quad (175)$$

$$\delta S_- = H^+ T^\dagger E_X (\mathbf{1} - S_-) + (\mathbf{1} - S_-)E_X^\dagger TH^+. \quad (176)$$

A conservative operator-norm bound is $\|\delta T\|_2 \leq 3\|E_X\|_2/\sigma_{\min,+}(X)$. Loss of a uniform bound near rank loss does not imply every particular perturbation diverges. Standard polar theory provides the underlying matrix framework [4].

A.2 Transport, metric and analysis

Transport variations use the three-term product rule, for example

$$\delta \rho_F^{\text{sel}} = (\delta T_{\text{sel}})\rho_F T_{\text{sel}}^\dagger + T_{\text{sel}}(\delta \rho_F)T_{\text{sel}}^\dagger + T_{\text{sel}}\rho_F(\delta T_{\text{sel}})^\dagger. \quad (177)$$

The same applies to unnormalized witness. With $m = m_{\text{sel}}$,

$$\delta m = \text{tr}[(\delta S_{\text{sel}}^-)\rho_W + S_{\text{sel}}^- \delta \rho_W], \quad \delta \rho_W^{\text{sel}} = \delta \rho_W^{\text{un,sel}}/m - \rho_W^{\text{sel}} \delta m/m. \quad (178)$$

$\delta C = -C\delta Y C$, $\delta B = C\delta Y C$, and $\delta G = \delta Y$. Choose a horizontal support frame $\delta Z = (\delta P_{\text{sel}})Z$. Then

$$\delta D = (\delta Z)^\dagger G Z + Z^\dagger \delta G Z + Z^\dagger G \delta Z, \quad (179)$$

$$\delta(D^{-1}) = -D^{-1}(\delta D)D^{-1}. \quad (180)$$

Differentiate $P_G = ZD^{-1}Z^\dagger G$ by the product rule. For a channel $L \in \{C, B\}$ set $H_L = P_G L$ and

$$\begin{aligned} \delta \Lambda_L = & \text{tr}[(\delta G)H_L \rho_F^{\text{sel}} H_L^\dagger + G(\delta H_L)\rho_F^{\text{sel}} H_L^\dagger \\ & + G H_L(\delta \rho_F^{\text{sel}})H_L^\dagger + G H_L \rho_F^{\text{sel}}(\delta H_L)^\dagger]. \end{aligned} \quad (181)$$

For $s = \Lambda_C + \Lambda_B$,

$$\delta \alpha = \frac{\Lambda_C \delta \Lambda_B - \Lambda_B \delta \Lambda_C}{s^2}, \quad \delta \beta = -\delta \alpha. \quad (182)$$

The analysis blocks obey $\delta A_C = (\delta \alpha)/(\sqrt{2}\alpha)P_G C + \sqrt{\alpha}[(\delta P_G)C + P_G \delta C]$, with the analogous B, β formula. Thus all analysis responses depend on the five input variations; support motion is not an extra free parameter.

A.3 Complex reduced-SVD derivative

Set $F_A = \delta A$, $M = U^\dagger F_A V$, $\Omega_U = U^\dagger \delta U$, $\Omega_V = V^\dagger \delta V$. These connections are skew-Hermitian and satisfy

$$M = \Omega_U \Sigma + \delta \Sigma - \Sigma \Omega_V. \quad (183)$$

The diagonal equations are

$$\delta \sigma_i = \Re M_{ii}, \quad (\Omega_U)_{ii} - (\Omega_V)_{ii} = i \Im M_{ii} / \sigma_i. \quad (184)$$

Only the common phase is free; both diagonal connections cannot generally be set to zero. For $i \neq j$,

$$(\Omega_U)_{ij} = \frac{\sigma_j M_{ij} + \sigma_i \overline{M_{ji}}}{\sigma_j^2 - \sigma_i^2}, \quad (\Omega_V)_{ij} = \frac{\sigma_i M_{ij} + \sigma_j \overline{M_{ji}}}{\sigma_j^2 - \sigma_i^2}. \quad (185)$$

The normal components are

$$(\mathbf{1} - UU^\dagger) \delta U = (\mathbf{1} - UU^\dagger) F_A V \Sigma^{-1}, \quad (\mathbf{1} - VV^\dagger) \delta V = (\mathbf{1} - VV^\dagger) F_A^\dagger U \Sigma^{-1}. \quad (186)$$

Finally $\delta Y_+ = \delta \Sigma \Sigma + \Sigma \delta \Sigma$ and

$$\delta K_+ = (\delta U)^\dagger (K \oplus K) U + U^\dagger (\delta K \oplus \delta K) U + U^\dagger (K \oplus K) \delta U. \quad (187)$$

A.4 R4, inheritance and moving seed complement

$\delta X_4 = (\delta V)^\dagger P_{\text{sel}} + V^\dagger \delta P_{\text{sel}}$ determines the polar response. On this ambient stratum $m_4 = 1$ identically, so $\delta m_4 = 0$; transport and frame responses still contribute. Apply the three-term product rule to $\rho_{W+} = T_4 \rho_W^{\text{sel}} T_4^\dagger$ and inherited formation.

With $\delta E = -\delta S_4^+$, choose $\delta Z_E = (\delta E) Z_E$. Then

$$\delta H_E = Z_E^\dagger [(\delta E) K_+ + \delta K_+ + K_+ \delta E] Z_E. \quad (188)$$

For the simple seed minimum define $R_\lambda = (H_E - \lambda \mathbf{1})^+$, which vanishes on its eigenvector. The derivatives are

$$\delta \lambda = \text{tr}(\pi \delta H_E), \quad (189)$$

$$\delta \pi = -R_\lambda (\delta H_E) \pi - \pi (\delta H_E) R_\lambda, \quad (190)$$

$$\delta \Pi_{\text{seed}} = (\delta E) \Pi_{\text{seed}} + \Pi_{\text{seed}} \delta E + Z_E (\delta \pi) Z_E^\dagger, \quad (191)$$

$$\delta \rho_{\text{seed}} = -\kappa_F [(\delta \lambda) \Pi_{\text{seed}} + \lambda \delta \Pi_{\text{seed}}]. \quad (192)$$

Together with inheritance these close the derivative chain. At threshold, rank or seed-degeneracy crossings, these fixed-stratum formulas cannot be used unchanged.

B Reproducible numerical record

All 40 declared complex preparations passed a complete nonterminal event. 20 included an admissible rank-one seed. All five terminal codes were exercised, including a replicated seed preparation that produced a multiple negative complement minimum. A successful two-dimensional seed trajectory reached full persistence and then continued for three seed-free events.

Quantity	Result
maximum R4 support residual	2.1678e-15
maximum second mass residual	2.5535e-15
maximum weighted load residual	1.7764e-15
maximum formation balance residual	3.9911e-14
maximum quotient covariance residual	7.9096e-14
tensor replication residual	1.6317e-15
maximum successor Gram eigenvalue	0.24989
driven fixed point	0.055962
driven 60 step max error	3.4694e-17

The constant-input experiment used $d = 0.15$, 100 initial scalar values in $[0, 0.84]$ and 60 events. Tensor replication used multiplicity three and a uniform auxiliary state. Random seed: 20261008; absolute numerical rank and gate tolerance: 10^{-12} . No empirical data are used.

Declarations

This is a manuscript draft prepared from the author’s supplied works with AI-assisted mathematical synthesis, drafting and computational checking. It requires author review before submission. No empirical dataset, human-participant study or animal experiment is reported. Funding and competing-interest declarations must be supplied by the author for publication; none are inferred here. The synthetic implementation and numerical record accompany the draft. The universality hypothesis and physical bridge proposals are not presented as experimentally established.

Reference scope and numbering

References 1–20 reproduce the bibliography of the supplied preprint and retain its numbering, so the inherited citations in Part I and the citations to the same works in Part II agree. The original preprint itself is Reference 21; the philosophical source is Reference 22. References 23–27 supply additional background for the new discussion. Duplicate works are listed once in this consolidated bibliography. The former bibliography at the end of Part I is replaced by this single consolidated list; the supplied original PDF remains available in the reproducibility archive.

The three precursor manuscripts are retained as bibliographic provenance reported by the supplied preprint; their full texts were not supplied for this synthesis. Established matrix, information, physical and biological literature supports the specifically cited background, not empirical validation of BFG or of its proposed bridge maps. The fixed-point and tensor-lift arguments in Part II are stated and proved here under their explicit assumptions.

References

- [1] M. T. Wende and T. W. Loker, “A Constructive Level-0 Foundation of the Balance–Field Equation (BFG): A Pre-Mathematical Balance Substrate for Emergent Order, Information, Process Time and Physics,” independent research manuscript, 11 March 2026.
- [2] M. T. Wende and L. K. C. Leighton, “The Recursive Conjugate Closure Extension of the Balance–Field Equation (BFG): Holomorphic/Anti-Holomorphic Recursive Geometry, Lift-Invariant Closure, and Stable Conscious Regimes,” BFG manuscript, 16 May 2026.

- [3] M. T. Wende and A. Pereira Jr., “From No-Supervenience to Conscious Artificial Systems: A Level-0 Balance—Field Reconstruction of the Missing Mediating Structure Between Function, Information, and Experience,” independent research preprint, 2 June 2026.
- [4] N. J. Higham, “Computing the Polar Decomposition—with Applications,” *SIAM Journal on Scientific and Statistical Computing* 7(4), 1160–1174 (1986). doi:10.1137/0907079.
- [5] N. J. Higham, *Functions of Matrices: Theory and Computation*, SIAM, Philadelphia, 2008, Chapter 8.
- [6] C. Davis and W. M. Kahan, “The Rotation of Eigenvectors by a Perturbation. III,” *SIAM Journal on Numerical Analysis* 7(1), 1–46 (1970). doi:10.1137/0707001.
- [7] T. Kato, *Perturbation Theory for Linear Operators*, 2nd ed., Springer, 1995. doi:10.1007/978-3-642-66282-9.
- [8] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed., Cambridge University Press, 2013. doi:10.1017/CBO9781139020411.
- [9] J.-P. Aubin, *Viability Theory*, Birkhäuser, 2009. doi:10.1007/978-0-8176-4910-4.
- [10] C. E. Shannon, “A Mathematical Theory of Communication,” *Bell System Technical Journal* 27(3), 379–423 and 27(4), 623–656 (1948). doi:10.1002/j.1538-7305.1948.tb01338.x; doi:10.1002/j.1538-7305.1948.tb00917.x.
- [11] K. Kraus, *States, Effects, and Operations: Fundamental Notions of Quantum Theory*, *Lecture Notes in Physics* 190, Springer, 1983. doi:10.1007/3-540-12732-1.
- [12] V. Gorini, A. Kossakowski, and E. C. G. Sudarshan, “Completely Positive Dynamical Semigroups of N-Level Systems,” *Journal of Mathematical Physics* 17, 821–825 (1976). doi:10.1063/1.522979.
- [13] J. D. Bekenstein, “Relation between Physical and Gravitational Geometry,” *Physical Review D* 48, 3641–3647 (1993). doi:10.1103/PhysRevD.48.3641.
- [14] S. Navas et al. (Particle Data Group), *Review of Particle Physics*, *Physical Review D* 110, 030001 (2024), and 2025 update. doi:10.1103/PhysRevD.110.030001. <https://pdg.lbl.gov/2025/>.
- [15] A. M. Turing, “The Chemical Basis of Morphogenesis,” *Philosophical Transactions of the Royal Society B* 237, 37–72 (1952). doi:10.1098/rstb.1952.0012.
- [16] H. R. Maturana and F. J. Varela, *Autopoiesis and Cognition: The Realization of the Living*, D. Reidel/Springer, 1980. doi:10.1007/978-94-009-8947-4.
- [17] K. Friston, “The Free-Energy Principle: A Unified Brain Theory?” *Nature Reviews Neuroscience* 11, 127–138 (2010). doi:10.1038/nrn2787.
- [18] Bureau International des Poids et Mesures (BIPM), *The International System of Units (SI)*, 9th ed., updated 2026. doi:10.59161/AUEZ1291.
- [19] National Institute of Standards and Technology, Quantum Measurement Division. <https://www.nist.gov/pml/quantum-measurement>. Accessed 7 October 2026.
- [20] Particle Data Group, 2025 Summary Tables. <https://pdg.lbl.gov/2025/tables/contentstables.html>. Accessed 7 October 2026.

- [21] M. T. Wende. *The Balance Field Equation as a Recursive World Formula: A finite operator model with geometric self feedback and explicit physical bridge requirements*. Supplied preprint, 6 October 2026, 68 pp.
- [22] M. T. Wende. *Die kosmische Ordnung: Eine kurze philosophische Reise durch das Universum*. German illustrated edition, 2026. Supplied EPUB.
- [23] NASA. *Forces*. <https://science.nasa.gov/universe/overview/forces/>. Accessed 8 October 2026. Used for the distinction between electromagnetic, strong and gravitational interactions.
- [24] NASA. *What Can We Learn from the Universe's Baby Picture?* <https://science.nasa.gov/universe/stories/quick-reads/what-can-we-learn-from-the-universes-baby-picture/>. Accessed 8 October 2026. Used for cooling and recombination.
- [25] J. E. Hutchinson. Fractals and Self-Similarity. *Indiana University Mathematics Journal* 30(5), 713–747 (1981). doi:10.1512/iumj.1981.30.30055.
- [26] K. Falconer. *Fractal Geometry: Mathematical Foundations and Applications*. 3rd ed., Wiley, 2014, Chapters 2, 3 and 9. ISBN 978-1-119-94239-9. Publisher record: <https://www.wiley-vch.de/de/fachgebiete/mathematik-und-statistik/fractal-geometry-978-1-119-94239-9>.
- [27] S. Banach. Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales. *Fundamenta Mathematicae* 3, 133–181 (1922). doi:10.4064/fm-3-1-133-181. Publisher record: <https://doi.org/10.4064/fm-3-1-133-181>.