Closing Edge Loops at Quantization Time: Helix Constraints for T-Mesh Quantization

Research note — draft, not peer reviewed

openclare project

18 October 2026

Abstract

Automatic quad remeshers produce edge loops that wind round a part of the shape as helices instead of closing into rings, a defect that makes such meshes awkward to edit by hand. Bommes et al. [4] identified these *q-helices* and remove them after meshing, by grid-preserving operators applied to the finished quad mesh. We take the opposite route and prevent them where the integer structure of the mesh is decided: in the quantization of a T-mesh. A straight walk across the rectangles of a T-mesh that comes back to its start defines a *turn*; we show that the pitch of the corresponding helix is a linear form in the integer lengths of the arcs, so that closing the helix is a linear constraint on the quantization. We add such constraints to a minimum-deviation-flow quantizer [10] through a least-change projection and targeted repair walks, select among a few ways of finding turns by the share of spiral edges they leave, and guard against folds and budget drift. On our bench the share of edges on spiral loops at 3,000 quads drops from 16.7% to 6.0% on twelve models; on a human body remeshed at the face count of the hand-made base mesh it drops from 12.1% to 1.1%, below the 2.5% of the hand-made mesh. We relate turns to the twist of flat cylinders in the theory of Jenkins–Strebel differentials [24], show with the holonomy of the flat metric that a pair of singularities of valences three and five joined by a staircase of arcs between two turns forbids closing both, and prove that on a sphere the shift of a turn has the parity of its length, so that a helix of odd pitch closes only if its length changes. Exact computations on the lattice of closed quantizations of the human body show that whole lengths closing nearly every ring exist, but not at the size asked for. Tested on the T-mesh of that body, the propositions hold on every turn and annulus they apply to; the conflicts between two turns are not, as we had guessed, pairs on one trace, but staircases between a pair of valences three and five, exchanges of two vertices of valence three, or caps, and the one turn that cannot close on its own is held open by a separatrix that winds round a wrist. On seven other models the Burgers sum reproduces the shift of 3,600 of 3,604 rings, yet moving a singular vertex in the parametrization barely moves the shift: the map moves the other cone points instead. With Fox's free differential calculus we write any combination of shifts as a sum over the singular vertices of twisted winding numbers times Burgers vectors; winding numbers alone do not suffice, on the body as in principle, but they do modulo two, which gives a checkerboard rule for any combination of turns. Turned around, the obstructions bound from below the number of helices of any quantization: on the body, the separation of the singularities alone forces one to nine of the turns held for closing to stay helices, and this is attained, where the engine leaves 13 to 123. We list the open problems that a complete treatment would need to solve.

1 Introduction

Field-aligned quad remeshing has reached a level where element shape and alignment are rarely the limiting factor. What remains is structure: the way edge loops run across the surface. Artists expect a loop round an arm, a leg or a neck to be a ring. Automatic methods often

produce a helix instead: the loop goes round, comes back one or a few rows further, and winds along the whole part. The problem is old and well identified: Bommes et al. showed in 2011 that helical configurations are what keeps the base complex of an otherwise good quad mesh dense [4].

This note documents an approach developed in the open-source remesher openclare, and positions it against the literature we could check. Our contributions are:

1. *Turns* of a T-mesh and their *shift*: a straight walk across the rectangles of a T-mesh that returns to its starting side, and a linear form in the arc lengths whose integer value is the pitch of the helix the walk becomes in the quad mesh (Section 3).

2. Closing turns within T-mesh quantization: a least-change projection of the target lengths, an integer search, and repair walks judged on the turns, with safeguards on folds, distortion and budget (Section 4).

3. A classification of the edge loops of a quad mesh into closed, spiral, open and border loops, with spirals of order up to four (Section 5), and a bench that holds automatic remeshers against meshes modelled by hand (Section 6).

4. Obstructions to closing: the shift of a turn is the Burgers vector of the singularities on one side, composed in the order in which a walk round a cutting tree first meets them; a pair of valences three and five, or two vertices of equal valence exchanged between two turns, joined by a staircase of arcs, forces one of the turns to be a helix; a separatrix that winds once round and lands beside its own vertex does the same to a single turn; a checkerboard rule gives the parity of the shift between two turns; on a sphere the shift of every turn has the parity of its length; and a turn whose shift is a non-negative form in the lengths can never close (Section 8).

5. A test of these statements on the T-mesh of a human body, and a description of its conflicts between turns (Sections 8.2 and 8.3); a test of the Burgers sum on seven other models, and why moving singularities in the parametrization does not set the shifts (Section 8.5).

6. A formula for any combination of shifts, by Fox calculus, with twisted winding numbers in place of winding numbers, a checkerboard rule for any combination, and a proof that winding numbers alone do not determine the shifts (Section 8.4).

7. A lower bound on the number of helices any quantization leaves among a set of turns, with chains of dislocations as a corollary, measured against the engine (Section 10).

8. An exact look at whole lengths: the balances are the incidence matrix of a bidirected graph, whose free arcs parametrize the lattice of balanced lengths; whole lengths closing nearly all rings exist but cost too much deviation, and the turns, not the rounding, decide what closes (Section 9).

9. Machine-checked proofs, in Lean with Mathlib, of the algebraic cores of the propositions of Sections 8 to 10, with the topological facts as explicit hypotheses; they found a few gaps, corrected here, and show what holds for other rotation orders (Section 10.1).

The results are preliminary: one seed per model, one implementation, and heuristics where an exact formulation is still missing (Section 11). Statements are *proved* when they come with a proof, *measured* when they come from exact arithmetic or solver output on the systems openclare exports, and *conjectured* otherwise; we say which.

2 Background and related work

Counting singularities. On a closed surface of genus g, the valences $k(v)$ of a quad mesh satisfy $\sum_v (4-k(v)) = 4\chi = 8(1-g)$. With only valences 3 and 5, $n_3 - n_5 = 8(1-g)$: a sphere or a closed human body needs at least eight vertices of valence three. This bound says how many singular vertices are needed, not where they go; a hand-made human base mesh has hundreds.

Quad meshes as quartic differentials. Lei et al. [14] show that a closed quad mesh induces a Riemann surface and a meromorphic quartic differential $\omega = (dz)^4$ whose zeros and poles are the singular vertices, a vertex of valence k being of order $k-4$; conversely a quartic differential with finite trajectories induces a quad mesh. The divisor of singularities then satisfies the Abel–Jacobi condition, and a divisor satisfying it is the divisor of a meromorphic quartic differential. Zheng et al. [25] optimize the singularity configuration by integer programming to satisfy the condition. Two points matter here. First, the condition is stated with holomorphic one-forms, which a surface of genus zero does not have; our reading is that in genus zero it adds nothing to the degree condition (this is an inference, not a statement of the papers). Second, the paper lists as an open problem the guarantee that all trajectories are *finite*. A trajectory that is not finite is precisely an edge loop that does not close: the helices of this note.

Closed trajectories. A quadratic differential all of whose non-critical horizontal trajectories are closed is a Jenkins–Strebel differential [24]; the complement of its critical graph is a union of flat cylinders, each with a height and a *twist*, the shift with which the cylinder is glued back after a cut along its core curve. In a quad mesh, a ring of quads is such a cylinder; a helix is a cylinder whose gluing is shifted by a whole number of rows. Section 3 makes this precise at the level of a T-mesh.

Quad meshes as flat metrics. Declaring every quad a unit square turns a quad mesh into a flat metric with cone points at its irregular vertices. Chen et al. [8] characterize the metrics that arise this way by four conditions: Gauss–Bonnet, rotational holonomy in multiples of a quarter-turn, alignment with the border, and a *finite geodesic lamination*: the straight lines of the field are finite loops. On a surface of genus zero the holonomy condition follows from Gauss–Bonnet (their Lemma 3.9). The translational part of the holonomy is not discussed, and the finiteness condition is assumed, not constructed. Lei et al. [15] show that the singularity configurations of quad meshes are a strict subset of those of cross fields, and explain in this way why no quad mesh of the torus has exactly one vertex of valence three and one of valence five.

Dislocations. Izmestiev et al. [11] give geometric proofs of such non-existence results, among them the torus with one vertex of valence three and one of valence five, first proved combinatorially by Barnette, Jucovič and Trenkler in 1971 (as cited there). Their tool is the holonomy of the developing map of the flat metric, a rigid motion per loop. When the rotation of a loop is the identity, its translation is the *Burgers vector* of crystallography; two opposite cone points at distinct positions always have a non-zero one, which is the dislocation a pair of valences three and five makes in the square grid. Section 8 uses this tool for the shifts of turns.

Helices in quad meshes. Bommes et al. [4] define a *q-helix* as a dual parametric line of a quad mesh with an integer *pitch* h and *turn length* τ: walking τ steps along it is equivalent to h side-steps. They detect minimal q-helices ($h \leq \tau$, no sub-helix of smaller pitch) by the self-intersections of orthogonal dual lines, and remove them greedily with *grid-preserving operators*: closed dual paths of shift and collapse steps, found by a graph search, that keep the mesh all-quad and its singularities in place. Tangential smoothing then repairs the distortion the shifts introduce. Their experiments remove most helices (from 19–52 down to 1–23 per model) and

coarsen the base complex accordingly. They note that aligning pairs of singularities directly in the parametrization, as Myles et al. do for T-meshes [19], is ambiguous to automate. The approach works on a finished mesh; it does not touch the parametrization or its quantization.

T-mesh quantization. Integer-grid maps [3] and their quantization decide the integer structure of a quad mesh. Campen et al. quantize via the integer lengths of the arcs of a T-mesh traced in a seamless parametrization [7]. Lyon et al. solve a constrained version as an integer linear program, bounding the directional deviation of the separatrices of the resulting layout [18] (we could only read the abstract). Heistermann et al. cast quantization as a minimum-deviation flow on a bi-directed graph and give a fast approximate solver and an exact refinement [10]; openclare follows their formulation. To our knowledge none of these constrains the helices: the lengths are chosen for consistency, validity and closeness to the parametrization, and the loops close or spiral as a side effect.

Periods round a tube. QuadCover [12] lifts the cross field to a branched covering where it becomes a vector field, and integrates it there. The stripe parameterization of tubular surfaces [13], built on it, measures the jump ("gap") of the parameter across each cut path, which is constant along the path, and rounds it to the nearest whole number; the grid lines are continuous exactly when every gap is whole. The gap of the coordinate across a loop round a tube is what we call the shift of a turn: rounded to zero it gives rings, rounded to a non-zero whole number a helix of that pitch. That paper treats stripes spiralling round the tube as a legitimate outcome, and notes that rounding the gaps for least distortion is a hard combinatorial problem that it does not solve exactly.

Layouts traced from the field. Myles et al. [20] trace the field into a partition into quadrilateral patches and adjust the layout until its lengths are consistent; Pietroni et al. [21] select traced separatrices by a binary linear program; Razafindrazaka et al. [22, 23] obtain a layout as a minimum-weight perfect matching on a graph of the singularities. Campen surveys these methods [5]. They build coarse layouts; whether the dense mesh made from the layout spirals is, as far as the abstracts we read state, not among their criteria.

Lower bounds. We found no result that bounds from below the number of helices of a quad mesh, or the number of its loops that do not close, for a given singularity configuration. The nearest results are existence and non-existence theorems: Strebel's existence of Jenkins–Strebel differentials with prescribed cylinders on a fixed Riemann surface [24], the non-existence results above [11, 15], and the finiteness condition of [8], which the open problem of [14] asks to guarantee.

Loops by construction. Dual loops meshing [6] builds a layout from loops traced on the surface, which close by construction; it produces coarse layouts rather than dense field-aligned meshes.

Measuring loops. QuadGPT [17] counts *quad rings* (closed loops of faces) and *quad lines* (open strips) and uses the share of faces in rings and the mean length of lines as reward signals for preference optimization; it does not report them as evaluation metrics, nor does it tell spirals from open lines.

3 Turns of a T-mesh

Setting. A T-mesh is a partition of the surface into rectangles, the *patches*, whose sides are made of *arcs*; nodes where a trace stops on another are T-junctions, and the singular vertices of

the field are nodes. Each patch P has local coordinates with its first corner at the origin and its sides along the axes. A quantization assigns every arc a a length $\ell_a \in \mathbb{Z}_{\geq 0}$ such that opposite sides of every patch have equal length; the quad mesh is then read off as the integer grid of each patch.

Definition 1 (Turn)**.** *A* turn γ *is a straight walk across patches: it enters a patch through a side, leaves through the opposite side at the same distance from the corresponding corner, enters the neighbouring patch across the arc it reaches, and so on, until it comes back into its first patch through its first side, with the four quarter-turns of the crossings summing to zero.*

Crossing from patch P to patch Q across an arc composes a quarter-turn rotation of the local frame with a translation whose components are signed sums of arc lengths along the rims of P and Q: the position of a rim point is the sum of the lengths of the arcs between it and the first corner.

Proposition 1 (The shift of a turn is linear)**.** *For a turn γ there are integer coefficients c_a, independent of the lengths, such that after a full turn a point of the first side has moved across the walking direction by*

$$s_\gamma(\ell) = \sum_a c_a \, \ell_a.$$

For integer lengths, the dual line of the quad mesh that follows γ comes back $s_\gamma(\ell)$ rows beside itself: it closes into a ring if and only if $s_\gamma(\ell) = 0$, and is otherwise a helix of pitch $|s_\gamma(\ell)|$ in the sense of [4].

Sketch. Each crossing is a map $x \mapsto R_k x + t$ with R_k a rotation by k quarter-turns and t a vector whose entries are sums of $\pm\ell_a$. The composition along γ has total rotation zero, hence is a translation $T_\gamma(\ell)$, linear in ℓ with integer coefficients. Its component along the walking direction does not move a point of the first side off that side; its component across is $s_\gamma(\ell)$. The walk crosses no singularity (singularities are nodes, never inside a patch), so the quad mesh is a regular grid along it, and a dual line at distance o from the first corner comes back at distance $o + s_\gamma(\ell)$. □

The component of $T_\gamma(\ell)$ along the walking direction is the *length* $\alpha_\gamma(\ell)$ of the turn, the distance walked before coming back to the first side; it is also a linear form in ℓ with integer coefficients, and for whole lengths it is the number of quads of the strip in one turn.

In openclare (`quantization/tubes.rs`) the shift is built once per turn as an expression in the arc lengths, by composing the rim expressions of the crossed patches; evaluating it with the real lengths of the parametrization gives the *real shift*, the twist the field itself would give the cylinder.

Definition 2 (Strip)**.** *For whole lengths ℓ balancing every patch, the quad mesh is, in each patch, the integer grid of its sides. A* strip *is a maximal sequence of quads, each sharing with the next the edge opposite the one it shares with the previous.*

Proposition 2 (Completeness at given lengths)**.** *Let ℓ be whole lengths balancing every patch. Every strip of the quad mesh that crosses a side σ of a patch P at the quad edge $[k, k+1]$ of σ is, as long as it stays in the regular region, the walk of Definition 1 taken* with the lengths ℓ *from σ at offset $k + \frac{1}{2}$. In particular, walking from every half-integer of every side finds every strip of the quad mesh that comes back to the side it left; by Proposition 1 it is a helix exactly when $s_\gamma(\ell) \neq 0$.*

Proof. Inside P, the quads of the strip are the unit squares between the grid lines at k and $k+1$ across σ; their dual line is the straight line at $k + \frac{1}{2}$, which reaches the opposite side at the same distance from the corresponding corner, which is the move of the walk. At the arc where it leaves P, the crossing map to the neighbouring patch is a quarter-turn followed by a translation

whose components are sums of $\pm\ell_a$, hence integers: it maps the grid of P onto the grid of the neighbour, and the midpoint of a quad edge onto the midpoint of a quad edge. Every node lies at a whole distance along the sides, so a walk at a half-integer offset never meets a node, and the arc it crosses is determined. The strip and the walk thus continue together, patch after patch. A strip only meets singular vertices at its sides, never inside, since they are corners of patches; the regular region is where the grid coordinates of the patches glue by such maps. □

The sampled walks of Section 4 start at fixed shares of the real side lengths. Proposition 2 says where the helices of a given mesh are *all* found: at the half-integers of the whole lengths. This suggests closing by *cuts*, in the sense of cutting planes:

1. quantize; let ℓ be the whole lengths;
2. walk every strip at ℓ (Proposition 2); among the ways round that come back, select those meant to be rings;
3. add their shift expressions to the closing constraints, keeping those already closed, as long as the bounds of separation allow it;
4. solve again in whole numbers; repeat until no selected way is a helix.

Remark 1. *A turn round a part of the shape whose real shift is far from zero is not a rounding artefact: the field winds round the part. We only hold turns whose real shift is a small share of their length, the* lean*; a turn round a group of singularities, by contrast, may legitimately come back shifted.*

4 Closing turns in quantization

The quantizer of openclare minimizes, as in [10], a weighted deviation $\sum_a w_a\,|\ell_a - \lambda\, r_a|$ of the integer lengths from the scaled real lengths λr_a, subject to the balance of each patch, by cancelling improving walks in a strip network; the scale λ is searched for the number of quads asked for, and arcs that would merge two singularities are held open. Closing turns is a side constraint that is not a flow constraint. We handle it in three steps.

Finding turns. From points at given shares along each side of each patch, we walk straight across patches and keep the walks that return. A turn is *held* if its real shift is above a small threshold and its lean below a bound, and if its length is below a multiple of the side of the T-mesh (the side of a square of its area). Several *ways* are tried: starting shares $\{0.5\}$ or $\{0.2, 0.5, 0.8\}$, lean bounds 0.05 or 0.12, length bounds 2 or 6 sides.

Least-change projection. The target lengths are changed as little as possible, in the weighted least squares sense, for every held turn to have zero shift and every patch to stay balanced:

$$\min_x \sum_a \frac{(x_a - r_a)^2}{w_a} \quad \text{s.t.} \quad Ax = 0, \quad s_\gamma(x) = 0 \;\; \forall \gamma \in H,$$

solved by a sparse Cholesky factorization of $AW^{-1}A^\top$. If the system cannot be met, the held turn left with the largest shift is let go and the projection repeated.

Integer search and repair. The integer search of the quantizer runs on the projected targets. Rounding leaves some turns shifted by one or a few rows. Repair walks are then taken one at a time: a walk is guided by a bias along the arcs of the shifted turns, and accepted if it reduces the total shift, costs little in deviation, keeps the structure intact and the number of quads within tolerance. A second, *aimed* repair treats each held turn still shifted on its own: the bias is put on its arcs alone, and a walk is accepted if it shifts that turn less and the others no more.

Selection and safeguards. Each way gives a quantization. Its quad mesh connectivity is built and the share of edges on spiral loops measured (Section 5); the way with the fewest spirals is kept, provided its number of quads is within tolerance and its mean deviation from the real lengths at most 1.5 times that of the plain search. The ways are independent and run in parallel. After meshing, if the kept quantization makes a face fold over the surface, the plain quantization is meshed as well and the mesh with fewer folds kept.

5 Measuring loops

An edge loop is followed straight through every regular vertex (valence four, interior), each edge belonging to one loop. A loop is *closed* if it comes back to its start; *border* if it runs from border to border; *open* if it stops at an irregular vertex or a non-quad face; and *spiral*, whatever its ends, if a strip of quads leaving it across meets it again, the same way round, within $k_{\max} = 4$ quads. The last test catches helices of pitch up to four, including bundles of interleaved helices [4], while a ring is met again only round a whole torus or over a cap the other way round. We report the share of edges on each kind of loop, edges of borders left out.

6 Experiments

All numbers come from the openclare bench (repository `marcgardent/openclare`), one seed per model, measured on the output meshes against their sources. Models: twelve organic and mechanical models for development, a corpus of 31 models for the full bench, and two human bodies from Blender's Essentials library (CC0) whose base meshes, modelled by hand, serve as references.

Spirals on the twelve development models, 3,000 quads. Table 1 shows the successive steps. Fidelity (Hausdorff distance, per mil of the diagonal) and the mean corner deviation stay within about one per cent; the last step trades a little of both for a third fewer spiral edges.

	spiral edges	Hausdorff (‰)	corner deviation (deg)
no closing	16.7%	8.77	4.75
closing, one way kept by spirals	9.7%	8.57	4.85
+ leaning ways, aimed repair	9.1%	8.58	4.91
+ longer turns, deviation cap	6.0%	8.68	5.01

Table 1: Mean over twelve models; each line adds to the one above.

Against other remeshers. On the full bench at 3,000 quads, the mean share of spiral edges is 19.1% for Quad Remesher, 21.9% for QuadriFlow and 25.3% for AutoRemesher, against 12.3% for openclare with the first version of closing (and 17.3% without).

Against the hand-made mesh. The male body of the Essentials library has a base mesh of 10,590 quads modelled by hand, with 2.5% of spiral edges. Remeshed by openclare at that count before closing, it had 12.1% (one leg was a single helix from hip to ankle, its turns leaning more than the first lean bound allowed). With the method of Section 4: 10,467 quads, 1.1% spiral edges, 36.2% of loops closed against 24.7% for the hand-made mesh, a Hausdorff distance of 4.6‰ against 6.5‰, and a mean corner deviation of 4.1 degrees against 9.2.

The quad sphere. Not a result of closing but of the field. An icosphere remeshed at 3,000 quads had 32 irregular vertices (20 of valence three, 12 of valence five), the pattern of its own facets, because the field was pulled towards curvature directions that a sphere does not have; at 1,000 quads it had the eight poles but no structure (a base complex of 1,010 patches) and 87% of spiral edges. With directions not trusted where the two principal curvatures are within 10% of one another, both give the layout of a cube: eight poles of valence three, six patches, no spiral.

7 An exact look at the closing system

To see how far the projection of Section 4 is from what the constraints allow, openclare exports, for each way of finding turns, the system it works on: the real lengths of the 2,069 arcs of the T-mesh of the male body at 10,590 quads, the 1,552 balance equations of its patches, every turn found with its shift, and the lower bounds the integer search placed on arcs to keep singularities apart (30 to 36 arcs held at length one). We solve it with HiGHS through SciPy, in real numbers. The ways are numbered as exported: way $m + 2j + 4i$ starts from the shares $\{0.5\}$ ($m = 0$) or $\{0.2, 0.5, 0.8\}$ ($m = 1$), with the lean bound 0.05 ($j = 0$) or 0.12 ($j = 1$) and the length bound 2 ($i = 0$) or 6 ($i = 1$) sides. Way 2 is the one the engine keeps for this body.

Where the conflict lies (measured). All held turns can be closed together if the lengths are only asked to be non-negative. With the bounds of separation, they cannot. In four ways (0, 1, 2 and 4) one held turn cannot even be closed on its own: the same turn round the left wrist, of length 7.0, whose shift is a sign-definite form in two arc lengths (Section 8, Remark 4). Giving up the least total amount of the bounds needed to close all held turns, a linear program, shows that in ways 0, 2 and 4 a *single* arc must collapse, the arc of that wrist turn, 0.08 quad edges long; in way 6 a single other arc, at a foot; ways 1 and 3 need 9 and 11 arcs, round the eyes and the hands, and ways 5 and 7, which hold the most turns, 22 and 30, most of them at the feet. The conflict between closing loops and keeping singularities apart is thus real, but local.

How far the heuristic is. With every bound kept, the largest set of held turns that can be closed together is found by a mixed-integer program with one binary per turn (Table 2). The projection keeps far fewer: it removes, one at a time, the turn left most shifted by a least-squares projection whose negative lengths are clamped to zero, and a single clamped arc can make it drop dozens of turns that conflict with nothing.

way	held	kept by the projection	closable together	arcs to collapse
0	48	42	47	1
1	78	46	73	9
2	52	46	51	1
3	86	47	82	11
4	84	49	83	1
5	160	48	154	22
6	88	53	87	1
7	169	49	160	30

Table 2: Male body at 10,590 quads, real numbers. “Closable together”: the largest set of held turns closable with every bound of separation kept; “arcs to collapse”: the arcs whose bound must be given up to close all held turns, for the least total amount given up.

In whole numbers, the same mixed-integer program did not finish within fifteen minutes on

the two systems tried. Section 9 shows that a similar failure, on a larger set of rings, came from the way the problem was posed to the solver and not from an absence of whole solutions.

From reals to whole numbers. We then fed the engine better targets and measured what survives its integer search (Table 3). Aiming at the lengths of the linear program closes fewer loops in the end: the weighted L^1 objective puts the changes on few arcs, the deviation from the real lengths grows, and the variants with the fewest spirals are refused by the deviation cap. A projection that holds lengths at a floor instead of clamping them (an active-set method on the same Cholesky solver) keeps every held turn closed in real numbers; yet after the integer search 117 of 169 are shifted again, and the mesh spirals more. With the current projection, too, about 40% of the turns it kept are shifted after the integer search. The bottleneck is not the projection but the rounding: the integer search follows the targets but ignores the shifts.

targets of the integer search	turns closed in reals	still shifted after	spiral edges (mesh)
projection, clamped (current)	about 50	10–23 of them	1.1%
linear program, L^1	up to 160	—	11.9%
active set, floor 0	all held (up to 169)	117 of 169	3.3%
active set, floor half the real length	36–72	—	9.8%

Table 3: Male body at 10,590 quads, one run each; the mesh keeps the way with the fewest spirals, so its share is that of a single way.

8 When rings cannot all close

Section 7 found that closing all held turns conflicts with keeping singularities apart, and that the conflict comes down to one or a few arcs. This section gives a mechanism for such conflicts, with proofs. The tool is the holonomy of the flat metric, as used by Izmestiev et al. [11]; its ingredients are classical, and we know of no earlier statement of the propositions below in terms of helices. Section 8.2 tests them on the male body, and Section 8.3 describes its conflicts.

8.1 Turns and singularities

Flat metric and holonomy. In this section the surface Σ is closed, without border; a surface of genus zero is then a sphere. Several statements below fail on a sphere with borders (Remark 2). Lengths $\ell > 0$ balancing every patch make each patch a Euclidean rectangle. Glued along their arcs, whose lengths agree on both sides, the rectangles give the surface Σ a flat metric with cone points at the singular vertices: a vertex of valence k has angle $k\pi/2$ and curvature $(4-k)\pi/2$; a T-junction has angle 2π and is not a cone point. Let Σ° be Σ without its singular vertices, x_0 a point inside a patch, and R the quarter-turn in the positive sense of the orientation of Σ. Developing a loop γ of Σ° based at x_0 patch after patch, as in the proof of Proposition 1, gives a motion $\hat{h}(\gamma)\colon x \mapsto R^j x + t$ of the coordinates of the first patch, its *holonomy*: the chart φ of x_0, continued along γ, comes back as $\hat{h}(\gamma) \circ \varphi$. It depends only on the homotopy class of γ in Σ°, and $\hat{h}$ is a homomorphism of the fundamental group into the group of such motions [11], $\hat{h}(\alpha\beta) = \hat{h}(\alpha) \circ \hat{h}(\beta)$. A small loop round a singular vertex c of valence k, run with c on its right and reached along a path that develops it at the point p, has holonomy $x \mapsto R^{4-k}(x-p)+p$: turning clockwise round a cone point of angle $k\pi/2$ rotates the chart by $-k\pi/2$, which is $(4-k)\pi/2$ modulo a full turn. When the rotation of $\hat{h}(\gamma)$ is the identity, its translation is the *Burgers vector* of γ.

For a turn γ, let $\bar{\gamma}$ be the closed curve that follows the walk and, back in its first patch, returns to its starting point along a segment parallel to the first side, just inside the patch. Let e_1

be the walking direction and e_2 the direction along the first side. By the proof of Proposition 1, $\hat{h}(\bar{\gamma})$ is a translation and

$$s_\gamma(\ell) = \langle e_2,\ \hat{h}(\bar{\gamma}) \rangle.$$

Shifts, developed positions and Burgers vectors are linear forms in ℓ with integer coefficients, as long as the walks and paths cross the same patches. An identity between them that holds on an open set of balanced positive lengths therefore holds for all balanced lengths, and for whole lengths the developed positions of the nodes are points of $\mathbb{Z}^2$.

Lemma 1 (Gauss–Bonnet for turns)**.** *Let γ be a turn whose curve $\bar{\gamma}$ is simple. If $\bar{\gamma}$ bounds a disc $D \subset \Sigma$, the singular vertices inside D satisfy $\sum_{c \in D}(4 - k(c)) = 4$. If the curves of two turns are disjoint and bound an annulus A, then $\sum_{c \in A}(4 - k(c)) = 0$.*

Proof. For a region of a flat surface with cone points, bounded by straight segments, Gauss–Bonnet reads $\sum_c \kappa_c + \sum_i \theta_i = 2\pi\chi$, where κ_c are the curvatures of the cone points inside and θ_i the turning angles at the corners of the border. The curve $\bar{\gamma}$ is straight except at the two ends of its closing segment, where it turns by $+\pi/2$ and $-\pi/2$ (or not at all if the shift is zero); the turning angles sum to zero on each curve, seen from either side. With $\kappa_c = (4 - k(c))\pi/2$, $\chi(D) = 1$ and $\chi(A) = 0$ the two statements follow. □

On a surface of genus zero, such as a body, every simple closed curve bounds a disc on each side. Every turn with a simple curve thus splits the singular vertices into two groups of total index four each, consistently with the total $8(1 - g) = 8$ of Section 2: a fingertip beyond a ring holds the equivalent of four vertices of valence three. A loop round a group of singularities whose indices do not add up to four is never a turn, whatever the lengths. A side holding at most one singular vertex has total index 0 or $4 - k \neq 4$, so each side of a turn holds at least two.

Curves of turns need not be simple, and on the male body most of those held for closing are not (Section 8.2). The first case beyond simple curves is still determined.

Lemma 2 (One self-crossing)**.** *Let the surface have genus zero, and let the curve $\bar{\gamma}$ of a turn cross itself exactly once, transversally, with no singular vertex on it. It cuts the surface into three discs: two* lobes*, each bounded by one of the two loops of $\bar{\gamma}$, and a third region bounded by both. If the crossing lies on the walk, the totals of $4 - k$ over the singular vertices of the lobes and the third region are* 3*,* 3 *and* 2*; if it lies on the closing segment, they are* 2*,* 2 *and* 4*.*

Proof. A walk follows the two directions of the field, so it crosses itself at a right angle. Each loop leaves the crossing point X along one passage and comes back along the other, so its two rays at X are a quarter-turn apart. A loop is simple; it bounds on one side a disc containing the quadrant between its two rays, and the other loop, which meets it only at X, lies in the closure of the other side. That disc is a lobe, with a corner of angle $\pi/2$ at X; the two lobes take opposite quadrants, and the third region has the other two. The curve is a connected graph on the sphere with one vertex and two edges, so its three faces are discs. By Gauss–Bonnet as in Lemma 1, a disc whose border turns by the angles θ_i at its corners holds singular vertices of total index $4 - \sum_i \theta_i/(\pi/2)$. A corner of angle $\pi/2$ turns by $\pi/2$, which gives 3, 3 and 2 before the corners of the closing segment, at which the curve turns by $+\pi/2$ and $-\pi/2$ in succession. If the crossing lies on the walk, both of these corners are on one loop, every region sees both, and they cancel. Otherwise let the walk start at A along e_1 and come back at Z, and let it cross the closing segment between them. It crosses walking along $-e_1$: walking along e_1, it would enter the first patch through its first side there, and the turn would have ended. The loop through A then turns the same way at A as at X, and so does the loop through Z at Z and at X (a direct check of the four directions). Each lobe lies on the side of its corner of angle $\pi/2$ at X, so it sees a second convex corner and holds $4 - 2 = 2$; the third region holds the rest of the total 8, that is 4. □

Proposition 3 (The shift is set by the singularities)**.** *Let γ, γ_1, γ_2 be turns with simple curves.*

(a) Let $\bar\gamma$ bound a disc D holding the singular vertices $c_1,\dots,c_m$, and let x_0 be a point of $\bar\gamma$, the start of the walk for instance. Let $T\subset D$ be a tree whose vertices include $c_1,\dots,c_m$ and which meets $\bar\gamma$ only at x_0. Cut along T, the disc D becomes a disc Ω without cone point, which develops into the plane, with the frame (e_1,e_2) at x_0. Number the c_i in the order in which a walk round T from x_0, keeping T on its right, first reaches them, and let p_i be the developed position of c_i at that first visit. Then $s_\gamma(\ell)=\pm\langle e_2,B_D\rangle$, with

$$B_D=\sum_{i=1}^{m} Q_i\,(I-R_i)\,p_i,\qquad R_i=R^{4-k(c_i)},\qquad Q_i=R_mR_{m-1}\cdots R_{i+1},$$

the translation of $\rho_m\circ\cdots\circ\rho_1$, where $\rho_i(x)=R_i(x-p_i)+p_i$; the sign is $+$ if D lies on the right of $\bar\gamma$, and $-$ otherwise.

(b) Let $\bar\gamma_1$ and $\bar\gamma_2$ be disjoint and bound an annulus A, round which the two turns go the same way, and let a path κ inside A from $\bar\gamma_1$ to $\bar\gamma_2$ carry the walking direction of γ_2 onto that of γ_1 (the turns are parallel*); measure both shifts along the direction e_2 carried by κ. Then $s_{\gamma_1}(\ell)-s_{\gamma_2}(\ell)=\langle e_2,B_A\rangle$, where B_A is the Burgers vector of the border of the disc A cut along κ, given by the same sum over the singular vertices inside A, developed inside the cut disc.*

(c) In particular, parallel turns bounding an annulus without singular vertex have the same shift for every ℓ: a helix is a property of a whole cylinder of parallel turns, not of one of them.

Proof. (a) Orient each edge f of T away from x_0. Its two sides on the border of Ω are glued by a motion G_f: a point of f that develops at u on the side the walk round T follows first develops at $G_f(u)$ on the other. Let f end at the vertex v, from which the edges $f_1,\dots,f_r$ leave in the order of the walk. Keeping T on its right, the walk turns clockwise round v, so a small arc round v from the first side of f to the second turns clockwise through the whole angle at v and crosses $f_1,\dots,f_r$ on its way. Continuing the chart of Ω along this arc composes it with $G_{f_1}^{-1},\dots,G_{f_r}^{-1}$ at the crossings, and turning once round v composes it with the rotation $\rho_v(x)=R^{4-k(v)}(x-p_v)+p_v$, where p_v is the position of v at the first visit, the one the arc starts from. Comparing the two at the second side of f,

$$G_f=G_{f_r}\circ\cdots\circ G_{f_1}\circ\rho_v,$$

with ρ_v the identity if v is not singular. The same comparison at x_0, with the edges $f_1,\dots,f_r$ leaving x_0 and the arc round v replaced by $\bar\gamma$ run with D on its right, gives $\hat h(\bar\gamma)=G_{f_r}\circ\cdots\circ G_{f_1}$. Unfolding the recursion, $\hat h(\bar\gamma)$ is the composition of the ρ_{c_i}, each vertex before the vertices below it and the branches in the order of the walk: the order of first visit. Its rotation is $R^{\sum_i(4-k(c_i))}=R^4=I$ by Lemma 1, and expanding the composition gives B_D; run with D on its left, $\bar\gamma$ has the inverse holonomy, the translation by $-B_D$. The shift is the component along e_2. When T is a star of disjoint paths from x_0, the order is the one in which the paths leave x_0, as the walk meets them. (b) Take x_0 on $\bar\gamma_1$ at the start of κ. The loop $\lambda=\bar\gamma_1\,\kappa\,\bar\gamma_2^{-1}\kappa^{-1}$ is the border of the disc obtained by cutting A along κ. Developing along κ is a change of coordinates; conjugating the translation $\hat h(\bar\gamma_2)$ by it gives that translation read in the coordinates at x_0, where the walk of γ_2 runs along e_1 by hypothesis. Hence $\hat h(\lambda)$ is the translation by $\hat h(\bar\gamma_1)-\hat h(\bar\gamma_2)$, both read in the same coordinates, whose component along e_2 is the difference of the shifts, and part (a) applied to this disc, whose singular vertices have total index zero by Lemma 1, gives the sum. (c) Without singular vertex the sum is empty. □

The simplest configuration between two parallel turns is a pair of singular vertices of valences three and five, the defect that lets the number of edges round a limb change. It is a dislocation of the grid.

Proposition 4 (A dislocation between two turns)**.** *In the setting of Proposition 3(b), let the annulus A hold exactly two singular vertices, c_3 of valence three and c_5 of valence five, and let $d = p_5 - p_3 = d_1 e_1 + d_2 e_2$ be their separation, developed inside the cut disc. Then:*

(i) $s_{\gamma_1}(\ell) - s_{\gamma_2}(\ell) = \varepsilon_1 d_1 + \varepsilon_2 d_2$ *with signs* $\varepsilon_1, \varepsilon_2 \in \{+1, -1\}$ *that depend on the configuration but not on* ℓ*, and* $|B_A| = \sqrt{2}\,|d|$*. The two turns can close together only if* c_5 *lies, as seen from* c_3*, on one particular diagonal of the field.*

(ii) If c_3 *and* c_5 *lie on one straight line of the field inside* A*, of length* δ *(in a T-mesh: they are joined by a chain of arcs along one trace, and* δ *is the sum of their lengths), then* $|s_{\gamma_1}(\ell) - s_{\gamma_2}(\ell)| = \delta$*. The two turns can both be rings only if* $\delta = 0$*, that is if the two singularities merge; as long as the bounds of separation keep* $\delta \geq 1$*, one of them at least is a helix.*

(iii) For whole lengths, $s_{\gamma_1}(\ell) - s_{\gamma_2}(\ell) \equiv d_1 + d_2 \pmod 2$*: a pair at an odd number of edges from one another, counted along the two directions of the field, never lies between two rings.*

Proof. With $R_3 = R$ and $R_5 = R^{-1}$, the sum of Proposition 3 for the two possible orders is $B_A = R^{-1}(I - R)\,p_3 + (I - R^{-1})\,p_5 = (I - R^{-1})\,d$ or $B_A = R\,(I - R^{-1})\,p_5 + (I - R)\,p_3 = (I - R)(-d)$. In coordinates (e_1, e_2), $R(a, b) = (-b, a)$, so $(I - R^{-1})(d_1, d_2) = (d_1 - d_2,\ d_1 + d_2)$ and $(I - R)(-d_1, -d_2) = (-d_1 - d_2,\ d_1 - d_2)$; reversing the orientation of the border, which inverts the rotations, gives the two other sign patterns. (We checked the four cases symbolically.) In all of them both components of B_A are of the form $\pm d_1 \pm d_2$ and $|B_A|^2 = 2|d|^2$, which is (i); the component along e_2 vanishes exactly on the line $\varepsilon_1 d_1 + \varepsilon_2 d_2 = 0$, a diagonal. For (ii), choose κ away from the straight segment joining c_3 to c_5, which does not separate the two borders of A, and develop c_5 from c_3 along it: d is δ times one of $\pm e_1$, $\pm e_2$, so $|\varepsilon_1 d_1 + \varepsilon_2 d_2| = \delta$. For (iii), $d \in \mathbb{Z}^2$ and $\pm d_1 \pm d_2 \equiv d_1 + d_2$. □

Plain counting confirms (iii) independently. If both turns were rings, the along component of B_A would be the difference $n_1 - n_2$ of the numbers of edges round them, of the parity of $d_1 + d_2$; but an annulus made of quads has an even number of edges on its border, since $4F = 2E_{\text{inside}} + E_{\text{border}}$, so $n_1 + n_2$ is even. Proposition 6 below makes this a statement about each turn. Part (ii) is the mechanism the problem of feasibility in Section 11 guessed, and Proposition 7 extends it from straight chains to staircases. An earlier draft suggested that it explains the conflict found on the male body at a single arc of the left wrist (Section 7). It does not: that conflict involves a single turn, not two (Remark 4), and Proposition 9 gives its mechanism.

For more singular vertices between two turns, the parity part generalizes into a checkerboard rule.

Proposition 5 (Checkerboard rule)**.** *In the setting of Proposition 3(b), with whole lengths, colour the points* $p \in \mathbb{Z}^2$ *of the cut disc black or white by the parity of* $p_1 + p_2$ *(either colour may be called white). Then* $s_{\gamma_1}(\ell) - s_{\gamma_2}(\ell)$ *has the parity of the number of singular vertices of odd valence inside* A *that develop onto white points. In particular, two rings bound an annulus only if that number is even.*

Proof. In the sum of Proposition 3, a vertex of even valence has $R_i = I$ or $R_i = -I$, and $(I - R_i)p_i \in \{0, 2p_i\}$ is even. A vertex of odd valence has $R_i = R^{\pm 1}$, and both components of $(I - R^{\pm 1})(a, b) = (a \pm b,\ b \mp a)$ have the parity of $a + b$. Quarter-turns keep the property "both components have the parity σ", and sums add the σ. So both components of B_A have the parity of the number of odd vertices on white points. The count does not depend on the coordinates: a quarter-turn keeps colours, and a whole translation t changes the colour of every point if $t_1 + t_2$ is odd, which keeps the parity of the count because the number of odd vertices inside A is even, their indices $4 - k$ being odd and summing to zero with the even ones (Lemma 1); for the same

reason the two colours give counts of the same parity. Developing a vertex along another path inside the cut disc moves it by a rotation $x \mapsto R_j(x - p_j) + p_j$ about another singular vertex, which keeps its colour. We checked the rule on 13,000 random configurations with sympy. □

Parity on a sphere. The checkerboard rule compares two turns. On a surface of genus zero each turn carries its own parity condition, which needs neither a simple curve nor the holonomy. Let B be the matrix of the balances of the patches, so that the balanced lengths are the solutions of $B\ell = 0$.

Proposition 6 (Parity of a turn on a sphere)**.** *Let the surface have genus zero and the T-mesh admit lengths that are positive and balance every patch (its real lengths do). For every turn γ and all whole lengths ℓ balancing every patch, of any sign,*

$$\alpha_\gamma(\ell) \equiv s_\gamma(\ell) \pmod 2.$$

Proof. First let ℓ be whole, balanced, and positive on every arc. The patches are then grids of unit squares glued along their arcs, a quad mesh of the sphere. Its graph is bipartite: over $\mathbb{Z}/2$, the cycles of a graph embedded in a sphere are sums of face boundaries, since $H_1(S^2; \mathbb{Z}/2) = 0$, and every face boundary has four edges, so every cycle has an even length. Fix a colouring of the vertices in two colours, adjacent vertices differing. In each patch P the grid is connected, so the colour of its point of local coordinates (i, j) is $i + j + \kappa_P \bmod 2$ for a constant κ_P. For a motion $M(x) = R^k x + t$ with $t \in \mathbb{Z}^2$, write $\pi(M) = t_1 + t_2 \bmod 2$; since a quarter-turn keeps the parity of $i + j$, the sum of the coordinates of $M(x)$ has the parity of that of x plus $\pi(M)$.

Let $P_0, P_1, \dots, P_N = P_0$ be the patches crossed by γ and D_k the motion that develops the coordinates of P_k into those of P_0, as in the proof of Proposition 1; D_0 is the identity and D_N the translation $T_\gamma(\ell)$. We show $\kappa_{P_k} \equiv \kappa_{P_0} + \pi(D_k)$ by induction. A vertex w of the arc crossed from P_k to P_{k+1} has coordinates y in P_k and y' in P_{k+1} with $D_k(y) = D_{k+1}(y')$; its colour is $y_1 + y_2 + \kappa_{P_k} \equiv (D_k y)_1 + (D_k y)_2 + \kappa_{P_0}$ by the induction hypothesis, and also $y'_1 + y'_2 + \kappa_{P_{k+1}}$; as $(D_{k+1}y')_1 + (D_{k+1}y')_2 \equiv y'_1 + y'_2 + \pi(D_{k+1})$, this gives $\kappa_{P_{k+1}} \equiv \kappa_{P_0} + \pi(D_{k+1})$. For $k = N$ the patch is P_0 again, so $\pi(T_\gamma(\ell)) \equiv 0$. In the frame (e_1, e_2) of the first side, which is that of P_0 up to a quarter-turn and signs (the coordinates swap when the first side is along the first axis of P_0), $T_\gamma(\ell) = (\alpha_\gamma(\ell), \pm s_\gamma(\ell))$; a quarter-turn keeps the parity of the sum of the coordinates, hence $\alpha_\gamma(\ell) + s_\gamma(\ell)$ is even.

For general whole balanced ℓ: the balanced lengths form a rational subspace, positive real balanced lengths exist, and positivity is an open condition, so there are positive rational balanced lengths, and, multiplying by a common denominator, positive whole ones p. For N large, $\ell + Np$ is whole, balanced and positive, and $\alpha_\gamma + s_\gamma$, being linear, takes at $\ell = (\ell + Np) - Np$ the difference of two even values. □

The proof uses only that the quad graph is bipartite, so it fails in higher genus: on a torus, a 3×3 grid of quads has rings of three quads. On a sphere it gives at once:

Corollary 1. *On a surface of genus zero:*

(a) every ring of quads has an even number of quads;

(b) a helix of odd pitch winds round an odd number of quads per turn;

(c) closing a helix of odd pitch changes the length of its walk by an odd number of edges: a repair that keeps the length of a turn cannot close it if its pitch is odd, and changing the pitch by an odd amount always changes the length.

Proof. (a) By Proposition 2 a ring is the strip of a walk with $s = 0$, so α is even. (b) and (c) are the proposition read before and after the change. □

Part (a) also follows directly from the bipartite graph: either border of a ring is a cycle of the quad graph with as many edges as the ring has quads. The proposition is consistent with the earlier statements and refines them. For a turn whose curve bounds a disc D, the argument of the checkerboard rule applied to Proposition 3(a) shows that both components of $B_D = (\alpha_\gamma, s_\gamma)$ have the parity of the number of vertices of odd valence of D on white points: they are equal, which is Proposition 6 for that turn. For two parallel turns, the proposition applied to each gives $s_{\gamma_1} - s_{\gamma_2} \equiv \alpha_{\gamma_1} - \alpha_{\gamma_2}$, the two components of B_A, which the checkerboard rule and Proposition 4(iii) evaluate from the singularities: two rings ($s_{\gamma_1} = s_{\gamma_2} = 0$) have even lengths, so no 3–5 pair at odd separation, and no odd number of odd vertices on white points, lies between them. On the eight systems exported for the male body (Section 7), the proposition holds on all 1,838 ways round found by walking every strip of the meshes (measured).

Remark 2 (Borders; a second proof)**.** *The proposition and part (a) of the corollary need the surface to be closed: on a sphere with two border loops, an open cylinder of three quads round has a ring of three quads. The proof uses $H_1(S^2; \mathbb{Z}/2) = 0$, which borders break. On a closed sphere there is a second proof, which needs neither the bipartite graph nor positive lengths, and which the formalization of Section 10.1 found. Every loop of Σ° is a product of the loops λ_i round the singular vertices (Section 8.4), whose holonomies are quarter-turns about the points p_i. For whole lengths these are points of $\mathbb{Z}^2$, and a quarter-turn about a whole point keeps the parity of $p_1 + p_2$. So the holonomy of every loop keeps that parity, and its translation, the image of the origin, has an even sum of coordinates: for a turn, $\alpha_\gamma + s_\gamma$ is even.*

How many independent rings. Rings that do not cross one another are disjoint simple closed curves of Σ°, none bounding a disc with fewer than two singular vertices (Lemma 1). On a surface of genus zero with n singular vertices there are at most $n - 3$ such curves pairwise non-homotopic, the number of curves of a decomposition of the n-punctured sphere into pairs of pants; disjoint homotopic ones bound an annulus without singular vertex and share their shift by Proposition 3(c). The rings of a quad mesh that do not cross therefore fall into at most $n-3$ cylinders, the cylinders of the Jenkins–Strebel picture, and their closing conditions have rank at most $n - 3$.

Remark 3 (Whole periods are automatic; rings need zero)**.** *For whole lengths, every shift is a whole number: the period of the integer grid along a loop is always whole, which is what quantization guarantees and what [13] enforces by rounding. Closing a ring is the stronger requirement that this period be zero. The real shift of the parametrization is the twist of the field's cylinder; it tells which whole value rounding is likely to pick, not whether zero is reachable.*

Remark 4 (Sign-definite turns; Farkas certificates)**.** *Say that a linear form f in the lengths equals g* modulo the balances *if $f - g$ is a combination of the rows of B, so that $f(\ell) = g(\ell)$ for every balanced ℓ. If the shift of a turn equals, modulo the balances, a form $\sum_a c_a \ell_a$ with every $c_a \geq 0$, then for all balanced $\ell \geq 0$ the shift $s_\gamma(\ell)$ is never negative, and it is zero only if every arc with $c_a > 0$ has length zero. If one of those arcs is held open by a bound of separation, the turn is a helix under every quantization meeting the bounds; it should never be held, and one linear program per turn finds such turns. More generally, by Farkas' lemma (with $x = b + z$, $z \geq 0$), the system $B\ell = 0$, $s_\gamma(\ell) = 0$ for $\gamma \in H$, $\ell \geq b$ has no solution exactly when some combination $\sum_{\gamma \in H} u_\gamma s_\gamma$ equals, modulo the balances, a form $\sum_a c_a \ell_a$ with $c \geq 0$ and $\sum_a c_a b_a > 0$; and an arc a has length zero in every solution with $\ell \geq 0$ exactly when some such combination has $c_a > 0$. The data being integral, the multipliers u_γ can be taken whole.*

On the male body this is what happens at the left wrist (measured; the certificate checked in exact rational arithmetic for way 0): in ways 0, 1, 2 and 4 one held turn, of length 7.0, has a shift equal, modulo the balances, to $-(\ell_{1921} + \ell_{1937})$, where arc 1937, 0.08 edges long, is held at length at least one. That turn cannot be closed on its own. A sentence of an earlier

draft of Section 7 stated that every held turn can be closed on its own; it is false for this body. Section 8.3 shows where this form comes from.

Remark 5 (Whole solutions exist at some size)**.** *The balance and closing constraints are homogeneous and the bounds of separation are lower bounds $\ell \geq b$ with $b \geq 0$. If they have a real solution they have a rational one, a vertex of a rational polyhedron, and multiplying it by the common denominator N gives whole lengths $N\ell \geq Nb \geq b$ that satisfy them all. Closing in whole numbers is thus never impossible in itself; what can fail is closing* at a given size *(number of quads) or within a given deviation from the targets. Section 9 measures this on the male body.*

8.2 Measured on the male body

The statements above concern the T-mesh, which the export of Section 7 first lacked. We exported it with every turn, for the eight ways of the male body at 10,590 quads: its 1,295 nodes (1,034 T-junctions and 261 singular vertices, 133 of valence three, 127 of valence five and one of valence two, of total index 8) with their corner angles, its 2,069 arcs with their traces, its 776 patches, and for each turn the arcs it crosses. A turn is drawn as a curve by straight chords across the patches it crosses, and the regions its curve cuts the surface into are found from the chords. Identities between quantities of this section are identities between linear forms in the arc lengths modulo the balances, checked on a basis of the balanced lengths in floating point; parities are evaluated at the whole lengths the engine found. Counts are summed over the eight ways, a turn found in several ways counting once in each (measured throughout).

What holds. For each of the 1,299 turns, the crossing maps composed along the walk give a translation whose component across is the exported shift (Proposition 1). Of these turns, 718 have a simple curve; each splits the 261 singular vertices into two groups of total index 4 (Lemma 1). On each side of each of them, with a tree of cuts found by breadth-first search, the holonomy of the curve equals the composition of Proposition 3(a), in the order of first visit, in both components: 1,436 discs holding 4 to 257 singular vertices, no exception. The order matters: taking the vertices in the order of their last visit fails on 150 of the 158 discs of way 2. The checkerboard rule (Proposition 5) holds on the 1,436 discs, in the form given after Corollary 1, and on the 30,148 annuli between two disjoint simple turns, whose singular vertices have total index 0 in every case, as Lemma 1 says. The 294 pairs of disjoint turns with no singular vertex between them have equal shifts (Proposition 3(c)). Ten annuli hold exactly one vertex of valence three and one of valence five; on all ten the identity of Proposition 4(i) holds, and in none of them do the two vertices lie on one trace, so part (ii) never applies at the real lengths. The parity of Proposition 6 holds for the 1,299 turns.

Most held turns are not simple. Of the 1,299 turns, 101 cross themselves once, as in Lemma 2: 97 split the singular vertices as $(3, 3, 2)$, and 4 as $(2, 2, 4)$, all four crossing their closing segment. The other 480 cross themselves at least twice, up to 107 times. Of the 765 turns held for closing, only 356 are simple: long turns cross themselves many times, and the statements about simple curves and annuli say nothing about them. For such curves we use winding numbers: a combination $\sum_\gamma u_\gamma \bar{\gamma}$ of curves winds round each region of their union a whole number of times. This is its class in the homology of Σ°; it tells round which singular vertices the combination goes, but not its holonomy, which is not commutative.

8.3 Conflicts between turns

Conflicts are local, but not always pairs. An earlier draft conjectured that whenever a set of turns, each closable on its own, cannot be closed together within the bounds of separation, it contains two parallel turns whose annulus forbids it, as in Proposition 4. This is false. We

searched the held turns of the eight exported systems for *irreducible* infeasible sets: sets that cannot be closed together within the bounds while every proper subset can (Table 4). For each, the Farkas multipliers of Remark 4 were made rational and the identity checked in exact arithmetic, and for each turn left out, whole lengths closing the others within the bounds were found by HiGHS and checked in exact integer arithmetic (measured).

ways	size	turns	multipliers	bounded arcs	exact check
0, 1, 2, 4	1	42 (way 0)	-1	1937	way 0
1, 3, 5, 7	2	54, 61 (way 1)	$1, -1$	1587	—
1, 3	2	45, 49 (way 1)	$1, -1$	1069	—
1, 3, 7	2	26, 37 (way 1)	$-1, 1$	619	—
5, 7	2	58, 94 (way 5)	$1, -1$	858	—
5	2	49, 71	$-1, 1$	619, 625	—
6	2	57, 58	$1, 1$	1241	—
7	2	8, 15	$1, 1$	191	—
1	3	36, 37, 39	$1, 1, -1$	619	yes
1	4	40, 56, 59, 61	$1, 1, -1, -1$	1587	yes
1	4	66, 71, 75, 76	$-\frac{7}{6}, -\frac{5}{6}, -\frac{1}{3}, -\frac{1}{2}$	1967, 1980, 1998	yes
3	4	73, 78, 82, 83	$-\frac{7}{4}, -\frac{5}{4}, -\frac{1}{2}, -\frac{3}{4}$	1980, 1998	yes
5	7	59, 91, 96, 98, 99, 157, 158	$2, 4, 1, -4, -1, 1, -1$	1036	yes
7	7	112–116, 145, 147	$8, \frac{1}{2}, 1, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}$	1157	yes
5	15	0, 2–9, 11, 14, 16–18, 64	whole, up to 23	17	14 of 15

Table 4: Irreducible infeasible sets of held turns on the male body, bounds of separation kept. Turn numbers are those of the first way listed; "bounded arcs": arcs held at length at least one in the support of the non-negative form. Rows without an exact check come from the floating-point linear program; their turns are each closable alone, so a pair is irreducible. In the last row, the whole lengths closing the set without turn 3 were not found within the time limit.

The smallest counterexample has three turns, in way 1 near the right eye, of lengths 54.9, 52.8 and 108.1:

$$s_{36} + s_{37} - s_{39} = \ell_{604} + \ell_{619} + \ell_{620} + \ell_{637} \quad \text{modulo the balances,}$$

with $\ell_{619} \geq 1$ imposed, so no lengths meeting the bounds close all three, while every two of them close together in whole numbers. Sets of four and seven turns behave alike, with the same signature: a combination of the shifts equals a non-negative form containing a bounded arc. Every set found is infeasible in real numbers already: none of these obstructions is one of parity. In the conflicts between two turns, the sum or difference of the shifts equals a sum of two to eight arc lengths with coefficient one, one of them bounded; the two turns have nearly equal lengths (39.7 and 40.0, 52.4 and 52.8, 277.5 and 278.0, ...) and lie in clusters of singularities, at the eyes, the hands and the feet. An earlier draft read this as Proposition 4(ii): a vertex of valence three and one of valence five joined along one trace by a chain holding the bounded arc. The T-mesh refutes it: we examined 13 conflicts between two turns (every row of two turns in Table 4 except the pair 49, 71 of way 5), and in none of them are the turns separated by such a pair on one trace. What separates them is a staircase.

Proposition 7 (Staircases)**.** *Let γ_1, γ_2 be turns and c, c' singular vertices such that*

$$s_{\gamma_1}(\ell) - s_{\gamma_2}(\ell) = \varepsilon_1 d_1 + \varepsilon_2 d_2$$

for all balanced ℓ, with $\varepsilon_1, \varepsilon_2 \in \{+1, -1\}$ and $d = p_{c'} - p_c = d_1 e_1 + d_2 e_2$ the separation of c' from c developed along a path π from c to c' (Propositions 4(i) and 8 give such identities). Let σ be a path from c to c' along arcs $a_1, \ldots, a_r$, through no other singular vertex, and homotopic to π with its ends fixed in Σ without the other singular vertices. Developed along σ, each arc a_i runs along

a direction $v_i \in \{\pm e_1, \pm e_2\}$*. If* $\varepsilon_1 v_{i,1} + \varepsilon_2 v_{i,2} = 1$ *for every* i*, so that* σ *is a* staircase*, using two directions a quarter-turn apart and climbing across the diagonal on which the two turns could close, then*

$$s_{\gamma_1}(\ell) - s_{\gamma_2}(\ell) = \sum_{i=1}^{r} \ell_{a_i}$$

for all balanced ℓ*. Both turns are rings only if every arc of* σ *has length zero. If this sum equals, modulo the balances, a form with non-negative coefficients containing an arc held at length at least one, one of the two turns is a helix under every quantization meeting the bounds.*

Proof. Developing along σ adds, arc after arc, the vector $\ell_{a_i} v_i$: the nodes σ passes are T-junctions, where the flat metric has no cone point. As σ and π are homotopic with fixed ends away from the other cone points, they develop c' at the same place relative to c, so $d = \sum_i \ell_{a_i} v_i$ and $\varepsilon_1 d_1 + \varepsilon_2 d_2 = \sum_i \ell_{a_i}$, first for positive balanced lengths, then for all by linearity. If both shifts vanish, this sum of non-negative lengths is zero. □

The proof uses only $\varepsilon_1 v_{i,1} + \varepsilon_2 v_{i,2} = 1$ for each step: neither that the steps run along the axes nor that $\varepsilon_1, \varepsilon_2 = \pm 1$, so it applies to any path on whose steps the linear form of the shift difference is one.

Proposition 4(ii) is the case of a straight σ, along one direction. The second source of such identities is an exchange of two vertices of equal valence.

Proposition 8 (Exchanging two vertices)**.** *Let* γ_1*,* γ_2 *be turns, and* κ *a path from the start of* $\bar{\gamma}_1$ *to that of* $\bar{\gamma}_2$ *that carries the walking direction of* γ_2 *onto that of* γ_1*, as in Proposition 3(b); let* $\lambda = \bar{\gamma}_1 \, \kappa \, \bar{\gamma}_2^{-1} \kappa^{-1}$*. Suppose that* λ *is homotopic in* Σ° *to* $\lambda_c \lambda_{c'}^{-1}$*, where* λ_c *and* $\lambda_{c'}$ *are small loops run with* c*,* c' *on their right round two singular vertices of the same valence* k*, reached along paths that develop them at* p_c *and* $p_{c'}$*. Then*

$$s_{\gamma_1}(\ell) - s_{\gamma_2}(\ell) = \langle e_2, \ (I - R^{4-k})(p_c - p_{c'}) \rangle.$$

For valence three or five this is $\pm d_1 \pm d_2$ *with* $d = p_c - p_{c'}$*, as for a dislocation, and Proposition 7 applies.*

Proof. As in the proof of Proposition 3(b), $\hat{h}(\lambda)$ is the translation by $\hat{h}(\bar{\gamma}_1) - \hat{h}(\bar{\gamma}_2)$, read in the coordinates at the start of γ_1. With $j = 4 - k$,

$$\hat{h}(\lambda_c) \circ \hat{h}(\lambda_{c'})^{-1}(x) = R^j \big(R^{-j}(x - p_{c'}) + p_{c'} - p_c\big) + p_c = x + (I - R^j)(p_c - p_{c'}),$$

and the shifts are the components along e_2. For $j = \pm 1$ both components of $(I - R^j)d$ are of the form $\pm d_1 \pm d_2$, as in the proof of Proposition 4. □

Table 5 lists the six configurations the 13 conflicts between two turns fall into. Configuration A is a dislocation: the two turns are simple and disjoint, and the annulus between them holds one vertex of valence three and one of valence five, 0.270 and 0.720 edges apart across and along, developed in the cut annulus. The difference of the shifts is $d_1 - d_2$ up to sign, as Proposition 4(i) says, and the staircase of two arcs joining the pair has length 0.990, the difference of the real shifts -0.955 and -1.945 of the two turns, identically on balanced lengths; modulo the balances it is a non-negative form containing the held arc 619. At the whole lengths the engine found, the two vertices are one edge apart on one grid line, the case of Proposition 4(ii) with $\delta = 1$. In configurations B, C and D the two turns cross. The combination of their curves winds $+1$ round a region holding one vertex of valence three, -1 round another holding another such vertex, and round no other singular vertex: in homology, an exchange of two vertices of valence three. The combination of the shifts is the length of a staircase joining them, and the held arc lies on it (D) or follows from it by the balances (B, C). The homotopy that Proposition 8 assumes was not checked; the identity it leads to with Proposition 7 was. In configurations E

and F, the combination winds once round singular vertices of total index four, like a turn round a cap; a bounded search found no staircase, nor sum of two, equal to the combination, and we leave them undecided.

	conflict	curves	between the turns	staircase	mechanism
A	$s_{37} - s_{26}$, ways 1, 3, 7; right eye	disjoint, simple; 52.4 and 52.8	annulus: 78 (val. 3), 76 (val. 5)	2 arcs, 0.990; held arc 619 by the balances	dislocation
B	$s_{45} - s_{49}$, ways 1, 3; right hand	simple; 39.7 and 40.0; 2 crossings	lenses: 102, 103 (val. 3)	4 arcs, 1.731; held arc 1069 by the balances	exchange
C	$s_{58} - s_{94}$, ways 5, 7; right hand	2 self-crossings each; 278.0 and 277.5; 6 crossings	109, 118 (val. 3)	3 arcs, 2.203; held arc 858 by the balances	exchange
D	$s_{57} + s_{58}$, way 6; left foot	figure-eights; 236.4 and 236.9; 4 crossings	155, 156 (val. 3)	2 arcs, 1.452, one of them held	exchange
E	$s_{54} - s_{61}$, ways 1, 3, 5, 7; left eye	simple and figure-eight; 59.5 and 71.9; 2 crossings	11 vertices of index 3, and 207 (val. 3)	none found	cap of index 4, undecided
F	$s_8 + s_{15}$, way 7; right foot	2 and 1 self-crossings; 251.5 and 247.4; 4 crossings	14, 23, 24, and 15 (val. 3)	none found	cap of index 4, undecided

Table 5: The conflicts between two turns on the male body, by configuration (measured). Turn numbers are those of the first way listed, signs those of the exported shifts; "between the turns": the singular vertices round which the combination of the two curves winds, numbered as nodes of the T-mesh; "staircase": a staircase between them whose length equals the combination of the shifts modulo the balances, and where the held arc of the certificate lies. A: Proposition 4(i) with Proposition 7; B–D: Proposition 8 with Proposition 7.

A separatrix that winds. The one turn that cannot be closed on its own is of another kind.

Proposition 9 (A separatrix that winds)**.** *Let σ and τ be separatrices of a singular vertex v such that σ, followed from v, stops on τ at a point q at distance $\delta(\ell)$ from v along τ, a sum of arc lengths. Let λ be the loop that runs along σ from v to q and back along τ to v, pushed off v into the sector it turns through. If λ is freely homotopic in Σ° to the curve $\bar{\gamma}$ of a turn γ, the walking direction of γ going to that of σ, then $s_\gamma(\ell) = \pm\delta(\ell)$ for all balanced ℓ. The turn is a ring only if $\delta = 0$, that is if σ comes back onto its own vertex; if δ equals, modulo the balances, a non-negative form containing an arc held at length at least one, γ is a helix under every quantization meeting the bounds.*

Proof. Develop λ from a point of σ next to v, with v at the origin and σ along e_1. The separatrix σ is straight and stops on τ at a right angle, so the return along τ is a segment of length δ along $\pm e_2$, like the closing segment of a turn. Being homotopic to the curve of a turn, λ has a translation as holonomy; it maps v, developed at the start, onto v developed at the end, which lies at distance δ from q across σ. Its component across is thus $\pm\delta$. A free homotopy conjugates the holonomy by a motion; the conjugate of a translation is the translation by the rotated vector, and the rotation is the identity since it carries the walking direction of γ onto that of σ. □

At the left wrist the held turn (42 in way 0, length 6.99, real shift -0.305, whole shift -1) is simple, and its small side holds four vertices of valence three, 226, 243, 244 and 245,

of total index 4: a cap. Proposition 3(a) on that side gives the shift -0.305 from their four positions, but as a form in the lengths of the arcs inside the cap it has no sign. The sign comes from outside. The vertex 246, of valence five, lies beyond the cap; one of its separatrices, the trace 964, runs $0.066 + 6.928 = 6.994$ edges once round the wrist, parallel to the turn, and stops on another of its separatrices, the trace 966, at the T-junction 1099, at the distance $\delta = \ell_{1920} + \ell_{1921} = 0.075 + 0.230 = 0.305$ from 246. The shift of the turn is $-\delta$ modulo the balances, and ℓ_{1920} equals, modulo the balances, the length of arc 1937, which starts at the vertex 248 of valence five and is held at length at least one: whence the certificate $-(\ell_{1921} + \ell_{1937})$ of Remark 4, and $s \leq -1$ at every admissible whole length (measured). A ring would need the separatrix to come back onto its own vertex, which the bounds of separation forbid. We checked the identity, not the homotopy that Proposition 9 assumes.

Conflicts of more than two turns. The three turns at the right eye are a dislocation between three turns. The turns 36 and 37 are simple; 39 is a figure-eight; each two of them cross twice, and together they cut the surface into five regions. The cycle $\bar{\gamma}_{36} + \bar{\gamma}_{37} - \bar{\gamma}_{39}$ winds -1 round the one region bordered by all three curves, which holds the vertices 79 of valence three and 77 of valence five, and round no other singular vertex: in homology, 39 is the sum of 36 and 37 and a loop round the pair. The combination $s_{36} + s_{37} - s_{39}$ is the length of a staircase of three arcs from 79 to 77 that includes the held arc 619, $1.367 = 1.712 - 0.955 + 0.610$ at the real lengths (measured). The set of four turns at the eyes (40, 56, 59, 61 of way 1, 72 to 167 long, crossing themselves up to 17 times and each other up to 28 times) winds $+1$ round the region holding 198 (valence three) and 199 (valence five), -1 round 207 and $+1$ round 202 (both valence three), and round no other singular vertex: in homology, a dislocation and an exchange, the held arc 1587 starting at 199; a bounded search did not write the combination as a sum of two staircases.

The other sets are not described. At the left hand (66, 71, 75, 76 of way 1, the same curves as 73, 78, 82, 83 of way 3), 76 is a figure-eight round three vertices of valence three, 29.9 long, and the others are 96 to 140 long, crossing themselves 13 to 28 times and each other 10 to 46 times; relative to the rest of the body, the weighted cycle winds round 18 of the 22 regions, with windings from $-\frac{1}{2}$ to $\frac{4}{3}$ in the normalization of Table 4. The set of seven in way 5 contains the two turns of configuration B and winds ± 4 round the two vertices they exchange; its other turns, up to 398 long, go round parts of both hands. The set of seven in way 7, at the left foot, contains a turn of length 4.92 round a cap of four vertices of valence three, whose multiplier is 16 against ± 1 or 2 for the six others, once scaled to whole numbers.

The fractional multipliers are an artefact of normalization. The linear program that finds the certificate asks the bounded arcs to weigh one in total: at the left hand the form has three bounded arcs of coefficient $\frac{1}{3}$ in way 1 and two of coefficient $\frac{1}{2}$ in way 3. Scaled to whole numbers, both ways give the same primitive multipliers,

$$7\, s_{66} + 5\, s_{71} + 2\, s_{75} + 3\, s_{76} = -2F \quad \text{modulo the balances,}$$

with F a form of non-negative whole coefficients 1, 2 and 5 containing the bounded arcs (measured). With these weights the weighted cycle winds alike round the rest of the body (236 singular vertices) and round the region of ten singular vertices of total index -2. Section 8.4 explains part of these weights: why $-2F$ is even, and which dislocation carries the coefficient 5.

Conjecture 1 (Conflicts are round a cluster)**.** *On a surface of genus zero, let H be an irreducible set of turns that cannot be closed together within the bounds of separation. Then the curves of the turns of H are, up to the cylinders of Proposition 3(c), the $k \geq 1$ borders of one planar region round a cluster of singularities, and a combination of their shifts that certifies the conflict is given by the holonomy relation of these borders: a form with non-negative coefficients containing an arc held open.*

An earlier draft proposed Conjecture 1 on the evidence of the certificates alone. The T-mesh confirms it for one turn (the wrist, a cap, though the form comes from a separatrix outside the region) and for the annulus of configuration A. It holds in a modified form for the three turns at the right eye: the curves are not the borders of a pair of pants, 39 being a figure-eight, but their combination winds round one region holding one pair of valences three and five, and the certificate is the staircase of that pair. As stated, it is refuted by the conflicts made of crossing turns: configurations B to F, and the sets of four and seven turns, whose curves cross each other up to tens of times and bound no single planar region. What the confirmed cases share is local, and suggests a refined statement, which we leave as a conjecture.

Conjecture 2 (Conflicts are sums of local dislocations)**.** *On a surface of genus zero, let H be an irreducible set of turns that cannot be closed together within the bounds of separation. Then some combination $\sum_{\gamma\in H} u_\gamma s_\gamma$ with whole multipliers that certifies the conflict (Remark 4) equals, modulo the balances, a sum with non-negative whole weights of*

(i) lengths of staircases (Proposition 7) between a vertex of valence three and one of valence five, or between two vertices of equal valence exchanged as in Proposition 8, and

(ii) shifts of caps: sign-definite forms carried by curves that wind once round singular vertices of total index four, as the wrist turn and its separatrix (Proposition 9),

one term at least containing an arc held open.

It holds, with a single term and up to the homotopies we did not check, for the wrist, configurations A to D and the three turns at the right eye. The set of four at the eyes fits it in homology (a dislocation and an exchange), and the set of seven at the left foot gives most of its weight to a turn round a cap. Configurations E and F, the left hand and the set of seven of way 5 are not yet written in this form; they, and the set of fifteen turns of way 5, which we did not examine, are the cases that could refute it.

8.4 Any combination of turns

The conflicts above were described by the winding numbers of a weighted cycle $\sum_\gamma u_\gamma \bar\gamma$. Winding numbers are its class in the homology of Σ°, which is commutative, while the holonomy is not; so they cannot carry the whole holonomy, and Proposition 3 needed an order of first visit. The bookkeeping that keeps the order is the free differential calculus of Fox [9]. It turns Proposition 3 into a formula for any combination of turns.

Setting. Let Σ have genus zero and singular vertices $c_1, \dots, c_n$. Fix a point x_0 inside a patch, with the frame of that patch, and disjoint paths π_i from x_0 to c_i; let λ_i be the loop that follows π_i, goes round c_i with c_i on its right, and comes back. The fundamental group of Σ° is generated by the λ_i, with a single relation, the product of all of them in the order in which the paths leave x_0. Let p_i be the developed position of c_i at the end of π_i and $R_i = R^{4-k(c_i)}$, so that $\hat h(\lambda_i) = \rho_i$, $\rho_i(x) = R_i(x - p_i) + p_i$. For a loop γ at x_0 write $\hat h(\gamma)(x) = \rho(\gamma)x + \tau(\gamma)$; since $\hat h$ is a homomorphism, $\tau(\alpha\beta) = \tau(\alpha) + \rho(\alpha)\,\tau(\beta)$. Let F be the free group on letters $x_1, \dots, x_n$, mapped onto the fundamental group by $x_i \mapsto \lambda_i$, and $\partial/\partial x_i$ the Fox derivatives of its group ring: $\partial x_j/\partial x_i = \delta_{ij}$ and $\partial(uv)/\partial x_i = \partial u/\partial x_i + u\,\partial v/\partial x_i$. Extend ρ linearly to the group ring; its values are the integer matrices $a_0 I + a_1 R + a_2 R^2 + a_3 R^3$, and the augmentation ϵ maps such an element to $a_0 + a_1 + a_2 + a_3$, the sum of the coefficients of the group ring element.

Proposition 10 (Fox–Burgers formula)**.** *For every word w of F,*

$$\tau(w) = \sum_{i=1}^{n} \rho\Big(\frac{\partial w}{\partial x_i}\Big)(I - R_i)\, p_i.$$

Let $\gamma_1, \dots, \gamma_m$ be turns, κ_r a path from x_0 to the start of $\bar\gamma_r$, f_r the direction e_2 of γ_r carried back to x_0 along κ_r (one of $\pm e_1$, $\pm e_2$), and w_r a word for $\kappa_r\bar\gamma_r\kappa_r^{-1}$. For multipliers u_r define the twisted winding *of the weighted cycle round c_i,*

$$W_i = \sum_{r=1}^{m} u_r\, \rho\Big(\frac{\partial w_r}{\partial x_i}\Big)^{\top} f_r \in \mathbb{Z}^2 \ \textit{(for whole } u_r\textit{)}.$$

Then, for all balanced lengths,

$$\sum_{r=1}^{m} u_r\, s_{\gamma_r}(\ell) = \sum_{i=1}^{n} \langle W_i,\ (I - R_i)\, p_i\rangle, \qquad \sum_{i=1}^{n} (I - R_i)^{\top} W_i = 0,$$

so the sum does not change when all the p_i are moved by one translation: it is a combination of separations $p_i - p_j$. The ordinary winding number of $\sum_r u_r\bar\gamma_r$ round c_i is, up to a constant common to all i, $w_i = \sum_r u_r\, \epsilon(\partial w_r/\partial x_i)$. The words w_r are defined up to the relation, and the W_i accordingly: inserting a conjugate $g\varrho g^{-1}$ of the relation ϱ in w_r, after a prefix a of w_r, adds $u_r\rho(ag\,\partial\varrho/\partial x_i)^{\top} f_r$ to every W_i (with a empty, $u_r\rho(g\,\partial\varrho/\partial x_i)^{\top} f_r$), and leaves the sum unchanged.

Proof. The map $w \mapsto \tau(w)$ and the right-hand side are both maps $F \to \mathbb{R}^2$ with $\phi(uv) = \phi(u) + \rho(u)\phi(v)$: the first by the rule above, the second by the rule of the Fox derivatives and the multiplicativity of ρ. They agree on the letters, where $\partial x_j/\partial x_i = \delta_{ij}$ and $\tau(\lambda_i) = (I - R_i)p_i$, and on the empty word, so they agree everywhere by induction on the length of words (for x_i^{-1}, use $0 = \phi(x_i x_i^{-1})$). For a turn, $\hat h(\kappa_r\bar\gamma_r\kappa_r^{-1})$ is the conjugate of the translation $\hat h(\bar\gamma_r)$ by $\hat h(\kappa_r)$, the translation by M_rT_r, where T_r is the translation of $\bar\gamma_r$ in its own frame and M_r the rotation of $\hat h(\kappa_r)$. Hence $s_{\gamma_r} = \langle e_2, T_r\rangle = \langle M_re_2, M_rT_r\rangle = \langle f_r, \tau(w_r)\rangle$, and $\langle f, \rho(D)\, b\rangle = \langle \rho(D)^{\top} f, b\rangle$ gives the formula. The fundamental formula of Fox calculus, $\sum_i (\partial w/\partial x_i)(x_i - 1) = w - 1$, maps under ρ to $\sum_i \rho(\partial w_r/\partial x_i)(R_i - I) = \rho(w_r) - I = 0$, the rotation of a turn being trivial; transposed and summed with the u_rf_r, it is the second identity. Finally $\epsilon(\partial w/\partial x_i)$ is the exponent sum of x_i in w, which is the class of w in the homology of Σ° written on the basis of the λ_i, defined up to the relation, in which each λ_i appears once: up to a common constant, the winding numbers. □

For a simple turn the word is a product of conjugates of the x_i, each once, and the formula is Proposition 3(a), with $\rho(\partial w/\partial x_i) = Q_i$. In general the twisted winding W_i is an integer vector, not a number: the cycle may go round c_i several times, each time with another rotation before it. Reduced modulo 2, the rotations disappear.

Corollary 2 (Checkerboard rule for any combination)**.** *On a surface of genus zero, let ℓ be whole and balanced and the multipliers u_r whole. Colour each singular vertex c_i by the parity χ_i of $p_{i,1} + p_{i,2}$; it does not depend on the paths π_i, and for positive lengths it is the colour of c_i in the bipartition of the quad mesh. Then*

$$\sum_r u_r\, s_{\gamma_r}(\ell) \equiv \sum_r u_r\, \alpha_{\gamma_r}(\ell) \equiv \sum_{i:\ k(c_i)\ \textit{odd}} w_i\, \chi_i \pmod 2,$$

with w_i the winding numbers of $\sum_r u_r\bar\gamma_r$, whatever the common constant and whatever the orientation of each $\bar\gamma_r$. In particular, a combination whose cycle winds an even number of times round every vertex of odd valence takes even values at every whole length.

Proof. Write $\rho(D) = \sum_k a_k R^k$, with $\sum_k a_k = \epsilon(D)$. Since $(R - I)^2 = -2R$, $R^k(I - R^{\pm 1}) \equiv I - R^{\pm 1}$ modulo 2, so $\rho(D)(I - R_i) \equiv \epsilon(D)(I - R_i)$ for $k(c_i)$ odd, and $\rho(D)(I - R_i) \equiv 0$ for $k(c_i)$ even, where $I - R_i \in \{0, 2I\}$. For $f \in \{\pm e_1, \pm e_2\}$ and $p \in \mathbb{Z}^2$, $\langle f, (I - R^{\pm 1})p\rangle \equiv p_1 + p_2$. This gives the value modulo 2 of the sum of Proposition 10, and the same computation with the direction e_1 of each turn gives that of $\sum_r u_r\alpha_{\gamma_r}$. A rotation about a singular vertex at a whole point, or a quarter-turn, keeps the parity of $p_1 + p_2$, so χ_i does not depend on the paths;

a whole translation of the origin flips all colours, which adds $\sum_{k(c_i)\text{ odd}} w_i$, even because the rotation of the cycle is trivial ($\sum_i w_i(4-k(c_i)) \equiv 0 \pmod 4$). Adding a constant q to all w_i adds $q\sum_{k(c_i)\text{ odd}} \chi_i$, which is even: it is the value modulo 2 of the relation, a loop with zero translation. Reversing a turn changes the sign of its winding numbers, which does not matter modulo 2. For positive whole lengths the developed parity is a proper colouring of the quad graph, as in the proof of Proposition 6. □

The corollary contains the checkerboard rule of Proposition 5 (winding one inside the annulus, zero outside), Proposition 4(iii), and, on a sphere, Proposition 6: for a single turn s and α have the same parity because both are congruent to the same sum.

Proposition 11 (Winding numbers do not suffice)**.** *Let c_a, c_b be singular vertices of odd valence developed at distinct points $p_a \neq p_b$. The loop $\lambda_a\lambda_b\lambda_a^{-1}\lambda_b^{-1}$ winds zero times round every singular vertex, and its holonomy is the translation by $(I-R_a)(I-R_b)(p_a-p_b)$, of length $2|p_a-p_b| \neq 0$. No function of the winding numbers gives the holonomy of the loops with trivial rotation.*

Proof. For $w = x_a x_b x_a^{-1} x_b^{-1}$, $\partial w/\partial x_a = 1 - x_a x_b x_a^{-1}$ and $\partial w/\partial x_b = x_a - w$, whose images under ρ are $I - R_b$ and $R_a - I$, the rotations commuting. By Proposition 10, $\tau(w) = (I-R_b)(I-R_a)p_a + (R_a-I)(I-R_b)p_b$. For odd valences $I-R_a$ and $I-R_b$ are $\sqrt{2}$ times a rotation. The exponent sums of w are zero. □

Measured on the male body. We implemented Proposition 10 on the T-mesh of Section 8.2. The patches are developed along a breadth-first tree of patches; the arcs this tree does not cross form a spanning tree of the 1,295 nodes, and the surface cut along it is a disc with one chart. Crossing an arc of the tree changes the chart by a motion; rooted at a singular vertex, every such motion is the composition, in the cyclic order round the node, of the motions of the arcs below it and of the rotation about the node, the recursion in the proof of Proposition 3(a). It holds at every node of the tree with one convention of order, found among eight, the places p_i being corners of the nodes in the chart. The holonomy of every turn is then a word in the rotations, and the formula of Proposition 10 gives its translation, in both components, for all 1,299 turns of the eight ways (identities modulo the balances; measured throughout this paragraph). Corollary 2 holds, at the whole lengths of the engine, on all 1,299 turns, 718 simple, 101 with one self-crossing and 480 with more, and on the 14 infeasible sets of Table 4 with whole multipliers (all but the set of fifteen). For the set at the left hand, with multipliers $(7, 5, 2, 3)$, the cycle winds an even number of times round every vertex of odd valence: the corollary is why its certificate $-2F$ is even at every whole length.

Turns that wind alike have, on this body, the same shift up to sign: the 1,299 turns give 293 distinct pairs of winding numbers and shift, and no two with the same winding numbers have shifts that differ by more than a sign; the 294 relations between two or three winding vectors with coefficients ±1, ±2 hold for the shifts, for some choice of signs. Proposition 11 is nevertheless not idle: the turns 18 of way 0 (7 self-crossings), 22 of way 0 (simple), 101 of way 4 (4) and 65 of way 5 (17) satisfy $w_{18} = w_{22} + w_{101} + w_{65}$ exactly, round every singular vertex, while no choice of quarter-turn frames makes their translations satisfy the same relation (the smallest residual has norm 0.25 on an orthonormal basis of the balanced lengths). A "winding Burgers formula" with numbers for weights is thus false, on the body as in principle; it holds modulo 2 (Corollary 2), and over the reals with the twisted windings.

The conjecture in these terms. Since $\sum_i (I-R_i)^\top W_i = 0$, the vertices with non-zero twisted winding can be grouped so that $\sum_{i\in G}(I-R_i)^\top W_i = 0$ in each group G; each group then contributes a combination of separations of its own vertices. A group of a vertex of valence three and one of valence five contributes $\langle V, p_c - p_{c'}\rangle$ with $V = (I-R_c)^\top W_c$, a vector whose coordinates have equal parity, that is $a(d_1+d_2) + b(d_1-d_2)$ with a, b whole and $d = p_c - p_{c'}$: the diagonal

forms of Proposition 4, which a staircase makes sign-definite (Proposition 7). An exchange is a group of two vertices of equal valence, a cap a group of total index four. So Conjecture 2 says that the twisted windings of an irreducible certificate, in a suitable development, split into such groups, each with a form of the sign of the certificate. At the left hand, developed from the patch of turn 76, twelve vertices have a non-zero twisted winding. The vertices 231 (valence three) and 228 (valence five) have $W = (-5, -5)$ and $(-5, 5)$; their group contributes, modulo the balances, exactly $10\,\ell_{1805}$ to $-(7s_{66} + 5s_{71} + 2s_{75} + 3s_{76}) = 10\,\ell_{1805} + 2\ell_{1817} + 4\ell_{1853} + 2(\ell_{1870} + \ell_{1879} + \ell_{1969} + \ell_{1998} + \ell_{2029})$. A separatrix of 231 (trace 915, 7.76 edges) stops on a separatrix of 228 at the T-junction 1142, and arc 1805 joins that junction to 228, 0.063 edges long: ℓ_{1805} is the offset of the pair across the line of the field that nearly joins them, the case of Proposition 4(ii) with $\delta = \ell_{1805}$. The coefficient 5 of F is this dislocation, round which the weighted cycle goes five times in the twisted sense. The ten other vertices, six of valence three and four of valence five, form a group of total index 2 whose form has the sign of the certificate, but an exhaustive search finds no split of them into smaller groups whose forms all have that sign, in this development. The sets of seven are less clear. In way 5, 24 vertices have a non-zero twisted winding; the pair 102, 103 exchanged in configuration B has $W = \pm(4, -4)$ and a form of the sign of the certificate, the pair 109, 118 of configuration C has $W = \pm(4, 0)$ and a form of the opposite sign, and the other groups found have no sign. In way 7 the twisted windings we compute are non-zero at 257 of the 261 vertices, nearly all ± 2 in one coordinate: the long turns wind round most of the body, and we did not remove the part that is a multiple of the relation. Conjecture 2 remains open; Proposition 10 reduces it to a statement about how the twisted windings of a certificate group into clusters.

8.5 Moving singularities in the parametrization

Proposition 3 suggests a lever before quantization: move the singular vertices so that the Burgers sum of each ring vanishes. We tested it on the seamless maps of the development models (measured throughout this subsection; the experiment is a patch of the engine, not part of it).

The Burgers sum on other models. For every ring of the seamless map that leans by at most 0.12, we computed the sum of Proposition 3(a) over the singular vertices of its smaller side, cut open along a tree of faces and ordered by first visit along its border. On seven closed surfaces of genus zero (blob, rock, creature, spot, pig, ogre, blub), 3,600 of the 3,604 rings that bound a disc are reproduced to $10^{-5}(1 + L)$, L the length of the ring; the reading fails only where a side is not a disc (borders on suzanne, handles on bob, part and cage).

Moving a singularity does not move the shift. The linearization $\delta s = \langle e_2, Q_i(I - R_i)\,\delta p_i \rangle$ assumes the other singular vertices keep their developed positions. They do not: on the blob at 3,000 quads, moving one pole of a cap one edge changes the shift of the cap ring (length 30.7, shift -1.065) by -0.13 with a local re-solve of the map, -0.02 with a global one, where the sum predicts -0.60; walking it six edges away saturates at -0.18; in the map, the three other poles move by 0.3 to 0.5 and absorb the rest. Over twelve moves per model, the measured change is 0.12, 0.03 and 0.01 of the predicted one on the blob, the creature and the pig. Rotating the field rigidly by $\alpha = |s|/L$ on the smaller side instead moves the shift by 26%, 46%, 64% and 87% of αL as the rotated region grows by 0, 10, 20 and 40 rings of vertices, at almost no cost in distortion (0.3362 to 0.3371). By Proposition 3(c) the shift is a property of the whole cylinder: shearing a long cylinder is expensive in the least-squares map, displacing a cone point in it is not, so the map keeps the shift the field gives the cylinder and moves its cone points instead. The identity holds exactly; it is the developed positions of the cone points that the field sets, not their places on the surface.

Two levers that fail. Cancelling the pairs of valences three and five of a twisted annulus changes the shift by its $\pm d_1 \pm d_2$ (Proposition 4), but raised the spiral edges of the blob at 3,000 quads from 11.6% to 23–36%. Rotating the field cylinder by cylinder to bring the shifts to zero reduced the shifts of the map but moved the twist onto rings of 90 to 200 edges, whose sensitivity grows with their length: over 36 runs (12 models, 3 sizes), 2 improved and 4 regressed, the mean gain coming from a single run, which we read as chaos rather than progress. Our reading, a conjecture: moving singularities would act through the Burgers sum only in a map whose scale is free to follow them, a cone metric; with a map of fixed scale the lever is the orientation of the field along whole cylinders, long rings must be handled jointly, and the conflicts are better attacked in quantization, where the staircases and dislocations of this section live.

9 Whole lengths

This section returns to the male body of Section 7, now in whole numbers. The propositions are proved; the numbers are measured, with exact integer or rational arithmetic where we say so, and otherwise from HiGHS through SciPy, stopped at a time limit (best solution found and proven bound).

Exact whole lengths. Closing exactly, in whole numbers, the turns found closable together with the bounds kept is feasible: HiGHS solves it in about two minutes for three of the four ways tried (the fourth, with 87 turns, runs out of ten minutes). The solutions leave 1 to 4 held turns shifted where the engine leaves 13 to 44, and fewer of the turns not held as well (5 against 9, 5 against 11, 0 against 27), at a deviation from the real lengths 4% to 62% higher. Yet the quad meshes they make spiral *more*: 12.2% of their edges against 8.8% for the first way, 3.7% against 1.1% for the second, and the third breaks the structure. Most of the loops that spiral are thus not turns our search finds: closing every turn it finds is not enough.

Closing by cuts. Walking every strip of the mesh the engine makes (Proposition 2) finds about 220 ways round per way of closing; each strip is found twice, once in each direction, with opposite shift expressions. Of these, 12 to 24 come back shifted although the field itself shifts them by less than half a row: the sampled walks of Section 4 had found only 4 to 10 of those, and held 1 to 5. A first selection of rings by that criterion (the field's shift under half a row) proved too narrow: the twisted thigh, shifted 1.65 rows a turn by the field, is a ring a modeller wants closed. Closing those rings by cuts converges (12, then 14, then 6 shifted over three rounds) but leaves the others free, and the mesh spirals more than with the engine (18% of the edges of way 2 against 1.1%). Selecting rings by their lean instead, as the engine does, gives 222 of the 230 ways round of way 2; 218 of them (109 strips) can be closed together in real numbers with every bound of separation kept. Posed directly, HiGHS found no whole solution within ten minutes, nor in a later run of thirty. The rest of this section explains why, and what that set of rings really costs.

The balances form a bidirected graph. The matrix B has two rows per patch, one per direction, 1,552 rows for the 776 patches, and one column per arc. An arc lies on a side of each of the two patches it borders, so its column has exactly two non-zero entries, each $+1$ or -1 according to the side of the patch it lies on (measured on all eight systems; in way 2, 958 arcs have signs $+/-$, 557 have $-/-$ and 554 have $+/+$). In the language of signed graphs, B is the incidence matrix of a *bidirected graph* G whose nodes are the balance equations and whose edges are the arcs. Call a cycle of G *balanced* if values ± 1 on its edges have a zero balance at each of its nodes, and *unbalanced* otherwise.

Lemma 3 (Handshake)**.** *For every $x \in \mathbb{Z}^n$, $\sum_r (Bx)_r$ is even.*

Proof. It is $\sum_a x_a \sum_r B_{ra}$, and each column sums to 0 or ± 2. □

So the rows of B are dependent modulo 2, although they are independent over the reals: a system $Bx = d$ with $\sum_r d_r$ odd has real solutions but no whole one. Consistently, integer elimination of B ends with one row whose entries have greatest common divisor 2, all other pivots being units (measured in all eight ways). This does not affect the balanced lengths:

Proposition 12 (Free arcs)**.** *Let G be connected and contain an unbalanced cycle. Let T be the edges of a spanning tree of G and one more edge closing an unbalanced cycle, and F the other edges. Then B has full row rank, and every $x_F \in \mathbb{Z}^F$ extends to exactly one $x \in \mathbb{Z}^n$ with $Bx = 0$. The whole balanced lengths thus form a lattice isomorphic to $\mathbb{Z}^F$, of dimension $n - m$ for m balances, with a basis of vectors z_f, $f \in F$, each supported on $T \cup \{f\}$ with $z_f(f) = 1$ and entries in $\{0, \pm 1, \pm 2\}$.*

Proof. Such a T exists: the sign of a cycle is a homomorphism from the cycle space of G over $\mathbb{Z}/2$, of which the fundamental cycles of a spanning tree are a basis, so if all of them were balanced every cycle would be. T has m edges. If $B_T y = 0$, a node met by a single edge of T forces that edge to zero; removing such edges repeatedly leaves the unbalanced cycle, on which each value fixes the next up to sign, and going round, since the cycle is unbalanced, forces them all to zero. So the square matrix B_T is invertible, B has full row rank, and x_F has exactly one real extension; it remains to see that it is whole. For $f \in F$, the graph $T \cup \{f\}$ has a one-dimensional space of balanced values. Removing the edges met at a node of degree one leaves either three paths between two nodes, whose three cycles are the unbalanced cycle of T, one more unbalanced and one balanced (the number of unbalanced cycles of such a graph is even), or two cycles joined by a path, possibly of length zero. In the first case the balanced cycle contains f, since the cycle of T does not, and carries values ± 1. In the second, the cycle through f either is balanced and carries values ± 1, or is unbalanced, and then values ± 1 on both cycles and ± 2 on the joining path are balanced. Either way there is a whole balanced z_f supported on $T \cup \{f\}$ with $z_f(f) = \pm 1$; changing its sign if needed, $x = \sum_{f \in F} x_f z_f$ is the whole extension. □

On the male body G is connected with an unbalanced cycle, B has rank 1,552, and the lattice of balanced lengths has dimension $2{,}069 - 1{,}552 = 517$ (measured in all eight ways). Balances alone thus never obstruct whole lengths: 517 arcs can be given any whole lengths and the others follow. With the shifts of the 218 rings of way 2 the rank grows by 105 and the lattice of closed quantizations has dimension 412; integer elimination still uses unit pivots only, with one row of greatest common divisor 2 (the shift of one ring is even as soon as the balances and the rings before it hold). Adding the shifts of the held turns instead gives dimensions 469, 439, 465, 431, 435, 363 and 431 in ways 0 to 6; in ways 4, 5 and 6 elimination needs steps of Euclid's algorithm, so turn rows can destroy the free-arc structure.

Integer solutions exist. By Remark 5, whole lengths closing the 218 rings within the bounds exist. HiGHS finds some in one second once it is told which arcs are zero in *every* real solution: closing the 218 rings with $\ell \geq 0$ forces 171 arcs to zero (measured), the implicit equalities of the cone, each certified by a non-negative combination of shifts as in Remark 4. There are 23 distinct certificates: 13 single rings whose shift is sign-definite (round fingers and toes), 4 pairs with multipliers $(1, 1)$ (left foot, right foot, head, top of the head), and 6 combinations of four to six rings (torso, neck). Given as upper bounds, these zeros make the problem easy; left implicit, the solver works on a face of lower dimension than the problem without knowing it, which is what the ten- and thirty-minute failures were. They are not an integer obstruction.

The difficulty is their size. The 171 arcs forced to zero carry 350.8 quad edges of target length, the longest 10.1. The real lengths closing the 218 rings nearest the targets, in the weighted deviation, make 3,960 quads instead of the 10,496 of the targets. With the deviation the engine measures, the mean of $|\ell_a - \lambda r_a|$, a linear program shows that no real lengths closing the 218 rings have a mean deviation below 0.817, nor, after dropping 7 strips (14 ways round) so that every arc with a target of at least 1.5 stays open, below 0.758; balanced lengths without rings reach 0.085. A lower bound in real numbers holds for whole numbers too. The engine refuses quantizations whose deviation exceeds 1.5 times that of the plain search, $1.5 \times 0.316 = 0.474$. So the 218 rings cannot close at the size asked for within the engine's tolerance, whatever the rounding: the obstruction is in the reals.

The 190 rings that fit. A subset of 190 of the 218 ways round (chosen greedily, so 190 is only a lower bound on what fits) closes in real numbers at a mean deviation of 0.399, keeping the number of quads (10,502) at that of the targets to first order: the linear form $\nabla q(t) \cdot \ell \geq \nabla q(t) \cdot t$, where $q(\ell)$ is the number of quads, a quadratic function of the lengths, and t the targets. In whole numbers we followed the reformulation of Aardal, Hurkens and Lenstra [1]: a basis M of the lattice of closed quantizations, with the 74 implicit zeros of this set imposed first (dimension 407), reduced by LLL [16] for a weighted inner product, which leaves entries in $[-2, 2]$; then $\ell = My$ with y whole, so that no equation is left, and a mixed-integer program over y. Babai's nearest plane [2] from the real solution gives a mean deviation of 0.586 but 10 arcs below their bounds. HiGHS, stopped after 1,200 seconds, gives whole lengths closing all 190 rings within every bound, with a mean deviation of 0.619 and 10,568 quads, and proves that no whole lengths do better than 0.536 under the same constraints, the first-order proxy of the number of quads included. Against 0.399 in real numbers, this rounding gap is the one integer obstruction we found, and it is not a parity condition: every infeasible set of Table 4 is infeasible in real numbers already.

Which turns, not rounding. Given to the engine in place of the lengths of way 2, these whole lengths close all 190 rings, with 10,568 quads and 2.5% of spiral edges against 1.1% for the engine's own lengths (the share of that way alone: the deviation cap would refuse it). At the new lengths the strips are different walks: walking all of them finds 230 ways round, 156 of them shifted (168 at the engine's lengths), and of the 214 that lean less than 0.12, 142 are shifted (162 of 222 at the engine's lengths). Closing the rings of one set of lengths moves the strips, and other loops spiral. On this body, then, whole lengths are not what is missing: rounding costs between 0.137 (proven) and 0.220 (found) of mean deviation above the real optimum, and it is the choice of the turns to close, at the lengths where they are the strips of the mesh, that decides how much the mesh spirals.

10 Helices that every quantization keeps

The obstructions of Section 8 say that a set of turns cannot close together. Turned around, they bound from below the number of helices that any quantization must leave, and so tell how far a heuristic such as the one of Section 4 is from the best possible. We are not aware of an earlier lower bound of this kind (Section 2).

Theorem 1 (Forced helices)**.** *Let H be a finite set of turns of a T-mesh and P a non-empty set of lengths containing every quantization considered: for instance the balanced lengths meeting the bounds of separation, $\ell \geq b$, possibly with a cap $\frac{1}{n}\sum_a |\ell_a - t_a| \leq D$ on the mean deviation from the targets t, or with a given number of quads. Let*

$$\tau_P(H) = \min\{\, |X| : X \subseteq H, \text{ the turns of } H \setminus X \text{ can be closed together by real lengths in } P \,\}.$$

Then for every $\ell \in P$, whole or not, at least $\tau_P(H)$ turns of H have $s_\gamma(\ell) \neq 0$. If H contains k pairwise disjoint subsets none of which can be closed within P, then $\tau_P(H) \geq k$.

Proof. At a whole $\ell \in P$, the turns of H with zero shift are closed by ℓ itself, which is real and lies in P; so the shifted turns form a set X as in the definition, and $|X| \geq \tau_P(H)$. A set X whose removal leaves a closable set meets each of the k subsets, which are disjoint, so $|X| \geq k$. □

The theorem is elementary; its content is in what makes subsets infeasible, which Section 8 relates to the singularities. The proof uses neither convexity, nor linearity of the shifts, nor whole lengths; convexity is what lets $\tau_P(H)$ be bounded by linear programs, as below. An earlier draft asked P to be convex. The bounds of separation are what make the shifts of a dislocation non-zero, and the statement becomes one about parallel rings.

Corollary 3 (Chains of dislocations)**.** *Let $\gamma_0, \gamma_1, \ldots, \gamma_m$ be turns and suppose that for $i = 1, \ldots, m$*

$$s_{\gamma_{i-1}}(\ell) - s_{\gamma_i}(\ell) = \varepsilon_i \, F_i(\ell) \quad \text{for all balanced } \ell,$$

with $\varepsilon_i \in \{+1, -1\}$ and forms F_i that are positive on the whole points of P; for instance F_i the length of a staircase (Proposition 7) between a vertex of valence three and one of valence five lying between γ_{i-1} and γ_i (Propositions 4 and 3(b)), the staircase containing an arc held at length at least one. Let q be the number of indices $i < m$ with $\varepsilon_{i+1} \neq \varepsilon_i$. Then for every whole $\ell \in P$ at most $q + 1$ of the turns γ_i are rings: at least $m - q$ are helices. In particular $m + 1$ nested parallel turns separated by m dislocations whose staircases all climb the same way leave at least m helices.

Proof. If γ_i and γ_j, $i < j$, are both rings, then $\sum_{k=i+1}^{j} \varepsilon_k F_k(\ell) = s_{\gamma_i}(\ell) - s_{\gamma_j}(\ell) = 0$; the $F_k(\ell)$ being positive, the signs $\varepsilon_{i+1}, \ldots, \varepsilon_j$ are not all equal, so a change of sign lies between i and j. Between two consecutive rings of the chain lies a change of sign of its own, so there are at most $q + 1$ rings. (Equivalently, in Theorem 1, every pair of rings of a monotone stretch is an infeasible pair.) □

Several chains, or disjoint conflicts of any of the kinds of Section 8.3, add up by the second part of the theorem. The corollary is stated with signs ε_i and forms F_i that the singular vertices between the turns determine: by Proposition 3(b), $s_{\gamma_{i-1}} - s_{\gamma_i}$ is the across component of the Burgers vector of the annulus between them, computed from the positions and valences of its singular vertices alone.

Remark 6 (What the bound counts)**.** *The theorem counts turns with a non-zero shift. At the lengths where a turn is a strip of the mesh (Proposition 2) a non-zero shift is a helix; at other lengths the same walk may not be a strip. One spiral strip may also carry several turns of H, and the measure of Section 5 counts edge loops that spiral within four quads, which is yet another count. The bound is therefore a bound on turns, and is compared below with the turns the engine leaves shifted.*

Measured on the male body. For each way of Section 7 we took for H its held turns, each up to sign once, and for P the balanced lengths meeting the bounds of separation, then also the cap $D = 0.474$ on the mean deviation, the engine's own (1.5 times the 0.316 of its plain search), with the targets of that way. Lower bounds on $\tau_P(H)$ come from a greedy packing of disjoint irreducible infeasible subsets, each found by a deletion filter and certified infeasible by HiGHS; upper bounds from the largest closable subset a mixed-integer program finds, checked feasible. We also took for H the strips of the engine's own mesh for way 2 (the way it keeps) whose real shift leans less than 0.12, the rings of Section 9: 222 ways round, 111 strips. Table 6 compares with what the engine leaves shifted at its whole lengths.

In every way the bounds of separation alone force very few helices: one turn in ways 0, 2, 4 and 6, the turn round the left wrist of Section 8.3, and four to nine in the ways that hold the

way	turns	forced, bounds	forced, bounds and cap	left shifted by the engine
0	48	1	1	13
1	78	5	5	39
2	52	1	1	16
3	86	4	4	44
4	84	1	1	49
5	160	6	7–17	115
6	88	1	1	44
7	169	8–9	8–26	123
2, strips of the mesh	111	2	3–12	81

Table 6: Male body at 10,590 quads: turns of each family that every quantization in P leaves shifted (Theorem 1), with P given by the bounds of separation, then also by the engine's cap on the mean deviation; a single number is exact (packing and mixed-integer program agree); a range gives the larger of the packing and the solver's proven bound, and the best subset found within the time limit (120 to 1,800 seconds). Last column: turns of the family with a non-zero shift at the engine's whole lengths for that way. Turns are counted up to sign; "strips": the ways round of the engine's mesh for way 2 leaning less than 0.12, two per strip.

most turns, where the packings are the irreducible sets of Table 4. The bounds are attained: the mixed-integer program closes all the other turns together (in way 7 its optimum, 160 of 169, is computed with every length at most 1,000 above its target, which the packing does not need), and Section 9 found whole lengths leaving only 1 to 4 held turns shifted. The engine leaves 13 to 123. Of the 111 strips of its own mesh that lean like rings, at least 2 must be helices under every quantization meeting the bounds, one of them round the wrist; the engine leaves 81 shifted, and the measure of Section 5 counts on that mesh 1.1% of spiral edges (1.15% to 1.59%, in 8 to 9 spiral loops, over recent runs of the engine). The engine's cap on the deviation does not raise the proven bound in seven of the eight ways; it adds one forced turn in way 5 (an infeasible set of 72 turns), and among the strips it raises the bound to 3, proven by the solver's dual bound after 1,800 seconds, against at most 12: with the cap kept, 99 of the 111 strips can be closed together in real numbers.

So on this body the singular configuration and its separation explain only a small part of the helices the engine leaves: the turns the theorem forces are one to a few per way, and most of the shifted turns of the engine could be closed by other lengths in P. What keeps them open in practice is the size and the deviation, as Section 9 found, and the heuristic, which keeps far fewer turns than can be closed (Table 2). The theorem holds with the number of quads fixed, a non-convex constraint; how to compute or certify $\tau_P(H)$ for such a P is left open.

Status of Sections 8 to 10. *Proved:* Lemmas 1, 2 and 3; Propositions 3 (with the order of first visit made precise), 4, 5, 6, 7, 8, 9, 10, 11 and 12, Corollary 2, Theorem 1 and Corollary 3, each under its hypotheses; Remark 4, and the existence of whole solutions when real ones exist. *Machine-checked:* the algebraic cores of most of these, with the topological steps as hypotheses (Section 10.1). *Measured:* on the T-mesh of the male body, every statement of Section 8.1 wherever it applies (Section 8.2); the configurations of the conflicts (Table 5), the staircases, the separatrix at the wrist and the windings of the larger sets, as identities modulo the balances, without the homotopies; the Fox–Burgers formula on all 1,299 turns and the checkerboard rule for combinations on all of them and on 14 infeasible sets; the Burgers sum on 3,600 of 3,604 disc-bounding rings of seven other models, and the response of the shift to moving a singular vertex or rotating the field (Section 8.5); the four turns whose winding numbers are related while their translations are not; the dislocation of weight 5 at the left hand; the structure and ranks of the eight systems, the 171 implicit zeros and their certificates, the bounds 0.817, 0.758

and 0.536, the infeasible sets of Table 4 (exactly where marked), the forced helices of Table 6, the meshes. *Refuted:* that the conflicts between two turns are pairs of valences three and five on one trace (0 of 13); Conjecture 1 as stated; and a formula for combinations of shifts in the winding numbers alone (in principle and on the body). *Conjectured:* Conjecture 2, reduced by Proposition 10 to the grouping of twisted windings; the left hand is half explained, the sets of seven and configurations E and F are not described; that moving singularities in a map of fixed scale cannot set the shifts (Section 8.5) is a reading of measurements.

10.1 What is machine-checked

The algebraic cores of the propositions of Sections 8 to 10 are proved in Lean 4 with Mathlib, in the folder `docs/research/closing-helices/lean` of the repository, pinned to Mathlib `v4.35.0` and the toolchain `leanprover/lean4:v4.35.0`; `make proofs` fetches Mathlib compiled and checks them, in about thirty seconds once the cache is there. The topological facts the proofs use, that a loop develops into a given motion, that a curve bounds a disc, that the fundamental group of Σ° is presented by the loops λ_i, that the quad graph of a sphere is bipartite, are not formalized: each is an explicit hypothesis of the theorem that uses it, and no proof is left open. The folder's `README.md` maps each statement to its Lean theorems.

What is checked.

- For motions whose linear parts lie in any ring acting on any module: the translation of $\rho_m \circ \cdots \circ \rho_1$ is $\sum_i Q_i(I - R_i)p_i$ with $Q_i = R_m \cdots R_{i+1}$ (Proposition 3(a)); the recursion of the gluing maps of a tree of cuts unfolds into the order of first visit, and the order of last visit gives another vector already for two vertices; the holonomy of the border of a cut annulus (Proposition 3(b)); exchanges (Proposition 8); the conjugation of Proposition 9; the linearity of Proposition 1.
- Over the reals: the four orders and orientations of a dislocation, the forms $\pm d_1 \pm d_2$ and $|B_A|^2 = 2|d|^2$, straight chains (Proposition 4(i), (ii)); staircases (Proposition 7); the positivity of a non-negative form with a held arc (Remark 4, first part).
- At whole lengths: the parities of $(I - R^{\pm 1})p$ and $(I - R^2)p$; the checkerboard rule (Proposition 5); the induction of the proof of Proposition 6, given a 2-colouring of the quad graph, and the second proof of Remark 2.
- The Fox–Burgers formula on words, which is well defined on the free group, the fundamental formula, the twisted windings and $\sum_i (I - R_i)^\top W_i = 0$, the insertion of the relation (Proposition 10); the commutator (Proposition 11); Corollary 2, with the two arguments of its proof on the colour of the origin and the constant of the winding numbers.
- Lemma 3, Theorem 1 and Corollary 3.

Not checked: Lemmas 1 and 2, Proposition 2, Proposition 12, which needs a theory of bidirected graphs, Farkas' lemma in Remark 4, and every measured statement.

Gaps found. Formalizing found no error of sign or convention, but a few gaps, corrected in this version: *genus zero* must mean a closed sphere in Section 8, and the parity of a turn fails with borders (Remark 2); inserting the relation in a word after a prefix a brings $\rho(a)$ into the change of the twisted windings (Proposition 10); Theorem 1 needs P non-empty, for $\tau_P(H)$ to be defined, but neither convex nor of whole lengths, so it holds for quantizations of a given size; the frame of the first side is that of the first patch up to a quarter-turn, not only up to signs (Proposition 6); the edge closing an unbalanced cycle of Proposition 12 exists, which the proof did not say; Proposition 7 holds for any path on whose steps the form of the shift difference is one.

Other rotation orders. Nothing in the Burgers sum, the order of first visit or the Fox–Burgers formula uses that R has order four: they hold for the rotations of order six of triangle and hexagonal meshes, for cone metrics of any angles, and for motions of space. What changes is parity. When every rotation is a power z^{j_i} of one rotation z, every Fox derivative is congruent to its exponent sum modulo $1-z$, and $1-z^j$ to $j(1-z)$ modulo $(1-z)^2$, so

$$\tau(w) \equiv (1-z)\sum_i \epsilon\Big(\frac{\partial w}{\partial x_i}\Big) j_i\, p_i \pmod{(1-z)^2},$$

proved in any commutative ring. For quarter-turns, $(1-z)^2 = -2z$, and this is Corollary 2, with $j_i = 4 - k(c_i)$. For rotations of order three, $(1-z)^2 = -3z$: a rule modulo three, with three colours. For rotations of order six, $1-z$ is a unit: there is no colouring rule on triangle or hexagonal meshes. A dislocation of order n has $|B| = 2\sin(\pi/n)\,|d|$: $\sqrt{2}\,|d|$ for quads, $|d|$ for a pair of valences five and seven on a triangle mesh; the forms $\pm d_1 \pm d_2$ are particular to quads. Theorem 1, Corollary 3 and Lemma 3 do not depend on the geometry at all.

Invariants for the engine. The checked identities give tests the engine can run on its own quantities: on a closed surface of genus zero, the two components of the translation of every turn, its length and its shift, have the same parity at the whole lengths of the integer search, so every ring has an even number of quads round it; the coefficients of every shift expression are whole; every column of the balances has two entries ± 1, so a balance system with an odd total right-hand side has no whole solution; where the developed cone points are known, the translation of a turn bounding a disc is the Burgers sum in the order of first visit, and its two components have the parity of the number of vertices of odd valence on points of odd colour; a turn whose shift is a non-negative form with a held arc, or both turns of a staircase with a held arc, are never closed; and the number of held turns left shifted is at least the number of disjoint infeasible sets found. The README of the folder lists them with the engine's names.

11 Open problems

Open problem 1 (Exact formulation)**.** *Minimize the deviation of the quantization subject to balance, validity and $s_\gamma(\ell) = 0$ for a set of turns. This is a minimum-deviation flow with linear side constraints; how much of the exact machinery of [10] carries over is open. Section 7 shows that the linear relaxation already closes almost all held turns.*

Open problem 2 (Closing in whole numbers)**.** *The solutions of the balance and shift equations in whole numbers form a lattice, and Section 9 searched it through a reduced basis and the reformulation of [1]: whole lengths closing every ring asked for are found, but in minutes, with the number of quads held only to first order, and at a deviation* 0.14 *to* 0.22 *above the real optimum. Can the integer search of the quantizer move within this lattice at the speed of a flow solver, with the number of quads held exactly? This matters once the right turns are chosen; Section 9 suggests it is not, on its own, where the loops are lost.*

Open problem 3 (Which turns)**.** *This is, on the evidence of Section 9, the central problem: closing rings changes the lengths, the lengths change the strips, and the strips of the new mesh are other walks. The straight walks that return are infinite in number; we sample them, and Section 9 shows the sample misses most of the loops that spiral. Is there a finite set of turns, a basis in some homological sense of the regular region of the T-mesh, whose closing implies that every loop of the quad mesh closes? Can the loops of the quad mesh be read off the T-mesh and its lengths directly, so that every one is accounted for?*

Open problem 4 (Feasibility)**.** *On a human torso, some held turns cannot be closed together: the projection lets them go. Our hypothesis was that pairs of singularities between two rings*

force a dislocation, and that closing all rings needs the poles to face one another, which the separation of singularities forbids at the scale of the T-mesh; Propositions 4 and 7 prove this for a single pair joined by a straight chain or a staircase. A capped cylinder shows the same conflict: its layout of six patches exists, but only with arcs our quantizer will not bring to zero. On the male body the conflict comes down to one arc in four of the eight ways, to 9 to 30 arcs in the others (Section 7), and to irreducible sets of one to fifteen turns; the smallest are dislocations, exchanges of two vertices, caps and a winding separatrix (Section 8.3). When is a set of turns simultaneously closable, in terms of the singularity configuration? Conjecture 2 proposes that the obstructions are always sums of such local ones; Proposition 10 reduces it to how the twisted windings of a certificate group into clusters.

Open problem 5 (Lower bounds with the size)**.** *Theorem 1 bounds the helices any quantization leaves within a set of lengths, and on the male body the bounds of separation force only one to nine turns. The theorem holds for quantizations of a given size, but the number of quads is a quadratic, not convex, function of the lengths, and the linear programs that bound $\tau_P(H)$ no longer apply. How can $\tau_P(H)$ be bounded for quantizations of a given size, and how close does it come to what the engine leaves?*

Open problem 6 (Finite trajectories)**.** *In the language of [14]: given a singularity configuration on a surface of genus zero, when does a quartic differential with all trajectories finite exist, and how close to a given cross field can it be chosen? Our turns are its discrete, integer version.*

Towards proofs. The algebraic cores of Sections 8 to 10 are machine-checked (Section 10.1); their topological hypotheses are not. Proposition 1 is elementary but its statement over a T-mesh with T-junctions is easy to get wrong. A formal model of an abstract complex of rectangles with integer side lengths, its turns and its developing map would turn those hypotheses into theorems, and could carry the sufficiency question of the third problem for special classes of layouts.

Reproducibility. The method is implemented in openclare, Rust, branches `close-quantized` (closing, measure) and `quadsphere` (confidence floor, merged traces); the bench is driven by `bench/run.py` and the marks by `scripts/scores.py`. The export of the T-mesh used in Section 8.2 is an experimental hook of the branch `rd-exact`, not part of the engine. The complete walk of the strips and the cut rounds are on the branch `complete-helices`; the Lean proofs are in `docs/research/closing-helices/lean` and replay with `make proofs`.

AI assistance

This note and the software it describes were made by a human author working with an AI system. Following the practice recommended by ACM and by Nature, we state how the work was divided. No AI system is an author of this note.

The human author chose the problem and its aims: edge loops that close into rings, the quad sphere with the eight poles of a cube as a target, and, as an artist, a mesh with few singular vertices and few spirals rather than perfectly square quads, always made of quads only. He set the direction of the research and the order of the work, decided which results to pursue, which changes to merge into the engine and which to drop, authorized the tools (LaTeX, Lean and Mathlib), and judged the results through the measurements, pictures and drafts presented to him.

The AI system , Claude Opus 5.5 (Anthropic), used through Claude Code in October 2026, with several sub-agents of the same model working in parallel, did the rest. It wrote the changes to the engine and the experiment scripts, ran the experiments and the bench, carried out the exact computations (SymPy, HiGHS through SciPy), searched the literature, stated and proved the propositions, wrote the Lean formalization and drafted the text of this note. References were checked for existence, and the note attached to each entry of the bibliography says how much of the work was read: the abstract only, a part, or all of it.

What was checked, and how. The algebraic cores of Sections 8 to 10 are machine-checked in Lean (Section 10.1), using only Lean's standard axioms. The topological statements that Lean takes as hypotheses, and the proofs given only on paper, were checked by AI agents rereading them and testing them numerically on the data (Section 8.2), not by a human expert. Every measured figure can be reproduced from the scripts and branches named above. The note is a draft and has not been peer reviewed. Errors remain possible, and the human author, who publishes the note, is responsible for its content.

References

[1] Karen Aardal, Cor A. J. Hurkens, and Arjen K. Lenstra. Solving a system of linear Diophantine equations with lower and upper bounds on the variables. *Mathematics of Operations Research*, 25(3):427–442, 2000. doi: 10.1287/moor.25.3.427.12219. Abstract read.

[2] László Babai. On Lovász' lattice reduction and the nearest lattice point problem. *Combinatorica*, 6(1):1–13, 1986. doi: 10.1007/BF02579403. Abstract read.

[3] David Bommes, Henrik Zimmer, and Leif Kobbelt. Mixed-integer quadrangulation. *ACM Transactions on Graphics*, 28(3), 2009. Proc. SIGGRAPH 2009.

[4] David Bommes, Timm Lempfer, and Leif Kobbelt. Global structure optimization of quadrilateral meshes. *Computer Graphics Forum*, 30(2):375–384, 2011. Proc. Eurographics 2011. Read in full.

[5] Marcel Campen. Partitioning surfaces into quadrilateral patches: A survey. *Computer Graphics Forum*, 36(8):567–588, 2017. doi: 10.1111/cgf.13153. Abstract read.

[6] Marcel Campen, David Bommes, and Leif Kobbelt. Dual loops meshing: Quality quad layouts on manifolds. *ACM Transactions on Graphics*, 31(4), 2012. doi: 10.1145/2185520.2185606. Proc. SIGGRAPH 2012. Abstract read.

[7] Marcel Campen, David Bommes, and Leif Kobbelt. Quantized global parametrization. *ACM Transactions on Graphics*, 34(6):1–12, 2015. doi: 10.1145/2816795.2818140. Proc. SIGGRAPH Asia 2015. Abstract read.

[8] Wei Chen, Xiaopeng Zheng, Jingyao Ke, Na Lei, Zhongxuan Luo, and Xianfeng Gu. Quadrilateral mesh generation I: Metric based method. *Computer Methods in Applied Mechanics and Engineering*, 356:652–668, 2019. Theorems 3.7, 3.8 and Lemma 3.9 of the arXiv version read.

[9] Ralph H. Fox. Free differential calculus. I: Derivation in the free group ring. *Annals of Mathematics*, 57(3):547–560, 1953. doi: 10.2307/1969736. Cited for the definition of the derivatives and their fundamental formula, which are standard.

[10] Martin Heistermann, Jethro Warnett, and David Bommes. Min-deviation-flow in bi-directed graphs for T-mesh quantization. *ACM Transactions on Graphics*, 42(4), 2023. doi: 10.1145/3592437. Proc. SIGGRAPH 2023. Abstract read; our quantizer follows its formulation.

[11] Ivan Izmestiev, Robert B. Kusner, Günter Rote, Boris Springborn, and John M. Sullivan. There is no triangulation of the torus with vertex degrees 5, 6, ..., 6, 7 and related results: geometric proofs for combinatorial theorems. *Geometriae Dedicata*, 166:15–29, 2013. doi: 10.1007/s10711-012-9782-5. Read in full (arXiv v2): holonomy of the developing map, Burgers vector, no 3,5-quadrangulation of the torus.

[12] Felix Kälberer, Matthias Nieser, and Konrad Polthier. QuadCover – surface parameterization using branched coverings. *Computer Graphics Forum*, 26(3):375–384, 2007. doi: 10.1111/j.1467-8659.2007.01060.x. Proc. Eurographics 2007. Abstract read.

[13] Felix Kälberer, Matthias Nieser, and Konrad Polthier. Stripe parameterization of tubular surfaces. In *Topological Methods in Data Analysis and Visualization: Theory, Algorithms, and Applications*, Mathematics and Visualization, pages 13–26. Springer, 2011. doi: 10.1007/978-3-642-15014-2_2. Preprint read in full (Sections 3–4: integer gaps along cut paths, rounded).

[14] Na Lei, Xiaopeng Zheng, Zhongxuan Luo, Feng Luo, and Xianfeng Gu. Quadrilateral mesh generation II: Meromorphic quartic differentials and Abel-Jacobi condition. *Computer Methods in Applied Mechanics and Engineering*, 366, 2020. Main theorems read (Thm. 4.7–4.13).

[15] Na Lei, Yiming Zhu, Xiaopeng Zheng, Hang Si, Zhongxuan Luo, and Xianfeng Gu. Why cross fields are not equivalent to quadrilateral meshes. *Computer Methods in Applied Mechanics and Engineering*, 417:116442, 2023. doi: 10.1016/j.cma.2023.116442. Abstract read.

[16] Arjen K. Lenstra, Hendrik W. Lenstra, Jr., and László Lovász. Factoring polynomials with rational coefficients. *Mathematische Annalen*, 261:515–534, 1982. doi: 10.1007/BF01457454. Bibliographic record checked.

[17] Jian Liu, Chunshi Wang, Song Guo, Haohan Weng, Zhen Zhou, Zhiqi Li, Jiaao Yu, Yiling Zhu, Jing Xu, Biwen Lei, Zhuo Chen, and Chunchao Guo. QuadGPT: Native quadrilateral mesh generation with autoregressive models, 2025. Appendix B.2 read: quad rings and quad lines as reward signals.

[18] Max Lyon, Marcel Campen, and Leif Kobbelt. Quad layouts via constrained T-mesh quantization. *Computer Graphics Forum*, 40(2):305–314, 2021. doi: 10.1111/cgf.142634. Proc. Eurographics 2021. Abstract read; the full text was not openly accessible.

[19] Ashish Myles, Nico Pietroni, Denis Kovacs, and Denis Zorin. Feature-aligned T-meshes. *ACM Transactions on Graphics*, 29(4):1–11, 2010. As cited by [4].

[20] Ashish Myles, Nico Pietroni, and Denis Zorin. Robust field-aligned global parametrization. *ACM Transactions on Graphics*, 33(4):135:1–135:14, 2014. doi: 10.1145/2601097.2601154. Proc. SIGGRAPH 2014. Abstract read.

[21] Nico Pietroni, Enrico Puppo, Giorgio Marcias, Roberto Scopigno, and Paolo Cignoni. Tracing field-coherent quad layouts. *Computer Graphics Forum*, 35(7):485–496, 2016. doi: 10.1111/cgf.13045. Proc. Pacific Graphics 2016. Abstract read.

[22] Faniry H. Razafindrazaka and Konrad Polthier. Optimal base complexes for quadrilateral meshes. *Computer Aided Geometric Design*, 52–53:63–74, 2017. Bibliographic data checked (DBLP); text not read.

[23] Faniry H. Razafindrazaka, Ulrich Reitebuch, and Konrad Polthier. Perfect matching quad layouts for manifold meshes. *Computer Graphics Forum*, 34(5):219–228, 2015. doi: 10.1111/cgf.12710. Proc. SGP 2015. Abstract read.

[24] Kurt Strebel. *Quadratic Differentials*. Ergebnisse der Mathematik und ihrer Grenzgebiete. Springer, 1984.

[25] Xiaopeng Zheng, Yiming Zhu, Na Lei, Zhongxuan Luo, and Xianfeng Gu. Quadrilateral mesh generation III: Optimizing singularity configuration based on Abel-Jacobi theory, 2020. Abstract read.