

THÈSE POUR OBTENIR LE GRADE DE DOCTEUR DE L'UNIVERSITÉ DE MONTPELLIER

En informatique

École doctorale I2S

Unité de recherche LIRMM

Génération procédurale d'environnements sous-marins Procedural generation of underwater environments

Présentée par **Marc HARTLEY**
le [XX mois année]

Sous la direction de **Christophe FIORIO, Noura FARAJ**
et **Karen GODARY-DEJEAN**

Devant le jury composé de

[Prénom NOM, Titre, Affiliation]
[Prénom NOM, Titre, Affiliation]

[Statut jury]
[Statut jury]



Contents

Table of Contents

	v
1 Introduction	1
1.1 Contributions and outlines	1
1.1.1 Semantics	2
1.1.2 Modeling	2
1.1.3 Amplification	2
1.2 Procedural generation	2
1.2.1 Notable examples in the entertaining industry	5
1.2.2 Parallelization and GPU utilisation	6
1.3 Procedural terrain generation	7
1.3.1 Terrain representations	7
1.3.2 Terrain generation	10
1.3.3 User interaction [TODO]	12
1.3.4 Fluid simulations	13
1.3.5 Vegetation generation	15
1.4 Underwater landscapes	15
1.4.1 Main differences with aerial terrains	16
1.4.2 Coral reefs (biological aspects)	17
1.5 Prototype creation	24
 I Semantic Representation	 25
2 Generation de terrain sémantique	29
2.1 Introduction	30
2.2 Related works	32
2.3 Description of the method	33
2.3.1 Pipeline	33
2.3.2 Environmental objects \mathcal{O}	35
2.3.3 Environmental attributes \mathcal{E}	35
2.4 Placement of environmental objects in an environment	36
2.4.1 Fitness function ω	37
2.4.2 Skeleton fitting function Γ	38

2.5	Environmental modifiers	41
2.5.1	Environmental material modifiers	41
2.5.2	Height modifiers	41
2.5.3	Influence on water currents	42
2.5.4	Environment stability	42
2.6	User interactions	43
2.6.1	Direct interactions with the environmental objects	43
2.6.2	Indirect interaction with environmental objects	44
2.7	Results and discussion	46
2.8	Conclusion	48
II	Modelisation	51
3	Graphical representation of environmental objects	55
4	Automatic Generation of Coral Islands	57
4.1	Introduction	57
4.1.1	Multiple theories	58
4.1.2	Darwinian theory	60
4.1.3	Overview	60
4.2	Related works	61
4.3	Example generation	61
4.3.1	Closed form of coral growth	62
4.3.2	Labeling of the map	62
4.3.3	Automation	63
4.4	cGAN	63
4.4.1	Definition of cGAN	63
4.4.2	Why cGAN?	63
4.4.3	Training	63
4.4.4	Model usage	64
5	Generation of karst networks	65
5.1	Introduction	65
5.2	Related works	67
5.3	Space colonization	67
5.4	Our method	67
5.5	Modeling	68
5.6	User control	68
5.7	Results	69
5.8	Conclusion	69
III	Erosion Simulation	71
6	Érosion par particules	75
6.1	Introduction	76
6.2	State of the art	78
6.2.1	Terrain representations	78
6.2.2	Erosion processes	79

6.3	Particle erosion	79
6.3.1	Overview	80
6.3.2	Erosion process	80
6.3.3	Transport	81
6.4	Our erosion method	83
6.4.1	Application on height fields	84
6.4.2	Application on layered terrains	85
6.4.3	Application on implicit terrains	85
6.4.4	Application on voxel grids	86
6.5	Results	87
6.5.1	Rain	88
6.5.2	Coastal erosion	88
6.5.3	Rivers	89
6.5.4	Landslide	89
6.5.5	Karsts	89
6.5.6	Wind	90
6.5.7	Underwater currents	90
6.5.8	Multiple phenomena	90
6.6	Comparisons	91
6.6.1	Coastal erosion on implicit terrain representation	92
6.6.2	Wind erosion on voxel grid representation	92
6.6.3	Hydraulic erosion on height field representation	93
6.6.4	Wind erosion on stacked materials representation	94
6.7	Discussion	94
6.7.1	Realism	94
6.7.2	Usage of velocity fields	95
6.7.3	Performances	95
6.8	Conclusion	96
IV		99
Conclusion		101
References		101
Glossary		113

Abstract

This thesis, entitled "*Procedural terrain generation for underwater environments*", as its name suggests, focuses on the topic of procedural terrain generation with the specificity to tackle the underwater environments. Terrain generation is still an open research area in computer graphics as, with the emergence of simulation, rendering, and interaction techniques, the entire field is experiencing increased collaboration with terrain experts. Virtualization of the physical world helps users to see and manipulate it, allowing for a better understanding of it and bringing together numerous experts, gradually breaking down the boundaries of scientific disciplines. Terrain science brings together, in almost a direct manner, geologists, oceanologists, physicists, meteorologists, biologists, roboticists, computer scientists, 3D artists, and more. We focused our work on the inclusion of the user in the generation process of virtual worlds through fast and controllable algorithms, with the possibility to dive underwater.

This thesis is divided into three parts, guiding the user from the design of a landscape, its 3D modeling, until its finalization, step by step. Firstly, we will propose a formalization of terrain designing, allowing for the conception of environments in a semantic sense, abstracting away from 3D geometry and data structures.

Secondly, we will see how to give 3D form to these environments. We will propose new models for the generation and modeling of coral islands and karst networks.

In the third part, we will focus on adding realism to 3D terrains through physical simulations of erosion processes. We will demonstrate a flexible and controllable method to imitate the long-term effects of water and wind on both terrestrial and marine landscapes.

Keywords: terrain representation, procedural generation, physical simulations, user interaction

CHAPTER 1

Introduction

Over the past 50 years, terrain generation has emerged as an increasingly active domain inside the field of computer graphics. As the demand for more realistic, faster, and automated processes has grown, terrain generation techniques have evolved to meet these challenges. As these objectives get progressively achieved, a new trend has started to emerge in which focus is pushed toward user control. The focus of this work is on one specific branch of terrain generation: the interactive creation and modeling of landscapes.

[TODO]

What sets this work apart is its emphasis on underwater landscapes or "seascapes", an area still overviewed but with an important relevance in marine biology, oceanography, and underwater robotics. Furthermore, the extension of these techniques to new areas of the entertainment industry, such as video games and cinema, presents new opportunities for innovation.

[TODO]

The central question guiding this research is: "How can we efficiently guide the user in the creation of virtual content along the production process line to maintain as much control as possible over the final product?" This concept of "guiding" rather than "replacing" the user is, from my point of view, fundamental, as no machine can truly know better than a human what the final product should look like. The goal is to present algorithms that can be used flexibly within a production pipeline, adapting to many terrain representation, fluid solver, landscape type, user's hardware, or objective, whether the final use is real-time rendering, realism, or animation.

Maintaining control over the creative process is essential. Each generated result should feel unique, meet the user's expectations, be explainable, and be easily correctible without requiring a complete regeneration. This approach ensures that the user remains at the center of the creative process, with the tools and algorithms serving to enhance their objectives, rather than replace them.

1.1 Contributions and outlines

This research follows a chronological order of terrain generation, beginning from the development of an abstract representation that bridges the gap between computer science

and Earth science. This approach offers a generalization of the desired landscape type, with a particular focus on underwater environments, but also applicable to terrestrial landscapes.

A second key contribution is the proposal of new types of landscape features, including karst networks and coral islands, which maintain the notion of sparseness through the use of implicit volumes.

Finally, a new particle-based erosion simulation method is introduced. This method is agnostic to the terrain representation, making it lightweight, fast, and easy to implement.

The overarching aim is to provide maximum control to the user, not only during the generation process but also in making corrections to details upstream, ensuring that the final product aligns with the user's vision.

1.1.1 Semantics

[TODO] This interdisciplinary approach enriches the modeling process, ensuring that the generated landscapes reflect real-world biological and geological phenomena.

1.1.2 Modeling

[NOT GOOD AT ALL, TO REDO]

The generation of certain landscape elements, such as karst networks and coral islands, remains relatively new. While karsts have been explored in previous research (Axel Paris), the representation of coral islands, particularly in the context of Darwin's theories, is an innovative contribution. Karst networks are represented in a highly user-friendly manner, using a directed acyclic graph structure, which closely resembles a tree structure. This method is enhanced by a fractal generation approach with cycles, allowing for iterative generation of complex landscapes. The modeling of coral islands is based on both historical observations and travel journals.

1.1.3 Amplification

To increase the realism of the generated landscapes, details are added through a particle-based erosion method. This method is designed to be generalizable for flexibility, and its implementation is oriented towards speed and parallelization.

These contributions collectively aim to advance the field of terrain generation, providing new tools and methods that offer both flexibility and control to users, while also pushing the boundaries of what is possible in the simulation and modeling of natural landscapes.

Due to the disparities of works for the different topics proposed in this thesis, we will introduce the topic, present a state of the art and develop our methods in each chapter.

1.2 Procedural generation

Procedural generation is a powerful technique for creating data algorithmically, rather than manually. This method is massively used in areas such as computer graphics, simulations, and game development. Essentially, it involves using predefined rules or algorithms to generate complex structures or systems. For example, it can be employed to create music, textures, or virtual ecosystem.

The history of procedural generation can be traced back to the mid-20th century, rooted in mathematical and algorithmic theories. During this period, foundational concepts such as

randomness, fractal geometry, and noise functions were introduced, laying the groundwork for modern procedural techniques.

One of the main benefits of procedural generation is its ability to generate content with minimal storage requirements. Content is typically generated in real-time or on-the-fly, rather than being explicitly stored, which allows for the creation of large amount of content without requiring significant storage space. Additionally, procedural generation is also advantageous for producing diverse content quickly, enabling the creation of vast amounts of data that may require only minor adjustments or improvements. In practical applications, procedural generation is widely used across various fields to create dynamic, diverse, and detailed content efficiently. In video games, it enables the automatic creation of expansive game worlds, complex level designs, and varied character animations. For example, procedural generation can produce entire game levels with unique layouts for each playthrough, enhancing replayability, or generate diverse character behaviors that adapt to in-game environments, making interactions more dynamic and lifelike.

Beyond video games, procedural generation plays a significant role in other industries as well. In the film industry, it is used to generate intricate visual effects, such as realistic smoke, fire, and water simulations, which would be labor-intensive to create manually (Fedkiw and Jensen, 2001; Hädrich et al., 2021; Stam, 2003b; Xing et al., 2016). Facial animation also benefits from procedural techniques, where algorithms can generate subtle, lifelike expressions that enhance the realism of CGI characters [ADD REFERENCES]. Crowd simulations are another critical application, with procedural methods used to populate scenes with thousands of unique, animated characters, each with distinct appearances and behaviors (Narain, 2011; Park, 2010; Yersin et al., 2009).

In simulations, procedural generation is integral to creating accurate and scalable representations of real-world environments. This includes the generation of complex ecosystems for environmental simulations, detailed urban layouts for city planning, and even diverse astronomical phenomena for space simulations [ADD REFERENCES]. Additionally, procedural generation allows researchers to isolate and control variables within these simulations, enabling them to test different scenarios and observe outcomes in a controlled, repeatable manner.

The integration of algorithms and data is a key aspect of procedural generation. Mathematical models and noise functions, such as Perlin or Simplex noise, are often used to create the foundational structures in procedural content. These algorithms can generate everything from the random yet cohesive textures of terrain to the complex, non-repetitive patterns seen in natural phenomena or digital art (Vergne et al., 2011). This might involve generating natural features like landscapes, complex textures and mesostructures on textile objects (É. Michel and Boubekeur, 2023), or even entire ecosystems. Incorporating elements of real-world phenomena, such as erosion patterns in terrain generation, can make the environments more believable and interactive [REPLACE WITH EROSION PATTERNS ON OBJECTS GENERATION I.E. CRACKS, BUMPS, MELTING].

The flexibility of these algorithms not only allows for the creation of complex content but also enables user-driven customization that can influence or guide the content generation process, leading to user-specific outcomes. This feature, combined with the scalability and efficiency of procedural methods, means that large amounts of data can be produced with relatively low computational and storage costs compared to manually crafted content. A urban layout can, for example, be constraint to certain areas (Zhou et al., 2021), to satisfy inhabitants accessibility needs (Lima et al., 2022), to minimize energy consumption (Shi et al., 2017) or even solar irradiations (Vermeulen et al., 2015).

Procedural generation can be categorized into two main types: deterministic and stochas-

tic systems.

Deterministic systems produce outputs that are entirely predictable and repeatable given the same initial parameters and input conditions. This means that running the same algorithm with identical inputs multiple times generates the exact same output. Such predictability is useful in scenarios where consistency and reliability are important. For example, in engineering simulations, deterministic systems ensure that specific results can be consistently reproduced, which is essential for validation and testing. Similarly, in video game development, deterministic algorithms might be employed to generate identical game worlds across different sessions, ensuring that all players experience the same environment under the same conditions, and that no unexpected obstacle blocks the character. This consistency is also beneficial in content creation pipelines, where artists or designers need to fine-tune specific aspects of generated content. Since the output is repeatable, they can iteratively adjust parameters and observe how these changes affect the final result, making it easier to achieve the desired outcome.

On the other hand, stochastic systems introduce elements of randomness into the generation process, resulting in varied outputs even when the same initial parameters are used. This randomness is particularly important in applications where variation and uniqueness are desired. In video games, for instance, stochastic systems might be utilized to create a wide variety of levels, landscapes, or in-game items, ensuring that each playthrough feels different and offers a new experience. Beyond that, stochastic processes are also used in fields like procedural music generation, where each composition can be unique, providing a unique listening experience every time.

Recognizing the strengths and limitations of both deterministic and stochastic systems, many applications integrate these approaches into hybrid systems. In such systems, deterministic algorithms might be used to establish the overall structure, ensuring consistency and adherence to specific rules or constraints. Then stochastic elements are introduced to fill in details and add variability, creating dynamic content that remains coherent. For example, in procedural architecture, a deterministic system might generate the building's foundational layout, while stochastic processes add varied interior details, making each design unique yet structurally coherent. This hybrid approach can provide a balance between consistency and variability, making it possible to create content that is both reliable and dynamic.

In rule-based systems, content is generated by following a set of predefined rules, which can range from simple conditions to complex algorithms. These rules dictate how different elements interact, ensuring that the output adheres to specific constraints. For instance, in procedural city generation, rules might govern the placement of buildings, roads, and parks to create a realistic urban layout. In this way, rule-based systems ensure that the generated content is both structured and coherent. Other examples of rule-based systems, in game design, might be used to generate levels that adhere to specific gameplay mechanics, such as ensuring that obstacles are placed in a way that challenges the player without making the game impossible. In architecture, rules might dictate the structural integrity of a building, ensuring that generated designs are not only aesthetically pleasing but also feasible.

(Temuçin et al., 2020)

Typically, rule-based systems involves iterative processes, where initial outputs are continuously refined or adjusted based on additional rules or feedback. This iterative approach allows for the progressive enhancement of the content, where each cycle of generation can introduce new elements or improve upon previous iterations, resulting in a more polished and coherent final output. Those systems are mainly considered deterministic but including a slight randomness may increase the variability and provide a more natural feeling.

Procedural generation significantly help reducing manual efforts, which becomes un-

avoidable for the creation of large assets or a diversity of variations of content. Including control over the algorithms in play allows the user to adapt the results to its needs while minimizing the required storage.

1.2.1 Notable examples in the entertaining industry

As procedural generation techniques matured, they found early applications in video games and interactive media. One of the first notable examples was the game *Rogue* (1980), which utilized procedural generation to create its dungeon levels. In *Rogue*, each dungeon was generated anew each time a player started a game, ensuring that no two playthroughs were alike. This approach not only enhanced replayability but also minimized the amount of storage needed, as the game did not store pre-made levels but rather generated them on-the-fly.

Another milestone in procedural generation was the game *Elite* (1984), which utilized procedural algorithms to generate a massive universe containing 2048 unique planets, each with its own name, economy, and location in space. This vast universe was generated using only about 20KB of memory, thanks to the efficiency of the procedural techniques employed. The planets' attributes were generated using deterministic algorithms, ensuring that each planet's characteristics were consistent across different game sessions.

Dwarf Fortress (2006) took procedural generation to new heights by generating not just terrain, but entire worlds, complete with histories, civilizations, and ecosystems. The game uses complex algorithms to simulate thousands of years of history, resulting in a deeply detailed and dynamic world. The procedural generation in *Dwarf Fortress* goes beyond mere randomization; it involves creating a coherent and interconnected world where every element, from geography to political structures, is generated algorithmically.

Spore pushed the boundaries of procedural content generation by procedurally generating creatures, planets, and even entire galaxies. Unlike traditional games, *Spore* did not store textures, music, or animations. Instead, it generated these elements on-the-fly, allowing for an almost infinite level of variety and creativity.

Minecraft (2011), revolutionized procedural generation in gaming by creating vast, open worlds composed of blocks, with different biomes, caves, and structures all generated procedurally. The game's world is generated using a combination of Perlin noise and other algorithms to create varied and dynamic landscapes that players can explore indefinitely. *Minecraft*'s use of procedural generation allowed for an infinite world size, limited only by the computing power available.

Procedural generation's impact extends beyond gaming into simulations, scientific research, and computer graphics. In scientific simulations, procedural techniques are used to generate terrains for geological studies, allowing researchers to model and analyze various scenarios, such as modeling natural disaster (Hädrich et al., 2021; Hudák and Ďuríkovič, 2011; Valette et al., 2005), animal behaviours (Ecormier-Nocca et al., 2021; Wampler and Popovi, 2009), robot simulations (Lejeune, 2021; Louis et al., 2019), military simulations (Dam et al., 2019), etc... Procedural generation is also used in fluid dynamics simulations to create realistic water flows, weather patterns, and other environmental phenomena.

In computer graphics and animation, procedural generation is essential for creating complex visual effects, such as realistic landscapes, textures (Gobron and Chiba, 2001), and particle systems in movies. Procedural methods enable the efficient generation of vast and detailed environments that would be impractical to model manually. For example, in movies like *The Lord of the Rings* (2001) and *Avatar* (2009), procedural techniques were used to generate large natural landscapes, facial animations, crowds simulations, etc.

1.2.2 Parallelization and GPU utilisation

The exponential increase in computing power over the past few decades allowed exponential advancements in procedural generation, particularly for real-time content creation and complex simulations. The evolution from single-core processors to multi-core CPUs (Central Processing Unit), and more significantly, the rise of powerful Graphics Processing Units (GPUs) capable of massive parallel processing, has transformed how procedural algorithms are implemented.

GPUs are perfect for procedural generation due to their architecture, which allows them to execute thousands of threads in parallel. Unlike CPUs, which are optimized for sequential processing and are typically limited to a few cores, GPUs contain thousands of smaller, simpler cores designed for handling multiple simple tasks simultaneously. This parallelization capability is essential for procedural generation, where large data grids need to be processed efficiently.

In procedural generation, many operations can be performed independently on different parts of the data, making them ideal candidates for parallel execution. For example, when generating a large terrain, each point on the terrain grid can be computed independently based on noise functions or subdivision algorithms. By distributing these computations across many parallel threads, GPUs can dramatically accelerate the generation process, allowing for real-time updates and interactions in complex environments.

GPGPU (General-purpose Processing on GPU) refers to the use of GPUs for performing computations that are traditionally handled by CPUs. Originally designed for rendering graphics, GPUs have evolved to support general-purpose computing tasks through frameworks like CUDA (Compute Unified Device Architecture) from NVIDIA and OpenCL (Open Computing Language). These frameworks allow developers to write programs that can harness the parallel processing power of GPUs for a wide range of computational tasks beyond just graphics rendering.

In procedural generation, GPGPU is employed to accelerate various algorithms that require intensive computation. For example, noise functions like Perlin or Simplex noise can be computed in parallel across the entire terrain grid or texture space, enabling the real-time generation of complex patterns and surfaces. Or techniques such as the Diamond-Square algorithm or midpoint displacement can be implemented on the GPU, where each refinement step of the grid can be processed simultaneously for different segments of the terrain. Finally, procedural simulations of fluid dynamics, particle systems, and other physical phenomena can be executed on the GPU, where the interactions of thousands or even millions of particles are computed in parallel, resulting in realistic, real-time simulations.

The integration of machine learning, particularly deep learning techniques, with procedural generation represents a new step forward in the domain. Machine learning models, such as Generative Adversarial Networks (GANs) and Variational Autoencoders (VAEs), have been leveraged to enhance the quality and diversity of procedurally generated content.

GANs are composed of two neural networks, a generator and a discriminator, that are trained together in a competitive manner. The generator creates new data samples, while the discriminator evaluates them against real data. Through this process, the generator learns to produce increasingly realistic outputs. In procedural generation, GANs can be trained on datasets of real-world terrains, textures, or architectural styles, and then used to generate new, highly detailed and realistic content that closely mimics the training data. For instance, GANs can create entire landscapes that are indistinguishable from real-world environments, adding a layer of realism that traditional procedural techniques.

VAEs are another type of deep learning model used in procedural generation. They work

by encoding input data into a latent space, where similar inputs are mapped close to each other, and then decoding from this space to generate new data. VAEs are particularly useful for generating content that needs to maintain a certain level of coherence or adhere to specific stylistic constraints. For example, VAEs can generate new character models or textures that fit within a desired aesthetic, providing a high degree of control over the final output.

[TALK ABOUT TRANSFORMERS ALSO]

1.3 Procedural terrain generation

1.3.1 Terrain representations

Terrain refers to the physical features and configuration of a specific area of land. It includes the elevation, slope, and the overall topography, such as mountains, valleys, and plains. Terrain is often used to describe the surface characteristics of the land, focusing on the natural contours and the geographical aspects that define a region's physical form.

While the term "terrain" describes the physical characteristics of land, it does not include the natural elements that shape an area's identity. Elements like vegetation, water bodies, and climatic conditions, such as snow cover, are essential to how we perceive and understand a landscape. Therefore, when discussing procedural generation in virtual environments, "landscape generation" is a more fitting term, as it integrates these natural elements along with the topographical features.

In addition to "terrain generation," other terms such as "landscape generation," "world generation," and "environment generation" can be used to describe the creation of virtual landscapes. These terms are interchangeable and can all refer to the process of generating physical terrain along with natural and artificial elements. However, by convention and for simplicity, the term "terrain generation" is most commonly used in the field. Despite its original focus on the physical features of the land, "terrain generation" has evolved to encompass a broader range of environmental elements, making it a convenient and widely accepted term for describing the comprehensive process of creating virtual environments.

A terrain can be represented in various ways, each of them suited for a given application of which we give an brief overview, more details can be found in (E. Galin et al., 2019).

Elevation models

Elevation models are a fundamental approach in terrain representation, widely used in procedural generation due to their simplicity and efficiency. These models define the terrain as a function $h : \mathbb{R}^2 \mapsto \mathbb{R}$, where each point in a 2D plane is mapped to an elevation value. This approach is particularly effective for representing terrains where the elevation is the only varying factor, such as hills, valleys, and plateaus, and it is best suited for terrains without complex 3D features like overhangs or caves. While we visualize elevation models in three dimensions, they are mathematically considered two-dimensional functions. In the domain of terrain generation, we will name them 2.5D models.

Elevation models are widely used in industries where large-scale terrain representation is crucial. In video games, they provide the foundation for creating vast open-world environments. In geographic information systems (GIS) and remote sensing, height fields are used to represent real-world terrain data, offering a practical means of visualizing and analyzing geographical features. The ability to manipulate and control terrain features procedurally makes elevation models a common choice for applications that require efficient terrain generation and rendering.

They offer a powerful method for representing terrains in procedural generation, combining simplicity with flexibility. While they have limitations in representing complex 3D structures, their efficiency and compatibility with existing algorithms make them indispensable in a variety of applications.

Implicit height fields

Implicit height fields represent the terrain as a mathematical function that provides a height value at any given point in the domain. These functions can be procedural or closed-form expressions, allowing for compact storage and infinite precision in theory. The elevation function allows for easy manipulation of terrain features, making it ideal for generating terrains that require smooth, continuous surfaces. However, the primary disadvantage is the computational complexity involved in evaluating the function, especially for large or highly detailed terrains. The challenge lies in constructing functions that can realistically represent large-scale terrains with complex landforms.

Discrete height fields

Discrete height fields, or explicit height fields, are one of the most prevalent methods for terrain representation. These models consist of a 2D grid where each cell contains a height value, representing the elevation at that point. Height fields are particularly advantageous because they are simple to implement and are directly compatible with many rendering techniques and hardware, but also due to their closeness with image processing, a domain studied for many decades now.

The main advantage of height fields is their ability to handle large datasets efficiently, providing a balance between memory usage and detail. However, they are limited by their inability to represent terrains with overhangs or caves, as each point on the grid can only hold a single elevation value. Additionally, height fields often require interpolation methods, such as bi-linear or bi-cubic interpolation, to reconstruct a continuous surface from the discrete grid points.

Volumetric models

Volumetric models represent a more complex approach to terrain modeling, allowing for the depiction of 3D features that go beyond the simple surface-based representation provided by elevation models. These models capture not only the surface of the terrain but also its internal structure, making them ideal for representing terrains with overhangs, caves, and other subsurface features.

Volumetric models, including layered materials, voxel grids, and implicit models, are essential in applications where terrain complexity and detail are primordial. In geological simulations, these models allow for accurate representation of subsurface structures and processes. Voxel models are widely used in games that require dynamic terrain deformation, providing a rich interactive environment for players. Implicit models are favored in situations where smooth, continuous surfaces are needed [FIND OTHER USE CASES].

Implicit volumetric models

Implicit volumetric models describe the terrain's shape and features using an implicit function. The terrain is represented by a mathematical function $f : \mathbb{R}^3 \mapsto \mathbb{R}$ that determines the terrain surface by evaluating to an isovalue, often zero. This function provides a continuous representation of the terrain, with points inside the terrain returning positive values and while points in the air evaluate to negative values. It allows for the seamless representation

of complex terrain features, including caves, overhangs, and varying geological structures, which are impossible to represent with elevation models.

One of the key advantages of implicit models is their ability to produce smooth surfaces without the need for discrete polygonal meshes, which can result in realistic and natural-looking terrains. However, the computational complexity of evaluating the implicit function, especially for large terrains, can be a significant drawback. Additionally, converting an implicit surface into a mesh for rendering can be challenging and resource-intensive. (de Araújo et al., 2015)

Layered models

Layered models are a type of volumetric representation that encode different material layers within the terrain and are defined by a function $\mu : \mathbb{R}^3 \mapsto \mathcal{M}$, where \mathcal{M} denotes the material type at any given point in 3D space. This allows for a detailed representation of the terrain's internal composition, which can be crucial for applications requiring realistic geological simulations. Each layer is defined by its thickness or elevation, and multiple layers can be stacked to represent complex geological formations. These layers might include materials like bedrock, sand, soil, or water, each contributing to the overall structure of the terrain. Layered models are particularly useful in simulations that involve processes like erosion or sedimentation, where the interaction between different material layers affects the physical process.

The primary advantage of layered models is their ability to represent a stratified terrain with distinct material properties, which can be manipulated individually. This makes them well-suited for simulations that require detailed geological accuracy. However, they are more complex to implement than simple elevation models and require additional computational resources to manage the interactions between layers.

Voxel grid models

Voxel grids are a common method for representing 3D terrains in procedural generation, offering the ability to capture complex internal structures and features that are difficult or impossible to represent with surface-based models. In a voxel grid, the 3D space is divided into a regular grid of small, cube-shaped elements called voxels (volumetric pixels). Each voxel holds information about the material or properties of the terrain at that specific point in space. This approach allows for detailed modeling of features such as caves, tunnels, overhangs, and intricate underground networks. The regular grid structure allows for the use of image processing-oriented algorithms.

There are three primary types of voxel grids used in terrain representation: binary voxel grids, material voxel grids and density voxel grids. Each has distinct characteristics, advantages, and limitations, making them suitable for different applications.

Binary voxel grids Binary voxel grids are the simplest form of voxel representation. In these grids, defined $f : \mathbb{R}^3 \mapsto [0, 1]$, each voxel is either "filled" or "empty," representing the presence or absence of material. This binary state is typically represented by a 1 (filled) or 0 (empty). Binary voxel grids are straightforward to implement and require much less memory compared to more complex voxel representations, making them ideal for applications where the primary concern is whether a space is occupied or not.

The simplicity of binary voxel grids is one of their main advantages. They are easy to understand and visualize, with each voxel requiring only a single bit of information to represent its state. Additionally, because only a binary state is stored, these grids can be memory-efficient when combined with compression techniques like Sparse Voxel Octrees (SVOs) (Laine and Karras, 2010) or voxel Directed Acyclic Graphs (DAG) (Careil et al., 2020;

Villanueva et al., 2017). The simplicity of the data structure also allows for quick processing, making binary voxel grids suitable for real-time applications where performance is required. However, the binary nature of these grids limits their ability to represent variations in material density or properties, or even smoothness, resulting in less detailed terrain models. This can lead to hard, blocky edges in the terrain, which may appear unnatural without additional smoothing or processing.

Material voxel grids Material voxel grids, defined as $\mu : \mathbb{R}^3 \mapsto \mathcal{M}$, are commonly used in applications where simple occupancy information is sufficient. For example, voxel-based games like Minecraft utilize material grids to create terrains composed of solid blocks with clear boundaries. These grids are also employed in scientific simulations where the primary concern is the presence or absence of materials, rather than detailed material properties.

Density voxel grids Finally, density voxel grids allow each voxel to store a range of values, representing varying degrees of material presence with $f : \mathbb{R}^3 \mapsto \mathbb{R}$. Instead of a simple discrete state, a density voxel grid assigns a continuous value to each voxel, which can represent material density, opacity, or other properties. This added complexity enables density voxel grids to represent subtle variations in terrain, such as gradual changes in material density or smooth transitions between solid and empty spaces, allowing for more realistic and natural-looking terrain models.

Density voxel grids are often used in high-fidelity simulations where detail and realism are essential. They are found in applications such as medical imaging, scientific visualizations, and advanced terrain modeling for films and visual effects. These grids are also employed in procedural terrain generation systems that require smooth and natural transitions between different terrain features, such as caves, cliffs, and eroded landscapes.

1.3.2 Terrain generation

Procedural generation encompasses a diverse range of techniques used to create complex, natural-looking content, particularly in terrain generation. These techniques span from mathematical models to advanced simulations and can be broadly categorized into methods for large-scale terrain generation and procedural landform generation.

Noise functions are foundational to procedural terrain generation, providing the basis for creating coherent, natural patterns across terrains. Perlin noise, developed by Ken Perlin in 1983, is a gradient-based noise function that produces smooth, continuous patterns, ideal for generating natural-looking textures and landscapes. Simplex noise, introduced by Perlin in 2001, improves upon Perlin noise by being more computationally efficient and reducing directional artifacts, especially in higher-dimensional spaces. Worley noise, another noise variant, generates patterns based on the distance between points, producing cellular structures useful for simulating natural textures like stone or marble, and for modeling phenomena like cloud formations. Fractional Brownian motion (fBm) builds on these noise functions by summing noise at different scales and amplitudes, creating terrains with varied features, from gentle hills to rugged mountains. Techniques like ridge noise focus on generating sharp features such as crests and ridges, while domain warping applies distortions to prevent grid artifacts and simulate erosion effects without the need for full physical simulations.

Cellular automata offer a rule-based approach to procedural generation, particularly useful for simulating complex systems and natural processes. Originating in the 1940s, these grid-based models evolve based on rules applied to neighboring cells. In terrain generation, cellular automata are commonly employed to simulate cave systems, forests, and other structured environments. Notable examples include Conway's Game of Life, which

demonstrates how simple rules can lead to complex, self-organizing patterns, and Langton's Ant, which shows how basic rules can produce intricate behaviors. Cellular automata can be generalized to continuous time and space fields, making them versatile tools for modeling a wide range of natural phenomena, from fluid dynamics (Boldea, 2009; Cattaneo and Jocher, 2005; Menshutina et al., 2020) to vegetation patterns (Greene, 1989).

Large-scale terrain generation involves creating expansive terrains that exhibit natural-looking landscapes over large areas. The main method for generating a large-scale terrain is the employ of the noise functions, but other techniques like subdivision schemes iteratively refine an initial coarse terrain by subdividing it and adding fractal detail. While effective in generating self-similar terrains, these methods may introduce artifacts such as directional inconsistencies, which require further refinement. Faulting is another technique, simulating tectonic activity by introducing random vertical faults into a flat terrain. This method creates realistic elevation changes by displacing points on either side of the fault line, with smooth step functions ensuring gradual transitions between displaced and non-displaced areas. Both methods are often used for generating the broad, natural features characteristic of large-scale terrains, although they may lack the precision needed for detailed landform creation. Large-scale methods are mostly defined in 2.5D. Gravity playing a large role in the shape of such environment, the surface of the terrain contains few concavities. Such cases, like caves, arches or overhangs, are mainly considered in the feature-scale.

Procedural landform generation can be focused on creating specific terrain features, such as rivers, cliffs, and canyons, with greater precision and control. Controlled subdivision enhances traditional fractal methods by integrating user-defined features, such as river networks, directly into the terrain. This allows for the creation of landscapes that adhere to specific topological and hydrological requirements. Feature-based construction techniques further refine this approach by using interactive sketch-based interfaces, where users can define terrain features through control curves. These curves guide the multi-resolution deformation process, enabling the creation of detailed and complex landforms like layered terraces and isolated mesas. By blending different types of curves, such as elevation profiles and cross-sectional shapes, users can produce terrains that are both realistic and tailored to specific needs.

Neural networks represent a more recent advancement in procedural generation, particularly in terrain and texture generation. Generative models like Generative Adversarial Networks (GANs) and Variational Autoencoders (VAEs) are used to create realistic terrains and photorealistic images. GANs operate by having two neural networks (a generator and a discriminator) compete to produce outputs that are increasingly realistic. VAEs, on the other hand, encode and decode data to and from a latent space, generating new, similar instances of terrains or textures. These models are powerful tools for adding detail and variety to procedurally generated environments, enhancing the realism and complexity of the final output.

Physical simulations use models of real-world processes to generate and refine natural features in terrains. These include simulations of fluid dynamics, erosion, sediment transport, and other geological and environmental processes. While these simulations are often used to improve the realism of terrains rather than for initial creation, they are crucial for adding realistic details such as river erosion, weathering, and sediment deposition. Erosion models mimic natural processes like water flow and wind, shaping the terrain over time, while sediment transport models simulate the movement and deposition of sediment, contributing to the natural evolution of the landscape.

By combining these diverse techniques, procedural generation enables the creation of vast, detailed, and realistic terrains with minimal input data. This approach balances compu-

tational efficiency with the complexity required to simulate natural landscapes, making it a powerful tool in terrain modeling and simulation. Whether generating expansive landscapes or intricate landforms, procedural generation techniques provide the flexibility and precision needed to create dynamic and believable environments across various applications, from video games to scientific simulations.

1.3.3 User interaction [TODO]

User interaction affects how users balance realism, speed, and control over generated content.

Realism refers to the extent to which generated content accurately represents real-world characteristics, such as visual details, physical processes, and natural patterns. This is especially important in simulations, visualizations, and training environments where plausibility [FIND BETTER TERM] is critical. Techniques to enhance realism include physical simulations, which model processes like erosion and sediment transport, and the integration of expert knowledge from fields such as geology and ecology. However, achieving realism can be challenging due to the complexity of detailed simulations and the need for specialized expertise, which can demand substantial computational resources and inputs.

Speed is about the efficiency of content generation within acceptable time constraints. While no official categorisation has been set, James Gain proposed to describe levels of speed based on response time:

- Real-time: generation in less than 30 milliseconds, essential for interactive applications like VR environments.
- Interactive: generation in under 3 seconds, suitable for user-driven customization in games and simulations.
- Near-interactive: generation in less than 5 minutes, applicable for larger-scale simulations where some delay is acceptable.
- Long-term: generation that takes longer, often used for precomputed content or offline rendering.

Optimizing speed involves using efficient algorithms and parallel processing. Almost all recent works achieve fast execution time thanks to high parallelisation on GPU. Other types of algorithm may rely on a refinement paradigm, generating a coarse result at first and then iteratively adding finer details, such that the global output shape is available long time before the real final result.

Control refers to how much users can influence or direct the procedural generation process to meet their specific needs or preferences. This includes parameters fine-tuning, allowing users to modify parameters like the noise functions parameters or the simulation parameters, and artistic control tools, made for artists and designers, to guide the generation process with the use of masks and brushes, allowing for combination and mixing of different algorithms.

Managing diverse and sometimes conflicting user expectations, such as balancing creative freedom with the demand for realistic outcomes, requires careful consideration. Additionally, it is essential to balance the complexity of procedural systems to ensure they remain user-friendly, avoiding overwhelming users or producing results that feel artificial or inconsistent.

1.3.4 Fluid simulations

Fluid simulations are an essential component of procedural terrain generation, particularly for modeling the behavior of water and its interactions with the terrain. These simulations are crucial for creating realistic and dynamic environmental models that accurately reflect natural phenomena, such as river flow, coastal erosion, sediment transport, and overall hydrology.

At the core of fluid simulations are the Navier-Stokes equations, which describe the motion of fluid substances. These equations account for various forces acting on a fluid, such as pressure, viscosity, and external forces like gravity.

$$\rho \left(\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} \right) = -\nabla p + \mu \nabla^2 \mathbf{u} + \mathbf{f} \quad (1.1)$$

Where $\mathbf{u} = (u, v, w)$ is the fluid velocity vector, with components u , v , and w in the x -, y -, and z -directions, respectively, t is time, ρ is the fluid density, p is the pressure field within the fluid, μ is the dynamic viscosity of the fluid, ∇p is the pressure gradient force, $\mu \nabla^2 \mathbf{u}$ is the viscous diffusion term, representing the effects of internal friction within the fluid and finally, \mathbf{f} represents external body forces, such as gravity.

For an incompressible fluid like water, the Navier-Stokes equations can be simplified by assuming that the fluid density remains constant, leading to the incompressibility condition:

$$\nabla \cdot \mathbf{u} = 0 \quad (1.2)$$

This condition ensures that the volume of fluid elements remains unchanged as they move through space, which is crucial for accurately simulating water flow.

The Navier-Stokes equations are highly non-linear, which means that even small changes in the input conditions can lead to significant and unpredictable changes in the fluid's behavior. The non-linearity makes solving these equations a expensive computational problem, as it requires iterative methods to find stable solutions. Each step of the simulation must balance the forces acting on the fluid, such as pressure and viscosity, while ensuring that the solution remains physically accurate and stable over time. This process becomes particularly demanding when aiming for realistic, high-resolution simulations where fine details like turbulence and surface tension must be accurately captured. In this case we require dense computational grids or a large number of particles, and these elements must be updated frequently to maintain the realism of the simulation, which exponentially increase the computations and the memory used.

The computation cost is further amplified when simulations are performed in 3D space. Unlike 2D simulations, where calculations are confined to a plane, 3D simulations must account for the full complexity of fluid movement in all directions. This increases the number of computations required, making real-time applications like as video games and interactive simulations almost incapable to use them.

Different fluid solvers have been developed to approximate the solutions to the Navier-Stokes equations, each with its strengths and weaknesses. These solvers vary in how they represent the fluid (the Lagrangian approach using particles or the Eulerian approach with discrete grids, or a combination of both) and how they handle the computational trade-offs between accuracy, stability, and performance. The choice of a fluid solver depends on the specific requirements of the simulation, such as the need for real-time performance, the level of detail required, or the types of fluid behavior being modeled.

Marker-And-Cell (MAC) Method

The Marker-And-Cell (MAC) method is one of the earliest and most fundamental grid-based techniques for simulating incompressible fluid flows. In the MAC method, the fluid's velocity components are stored at the faces of grid cells, while pressure values are stored at the cell centers. Marker particles are used to track the fluid's free surface, ensuring that fluid interfaces and boundaries are accurately captured. The MAC method excels at maintaining the incompressibility condition of fluids and accurately modeling pressure fields, which are important for realistic fluid dynamics. Because of its emphasis on accuracy and detailed pressure modeling, MAC is more oriented toward realism, especially in scenarios where the precise behavior of fluids is essential. However, its grid-based nature can lead to challenges in handling complex geometries and fine details at fluid boundaries, as the resolution is limited by the grid size. Moreover, the method can be computationally intensive, especially for high-resolution grids, making it less ideal for real-time applications.

Stable Fluids

Building on the grid-based approach of the MAC method, the Stable Fluids method addresses key stability issues that arise in fluid simulations (Stam, 1999). Stable Fluids uses a semi-Lagrangian advection scheme and implicit solvers to achieve stability even with large time steps, which is particularly advantageous in real-time applications like video games or interactive simulations. This method is designed with performance in mind, allowing for the simulation of fluid motion without the numerical dissipation that can degrade the accuracy of results over time, which was a common problem in other grid-based methods. While Stable Fluids prioritize performance and stability, particularly in real-time environments, the use of implicit solvers can introduce smoothing effects that reduce the sharpness of fine details in the fluid's motion. Additionally, the method's reliance on grid resolution still limits its ability to represent highly detailed surface interactions, making it a good compromise between realism and performance.

Particle-In-Cell (PIC) Method

The Particle-In-Cell (PIC) method represents a hybrid approach that combines the strengths of particle-based and grid-based methods. Fluid properties are stored on a grid, but the fluid's motion is tracked using particles, which carry velocity and position information only. The particles interact with the grid to update the fluid's velocity field, and the grid provides a stable mean for solving the fluid dynamics equations. PIC is particularly useful for capturing the large-scale motion of fluids, benefiting from the grid's stability while using particles to track detailed fluid behavior. However, the method leans more toward performance over accuracy due to its hybrid nature and ability to handle large-scale fluid movements efficiently. A major downside of PIC is its tendency toward numerical dissipation, where the fluid's kinetic energy is artificially reduced over time, leading to a loss of fine details, especially in turbulent flows. This dissipation occurs because the interpolation between particles and the grid tends to average out small-scale variations in velocity, making PIC less suitable for scenarios where high fidelity and detail are required.

Fluid-Implicit Particle (FLIP) Method

The Fluid-Implicit Particle (FLIP) method is an evolution of the PIC method designed to address its numerical dissipation problem (Brackbill et al., 1988). FLIP retains the grid-

particle hybrid approach but modifies how particle velocities are updated. Instead of fully relying on the grid's velocity field, FLIP updates particle velocities by applying only the changes in the grid's velocity field, preserving the fluid's kinetic energy and maintaining the detail in small-scale motions. This approach significantly reduces numerical dissipation, making FLIP more oriented toward realism, particularly in simulations that require detailed fluid dynamics like splashing, swirling, and other turbulent effects. However, FLIP is more computationally expensive than PIC and can suffer from instability issues, particularly when dealing with highly turbulent flows. The increased computational cost and the need for careful tuning of simulation parameters to maintain stability make FLIP less ideal for performance-focused applications but suitable for scenarios where high fidelity and detailed fluid behavior are essential.

Smoothed Particle Hydrodynamics (SPH)

Smoothed Particle Hydrodynamics (SPH) (Müller et al., 2003) is a purely particle-based method that models fluids using discrete particles, each representing a small volume of fluid. These particles interact with each other based on smoothing kernels, which define the influence of one particle on its neighbors over a certain distance (Koschier et al., 2022). SPH is particularly well-suited for simulating free-surface flows, such as waves, splashes, and other fluid phenomena where the interaction between fluid particles is complex and highly dynamic. Due to its ability to handle complex boundaries and fluid interfaces naturally, SPH is much more focused on realism, especially in scenarios that require accurate computation of fluid dynamics and interactions. However, SPH is computationally intensive, especially for high-resolution simulations where a large number of particles is required. Additionally, SPH can suffer from stability issues, such as particle clumping or excessive smoothing, which can detract from the realism of the simulation if not carefully managed. These factors make SPH better suited for applications where realism is prioritized over performance, particularly in high-fidelity simulations used in film and scientific research, but not for real-time applications.

We can find improvements (Roose et al., 2011), can represent may elements (Iwasaki et al., 2010), either fluid or sediments (Lenaerts and Dutré, 2009), so it is often used to simulate accurately water bodies (Nikeghbali and Omidvar, 2018).

Some other models exist, and are often proposed in OpenFOAM [CITE SIMPLE, OPEN-FOAM, ETC] (Caretto et al., 1973)

[BLA BLA Now some Neural networks, deep learning bla bla (Tompson et al., 2017) or some cellular automata (Boldea, 2009; Cattaneo and Jocher, 2005)]

1.3.5 Vegetation generation

Bla bla...

1.4 Underwater landscapes

- Use "seascape" instead of "landscape" and "aerial landscapes" for above water.

** Not official terms, but may be very useful for the rest of this thesis..!

- Describe main differences with landscape generation

- Describe coral islands ecosystems.

1.4.1 Main differences with aerial terrains

In the field of terrain generation, particularly when it comes to modeling underwater landscapes, the challenge is to accurately represent complex environments that include both geological and biological features. The underwater world is filled with diverse structures, ranging from vast coral reefs to intricate cave systems, each requiring specialized techniques to model their unique characteristics.

3D Data for underwater landscapes

One of the most challenging aspects of underwater landscape modeling is capturing the complexity of its structure. Taking the example of coral reefs, these environments feature highly intricate 3D structures, including branching corals, coral mounds, and encrusting forms. These formations are not just surface-level features; they extend into the terrain with voids and cavities, such as caves, grottos, and karst networks, which add layers of complexity to their representation. Accurate modeling of coral reefs demands that the porous and irregular nature of these formations is represented in high detail, which is a significant challenge due to the variability of their structures.

In addition to biological features, underwater landscapes are shaped by geological processes that must also be accurately represented. For example, sedimentary layers and underwater geological formations like seamounts and ridges are important components of the terrain.

To gather the data needed for such detailed modeling, advanced measurement techniques are employed. Sonar, including multi-beam echo sounders, and LIDAR (Light Detection and Ranging) are commonly used to capture detailed 3D data of underwater terrains. These technologies allow for the mapping of areas that are otherwise difficult to access, providing the foundational data necessary for accurate terrain generation. Remote sensing technologies also play a major role, enabling the collection of data over large and remote areas, further enhancing the model's accuracy, but represent a significant challenge as the provision of humans and hardware in these areas is far from easy to achieve.

Interdisciplinary data integration

Accurately modeling underwater landscapes requires more than just geological data; it necessitates an interdisciplinary approach that integrates biological and ecological data. Geological validation is an important step in this process, involving collaboration with geologists and marine scientists to ensure that the models are not only scientifically accurate but also reflective of real-world conditions. This validation process ensures that the generated models accurately depict the geological and biological features present in underwater environments.

Biological data is also essential, particularly when modeling coral reefs and other marine ecosystems. Incorporating data on coral species, marine biodiversity, and ecosystem dynamics allows for the creation of realistic and biologically accurate representations. Additionally, the impact of biological processes, such as coral growth patterns and marine erosion, must be considered, as these factors significantly influence the terrain shape over time.

However, one of the major challenges in modeling underwater landscapes is the scarcity of high-resolution data, especially in remote, protected or less studied areas. This data scarcity makes it difficult to create detailed and accurate models. Another challenge lies in the integration of diverse data types, geological, biological, hydrological, etc, into a cohesive and comprehensive model. Effective integration requires complex techniques to ensure that all aspects of the underwater environment are accurately represented.

Multi-scale modeling

Underwater landscapes are characterized by features that vary greatly in scale, from vast underwater mountains and ridges to small-scale elements like individual coral polyps and marine vegetation. This diversity in scale necessitates multi-scale modeling techniques that can accommodate both large and small features within the same model.

Paris, *Modeling and simulating virtual terrains* (2023) classified the different geological structures in four different spatial scales: the *megascale* describes landforms spread on more than 100km such as continents or mountain ranges, the *macroscale* between 1km and 100km like river or cave networks, the *mesoscale* ranging from 10m to 1km contains most complex structures (arches, ravines, cliffs, ...) and finally the *microscale* for features between 10cm and 10m, like sand ripples, ventifacts, rocks, etc. Generally, we consider as *large-scale* features that can be seen from far away and *small-scale* elements for which somebody has to be too close to observe it.

Large-scale features, such as underwater mountains and ridges, define the macrostructure of the landscape. These must be integrated seamlessly with small-scale features, such as detailed sedimentary textures and small coral formations, to ensure that the terrain is represented accurately across different scales.

[TODO]

The procedural generation of underwater landscapes involves the integration of advanced 3D data collection techniques, interdisciplinary collaboration, and multi-scale modeling approaches. Modeling underwater environments, from the complex structures of coral reefs to the accurate representation of geological and biological features, requires innovative solutions that combine geological plausibility with visual realism.

1.4.2 Coral reefs (biological aspects)

Coral reefs have intrigued humans for centuries, with early observations by ancient civilizations often misunderstanding these structures as mysterious sea plants or rocks. The study of coral reefs gained scientific momentum during the era of exploration, particularly through the work of figures like Captain James Cook and Charles Darwin. Darwin's theory of atoll formation, developed during his voyage on the HMS Beagle, was a significant milestone, proposing that atolls form around sinking volcanic islands as coral growth keeps pace with subsidence.

The 20th century brought technological advancements, such as scuba diving and sonar, which allowed scientists to directly observe and map coral reefs, deepening our understanding of their biology and ecological significance. In recent decades, tools like remote sensing and genetic analysis have further expanded our knowledge, highlighting the critical role of coral reefs in marine biodiversity and their vulnerability to environmental changes.

Anatomy of coral reefs

Coral polyps

Coral polyps are the essential building blocks of coral reefs. These tiny, soft-bodied organisms, just a few millimeters in diameter, form the vast structures of coral reefs. Each polyp consists of a sac-like body with a central mouth surrounded by tentacles equipped with stinging cells (nematocysts) for defense and capturing prey.

The body of the polyp is anchored to a hard surface, which it helps create by secreting calcium carbonate CaCO_3 . This secretion forms a protective exoskeleton around the polyp,

allowing it to attach securely to the substrate or to other polyps in a colony. This secretion process creates the coral's hard structure, which becomes the foundation of the reef.

Reproduction in coral polyps occurs in two main ways: asexual and sexual reproduction. Asexually, polyps reproduce through a process called budding, where new polyps form from the sides of an existing polyp. These new polyps remain attached, leading to the growth of a larger coral colony. In sexual reproduction, polyps release eggs and sperm into the water during spawning events, where fertilization occurs externally. The resulting larvae settle onto a suitable substrate, developing into new polyps and eventually forming new colonies.

Coral polyps are in a symbiotic relationship with zooxanthellae, a type of photosynthetic algae that live within the tissues of the coral polyps, providing them with vital energy through photosynthesis, fueling its growth and calcification.

In return, the coral provides the zooxanthellae with protection and access to sunlight. This mutualistic relationship is highly efficient, allowing corals to thrive in the nutrient-poor waters of tropical oceans.

However, this relationship is fragile. When corals are exposed to environmental stressors such as elevated water temperatures, they may expel their zooxanthellae in a phenomenon known as coral bleaching. Without the algae, the coral loses its primary energy source, leading to a whitening of the coral and, if conditions do not improve in the next weeks, eventual death.

Coral colonies

Coral polyps, though individually small and simple, have the remarkable ability to form large, complex colonies that create the structure of coral reefs. These colonies exhibit a variety of growth patterns, each of which plays a different role in the overall architecture and ecological function of coral reefs.

Once established, coral colonies can develop into a wide range of growth forms, each adapted to specific environmental conditions and with unique functions to contribute to the reef shape.

Massive corals, like those in the genus *Porites*, grow in boulder-like forms. They tend to grow more slowly than other types but are much more resistant to physical damage from waves and storms. They contribute to the long-term stability and resilience of coral reefs. Massive corals can live for hundreds of years, with some colonies growing to several meters in diameter. Their dense, solid structure provides a stable and enduring foundation for the reef, making them useful for the long-term stability of the reef ecosystem. Their robust skeletons can withstand strong wave action, storms, and other environmental stresses that would easily damage more delicate coral types. As they grow and expand, massive corals create the core of the reef, providing a solid base that other corals and marine organisms can build upon. Their presence is vital for maintaining the reef's overall integrity and resilience.

In contrast, branching corals, such as those in the genus *Acropora* (e.g., staghorn corals), look like a tree structure with numerous branches. These corals grow rapidly, extending outward to maximize exposure to sunlight. The dense thickets formed by branching corals provide essential habitat for a wide variety of marine species, from small fish to invertebrates. This growth form is particularly present in shallow, well-lit waters where competition for space and light is intense. This rapid growth allows them to quickly occupy space and expand the reef's surface area. The three-dimensional architecture of branching corals provides important habitats for a variety of marine life, including small fish and invertebrates, offering them protection from predators and strong currents. However, branching corals are relatively fragile and can be easily damaged by storms, waves, or human activity. Their ability to regenerate through fragmentation allows them to recover quickly from physical damage, but they do not contribute significantly to the reef's overall structural strength.

Similarly, foliose corals like in the genus *Astreopora*, which form leaf-like structures, add another dimension to reef complexity. These corals grow in overlapping layers, creating sheltered environments within the reef that are ideal for smaller marine organisms. Foliose corals can be particularly abundant in areas with moderate water movement, where their shape helps capture food particles from the water.

Corals that grow in flat, horizontal formations are known as table corals or plate coral, such as *Acropora*, can spread out to cover large areas, creating expansive, flat surfaces that provide important habitats for animals and other coral colonies. Their growth form allows them to efficiently capture sunlight, making them well-suited for shallow reef environments.

Some corals, known as encrusting corals like in the genus *Cyphastrea*, grow as a thin layer over hard substrates, including rocks, dead coral skeletons, and other surfaces. They are particularly adept at covering large areas of the reef, often outcompeting other organisms for space. Their growth form helps to bind loose sediments and rubble together, contributing to the stabilization of the reef's foundation. Although encrusting corals do not add significant height or volume to the reef, they are critical for cementing the reef structure, preventing erosion, and enhancing the overall stability of the reef. By filling in gaps between other coral colonies and consolidating the reef framework, encrusting corals help maintain the reef's structural cohesion, which is essential for its longevity.

The position of different coral types within the reef varies depending on factors such as water depth, wave exposure, and light availability. Massive corals are typically found on the reef slope, where the water is deeper and more stable. Encrusting corals thrive in the less turbulent waters of the deep reef, where they can spread across stable surfaces. Branching corals, such as *Acropora* species, are often found on the reef crest and fore reef, where they are exposed to strong wave action and plenty of sunlight. Their fast growth and ability to regenerate quickly make them well-suited for these dynamic environments. Foliose corals are often found on the reef flat and in the back reef areas, where the water is calmer and light levels are moderate. Table corals are also commonly found in shallow reef environments, where they can maximize their exposure to sunlight.

These varied growth patterns not only shape the physical structure of coral reefs but also create habitats that support immense biodiversity, making coral reefs, even if representing less than 0.1% of the ocean area, some of the most biologically diverse ecosystems on Earth, with about 25% of the marine species.

Reproduction

The formation and expansion of coral colonies are driven by a combination of sexual and asexual reproductive strategies, each of which plays a vital role in the growth, survival and spread of coral species.

Asexual reproduction is the primary mechanism by which coral colonies expand and repair themselves. The most common form of asexual reproduction is budding, where new polyps are produced from the body of an existing polyp. These new polyps remain attached to the parent polyp, leading to the growth of a larger, stronger, interconnected colony. The shared calcium carbonate skeleton secreted by these polyps provides the structural foundation for the entire colony.

Another important method of asexual reproduction is fragmentation, in which parts of the coral break off, often due to physical disturbances such as storms or strong currents, emit fragments with polyps that can then reattach to a suitable substrate and form new colonies further away. Species like *Acropora* spp. and *Pocillopora* spp. are known for their ability to reproduce through fragmentation, making them prolific in dynamic environments. Similarly, polyp bail-out, where individual polyps detach from the colony in response to environmental stress allows corals to survive adverse conditions and spread its colonies.

Species such as *Fungia* spp. and *Scleractinia* spp. have been observed to reproduce through polyp bail-out.

Sexual reproduction is essential for generating new colonies and ensuring genetic diversity within coral populations. Many coral species, particularly those that are broadcast spawners, engage in mass spawning events. During these events, numerous corals simultaneously release eggs and sperm into the water column, where fertilization occurs externally. The resulting larvae (planulae), are carried by ocean currents, sometimes over great distances, before settling onto a suitable substrate where they can develop into new polyps and establish a new colony. The precise synchronization of these spawning events, while not fully understood, is thought to be influenced by environmental elements such as lunar cycles, water temperature, and daylight length. Species such as *Acropora* spp. and *Porites* spp. are typical broadcast spawners, relying on this strategy to disperse their offspring and colonize new areas of the reef.

Some coral species are brooders, where fertilization occurs internally within the polyp. The larvae are retained inside the polyp until they are more developed and ready to settle soon after being released. This strategy often results in the larvae settling near the parent colony, contributing to the growth of dense coral clusters. *Pocillopora* spp. and *Stylophora* spp. are examples of brooding corals that employ this strategy, leading to the formation of tightly clustered colonies that enhance the local complexity of the reef.

Together, these reproductive strategies ensure the survival, expansion, and genetic diversity of coral populations. They enable coral reefs to adapt to changing environmental conditions, recover from disturbances by colonizing new areas and reinforce existing structures.

Corals, animals and vegetation interactions

Coral reefs are dynamic ecosystems where corals, algae, and reef-dwelling animals are constantly interacting, with each playing a vital role in the health and stability of the reef. The competition between corals and algae is a fundamental aspect of these interactions. Both corals and algae compete for the same resources: sunlight, nutrients, and substrate. However, their growth strategies differ significantly. Corals are slow-growing, reef-building organisms that provide the structural foundation of the reef, while algae are generally fast-growing and can quickly colonize available space, especially when nutrient levels are elevated due to pollution or when populations of herbivores, such as fish and sea urchins, are reduced due to overfishing or diseases. This algal overgrowth can suffocate coral colonies, blocking sunlight and hindering coral growth and reproduction. A shift in the ecosystem from a coral-dominated system to an algal-dominated significantly reduces the structural complexity of the reef, as algae do not build the rigid frameworks that corals do. The loss of coral cover and the dominance of algae decrease the reef's ability to provide habitat for marine life, leading to a decline in biodiversity and altering the dynamics of the entire ecosystem.

In addition to controlling algae, certain reef animals contribute to coral health through bioerosion, a process where organisms like parrotfish and certain invertebrates break down dead coral structures, which help recycle calcium carbonate within the reef ecosystem. The fine sediment produced by bioerosion can fill in gaps between living coral colonies, stabilizing the reef and providing a more solid foundation for new coral growth. Fishes also contribute to coral preservation through nutrient cycling. As fishes move through the reef and excrete waste, they release nutrients that can be utilized by corals and other reef organisms. These nutrients can be absorbed by corals to grow, especially in nutrient-poor environments where external nutrient sources are limited. Furthermore, the physical presence of fish within the

reef can enhance water circulation, which helps distribute nutrients and oxygen throughout the reef, supporting coral health and resilience.

Corals have evolved various defense mechanisms to protect themselves from algal overgrowth and other threats. Some corals produce chemical compounds that deter algal growth or attract herbivorous fish that feed on algae. Additionally, certain coral species engage in mutualistic relationships with small fish or invertebrates that help defend the coral from algal encroachment and predation.

Environmental changes, particularly those driven by climate change, can disrupt the balance between corals and algae. Warmer ocean temperatures and acidification by increased CO₂ can increase the frequency and severity of coral bleaching events, weakening corals and giving algae a competitive advantage. Additionally, nutrient enrichment from agricultural runoff, sewage, and other sources can lead to algal domination on coral reefs.

Fish and other marine organisms derive significant benefits from living in coral reef habitats, which are essential for their survival, growth, and reproduction. Coral reefs offer a complex three-dimensional structure that provides shelter and protection from predators, particularly for juvenile fish. This protective environment increases the chances of survival for many species.

Additionally, coral reefs serve as critical breeding and nursery grounds. The sheltered environments within the reef are ideal for spawning and raising young, offering protection to eggs and larvae. Coral reefs also facilitate important social interactions, such as territory establishment and mating behaviors, which are vital for reproduction and species survival.

Lastly, some fish species engage in mutualistic relationships with corals, such as clownfish with anemones, gaining protection while providing benefits like nutrient exchange or cleaning services. These interactions underscore the interdependence of fish and corals within the reef ecosystem.

Coastal impact

Coral reefs serve as vital natural barriers for coastlines, acting as a first line of defense against oceanic forces. These complex structures, composed of the reef crest, fore reef, and back reef, significantly reduce incoming wave energy, thereby protecting coastal ecosystems and communities from storms, waves, and rising sea levels. Coral reefs are especially important for densely populated coastal regions, where they provide crucial protection, as evidenced by the percentage of coastlines safeguarded by reefs globally.

[ADD PROFILE IMAGE FOR FORE-REEF, CREST AND BACK REEF]

A key protective function of coral reefs is their ability to dissipate wave energy. As waves approach the shore, the reef crest acts as a primary barrier, breaking and reducing their intensity. The slope and structural complexity of branching corals contribute to slowing down and dispersing wave energy. This reduction in wave energy directly impacts the shoreline, by reducing coastal erosion and preserving beaches. The stability provided by reefs extends to adjacent coastal ecosystems, such as mangroves and seagrasses, which are also vital for shoreline protection. Together, these ecosystems form a multi-layered defense system where each component enhances the protective functions of the others.

Storm surges have the potential to cause severe coastal flooding. Coral reefs act as natural buffers, reducing the height and energy of these surges before they reach the coast. However, storms also affect the physical structure of the reef itself, with different types of corals responding differently to storm impacts. In regions where coral reefs have remained healthy, communities have benefited from reduced damage during storms, whereas areas with degraded reefs have experienced significant flooding and destruction.

Coral reefs also provide long-term protection against sea-level rise through their ability to grow vertically by depositing calcium carbonate. The sediment stabilization and the protection of mangroves and seagrass areas reduces the erosion on the coasts, the loss of land and flooding. The ability of a reef to respond to sea-level rise is often described in terms of "keep up," "catch up," and "give up." Reefs that "keep up" grow vertically at a rate that matches or exceeds the rate of sea-level rise, ensuring that they continue to provide effective coastal protection. Reefs that "catch up" grow slower than the current rate of sea-level rise but may eventually match it, potentially restoring their protective capabilities over time. On the other hand, reefs that "give up" are unable to grow fast enough to keep up with rising sea levels, leading to their submersion and loss of protective function.

Coral reefs' shape

In Darwin and Jackson, *The Structure and Distribution of Coral Reefs: Being the First Part of the Geology of the Voyage of the 'Beagle,' under the Command of Capt. Fitzroy, R. N., during the Years 1832 to 1836* (1842), Darwin outlined his atoll reefs formation theory, inspired by his Beagle voyage. He proposed that Earth's crust movements under the oceans created atolls. Darwin described a three-stage process in atoll formation: a fringing reef forms around a sinking volcanic island, evolving into a barrier reef, and finally an atoll reef as subsidence continues.

Fringing reefs

Fringing reefs are the most common and widespread type of coral reef, typically found in close proximity to the shore, forming a narrow band that runs parallel to the coastline. These reefs are directly attached to the land, often with no significant separation between the reef and the shore, or they may be separated by a shallow, narrow lagoon. The proximity of fringing reefs to the coast makes them easily accessible, but it also exposes them to various environmental stressors from both the land and the sea.

The structure of fringing reefs is defined by several key components. The reef flat is the area closest to the shore, characterized by shallow waters where corals grow in a relatively calm environment. The reef flat often extends out to the reef crest, which is the highest point of the reef and is typically exposed at low tide. The reef crest is subjected to the full force of incoming waves, making it a zone of high energy where only the hardiest coral species can thrive. Beyond the reef crest, the reef slope descends into deeper water, where the diversity and density of coral species often increase. The reef slope provides a habitat for a wide variety of marine organisms, taking advantage of the increased light availability and water movement.

Fringing reefs form in areas where the environmental conditions are favorable for coral growth, including clear, shallow waters with high light availability, warm temperatures, and moderate wave action. These conditions allow corals to establish themselves close to the shore, where they can benefit from the nutrients carried by coastal currents. However, the proximity of fringing reefs to land also makes them more vulnerable to the impacts of sedimentation, freshwater runoff, and pollution from human activities. Sedimentation can smother corals, blocking the light they need for photosynthesis, while excess nutrients from runoff can lead to algal blooms that compete with corals for space and resources.

Within fringing reefs, there is a distinct pattern of zonation, where different coral species and marine organisms are distributed according to the environmental conditions across the reef. On the reef flat, where conditions are relatively calm, corals tend to be more resilient to temperature fluctuations and sedimentation. On the reef crest, the coral species are more adapted to withstand the wave action and exposure during low tides. The reef slope, with

its deeper waters and more stable conditions, often supports a greater diversity of coral species, including those that form large, complex structures that provide habitat for a variety of marine life.

Barrier reefs

These reefs typically develop from fringing reefs that have grown over long periods, during which the coastline has either subsided or sea levels have risen. As the land subsides or the sea level rises, the original fringing reef is gradually separated from the shore, forming a lagoon between the reef and the coastline. The depth and size of the lagoon, as well as the height of the reef crest, are influenced by factors such as wave action, currents, and the rate of coral growth. The lagoon often contains patch reefs, seagrass beds, and sandy areas. The zonation within barrier reefs creates distinct ecological niches, with different species of corals, fish, and invertebrates adapted to the specific conditions found in each zone. The fore reef slope, with its high-energy environment, supports species that are resilient to strong waves and currents. In contrast, the more sheltered back reef and lagoon provide habitats for species that prefer calmer waters and stable conditions.

Barrier reefs also play a role in the connectivity between marine ecosystems. They act as a bridge between the open ocean and the coastal environments, facilitating the movement of marine species across different habitats. Many species of fish and invertebrates migrate between the reef and the lagoon during different life stages, using the barrier reef as a nursery or feeding ground.

Atolls

The formation of atolls is a process that unfolds over millions of years. As the island gradually subsides, mainly due to tectonic activity, the coral continues to grow upward toward the sunlight. Eventually, the island disappears entirely beneath the ocean's surface, leaving behind a ring-shaped coral reef that encircles the central lagoon. This theory of atoll formation was first proposed by Charles Darwin and remains one of the most widely accepted explanations for the development of atolls.

The lagoon varies greatly in depth, ranging from shallow areas with sandy bottoms to deeper sections where patch reefs and seagrass beds may develop. The sheltered waters of the lagoon provide a calmer environment compared to the outer reef slope, allowing a different community of marine species to develop. These species often include a mix of corals, fish, and invertebrates that are weakly tolerant of the high-energy conditions. The lagoon may also contain small coral islands, known as motus, which form from the accumulation of coral debris and sand.

The outer reef slope, with its exposure to the open ocean, is home to species that are adapted to strong currents and waves, including large predatory fish and robust coral species. In contrast, the sheltered lagoon provides a haven for more delicate species, such as seagrasses, small fish, and invertebrates, which rely on the calmer waters for feeding and reproduction. This diversity of habitats within a relatively small area makes atolls particularly important for marine biodiversity.

Other types of coral reefs

Patch reefs are smaller, isolated coral formations that occur within lagoons, between larger reef systems, or in shallow coastal waters. These reefs are typically found in the sheltered environments of lagoons, where conditions are favorable for coral growth but do not support the development of larger reef structures.

Patch reefs are characterized by their simplicity and isolation compared to their larger counterparts. These small, often circular or irregularly shaped coral assemblages form in

areas where the physical and environmental conditions, such as water depth, light availability, and wave action, allow for the localized growth of coral colonies.

The formation of patch reefs is influenced by a combination of biological and physical factors. Coral larvae settle in suitable areas where they can establish colonies and grow over time. They often develop in locations where the water is clear, and the substrate is stable, such as on sandy bottoms or rocky outcrops.

1.5 Prototype creation

- C++23 and Qt5.12
- OpenGL 4.6 and GLSL
- Marching Cubes on geometry shader
 - ** Bad idea, but justify why
- Renderings:
 - ** With the prototype:
 - *** Marching Cubes on geometry shader
 - *** Triplanar texture
 - *** Real-time results
 - *** Textures based on materials
 - ** With Unreal Engine 5:
 - *** Static meshes
 - *** Added procedural vegetation with plugin [PLUGIN NAME] and ocean with [PLUGIN NAME]
 - ** With Blender 4.1:
 - *** Static meshes
 - *** Easier script usage
 - ...

Part I

Semantic Representation

Abstract

This chapter introduces a novel method for procedural terrain generation, which leverages a sparse representation of environmental features to produce landscapes that are lightweight, plausible and adaptable to user desires. The method differs from traditional terrain generation approaches by emphasizing multi-scale user interaction and incorporating expert knowledge to model the evolution of terrain features over time. By representing terrain features as discrete entities, or "environmental objects", the method enables dynamic interaction between these entities and their surrounding environment, represented through continuous scalar and vector fields. The generation process is iterative and allows for user-guided modifications at any iteration, including the introduction of environmental events that can influence the terrain's evolution. The proposed approach is particularly flexible, capable of generating both terrestrial and underwater landscapes with a focus on large-scale plausibility and detailed, localized feature representation.

CHAPTER 2

Generation de terrain sémantique



Figure 2.1: Our method can produce different scenes including coral islands and canyons at multiple scales using environmental objects to represent terrain features.

Contents

2.1	Introduction	30
2.2	Related works	32
2.3	Description of the method	33
2.3.1	Pipeline	33
2.3.2	Environmental objects \mathcal{O}	35
2.3.3	Environmental attributes \mathcal{E}	35
2.4	Placement of environmental objects in an environment	36
2.4.1	Fitness function ω	37
2.4.2	Skeleton fitting function Γ	38
2.5	Environmental modifiers	41
2.5.1	Environmental material modifiers	41
2.5.2	Height modifiers	41
2.5.3	Influence on water currents	42
2.5.4	Environment stability	42
2.6	User interactions	43
2.6.1	Direct interactions with the environmental objects	43
2.6.2	Indirect interaction with environmental objects	44
2.7	Results and discussion	46

2.8 Conclusion 48
[Back to summary](#)

2.1 Introduction




Figure 2.2: Four examples of topographic maps used in earth science. From left to right and top to bottom: Sedimentary distribution other Madagascar island (Pratt et al., 2017), evolution of coastlines at Bideford, UK and Scalby, UK over the last 6000 years and 2000 years respectively (BP = Before Present) (Shadrick et al., 2022), localization of key parameters of Perth Submarine Canyon, Autralia (Huang et al., 2014), and geological features distribution of karstic landscape at Qasr Lybia, Libya (Amawy and Muftah, 2009)

Topographic maps are very useful tools for biologists, geologists or even oceanologists (which would call them "bathymetric charts"). These maps are displayed in 2D but provide 3D information about the altitude (or depth), but can also use symbology to represent the important elements that need to be visible. Map symbols are important in order to extract as

much information as possible from a 2D object. In cartography, map symbols are defined as geometric primitives such as points, polylines, polygons, and (rarely) polyhedrons. These symbols are a simplification of the content of an environment, or an abstraction of the 3D shape of the terrain features. This is useful to understand the relationship between the different features, which may enable to deduce physical rules in the evolution of a terrain.

In such way, geologists can study the distribution of peaks in a mountain range, the location of soil types in an area, which in turn allow to deduce possible locations of karst networks, for example. Using the same tools, a biologist may interpret the effect of natural or artificial reefs on coastal erosion, or understand better the interactions inside an ecosystem. Oceanologists may deduce the formation of canyons and fans from old river systems.

By simplifying the surface details in maps or models, we can concentrate more effectively on gaining a deep understanding of the underlying processes that shape the terrain. This focused understanding allows us to apply the insights we gain to a wide variety of terrain types. Essentially, this approach enables us to generalize our findings and apply them across diverse geographical landscapes, facilitating broader and more versatile applications of our knowledge.

Parallelly, terrain artists most of the time sketch the global shape of the terrain they will model beforehand, such that they can check, before the modeling part, that the consistency and plausibility of the terrain will be valid. Looking at a simplified map before starting the modeling step allows the designer to modify the overall shape of the terrain, at a large scale, before the 3D geometry comes into play, generating too much control points or vertices to be able to deal with.

Starting from an initial configuration or providing conditions on the desired output terrain, the algorithm we propose will let the different terrain features evolve as a multi-agent system in which the user can apply modifications or new constraints on the state of the environment. The resulting configuration is an environment conform to the constraints given by the user over the distribution of features present in the scene.

As described in the previous chapter, most terrain generation algorithm use the geometry of an initial terrain surface to iteratively apply changes such as noise-based algorithms or erosion simulation. While the introduction of information about some environment variables or properties of the ground may influence the result of an algorithm, the control of the global shape of the terrain is lost as it treat locally the surface, without knowing which feature a certain point on the surface lies on.

The question which led to our solution is the multi-scale user interaction: "Is it possible to provide an interaction mean for terrain generation allowing the user to interact with a small structure like a rock in the same manner as with a large structure like a mountain?". In this work, we want the user to be able to have a large scale representation of the terrain in order to generate a landscape that satisfies his needs while keeping the possibility to apply large modifications. In discussion with robotician users, we realise that we want to create a large landscape that can contain interesting configurations, select a smaller region that have features in a disposition that fits its requirements and then refine again the given region. In the optic of generating a large scale terrain in which we could focus the generation effort in a certain region, we wished to be able to see a coarse representation that can be computed quickly.

We aim for our terrain generation method to be versatile enough to handle both terrestrial and aquatic environments. This dual capability would allow the method to be applicable to landscapes above the water level, such as mountains and valleys, as well as to submerged terrains, such as oceanic canyons and coral landscapes.

Because many geographical terms and computer science terms are deceptive cognates, we

will try to find middle ground in the naming of our introduced structures to avoid as much as possible any ambiguity between the research fields.

2.2 Related works

Procedural terrain generation has been heavily studied for the last 40 years (E. Galin et al., 2019). Researches in this topic try to find new solutions to compromise between realism, user control and efficiency (Gain et al., 2009). Using fractal noise parametrized to resemble real landscape has been an important first step (Musgrave et al., 1989) as it's a fast and light solution to generate procedurally the appearance of mountains. The lack of user control pushed newer works toward the use of controlled noise by including real DEM in the process through learning (Brosz et al., 2007; Kapp et al., 2020), while the rise of deep learning technologies gave higher control to the user through sketches (É. Guérin et al., 2017; Talgorn and Belhadj, 2018).

By including expert knowledge of tectonic process and subsurface geology, some algorithms tend to get more realistic (Cortial et al., 2019; E. Michel et al., 2015; Patel et al., 2021).

While these algorithms are able to generate large-scale landscapes, the finer details of the terrain is often computed by the use of erosion simulation (Cordonnier et al., 2023; Paris et al., 2019b; Schott et al., 2023). This process can be expensive in time but results in more plausible surfaces.

All the algorithms aim to reproduce plausible relief in terrestrial landscapes, mostly limited to alpine landscapes, but a lack of research can be found in almost all other biomes. Underwater landscapes generation, for example, has been almost completely absent from literature for many reasons: the difficulty of accessing the area, the lack of visibility under water and the complex physics of underwater geology and biology make the algorithms adapted for this environment scarce.

The majority of the ocean floor can be represented as a fractal terrain (Mareschal, 1989). While stochastic noise can be sufficient to model the ocean floor, this process won't cover areas with the biggest biomass, near shallower waters such as near coasts and islands.

Due to the impossibility to observe the large-scale and the small-scale of underwater environments, some works related to geology model large structures like the profile shape of the coral reef (Bosscher and Schlager, 1992), simulate its surface growth (Li, 2021), or use procedural algorithms for single coral colonies (Abela et al., 2015). We however don't have a mix of the different scales, and neither methods take into account the environment such as the topography or the interaction of different terrain features. This is mainly due to the fact that the evolution time for each scale varies from a span of weeks to thousands of years.

In an ecosystem, every element within the system has an impact on its surroundings, and accurately simulating physical properties like shading, heat, and humidity can require immense computational power. To address this, instead of simulating these properties individually for each object, they can be represented as scalar fields that span the entire scene. These scalar fields, which may represent temperature, humidity, or occlusion, are locally influenced by nearby objects and terrain features, such as trees casting shadows or buildings radiating heat. This approach, as described in (Grosbellet et al., 2016; É. Guérin et al., 2016), allows for the environment's physical properties to be computed in a more efficient manner. By simplifying the complex interactions into these localized scalar fields, their method provides a scalable system that can handle large and detailed scenes. This system effectively renders scene details, such as snow accumulation or leaf distribution, in a way that

is visually plausible, ensuring realistic results without the need for exhaustive simulations. In a similar approach, parallelly Wejchert and Haumann (Wejchert and Haumann, 1991) and Sims (**Sims1990**) represent wind flow as a composition of local vector fields, known as "flow primitives", which can be combined to simulate complex fluid behavior. This method simplifies the computation by avoiding full fluid simulations, instead offering a lightweight model that allows for intuitive user control over the flow dynamics. By combining these approaches, we can create dynamic ecosystems where environmental effects and fluid flows interact in a plausible manner, while requiring a low computation effort and preserving a high user control.

2.3 Description of the method

In this section, we present the pipeline and processes that underpin our method. Our approach introduces the concept of environmental objects, simplified terrain features that interact with their surroundings to simulate complex ecosystem dynamics. These environmental objects, which can represent natural features such as trees, rivers, or rocks, influence and are influenced by scalar fields like temperature, humidity, and elevation. The method is structured into several key phases: initialization, where the foundational elements of the terrain and environmental objects are set up; an iterative generation process, where these environmental objects are instantiated and interact with their environment; and finally, the production of a sparse representation of the scene's features. This section details each of these phases, explaining how the system dynamically adapts to user input and environmental changes.

2.3.1 Pipeline

Initialization

The generation of the terrain is initialized using an initial height field h and an initial water level \mathcal{L} . During this chapter we will include many features depending on altitude or depth, so we will use the shorthand notations $\mathcal{H} = h - \mathcal{L}$ and $\mathcal{D} = -\mathcal{H}$. The height field provides variation on the altitude, which can influence the generation process of the scene.

The list of available environmental objects \mathcal{O} , representing the different features that can be present in the scene, are provided with their properties: type, size, generation rules and effects on the environmental attributes (Section 2.3.2). A target list of environmental objects $\hat{\mathcal{O}}$ can be defined to control the final result of the generation.

Finally, different environmental materials are defined with their properties such as diffusion speed, mass, decay rate and influence from the water currents. An initial environment configuration resulting from the initial height field \mathcal{H} and water level \mathcal{L} , the environmental materials distribution \mathcal{M} (represented as scalar fields $\mathcal{M} : \mathbb{R}^2 \mapsto \mathbb{R}$) and water currents \mathcal{W} (as a vector field $\mathcal{W} : \mathbb{R}^2 \mapsto \mathbb{R}^2$) will be used by the environmental objects of the scene to simulate their growth and spawn at the most probable position. The environmental attributes are noted $\mathcal{E} = (\mathcal{D}, \mathcal{W}, \mathcal{L}, \mathcal{M})$ (Section 2.3.3).

The definition of environmental objects' properties and environmental attributes is done with field experts, providing the pertinent parameters required to model the evolution of the terrain features using expert knowledge (Section ??).

The generation phase starts with an initial set of environmental objects present in the scene, which can optionally be pre-filled.




Figure 2.3: Overview of the pipeline of the method. The user provides as input an initial height field and sets the water level, as well as a definition of the environmental materials properties and environmental objects properties that will be used in the iterative process. These inputs are initialized as an initial set of environmental objects and scalar fields that represents the environmental attributes. In the iterative loop, new environmental objects are instantiated using the current state of the environment at their optimal position. The existing environmental objects in the terrain reevaluate their fitness function to grow or die and update the environmental attributes locally. At each iteration, geomorphic events can update the environmental attributes, while the user can interact directly with the environmental objects. The result of the whole process is a set of environmental objects which is a sparse representation of the features of the scene.

Generation process

Once the initialization phase is done, the generation begins. The generation process is incremental and its main loop is composed of two different steps: the instantiation of new environmental objects then the update of the environment.

Instantiation

At each iteration, new environmental objects can be created at their most fitting locations if possible. The generation rules provided in the initialization phase are used to find an optimal position from stochastic sampling (Section 2.4). All environmental objects are evaluating their state analytically using a fitness function and a skeleton fitting function provided as input (Section 2.4).

Environment update

Once the instantiation step is done, the environmental attributes are updated by each environmental object through environmental modifiers, which deposit and absorb some of the environmental materials \mathcal{M} (Section 2.5) while modifying the water currents \mathcal{W} (Section 2.5.3) and the height field \mathcal{H} around them. Finally, water currents and terrain slope displace environmental materials of the terrain until reaching a dynamic equilibrium in the environment at each iteration.

During the generation process, the user can alter directly the distribution and shapes of the environmental objects (Section 2.6.1) and perturb the generation process by planning geomorphic events that have impacts on the environmental attributes (Section 2.6.2).

Output

The output of our system is a set of environmental objects disposed in the plane. We do not provide the 3D representation of the environmental objects in this chapter, letting the user define the rendering method. The figures used in the chapter use a mix of implicit surfaces and triangular meshes.

2.3.2 Environmental objects \mathcal{O}

A geographical feature, also called object or entity, is defined as a discrete phenomenon located at or near the Earth's surface, relevant in geography and geographic information science (GIScience). It represents geographic information that can be depicted in maps, geographic information systems (GIS), and other forms of geographic media. This term includes both natural and human-made objects, ranging from tangible items like buildings or trees to intangible concepts like neighborhoods or savanna. Features are distinct entities with defined boundaries, differentiating them from continuous geographic masses or processes occurring over time. They can be categorized as natural features, such as ecosystems, biomes, water bodies, and landforms, or artificial features, such as settlements, administrative regions, and engineered constructs. Geographic features are described by characteristics including identity, existence, classification, relationships with other features, location, attributes, and temporal aspects. Information about these features is stored in geographic databases using models like GIS datasets, which organize and represent these features in structured formats. As the term "feature" is overused in computer science, we will use the term environmental object in this work.

Each environmental object is shaped with a simple geometric shape called a "skeleton" that defines where it is located and how it fits into the environment. The skeleton can either be, as used in cartography, a point, a curve or a region. We will then refer to our environmental objects as point-based, curve-based or region-based, respectively. These environmental objects interact with the environment \mathcal{E} by changing local conditions using the environmental modifiers. For example, the presence of a river might increase the moisture in the surrounding area, while a mountain might induce more rockiness in the soil composition. They can also absorb changes from the environment, such as a forest taking in humidity from the air. The placement of environmental object is determined by a fitness function ω , which evaluates how suitable a location is based on the environmental attributes. Once a suitable location is found, the skeleton fitting function optimizes the shape and position of the entity to fit as best as possible into the environment Γ .

2.3.3 Environmental attributes \mathcal{E}

In geography, a "field" refers to a continuous spatial phenomenon across a region where each point has a specific value of a variable, unlike discrete objects with distinct boundaries. Fields can be scalar, representing a single value at every point (like temperature or elevation), or vector, representing quantities with magnitude and direction (like wind velocity). Examples include topographic fields for elevation, climatic fields for temperature or precipitation, and magnetic fields for magnetic forces. Because of the ambiguous nature of the term "field" with mathematics and computer science, we will define the geographic fields as environmental attribute.

In an ecosystem simulation, each actor of the ecosystem has an impact on all other actors, which results in an exponentially growing computation effort as the number of elements

of the terrain increase. We avoid this problem by considering the environmental attributes as a proxy to allow any environmental object to interact with any other one. Each of the environmental object have a local impact on the environmental attributes without knowledge of neighboring environmental objects. This modification of the environmental attributes are presented as the effect of environmental modifiers defined for each environmental object.

In this work, we have integrated vector environmental attributes (e.g., water currents \mathcal{W}) and scalar environmental attributes (e.g., altitude \mathcal{H} , water level \mathcal{L} , and various material properties \mathcal{M}) under the unified term "environment," denoted as $\mathcal{E} = (\mathcal{H}, \mathcal{W}, \mathcal{L}, \mathcal{M})$. The environmental materials represent abstract quantities such as the availability of sand, salt, moisture, or rocks at each point. It is important to emphasize that these materials are not to be visualized as physical layers stacked on the terrain surface. Instead, they should be understood as conceptual resource distributions that influence the environment and the behavior of environmental objects, rather than as something directly observable in the scene.

Environmental modifiers \mathcal{E}^+

The environment determine if a environmental object does belong at a certain position. When a environmental object is placed, its surrounding environmental attributes can be affected though environmental modifiers noted $\mathcal{E}^+ = (\mathcal{H}^+, \mathcal{W}^+, \emptyset, \mathcal{M}^+)$ defining a change of height \mathcal{H}^+ , changes in the water currents \mathcal{W}^+ and environmental material alteration \mathcal{M}^+ .

Environmental objects are subject to altitude conditions. However, to maintain a clear distinction between semantic modeling and 3D modeling, we do not compute the exact physical shape or detailed height field of each environmental object. Instead, we define a coarse height function, a parametric representation derived from the skeleton to provide a rough estimate of the changes in elevation around it. This simplified model of how the environmental object influences the surrounding terrain's altitude allows us to cheaply evaluate the potential presence of new entities that have altitude-dependent conditions without the need to perform expensive computations to generate a detailed height map of the entire terrain.

Altering the vector field of the water currents \mathcal{W} is done by the composition of the effect \mathcal{W}^+ of each object at a position \mathbf{p} as introduced in Wejchert and Haumann, *Animation aerodynamics* (1991), while we use the formulation of Kelvinlets (Goes and James, 2017) in the computation of effect of each environmental object.

Each environmental object has intrinsic environmental materials that can be seen as "spreading" and "absorbed" around its skeleton over time. A coral reef may produce coral polyps and at the same time reduce the water currents. It grows thanks to the deposition of limestone from coral colonies. In our model, the colonies affect the environmental attributes through the deposition of environmental material $\mathcal{M}_{\text{limestone}}^+$, which in turn, is absorbed by the coral reef, without a direct exchange between the two environmental objects.

The alteration of a scalar environmental attribute is done by adding or removing some amount around the skeleton of the environmental object and diffusing it in the space, influenced by the water currents. We consider the system to be steady state, garantied by the introduction of a decay rate $k > 0$ in the computation of the diffusion and advection.

2.4 Placement of environmental objects in an environment

At each iteration of our algorithm, we want our environmental objects to be at plausible positions. We do not guaranty a temporal continuity between iterations as in Ecormier-Nocca

et al., *Authoring consistent landscapes with flora and fauna* (2021), so the objective is to add new environmental object in order to satisfy the users wishes, while conserving the plausibility of the scene. Rather than "forming" these environmental objects, our method "reveals" them, much like a paleontologist uncovers fossils during an excavation. A paleontologist does not dig randomly across the Earth to find fossils; instead, they analyze the geological context to identify the most likely locations. Similarly, our method observes the environment and estimates where certain elements are likely to exist. For example, in hydrology, if a river appears to originate from nowhere, it might suggest the presence of a karstic river system upstream. In urban planning, if many roads converge at a certain point, it is reasonable to expect that a city is located there. This approach ensures that Semantic Terrain Entities are revealed in positions that are contextually appropriate and coherent within the landscape.

We will follow the same intuition using a fitness function for each of the environmental object that may be spawn in the terrain. The fitness function defined $\omega : \mathcal{E} \mapsto \mathbb{R}$ provides a score indicating how well the environmental object may fit in this position. Evaluating this function at multiple position results in an approximation of the fitness map of the entity. Once the most probable position is found, we can find the most plausible shape of the environmental object using the skeleton fitting function Γ .

For this task, we place a new elements required at the most plausible position using the analysis of the fitness function of each environmental object. We know that each environmental object will modify the environment surrounding, which may make previously instantiated environmental objects unfitted. Knowing this, the goal is to add the new element at the position that will change the least the stability of the system.

Genetic algorithms or Depth First Search algorithms could be used to try many possibilities until a local or global minimum could be found, but this would require a large processing power. Naive genetic algorithms would place a environmental object at a certain position at each iteration and evaluate the stability of the environment, repeating this operation while varying slightly the position of the environmental objects or the type of environmental object instantiated at each iteration, resulting in way too much computation to stay interactive. The Depth First Seach algorithms would require to compute all the possible combinations of environmental objects and positions which, given the fact that we want a continuous position in order to work multi-scale, would require to compute an incredibly high amount of possible configurations in order to find a plausible situation, on average. We will work with an evolutionary algorithm to find a compromise between fast computation and a satisfying result.

Our placing algorithm is done in two steps: first, it identifies the global location where a environmental object best fits within the environment \mathcal{E} using its fitness function ω , which evaluates the suitability of each point \mathbf{p} based on the environmental attributes \mathcal{E}_p , including factors such as altitude $\mathcal{H}(\mathbf{p})$, water current velocity and direction $\mathcal{W}(\mathbf{p})$, water level $\mathcal{L}(\mathbf{p})$, and the availability of environmental material $\mathcal{M}(\mathbf{p})$. Second, once the most suitable location is identified, the algorithm determines the most plausible shape of the environmental object using the skeleton fitting function Γ .

To ease the reading the functions $\omega(\mathcal{E}(\mathbf{p}))$ and $\Gamma(\mathcal{E}(\mathbf{p}))$ with $\mathcal{E}(\mathbf{p})$ the environmental attributes at the point \mathbf{p} of the terrain are simplified to $\omega(\mathbf{p})$ and $\Gamma(\mathbf{p})$ respectively.

2.4.1 Fitness function ω

"Darwinian fitness" refers to an organism's ability to survive and reproduce in its environment. It is a measure of how well-suited an organism is to its surroundings, and those with higher fitness are more likely to pass on their genes to the next generation. In a similar

vein, the fitness function of a environmental object evaluates its suitability at a location in the terrain, extending the meaning to living features like forests and corals and non-living elements such as rivers, mountains, karsts, etc.

The fitness function is constructed by evaluating several environmental variables at a given location \mathbf{p} . These include altitude ($\mathcal{H}(\mathbf{p})$) and its gradient ($\nabla \mathcal{H}(\mathbf{p})$), the availability and gradient of various materials ($\mathcal{M}_i(\mathbf{p})$ and $\nabla \mathcal{M}_i(\mathbf{p})$), water current characteristics ($\mathcal{W}(\mathbf{p})$), and water level ($\mathcal{L}(\mathbf{p})$). Altitude and its gradient can influence the placement of objects like rivers, which may prefer lower elevations, or forests, which might thrive on slopes. Each material, such as limestone, sand, or clay, is considered separately, and their availability and gradients at a location are crucial for determining the suitability of different environmental objects, such as a coral reef requiring specific substrates. The velocity and direction of water currents are essential for placing aquatic features or determining where erosion might occur, while the proximity to water influences the likelihood of placing wetlands, lakes, or other hydrological features.

These environmental variables are typically combined in the fitness function, often expressed as a weighted sum, but not restricted to this form:

$$\omega(\mathbf{p}) = w_1 \cdot \mathcal{H}(\mathbf{p}) + w_2 \cdot \nabla \mathcal{H}(\mathbf{p}) + \sum_i (w_{3,i} \cdot \mathcal{M}_i(\mathbf{p}) + w_{4,i} \cdot \nabla \mathcal{M}_i(\mathbf{p})) + w_5 \cdot \mathcal{W}(\mathbf{p}) + w_6 \cdot \mathcal{L}(\mathbf{p})$$

Here, $w_1, w_2, w_{3,i}, w_{4,i}, w_5, w_6$ are weights that reflect the relative importance of each factor for the specific environmental object being considered.

Different types of environmental objects require different criteria for their fitness evaluation. For example, a river might prioritize lower altitude and proximity to water currents, while a forest might prioritize higher altitude and specific material availability. The flexibility of the fitness function allows it to be customized for each environmental object, ensuring that the generated terrain remains coherent and realistic.

2.4.2 Skeleton fitting function Γ

The seed point of a spawning environmental object is defined at the point \mathbf{p} satisfying $\arg \max_{\mathbf{p}} \omega(\mathbf{p})$.

Point-based skeleton

The spawning position of a punctual environmental object is found at the local maxima of the skeleton fitting function from the seed point. While the skeleton fitting function doesn't have to be identical to the fitness function, we usually use $\Gamma = \omega$ for point-based environmental objects. If the two functions are different, the optimisation process simply follows the field's gradient $\nabla \Gamma$ until the local maxima is reached.

Curve-based skeleton

The skeleton of a curve-based environmental object is determined by the shape that fits the most given the environment it is added to. Using a modified version the Active Contours algorithm (Kass et al., 1988), we can minimize the energy E for the parametric curve C given

$E(C) = E_{\text{internal}} + E_{\text{external}} + E_{\text{shape}} + E_{\text{gradient}}$ with

$$E_{\text{internal}} = \alpha_i \left(\int_C \|C'(s)\|^2 ds + \int_C \|C''(s)\|^2 ds \right) \quad (2.1)$$

$$E_{\text{external}} = \alpha_e \int_C \Gamma(C(s)) ds \quad (2.2)$$

$$E_{\text{shape}} = \alpha_s \left(L - \int_C ds \right)^2 \quad (2.3)$$

$$E_{\text{gradient}} = \alpha_g \int_C \langle C'(s), \Delta\Gamma(C(s)) \rangle ds \quad (2.4)$$

In this configuration, E_{internal} induce a smooth continuity of the curve by reducing the spacing of each point while reducing the curvature. Another energy, E_{external} integrate the skeleton fitting function over the curve, often seen as an attractor of the points, that tries to descent the gradient to find local minima. At the same time, E_{shape} apply constraints on the curve shape, which, in this case, is to target a specific length L . As such, the curve search for an optimized shape given constraints in E_{shape} for minimizing E_{external} . We introduced a new term E_{gradient} in the energy computation that push the points of the curve in the direction of the slope of the skeleton fitting function, also providing an orientation for the curve.

If a steep coast can be found where the terrain slope is important near the water level, we can define $\Gamma = |\mathcal{H} + 1| / \|\Delta\mathcal{H}\|$, but no orientation is needed, thus we set $\alpha_g = 0$. In this case, the curve will follow a path at the water level and spread its extremities over areas with a steep slope. A river may also be symbolized as a parametric curve, but we need to add information about the direction and magnitude of the slope. As such, we can use $\Gamma = \mathcal{H}$ which forces the direction of the curve to fit with the terrain slope. The introduction of the gradient component provides also an orientation to shapes.

Internal energy

The internal energy, already introduced in Kass et al., *Snakes: Active contour models* (1988) is composed of two components imposing penalties on the local properties of the points of the curve. The first derivative forces the points along the curve to be evenly spaced by minimizing the curve tension, while the second derivative restrict it from forming sharp corners. As our aim is to represent natural elements, we rarely find sharp elements and thus, keep the original definition from the Snake formulation.

External energy

The external energy is also present in the original work. Using an external scalar field, each point of the curve is forced to follow the steepest slope of the field.

Shape energy

We introduced the shape energy, an energy defined on the whole curve to apply constraints on its final shape. As many natural features have a given dimension, we may add a constraint on the lenght of the skeleton.

Gradient energy

In our application, having information about the orientation of environmental objects may be essential. For this purpose, we introduced a gradient energy component in the formulation. The equation impose that the direction of the curve at any point should be directed towards the gradient of the scalar field. As the external energy pushes points toward the lowest point of the scalar field, the gradient field restrict the gradient descent for the global curve into a specific way, which may feel more natural.

During the optimization process, the gradient of the scalar field is already evaluated by the external energy optimization, so the addition of the gradient component is almost free.

Region-based skeleton

Region-based environmental objects follows the same process than curve-based environmental objects to define their skeleton, at the exception of the gradient energy E_{gradient} which have null value on closed shapes.

The resulting energy E to minimize for a closed region whose borders are defined by the curve C can then be expressed as $E(C) = E_{\text{internal}} + E_{\text{external}} + E_{\text{shape}}$.

The internal energy is expressed identically as for curve-based environmental objects. The external energy however, has to be modified to take into account the interior of the region instead of only the borders. We will use the idea of the Chan-Vese algorithm, differentiating the energy value for the inside Ω and the borders C of the region (Chan2001; Getreuer, 2012).

By adding a factor for the inside λ_1 and a factor for the borders λ_2 , we can add weight depending on the required optimization. We can then define the external energy component E_{external} as:

$$E_{\text{external}} = \lambda_1 \int_{\Omega} \Gamma(s) ds + \lambda_2 \int_C \Gamma(C(s)) ds \quad (2.5)$$

We see that the definition of the energy formulation of the curve-based environmental objects is a specialization of the region-based, with $\lambda_1 = 0$ and $\lambda_2 = \alpha_e$

The use of external forces on the skeleton can be useful as some landscape features are easier to define by what they contain, while other by what they separate. For example, a lagoon is formed by coral reefs surrounding it, while a forest is formed by the climatic conditions inside it that are propice for trees. A lagoon may then be defined with $\lambda_1 = 0$ and $\Gamma_{\text{lagoon}} = -\mathcal{M}_{\text{coral limestone}}$ while a forest sets $\lambda_2 = 0$ and $\Gamma_{\text{forest}} = -\mathcal{M}_{\text{humidity}} + \mathcal{M}_{\text{shade}} + |\mathcal{M}_{\text{temperature}} - 10|^2$.

Finally, the shape constraint energy E_{shape} can target an area A , for example.

$$E_{\text{shape}} = \left(A - \int_{\Omega} ds \right)^2 \quad (2.6)$$

In our implementation, each environmental object's skeleton is a connected component as we define the boundaries by the connected curve C , but can be convex or concave. An infinite penalty is added for the curve self-intersection.

At each iteration, environmental objects are interrogated to verify if they are still fitted to belong at their position. We first check that the fitness is above a given threshold $\omega > T_{\omega}$. If not, we remove the environmental object from the scene. On the other case, we can improve the shape of the skeleton by minimizing again their skeleton fitting function. As the number of environmental objects become important in the scene, the reajustment time for the skeletons becomes important. For environmental objects that have been present for multiple iterations, we consider less iterations and an increased convergence threshold up to a point where the features are static.

2.5 Environmental modifiers

2.5.1 Environmental material modifiers

The environment is composed of a scalar field for each of the possible material that can be found in the terrain. The scalar fields represents the availability of the material at any point, but not a height field. Each material is defined with a mass m , a fluid velocity factor ν , a diffusion rate D and finally a decay rate k .

Each environmental object in the terrain is a source and a sink of environmental materials. It is the main mean of communication between environmental objects as it allows them to interact with their surrounding environment. We define the amount of deposited material with D_M and A_M the amount of material deposited and absorbed by the environmental object and $\gamma(t) \in [0, 1]$ a factor related with the current state of the environmental object, which state that more material will be displaced when the environmental object is fully formed than when it was just spawn:

$$\int_0^t \gamma(t) (D_M - A_M) dt$$

The deposition and absorption around an environmental object is defined using the Gaussian kernel distance computation from the skeleton.

The scalar field for the material M is displaced by using a warp operator Φ , taking into account the water flow \mathcal{W} and the terrain slope ∇H . We unified the warp with m the mass of the material and ν a influence factor of the fluid on the material:

$$\Phi(\mathbf{p}, t) = m \nabla H(\mathbf{p}, t) + \nu \mathcal{W}(\mathbf{p}, t)$$

The environmental materials are also dispersed at a diffusion rate D , for which we can use the advection-diffusion-reaction equation to evaluate the distribution after a time t

$$\frac{\partial M}{\partial t} \Phi \nabla M = D \nabla^2 M - k M \quad (2.7)$$

We solve Equation (2.7) numerically using Euler integration

$$\begin{aligned} M(\mathbf{p}, t + dt) &= M(\mathbf{p}, t) + dt(D \nabla^2 M(\mathbf{p}, t) - k M(\mathbf{p}, t) \\ &\quad - \Phi(\mathbf{p}, t) \nabla M(\mathbf{p}, t)) \end{aligned} \quad (2.8)$$

The introduction of the decay rate $k \in]0; 1]$ in the equation allows for the reach of a steady-state, where we can consider the simulation stable. As the user updates the state of the simulation manually, we observe the reach of this steady state before continuing the iterative steps.

2.5.2 Height modifiers

The computation of the height $H(\mathbf{p})$ is done using the coarse height function of each environmental object affecting a point \mathbf{p} .

2.5.3 Influence on water currents

We define our water currents as a vector field defined as

$$\mathcal{W}(\mathbf{p}) = \mathcal{W}_{\text{user}}(\mathbf{p}) + \mathcal{W}_{\text{simulation}}(\mathbf{p}) + \mathcal{W}_{\text{objects}}^+(\mathbf{p})$$

With $\mathcal{W}_{\text{user}}$ a user-defined vector field, $\mathcal{W}_{\text{simulation}}$ an analytical solution inspired by a wind flow simulation (Paris et al., 2019a), and $\mathcal{W}_{\text{objects}}^+$ the water flow alteration computed from the environmental objects. The component $\mathcal{W}_{\text{simulation}}$ is terrain-induced. Given an input flow direction a , we modify the vector field by warping it with the terrain gradient smoothed at multiple scales :

$$\mathcal{W}_{\text{simulation}}(\mathbf{p}) = \sum_{i=0}^{i=n} c_i \Phi_i \cdot v$$

with $v = (a(1 + k_w \mathcal{D}(\mathbf{p}))$ and $\mathcal{D}(\mathbf{p})$ the depth at point \mathbf{p} and k_w a scaling factor, used to simulate the Venturi effects. $\Phi_i \cdot v$ is the warping operator at scale i with a coefficient c_i defined as

$$\Phi_i \cdot v = (1 - \alpha)v + \alpha k_i \nabla \tilde{h}_i^\perp(\mathbf{p}) \quad \alpha = \|\nabla \tilde{h}_i(\mathbf{p})\|$$

with k_i a deviation coefficient, α the slope of the smoothed terrain and $\nabla \tilde{h}_i^\perp(\mathbf{p})$ the orthogonal vector of the smoothed terrain. As proposed by the authors, we used two scaling levels $n = 2$ with gaussian kernels of radii 200m and 50m with weights 0.8 and 0.2 and deviation coefficients k_0 and k_1 of 30 and 5.

$\mathcal{W}_{\text{objects}}^+$ is a deformation field defined as the accumulation of flow primitives (Wejchert and Haumann, 1991). Kelvinlets are applied on each environmental objects to deflect the water flow. We use the scale and grab formulations of the regularized Kelvinlets brushes (Goes and James, 2017), denoted as $s_\varepsilon(r)$ and $g_\varepsilon(r)$ respectively to simulate obstruction and diversion, are defined as

$$s_\varepsilon(r) = (2b - a) \left(\frac{1}{r_\varepsilon^3} + \frac{1}{2r_\varepsilon^5} \right) (sr)$$

$$g_\varepsilon(r) = \left[\frac{a - b}{r_\varepsilon} I + \frac{b}{r_\varepsilon^3} rr^t + \frac{ae^2}{2r_\varepsilon^3} \mathbf{I} \right] \mathbf{F}$$

with $a = \frac{1}{4\pi\mu}$ and $b = \frac{a}{4(1-v)}$ provided μ a shear modulus and v a Poisson ratio provided for each Kelvinlet, $r = \mathbf{p} - \mathbf{q}$ for \mathbf{p} the evaluation position and \mathbf{q} the center point of the Kelvinlet, $r_\varepsilon = \sqrt{\|r\|^2 + \varepsilon^2}$ the regularized distance, ε a radial scale for the deformation field, s a scaling factor and \mathbf{F} the force vector of the grab operation. Deformations defined on curves use $\mathbf{q} = C(\mathbf{p})$ with $C(\mathbf{p})$ the closest point on the curve from the point \mathbf{p} and $f = C'(\mathbf{p})$. We can then define $u_o(\mathbf{p}) = s_\varepsilon(\mathbf{q} - \mathbf{p}) + g_\varepsilon(\mathbf{p} - \mathbf{q})$. Finally, we can retrieve the velocity field from the objects:

$$\mathcal{W}_{\text{objects}}^+(\mathbf{p}) = \sum_{o \in \mathcal{O}} \lambda_o u_o(\mathbf{p})$$

2.5.4 Environment stability

Environmental objects and environmental materials are inspired by the ecological concept of biogeocoenosis, which describes the relationship between living organisms (biotic)

and their non-living environment (abiotic). These interactions closely mirror the processes observed in natural ecosystems, where biotic and abiotic components continuously influence each other, leading to a balanced and stable environment. Gubanov and Degermendzhy [CITATION] describe biogeocoenosis as almost-closed systems, with many parallels possible with thermodynamics such as the concept of dynamic equilibrium. In a biogeocoenosis, the various processes, such as the spread of nutrients, the growth of vegetation, or the erosion of soil, tend toward a state of balance.

Similarly, in our method, the interaction between environmental objects and environmental materials is modeled using a reaction-diffusion-advection framework. This model captures how environmental materials are spread and absorbed within the terrain. Thanks to the decay rate of the environmental materials, the system progresses toward a dynamic equilibrium, where the distribution of environmental materials stabilizes. This stabilization reflects a state of balance where the rates of spreading and absorption and decay are equal, and the environmental attributes become coherent across the landscape. For example, a river might spread sediments downstream, which are gradually absorbed by the surrounding terrain, leading to a stable sediment distribution over time. Similarly, moisture emitted by a forest will eventually reach a balance with the surrounding soil and atmosphere, creating a stable, humid environment.

The only factor that can disrupt this equilibrium is a change in the environmental attributes. Changes such as an increase in temperature, a shift in wind patterns, or the introduction of a new environmental object can alter the balance, leading to a new phase of interaction and stabilization.

2.6 User interactions

The user can guide the generation process. The use of simple shapes as environmental objects facilitate the edition of the simulation, as we can interactively add, remove or modify environmental objects, or focus the generation process in a restricted area. Interaction with the environmental attributes is also provided as geomorphic events, that the user can invoke during the simulation. While the direct interactions on the environmental objects are instantaneous, as the geomorphic events are active on a given duration.

2.6.1 Direct interactions with the environmental objects

The interactive nature of our simulation enables the user to modify the state of the terrain by manipulating directly the environmental objects of the scene. We assume the modifications applied between two iterations of the simulation.

Translating an environmental object is trivial, we simply require to evaluate the state of the environmental objects at a translated position. The deformation of environmental objects can be applied on curve and region environmental objects by updating the control points of the skeleton and recomputing the resulting implicit surfaces. The evaluation positions used for region environmental objects are displaced by applying a cage deformation of the 2D shape using the Green coordinates of points in the shape. After the alteration of the region, evaluation points should be keeping a similar distribution than before, avoiding unexpected results during the interaction. By modifying an environmental object, the environmental attributes may change, which can result in the destruction of the now incompatible environment objects in the scene (Figure 2.4).



Figure 2.4: Starting from a coral colony developed around a canyon (*left*), the user edits the shape of the canyon, resulting in a different configuration of the scene, killing the corals that ends too deep in the water (*center*) and the development and growth of new corals at the previous location of the canyon (*right*).

As long as a non-zero fitness function is defined in the terrain, new environmental objects can be forced by the user at any point of the simulation.

Control over the region of the terrain that should be updated can be given by adjusting all fitness functions through a scalar field $\lambda : \mathbb{R}^2 \mapsto \mathbb{R}$ such that the fitness function $\omega(\mathbf{p})$ of any new environmental object is evaluated as $\omega^*(\mathbf{p}) = \lambda \mathbf{p} \omega(\mathbf{p})$. This is especially useful in the planning of robotic simulations as we can first generate the overall shape of our terrain and secondly focus the generation process around the areas that may be visited by the robot, avoiding useless simulations and computer power. Figure 2.8 shows an example of colonization of the coral polyps that we limited manually into an annulus.

Our water current simulation is modeled as a simple vector field. As such, the user is able to interact with it at any moment of the simulation, allowing for the death of sensible environmental objects while it will guide the simulation into a new landscape. By modifying the water currents, the user also modifies the transport rate of environmental materials at this position. The modification of currents is given as a stroke, a parametric curve C for which we evaluate $\Delta \mathcal{W}_{\text{user}}(\mathbf{p})$ just as for curved environment objects (Section 2.5.3).

2.6.2 Indirect interaction with environmental objects

A configuration file can define in advance the different geomorphic events that should be triggered during the simulation. This can be useful to generate landscapes that are close to some existing locations. Multiple geomorphic events can be triggered either as sudden or continuous environmental changes. These changes play a huge role in the morphology of landscapes. We define geomorphic events with a starting point and an ending point, such that at any time of the simulation we can compute the progress of the geomorphic event as $t_e \in [0, 1]$.

Water level changes are important geomorphic events that shape the underwater landscapes. As previously submerged environmental objects get elevated above water level, flora and fauna terrain features dry and die. Deprived from the living part of the features, everything is more affected by terrestrial erosion. By updating the value of the depth \mathcal{D} evaluated in the fitness functions, any environmental object that is sensible to the depth will be impacted automatically, that may be causing death (Figure 2.5). The modification of the water level is defined as

$$\mathcal{D}(\mathbf{p}) = \mathcal{D}_0(\mathbf{p}) + \sum_{e \in \text{events}} \Delta \mathcal{D}_e t_e$$



Figure 2.5: Lowering the water level by a few meters caused most of the coral objects to satisfy $\omega \leq 0$, causing their death. Since the water level (blue) decrease slowly, new coral objects spawn progressively at a lower altitude.



Figure 2.6: Simulating subsidence on a part of the terrain (brown area) cause the depth value to change locally, resulting in the death of coral objects that find themselves too deep to survive. Here two subsidence geomorphic events are triggered in parallel.

with $\Delta\mathcal{D}_e$ the amount of water rising or lowering during an geomorphic event. We assumed a linear evolution of the water level during an geomorphic event. This allows to evaluate the depth at any point in space and in time.

Subsidence and uplift are the main geomorphic events that create or destroy islands in the long term. These geomorphic events are simulated as a simple factor on the height field of the generated terrain (Figure 2.6). Subsidence is not always uniform in the terrain. As such, the user can provide a position \mathbf{q} at which the subsidence is the strongest, the amount of subsidence applied $\Delta\mathcal{H}_e$ and a standard deviation σ for which we can then compute at any point in space and time of the simulation the height of the terrain

$$\mathcal{H}(\mathbf{p}) = \mathcal{H}_0(\mathbf{p}) \cdot \sum_{e \in \text{events}} \frac{G(\|\mathbf{p} - \mathbf{q}\|)}{G(0)} \Delta\mathcal{H}_e t_e$$

with $G(x)$ the Gaussian function

$$G(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{x^2}{2\sigma^2}\right)$$

Storms are factors of the geomorphology of coral reefs (Oron et al., 2023; Vila-Concejo and Kench, 2016) and coasts (Cowart et al., 2010; Domínguez et al., 2005). Due to the extreme




Figure 2.7: The result of a storm localized on one side of the island (red area) modifies the result of the evaluation of environmental objects around its epicenter for a short period of time. Most of the coral objects died from the geomorphic event, except few environmental objects less sensible to water currents strength.

wind and wave velocities coasts are highly eroded in a short time period and the more fragile corals near the water surface are broken, possibly causing breaches in the reefs and spreading polyps in the currents direction. While there are many factors at play to understand the apparition of storms and the hydrodynamics affecting it, we simplified the model of storms to the user as a single epicenter \mathbf{q} with a wind velocity v_{wind} and a standard deviation σ representing the spread around the epicenter (Figure 2.7). The computation of water currents are then computed as

$$\mathcal{W}_{\text{user}}(\mathbf{p}) = \mathcal{W}_{\text{user}}^*(\mathbf{p}) + \sum_{e \in \text{events} | t_e \in [0,1]} v_{\text{wind}} \frac{G(\|\mathbf{p} - \mathbf{q}\|)}{G(0)}$$

In this case, we did not include the linear factor t_e as storms are usually conserving a constant force for the time of the few weeks or months of their occurrence.

with ΔT_e the change of heat during an geomorphic event, T_0 the temperature at the water surface, and c a very small factor.

The framework can easily be extended as the geomorphic event system stays similar for all geomorphic events. Including higher level simulations in the geomorphic event system can be added, such as the simulation of tectonic activity, the use of fluid dynamics for tsunami geomorphic events, the integration of human activity, ...

2.7 Results and discussion

Our method provides a way to generate scenes at different scales. We demonstrate this capacity with the generation of a large scene of an island (Figure 2.1) after what we focused the generation process in a canyon (Figure 2.9), then a small-scale visualization of coral colonies (Figure 2.8). In the examples, we rendered the environmental objects as a implicit




Figure 2.8: Three colonies of coral (red, blue, green) restricted to an annulus the middle section of the terrain fighting for the space.




Figure 2.9: Evolution of a canyon scene at different iterations of the simulation. The apparition of an arch causes the spawning of rocks, pebbles, and finally some deposition of sand at the bottom of the canyon, spawning ripples.

tree or as individual meshes. The island, lagoons, reefs, canyons and sand ripples as implicit surfaces

A canyon scene can be generated using our method. The water flow is affected by the curve of the canyon such that the currents are oriented in the direction of the curve's tangent. In this example, we force the position of arches to be inside the canyon. The arches deposits a material "rock deposit", which is the main element of the fitness function of the Rock object. The "rock deposit" is slightly affected by water currents, but its mass make it highly affected by gravity. As such, rocks will spawn underneath arches. In reality, an arch is often created as part of a large coral boulder that sees the calcareous bottom part detached by the water currents, often resulting in an arch surrounded by big rocks and smaller rocks from the erosion of the first rocks. As such, we define an environmental object "Arch" with a fitness function $\omega_{arch}(\mathbf{p}) = 5 - d(canyon - \mathbf{p}) * \|\mathcal{W}(\mathbf{p})\|$, an environmental object "Rock" using $\omega_{rock}(\mathbf{p}) = \mathcal{M}_{rock_deposit}(\mathbf{p})$ and Pebble using $\omega_{pebble}(\mathbf{p}) = \mathcal{M}_{smaller_rock_deposit}(\mathbf{p})$. Finally, sand ripples are simply described as curves appearing where there is a lot of sand available: $\omega_{ripple}(\mathbf{p}) = \mathcal{M}_{sand}(\mathbf{p})$. Following these simple rules, Figure 2.9 shows the emergence of details in the scene.

In this example we defined three different types of corals, coralA, coralB and coralC, to illustrate the possibility to model behaviours from the choice of fitness functions. Each of the coral types deposits a material "coral polyp" and "coral polyp A" ("coral polyp B" and "coral polyp C" respectively). By considering a fitness function that minimize the ratio $\frac{\text{coral polyp}}{\text{coral polyp A}}$, we can see an emergent behavior of the three types of coral fighting for the space colonization. Figure 2.8 shows the result of this simulation at three different interations. At the border between two colonies, none of the colonies make progression due to the amount of coral polyp specific from the other colony.

The proposed method aims to generate plausible landscapes using simplified versions

of the evolution of an ecosystem and of the 3D representation. The biological realism of the result is highly correlated to the amount of simplification and assumptions, while the visual realism is completely dependent to the geometric functions used for the 3D modeling of the environmental objects. While proposing a flexible method that propose a generic approach for terrain generation, a close collaboration with fields experts and with graphists is needed to achieve optimal results.

Most simulation algorithm's quality depends on the size of the time step used, but with the introduction of a decay rate in the environmental materials properties, we limit the influence of time steps by considering that steady-state are reachable. The material deposition and absorption on punctual environmental objects can be seen as a Dirac function δ centered at their position resulting in the advantage that material displacement function can use the definition of the diffusion equation instead of the advection-diffusion-reaction equation. This equation allowing us to evaluate the state of the material M without intermediate steps, but this is not applicable with curve- and region-based environmental objects.

2.8 Conclusion

We have proposed a method to generate terrains procedurally using sparse representations. This representation, the environmental objects, enables to introduce expert knowledge by the mean of the fitness functions that rule the environmental objects life cycle, but also to integrate the user in the loop during the generation process. We reduced the terrain resolution limitations by defining the environment objects as parametric features. Thanks to the sparse representation based on single points, curves and regions, we allow for direct manipulation of the environmental objects of the scene by the user which, thanks to the environment steady state consideration, also enables to include these interactions in the automatic simulation process. Integrating environmental properties in the fitness function of environmental objects allows the user to guide the generation through geomorphic events. Our method enables each environmental object of the scene to influence the environment locally, reducing the need of computations while also retrieving environmental attributes locally, which result in a parallelizable life-like simulation process. The genericity of the environment properties definitions should be sufficient for plausible generation of other landscape types as long as expert knowledge can be translated to environmental object's formalism.

We limited our work to the use of 2D scalar fields as they are more easily differentiable, interpretable and lighter than volumetric representations. However, future works include using 3D representations of the terrain and the environment to generate 3D terrains, including cavities, sub-terrestrial areas and the interior of coral structures.



Figure 2.10: A simple coral island is generated using an island, a lagoon, reefs coral polyps, beaches, trees and algae environmental objects. Trees appear on beaches and algae grow in the lagoon's sand.

Part II

Modelisation

Abstract

- Methods completely different
- ** Physical phenomena are different
- ** => Simulation/generation methods must be specific for one landscape
- Methods are developed at different instants of the thesis work,
- ** We will see different procedural generation domains:
 - *** Analytical solutions (coral islands #1)
 - *** Deep Neural Networks (coral islands #2)
 - *** 3D user interaction (karst networks)
- Deep Learning tends to replace procedural methods in 2D domain,
- ** But still too complex for 3D models
 - *** (lack of data, lack of research interest for now)
- ** Still requires a lot of data, which, is (while not easy) possible using aerial images, but is way too sparse with underwater landscapes or underground biomes.
- We want to control the area in which an element is modeled in the terrain
 - ** Because that's how we define them in the previous chapter.
 - ** Thus we cannot use simple random noise methods => no bounds
 - *** Only solution is to use falloff maps, but meh...
- As such, we want to keep the skeleton of our elements using primitives (points, curves, regions)
 - ** => Much easier to add constraints / manipulate primitives in a "procedural generation" way.
- Warning: usage of Deep Learning is at the limit of "procedural generation" and is not considered as part of it by the whole community.
(Complete with the Reddit poll).
- ...

CHAPTER 3

Graphical representation of environmental objects

- Implicit surfaces
- Meshes
- ...

CHAPTER 4

Automatic Generation of Coral Islands

Contents

4.1	Introduction	57
4.1.1	Multiple theories	58
4.1.2	Darwinian theory	60
4.1.3	Overview	60
4.2	Related works	61
4.3	Example generation	61
4.3.1	Closed form of coral growth	62
4.3.2	Labeling of the map	62
4.3.3	Automation	63
4.4	cGAN	63
4.4.1	Definition of cGAN	63
4.4.2	Why cGAN?	63
4.4.3	Training	63
4.4.4	Model usage	64

[Back to summary](#)

4.1 Introduction

- Definition of coral islands
- ** Different types of coral islands
- *** Here, volcanic islands
- Presentation of corals
- Difference from regular landscapes
- ** Concept of corals
- *** Long-term evolution (island) and short-term evolution (corals)
- **** Geological process of island subsidence
- ...

4.1.1 Multiple theories

- Complexity of coral reef ecosystems
- ** Biological diversity
 - *** Varied species of corals and their differing growth patterns
 - *** Interaction with marine flora and fauna
- ** Environmental factors
 - *** Influence of water temperature, salinity, and light availability
 - *** Impact of ocean currents and wave action
- Geological processes
 - ** Dynamic nature of earth's crust
 - *** Plate tectonics and volcanic activity
 - *** Subsidence and uplift processes
 - ** Variations in sea levels
 - *** Historical fluctuations due to glacial and interglacial periods
 - *** Current sea level rise and its impact on coral reefs
- Historical and technological context
 - ** Historical developments in marine science
 - *** Early exploration and observations by naturalists like Darwin and Wallace
 - *** Advances in geological and oceanographic methods
 - ** Technological advancements
 - *** Development of deep-sea exploration tools
 - *** Use of sonar mapping, core sampling, and seismic surveys
- Limitations of early theories
 - ** Inadequate data and observation
 - *** Limited access to deep-sea environments in the 19th and early 20th centuries
 - *** Reliance on surface observations and anecdotal evidence
 - ** Evolving scientific understanding
 - *** Changes in scientific paradigms and theories over time
 - *** Integration of new findings and methodologies
- Different perspectives and disciplines
 - ** Geological vs. biological perspectives
 - *** Focus on geological processes like subsidence and sea level change
 - *** Emphasis on biological factors such as coral growth and reproduction
 - ** Interdisciplinary approaches
 - *** Collaboration between geologists, marine biologists, and oceanographers
 - *** Diverse methodologies leading to different interpretations
- Regional and case-specific variations
 - ** Geographical differences
 - *** Variations in reef types and formations across different regions
 - *** Influence of local environmental conditions and geological settings
 - ** Case studies of specific islands
 - *** Unique formation histories of individual atolls and coral islands
 - *** Examples from the Pacific, Indian, and Atlantic Oceans
- Ongoing research and discoveries
 - ** New findings and theories
 - *** Continuous exploration leading to new hypotheses and models
 - *** Advances in climate science impacting understanding of historical sea levels
 - ** Reevaluation of existing theories

- *** Critical assessment of long-standing theories with new data
- *** Adaptation and refinement of theories over time

Theory 1: Subsidence theory

- Origin and proponents
 - ** Charles Darwin (The Structure and Distribution of Coral Reefs, 1842)
 - ** Alfred Russel Wallace
- Mechanism
 - ** Initial formation around a volcanic island (fringing reef)
 - ** Gradual sinking of the volcanic island (subsidence)
 - ** Transition to a barrier reef with a lagoon
 - ** Complete submersion of the volcanic island leading to atoll formation
- Supporting evidence
 - ** Observations from the HMS Beagle voyage
 - ** Modern geological surveys and core samples

Theory 2: Growth on submarine mountains theory

- Origin and proponents
 - ** John Murray
 - ** Alexander Agassiz
- Mechanism
 - ** Coral colonization on underwater mountains (guyots)
 - ** Vertical growth of corals towards the sea surface
 - ** Formation of atolls without the need for subsidence
- Supporting evidence
 - ** Studies on deep-sea coral formations
 - ** Distribution of guyots and seamounts in tropical regions

Theory 3: Sea level change theory

- Origin and proponents
 - ** Reginald Daly (The Coral Reef Problem, 1915)
- Mechanism
 - ** Impact of glacial and interglacial periods on sea levels
 - ** Lowered sea levels exposing coral reefs, allowing vertical growth
 - ** Rising sea levels creating conditions for atoll formation
- Supporting evidence
 - ** Geological records of sea level changes
 - ** Correlation with coral reef growth periods

Theory 4: Erosion and sedimentation theory

- Origin and proponents
 - ** Maurice Ewing
 - ** William Donn
- Mechanism
 - ** Erosion of coral reefs by waves and currents
 - ** Accumulation of coral debris forming islands

- ** Continuous process of erosion and sediment deposition
- Supporting evidence
- ** Sediment analysis around coral reefs
- ** Observations of island formation from coral rubble

Theory 5: Platform reef theory

- Origin and proponents
- ** William Morris Davis
- Mechanism
- ** Coral growth on stable continental or insular platforms
- ** Formation of reef structures (platform reefs)
- ** Development into coral islands when conditions are favorable
- Supporting evidence
- ** Studies on platform reefs and their distribution
- ** Geological stability analysis of coral platforms

Comparative analysis

- Similarities and differences
- ** Common themes: coral growth, geological activity, sea level changes
- ** Differences in primary mechanisms (subsidence vs. growth on seamounts)
- Complementary aspects
- ** Integration of multiple theories for a comprehensive understanding
- ** Case studies showing multiple processes at work

Modern advances and technologies

- Geological surveys
- ** Core sampling techniques
- ** Seismic and sonar mapping
- Environmental studies
- ** Impact of climate change on coral growth and sea levels
- ** Conservation efforts and their implications for island formation

4.1.2 Darwinian theory

- Several theories
- Difficulty in studying environments
- ** Use of observations
- Theory refuted by (Droxler and Jorry, 2021)
- ** But it's too early to judge
- ** Practical for our case.
- ...

4.1.3 Overview

- Proposed tool
- ** Sketching the island
- ** Sketching the profile

- ** Wind simulation
- Automation of examples
- ** Data augmentation
- ...

4.2 Related works

- Perlin + falloff
- ** But lacks control
- Uplift [SCHOTT UPLIFT] (Cordonnier et al., 2016; 2017a)
- ** But we propose a method that limits the required interaction
- Sketching [PEYTAVIE SKETCHING, EMILIEN FIRST PERSON] (Gain et al., 2009)
- ** Added radial constraint to simplify interaction
- Geological modeling (Patel et al., 2021)
- ** For us, it's hard to know what's happening underground (?)
- Unlike the literature, integration of the underwater part
- ** Integration of observation data into our process
- *** -> Typical shape and profile of an island
- ** Difficult to integrate physical simulations due to uncertainty
- Layer-based generation could improve the realism of examples by considering layer/environment interactions.
- ...

4.3 Example generation

- We use Darwin's theory in our case because generation is relatively simple.
- 2 steps:
 - ** Generation of the island
 - ** Generation of the reef
- Numerous assumptions:
 - ** Island has a relatively round shape
 - ** Coral grows to a constant height around the island
 - ** All "features" are radial
 - ** Deformations of islands caused by winds and waves
 - ** Islands are independent of each other
 - ** The profile is relatively identical all around the island
- ...

Input

- Sketching of the island from above + profile view
- Wind strength
- Water level
- Subsidence rate
- ...

"Simulation"

- Subsidence calculated by simple scaling
- ** Note: No geological consistency
- *** Here, using a zero level to scale in Z
- *** No consideration of different soil materials
- Reef growth
- ** Does not consider any weather events
- ** Considers the "keep up" strategy (as opposed to "give up" and "catch up") [large simplification]
- Wind deformation
- ** using directly $f(\mathbf{p}) = f(\Phi(\mathbf{p}))$.
- ...

Output

- Height map of the island's surface
- Island zones and coral zones
- Possibility to recalculate ground height and coral height
- ...

4.3.1 Closed form of coral growth

- Proposes a closed-form solution for surface calculation
- Calculation of ground height:
 - ** Calculation of height by revolution around the origin point
 - ** Deformation of the revolution profile using the top-down sketch
 - ** Deformation of the height field by the wind map.
- Calculation of growth:
 - ** On the initial heightmap,
 - *** Any surface $z_{min} < h(p) < z_{max}$ becomes coral
 - *** Calculation of the "low" contour $h(p) = z_{min}$ and "high" contour $h(p) = z_{max}$
 - *** ...
- Deformation of the map using the wind map:
 - ** ...
 - ...

4.3.2 Labeling of the map

- Using top-down sketch:
 - ** Features "Mountain", "Island borders", "Beach" are radial (θ, r) , so we can label each point of the map as the "next" feature
- Using coral simulation information:
 - ** Provides the labels "Lagoon" and "Reef {begin, peak, end}".
 - The height map is directly associated to the feature map
 - This is perfect to feed a cGAN.
 - ...

4.3.3 Automation

- Take advantage of the radial nature of the features
- Some features can be optional (mountains)
- Deformation of the feature lines
- ** Influence on the radius: $\tilde{r}(\theta) = r(\theta) + \eta(\theta)$ with η continuous noise function 2π -periodic.
- ...

4.4 cGAN

- Conditional GAN: a type of Generative Adversarial Network (GAN) where the generation process is conditioned on additional information.
- ...

4.4.1 Definition of cGAN

- Two networks: consists of two neural networks, a generator and a discriminator, which compete against each other.
- ** Generator: takes both random noise and additional information (like class labels or data) to produce synthetic data.
- ** Discriminator: evaluates whether a given data instance is real (from the actual dataset) or fake (produced by the generator), while also considering the additional information.
- ** Additional information: this can be labels, data from other modalities, or any other contextual information that guides the generation process.
- Training process: the generator tries to create realistic data to fool the discriminator, while the discriminator tries to correctly classify data as real or fake based on both the data and additional information.
- Objective: the goal is to improve the generator's ability to produce realistic data that matches the given conditions and to enhance the discriminator's ability to distinguish between real and fake data.
- Applications: used in various domains including image-to-image translation, text-to-image synthesis, and other tasks where generating data based on specific conditions is required.
- ...

4.4.2 Why cGAN?

- Flexibility of input
- Moving beyond the "radial" input condition
- Output even for "inconsistent" data (e.g., ocean in an island)
- No math, geometry, geology, or complicated things to master (hehe)
- ...

4.4.3 Training

- ...

Use of synthetic data

- + Problem with synthetic data
- ...

Data augmentation

- ...

4.4.4 Model usage

- ...

Generation from sketch

- ...

Interactive times

- ...

Realism

- ...

CHAPTER 5

Generation of karst networks

Contents

5.1	Introduction	65
5.2	Related works	67
5.3	Space colonization	67
5.4	Our method	67
5.5	Modeling	68
5.6	User control	68
5.7	Results	69
5.8	Conclusion	69

Back to summary

- ...

5.1 Introduction

- Definition: complex subterranean systems formed primarily in soluble rocks like limestone, dolomite, and gypsum.
- Importance: significant for water resources, unique ecosystems, and land use management.
- Formation and development
 - ** Chemical weathering
 - *** Study the role of rainwater acidity (carbonic acid) in dissolving carbonate minerals.
 - ** Underground drainage
 - *** Analyze the development of subterranean drainage systems, including caves, tunnels, and sinkholes.
 - ** Speleogenesis
 - *** Examine the process of cave formation, especially at or below the water table.
- Key features of karst networks
 - ** Caves and caverns
 - *** Document major cave systems and their formation processes.
 - ** Sinkholes (dolines)
 - *** Identify areas prone to sinkhole formation and study their causes.

- ** Disappearing streams and springs
- *** Trace the path of streams that vanish into the ground and re-emerge.
- ** Karst valleys
- *** Investigate valleys formed by the collapse of underground voids.
- ** Stalactites and stalagmites
- *** Study the formation of these features within caves.
- Hydrology of karst networks
- ** Aquifers
- *** Assess the productivity and structure of karst aquifers.
- ** Groundwater flow
- *** Model the rapid and turbulent flow of water through karst systems.
- ** Vulnerability to contamination
- *** Evaluate the risks and sources of contamination in karst aquifers.
- Ecology of karst environments
- ** Unique habitats
- *** Research cave-dwelling species (troglobites) and their adaptations.
- ** Biodiversity hotspots
- *** Identify and document biodiversity hotspots in karst regions.
- Human interaction with karst networks
- ** Water resources
- *** Study the dependence of regions on karst aquifers for freshwater.
- ** Tourism
- *** Assess the impact of tourism on karst landscapes and caves.
- ** Land use challenges
- *** Develop guidelines for construction and land use planning in karst regions.
- ** Environmental concerns
- *** Identify and mitigate pollution and land degradation impacts.
- Examples of karst regions
- ** The Mammoth Cave System (USA): world's longest cave system.
- ** The Guilin Karst (China): unique limestone peaks and river systems.
- ** The Dinaric Karst (Balkans): extensive cave systems and karst phenomena.
- Research and study in karst science
- ** Speleology
- *** Promote the scientific study of caves and karst features.
- ** Geohydrology
- *** Conduct studies on water flow in karst systems for resource management.
- ** Geomorphology
- *** Research the development of karst landforms over geological timescales.
- Action steps
- ** Field surveys and mapping
- *** Conduct detailed field surveys and create maps of karst features.
- ** Hydrological studies
- *** Implement hydrological modeling and water quality testing.
- ** Ecological research
- *** Carry out biodiversity assessments and ecological studies in karst habitats.
- ** Human impact analysis
- *** Study the impact of human activities on karst systems and develop mitigation strategies.
- ** Education and outreach
- *** Educate local communities and stakeholders about the importance of karst networks and

sustainable practices.

- Goals

- ** Conservation

- *** Preserve unique karst ecosystems and landscapes.

- ** Sustainable development

- *** Ensure that human activities in karst regions are sustainable and minimize environmental impacts.

- ** Scientific advancement

- *** Promote research and understanding of karst processes and features.

5.2 Related works

- State of the art:

- ** (Paris et al., 2021)

- *** In our case, we do not rely on a graph and path finder, which allows us to compute sections of the karst on the fly (no need to know the whole network to find paths)

- ** (Pytel and Mann, 2015)

- *** Based on voxels, looks plausible, with a low number of parameters, but take way too long (10 to 20 minutes per generation)

- *** Which makes it unfit for user interaction

- ** (Collon et al., 2015; 2017)

- With some imagination, we can see the shape of trees:

- ** [Prusinkiewicz : ANY]

- ** (Runions, 2008)

- Or miscellaneous networks:

- ** (Fernandes and Fernandes, 2018; É. Galin et al., 2010)

- Maybe even look for lichen or ant colonies

- ...

5.3 Space colonization

- Not really a tree, but...

- ** Some networks are branching

- ** Some have low number of cycles

- SC is easy to manipulate for an user

- For these reasons, we will consider the use of SC.

- Description of the algorithm:

- ** Definition of a root and sinks

- ** Evolution from the root toward closest sinks

- ** Branching [WHEN NEEDED]

- ...

5.4 Our method

- Based on classic SC

- ** Merging branches

- ** Closing paths based on angle and distances to create cycles




Figure 5.1: Classification of tunnel shapes

- Adding width to the branches
- ** Destroying paths when width too small
- Leaves as chambers/cavities
- ...

5.5 Modeling

- Tunnels:
- ** The output of SC is a set of paths
- ** (Paris et al., 2021) provides a classification of tunnel shapes (5.1).
- ...

5.6 User control

- Importance of keeping user control
- ** Shape of a karst is close to randomness
- ** Want to be predictable
- Manipulation of control points
- ** Source
- ** Sink
- Paths tortuosity
- Inclusion of soil properties in the formation of paths, using the gradient of:
 - ** Humidity,
 - ** Porosity
- Real-time editing
- ** Allows for precise manipulation
- ...

5.7 Results

- ...

5.8 Conclusion

- ...

Part III

Erosion Simulation

Abstract

- Work carried out at the beginning of the thesis
 - Representation choice still uncertain
 - Search for abstraction of the representation
- ** Drives the generalization of the erosion system

CHAPTER 6

Érosion par particules

Contents

6.1	Introduction	76
6.2	State of the art	78
6.2.1	Terrain representations	78
6.2.2	Erosion processes	79
6.3	Particle erosion	79
6.3.1	Overview	80
6.3.2	Erosion process	80
6.3.3	Transport	81
6.4	Our erosion method	83
6.4.1	Application on height fields	84
6.4.2	Application on layered terrains	85
6.4.3	Application on implicit terrains	85
6.4.4	Application on voxel grids	86
6.5	Results	87
6.5.1	Rain	88
6.5.2	Coastal erosion	88
6.5.3	Rivers	89
6.5.4	Landslide	89
6.5.5	Karsts	89
6.5.6	Wind	90
6.5.7	Underwater currents	90
6.5.8	Multiple phenomena	90
6.6	Comparisons	91
6.6.1	Coastal erosion on implicit terrain representation	92
6.6.2	Wind erosion on voxel grid representation	92
6.6.3	Hydraulic erosion on height field representation	93
6.6.4	Wind erosion on stacked materials representation	94
6.7	Discussion	94
6.7.1	Realism	94
6.7.2	Usage of velocity fields	95



Figure 6.1: Applying shading and textures on the generated geometry can produce a plausible aspect of a coast eroded by waves on a long timespan, or a desertic landscape eroded by wind, or a mountainous area flatten by thermal erosion.

6.7.3 Performances	95
6.8 Conclusion	96

[Back to summary](#)

6.1 Introduction

Automated terrain generation is a key component of natural scene digital modeling for animated movies and video games. A standard approach is to first generate a base terrain geometry using noise to define the height on the input domain (Musgrave et al., 1989; Olsen, 2004; Roudier, 1993), the result will most likely lack realism and feel synthetic. Erosion simulation algorithms are applied, to simulate thousands of years of ageing by reproducing physical phenomena - i.e. effects of the elements (rain, wind, running water...) - affecting the terrain making it more believable (E. Galin et al., 2019; Smelik et al., 2009; Stachniak and Stuerzlinger, 2005).

The process of terrain alteration caused by the effect of water, air, or any other element - natural or not - over time is usually performed in three steps (Neidhold et al., 2005): **detachment** - pieces of the ground of variable dimensions, ranging from complete ledges to grains of sand, are removed from the terrain depending on the simulated meteorological phenomenon - **transport** - pieces of ground fallen from their initial position are moved to a different one (e.g. a cornice falls down a slope or a grain of sand is thrown into the air) - and **deposition** - transported pieces of land are accumulated at a new part of the landscape. Various phenomena can cause these alterations: **thermal erosion** (bursting of rocks caused by expansion of water under frost, then falling of debris to the bottom of a slope), **hydraulic erosion** (detachment caused by the impact of water particles on surfaces and the transport of sediments by the flow of runoff), **wind erosion** (fine particles carried away in the wind and hit surfaces on their way, creating new fine particles which then also fly away), **chemical erosion** (chemical decomposition of rocks caused by rainwater or other fluids), other exceptional phenomena such as avalanches, animals, lightning, etc... modify the terrain (Argudo et al., 2020; Cordonnier et al., 2017b; 2017a; 2018; 2023).

In practice, the core idea to simulate erosion is to add or remove material from the terrain at given positions on the interface between the terrain and fluid eroding it (e.g. air or water). Hence, the two major problems to tackle are: how to locally alter the terrain geometry for material detachment and deposition and where to perform these alteration given the properties of the environment (terrain slope, fluid density and velocity). A terrain is more than often represented in 2.5D using a 2D image called a **heightmap** which grey scale values define terrain elevation. While being the major terrain representation, only a limited number of environments can be modeled. Indeed, natural landscapes are intrinsically 3D (overhangs,




Figure 6.2: Our method require a base geometry, a small number of parameters for the particles and the medium used for the erosion simulation. It can be easily adapted to be compatible with different mediums and terrain representations.

cavities or geological structures such as arches or gobelins), this is particularly true for underwater environments generation. Alternate representation such as voxel grids, material layers or implicit surfaces can be used. A wide variety of method have been proposed to simulate natural erosion phenomena on heightmaps as the partial differential equations to model erosion can be discretized and solved in 2D and the material detachment and deposition at a given point of the terrain surface can be easily performed by elevating or lowering the ground level i.e. changing locally pixel intensities. For volumetric representations, the alteration of the terrain is not as trivial. To define where to perform the erosion process the local slope variations are more than often used combined with eroding medium information. This fluid can be simulated using particle systems, Smoothed Particle Hydrodynamics (SPH) (Krištof et al., 2009) or approximated using a simple vector field. Proposed methods offer a specific erosion effect tailored to a single terrain representation and fluid simulation.

In this work we propose an approach to simulate a large part of the geomorphological and meteorological phenomena present in the literature of terrain generation (including 3D and volumetric effects). We introduce a generalized algorithm performing the three stages of erosion on surface and volume representations alike, and expose very few intuitive parameters to be adjusted by the user (Figure 6.2). We propose to tackle separately the material variation and the fluid simulation. Our method relies on a particle system to simulate eroding agents, each thrown particle will collide with the terrain, perform terrain alteration at the collision point and transport material along its path. Their motion is computed using simple particle physics accounting for the medium density and particle properties (buoyancy and gravity forces). We consider each particle as independent, hence, they do not interact with each other, no collision detection or response. This simplification allows for efficient parallel computation. When more accuracy or control is needed, we propose to provide a vector field used to modify the particle speed at each time step. The nature of this vector field is flexible, it can be computed using a more or less accurate fluid simulation (SPH, FLIP,...) or be manually defined by the user. We propose a particle-based strategy for material alteration that can be applied on surface and volumetric representation.

The main contributions of this paper are:

- a generalized particle-based algorithm performing the three stages of erosion on surface and volume representations,
- decoupling the erosion system from the fluid simulation, making the process more flexible in its usage and implementation and opening the door for richer effects that can easily be produced.

6.2 State of the art

In this section, we first present the major terrain representations (height fields, layered representations, voxel grids, and scalar functions) and a subset of the major simulated phenomena used to erode terrains. We highlight the fact that, in the literature, a specific erosion method tailored to a given terrain representation is proposed for given phenomena which might lead to limitation in term of terrain modeling. Indeed, changing representation costs information and precision loss.

6.2.1 Terrain representations

A terrain can be represented in various ways, each of them suited for a given application of which we give an brief overview, more details can be found in (E. Galin et al., 2019).

Height Fields

Height fields represents the surface of the terrain by defining the elevation at each point in a 2D grid. This representation is simple, regular, and fast to process allowing for easy manipulation, such as raising or lowering the terrain (Emilien et al., 2015; Gain et al., 2009).

Layered Representations

Layered representations are an extension of height fields using a 2D grid where each cell represents a stack of different materials instead of a simple height (Beneš and Forsbach, 2001; Peytavie et al., 2009) allowing for memory efficient representation of volumetric terrains. To create complex structures, arches or caves, solid materials can be transformed into more granular ones, that can be stabilized (Peytavie et al., 2009).

Voxel Grids

Voxel grids are regular, uniform volumetric grids that encode information on the presence or absence of material for each 3D point in the domain. Voxel grids are advantageous due to their regularity (Dey et al., 2018) and ability to represent volumetric models at the cost of high memory footprint, which has limited their use in terrain generation (Ito et al., 2003; Kaufman et al., 1993; Lengyel, 2010). We consider two voxel grid representations : *density-voxel* grids for which each voxel contains a scalar value, for instance the occupation percentage (Eisemann and Decoret, 2008) and *binary voxel* grids that can be seen as a mask containing the presence of material information.

Implicit terrains

Implicit terrains represent landscapes as an implicit surface defined by a scalar function. This allows for high definition large terrain modeling. The application of combinable scalar function overlays (É. Guérin et al., 2016) or the definition of user-defined gradients (E. Guérin et al., 2022) can be used to create complex terrain features. Altering a implicitly defined surface is a challenging task hence limited option exist for erosion simulation (Paris et al., 2019b).

6.2.2 Erosion processes

Erosion processes play a crucial role in shaping landscapes over time. We present different kind of erosion and how they apply to given terrain representations. Note that using existing methods all erosion methods can not be used on all representation.

Thermal erosion

Thermal erosion is driven by large temperature shifts, transferring material based on slope thresholds. The process is iterative, redistributing material until slopes stabilize. It can be computed efficiently on height fields and layered terrains due to their manipulable height nature (Beneš and Forsbach, 2001; Musgrave et al., 1989; Peytavie et al., 2009). However, its application on voxel grids is challenging due to limited Z-axis resolution.

Hydraulic erosion

Hydraulic erosion stems from water movement, eroding and depositing sediment based on water flow intensity. 2.5D terrains are widely studied for this simulation, using either water slope velocities (Neidhold et al., 2005) or water simulations for erosion effects (Mei et al., 2007). For smaller scales, 3D fluid simulations on voxel grids have been proposed (Beneš et al., 2006). Kristof et al (Krištof et al., 2009) used SPH (Smoothed Particle Hydrodynamics) for meshless erosion simulations on various terrains. Their method involves numerous particle interactions, demanding significant computational power. Our approach draws inspiration from this but enhances efficiency by removing certain particle interactions.

Wind erosion

Wind erosion shifts material through wind force, notably impacting areas with fine surface particles like deserts. It has been modeled on discrete height fields (Paris et al., 2019a; Roa and Benes, 2004) by mimicking sand's wind-driven trajectory and using thermal erosion for corrections. This process is simulated by iteratively displacing small amounts of matter, which make it less suitable for representations with discrete height resolution.

Erosion by other forces

Erosion comes in many forms. These includes influences like glaciers, snow, tectonic movements, and fauna, each introducing distinctive terrain patterns, enriching its intricacy (Argudo et al., 2020; Cordonnier et al., 2016; 2017b; 2017a; 2018). However, most methods are tailored by a given terrain representation, often the height fields, and might not be applicable to other representations due to their intrinsic properties.

6.3 Particle erosion

Erosion occurs in three stages: material detachment, transport and deposition (respectively in red, black and green in Figure 6.3). In our approach, particles move through the medium following its flow (i.e. wind in air or currents in water) and then absorb or deposit a small amount of material upon contact with the land surface, effectively fulfilling the three stages of erosion.





Figure 6.3: Three steps of the erosion process from the sediment point of view: detachment from its original location - dotted red circle -, transport in a fluid - dotted black circle -, deposition at a new location - dotted green circle.

6.3.1 Overview

Particles are transported through the medium and can pass through several different media. Each medium is defined by a density and a flow. Consider, for example, water density to be 1000 kg m^{-3} and that of air to be 1 kg m^{-3} . The gravity applied to the particles is then very different between open and submerged environments due to the difference in buoyancy, while the process remains similar. Using a pre-calculated flow field to guide particle movement simplifies the simulation by treating particles as independent entities, eliminating the need for inter-particle calculations. This not only reduces significantly the overall execution time but also offers users high flexibility over the quality of the simulation and simplify the implementation.

6.3.2 Erosion process

Every time the particle hits the ground, a given amount $q_{\text{detachment}}$ of sediment is detached from the ground (red arrows) while another amount q_{deposit} of sediments is deposited at this location (green arrows). Our erosion model is based on the work of Wojtan et al where regular 3D grids are used to estimate the fluid velocity and sediment transport (Wojtan et al., 2007). In the spirit of (Krištof et al., 2009), we transposed their method into a particle-based erosion simulation, but, in our proposition, we decouple the particle system from the fluid simulation, making the process more flexible and opening the door for richer effects that can easily be produced.



Detachment

As a particle approaches the surface of the terrain, its motion applies friction at the interface between fluid and ground, causing bedrock to dislocate microscopic parts, that we call abrasion. We use pseudoplastics model to approximate the amount of matter removed due to the shear forces while considering the physical properties of the fluid and the ground (Wojtan et al., 2007).

The shear rate θ is approximated by the relative velocity of the fluid to the solid boundary v_{rel} over a short distance l . We approximate the shear stress τ at the solid boundary by a power-law:

$$\tau = K\theta^n \quad (6.1)$$

where $\theta = v_{rel}/l$, K is the shear stress constant (often set to 1) and $n \in [0, 1]$ is the flow behaviour index. Shear-thinning models typically assume n close to $\frac{1}{2}$, which is why we used this value as a constant.

We can then compute the erosion rate ε at any contact point between a fluid and a solid boundary using Equation (6.1) by

$$\varepsilon = K_\varepsilon (\tau - \tau_{\text{critical}})^a \quad (6.2)$$

with $K_\varepsilon \in [0, 1]$ a user-defined erosion constant, τ_{critical} the critical shear stress value for which the matter starts to behave like a fluid and a a power-law constant, typically considered as $a = 1$.

In our method, the eroded quantity is approximated as the material contained in the half sphere, of radius R , in the normal opposite direction at the particle impact point (Figure 6.6). We then use Equation (6.2):

$$q_{\text{detachment}} = \varepsilon \frac{2\pi R^3}{3} \quad (6.3)$$

to get the final eroded amount $q_{\text{detachment}}$. The particle is also defined by a maximal amount of sediments that can be contained in its volume before being saturated noted C_{max} . Note that this constant will be used for the settling velocity computation Equation (6.7).

Deposition

The eroded sediments are considered in suspension in a fluid and are affected by its velocity. A fluid particle then transports the sediments in its flow until gravity settles it onto the ground again. The effect of gravity is modeled by a settling velocity w_s defined in Eq Equation (6.7). We consider that the amount of sediment settled is proportional to the norm of the settling velocity as proposed in (Wojtan et al., 2007) with $\omega \in [0, 1]$:

$$q_{\text{deposit}} = \omega \|w_s\|. \quad (6.4)$$

6.3.3 Transport

Our simulation is computed by integrating the full trajectory of multiple particles at each iteration unlike most other erosion methods. This allows to constantly have a terrain in a plausible state, while giving the possibility to increase the aging effect by running more iterations. Note that, reducing progressively the overall erosion strength can be used as a strategy to adapt the computation time to a chosen level-of-details.

We first present how to compute the particle speed using particle's physics then how to add optional medium velocity field to add a fluid simulation or user control.

Particle's physics

From its independence with other particles: we consider each particle following Newton's laws of motion.

First, we define the external forces \vec{F}_{ext} applied on each particle, we consider gravity and buoyancy. We calculate the buoyancy force $\vec{F}_{\text{buoyancy}} = -\rho_{\text{fluid}} V \vec{g}$ with ρ_{fluid} the density of the fluid, V the volume of the particle and \vec{g} the gravitational acceleration, but we can also calculate the force of gravity $\vec{F}_{\text{gravity}} = m_{\text{particle}} \vec{g}$ with m_{particle} the mass of the particle. We



Figure 6.4: The coefficient of restitution affects the amount of energy absorbed from the particle when hitting the ground. Here, rain is applied on an initial slope (yellow). Only two particles are displayed, with a high (blue) and low (red) coefficient of restitution. The resulting slope after erosion is displayed in blue and red (right).

then have the final external force $\vec{F}_{\text{ext}} = \vec{F}_{\text{gravity}} + \vec{F}_{\text{buoyancy}} = m_{\text{particle}}\vec{g} - v_{\text{fluid}}V\vec{g}$ knowing the density of an object $\rho_{\text{particle}} = \frac{m_{\text{particle}}}{V}$, we have:

$$\vec{F}_{\text{ext}} = V\vec{g}(\rho_{\text{particle}} - \rho_{\text{fluid}}). \quad (6.5)$$

The particle velocity v can be integrated from Equation (6.5) by:

$$v = \int \vec{F}_{\text{ext}} dt + w_s + v_0, \quad (6.6)$$

with w_s the settling speed of sediments in a fluid with a viscosity μ given by Stoke's Law (Stokes, 1850):

$$w_s = \frac{2}{9}\vec{g}R^2 \frac{(\rho_{\text{particle}} - \rho_{\text{fluid}})}{\mu} f(C). \quad (6.7)$$

We use the Richardson-Zaki relation as the hindered settling coefficient:

$$f(C) = 1 - \left(\frac{C}{C_{\max}} \right)^n$$

with C and C_{\max} respectively the fraction of volume of sediments contained and the maximal fraction of sediments the particle can contain, and n an exponent typically 4–5.5, which we set to 5 (Richardson and Zaki, 1954; Wojtan et al., 2007).

Finally, the particle position can be integrated as:

$$\mathbf{p} = \int v dt + \mathbf{p}_0.$$

When the particle hits the ground, a coefficient of restitution affects its behaviour by reducing its velocity post-collision. This value depends on ground material as it is influenced mainly by the material's particle shape, coefficient of friction and density (Yan et al., 2020). Less bouncy particles lose speed quickly and settle down sooner, forming a steeper pile (Figure 6.4 blue), or a higher talus angle like chalk. On the other hand, more bouncy particles disperse more widely upon hitting a surface, resulting in a gentler accumulation like clay (Figure 6.4 red).

Velocity field

In our model, we allow the user to add a velocity field to the environment that influences particles motion. This velocity field can be the result of a complex fluid simulation, a uniform

vector field, or an artistic motion field. We modify Equation (6.6) such that the particle's speed will be influenced by the velocity field as follows:

$$v = \int \vec{F}_{\text{ext}} dt + w_s + \alpha v_{\text{fluid}} + v_0, \quad (6.8)$$

with v_{fluid} medium velocity field modulated by $\alpha \in [0, 1]$.

Our particle system can model intricate scenarios, like the erosion caused by water currents on the seabed or aeolian erosion. The velocity field remains static during the erosion, which may cause inconsistencies in the fluid velocity field. However, minor changes can be overlooked to maintain a balance between realism and computational efficiency (Tychonievich and Jones, 2010). We offer several velocity improvement methods:

- **Fluid simulation refinement:**

Many erosion systems incorporate fluid simulation, requiring regular updates for erosion and velocity (Krištof et al., 2009; Wojtan et al., 2007). Our method can use fluid simulations with multi-resolution refinement, with the possibility to focus the velocity field adjustments near the updated boundaries of the surface (Roose et al., 2011).

- **Particle velocities in fluid simulation:**

With a Lagrangian fluid simulation relying on particle systems (Koschier et al., 2022), our particle velocities can be incorporated in its computation. This approach is only a provisional solution due to potential parameter mismatches with main fluid simulation.

- **Velocity field diffusion:**

Given the minor changes to the surface level at each erosion iteration, which reflect the gradual alterations in terrain surface, we can estimate that the velocity at a fixed point transitioning between the inside and outside of the terrain closely mirrors the velocities observed in its surrounding area. In this context, we can simply interpolate the velocity field at any transitioning point. This simple method, as used in Figure 6.8, allows us to find a balance between achieving realistic flow simulations and maintaining computational efficiency.

6.4 Our erosion method

In this section, we describe how to apply detachment and deposition to different terrain representations with our method (Figure 6.5). We cover the most commonly used representations namely height fields, layered terrains, voxel grid and implicit surfaces, note that our work could be extended to additional representations. Two conditions need to be satisfied for a representation to be eligible for our erosion method: being able to evaluate the intersection of a particle with the ground and compute the normal of the terrain at this point. To the best of our knowledge, all representation do.

We use Verlet integration for the particle's physics (Verlet, 1967), with low error rate and stability even for high dt , reducing computation time for negligible imprecision (Baraff and Witkin, 1998; Swope et al., 1982).

For all the representations, the amount of material absorbed by the particle, i.e. the erosion value $q_{\text{detachment}}$ from Equation (6.3), is taken around the particle at a radius R , meaning that the modification of the terrain by a particle at position c will only occur for the positions




Figure 6.5: Our overall pipeline: our erosion process compute matter displacement of a terrain using an arbitrary representation as long as intersections between particles and the ground can be detected. An optional velocity field, provided by the user, guides the particles trajectories. We propose surface alteration methods to apply the erosion to the terrain in a coherent way between possible representations.




Figure 6.6: Illustration of the material detachment in the (half-)sphere at contact point C (cross) on different representations. (height field) When $Q < 0$ material detachment happen in the bottom scaled half sphere of the particle's contact with the ground, while the deposition is applied on the upper half sphere of volume when $Q > 0$. Unlike the height field, for 3D terrains detachment and deposit are applied in the full sphere around the contact point.

\mathbf{p} satisfying $\|\mathbf{p} - \mathbf{c}\| < R$. At the same time, the amount q_{deposit} from Equation (6.4) is deposited, resulting in a change $Q = q_{\text{deposit}} - q_{\text{detachment}}$

In our simulation, while the dynamics are informed by physical principles, the particle size is conceptualized within a dimensionless framework. This provides the flexibility to adapt our results to various real-world scales, ensuring the applicability of our model across diverse scenarios. Note that, for a 2.5D terrain, we can consider that half of the sphere surrounding the particle is affected which has a volume of $V_{2.5D} = \frac{2\pi R^3}{3}$ while a 3D terrain is affected by the full sphere $V_{3D} = \frac{4\pi R^3}{3}$ (as illustrated Figure 6.6). In the following sections, we will describe the strategies used to modify the amount of matter for different representations.

6.4.1 Application on height fields

On a height field defined by $h(\mathbf{p}) = z$, the intersection point with the surface is verified at $\mathbf{p}_z = h(\mathbf{p})$, and the normal can be computed at the intersection point.

For this representation, the half sphere is scaled in the z direction to fit $\alpha V = Q$ using $\alpha = \frac{Q}{V}$. We then can decrease the height $h'(\mathbf{p})$ at all points \mathbf{p} by the height of the scaled half sphere at position \mathbf{p} . Given the height of the scaled half sphere of center \mathbf{c} and the distance of the particle to the center $d = \|\mathbf{p} - \mathbf{c}\|$ by $h_{\text{half sphere}}(\mathbf{p}) = \alpha \sqrt{R^2 - d^2}$ for all \mathbf{p} such that $d \leq R$

the radius around a particle.

This change of height can be sampled at all points of the 2D grid by reducing the height by

$$\Delta h(\mathbf{p}) = \frac{\sqrt{R^2 - d^2}}{\alpha} = \frac{Q}{\frac{2}{3}\pi R^3} \sqrt{R^2 - d^2} \quad (6.9)$$

The height at each point after an erosion is then computed as $\tilde{h}(\mathbf{p}) = h(\mathbf{p}) + \Delta h(\mathbf{p})$.

6.4.2 Application on layered terrains

Layered terrains are defined as $\mu : \mathbb{R}^3 \mapsto \mathbb{N}$ assigning a discrete material index μ for any point in space (Beneš and Forsbach, 2001; Peytavie et al., 2009). In the original work, outer borders stack elements of the terrain are transformed into density-voxels to enable global erosion through height changes. We enable the erosion/deposition process directly on the layers hence removing the need for representation changes.

When intersecting the terrain, the amount eroded for each material stack should be the integration of the volume of the intersection between the sphere surrounding the particle and the cubicle represented by the stack. Since there is no easy solution (Jones and Williams, 2017), we approximate the volume of the stack we need to alter using the previously defined height field equation Equation (6.9). At a distance d from the particle, the height is defined as:

$$H(d) = \frac{|Q|}{\frac{2}{3}\pi R^3} \sqrt{R^2 - d^2}. \quad (6.10)$$

If $Q > 0$ (more deposition is applied than detachment) then we transform the materials in the stack contained in the sphere to become ground material. For $Q < 0$ the materials are transformed in background material.

6.4.3 Application on implicit terrains

Implicit terrain are defined using a function $f(\mathbf{p})$ and its variation resulting from the erosion process using $\Delta f(\mathbf{p})$. We propose a strategy to compute $\Delta f(\mathbf{p})$ at any point of the sphere surrounding the erosion point based on metaball primitives. At each contact point a metaball is added to create a hole or a bump in the terrain. A metaball is defined as:

$$\Delta f(\mathbf{p}) = \frac{3Q}{\pi} \frac{(1-d)}{R} \quad (6.11)$$

with d the distance of the point \mathbf{p} to the sphere center. For all point \mathbf{p} for which $d \geq R$, $\Delta f(\mathbf{p}) = 0$ (see ??).

As they are the most commonly used representations, we propose a formulation to erode implicit terrains defined by Signed Distance Functions (SDF) and by gradient or vector fields.

Signed Distance Functions

Considering SDF, the terrain is defined as the 0-set of the signed distance function $f : \mathbb{R}^3 \mapsto \mathbb{R}$, hence, for $f(\mathbf{p}) = 0$, the inside as $f(\mathbf{p}) < 0$ and outer-part (i.e. air or water) as $f(\mathbf{p}) > 0$.

The particle erosion applies at impact points at discrete positions, so we propose to add or




Figure 6.7: Our erosion method is applied iteratively on a completely synthetic island, the terrain is altered to obtain a plausible shape by forming rills. The use of particles with hydraulic densities dropped from the sky results in a strong erosion on the sides of the mountains, and the particles that slide to the sea are mainly drifting offshore resulting in the formation of small beaches and a weaker erosion on the bottom of the water body. Repeating the process causes the island height to decrease progressively up to the point where only the submerged part of the terrain is sheltered from erosion.

subtract metaballs defined using equation Equation (6.11) to respectively deposit or erode material using a composition tree:

$$\text{metaball}(\mathbf{p}) = -\Delta f(\mathbf{p}).$$

Now the eroded terrain function $\tilde{f}(\mathbf{p})$ will be evaluated at each point \mathbf{p} from the initial terrain value $f(\mathbf{p})$, the erosion function $\text{metaball}(\mathbf{p})$ and the composition function $g(f_1, f_2)$:

$$\tilde{f}(\mathbf{p}) = g(f(\mathbf{p}), \text{metaball}(\mathbf{p})).$$

As a metaball is added for each particle bounce on the terrain space partitioning optimization algorithms such as k-d trees, BSP trees or BVH can easily be used to improve performances.

Other implicit terrains

are present in the literature, notably a 2.5D representation based on the surface gradient (E. Guérin et al., 2022) and a 3D representation based on curves (Becher et al., 2017) for which the trajectory of each particle projected to the closest surface could be used to define the alteration of the terrain.

In the case of gradient-based representation, we propose to use the partial derivative from the equation of the 2D scalar fields Equation (6.9) that gives:

$$\nabla h' = -\frac{Q}{\frac{2}{3}R^3} \frac{1}{\sqrt{R^2 - d^2}} \vec{CP} \quad (6.12)$$

with \vec{CP} the vector from the position \mathbf{p} to evaluate to the center of the erosion point c . Now the new gradient field can be computed as:

$$\nabla \tilde{h}(\mathbf{p}) = \nabla h(\mathbf{p}) + \nabla h'(\mathbf{p}).$$

6.4.4 Application on voxel grids

We consider two of the voxel grids representations: density-voxel grids and binary voxel grids for which we present our material alternation strategy.

Density voxels

We consider "density-voxel" grids defined on $f : \mathbb{Z}^3 \mapsto [-1, 1]$ for which a voxel is full for $f(\mathbf{p}) = 1$, partially full for $-1 < f(\mathbf{p}) < 1$ or empty for $f(\mathbf{p}) \leq -1$. This definition allows us to erode them smoothly. Since this kind of grid is a discretization of a scalar function, We could directly use Equation (6.11), as described previously, but we take advantage of the discrete nature of the representation to avoid expensive computation.

We apply the erosion from a particle at position c on all points \mathbf{p} in the volume proportionally to the distance from the center of the sphere $d = ||\mathbf{p} - c||$ to find an approximation to the real erosion value per voxel $Q_{approx} = Q \frac{1-d}{R}$. Using their discrete nature, we rectify this value to sum up the total erosion value to Q by dividing each value by the sum of the distances. We now consider eroding the "empty" voxels since their density can drop until -1 . We then have for all surrounding voxels:

$$\Delta f(\mathbf{p}) = Q \frac{(1 - \frac{d}{R})}{\sum (1 - \frac{d}{R})}. \quad (6.13)$$

Resulting voxel value is computed as $\tilde{f}(\mathbf{p}) = f(\mathbf{p}) + \Delta f(\mathbf{p})$. In our implementation, when $f(\mathbf{p}) > 1$, we simply transport the density excess to the above voxel, giving it a very close analogy to height fields as long as $|\Delta f| < 1$.

Binary voxels

The terrain can be represented using an occupancy function as $f : \mathbb{Z}^3 \mapsto \{0, 1\}$ where a voxel $f = 1$ defines the ground and $f = 0$ the background.

We propose to apply particle erosion by assigning voxels a number of hits, and transform them as air or as ground when this number reaches a critical value C that is proportional to the particle's strength parameter K_e (Beardall et al., 2010).

On a hit, all voxels in a radius R receive a hit number:

$$\Delta hits = \lfloor \alpha \Delta f \rfloor \quad (6.14)$$

with Δf the erosion per voxel computed using Equation (6.13) and α a coefficient high enough to obtain values above 1.

All voxels with $\#hits > C$ are transformed to background and voxels with $\#hits < -C$ are transformed to ground.

Note that, a binary voxel grid can also be transformed into a density-voxel grid to be eroded smoothly.

Our formulation for height fields Equation (6.9), can be used to erode 2D scalar field-based representations. Similarly, our proposition for SDF Equation (6.11) enables erosion for continuous 3D scalar fields and voxels Equation (6.13) for discrete 3D scalar fields respectively.

6.5 Results

Our erosion process enables the simulation of a wide range of erosion effects on the major terrain representations alike. In this section, we present applications that demonstrate the versatility of our method by changing the particle's effect size, quantity, density, maximum capacity, deposition factor and the velocity fields. The results of each process are presented in Figure 6.14, parameters used are available at Table 6.1. It is important to note that all erosion

Name	Rep.	Dimensions	Res	#P	#N	R	COR	ρ_{particle}	C_{factor}	ε	ω	Vel f
Rain	H	100x100	20	100	10	1.0	1.0	1000	10.0	2.5	0.3	None
Coastal	DV	100x100x30	10	80	3	5	0.1	500	10.0	5.0	0.5	Uniform
Meanders	I	N/A	N/A	10	20	5.0	1.0	1000	1.0	1.0	1.0	(¹)
River	H	100x100	5	100	50	1.5-5	0.5	900	0.1	1.0	1.0	None
Landslide	H	100x100	20	200	10	2.5	0.2	500	0.1	1.0	1.0	None
Volcano	DV	100x100x40	50	150	30	1.0	5.0	2000	1.0	1.0	5.0	None
Karst	BV	100x100x50	2	1000	40	5	0.5	500	10.0	5.0	0.5	Uniform
Tunnel	DV	100x100x50	1	100	100	2.5	0.1	500	1.0	1.0	1.0	None
Wind	DV	100x100x50	0.2	100	10	1.5	0.9	1.5	1.0	1.0	1.0	(Paris e)
Underwater	H	100x100	10	100	50	2.5	0.9	1000	1.0	1.0	1.0	(Stam, 2

Table 6.1: Parameters used for the generation of the terrains presented in Figure 6.14, with "Rep" the representation (H: Heightmap, DV: Density-voxels, BV: Binary voxels, I: Implicit) "Res" the resolution in meter per voxel or cell, #P the number of particles per iteration, #N the number of iterations, R the particles radius (in voxel or cell unit), COR the coefficient of restitution, ρ_{particle} the particle density in kg m^{-3} , C_{factor} , ε and ω respectively the capacity, erosion and deposition factors, "Vel field" the type of velocity field used and t the computation time of the simulation in seconds on CPU.

(¹) The velocity field is a vector field defined as $v_{\text{fluid}}(\mathbf{p}) = [0 \sin(\mathbf{p}_x) 0]^T$.

examples presented in this section are available for any 3D terrain representation. However, we cannot create volumetric structure, such as overhangs, using 2.5D representations (height fields).

Environment density ρ_{fluid} is set to 1 kg m^{-3} above water level (terrain blue part) and to 1000 kg m^{-3} below it. Velocity field's refinement is done by using the presented diffusion strategy.

6.5.1 Rain

Hydraulic erosion from rain is the most common process used in terrain generation. In this case, particles are seen as water droplets falling from the sky and rolling downhill due to the gravitational force of Earth. No velocity field is required from fluid simulation. These parameters result in a detailed geometry of the rills on the side of mountains that quickly emerge and deposit many sediments in the valley. We demonstrate the result of rain erosion in *Figure 6.14: Rain* with a computation time of 4 seconds.

Using this erosion parameters in combination with water bodies results in different outcomes (Figure 6.7). The terrain above water is directly affected by the erosion process while particles colliding with the underwater part of the terrain are slowed down and filled with sediments, leading to mainly apply deposition. The result is a typical hydraulic erosion on mountains and the formation of slopes and beaches near water level.

6.5.2 Coastal erosion

Waves repeated motion creates coastal erosion, that can be seen as cliffs with holes at the water level.

We apply a uniform velocity field in the water pointing towards the coast to simulate waves and emit particles from the water area with a large size, a density between air and water densities, a high capacity factor and a low deposition factor ω . Using these parameters, the

erosion process is focused at the interface of air and water, and apply a coarse detachment while depositing a very small quantity of sediments, simulating the corrosive effect of water on limestone.

This effect can only be simulated on 3D terrain representations, but will create cliffs on a 2D representation. *Figure 6.14: Coastal* presents the result of coastal erosion on a density-voxel grid that creates overhangs around sea level using a small amount of particles. Note that, the same effect using an alternate implicit representation based on SDF is displayed in Figure 6.10. A shaded version of this effect is presented in Figure 6.1.

6.5.3 Rivers

Given a source point, we generate particles that run downhill, simulating the formation of a river. More complex erosion simulation using fluid simulations like SPH (Krištof et al., 2009) would create realistic results at the cost of high processing time. Our method offers the flexibility to be applied either with a velocity field (simple, used given or resulting from a fluid simulation) or without allowing for simplicity and efficiency.

When provided with a hand-made or procedural velocity field, our particle system can reproduce simple river meanders (*Figure 6.14: Meanders*).

Figure 6.14: River presents a river that has been modeled by emitting water particles with different sizes that ranges from 1.5 m to 5 m, a high coefficient of restitution and a low capacity factor. Random sizes are used to simulate a river for which the flow rate had fluctuated over formation time, while the low capacity ensure that the banks of the river stays smooth. A high coefficient of restitution is a strategy that let the particles flow with low friction, approaching a water behaviour. Our particles are affected only by gravity, without fluid simulation.

6.5.4 Landslide

are mainly caused by large amount of water saturating the ground and flowing downhill, transporting matter in its path.

By using water particles with a medium size, a low coefficient of restitution and a low capacity factor but a high deposition factor ω , they transport sediments on short distances as the velocity quickly drops to 0, and ground material is completely spread along its path since it is easier to deposit the same amount of sediment than the eroded amount at each collision point. Reducing the density of the particle simulates a rise of viscosity in the settling velocity formula, increasing again the quantity of matter to deposit at contact with the ground. By this means, we can simulate landslides as illustrated on *Figure 6.14: Landslide*. A smoother surface is resulting, compared to the rain erosion as the rills are filled with sediments as soon as they begin to form. By setting the initial capacity of the particle equal to 10% of its max capacity, the mass of the terrain increases, simulating a volcano eruption as illustrated on *Figure 6.14: Volcano*.

6.5.5 Karsts

networks are created over hundreds of years from the corrosion of water on the limestone in the ground. A limited number of methods have been proposed for the procedural generation of karsts (Paris et al., 2021).

By reducing the deposition factor ω , the particles simulate corrosion (without mass conservation). We can use the same particle parameters than the coastal erosion (big size, a density between air and water densities, a high capacity factor and a low ω) and optionally provide

a 3D shear stress map. The karst will automatically follow the softest materials, which is geologically coherent as given in example in *Figure 6.14: Karst*, where we can observe a "pillar" that is formed in the center, and thus the karst forms two corridors that finally merge partially. Underground results are only available for representations allowing 3D structures. Another underground terrain simulation is shown in *Figure 6.14: Tunnel* in which a water runoff is eroding a tunnel without the use of a fluid simulation. Here, when particles bounce often on the terrain surface, the coefficient of restitution may be seen as a viscosity parameter.

6.5.6 Wind

erosion is a significant process in deserts shaping since there are no obstacles on the airflow path. Air particles can reach high velocities, transporting sand over long distances forming either dunes or are blasted into rocks, eroding into goblins.

By setting the density of our particles close to 1 kg m^{-3} , two erosion simulations can be applied at once. Air particles follow closely the flowfield given by the user in air. This flowfield can be given from a complex simulation, a user-defined wind rose (Paris et al., 2019a) or a random flowfield with a general direction.

The generation of the different sand structures depends on the velocity field provided, and a simple field will easily generate linear dunes. On contact with a rock block, the simulation will automatically erode block borders, creating shapes looking like gobelins.

Figure 6.14: Wind gives an example of wind erosion on a flat surface with rock columns being eroded. Given a strong 2D velocity field computed by the high wind simulation proposed in (Paris et al., 2019a) is used on light particles, the simulation is fast thanks to the low number of collisions each particle has with the ground.

6.5.7 Underwater currents

Procedural generation of underwater 3D terrains has received little attention. The difference between the underwater and the surface rely on the buoyancy force that is much stronger, meaning that the water flow has a much more impacting effect on erosion than wind. Taking into account the density of the environment and the velocity field of water in our formulas are the keys to be able to apply any erosion in this environment. Our method works in a water environment by giving at least water density to particles. Given a velocity field describing underwater currents from a complex simulation or from a sketch, the particle system erodes the terrain.

In the example presented in *Figure 6.14: Underwater*, the velocity field is given by a simple 3D fluid simulation (Stam, 1999) applied on the terrain.

A complex water flow simulation is computed using SIMPLE (Caretto et al., 1973) fluid simulation with OpenFOAM. The resulting erosion can then follow complex water movement and erode the terrain at the most affected parts of the 3D terrain as the trajectories of the particles (green) is highly affected by the fluid velocity (blue). The density of the particles and the environment being close, the buoyancy cancels most of the gravity force, leaving the velocity of the particles computed by the fluid velocity v_{fluid} and settling velocity w_s from Equation (6.8) (Figure 6.8).

6.5.8 Multiple phenomena

A terrain eroded with multiple erosion phenomena applied on a 500x500x50 density-voxel grid is illustrated in Figure 6.9. Here, water-density particles are applying rain on the terrain




Figure 6.8: A complex water flow simulation is computed using OpenFOAM. Particle trajectories (green) are highly affected by the fluid velocity (blue). Most the terrain exposed surfaces is eroded (bottom).

while the coasts of the river are being eroded thanks to a velocity field defined at the water level. The velocity field defined in the air mainly affects particles with air-density, such that wind erosion can be applied at the same time. The computation of these effects took 7 seconds on CPU.




Figure 6.9: Multiple erosion types can be combined. On an initial synthetic 500x500x50 density voxel grid, the a wind erosion is applied on the surface of the terrain while hydraulic erosion shapes the rills and the base of the mountains. A water current digs its borders and spreads sediments at the bottom.

6.6 Comparisons

In the following section, we compare our method with existing ones to show that while we are versatile on the terrain representation, we are also able to reproduce various effects without applying specific algorithms. The other works are displayed in blue to distinguish them from ours.



Figure 6.10: The algorithm proposed by (Paris et al., 2019b) allows for the simulation of coastal erosion (left) that we can reproduce almost identically by allowing our particles to collide only once with the ground and applying only erosion (center). If we apply our erosion with the full tracking of our particles and using deposition, we can achieve more diverse results (right).

6.6.1 Coastal erosion on implicit terrain representation

Paris et al., *Terrain Amplification with Implicit 3D Features* (2019) present an erosion simulation method applied to implicit terrains able to create coastal erosion, karsts and caves by adding negative sphere primitive in the terrain's construction tree. The positions of the spheres are determined using a Poisson disk sampling at the weakest terrain area defined by the Geology tree of their model. They are simulating the corrosion effect of water on the rocks. Our work is also able to approximate this phenomena by defining the position of these sphere primitives at the position where the water particles hit the surface. While the computation time of the positions of the sphere is higher due to the fact that we are evaluating the position of our particles at every time step in the implicit model (which could be improved by the triangulation of the implicit surface, or better, a dynamic triangulation), the distribution of our erosion primitives is based on a physical model instead of a mathematical model, meaning that we can integrate more easily the direction and strength of the waves for example. The management of their sphere primitives can be replicated with our method by considering that a particle exists until a collision occurs, at which point it disappears. Their method is not conserving the mass of the terrain, which is acceptable for the corrosion simulation, but limits its validity for other erosion simulations. In our method, the particle can be tracked until it settles, ensuring mass conservation (Figure 6.10);

6.6.2 Wind erosion on voxel grid representation

Beardall et al., *Directable weathering of concave rock using curvature estimation* (2010) propose a weathering erosion on voxel grids by approximating and eroding continuously the most exposed voxels. When a solid voxel is decimated, it is considered deposit and is displaced down the slope until a minimal talus angle in the terrain is reached and if the deposition is eroded again, it disappears. Our work is able to reproduce their algorithm by sending our particles from a close distance to the terrain surface. By doing so, we reproduce the erosion process as much as the deposition process since the air particles, filled with sediments, is falling automatically towards the local minimum of the erosion point. Just like in their work, we can easily define the resistance value of the materials to add diversity in the results. By adding the possibility of a wind field, even a very simple uniform vector field, to the simulation, we naturally add the wind shadowing effect that protects a gobelin surrounded by bigger gobelins, and also allows the deposit slope to fit more closely to the wind direction (Figure 6.11).




Figure 6.11: The algorithm proposed by (Beardall et al., 2010) allows for an efficient simulation of the spheroidal erosion, making the creation of gobelins on voxel grids in a plausible way (left). Our algorithm naturally erodes the most exposed areas of the terrain when particles are affected by the wind (right).

6.6.3 Hydraulic erosion on height field representation

Mei et al., *Fast hydraulic erosion simulation and visualization on GPU* (2007) integrate and adapt to the GPU the pipe model proposed in (O'Brien and Hodgins, 1995) for the fluid simulation. This simulation is simple but efficient enough to approximate the Shallow-Water equations in real time and use the speed of columns of water to compute the erosion and deposition rate on the 2D grid of the terrain at each time step. Using columns of water even allows the flow to overpass small bumps on the terrain over time. Our method initially rely on a stable fluid flow that is consistent during the whole life time of a particle, but by refining the simulation at each time step instead of at the end of the particles lifetime, our erosion model is able to reproduce this effect, allowing the terrain to have a single batch of fluid going through it. Our method can be seen as a generalization of Mei et al. that can then be used on more than discrete 2D grids (Figure 6.12).




Figure 6.12: While our resulting geometry on the hydraulic erosion (bottom) is less smoothed than the one proposed proposed by (Mei et al., 2007) (top), our method allows the application on more terrain representations than the height fields only.

6.6.4 Wind erosion on stacked materials representation

Paris et al., *Desertscape Simulation* (2019) simulate the effect of wind over sand fields defined on stacked materials, creating dune structures, even taking into account obstacles like (Roa and Benes, 2004) and different material layers like vegetation (Cordonnier et al., 2017a) that are not affected by abrasion(Paris et al., 2019a). A wind field simulation is required to produce results, and while (Roa and Benes, 2004) and (Onoue and Nishita, 2000) consider a uniform vector field, this work consider a dynamic vector multi-scaled warped field from the terrain height. The sand grains then apply multiple moves: sand lift, bounces, reptation and avalanching. Once the sand is lifted by the wind, the trajectory of the grains can be seen as the displacement of particles, fitting completely with our model as illustrated Figure 6.13.




Figure 6.13: The algorithm from Paris 2020 allow the generation of desertscapes (top), which we can (at least partially) reproduce with our erosion simulation (bottom). The different effects are achieved by affecting the wind direction and strength.

6.7 Discussion

This work is a generalization of erosion that is applicable to any terrain representation. In practice, while similar particle physics is used on different terrain representations, using similar parameters does not ensure resulting in the same eroded terrain. Surfaces and normals being approximated differently have rippling effect on particle trajectories. Note that, not all effects can be applied to all representations, for instance, karsts generation on 2.5D data structures.

6.7.1 Realism

Realism of the erosion simulation is highly correlated to the size and quantity of particles used and their distribution. Using too few or distributing them too sparsely will result in a terrain that is unrealistic since the alteration will have localized effects, breaking process homogeneity.

The resolution is also limited by the number and size of the particles, which can be problematic on implicit terrains that can theoretically have a infinite resolution.

Our method allows to perform erosion on implicit terrains. However, in its current form, our algorithm is time expensive on implicit representations since a large number of primitives are added in the composition tree. Using skeletons-defined primitives (Hong, 2013; Rigaudi  re et al., 2000) from particles trajectories and erosion/deposition values could be a solution to optimize the computation time.

6.7.2 Usage of velocity fields

In our erosion algorithm, we simplify particle physics to enhance computational efficiency and facilitate parameterization. We use the velocity field from fluid simulations to approximate particle velocities. Sediment mass is harnessed to compensate for this approximation, allowing compatibility with various fluid simulation algorithms. Velocity fields can be recomputed at a frequency meeting the applications needs, ranging from "classic erosion simulation" (recomputed at each time step) to "simple simulation" (never recomputed). We addressed provisional adjustments to mitigate discrepancies when terrain changes due to erosion are not reflected in a static velocity field in section "3.3 Transport". However, it is important to note that these are expedient solutions and may not fully capture precise dynamics of an evolving terrain.

6.7.3 Performances

To facilitate parallelization, we intentionally overlook particle interactions and sediment exchanges, albeit at the expense of achieving smoother results. Surface collisions are simplified to basic bounces with a damping parameter instead of relying on complex particles and ground properties (Young's modulus, friction, material, ...)(Yan et al., 2020), further easing the parameterization process. However, these simplifications, combined with the inherent discrete nature of particles, as opposed to the continuous nature of erosion, result in a correlation between realism and particle count.

The performance of our method is influenced by the time required for collision detection. Consequently, we mainly observe better performances with explicit terrain models than with implicit models.

Particle's atomicity

While we can replicate various effects, the "fan" shape commonly observed in natural erosion patterns is not perfectly represented. This limitation arises because we do not account for the splitting of a particle, a process that significantly influences the multidirectional dislocation and trajectory of individual particles (Ranz et al., 1960). Additionally, we acknowledge an issue where particles may collide with the ceiling and the deposition is stuck. While a potential resolution involves splitting particles upon impact rather than simply depositing sediments, this introduces complexities to the parallelization layer of the method. Allowing particles to split introduces unpredictability in the total number of particles that will exist in the simulation. This unpredictability can complicate the use of multi-threading. Future works includes finding a data structure allowing this splitting efficiently, leading to more realistic erosion patterns.

Simulation with multiple materials

One aspect we haven't addressed is a layered terrain with multiple materials. In the native way our method is done, we do not consider the transport of different materials (all

sediments are considered as sand), but by storing a list of the different materials and the quantity transported by each particle, the same simulation process could be done at the cost of some memory and performance overhead.

Another possible adaptation of the erosion strategy for material voxels is to extend the erosion computation from binary voxels by define transformation rules from one material to another when a voxel is eroded a number $\#hits < -C$ or $\#hits > C$. For example, the material "clay" may transform to "sand" when eroded or to "rock" when many depositions occurred.

6.8 Conclusion

We introduced a flexible particle-based erosion system that is easy to use and simple to implement. We have presented how to adapt the process for various terrain representations and generate a variety of erosion phenomenon due to rain, wind, water bodies... by adjusting intuitive parameters hence generate automatically realistic 2.5D and 3D terrains. The use of external velocity fields provides a high flexibility i.e. using the simulations that best fits the user's needs (precision, control, implementation efficiency...). Our method can also be applied to underwater environments with identical physics simulation since our erosion method can be applied on 3D representations. Erosion algorithms are often limited to the use of height fields, but by finding more generalized methods, we can go toward a global use of 3D terrains, which can offer richer and more diverse landscapes.




Figure 6.14: Erosion processes results on various representations presented in section 6.5. Used parameters used are detailed in Table 6.1.

Part IV

Conclusion

- ...

References

- Abela, R., Liapis, A., & Yannakakis, G. N. (2015). A constructive approach for the generation of underwater environments. *Proceedings of the FDG workshop on Procedural Content Generation in Games* (cit. on p. 32).
- Amawy, E. A., & Muftah, A. M. (2009). Karst development and structural relationship in the tertiary rocks of the al jabal al akhdar, ne libya: A case study in qasr libya area. *3rd International Symposium Karst Evolution in the South Mediterranean Area*, 14, 173–189 (cit. on p. 30).
- Argudo, O., Galin, E., Peytavie, A., Paris, A., & Guérin, E. (2020). Simulation, modeling and authoring of glaciers. *ACM Transactions on Graphics*, 39, 1–14. <https://doi.org/10.1145/3414685.3417855> (cit. on pp. 76, 79).
- Baraff, D., & Witkin, A. (1998). Large steps in cloth simulation. *Proceedings of the 25th Annual Conference on Computer Graphics and Interactive Techniques, SIGGRAPH 1998*, 43–54. <https://doi.org/10.1145/280814.280821> (cit. on p. 83).
- Beardall, M., Butler, J., Farley, M., & Jones, M. D. (2010). Directable weathering of concave rock using curvature estimation. *IEEE Transactions on Visualization and Computer Graphics*, 16, 81–94. <https://doi.org/10.1109/TVCG.2009.39> (cit. on pp. 87, 92, 93).
- Becher, M., Krone, M., Reina, G., & Ertl, T. (2017). Feature-based volumetric terrain generation. *Proceedings - I3D 2017: 21st ACM SIGGRAPH Symposium on Interactive 3D Graphics and Games*. <https://doi.org/10.1145/3023368.3023383> (cit. on p. 86).
- Beneš, B., & Forsbach, R. (2001). Layered data representation for visual simulation of terrain erosion. *Proceedings - Spring Conference on Computer Graphics, SCCG 2001*, 80–86. <https://doi.org/10.1109/SCCG.2001.945341> (cit. on pp. 78, 79, 85).
- Beneš, B., Těšínský, V., Hornyš, J., & Bhatia, S. K. (2006). Hydraulic erosion. *Computer Animation and Virtual Worlds*, 17, 99–108. <https://doi.org/10.1002/cav.77> (cit. on p. 79).
- Boldea, C.-R. (2009). A particle cellular automata model for fluid simulations. *Math. Comp. Sci. Ser*, 36, 35–41 (cit. on pp. 11, 15).
- Bosscher, H., & Schlager, W. (1992). Computer simulation of reef growth. *Sedimentology*, 39, 503–512. <https://doi.org/10.1111/j.1365-3091.1992.tb02130.x> (cit. on p. 32).

- Brackbill, J. U., Kothe, D. B., & Ruppel, H. M. (1988). Flip: A low-dissipation, particle-in-cell method for fluid flow. *Computer Physics Communications*, 48, 25–38. [https://doi.org/10.1016/0010-4655\(88\)90020-3](https://doi.org/10.1016/0010-4655(88)90020-3) (cit. on p. 14).
- Brosz, J., Samavati, F. F., & Sousa, M. C. (2007). Terrain synthesis by-example. *Communications in Computer and Information Science*, 4 CCIS, 58–77. https://doi.org/10.1007/978-3-540-75274-5_4 (cit. on p. 32).
- Careil, V., Billeter, M., & Eisemann, E. (2020). Interactively modifying compressed sparse voxel representations. *Computer Graphics Forum*, 39, 111–119. <https://doi.org/10.1111/cgf.13916> (cit. on p. 9).
- Caretto, L. S., Gosman, A. D., Patankar, S. V., & Spalding, D. B. (1973). Two calculation procedures for steady, three-dimensional flows with recirculation. *Proceedings of the Third International Conference on Numerical Methods in Fluid Mechanics*, 19, 60–68 (cit. on pp. 15, 90).
- Cattaneo, G., & Jocher, U. (2005). Cellular automata for 2d and 3d fluid-dynamics simulations. (Cit. on pp. 11, 15).
- Collon, P., Bernasconi, D., Vuilleumier, C., & Renard, P. (2017). Statistical metrics for the characterization of karst network geometry and topology. *Geomorphology*, 283, 122–142. <https://doi.org/10.1016/j.geomorph.2017.01.034> (cit. on p. 67).
- Collon, P., Steckiewicz-Laurent, W., Pellerin, J., Laurent, G., Caumon, G., Reichart, G., & Vaute, L. (2015). 3d geomodelling combining implicit surfaces and voronoi-based remeshing: A case study in the lorraine coal basin (france). *Computers and Geosciences*, 77, 29–43. <https://doi.org/10.1016/j.cageo.2015.01.009> (cit. on p. 67).
- Cordonnier, G., Braun, J., Cani, M.-P., Benes, B., Galin, É., Peytavie, A., & Guérin, É. (2016). Large scale terrain generation from tectonic uplift and fluvial erosion. *Computer Graphics Forum*, 35, 165–175. <https://doi.org/10.1111/cgf.12820> (cit. on pp. 61, 79).
- Cordonnier, G., Cani, M.-P., Beneš, B., Braun, J., & Galin, É. (2017a). Sculpting mountains: Interactive terrain modeling based on subsurface geology. *IEEE Transactions on Visualization and Computer Graphics*, 24. <https://doi.org/10.1109/TVCG.2017.2689022> (cit. on pp. 61, 76, 79, 94).
- Cordonnier, G., Ecormier-nocca, P., Galin, É., Gain, J., Beneš, B., & Cani, M.-P. (2018). Interactive generation of time-evolving, snow-covered landscapes with avalanches. *Computer Graphics Forum*, 37, 497–509. <https://doi.org/10.1111/cgf.13379> (cit. on pp. 76, 79).
- Cordonnier, G., Galin, É., Gain, J., Beneš, B., Guérin, É., Peytavie, A., & Cani, M.-P. (2017b). Authoring landscapes by combining ecosystem and terrain erosion simulation. *ACM Transactions on Graphics*, 36. <https://doi.org/10.1145/3072959.3073667> (cit. on pp. 76, 79).
- Cordonnier, G., Jouvet, G., Peytavie, A., Braun, J., Cani, M.-P., Benes, B., Galin, E., Guérin, E., & Gain, J. (2023). Forming terrains by glacial erosion. *ACM Transactions on Graphics*, 42, 14. [\\"{i}](https://doi.org/10.1145/3592422) (cit. on pp. 32, 76).
- Cortial, Y., Peytavie, A., Galin, É., & Guérin, É. (2019). Procedural tectonic planets. *Eurographics Computer Graphics Forum*, 38, 1–11. <https://doi.org/10.1111/cgf.13614> (cit. on p. 32).

- Cowart, L., Walsh, J. P., & Corbett, D. R. (2010). Analyzing estuarine shoreline change: A case study of cedar island, north carolina. *Journal of Coastal Research*, 265, 817–830. <https://doi.org/10.2112/jcoastres-d-09-00117.1> (cit. on p. 45).
- Dam, P., Duarte, F., & Raposo, A. (2019). Terrain generation based on real world locations for military training and simulation. *Brazilian Symposium on Games and Digital Entertainment, SBGAMES, 2019-Octob*, 173–181. <https://doi.org/10.1109/SBGames.2019.00031> (cit. on p. 5).
- Darwin, C., & Jackson, C. (1842, September). The structure and distribution of coral reefs: Being the first part of the geology of the voyage of the 'beagle,' under the command of capt. fitzroy, r. n., during the years 1832 to 1836. Cambridge University Press. <https://doi.org/10.2307/1797986> (cit. on p. 22).
- de Araújo, B. R., Lopes, D. S., Jepp, P., Jorge, J. A., & Wyvill, B. (2015). A survey on implicit surface polygonization. *ACM Computing Surveys*, 47, 1–39. <https://doi.org/10.1145/2732197> (cit. on p. 9).
- Dey, R., Doig, J. G., & Gatzidis, C. (2018). Procedural feature generation for volumetric terrains using voxel grammars. *Entertainment Computing*, 27, 128–136. <https://doi.org/10.1016/j.entcom.2018.04.003> (cit. on p. 78).
- Domínguez, L., Anfuso, G., & Gracia, F. J. (2005). Vulnerability assessment of a retreating coast in sw spain. *Environmental Geology*, 47, 1037–1044. <https://doi.org/10.1007/s00254-005-1235-0> (cit. on p. 45).
- Droxler, A. W., & Jorry, S. J. (2021). The origin of modern atolls : Challenging darwin's deeply ingrained theory. *Annual Review of Marine Science*. <https://doi.org/10.1146/annurev-marine-122414-034137> (cit. on p. 60).
- Ecormier-Nocca, P., Cordonnier, G., Carrez, P., Moigne, A. M., Memari, P., Benes, B., & Cani, M. P. (2021). Authoring consistent landscapes with flora and fauna. *ACM Transactions on Graphics*, 40. <https://doi.org/10.1145/3450626.3459952> (cit. on pp. 5, 36).
- Eisemann, E., & Decoret, X. (2008). Single-pass gpu solid voxelization for real-time applications. *Proceedings - Graphics Interface*, 73–80 (cit. on p. 78).
- Emilien, A., Poulin, P., Cani, M.-P., & Vimont, U. (2015). Interactive procedural modelling of coherent waterfall scenes. *Computer Graphics Forum*, 34, 22–35. <https://doi.org/10.1111/cgf.12515> (cit. on p. 78).
- Fedkiw, R., & Jensen, H. W. (2001). Visual simulation of smoke. *SIGGRAPH 2001 Conference Proceedings, Annual Conference Series*, 15–22 (cit. on p. 3).
- Fernandes, G. D., & Fernandes, A. R. (2018). Space colonisation for procedural road generation. *2018 International Conference on Graphics and Interaction (ICGI)*, 1–8. <https://doi.org/10.1109/ITCGI.2018.8602928> (cit. on p. 67).
- Gain, J., Marais, P., & Straßer, W. (2009). Terrain sketching. *Proceedings of I3D 2009: The 2009 ACM SIGGRAPH Symposium on Interactive 3D Graphics and Games*, 1, 31–38. <https://doi.org/10.1145/1507149.1507155> (cit. on pp. 32, 61, 78).
- Galin, E., Guérin, E., Peytavie, A., Cordonnier, G., Cani, M.-P., Benes, B., & Gain, J. (2019). A review of digital terrain modeling. *Computer Graphics Forum*, 38, 553–577. <https://doi.org/10.1111/cgf.13657> (cit. on pp. 7, 32, 76, 78).

- Galin, É., Peytavie, A., Maréchal, N., & Guérin, É. (2010). Procedural generation of roads. *Computer Graphics Forum*, 29, 429–438. <https://doi.org/10.1111/j.1467-8659.2009.01612.x> (cit. on p. 67).
- Getreuer, P. (2012). Chan-vese segmentation. *Image Processing On Line*, 2, 214–224. <https://doi.org/10.5201/ipol.2012.g-cv> (cit. on p. 40).
- Gobron, S., & Chiba, N. (2001). Crack pattern simulation based on 3d surface cellular automata. *Visual Computer*, 17, 287–309. <https://doi.org/10.1007/s003710100099> (cit. on p. 5).
- Goes, F. D., & James, D. L. (2017). Regularized kelvinlets: Sculpting brushes based on fundamental solutions of elasticity. *ACM Transactions on Graphics*, 36, 401–411. <https://doi.org/10.1145/3072959.3073595> (cit. on pp. 36, 42).
- Greene, N. (1989). Voxel space automata: Modeling with stochastic growth processes in voxel space. (Cit. on p. 11).
- Grosbellet, F., Peytavie, A., Guérin, É., Galin, É., Mérillou, S., & Benes, B. (2016). Environmental objects for authoring procedural scenes. *Computer Graphics Forum*, 35, 296–308. <https://doi.org/10.1111/cgf.12726> (cit. on p. 32).
- Guérin, E., Peytavie, A., Masnou, S., Digne, J., Sauvage, B., Gain, J., & Galin, E. (2022). Gradient terrain authoring. *Computer Graphics Forum*, 41, 85–95. <https://doi.org/10.1111/cgf.14460> (cit. on pp. 78, 86).
- Guérin, É., Digne, J., Galin, É., & Peytavie, A. (2016). Sparse representation of terrains for procedural modeling. *Computer Graphics Forum*, 35, 177–187. <https://doi.org/10.1111/cgf.12821> (cit. on pp. 32, 78).
- Guérin, É., Digne, J., Galin, É., Peytavie, A., Wolf, C., Beneš, B., & Martinez, B. (2017). Interactive example-based terrain authoring with conditional generative adversarial networks. *ACM Transactions on Graphics*, 36. <https://doi.org/10.1145/3130800.3130804> (cit. on p. 32).
- Hädrich, T., Banuti, D. T., Pałubicki, W., Pirk, S., & Michels, D. L. (2021). Fire in paradise. *ACM Transactions on Graphics*, 40. <https://doi.org/10.1145/3450626.3459954> (cit. on pp. 3, 5).
- Hong, Q. (2013). A skeleton-based technique for modelling implicit surfaces. *Proceedings of the 2013 6th International Congress on Image and Signal Processing, CISIP 2013*, 2, 686–691. <https://doi.org/10.1109/CISP.2013.6745253> (cit. on p. 95).
- Huang, Z., Nichol, S. L., Harris, P. T., & Caley, M. J. (2014). Classification of submarine canyons of the australian continental margin. *Marine Geology*, 357, 362–383. <https://doi.org/10.1016/j.margeo.2014.07.007> (cit. on p. 30).
- Hudák, M., & Ďuríkovič, R. (2011). Terrain models for mass movement erosion. *Theory and Practice of Computer Graphics 2011, TPCG 2011 - Eurographics UK Chapter Proceedings*, 9–16. <https://doi.org/10.2312/LocalChapterEvents/TPCG/TPCG11/009-016> (cit. on p. 5).
- Ito, T., Fujimoto, T., Muraoka, K., & Chiba, N. (2003). Modeling rocky scenery taking into account joints. *Proceedings of Computer Graphics International Conference, CGI, 2003-Janua*, 244–247. <https://doi.org/10.1109/CGI.2003.1214475> (cit. on p. 78).

- Iwasaki, K., Uchida, H., Dobashi, Y., & Nishita, T. (2010). Fast particle-based visual simulation of ice melting. *Computer Graphics Forum*, 29, 2215–2223. <https://doi.org/10.1111/j.1467-8659.2010.01810.x> (cit. on p. 15).
- Jones, B. D., & Williams, J. R. (2017). Fast computation of accurate sphere-cube intersection volume. *Engineering Computations*, 34, 1204–1216. <https://doi.org/10.1108/EC-2016-0052> (cit. on p. 85).
- Kapp, K., Gain, J., Guérin, E., Galin, E., & Peytavie, A. (2020). Data-driven authoring of large-scale ecosystems. *ACM Transactions on Graphics*, 39. <https://doi.org/10.1145/3414685.3417848> (cit. on p. 32).
- Kass, M., Witkin, A., & Terzopoulos, D. (1988). Snakes: Active contour models. *International Journal of Computer Vision*, 1, 321–331. <https://doi.org/10.1007/BF00133570> (cit. on pp. 38, 39).
- Kaufman, A., Cohen, D., & Yagel, R. (1993). Volume graphics. *Computer*, 26, 51–64. <https://doi.org/10.1109/MC.1993.274942> (cit. on p. 78).
- Koschier, D., Bender, J., Solenthaler, B., & Teschner, M. (2022). A survey on sph methods in computer graphics. *Computer Graphics Forum*, 41, 737–760. <https://doi.org/10.1111/cgf.14508> (cit. on pp. 15, 83).
- Krištof, P., Beneš, B., Křivánek, J., & Šťava, O. (2009). Hydraulic erosion using smoothed particle hydrodynamics. *Computer Graphics Forum*, 28, 219–228. <https://doi.org/10.1111/j.1467-8659.2009.01361.x> (cit. on pp. 77, 79, 80, 83, 89).
- Laine, S., & Karras, T. (2010). Efficient sparse voxel octrees. *Proceedings of the 2010 ACM SIGGRAPH symposium on Interactive 3D Graphics and Games*, 55–63. <https://doi.org/10.1145/1730804.1730814> (cit. on p. 9).
- Lejeune, C. (2021). Génération et analyse de tests pour les systèmes autonomes, 170 (cit. on p. 5).
- Lenaerts, T., & Dutré, P. (2009). Mixing fluids and granular materials. *Computer Graphics Forum*, 28, 213–218. <https://doi.org/10.1111/j.1467-8659.2009.01360.x> (cit. on p. 15).
- Lengyel, E. (2010). Voxel-based terrain for real-time virtual simulations, 148 (cit. on p. 78).
- Li, W. (2021). Procedural modeling of the great barrier reef. *Lecture Notes in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics)*, 13017 LNCS, 381–391. https://doi.org/10.1007/978-3-030-90439-5_30 (cit. on p. 32).
- Lima, F. T., Brown, N. C., & Duarte, J. P. (2022). A grammar-based optimization approach for walkable urban fabrics considering pedestrian accessibility and infrastructure cost. *Environment and Planning B: Urban Analytics and City Science*, 49, 1489–1506. <https://doi.org/10.1177/23998083211048496> (cit. on p. 3).
- Louis, S., Andreu, D., Godary-dejean, K., Lapierre, L., Louis, S., Andreu, D., Godary-dejean, K., Lapierre, L., Simulator, H. I. L., Ouis, S. L., Ndreu, D. A., Ejean, K. G. O., & Apierre, L. L. (2019). Hil simulator for auv with contract. *CAR: Control Architectures of Robots* (cit. on p. 5).

- Mareschal, J. C. (1989). Fractal reconstruction of sea-floor topography. *Pure and Applied Geophysics PAGEOPH*, 131, 197–210. <https://doi.org/10.1007/BF00874487> (cit. on p. 32).
- Mei, X., Decaudin, P., & Hu, B. G. (2007). Fast hydraulic erosion simulation and visualization on gpu. *Proceedings - Pacific Conference on Computer Graphics and Applications*, 47–56. <https://doi.org/10.1109/PG.2007.27> (cit. on pp. 79, 93).
- Menshutina, N. V., Kolnoochenko, A. V., & Lebedev, E. A. (2020). Cellular automata in chemistry and chemical engineering. <https://doi.org/10.1146/annurev-chembioeng> (cit. on p. 11).
- Michel, E., Emilien, A., & Cani, M.-P. (2015). Generation of folded terrains from simple vector maps. *Eurographics 2015 short paper proceedings*, 4–8. <https://doi.org/10.2312/egsh.20151019> (cit. on p. 32).
- Michel, É., & Boubekeur, T. (2023). Mesogen: Designing procedural on-surface stranded mesostructures. *Proceedings - SIGGRAPH 2023 Conference Papers*. <https://doi.org/10.1145/3588432.3591496> (cit. on p. 3).
- Müller, M., Charypar, D., & Gross, M. (2003). Particle-based fluid simulation for interactive applications. *Proceedings of the 2003 ACM SIGGRAPH/Eurographics Symposium on Computer Animation, SCA 2003* (cit. on p. 15).
- Musgrave, F. K., Kolb, C. E., & Mace, R. S. (1989). The synthesis and rendering of eroded fractal terrains. *Proceedings of the 16th Annual Conference on Computer Graphics and Interactive Techniques, SIGGRAPH 1989*, 41–50. <https://doi.org/10.1145/74333.74337> (cit. on pp. 32, 76, 79).
- Narain, R. (2011). Visual modeling and visualisation of multiscale phenomena (cit. on p. 3).
- Neidhold, B., Wacker, M., & Deussen, O. (2005). Interactive physically based fluid and erosion simulation. *Natural Phenomena*, 25–32 (cit. on pp. 76, 79).
- Nikeghbali, P., & Omidvar, P. (2018). Application of the sph method to breaking and undular tidal bores on a movable bed. *Journal of Waterway, Port, Coastal, and Ocean Engineering*, 144. [https://doi.org/10.1061/\(asce\)ww.1943-5460.0000424](https://doi.org/10.1061/(asce)ww.1943-5460.0000424) (cit. on p. 15).
- O'Brien, J. F., & Hodgins, J. K. (1995). Dynamic simulation of splashing fluids. *Proceedings Computer Animation, CA 1995*, 198–205. <https://doi.org/10.1109/CA.1995.393532> (cit. on p. 93).
- Olsen, J. (2004). Realtime procedural terrain generation. *Department of Mathematics And Computer Science* (..., 20 (cit. on p. 76).
- Onoue, K., & Nishita, T. (2000). A method for modeling and rendering dunes with wind-ripples. *Proceedings - Pacific Conference on Computer Graphics and Applications, 2000-Janua*, 427–428. <https://doi.org/10.1109/PCCGA.2000.883978> (cit. on p. 94).
- Oron, S., Akkaynak, D., Tchernov, B. N. G., & Shaked, Y. (2023). How monster storms shape fringing reefs: Observations from the 2020 middle east cyclone. *Ecosphere*, 14. <https://doi.org/10.1002/ecs2.4602> (cit. on p. 45).
- Paris, A., Guérin, E., Peytavie, A., Collon, P., & Galin, E. (2021). Synthesizing geologically coherent cave networks. *Computer Graphics Forum*, 40, 277–287. <https://doi.org/10.1111/cgf.14420> (cit. on pp. 67, 68, 89).

- Paris, A., Peytavie, A., Guérin, E., Argudo, O., & Galin, E. (2019a). Desertscape simulation. *Computer Graphics Forum*, 38, 47–55. <https://doi.org/10.1111/cgf.13815> (cit. on pp. 42, 79, 90, 94).
- Paris, A. (2023). Modeling and simulating virtual terrains (cit. on p. 17).
- Paris, A., Galin, E., Peytavie, A., Guérin, E., & Gain, J. (2019b). Terrain amplification with implicit 3d features. *ACM Transactions on Graphics*, 38, 1–15. <https://doi.org/10.1145/3342765> (cit. on pp. 32, 78, 88, 92).
- Park, M. J. (2010). Guiding flows for controlling crowds. *Visual Computer*, 26, 1383–1391. <https://doi.org/10.1007/s00371-009-0415-4> (cit. on p. 3).
- Patel, D., Natali, M., Lidal, E. M., Parulek, J., Brazil, E. V., & Viola, I. (2021). Modeling terrains and subsurface geology. *Interactive Data Processing and 3D Visualization of the Solid Earth*, 1–43. <https://doi.org/10.1007/978-3-030-90716-7\1> (cit. on pp. 32, 61).
- Peytavie, A., Galin, E., Grosjean, J., & Merillou, S. (2009). Arches: A framework for modeling complex terrains. *Computer Graphics Forum*, 28, 457–467. <https://doi.org/10.1111/j.1467-8659.2009.01385.x> (cit. on pp. 78, 79, 85).
- Pratt, M. J., Wysession, M. E., Aleqabi, G., Wiens, D. A., Nyblade, A. A., Shore, P., Rambolamana, G., Andriampenomanana, F., Rakotondraibe, T., Tucker, R. D., Barruol, G., & Rindraharisaona, E. (2017). Shear velocity structure of the crust and upper mantle of madagascar derived from surface wave tomography. *Earth and Planetary Science Letters*, 458, 405–417. <https://doi.org/10.1016/j.epsl.2016.10.041> (cit. on p. 30).
- Pytel, A., & Mann, S. (2015). Procedural modeling of cave-like channels. (Cit. on p. 67).
- Ranz, W. E., Talandis, G. R., & Guterman, B. (1960). Mechanics of particle bounce. *AIChE Journal*, 6, 124–127. <https://doi.org/10.1002/aic.690060123> (cit. on p. 95).
- Richardson, J. F., & Zaki, W. N. (1954). The sedimentation of a suspension of uniform spheres under conditions of viscous flow. *Chemical Engineering Science*, 3 (cit. on p. 82).
- Rigaudière, D., Gesquière, G., & Faudot, D. (2000). Shape modelling with skeleton based implicit primitives. *Methods* (cit. on p. 95).
- Roa, T., & Benes, B. (2004). Simulating desert scenery. *Winter School of Computer Graphics SHORT communication Papers Proceedings*, 17–22 (cit. on pp. 79, 94).
- Roose, D., Leuven, K. U., & López, Y. R. (2011). Dynamic refinement for fluid flow simulations with sph particle refinement for fluid flow simulations with sph. (Cit. on pp. 15, 83).
- Roudier, P. (1993). Synthèse de paysages réalistes par simulation de processus d'érosion (cit. on p. 76).
- Runions, A. (2008). Modeling biological patterns using the space colonization algorithm. *A Thesis submitted to the faculty of graduate studies for degree of master of science*, 74 (cit. on p. 67).
- Schott, H., Paris, A., Fournier, L., Guérin, E., & Galin, E. (2023). Large-scale terrain authoring through interactive erosion simulation. *ACM Transactions on Graphics*, 42, 1–15. <https://doi.org/10.1145/3592787> (cit. on p. 32).

- Shadrick, J. R., Rood, D. H., Hurst, M. D., Piggott, M. D., Hebditch, B. G., Seal, A. J., & Wilcken, K. M. (2022). Sea-level rise will likely accelerate rock coast cliff retreat rates. *Nature Communications*, 13. <https://doi.org/10.1038/s41467-022-34386-3> (cit. on p. 30).
- Shi, Z., Fonseca, J. A., & Schlueter, A. (2017). A review of simulation-based urban form generation and optimization for energy-driven urban design. *Building and Environment*, 121, 119–129. <https://doi.org/10.1016/j.buildenv.2017.05.006> (cit. on p. 3).
- Smelik, R. M., Kraker, K. J. D., Groenewegen, S. A., Tutenel, T., & Bidarra, R. (2009). A survey of procedural methods for terrain modelling. *Proceedings of the CASA workshop on 3D advanced media in gaming and simulation (3AMIGAS)* (cit. on p. 76).
- Stachniak, S., & Stuerzlinger, W. (2005). An algorithm for automated fractal terrain deformation. In *Proceedings of Computer Graphics and Artificial Intelligence*, 64–76 (cit. on p. 76).
- Stam, J. (1999). Stable fluids. *Proceedings of the 26th Annual Conference on Computer Graphics and Interactive Techniques, SIGGRAPH 1999*, 121–128. <https://doi.org/10.1145/311535.311548> (cit. on pp. 14, 90).
- Stam, J. (2003a). Flows on surfaces of arbitrary topology. *ACM Transactions on Graphics*, 22, 724–731. <https://doi.org/10.1145/882262.882338> (cit. on p. 88).
- Stam, J. (2003b). Real-time fluid dynamics for games. *Proceedings of the Game Developer Conference*, 18, 17 (cit. on p. 3).
- Stokes, G. G. (1850). On the effect of the internal friction of fluids on the motion of pendulums. Pitt Press Cambridge. <https://doi.org/10.1017/CBO9780511702266.002> (cit. on p. 82).
- Swope, W. C., Andersen, H. C., Berens, P. H., & Wilson, K. R. (1982). A computer simulation method for the calculation of equilibrium constants for the formation of physical clusters of molecules: Application to small water clusters. *The Journal of Chemical Physics*, 76, 637–649. <https://doi.org/10.1063/1.442716> (cit. on p. 83).
- Talgorn, F. X., & Belhadj, F. (2018). Real-time sketch-based terrain generation. *ACM International Conference Proceeding Series*, 13–18. <https://doi.org/10.1145/3208159.3208184> (cit. on p. 32).
- Temuçin, M. B., Kocabas, İ., & Oğuz, K. (2020). Using cellular automata as a basis for procedural generation of organic cities. *European Journal of Engineering Research and Science*, 5, 116–120. <https://doi.org/10.24018/ejers.2020.5.12.2293> (cit. on p. 4).
- Tompson, J., Schlachter, K., Sprechmann, P., & Perlin, K. (2017). Accelerating eulerian fluid simulation with convolutional networks. *34th International Conference on Machine Learning, ICML 2017*, 7, 5258–5267 (cit. on p. 15).
- Tychonievich, L. A., & Jones, M. D. (2010). Delaunay deformable mesh for the weathering and erosion of 3d terrain. *Visual Computer*, 26, 1485–1495. <https://doi.org/10.1007/s00371-010-0506-2> (cit. on p. 83).
- Valette, G., Herbin, M., Lucas, L., & Léonard, J. (2005). A preliminary approach of 3d simulation of soil surface degradation by rainfall. *Natural Phenomena*, 41–50 (cit. on p. 5).
- Vergne, R., Vanderhaeghe, D., Chen, J., Barla, P., Granier, X., & Schlick, C. (2011). Implicit brushes for stylized line-based rendering. *Computer Graphics Forum*, 30, 513–522. <https://doi.org/10.1111/j.1467-8659.2011.01892.x> (cit. on p. 3).

- Verlet, L. (1967). Computer "experiments" on classical fluids. i. thermodynamical properties of lennard-jones molecules. *Physical Review*, 159, 98–103. <https://doi.org/10.1103/PhysRev.159.98> (cit. on p. 83).
- Vermeulen, T., Knopf-Lenoir, C., Villon, P., & Beckers, B. (2015). Urban layout optimization framework to maximize direct solar irradiation. *Computers, Environment and Urban Systems*, 51, 1–12. <https://doi.org/10.1016/j.compenvurbsys.2015.01.001> (cit. on p. 3).
- Vila-Concejo, A., & Kench, P. (2016, August). Storms in coral reefs. Wiley Blackwell. <https://doi.org/10.1002/9781118937099.ch7> (cit. on p. 45).
- Villanueva, A. J., Marton, F., & Gobbetti, E. (2017). Symmetry-aware sparse voxel dags (ssvdags) for compression-domain tracing of high-resolution geometric scenes. *Journal of Computer Graphics Techniques*, 6, 1–30 (cit. on p. 10).
- Wampler, K., & Popovi, Z. (2009). Optimal gait and form for animal locomotion. *ACM Transactions on Graphics*, 28. <https://doi.org/10.1145/1531326.1531366> (cit. on p. 5).
- Wejchert, J., & Haumann, D. (1991). Animation aerodynamics. *Proceedings of the 18th Annual Conference on Computer Graphics and Interactive Techniques, SIGGRAPH 1991*, 25, 19–22. <https://doi.org/10.1145/122718.122719> (cit. on pp. 33, 36, 42).
- Wojtan, C., Carlson, M., Mucha, P. J., & Turk, G. (2007). Animating corrosion and erosion. *Natural Phenomena*, 15–22 (cit. on pp. 80–83).
- Xing, J., Kazi, R. H., Grossman, T., Wei, L. Y., Stam, J., & Fitzmaurice, G. (2016). Energy-brushes: Interactive tools for illustrating stylized elemental dynamics. *UIST 2016 - Proceedings of the 29th Annual Symposium on User Interface Software and Technology*, 755–766. <https://doi.org/10.1145/2984511.2984585> (cit. on p. 3).
- Yan, P., Zhang, J., Kong, X., & Fang, Q. (2020). Numerical simulation of rockfall trajectory with consideration of arbitrary shapes of falling rocks and terrain. *Computers and Geotechnics*, 122. <https://doi.org/10.1016/j.compgeo.2020.103511> (cit. on pp. 82, 95).
- Yersin, B., Maim, J., Pettré, J., & Thalmann, D. (2009). Crowd patches. *Proceedings of the 2009 symposium on Interactive 3D graphics and games*, 207–214. <https://doi.org/10.1145/1507149.1507184> (cit. on p. 3).
- Zhou, X., Benes, B., Chang, P., & Cani, M. P. R. (2021). Urban brush: Intuitive and controllable urban layout editing. *UIST 2021 - Proceedings of the 34th Annual ACM Symposium on User Interface Software and Technology*, 796–814. <https://doi.org/10.1145/3472749.3474787> (cit. on p. 3).

Glossary

Binary voxel A voxel that can represent two states (e.g., filled or empty), used for creating binary volumetric models where each voxel is either solid or not..

Cellular automata A model used in computational systems where cells in a grid evolve rules based on the states of neighboring cells, applicable in various scientific fields..

Coarse height function Simplified function representing the height of a environmental object in the 2D plane, useful to estimate the altitude of a point \mathbf{p} of the terrain $\mathcal{H}(\mathbf{p})$ as a combination of the coarse height function of all environmental objects close to \mathbf{p} .. 36, 41, , 103

Computer graphics The field of computing dedicated to generating and managing visual content and images through software and hardware..

Control curve Parametric curve used in computer graphics and modeling to guide the transformation or manipulation of objects, allowing for precise control over the resulting shape..

Controlled subdivision The deliberate manipulation of subdivision processes in modeling to achieve specific forms or structures..

Coral island An island formed from coral debris and associated organic material, found in tropical ocean regions..

Deep learning A subset of machine learning involving neural networks with many layers, used for tasks like image recognition, speech recognition, and natural language processing..

Density voxel A voxel that represents the density of material at a point, used in simulations where variable material density is needed..

Diamond-square algorithm An algorithm used for generating fractal landscapes or terrains through recursive subdivision, providing a simple method for creating rough, naturally appearing terrains..

Discrete height field A specific form of terrain representation where the terrain's elevation is captured at discrete points, often organized in a grid..

Domain warping A technique in procedural generation where the input coordinates to a function or texture are distorted, used to create more organic and varied patterns..

Ecosystem A biological community of interacting organisms and their physical environment, studied in ecology to understand the relationships and dynamics within natural systems..

Elevation terrain model A model representing the elevation or altitude of various points on the Earth's surface, used in geographic studies..

Environmental attribute Value of a geographic field at one point in space and time.. iii, 29, 33–37, 43, 48, , 104, 105, 107

Environmental material Material available.. iv, 29, 33, 34, 36, 37, 41–44, 48, , 104, 107

Environmental modifier Description of a modification of a environmental attribute around a environmental object. It can represent the spread or the absorption of material around it, or the deformation of water currents.. iv, 29, 34–36, 41,

Environmental object Geographic feature represented sparsly which describe a landscape. We represent it with a skeleton, generation rules and environmental materials.. iii, iv, 27, 29, 33–44, 46–49, , 103–105, 107

Erosion simulation The computational imitation of geological erosion processes in digital environments to create more realistic landscapes in games and simulations..

Faulting A process that simulates geological faults, creating abrupt changes in elevation to mimic earth's tectonic movements..

Feature-based construction technique The integration and manipulation of specific terrain features such as ridges, valleys, cliffs, and riverbeds, directly into the terrain model. It allows for precise control over the placement and characteristics of these features, enhancing the realism and complexity of the generated landscape..

Fine tuning The process of making small adjustments to a model or algorithm to optimize its performance or output quality..

Fitness function Function affected to a environmental object that, given the environmental attribute at a (x, y) position in the space, returns a score describing how well this environmental object may survive.. iii, 29, 33–35, 37, 38, 44, 47, 48, , 105, 107

Fluid simulation A computational method to model fluid behaviors in graphics, gaming, and simulation, often using algorithms that calculate the motion and interaction of liquids and gases..

Fluid solver A software or algorithm designed to calculate and predict the behavior of fluids within a simulation environment by solving equations relative to fluid dynamics..

Fractal geometry The study of shapes and patterns that repeat at increasingly fine scales, used in procedural generation to create complex, self-similar textures and structures..

Fractional Brownian motion A random motion whose changes are self-similar and depend on a scaling factor, used in graphics to generate landscapes, clouds, and other natural phenomena..

Generation rule Composed of a fitness function and skeleton fitting function, the optimisation of the generation rule of an environmental object through the maximization of its components describe where and how new elements can be added to a semantic terrain.. 34, , 104, 105

Generative adversarial network A class of machine learning systems where two neural networks contest with each other in a game (as in "adversarial"), typically used to generate realistic synthetic data..

Geographic information science A scientific discipline that encompasses the theoretical aspects, methodologies, and applications of spatial data, focusing on the study and improvement of geographic information systems (GIS) and related technologies. It deals with the principles behind collecting, analyzing, and visualizing geographic data, including spatial analysis, data management, and the integration of various data forms..

Geographic information system A system designed to capture, store, manipulate, analyze, manage, and present spatial or geographic data..

Geomorphic event Modification of a environmental attribute in an interval of time with or without spatial bounds.. 33, 34, 43–46, 48,

Gradient noise function A type of noise used in computer graphics to generate smoothly varying random patterns, often used for textures and terrain modeling..

Implicit height field A field representing variations in height across a surface, defined through mathematical functions rather than discrete measurements..

Implicit surface A surface in 3D space defined by an equation, used in computer graphics to generate complex geometrical shapes procedurally..

Implicit volume A type of 3D model defined mathematically by a function that describes whether points in space are inside or outside the volume..

Karst network Complex underground drainage systems and structures found in soluble rock landscapes, important in geological studies..

Landscape feature Distinct attributes or formations within a terrain, such as hills, valleys, or rivers, that define the character of the landscape..

Large-scale terrain Refers to large digital terrains used in simulations and games that require extensive geographical details and can often be explored interactively..

Latent space A representation in machine learning where inputs are converted to a set of abstract dimensions, used to generate new data instances with similar properties..

Layered material terrain model A representation that incorporates different material layers to depict the geological stratification of an area..

Level of detail Amount of graphical data or information displayed in a scene or image. Higher LOD means more detail, while lower LOD means less detail. This concept helps manage computational resources, where less important or distant objects have lower LOD to improve performance..

Machine learning The study of algorithms and statistical models that computer systems use to perform specific tasks without explicit instructions, relying on patterns and inference..

Material voxel A voxel that carries information about the material type at its location..

Multi-fractal noise Noise generated with fractal properties that vary spatially, used to create more complex, varied textures and terrains..

Neural network A computing system inspired by the structure and functions of human brain neural networks, designed to recognize patterns and solve specific problems by learning from data. It consists of layers of interconnected nodes (neurons) that can learn to perform tasks by considering examples, generally without being programmed with any task-specific rules..

Noise function A function that generates a pattern of noise used to add surface detail or randomness to computer graphics, often used in texture and terrain generation..

Offline rendering The process of generating a high-quality image or animation frame by allowing the computer more processing time, typically used in film and animation production..

Parallelization The process of dividing computational tasks into multiple components that can be processed simultaneously, improving performance and speed, especially in graphics and simulation..

Parameter space Set of adjustable parameters that can be modified to control and influence the generation process of content or assets. These parameters can include properties like size, shape, density, or other relevant factors, depending on the specific procedural algorithm being used. By adjusting these parameters, creators can achieve a wide variety of outcomes and patterns, enabling the generation of diverse and customizable content such as textures, levels, or 3D models..

Procedural generation A method for creating data algorithmically rather than manually, used across various fields to automate content creation..

Real-time rendering The process of generating images from 3D models at a fast enough rate to provide interaction and movement, as seen in video games and simulations..

Regeneration The process of re-creating or updating digital content dynamically, often used in procedural generation to modify terrains or environments based on user interaction or changes in parameters..

Ridge noise A type of noise function used in procedural terrain generation to create ridged, mountainous landscapes with sharp crests and rugged features..

Seascape An underwater environment or ecosystem, important in marine biology and underwater photography..

Simplex noise A method for constructing a gradient noise similar to Perlin noise but with fewer directional artifacts and lower computational overhead. It is used in higher-dimensional graphics for more efficient and visually appealing results..

Skeleton Simplified shape of a environmental object, as it could be symbolized in a map: as a point, a curve or a region.. 35, 36, 38–40,, 104

Skeleton fitting function Function affected to a environmental object that, given the local environmental attribute and the shape of the skeleton of the environmental object, returns a score describing how well this environmental object fits. It may be seen as a refinement of the fitness function.. iii, 29, 34, 35, 37–40,, 105

Sketch-based interface An interface design that allows users to interact with digital systems through sketching or drawing..

Small-scale terrain Refers to small digital terrains with more detailed segments of terrain used in close-up views or where high levels of detail are required for visual effects or analysis..

Smooth step A mathematical function used in graphics programming to interpolate smoothly between two values with zero derivatives at both ends, useful in shading and texture mapping, but also in blending together two different functions..

Smoothed Particle Hydrodynamics (SPH) A particle-based method for simulating the physics of fluids and gases, using particles to approximate the properties and motion of the fluid..

Smoothing kernel A function used in particle-based simulations (like SPH) to smooth out values across particles, enhancing stability and realism in fluid simulations..

Sparse voxel octree An efficient data structure that stores voxel data in a tree hierarchy, reducing memory usage by only storing data for regions that contain relevant information..

Sparseness A property of data or procedural models where significant portions of the space contain no actual data or minimal features, often used to optimize storage and processing..

Steady state A system is at steady state when internal properties do not change overtime once a dynamic equilibrium has been reached. The environmental materials emission from environmental objects can be seen as a thermodynamic system which achieve thermodynamic equilibrium over time.. 36,

Subdivision scheme A technique in computer graphics used to refine a mesh by splitting each polygonal face into smaller faces, improving smoothness and detail. This method allows for progressively higher levels of detail in 3D models and images..

Terrain generation The process of algorithmically creating landforms in virtual environments, often using mathematical models and noise functions to simulate natural terrains..

Terrain representation Different methods used to depict geographical features in two-dimensional and three-dimensional formats..

Texture The visual appearance or feel of a surface. In computer graphics, textures are bitmap images applied to the surface of a 3D model to add color, detail, and material properties, simulating the look of real-world materials..

Variational autoencoder A type of neural network that aims to encode input data into a latent representation, then reconstruct the input data from this representation, used in unsupervised learning tasks..

Virtual terrain Digitally simulated landscape used in video games, simulations, and visualizations, which mimics real-world or imaginary terrains..

Volumetric terrain model A 3D representation of terrain that includes volume, allowing for the depiction of complex underground structures and features..

Voxel The 3D equivalent of a pixel, representing a value on a regular grid in three-dimensional space, used for constructing volumetric environments and objects..

Voxel directed acyclic graph A data structure that organizes voxels in a hierarchical, non-cyclic manner, enabling efficient storage and manipulation of volumetric data in graphics and spatial analysis..

Voxel grid terrain model A terrain representation based on a grid of voxels, each storing information about the terrain's properties at that point..

Warp function $\Phi : \mathbb{R}^n \mapsto \mathbb{R}^n$ is a function that, given a n -dimensional point \mathbf{p} returns another point \mathbf{q} . The function is usually bijective..

Warping The process of distorting a field to modify structural and visual properties..

Worley noise A noise function that generates a cellular pattern, used in graphics to create textures that mimic certain natural materials like stone or wood..