

1. The field of complex numbers
--

introduction

The field of complex numbers

Let:

$$\begin{aligned} \cdot \quad + : \mathbb{R}^2 \times \mathbb{R}^2 &\longrightarrow \mathbb{R}^2 \\ \cdot \quad ((a, b), (c, d)) &\longmapsto (a + c, b + d) \\ \cdot \quad \cdot : \mathbb{R}^2 \times \mathbb{R}^2 &\longrightarrow \mathbb{R}^2 \\ \cdot \quad ((a, b), (c, d)) &\longmapsto (ac - bd, ad + bd) \end{aligned}$$

We name the field of complex numbers to:

$$\cdot (\mathbb{R}^2, +, \cdot)$$

We denote:

$$\cdot (\mathbb{R}^2, +, \cdot) : \mathbb{C}$$

$$\cdot (0, 1) \in \mathbb{C} : i$$

$$\cdot (a, b) \in \mathbb{C} : a + bi$$

$$\cdot \pi_1(\mathbb{C}) : Re$$

$$\cdot \pi_2(\mathbb{C}) : Im$$

Conjugation

We name complex conjugation to:

$$\begin{aligned} \cdot \quad f : \mathbb{C} &\longrightarrow \mathbb{C} \\ \cdot \quad (a, b) &\longmapsto (a, -b) \end{aligned}$$

We denote:

$$\cdot f((a, b)) : \overline{(a, b)}$$

Norm

We name complex norm to:

$$\begin{array}{lcl} f : \mathbb{C} & \longrightarrow & \mathbb{R} \\ (a, b) & \longmapsto & \sqrt{a^2 + b^2} \end{array}$$

We denote:

$$\cdot f((a, b)) : |(a, b)|$$

Polar transformation

We name polar transformation to:

$$\begin{array}{lcl} f : \mathbb{C} & \longrightarrow & \mathbb{R}^+ \times [0, 2\pi) \\ (a, b) & \longmapsto & (\sqrt{|(a, b)|}, \arctan(\frac{b}{a})) \end{array}$$

We denote:

$$\cdot f((a, b)) : (r, \theta)$$

Unit sphere projection

We name unit sphere projection to:

$$\begin{array}{lcl} f : \mathbb{C} & \longrightarrow & S^1 \\ z & \longmapsto & \frac{z}{|z|} \end{array}$$

We denote:

$$\cdot f(z) : \pi(z)$$

Roots of unity

Let:

$$\cdot z \in \mathbb{C}$$

Then, z is a root of unity if:

$$\cdot \exists n \in \mathbb{N} :$$

$$z^n = 1$$

We denote:

$$\cdot \{z \in \mathbb{C} \mid z \text{ root of unity} \} : S^1$$

Disk

Let:

$$\cdot p \in \mathbb{C}$$

$$\cdot r \in \mathbb{R}^+ \setminus \{0\}$$

We name Disk centered in p and radius r to:

$$\cdot \{z \in \mathbb{C} \mid |z - p| < r\}$$

We denote:

$$\cdot \{z \in \mathbb{C} \mid |z - p| < r\} : D^1$$

Component decomposition

Let:

$$f : \mathbb{C} \rightarrow \mathbb{C}$$

We name real component of f to:

$$\begin{aligned} f_{Re} : \mathbb{C} &\longrightarrow \mathbb{R} \\ z &\longmapsto Re(f(z)) \end{aligned}$$

We name imaginary component of f to:

$$\begin{aligned} f_{Im} : \mathbb{C} &\longrightarrow \mathbb{R} \\ z &\longmapsto Im(f(z)) \end{aligned}$$

We name component decomposition of f to:

$$\begin{aligned} f_{\mathbb{R}^2} : \mathbb{C} &\longrightarrow \mathbb{R}^2 \\ (x, y) &\longmapsto (f_{Re}(x + yi), f_{Im}(x + yi)) \end{aligned}$$