

Complex Analysis

Martin Azpillaga

March 5, 2014

Contents

I	Definitions	7
1	The field of complex numbers	7
	The field of complex numbers	9
	Conjugation	9
	Norm	9
	Polar transformation	10
	Unit sphere projection	10
	Roots of unity	10
	Disk	11
	Component decomposition	11
2	Holomorphic functions	12
	Incremental quotient	14
	Holomorphic function	14
	Cauchy-Riemman equations	15
	Conformal	16
	Power series	17
	Convergence radius	18
	Absolutely convergent	18
	Numeric series	19
II	Propositions	19
1	The field of complex numbers	21
2	Holomorphic functions	23
	Cauchy-Riemman	24
	tangent venvtor	25

arc images	26
Holomorphic functions are conform	27
Convergence of complex series	28
Absolutely convergent are convergent	29
Series and norm	30
Root test	31
Quotient test	32
Power series theorem	33
Power series theorem	35
functions associated to series are \mathcal{C}^∞	38
III Examples	39
1 Holomorphic functions	39
2 Holomorphic functions	41
Conjugation	42
Quadratic norm	42
Non preserving angles function	43
Exponential	44
Geometric series	44
Series not centered in 0	45
Radius of convergence without quotient test	46
IV Problems	46
1 The field of complex numbers	50
entity	51
2 Holomorphic functions	51
3. Cauchy-Riemman	52
B.2 a)	53
Preservation of angles	54
V Tasks	56
1 1st laboratory	57
Existence of holomorphic functions	58

block name	5
<hr/>	
Constant tests	61
Real part of holomorphic functions	63
2 2nd laboratory	65
Power series	66

Block I

Definitions

1. The field of complex numbers
--

introduction

The field of complex numbers

Let:

$$\begin{aligned} \cdot \quad & + : \mathbb{R}^2 \times \mathbb{R}^2 \longrightarrow \mathbb{R}^2 \\ & ((a, b), (c, d)) \longmapsto (a + c, b + d) \\ \cdot \quad & \cdot : \mathbb{R}^2 \times \mathbb{R}^2 \longrightarrow \mathbb{R}^2 \\ & ((a, b), (c, d)) \longmapsto (ac - bd, ad + bd) \end{aligned}$$

We name the field of complex numbers to:

$$\cdot (\mathbb{R}^2, +, \cdot)$$

We denote:

$$\cdot (\mathbb{R}^2, +, \cdot) : \mathbb{C}$$

$$\cdot (0, 1) \in \mathbb{C} : i$$

$$\cdot (a, b) \in \mathbb{C} : a + bi$$

$$\cdot \pi_1(\mathbb{C}) : Re$$

$$\cdot \pi_2(\mathbb{C}) : Im$$

Conjugation

We name complex conjugation to:

$$\begin{aligned} \cdot \quad & f : \mathbb{C} \longrightarrow \mathbb{C} \\ & (a, b) \longmapsto (a, -b) \end{aligned}$$

We denote:

$$\cdot f((a, b)) : \overline{(a, b)}$$

Norm

We name complex norm to:

$$\begin{aligned} f : \mathbb{C} &\longrightarrow \mathbb{R} \\ (a, b) &\longmapsto \sqrt{a^2 + b^2} \end{aligned}$$

We denote:

$$\cdot f((a, b)) : |(a, b)|$$

Polar transformation

We name polar transformation to:

$$\begin{aligned} f : \mathbb{C} &\longrightarrow \mathbb{R}^+ \times [0, 2\pi) \\ (a, b) &\longmapsto (\sqrt{|(a, b)|}, \arctan(\frac{b}{a})) \end{aligned}$$

We denote:

$$\cdot f((a, b)) : (r, \theta)$$

Unit sphere projection

We name unit sphere projection to:

$$\begin{aligned} f : \mathbb{C} &\longrightarrow S^1 \\ z &\longmapsto \frac{z}{|z|} \end{aligned}$$

We denote:

$$\cdot f(z) : \pi(z)$$

Roots of unity

Let:

$$\cdot z \in \mathbb{C}$$

Then, z is a root of unity if:

$$\cdot \exists n \in \mathbb{N} :$$

$$z^n = 1$$

We denote:

$$\cdot \{z \in \mathbb{C} \mid z \text{ root of unity} \} : S^1$$

Disk

Let:

$$\cdot p \in \mathbb{C}$$

$$\cdot r \in \mathbb{R}^+ \setminus \{0\}$$

We name Disk centered in p and radius r to:

$$\cdot \{z \in \mathbb{C} \mid |z - p| < r\}$$

We denote:

$$\cdot \{z \in \mathbb{C} \mid |z - p| < r\} : D^1$$

Component decomposition

Let:

$$f : \mathbb{C} \rightarrow \mathbb{C}$$

We name real component of f to:

$$\begin{aligned} f_{Re} : \mathbb{C} &\longrightarrow \mathbb{R} \\ z &\longmapsto Re(f(z)) \end{aligned}$$

We name imaginary component of f to:

$$\begin{aligned} f_{Im} : \mathbb{C} &\longrightarrow \mathbb{R} \\ z &\longmapsto Im(f(z)) \end{aligned}$$

We name component decomposition of f to:

$$\begin{aligned} f_{\mathbb{R}^2} : \mathbb{C} &\longrightarrow \mathbb{R}^2 \\ (x, y) &\longmapsto (f_{Re}(x + yi), f_{Im}(x + yi)) \end{aligned}$$

2. Holomorphic functions**Incremental quotient**

Let:

$$\cdot \mathcal{U} \subset \mathbb{C} \text{ open}$$

$$\cdot f : \mathcal{U} \rightarrow \mathbb{C}$$

$$\cdot p \in \mathcal{U}$$

We name incremental quotient of f in p to:

$$\cdot \lim_{z \rightarrow p} \frac{f(z) - f(p)}{z - p}$$

We denote:

$$\cdot f'(p)$$

Holomorphic function

Let:

$$\cdot \mathcal{U} \subset \mathbb{C} \text{ open}$$

$$\cdot f : \mathcal{U} \rightarrow \mathbb{C}$$

$$\cdot p \in \mathcal{U}$$

Then, f is holomorphic over p if:

$$\cdot \exists f'(p)$$

Then, f is holomorphic over U if:

$$\cdot \forall p \in \mathcal{U} :$$

$$\exists f'(p)$$

We denote:

$$\cdot \{f : \mathcal{U} \rightarrow \mathbb{C} \mid f \text{ holomorphic over } \mathcal{U}\} : \mathcal{H}(\mathcal{U})$$

$$\cdot f \in \mathcal{H}(\mathbb{C}) : f \text{ entire}$$

Cauchy-Riemman equations

Let:

$$\begin{aligned} \cdot & u, v : \mathbb{R}^2 \rightarrow \mathbb{R} \\ \cdot & f : \mathbb{R}^2 \longrightarrow \mathbb{R}^2 \\ \cdot & (x, y) \longmapsto (u((x, y)), v((x, y))) \end{aligned}$$

Then, f is satisfies the Cauchy-Riemman equations if:

$$\begin{aligned} \cdot & \exists u_x, u_y, v_x, v_y \\ \cdot & u_x = v_y \\ \cdot & u_y = -v_x \end{aligned}$$

We denote:

$$\begin{aligned} \cdot & u_x + iv_x : f_x \\ \cdot & u_y + iv_y : f_y \end{aligned}$$

Conformal

Let:

$$\cdot \mathcal{U} \subset \mathbb{C} \text{ open}$$

$$\cdot f : \mathcal{U} \rightarrow \mathbb{C}$$

$$\cdot z \in \mathcal{U}$$

Then, f is conformal in z if:

$$\cdot \exists c \in \mathbb{C} :$$

$$\forall I \subset \mathbb{R} \quad 0 \in I :$$

$$\forall \gamma : I \rightarrow \mathbb{R}^2 \quad \gamma \text{ differentiable} \wedge \gamma(0) = z \wedge \gamma'(0) \neq$$

$0 :$

$$\frac{(f \circ \gamma)'(0)}{\gamma'(0)} = c$$

Then, f is conformal if:

$$\cdot \forall z \in \mathcal{U} :$$

$$f \text{ conformal in } z$$

Power series

Let:

$$\cdot \sum_{n \geq 0} a_n f_n \text{ complex valued sequence}$$

Then, $\sum_{n \geq 0} a_n f_n$ is a power series if:

$$\cdot \forall n \in \mathbb{N} :$$

$$\begin{array}{ccc} f_n : \mathbb{C} & \longrightarrow & \mathbb{C} \\ z & \longmapsto & (z - a)^n \end{array}$$

Convergence radius

Let:

$$\cdot \text{statements}$$

.

Then, *item* is a/an entity if:

$$\cdot \text{conditions}$$

.

We denote:

$$\cdot \text{property} : \text{notation}$$

.

Absolutely convergent

Let:

· $\sum c_n n[0]$ series

Then, $\sum c_n n[0]$ is absolutely convergent if:

· $\sum |c_n| n[0]$ convergent

Numeric series

Let:

· $(c_k)_{k \in \mathbb{N}}$

Then, *item* is a/an entity if:

· *conditions*

.

We denote:

· *property : notation*

.

Block II

Propositions

1. The field of complex numbers
--

introduction

go

2. Holomorphic functions**Cauchy-Riemman**

Let:

$$\cdot f : \mathbb{R}^2 \rightarrow \mathbb{R}^2 \text{ satisfies CR}$$

Then, holds:

$$\cdot f_x = f'(z)$$

$$\cdot f_y = -if_y$$

$$\cdot \partial_{f(z)} z = 0$$

Demonstration:

$$f_x = u_x + iv_x = a + bi = f'(z)$$

$$f_y = i(a + ib) = if'(z)$$

$$f'(z) = -if_y$$

notation

tangent vector

Let:

γ differentiable plane arc $\gamma : I \rightarrow \mathbb{R}^2, \forall t \in I :$

$$\gamma'(t) \neq 0$$

Then, holds:

$\gamma'(t)$ tangent to γ

Demonstration:

demonstration

arc images

Let:

$$\cdot f \in \mathcal{H}(\mathcal{U})$$

$$\cdot \gamma \text{ differentiable plane arc} \quad \gamma \subset \mathcal{U}$$

$$\cdot \sigma = f(\gamma)$$

$$\cdot z_0 = \gamma(0)$$

Then, holds:

$$\cdot \sigma' = f'(\gamma)\gamma'$$

$$\cdot \gamma'(0) \neq 0 \rightarrow f'(z_0) \neq 0$$

$$\cdot \sigma'(0) = f'(z_0)\gamma'(0)$$

$$\cdot |\sigma'(0)| = |f'(z_0)| |\gamma'(0)|$$

$$\cdot \arg \sigma'(0) = \arg \gamma'(0) + \arg f'(z_0)$$

$$\cdot f \text{ aplica una homotecia mas una rotacion constante a todos los}$$

vectores tangentes que salen de z_0

Demonstration:

obvio

Holomorphic functions are conform

Let:

$$f : \mathcal{U} \rightarrow \mathbb{C}$$

.

Then, holds:

$$f \text{ holomorph in } z \iff f'(z) \neq 0 \leftrightarrow f \text{ conform}$$

Demonstration:

\rightarrow):

already seen

\leftarrow):

too hard

Convergence of complex series

Let:

$$\cdot \sum_{n \geq 0} c_n \text{ complex series}$$

Then, holds:

$$\cdot \sum_{n \geq 0} c_n \text{ convergent} \leftrightarrow \sum_{n \geq 0} \text{Re} c_n \text{ convergent} \wedge \sum_{n \geq 0} \text{Im} c_n \text{ convergent}$$

Demonstration:

demonstration

Absolutely convergent are convergent

Let:

$$\cdot \sum c_n n[0] \text{ absolutely convergent}$$

Then, holds:

$$\cdot \sum c_n n[0] \text{ convergent}$$

Demonstration:

$$S_k := \sum c_n n[0][k]$$

$$\forall m \in \mathbb{N} \quad m < k :$$

$$|S_k - S_m| = |\sum c_n n[m+1][k]| \leq \sum |c_n| n[m+1][k]$$

$$\leq \sum |c_n| n[m+1] \xrightarrow{n} 0$$

$$|S_k - S_m| \xrightarrow{n} 0 \rightarrow (S_k)_k \text{ convergent} \rightarrow \sum c_n n[0] \text{ convergent}$$

gent

Series and norm

Let:

$$\cdot \sum c_n n[0] \text{ convergent}$$

Then, holds:

$$\cdot |c_n| \xrightarrow{n} 0$$

Demonstration:

$$\sum c_n n[0] \text{ convergent} \leftrightarrow (S_n)_n \text{ convergent}$$

$$\rightarrow \text{Cauchy } |S_n - S_m| \xrightarrow{n} 0 \text{ por n y m} \rightarrow |S_n - S_{n-1}| \xrightarrow{n} 0$$

$$\rightarrow |c_n| \xrightarrow{n} 0$$

Root test

Let:

$$\begin{aligned} & \cdot \sum_{n \geq 0} c_n \text{ real series} \\ & \cdot l \in \mathbb{R} \quad \text{, } \lim_{k \rightarrow \infty} |c_k|^{\frac{1}{k}} = l \end{aligned}$$

Then, holds:

$$\begin{aligned} & \cdot l > 1 \rightarrow \sum_{n \geq 0} c_n \notin \mathbb{R} \\ & \cdot l < 1 \rightarrow \sum_{n \geq 0} c_n \in \mathbb{R} \end{aligned}$$

Demonstration:

demonstration

Quotient test

Let:

$$\begin{aligned} & \cdot \sum_{n \geq 0} c_n \text{ real series} \\ & \cdot l \in \mathbb{R} \quad \text{„} \end{aligned}$$

Then, holds:

$$\begin{aligned} & \cdot \exists l \in \mathbb{R} : \\ & \quad \lim_k \frac{c_{k+1}}{c_k} = l \\ & \cdot \overline{\lim}_{c_k} |c_k|^{\frac{1}{k}} k = l \end{aligned}$$

Power series theorem

Let:

$$\cdot \sum_{n \geq 0} a_n c^n \text{ power series}$$

Then, holds:

$$\cdot |z - a| < R \rightarrow \text{absolutely convergent}$$

$$\cdot |z - a| > R \rightarrow \text{divergent}$$

$$\cdot \text{convergent in } D(a, R)$$

$$\cdot \begin{array}{ccc} f : \mathbb{C} & \longrightarrow & \mathbb{C} \\ z & \longmapsto & \sum_{n \geq 0} c_n (z - a)^n \in \mathcal{H}(D(a, R)) \end{array}$$

$$\cdot \forall z \in \mathbb{C} :$$

$$f'(z) = \sum_{n \geq 0} n c_n (z - a)^{n-1} \text{ convergent}$$

$$\cdot \text{convergence radius of } f' = \text{convergence radius of } f$$

Demonstration:

$$\forall z \in \mathbb{C} \quad \text{,, } |z - a| < R :$$

$$\begin{array}{l} \text{Root test over } \sum_{n \geq 0} |c_n| |z - a|^n \text{ rlimit}(|c_n| |z - a|^n)^{\frac{1}{n}} = |z - a| \text{ rlimit} |c_n|^{\frac{1}{n}} = \frac{|z - a|}{R} < 1 \text{ Root test} \rightarrow \text{absolutely convergent} \end{array}$$

$$\forall z \in \mathbb{C} \quad \text{,, } |z - a| < R :$$

$$\forall \rho \in \mathbb{R} \quad \text{,, } |z - a| < \rho < R :$$

$$\frac{1}{\rho} < \frac{1}{R}$$

$$\text{rlimit} |c_n|^{1/n} = \frac{1}{R} \text{ exists partial of } |c_n|^{1/n}$$

$$|c_n| |z - a|^n > \frac{|z - a|^n}{\rho^n} \text{ no } \xrightarrow{n} 0$$

General term test \rightarrow divergent

Power series theorem

Let:

.

Then, holds:

· *Part II, III, IV*

Demonstration:

Follow 3 steps

Step 1 : Uniform convergence in compacts of $D(a, R)$:

$$\begin{aligned} g_n : \mathbb{C} &\longrightarrow \mathbb{C} \\ z &\longmapsto c_n(z-a)^n \end{aligned}$$

$$\forall \rho \in \mathbb{R} \quad \text{,, } \rho < R :$$

$$\forall z \in \overline{D(a, \rho)} :$$

$$|g_n(z)| = |c_n||z-a|^n \leq |c_n|\rho^n$$

$$M_n := |c_n|\rho^n$$

$$\rho \limsup_{n \rightarrow \infty} (|c_n|\rho^n)^{1/n} = \rho \limsup_{n \rightarrow \infty} |c_n|^{1/n} \rho = \frac{\rho}{R} < 1$$

$$\text{Root test} \rightarrow \sum_{n \geq 0} M_n \text{ convergent}$$

$$\text{M-Weierstrass} \rightarrow \sum_{n \geq 0} c_n(z-a)^n \text{ uniformly convergent}$$

over compacts of $D(a, \rho)$

$$\sum_{n \geq 0} c_n(z-a)^n \text{ uniformly convergent over compacts of } D(a, r)$$

$$f(z) := \sum_{n \geq 0} g_n(z)$$

g uniformly convergent $\rightarrow f$ continuous

Step 2 : IV. R' :

$$\tilde{f}(z) := \sum_{n \geq 1} n c_n (z - a)^{n-1}$$

$$\tilde{f}(z) = \sum_{n \geq 0} (n+1) c_{n+1} (z - a)^n$$

$$\frac{1}{R'} = rlimit(n+1) |c_{n+1}|^{1/n} n = rlimit(n+1)^{1/n} |c_{n+1}|^{1/n} n =$$

$$rlimit(|c_{n+1}|^{1/n+1})^{\frac{n+1}{n}} n = \frac{1}{R}$$

$$R' = R$$

Step 3 : III :

\tilde{f} well defined in $D(a, R)$

$\forall z_0 \in D(a, R) :$

$$\left| \frac{f(z) - f(z_0)}{z - z_0} - \tilde{f}(z_0) \right| \xrightarrow{n} 0?$$

$\forall n \in \mathbb{N} :$

$$S_n(z) := \sum_{k=0}^n f_k(z)$$

$$R_n(z) := \sum_{k \geq n+1} f_k(z)$$

$$f(z) = S_n(z) + R_n(z)$$

$$\tilde{f}(z) = \tilde{S}_n(z) + \tilde{R}_n(z)$$

$$R_n(z), \tilde{R}_n(z) \xrightarrow{n} 0$$

$$\forall \varepsilon \in \mathbb{R}^+ :$$

$$\left| \frac{f(z)-f(z_0)}{z-z_0} \right| = \left| \frac{S_n(z)-S_n(z_0)}{z-z_0} + \frac{R_n(z)-R_n(z_0)}{z-z_0} - \tilde{S}_n(z_0) - \tilde{R}_n(z_0) \right|$$

$$\forall \rho \in \mathbb{R} \quad \text{,, } |z_0 - a| < \rho < R :$$

$$|\tilde{R}_n(z_0)| \leq \sum_{k \geq n+1} k |c_k| \rho^{k-1} < \frac{\varepsilon}{3} (n \geq n_1)$$

$$\begin{aligned} \left| \frac{R_n(z)-R_n(z_0)}{z-z_0} \right| &\leq \sum_{k \geq n+1} |c_k| \left| \frac{(z-a)^k - (z_0-a)^k}{z-z_0} \right| \\ \frac{a^k - b^k}{a-b} &= a^{k-1} - a^{k-2}b + \dots + b^{k-1} \\ &\leq \sum_{k \geq n+1} |c_k| (|z-a|^{k-1} + |z-a|^{k-2}|z_0-a| + \dots + \\ &\quad |z_0-a|^{k-1} \end{aligned}$$

$$|z-a|, |z_0| < \rho$$

$$\begin{aligned} &\leq \sum_{k \geq n+1} |c_k| k \rho^{k-1} < \frac{\varepsilon}{3} (n \geq n_1) \\ \left| \frac{S_n(z)-S_n(z_0)}{z-z_0} - \tilde{S}_n(z_0) \right| &< \frac{\varepsilon}{3} (S'_n = \tilde{S}_n) (n \geq n_2) \end{aligned}$$

$$\forall n \in \mathbb{N} \quad \text{,, } n \geq \max(n_1, n_2) :$$

$$\left| \frac{f(z)-f(z_0)}{z-z_0} \right| < \varepsilon$$

functions associated to series are \mathcal{C}^∞

Let:

$$\cdot f(z) = \sum c_n (z-a)^n n[0] \text{ series}$$

$$\cdot R \text{ radius of convergence of } f$$

Then, holds:

$$\cdot f \in \mathcal{C}^\infty \text{ over } D(a, R)$$

$$\cdot \forall n \in \mathbb{N} :$$

$$f^{(n)} \in \mathcal{H}(D(a, R))$$

$$\cdot c_k = \frac{f^{(k)}}{k!}$$

$$\cdot \text{series associated to } f \text{ is unique}$$

Demonstration:

$$f^{(k)}(z) = \sum n(n-1) \dots (n-k+1) c_n (z-a)^{n-k} n[k]$$

$$f^{(k)}(a) = k! c_k$$

$$c_k = \frac{f^{(k)}(a)}{k!}$$

Block III

Examples

1. Holomorphic functions

introduction

go

2. Holomorphic functions**Conjugation**

Let:

$$\begin{aligned} \bar{a} : \mathbb{C} &\longrightarrow \mathbb{C} \\ z &\longmapsto \bar{z} \end{aligned}$$

Then, \bar{a} is not holomorphic :

$$u_x = 1$$

$$u_y = 0$$

$$v_x = 0$$

$$v_y = -1$$

$$\forall z \in \mathbb{C} :$$

$$-1 \neq 1 \rightarrow f \text{ not holomorphic in } z$$

Quadratic norm

Let:

$$\begin{aligned} f : \mathbb{C} &\longrightarrow \mathbb{C} \\ z &\longmapsto |z|^2 \end{aligned}$$

$\cdot f_{\mathbb{R}^2}$ component decomposition of f

Then, f is holomorphic in 0:

f differentiable in \mathbb{R}^2 polynomial

$\forall z \in \mathbb{C} :$

$$u_x(x, y) = 2x$$

$$u_y(x, y) = 2y$$

$$v_x(x, y) = 0$$

$$v_y(x, y) = 0$$

$$u_x = v_y \leftrightarrow x = 0$$

$$u_y = -v_x \leftrightarrow y = 0$$

f holomorphic function in $z \leftrightarrow z = 0$

Non preserving angles function

Let:

$$\cdot f(z) = z^2$$

Then, f is conform in $\mathbb{R} \setminus \{0\}$:

$$f(\{(x, 0) \in \mathbb{C} \mid x > 0\}) = \{(x, 0) \in \mathbb{C} \mid x > 0\}$$

$$f(\{(x, 0) \in \mathbb{C} \mid x < 0\}) = \{(x, 0) \in \mathbb{C} \mid x > 0\}$$

$$\text{ang}(A, B) = \pi \neq 0 = \text{ang}(f(A), f(B))$$

Exponential

Let:

$$\cdot a : 0$$

$$\cdot c_n : \frac{1}{n}$$

Then, $\sum_{n \geq 0} c_n (z - a)^n$ is convergent in D_1 :

$$\lim_n \frac{|c_n|}{|c_{n+1}|} = \lim_n \frac{n+1}{n} = 1 \rightarrow R = 1$$

$\mathbb{CH} \rightarrow D(0, 1)$ convergent

$\mathbb{C} \setminus D(0, 1)$ divergent

$$f' = f$$

Geometric series

Let:

$$\cdot a : 0$$

$$\cdot c_n : 0$$

Then, $\sum z^n n[0]$ is convergent in \mathbb{D} :

$$R = \frac{c_n}{c_{n+1}} = 1 \quad \text{Then, holds:}$$

$$\cdot \sum z^n n[0] = \frac{1}{1-z}$$

$$\cdot \sum n z^{n-1} n[0] = \frac{1}{(1-z)^2}$$

$$\cdot \sum \frac{z^{n+1}}{n+1} n[0] = -\log(1-z)$$

Demonstration:

$$\forall z \in \mathbb{D} :$$

$$\sum z^n n[0] \text{ geometric series}$$

$$\sum z^n n[0] = \frac{1}{1-z}$$

II differentiating

III integrating

Series not centered in 0

Let:

$$\cdot a : i$$

$$\cdot c_n : \frac{n+1}{5^{n+1}}$$

Then, *item* is a/an entity :

$$\sum \frac{n(z-i)^{n-1}}{5^n} n[1]$$

$$= \frac{1}{5} \sum n \frac{z-i}{5} n^{n-1} n[1] = \frac{1}{5} \sum n u^{n-1} n[1]$$

$$S(u) = \tilde{S}'(u)$$

$$\tilde{S}(u) = \frac{1}{5} \sum u^n n[1] = \frac{u}{5(1-u)}$$

$$S(u) = \frac{1}{5(1-u)^2}$$

$$S(z) = \frac{5}{(5+i-z)^2} \text{ over } D(i, 5)$$

Radius of convergence without quotient test

Let:

$$\cdot \sum \frac{(-1)^n}{n(n+1)} (z-2)^{n(n+1)} n[1]$$

Then, *R* is a/an entity :

$$\lim_{c_{n+1}} c_n \nexists$$

$$\lim_n \frac{1}{n(n+1)} \frac{1}{\frac{1}{n(n+1)}} = 1$$

ignore zeros

Block IV

Problems

PROBLEMES D'ANÀLISI COMPLEXA
2n quadrimestre del curs 2013-2014.

Llista 1: Els nombres complexos

B.1. Expressen en la forma $a + ib$ els següents nombres:

- | | | | |
|-----------------------|---------------------|-----------------|------------------|
| (a) $(2 + 3i)(4 + i)$ | (c) $\frac{1}{4+i}$ | (e) \sqrt{i} | (g) $\sqrt{9i}$ |
| (b) $(4 + 2i)^2$ | (d) $\frac{i}{4+i}$ | (f) $\sqrt{-i}$ | (h) $\sqrt{1+i}$ |

B.2. Si $z = x + iy$ trobeu les parts real i imaginària de les expressions següents:

- | | | | |
|-----------|----------------|-------------------|---------------------|
| (a) z^2 | (b) $z(z + 1)$ | (c) $\frac{1}{z}$ | (d) $\frac{1}{z-3}$ |
|-----------|----------------|-------------------|---------------------|

B.3. És cert que

- a) $\operatorname{Re}(z + w) = \operatorname{Re} z + \operatorname{Re} w$?
- b) $\operatorname{Re}(zw) = (\operatorname{Re} z)(\operatorname{Re} w)$?
- c) $\operatorname{Re}\left(\frac{z}{w}\right) = \frac{\operatorname{Re} z}{\operatorname{Re} w}$?

B.4. Trobeu la forma polar dels nombres següents i dibuixeu-los.

- | | | | |
|------------------------|----------------------|---------------|--------------|
| (a) $3(1 + \sqrt{3}i)$ | (b) $2\sqrt{3} - 2i$ | (c) $-2 + 2i$ | (d) $-1 - i$ |
|------------------------|----------------------|---------------|--------------|

B.5. Sigui $(x + iy)/(x - iy) = a + ib$. Proveu que $a^2 + b^2 = 1$.

B.6. Proveu que si $p(z)$ és un polinomi amb coeficients reals i z és un zero de p llavors \bar{z} també ho és.

B.7. Descriu els conjunts del pla que satisfan (recordeu que $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$.)

- | | | |
|---|-------------------------------------|-------------------------|
| (a) $\operatorname{Im} \frac{z-a}{z} = 0, a \in \mathbb{C}^*$ | (b) $ z = \operatorname{Re} z + 1$ | (c) $ z - 2 > z - 3 $ |
|---|-------------------------------------|-------------------------|

SOL. B.1. a) $5 + 14i$; b) $12 + 16i$; c) $4/17 - i/17$; d) $1/17 + 4i/17$; e) $\pm\sqrt{2}/2(1 + i)$; f) $\pm\sqrt{2}/2(1 - i)$; g) $\pm 3\sqrt{2}/2(1 + i)$; h) $\pm 2^{1/4}(\cos(\pi/8) + i \sin(\pi/8))$.

B.2 a) $x^2 - y^2 + 2ixy$; b) $x^2 - y^2 + x + i(y + 2xy)$; c) $(x - iy)/(x^2 + y^2)$; d) $(x - 3 - iy)/((x - 3)^2 + y^2)$.

B.3 a) si. b) no. c) no.

B.4 a) $6(\cos(\pi/3) + i \sin(\pi/3))$; b) $4(\cos(\pi/6) - i \sin(\pi/6))$; c) $2\sqrt{2}(\cos(\pi/4) + i \sin(\pi/4))$; d) $\sqrt{2}(\cos(3\pi/4) - i \sin(3\pi/4))$.

B.6 Conjugueu tot el polinomi.

B.7 a) Recta que passa per 0 i a ; b) Paràbola horitzontal $x = (1/2)(y^2 - 1)$; c) $\{\operatorname{Re} z > 3/2\}$.

1. Expressen en la forma $a + ib$ els següents nombres:

- | | | | |
|-----------------------|--|---------------------------------------|------------------------------|
| (a) $\frac{1}{i}$ | (c) $\frac{1}{2+i} + \frac{1}{2-i}$ | (e) $\left(\frac{2+i}{3-2i}\right)^2$ | (g) $\sqrt[4]{-i}$ |
| (b) $\frac{1+i}{1-i}$ | (d) $\frac{1}{2+i} + \frac{4-2i}{3+i}$ | (f) $(1+i)^{100} + (1-i)^{100}$ | (h) $(3 + 4i)^{\frac{1}{2}}$ |

2. Si $z = x + iy$ on $x, y \in \mathbb{R}$, trobeu les parts real i imaginària de:

PROBLEMES D'ANLISI COMPLEXA
2n quadrimestre del curs 2013-2014

Llista 2: Funcions de variable complexa i equacions de Cauchy-Riemann

B.1. Trobeu els punts on la funció f és derivable (en el sentit complex), en els següents casos, i calculeu la derivada.

(a) $\cos |z|^2$

(c) e^{iz}

(e) $\frac{1}{(z-1)^2(z^2+2)}$

(b) $|z|^4$

(d) $z + \frac{1}{z}$

(f) $\frac{1}{(z+\frac{1}{z})^2}$

Solució: (a) \emptyset ; (b) \emptyset ; (c) \mathbb{C} ; $f'(z) = ie^{iz}$; (d) $\mathbb{C} \setminus \{0\}$; $f'(z) = 1 - \frac{1}{z^2}$; (e) $\mathbb{C} \setminus \{1, \pm\sqrt{2}i\}$; (f) \mathbb{C} .

B.2. Determineu si aquestes funcions poden ser la part real d'una funció holomorfa, i en cas que ho siguin calculeu la part imaginària.

(a) $e^x \cos y$

(b) $x^3 + 6xy^2$

(c) $\log(x^2 + y^2)$

Solució: (a) $e^x \sin y$; $f(z) = e^z$; (b) No ho és; (c) $2 \arctan(y/x)$; $f(z) = \log(z^2)$.

B.3. Sigui f una funció holomorfa en un obert $\Omega \subset \mathbb{C}$ i $z_0 \in \Omega$ tal que $f'(z_0) \neq 0$. Quin angle formen les corbes $\operatorname{Re} f(z) = \operatorname{Re} f(z_0)$ i $\operatorname{Im} f(z) = \operatorname{Im} f(z_0)$ en un punt z_0 ?

Solució: $\pi/2$.

1. Trobeu els punts on la funció f és derivable (en el sentit complex), en els següents casos:

(a) $f(z) = |z|$

(d) $f(z) = z + z\bar{z}$

(b) $\cosh x \cos y + i \sinh x \sin y$

(c) $f(z) = \operatorname{Re} z$

(e) $f(z) = \operatorname{Im} e^{\bar{z}} + i \operatorname{Re} e^z$

2. Sigui $\Omega \subset \mathbb{C}$ un obert, $z_0 \in \Omega$ i $f : \Omega \rightarrow \mathbb{C}$ una funció.

a) Identificant \mathbb{R}^2 amb \mathbb{C} de la forma habitual, demostreu que si f és diferenciable en z_0 , llavors

$$Df(z_0)(z) = \frac{\partial f}{\partial z}(z_0) \cdot z + \frac{\partial f}{\partial \bar{z}}(z_0) \cdot \bar{z} \quad (z \in \mathbb{C}),$$

on

$$\frac{\partial f}{\partial z} = \frac{1}{2} \left(\frac{\partial f}{\partial x} - i \frac{\partial f}{\partial y} \right), \quad \frac{\partial f}{\partial \bar{z}} = \frac{1}{2} \left(\frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right).$$

b) Proveu que f és holomorfa en Ω si, i només si, f és diferenciable i $\frac{\partial f}{\partial \bar{z}} = 0$ en Ω . En tal cas, $f' = \frac{\partial f}{\partial z}$.

3. Demostreu que si f és diferenciable en un obert de \mathbb{C} , llavors

$$\overline{\frac{\partial f}{\partial z}} = \frac{\partial \bar{f}}{\partial \bar{z}} \quad \text{i} \quad \overline{\frac{\partial f}{\partial \bar{z}}} = \frac{\partial \bar{f}}{\partial z}.$$

1. The field of complex numbers
--

introduction

entity

Let:

· *statements*

·

Then, *item* is a/an entity if:

· *conditions*

·

We denote:

· *property : notation*

·

2. Holomorphic functions

3. Cauchy-Riemann

Let:

$$\cdot f \in \mathcal{H}(\mathbb{C}) \quad \text{,,} \quad \operatorname{Re} f + i \operatorname{Im} f = c_a$$

Show that:

$$\cdot \exists a' \in \mathbb{C} :$$

$$f = c_{a'}$$

Demonstration:

u, v real components of f

$$u(x, y) + v(x, y) = a$$

differentiate respect x and y

$$u_x + v_x = 0$$

$$u_y + v_y = 0$$

f holomorphic $\rightarrow f$ CR

$$u_x - u_y = 0$$

$$u_y + u_x = 0$$

$$u_x, u_y, v_x, v_y = 0$$

$$\exists a_1 \in \mathbb{R} :$$

$$u = c_{a_1}$$

$$\exists a_2 \in \mathbb{R} :$$

$$v = c_{a_2}$$

$$f = c_{(a_1, a_2)}$$

B.2 a)

Let:

$$\begin{aligned} u : \mathbb{R}^2 &\longrightarrow \mathbb{R}^2 \\ (x, y) &\longmapsto \exp(y) \cos(x) \end{aligned}$$

Show that:

$$\exists f \in \mathcal{H}(\mathbb{C}) :$$

u real component of f

Demonstration:

$$\text{lab } 1 \rightarrow u_{xx} + v_{yy} = 0$$

$$u_x = \exp(x) \cos(y)$$

$$u_{xx} = \exp(x) \cos(y)$$

$$u_y = -\exp(x) \sin(y)$$

$$u_{yy} = -\exp(x) \cos(y)$$

ok

Calculate v using CR

$$v_y = u_x = \exp(x) \cos(y)$$

$$v(x, y) = \int_{\mathbb{C}} \exp(x) \sin(y) dy = \exp(x) \sin(y) + \phi(x)$$

$$v_x = \partial_v x = \exp(x) \sin(y) + \phi'(x)$$

$$-u_y = \exp(x) \sin(y) + \phi'(x)$$

$$\text{CR} \rightarrow \phi'(x) = 0$$

$$\forall c \in \mathbb{R} :$$

$$\phi(x) = c \text{ ok}$$

$$v(x, y) = \exp(x) \sin(y)$$

Preservation of angles

Let:

$$\cdot \gamma_1, \gamma_2 \text{ plane arcs} \quad \parallel \quad \gamma_1(0) = \gamma_2(0)$$

Then, holds:

$$\cdot \text{angle of } \gamma_1'(0) \text{ and } \gamma_2'(0) = \text{angle } \sigma_1'(0), \sigma_2'(0)$$

Demonstration:

rotations and homotecies let angles invariant

Block V

Tasks

1. 1st laboratory
Existence of holomorphic functions

Let:

$$\cdot f \in \mathcal{H}(\mathbb{D})$$

Study:

$$\cdot \exists f \in \mathcal{H}(\mathbb{D}) :$$

$$\forall n \in \mathbb{N} \quad n \geq 2 :$$

$$a) f\left(\pm \frac{1}{n}\right) = \frac{1}{2n+1}$$

$$b) f\left(\pm \frac{1}{n}\right) = \frac{1}{n^2}$$

$$c) \left|f\left(\frac{1}{n}\right)\right| = \frac{1}{\log(n+1)}$$

$$d) \left|f\left(\frac{1}{n}\right)\right| = \frac{n}{n+1}$$

Demonstration:

a):

$$E_1 := \left\{ +\frac{1}{n} + 0i \in \mathbb{C} \mid n \in \mathbb{N} \right\}$$

$$E_2 := \left\{ -\frac{1}{n} + 0i \in \mathbb{C} \mid n \in \mathbb{N} \right\}$$

$$\lim_{E_1} \frac{f(z) - f(0)}{z - 0} = \lim_n \frac{f\left(\frac{1}{n}\right) - f(0)}{\frac{1}{n}} = \frac{1}{2} - \lim_n \frac{f(0)}{\frac{1}{n}}$$

$$\lim_{E_1} \frac{f(z) - f(0)}{z - 0} \begin{cases} = \frac{1}{2} & f(0) = 0 \\ \notin \mathbb{C} & f(0) \neq 0 \end{cases}$$

Case $f(0) = 0$:

$$\lim_{E_2} \frac{f(z) - f(0)}{z - 0} = \lim_n \frac{f\left(-\frac{1}{n}\right) - f(0)}{-\frac{1}{n}} = -\frac{1}{2} \neq \lim_{E_1} \frac{f(z) - f(0)}{z - 0}$$

$$\nexists f \in \mathcal{H}(0) \quad \text{„} f \text{ satisfies } a)$$

In particular:

$$\nexists f \in \mathcal{H}(\mathbb{D}) \quad \text{„} f \text{ satisfies } a)$$

b):

$$\begin{aligned} f : \mathbb{C} &\longrightarrow \mathbb{C} \\ z &\longmapsto z^2 \end{aligned}$$

$$\forall n \in \mathbb{N} \quad \text{„} n \geq 2 :$$

$$f\left(\pm \frac{1}{n}\right) = \frac{1}{n^2}$$

f satisfies b)

$$\begin{aligned} \bar{f} : \mathbb{R}^2 &\longrightarrow \mathbb{R}^2 \\ (x, y) &\longmapsto (u(x, y), v(x, y)) = (x^2 - y^2, 2xy) \\ \bar{f} \in \text{Pol}(\mathbb{R}^2) &\rightarrow \bar{f} \text{ differentiable in } \mathbb{R}^2 \end{aligned}$$

$$\forall (x, y) \in \mathbb{R}^2 :$$

$$\partial_x u(x, y) = 2x = \partial_y v(x, y)$$

$$\partial_y u(x, y) = -2y = -\partial_x v(x, y)$$

f satisfies CR

$$\therefore f \in \mathcal{H}(\mathbb{R}^2)$$

In particular:

$$f \in \mathcal{H}(\mathbb{D})$$

c):

Suppose $\exists f \in \mathcal{H}(\mathbb{D}) \quad \text{„} f \text{ satisfies } c)$

$$f \in \mathcal{C}^0(\mathbb{D}) \rightarrow f(0) = f\left(\lim_n \frac{1}{n}\right) = \lim_n f\left(\frac{1}{n}\right) = 0$$

$$\left| \lim_{E_1} \frac{f(z) - f(0)}{z - 0} \right| = \lim_n \frac{\left| f\left(\frac{1}{n}\right) \right|}{\frac{1}{n}} \notin \mathbb{C}$$

$f \notin \mathcal{H}(0)$ absurd

d):

$$\begin{array}{ccc} f : \mathbb{C} & \longrightarrow & \mathbb{C} \\ z & \longmapsto & \frac{1}{z+1} \end{array}$$

$$\forall n \in \mathbb{N} \quad n \geq 2 :$$

$$\left| f\left(\frac{1}{n}\right) \right| = \frac{1}{\frac{1}{n}+1} = \frac{n}{n+1}$$

f satisfies d)

$$\begin{array}{ccc} \bar{f} : \mathbb{R}^2 & \longrightarrow & \mathbb{R}^2 \\ (x, y) & \longmapsto & (u(x, y), v(x, y)) = \left(\frac{x+1}{(x+1)^2+y^2}, \frac{-y}{(x+1)^2+y^2} \right) \end{array}$$

$$\bar{f} \in \text{Rat}(\mathbb{R}^2) \wedge \forall (x, y) \in \mathbb{R}^2 :$$

$$(x+1)^2 + y^2 \neq 0$$

\bar{f} differentiable in \mathbb{R}^2

$$\forall (x, y) \in \mathbb{R}^2 :$$

$$\partial_x u(x, y) = \frac{y^2 - (x+1)^2}{((x+1)^2 + y^2)^2} = \partial_y v(x, y)$$

$$\partial_y u(x, y) = \frac{-2y(x+1)}{((x+1)^2 + y^2)^2} = -\partial_x v(x, y)$$

f satisfies CR

$$\therefore f \in \mathcal{H}(\mathbb{R}^2)$$

In particular:

$$f \in \mathcal{H}(\mathbb{D})$$

Constant tests

Let:

$$\cdot \Omega \subset \mathbb{C} \text{ region}$$

$$\cdot f \in \mathcal{H}(\Omega)$$

Then, holds:

$$\cdot f_{Re} = 0 \vee f_{Im} = 0 \rightarrow f \in \text{Cst}(\Omega)$$

$$\cdot |f| \in \text{Cst}(\Omega) \rightarrow f \in \text{Cst}(\Omega)$$

$$\cdot \text{Im} f \text{ circumference} \rightarrow f \in \text{Cst}$$

Demonstration:

$$f_{Re} = 0 \vee f_{Im} = 0:$$

$$u := f_{Re}$$

$$v := f_{Im}$$

$$f \in \mathcal{H}(\Omega) \rightarrow f \text{ satisfies CR in } \Omega$$

$$\partial_x u = \partial_y v = 0$$

$$\partial_y u = -\partial_x v = 0$$

Null diferencial test:

$$\Omega \text{ connex} \rightarrow u, v \in \text{Cst}$$

$$u, v \in \text{Cst} \rightarrow f \in \text{Cst}$$

$$|f| \in \text{Cst}(\Omega):$$

$$\begin{aligned} |f| : \mathbb{R}^2 &\longrightarrow \mathbb{R} \\ (x, y) &\longmapsto \sqrt{u(x, y)^2 + v(x, y)^2} \end{aligned}$$

$$|f| \in \text{Cst} \rightarrow \exists a \in \mathbb{R} :$$

$$\sqrt{u(x, y)^2 + v(x, y)^2} = a$$

$$u(x, y)^2 + v(x, y)^2 = a^2$$

$$2\partial_x u(x, y) + 2\partial_x v(x, y) = 0$$

$$2\partial_y u(x, y) + 2\partial_y v(x, y) = 0$$

$$f \in \mathcal{H}(\Omega) \rightarrow f \text{ satisfies CR in } \Omega$$

$$2\partial_y v(x, y) + 2\partial_x v(x, y) = 0$$

$$-2\partial_x v(x, y) + 2\partial_y v(x, y) = 0$$

$$+ : 4\partial_y v(x, y) = 0 \rightarrow \partial_y v(x, y) = 0$$

$$- : 4\partial_x v(x, y) = 0 \rightarrow \partial_x v(x, y) = 0$$

Null differential test:

$$\Omega \text{ connex} \rightarrow u, v \in \text{Cst}$$

$$u, v \in \text{Cst} \rightarrow f \in \text{Cst}$$

$\text{Im}(f)$ circumference :

$$\exists (x_0, y_0) \in \mathbb{R}^2, r \in \mathbb{R}^+ :$$

$$\text{Im}(f) = C_r(x_0, y_0)$$

$$\begin{aligned} \bar{f} : \mathbb{R}^2 &\longrightarrow C_r(x_0, y_0) \\ (x, y) &\longmapsto (r \cos(x - x_0), r \sin(y - y_0)) \end{aligned}$$

$$\forall (x, y) \in \Omega :$$

$$|\bar{f}|(x, y) = \sqrt{r^2(\cos^2(x - x_0) + \sin^2(y - y_0))} = r$$

$$|f| \in \text{Cst} \rightarrow f \in \text{Cst}$$

Real part of holomorphic functions

Let:

$$\cdot \Omega \subset \mathbb{R}^2 \text{ region}$$

$$\cdot u \in \mathcal{C}^2(\Omega) \quad \text{,,} \quad \exists f \in \mathcal{H}(\Omega) :$$

$$f_{Re} = u$$

Show that:

$$\cdot \partial_{xx}u + \partial_{yy}u = 0$$

Study:

$$\cdot \exists f \in \mathcal{H}(\Omega) :$$

$$a) f_{Re}(x, y) = x^2 + y^2$$

$$b) f_{Re}(x, y) = x(x+1) - y^2$$

$$c) \forall \alpha \in \mathbb{R} :$$

$$f_{Re} = y^3 + \alpha x^2 y \wedge \Omega = \mathbb{C}$$

Demonstration:

$$\partial_{xx}u + \partial_{yy}u = 0:$$

$$u := f_{Re}, v := f_{Im}$$

$$f \in \mathcal{H}(\Omega) \rightarrow f \text{ satisfies CR in } \Omega$$

$$\partial_x u = \partial_y v \rightarrow \partial_{xx}u = \partial_{xy}v$$

$$\partial_y u = -\partial_x v \rightarrow \partial_{yy}u = -\partial_{xy}v$$

$$\therefore \partial_{xx}u + \partial_{yy}u = 0$$

$$f_{Re}(x, y) = x^2 + y^2:$$

$$\partial_{xx}u + \partial_{yy}u = 4 \neq 0$$

$$\nexists f \in \mathcal{H}(\Omega) \quad \text{,} \quad f_{Re}(x, y) = x^2 + y^2$$

$$f_{Re}(x, y) = x(x+1) - y^2:$$

$$\begin{aligned} \bar{f} : \Omega &\longrightarrow \mathbb{R}^2 \\ (x, y) &\longmapsto (u(x, y), v(x, y)) = (x(x+1) - y^2, 2xy + y) \end{aligned}$$

$$\bar{f} \in \text{Pol} \rightarrow \bar{f} \text{ differentiable in } \Omega$$

$$\forall (x, y) \in \Omega :$$

$$\partial_x u(x, y) = 2x + 1 = \partial_y v(x, y)$$

$$\partial_y u(x, y) = -2y = -\partial_x v(x, y)$$

$$f \text{ satisfies CR in } \Omega$$

$$f \in \mathcal{H}(\Omega) \wedge f_{Re} = u$$

$$f_{Re}(x, y) = y^3 + \alpha x^2 y:$$

$$f \text{ has to satisfy CR in } \mathbb{C}:$$

$$\forall (x, y) \in \mathbb{R}^2 :$$

$$\partial_y v(x, y) = \partial_x u(x, y) = 2\alpha xy$$

$$\partial_x v(x, y) = -\partial_y u(x, y) = -3y^2 - \alpha x^2$$

$$v(x, y) = \alpha xy^2 + c(x)$$

$$v(x, y) = -3xy^2 - \frac{\alpha}{3}x^3 + c(y)$$

$$\alpha = -3, \quad c(x) = x^3, \quad c(y) = 0$$

$$v(x, y) = -3xy^2 + x^3$$

2. 2nd laboratory

Power series

Study:

$$\cdot \sum_{n \geq 1} n(n+1)z^n$$

Demonstration:

Naming:

R radius of convergence of the series

$\forall n \in \mathbb{N} :$

$$c_n := n(n+1)$$

Convergence domain:

$$\lim_n \frac{c_n}{c_{n+1}} = \lim_n \frac{n(n+1)}{(n+1)(n+1)} = 1$$

Quotient test:

$$R^{-1} = \overline{\lim}_n |c_n|^{\frac{1}{n}} = 1 \rightarrow R = 1$$

Cauchy-Hadamard theorem:

$\sum_{n \geq 1} n(n+1)z^n$ convergent over \mathbb{D}

$\sum_{n \geq 1} n(n+1)z^n$ divergent over $\mathbb{C} \setminus \overline{\mathbb{D}}$

$\forall K \subset \mathbb{D} \quad \parallel K \text{ compact} :$

$\sum_{n \geq 1} n(n+1)z^n$ uniformly convergent over K

Define

$$\begin{aligned} f : \mathbb{D} &\longrightarrow \mathbb{C} \\ z &\longmapsto z \sum_{n \geq 1} n(n+1)z^{n-1} \end{aligned}$$

Sum:

$$\begin{aligned} g : \mathbb{D} &\longrightarrow \mathbb{C} \\ z &\longmapsto \sum_{n \geq 1} n(n+1)z^{n-1} \end{aligned}$$

UCI theorem:

$$\int_0^z g(t) dt = \sum_{n \geq 1} (n+1)z^n$$

$$\begin{aligned} h : \mathbb{D} &\longrightarrow \mathbb{C} \\ z &\longmapsto \sum_{n \geq 1} (n+1)z^n \end{aligned}$$

$$\int_0^z h(t) dt = \sum_{n \geq 1} z^{n+1} = \sum_{n \geq 0} z^n = \frac{1}{1-z}$$

$$h(z) = \partial_z \frac{1}{1-z} = \frac{1}{(1-z)^2}$$

$$g(z) = \partial_z h(z) = \frac{2}{(1-z)^3}$$

$$f(z) = \frac{2z}{(1-z)^3}$$

Application:

In particular:

$$\sum_{n \geq 1} (-1)^n \frac{n(n+1)}{2^n} = f\left(-\frac{1}{2}\right) = \frac{-2^3}{3^3}$$

Bibliography

- [1] Joaquim Ortega-Cerda. Funcions analtiques. "https://campusvirtual2.ub.edu/pluginfile.php/501998/mod_resource/content/1/APUNTS/FA.pdf", February 2005.