

# Block I

# Laboratory

**1. Fixed points cardinality**
**II. Martin Azpillaga**

Let:

- $f : [0, 1] \rightarrow [0, 1] \in \mathcal{C}^2([0, 1])$
- $f(1) < 1$
- $f'' > 0 \in [0, 1]$

Show that:

- $\# \{x \in [0, 1] \mid f(x) = x\} = 1$

Demonstration:

- $\# \{x \in [0, 1] \mid f(x) = x\} \geq 1$ :

Case  $f(0) = 0$ :

0 fixed point

Case  $f(0) > 0$ :

$$\begin{array}{ccc} g : [0, 1] & \longrightarrow & [-1, 1] \\ x & \longmapsto & f(x) - x \end{array} \in \mathcal{C}^2([0, 1])$$

$$g(0) = f(0) - 0 > 0$$

$$g(1) = f(1) - 1 < 0$$

Bolzano's theorem:

$$\exists x \in (0, 1) :$$

$$g(x) = 0$$

$$f(x) = x$$

- $\# \{x \in [0, 1] \mid f(x) = x\} \leq 1$ :

$$g'' > 0 \text{ over } [0, 1]$$

Rolle's theorem:

$$\# \{x \in (0, 1) \mid g'(x) = 0\} \leq 1$$

$$\# \{x \in (0, 1) \mid g(x) = 0\} \leq 2$$

$$\# \{x \in (0, 1) \mid f(x) = x\} \leq 2$$

$$f'' > 0 \text{ over } [0, 1]$$

Monotonicity test:

$$f' \text{ increasing in } [0, 1]$$

$$\forall a < b \in [0, 1] \quad f(a) = a, f(b) = b :$$

Mean Value Theorem:

$$\exists c \in (a, b) :$$

$$f'(c) = \frac{f(b) - f(a)}{b - a} = 1$$

$$\exists d \in (b, 1) :$$

$$\begin{aligned}
 f'(d) &= \frac{f(1)-f(b)}{1-b} < 1 \\
 f' \text{ increasing} &\rightarrow f'(c) < f'(b) < f'(d) \\
 1 < f'(b) < 1 &\text{ absurd} \\
 \therefore \# \{x \in [0, 1] \mid f(x) = x\} &= 1
 \end{aligned}$$

### III. Bifurcation Theory

#### Bifurcation diagram

Let:

$$\begin{aligned}
 \cdot \quad f : \mathbb{R}^2 &\longrightarrow \mathbb{R} \\
 \quad (a, x) &\longmapsto x^3 - 3x^2 + (5-a)x - 2 + a \\
 \cdot \quad \forall a \in \mathbb{R} : \\
 \quad f_a : \mathbb{R} &\longrightarrow \mathbb{R} \\
 \quad x &\longmapsto x^3 - 3x^2 + (5-a)x - 2 + a
 \end{aligned}$$

Study:

$$\cdot \text{ Bifurcations of } (\mathbb{R}, \mathbb{N}, \{f_a\}_{a \in \mathbb{R}})$$

Start:

Fixed points:

$$f_a(x) = x \Leftrightarrow x^3 - 3x^2 + (4-a)x - 2 + a = 0 \Leftrightarrow (x-1)(x^2 - 2x + 2 - a) = 0$$

$$x^2 - 2x + 2 - a = 0 \Leftrightarrow x = \pm \sqrt{a-1}$$

$$\forall a \in \mathbb{R} \quad \text{,, } a \leq 1 :$$

$$\text{Fix}(f_a) = \{1\}$$

$$\forall a \in \mathbb{R} \quad \text{,, } a > 1 :$$

$$\text{Fix}(f_a) = \{1, \pm \sqrt{a-1}\}$$

Stability:

$$\partial_x f(a, x) = 3x^2 - 6x + 5 - a$$

$$\partial_{x^2} f(a, x) = 6x - 6$$

$$\partial_{x^3} f(a, x) = 6$$

$$|\partial_x f(a, 1)| < 1 \leftrightarrow |2 - a| < 1 \leftrightarrow a \in (1, 3)$$

$$\partial_{x^2} f(1, 1) = 0, \quad \partial_{x^3} f(1, 1) > 0$$

$$\partial_{x^2} f(3, 1) = 0, \quad \partial_{x^3} f(3, 1) > 0$$

$$\forall a \in \mathbb{R} \quad \text{,, } a \leq 1 \vee a \geq 3 :$$

1 repulsive

$$\forall a \in \mathbb{R} \quad \text{,, } a \in (1, 3) :$$

1 attractive

$$\forall a \in \mathbb{R} \quad \text{,, } a > 1 :$$

$$|\partial_x f(a, \pm\sqrt{a-1})| = |2a-1| > 1$$

$\pm\sqrt{a-1}$  repulsive

Pitchfork bifurcation at 1 :

$$\partial_a f(1, 1) = 1 - 1 = 0$$

$$\partial_{x^2} f(1, 1) = 6 - 6 = 0$$

$$\partial_{ax} f(1, 1) = -1 \neq 0$$

$$\partial_{x^3} f(1, 1) = 6 \neq 0$$

Period-doubling bifurcation at 3 :

$$\partial_a f^2(3, 1) = \partial_a f(3, f(3, 1)) \partial_a f(3, 1) = (1 - 1)(1 - 1) = 0$$

$$\partial_{x^2} f^2(3, 1) = \partial_{x^2} f(3, f(3, 1)) \partial_{x^2} f(3, 1) = 0$$

$$\partial_{ax}f^2(3,1) = \partial_{ax}f(3,f(3,1))\partial_{ax}f(3,1) \neq 0$$

$$\partial_{x^3}f^2(3,1) = \partial_{x^3}f(3,f(3,1))\partial_{x^3}f(3,1) \neq 0$$

## Source Code

```

#include "stdio.h"
#include "stdlib.h"
#include "math.h"
#include "string.h"

void plot( char *input_file , char *output_file )
{
    FILE *gnuplot;
    gnuplot = popen("gnuplot", "w");
    if( output_file )
    {
        fprintf(gnuplot, "set term svg\n");
        fprintf(gnuplot, "set out \"%s\" \n", output_file );
    }
    fprintf(gnuplot, "plot \"%s\" with dots\n", input_file);
    fflush(gnuplot);
    fclose(gnuplot);
}

double example_function( double param, double point )
{
    return pow(point,3) - 3*pow(point,2) + (5-param)*point - 2 + param;
}

void bifurcation_diagram( int param_min, int param_max, double param_step,
    int point_min, int point_max, int num_points,
    double (*f)(double,double), int num_iter, int tolerancy)
{
    FILE* file;
    double param, point;
    int i,j;

    srand(time(NULL));
    file = fopen("data.dat", "w");

    for ( param = param_min; param < param_max; param += param_step )
    {
        for ( i = 0; i < num_points; i++ )
        {
            point = point_min + ((double) rand() / (double) RAND_MAX) * (point_max -
                point_min);

            for ( j = 0; j < num_iter && abs(point) < tolerancy; j++ )
            {
                point = (*f)(param, point);
            }

            if(abs(point) < tolerancy)
            {
                fprintf(file, "%lf %lf\n", param, point);
            }
        }
    }
    plot( "data.dat", "graph.svg" );
}

```

```

int main(int argc, char const *argv[])
{
    bifurcation_diagram( 0, 5, 10e-3, 0, 5, 100, &example_function, 100, 10e1);

    return 0;
}

```



