

# Machine Learning: Homework #2

Due on November 06, 2017 at 10:00am

*Professor Dr. Stephan Guennemann*

Marc Meissner

## Problem 1

A secret government agency has developed a scanner which determines whether a person is a terrorist. The scanner is fairly reliable; 95% of all scanned terrorists are identified as terrorists, and 95% of all upstanding citizens are identified as such. An informant tells the agency that exactly one passenger of 100 aboard an airplane in which you are seated is a terrorist. The agency decides to scan each passenger and the shifty looking man sitting next to you is tested as 'TERRORIST'. What are the chances that this man is a terrorist? Show your work!

### Solution

Let  $S$  symbolize the correct result of the scan and  $T$  the fact that the subject is a terrorist. According to the description, the following holds:

$$P(S|T) = P(\bar{S}|\bar{T}) = 0.95, P(S|\bar{T}) = 0.05, P(T) = 0.01 \text{ and } P(\bar{T}) = 0.99$$

We can then acquire the solution by Bayes Law:

$$P(T|S) = \frac{P(S|T) \cdot P(T)}{P(S|T) \cdot P(T) + P(S|\bar{T}) \cdot P(\bar{T})} = \frac{19}{118} \approx 0.161$$

The probability is only 16.1%. But since the man is looking shifty, he probably is the terrorist, anyway.

## Problem 2

A fair coin is tossed twice. Whenever it turns up heads, a red ball is placed into a box, otherwise a white ball. Afterwards, balls are drawn from the box three times in succession (placing the drawn ball back into the box every time). It is found that on all three occasions a red ball is drawn. What is the probability that both balls in the box are red? Show your work!

### Solution

Let  $RR$ ,  $WW$  and  $RW$  be the three possible combinations of colored balls in the box. Let  $D3R$  denote the event that three red balls have been drawn. Then:

$$P(RR) = P(WW) = \frac{1}{4}, P(RW) = \frac{1}{2}$$

$$P(D3R|RR) = 1, P(D3R|WW) = 0, P(D3R|RW) = \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{8}$$

Bayes Law leads to the solution:

$$P(RR|D3R) = \frac{P(D3R|RR) \cdot P(RR)}{P(D3R|RR) \cdot P(RR) + P(D3R|RW) \cdot P(RW)} = \frac{4}{5} = 0.8$$

## Problem 3

A fair coin is flipped until heads shows up for the first time. What is the expected number of tails  $T$  and the expected number of heads  $H$  in any one run of this experiment? Show your work.

**Solution****Problem 4**

Calculate mean and variance of a uniform random variable  $X$  on the interval  $[a, b]$ ,  $a < b$  with probability density function

$$p(x) = \begin{cases} \frac{1}{b-a} & a \leq x \leq b. \\ 0 & \text{elsewhere.} \end{cases} \quad (1)$$

**Solution**

Just apply the formulas/definitions of mean and variance:

$$E[X] = \int_{-\infty}^{\infty} x \cdot p(x) dx = \frac{1}{b-a} \cdot \int_a^b x dx = \frac{1}{2} \cdot \frac{b^2 - a^2}{b-a} = \frac{a+b}{2}$$

$$E[X^2] = \frac{1}{b-a} \cdot \int_a^b x^2 dx = \frac{1}{3 \cdot (b-a)} \cdot [b^3 - a^3] = \frac{b^2 + ab + a^2}{3}$$

$$\text{Var}[X] = E[X^2] - (E[X])^2 = \frac{b^2 + ab + a^2}{3} - \frac{b^2 + ab + a^2}{4} = \frac{(b-a)^2}{12}$$

**Problem 5**

Let  $X$  and  $Y$  be random variables with joint density  $p(x, y)$ . Prove the tower properties:

$$E[X] = E_Y[E_{X|Y}[X]],$$

$$\text{Var}[X] = E_Y[\text{Var}_{X|Y}[X]] + \text{Var}_Y[E_{X|Y}[X]].$$

$E_{X|Y}[X]$  and  $\text{Var}_{X|Y}[X]$  denote the expectation and variance of  $X$  under the conditional density  $p(x|y)$ .

**Solution****Problem 6**

Prove eq. (1). You may assume that the  $X_i$  have finite variance  $\text{Var}[X_i]$ . You may further use Markov's and Chebyshev's inequalities without proof.

**Solution**

$$E[X] = \frac{1}{n} \cdot \sum_{i=1}^n X_i = \mu$$

$$\text{Var}[X] = \text{Var}\left[\frac{1}{n} \cdot \sum_{i=1}^n X_i\right] = \sum_{i=1}^n \text{Var}\left[\frac{X_i}{n}\right] = n \cdot \frac{\sigma^2}{n^2} = \frac{\sigma^2}{n}$$

$$P(|X - E[X]| > \epsilon) = P\left(\left|\frac{1}{n} \cdot \sum_{i=1}^n X_i - E[X]\right| > \epsilon\right) \leq \frac{\sigma^2}{\epsilon^2 \cdot n}$$