RIASSUNTO TESI

Titolo:	The Mori-Zwanzig formalism as a tool to systematically derive hybrid coarse-grained models
Autore:	Marco Spinaci
Relatore interno:	Lorenzo Rovigatti
Data della sessione di laurea:	16/12/2019

My Master thesis project deals with the Mori-Zwanzig (MZ) formalism as a tool to systematically derive hybrid coarse-grained (CG) models. We consider two resolution levels for a given molecular system. The first one, at high resolution, is the fine-grained (FG), while the second one, at low resolution, is the CG system. This project is based on the work of N. Di Pasquale *et al.* [1], in which the authors derived a coarse-grained model for hybrid systems, through the MZ projection, with a theoretical discussion supported by numerical simulations in the NVE ensemble. The MZ formalism allows to study the evolution of a dynamical system through the projection on a subset of its degrees of freedom. This operation can be done through various projection operator techniques [2].

The first part of the thesis is focused on the study and detailed derivation of the equations governing the coarse-grained dynamics, obtained through MZ formalism. The equations of motion we get are described by a Generalized Langevin Equation (GLE), see [1]. Which is an equation that allows to describe a particle, in a system close to equilibrium, subject to a potential and coupled to a general thermal bath in which memory effects are present.

The second part of the thesis is focused on the study of numerical methods for Langevin dynamics and for the constrained Langevin dynamics. These methods are used to numerically implement full and the constrained dynamics in the canonical ensemble. The integrators applied were developed by Leimkuhler *et al.* [3, 4], those methods are splitting integrators that decompose the stochastic vector field which drives the Langevin Dynamics into simpler vector fields which can be solved exactly. To sample the full dynamics we used the BAOAB method [3]. Instead, to sample the constrained dynamics we used a geodesic-BAOAB method [4].

The third and final part of the thesis is on the numerical experiments in the N,V,T ensemble. We modified the open-source parallel code, written in Julia 0.6, developed by Di Pasquale *et al.* [1], introducing the integrators studied in the previous part. This code allows to solve the full, constrained and coarse-grained dynamics, as well as all the components needed for the CG system. The FG system that we consider is a periodic chain of M^{FG} "atoms" of two different species, which are assumed to have different masses and stiffness, in a repeating pattern. We assumed that the chemical structure of the molecular system is fixed, i.e. no reactions occur. Each "atom" is mapped to a fixed (unique) particle in the CG system, called bead, so we considered a fully coarse-grained system. We performed the simulation for four different systems with different parameter and for different dynamics: the FG dynamics, the orthogonal dynamics (or constrained dynamics), the deterministic coarse-grained dynamics and the Markovian Mori-Zwanzig dynamics.

In all cases simulated no significant scale separation between the dynamics of the beads and individual atoms was observed. Nevertheless, a comparison was made between the true and approximate dynamics, and as expected, systems with 'more rigid' beads, having greater internal stiffness, showed much weaker memory effects. Furthermore we observed that the Langevin dynamics does not appear to be appropriate for generic CG systems, and that the preservation of more complex dynamical properties, such as the mean-square displacement of the inter bead distance in time, requires instead the strict validity of this assumption or a correct parametrisation and implementation of the memory term.

References

[1] N. Di Pasquale, T. Hudson, and M. Icardi. Systematic derivation of hybrid coarse-grained models. *Physical Review E*, 99:013303, Jan 2019.

[2] H. Grabert. *Projection operator techniques in nonequilibrium statistical mechanics*. Springer tracts in modern physics. Springer-Verlag, 1982.

[3] B. Leimkuhler and C. Matthews. Rational construction of stochastic numerical methods for molecular sampling. *Applied Mathematics Research eXpress*, 2013(1):34–56, 2013.

[4] B. Leimkuhler and C. Matthews. Efficient molecular dynamics using geodesic integration and solvent-solute splitting. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 472(2189), 2016.