

# The doctrinal Herbrand’s theorem and its Stone dual

Marco Abbadini

Catholic University of Louvain, Louvain-la-Neuve, Belgium.

`marco.abbadini@uclouvain.be`

Herbrand’s theorem (in its weak form) says that if  $\exists x \phi(x)$  is provable (with  $\phi(x)$  quantifier-free), then some finite disjunction  $\phi(t_1) \vee \dots \vee \phi(t_n)$  of its ground instances is already provable.

In [AG], we obtained Herbrand’s theorem within the formalism of first-order Boolean doctrines. We recall that *first-order Boolean doctrines* are the analogue, for classical first-order logic, of Boolean algebras for classical propositional logic: given a first-order theory  $\mathcal{T}$ , one considers formulas (indexed by their contexts of free variables) modulo  $\mathcal{T}$ -provable equivalence, and obtains a first-order Boolean doctrine. Formally, a *first-order Boolean doctrine* over a category  $\mathbf{C}$  with finite products is a functor  $\mathbf{P}: \mathbf{C}^{\text{op}} \rightarrow \mathbf{BA}$  in which the reindexings along the product projections have right and left adjoints satisfying the Beck-Chevalley condition.

Recall now that a (not necessarily first-order) *Boolean doctrine* over a category  $\mathbf{C}$  with finite products is simply a functor  $\mathbf{P}: \mathbf{C}^{\text{op}} \rightarrow \mathbf{BA}$ ; a Boolean doctrine models the class of quantifier-free formulas modulo a universal first-order theory, indexed by their contexts. Any Boolean doctrine over a small category admits a quantifier completion  $\mathbf{P} \hookrightarrow \mathbf{P}^{\forall\exists}$ , i.e., a universal morphism into a *first-order* Boolean doctrine, which freely adds first-order quantifiers to  $\mathbf{P}$ . Herbrand’s theorem gives an explicit description of the fragment of  $\mathbf{P}^{\forall\exists}$  consisting of formulas of quantifier-alternation depth at most 1.

In this talk, I will present joint work in progress with Francesca Guffanti, in which we develop the Stone dual of this description. Composing a first-order Boolean doctrine  $\mathbf{P}: \mathbf{C}^{\text{op}} \rightarrow \mathbf{BA}$  with Stone duality  $\mathbf{BA}^{\text{op}} \cong \mathbf{Stone}$  yields Joyal’s notion of polyadic space, namely a functor  $\mathbf{E}: \mathbf{C} \rightarrow \mathbf{Stone}$  in which product projections are sent to open maps, and certain squares are epi-pullbacks. Just as a Stone space captures the space of models of a classical propositional theory, a polyadic space captures the space of models of a classical first-order theory, up to elementary equivalence.

For the Stone dual of Herbrand’s theorem, we start from a Stone-valued functor  $\mathbf{E}: \mathbf{C}^{\text{op}} \rightarrow \mathbf{Stone}$ , representing the space of models of a universal theory modulo agreement on quantifier-free formulas. We then give an explicit description—entirely in terms of  $\mathbf{E}$ —of the Stone-valued functor capturing the space of models of the same theory modulo agreement on formulas of quantifier-alternation depth at most 1. In other words, we identify the first nontrivial layer of the quantifier completion on the Stone/polyadic side, thus obtaining the Stone dual of the doctrinal Herbrand theorem.

## References

- [AG] M. Abbadini and F. Guffanti. Freely adding one layer of quantifiers to a Boolean doctrine. Preprint at <https://arxiv.org/abs/2410.16328>.