



LIP Internship: *Tensor Networks for in-medium particle dynamics*

Compendium of problems

Author: Marco Leitão^a

Contents

I	Numerical Methods Review	1
II	Tensor Networks	5

^aEmail: mleitao@lip.pt

I. Numerical Methods Review

The following chapter is a review of numerical methods. It is recommended to explore it in order to refresh your knowledge on the subject, but it is not strictly required. Problems 1-3 should be doable with basic computational physics knowledge (possibly except 3.ix). Problem 4 is more advanced and requires some research on your part, but it is a very interesting problem that can give you a taste of what is done in some areas of modern physics research.

Problem 1. Numerical errors in derivatives

The total error in computing a numerical derivative is $E = E_{\text{acc}} + E_{\epsilon}$, where E_{acc} is the error associated to the analytical Taylor expansion on the resolution step (truncation error) and E_{ϵ} the error associated to the numerical precision.

In this exercise, your task is to perform the numerical analysis of the errors for the *forward difference derivative*, which has an accuracy error of $\mathcal{O}(h)$, and the *central difference second derivative*, which is of $\mathcal{O}(h^2)$. You should be able to derive all the formulas, including the Taylor expansions, by yourself (you can check literature for help). Plot the error total as a function of h (in full log-axis), for a function of your choice, and compare the numerical error of both methods. What can you conclude?

Problem 2. Lotka-Volterra prey-predator problem

A classical model for the population of prey and predators in a given ecosystem are the Lotka-Volterra equations. This is a coupled set of two differential equations, given by

$$\begin{aligned}\frac{dx}{dt} &= \alpha x - \beta xy \\ \frac{dy}{dt} &= \theta xy - \gamma y,\end{aligned}\tag{I.1}$$

where x and y correspond to the number of preys and the number of predators, respectively. The constant α corresponds to the growth rate of the preys, β is the predation rate, θ the growth rate of predators per prey eaten and γ the predator death rate.

You are free to explore this problem however you want, by solving the ODEs for different combinations of variables and initial conditions, using your preferred method. Here are some suggested studies:

- i) Find the equilibrium points of the system and study their stability;
- ii) Plot the phase space of the system, i.e., plot x vs. y , for different configurations;
- iii) Do some numerical analysis;
- iv) Explore variations of the problem, e.g., add a new predator higher in the *food chain*, add a quadratic term to the equations, etc.;

- v) Do animations and/or get nice and informative plots (this is quite important). Tip: use *matplotlib* for your plots (you can write a script that imports your results from, for instance, your C++ routine, and builds plots with that data).

Problem 3. Lennard-Jones potential

The interaction of two atoms can be modeled using the Lennard-Jones Potential, expressed by

$$V(r) = 4\epsilon \left[\left(\frac{\sigma}{r} \right)^{12} - \left(\frac{\sigma}{r} \right)^6 \right], \quad (\text{I.2})$$

where r is the distance between the two atoms, ϵ is the depth of the potential and σ is the distance at which the potential is zero.

In the file `lennard_jones.csv`, you can find experimental data with uncertainties for the measurement of the potential of between a pair of atoms. Solve the following questions:

- i) Read the data, *fit* the data to the Lennard-Jones potential in Eq.(I.2), thus extracting the parameters ϵ and σ . You should explore techniques and/or libraries to fit the data with uncertainties;
- ii) Plot the data and the fitted curve, both with uncertainties;
- iii) Perform an analytical harmonic oscillator (HO) approximation around the minimum of the potential ($V(r) \sim r^2$) (*small oscillations*).
- iv) Consider that the motion of the atoms are approximately classical and one dimensional, ie, described by the Lagrangian

$$L = \frac{1}{2}\mu\dot{r}^2 - V(r), \quad (\text{I.3})$$

where μ is the reduced mass of the atoms (use $\mu = 1$). Use the Euler-Lagrange equations to derive the equation of motion for the distance between the atoms. Then, solve analytically the equations of motion for small oscillations (HO approximation).

- v) Implement a numerical routine to solve the equation of motion with a general potential $V(r)$. You can use RK methods, other method such as leap-frog. Then, use your numerical method to solve the equation of motion for the HO approximated potential, for adequate initial conditions. Solving numerically a limiting case of a problem that has analytical solution is a good way to benchmark your method. Finally, after making sure that the method matches the analytical predictions, you should solve the full case for $V(r)$. Do the simulation for various initial conditions and compare them with the HO case. Study the regimes of compatibility and discrepancies between the full solution and the HO.
- vi) Verify that your numerical method conserves the total energy. You may find minor discrepancies due to numerical errors. Thus, try to do some numerical analysis to estimate the error of your method and verify if it is compatible with the discrepancies you find in the energy conservation.

- vii) At the beginning you were given experimental data with uncertainties, and then fit into the Lennard-Jones potential. Now, you will plan and do the extraction of the potential yourself based on numerical realizations. Starting from the equations of motion for parameters of your choice, simulate several motions of the atoms, for different initial conditions, and store all the positions registered. This data is what you are going to work with. Then, from that data, you should already be able to extract the potential, based on the connection between the potential and the equations of motion. *Hint*: find the accelerations from the recorded positions.
- viii) Explore variations of the problem. For instance, do animations, let the atoms rotate in polar coordinates (this changes the Lagrangian), add an external magnetic field, etc.
- ix) Finally, let's do some Quantum Mechanics! Consider the Lennard-Jones potential. The Hamiltonian operator for the system is given by

$$H = -\frac{\hbar^2}{2\mu} \frac{\partial^2}{\partial r^2} + V(r), \quad (\text{I.4})$$

where $\hbar$ is the reduced Planck constant (take $\hbar = 1$ for the sake of numerics). The time-dependent Schrödinger equation for the system is hence given by

$$i\hbar \frac{\partial \psi(r, t)}{\partial t} = \left(-\frac{\hbar^2}{2\mu} \frac{\partial^2}{\partial r^2} + V(r) \right) \psi(r, t), \quad (\text{I.5})$$

where $\psi(r, t)$ is the wave function of the system (if you are not familiar with the Schrödinger equation, you can check the literature/Wikipedia).

In quantum mechanics, everything is probabilistic, so one needs to talk about distributions and expectation values of observables (e.g. positions, momenta, etc.), rather than their exact values. The expectation value of the position is given by

$$\langle r(t) \rangle = \int_{-\infty}^{\infty} r |\psi(r, t)|^2 dr, \quad (\text{I.6})$$

where $|\psi(r, t)|^2$ is the probability density of finding the particle at position r at time t . Your task is to solve the Schrödinger equation using the Chebyshev polynomial method, starting with a gaussian pulse around 0 on the average initial position, and compute the expectation value of the position as a function of time, using the same parameters you found in i) and $\mu = 1$. Compare with the classical case and discuss the results.

Problem 4. Recreating phase transitions in the early universe

In the beginning, the universe was in a highly symmetric state, with a balance of matter and anti-matter, and in a unified picture of interactions. However, as the universe evolved, these symmetries started to *break*, evolving into a different phase and eventually leading to the universe we know today. These transitions were driven by what we call *fields*, which are the fundamental entities, such as the Higgs field, the electromagnetic field, etc. Mathematically, they are

described functions of the position and time, $\phi(\mathbf{r}, t)$.

Near a phase transition, these fields are approximately ruled by what we call a ϕ^4 theory. This gives us the possibility of studying the phase transitions in the early universe using other simpler systems that exhibit similar phase transitions near critical points, for instance thin ferromagnetic films. Thus, we can say, at some extent, that ferromagnetic films constitute an *analogue system* of the early universe.

In a ferromagnetic system, the field that describes the system is the magnetization $m(\mathbf{r}) \in [-1, 1]$. For such system around the critical point, the Free Energy takes the form

$$F = \int d^d x \left[\frac{1}{2} (\nabla m)^2 + T m^2 + \lambda(T) m^4 + h m \right], \quad (\text{I.7})$$

where T is the temperature, λ is a constant, h is an external magnetic field and d the dimensions of the system.

In this problem, you will use the *Hybrid Monte Carlo* (HMC) method to simulate this system. You should do a small research on the HMC method procedure (ask for references if needed!) and work in team, in order to establish and plan the methodology to attack this problem. For this method, you will need the pseudo-Hamiltonian of the system in a discretized lattice, which can be written as

$$H = a^d \sum_i \left[\frac{1}{2} \pi_i^2 + \frac{1}{2a^2} \sum_{j, \text{neighbor}} (m_i - m_j)^2 + T m_i^2 + \lambda m_i^4 + h m_i \right], \quad (\text{I.8})$$

where m_i is the field at site i , π_i is the conjugate momentum of the field, and the summation in j is done over **immediate neighbors**. The quantity a corresponds to the lattice spacing, and you can set $a = 1$. The goal of this problem is to study, from simulations, the behaviour of the average value of the magnetization $\langle m \rangle$ in terms of the temperature T and the magnetic field h , in a lattice of **2 dimensions**. Some suggestions:

- i) Use the HMC method to sample evolve the system until the thermal equilibrium (constant $\langle m \rangle$ over time), as described above;
- ii) Start by exploring $h = 0$ and study what happens to $\langle m \rangle$ when varying λ and T . You should be able to find a critical temperature T_c where the system undergoes a phase transition. What is the behaviour of $\langle m \rangle$ around T_c ? What is the relation between this temperature and λ (what is λ in such transitions)?
- iii) Check David Tong's Statistical Field Theory, Chapter 1, for analytical study of the Ising Model, and the CompPhysics 24/25 IST first homework. You can use this as a reference to compare your results and to understand how to build a phase diagram;
- iv) Do a detailed analysis, both from a physical and a numerical point of view.
- v) Establish and divide tasks among the team members.
- vi) BONUS: can you derive the discretized pseudo-Hamiltonian?

With this problem, you are doing your very first state-of-the-art *lattice gauge theory* simulation!

II. Tensor Networks

Problem 1. Memory

Consider the following tensors. Indicate the vector space they belong to and estimate the memory required to completely store them. Evaluate whether it is possible to store them in typical RAM memory, L2 cache, and L1 cache. Whenever possible, implement them in your computer and check explicitly the memory used by them (e.g. in numpy, you can use the `.nbytes` attribute of the array). Use floating point precision of 64 bits (8 bytes) *whenever necessary*.

- i) $T_k = \{k : k \in \mathbb{N}, 1 \leq k \leq 100000\}$
- ii) $T_k = \{dx \times k : k \in \mathbb{N}, 1 \leq k \leq 100000\}$ with $dx = 10^{-7}$.
- iii) $T_{kl} = \frac{kl}{N}$, for $1 \leq k, l \leq N = 1000$.
- iv) $T_{klm} = \sin(klm)$, for $1 \leq k, l, m \leq N = 1000$.
- v) $T_{klm} = e^{iklm}$, for $1 \leq k, l, m \leq N = 1000$.
- vi) $T_{k_1 \dots k_6} = e^{ik_1 \dots k_6}$, for $1 \leq k_1, \dots, k_6 \leq N = 32$.
- vii) $T_{k_1 \dots k_6} = \frac{k_1 k_2 k_3 e^{ik_4 k_5 k_6}}{N^3}$, for $1 \leq k_1, \dots, k_6 \leq N = 32$.
- viii) $T_{k_1 \dots k_8} = e^{i(k_1 + \dots + k_8)}$, for $1 \leq k_1, \dots, k_8 \leq N = 32$.
- ix) $T_{k_1 \dots k_8} = k_1 \dots k_8$, for $1 \leq k_1, \dots, k_8 \leq N = 16$.

Problem 2. Factorization

Which tensors in Problem 1 can be exactly factorized into a product of lower-rank tensors? For those that can, write down the factorization explicitly. How does this affect the memory requirements?





Problem 3. Drawing diagrams

Represent in graph diagrams the following tensor decomposition. Indicate the vector space of each tensor appearing in the diagram and the resulting tensor after contraction. Identify which decompositions correspond to tensor trains (TT).

- i) $T_{kl} = \sum_{a=1}^r A_{ka} B_{al}$, for $1 \leq k, l \leq N$.
- ii) $T_{klm} = \sum_{a=1}^{r_1} \sum_{b=1}^{r_2} A_{ka} B_{alb} C_{bm}$, for $1 \leq k \leq N_1, 1 \leq l \leq N_2, 1 \leq m \leq N_3$.
- iii) $T_{klm} = \sum_{a=1}^{r_1} \sum_{b=1}^{r_2} \sum_{c=1}^{r_3} A_{kab} B_{alc} C_{bcm}$, for $1 \leq k \leq N_1, 1 \leq l \leq N_2, 1 \leq m \leq N_3$.

Problem 4. Reading diagrams

Provide the explicit tensor decomposition corresponding to the following diagrams. Indicate the vector space of each tensor appearing in the diagram and the resulting tensor after contraction. Assume all contractions to be of rank r and all free indices to be of dimension N . Identify which decompositions correspond to tensor trains (TT).



Problem 5. TT/SVD decomposition

Implement a numerical routine that performs a TT/SVD decomposition of an arbitrary tensor. The algorithm must be able to take as input a tensor of arbitrary dimensions, and return the TT decomposition of the tensor. The routine should also allow for a specified accuracy ϵ to control the truncation of singular values during the decomposition.

Problem 6. TT decomposition of tensors

Consider the following tensors. Implement a numerical routine that performs a TT decomposition of them up to a chosen accuracy ϵ . Evaluate the compression ratio (ratio between the memory required by the original tensor and the TT decomposition) as a function of the chosen accuracy ϵ , as well as the ranks in the Tensor Train. You should be able to store all the given tensors in your memory (most of them are of ~ 1 GB for double precision).

i) $T_{klm} = (k + l)/m$, for $1 \leq k, l, m \leq N = 500$.

ii) $T_{klmn} = \sin(klmn)$, for $1 \leq k, l, m, n \leq N = 100$.

iii) $T_{klmn} = e^{i(kl+lm+mn)}$, for $1 \leq k, l, m, n \leq N = 75$.

iv) $T_{i_1 i_2 i_3 i_4 i_5 i_6} = i_1 i_2 i_3 \ln(1 + i_1 i_2 i_3 i_4 i_5 i_6)$, for $1 \leq i_1, i_2, i_3, i_4, i_5, i_6 \leq N = 24$.