How much does $\partial C_\ell/\partial \omega_m$ depend on the other parameters?

At the benchmark-1 point, $\partial C_\ell/\partial \omega_m$ recomputed with each ΛCDM parameter θ_i shifted by ±3%; change measured with Planck-like noise weighting (33 μK-arcmin, 7′ beam, $f_{\rm sky} = 0.57$).

	change in $\partial C_\ell/\partial \omega_m$, θ_i shifted ±3%				same change at a
shifted θ_i	TT	TE	EE	$\phi\phi$	1σ shift (TT/TE/EE)
h	10.5%	11.0%	8.9%	1.7%	$\sim 3\%$
ω_b	4.6%	4.5%	3.4%	1.3%	$\sim 1\%$
A_s	2.8%	2.7%	2.8%	3.0%	$\sim 0.8\%$
n_s	2.0%	1.6%	1.7%	5.3%	$\sim 0.3\%$
τ	0.6%	0.8%	0.9%	0.1%	$\sim 1.6\%$

For a 1σ shift in any other parameter, $\partial C_\ell/\partial \omega_m$ changes by at most $\sim 3\%$ (from h). A ±3% step is a much larger move than 1σ for h, ω_b, A_s, n_s, and a smaller one for τ, hence its larger 1σ value.