

Probability Theory and Statistics

Exercise 4 solutions

09.29. – 10.03.

Expected value, Expected value of transformed random variables, deviation

1. Let X be the maximum of the two dice (a random variable). We can compute the expected value of X as follows:

$$\mathbb{E}(X) = \sum_{k=1}^6 k \cdot \mathbb{P}(X = k).$$

The maximum is $X = 1$ if both dice show 1 – there is only one such outcome, so $\mathbb{P}(X = 1) = 1/36$ (there are 36 possible outcomes in total). $\mathbb{P}(X = 2) = 3/36$, since there are three favorable cases: one die shows 1 and the other shows 2 (in either order), or both dice show 2. Proceeding similarly for each $k \in \{1, 2, \dots, 6\}$, we obtain

$$\mathbb{E}(X) = 1 \cdot \frac{1}{36} + 2 \cdot \frac{3}{36} + 3 \cdot \frac{5}{36} + 4 \cdot \frac{7}{36} + 5 \cdot \frac{9}{36} + 6 \cdot \frac{11}{36} = \frac{1}{36} \sum_{i=1}^6 i(2i-1) = 4.4722.$$

2. Let X be the number of flips needed. Note that $\text{Ran}(X) = \{2, 3, 4, \dots\}$ (since to get two equal consecutive outcomes, at least two flips are needed, but any larger number is also possible). Define $Y := X - 1$. Then Y is geometrically distributed with parameter $1/2$.

Indeed, for $k \geq 1$, if the experiment has not ended after the k -th flip, then it ends at the $(k+1)$ -th flip exactly when the $(k+1)$ -th outcome equals the k -th, which has probability $1/2$, independent of the previous outcomes. Thus for all $k \geq 1$,

$$\mathbb{P}(Y > k+1 | Y > k) = 1/2.$$

Since $\mathbb{P}(Y > 0) = 1$, the multiplication rule gives

$$\mathbb{P}(Y > k+1) = (1/2)^{k+1},$$

so

$$\mathbb{P}(Y = k) = \mathbb{P}(Y > k-1) - \mathbb{P}(Y > k) = (1/2)^{k-1} - (1/2)^k = (1/2)^k = (1/2)^{k-1} \cdot (1-1/2).$$

Thus Y is indeed $\text{Geo}(1/2)$. Therefore, by linearity of expectation,

$$\mathbb{E}(X) = \mathbb{E}(Y+1) = \mathbb{E}(Y)+1 = 2+1 = 3, \quad \mathbb{D}^2(X) = \mathbb{D}^2(Y) = \frac{1-p}{p^2} = 2, \quad \mathbb{D}(X) = \sqrt{2}.$$

Remark: On practice class at least one student suggested the following alternative solution. The sample space may be taken as the set of finite F – H sequences ending with two identical outcomes but containing no two consecutive identical ones before, e.g. $FF, HH, FHH, HFF, FHFF, HFFH, \dots$. For each $k = 2, 3, 4, \dots$ there are exactly 2 such sequences of length k . The value X is precisely the length of the outcome. A sequence of length k has probability $(1/2)^k$, hence $\mathbb{P}(X = k) = 2(1/2)^k = (1/2)^{k-1}$ for $k = 2, 3, 4, \dots$. From this, $\mathbb{E}(X)$ can be computed directly as in the case of the geometric distribution, and similarly for $\mathbb{D}(X)$.

3. As we have already seen earlier, the distribution of the number of defective bulbs is binomial with parameters $n = 100$ and $p = 0.01$. Thus $\mathbb{E}(X) = np = 100 \cdot 0.01 = 1$.

4. Let X denote the number of points. Then X is geometrically distributed, since we repeat until the first success (a point inside the unit circle). So $X \sim \text{Geo}(p)$, where

$$p = \frac{\text{area of circle}}{\text{area of square}} = \frac{\pi \cdot 1^2}{4} = \frac{\pi}{4} \approx 0.7854.$$

Therefore

$$\mathbb{E}(X) = \frac{1}{p} = \frac{4}{\pi} \approx 1.2732.$$

5. Let X be the number of broken telephones on a given day. Since the number of telephones is unbounded and each fails independently with small probability, X has a Poisson distribution: $X \sim \text{Poi}(\lambda)$. The problem states

$$\mathbb{P}(X = 0) = \frac{12}{360} = \frac{1}{30}.$$

But for a Poisson variable $\mathbb{P}(X = 0) = e^{-\lambda}$, hence $\lambda = -\ln(1/30) \approx 3.401$. Thus $\mathbb{E}(X) = \lambda = 3.401$.

Now let Y be the number of days (out of 360) with at least 2 failures. Then

$$\mathbb{P}(X \geq 2) = 1 - \mathbb{P}(X \leq 1) = 1 - (e^{-\lambda} + \lambda e^{-\lambda}) \approx 0.8533.$$

So $Y \sim \text{Bin}(360; 0.8533)$ and $\mathbb{E}(Y) = 360 \cdot 0.8533 \approx 307.18$.

6. (a) Let X be the number of defective bulbs on that day. Then $X \sim \text{Bin}(n = 10, p = 1/20)$. Thus $\mathbb{E}(X) = np = 1/2$, $\mathbb{D}^2(X) = np(1 - p) = 10 \cdot 1/20 \cdot 19/20 = 19/40 = 0.475$.

(b) The required probability is

$$\mathbb{P}(4 < X < 8) = \mathbb{P}(X = 5) + \mathbb{P}(X = 6) + \mathbb{P}(X = 7),$$

which equals

$$\binom{10}{5}(1/20)^5(19/20)^5 + \binom{10}{6}(1/20)^6(19/20)^4 + \binom{10}{7}(1/20)^7(19/20)^3 \approx 0.0000637.$$

(c) For large n , the number of defective bulbs can be approximated by a Poisson random variable Y with parameter $\lambda = \lim_{n \rightarrow \infty} n(1/20) = 1/2$. Hence

$$\mathbb{P}(Y = 3) = \frac{\lambda^3}{3!} e^{-\lambda} = \frac{(1/2)^3}{6} e^{-1/2} \approx 0.0126.$$

7. $\mathbb{E}((X - 3)^2) = \mathbb{E}(X^2 - 6X + 9) = \mathbb{E}(X^2) - 6\mathbb{E}(X) + 9$. We know $\mathbb{E}(X) = 3.5 = 21/6$ and $\mathbb{E}(X^2) = 91/6$, so

$$\mathbb{E}((X - 3)^2) = \frac{91}{6} - \frac{126}{6} + \frac{54}{6} = \frac{19}{6} = 3\frac{1}{6}.$$

8. a) $\mathbb{P}(Y = 1) = 5/9$, $\mathbb{P}(Y = -1) = 4/9$. b) $\mathbb{E}(Y) = 1/9$. c) $\mathbb{E}((-1)^X) = 1/9$.

9. $X \sim \text{Bin}(n = 3, p = 1/4)$. Then

$$\mathbb{E}(Y) = \mathbb{E}(X^2) = \sum_{k=0}^3 k^2 \mathbb{P}(X = k) = 9/8.$$

So $\mathbb{E}(Z) = \mathbb{E}(X^2 + X + 1) = \mathbb{E}(X^2) + \mathbb{E}(X) + 1 = 9/8 + 3/4 + 1 = 23/8$. Also $\mathbb{D}^2(X) = np(1 - p) = 9/16$, hence $\mathbb{D}(X) = 3/4$.

10. The pmf is

$$\mathbb{P}(X = k) = \binom{k+2}{2} p^3 (1-p)^k, \quad k = 0, 1, 2, \dots$$

Then

$$\mathbb{E}\left(\frac{1}{(X+1)(X+2)}\right) = \sum_{k=0}^{\infty} \frac{1}{(k+1)(k+2)} \binom{k+2}{2} p^3 (1-p)^k = \frac{p^2}{2}.$$

11. Sample space size 27. The distribution of X (number of surviving deer):

$$\mathbb{P}(X = 0) = 2/9, \quad \mathbb{P}(X = 1) = 2/3, \quad \mathbb{P}(X = 2) = 1/9.$$

Thus $\mathbb{E}(X) = 8/9$, $\mathbb{D}^2(X) = 26/81$, $\mathbb{D}(X) \approx 0.567$.

12. $\mathbb{D}^2(2X + 1) = 4\mathbb{D}^2(X) = 4\lambda$.

13. (a) $\mathbb{E}((2 + X)^2) = 14$. (b) $\mathbb{D}^2(4 + 3X) = 45$.

14. With $p = 1/3$, $\mathbb{E}(X) = 3$, $\mathbb{D}^2(X) = 6$. Then

$$\mathbb{E}((3 - X)^2) = 6, \quad \mathbb{D}(5 - 2X) = \sqrt{24} \approx 4.899.$$