

Decision Making under Model Uncertainty

Marco Cattaneo
Department of Statistics, LMU Munich
`cattaneo@stat.uni-muenchen.de`

April 24, 2009

past and present research

- ▶ foundations of statistics
(reasoning and decision making under uncertainty)

past and present research

- ▶ foundations of statistics
(reasoning and decision making under uncertainty)
- ▶ PhD with Frank Hampel at ETH Zurich
(November 2002 – March 2007):
Statistical Decisions Based Directly on the Likelihood Function

past and present research

- ▶ foundations of statistics
(reasoning and decision making under uncertainty)
- ▶ PhD with Frank Hampel at ETH Zurich
(November 2002 – March 2007):
Statistical Decisions Based Directly on the Likelihood Function
- ▶ complex uncertainty:
simple randomness is superposed by non-stochastic aspects of
uncertainty (model uncertainty)

past and present research

- ▶ foundations of statistics
(reasoning and decision making under uncertainty)
- ▶ PhD with Frank Hampel at ETH Zurich
(November 2002 – March 2007):
Statistical Decisions Based Directly on the Likelihood Function
- ▶ complex uncertainty:
simple randomness is superposed by non-stochastic aspects of
uncertainty (model uncertainty)
- ▶ Postdoc with Thomas Augustin at LMU Munich
(SNSF Research Fellowship, October 2007 – March 2009):
*Decision making on the basis of a probabilistic-possibilistic
hierarchical description of uncertain knowledge*

decision making

- ▶ models of reality: $m \in \mathcal{M}$

decision making

- ▶ models of reality: $m \in \mathcal{M}$
- ▶ consequences of decisions: $c \in \mathcal{C}$

decision making

- ▶ models of reality: $m \in \mathcal{M}$
- ▶ consequences of decisions: $c \in \mathcal{C}$
- ▶ **decisions:** $d : \mathcal{M} \longrightarrow \mathcal{C}$

decision making

- ▶ models of reality: $m \in \mathcal{M}$
- ▶ consequences of decisions: $c \in \mathcal{C}$
- ▶ **decisions:** $d : \mathcal{M} \longrightarrow \mathcal{C}$
- ▶ simplifying assumption: $\mathcal{C} = [0, \infty)$,
 $d(m)$ is the loss incurred by making decision d , according to model m

decision making

- ▶ models of reality: $m \in \mathcal{M}$
- ▶ consequences of decisions: $c \in \mathcal{C}$
- ▶ **decisions:** $d : \mathcal{M} \longrightarrow \mathcal{C}$
- ▶ simplifying assumption: $\mathcal{C} = [0, \infty)$,
 $d(m)$ is the loss incurred by making decision d , according to model m
- ▶ **minimax** decision criterion:

$$\text{minimize} \quad \sup_{m \in \mathcal{M}} d(m)$$

statistical information

- ▶ P_m describes the probabilistic forecasts associated with model $m \in \mathcal{M}$

statistical information

- ▶ P_m describes the probabilistic forecasts associated with model $m \in \mathcal{M}$
- ▶ if the event A describes the observed data, then a **likelihood** function $l : \mathcal{M} \rightarrow [0, \infty)$ with

$$l(m) \propto P_m(A) \quad \text{for all } m \in \mathcal{M}$$

describes the (relative) forecasting ability of the models in \mathcal{M}

statistical information

- ▶ P_m describes the probabilistic forecasts associated with model $m \in \mathcal{M}$
- ▶ if the event A describes the observed data, then a **likelihood** function $l : \mathcal{M} \rightarrow [0, \infty)$ with

$$l(m) \propto P_m(A) \quad \text{for all } m \in \mathcal{M}$$

describes the (relative) forecasting ability of the models in \mathcal{M}

- ▶ **maximum likelihood** decision criterion:

$$\text{minimize } d(m_{ML}),$$

where m_{ML} is the only absolute maximum point of l

statistical information

- ▶ P_m describes the probabilistic forecasts associated with model $m \in \mathcal{M}$
- ▶ if the event A describes the observed data, then a **likelihood** function $l : \mathcal{M} \rightarrow [0, \infty)$ with

$$l(m) \propto P_m(A) \quad \text{for all } m \in \mathcal{M}$$

describes the (relative) forecasting ability of the models in \mathcal{M}

- ▶ **maximum likelihood** decision criterion:

$$\text{minimize } d(m_{ML}),$$

where m_{ML} is the only absolute maximum point of l

- ▶ **Bayesian** decision criterion with prior probability measure π on \mathcal{M} :

$$\text{minimize } \int d(m) l(m) d\pi(m)$$

nonadditive measures and integrals

- ▶ in the likelihood approach to statistics, the likelihood function is usually extended to the nonadditive measure λ on \mathcal{M} defined by

$$\lambda(M) = \sup_{m \in M} l(m) \quad \text{for all } M \subseteq \mathcal{M}$$

nonadditive measures and integrals

- ▶ in the likelihood approach to statistics, the likelihood function is usually extended to the nonadditive measure λ on \mathcal{M} defined by

$$\lambda(M) = \sup_{m \in M} l(m) \quad \text{for all } M \subseteq \mathcal{M}$$

- ▶ λ is a completely maxitive measure (**possibility** measure):

$$\lambda\left(\bigcup_{M \in \mathcal{S}} M\right) = \sup_{M \in \mathcal{S}} \lambda(M) \quad \text{for all } \mathcal{S} \subseteq 2^{\mathcal{M}}$$

nonadditive measures and integrals

- ▶ in the likelihood approach to statistics, the likelihood function is usually extended to the nonadditive measure λ on \mathcal{M} defined by

$$\lambda(M) = \sup_{m \in M} l(m) \quad \text{for all } M \subseteq \mathcal{M}$$

- ▶ λ is a completely maxitive measure (**possibility** measure):

$$\lambda\left(\bigcup_{M \in \mathcal{S}} M\right) = \sup_{M \in \mathcal{S}} \lambda(M) \quad \text{for all } \mathcal{S} \subseteq 2^{\mathcal{M}}$$

- ▶ **likelihood-based** decision criterion:

$$\text{minimize} \quad \int d(m) \, d\lambda(m)$$

likelihood-based statistical decisions

- ▶ the likelihood-based decisions share the properties of the likelihood-based inferences, in particular:
 - ▶ asymptotic optimality (consistency)
 - ▶ asymptotic efficiency
 - ▶ parametrization invariance
 - ▶ equivariance

likelihood-based statistical decisions

- ▶ the likelihood-based decisions share the properties of the likelihood-based inferences, in particular:
 - ▶ asymptotic optimality (consistency)
 - ▶ asymptotic efficiency
 - ▶ parametrization invariance
 - ▶ equivariance
- ▶ a simple property (sure-thing principle) characterizes the likelihood-based decision criterion based on the Shilkret integral: the **MPL** (Minimax Plausibility-weighted Loss) decision criterion:

$$\text{minimize} \quad \int^S d(m) d\lambda(m) = \sup_{m \in \mathcal{M}} d(m) l(m)$$

example: estimation of variance components

- ▶ estimation of the variance components in the 3×3 random effect one-way layout, under normality assumptions and weighted squared error loss

$$X_{ij} = \mu + \alpha_i + \varepsilon_{ij} \quad \text{for all } i, j \in \{1, 2, 3\}$$

example: estimation of variance components

- ▶ estimation of the variance components in the 3×3 random effect one-way layout, under normality assumptions and weighted squared error loss

$$X_{ij} = \mu + \alpha_i + \varepsilon_{ij} \quad \text{for all } i, j \in \{1, 2, 3\}$$

- ▶ normality assumptions:

$$\alpha_i \sim \mathcal{N}(0, v_a), \quad \varepsilon_{ij} \sim \mathcal{N}(0, v_e), \quad \text{all independent}$$

$$\Rightarrow X_{ij} \sim \mathcal{N}(\mu, v_a + v_e) \quad \text{dependent, } \mu \in \mathbb{R}, \quad v_a, v_e \in (0, \infty)$$

example: estimation of variance components

- ▶ estimation of the variance components in the 3×3 random effect one-way layout, under normality assumptions and weighted squared error loss

$$X_{ij} = \mu + \alpha_i + \varepsilon_{ij} \quad \text{for all } i, j \in \{1, 2, 3\}$$

- ▶ normality assumptions:

$$\alpha_i \sim \mathcal{N}(0, v_a), \quad \varepsilon_{ij} \sim \mathcal{N}(0, v_e), \quad \text{all independent}$$

$$\Rightarrow X_{ij} \sim \mathcal{N}(\mu, v_a + v_e) \quad \text{dependent, } \mu \in \mathbb{R}, \quad v_a, v_e \in (0, \infty)$$

- ▶ probabilistic models:

$$\mathcal{M} = \{m_{\mu, v_a, v_e} : \mu \in \mathbb{R}, \quad v_a, v_e \in (0, \infty)\}$$

example: estimation of variance components

- estimates \hat{v}_e and \hat{v}_a of variance components v_e and v_a are functions of

$$SS_e = \sum_{i=1}^3 \sum_{j=1}^3 (x_{ij} - \bar{x}_{i.})^2 \quad \text{and} \quad SS_a = 3 \sum_{i=1}^3 (\bar{x}_{i.} - \bar{x}_{..})^2,$$

where

$$\bar{x}_{i.} = \frac{1}{3} \sum_{j=1}^3 x_{ij}, \quad \bar{x}_{..} = \frac{1}{9} \sum_{i=1}^3 \sum_{j=1}^3 x_{ij},$$

$$\frac{SS_e}{v_e} \sim \chi_6^2, \quad \text{and} \quad \frac{\frac{1}{3} SS_a}{v_a + \frac{1}{3} v_e} \sim \chi_2^2$$

example: estimation of variance components

- estimates \hat{v}_e and \hat{v}_a of variance components v_e and v_a are functions of

$$SS_e = \sum_{i=1}^3 \sum_{j=1}^3 (x_{ij} - \bar{x}_{i.})^2 \quad \text{and} \quad SS_a = 3 \sum_{i=1}^3 (\bar{x}_{i.} - \bar{x}_{..})^2,$$

where

$$\bar{x}_{i.} = \frac{1}{3} \sum_{j=1}^3 x_{ij}, \quad \bar{x}_{..} = \frac{1}{9} \sum_{i=1}^3 \sum_{j=1}^3 x_{ij},$$


$$\frac{SS_e}{v_e} \sim \chi_6^2, \quad \text{and} \quad \frac{\frac{1}{3} SS_a}{v_a + \frac{1}{3} v_e} \sim \chi_2^2$$

- decisions (estimates $\hat{v}_e, \hat{v}_a \in (0, \infty)$ with invariant loss):


$$d_{\hat{v}_e}(m_{\mu, v_a, v_e}) = 3 \frac{(v_e - \hat{v}_e)^2}{v_e^2} \quad \text{and} \quad d_{\hat{v}_a}(m_{\mu, v_a, v_e}) = \frac{(v_a - \hat{v}_a)^2}{(v_a + \frac{1}{3} v_e)^2}$$

example: estimation of variance components

$$\frac{\widehat{v_e}}{SS_a + SS_e}$$




$$\frac{\widehat{v_a}}{SS_a + SS_e}$$




example: estimation of variance components

$$3 \frac{E[(\hat{v}_e - v_e)^2]}{v_e^2}$$



$$\frac{E[(\hat{v}_a - v_a)^2]}{(v_a + \frac{1}{3} v_e)^2}$$



present and future research

- ▶ comparison/combination with other approaches in various applications

present and future research

- ▶ comparison/combination with other approaches in various applications
- ▶ application to robustness problems
(relaxation of i.i.d. assumption, robust likelihood)

present and future research

- ▶ comparison/combination with other approaches in various applications
- ▶ application to robustness problems
(relaxation of i.i.d. assumption, robust likelihood)
- ▶ application to imprecise probability theory
(better updating, possibilistic previsions, convex set of non-normalized measures)

present and future research

- ▶ comparison/combination with other approaches in various applications
- ▶ application to robustness problems
(relaxation of i.i.d. assumption, robust likelihood)
- ▶ application to imprecise probability theory
(better updating, possibilistic previsions, convex set of non-normalized measures)
- ▶ application to graphical models
(probabilistic and non-probabilistic aspects of uncertainty)

present and future research

- ▶ comparison/combination with other approaches in various applications
- ▶ application to robustness problems
(relaxation of i.i.d. assumption, robust likelihood)
- ▶ application to imprecise probability theory
(better updating, possibilistic previsions, convex set of non-normalized measures)
- ▶ application to graphical models
(probabilistic and non-probabilistic aspects of uncertainty)
- ▶ application to financial risk measures
(derivation and interpretation of convex risk measures)