

# **A Continuous Updating Rule for Imprecise Probabilities**

Marco Cattaneo

Department of Statistics, LMU Munich

IPMU 2014, Montpellier, France

18 July 2014

## previsions

- ▶ let  $\Omega$  be a nonempty set (of possible states of the world) and  $\mathcal{L}$  be the set of all bounded functions  $X : \Omega \rightarrow \mathbb{R}$  (describing an uncertain quantity)

# previsions

- ▶ let  $\Omega$  be a nonempty set (of possible states of the world) and  $\mathcal{L}$  be the set of all bounded functions  $X : \Omega \rightarrow \mathbb{R}$  (describing an uncertain quantity)
- ▶ let  $\mathcal{P}$  be the set of all **linear previsions**  $P : \mathcal{L} \rightarrow \mathbb{R}$ , i.e., all expectation functionals  $P : X \rightarrow \int X d\pi$  corresponding to some finitely additive probability measure  $\pi$  on  $\Omega$  (Bhaskara Rao and Bhaskara Rao, 1983)

# previsions

- ▶ let  $\Omega$  be a nonempty set (of possible states of the world) and  $\mathcal{L}$  be the set of all bounded functions  $X : \Omega \rightarrow \mathbb{R}$  (describing an uncertain quantity)
- ▶ let  $\mathcal{P}$  be the set of all **linear previsions**  $P : \mathcal{L} \rightarrow \mathbb{R}$ , i.e., all expectation functionals  $P : X \rightarrow \int X d\pi$  corresponding to some finitely additive probability measure  $\pi$  on  $\Omega$  (Bhaskara Rao and Bhaskara Rao, 1983)
- ▶ let  $\underline{\mathcal{P}}$  be the set of all **coherent lower previsions**  $\underline{P} : \mathcal{L} \rightarrow \mathbb{R}$ , i.e., all functionals of the form  $\underline{P} = \min \mathcal{M}(\underline{P})$  with  $\mathcal{M}(\underline{P}) = \{P \in \mathcal{P} : P \geq \underline{P}\}$  (Walley, 1991)

# previsions

- ▶ let  $\Omega$  be a nonempty set (of possible states of the world) and  $\mathcal{L}$  be the set of all bounded functions  $X : \Omega \rightarrow \mathbb{R}$  (describing an uncertain quantity)
- ▶ let  $\mathcal{P}$  be the set of all **linear previsions**  $P : \mathcal{L} \rightarrow \mathbb{R}$ , i.e., all expectation functionals  $P : X \rightarrow \int X d\pi$  corresponding to some finitely additive probability measure  $\pi$  on  $\Omega$  (Bhaskara Rao and Bhaskara Rao, 1983)
- ▶ let  $\underline{\mathcal{P}}$  be the set of all **coherent lower previsions**  $\underline{P} : \mathcal{L} \rightarrow \mathbb{R}$ , i.e., all functionals of the form  $\underline{P} = \min \mathcal{M}(\underline{P})$  with  $\mathcal{M}(\underline{P}) = \{P \in \mathcal{P} : P \geq \underline{P}\}$  (Walley, 1991)
- ▶ the upper prevision conjugate to  $\underline{P} \in \underline{\mathcal{P}}$  is the functional  $\bar{P} = \max \mathcal{M}(\underline{P})$

# previsions

- ▶ let  $\Omega$  be a nonempty set (of possible states of the world) and  $\mathcal{L}$  be the set of all bounded functions  $X : \Omega \rightarrow \mathbb{R}$  (describing an uncertain quantity)
- ▶ let  $\mathcal{P}$  be the set of all **linear previsions**  $P : \mathcal{L} \rightarrow \mathbb{R}$ , i.e., all expectation functionals  $P : X \rightarrow \int X d\pi$  corresponding to some finitely additive probability measure  $\pi$  on  $\Omega$  (Bhaskara Rao and Bhaskara Rao, 1983)
- ▶ let  $\underline{\mathcal{P}}$  be the set of all **coherent lower previsions**  $\underline{P} : \mathcal{L} \rightarrow \mathbb{R}$ , i.e., all functionals of the form  $\underline{P} = \min \mathcal{M}(\underline{P})$  with  $\mathcal{M}(\underline{P}) = \{P \in \mathcal{P} : P \geq \underline{P}\}$  (Walley, 1991)
- ▶ the upper prevision conjugate to  $\underline{P} \in \underline{\mathcal{P}}$  is the functional  $\overline{P} = \max \mathcal{M}(\underline{P})$
- ▶  $P, \underline{P}, \overline{P}$  denote probabilities as well as previsions: e.g.,  $P(A) = P(I_A)$

# updating

- ▶ when an event  $B \subset \Omega$  is observed, the previsions must be updated (to avoid trivial results,  $|B| > 1$  is assumed)

# updating

- ▶ when an event  $B \subset \Omega$  is observed, the previsions must be updated (to avoid trivial results,  $|B| > 1$  is assumed)
- ▶ **Bayesian updating:** each  $P \in \mathcal{P}$  with  $P(B) > 0$  is updated to

$$P(\cdot | B) : X \mapsto P(X | B) = \frac{P(X I_B)}{P(B)}$$



# updating

- ▶ when an event  $B \subset \Omega$  is observed, the previsions must be updated (to avoid trivial results,  $|B| > 1$  is assumed)
- ▶ **Bayesian updating:** each  $P \in \mathcal{P}$  with  $P(B) > 0$  is updated to

$$P(\cdot | B) : X \mapsto P(X | B) = \frac{P(X I_B)}{P(B)}$$

- ▶ **regular extension:** each  $\underline{P} \in \underline{\mathcal{P}}$  with  $\overline{P}(B) > 0$  is updated to

$$\underline{P}(\cdot | B) : X \mapsto \underline{P}(X | B) = \inf_{P \in \mathcal{M}(\underline{P}) : P(B) > 0} P(X | B)$$

## example

- ▶ Piatti et al. (2009) studied exchangeable sequences of binary experiments  $X_1, X_2, \dots \in \{0, 1\}$  such that the realization of each  $X_i$  can be observed incorrectly with a known probability  $\varepsilon$  (the errors of observation are independent, conditional on the realizations of  $X_1, X_2, \dots$ )

## example

- ▶ Piatti et al. (2009) studied exchangeable sequences of binary experiments  $X_1, X_2, \dots \in \{0, 1\}$  such that the realization of each  $X_i$  can be observed incorrectly with a known probability  $\varepsilon$  (the errors of observation are independent, conditional on the realizations of  $X_1, X_2, \dots$ )
- ▶ event  $B$ : (possibly incorrect) observation of 7 times “1” and 2 times “0” in the first 9 experiments

## example

- ▶ Piatti et al. (2009) studied exchangeable sequences of binary experiments  $X_1, X_2, \dots \in \{0, 1\}$  such that the realization of each  $X_i$  can be observed incorrectly with a known probability  $\varepsilon$  (the errors of observation are independent, conditional on the realizations of  $X_1, X_2, \dots$ )
- ▶ event  $B$ : (possibly incorrect) observation of 7 times “1” and 2 times “0” in the first 9 experiments
- ▶ Bayesian conjugate model  $P$  with Jeffreys’ prior, and IDM model  $\underline{P}$  with hyperparameter  $s = 1$ :

	$\varepsilon = 0$	$\varepsilon = 10^{-6}$
$P(X_1) = 0.5$	$P(X_{10}   B) = 0.75$	$P(X_{10}   B) \approx 0.750$
$\bar{P}(X_1) = 1$	$\bar{P}(X_{10}   B) = 0.8$	$\bar{P}(X_{10}   B) = 1$
$\underline{P}(X_1) = 0$	$\underline{P}(X_{10}   B) = 0.7$	$\underline{P}(X_{10}   B) = 0$

## continuity

- ▶ metric on  $\mathcal{P}$  (dual norm =  $2 \times$  total variation distance):

$$d(P, P') = \sup_{X: \Omega \rightarrow [-1, 1]} |P(X) - P'(X)| = 2 \sup_{A \subseteq \Omega} |P(A) - P'(A)|$$

## continuity

- ▶ metric on  $\mathcal{P}$  (dual norm =  $2 \times$  total variation distance):

$$d(P, P') = \sup_{X: \Omega \rightarrow [-1, 1]} |P(X) - P'(X)| = 2 \sup_{A \subseteq \Omega} |P(A) - P'(A)|$$

- ▶ metric on  $\underline{\mathcal{P}}$  (Hausdorff distance = “dual norm”):

$$d(\underline{P}, \underline{P}') = d_H(\mathcal{M}(\underline{P}), \mathcal{M}(\underline{P}')) = \sup_{X: \Omega \rightarrow [-1, 1]} |\underline{P}(X) - \underline{P}'(X)|$$

## continuity

- ▶ metric on  $\mathcal{P}$  (dual norm =  $2 \times$  total variation distance):

$$d(P, P') = \sup_{X: \Omega \rightarrow [-1, 1]} |P(X) - P'(X)| = 2 \sup_{A \subseteq \Omega} |P(A) - P'(A)|$$

- ▶ metric on  $\underline{\mathcal{P}}$  (Hausdorff distance = “dual norm”):

$$d(\underline{P}, \underline{P}') = d_H(\mathcal{M}(\underline{P}), \mathcal{M}(\underline{P}')) = \sup_{X: \Omega \rightarrow [-1, 1]} |\underline{P}(X) - \underline{P}'(X)|$$

- ▶ the Bayesian updating  $\{P \in \mathcal{P} : P(B) > 0\} \rightarrow \{P \in \mathcal{P} : P(B) = 1\}$  is **continuous**

# continuity

- ▶ metric on  $\mathcal{P}$  (dual norm =  $2 \times$  total variation distance):

$$d(P, P') = \sup_{X: \Omega \rightarrow [-1, 1]} |P(X) - P'(X)| = 2 \sup_{A \subseteq \Omega} |P(A) - P'(A)|$$

- ▶ metric on  $\underline{\mathcal{P}}$  (Hausdorff distance = “dual norm”):

$$d(\underline{P}, \underline{P}') = d_H(\mathcal{M}(\underline{P}), \mathcal{M}(\underline{P}')) = \sup_{X: \Omega \rightarrow [-1, 1]} |\underline{P}(X) - \underline{P}'(X)|$$

- ▶ the Bayesian updating  $\{P \in \mathcal{P} : P(B) > 0\} \rightarrow \{P \in \mathcal{P} : P(B) = 1\}$  is **continuous**
- ▶ the regular extension  $\{\underline{P} \in \underline{\mathcal{P}} : \overline{P}(B) > 0\} \rightarrow \{\underline{P} \in \underline{\mathcal{P}} : \underline{P}(B) = 1\}$  is **not** continuous: it has discontinuities at points  $\underline{P} \in \underline{\mathcal{P}}$  with  $\overline{P}(B) > \underline{P}(B) = 0$



# updating

- ▶ when an event  $B \subset \Omega$  is observed, the previsions must be updated (to avoid trivial results,  $|B| > 1$  is assumed)
- ▶ **Bayesian updating:** each  $P \in \mathcal{P}$  with  $P(B) > 0$  is updated to

$$P(\cdot | B) : X \mapsto P(X | B) = \frac{P(X I_B)}{P(B)}$$

- ▶ **regular extension:** each  $\underline{P} \in \underline{\mathcal{P}}$  with  $\bar{P}(B) > 0$  is updated to

$$\underline{P}(\cdot | B) : X \mapsto \underline{P}(X | B) = \inf_{P \in \mathcal{M}(\underline{P}) : P(B) > 0} P(X | B)$$

- ▶  **$\alpha$ -cut rule**, where  $\alpha \in (0, 1)$ : each  $\underline{P} \in \underline{\mathcal{P}}$  with  $\bar{P}(B) > 0$  is updated to

$$\underline{P}_\alpha(\cdot | B) : X \mapsto \underline{P}_\alpha(X | B) = \min_{P \in \mathcal{M}(\underline{P}) : P(B) \geq \alpha \bar{P}(B)} P(X | B)$$

## example

- ▶ Piatti et al. (2009) studied exchangeable sequences of binary experiments  $X_1, X_2, \dots \in \{0, 1\}$  such that the realization of each  $X_i$  can be observed incorrectly with a known probability  $\varepsilon$  (the errors of observation are independent, conditional on the realizations of  $X_1, X_2, \dots$ )
- ▶ event  $B$ : (possibly incorrect) observation of 7 times “1” and 2 times “0” in the first 9 experiments
- ▶ Bayesian conjugate model  $P$  with Jeffreys’ prior, and IDM model  $\underline{P}$  with hyperparameter  $s = 1$  (and  $\alpha = 0.001$ ):

	$\varepsilon = 0$	$\varepsilon = 10^{-6}$
$P(X_1) = 0.5$	$P(X_{10}   B) = 0.75$	$P(X_{10}   B) \approx 0.750$
$\overline{P}(X_1) = 1$	$\overline{P}(X_{10}   B) = 0.8$	$\overline{P}(X_{10}   B) = 1$
$\underline{P}(X_1) = 0$	$\underline{P}(X_{10}   B) = 0.7$	$\underline{P}(X_{10}   B) = 0$
$\overline{P}(X_1) = 1$	$\overline{P}_\alpha(X_{10}   B) \approx 0.800$	$\overline{P}_\alpha(X_{10}   B) \approx 0.800$
$\underline{P}(X_1) = 0$	$\underline{P}_\alpha(X_{10}   B) \approx 0.700$	$\underline{P}_\alpha(X_{10}   B) \approx 0.700$

# continuity

- ▶ metric on  $\mathcal{P}$  (dual norm =  $2 \times$  total variation distance):

$$d(P, P') = \sup_{X: \Omega \rightarrow [-1, 1]} |P(X) - P'(X)| = 2 \sup_{A \subseteq \Omega} |P(A) - P'(A)|$$

- ▶ metric on  $\underline{\mathcal{P}}$  (Hausdorff distance = “dual norm”):

$$d(\underline{P}, \underline{P}') = d_H(\mathcal{M}(\underline{P}), \mathcal{M}(\underline{P}')) = \sup_{X: \Omega \rightarrow [-1, 1]} |\underline{P}(X) - \underline{P}'(X)|$$

- ▶ the Bayesian updating  $\{P \in \mathcal{P} : P(B) > 0\} \rightarrow \{P \in \mathcal{P} : P(B) = 1\}$  is **continuous**
- ▶ the regular extension  $\{\underline{P} \in \underline{\mathcal{P}} : \overline{P}(B) > 0\} \rightarrow \{\underline{P} \in \underline{\mathcal{P}} : \underline{P}(B) = 1\}$  is **not** continuous: it has discontinuities at points  $\underline{P} \in \underline{\mathcal{P}}$  with  $\overline{P}(B) > \underline{P}(B) = 0$
- ▶ the  $\alpha$ -cut rule  $\{\underline{P} \in \underline{\mathcal{P}} : \overline{P}(B) > 0\} \rightarrow \{\underline{P} \in \underline{\mathcal{P}} : \underline{P}(B) = 1\}$  is **continuous**, for all  $\alpha \in (0, 1)$

## conclusion

- ▶ the Bayesian updating of linear previsions is continuous

## conclusion

- ▶ the Bayesian updating of linear previsions is continuous
- ▶ the regular extension, as well as the natural extension and any other updating rule that is **coherent** in the sense of Walley (1991):

# conclusion

- ▶ the Bayesian updating of linear previsions is continuous
- ▶ the regular extension, as well as the natural extension and any other updating rule that is **coherent** in the sense of Walley (1991):
  - ▶ generalizes Bayesian updating to lower previsions

## conclusion

- ▶ the Bayesian updating of linear previsions is continuous
- ▶ the regular extension, as well as the natural extension and any other updating rule that is **coherent** in the sense of Walley (1991):
  - ▶ generalizes Bayesian updating to lower previsions
  - ▶ is **discontinuous** when  $\underline{P}(B) = 0$ , and this leads to robustness issues

# conclusion

- ▶ the Bayesian updating of linear previsions is continuous
- ▶ the regular extension, as well as the natural extension and any other updating rule that is **coherent** in the sense of Walley (1991):
  - ▶ generalizes Bayesian updating to lower previsions
  - ▶ is **discontinuous** when  $\underline{P}(B) = 0$ , and this leads to robustness issues
- ▶ the  $\alpha$ -cut rule:



## conclusion

- ▶ the Bayesian updating of linear previsions is continuous
- ▶ the regular extension, as well as the natural extension and any other updating rule that is **coherent** in the sense of Walley (1991):
  - ▶ generalizes Bayesian updating to lower previsions
  - ▶ is **discontinuous** when  $\underline{P}(B) = 0$ , and this leads to robustness issues
- ▶ the  $\alpha$ -cut rule:
  - ▶ generalizes Bayesian updating, and corresponds to regular/natural extension (and any other coherent updating rule) when  $\underline{P}(B) \geq \alpha \bar{P}(B) > 0$

# conclusion

- ▶ the Bayesian updating of linear previsions is continuous
- ▶ the regular extension, as well as the natural extension and any other updating rule that is **coherent** in the sense of Walley (1991):
  - ▶ generalizes Bayesian updating to lower previsions
  - ▶ is **discontinuous** when  $\underline{P}(B) = 0$ , and this leads to robustness issues
- ▶ the  $\alpha$ -cut rule:
  - ▶ generalizes Bayesian updating, and corresponds to regular/natural extension (and any other coherent updating rule) when  $\underline{P}(B) \geq \alpha \bar{P}(B) > 0$
  - ▶ is **continuous**

# conclusion

- ▶ the Bayesian updating of linear previsions is continuous
- ▶ the regular extension, as well as the natural extension and any other updating rule that is **coherent** in the sense of Walley (1991):
  - ▶ generalizes Bayesian updating to lower previsions
  - ▶ is **discontinuous** when  $\underline{P}(B) = 0$ , and this leads to robustness issues
- ▶ the  $\alpha$ -cut rule:
  - ▶ generalizes Bayesian updating, and corresponds to regular/natural extension (and any other coherent updating rule) when  $\underline{P}(B) \geq \alpha \overline{P}(B) > 0$
  - ▶ is **continuous**
  - ▶ allows **vacuous priors** in statistical analyses, and has been successfully applied (Antonucci et al., 2012; Cattaneo and Wiencierz, 2012; Destercke, 2013)

# conclusion

- ▶ the Bayesian updating of linear previsions is continuous
- ▶ the regular extension, as well as the natural extension and any other updating rule that is **coherent** in the sense of Walley (1991):
  - ▶ generalizes Bayesian updating to lower previsions
  - ▶ is **discontinuous** when  $\underline{P}(B) = 0$ , and this leads to robustness issues
- ▶ the  $\alpha$ -cut rule:
  - ▶ generalizes Bayesian updating, and corresponds to regular/natural extension (and any other coherent updating rule) when  $\underline{P}(B) \geq \alpha \overline{P}(B) > 0$
  - ▶ is **continuous**
  - ▶ allows **vacuous priors** in statistical analyses, and has been successfully applied (Antonucci et al., 2012; Cattaneo and Wiencierz, 2012; Destercke, 2013)
  - ▶ lies between regular extension (limit  $\alpha \rightarrow 0$ ) and a generalization of Dempster's rule of conditioning (limit  $\alpha \rightarrow 1$ ), and is strictly related to other proposals (Moral, 1992; Gilboa and Schmeidler, 1993; Cano and Moral, 1996)

# conclusion

- ▶ the Bayesian updating of linear previsions is continuous
- ▶ the regular extension, as well as the natural extension and any other updating rule that is **coherent** in the sense of Walley (1991):
  - ▶ generalizes Bayesian updating to lower previsions
  - ▶ is **discontinuous** when  $\underline{P}(B) = 0$ , and this leads to robustness issues
- ▶ the  $\alpha$ -cut rule:
  - ▶ generalizes Bayesian updating, and corresponds to regular/natural extension (and any other coherent updating rule) when  $\underline{P}(B) \geq \alpha \bar{P}(B) > 0$
  - ▶ is **continuous**
  - ▶ allows **vacuous priors** in statistical analyses, and has been successfully applied (Antonucci et al., 2012; Cattaneo and Wiencierz, 2012; Destercke, 2013)
  - ▶ lies between regular extension (limit  $\alpha \rightarrow 0$ ) and a generalization of Dempster's rule of conditioning (limit  $\alpha \rightarrow 1$ ), and is strictly related to other proposals (Moral, 1992; Gilboa and Schmeidler, 1993; Cano and Moral, 1996)
  - ▶ is not iteratively consistent (i.e., updates do not commute), but this can be remedied by recording the whole likelihood function (Cattaneo, 2008, 2009)

# conclusion

- ▶ the Bayesian updating of linear previsions is continuous
- ▶ the regular extension, as well as the natural extension and any other updating rule that is **coherent** in the sense of Walley (1991):
  - ▶ generalizes Bayesian updating to lower previsions
  - ▶ is **discontinuous** when  $\underline{P}(B) = 0$ , and this leads to robustness issues
- ▶ the  $\alpha$ -cut rule:
  - ▶ generalizes Bayesian updating, and corresponds to regular/natural extension (and any other coherent updating rule) when  $\underline{P}(B) \geq \alpha \overline{P}(B) > 0$
  - ▶ is **continuous**
  - ▶ allows **vacuous priors** in statistical analyses, and has been successfully applied (Antonucci et al., 2012; Cattaneo and Wiencierz, 2012; Destercke, 2013)
  - ▶ lies between regular extension (limit  $\alpha \rightarrow 0$ ) and a generalization of Dempster's rule of conditioning (limit  $\alpha \rightarrow 1$ ), and is strictly related to other proposals (Moral, 1992; Gilboa and Schmeidler, 1993; Cano and Moral, 1996)
  - ▶ is not iteratively consistent (i.e., updates do not commute), but this can be remedied by recording the whole likelihood function (Cattaneo, 2008, 2009)
- ▶ the relative importance of properties like coherence, continuity, and ability of using vacuous priors depends on the application field and on the exact interpretation of imprecise probabilities

# references

- Antonucci, A., Cattaneo, M., and Corani, G. (2012). Likelihood-based robust classification with Bayesian networks. In *IPMU '12*. Vol. 3. Springer, 491–500.
- Bhaskara Rao, K. P. S., and Bhaskara Rao, M. (1983). *Theory of Charges*. Academic Press.
- Cano, A., and Moral, S. (1996). A genetic algorithm to approximate convex sets of probabilities. In *IPMU '96*. Vol. 2. Proyecto Sur, 859–864.
- Cattaneo, M. (2008). Fuzzy probabilities based on the likelihood function. In *SMPS '08*. Springer, 43–50.
- Cattaneo, M. (2009). A generalization of credal networks. In *ISIPTA '09*. SIPTA, 79–88.
- Cattaneo, M., and Wiencierz, A. (2012). Likelihood-based Imprecise Regression. *Int. J. Approx. Reasoning* 53, 1137–1154.
- Destercke, S. (2013). A pairwise label ranking method with imprecise scores and partial predictions. In *ECMLPKDD '13*. Vol. 2. Springer, 112–127.
- Gilboa, I., and Schmeidler, D. (1993). Updating ambiguous beliefs. *J. Econ. Theory* 59, 33–49.
- Moral, S. (1992). Calculating uncertainty intervals from conditional convex sets of probabilities. In *UAI '92*. Morgan Kaufmann, 199–206.
- Piatti, A., Zaffalon, M., Trojani, F., and Hutter, M. (2009). Limits of learning about a categorical latent variable under prior near-ignorance. *Int. J. Approx. Reasoning* 50, 597–611.
- Walley, P. (1991). *Statistical Reasoning with Imprecise Probabilities*. Chapman and Hall.