

Introduction to Differentiable Manifolds

original notes by Anna Kiesenhofer
modified by Marcos Cossarini

EPFL, autumn 2021
date of this version: 2021-11-23

Contents

0.1	Practical remarks about the course	3
0.1.1	Content for the exam	3
1	Manifolds	4
1.1	Topological manifolds	4
1.2	Differentiable manifolds	7
1.2.1	One-step \mathcal{C}^k manifolds*	9
1.3	Differentiable maps	10
1.4	Partitions of unity	12
1.4.1	Paracompactness*	13
1.4.2	Applications	14
2	Tangent vectors	16
2.0.1	Differential of a \mathcal{C}^k map between manifolds	18
2.0.2	Pushforward and tangent operators	24
2.0.3	Classical notation for tangent vectors	26
2.0.4	General derivations*	26
3	Submanifolds	27
3.1	How to recognize an embedding	28
3.2	Constant rank theorem	29
3.3	Some properties of immersions and embeddings	32
3.3.1	Local properties: slices and retractions	32
3.3.2	The initial property	33
3.3.3	Consequences of the initial property	34
3.4	How to recognize an embedded submanifold	34
3.5	How to produce a submanifold as a level set	35
3.6	Whitney's theorem*	36
4	Vector bundles and vector fields	37
4.1	The tangent bundle	37
4.2	Sections, frames, and trivial bundles	38
4.3	Vector fields	40
4.4	Integral curves and flow of a vector field	40
5	Covectors and 1-forms	43
5.1	Vectors and covectors; superindices and subindices	43
5.2	Cotangent vectors	43
5.3	Covector fields	44
5.4	Contraction of a 1-form with a vector field	45
5.5	Differential of a function	45
5.6	Pullback of a 1-form	45
5.7	Integration of a 1-form along a curve	46
5.8	Integration along curves via pullback	47

Main references

- [Lee10] John M. Lee. *Introduction to Topological Manifolds*. Springer, 2010.
- [Lee13] John M. Lee. *Introduction to Smooth Manifolds*. Second edition. Springer, 2013.

Both books have online versions available in the library.

0.1 Practical remarks about the course

0.1.1 Content for the exam

Studying the content under a heading marked by an asterisk is not mandatory by itself. You should read it only if you find it interesting or helpful for understanding the rest.

1 Manifolds

The goal of this course is to extend differential and integral calculus from Euclidean space \mathbb{R}^n to all *differentiable manifolds* such as the n -sphere, the n -torus, etc. Roughly speaking, a differentiable manifold is a space that

- is endowed with a certain topology,
- has, in addition, a *differentiable structure* that allows us to distinguish whether a map is differentiable or not, rather than just continuous, and
- locally looks like Euclidean space \mathbb{R}^n .

1.1 Topological manifolds

[Lee13], Chapter 1 and [Lee10], Chapter 2

Let us postpone the question of differentiability and focus on topology. As said, we want to study spaces that “locally look like” Euclidean space \mathbb{R}^n .

Definition 1.1.1 (Locally Euclidean space). Let $n \in \mathbb{N} = \{0, 1, \dots\}$. A topological space M is **locally Euclidean** of dimension n at a point $p \in M$ if the point p has an open neighborhood that is homeomorphic to an open subset of \mathbb{R}^n . If this holds for all points $p \in M$, we say that M is locally Euclidean of dimension n .

A typical example is the circle: it is locally Euclidean of dimension 1 but not globally homeomorphic to any subset of \mathbb{R} .

Example 1.1.2. The **circle** $\mathbb{S}^1 = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}$ (with the subspace topology) is locally Euclidean of dimension 1: let $(x_0, y_0) \in \mathbb{S}^1$, wlog $y_0 > 0$, then $U := (\mathbb{R} \times \mathbb{R}^+) \cap \mathbb{S}^1$ is an open subset of \mathbb{S}^1 containing (x_0, y_0) and homeomorphic to $(-1, 1)$ via the map $U \rightarrow (-1, 1)$ that sends $(x, y) \mapsto x$.

We will see more examples later on (see e.g. Examples 1.1.11 below). Let us make some general comments.

Remark 1.1.3. If a space M is locally Euclidean of dimension 0, then every point has a neighborhood homeomorphic to $\mathbb{R}^0 = \{0\}$, i.e. a point. In other words, M is a discrete topological space.

Remark 1.1.4. In the definition of locally Euclidean space, we could have replaced “...homeomorphic to an open subset of \mathbb{R}^n ” by “...homeomorphic to \mathbb{R}^n ”. (Exercise.)

Remark 1.1.5. Brouwer’s theorem of *invariance of domain* implies¹ that if two nonempty open sets $U \subset \mathbb{R}^m$, $V \subset \mathbb{R}^n$ are homeomorphic, then $m = n$. It follows that the dimension of a locally Euclidean space at each point can be defined unambiguously. Furthermore, it is easy to prove that the dimension is constant throughout each connected component. Thus the only way to get a locally Euclidean space of mixed dimensions is to make a disjoint union of components of different dimensions. Anyway, in the definition of topological manifold (see below) we will not admit this kind of spaces.

¹The actual theorem says that if $U \subseteq \mathbb{R}^m$ is an open set, then any injective continuous map $f : U \rightarrow \mathbb{R}^m$ is an open map.

For the definition of *topological manifold* we demand some further topological properties that ensure that the space is topologically “well-behaved”. (For instance, we want the limit of every sequence to be unique.)

Definition 1.1.6. A **topological manifold of dimension n** , or **topological n -manifold**, is a topological space M that is locally Euclidean of dimension n , Hausdorff² and second countable.³

A **topological manifold** is a topological space that is a topological n -manifold for some n .

Side note: Make sure you are familiar with some basic definitions from topology such as Hausdorff, second countable, connected and compact spaces, and the construction of subspace, product, coproduct and quotient topologies. Chapters 2 and 3 in [Lee10] provides a succinct overview of everything we need.

Remark 1.1.7. The conditions of Hausdorff resp. second countable in Definition 1.1.6 are not redundant. For example, the *line with two origins* (see Exercises) is a locally Euclidean, second countable space that is not Hausdorff. The *long line* and the *Prüfer surface* (see Wikipedia if interested) are locally Euclidean of dimension 1 and 2 respectively, Hausdorff, and connected, but not second countable.

The homeomorphisms that locally identify a topological manifold with Euclidean space are called *charts*:

Definition 1.1.8 (Coordinate charts). Let M be a topological n -manifold. A **chart** (or **coordinate chart**) for M is a homeomorphism $\varphi : U \rightarrow V$, where $U \subseteq M$ and $V \subseteq \mathbb{R}^n$ are open sets. Its inverse φ^{-1} is a **local parametrization** of M . An **atlas** for M is a collection of charts whose domains cover M .

For the moment, we can see an atlas simply as a way of showing that a space is locally Euclidean.

Remark 1.1.9. Some authors define a chart for M as a *pair* (φ, U) or even a *triple* (φ, U, V) where $U \subseteq M$ and $V \subseteq \mathbb{R}^n$ are open sets and $\varphi : U \rightarrow V$ is a homeomorphism. Here, instead, we consider the sets U and V as part of the function φ , namely, its domain $\text{Dom}(\varphi)$ and codomain $\text{Cod}(\varphi)$, thus there’s no need not specify them separately.⁴ From this point of view, the letters “ U ”, “ V ” are just shorter names for the sets $\text{Dom}(\varphi)$, $\text{Cod}(\varphi)$.

Convention: When talking about subsets (resp. quotients, products, disjoint unions) of topological spaces we’ll assume that they are endowed with the subspace (resp. quotient, product, coproduct) topology unless otherwise stated.

Using this convention, let us mention some easy ways to construct new topological manifolds from old ones.

Proposition 1.1.10 (New manifolds from old). *The properties of being Hausdorff or second countable are preserved by taking subspaces, finite products and countable coproducts. In consequence:*

²Recall that a topological space X is **Hausdorff** if every two different points $x, y \in X$ have disjoint neighborhoods.

³Recall that a topological space X is **second countable** if its topology admits a countable base. A **base** for a topology is a family \mathcal{B} of open sets such that every open set is a union of some sets of \mathcal{B} .

⁴Formally, a function f is a triple $f = (X, Y, \Gamma)$ where X, Y are sets (called the **domain** and **codomain** of f , and denoted $\text{Dom}(f)$ and $\text{Cod}(f)$), and Γ is a subset of $X \times Y$ (called the **graph** of f , denoted $\text{Gra}(f)$), such that for each $x \in X$ there is a unique $y \in Y$ (called the **image** of x by f , denoted $f(x)$) such that $(x, y) \in \Gamma$.

- An open subset of a topological n -manifold is a topological n -manifold.
- A disjoint union $M = \coprod_i M_i$ of countably many topological n -manifolds M_i is a topological n -manifold.
- A product $M = \prod_i M_i$ of finitely many topological manifolds M_i is a topological manifold of dimension $\dim(M) = \sum_i \dim(M_i)$.

Proof. Exercise. □

Example 1.1.11 (Examples of topological manifolds).

- Of course any open subset of \mathbb{R}^n is a topological manifold.
- An example of topological n -manifold is the **graph**

$$\Gamma_f := \{(x, f(x)) \mid x \in U\} \subseteq U \times \mathbb{R}^m$$

of a continuous function $f : U \rightarrow \mathbb{R}^m$, where $U \subset \mathbb{R}^n$ is an open set. Indeed, it is homeomorphic to U via the **graph parametrization**

$$\begin{array}{ccc} U & \xrightarrow{\quad \Gamma_f \quad} & \Gamma_f \\ x & \mapsto (x, f(x)), & \text{whose inverse is the projection} \\ & & \Gamma_f \rightarrow U \\ & & (x, y) \mapsto x. \end{array}$$

- The sphere $\mathbb{S}^n \subset \mathbb{R}^{n+1}$ is a topological manifold. Being a subset of \mathbb{R}^{n+1} it is Hausdorff and second countable. A possible choice of atlas is given by the so-called **graph coordinates**: Cover \mathbb{S}^n by the $2(n+1)$ open sets $U_i^\pm := \{x \in \mathbb{R}^n \mid \pm x_i > 0\}$, then $\mathbb{S}^n \cap U_i^\pm$ is homeomorphic to the open unit n -ball \mathbb{B}^n via the projection⁵

$$\begin{array}{ccc} \varphi_i^\pm : & \mathbb{S}^n \cap U_i^\pm & \rightarrow \mathbb{B}^n \\ & (x_0, \dots, x_n) & \rightarrow (x_0, \dots, \widehat{x}_i, \dots, x_n). \end{array}$$

The maps φ_i^\pm are coordinate charts for \mathbb{S}^n ; we call them *graph coordinates*. Locally this is a special case of the previous item (b): each set $\mathbb{S}^n \cap U_i^\pm$ is (up to permutation of coordinates) the graph of the continuous function on the unit n -ball $\mathbb{B}^n(0)$:

$$\mathbb{B}^n \rightarrow \mathbb{R} : y \mapsto \pm \sqrt{1 - \sum_i y_i^2}.$$

- Real projective space \mathbb{P}^n is a topological n -manifold (exercise).
- The torus $\mathbb{T}^n = \mathbb{R}^n / \mathbb{Z}^n$ is a topological n -manifold (exercise).
- More generally, If M is a topological n -manifold and G is a group of homeomorphisms of M that acts properly discontinuously and without fixed points, then the quotient space M/G is a topological n -manifold.
- If M is a topological n -manifold and $\pi : N \rightarrow M$ is a covering map (with N connected), then N is a topological n -manifold. In particular, the universal covering space of any connected topological manifold is a topological manifold.

We will not prove the following result (although it can be done elementarily).

Theorem 1.1.12 (Classification of topological 1-manifolds). *Every connected topological 1-manifold is homeomorphic to either \mathbb{S}^1 (if it is compact) or to \mathbb{R} (if it is not compact).*

⁵The hat on \widehat{x}_i means that we omit the respective coordinate x_i .

1.2 Differentiable manifolds

Our next goal is to define a kind of spaces and maps called *differentiable manifolds* and *differentiable maps* (or, more precisely \mathcal{C}^k manifolds and maps) with which we can actually do differential calculus. Topological manifolds do not have enough structure because a topology does not allow us to determine whether a function is differentiable or not; it only distinguishes continuous functions. Differentiable manifolds should be locally equivalent to Euclidean open sets (where we already have a well defined notion of \mathcal{C}^k maps; see below), but at the global level they should be allowed to have a more interesting topology. In particular, the sphere, torus, projective space, etc. should become differentiable manifolds.

Before defining \mathcal{C}^k manifolds, let us set up some terminology for \mathcal{C}^k maps in \mathbb{R}^n .

Definition 1.2.1 (Euclidean open sets and Euclidean \mathcal{C}^k maps). A **Euclidean open set** is an open subset of some Euclidean space \mathbb{R}^n .

Let $k \in \{0, 1, \dots, \infty\}$. A function $f : U \rightarrow V$ between Euclidean open sets is \mathcal{C}^k at a point $p \in U$ if its partial derivatives of order $\leq k$ are defined in a neighborhood of p and continuous at p . We say that f is \mathcal{C}^k (and we call it a **Euclidean \mathcal{C}^k map**) if it is \mathcal{C}^k at all points $p \in U$.

An **Euclidean \mathcal{C}^k isomorphism** is an Euclidean \mathcal{C}^k map that has an Euclidean \mathcal{C}^k inverse.

Note that every Euclidean \mathcal{C}^k map is continuous because the function f itself is a partial derivative (of order 0) of f . In fact, a \mathcal{C}^0 map is the same thing as a continuous map.

We are now ready to define \mathcal{C}^k manifolds. The key to turn a topological manifold into a \mathcal{C}^k manifold is to choose an appropriate atlas.

Definition 1.2.2 (\mathcal{C}^k manifolds). Let M be a topological n -manifold and $k = 0, \dots, \infty$. Two charts φ, ψ for M , with respective domains $U, V \subseteq M$, are **\mathcal{C}^k -compatible** if the **transition map** from φ to ψ , that is, the homeomorphism

$$\psi \circ \varphi^{-1} : \varphi(U \cap V) \rightarrow \psi(U \cap V),$$

is a \mathcal{C}^k isomorphism (i.e. itself and its inverse are both Euclidean \mathcal{C}^k maps).

A **\mathcal{C}^k -consistent atlas** (or **\mathcal{C}^k atlas**, for short) is an atlas for M whose charts are \mathcal{C}^k compatible with each other. Two \mathcal{C}^k atlases for M are **\mathcal{C}^k -equivalent** if their union is \mathcal{C}^k -consistent. A **\mathcal{C}^k structure** on M is a maximal \mathcal{C}^k atlas, i.e., a \mathcal{C}^k atlas that is not contained in any other strictly larger \mathcal{C}^k atlas. A **\mathcal{C}^k manifold** is a topological manifold M endowed with a \mathcal{C}^k structure \mathcal{A} . (More formally, the \mathcal{C}^k manifold is the pair (M, \mathcal{A}) .)

Note that a \mathcal{C}^0 manifold is the same thing as a topological manifold. A \mathcal{C}^k manifold with $k \geq 1$ is called a **\mathcal{C}^k -differentiable manifold**. A **smooth manifold** is a \mathcal{C}^∞ manifold.

Remark 1.2.3 (Domains and codomains of functions). To be precise, the transition map that we wrote as $\psi \circ \varphi^{-1}$ should actually be defined as $\psi|_{\varphi(U \cap V)} \circ (\varphi|_{U \cap V})^{-1}$, using the *restricted* charts

$$\varphi|_{U \cap V}^{\varphi(U \cap V)} : U \cap V \rightarrow \varphi(U \cap V), \quad \psi|_{U \cap V}^{\psi(U \cap V)} : U \cap V \rightarrow \psi(U \cap V).$$

In general we will not write the restrictions explicitly because it is cumbersome. When we compose functions, it should be understood that the resulting composite

function is defined in principle at all points where it is possible. (Maybe no points at all!)

We may further restrict a function by specifying a reduced domain or codomain. On the other hand, we shall never specify a domain containing points where the function is not defined, nor a codomain that does not contain the image of the specified domain. Thus a function “ $f : A \rightarrow B$ ” always has domain A and codomain B .

The next proposition shows that it suffices to give any \mathcal{C}^k -consistent atlas (not necessarily a maximal one) to determine a \mathcal{C}^k structure.

Proposition 1.2.4 (\mathcal{C}^k atlas defines \mathcal{C}^k structure). *For a fixed topological manifold M , each \mathcal{C}^k atlas \mathcal{A} is contained in a unique maximal \mathcal{C}^k atlas $\overline{\mathcal{A}}$, which consists of all charts for M that are \mathcal{C}^k -compatible with those of \mathcal{A} . Any other \mathcal{C}^k atlas \mathcal{B} for M is equivalent to \mathcal{A} if and only if it is contained in $\overline{\mathcal{A}}$.*

Proof. Let \mathcal{A} be any \mathcal{C}^k atlas for M . Define

$$\overline{\mathcal{A}} := \{\varphi \text{ chart for } M \text{ that is } \mathcal{C}^k \text{ compatible with all charts } \theta \in \mathcal{A}\}.$$

Clearly $\overline{\mathcal{A}}$ contains \mathcal{A} . We claim that $\overline{\mathcal{A}}$ is a \mathcal{C}^k atlas. To prove this we have to show that if $\varphi, \psi \in \overline{\mathcal{A}}$ are charts with respective domains $U, V \subseteq M$, then the transition map $\varphi \circ \psi^{-1} : \psi(U \cap V) \rightarrow \varphi(U \cap V)$ is \mathcal{C}^k . Take any point $\psi(p) \in \psi(U \cap V)$ and let $\theta \in \mathcal{A}$ be a chart whose domain W contains the point $p \in U \cap V$. Then $\psi(U \cap V \cap W)$ is an open neighborhood of $\psi(p)$ and we can write the restriction⁶

$$\varphi \circ \psi^{-1} : \psi(U \cap V \cap W) \rightarrow \varphi(U \cap V \cap W)$$

as the composition $(\varphi \circ \theta^{-1}) \circ (\theta \circ \psi^{-1})$, which is \mathcal{C}^k because $\varphi \circ \theta^{-1}$ and $\theta \circ \psi^{-1}$ are \mathcal{C}^k by assumption. This proves that $\varphi \circ \psi^{-1}$ is \mathcal{C}^k in a neighborhood of $\psi(p)$, but the same reasoning is valid at any point of $\psi(U \cap V)$, therefore $\varphi \circ \psi^{-1}$ is \mathcal{C}^k .

Finally, from the definition of $\overline{\mathcal{A}}$ it is clear that it is maximal, and that any atlas \mathcal{B} is equivalent to \mathcal{A} if and only if it is contained in $\overline{\mathcal{A}}$. In particular, any atlas \mathcal{B} containing \mathcal{A} is equivalent to \mathcal{A} (by definition of equivalent atlases), and is therefore contained in $\overline{\mathcal{A}}$. Therefore a maximal atlas containing \mathcal{A} is contained in $\overline{\mathcal{A}}$, but in fact it must be equal to $\overline{\mathcal{A}}$ (by maximality). We conclude that $\overline{\mathcal{A}}$ is the unique maximal \mathcal{C}^k atlas containing \mathcal{A} . \square

In consequence, given a topological manifold M and some \mathcal{C}^k atlas \mathcal{A} on M we can speak without ambiguity of *the* \mathcal{C}^k structure $\overline{\mathcal{A}}$ determined by \mathcal{A} .

Remark 1.2.5. For practical purposes the concept of a maximal \mathcal{C}^k atlas is not really important. We usually work with a smaller \mathcal{C}^k atlas and this is all we need e.g. for checking that a function is \mathcal{C}^k (see next section). (In fact, we could have defined a \mathcal{C}^k structure on M as an equivalence class of \mathcal{C}^k atlases, rather than as a maximal \mathcal{C}^k atlas.) In general we won't give any name to the maximal atlas and we'll just speak about “a \mathcal{C}^k manifold M ” with the maximal \mathcal{C}^k atlas \mathcal{A} being implicit.

Also, when we say “a \mathcal{C}^k chart” or simply “a chart” of M , we mean a chart $\varphi \in \mathcal{A}$. In the rare case that we may need a \mathcal{C}^l chart φ with $l \leq k$ (which means φ is only \mathcal{C}^l compatible with the charts of \mathcal{A}), we will say it explicitly. In particular, a “topological chart” is a \mathcal{C}^0 chart, i.e. a homeomorphism.

⁶see Remark 1.2.3

Remark 1.2.6. Every \mathcal{C}^k manifold is automatically a \mathcal{C}^l manifold for every $l \leq k$, because every \mathcal{C}^k atlas is a \mathcal{C}^l atlas. In the other direction, Whitney (*Differentiable Manifolds*, 1936) proved that every \mathcal{C}^k structure contains a (non unique!) \mathcal{C}^l structure for any $l > k$. The proof is reproduced in Munkres' *Elementary Differential Topology* and in Hirsch's *Differential Topology*.

Example 1.2.7 (Examples of smooth manifolds).

1. \mathbb{R}^n (with the atlas consisting of the single chart $\text{id}_{\mathbb{R}^n}$) is a smooth manifold. In general, any topological manifold endowed with a single-chart atlas is automatically a smooth manifold. For example, the graph Gra_f of any *continuous* (sic) function $f : U \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$ as described in Example 1.1.11, endowed with the projection chart, is a smooth manifold.
2. Any open subset U of a \mathcal{C}^k manifold M has a natural \mathcal{C}^k structure consisting of the \mathcal{C}^k charts of M whose domain is contained in U . (Exercise.) We will also see in the exercises that finite products of \mathcal{C}^k manifolds have a natural \mathcal{C}^k structure.
3. The sphere $\mathbb{S}^n \subset \mathbb{R}^{n+1}$ is a smooth manifold. Indeed, the atlas given by the graph coordinates (Example 1.1.11) is smooth. To see this, we compute the transition functions (wlog $i < j$):

$$\begin{aligned} \varphi_i^+ \circ (\varphi_j^\pm)^{-1} : \varphi_j^\pm(U_j^\pm \cap U_i^+ \cap \mathbb{S}^n) &\rightarrow \varphi_i^\pm(U_j^\pm \cap U_i^+ \cap \mathbb{S}^n) \\ (y_0, \dots, y_{n-1}) &\mapsto (y_0, \dots, \hat{y}_i, \dots, y_{j-1}, \pm \sqrt{1 - \sum_i (y_i)^2}, y_j, \dots, y_{n-1}) \end{aligned}$$

and a similar formula works if we replace φ_i^+ by φ_i^- . Hence all the transition maps are smooth.

4. More generally, any subset M of \mathbb{R}^k given as the *regular level set* of a smooth map $F : \mathbb{R}^k \rightarrow \mathbb{R}^\ell$ is a $k - \ell$ dimensional smooth manifold “in a natural way”⁷. (Being a level set means $M = F^{-1}(\{c\})$ for some $c \in \mathbb{R}^\ell$ and being a regular level set means that, moreover, the Jacobian $D|_p F$ is surjective for all $p \in M$.) You can prove this quite easily using the implicit function theorem and writing M locally as a graph of smooth functions (analogous to the graph coordinates for the sphere). We will show a more general statement later on when discussing submanifolds (Chapter 3).
5. Projective space \mathbb{P}^n is naturally a smooth manifold; see exercises.
6. The torus $\mathbb{T}^n = \mathbb{R}^n / \mathbb{Z}^n$ is naturally a smooth manifold. (Exercise.)
7. On the topological 1-manifold $M = \{(x, y) \in \mathbb{R}^2 \mid y = x^3\}$, each of the two projections $\pi_0, \pi_1 : M \rightarrow \mathbb{R}$ (given by $\pi_0(x, y) = x$ and $\pi_1(x, y) = y$) is a chart defined on the whole manifold M , but these two charts are not \mathcal{C}^k compatible for any $k \geq 1$. (The transition functions are $\pi_1 \circ (\pi_0)^{-1} : x \mapsto x^3$ and its inverse $\pi_0 \circ (\pi_1)^{-1} : y \mapsto \sqrt[3]{y}$, which is not differentiable.) Thus these charts determine two different \mathcal{C}^k structures on M .

1.2.1 One-step \mathcal{C}^k manifolds*

Here it is convenient to work with local parametrizations rather than charts. Recall that an (n -dimensional) **local parametrization** of a topological space M is an homeomorphism from an open subset U of \mathbb{R}^n to an open subset of M or, equivalently, an open injective map $U \rightarrow M$. A family \mathcal{A} of such maps whose images cover M is called a **parametrization atlas** or **inverse atlas** for M , and such an atlas is \mathcal{C}^k if the transition map $\psi^{-1} \circ \phi$ is \mathcal{C}^k for each $\phi, \psi \in \mathcal{A}$. Equivalently, a local parametrization is the inverse of a chart, and \mathcal{A} is a \mathcal{C}^k parametrization atlas iff $\mathcal{A}^{-1} := \{\varphi^{-1} : \varphi \in \mathcal{A}\}$ is a \mathcal{C}^k atlas.

⁷In particular, the inclusion map $M \hookrightarrow \mathbb{R}^k$ is smooth.

Note that if M is a \mathcal{C}^k manifold, then the topology of M is determined by the \mathcal{C}^k structure. Indeed, if \mathcal{A} is a \mathcal{C}^k parametrization atlas, then

$$\text{a set } U \subseteq M \text{ is open iff } \phi^{-1}(U) \text{ is open in } \text{Dom}(\phi) \subseteq \mathbb{R}^n \text{ for all } \phi \in \mathcal{A}. \quad (1.1)$$

In other words, the topology of M is the *final topology* induced by the family \mathcal{A} .⁸

We can ask then, given a set M and a collection \mathcal{A} of functions $\phi : U_\phi \subseteq \mathbb{R}^n \rightarrow M$, whether there exists a topology on M such that \mathcal{A} is a \mathcal{C}^k parametrization atlas of M . Such a topology is necessarily unique by (1.1). This provides a way to define a \mathcal{C}^k manifold without constructing first a topological space.

Proposition 1.2.8 (One-step manifolds). *Consider a set M endowed with a family \mathcal{A} of injections $\phi : U_\phi \subseteq \mathbb{R}^n \rightarrow M$ satisfying the the following properties:*

(Denote $\tilde{U}_\phi = \phi(U_\phi)$ the image of each $\phi \in \mathcal{A}$.)

- (1) For all $\phi, \psi \in \mathcal{A}$, the set $U_\phi^\psi := \phi^{-1}(\tilde{U}_\psi)$ is open in \mathbb{R}^n ,*
- (2) and the transition map $\psi^{-1} \circ \phi : U_\phi^\psi \rightarrow U_\psi^\phi$ is \mathcal{C}^k .*
- (3) Countably many of the sets \tilde{U}_ϕ cover M .*
- (4) For every two distinct points $p, q \in M$ there exist maps $\phi, \psi \in \mathcal{A}$ containing p, q in their respective images, and open neighborhoods $V \subseteq U_\phi, W \subseteq U_\psi$ of $\phi^{-1}(p), \psi^{-1}(q)$ respectively, such that $\phi(V) \cap \psi(W) = \emptyset$.*

Then the set M endowed with the final topology w.r.t. \mathcal{A} is a topological n -manifold admitting \mathcal{A} as a \mathcal{C}^k parametrization atlas.

Remarks:

- The hypothesis implies that each set $U_\phi \subseteq \mathbb{R}^n$ is open, because it is equal to U_ϕ^ϕ .
- Each transition map $\psi^{-1} \circ \phi : U_\phi^\psi \rightarrow U_\psi^\phi$ is a \mathcal{C}^k isomorphism because it is \mathcal{C}^k and bijective, with inverse $\phi^{-1} \circ \psi$, which is also \mathcal{C}^k .

Proof. We endow M with the final topology w.r.t. \mathcal{A} , so that a set $U \subseteq M$ is open if and only if $\psi_\alpha^{-1}(U)$ is open in U_α for all α . This is clearly a topology on M .

We claim that each map $\phi \in \mathcal{A}$ is open. Indeed, if $W \subseteq U_\phi$ is an open set, then $\phi(W)$ is open in M because for any $\psi \in \mathcal{A}$, the set $\psi^{-1}(\phi(W)) = (\psi^{-1} \circ \phi)(W \cap U_\phi^\psi)$ is open in U_ψ^ϕ since $\psi^{-1} \circ \phi$ is a homeomorphism $U_\phi^\psi \rightarrow U_\psi^\phi$.

It follows that each $\phi \in \mathcal{A}$ is a homeomorphism onto its image, because it is open and injective. Therefore the maps ϕ are n -dimensional local parametrizations of M ensuring that M is locally Euclidean of dimension n . Moreover, M is a topological manifold because it is second countable by (3) and Hausdorff by (4). Finally, the parametrization atlas \mathcal{A} is \mathcal{C}^k -consistent by (2). \square

An example of manifold that can be constructed in this way is the Grassman manifold $G_k(V)$ of k -subspaces of $V \simeq \mathbb{R}^n$; see [Lee13, p. 1.36].

1.3 Differentiable maps

We are about to define \mathcal{C}^k maps between \mathcal{C}^k manifolds. The plan is to reduce the question of differentiability to the case of a map between Euclidean open sets. We will do so by using charts.

In general, when studying a map $f : M \rightarrow N$ between manifolds, charts allow us to *locally* express f as a map between subsets of Euclidean space.

⁸The **final topology** on a set M induced by a family of maps $\phi : U_\phi \rightarrow M$, where U_ϕ are topological spaces, is the topology defined by the formula (1.1). Exercise: check that it is indeed a topology.

Definition 1.3.1 (Local expression of a map). Let M, N be \mathcal{C}^k manifolds and let $f : M \rightarrow N$ be any function (not necessarily continuous). A **local expression** (or **coordinate representation**) of f at some point $p \in M$ is a composite map

$$f|_{\varphi}^{\psi} := \psi \circ f \circ \varphi^{-1},$$

where φ and ψ are charts of M and N whose domains U, V contain the points p and $f(p)$ respectively. The composite is defined at all possible points; see Remark 1.2.3.

Local expressions allow us, for instance, to determine whether a function is continuous or not.

Remark 1.3.2. Exercise: Show that f is continuous at p if and only if the local expression $f|_{\varphi}^{\psi}$ at p is defined in a neighborhood of $\varphi(p)$ and is continuous at $\varphi(p)$. This fact holds however we choose the charts φ, ψ (provided their domains contain the points $p, f(p)$ respectively, of course).

The idea is to use local expressions to *define* whether a map is \mathcal{C}^k or not.

Definition 1.3.3 (\mathcal{C}^k maps between manifolds). We say a function $f : M \rightarrow N$ between \mathcal{C}^k manifolds is \mathcal{C}^k at a point $p \in M$ if there exists a local expression $f|_{\varphi}^{\psi}$ of f at p such that

$$f|_{\varphi}^{\psi} \text{ is defined in a neighborhood of } \varphi(p) \text{ and is } \mathcal{C}^k \text{ at the point } \varphi(p). \quad (1.2)$$

If this holds for all points $p \in M$, we say that f is a \mathcal{C}^k map. The set of \mathcal{C}^k maps $M \rightarrow N$ is denoted $\mathcal{C}^k(M, N)$. A \mathcal{C}^k **isomorphism** is a \mathcal{C}^k map that has a \mathcal{C}^k inverse.

If $k \geq 1$, a \mathcal{C}^k map is also called \mathcal{C}^k -**differentiable**, and a \mathcal{C}^k isomorphism is also called a \mathcal{C}^k **diffeomorphism** (or a \mathcal{C}^k **diffeo**, for short).

In fact the condition (1.2), that determines whether f is \mathcal{C}^k or not, does not depend on how we choose the charts φ, ψ .

Proposition 1.3.4. *If f is \mathcal{C}^k at the point p , then every local expression $f|_{\varphi}^{\psi}$ of f at p satisfies (1.2).*

Proof. Assume that the local expression $f|_{\varphi}^{\psi}$ satisfies (1.2). This implies that f is continuous at p , by Remark 1.3.2. Consider a second local expression $f|_{\tilde{\varphi}}^{\tilde{\psi}}$ of f at p . This new local expression is defined on some neighborhood of $\tilde{\varphi}(p)$ (again by Remark 1.3.2), and is related to the old one by the **chart-change formula**

$$f|_{\tilde{\varphi}}^{\tilde{\psi}} \equiv (\psi \circ \tilde{\psi}^{-1}) \circ f|_{\varphi}^{\psi} \circ (\varphi \circ \tilde{\varphi}^{-1}), \quad (1.3)$$

which holds at all points where the two expressions are defined (in particular, in some neighborhood of $\tilde{\varphi}(p)$). Since the transition maps $(\psi \circ \tilde{\psi}^{-1})$ and $(\varphi^{-1} \circ \tilde{\varphi})$ are \mathcal{C}^k , we conclude that $f|_{\tilde{\varphi}}^{\tilde{\psi}}$ is \mathcal{C}^k at the point $\tilde{\varphi}(p)$. \square

Example 1.3.5. 1. The identity map of any \mathcal{C}^k manifold is a \mathcal{C}^k map. (Exercise.)
 2. A composite map $g \circ f$ is \mathcal{C}^k at a point p if f is \mathcal{C}^k at p and g is \mathcal{C}^k at $f(p)$. (Exercise.)
 3. If M is a \mathcal{C}^k manifold, then every \mathcal{C}^k chart of M , as well as its inverse, are \mathcal{C}^k maps. (Exercise.)

A \mathcal{C}^k structure on a topological manifold M allows us to determine which maps that go to M are \mathcal{C}^k . But the reciprocal property also holds: if we know which maps to M are \mathcal{C}^k , this information determines the \mathcal{C}^k structure of M .

Proposition 1.3.6. *Let $\mathcal{A}_0, \mathcal{A}_1$ be two \mathcal{C}^k atlases on a topological manifold M , defining two \mathcal{C}^k manifolds $M_i = (M, \overline{\mathcal{A}_i})$. Then the two atlases \mathcal{A}_i are equivalent if and only if the following property holds:*

For every function $f : N \rightarrow M$ (where N is a \mathcal{C}^k manifold), the function f is \mathcal{C}^k as a map $N \rightarrow M_0$ if and only if it is \mathcal{C}^k as a map $N \rightarrow M_1$.

Proof. Exercise. □

1.4 Partitions of unity

In this section we develop *partitions of unity*, a tool often used to turn a local construction, obtained by working in coordinates, into a global one. (Don't worry if this sounds vague; we will see examples later on.) Existence of partitions of unity on a manifold is easier to prove on a compact manifold; we will deal first with this case. The general proof relies on a topological property of manifolds called *paracompactness*, which is in turn a consequence of being Hausdorff, second countable and locally compact.

Recall that the **support** of a continuous function $\eta : M \rightarrow \mathbb{R}$ is the closed set

$$\text{supp}(\eta) := \overline{\{p \in M \mid \eta(p) \neq 0\}}.$$

Definition 1.4.1. A \mathcal{C}^k **partition of unity** (or **POU**, for short) on a \mathcal{C}^k manifold M is a family $(\eta_i)_i$ of \mathcal{C}^k functions $\eta_i : M \rightarrow [0, +\infty)$ satisfying

- Sum condition: $\sum_i \eta_i(x) = 1$ for every $x \in M$,
- Local finiteness: Every point of M has a neighborhood where all except finitely many of the functions η_i vanish.

A partition of unity $(\eta_i)_i$ is **subordinate** to an open cover \mathcal{U} of M if every η_i has its support contained in some open set $U \in \mathcal{U}$.

Theorem 1.4.2 (Existence of Partitions of Unity). *For any open cover \mathcal{U} of a \mathcal{C}^k manifold M there exists a partition of unity $(\eta_i)_i$ subordinate to \mathcal{U} such that the functions η_i have compact support (in fact, their supports are closed coordinate balls; see definition below).*

Another version of the theorem is the following.

Corollary 1.4.3 (Existence of Partitions of Unity, alternate form). *For any open cover $\mathcal{U} = \{U_j\}_{j \in J}$ of a \mathcal{C}^k manifold M there exists a partition of unity $(\xi_j)_{j \in J}$ such that $\text{supp}(\xi_j) \subseteq U_j$ for all j .*

This version of the theorem can be deduced from the first one (exercise). Note that the functions g_j may not have compact support in this case.

For the proof of Theorem 1.4.2 we will use *bump functions*.

Lemma 1.4.4 (Bump functions on Euclidean space). *For any numbers $0 < a < b$ there exists a smooth **bump function** $h : \mathbb{R}^n \rightarrow [0, 1]$ satisfying $h(x) = 1$ iff $\|x\| \leq a$ and $h(x) = 0$ iff $\|x\| > b$.*

Proof. It suffices to let $h(x) = g(\|x\|)$, where $g : \mathbb{R} \rightarrow [0, 1]$ is a smooth **cutoff function** satisfying $g(t) = 1$ iff $t < a$ and $g(t) = 0$ iff $t > b$. This cutoff function, in turn, may be defined as

$$g(t) = \frac{f(b-t)}{f(b-t) + f(t-a)},$$

where $f : \mathbb{R} \rightarrow [0, +\infty)$ is a smooth function such that $f(t) > 0$ iff $t > 0$. Such a function f may be given e.g. by the formula

$$f(t) = \begin{cases} e^{-\frac{1}{t}} & t > 0 \\ 0 & t \leq 0 \end{cases}.$$

□

We may use charts to transport these bump functions from Euclidean space to any manifold. The resulting bump function will be supported on a closed *coordinate ball*.

Definition 1.4.5. A **closed coordinate ball** in a \mathcal{C}^k manifold M is the preimage $\varphi^{-1}(\overline{B})$ of a closed Euclidean ball \overline{B} by a \mathcal{C}^k chart φ containing \overline{B} in its codomain.

Proof of Thm. 1.4.2 for compact manifolds. Let M be a compact \mathcal{C}^k manifold and let \mathcal{U} be an open cover of M . In this case we'll obtain a *finite* \mathcal{C}^k partition of unity $(\eta_i)_i$ subordinate to \mathcal{U} . In fact, it is sufficient to find finitely many \mathcal{C}^k functions $\tilde{\eta}_i : M \rightarrow [0, +\infty)$, each of which has its support $\text{supp}(\tilde{\eta}_i)$ contained in some $U \in \mathcal{U}$, and such that $\sum_i \tilde{\eta}_i(x) > 0$ for all x . The functions η_i can then be obtained by dividing each $\tilde{\eta}_i$ by the strictly positive \mathcal{C}^k function $\tilde{\eta} = \sum_i \tilde{\eta}_i$.

To construct the functions η_i we proceed as follows. Each point of a set $U \in \mathcal{U}$ is contained in the interior of some closed coordinate ball $D \subseteq U$. Thus there is a family of closed coordinate balls, each of them contained in some $U \in \mathcal{U}$, whose interiors cover M . By compactness, we may take a finite subfamily of balls D_i whose interiors still cover M . Write each ball D_i as $\varphi_i^{-1}(\overline{B}_i)$, where B_i is an open Euclidean ball and φ_i is a \mathcal{C}^k chart containing \overline{B}_i in its codomain. Then let $h_i : \mathbb{R}^n \rightarrow [0, +\infty)$ be a \mathcal{C}^k bump function that is supported on the closed ball \overline{B}_i and strictly positive on the interior B_i . Finally, define a \mathcal{C}^k function $\tilde{\eta}_i : M \rightarrow [0, 1]$ by the formula

$$\tilde{\eta}_i = \begin{cases} h_i \circ \varphi_i & \text{on } \text{Dom}(\varphi_i) \\ 0 & \text{on } M \setminus D_i \end{cases}$$

This function is supported on D_i , which is contained in some $U \in \mathcal{U}$, thus the function family $(\tilde{\eta}_i)$ is subordinate to \mathcal{U} . In addition, $\tilde{\eta}_i$ is strictly positive on B_i , and since the balls B_i cover M , we conclude that $\sum_i \tilde{\eta}_i(x) > 0$ for all $x \in M$, as required. □

1.4.1 Paracompactness*

Definition 1.4.6 (Paracompact space). Let X be a topological space.

- An **open cover** of X is a family \mathcal{U} of open sets whose union is X .
- Another open cover \mathcal{V} **refines** \mathcal{U} if every $V \in \mathcal{V}$ is contained in some $U \in \mathcal{U}$.
- The space X is **paracompact** if every open cover \mathcal{U} admits as refinement some open cover \mathcal{V} that is locally finite.
- (A family of subsets of X is **locally finite** if every point $x \in X$ has a neighborhood that intersects only finitely many sets of the family.)

Proposition 1.4.7. *If a topological space X is Hausdorff, second countable and locally compact, then it is paracompact.*

In fact, given any topological base \mathcal{B} of X , every open cover \mathcal{U} has a locally finite refinement \mathcal{V} consisting of open sets $B \in \mathcal{B}$ with their closures \overline{B} contained in some $U \in \mathcal{U}$.

Proof.

Lemma 1.4.8. *X admits an exhaustion by compact sets, i.e. a sequence $(K_i)_{i \in \mathbb{N}}$ of compact sets that cover X and satisfy $K_i \subseteq \text{Int}(K_{i+1})$.*

Proof of lemma. Let $(V_j)_{j \in \mathbb{N}}$ be a countable open cover of M where each V_j has compact closure. Let $K_0 = \emptyset$, and define inductively for each $i \in \mathbb{N}$ an integer $j_i \geq i$ such that the sets $(V_j)_{j < j_i}$ cover K_i , and a compact set $K_{i+1} = \overline{V_{j_i}} \cup \bigcup_{j < j_i} \overline{V_j}$. These compact sets K_i cover M and satisfy $K_i \subseteq \text{Int}(K_{i+1})$. \square

Consider the compact sets $L_i = K_i \setminus \text{Int}(K_{i-1})$ and their respective open neighborhoods $W_i = \text{Int } K_{i+1} \setminus K_{i-2}$.

Each point $x \in L_i$ is contained in some open $U_x \in \mathcal{U}$ and has a basic neighborhood $B_x^i \in \mathcal{B}$ such that $\overline{B_x^i} \subseteq W_i \cap U$.

Take a finite subfamily $(B_{x_j}^i)_j$ that covers L_i . Doing this for each i we obtain a family of basics $(B_{x_j}^i)_{i,j}$ that cover M . Their closures satisfy $\overline{B_{x_j}^i} \subseteq U_{x_j}$, which gives the subordination condition, and $\overline{B_{x_j}^i} \subseteq W_i$, which ensures local finiteness since every point $x \in L_i$ is contained in at most three sets W_ℓ , namely, those with $|\ell - i| \leq 1$. \square

Using paracompactness, we can prove existence of partitions of unity without the hypothesis of compactness.

Proof of Thm. 1.4.2. Let M be a manifold and \mathcal{U} an open cover.

Let \mathcal{B} be the topological base of M consisting of the interiors of closed coordinate balls. By Proposition 1.4.7, there exists a family of closed coordinate balls D_i , each contained in some open set $U \in \mathcal{U}$, whose interiors cover M .

The proof finishes as in the compact case. We take for each i a \mathcal{C}^k function $\tilde{\eta}_i : M \rightarrow [0, 1]$ that is strictly positive on $\text{Int}(D_i)$ and supported on D_i . Then the functions $\eta_i = \frac{\tilde{\eta}_i}{\sum_j \tilde{\eta}_j}$ form a \mathcal{C}^k partition of unity $(\eta_i)_i$ that is subordinate to the open cover \mathcal{U} . \square

1.4.2 Applications

Corollary 1.4.9 (Bump functions). *If M is a \mathcal{C}^k manifold, $A \subset M$ a closed set and $U \subseteq M$ an open neighborhood of A , then there exists a \mathcal{C}^k function $\eta : M \rightarrow [0, 1]$ such that $\eta \equiv 1$ on A and $\text{supp}(\eta) \subset U$.*

We call η a *bump function* for A supported in U .

Proof. Just take the open cover $\{V_0 = U, V_1 = M \setminus A\}$ of M and a partition of unity $(\eta_i)_{i=0,1}$ satisfying $\text{supp } \eta_i \subseteq V_i$, then set $\eta = \eta_0$. \square

We have already defined differentiability for function $M \rightarrow N$ between \mathcal{C}^k manifolds. For functions on closed sets we make the following definition:

Definition 1.4.10. Let $f : A \rightarrow N$ be a function where $A \subseteq M$ is a closed set and M, N are \mathcal{C}^k manifolds. We say that f is \mathcal{C}^k if it can be extended to a \mathcal{C}^k function defined on an open neighborhood of A .

As a corollary of the existence of bump functions we can extend a smooth function on a closed set to a smooth function on the whole manifold:

Corollary 1.4.11 (Extension lemma). *Let $f : A \rightarrow \mathbb{R}$ be \mathcal{C}^k , where $A \subseteq M$ is a closed subset of a \mathcal{C}^k manifold M , and let $U \subseteq M$ be an open set containing A . Then there exists a \mathcal{C}^k function $\tilde{f} : M \rightarrow \mathbb{R}$ such that $\tilde{f}|_A = f$ and $\text{supp } \tilde{f} \subseteq U$.*

Proof. By definition f can be extended to a \mathcal{C}^k function (say, also called f) on some open set $W \supseteq A$; wlog $W \subseteq U$. We take a \mathcal{C}^k bump function η for A supported in W . Then ηf has support in W and therefore extending the function by 0 outside W we obtain a \mathcal{C}^k function \tilde{f} on M with the desired properties. \square

2 Tangent vectors

Recall that if a map $f : U \rightarrow V$ between Euclidean open sets $U \subseteq \mathbb{R}^m$, $V \subseteq \mathbb{R}^n$ is \mathcal{C}^1 at a point $p \in U$, then there exists a unique linear transformation $D_p f : \mathbb{R}^m \rightarrow \mathbb{R}^n$, called the **differential transformation** of f at p , which gives a first-order approximation

$$f(p + v) = f(p) + D_p f(v) + r_p(v)$$

where $\frac{r_p(v)}{\|v\|} \rightarrow 0$ as $v \rightarrow 0$.

To define the differential of a map between \mathcal{C}^k manifolds, we need a notion of tangent space.

Definition 2.0.1. Let M be an n -dimensional \mathcal{C}^k manifold with $k \geq 1$. A **co-ordinatized tangent vector** on M is a triple (p, φ, v) where $p \in M$ is a point, φ is a \mathcal{C}^k chart of M defined at p , and $v \in \mathbb{R}^n$ is a vector in Euclidean space. A **tangent vector** on M is the equivalence class $[p, \varphi, v]$ of a coordinatized tangent vector (p, φ, v) under the equivalence relation

$$(p, \varphi, v) \sim (\tilde{p}, \tilde{\varphi}, \tilde{v}) \iff \tilde{p} = p \quad \text{and} \quad \tilde{v} = D_{\varphi(p)}(\tilde{\varphi} \varphi^{-1})(v)$$

The set TM of tangent vectors is the **tangent bundle** of M , and there is a canonic projection map $\pi_{TM} : TM \rightarrow M$ sending $[p, \varphi, v] \mapsto p$.

The **tangent space** at a point $p \in M$ is the set $T_p M := \pi^{-1}(p)$. It is a vector space with vector addition

$$[p, \varphi, v] + [p, \varphi, w] := [p, \varphi, v + w]$$

and vector scaling

$$\lambda [p, \varphi, v] := [p, \varphi, \lambda v] \quad \text{for } \lambda \in \mathbb{R}.$$

Remark 2.0.2. 1. \sim is indeed an equivalence relation. (Exercise.)

2. Fixed a point $p \in M$ and a \mathcal{C}^k chart φ defined on p , the function $\iota : \mathbb{R}^n \rightarrow T_p M$ sending $v \mapsto [p, \varphi, v]$ is a bijection. (Exercise.)

3. A tangent vector $X = [p, \varphi, v] \in T_p M$ can be considered as a function

$$\begin{aligned} \{\text{charts of } M \text{ def. at } p\} &\rightarrow \mathbb{R}^n \\ \psi &\mapsto D_{\varphi(p)}(\psi \varphi^{-1})(v). \end{aligned}$$

The vector $D_{\varphi(p)}(\psi \varphi^{-1})(v)$ is the only $w \in \mathbb{R}^n$ such that $[p, \psi, w] = X$.

4. Vector addition and scaling are well defined and make $T_p M$ a vector space isomorphic to \mathbb{R}^n . (Exercise.)

Remark 2.0.3. If $U \subseteq \mathbb{R}^n$ is an open set (considered as a smooth manifold), we identify $TU \equiv U \times \mathbb{R}^n$ by the bijection

$$(p, v) \in U \times \mathbb{R}^n \mapsto [p, \text{id}_U, v] \in TU.$$

Thus for each $p \in U$ we have $T_p U \equiv \{p\} \times \mathbb{R}^n \equiv \mathbb{R}^n$, and for $p \in U$ and $v \in \mathbb{R}^n$ we write

$$v|_p := [p, \text{id}_U, v] \in T_p U. \tag{2.1}$$

Coordinate base for the tangent space We can construct a base of the tangent space $T_p M$ as follows. Consider the canonic base $(e_i)_i$ of \mathbb{R}^n , and take a chart φ defined at p . Then the vectors

$$\left. \frac{\partial}{\partial \varphi^i} \right|_p := [p, \varphi, e_i],$$

called the **coordinate vectors** at p associated to the chart φ , form a base of $T_p M$.

If ψ is another chart defined at p , then this chart determines a second base consisting of vectors $\left. \frac{\partial}{\partial \psi^j} \right|_p$. This second base is related to the first one by the formula

$$\left. \frac{\partial}{\partial \varphi^i} \right|_p = \sum_j \left. \frac{\partial \psi^j}{\partial \varphi^i} \right|_{\varphi(p)} \left. \frac{\partial}{\partial \psi^j} \right|_p.$$

where $\left. \frac{\partial \psi^j}{\partial \varphi^i} \right|_{\varphi(p)}$ is the partial derivative of $\psi \circ \varphi^{-1}$ that appears as the coefficient (j, i) of the matrix expression of the linear map $D_{\varphi(p)}(\psi \circ \varphi^{-1}) : \mathbb{R}^n \rightarrow \mathbb{R}^n$. To see this we compute as follows:

$$\begin{aligned} \left. \frac{\partial}{\partial \varphi^i} \right|_p &= [p, \varphi, e_i] = [p, \psi, D_{\varphi(p)}(\psi \circ \varphi^{-1})e_i] \\ &= \left[p, \psi, \sum_j \left. \frac{\partial \psi^j}{\partial \varphi^i} \right|_{\varphi(p)} e_j \right] \\ &= \sum_j \left. \frac{\partial \psi^j}{\partial \varphi^i} \right|_{\varphi(p)} [p, \psi, e_j] \\ &= \sum_j \left. \frac{\partial \psi^j}{\partial \varphi^i} \right|_{\varphi(p)} \left. \frac{\partial}{\partial \psi^j} \right|_p. \end{aligned}$$

Remark 2.0.4. In concrete examples it is common to use more intuitive symbols for φ^i , e.g. the polar coordinates (r, φ) or the spherical coordinates (r, ϕ, θ) . The standard coordinates (x^0, \dots, x^{n-1}) on \mathbb{R}^n are usually written (x, y) for $n = 2$ and (x, y, z) for $n = 3$. Bear in mind that an expression like (r, φ) can mean either the map (chart) or the coordinates of a particular point; see example below.

Example 2.0.5 (Polar coordinates). Let $W := \mathbb{R}^+ \times (0, 2\pi)$. The map

$$\Psi : W \rightarrow \mathbb{R}^2 : (r, \varphi) \mapsto (r \cos \varphi, r \sin \varphi)$$

is a diffeomorphism onto its image $U := \Psi(W) = \mathbb{R}^2 \setminus (\mathbb{R}_{\geq 0} \times \{0\})$. Its inverse $\Psi^{-1} : U \rightarrow W$ is therefore a smooth chart for \mathbb{R}^2 . The components of Ψ^{-1} are usually written (r, φ) and called *polar coordinates*. On the other hand, we have the standard coordinates (x, y) (i.e. the identity map) on \mathbb{R}^2 . Take a point $p = (x, y) \in U$ and let $(r, \varphi) = \Psi^{-1}(x, y)$ be its polar coordinates. The polar coordinate vectors $\left. \frac{\partial}{\partial r} \right|_p, \left. \frac{\partial}{\partial \varphi} \right|_p$ can be expressed as a linear combination of the standard coordinate vectors $\left. \frac{\partial}{\partial x} \right|_p, \left. \frac{\partial}{\partial y} \right|_p$, using the change of coordinates formula:

$$\begin{aligned} \frac{\partial}{\partial r} &= \frac{\partial x}{\partial r} \frac{\partial}{\partial x} + \frac{\partial y}{\partial r} \frac{\partial}{\partial y} = \cos(\varphi) \frac{\partial}{\partial x} + \sin(\varphi) \frac{\partial}{\partial y} \\ \frac{\partial}{\partial \varphi} &= \frac{\partial x}{\partial \varphi} \frac{\partial}{\partial x} + \frac{\partial y}{\partial \varphi} \frac{\partial}{\partial y} = -r \sin(\varphi) \frac{\partial}{\partial x} + r \cos(\varphi) \frac{\partial}{\partial y}. \end{aligned}$$

Here we do certain standard abuses of notation. First, the letters r, φ, x, y represent functions when preceded by a ∂ , otherwise they are numbers (obtained by evaluating these functions at the point p). Second, we have omitted the “ $|_p$ ” on the vectors and the evaluation at (r, φ) for the partial derivatives.

2.0.1 Differential of a \mathcal{C}^k map between manifolds

Now that we have tangent spaces, we can define the differential of a \mathcal{C}^k map.

Definition 2.0.6. The **differential transformation** (or **differential**, for short) of a map $f : M \rightarrow N$ that is \mathcal{C}^k at a point $p \in M$ is the linear operator

$$\begin{aligned} D_p f : T_p M &\rightarrow T_{f(p)} N \\ [p, \varphi, v] &\mapsto [f(p), \psi, D_{\varphi(p)} f|_{\varphi}^{\psi}(v)]. \end{aligned}$$

where φ, ψ are charts of M, N defined at the points $p, f(p)$ respectively, and $f|_{\varphi}^{\psi} = \psi \circ f \circ \varphi^{-1}$ is the local expression of f with respect to the charts φ, ψ .

If f is \mathcal{C}^k everywhere, the union of the maps $D_p f$ for all $p \in M$ is a map $Df : TM \rightarrow TN$, also denoted f_* and called the **pushforward** by f .

Note that $D_p f$ is well defined (independent of φ, ψ) and linear. To see that it is well defined, we compute twice

$$\begin{aligned} D_p f : X = [p, \varphi, v] &\mapsto Y = [f(p), \psi, w = D_{\varphi(p)} f|_{\varphi}^{\psi}(v)]. \\ D_p f : \tilde{X} = [p, \tilde{\varphi}, \tilde{v}] &\mapsto \tilde{Y} = [f(p), \tilde{\psi}, \tilde{w} = D_{\tilde{\varphi}(p)} f|_{\tilde{\varphi}}^{\tilde{\psi}}(\tilde{v})]. \end{aligned}$$

and verify that $X = \tilde{X}$ implies $Y = \tilde{Y}$. That is, we must check that $\tilde{v} = D_{\varphi(p)}(\tilde{\varphi}\varphi^{-1})(v)$ (or, equivalently, $v = D_{\tilde{\varphi}(p)}(\varphi\tilde{\varphi}^{-1})(\tilde{v})$) implies $\tilde{w} = D_{\psi(p)}(\tilde{\psi}\psi^{-1})(w)$. This follows from the equation $f|_{\tilde{\varphi}}^{\tilde{\psi}} = (\tilde{\psi}\psi^{-1})f|_{\varphi}^{\psi}(\varphi\tilde{\varphi}^{-1})$ by the chain rule (in its Euclidean version).

The linearity of $D_p f$ follows from the linearity of $D_{\varphi(p)} f|_{\varphi}^{\psi}$.

The chain rule has the following version for maps between manifolds.

Proposition 2.0.7 (Chain rule). *If $f : M \rightarrow N$ is \mathcal{C}^k at some point p and $g : N \rightarrow L$ is \mathcal{C}^k at $f(p)$, then $g \circ f$ is \mathcal{C}^k at p and has differential*

$$D_p(g \circ f) = D_{f(p)}g \circ D_p f.$$

In particular, for a diffeo f , the differential $D_p f$ is a linear isomorphism whose inverse is $D_{f(p)}(f^{-1})$. (Exercise.)

Example 2.0.8 (Derivative of a chart). Let M be an n -dimensional \mathcal{C}^k -manifold, let $p \in M$, and let $\varphi : U \rightarrow V$ be a chart of M defined at p . Then we have

$$D\varphi\left(\frac{\partial}{\partial\varphi^i}\Big|_p\right) = e_i|_{\phi(p)} \in T_{\phi(p)}\mathbb{R}^n.$$

To see this, using the local expression $\varphi|_{\varphi}^{\text{id}_V} = \text{id}_V \circ \varphi \circ \varphi^{-1} = \text{id}_V$ we compute

$$\begin{aligned} D\varphi\left(\frac{\partial}{\partial\varphi^i}\Big|_p\right) &= D\varphi[p, \varphi, e_i] \\ &= [\varphi(p), \text{id}_V, D_{\varphi(p)}(\varphi|_{\varphi}^{\text{id}_V})(e_i)] = [\varphi(p), \text{id}_V, e_i] = e_i|_p \in T_p\mathbb{R}^n. \end{aligned}$$

Example 2.0.9 (Velocity of a curve). For a differentiable curve $\gamma : I \subseteq \mathbb{R} \rightarrow M$ on a manifold M , we define its **velocity vector** at an instant $t \in I$ as the vector $\gamma'(t) := D_t\gamma(1|_t) \in T_{\gamma(t)}M$ where $1|_t$ represents the element $[t, \text{id}_I, 1]$ of $T_t I$ according to the identification $TI \cong I \times \mathbb{R}$ given in Remark 2.0.3.

Exercise: Show that for any vector $X \in TM$ there is a curve $\gamma : (-\varepsilon, \varepsilon) \rightarrow M$ such that $\gamma'(0) = X$.

Tangent vectors as derivations

Definition 2.0.10. A **derivation** on a C^k differentiable manifold M at a point $p \in M$ is a linear function $D : C^k(M, \mathbb{R}) \rightarrow \mathbb{R}$ satisfying the Leibniz identity

$$D(fg) = D(f)g(p) + f(p)D(g).$$

The set $Der_p M$ of derivations on M at p is a vector space with the operations

$$\begin{aligned}(D + E)(f) &:= D(f) + E(f), \\ (\lambda D)(f) &:= \lambda(D(f))\end{aligned}$$

defined for $D, E \in Der_p M$ and $\lambda \in \mathbb{R}$.

Each vector $X \in T_p M$ induces a derivation $D_X \in Der_p M$ defined by the formula

$$D_X(f) := D_p f(X) \in T_{f(p)} \mathbb{R} \equiv \mathbb{R}.$$

(Here we use the identification of Remark 2.0.3).

Proposition 2.0.11. *The map $\nu_p : X \in T_p M \mapsto D_X \in Der_p M$ is a linear injection. (Exercise.)*

Therefore we may identify a tangent vector X with the derivation D_X and write $X(f) := D_X(f)$. In fact, the map ν is bijective if M is smooth (see e.g. [Lee13, Prop. 3.2]). Therefore in some books (e.g. [Lee13]), the tangent space $T_p M$ of a smooth manifold M is *defined* as the vector space of derivations at p . Here we do not use that definition because it does not work for non-smooth manifolds (i.e. C^k manifolds with $k < \infty$).

Tangent vectors as derivations

In this chapter we give an alternative definition of the “tangent space” using derivations. This second definition is equivalent to the one given in the previous chapter, but is more similar to the one used in most books (e.g. Lee’s book [Lee13]).

In more detail, for a point p of a differentiable manifold M , we will consider first a vector space $\text{Der}_p M$ of *derivation operators* (or *derivations*, for short) and then we will define the tangent space $T_p M$ as a certain subspace of $\text{Der}_p M$. Tangent vectors are the derivations that correspond (via a chart) to a derivation defined by a vector of \mathbb{R}^n . Equivalently, tangent vectors are those derivations that can be expressed as the velocity vector of a curve.

The objects that we will define are analogous to those defined in the previous chapter (except that the differential of a function will be called “tangent map”). The plan is the following:

- (1) For each \mathcal{C}^k -differentiable n -manifold M and each point $p \in M$, define a vector space $T_p M \simeq \mathbb{R}^n$, called the *tangent space* of M at p .
- (2) For each \mathcal{C}^k map $f : M \rightarrow N$ and each point $p \in M$, define a linear map $T_p f : T_p M \rightarrow T_p N$ called the *tangent map* of f at p .

If the manifolds M, N are open subsets of $\mathbb{R}^m, \mathbb{R}^n$ respectively, then the map $T_p f$ will be essentially equivalent to the **differential transformation** studied in calculus courses, that is, the unique linear map $D_p f : \mathbb{R}^m \rightarrow \mathbb{R}^n$ which gives the first-order approximation

$$f(p + v) = f(p) + D_p f(v) + r(v)$$

where $\frac{r(v)}{\|v\|} \rightarrow 0$ as $v \rightarrow 0$.

Motivation for definition of tangent vectors Let M be a \mathcal{C}^k manifold M with $k \geq 1$ and let $p \in M$. We want to have a vector space $T_p M$ of vectors tangent to M at p . How can we define the notion of a “tangent vector”?

For example, let $\gamma : I \subseteq \mathbb{R} \rightarrow M$ be a differentiable curve such that $\gamma(0) = p$. This curve should have a “velocity vector” X at the instant $t = 0$, which should be vector tangent to M at p . This object X should tell us the *direction and speed* of movement of $\gamma(t)$ at time $t = 0$. How can we measure this “direction and speed”?

Idea: take a function $h \in \mathcal{C}^k(M, \mathbb{R})$. The composite $h \circ \gamma$ is a function $I \rightarrow \mathbb{R}$ that is differentiable at $t = 0$, and the number $(h \circ \gamma)'(0)$ tells us something about how γ is moving at $t = 0$. If we know this number for all functions h , let us agree that we know the velocity vector X completely. Therefore we propose to *define* the velocity vector X simply as the function $h \mapsto (h \circ \gamma)'(0)$. This function has certain algebraic properties that make it a *derivation*.

Definition 2.0.12. A **derivation operator** (or **derivation**, for short) on a \mathcal{C}^k -differentiable manifold M at a point $p \in M$ is a function $X : \mathcal{C}^k(M, \mathbb{R}) \rightarrow \mathbb{R}$ that is \mathbb{R} -linear, that is, such that

$$X(a \cdot g + b \cdot h) = a \cdot X(g) + b \cdot X(h) \quad \text{for } g, h \in \mathcal{C}^k(M, \mathbb{R}) \text{ and } a, b \in \mathbb{R},$$

and satisfies the **Leibniz identity**

$$X(g \cdot h) = X(g) \cdot h(p) + g(p) \cdot X(h). \tag{2.2}$$

The set $\text{Der}_p(M)$ of derivations on M at p is a real vector space: for $X, Y \in \text{Der}_p M$ and $a, b \in \mathbb{R}$ we define the derivation¹ $aX + bY \in \text{Der}_p M$ by the formula

$$(a \cdot X + b \cdot Y)(h) := a \cdot X(h) + b \cdot Y(h).$$

We also define the set $\text{Der } M := \coprod_{p \in M} \text{Der}_p M$.

Before getting to the derivations that we are interested in (namely, those we'll call "tangent vectors"), let us prove some properties of general derivations.

Proposition 2.0.13. *Consider any derivation $X \in \text{Der}_p M$ at a point p of a \mathcal{C}^k -differentiable manifold M . Then for all functions $g, h \in \mathcal{C}^k(M, \mathbb{R})$ we have:*

- (a) *If h is constant, then $X(h) = 0$.*
- (b) *If both g and h vanish at p (i.e. $g(p) = h(p) = 0$), then $X(g \cdot h) = 0$.*
- (c) *Locality: If $g \equiv h$ on a neighborhood of p , then $X(g) = X(h)$.*

Proof. To prove (a), since X is linear, it suffices to show that $X(h) = 0$ if $h \equiv 1$. And indeed, in this case we have

$$X(h) = X(h^2) = X(h) \cdot h(p) + h(p) \cdot X(h) = 2X(h)$$

which implies $X(h) = 0$.

Fact (b) follows from the Leibniz identity.

To prove (c), since X is linear, it suffices to show that $X(f) = 0$ if $f \equiv 0$ in some open neighborhood U of p . Let $\eta : \mathcal{C}^k(M, \mathbb{R})$ be a bump function that is constantly 1 on the closed set $M \setminus U$ and whose support is contained in the open set $M \setminus \{p\}$. Note that $f \cdot \eta = f$, therefore $X(f) = X(f \cdot \eta) = 0$ since $f(p) = \eta(p) = 0$. \square

One first example of derivation is the one we already mentioned: velocity vectors.

Definition 2.0.14. The **velocity vector** of a curve $\gamma : I \subseteq \mathbb{R} \rightarrow M$ at an instant $t \in I$ (where γ is differentiable) is the derivation $\text{Vel}_\gamma(t) \in \text{Der}_{\gamma(t)} M$ defined by

$$\begin{aligned} \text{Vel}_\gamma(t) : \mathcal{C}^k(M, \mathbb{R}) &\rightarrow \mathbb{R} \\ h &\mapsto (h \circ \gamma)'(t). \end{aligned}$$

Let us check that the function $X = \text{Vel}_\gamma(t_0)$ is indeed a derivation on M at the point $p = \gamma(t_0)$. For a function $h \in \mathcal{C}^k(M, \mathbb{R})$ we denote $h_\gamma := h \circ \gamma \in \mathcal{C}^k(I, \mathbb{R})$. The fact that X satisfies the Leibniz identity for functions $g, h \in \mathcal{C}^k(M, \mathbb{R})$ follows from the Leibniz identity for the single-variable functions $g_\gamma, h_\gamma : I \rightarrow \mathbb{R}$ (that you already know from calculus):

$$\begin{aligned} X(g \cdot h) &= (g \cdot h)'_\gamma(t_0) = (g_\gamma \cdot h_\gamma)'(t_0) = g'_\gamma(t_0) \cdot h_\gamma(t_0) + g_\gamma(t_0) \cdot h'_\gamma(t_0) \\ &= X(g) \cdot h(p) + g(p) \cdot X(h) \end{aligned}$$

The fact that X is \mathbb{R} -linear can be proven in a similar way (exercise).

Remark 2.0.15. The velocity vector $\text{Vel}_\gamma(t)$ is sometimes denoted $\gamma'(t)$, but it is different from the usual $\gamma'(t)$ defined for a curve in \mathbb{R}^n , because it also contains the information of the point $\gamma(t)$ where the derivation is located.

Exercise: If γ, β are differentiable curves in \mathbb{R}^n , show that $\text{Vel}_\gamma(t) = \text{Vel}_\beta(t)$ if and only if $\gamma(t) = \beta(t)$ and $\gamma'(t) = \beta'(t)$.

¹Exercise: check that $a \cdot X + b \cdot Y$ is indeed a derivation and convince yourself that these operations make $\text{Der}_p M$ a vector space.

Another kind of derivations are *vectorial derivations* (also known as *directional derivations*²). Before considering general manifolds, let us recall how these derivation operators are defined on an open subset of \mathbb{R}^n .

Definition 2.0.16. A **vectorial derivation** at a point p of an open set $U \subseteq \mathbb{R}^n$ is a derivation $\delta_{p,v} \in \text{Der}_p U$ determined by a vector $v \in \mathbb{R}^n$ by the formula

$$\begin{aligned} \delta_{p,v} : \mathcal{C}^k(U, \mathbb{R}) &\rightarrow \mathbb{R} \\ h &\mapsto D_p h(v), \end{aligned} \quad (2.3)$$

where $D_p h : \mathbb{R}^n \rightarrow \mathbb{R}$ is the differential operator of h at the point p .

Note that $\delta_{p,v}$ is a derivation; in fact, it is the velocity vector at time $t = 0$ of any curve $\gamma : (-\varepsilon, \varepsilon) \rightarrow U$ satisfying $\gamma'(0) = p$ and $\gamma'(0) = v$. (For example, the curve $\gamma(t) = p + tv$.) Indeed, if γ is such a curve, then for any function $h \in \mathcal{C}^k(U, \mathbb{R})$ we have

$$\delta_{p,v}(h) = D_p h(v) = D_p h(\gamma'(0)) = (h \circ \gamma)'(0) = \text{Vel}_\gamma(0)(h).$$

The definition of vectorial derivation can be adapted to a general manifold by using a chart.

Definition 2.0.17. A **vectorial derivation** on a \mathcal{C}^k differentiable n -manifold M at a point $p \in M$ is a derivation of the form

$$\begin{aligned} \delta_{p,\phi,v} : \mathcal{C}^k(M, \mathbb{R}) &\rightarrow \mathbb{R} \\ h &\mapsto D_{\phi(p)}(h \circ \phi^{-1})(v) \end{aligned} \quad (2.4)$$

where ϕ is a chart defined at p and $v \in \mathbb{R}^n$. The triple (p, ϕ, v) is called a **coordinate expression** of $\delta_{p,\phi,v}$.

A **tangent vector** to M at p is the same thing as a vectorial derivation on M at p .

The **tangent space** of M at p is the set $T_p M \subseteq \text{Der}_p M$ of vectors tangent to M at p . (We will show that $T_p M$ is in fact an n -dimensional subspace of $\text{Der}_p M$.)

The **tangent bundle** of M is the set $TM := \coprod_{p \in M} T_p M$.

Exercise 2.0.18. Prove that tangent vectors are derivations indeed. In fact, tangent vectors are the same thing as velocity vectors. That is, every tangent vector is the velocity vector of some curve, and every velocity vector can also be expressed as a vectorial derivation.

Remark 2.0.19 (Domain of tangent vectors). Although the “official” domain of a tangent vector $X \in T_p M$ is the set $\mathcal{C}^k(M, \mathbb{R})$, in fact we can apply X to any real-valued h that is defined on a neighborhood U of p and is differentiable at p . The number $X(h)$ is defined using the same formula (2.4) that we would use for a function $h \in \mathcal{C}^k(M, \mathbb{R})$.

Remark 2.0.20 (Locality of tangent vectors). It is also clear from the definition that the number $X(h)$ depends only on the values of h near p . Thus if two functions g, h coincide on a neighborhood of p , then $X(g) = X(h)$. (We have already shown in Proposition 2.0.13 that this is a property of general derivations, but in the case of vectorial derivations it is especially evident.)

²I don't like saying “directional” because it suggests that the derivation depends only on the direction of the vector, and not on its size.

Let us show that the tangent space $T_p M$ of an n -dimensional differentiable manifold M at any point $p \in M$ is an n -dimensional vector space, as promised. We defined $T_p M$ as a subset of $\text{Der}_p M$, which is itself a vector space (with the obvious operations that we already defined). Thus we need just show that $T_p M$ is an n -dimensional subspace of $\text{Der}_p M$, and we will do so by showing that $T_p M$ is the image of an injective linear transformation $\mathbb{R}^n \rightarrow \text{Der}_p M$.

Proposition 2.0.21. *Let M be a C^k -differentiable manifold and let $p \in M$. Then for any chart ϕ of M defined at p , the map*

$$\begin{aligned} \iota_{p,\phi} : \mathbb{R}^n &\rightarrow \text{Der}_p M \\ v &\mapsto \delta_{p,\phi,v} \end{aligned}$$

is a linear isomorphism onto its image, which is the tangent space $T_p M$. In consequence, $T_p M$ is an n -dimensional subspace of $\text{Der}_p M$.

In other words, *any single* chart ϕ defined at p allows us to express *all* tangent vectors $X \in T_p M$ in the form $X = \delta_{p,\phi,v}$, with the vector $v \in \mathbb{R}^n$ being unique for each X .

Proof. To show that $\iota_{p,\phi}$ is an isomorphism we develop some formulas that will be useful elsewhere.

Lemma 2.0.22. *Any tangent vector $X = \delta_{p,\phi,v} \in T_p M$ applied to a component function ϕ^i of the chart ϕ gives*

$$X(\phi^i) = v^i. \quad (2.5)$$

In consequence, $\iota_{p,\phi}$ is injective.

Proof of lemma. Note first that $\phi^i = \pi^i \circ \phi$, where $\pi^i : \mathbb{R}^n \rightarrow \mathbb{R}$ is the projection on the i -th coordinate, defined by $\pi^i(x) = x^i$. Using this, we can compute

$$X(\phi^i) = D_p(\phi^i \circ \phi^{-1})(v) = D_p(\pi^i)(v) = \pi^i(v) = v^i.$$

Here we used the fact that $D_p(\pi^i) = \pi^i$ since π^i is linear.

Note that the functions ϕ^i are not in the “official” domain $\mathcal{C}^k(M, \mathbb{R})$ of the operator $\delta_{p,\phi,v}$, because they are defined on $U \subseteq M$ (the domain of ϕ) rather than on the whole manifold M . However, we can construct functions $\tilde{\phi}^i \in \mathcal{C}^k(M, \mathbb{R})$ that coincide with ϕ^i on a neighborhood of p . (For example, using bump functions; recall Lemma 1.4.11.) Using the functions $\tilde{\phi}^i$ instead of ϕ^i we obtain an analogous formula $X(\tilde{\phi}^i) = v^i$. This formula allows us to recover the components of the vector $v \in \mathbb{R}^n$ from the derivation $\delta_{p,\phi,v}$. It follows that the map $\iota_{p,\phi} : v \mapsto \delta_{p,\phi,v}$ is injective. \square

Lemma 2.0.23. *For any two charts ϕ, ψ defined at p , and for any two vectors $v, w \in \mathbb{R}^n$, we have*

$$\begin{aligned} \delta_{p,\phi,v} = \delta_{p,\psi,w} &\iff w = D_{\phi(p)}(\psi \circ \phi^{-1})(v) \\ &\iff v = D_{\psi(p)}(\phi \circ \psi^{-1})(w) \end{aligned} \quad (2.6)$$

In consequence, the map $\iota_{p,\phi}$ is surjective onto $T_p M$.

Proof of lemma. Note first that the two equations on the right are equivalent since the linear transformations $D_{\phi(p)}(\psi \circ \phi^{-1})$ and $D_{\psi(p)}(\phi \circ \psi^{-1})$ are inverse of each other.

Suppose first that these equations hold. Then for any function $h \in \mathcal{C}^k(M, \mathbb{R})$ we have

$$\begin{aligned}\delta_{p,\psi,w}(h) &= D_{\psi(p)}(h \circ \psi^{-1})(w) \\ &= D_{\psi(p)}(h \circ \psi^{-1})(D_{\phi(p)}(\psi \circ \phi^{-1})(v)) \\ &= D_{\phi(p)}(h \circ \phi^{-1})(v) \\ &= \delta_{p,\phi,v}(h).\end{aligned}$$

It follows that $\iota_{p,\phi}$ is surjective onto $T_p M$, since any vector $X \in T_p M$ is of the form $X = \delta_{p,\psi,w}$ for some chart ψ and some vector $w \in \mathbb{R}^n$, and we can obtain $\iota_{p,\phi}(v) = X$ by putting $v = D_{\psi(p)}(\phi \circ \psi^{-1})(w)$.

Finally, suppose that $\delta_{p,\phi,v} = \delta_{p,\psi,w}$. We have already shown that this equation holds when $v = D_{\psi(p)}(\phi \circ \psi^{-1})(w)$, and it cannot hold for any other value of v because the map $\iota_{p,\phi} : v \mapsto \delta_{p,\phi,v}$ is injective by Lemma 2.0.22. We conclude that $v = D_{\psi(p)}(\phi \circ \psi^{-1})(w)$. \square

This finishes the proof that $\iota_{p,\phi}$ is an isomorphism, and hence $T_p M \simeq \mathbb{R}^n$. \square

Remark 2.0.24. Lemma 2.0.23 shows that a tangent vector $X \in T_p M$ can be thought of as an equivalence class of triples (p, ϕ, v) (where ϕ is a chart of M and $v \in \mathbb{R}^n$) under the equivalence relation

$$\begin{aligned}(p, \phi, v) \sim (p, \psi, w) &\iff w = D_{\phi(p)}(\psi \circ \phi^{-1})(v) \\ &\iff v = D_{\psi(p)}(\phi \circ \psi^{-1})(w)\end{aligned}$$

This is another way to define tangent vectors, equivalent to ours.

Example 2.0.25 (Tangent bundle of an open subset of \mathbb{R}^n). Recall that any open set $U \subseteq \mathbb{R}^n$ is naturally a \mathcal{C}^k differentiable manifold with an atlas consisting of the single chart id_U . This single chart is sufficient to express all tangent vectors, by Proposition 2.0.21. That is, all tangent vectors $X \in TU$ are of the form $X = \delta_{p,\text{id}_U,v}$, with $p \in U$ and $v \in \mathbb{R}^n$. Thus there is a natural identification³

$$\begin{aligned}TU &\equiv U \times \mathbb{R}^n \\ \delta_{p,\text{id}_U,v} &\leftrightarrow (p, v).\end{aligned}$$

Note also that the operator $\delta_{p,\text{id}_U,v}$ coincides with the operator $\delta_{p,v}$ of Definition 2.0.17, therefore the two definitions of “vectorial derivation” (2.0.17 and 2.0.16) are equivalent for an open set $U \subseteq \mathbb{R}^n$.

2.0.2 Pushforward and tangent operators

Any differentiable map between manifolds induces a map on derivations called the **pushforward operator**.

Definition 2.0.26 (Pushforward of derivations). The **pushforward operator** of a \mathcal{C}^k -differentiable map $f : M \rightarrow N$ at some point $p \in M$ is the linear map

$$\begin{aligned}f_*|_p : \text{Der}_p M &\rightarrow \text{Der}_{f(p)} N \\ X &\mapsto (h \in \mathcal{C}^k(N, \mathbb{R}) \mapsto X(h \circ f) \in \mathbb{R}).\end{aligned}$$

The union of the operators $f_*|_p$ over all points $p \in M$ is the **pushforward map**

$$f_* := \coprod_{p \in M} f_*|_p : \text{Der } M \rightarrow \text{Der } N.$$

³We write $A \equiv B$ when there is a natural isomorphism between two objects A, B .

Example 2.0.27. Each derivation $\delta_{p,\phi,v}$ of Definition 2.0.17 is the image of the derivation $\delta_{\phi(p),v}$ of Definition 2.0.16 by the pushforward operator of the map ϕ^{-1} at the point $\phi(p)$. (Check it!)

The **tangent operator** of a differentiable map is obtained by restricting the pushforward operator to tangent vectors.

Definition 2.0.28. The **tangent operator** of a \mathcal{C}^k map $f : M \rightarrow N$ at a point $p \in M$ is the map

$$T_p f : T_p M \rightarrow T_{f(p)} N$$

obtained by restricting the pushforward map $f_*|_p : \text{Der}_p M \rightarrow \text{Der}_{f(p)} N$.

The **tangent map** of f is the union of the tangent operators $T_p f$ over all points $p \in M$,

$$Tf := \coprod_{p \in M} T_p f : TM \rightarrow TN.$$

To show that the tangent operator $T_p f$ is well defined, we have to verify that the pushforward operator $f_*|_p$ indeed maps tangent vectors to tangent vectors (or, equivalently, it maps velocity vectors to velocity vectors).

Proposition 2.0.29 (Local expression of the tangent operator). *If $f : M \rightarrow N$ is differentiable at a point $p \in M$, then for any tangent vector $\delta_{p,\phi,v} \in T_p M$ and any chart ψ of N defined at $f(p)$ we have*

$$f_*|_p(\delta_{p,\phi,v}) = \delta_{\phi(p),\psi,w} \quad \text{where} \quad w = D_{\phi(p)}(f|_{\phi}^{\psi})(v). \quad (2.7)$$

Proof. For any $h \in \mathcal{C}^k(N, \mathbb{R})$ we have

$$\begin{aligned} (f_*|_p \delta_{p,\phi,v})(h) &= \delta_{p,\phi,v}(hf) \\ &= D_{\phi(p)}(hf\phi^{-1})(v) \\ &= D_{\psi(f(p))}(h\psi^{-1})(D_{\phi(p)}(\psi f \phi^{-1})(v)) \\ &= \delta_{f(p),\psi,w}(h). \end{aligned}$$

□

Proposition 2.0.30 (Pushforward of velocity vectors; exercise). *Let $f : M \rightarrow N$ be a \mathcal{C}^k -differentiable map, and let $\gamma : I \subseteq \mathbb{R} \rightarrow M$ be a curve that is differentiable at some instant $t_0 \in I$. Then $f_*(\text{Vel}_{\gamma}(t_0)) = \text{Vel}_{f \circ \gamma}(t_0)$.*

Remark 2.0.31. The tangent transformation $T_p f : T_p M \rightarrow T_{f(p)} N$ can in fact be defined for any function $f : U \subseteq M \rightarrow N$ that is defined on a neighborhood U of p and is differentiable at p . This follows from Remarks 2.0.19 and 2.0.20 (check it!).

Example 2.0.32. For a \mathcal{C}^k differentiable map $f : U \rightarrow V$, where $U \subseteq \mathbb{R}^m$, $V \subseteq \mathbb{R}^n$ are open sets, the tangent operator $T_p f : T_p U \rightarrow T_p V$ is essentially equivalent to the differential $D_p f : \mathbb{R}^m \rightarrow \mathbb{R}^n$ because for all vectors $X \in T_p U$ (which can be written as $X = \delta_{p,v}$ with $v \in \mathbb{R}^m$) we have

$$T_p f(X) = T_p f(\delta_{p,v}) = \delta_{p,w} \quad \text{where} \quad w = D_p f(v).$$

The chain rule that you know from calculus also holds for maps between manifolds.

Proposition 2.0.33 (Chain rule; exercise). *Let $f : M \rightarrow N$ and $g : N \rightarrow L$ be maps between differentiable manifolds.*

1. If f and g are differentiable at points $p \in M$, $f(p) \in N$ respectively, then the composite map $g \circ f$ is differentiable at p , and

$$T_p(g \circ f) = T_{f(p)}g \circ T_p f. \quad (2.8)$$

2. For any point $p \in M$ we have $T_p \text{id}_M = \text{id}_{T_p M}$.
3. If f is a diffeomorphism, then $T_p f$ is a linear isomorphism with inverse $T_{f(p)}(f^{-1})$.

2.0.3 Classical notation for tangent vectors

Here we could do the same as in the last chapter.

Given a chart ϕ of M at p , define a basis of $T_p M$ consisting of vectors $\frac{\partial}{\partial \phi^i} \Big|_p := \delta_{p, \phi, e_i}$.

Note that for any function $f \in \mathcal{C}^k(M, \mathbb{R})$ $\frac{\partial}{\partial \phi^i} \Big|_p f = \frac{\partial}{\partial x^i} \Big|_{\phi(p)} f|_{\phi(x)}$.

The vector $\frac{\partial}{\partial \phi^i} \Big|_p$ is the velocity vector at $t = 0$ of a curve $\gamma_i(t)$ that satisfies $\phi^j(\gamma_i(t)) = \phi^j(p) + \delta_i^j t$. That is, along this curve $\gamma_i(t)$ the i -th coordinate increases at unit speed while the other coordinates remain constant.

One can prove that $\frac{\partial}{\partial \phi^i} \Big|_p (\phi^j) = \delta_i^j$ and also give the coordinate change formula.

2.0.4 General derivations*

If M is a smooth manifold, then the space $\text{Der}_p M$ contains no other derivations than the tangent vectors (see exercises). Thus one can define the tangent space $T_p M$ of a smooth manifold M simply as the space $\text{Der}_p M$ of derivations. However, if M is a \mathcal{C}^k manifold with $k < \infty$, then the space $\text{Der}_p M$ is infinite dimensional (see exercises).

3 Submanifolds

Let us start with a short motivation. The “bent line” in \mathbb{R}^2 , defined as $C = \{(x, y) \in \mathbb{R}^2 \mid xy = 0, x \geq 0, y \geq 0\}$, is intuitively not “smoothly contained” in \mathbb{R}^2 . It is topological submanifold of \mathbb{R}^2 because, when endowed with the subspace topology, it is homeomorphic to \mathbb{R} . Moreover, it follows that C can be given the structure of a smooth manifold (using a single-chart atlas), but the smooth structure does not “agree” with the ambient one; the bent line is not a *smooth submanifold* of \mathbb{R}^2 . To make this precise we need a few definitions:

Definition 3.0.1. Let $f : M \rightarrow N$ be a \mathcal{C}^k -differentiable map between manifolds of respective dimensions m, n .

We say that f is **submersive** at a point $p \in M$ if $T_p f$ is surjective. If this holds for all points $p \in M$, then f is a **submersion**.

We say that f is **immersive** at a point $p \in M$ if $T_p f$ is injective. If this holds for all points $p \in M$, then f is an **immersion**.

We say that f is an **embedding** if it is an immersion and also a topological embedding (i.e. a homeomorphism onto its image, the latter being endowed with the subspace topology). Its image is called a \mathcal{C}^k **embedded submanifold** of N (or a \mathcal{C}^k **embedded m -submanifold**, to highlight its dimension).

The **rank** of f at a point $p \in M$, denoted $\text{rank}_p f$, is the rank of the tangent operator $T_p f$. If f has the same rank k at all points (“constant rank”) then we write $\text{rank } f = k$.

So to sum up, a \mathcal{C}^k map $f : M \rightarrow N$ is a submersion iff $\text{rank } f = n = \dim N$ and is an immersion iff $\text{rank } f = m = \dim M$. In both cases the tangent operator $T_p f$ has maximal rank for all points $p \in M$.¹ Note that the set of points where f has maximal rank is open because the rank is an upper semicontinuous function: if $k = \text{rank}_p f$, then there is a neighborhood of p where f has $\text{rank} \geq k$.

Example 3.0.2. The **standard immersion** $\iota_k^n : \mathbb{R}^k \rightarrow \mathbb{R}^n$, defined for $k \leq n$, is the map

$$\iota_k^n : (x^0, \dots, x^{k-1}) \mapsto (x^0, \dots, x^{k-1}, 0, \dots, 0).$$

This is also a smooth embedding.

The **standard submersion** $\pi_m^k : \mathbb{R}^m \rightarrow \mathbb{R}^k$, defined for $m \geq k$, is the map

$$\pi_m^k : (x^0, \dots, x^{m-1}) \mapsto (x^0, \dots, x^{k-1}).$$

Composing these we obtain the **standard map of rank k**

$$\iota_k^n \circ \pi_m^k : \mathbb{R}^m \rightarrow \mathbb{R}^n : (x^0, \dots, x^{m-1}) \mapsto (x^0, \dots, x^{k-1}, 0, \dots, 0).$$

Example 3.0.3. • A smooth curve $\gamma : J \rightarrow M$, where $J \subset \mathbb{R}$ is an open interval and M is a smooth manifold, is an immersion iff $\gamma'(t) \neq 0$ for all $t \in J$. For example, the curve on \mathbb{R}^2 given by $t \mapsto (\cos t, \sin t)$, $t \in \mathbb{R}$, is an immersion, but the curve on \mathbb{R}^2 given by $t \mapsto (t^2, t^3)$, $t \in \mathbb{R}$, is not.

¹A linear map $V \simeq \mathbb{R}^m \rightarrow W \simeq \mathbb{R}^n$ has **maximal rank** if its rank is equal to $\min\{m, n\}$, which is the maximum rank that a linear map $V \rightarrow W$ can have.

- The *bent line* $C = \{(x, y) \in \mathbb{R}^2 \mid xy = 0, x, y \geq 0\}$ is not an embedded submanifold. In fact, it is not the image of an immersion. (Exercise.)
- The inclusion $\mathbb{S}^n \hookrightarrow \mathbb{R}^{n+1}$ is a smooth embedding (where we endow \mathbb{S}^n and \mathbb{R}^{n+1} with the standard smooth structures).
- Let $M = M_0 \times M_1$ be a product of smooth manifolds. Then the canonical projections $\pi^i : M \rightarrow M_i$ are submersions.
- “The figure 8”: Let $J = (-\frac{\pi}{2}, \frac{3\pi}{2})$ and consider the curve

$$\gamma : J \rightarrow \mathbb{R}^2, \quad \gamma(t) = (\sin(2t), \cos t).$$

It is easy to verify that γ is an injective immersion. However, it is not a homeomorphism onto its image $N := \gamma(J) \subset \mathbb{R}^2$ (endowed with the subspace topology): removing e.g. the point $(0, 1) = \gamma(0)$ the set $N \setminus \{(0, 1)\}$ is connected, but its preimage under γ is $(-\frac{\pi}{2}, 0) \cup (0, \frac{3\pi}{2})$ hence disconnected. Therefore γ is not an embedding.

- *Irrational line on the torus*: Consider the curve in the torus $f : \mathbb{R} \rightarrow \mathbb{T}^2$ defined by $f(t) = [t, \alpha t]$, where $\alpha \in \mathbb{R}$ is an irrational number. This map is an injective immersion, but is not an embedding since one can find a divergent sequence of numbers $t_i \in \mathbb{R}$ such that $f(t_i)$ converges to some point in the image of f . (See [Lee13, Example 4.20] for more details.)

Example 3.0.4 (Graphs as submanifolds). The **graph** of a \mathcal{C}^k map $f : M \rightarrow N$, defined as the set $\text{Gra}_f = \{(x, f(x)) : x \in M\}$, is a submanifold of $M \times N$. Indeed, it is the image of the **graphing map** of f , i.e. the map $g : M \rightarrow M \times N$ defined by $g(x) = (x, f(x))$, which is an embedding.

To see that g is an immersion we note that for any point $p \in M$, the tangent operator $T_p g : T_p M \rightarrow T_{p, f(p)}(M \times N) \equiv T_p M \times T_{f(p)} N$ maps every nonzero vector $v \in T_p M$ to the vector $T_p g(v) = (v, T_p f(v))$, which is nonzero because its first component is nonzero.

To see that the map g is a topological embedding we note that it admits a continuous retraction, namely, the projection map $\pi : M \times N \rightarrow M : (x, y) \mapsto x$. (A **retraction** of g is a left inverse, i.e. a map ρ satisfying $\rho \circ g = \text{id}$.) Equivalently, the map $g|_{\text{Gra}_f} : M \rightarrow \text{Gra}_f$ is a homeomorphism because it admits as inverse the continuous map $\pi|_{\text{Gra}_f} : \text{Gra}_f \rightarrow M$.

3.1 How to recognize an embedding

Suppose we are given a differentiable map $f : M \rightarrow N$ between manifolds, and we have already checked that it is an injective immersion. How can we prove that it is an embedding? This is essentially a topological question: how can we show that a given injective map² is actually a topological embedding? Here are some ways to do so:

Proposition 3.1.1. *An injective \mathcal{C}^r immersion $f : M \rightarrow N$ is an embedding in any of the following cases:*

- f is an open map.*
- $\dim M = \dim N$.*
- f is a closed map.*
- The domain M is compact.*
- f is a proper map.*

²A **map** between topological spaces is a continuous function.

(f) The image $f(M)$ has an open neighborhood W such that the map $f|_W : M \rightarrow W$ is closed or proper.

Proof. (a) In this case f sends open subsets of M to open subsets of N , which are clearly also open in the image $f(M)$. Thus f is an open (and bijective) map onto its image, hence an homeomorphism onto its image.

(b) In this case f is a local diffeomorphism by the IFT, hence an open map.

(c) Similar to (a), replacing “open” by “closed”.

(d) A map from a compact space to a Hausdorff space is closed.

(e) A proper map is closed if its codomain is Hausdorff and locally compact; see e.g. [Lee13, A.57] or [Lee10, p. 4.95].

(f) In this case f is the composite of two embeddings: the closed embedding $f|_W$, followed by the inclusion map $\iota_W : W \rightarrow M$, which is an open embedding. \square

Remark 3.1.2. The last case (f) is interesting because it gives a general criterion: for every embedding $f : M \rightarrow N$ there exists an open neighborhood W of $\text{Img}(f)$ such that $f|_W : M \rightarrow W$ is closed. (Exercise.)

A different, arguably more direct way to check that a \mathcal{C}^k map $f : M \rightarrow N$ is an embedding is by constructing a \mathcal{C}^k **retraction** (a.k.a left inverse) of f , that is, a \mathcal{C}^k map $g : N \rightarrow M$ satisfying $g \circ f = \text{id}_M$. In fact, the retraction need not be defined everywhere; an open neighborhood of the image suffices.

Proposition 3.1.3. *Let $f : M \rightarrow N$ be a \mathcal{C}^k map. Then f is an embedding if it admits a \mathcal{C}^k **neighborhood retraction**, that is, a \mathcal{C}^k map $g : U \rightarrow M$, where $U \subseteq N$ is an open neighborhood of the image $f(M)$, such that $g \circ f = \text{id}_M$.*

Proof. Suppose $f : M \rightarrow N$ has a neighborhood retraction $g : U \rightarrow M$. Then f is clearly injective. Its tangent operator $T_p f$ at any point $p \in M$ is injective as well because

$$T_p \text{id}_M = T_p(g \circ f) = T_{f(p)}g \circ T_p f$$

is an isomorphism. Therefore f is an immersion. Finally, f is a homeomorphism onto its image $f(M)$ because it admits as inverse the continuous map $g|_{f(M)}$. \square

Note that this is the argument we used in Example 3.0.4 to prove that the graph of a \mathcal{C}^r map $f : M \rightarrow N$ is a \mathcal{C}^r submanifold of $M \times N$. The graphing map $g : M \rightarrow M \times N : x \mapsto (x, f(x))$ admits as a retraction the projection $\pi : M \times N \rightarrow M : (x, y) \mapsto x$.

Remark 3.1.4. This method for showing that a map is a submanifold is also general in the sense that every embedding $f : M \rightarrow N$ admits a neighborhood retraction (defined on a so-called *tubular neighborhood* of the submanifold $f(M)$). We won’t prove that, but we’ll show how to produce local neighborhood retractions, which are sufficient to prove that a given subset of a manifold is an embedded submanifold (see Propositions 3.3.1 and 3.4.1 below).

3.2 Constant rank theorem

Many theorems about submanifolds are based on the inverse function theorem.

Theorem 3.2.1 (Inverse Function Theorem, or IFT). *Let $f : M \rightarrow N$ be a \mathcal{C}^k map, where M, N are n -manifolds, and let $p \in M$ be a point where the tangent operator $T_p f$ is invertible. Then f is a **local diffeomorphism** at p , i.e. there exists respective open neighborhoods U, V of $p, f(p)$ such that $f|_U^V$ is a \mathcal{C}^k diffeomorphism.*

In case that M, N are open subsets of \mathbb{R}^n , this is just the inverse function theorem that you know from calculus. The general case can be deduced by using charts. (Exercise.)

A useful generalization (and corollary) of the IFT is the *constant rank theorem*.

Theorem 3.2.2 (Constant Rank Theorem, or CRT). *Let $f : M \rightarrow N$ be \mathcal{C}^r map that has rank k at some point $p \in M$, and let $m = \dim M$ and $n = \dim N$. Suppose, further that f has constant rank k in a neighborhood of p . Then f admits at the point p a local expression $\tilde{f} = f|_{\tilde{\phi}}$ of the form*

$$\begin{aligned} \tilde{f} : \tilde{U} \subseteq \mathbb{R}^m &\rightarrow \tilde{V} \subseteq \mathbb{R}^n \\ x &\mapsto (x^0, \dots, x^{k-1}, 0, \dots, 0). \end{aligned}$$

where $\phi : U \rightarrow \tilde{U}$, $\psi : V \rightarrow \tilde{V}$ are charts of M, N defined and centered at p , $f(p)$ respectively. (This means that $\phi(p) = 0_{\mathbb{R}^m}$ and $\psi(f(p)) = 0_{\mathbb{R}^n}$.) Moreover, one can assume that the chart images are product sets $\tilde{U} = \tilde{W} \times \tilde{U}'$, $\tilde{V} = \tilde{W} \times \tilde{V}'$, for some open sets $\tilde{W} \subseteq \mathbb{R}^k$, $\tilde{U}' \subseteq \mathbb{R}^{m-k}$ and $\tilde{V}' \subseteq \mathbb{R}^{n-k}$.

The hypothesis of constant rank near p is satisfied automatically in the following cases:

- If f is submersive at p (i.e. if $n = k$): In this case the chart ψ may be obtained by restricting an arbitrary chart. Equivalently, if N is an open subset of \mathbb{R}^k , one can take $\psi = \text{id}_V$, where $V \subseteq N$ is an open neighborhood of $f(p)$.
- If f is immersive at p (i.e. if $m = k$): In this case the chart ϕ may be obtained by restricting an arbitrary chart. Equivalently, if M is an open subset of \mathbb{R}^k one can take $\phi = \text{id}_U$, where $U \subseteq M$ is an open neighborhood of p .

This theorem says that any map of constant rank (in particular, any immersion or submersion) is locally equivalent to the standard map given in Example 3.0.2 above. If $T_p f$ does not have maximum rank (i.e. if $\text{rank}_p f =: k < m, n$), then the hypothesis of having constant rank k near p is hard to satisfy because the rank is lower semicontinuous but not upper semicontinuous in general. We will prove the constant rank theorem only in the cases $k = m$ and $k = n$, which are the most useful.

Example 3.2.3. If $f : V \rightarrow W$ is an *linear* map of rank k between topological spaces $V \simeq \mathbb{R}^m$ and $W \simeq \mathbb{R}^n$, then there are *linear* isomorphisms $\phi : V \rightarrow \mathbb{R}^m$, $\psi : W \rightarrow \mathbb{R}^n$ such that $\psi \circ f \circ \phi^{-1}$ is a standard map of rank k . (Exercise.)

Before proving the constant rank theorem, let us discuss the situation. We use the following terminology and notation for a point $x = (x^i)_{0 \leq i < m} \in \mathbb{R}^m$. Its **horizontal part** is the tuple of its first k -coordinates,

$$x^{\text{hor}} := (x^0, \dots, x^{k-1}) = (x^i)_{0 \leq i < k} \in \mathbb{R}^{[0,k)} = \mathbb{R}^k,$$

and its **vertical part** is the tuple of the remaining coordinates

$$x^{\text{ver}} := (x^k, \dots, x^{m-1}) = (x^i)_{k \leq i < m} \in \mathbb{R}^{[k,m)} \equiv \mathbb{R}^{m-k}.$$

Similarly, for a point $y \in \mathbb{R}^n$, its horizontal and vertical parts are $y^{[0,k)}$ and $y^{[k,n)}$.

From the constant rank theorem we conclude that the map $f|_U^V$ has the following features: its image is a k -submanifold of V , and each fiber (i.e. the preimage of a point of the image) is an $(m - k)$ -submanifold of U . These features are evident in the local expression $f|_{\tilde{\phi}}^{\psi}$, but they cannot be created by the charts, therefore they must be already present in the function $f|_U^V$.

The main job of the chart ϕ is to straighten the fibers of $f|_U^V$ so that they appear in the chart image $\tilde{U} \subseteq \mathbb{R}^m$ as the vertical slices $\{x^{[0,k]}\} \times \tilde{U}'$. For this task, the first k coordinates must be chosen carefully, so that they are constant on the fibers of f . This straightening is not necessary if $k = m$ (because the fibers are points), hence we may choose ϕ arbitrarily in this case.

Similarly, the main job of the chart ψ is to straighten the image of $f|_U^V$ so that it appears in the chart image $\tilde{V} \subseteq \mathbb{R}^m$ as the horizontal null plane $\tilde{W} \times \{0_{\mathbb{R}^{[k,n]}}\}$. For this task, the last $n - k$ coordinates are the most important; they must vanish on the image of $f|_U^V$. This straightening is not necessary if $k = n$ (because the image is an open set, not a submanifold of smaller dimension), hence we may choose ψ arbitrarily in this case.

Finally, we must ensure that the local expression $f|_\phi^\psi$ maps each vertical fiber $\{x^{hor}\} \times \tilde{U}'$ to the right point $(x^{hor}, 0, \dots, 0)$ of the image. (To see that something is needed consider a map $f : \mathbb{R}^m \rightarrow \mathbb{R}^n$ of the form $f(x) = (\theta(x^{hor}), 0_{\mathbb{R}^{n-k}}) \in \mathbb{R}^n$, where $\theta : \mathbb{R}^k \rightarrow \mathbb{R}^k$ is a diffeomorphism. This map f has constant rank k but it is not in the standard form, even though its fibers are the vertical slices and its image is the horizontal null plane.) This third job can be done either by ϕ or by ψ , but it must be done by one of the two charts. This is why in the case $k = m = n$ we cannot choose both ϕ and ψ arbitrarily.

Proof of the constant rank theorem (for submersions and immersions). Since the theorem is local, by taking coordinates we may assume that M, N are open subsets of $\mathbb{R}^m, \mathbb{R}^n$ respectively and that $p = 0_{\mathbb{R}^m}$ and $f(p) = 0_{\mathbb{R}^n}$. Each point $x \in M$ is a tuple of coordinates $x = (x^i)_{i \in [0,m]}$, and, similarly a point $y \in N$ is of the form $y = (y^j)_{j \in [0,n]}$. Since $D_p f$ has rank k , we may assume wlog (by permutting the indices, if necessary) that the $k \times k$ matrix

$$\left(\frac{\partial y^j}{\partial x^i} \Big|_p \right)_{i \in [0,k], j \in [0,k]}, \text{ where } y = f(x), \text{ is invertible.} \quad (3.1)$$

Note that if $k = m$, then no permutation of the i indices is needed, and if $k = n$, then no permutation of the j indices is needed.

Case f submersive at p (i.e. $k = n$): We define on M near p a coordinate system ϕ by the formula

$$\phi^j(x) = u^j = \begin{cases} y^j & \text{if } j \in [0, k) \\ x^j & \text{if } j \in [k, m) \end{cases}$$

where $y = f(x)$. The first k coordinates are defined in the only possible way that ensures that the map $f|_\phi^{\text{id}_V} = f \circ \phi^{-1}$ has the right form, sending $u \mapsto (u^j)_{j \in [0,k]}$. The remaining $m - k$ coordinates are just meant to complete the coordinate system. To verify that this correspondence $x \mapsto u$ really defines a chart ϕ on a neighborhood of p , we check that its differential $D_p \phi$ is invertible. And indeed, the matrix

$$\left(\frac{\partial u^j}{\partial x^i} \Big|_{x=p} \right)_{\substack{j \in m \\ i \in m}} = \left(\begin{array}{c|c} \left(\frac{\partial y^j}{\partial x^i} \Big|_{x=p} \right)_{\substack{j \in k \\ i \in k}} & * \\ \hline 0 & I_{m-k} \end{array} \right)$$

is invertible because the block $\left(\frac{\partial y^j}{\partial x^i} \Big|_{x=p} \right)_{\substack{j \in k \\ i \in k}}$ by the hypothesis (3.1). Thus by the

IFT, there are respective open neighborhoods $U \subseteq M, \tilde{U} \subseteq \mathbb{R}^m$ of $p, \phi(p)$, such that the restriction $\phi|_{\tilde{U}}$ is a diffeomorphism. Moreover, by shrinking the sets U, \tilde{U}

we can ensure that $\tilde{U} = \tilde{W} \times \tilde{U}'$, where $\tilde{W} \subseteq \mathbb{R}^k$ and $\tilde{U}' \subseteq \mathbb{R}^{m-k}$ are open sets. On the manifold N we choose the chart $\psi = \text{id}_{\tilde{W}}$.

Note: The fact that we were able to put $\psi = \text{id}_{\tilde{W}}$ means that when M is a general manifold (instead of an open subset of \mathbb{R}^m) we can obtain ψ as the restriction of an arbitrary chart.

Case f immersive at p (i.e. $k = m$): Note that the horizontal part of f , i.e. the map $\theta : x \mapsto f(x)^{ver}$, is a diffeomorphism when restricted to a suitable domain and codomain (respective open neighborhoods of p and $f(p)^{hor}$ in \mathbb{R}^k). This follows from condition (3.1) by the IFT.

We define on N near $f(p)$ a coordinate system ψ by the formula

$$\begin{cases} \phi^{hor}(y) = \underline{x} \\ \phi^{ver}(y) = y^{ver} - f^{ver}(\underline{x}) \end{cases}$$

where $\underline{x} = \theta^{-1}(y^{hor})$. For a point $y = f(x)$ we have

$$\underline{x} = \theta^{-1}(y^{hor}) = \theta^{-1}(f^{hor}(x)) = \theta^{-1}(\theta(x)) = x,$$

therefore the coordinates $\phi(y)$ have the expected behaviour: the horizontal part $\phi(y)^{hor}$ equals x , while the vertical part vanishes:

$$\phi(y)^{ver} = f(x)^{ver} - f^{ver}(\underline{x}) = 0.$$

Therefore the map $f|_{\text{id}}^\psi = \psi \circ f$ sends $x \mapsto (x, 0)$, as required.

To ensure that ψ is a chart on a neighborhood of $f(p)$, we verify that its differential $D_{f(p)}\phi$ is invertible. And indeed, its matrix of partial derivatives

$$\left(\frac{\partial v^i}{\partial y^j} \Big|_{y=f(p)} \right)_{\substack{i \in n \\ j \in n}} = \left(\begin{array}{c|c} \left(\frac{\partial x^i}{\partial y^j} \Big|_{y=f(p)} \right)_{\substack{i \in k \\ j \in k}} & 0 \\ \hline * & I_{n-k} \end{array} \right),$$

is invertible. This follows because the matrix $\left(\frac{\partial x^i}{\partial y^j} \Big|_{y=f(p)} \right)_{i \in k, j \in k}$, which represents the linear transformation $D_{f(p)}\theta^{-1} = (D_p\theta)^{-1}$, is invertible. Thus by the IFT, there are respective open neighborhoods $V \subseteq N$, $\tilde{V} \subseteq \mathbb{R}^n$ of $f(p)$, $\psi(f(p))$, such that the restriction $\phi|_{\tilde{V}}$ is a diffeomorphism. Moreover, by shrinking the sets \tilde{V} , V we can ensure that $\tilde{V} = \tilde{W} \times \tilde{V}'$, where $\tilde{W} \subseteq \mathbb{R}^k$ and $\tilde{V}' \subseteq \mathbb{R}^{n-k}$ are open sets. On the manifold M we choose the chart $\psi = \text{id}_{\tilde{W}}$.

General case* (i.e. $k \leq m, n$) Here one can combine the two constructions. If interested, you can take it as an exercise or read the proof e.g. on p. 42 of Spivak's "Comprehensive introduction to Differential Geometry", volume 1. \square

3.3 Some properties of immersions and embeddings

3.3.1 Local properties: slices and retractions

The following corollary of the constant rank theorem says that for any immersion $f : M \rightarrow N$, if we restrict f to a sufficiently small neighborhood U of any point $p \in M$, its image $f(U)$ looks like a "slice" of some open set $V \subseteq N$. This implies that f has a local retraction and is locally an embedding.

Lemma 3.3.1 (Slice property (and local retractions) for immersions). *Let N be an n -manifold and let $f : M \rightarrow N$ be an immersion with $\dim M = k$. Then for each point p of M there exists a chart $\psi : V \rightarrow \tilde{V}$ of N and an open neighborhood U of p such that*

$$f(U) = \{q \in V : \psi^k(q) = \cdots = \psi^{n-1}(q) = 0\},$$

and such that the map $f|_U^V$ is an embedding and has a \mathcal{C}^k retraction $\rho : V \rightarrow U$.

Proof. By the constant rank theorem, the map f admits at the point p a local expression $\tilde{f} = f|_\phi^\psi$ of the form $\tilde{f}(x) = (x^0, \dots, x^{k-1}, 0, \dots, 0)$, where $\phi : U \rightarrow \tilde{U}$, $\psi : V \rightarrow \tilde{V}$ are charts of M, N defined at the points $p, f(p)$ respectively. Moreover, we can assume that $\tilde{V} = \tilde{U} \times \tilde{V}'$, where $\tilde{V}' \subseteq \mathbb{R}^{n-k}$ is an open neighborhood of $0_{\mathbb{R}^{n-k}}$. It follows that $\tilde{f}(\tilde{U}) = \tilde{U} \times \{0_{\mathbb{R}^{n-k}}\}$ and hence

$$f(U) = \psi^{-1}(\tilde{U} \times \{0_{\mathbb{R}^{n-k}}\}) = \{q \in V : \psi^k(q) = \cdots = \psi^{n-1}(q) = 0\}.$$

To construct the retraction ρ , consider the projection map

$$\begin{aligned} \pi : \tilde{V} = \tilde{U} \times \tilde{V}' &\rightarrow \tilde{U} \\ (x, y) &\mapsto x. \end{aligned}$$

There is a unique map $\rho : V \rightarrow U$ that has local expression $\rho|_\psi^\phi = \pi$, namely, the map $\rho = \phi^{-1} \circ \pi \circ \psi : V \rightarrow U$. This map ρ is a \mathcal{C}^r retraction of $f|_U^V$. The existence of this retraction implies that $f|_U^V$ is an embedding by Proposition 3.1.3. \square

We have shown that $f(U)$ is a slice of V , but note that the set $f(M \setminus U)$ may also intersect the set V . (Exercise: Find an example (i.e. an immersion f and a point p) such that this happens for any V .) However, this situation can be avoided if f is an embedding.

Proposition 3.3.2 (Slice property for embedded submanifolds). *Let S be the image of a \mathcal{C}^r embedding $f : M \rightarrow N$, and let $k = \dim M$ and $n = \dim N$. Then each point of S is contained in the domain V of a chart ψ that is k -**sliced** by S , meaning that*

$$S \cap V = \{q \in V : \phi^k(q) = \cdots = \phi^{n-1}(q) = 0\}. \quad (3.2)$$

Proof. Take a point $q_0 = f(p_0) \in S$. By Lemma 3.3.1, there exists a chart ψ of M with domain V' and a neighborhood U of p_0 such that $f(U) = \{q \in V' : \psi^k(q) = \cdots = \psi^{n-1}(q) = 0\}$. Since $f(U)$ is open in S , we can write $f(U) = S \cap V''$ using some open set $V'' \subseteq M$. But since $f(U) \subseteq V'$, we have $f(U) = S \cap V$, and therefore

$$S \cap V = f(U) = \{q \in V : \psi^k(q) = \cdots = \psi^{n-1}(q) = 0\}.$$

\square

3.3.2 The initial property

Recall that topological embeddings have the following *initial property*.

Proposition 3.3.3 (Initial property of topological embeddings). *If a continuous map $f : X \rightarrow Y$ between topological spaces is an embedding, then any function $h : Z \rightarrow X$ (where Z is a topological space) is continuous if the composite $f \circ h$ is continuous.*

A \mathcal{C}^r immersion has a similar property that allows us to show that a continuous function is \mathcal{C}^r .

Proposition 3.3.4 (Initial property of immersions and embeddings). *Let $f : M \rightarrow N$ be a \mathcal{C}^r -differentiable map, and let L be another \mathcal{C}^r manifold.*

- (a) *If f is an immersion, then any continuous map $h : L \rightarrow M$ is \mathcal{C}^r if the composite $f \circ h$ is \mathcal{C}^r .*
- (b) *If f is an embedding, then any function $h : L \rightarrow M$ is \mathcal{C}^r if the composite $f \circ h$ is \mathcal{C}^r .*

The proof is based on the fact that an immersion admits local retractions.

Proof. (a) Suppose that f is an immersion and that $h : L \rightarrow M$ is continuous at some point $z \in L$, and the composite $f \circ h$ is \mathcal{C}^r at some point $z \in L$. Let us show that h is \mathcal{C}^r at z .

Let $U \subseteq M$ and $V \subseteq N$ be open neighborhoods of the points $h(z)$ and $f(h(z))$ such that $f(U) \subseteq V$ and the map $f|_U^V$ has a \mathcal{C}^r retraction $\rho : V \rightarrow U$. Thus we have $\rho \circ f|_U^V = \text{id}_U$. Note that the set $W = h^{-1}(U)$ is a neighborhood of x since h is continuous at x . Thus we can write

$$h|_W^U = \text{id}_U \circ h|_W^U = (\rho \circ f|_U^V) \circ h|_W^U = \rho \circ (f \circ h)|_W^V.$$

This factorization shows that h is \mathcal{C}^r at the point z , since $f \circ h$ is \mathcal{C}^r at z and ρ is \mathcal{C}^r everywhere.

(b) Now suppose that f is a \mathcal{C}^r embedding and that $h : L \rightarrow M$ is any function such that the composite $f \circ h$ is \mathcal{C}^r . Since f is a topological embedding, the function h is continuous by Proposition 3.3.3. Since f is an immersion, the map h is \mathcal{C}^r by part (a). \square

3.3.3 Consequences of the initial property

- If two embeddings $f_0 : M_0 \rightarrow N$, $f_1 : M_1 \rightarrow M$ have the same image, then they are equivalent in the sense that there is a diffeomorphism $g : M_0 \rightarrow M_1$ such that $f_1 \circ g = f_0$. (Exercise.)
- If S is an embedded submanifold of M , then there is a unique topology and smooth structure on S such that the inclusion map $S \rightarrow M$ is an embedding. (Exercise.)

This allows us to define the tangent space of an embedded submanifold.

Definition 3.3.5. The **tangent space** of an embedded submanifold S of a differentiable manifold M is the subspace $T_p S \subseteq T_p M$ defined as the image $T_p S = \text{Im}(T_{f^{-1}(p)} f)$ of the differential of any embedding $f : N \rightarrow M$ whose image is S .

This subspace is well defined (i.e. independent of f). This follows from the fact that all the embeddings with image S are equivalent. (Exercise.)

3.4 How to recognize an embedded submanifold

Sometimes we are just given a subset S of a manifold M , and we have to determine whether it is an embedded submanifold or not. There are several ways to do so.

Proposition 3.4.1. *Let S be a subset of a \mathcal{C}^r manifold M , and let $k \leq n = \dim M$. The following are equivalent:*

- (a) *S is a \mathcal{C}^r -embedded k -submanifold of M .*
- (b) *Local embedded submanifold: Each point of S has an open neighborhood W in M such that the set $S \cap W$ is a \mathcal{C}^r -embedded k -submanifold of W .*

- (c) *Local slice:* Each point of S is contained in the domain W of a chart ψ of M that is k -sliced by S .
- (d) *Local retract:* Each point of S has an open neighborhood W in M such that the set $S \cap W$ is the image of a \mathcal{C}^r map $\phi : U \rightarrow W$ (where $U \subseteq \mathbb{R}^k$ is an open set), which in turn admits a \mathcal{C}^r retraction $\rho : W \rightarrow U$.

Proof. (a) \Rightarrow (b) Trivial: take $W = M$.

(a) \Rightarrow (c) This is Proposition 3.3.2.

(b) \Rightarrow (c) Same proof as (a) \Rightarrow (c).

(c) \Rightarrow (d) Let p be a point of S , and let $\psi : W \rightarrow \widetilde{W}$ be a chart of M , defined at p , that is k -sliced by S . By shrinking the sets W and \widetilde{W} , we may assume that $\widetilde{W} = \widetilde{U} \times \widetilde{V}$, for some open sets $\widetilde{U} \subseteq \mathbb{R}^k$ and $\widetilde{V} \subseteq \mathbb{R}^{n-k}$. The slice condition implies that

$$S \cap W = \psi^{-1}(\widetilde{U} \times \{0\}).$$

Consider the projection map $\pi : \widetilde{U} \times \widetilde{V} \rightarrow \widetilde{U} : (x, y) \mapsto x$ and its section $\sigma : \widetilde{U} \rightarrow \widetilde{U} \times \widetilde{V} : x \mapsto (x, 0)$. Note that $\pi \circ \sigma = \text{id}_{\widetilde{U}}$.

Now we use the diffeomorphism $\psi : W \rightarrow \widetilde{W}$ to transport the maps π and ρ from the Euclidean set $\widetilde{W} \subseteq \mathbb{R}^n$ to the manifold subset $W \subseteq M$. That is, we define the \mathcal{C}^r maps $\phi = \psi^{-1} \circ \sigma : \widetilde{U} \rightarrow W$ and $\rho = \pi \circ \psi : W \rightarrow \widetilde{U}$. Clearly ρ is a retraction of ϕ , that is, we have $\rho \circ \phi = \text{id}_{\widetilde{U}}$. The image of ϕ is the set

$$\text{Img}(\phi) = \phi^{-1}(\text{Img}(\sigma)) = \phi^{-1}(\widetilde{U} \times \{0\}) = S \cap W.$$

(d) \Rightarrow (a) Suppose S is covered by open sets W_i such that each set $V_i := S \cap W_i$ is the image of a \mathcal{C}^r maps $\phi_i : U_i \rightarrow W_i$ admitting a retraction $\rho_i : W_i \rightarrow U_i$.

Let us show that S admits a structure of k -dimensional \mathcal{C}^r manifold such that the inclusion map $\iota_S : S \rightarrow M$ is a \mathcal{C}^r embedding. The only topology we may put on S (so that ι_S a topological embedding) is the subspace topology. This topology is Hausdorff and second countable.

Let us construct a \mathcal{C}^r atlas on S . The maps ϕ_i and ρ_i of course satisfy $\rho_i \circ \phi_i = \text{id}_{U_i}$. Composing in the other order we get a map $\phi_i \circ \rho_i$ that fixes the image of ϕ_i , which is the set V_i . Therefore the continuous maps

$$\overline{\phi}_i := \phi_i|_{V_i} : U_i \rightarrow V_i, \quad \overline{\rho}_i := \rho_i|_{V_i} : V_i \rightarrow U_i$$

are inverse of each other. The maps $\overline{\rho}_i$ form a \mathcal{C}^r atlas of S since the transition maps

$$\overline{\rho}_i \circ \overline{\rho}_j^{-1} = \overline{\rho}_i \circ \overline{\phi}_j = \rho_i \circ \phi_j$$

are \mathcal{C}^r .

The inclusion $\iota_S : S \rightarrow M$ is an immersion because composing with each local parametrization $\overline{\phi}_i$ we obtain a map $\iota_S \circ \overline{\phi}_i = \phi_i$ that is a \mathcal{C}^k immersion, since it admits a \mathcal{C}^k retraction. \square

Example 3.4.2. A subset $S \subseteq \mathbb{R}^n$ that can be locally expressed as a graph of a \mathcal{C}^r function (with some $n - k$ coordinates expressed as a function of the other k coordinates) is a k -submanifold of \mathbb{R}^n because it satisfies condition (d), since graphing maps admit projections as retractions. This is essentially what we used to construct the manifold structure on the sphere \mathbb{S}^{n-1} .

3.5 How to produce a submanifold as a level set

Definition 3.5.1. Let $f : M \rightarrow N$ be a \mathcal{C}^k -differentiable map. A point $q \in N$ is called a **regular value** of f if $T_p f$ is surjective for all points $p \in f^{-1}(q)$.

Theorem 3.5.2 (Regular preimage theorem). *Let c be a regular value of a C^r -differentiable map $f : M \rightarrow N$, and let $m = \dim M$, $n = \dim N$. Then the set $S = f^{-1}(c)$ is a C^r -embedded submanifold of M of dimension $k = m - n$. Its tangent space at any point $p \in S$ is $T_p S = \text{Ker}(T_p f)$.*

Proof. We will use the constant rank theorem to show that S fulfills the “local slice” condition of Proposition 3.4.1, therefore S is a submanifold.

At a point $p_0 \in S$ the map f is submersive, therefore by the CRT it admits at the point p_0 a local expression $\tilde{f} = f|_{\tilde{\phi}}^\psi$ of the form $\tilde{f}(x) = (x^k, \dots, x^{m-1})$, where $\phi : U \rightarrow \tilde{U}$, $\psi : V \rightarrow \tilde{V}$ are charts of M , N defined and centered at the points p_0 , $f(p_0) = c$ respectively. In addition, we may assume that $\tilde{U} = \tilde{V} \times \tilde{U}'$ where $U' \subseteq \mathbb{R}^{m-n}$ is an open set.

Note that $\psi(c) = 0$. Thus for a point $p \in U$, denoting $x = \phi(p) \in \mathbb{R}^m$, we have $p \in S$ iff $f(p) = c$ iff $\tilde{f}(x) = 0$ iff the last n coordinates x^k, \dots, x^{m-1} vanish. In other words, the chart ϕ is k -sliced by $S = f^{-1}(c)$. Since we can find such a chart ϕ for any point $p_0 \in S$, we conclude that S is a C^r -embedded k -submanifold of M .

Tangent space: exercise. □

3.6 Whitney's theorem*

Theorem 3.6.1 (Whitney). *Every compact C^k manifold M can be C^k embedded in \mathbb{R}^m for some m .*

Proof. Let $n = \dim M$, and denote $B(r) = \{x \in \mathbb{R}^n : \|x\| < r\}$ for $r > 0$. Since M is compact, there is a finite family $(\phi_i)_{i \in N}$ of charts $\phi_i : U_i \rightarrow B(2)$ such that the open sets $B_i := \phi_i^{-1}B(1)$ cover M .

For each i , take a C^k bump function $\eta_i : M \rightarrow [0, 1]$ with support $\text{supp } \eta_i \subseteq U_i$ such that $\eta_i \equiv 1$ on B_i , and define a function $f_i : M \rightarrow \mathbb{R}^n$ by

$$f_i(x) = \begin{cases} \eta_i(x) \phi_i(x) & \text{if } x \in U_i \\ 0 & \text{if } x \notin \text{supp}(\eta_i) \end{cases}$$

Finally, define a C^k map $f : M \rightarrow \mathbb{R}^{Nn+N}$ by

$$f(x) = ((f_i(x))_{i \in N}, (\eta_i(x))_{i \in N}).$$

We claim that f is an embedding.

f is an immersion because each f_i is immersive on B_i , since it coincides with the chart ϕ_i .

f is injective: suppose that $f(x) = f(y)$. Take some i such that $x \in B_i$. Then $\eta_i(x) = 1$, and therefore $\eta_i(y) = 1$. But then $x, y \in U_i$ and we have $\phi_i(x) = f_i(x) = f_i(y) = \phi_i(y)$, which implies that $x = y$.

Since f is an injective map with compact domain (and Hausdorff codomain), it is a topological embedding. □

4 Vector bundles and vector fields

So far we have looked at each of the vector spaces $T_p M$, where $p \in M$, separately. Their union $TM := \bigcup_{p \in M} T_p M$ is called the **tangent bundle** of M . It has a special structure that we will encounter several times and so it is worthwhile introducing the following definition:

Definition 4.0.1. Let M be a C^r manifold. A C^r **vector bundle** of rank k over M is a C^r manifold E together with a C^r map $\pi = \pi_E : E \rightarrow M$ (the **projection map**) satisfying

- (a) For each $p \in M$, the set $\pi^{-1}(p) =: E_p$ (called the **fibre** of E over p) is a k -dimensional vector space.
- (b) Each point of M is contained in an open set $U \subseteq M$ such that there is a C^r diffeomorphism

$$\Phi : U \times \mathbb{R}^k \rightarrow \pi^{-1}(U)$$

(called a **local trivialization** of E over U) such that for each $p \in U$, Φ maps $\{p\} \times \mathbb{R}^k$ to E_p , and the restriction $\Phi_p : \{p\} \times \mathbb{R}^k \cong \mathbb{R}^k \rightarrow E_p$ is a vector space isomorphism.

We call E the **total space** of the bundle and M the **base space**.

Note: When talking about a vector bundle (E, π) , sometimes we omit mentioning the projection map π and we just say that E is a vector bundle.

If (E, π) and (E', π') are smooth vector bundles over the same smooth manifold M , a **vector bundle map** (or **vector bundle morphism**) $(E, \pi) \rightarrow (E', \pi')$ is a smooth map $f : E \rightarrow E'$ such that $\pi' \circ f = \pi$ and with the property that for each $p \in M$ the restricted map $f|_{E_p} : E_p \rightarrow E'_p$ is linear. A **vector bundle isomorphism** is a bundle map that has an inverse.

The product $M \times \mathbb{R}^k$ with the standard projection $\pi : M \times \mathbb{R}^k \rightarrow M$ is called **the trivial vector bundle** of rank k over M . (This is indeed a vector bundle since one can take a single local trivialization: the identity map.) A vector bundle of rank k over M is called **trivial** if it is isomorphic to *the* trivial bundle $M \times \mathbb{R}^k$.

4.1 The tangent bundle

By this point you are certainly suspecting that the tangent bundle TM of a differentiable manifold is a vector bundle. First, let us show that it is a smooth manifold.

Lemma 4.1.1. *Let M be a C^{k+1} manifold. Then the tangent bundle TM has a natural topology and C^k structure making it a $2n$ -dimensional manifold. The natural projection $\pi : TM \rightarrow M$, which sends $X \in T_p M$ to p , is a C^k map.*

Proof. We define the maps that will become our charts as follows: Given a local parametrization $\varphi : \tilde{U} \subseteq \mathbb{R}^n \rightarrow U \subseteq M$, let

$$\begin{aligned} \Phi : \tilde{U} \times \mathbb{R}^n &\rightarrow \pi^{-1}(U) \\ (x, v) &\mapsto [\varphi(x), \varphi^{-1}, v], \end{aligned}$$

or equivalently, in terms of the coordinate vectors of the chart $\psi = \varphi^{-1}$, $\Phi(x, v) = \sum_i \frac{\partial}{\partial \psi^i} \Big|_{\varphi(x)}$. This function Φ is a bijection onto its image $\pi^{-1}(U)$. We define on TM a topology as follows:

A set $A \subseteq TM$ is open \iff for each φ , the set $\Phi^{-1}(A) \subseteq \tilde{U} \times \mathbb{R}^n$ is open

You can check that this is indeed a topology that is Hausdorff and second countable. Each map Φ is a homeomorphism from an open subset of \mathbb{R}^n onto its image, which is an open subset of M . Hence TM is a topological manifold of dimension $2n$.

We show that the collection of maps Φ obtained as above from local parametrizations φ of M is a parametrization atlas for TM . Let $\varphi : \tilde{U} \rightarrow U$, $\psi : \tilde{V} \rightarrow V$ be smooth local parametrizations of M and let Φ, Ψ be the corresponding parametrizations of TM . Then the transition map is

$$\begin{aligned} \Psi^{-1} \circ \Phi : \varphi^{-1}(U \cap V) \times \mathbb{R}^n &\rightarrow \psi^{-1}(U \cap V) \times \mathbb{R}^n \\ (x, v) &\mapsto (\psi^{-1} \circ \varphi(x), D_x(\psi^{-1} \circ \varphi)(v)) \end{aligned}$$

which is smooth. This formula comes from the fact that if two tangent vectors $\Phi(x, v) = [\varphi(x), \varphi^{-1}, v]$ and $\Psi(y, w) = [\psi(y), \psi^{-1}, w]$ are equal, then $y = \psi^{-1}\varphi(x)$ and $w = D_x(\psi^{-1} \circ \varphi)(v)$.

To check that the projection map π is smooth, just note that its coordinate representation with respect to the local parametrizations φ and Φ is $\pi|_{\Phi}^{\varphi}(x, v) = x$. \square

The smooth structure defined in the previous proof is called the **standard smooth structure** for TM . (Similarly, if M is a \mathcal{C}^{r+1} manifold, then TM is naturally a \mathcal{C}^r manifold.)

Proposition 4.1.2. *The tangent bundle TM with the standard smooth structure and the standard projection $\pi : TM \rightarrow M$ is a smooth vector bundle of rank n over M .*

Proof. Each fibre $T_p M = \pi^{-1}(\{p\})$, $p \in M$, is a vector space. Given a smooth local parametrization $\varphi : \tilde{U} \subseteq \mathbb{R}^n \rightarrow U \subseteq M$, the map

$$\begin{aligned} U \times \mathbb{R}^n &\rightarrow \pi^{-1}(U) \\ (p = \varphi(x), v) &\mapsto [p, \varphi^{-1}, v] \end{aligned}$$

is a local trivialization of TM over U . \square

4.2 Sections, frames, and trivial bundles

Definition 4.2.1. Let (E, π) be a \mathcal{C}^k vector bundle over a manifold M . A **section** of a E is a continuous map $\sigma : M \rightarrow E$ such that $\pi \circ \sigma = \text{id}_M$. It is \mathcal{C}^k if it is \mathcal{C}^k as a map between manifolds.

If $U \subset M$ is open it is easy to verify that $E|_U := \pi^{-1}(U)$ is again a vector bundle with the restriction of π as its projection map. A (smooth) section of $E|_U$ is called a (smooth) **section of E over U** .

Definition 4.2.2. If $U \subset M$ is open, a **local frame** for E over U is an ordered k -tuple $(\sigma_0, \dots, \sigma_{k-1})$ where each σ_i is a smooth section of E over U such that $(\sigma_0(p), \dots, \sigma_{k-1}(p))$ is a base for the fibre E_p for each $p \in U$. It is called a **global frame** if $U = M$.

For the vector bundles we encounter in this lecture we usually write the value of a section σ at a point p as $\sigma_p := \sigma(p) \in E_p$. Addition of two sections σ, τ of E over an open subset $U \subset M$ is defined pointwise: $(\sigma + \tau)_p := \sigma_p + \tau_p$. We can also multiply a section over U by a function $f \in C^\infty(U)$ by defining the operation pointwise: $(f\sigma)_p := f(p)\sigma_p$. For those who know what it means: this makes the set of smooth sections of E over U a module over the ring $C^\infty(U)$.

Proposition 4.2.3. *A smooth vector bundle is trivial iff it has a global frame.*

Proof. Exercise. □

E.g. the tangent bundle $T\mathbb{R}^n$ is trivial, since it has a global frame consisting of the coordinate vector fields $p \mapsto \frac{\partial}{\partial x^i}|_p$.

Let us give an example of a non-trivial vector bundle.

Example 4.2.4. (Möbius band) The Möbius band E is the quotient space of \mathbb{R}^2 under the equivalence relation

$$(x, t) \sim (x', t') \iff x' - x = n \in \mathbb{Z} \text{ and } t' = (-1)^n t.$$

We endow E with the quotient topology and one can check that E is Hausdorff and second countable. We write $[x, t]$ for the equivalence class of $(x, t) \in \mathbb{R}^2$ in E .

We want to make E a vector bundle over the 1-torus $\mathbb{T}^1 = \mathbb{R}/\mathbb{Z}$. The projection map will be $\pi : E \rightarrow \mathbb{T}^1 : [x, t] \mapsto [x]$. Note that each fibre $E_t = \pi^{-1}[x] = \{[x, t] : t \in \mathbb{R}\}$ is a vector space with the operations $a[x, t] + a'[x, t'] = [x, at + a't']$.

Recall the smooth structure of the torus \mathbb{T}^1 . The quotient map $\kappa : \mathbb{R} \rightarrow \mathbb{T}^1 : x \mapsto [x]$ is open, and it is injective when restricted e.g. to any open interval $W \subseteq \mathbb{R}$ of length ≤ 1 . Consider a family of such intervals W_i whose images $U_i = \kappa(W_i)$ cover \mathbb{T}^1 (for example, $W_1 = (0, 1)$ and $W_0 = (-\frac{1}{2}, \frac{1}{2})$). Then the maps $\kappa|_{W_i} : W_i \rightarrow U_i$ form a smooth inverse atlas of \mathbb{T}^1 (i.e. its inverse functions $W_i \rightarrow U_i$ form a smooth atlas).

Let us give E a smooth structure. We cover E with open sets

$$V_i = \pi^{-1}(U_i) = \{[x, t] : x \in W_i, t \in \mathbb{R}\}$$

and we define homeomorphisms

$$\psi_i : W_i \times \mathbb{R} \rightarrow V_i : (x, t) \mapsto [x, t]$$

These maps form a smooth inverse atlas of E since the transition maps are smooth (check it!). Therefore E is a smooth manifold. In addition, (E, π) is a smooth vector bundle of rank 1 over \mathbb{T}^1 since it admits the local trivializations

$$\tilde{\psi}_i : U_i \times \mathbb{R} \rightarrow V_i : ([x], t) \mapsto [x, t].$$

We claim that E is not trivial. Indeed, suppose that E is trivial. Then there is a global frame i.e. a nowhere vanishing smooth section $\sigma : \mathbb{T}^1 \rightarrow E$.

Note that a smooth section σ is of the form $\sigma([x]) = [x, f(x)]$ for some function $f : \mathbb{R} \rightarrow \mathbb{R}$, and we claim that f is smooth. To see that f is smooth on each interval W_i , we express it as a composite of smooth functions

$$\begin{array}{ccccccc} f|_{W_i} : & W_i & \xrightarrow{\kappa} & U_i & \xrightarrow{\sigma} & V_i & \xrightarrow{\psi_i^{-1}} & W_i \times \mathbb{R} & \xrightarrow{\pi_1} & \mathbb{R} \\ & x & \mapsto & [x] & \mapsto & [x, t] & \mapsto & (x, t) & \mapsto & t. \end{array}$$

Also note that f is “antiperiodic” with period 1, i.e., it satisfies $f(x+1) = -f(x)$ because

$$[x, f(x)] = \sigma[x] = \sigma[x+1] = [x+1, f(x+1)] = [x, -f(x+1)].$$

On the other hand, f does not vanish if σ is a nonvanishing section. This is incompatible with antiperiodicity (by the intermediate value theorem), therefore there is no nonvanishing section σ .

4.3 Vector fields

Definition 4.3.1. A **tangent vector field** (or a **vector field**, for short) on a \mathcal{C}^{k+1} manifold M is a function $X : M \rightarrow TM$ such that

$$X_p \in T_p M \quad \text{for each } p \in M$$

where $X_p := X(p)$.

Its **component functions** with respect to a chart (U, ϕ) are the functions $X^i : U \rightarrow \mathbb{R}$ such that $X \equiv \sum_i X^i \frac{\partial}{\partial \phi^i}$ on U .

We say that X is \mathcal{C}^k at a point $p \in M$ if its component functions (w.r.t. any chart ϕ defined at p) are \mathcal{C}^k at the point p . (Exercise: it suffices to check just one chart.) We say that X is \mathcal{C}^k if it is \mathcal{C}^k at all points. The set of \mathcal{C}^k tangent vector fields on M is denoted $\mathfrak{X}(M)$.

Equivalently, in terms of vector bundles, a tangent vector field is a section of the tangent bundle $\pi : TM \rightarrow M$, i.e. a function $X : M \rightarrow TM$ such that $\pi \circ X = \text{id}_M$. It is \mathcal{C}^k if it is \mathcal{C}^k as a function $M \rightarrow TM$. (Exercise.)

Addition of two vector fields and multiplication by functions on M is defined in the natural way, i.e. pointwise: $(X + Y)_p = X_p + Y_p$, $(fX)_p = f(p)X_p$ where $f \in C^\infty(M)$.

Remark 4.3.2. A vector field over an open subset $U \subset M$ is a section of TM over U . In particular, given (U, φ) a \mathcal{C}^{k+1} chart, the so-called **coordinate vector fields**

$$\frac{\partial}{\partial \varphi^i} : U \rightarrow T_p M : p \mapsto \left. \frac{\partial}{\partial \varphi^i} \right|_p$$

are \mathcal{C}^k vector fields over U .

Example 4.3.3. Recall the polar coordinates (r, φ) defined on the open subset $U \subset \mathbb{R}^2$ (Example 2.0.5). The coordinate vector field $\frac{\partial}{\partial \varphi}$ is a smooth vector field on U . Writing the vector field in terms of the Euclidean coordinate vectors

$$\frac{\partial}{\partial \varphi} = -y \frac{\partial}{\partial x} + x \frac{\partial}{\partial y}$$

we see that $\frac{\partial}{\partial \varphi}$ extends to a smooth vector field on \mathbb{R}^2 , which we denote by the same symbol $\frac{\partial}{\partial \varphi}$. So while the coordinate φ cannot be extended to the whole space \mathbb{R}^2 as a smooth function, the corresponding coordinate vector field can be extended.

Example 4.3.4. On \mathbb{S}^1 we can define a nowhere vanishing vector field (exercise). Therefore, the vector bundle $T\mathbb{S}^1$ is trivial. The tangent bundle $T\mathbb{S}^2$ is *not* trivial: the *hairy ball theorem* says that there is no nowhere vanishing vector field on \mathbb{S}^2 .

4.4 Integral curves and flow of a vector field

Definition 4.4.1. An **integral curve** of a tangent vector field X on a manifold M is a \mathcal{C}^1 curve $\gamma : I \rightarrow M$ (where $I \subseteq \mathbb{R}$ is an open interval) such that

$$\gamma'(t) = X|_{\gamma(t)} \quad \text{for all } t \in I. \quad (4.1)$$

(Here we use the notation $\gamma'(t)$ for the velocity vector $\text{Vel}_t \gamma$.)

Equation (4.1) is an ordinary differential equation (ODE) on the manifold M . In a local chart, it boils down to solving the system of ODEs:

$$(\gamma^i)'(t) = X^i(\gamma(t)), \quad i = 0, \dots, n-1$$

in \mathbb{R}^n . Similarly to ODE's in \mathbb{R}^n , for manifolds there is a theorem of existence and uniqueness of solutions.

Proposition 4.4.2 (Existence and uniqueness of solutions for an ODE on a manifold). *If X is a \mathcal{C}^1 vector field, then for each point $p \in M$ there exists a unique maximal integral curve $\gamma_{X,p} : I_{X,p} \rightarrow M$ of X that satisfies the initial condition $\gamma_{X,p}(0) = p$. Any other integral curve of X satisfying the same initial condition is equal to $\gamma_{X,p}$, restricted to a subinterval of $I_{X,p}$.*

This proposition can be deduced from the corresponding proposition in \mathbb{R}^n . We will not study the proof.

Definition 4.4.3. The **flow** of a \mathcal{C}^1 vector field on a manifold M is the function

$$\begin{aligned} \Phi_X &= \coprod_{p \in M} I_{X,p} && \rightarrow && M \\ &(p, t) && \mapsto && \gamma_{X,p}(t). \end{aligned}$$

We also write $\Phi_X^t(p) := \Phi_X(p, t)$.

Proposition 4.4.4 (Differentiability w.r.t. initial conditions for an ODE on a manifold). *If X is a \mathcal{C}^k vector field with $k \geq 1$, the flow Φ_X is a \mathcal{C}^k map defined on an open subset of $M \times \mathbb{R}$.*

This theorem also follows from a corresponding theorem in \mathbb{R}^n , found e.g. in Hartman's ODE book, Thm. 4.1. Again, we skip the proof.

Definition 4.4.5. We say that a vector field X is **complete** if the flow Φ_X is defined over $M \times \mathbb{R}$, or equivalently, if every maximal integral curve of X is defined on \mathbb{R} .

Note that a vector field on a compact manifold (or, more generally, a vector field with compact support) is complete. (Exercise.)

Example 4.4.6. On Euclidean space \mathbb{R}^n , a constant vector field $X \equiv a$ has maximal integral curves $\gamma_{X,p}(t) = p + ta$, defined for all $t \in \mathbb{R}$. Hence X is complete.

Example 4.4.7. On the line \mathbb{R} consider the vector field X given by $X_p = p^2 \frac{\partial}{\partial \phi} \Big|_p$, where $\phi = \text{id}_{\mathbb{R}}$ is the identity map considered as a chart with just one component function $\phi^0 = \phi$. Let us find an integral curve γ with starting point $\gamma(0) = 1$. This means that we have to solve

$$\gamma'(t) = (\gamma(t))^2, \quad \gamma(0) = 1.$$

As is shown easily by separation of variables, the maximal solution is $\gamma(t) = \frac{1}{1-t}$, $t \in (-\infty, 1)$, i.e. it goes to infinity in a finite time and cannot be defined for all $t \in \mathbb{R}$. Therefore X is not complete.

The notion of *related vector fields* is useful for transporting solutions of ODE's from one manifold to another.

Definition 4.4.8. Let $f : M \rightarrow N$ be a smooth map. A vector field $X \in \mathfrak{X}(M)$ is **f -related** to a vector field $Y \in \mathfrak{X}(N)$ if $T_p f(X_p) = Y_{f(p)}$ for all $p \in M$.

The following facts follow from the definition (exercise):

- (a) X is f -related to Y if and only if $X_p(f \circ h) = Y_{f(p)}(h)$ for all functions $h \in \mathcal{C}^\infty(N, \mathbb{R})$ and all points $p \in M$.
- (b) If X is f -related to Y and γ is an integral curve of X , then $f \circ \gamma$ is an integral curve of Y .
- (c) If f is a local diffeo, for every vector field $Y \in \mathfrak{X}(N)$ there exists a unique $X \in \mathfrak{X}(M)$ that is f -related to Y . We denote $f^*Y := X$.

Thus if f is a diffeo, then f -relatedness is a bijection from $\mathfrak{X}(M)$ to $\mathfrak{X}(N)$.

In this case, if X is f -related to Y , we write $X = f^*Y$ and $Y = f_*X$.

Finally, a nice property of any vector field X is that its aspect near any point p where $X_p \neq 0$ is quite standard:

Exercise 4.4.9. If X is a smooth vector field on a manifold M and $p \in M$ is a point where $X_p \neq 0$, then there exists a chart (U, ϕ) of M defined at p such that $X|_U = \frac{\partial}{\partial \phi^0}$.

5 Covectors and 1-forms

5.1 Vectors and covectors; superindices and subindices

In differential geometry we often use subindices and superindices as follows.

Consider a vector space $V \simeq \mathbb{R}^n$ and its dual space V^* . (Typically V is the tangent space $T_p M$ of a manifold M .) Let $\mathcal{B} = (E_i)_i$ and $\mathcal{B}^* = (E^j)_j$ be bases of V and V^* that are dual to each other, which means that

$$E^j(E_i) = \delta_i^j \quad \text{for all } i, j.$$

For a vector $X \in V$ we denote X^i its coordinates or components with respect to the base \mathcal{B} , and for a covector $\xi \in V^*$ we denote ξ_j its coordinates with respect to the base \mathcal{B}^* , so that

$$X = \sum_i X^i E_i, \quad \xi = \sum_j \xi_j E^j, \quad \text{and thus} \quad \xi(X) = \sum_k \xi_k X^k.$$

(Note that we always sum over an index that appears exactly twice: once as subindex, once as superindex.) We can obtain the coordinates of X and ξ using the formulas

$$X^i = E^i(X) \quad \text{and} \quad \xi_j = \xi(E_j). \quad (5.1)$$

To do: Picture for $V = \mathbb{R}^2$, $X = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$, $\xi = (1 \ 2)$.

A nonzero covector $\xi \in V^*$ is represented by drawing the hyperplane $\{v \in V : \xi(v) = 1\}$.

5.2 Cotangent vectors

Definition 5.2.1. The **cotangent space** of a differentiable manifold M at a point $p \in M$ is the dual of the tangent space:

$$T_p^* M = (T_p M)^*$$

i.e. its elements (called **cotangent vectors**, or **covectors**) are the linear functions from $T_p M$ to \mathbb{R} .

We can obtain a covector by differentiating a function at a point.

Definition 5.2.2. The **differential** of a function $h : M \rightarrow \mathbb{R}$ at a point $p \in M$ where h is \mathcal{C}^1 is the covector $dh|_p \in T_p^* M$ given by

$$dh|_p(X) = X(h) \quad \text{for all vectors } X \in T_p M.$$

Example 5.2.3 (Differential of chart components). If φ is a chart of M defined at $p \in M$, the differential of a chart component φ^i applied to a coordinate vector $\frac{\partial}{\partial \varphi^j}$ gives

$$d\varphi^i|_p \left(\frac{\partial}{\partial \varphi^j} \Big|_p \right) = \frac{\partial}{\partial \varphi^j} \Big|_p \varphi^i = \delta_j^i. \quad (5.2)$$

This shows that the covectors $d\varphi^i|_p$ form a basis of $T_p^* M$ that is dual to the basis of $T_p M$ consisting of the coordinate vectors $\frac{\partial}{\partial \varphi^j} \Big|_p$.

Definition 5.2.4. The **coefficients** or **components** of a covector $\xi \in T_p^*M$ with respect to the chart φ are the real numbers ξ_i such that

$$\xi = \sum_i \xi_i d\phi^i|_p.$$

These coefficients can be obtained by evaluating ξ on the coordinate vectors:

$$\xi_j = \xi \left(\frac{\partial}{\partial \varphi^j} \Big|_p \right); \quad (5.3)$$

this formula follows from the duality relation (5.2).

Proposition 5.2.5 (Transformation law for covector coefficients). *Let $(\xi^i)_i, (\tilde{\xi}^j)_j \in \mathbb{R}^n$ be the tuples coefficients of a covector $\xi \in T_p^*M$ with respect to two charts $\varphi, \tilde{\varphi}$. These tuples are related by the formula*

$$\tilde{\xi}_j = \sum_i \frac{\partial \varphi^i}{\partial \tilde{\varphi}^j} \Big|_p \xi_i. \quad (5.4)$$

Proof.

$$\begin{aligned} \tilde{\xi}_j &= \xi \left(\frac{\partial}{\partial \tilde{\varphi}^j} \Big|_p \right) = \xi \left(\sum_i \frac{\partial \varphi^i}{\partial \tilde{\varphi}^j} \Big|_p \frac{\partial}{\partial \varphi^i} \Big|_p \right) \\ &= \sum_i \frac{\partial \varphi^i}{\partial \tilde{\varphi}^j} \Big|_p \xi \left(\frac{\partial}{\partial \varphi^i} \Big|_p \right) \\ &= \sum_i \frac{\partial \varphi^i}{\partial \tilde{\varphi}^j} \Big|_p \xi_i. \end{aligned}$$

□

5.3 Covector fields

Definition 5.3.1. The **cotangent bundle** T^*M of a differentiable manifold M is the disjoint union of the cotangent spaces: $T^*M = \coprod_{p \in M} T_p^*M$.

This set has a natural structure of vector bundle over M ...

Definition 5.3.2. A **covector field** or **1-form** on a \mathcal{C}^{k+1} manifold M is a function $\xi : M \rightarrow T^*M$ such that $\xi_p \in T_p^*M$ for all $p \in M$. Its **component functions** with respect to a chart (U, φ) are the functions $\xi_j : U \rightarrow \mathbb{R}$ such that

$$\xi_p = \sum_j \xi_j(p) d\varphi^j|_p \quad \text{for all } p \in U.$$

Note that $\xi_j(p) = \xi|_p \left(\frac{\partial}{\partial \varphi^j} \Big|_p \right)$ for all $p \in U$.

A covector field ξ is \mathcal{C}^k at a point $p \in M$ if its component functions (w.r.t. any chart that is defined at p)¹ are \mathcal{C}^k at the point p , and ξ is \mathcal{C}^k on M if it is \mathcal{C}^k at all points of M .

We denote $\Omega^1(M)$ the set of \mathcal{C}^k covector fields on a \mathcal{C}^{k+1} manifold M . Note that $\Omega^1(M)$ is a module over the ring $\mathcal{C}^k(M)$, i.e. we can obtain a new covector field by adding two covector fields, or by multiplying a covector field $\xi \in \Omega^1(M)$ by a function $h \in \mathcal{C}^k(M)$.

¹It doesn't matter which chart we use; this follows from the transformation law.

5.4 Contraction of a 1-form with a vector field

The **contraction** of a 1-form $\xi \in \Omega^1(M)$ with a vector field $X \in \mathfrak{X}(M)$ is the function $\langle \xi, X \rangle$, also denoted $\iota_X \xi$, defined by

$$\langle \xi, X \rangle(p) = \xi|_p(X|_p) \quad \text{for all } p \in M.$$

Thus the components of ξ w.r.t. a chart ϕ are $\xi_i = \langle \xi, \frac{\partial}{\partial \phi^i} \rangle$.

5.5 Differential of a function

An example of covector field is the differential of a function.

Definition 5.5.1. The **differential** of a function $h \in \mathcal{C}^{k+1}(M)$ is the 1-form $dh \in \Omega^1(M)$ defined by

$$dh|_p(X) = X(h) \quad \text{for all points } p \in M \text{ and all vectors } X \in T_p M.$$

A 1-form is **exact** if it is the differential of some function.

On a chart (U, ϕ) we can express dh as

$$dh|_U = \sum_i \frac{\partial h}{\partial \phi^i} d\phi^i$$

since the components of dh are $dh\left(\frac{\partial}{\partial \phi^i}\right) = \frac{\partial h}{\partial \phi^i}$ by (5.3). This shows that dh is \mathcal{C}^k if h is \mathcal{C}^{k+1} .

Example 5.5.2 (Differential of coordinate functions). If (U, φ) is a chart on M , each coordinate differential $d\varphi^i$ is a \mathcal{C}^k covector field since its component functions $\langle d\varphi^i, \frac{\partial}{\partial \varphi^j} \rangle \equiv \delta_j^i$ are constant.

In the exercises we prove some basic properties of the differential:

- Proposition 5.5.3.** (i) For $f, g \in C^\infty(M)$, $a, b \in \mathbb{R}$ we have: $d(a \cdot f + b \cdot g) = a \cdot df + b \cdot dg$, $d(f \cdot g) = f \cdot dg + g \cdot df$, $d(\frac{f}{g}) = \frac{g \cdot df - f \cdot dg}{g^2}$ (on the set where $g \neq 0$)
- (ii) Chain rule: If $h : \mathbb{R} \rightarrow \mathbb{R}$ is a smooth function then $d(h \circ f) = (h' \circ f) \cdot df$.
- (iii) If $df = 0$ then f is constant on each connected component of M .

5.6 Pullback of a 1-form

Definition 5.6.1. The **pullback** of a covector field $\xi \in \Omega^1(N)$ by a \mathcal{C}^1 map $f : M \rightarrow N$ is the covector field $f^*\xi \in \Omega^1(M)$ defined by

$$(f^*\xi)|_p = (T_p f)^*\xi|_{f(p)} \quad \text{for all } p \in M,$$

that is,

$$(f^*\xi)|_p(X) = \xi(T_p f X) \quad \text{for all } X \in T_p M.$$

Proposition 5.6.2. Pullback commutes with differentiation. That is, for any \mathcal{C}^1 map $f : M \rightarrow N$ and any function $h \in \mathcal{C}^1(N, \mathbb{R})$, we have $f^*(dh) = d(f^*h)$ where $f^*h = h \circ f$.

Also, pullback commutes with the module operations on Ω^1 , i.e. we have

$$f^*(\theta + \xi) = f^*\theta + f^*\xi \quad \text{and} \quad f^*(h \cdot \xi) = (f^*h) \cdot f^*\xi.$$

for covector fields $\xi, \theta \in \Omega^1(N)$ and a function $h \in \mathcal{C}^1(N, \mathbb{R})$.

Proof. To show that the 1-forms $f^*(dh)$ and $d(h \circ f)$ are equal, we apply them to a tangent vector $X \in T_p M$:

$$f^*(dh)(X) = dh(T_p f X) = (T_p f X)h = X(h \circ f) = d(h \circ f)(X).$$

The rest: exercise. □

5.7 Integration of a 1-form along a curve

Definition 5.7.1. The **integral** of a continuous 1-form $\xi \in \Omega^1(M)$ along a \mathcal{C}^1 curve $\gamma : I = [a, b] \rightarrow M$ in a differentiable manifold M is defined as

$$\int_{\gamma} \xi := \int_a^b \xi_{\gamma(t)}(\gamma'(t)) \, dt.$$

Example 5.7.2. Let us compute the integral of the 1-form $\xi = -y \, dx + x \, dy \in \Omega^1(\mathbb{R}^2)$ along the radius r circle $\gamma : t \in [0, 2\pi] \mapsto (r \cos t, r \sin t)$. The pullback of ξ by γ is a 1-form $\gamma^* \xi = f \, d_0$, where

$$\begin{aligned} f(t) &= \gamma^* \xi(\partial_0|_t) = \xi_{\gamma(t)}(\partial_0|_t) \\ &= \xi|_{\gamma(t)}(-r \sin(t) \partial_x + r \cos(t) \partial_y)|_{\gamma(t)} \\ &= (-r \sin(t) \, dx + r \cos(t) \, dy)(-r \sin(t) \partial_x + r \cos(t) \partial_y) \\ &= r^2 \end{aligned}$$

Thus

$$\int_{\gamma} \xi = \int_0^{2\pi} \gamma^* \xi = \int_0^{2\pi} f \, d_0 = \int_0^{2\pi} f(t) \, dt = 2\pi r^2.$$

Theorem 5.7.3 (1-dimensional Stokes). *The integral along a \mathcal{C}^1 curve $\gamma : [a, b] \rightarrow M$ of the differential of a function $h \in \mathcal{C}^1(M, \mathbb{R})$ is*

$$\int_{\gamma} dh = h(\gamma(b)) - h(\gamma(a)).$$

Proof.

$$\begin{aligned} \int_{\gamma} dh &= \int_a^b dh(\gamma'(t)) \, dt \\ &= \int_a^b (h \circ \gamma)'(t) \, dt = h(\gamma(b)) - h(\gamma(a)). \end{aligned}$$

□

In the exercises we prove the following property of integrals of 1-forms.

Proposition 5.7.4 (Reparametrization invariance of curve integrals). *If two \mathcal{C}^1 curves $\gamma : J \rightarrow M$, $\beta : I \rightarrow M$ are equivalent as oriented curves, in the sense that $\beta = \gamma \circ \tau$, for some positive (= increasing) diffeo $\tau : I \rightarrow J$, then*

$$\int_{\beta} \xi = \int_{\gamma} \xi \quad \text{for any 1-form } \xi \in \Omega^1(M).$$

Proof. The integrals we are comparing are

$$\int_{\gamma} \xi = \int_J g(t) \, dt \quad \text{and} \quad \int_{\beta} \xi = \int_I f(s) \, ds$$

where $g(t) = \xi_{\gamma(t)}(\gamma'(t))$ and $f(s) = \xi_{\beta(s)}(\beta'(s))$. Since $\beta = \gamma \circ \tau$, applying the chain rule we get

$$\begin{aligned} f(s) &= \xi_{\beta(s)}(\beta'(s)) = \xi_{\gamma(\tau(s))}((\gamma \circ \tau)'(s)) \\ &= \xi_{\gamma(\tau(s))}(\tau'(s) \cdot \gamma'(\tau(s))) \\ &= \tau'(s) \cdot \xi_{\gamma(\tau(s))}(\gamma'(\tau(s))) \\ &= \tau'(s) \cdot g(\tau(s)). \end{aligned}$$

Thus the two integrals are equal by the change of variables formula. \square

5.8 Integration along curves via pullback

There is an equivalent definition of the integral $\int_{\gamma} \xi$ using the pullback $\gamma^*\xi$.

Note that any 1-form $\theta \in \Omega^1(J)$ defined on an open interval $J \subseteq \mathbb{R}$ can be written as $\theta = \theta_0 d\phi$ where $\theta_0 : J \rightarrow \mathbb{R}$ is a function. This function θ_0 is the single component function of θ w.r.t. the chart ϕ and can be computed as

$$\theta_0 = \langle \phi, \frac{\partial}{\partial \phi} \rangle.$$

Definition 5.8.1. The **integral** of a 1-form $\theta \in \Omega^1(I)$ along a compact interval $I = [a, b] \subseteq \mathbb{R}$ is defined as

$$\int_a^b \theta := \int_a^b \theta_0(t) dt,$$

where $\theta_0 \in \mathcal{C}(I, \mathbb{R})$ is the component function of θ with respect to the standard coordinate $\phi = \text{id}_I$.

Proposition 5.8.2. *The integral of a continuous 1-form $\xi \in \Omega(M)$ over a \mathcal{C}^1 curve $\gamma : I = [a, b] \rightarrow M$ in a differentiable manifold M is*

$$\int_{\gamma} \xi := \int_I \gamma^* \xi.$$

Remark 5.8.3. The concept of 1-form on I is not yet defined because the interval I is not a manifold. (It is a *manifold-with-boundary*, but we have not yet defined this kind of objects.) For the moment we can just consider I as a subset of the manifold \mathbb{R} (or of any open interval I^+ that contains I). A 1-form on I is the restriction of a 1-form defined on \mathbb{R} (or on I^+). We can assume that γ is extended to some open interval $I^+ \supseteq I$, the 1-form $\gamma^*\xi$ can be extended to I^+ , so it is a 1-form on I .

In the exercises we prove that this second definition of $\int_{\gamma} \xi$ is equivalent to the previous one, and use this to give a second proof of the reparametrization invariance of line integrals.