Caterina Daidone, Marco Zanotti

University Milano-Bicocca





- 1. Introduction
- 2. Methods
- 3. Simulation & Results
- 4. Conclusions

## 1. Introduction

1. Introduction

1. Introduction

## 2. Methods


2. Methods

3. Simulation & Results

•00000




7 / 16



kappa = 5; epsilon = 0.05


kappa = 5; epsilon = 0.1


kappa = 5; epsilon = 0.2



Variogram Estimation: kappa = 1; epsilon = 0

000000





process	kappa	epsilon	nugget	sigmasq	phi	robust_nugget	robust_sigmasq	robust_phi
GRF	1	0.00	0.00	0.64	0.15	0.00	0.65	0.16
CGRF	5	0.05	0.00	0.64	0.14	0.00	0.65	0.14
CGRF	5	0.10	0.00	0.67	0.13	0.00	0.69	0.14
CGRF	5	0.20	0.00	0.68	0.12	0.00	0.69	0.13
CGRF	10	0.05	0.08	0.70	0.13	0.00	0.76	0.14
CGRF	10	0.10	0.25	0.68	0.14	0.05	0.84	0.15
CGRF	10	0.20	0.46	0.72	0.14	0.26	0.90	0.16



## 4. Conclusions

4. Conclusions

Cressie N. 1993. Statistics for Spatial Data. John Wiley and Sons Inc., New York

Thank you!

16 / 16