

# Relativistic Quantum Waves (Klein-Gordon Equation

Marcus Allen Denslow

2025-11-14

# Contents

<b>Chapter 1</b>	<b>Deriving the KG Equation</b>	<b>Page 2</b>
1.1	double deriving	2
1.2	Plugging in the new values	2
1.3	this is the Klein-Gordon Equation!	2
1.4	Replacing with Laplacian	3
1.5	d'Alembertian	3
<b>Chapter 2</b>	<b>Four-momentum Eigenstates</b>	<b>Page 4</b>
2.1	Klein-Gordon Plane Wave	4
2.2	Proof that plane waves $\psi = A \exp \left[ -ip \cdot x/\hbar \right]$ satisfy K.G.	4
<b>Chapter 3</b>	<b>Superposition</b>	<b>Page 5</b>
3.1	You can add together many states that solve K.G., and the resulting sum also solves K.G.	5
<b>Chapter 4</b>	<b>Group Velocity, and c Speed limit</b>	<b>Page 6</b>
4.1	calculating the group velocity of a Klein-Gordon Wave packet	6
4.2	Finding the group relation	6

# Chapter 1

## Deriving the KG Equation

### 1.1 double deriving

**Definition 1.1.1: Relativity: the mass shell**

$$p \cdot p = (mc)^2 \rightarrow (mc)^2 = \left(\frac{E}{c}\right)^2 - p_x^2 - p_y^2 - p_z^2$$

**Definition 1.1.2: Quantum: energy and momentum operators**

$$\hat{E} = i\hbar \frac{\partial}{\partial t}, \text{ so } \left(\frac{E}{c}\right)^2 \text{ becomes } -\frac{\hbar^2}{c^2} \frac{\partial^2}{\partial t^2}.$$

$$\hat{p} = -i\hbar \nabla, \text{ so } -p_x^2 \text{ becomes } \hbar^2 \frac{\partial^2}{\partial x^2}.$$

$$\text{likewise, } -p_y^2 \text{ becomes } \hbar^2 \frac{\partial^2}{\partial y^2} \text{ and } -p_z^2 \text{ becomes } \hbar^2 \frac{\partial^2}{\partial z^2}.$$

### 1.2 Plugging in the new values

we can now plugin these into the original equation:

$$\begin{aligned} (mc)^2 &= \left(\frac{E}{c}\right)^2 - p_x^2 - p_y^2 - p_z^2 \\ (mc)^2 &= -\frac{\hbar^2}{c^2} \frac{\partial^2}{\partial t^2} + \hbar^2 \frac{\partial^2}{\partial x^2} + \hbar^2 \frac{\partial^2}{\partial y^2} + \hbar^2 \frac{\partial^2}{\partial z^2} \\ \frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2} - \frac{\partial^2}{\partial z^2} + \left(\frac{mc}{\hbar}\right)^2 &= 0. \end{aligned}$$

### 1.3 this is the Klein-Gordon Equation!

$$\frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2} - \frac{\partial^2}{\partial z^2} + \left(\frac{mc}{\hbar}\right)^2 = 0 \quad (1.1)$$

$$\left[ \frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2} - \frac{\partial^2}{\partial z^2} + \left(\frac{mc}{\hbar}\right)^2 \right] \psi = 0 \quad (1.2)$$

## 1.4 Replacing with Laplacian

We know that  $-\frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2} - \frac{\partial^2}{\partial z^2} = \nabla^2$  -otherwise known as a Laplacian

So the function becomes

$$\left[ \frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \nabla^2 + \left( \frac{mc}{\hbar} \right)^2 \right] \psi = 0 \quad (1.3)$$

## 1.5 d'Alembertian

$\square =$  d'Alembertian

We can also rewrite  $\left( \frac{mc}{\hbar} \right)^2$  as  $\mu^2$

**Note:-**

We could also write  $\left( \frac{mc}{\hbar} \right)$  as just  $m$  as in this universe it would become  $\left( \frac{m \cdot 1}{1} \right)$  which is just the mass but this is the correct way to write it.

So our final equation becomes

$$[\square + \mu^2] \psi = 0 \quad (1.4)$$

## Chapter 2

# Four-momentum Eigenstates

### 2.1 Klein-Gordon Plane Wave

#### Definition 2.1.1: Klein-Gordon Plane Wave function

$$\psi = A \exp \left( -\frac{i}{\hbar} p \cdot x \right)$$

$$p = [E/c, \vec{p}], \quad x = [ct, \vec{x}]$$
$$A \in \mathbb{C}, \quad p^0 = \frac{E}{c} = 0 \pm \sqrt{|\vec{p}|^2 + m^2 c^2}$$

we can rewrite the original Equation as:

$$\psi = A \exp \left( \frac{i}{\hbar} (\vec{p} \cdot \vec{x} - Et) \right)$$
$$\psi = A \exp \left( \frac{i}{\hbar} \left( \vec{p} \cdot \vec{x} \pm c \sqrt{|\vec{p}|^2 + m^2 c^2} t \right) \right)$$

### 2.2 Proof that plane waves $\psi = A \exp \left[ -ip \cdot x / \hbar \right]$ satisfy K.G.

rewrite K.G.:  $[\square + \mu^3] \psi = 0 \rightarrow \square \psi = -\mu^2 \psi$

Does d'Alembertian do the same thing as multiplying by  $-\mu^2$ ?

$$\square = \square \left[ \exp \left[ -\frac{i}{\hbar} p \cdot x \right] \right] = \square \left[ \exp \left[ \frac{i}{\hbar} (\vec{p} \cdot \vec{x} - Et) \right] \right] \quad (2.1)$$

Definition of d'Alembertian:  $\square = \frac{1}{c^2} \frac{\partial}{\partial t^2} - \nabla^2$

$$\square = \left[ -\left( \frac{E}{\hbar c} \right)^2 + \left( \frac{|\vec{p}|}{\hbar} \right)^2 \right] \exp \left[ \frac{i}{\hbar} (\vec{p} \cdot \vec{x} - Et) \right] = -\left( \frac{mc}{\hbar} \right)^2 \psi \quad (2.2)$$

...

## Chapter 3

# Superposition

### 3.1 You can add together many states that solve K.G., and the resulting sum also solves K.G.

Another important concept we have to know about when working with the Klein-Gordon Equation is that any superposition of wave functions that satisfy the Klein-Gordon Equation, also satisfy the Klein-Gordon Equation. That lets us create a complex landscape starting with the simple basis set of functions

Let's say that you have two functions that satisfy the Klein-Gordon Equation, call them  $\psi_1$  and  $\psi_2$

$$[\square + \mu^2] \psi_1 = 0 \quad (3.1)$$

$$[\square + \mu^2] \psi_2 = 0 \quad (3.2)$$

Let's call their sum  $\psi_3$  so

$$\psi_3 = \psi_1 + \psi_2.$$

The question then arises: does  $\psi_3$  satisfy the Klein-Gordon Equation?

Well, yes it does. We can show this with:

$$\begin{aligned} [\square + \mu^2] \psi_3 &= [\square + \mu^2] (\psi_1 + \psi_2) \\ &= [\square + \mu^2] \psi_1 + [\square + \mu^2] \psi_2 = 0 + 0 = 0 \dots \text{also satisfies K.G..} \end{aligned}$$

The following reasoning does not just apply to the sum of two wave functions, but also scaling up a wave function or taking the arbitrary sum over many wave functions. Or even more usefully if you have a basis set of functions that satisfy the Klein-Gordon Equation, then you can make wave functions of that basis set by summing over them with some series of coefficients

Say we write a wavefunction  $\psi$  as a linear combination of  $\{\psi_n\}$

$$\psi = \sum_n C_n \psi_n, \quad C_n \in \mathbb{C} \quad (3.3)$$

Since each basis function in  $\psi_n \in \{\psi_n\}$  satisfies K.G....

$$[\square + \mu^2] \psi_n = 0 \rightarrow [\square + \mu^2] \sum_n C_n \psi_n = 0 \quad (3.4)$$

And that's how we actually use energy and momentum in Eigen-states most of the time.

## Chapter 4

# Group Velocity, and c Speed limit

### 4.1 calculating the group velocity of a Klein-Gordon Wave packet

In order to calculate the group velocity of a Klein-Gordon Wave packet, we first have to calculate something called the **DispersionRelation**.

The **DispersionRelation** is when **angularfrequency**  $\omega$  is written as a function of **angularwavenumber**.

General form of plane wave:

$$\psi = A \exp \left( i \left( \vec{k} \cdot \vec{x} - \omega t \right) \right) \quad (4.1)$$

Klein-Gordon plane wave:

$$\psi = A \exp \left( \frac{i}{\hbar} \left( \vec{p} \cdot \vec{x} \pm c \sqrt{|\vec{p}|^2 + m^2 c^2} t \right) \right) \quad (4.2)$$

$$\begin{aligned} \omega &= \frac{c}{\hbar} \sqrt{|\vec{p}|^2 + m^2 c^2}, & k &= |\vec{p}| / \hbar \\ |\vec{p}|^2 &= \hbar^2 k^2, & \rightarrow \omega &= \sqrt{(kc)^2 + \left( \frac{mc^2}{\hbar} \right)^2} \end{aligned}$$

### 4.2 Finding the group relation

Now we have the Dispersion Relation:  $\omega = \sqrt{(kc)^2 + \left( \frac{mc^2}{\hbar} \right)^2}$

The group velocity is then:  $v_g = \frac{d\omega}{dk}$

$$v_g = \frac{d}{dk} \left[ \sqrt{(kc)^2 + (mc^2/\hbar)^2} \right] \quad (4.3)$$

with  $k = |\vec{p}| / \hbar$  in mind:

$$v_g = \frac{kc}{\sqrt{k^2 + (mc/\hbar)^2}} \rightarrow v_g = \frac{c |\vec{p}|}{\sqrt{|\vec{p}|^2 + (mc)^2}} \quad (4.4)$$

Playing around with the Klein-Gordon equation, we have found a speed limit: the speed of light!