

# Relativistic Quantum Waves (Klein-Gordon Equation

Marcus Allen Denslow

2025-11-14

# Contents

<b>Chapter 1</b>	<b>Deriving the KG Equation</b>	<b>Page 2</b>
1.1	double deriving	2
1.2	Plugging in the new values	2

# Chapter 1

## Deriving the KG Equation

### 1.1 double deriving

**Definition 1.1.1: Relativity: the mass shell**

$$p \cdot p = (mc)^2 \rightarrow (mc)^2 = \left(\frac{E}{c}\right)^2 - p_x^2 - p_y^2 - p_z^2$$

**Definition 1.1.2: Quantum: energy and momentum operators**

$$\hat{E} = i\hbar \frac{\partial}{\partial t}, \text{ so } \left(\frac{E}{c}\right)^2 \text{ becomes } -\frac{\hbar^2}{c^2} \frac{\partial^2}{\partial t^2}.$$

$$\hat{p} = -i\hbar \nabla, \text{ so } -p_x^2 \text{ becomes } \hbar^2 \frac{\partial^2}{\partial x^2}.$$

$$\text{likewise, } -p_y^2 \text{ becomes } \hbar^2 \frac{\partial^2}{\partial y^2} \text{ and } -p_z^2 \text{ becomes } \hbar^2 \frac{\partial^2}{\partial z^2}.$$

### 1.2 Plugging in the new values

we can now plugin these into the original equation:

$$(mc)^2 = \left(\frac{E}{c}\right)^2 - p_x^2 - p_y^2 - p_z^2$$
$$(mc)^2 = -\frac{\hbar^2}{c^2} \frac{\partial^2}{\partial t^2} + \hbar^2 \frac{\partial^2}{\partial x^2} + \hbar^2 \frac{\partial^2}{\partial y^2} + \hbar^2 \frac{\partial^2}{\partial z^2}.$$