

Corregidum to “Novel whitening approaches in functional settings”

Marc Vidal* ^{1,2} and Ana M. Aguilera*²

¹*Ghent University, Belgium*

²*University of Granada, Spain*

1. Due to production errors, Equation 3 in pp. 3 is written as

$$\langle f, g \rangle_{\mathbb{M}} = \sum_{j=1}^{\infty} \lambda_j^{-1} \langle f, \gamma_j \rangle \langle g, \gamma_j \rangle = \left\langle \Gamma^{1/2\dagger} f, \Gamma^{1/2\dagger} g \right\rangle, \quad g \in \mathbb{M},$$

while it was originally written as

$$\langle f, g \rangle_{\mathbb{M}} = \sum_{j=1}^{\infty} \lambda_j^{-1} \langle f, \gamma_j \rangle \langle g, \gamma_j \rangle = \left\langle \Gamma^{1/2\dagger} f, \Gamma^{1/2\dagger} g \right\rangle \quad f, g \in \mathbb{M}.$$

2. In §3, the statement reads, “Then, we can use the inner product (3) to construct a space of isotropic functions (i.e., their covariance operator satisfies the identity), so that the space ends up having a certain Gaussian appearance...” The term *space of isotropic functions* is unclear in the current context. Since our whitening operators are mappings defined through elements over T , this does not necessarily imply that the realizations of \mathbb{X} are on the unit sphere $S = \{f \in \mathbb{M} \mid \|f\|^2 = 1\}$. The isotropy property would be satisfied when whitening the basis expansion coefficients in the direction of its transpose, assuming dependencies in a secondary domain exist. Therefore, one could use the following instead: “...to construct a *space of whitened functions* (i.e., their covariance operator satisfies the identity), so that the space ends up having a certain Gaussian appearance...”.
3. In §4, the sentence “As $2\text{tr}(\Gamma_{X\mathbb{X}})$ is the only dependence between the original and the whitened variable, the minimization problem can be reduced to the maximization of $\text{tr}(\Gamma_{X\mathbb{X}})$.” reads also as “.. is the only *dependent term*...”.
4. Note that, in §4 the term $\text{tr}(\Gamma_{\mathbb{X}})$ in the quadratic distances diverges (the trace of $\Gamma_{\mathbb{X}}$ is an infinite sum of ones). However, $\text{tr}(\Gamma_{\mathbb{X}})$ is not accounted for in the proof. In order for these distances to converge, one has to consider regularization, a weaker norm or finite space dependency. Furthermore, the operator $\Gamma_{X\mathbb{X}}$ coincides with $\Gamma^{1/2}$ if $\Psi \equiv \Gamma^{1/2\dagger}$. Note we only know that this operator belongs to the class of Hilbert Schmidt operators (from the trace property of the autocovariance operator), but this fact does not necessarily imply $\Gamma^{1/2}$ has finite trace. Hence, we further assume that under mild conditions, $\text{tr}(\Gamma^{1/2}) < \infty$ is satisfied.
5. In the Technical proofs (first paragraph), due to abuse of notation, in the sentence “Note that Condition 1 cannot be reached when $\langle X, \gamma_j \rangle^2 = \lambda_j$, or for $c_j \rightarrow c > 0$, $\langle X, \gamma_j \rangle^2 = \lambda_j c_{j\dots}$ ”, X stands for a deterministic function.

*For correspondence: marc.vidalbadia@ugent.be (M.V.), aaguiler@ugr.es (A.M.A)

CHARACTERIZATION OF $\mathcal{R} = \mathcal{V}^{1/2\dagger} \Gamma \mathcal{V}^{1/2\dagger}$

Suppose X is expanded as $X = \sum_{k=1}^{\infty} \langle X, e_k \rangle e_k$ and note the following:

$$\mathcal{V} = \sum_{k=1}^{\infty} \mathcal{P}_{e_k} \Gamma \mathcal{P}_{e_k} = \sum_{k=1}^{\infty} \langle \Gamma e_k, e_k \rangle \mathcal{P}_{e_k} = \sum_{k=1}^{\infty} E(\langle X, e_k \rangle^2) (e_k \otimes e_k) = \sum_{k=1}^{\infty} \eta_k (e_k \otimes e_k). \quad (1)$$

Now, consider the operator

$$\mathcal{R} \equiv E \left(\sum_{j=1}^{\infty} \frac{\langle X, e_j \rangle}{\eta_j^{1/2}} e_j \otimes \frac{\langle X, e_j \rangle}{\eta_j^{1/2}} e_j \right). \quad (2)$$

Observe that $\mathcal{V}^{1/2\dagger} \Gamma \mathcal{V}^{1/2\dagger} = \sum_{j=1}^{\infty} \eta_j^{-1/2} E(\mathcal{P}_{e_j} X \otimes \mathcal{P}_{e_j} X) \eta_j^{-1/2}$, where \mathcal{V}^{\dagger} is the Moore-Penrose inverse of $\mathcal{V} = \sum_{j=1}^{\infty} \eta_j \mathcal{P}_{e_j}$. This shows that $\mathcal{V}^{1/2\dagger} \Gamma \mathcal{V}^{1/2\dagger}$ is equivalent to the operator \mathcal{R} , as defined in [2](#). Note this operator bears resemblance to the classical correlation matrix in the multivariate setting.

REFERENCES

- Vidal, M. and Aguilera, M. (2023). Novel whitening approaches in functional settings. *Stat*, 12(1):e516.
 Vidal, M., Rosso, M., and Aguilera, A. M. (2021). Bi-smoothed functional independent component analysis of EEG data. *Mathematics*, 9:1243.