



Erfahrungen mit einem Computer-gestützten System zur automatischen Überwachung und Erfassung von nosokomialen Infektionen an Intensivstationen – MONI

**Alexander Blacky¹, Walter Koller², Andrea Rappelsberger³,
and Klaus-Peter Adlassnig^{3,4}**

¹ *VAMED-KMB, Accredited Inspection Unit for Sterilisation and Disinfection, Vienna*

² *Department of Hospital Epidemiology and Infection Control, Medical University of Vienna and Vienna General Hospital*

³ *Section for Medical Expert and Knowledge-Based Systems, Center for Medical Statistics, Informatics, and Intelligent Systems, Medical University of Vienna*


⁴ *Medexter Healthcare GmbH, Vienna, Austria*




ICU Wards at the Vienna General Hospital

- 13B1
- 13B2
- 13C1
- 13C2
- 13C3
- 13H1
- 13H3
- 13I1
- 13I2
- 13I3
- transplant ICU
- neurosurgery ICU
- NICU 9C
- NICU E10
- NICU E12





↑ ↑ ↓ ↓





Hospital in Europe
Link for Infection Control through
Surveillance

Surveillance of Nosocomial Infections in Intensive Care Units

Protocol

Version 6.1

(Based on Version 5.0 including technical amendments)

SEPTEMBER 2004



Project commissioned by the EC / DG SANCO/ F/ 4
Agreement Reference number: VS/1999/5235 (99CVF4-025)

ICU_v6_1

Page 1 of 51



TECHNICAL DOCUMENT

Surveillance of surgical site infections in European hospitals – HAISSI protocol

Protocol version 1.02

www.ecdc.europa.eu

INFECTION SITE: Symptomatic urinary tract infection

CODE: UTI-SUTI

DEFINITION: A symptomatic urinary tract infection must meet at least one of the following criteria:

Criterion 1: Patient has at least *one* of the following signs or symptoms with no other recognized cause: fever ($>38^{\circ}\text{C}$), urgency, frequency, dysuria, or suprapubic tenderness *and* patient has a positive urine culture, that is, $\geq 10^5$ microorganisms per cm^3 of urine with no more than two species of microorganisms.

Criterion 2: Patient has at least *two* of the following signs or symptoms with no other recognized cause: fever ($>38^{\circ}\text{C}$), urgency, frequency, dysuria, or suprapubic tenderness *and* at least *one* of the following:

- positive dipstick for leukocyte esterase and/or nitrate
- pyuria (urine specimen with ≥ 10 wbc/mm^3 or ≥ 3 $\text{wbc}/\text{high power field}$ of unspun urine)
- organisms seen on Gram stain of unspun urine
- at least *two* urine cultures with repeated isolation of the same

uropathogen (gram-negative bacteria or *S. saprophyticus*) with $\geq 10^2$ colonies/ml in nonvoided specimens

- $\leq 10^5$ colonies/ml of a single uropathogen (gram-negative bacteria or *S. saprophyticus*) in a patient being treated with an effective antimicrobial agent for a urinary tract infection
- physician diagnosis of a urinary tract infection
- physician institutes appropriate therapy for a urinary tract infection.

Criterion 3: Patient ≤ 1 year of age has at least *one* of the following signs or symptoms with no other recognized cause: fever ($>38^{\circ}\text{C}$), hypothermia ($<37^{\circ}\text{C}$), apnea, bradycardia, dysuria, lethargy, or vomiting *and* patient has a positive urine culture, that is, $\geq 10^5$ microorganisms per cm^3 of urine with no more than two species of microorganisms.

Criterion 4: Patient ≤ 1 year of age has at least *one* of the following signs or symptoms with no other recognized cause: fever ($>38^{\circ}\text{C}$), hypothermia ($<37^{\circ}\text{C}$), apnea, bradycardia, dysuria, lethargy, or vomiting *and* at least *one* of the following:

- positive dipstick for leukocyte esterase and/or nitrate
- pyuria (urine specimen with ≥ 10

wbc/mm^3 or >3 $\text{wbc}/\text{high power field}$ of unspun urine)

- organisms seen on gram stain or unspun urine
- at least *two* urine cultures with repeated isolation of the same uropathogen (gram-negative bacteria or *S. saprophyticus*) with $\geq 10^2$ colonies/ml in nonvoided specimens
- $\leq 10^5$ colonies/ml of a single uropathogen (gram-negative bacteria or *S. saprophyticus*) in a patient being treated with an effective antimicrobial agent for a urinary tract infection
- physician diagnosis of a urinary tract infection
- physician institutes appropriate therapy for a urinary tract infection.

COMMENTS:

- A positive culture of a urinary catheter tip is *not* an acceptable laboratory test to diagnose a urinary tract infection.
- Urine cultures must be obtained using appropriate technique, such as clean catch collection or catheterization.
- In infants, a urine culture should be obtained by bladder catheterization or suprapubic aspiration; a positive urine culture from a bag specimen is unreliable and should be confirmed by a specimen aseptically obtained by catheterization or suprapubic aspiration.

Bloodstream Infections (BSI)

CODE: BSI

BSI-A:

- 1 positive blood culture for a recognised pathogen

or

- Patient has at least one of the following signs or symptoms: fever ($>38^{\circ}\text{C}.$), chills, or hypotension and 2 positive blood cultures for a common skin contaminant (from 2 separate blood samples drawn within 48 hours).

skin contaminants = coagulase-negative staphylococci, *Micrococcus sp.*, *Propionibacterium acnes*, *Bacillus sp.*, *Corynebacterium sp.*

BSI-B: Patient has at least one of the following signs or symptoms: fever ($>38^{\circ}\text{C}.$), chills, or hypotension

And either

- 1 positive blood culture with a skin contaminant in patient with an intravascular line in place and in whom the physician instituted appropriate antimicrobial therapy.

or

- positive blood Antigen test (e.g. *H.influenzae*, *S.pneumoniae*, *N. meningitidis* or Group B *Streptococcus*)

Comment:

BSI-A is the definition used by the majority of NI surveillance networks in Europe. BSI-B extends this definition to the CDC definition of laboratory-confirmed bloodstream infection. Networks should specify in the network data (table icu_net, see 6.3.1) whether only BSI A or both BSI B and BSI A are included in the surveillance (i.e. networks using CDC definition of laboratory confirmed bloodstream infection [$\text{CDC}_{\text{LCBI}} = \text{BSI-A+B}$]). If this is the case, then BSI A and BSI B categories should be specified in the data collection.

• bloodstream infection with

- recognized pathogen
- clinical signs and growth of same skin contaminant from two separate blood samples
- clinical signs and growth skin contaminant from blood and intravascular line in place and AB Therapy
- clinical signs and positive antigen test from blood

Bloodstream infection with clinical signs and growth of same skin contaminant from two separate blood samples

- Patient has at least one of the following signs or symptoms: fever ($>38^{\circ}\text{C}.$), chills, or hypotension and 2 positive blood cultures for a common skin contaminant (from 2 separate blood samples drawn within 48 hours).

skin contaminants = coagulase-negative staphylococci, *Micrococcus sp.*, *Propionibacterium acnes*, *Bacillus sp.*, *Corynebacterium sp.*



BSI-A2

1
←

clinical_signs_of_BSI (t-1d, t, t+1d)

^

same_skin_contaminant_from_two_separate_blood_samples

Decomposition—clinical signs

clinical_signs_of_BSI (t-1d, t, t+1d)[yesterday, today, tomorrow]

=

clinical_signs_of_BSI (t-1d)

=

∨

clinical_signs_of_BSI (t)

=

∨

clinical_signs_of_BSI (t+1d)

=

fever (t-1d)

∨

hypotension (t-1d)

∨

leucopenia (t-1d)

∨

leucocytosis (t-1d)

∨

CRP increased (t-1d)

fever (t)

∨

hypotension (t)

∨

leucopenia (t)

∨

leucocytosis (t)

∨

CRP increased (t)

fever (t+1d)

∨

hypotension (t+1d)

∨

leucopenia (t+1d)

∨

leucocytosis (t+1d)

∨

CRP increased (t+1d)

Clinical signs—fever


fever (t-1d) \Leftarrow ...

fever (t) \Leftarrow

body temperature \uparrow

\vee

thermoregulation applied




fever (t+1d) \Leftarrow ...

Clinical Signs—CRP Increased

CRP increased (t-1d) \Leftarrow ...

CRP increased (t) \Leftarrow



data import

intensive care unit

[maximum value
of the day
e.g., 5 mg/dl]

CRP increased (t+1d) \Leftarrow ...

Decomposition—skin contaminant

same_skin_contaminant_from
two_separate_blood_samples ⇐

first blood culture

- coagulase-negative staphylococci
- *Micrococcus* sp.
- *Propionibacterium acnes*
- *Bacillus* sp.
- *Corynebacterium* sp.

^ (within 48 hours)

data import
microbiology ←

second blood culture

- coagulase-negative staphylococci
- *Micrococcus* sp.
- *Propionibacterium acnes*
- *Bacillus* sp.
- *Corynebacterium* sp.

Cockpit Surveillance at the infection control unit

Moni-IV - Surveillance

Patientendaten Surveillance Ausgabe Hilfe

Surveillance

☒ retrospektiv ☐ aktuell Datum: 2005-05-02 anzeigen ☐ Diagnosen ☐ aktuelle Param. ☒ alle Parameter drucken

Patienten

- 13B1
- 13B3
- 13C1
- 13C2
- 13H1
- 13H3
- 13I1
- 13I2
- 13I3
- Neurochir. ICU
- Transplant ICU

Parameter

Catheter-Associated Symptomatic Urinary Tract Infection completely fulfilled (100%)

Moni-IV - [Surveillance]

Patientendaten Surveillance Ausgabe Hilfe

☒ retrospektiv ☐ aktuell Datum: 2005-06-14 anzeigen

☐ Diagnosen ☐ aktuelle Param. ☒ alle Parameter drucken

Patienten

[2005-05-17 -]

- (2005-06-06 - 2005-06-08) (keine Mibil)
- (2005-06-01 - 2005-06-05) (keine Mibil)
- (2005-06-10 - 2005-06-12) (keine Mibil)
- (2005-06-10 - 2005-06-11) (keine Mibil)
- (2005-06-08 - 2005-06-09) (keine Mibil)
- (2005-06-03 - 2005-06-05) (keine Mibil)
- (2005-06-06 - 2005-06-07) (keine Mibil)
- (2005-06-06 - 2005-06-08) (keine Mibil)
- (2005-06-12 - 2005-06-13) (keine Mibil)
- (2005-06-13 -) (keine Mibil)
- (2004-06-09 -)
- (2004-06-21 -) (keine Mibil)
- (2005-06-03 - 2005-06-04) (keine Mibil)
- (2005-06-02 - 2005-06-03) (keine Mibil)
- (2005-06-02 - 2005-06-03) (keine Mibil)
- (2005-06-01 - 2005-06-03) (keine Mibil)
- (2005-04-18 - 2005-06-07)
- (2005-06-13 -) (keine Mibil)
- [2005-05-17 -]**
- (2005-03-23 - 2005-06-01) (Aufenthalt unterbr.!)
- (2005-05-27 - 2005-06-06) (keine Mibil)
- + Transplant ICU**

Parameter

- 2005-06-05 (NCHINT)
- 2005-06-04 (NCHINT)
- 2005-06-03 (NCHINT)
- 2005-06-02 (NCHINT)
- 2005-06-01 (NCHINT)
- Messwerte
- Interpretationen und Diagnosen
 - UTI-B-k:** 100 %
 - KiSeps (t-1d - t+1d): 100 %
 - KiSeps (t-1d): 80 % (2005-06-15 10:42:46)
 - KiSeps: 100 %
 - KiSeps (t+1d): 80 % (2005-06-15 10:42:46)
 - KiPneu (t-1d - t+1d): 47 % (2005-06-15 10:42:46)
 - KiPneu (t-1d): 47 % (2005-06-15 10:42:46)
 - KiHwI:** t-1d - t+1d: 100 %
 - KiHwI (t-1d): 80 % (2005-06-15 10:42:46)
 - KiHwI: 100 %
 - KiHwI (t+1d): 80 % (2005-06-15 10:42:46)
 - andere HwI-Befunde: 100 %
 - Pyurie:** t-1d - t+1d: 100 %
 - Harnkatheter:** t-2d - t: 100 %
 - Harnkatheter: 100 %
 - ZVK (t-2d - t): 100 %
 - ZVK: 100 %
 - Beatmung: 100 %
 - Schockindex (t-1d) - Schock: 92 % (2005-06-15 10:42:46)
 - Fieber:** t-1d: 80 % (2005-06-15 10:42:46)
 - Fieber (t+1d): 80 % (2005-06-15 10:42:46)
 - Thermoregulation (t-1d): 100 %
 - Thermoregulation (t+1d): 100 %
 - CRP:** 100 % (2005-06-15 10:42:46)
 - Hypotonie (t-1d): 80 % (2005-06-15 10:42:46)
 - Adaptierter-Murray-Score-Erhöhung (t-1d): 47 % (2005-06-15 10:42:46)
- 2005-05-31 (NCHINT)
- 2005-05-30 (NCHINT)

Backtracking of the Logical Chain of Reasoning patient has urinary catheter

Moni-IV

Patientendaten Surveillance Ausgabe Hilfe

zurück vor Erklärung

UTI-B-k 2005-06-01: [REDACTED] (17.5.2005 -): NCHINT

Regel: H3: UTI-B-k (kathassoz. sympt. Harnwegsinfektion) (Gewicht: 100)

Bezeichnung	Zutreffen %	Herkunft	ermittelt	Bemerkung
UTI-B-k	100	Moni-IV: Inferenz	2005-06-14 09:39:08	

Bedingungen:

Bezeichnung	Zutreffen %	Herkunft	ermittelt	Bemerkung
UND	100	Moni-IV: Inferenz	2005-06-14 09:39:08	
HarnKath (t-2d - t)	100	Import: CareVue	2005-06-14 09:17:43	2005-05-30 00:00:00 - Harnkatheter 1 Ch/Material 14 Silikon(2005-05-30 06:00:00 - H...
KH-Hwl (t-1d - t+1d)	100	Moni-IV: Inferenz	2005-06-14 09:39:08	
andere Hwl-Befunde	100	Moni-IV: Inferenz	2005-06-14 09:39:08	

Elevated CRP as a Clinical Sign present (100%)

Moni-IV Patientendaten Surveillance Ausgabe Hilfe

KliHWI 2005-06-01: [REDACTED] (17.5.2005 -); NCHINT

Regel: KliHWI: KliHWI (Gewicht: 100)

Bezeichnung	Zutreffen %	Herkunft	ermittelt	Bemerkung
KliHWI	100	Moni-IV: Inferenz	2005-06-14 09:39:08	

Bedingungen:

Bezeichnung	Zutreffen %	Herkunft	ermittelt	Bemerkung
ODER	100	Moni-IV: Inferenz	2005-06-14 09:39:08	
Fieber				
CRPEth	100	Moni-IV: Daten-Symbo...	2005-06-14 09:39:08	

Elevated CRP is Present

6 mg/dl is measured

Moni-IV

Patientendaten Surveillance Ausgabe Hilfe

zurück vpr Erklärung

CRP 2005-06-01: [REDACTED] (17.5.2005 -); NCHINT

Daten-Symbol-Konversion: Fuzzy-Set: CRPErh

Bezeichnung	Zutreffen %	Herkunft	ermittelt	Bemerkung
CRP	100	Moni-IV: Daten-Symbo...	2005-06-14 09:39:08	

quantitativer Wert:

Bezeichnung	Wert	Herkunft	ermittelt	Bemerkung
CRP	6 mg/dl	Import: CareVue	2005-06-14 09:17:43	2005-06-01 07:00:00 Serum: CRP: 6 mg/dl

Other Signs of Urinary Tract Infection

Pyurie

Moni-IV

Patientendaten

Surveillance

Ausgabe

Hilfe

zurück

vor

Erklärung

andere HWI-Befunde

2005-06-01: [REDACTED] (17.5.2005 -); NCHINT

Regel: HWIBef: andere HWI-Befunde (Gewicht: 100)

Bezeichnung	Zutreffen %	Herkunft	ermittelt	Bemerkung
andere HWI-Befunde	100	Moni-IV: Inferenz	2005-06-14 09:39:08	


Bedingungen:

Bezeichnung	Zutreffen %	Herkunft	ermittelt	Bemerkung
ODER	100	Moni-IV: Inferenz	2005-06-14 09:39:08	
Hamteststreifen				
Pyurie	100	Import: CareVue	2005-06-14 09:17:43	2005-06-01 06:00:00 - Ham: U-Ery/Leuko: 10 2005-06-02 06:00:00 - Ham: U-Ery/Leuko: 25
Gram-Färbung				
2HaKw>10 ²				
HaKw<10 ⁵				

Neues MONI : MONI ICU und MONI N-ICU „The Next Generation“

von 2008-04-01 bis 2008-05-04 Anzeigen Standardansicht Grafik ausblenden fixe Tagesbreite

Dep. 21



Stationen/Patienten

Dep. 21

Pat.	3623438
Pat.	3623446
Pat.	3444535
Pat.	3623434
Pat.	3444533
Pat.	3623444
Pat.	3623432



Berechnung



Surveillance



Übersicht



Statistik



Verwaltung




Hilfe



Abmelden

von 2008-05-01 bis 2008-05-05 Anzeigen Standardansicht Grafik ausblenden fixe Tabellenbreite

13c3



- Aufenthalt
- 10% - 50%
- 50% - 90%
- 90% - 100%

- Stationen/Patienten
- 13b1
 - 13b3
 - 13c1
 - 13c2
 - 13c3
-
- 13h1
 - 13h3
 - 13i1

2008-05-10	
2008-05-09	
2008-05-08	
2008-05-07	
2008-05-06	
2008-05-05	
2008-05-04	
2008-05-03	
2008-05-02	
2008-05-01	
Messwerte	
Interpretationen und Diagnosen	
CRI2 (generalisierte ZVK-assoz. Infektion)	80 %
Fieber	80 %
Hypotonie	80 %
klin. Anzeichen für Entzündung bei HWI	80 %
klin. Anzeichen für Entzündung bei Sepsis	80 %
CRP erhöht	37 %
Schock	100 %
max. Körpertemperatur	37,4 °C
Blutdruck-Profil (unterer Grenzwert)	86 mmHg
Blutdruck-Profil (oberer Grenzwert)	88,33 mmHg
Blutdruck-Profil (21:00 (Vortag) bis 3:00)	86,11 mmHg
Blutdruck-Profil (0:00 bis 6:00)	85,89 mmHg
Blutdruck-Profil (3:00 bis 9:00)	89,67 mmHg
Blutdruck-Profil (6:00 bis 12:00)	93,83 mmHg

CRI2 (generalisierte ZVK-assoz. Infektion)	
UND	80 %
NICHT	100 %
pos. Blutkultur	
klin. Anzeichen für Entzündung bei Sepsis	80 %
ODER	100 %
quant. Kultur Katheterspitze	100 %
semiquant. Kultur Katheterspitze	100 %
klin. Anzeichen für Entzündung bei Sepsis	
ODER	80 %
Fieber	80 %
Hypotonie	80 %
Leukopenie	
Leukocytose	
CRP erhöht	37 %
Hypotonie	
ODER	100 %
Blutdruck-Abfall	
Schock	100 %
Schock	
Schockindex	0,8



Berechnung



Surveillance



Übersicht



Statistik



Verwaltung



Hilfe



Abmelden

von 2008-05-01 bis 2008-05-05 Anzeigen Standardansicht Grafik einblenden fixe Tabellenbreite

Stationen/Patienten

13b1
13b3
13c1
13c2
13c3
13h1
13h3
13i1
NICU_E10



NICU_E9

2008-05-01

Messwerte

Interpretationen und Diagnosen

lab: Interleukin 8: Min. 262 mg/dl
lab: Glukose: Max. 226 mg/dl
BSI-1 (KISS) 100 %
BSI-1 (Alert) 100 %
1 klin. Anz. f. Sepsis (KISS) 100 %
1 klin. Anz. f. Sepsis (Alert) 26 %
2 Labor- und klin. Anz. f. Sepsis (KISS) 100 %
2 Labor- und klin. Anz. f. Sepsis (Alert) 100 %
neue Hyperglykämie (KISS) 100 %
neue Hyperglykämie (Alert) 26 %
laborchem. Zeichen für Pneumonie (KISS) 100 %
laborchem. Zeichen für Pneumonie (Alert) 100 %
laborchem. Zeichen für Entzündung (KISS) 100 %
laborchem. Zeichen für Entzündung (Alert) 100 %
Interleukin 8 erhöht 100 %
Hyperglykämie (KISS) 100 %
Hyperglykämie (Alert) 26 %

BSI-1 (KISS)

UND
mibi: pos. Blutku., Liquor: keine KNS 100 %
2 Labor- und klin. Anz. f. Sepsis (KISS) 100 %

2 Labor- und klin. Anz. f. Sepsis (KISS)

MIND 2 100 %
path. Körpertemp.
path. Herzfrequenz (Definition)
metabolische Azidose (KISS)
neue Hyperglykämie (KISS) 100 %
laborchem. Zeichen für Entzündung (KISS) 100 %

laborchem. Zeichen für Entzündung (KISS)

Interleukin 8 erhöht 100 %

Interleukin 8 erhöht

lab: Interleukin 8: Min. 262 mg/dl

lab: Interleukin 8: Min.

lab: Interleukin 8 262 mg/dl

lab: Interleukin 8 262 mg/dl

lab: Interleukin 8 262 mg/dl

MONI: Evaluierung

2 ICUs:	n	without HCAI	with HCAI
patients	89		
ICU stays	93	75	18*
LOS (days)	median:7 shortest:1 longest:40		
patient days	1005	918	88

* comprising 30 HCAI episodes: 12 with one, 1 with two, 4 with three and 1 with four distinct infection episodes during one stay

HCAI condition correctly or falsely identified by MONI and by the human expert

MONI reported:	HCAIs present	HCAIs absent
HCAIs present	26 (86.7%)	1* (1.3%)
HCAIs absent	4** (13.3%)	75 (98.7%)
human expert reported:		
HCAI present	12 (40%)	5*** (6.4%)
HCAI absent	18**** (60%)	73 (93.6%)

* A CRI episodes was falsely detected. The underlying cause was elevated CRI as a result of leukemia.

** 3 PN episodes and 1 CRI episode not reported due to missing import of electronic microbiology report

*** human expert misinterpreted or wrongly reported 4 UTI episodes and 1 BSI episode.

**** human expert missed 3 UTI episodes, 3 BSI episodes, 11 CRI episodes and 1 PN episode


Surveillance - Zeitbedarf

2 ICUs

alle ICUs

	Conventional surveillance	MONI-ICU surveillance
Time spent	82.5 h (100%)	12.5 h (15.2%)

**Erfolge
bei der
Infektions-
prävention**



**Aufwand
für die
Surveillance**



Herzlichen Dank!