

## Exercise 8.1

(a) Linearise around  $(0, 0)$ :

$$A := \left. \frac{df}{dx} \right|_{x=(0,0)} = \begin{pmatrix} 4x_1^3 + x_2 & x_1 \\ -2x_1 + x_2^2 & -2 + 2x_1x_2 \end{pmatrix} \Big|_{x=(0,0)} = \begin{pmatrix} 0 & 0 \\ 0 & -2 \end{pmatrix}.$$

We see that the eigenvalues are  $\lambda_1 = 0$  and  $\lambda_2 = -2$ . As  $\Re(\lambda_1) = 0$ , the fixed point  $(0, 0)$  is not hyperbolic and thus the theorem of Poincaré-Lyapunov is not applicable.

(b) The center subspace  $E^C(0, 0)$  can be obtained by

$$(A - \lambda_1 E)x = 0 \iff Ax = 0 \iff -2x_2 = 0 \iff x_2 = 0.$$

Therefore  $E^C(0, 0) = \{(x_1, x_2) \in \mathbb{R}^2 : x_2 = 0\} = \text{span}((1, 0))$ .

Idea:  $W^C(0, 0) = \{(x_1, h(x_1)) =: H(x_1) : |x_1| \leq \epsilon\}$ . Due to  $\Phi^t(W^C(0, 0)) \subset W^C(0, 0)$ , it follows

$$f(H(x_1)) \in T_{H(x_1)}W^C(0, 0) \implies \langle f(H(x_1)), N(x_1) \rangle = 0$$

with  $N(x_1) = \begin{pmatrix} 1 \\ h'(x_1) \end{pmatrix}^\perp = \begin{pmatrix} h'(x_1) \\ -1 \end{pmatrix}$ . So we obtain

$$\left\langle \begin{pmatrix} x_1^4 + x_1 h(x_1) \\ -2h(x_1) - x_1^2 + x_1 h^2(x_1) \end{pmatrix}, \begin{pmatrix} h'(x_1) \\ -1 \end{pmatrix} \right\rangle = 0. \quad (1)$$

Now we expand  $h(x_1)$  by Taylor Expansion up to degree 5:

$$\begin{aligned} h(x_1) &= h(0) + h'(0)x_1 + \frac{1}{2}h''(0)x_1^2 + \frac{1}{6}h^{(3)}(0)x_1^3 + \frac{1}{24}h^{(4)}(0)x_1^4 + O(x_1^5) \\ &= \frac{1}{2}ax_1^2 + \frac{1}{6}bx_1^3 + \frac{1}{24}cx_1^4 + O(x_1^5), \end{aligned}$$

where  $a, b, c$  are yet to be determined. Derivating the local approximation of  $h(x_1)$  yields

$$h'(x_1) = ax_1 + \frac{1}{2}bx_1^2 + \frac{1}{6}cx_1^3 + O(x_1^4)$$

Substituting in (1) gives

$$\left\langle \begin{pmatrix} x_1^4 + x_1 h(x_1) \\ -2h(x_1) - x_1^2 + x_1 h^2(x_1) \end{pmatrix}, \begin{pmatrix} ax_1 + \frac{1}{2}bx_1^2 + \frac{1}{6}cx_1^3 + O(x_1^4) \\ -1 \end{pmatrix} \right\rangle = 0$$

The first row gives a term  $O(x^5)$  and therefore

$$2h(x_1) + x_1^2 - \underbrace{x_1 h^2(x_1)}_{=O(x^5)} + O(x^5) = 0 \iff ax_1^2 + \frac{1}{3}bx_1^3 + \frac{1}{12}cx_1^4 + x_1^2 + O(x^5) = 0.$$

Now we get


$$0 = (a+1)x_1^2 + \frac{1}{3}bx_1^3 + \frac{1}{12}cx_1^4 \implies a = -1, b = c = 0.$$

Therefore,  $h(x_1) = -\frac{1}{2}x_1^2$  and

$$W^C(0, 0) = \left\{ \begin{pmatrix} 2x_1 \\ -x_1^2 \end{pmatrix} : |x_1| \leq \epsilon \right\}.$$

(c) Restricting the dynamical system locally on  $W^C(0,0)$  results in

$$\dot{x}_1 = x_1^4 + x_1(-0.5x_1^2) \iff \dot{x}_1 = x_1^4 - 0.5x_1^3.$$



We see that the fixed point  $(0,0)$  is stable. For positive  $x_1$  values and  $x_1 < 0.5$ ,  $\dot{x}_1$  is negative, which results in convergence to 0. Similarly, for negative  $x_1$  values,  $\dot{x}_1$  is positive, which results in convergence to 0.

## Exercise 8.2