

Bayesian learning




Observed Data: $\{x^{(i)}\}_{i=1}^n$

Parameter: $\boldsymbol{w} = \{\mu \in \mathbb{R}, \sigma^2 \in \mathbb{R}_{++}\}$

Model distribution: $p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w}) = \prod_{i=1}^n p(\boldsymbol{x}^{(i)}|\boldsymbol{w})$

Prior distribution: $p(w) \propto 1$






Observed Data: $\{x^{(i)}\}_{i=1}^n$ Parameter: $w = \{\mu \in \mathbb{R}, \sigma^2 \in \mathbb{R}_{++}\}$

Model distribution: $p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w}) = \prod p(\boldsymbol{x}^{(i)}|\boldsymbol{w})$

Prior distribution: $p(w) \propto 1$

Joint distribution: $p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w}) = p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n | \boldsymbol{w}) p(\boldsymbol{w})$

observed to be estimated!



We want to estimate parameter from observed data.



Observed Data: $\{x^{(i)}\}_{i=1}^n$ Parameter: $w = \{\mu \in \mathbb{R}, \sigma^2 \in \mathbb{R}_{++}\}$

Model distribution: $p(\{\boldsymbol{x}^{(i)}\}_{i=1}^{n}|\boldsymbol{w}) = \prod p(\boldsymbol{x}^{(i)}|\boldsymbol{w})$

Prior distribution: $p(w) \propto 1$

Joint distribution: $p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w}) = p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n | \boldsymbol{w}) p(\boldsymbol{w})$

observed to be estimated!

$$p(b|a)p(a) = p(a,b) = p(a|b)p(b)$$

$$a o \{\boldsymbol{x}^{(i)}\}_{i=1}^n,$$

 $b \rightarrow w$



Observed Data: $\{x^{(i)}\}_{i=1}^n$ Parameter: $w = \{\mu \in \mathbb{R}, \sigma^2 \in \mathbb{R}_{++}\}$

Model distribution: $p(\{\boldsymbol{x}^{(i)}\}_{i=1}^{n}|\boldsymbol{w}) = \prod p(\boldsymbol{x}^{(i)}|\boldsymbol{w})$

Prior distribution: $p(w) \propto 1$


Joint distribution: $p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w}) = p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n | \boldsymbol{w}) p(\boldsymbol{w})$



$$p(b|a)p(a) = p(a,b) = p(a|b)p(b)$$

$$a \to \{\boldsymbol{x}^{(i)}\}_{i=1}^n, \\ b \to \boldsymbol{w}$$

$$p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w}) = p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})$$





Observed Data: $\{x^{(i)}\}_{i=1}^n$ Parameter: $w = \{\mu \in \mathbb{R}, \sigma^2 \in \mathbb{R}_{++}\}$

Model distribution: $p(\{x^{(i)}\}_{i=1}^n|w) = \prod p(x^{(i)}|w)$

Prior distribution: $p(w) \propto 1$

Joint distribution: $p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w}) = p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n | \boldsymbol{w}) p(\boldsymbol{w})$


observed to be estimated!

$$p(b|a)p(a) = p(a,b) = p(a|b)p(b)$$

$$a \to \{\boldsymbol{x}^{(i)}\}_{i=1}^n,$$

 $b \to \boldsymbol{w}$

$$p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w})}_{\text{joint}} = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})}_{\text{likelihood}}$$





Observed Data: $\{x^{(i)}\}_{i=1}^n$ Parameter: $w = \{\mu \in \mathbb{R}, \sigma^2 \in \mathbb{R}_{++}\}$

Model distribution: $p(\{x^{(i)}\}_{i=1}^n|w) = \prod p(x^{(i)}|w)$

Prior distribution: $p(w) \propto 1$

Joint distribution: $p(\{x^{(i)}\}_{i=1}^n, w) = p(\{x^{(i)}\}_{i=1}^n | w) p(w)$

observed to be estimated!

$$p(b|a)p(a) = p(a,b) = p(a|b)p(b)$$

$$a \to \{\boldsymbol{x}^{(i)}\}_{i=1}^n, \\ b \to \boldsymbol{w}$$

 $\{x_i\}_{i=1}^n$

$$p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w})} = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w})}_{\text{posterior}} = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w})}_{\text{point}} = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i$$

must compute

given



$$p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w})}_{\text{joint}} = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})}_{\text{likelihood}}$$

Posterior distribution:

marginal likelihood:



$$p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w})}_{\text{joint}} = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})}_{\text{likelihood}}$$

Posterior distribution:
$$p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \frac{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})}{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n)}$$

marginal likelihood:



$$p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w})}_{\text{posterior}} = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w})}_{\text{point}} = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})}_{\text{point}} \underbrace{p(\boldsymbol{w})}_{\text{prior}}$$

Posterior distribution:
$$p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \frac{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})}{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n)}$$

marginal likelihood:

constant wrt w!



$$p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w})}_{\text{joint}} = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})}_{\text{likelihood prior}}$$

$$\begin{aligned} \text{Posterior distribution:} \quad & p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \frac{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})}{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n)} \\ \text{marginal likelihood:} \quad & p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \int p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})d\boldsymbol{w} \end{aligned}$$



$$p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w})}_{\text{joint}} = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})}_{\text{likelihood}} \underbrace{p(\boldsymbol{w})}_{\text{prior}}$$

Posterior distribution:
$$p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \frac{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})}{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n)}$$
 marginal likelihood:
$$p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \int p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})d\boldsymbol{w}$$
 Bayes estimator:
$$\widehat{\boldsymbol{w}} = \int \boldsymbol{w} \cdot p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)d\boldsymbol{w} = \langle \boldsymbol{w} \rangle_{p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)}$$
 posterior

Want to know the parameter $m{w}$ estimated from observation $\{m{x}^{(i)}\}_{i=1}^n$.



$$p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w})}_{\text{joint}} = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})}_{\text{likelihood}} \underbrace{p(\boldsymbol{w})}_{\text{prior}}$$

Posterior distribution:
$$p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \frac{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})}{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n)}$$
 marginal likelihood:
$$p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \int p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})d\boldsymbol{w}$$
 Bayes estimator:
$$\widehat{\boldsymbol{w}} = \int \boldsymbol{w} \cdot p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)d\boldsymbol{w} = \langle \boldsymbol{w} \rangle_{p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)}$$

Predictive distribution: $p(\boldsymbol{x}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \int p(\boldsymbol{x}, \boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n) d\boldsymbol{w}$

Want to predict a future x estimated from past observation $\{x^{(i)}\}_{i=1}^n$.



$$p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w})}_{\text{joint}} = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})}_{\text{likelihood}} \underbrace{p(\boldsymbol{w})}_{\text{prior}}$$



$$p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w})}_{\text{joint}} = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})}_{\text{likelihood}} \underbrace{p(\boldsymbol{w})}_{\text{prior}}$$

Posterior distribution:
$$p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \frac{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})}{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n)}$$
 marginal likelihood:
$$p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \int p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})d\boldsymbol{w}$$
 Bayes estimator:
$$\widehat{\boldsymbol{w}} = \int \boldsymbol{w} \cdot p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)d\boldsymbol{w} = \langle \boldsymbol{w} \rangle_{p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)}$$
 Predictive distribution:
$$p(\boldsymbol{x}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \int p(\boldsymbol{x},\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)d\boldsymbol{w}$$
 independent!
$$= \int p(\boldsymbol{x}|\boldsymbol{w},\{\boldsymbol{x}^{(i)}\}_{i=1}^n)p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)d\boldsymbol{w}$$



$$p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w})}_{\text{posterior}} = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n, \boldsymbol{w})}_{\text{point}} = \underbrace{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})}_{\text{pion}}$$

Posterior distribution:
$$p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \frac{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})}{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n)}$$
 marginal likelihood:
$$p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \int p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})d\boldsymbol{w}$$
 Bayes estimator:
$$\widehat{\boldsymbol{w}} = \int \boldsymbol{w} \cdot p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)d\boldsymbol{w} = \langle \boldsymbol{w} \rangle_{p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)}$$
 Predictive distribution:
$$p(\boldsymbol{x}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \int p(\boldsymbol{x},\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)d\boldsymbol{w}$$
 independent!
$$= \int p(\boldsymbol{x}|\boldsymbol{w},\{\boldsymbol{x}^{(i)}\}_{i=1}^n)p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)d\boldsymbol{w}$$

$$= \int p(\boldsymbol{x}|\boldsymbol{w})p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)d\boldsymbol{w}$$
 model distribution posterior




Bayesian learning

Observed Data: $\{x^{(i)}\}_{i=1}^n$

Parameter: $\boldsymbol{w} = \{\mu \in \mathbb{R}, \sigma^2 \in \mathbb{R}_{++}\}$

Model distribution: $p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w}) = \prod_{i=1}^n p(\boldsymbol{x}^{(i)}|\boldsymbol{w})$

Prior distribution: $p(w) \propto 1$



$$\begin{aligned} \text{Posterior distribution:} \quad p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n) &= \frac{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})}{p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n)} \\ \text{marginal likelihood:} \quad p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n) &= \int p(\{\boldsymbol{x}^{(i)}\}_{i=1}^n|\boldsymbol{w})p(\boldsymbol{w})d\boldsymbol{w} \end{aligned}$$

Bayes estimator:
$$\widehat{\boldsymbol{w}} = \int \boldsymbol{w} \cdot p(\boldsymbol{w} | \{\boldsymbol{x}^{(i)}\}_{i=1}^n) d\boldsymbol{w} = \langle \boldsymbol{w} \rangle_{p(\boldsymbol{w} | \{\boldsymbol{x}^{(i)}\}_{i=1}^n)}$$

Predictive distribution:
$$p(\boldsymbol{x}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n) = \int p(\boldsymbol{x}|\boldsymbol{w})p(\boldsymbol{w}|\{\boldsymbol{x}^{(i)}\}_{i=1}^n)d\boldsymbol{w}$$



Questions?