


Algorithms


Lecture 15: NP Completeness (Part 1)

Anxiao (Andrew) Jiang

CH 34. NP Completeness




CH 34. NP Completeness



Example of unsolvable problems:

Turing's Halting Problem

CH 34. NP Completeness



Example of unsolvable problems:

Turing's Halting Problem

Among solvable problems:

Polynomial-time

v.s.

Super-polynomial time
(Often exponential time)

CH 34. NP Completeness

Polynomial-time algorithm

Problem's input size: **n bits**


Polynomial time: $O(n)$, $O(n^2)$, $O(n^c)$

Polynomial-time solvable

P: the set of all problems that can be solved in polynomial time.

Instance of a problem

Worst-case time complexity




CH 34. NP Completeness

Shortest-Path Problem

Input: A directed graph $G=(V,E)$,
where every edge $e \in E$
has a weight $w(e) > 0$.
Let $s, t \in V$ be two nodes.

Output: A shortest path from s to t .

An instance:



Shortest-Path Problem can be solved in polynomial time. (We have learned it.)

CH 34. NP Completeness



Shortest-Path Problem

Input: A directed graph $G=(V,E)$,
where every edge $e \in E$
has a weight $w(e) > 0$.
Let $s, t \in V$ be two nodes.

Output: A shortest path from s to t .

Simple path: a path without cycles.

An instance:



CH 34. NP Completeness



Shortest-Path Problem

Input: A directed graph $G=(V,E)$,
where every edge $e \in E$
has a weight $w(e) > 0$.
Let $s, t \in V$ be two nodes.

Output: A shortest path from s to t .

Simple path: a path without cycles.

An instance:



simple path


CH 34. NP Completeness

~~Longest~~ ~~Shortest-Path~~ Problem

Input: A directed graph $G=(V,E)$,
where every edge $e \in E$
has a weight $w(e) > 0$.
Let $s, t \in V$ be two nodes.

Output: A ~~shortest~~ simple path from s to t .
longest

An instance:



Can the Longest Path Problem be solved in polynomial time?


CH 34. NP Completeness

~~Longest~~ ~~Shortest-Path~~ Problem

Input: A directed graph $G=(V,E)$,
where every edge $e \in E$
has a weight $w(e) > 0$.
Let $s, t \in V$ be two nodes.

Output: A ~~shortest~~ simple path from s to t .
longest

An instance:




Can the Longest Path Problem be solved in polynomial time?

No one knows. But if you do

CH 34. NP Completeness

Euler Tour:


Given a graph $G=(V,E)$, an Euler Tour is a cycle that passes every edge of G exactly once.



CH 34. NP Completeness

Euler Tour:


Given a graph $G=(V,E)$, an Euler Tour is a cycle that passes every edge of G exactly once.



CH 34. NP Completeness

Euler Tour:


Given a graph $G=(V,E)$, an Euler Tour is a cycle that passes every edge of G exactly once.



CH 34. NP Completeness

Euler Tour:


Given a graph $G=(V,E)$, an **Euler Tour** is a **cycle** that passes every **edge** of G **exactly once**.



CH 34. NP Completeness

Euler Tour:


Given a graph $G=(V,E)$, an Euler Tour is a cycle that passes every edge of G exactly once.



CH 34. NP Completeness

Euler Tour:


Given a graph $G=(V,E)$, an **Euler Tour** is a **cycle** that passes every **edge** of G exactly once.



CH 34. NP Completeness

Euler Tour:


Given a graph $G=(V,E)$, an Euler Tour is a cycle that passes every edge of G exactly once.



CH 34. NP Completeness

Euler Tour:


Given a graph $G=(V,E)$, an Euler Tour is a cycle that passes every edge of G exactly once.



CH 34. NP Completeness

Euler Tour:

Given a graph $G=(V,E)$, an **Euler Tour** is a **cycle** that passes every **edge** of G exactly once.



Euler Tour Problem:


Input: A graph $G=(V,E)$.

Question: Does G have an Euler Tour?

CH 34. NP Completeness

Euler Tour:

Given a graph $G=(V,E)$, an **Euler Tour** is a **cycle** that passes every **edge** of G exactly once.



Euler Tour Problem:

Input: A graph $G=(V,E)$.

Question: Does G have an Euler Tour?

Theorem: G has an Euler Tour if and only if G is connected and all its nodes have even degrees.

So the Euler Tour Problem is polynomial-time solvable.

CH 34. NP Completeness

How about we change “edge” to “node” below”


Hamiltonian cycle:

~~Euler Tour:~~

Hamiltonian cycle

node

Given a graph $G=(V,E)$, an ~~Euler Tour~~ is a cycle that passes every ~~edge~~ of G exactly once.



Euler Tour Problem:

Input: A graph $G=(V,E)$.

Question: Does G have an Euler Tour?

The Euler Tour Problem is polynomial-time solvable.

CH 34. NP Completeness

Hamiltonian cycle:

~~Euler Tour:~~

Hamiltonian cycle

node

Given a graph $G=(V,E)$, an ~~Euler Tour~~ is a cycle that passes every ~~edge~~ of G exactly once.


Hamiltonian cycle

~~Euler Tour Problem:~~

Input: A graph $G=(V,E)$.

Question: Does G have an ~~Euler Tour~~?

Hamiltonian cycle



The Euler Tour Problem is polynomial-time solvable.

Is the Hamiltonian cycle Problem polynomial-time solvable?

No one knows.

CH 34. NP Completeness

Boolean logic: AND operation : $0 \wedge 0 = 0, \quad 0 \wedge 1 = 0, \quad 1 \wedge 0 = 0, \quad 1 \wedge 1 = 1$

OR operation : $0 \vee 0 = 0, \quad 0 \vee 1 = 1, \quad 1 \vee 0 = 1, \quad 1 \vee 1 = 1$

NOT operation : $\bar{0} = 1, \quad \bar{1} = 0$

Boolean variables: $x_1, x_2, \dots, x_n \in \{0,1\}$

Boolean literal: x_i, \bar{x}_i

Boolean formula: $(x_1 \vee x_2 \vee x_3) \wedge (\bar{x}_1 \vee \bar{x}_2 \vee x_4) \wedge (x_1 \vee \bar{x}_4 \vee x_5) \wedge (x_2 \vee \bar{x}_3 \vee \bar{x}_5)$

clause

clause

clause

clause

CNF: Conjunctive Normal Form

CH 34. NP Completeness

2-CNF SAT Problem:

Input: A CNF formula with n variables and k clauses,
where each clause is the “OR” of 2 literals.

Question: Does there exist a solution to the variables that make the formula be true?

Instance:

$$(x_1 \vee x_3) \wedge (\bar{x}_2 \vee x_4) \wedge (x_1 \vee x_5) \wedge (\bar{x}_3 \vee \bar{x}_5)$$

$n=5$ variables

$k=4$ clauses

CH 34. NP Completeness

2-CNF SAT Problem:

Input: A CNF formula with n variables and k clauses, where each clause is the “OR” of 2 literals.

Question: Does there exist a solution to the variables that make the formula be **true**?
Satisfied

Instance:

$$(x_1 \vee x_3) \wedge (\bar{x}_2 \vee x_4) \wedge (x_1 \vee x_5) \wedge (\bar{x}_3 \vee \bar{x}_5)$$

A solution: $x_1 = 1, x_2 = 0, x_3 = 0, x_4 = 0, x_5 = 0$

CH 34. NP Completeness

2-CNF SAT Problem:

Input: A CNF formula with n variables and k clauses, where each clause is the “OR” of 2 literals.

Question: Does there exist a solution to the variables that make the formula be **true**?

Satisfied

Instance:

$$\begin{array}{cccc} 1 & 0 & 1 & 0 & 1 & 0 & 1 & 1 \\ (x_1 \vee x_3) \wedge (\bar{x}_2 \vee x_4) \wedge (x_1 \vee x_5) \wedge (\bar{x}_3 \vee \bar{x}_5) \end{array}$$

A solution: $x_1 = 1, x_2 = 0, x_3 = 0, x_4 = 0, x_5 = 0$

CH 34. NP Completeness

2-CNF SAT Problem:

Input: A CNF formula with n variables and k clauses,
where each clause is the “OR” of 2 literals.

Question: Does there exist a solution to the variables that make the formula be true?

The 2-CNF SAT Problem can be solved in polynomial time.

Instance:

$$(x_1 \vee x_3) \wedge (\bar{x}_2 \vee x_4) \wedge (x_1 \vee x_5) \wedge (\bar{x}_3 \vee \bar{x}_5)$$

$n=5$ variables

$k=4$ clauses

CH 34. NP Completeness

3-CNF SAT Problem:

Input: A CNF formula with n variables and k clauses, where each clause is the “OR” of **3** literals.

Question: Does there exist a solution to the variables that make the formula be true?

The 2-CNF SAT Problem can be solved in polynomial time.


How about 3-CNF SAT Problem? No one knows. But if you do ...

Instance: $(x_1 \vee x_2 \vee x_3) \wedge (\bar{x}_1 \vee \bar{x}_2 \vee x_4) \wedge (x_1 \vee \bar{x}_4 \vee x_5) \wedge (x_2 \vee \bar{x}_3 \vee \bar{x}_5)$

clause clause clause clause

n=5 variables k=4 clauses


CH 34. NP Completeness



P: the set of all problems that can be solved in polynomial time.

NP: the set of all problems with this property:
it takes polynomial time to verify the correctness of the solution.

CH 34. NP Completeness



P: the set of all problems that can be solved in polynomial time.

NP: the set of all problems with this property: it takes polynomial time to verify the correctness of the solution.


Two elements of an algorithm:

- 1) The process of finding a solution.
- 2) Verify the correctness of the solution (so it knows when to end).

CH 34. NP Completeness

$$P \subseteq NP$$

$$P \stackrel{?}{=} NP$$



P: the set of all problems that can be solved in polynomial time.

NP: the set of all problems with this property: it takes polynomial time to verify the correctness of the solution.

Two elements of an algorithm:

- 1) The process of finding a solution.
- 2) Verify the correctness of the solution (so it knows when to end).

CH 34. NP Completeness


$$P \subseteq NP$$

$$P \stackrel{?}{=} NP$$

Hamiltonian cycle Problem:

Input: A graph $G=(V,E)$.

Question: Does G have a Hamiltonian cycle?



Finding a Hamiltonian cycle
might be hard

P: the set of all problems
that can be solved in
polynomial time.

NP: the set of all problems
with this property:
it takes polynomial time
to verify the correctness
of the solution.

Two elements of an algorithm:

- 1) The process of finding a solution.
- 2) Verify the correctness of the solution
(so it knows when to end).

CH 34. NP Completeness


$$P \subseteq NP$$

$$P \stackrel{?}{=} NP$$

Hamiltonian cycle Problem:

Input: A graph $G=(V,E)$.

Question: Does G have a Hamiltonian cycle?



Finding a Hamiltonian cycle
might be hard

But given a Hamiltonian cycle,
it is easy to verify it is indeed
A Hamiltonian cycle.

P: the set of all problems
that can be solved in
polynomial time.

NP: the set of all problems
with this property:
it takes polynomial time
to verify the correctness
of the solution.

Two elements of an algorithm:

- 1) The process of finding a solution.
- 2) Verify the correctness of the solution
(so it knows when to end).

CH 34. NP Completeness

Optimization Problem: A problem where we look for a solution whose objective value is maximized or minimized.

CH 34. NP Completeness

Optimization Problem: A problem where we look for a solution whose objective value is maximized or minimized.

Decision Problem: A problem that asks a Yes/No question.

CH 34. NP Completeness

Optimization Problem: A problem where we look for a solution whose objective value is maximized or minimized.

Decision Problem: A problem that asks a Yes/No question.

Optimization Problem

Shortest-Path Problem

Input: A directed graph $G=(V,E)$,
where every edge $e \in E$
has a weight $w(e)$.
Let $s, t \in V$ be two nodes.

Output: A shortest path from s to t .

Decision Problem

Shortest-Path Problem

Input: A directed graph $G=(V,E)$,
where every edge $e \in E$
has a weight $w(e)$.
Let $s, t \in V$ be two nodes.

Let k be a real number.

Question: Is there a path from s to t
whose length is at most k ?

CH 34. NP Completeness

Optimization Problem: A problem where we look for a solution whose objective value is maximized or minimized.

Decision Problem: A problem that asks a Yes/No question.

Is there a sufficiently short path?

Optimization Problem

Shortest-Path Problem

Input: A directed graph $G=(V,E)$,
where every edge $e \in E$
has a weight $w(e)$.
Let $s, t \in V$ be two nodes.

Output: A shortest path from s to t .

Find the shortest path.

Decision Problem

Shortest-Path Problem

Input: A directed graph $G=(V,E)$,
where every edge $e \in E$
has a weight $w(e)$.
Let $s, t \in V$ be two nodes.
Let k be a real number.

Question: Is there a path from s to t
whose length is at most k ?

CH 34. NP Completeness

Optimization Problem: A problem where we look for a solution whose objective value is maximized or minimized.

Decision Problem: A problem that asks a Yes/No question.

Is there a sufficiently good solution?

Optimization Problem

Shortest-Path Problem

Input: A directed graph $G=(V,E)$,
where every edge $e \in E$
has a weight $w(e)$.
Let $s, t \in V$ be two nodes.

Output: A shortest path from s to t .

Find the best solution.

Decision Problem

Shortest-Path Problem

Input: A directed graph $G=(V,E)$,
where every edge $e \in E$
has a weight $w(e)$.
Let $s, t \in V$ be two nodes.
Let k be a real number.

Question: Is there a path from s to t
whose length is at most k ?

CH 34. NP Completeness

Optimization Problem: A problem where we look for a solution whose objective value is maximized or minimized.

Decision Problem: A problem that asks a Yes/No question.

Optimization Problem

Longest-Path Problem

Input: A directed graph $G=(V,E)$,
where every edge $e \in E$
has a weight $w(e)$.
Let $s, t \in V$ be two nodes.

Output: A longest simple path from s to t .

Decision Problem

Longest-Path Problem

Input: A directed graph $G=(V,E)$,
where every edge $e \in E$
has a weight $w(e)$.
Let $s, t \in V$ be two nodes.
Let k be a real number.

Question: Is there a path from s to t
whose length is at least k ?

CH 34. NP Completeness

Optimization Problem: A problem where we look for a solution whose objective value is maximized or minimized.

Decision Problem: A problem that asks a Yes/No question.

Is there a sufficiently long path?

Optimization Problem

Longest-Path Problem

Input: A directed graph $G=(V,E)$,
where every edge $e \in E$
has a weight $w(e)$.
Let $s, t \in V$ be two nodes.

Output: A longest simple path from s to t .

Find the longest path.

Decision Problem

Longest-Path Problem

Input: A directed graph $G=(V,E)$,
where every edge $e \in E$
has a weight $w(e)$.
Let $s, t \in V$ be two nodes.
Let k be a real number.

Question: Is there a path from s to t
whose length is at least k ?

CH 34. NP Completeness

Optimization Problem: A problem where we look for a solution whose objective value is maximized or minimized.

Decision Problem: A problem that asks a Yes/No question.

Is there a sufficiently good solution?

Optimization Problem

Longest-Path Problem

Input: A directed graph $G=(V,E)$,
where every edge $e \in E$
has a weight $w(e)$.
Let $s, t \in V$ be two nodes.

Output: A longest simple path from s to t .

Find the best solution.

Decision Problem

Longest-Path Problem

Input: A directed graph $G=(V,E)$,
where every edge $e \in E$
has a weight $w(e)$.
Let $s, t \in V$ be two nodes.
Let k be a real number.

Question: Is there a path from s to t
whose length is at least k ?

CH 34. NP Completeness

Optimization Problem: A problem where we look for a solution whose objective value is maximized or minimized.

Decision Problem: A problem that asks a Yes/No question.

Given an optimization problem, there is a corresponding decision problem.

CH 34. NP Completeness

Optimization Problem: A problem where we look for a solution whose objective value is maximized or minimized.

Decision Problem: A problem that asks a Yes/No question.

Given an optimization problem, there is a corresponding decision problem.

If an optimization problem can be solved in polynomial time, then the corresponding decision problem can also be solved in polynomial time.

CH 34. NP Completeness

Optimization Problem: A problem where we look for a solution whose objective value is maximized or minimized.

Decision Problem: A problem that asks a Yes/No question.

Given an optimization problem, there is a corresponding decision problem.

If an optimization problem can be solved in polynomial time, then the corresponding decision problem can also be solved in polynomial time.

If an optimization problem is “easy”, then the corresponding decision problem is also “easy”.

CH 34. NP Completeness

Optimization Problem: A problem where we look for a solution whose objective value is maximized or minimized.

Decision Problem: A problem that asks a Yes/No question.

Given an optimization problem, there is a corresponding decision problem.

If an optimization problem can be solved in polynomial time, then the corresponding decision problem can also be solved in polynomial time.

If an optimization problem is “easy”, then the corresponding decision problem is also “easy”.


If a decision problem is “hard”, then the corresponding optimization problem is also “hard”.

From now on, we focus on “Decision Problems” only.

CH 34. NP Completeness

Decision Problem: A problem that asks a Yes/No question.


Reduction



CH 34. NP Completeness

Decision Problem: A problem that asks a Yes/No question.


Reduction



CH 34. NP Completeness

Decision Problem: A problem that asks a Yes/No question.

Reduction



The mapping is from A to B (not B to A)


The mapping preserves the “Yes/No” answer.

CH 34. NP Completeness

Decision Problem: A problem that asks a Yes/No question.

Reduction: If there is a mapping from the instances of Problem A to the instances of Problem B, such that every “Yes” instance of Problem A is mapped to a “Yes” instance of Problem B, and every “No” instance of Problem A is mapped to a “No” instance of Problem B, then the mapping is called a “Reduction” from Problem A to Problem B.

We can also say
“Problem A is reduced
to Problem B”.



The mapping is from A to B (not B to A)


The mapping preserves the “Yes/No” answer.

CH 34. NP Completeness

Example:

Problem A: Does $ax + b = 0$
have an integer solution?

Problem B: Does $ax^2 + bx + c = 0$
have an integer solution?



The mapping is from A to B (not B to A)

The mapping preserves the “Yes/No” answer.

CH 34. NP Completeness

Example:

Problem A: Does $ax + b = 0$
have an integer solution?


Problem B: Does $ax^2 + bx + c = 0$
have an integer solution?

Mapping:

Instance of Problem A: $ax + b = 0$



Instance of Problem B: $0x^2 + ax + b = 0$



The mapping is from A to B (not B to A)

The mapping preserves the “Yes/No” answer.

CH 34. NP Completeness

Example:

Problem A: Does $ax + b = 0$
have an integer solution?

Problem B: Does $ax^2 + bx + c = 0$
have an integer solution?


Mapping:

Instance of Problem A: $ax + b = 0$



Instance of Problem B: $0x^2 + ax + b = 0$

The mapping is a reduction.



The mapping is from A to B (not B to A)

The mapping preserves the “Yes/No” answer.

CH 34. NP Completeness

Example:

Problem A: Does $ax + b = 0$
have an integer solution?

Problem B: Does $ax^2 + bx + c = 0$
have an integer solution?

Mapping:


Instance of Problem A: $ax + b = 0$



Instance of Problem B: $0x^2 + ax + b = 0$

The mapping is a reduction.

Polynomial-time reduction:
A reduction that takes polynomial time.




The mapping is from A to B (not B to A)

The mapping preserves the "Yes/No" answer.

CH 34. NP Completeness

If B is polynomial-time solvable,
then A is polynomial-time solvable.

Why?




CH 34. NP Completeness

If B is polynomial-time solvable,
then A is polynomial-time solvable.

In this sense,

Problem A is “easier” (or “no harder”),
Problem B is “harder” (or “no easier”).



We will use this property to prove the “hardness” of problems.