

MatematiCAL anal for economists

Anonymus fan of mat.anal

December 2, 2025

Contents

1	Numbers	3
1.1	Basic Classes of Numbers	3
2	Derivative	4
2.1	Basic derivatives	4
3	Taylor	7
3.1	Taylor's formula with the remainder term (and why is it needed? Without it, everything is obvious)	7

Preface

This textbook is designed to assist economics students studying the basic course of mathematical analysis. It summarizes the entire mathematical analysis course taught to economists in the best undergraduate economics program in Eastern Europe.

The lectures include only the essential material, ensuring that students who have achieved top honors in national economics Olympiads are not overburdened and can maintain their sense of superiority over the rest of the world. After all, they likely mastered all this material in kindergarten (or at the latest, by first grade). The division of topics into lectures corresponds well to the actual pace of the course, which spans an entire semester. Almost all statements in the course are self-evident, and their proofs are left to the reader as straightforward exercises.

Chapter 1

Numbers

1.1 Basic Classes of Numbers

First, let's introduce the definitions of the basic classes of numbers that we will constantly work with throughout the course.

Definition 1 *Numbers 1, 2, 3, ... are called natural numbers. The notation for the set of all natural numbers is \mathbb{N} .*

Definition 2 *A number is called an integer if it is equal to... but you don't need this because everything in economics is positive.*

Definition 3 *A number is called textitratational if it can be represented as something above a line and something below a line.*

Definition 4 *A number is called textitirrational if it is not rational.*

Obvious Fact 1 *The sum of all natural numbers equals $-1/12$.*

Kindergarten Example: If Vasya had 2 apples and Petya took 1 apple from him, how many apples does Vasya have left? The answer is obviously $-1/12$, as any advanced mathematician knows.

Chapter 2

Derivative

2.1 Basic derivatives

Definition 5 *The definition of derivative is omitted because it is obvious.*

Everything in this chapter is so obvious that no additional explanations will be provided - we'll immediately proceed to analyze an example from kindergarten.

Let's calculate a simple derivative:

$$(0.5 \cdot x)^2 + \cos(x^3) \tag{2.1}$$

According to the theorem (which number?) from paragraph ??:

$$\frac{df}{dx}(0.5 \cdot x) = 0.5 \tag{2.2}$$

It is obvious that:

$$\frac{df}{dx}((0.5 \cdot x)^2) = 2 \cdot 0.5 \cdot x \cdot 0.5 \tag{2.3}$$

Plus a constant:


$$\frac{df}{dx}(x^3) = 3 \cdot x^2 \tag{2.4}$$

According to the theorem (which number?) from paragraph ??:

$$\frac{df}{dx}(\cos(x^3)) = 3 \cdot x^2 \cdot -1 \cdot \sin(x^3) \tag{2.5}$$

A good, solid task?

$$\frac{df}{dx}((0.5 \cdot x)^2 + \cos(x^3)) = 2 \cdot 0.5 \cdot x \cdot 0.5 + 3 \cdot x^2 \cdot -1 \cdot \sin(x^3) \tag{2.6}$$



Let's calculate a simple derivative:

$$2 \cdot 0.5 \cdot x \cdot 0.5 + 3 \cdot x^2 \cdot -1 \cdot \sin(x^3) \quad (2.7)$$

Understanding this transformation is left to the reader as a simple exercise:

$$\frac{df}{dx}(0.5 \cdot x) = 0.5 \quad (2.8)$$

It is easy to see:

$$\frac{df}{dx}(2 \cdot 0.5 \cdot x) = 1 \quad (2.9)$$

It is common knowledge:

$$\frac{df}{dx}(2 \cdot 0.5 \cdot x \cdot 0.5) = 0.5 \quad (2.10)$$

A good, solid task?


$$\frac{df}{dx}(x^2) = 2 \cdot x \quad (2.11)$$

Let's imagine this household as:

$$\frac{df}{dx}(3 \cdot x^2) = 3 \cdot 2 \cdot x \quad (2.12)$$

Let's imagine this household as:

$$\frac{df}{dx}(x^3) = 3 \cdot x^2 \quad (2.13)$$



If this is not obvious to you, try attending a lecture for a change:

$$\frac{df}{dx}(\sin(x^3)) = 3 \cdot x^2 \cdot \cos(x^3) \quad (2.14)$$

A similar one can be proved:

$$\frac{df}{dx}(-1 \cdot \sin(x^3)) = -1 \cdot 3 \cdot x^2 \cdot \cos(x^3) \quad (2.15)$$

It is obvious that:

$$\frac{df}{dx}(3 \cdot x^2 \cdot -1 \cdot \sin(x^3)) = 3 \cdot 2 \cdot x \cdot -1 \cdot \sin(x^3) + 3 \cdot x^2 \cdot -1 \cdot 3 \cdot x^2 \cdot \cos(x^3) \quad (2.16)$$

If this is not obvious to you, try attending a lecture for a change:

$$\begin{aligned} \frac{df}{dx}(2 \cdot 0.5 \cdot x \cdot 0.5 + 3 \cdot x^2 \cdot -1 \cdot \sin(x^3)) &= 0.5 + 3 \cdot 2 \cdot x \cdot -1 \cdot \sin(x^3) \\ &\quad + 3 \cdot x^2 \cdot -1 \cdot 3 \cdot x^2 \cdot \cos(x^3) \end{aligned} \quad (2.17)$$

Chapter 3

Taylor

3.1 Taylor's formula with the remainder term (and why is it needed? Without it, everything is obvious)

Definition 6 *Taylor's formula is obvious, so no additional explanations will be given. Let's start straight with an example.*

At first the derivatives must be calculated:

Let's calculate a simple derivative:

$$\cos(\sin(x)) + x^3 \tag{3.1}$$

According to the theorem (which number?) from paragraph ??:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \tag{3.2}$$

It is easy to see:


$$\frac{df}{dx}(\cos(\sin(x))) = \cos(x) \cdot -1 \cdot \sin(\sin(x)) \tag{3.3}$$

It is easy to see:

$$\frac{df}{dx}(x^3) = 3 \cdot x^2 \tag{3.4}$$

Understanding this transformation is left to the reader as a simple exercise:

$$\frac{df}{dx}(\cos(\sin(x)) + x^3) = \cos(x) \cdot -1 \cdot \sin(\sin(x)) + 3 \cdot x^2 \tag{3.5}$$



Let's calculate a simple derivative:

$$\cos(x) \cdot -1 \cdot \sin(\sin(x)) + 3 \cdot x^2 \quad (3.6)$$

Let's imagine this household as:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.7)$$

According to the theorem (which number?) from paragraph ??:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.8)$$

Plus a constant:

$$\frac{df}{dx}(\sin(\sin(x))) = \cos(x) \cdot \cos(\sin(x)) \quad (3.9)$$

As already shown earlier:


$$\frac{df}{dx}(-1 \cdot \sin(\sin(x))) = -1 \cdot \cos(x) \cdot \cos(\sin(x)) \quad (3.10)$$

As already shown earlier:

$$\begin{aligned} \frac{df}{dx}(\cos(x) \cdot -1 \cdot \sin(\sin(x))) &= -1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) \\ &\quad + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) \end{aligned} \quad (3.11)$$

A similar one can be proved:

$$\frac{df}{dx}(x^2) = 2 \cdot x \quad (3.12)$$



Let's imagine this household as:

$$\frac{df}{dx}(3 \cdot x^2) = 3 \cdot 2 \cdot x \quad (3.13)$$

Understanding this transformation is left to the reader as a simple exercise:

$$\begin{aligned} \frac{df}{dx}(\cos(x) \cdot -1 \cdot \sin(\sin(x)) + 3 \cdot x^2) = & -1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \quad (3.14) \\ & \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + 3 \cdot 2 \cdot x \end{aligned}$$

Let's calculate a simple derivative:

$$-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + 3 \cdot 2 \cdot x \quad (3.15)$$

Understanding this transformation is left to the reader as a simple exercise:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.16)$$

A good, solid task?

$$\frac{df}{dx}(-1 \cdot \sin(x)) = -1 \cdot \cos(x) \quad (3.17)$$

Understanding this transformation is left to the reader as a simple exercise:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.18)$$

Plus a constant:

$$\frac{df}{dx}(\sin(\sin(x))) = \cos(x) \cdot \cos(\sin(x)) \quad (3.19)$$

A good, solid task?

$$\frac{df}{dx}(-1 \cdot \sin(\sin(x))) = -1 \cdot \cos(x) \cdot \cos(\sin(x)) \quad (3.20)$$

According to the theorem (which number?) from paragraph ??:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x))) &= -1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \\ &\quad \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) \end{aligned} \quad (3.21)$$

A similar one can be proved:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.22)$$

Should be known from school:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.23)$$

If this is not obvious to you, try attending a lecture for a change:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.24)$$

If you don't understand this obvious transformation, then you need to go into a program where they don't study mathematical analysis:

$$\frac{df}{dx}(\cos(\sin(x))) = \cos(x) \cdot -1 \cdot \sin(\sin(x)) \quad (3.25)$$

If this is not obvious to you, try attending a lecture for a change:


$$\begin{aligned} \frac{df}{dx}(\cos(x) \cdot \cos(\sin(x))) &= -1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \\ &\quad (3.26) \end{aligned}$$

Should be known from school:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \cos(x) \cdot \cos(\sin(x))) &= -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \\ &\quad \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \end{aligned} \quad (3.27)$$

A good, solid task?

$$\begin{aligned} \frac{df}{dx}(\cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) &= -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) \\ &\quad + \cos(x) \cdot -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) \\ &\quad + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \end{aligned} \quad (3.28)$$



As already shown earlier:

$$\begin{aligned}
 & \frac{df}{dx}(-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \\
 &= -1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \\
 & \quad -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)))
 \end{aligned} \tag{3.29}$$

As already shown earlier:

$$\frac{df}{dx}(2 \cdot x) = 2 \tag{3.30}$$

A similar one can be proved:

$$\frac{df}{dx}(3 \cdot 2 \cdot x) = 6 \tag{3.31}$$

If you don't understand this obvious transformation, then you need to go into a program where they don't study mathematical analysis:

$$\begin{aligned}
 & \frac{df}{dx}(-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + 3 \cdot 2 \cdot x) \\
 &= -1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + \cos(x) \cdot -1 \\
 & \quad \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + \cos(x) \cdot -1 \\
 & \quad \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + 6
 \end{aligned} \tag{3.32}$$

Let's calculate a simple derivative:

$$\begin{aligned}
 & -1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + \cos(x) \cdot -1 \\
 & \quad \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + \cos(x) \cdot -1 \\
 & \quad \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + 6
 \end{aligned} \tag{3.33}$$

It is easy to see:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.34)$$

It is easy to see:

$$\frac{df}{dx}(-1 \cdot \cos(x)) = -1 \cdot -1 \cdot \sin(x) \quad (3.35)$$

It is common knowledge:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.36)$$

Let's imagine this household as:

$$\frac{df}{dx}(\sin(\sin(x))) = \cos(x) \cdot \cos(\sin(x)) \quad (3.37)$$

Plus a constant:

$$\frac{df}{dx}(-1 \cdot \sin(\sin(x))) = -1 \cdot \cos(x) \cdot \cos(\sin(x)) \quad (3.38)$$

It is obvious that:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) &= -1 \cdot -1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \\ &\quad -1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) \end{aligned} \quad (3.39)$$

Should be known from school:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.40)$$

Understanding this transformation is left to the reader as a simple exercise:

$$\frac{df}{dx}(-1 \cdot \sin(x)) = -1 \cdot \cos(x) \quad (3.41)$$

If you don't understand this obvious transformation, then you need to go into a program where they don't study mathematical analysis:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.42)$$

It is easy to see:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.43)$$

If you don't understand this obvious transformation, then you need to go into a program where they don't study mathematical analysis:

$$\frac{df}{dx}(\cos(\sin(x))) = \cos(x) \cdot -1 \cdot \sin(\sin(x)) \quad (3.44)$$

A similar one can be proved:

$$\frac{df}{dx}(\cos(x) \cdot \cos(\sin(x))) = -1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \quad (3.45)$$

It is common knowledge:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \cos(x) \cdot \cos(\sin(x))) &= -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \\ &\quad \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \end{aligned} \quad (3.46)$$

A similar one can be proved:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \\ = -1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \\ \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \end{aligned} \quad (3.47)$$

If you don't understand this obvious transformation, then you need to go into a program where they don't study mathematical analysis:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \\ = -1 \cdot -1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \\ \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \\ \cdot -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \end{aligned} \quad (3.48)$$

It is common knowledge:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.49)$$

If you don't understand this obvious transformation, then you need to go into a program where they don't study mathematical analysis:

$$\frac{df}{dx}(-1 \cdot \sin(x)) = -1 \cdot \cos(x) \quad (3.50)$$

A similar one can be proved:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.51)$$

If you don't understand this obvious transformation, then you need to go into a program where they don't study mathematical analysis:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.52)$$

As already shown earlier:

$$\frac{df}{dx}(\cos(\sin(x))) = \cos(x) \cdot -1 \cdot \sin(\sin(x)) \quad (3.53)$$

A similar one can be proved:

$$\frac{df}{dx}(\cos(x) \cdot \cos(\sin(x))) = -1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \quad (3.54)$$

If this is not obvious to you, try attending a lecture for a change:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \cos(x) \cdot \cos(\sin(x))) &= -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \\ &\quad \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \end{aligned} \quad (3.55)$$

Plus a constant:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \\ = -1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \\ \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \end{aligned} \quad (3.56)$$

It is obvious that:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.57)$$

It is easy to see:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.58)$$

As already shown earlier:

$$\frac{df}{dx}(-1 \cdot \sin(x)) = -1 \cdot \cos(x) \quad (3.59)$$

It is obvious that:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.60)$$

According to the theorem (which number?) from paragraph ??:

$$\frac{df}{dx}(\cos(\sin(x))) = \cos(x) \cdot -1 \cdot \sin(\sin(x)) \quad (3.61)$$

A good, solid task?

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \sin(x) \cdot \cos(\sin(x))) &= -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \\ &\quad \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \end{aligned} \quad (3.62)$$

Let's imagine this household as:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.63)$$

If this is not obvious to you, try attending a lecture for a change:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.64)$$

It is obvious that:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.65)$$

Let's imagine this household as:

$$\frac{df}{dx}(\sin(\sin(x))) = \cos(x) \cdot \cos(\sin(x)) \quad (3.66)$$

It is common knowledge:

$$\frac{df}{dx}(-1 \cdot \sin(\sin(x))) = -1 \cdot \cos(x) \cdot \cos(\sin(x)) \quad (3.67)$$

Let's imagine this household as:

$$\begin{aligned} \frac{df}{dx}(\cos(x) \cdot -1 \cdot \sin(\sin(x))) &= -1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) \\ &\quad + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) \end{aligned} \quad (3.68)$$

Understanding this transformation is left to the reader as a simple exercise:

$$\begin{aligned} \frac{df}{dx}(\cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) &= -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\ &\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot \\ &\quad -1 \cdot \cos(x) \cdot \cos(\sin(x))) \end{aligned} \quad (3.69)$$

As already shown earlier:

$$\begin{aligned}
& \frac{df}{dx}(-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \\
&= -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \\
&\quad \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
&\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))
\end{aligned} \tag{3.70}$$

A similar one can be proved:

$$\begin{aligned}
& \frac{df}{dx}(-1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)))) \\
&= -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \\
&\quad \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
&\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))))
\end{aligned} \tag{3.71}$$

A similar one can be proved:

$$\begin{aligned}
& \frac{df}{dx}(\cos(x) \cdot -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)))) \\
&= -1 \cdot \sin(x) \cdot -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \\
&\quad + \cos(x) \cdot -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \\
&\quad + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
&\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))))
\end{aligned} \tag{3.72}$$

Let's imagine this household as:


$$\begin{aligned}
& \frac{df}{dx}(-1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + \cos(x) \cdot -1 \\
&\quad \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)))) = \\
&\quad -1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \\
&\quad \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + -1 \cdot \sin(x) \cdot \\
&\quad -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + \cos(x) \cdot -1 \\
&\quad \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \\
&\quad \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
&\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))))
\end{aligned} \tag{3.73}$$

According to the theorem (which number?) from paragraph ??:

$$\begin{aligned}
& \frac{df}{dx}(-1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \\
& \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + \cos(x) \cdot -1 \\
& \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)))) = \\
& \quad -1 \cdot -1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \\
& \quad \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \\
& \quad \cdot -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \\
& \quad + -1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \\
& \quad \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + -1 \cdot \sin(x) \cdot \\
& \quad -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + \cos(x) \cdot -1 \\
& \quad \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \\
& \quad \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
& \quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) \\
& \hspace{15em} (3.74)
\end{aligned}$$

It is obvious that:

$$\begin{aligned}
& \frac{df}{dx}(-1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \\
& \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + \cos(x) \cdot -1 \\
& \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + 6) = \\
& \quad -1 \cdot -1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \\
& \quad \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \\
& \quad \cdot -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \\
& \quad + -1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \\
& \quad \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + -1 \cdot \sin(x) \cdot \\
& \quad -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + \cos(x) \cdot -1 \\
& \quad \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \\
& \quad \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
& \quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) \\
& \hspace{15em} (3.75)
\end{aligned}$$



Let's calculate a simple derivative:

$$\begin{aligned}
& -1 \cdot -1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \\
& \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \\
& \cdot -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \\
& + -1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \\
& \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \\
& + -1 \cdot \sin(x) \cdot -1 \\
& \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \\
& + \cos(x) \cdot -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot \\
& -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
& \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))))
\end{aligned} \tag{3.76}$$

Let's imagine this household as:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \tag{3.77}$$

A good, solid task?

$$\frac{df}{dx}(-1 \cdot \sin(x)) = -1 \cdot \cos(x) \tag{3.78}$$

Plus a constant:

$$\frac{df}{dx}(-1 \cdot -1 \cdot \sin(x)) = -1 \cdot -1 \cdot \cos(x) \tag{3.79}$$

It is easy to see:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \tag{3.80}$$

It is easy to see:

$$\frac{df}{dx}(\sin(\sin(x))) = \cos(x) \cdot \cos(\sin(x)) \quad (3.81)$$

Should be known from school:

$$\frac{df}{dx}(-1 \cdot \sin(\sin(x))) = -1 \cdot \cos(x) \cdot \cos(\sin(x)) \quad (3.82)$$

If this is not obvious to you, try attending a lecture for a change:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot -1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x))) &= -1 \cdot -1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \\ &\quad \cdot -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) \end{aligned} \quad (3.83)$$

A similar one can be proved:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.84)$$

Should be known from school:

$$\frac{df}{dx}(-1 \cdot \cos(x)) = -1 \cdot -1 \cdot \sin(x) \quad (3.85)$$

A similar one can be proved:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.86)$$

It is easy to see:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.87)$$

Plus a constant:

$$\frac{df}{dx}(\cos(\sin(x))) = \cos(x) \cdot -1 \cdot \sin(\sin(x)) \quad (3.88)$$

A similar one can be proved:

$$\begin{aligned} \frac{df}{dx}(\cos(x) \cdot \cos(\sin(x))) &= -1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \\ &\quad \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \end{aligned} \quad (3.89)$$

Understanding this transformation is left to the reader as a simple exercise:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \cos(x) \cdot \cos(\sin(x))) &= -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \\ &\quad \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \end{aligned} \quad (3.90)$$

As already shown earlier:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \\ = -1 \cdot -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \cdot -1 \\ \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \end{aligned} \quad (3.91)$$

A good, solid task?

$$\begin{aligned} \frac{df}{dx}(-1 \cdot -1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \\ = -1 \cdot -1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \\ \cdot \cos(\sin(x)) + -1 \cdot -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \\ \cdot -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \end{aligned} \quad (3.92)$$

If you don't understand this obvious transformation, then you need to go into a program where they don't study mathematical analysis:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.93)$$

Understanding this transformation is left to the reader as a simple exercise:

$$\frac{df}{dx}(-1 \cdot \cos(x)) = -1 \cdot -1 \cdot \sin(x) \quad (3.94)$$

A similar one can be proved:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.95)$$

If you don't understand this obvious transformation, then you need to go into a program where they don't study mathematical analysis:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.96)$$

It is obvious that:

$$\frac{df}{dx}(\cos(\sin(x))) = \cos(x) \cdot -1 \cdot \sin(\sin(x)) \quad (3.97)$$

Understanding this transformation is left to the reader as a simple exercise:

$$\begin{aligned} \frac{df}{dx}(\cos(x) \cdot \cos(\sin(x))) = -1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \end{aligned} \quad (3.98)$$

If this is not obvious to you, try attending a lecture for a change:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \cos(x) \cdot \cos(\sin(x))) &= -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \\ &\quad \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \end{aligned} \quad (3.99)$$

Should be known from school:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \\ = -1 \cdot -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \cdot -1 \\ \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \end{aligned} \quad (3.100)$$

A similar one can be proved:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.101)$$

Let's imagine this household as:

$$\frac{df}{dx}(-1 \cdot \sin(x)) = -1 \cdot \cos(x) \quad (3.102)$$

Understanding this transformation is left to the reader as a simple exercise:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.103)$$

Let's imagine this household as:

$$\frac{df}{dx}(-1 \cdot \sin(x)) = -1 \cdot \cos(x) \quad (3.104)$$

Plus a constant:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.105)$$

Should be known from school:

$$\frac{df}{dx}(\cos(\sin(x))) = \cos(x) \cdot -1 \cdot \sin(\sin(x)) \quad (3.106)$$

If you don't understand this obvious transformation, then you need to go into a program where they don't study mathematical analysis:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \sin(x) \cdot \cos(\sin(x))) &= -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \\ &\quad \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \end{aligned} \quad (3.107)$$

It is common knowledge:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.108)$$

According to the theorem (which number?) from paragraph ??:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.109)$$

A similar one can be proved:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.110)$$

It is obvious that:

$$\frac{df}{dx}(\sin(\sin(x))) = \cos(x) \cdot \cos(\sin(x)) \quad (3.111)$$

It is easy to see:

$$\frac{df}{dx}(-1 \cdot \sin(\sin(x))) = -1 \cdot \cos(x) \cdot \cos(\sin(x)) \quad (3.112)$$

It is obvious that:

$$\begin{aligned} \frac{df}{dx}(\cos(x) \cdot -1 \cdot \sin(\sin(x))) &= -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \\ &\quad + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) \end{aligned} \quad (3.113)$$

As already shown earlier:

$$\begin{aligned} \frac{df}{dx}(\cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) &= -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\ &\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot \\ &\quad -1 \cdot \cos(x) \cdot \cos(\sin(x))) \end{aligned} \quad (3.114)$$

It is easy to see:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \\ = -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \\ \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\ \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \end{aligned} \quad (3.115)$$

If you don't understand this obvious transformation, then you need to go into a program where they don't study mathematical analysis:

$$\begin{aligned}
& \frac{df}{dx}(-1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)))) \\
&= -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \\
&\quad \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
&\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) \\
&\hspace{15em} (3.116)
\end{aligned}$$

A good, solid task?

$$\begin{aligned}
& \frac{df}{dx}(-1 \cdot \sin(x) \cdot -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)))) \\
&= -1 \cdot \cos(x) \cdot -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + \\
&\quad -1 \cdot \sin(x) \cdot -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \\
&\quad \quad + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
&\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) \\
&\hspace{15em} (3.117)
\end{aligned}$$

A similar one can be proved:

$$\begin{aligned}
& \frac{df}{dx}(-1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \\
&\quad \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)))) = \\
&\quad -1 \cdot -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \cdot -1 \\
&\quad \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + -1 \cdot \cos(x) \cdot \\
&\quad -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + -1 \cdot \sin(x) \\
&\quad \cdot -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \\
&\quad \quad \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
&\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) \\
&\hspace{15em} (3.118)
\end{aligned}$$

Plus a constant:

$$\begin{aligned}
\frac{df}{dx} & (-1 \cdot -1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \\
& \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \\
& -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)))) = \\
& -1 \cdot -1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \\
& \cdot \cos(\sin(x)) + -1 \cdot -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \\
& \cdot -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \\
& + -1 \cdot -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \cdot -1 \\
& \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + -1 \cdot \cos(x) \cdot \\
& -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + -1 \cdot \sin(x) \\
& \cdot -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \\
& \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
& \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))))
\end{aligned} \tag{3.119}$$

As already shown earlier:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \tag{3.120}$$

A similar one can be proved:

$$\frac{df}{dx}(-1 \cdot \cos(x)) = -1 \cdot -1 \cdot \sin(x) \tag{3.121}$$

Plus a constant:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \tag{3.122}$$

If you don't understand this obvious transformation, then you need to go into a program where they don't study mathematical analysis:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \tag{3.123}$$

Understanding this transformation is left to the reader as a simple exercise:

$$\frac{df}{dx}(\cos(\sin(x))) = \cos(x) \cdot -1 \cdot \sin(\sin(x)) \tag{3.124}$$

If you don't understand this obvious transformation, then you need to go into a program where they don't study mathematical analysis:

$$\frac{df}{dx}(\cos(x) \cdot \cos(\sin(x))) = -1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \tag{3.125}$$

It is easy to see:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \cos(x) \cdot \cos(\sin(x))) &= -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \\ &\quad \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \end{aligned} \quad (3.126)$$

It is obvious that:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \\ = -1 \cdot -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \cdot -1 \\ \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \end{aligned} \quad (3.127)$$

Let's imagine this household as:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.128)$$

It is obvious that:

$$\frac{df}{dx}(-1 \cdot \sin(x)) = -1 \cdot \cos(x) \quad (3.129)$$

Let's imagine this household as:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.130)$$

As already shown earlier:

$$\frac{df}{dx}(-1 \cdot \sin(x)) = -1 \cdot \cos(x) \quad (3.131)$$

Plus a constant:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.132)$$

It is obvious that:

$$\frac{df}{dx}(\cos(\sin(x))) = \cos(x) \cdot -1 \cdot \sin(\sin(x)) \quad (3.133)$$

Understanding this transformation is left to the reader as a simple exercise:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \sin(x) \cdot \cos(\sin(x))) &= -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \\ &\quad \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \end{aligned} \quad (3.134)$$

It is common knowledge:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.135)$$

It is common knowledge:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.136)$$

It is easy to see:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.137)$$

A similar one can be proved:

$$\frac{df}{dx}(\sin(\sin(x))) = \cos(x) \cdot \cos(\sin(x)) \quad (3.138)$$

Should be known from school:

$$\frac{df}{dx}(-1 \cdot \sin(\sin(x))) = -1 \cdot \cos(x) \cdot \cos(\sin(x)) \quad (3.139)$$

Understanding this transformation is left to the reader as a simple exercise:

$$\begin{aligned} \frac{df}{dx}(\cos(x) \cdot -1 \cdot \sin(\sin(x))) &= -1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) \\ &\quad + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) \end{aligned} \quad (3.140)$$

If you don't understand this obvious transformation, then you need to go into a program where they don't study mathematical analysis:

$$\begin{aligned} \frac{df}{dx}(\cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) &= -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\ &\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot \\ &\quad -1 \cdot \cos(x) \cdot \cos(\sin(x))) \end{aligned} \quad (3.141)$$

Let's imagine this household as:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \\ = -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \\ \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\ \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \end{aligned} \quad (3.142)$$

Should be known from school:

$$\begin{aligned}
& \frac{df}{dx}(-1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)))) \\
&= -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \\
&\quad \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
&\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) \\
&\hspace{15em} (3.143)
\end{aligned}$$

It is obvious that:

$$\begin{aligned}
& \frac{df}{dx}(-1 \cdot \sin(x) \cdot -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)))) \\
&= -1 \cdot \cos(x) \cdot -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + \\
&\quad -1 \cdot \sin(x) \cdot -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \\
&\quad \quad + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
&\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) \\
&\hspace{15em} (3.144)
\end{aligned}$$

A good, solid task?

$$\begin{aligned}
& \frac{df}{dx}(-1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \\
&\quad \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)))) = \\
&\quad -1 \cdot -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \cdot -1 \\
&\quad \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + -1 \cdot \cos(x) \cdot \\
&\quad -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + -1 \cdot \sin(x) \\
&\quad \cdot -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \\
&\quad \quad \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
&\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) \\
&\hspace{15em} (3.145)
\end{aligned}$$

Let's imagine this household as:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \tag{3.146}$$

It is obvious that:

$$\frac{df}{dx}(-1 \cdot \sin(x)) = -1 \cdot \cos(x) \tag{3.147}$$

Understanding this transformation is left to the reader as a simple exercise:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \tag{3.148}$$

It is obvious that:

$$\frac{df}{dx}(-1 \cdot \sin(x)) = -1 \cdot \cos(x) \quad (3.149)$$

It is easy to see:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.150)$$

Let's imagine this household as:

$$\frac{df}{dx}(\cos(\sin(x))) = \cos(x) \cdot -1 \cdot \sin(\sin(x)) \quad (3.151)$$

Understanding this transformation is left to the reader as a simple exercise:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \sin(x) \cdot \cos(\sin(x))) &= -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \\ &\quad \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \end{aligned} \quad (3.152)$$

If this is not obvious to you, try attending a lecture for a change:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.153)$$

As already shown earlier:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.154)$$

A good, solid task?

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.155)$$

A similar one can be proved:

$$\frac{df}{dx}(\sin(\sin(x))) = \cos(x) \cdot \cos(\sin(x)) \quad (3.156)$$

As already shown earlier:

$$\frac{df}{dx}(-1 \cdot \sin(\sin(x))) = -1 \cdot \cos(x) \cdot \cos(\sin(x)) \quad (3.157)$$

It is common knowledge:

$$\begin{aligned} \frac{df}{dx}(\cos(x) \cdot -1 \cdot \sin(\sin(x))) &= -1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) \\ &\quad + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) \end{aligned} \quad (3.158)$$

It is common knowledge:

$$\begin{aligned} \frac{df}{dx}(\cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) &= -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\ &\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot \\ &\quad -1 \cdot \cos(x) \cdot \cos(\sin(x))) \end{aligned} \quad (3.159)$$

According to the theorem (which number?) from paragraph ??:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \\ = -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \\ \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\ \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \end{aligned} \quad (3.160)$$

Should be known from school:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)))) \\ = -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \\ \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\ \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) \end{aligned} \quad (3.161)$$

Let's imagine this household as:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \sin(x) \cdot -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)))) \\ = -1 \cdot \cos(x) \cdot -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + \\ -1 \cdot \sin(x) \cdot -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \\ + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\ \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) \end{aligned} \quad (3.162)$$

As already shown earlier:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.163)$$

Should be known from school:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.164)$$

Plus a constant:

$$\frac{df}{dx}(-1 \cdot \cos(x)) = -1 \cdot -1 \cdot \sin(x) \quad (3.165)$$

According to the theorem (which number?) from paragraph ??:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.166)$$

As already shown earlier:

$$\frac{df}{dx}(\cos(\sin(x))) = \cos(x) \cdot -1 \cdot \sin(\sin(x)) \quad (3.167)$$

As already shown earlier:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \cos(x) \cdot \cos(\sin(x))) &= -1 \cdot -1 \cdot \sin(x) \cdot \cos(\sin(x)) + -1 \\ &\quad \cdot \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \end{aligned} \quad (3.168)$$

A similar one can be proved:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.169)$$

Let's imagine this household as:

$$\frac{df}{dx}(-1 \cdot \sin(x)) = -1 \cdot \cos(x) \quad (3.170)$$

If this is not obvious to you, try attending a lecture for a change:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.171)$$

Plus a constant:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.172)$$

If you don't understand this obvious transformation, then you need to go into a program where they don't study mathematical analysis:

$$\frac{df}{dx}(\sin(\sin(x))) = \cos(x) \cdot \cos(\sin(x)) \quad (3.173)$$

Should be known from school:

$$\frac{df}{dx}(-1 \cdot \sin(\sin(x))) = -1 \cdot \cos(x) \cdot \cos(\sin(x)) \quad (3.174)$$

Should be known from school:

$$\begin{aligned} \frac{df}{dx}(\cos(x) \cdot -1 \cdot \sin(\sin(x))) &= -1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) \\ &\quad + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) \end{aligned} \quad (3.175)$$

A similar one can be proved:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) &= -1 \cdot \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \\ &\quad -1 \cdot \sin(x) \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) \\ &\quad + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \end{aligned} \quad (3.176)$$

A similar one can be proved:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \\ = -1 \cdot -1 \cdot \sin(x) \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \cdot \cos(x) \cdot -1 \\ \cdot \sin(\sin(x)) + -1 \cdot \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \\ \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \end{aligned} \quad (3.177)$$

Understanding this transformation is left to the reader as a simple exercise:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.178)$$

Should be known from school:

$$\frac{df}{dx}(-1 \cdot \sin(x)) = -1 \cdot \cos(x) \quad (3.179)$$

It is common knowledge:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.180)$$

As already shown earlier:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.181)$$

It is common knowledge:

$$\frac{df}{dx}(\sin(\sin(x))) = \cos(x) \cdot \cos(\sin(x)) \quad (3.182)$$

As already shown earlier:

$$\frac{df}{dx}(-1 \cdot \sin(\sin(x))) = -1 \cdot \cos(x) \cdot \cos(\sin(x)) \quad (3.183)$$

As already shown earlier:

$$\begin{aligned} \frac{df}{dx}(\cos(x) \cdot -1 \cdot \sin(\sin(x))) &= -1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) \\ &\quad + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) \end{aligned} \quad (3.184)$$

As already shown earlier:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) &= -1 \cdot \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \\ &\quad -1 \cdot \sin(x) \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) \\ &\quad + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \end{aligned} \quad (3.185)$$

A good, solid task?

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.186)$$

According to the theorem (which number?) from paragraph ??:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.187)$$

If this is not obvious to you, try attending a lecture for a change:

$$\frac{df}{dx}(-1 \cdot \sin(x)) = -1 \cdot \cos(x) \quad (3.188)$$

If this is not obvious to you, try attending a lecture for a change:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.189)$$

If this is not obvious to you, try attending a lecture for a change:

$$\frac{df}{dx}(\sin(\sin(x))) = \cos(x) \cdot \cos(\sin(x)) \quad (3.190)$$

According to the theorem (which number?) from paragraph ??:

$$\frac{df}{dx}(-1 \cdot \sin(\sin(x))) = -1 \cdot \cos(x) \cdot \cos(\sin(x)) \quad (3.191)$$

It is easy to see:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x))) &= -1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \\ &\quad \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) \end{aligned} \quad (3.192)$$

Understanding this transformation is left to the reader as a simple exercise:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.193)$$

Understanding this transformation is left to the reader as a simple exercise:

$$\frac{df}{dx}(\cos(x)) = -1 \cdot \sin(x) \quad (3.194)$$

According to the theorem (which number?) from paragraph ??:

$$\frac{df}{dx}(\sin(x)) = \cos(x) \quad (3.195)$$

A good, solid task?

$$\frac{df}{dx}(\cos(\sin(x))) = \cos(x) \cdot -1 \cdot \sin(\sin(x)) \quad (3.196)$$

A similar one can be proved:

$$\begin{aligned} \frac{df}{dx}(\cos(x) \cdot \cos(\sin(x))) &= -1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \\ &\quad (3.197) \end{aligned}$$

Plus a constant:

$$\begin{aligned} \frac{df}{dx}(-1 \cdot \cos(x) \cdot \cos(\sin(x))) &= -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \\ &\quad \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \end{aligned} \quad (3.198)$$

Should be known from school:

$$\begin{aligned} \frac{df}{dx}(\cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) &= -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) \\ &\quad + \cos(x) \cdot -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) \\ &\quad + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \\ &\quad (3.199) \end{aligned}$$

It is easy to see:

$$\begin{aligned}
& \frac{df}{dx}(-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \\
&= -1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \\
&\quad \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \\
&\quad -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)))
\end{aligned} \tag{3.200}$$

Plus a constant:

$$\begin{aligned}
& \frac{df}{dx}(\cos(x) \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) \\
&= -1 \cdot \sin(x) \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \\
&\quad + \cos(x) \cdot (-1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) \\
&\quad \quad + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + \cos(x) \cdot -1 \\
&\quad \quad \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))))
\end{aligned} \tag{3.201}$$

A similar one can be proved:

$$\begin{aligned}
& \frac{df}{dx}(-1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
&\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) = \\
&\quad -1 \cdot \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \\
&\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) + -1 \cdot \sin(x) \\
&\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) + \cos(x) \\
&\quad \cdot (-1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \\
&\quad \quad \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + \cos(x) \cdot -1 \\
&\quad \quad \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))))
\end{aligned} \tag{3.202}$$

Plus a constant:

$$\begin{aligned}
& \frac{df}{dx}(-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \\
&\quad \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
&\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) = \\
&\quad -1 \cdot -1 \cdot \sin(x) \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \cdot \cos(x) \cdot -1 \\
&\quad \cdot \sin(\sin(x)) + -1 \cdot \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \\
&\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \\
&\quad + -1 \cdot \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \\
&\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) + -1 \cdot \sin(x) \\
&\quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) + \cos(x) \\
&\quad \cdot (-1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \\
&\quad \quad \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + \cos(x) \cdot -1 \\
&\quad \quad \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))))
\end{aligned} \tag{3.203}$$

A similar one can be proved:

$$\begin{aligned}
& \frac{df}{dx} (-1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \\
& \quad \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
& \quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) = -1 \\
& \quad \cdot (-1 \cdot -1 \cdot \sin(x) \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \\
& \quad \quad \cdot \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \\
& \quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) + -1 \\
& \quad \quad \cdot \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \\
& \quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) + -1 \\
& \quad \cdot \sin(x) \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \\
& \quad + \cos(x) \cdot (-1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \\
& \quad \quad \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + \cos(x) \cdot -1 \\
& \quad \quad \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)))) \\
& \hspace{15em} (3.204)
\end{aligned}$$

If this is not obvious to you, try attending a lecture for a change:

$$\begin{aligned}
& \frac{df}{dx} (\cos(x) \cdot -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot \\
& \quad -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
& \quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) = -1 \cdot \sin(x) \cdot \\
& \quad -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \\
& \quad \quad \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
& \quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) + \cos(x) \cdot \\
& \quad -1 \cdot (-1 \cdot -1 \cdot \sin(x) \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \\
& \quad \quad \cdot \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \\
& \quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) + -1 \\
& \quad \quad \cdot \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \\
& \quad \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) + -1 \\
& \quad \cdot \sin(x) \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \\
& \quad + \cos(x) \cdot (-1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \\
& \quad \quad \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + \cos(x) \cdot -1 \\
& \quad \quad \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)))) \\
& \hspace{15em} (3.205)
\end{aligned}$$



It is common knowledge:

$$\begin{aligned}
& \frac{df}{dx}(-1 \cdot \sin(x) \cdot -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \\
& + \cos(x) \cdot -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot \\
& -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
& \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) = -1 \cdot \cos(x) \cdot \\
& -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + -1 \cdot \sin(x) \\
& \cdot -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \\
& \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
& \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) + -1 \\
& \cdot \sin(x) \cdot -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \\
& + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
& \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) + \cos(x) \cdot \\
& -1 \cdot (-1 \cdot -1 \cdot \sin(x) \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \\
& \cdot \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \\
& \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) + -1 \\
& \cdot \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \\
& \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) + -1 \\
& \cdot \sin(x) \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \\
& + \cos(x) \cdot (-1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \\
& \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + \cos(x) \cdot -1 \\
& \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)))) \\
& \hspace{10em} (3.206)
\end{aligned}$$

Let's imagine this household as:

$$\begin{aligned}
& \frac{df}{dx}(-1 \cdot \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \\
& \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + -1 \cdot \sin(x) \cdot -1 \\
& \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + \cos(x) \cdot -1 \cdot (-1 \cdot \cos(x) \\
& \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \\
& + \cos(x) \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) = \\
& -1 \cdot -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \cdot -1 \\
& \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + -1 \cdot \cos(x) \cdot \\
& -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) + -1 \cdot \sin(x) \\
& \cdot -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \\
& \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
& \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) + -1 \\
& \cdot \cos(x) \cdot -1 \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x))) \\
& + -1 \cdot \sin(x) \cdot -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \\
& \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
& \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) + -1 \\
& \cdot \sin(x) \cdot -1 \cdot (-1 \cdot \cos(x) \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) \\
& + -1 \cdot \sin(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \\
& \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)))) + \cos(x) \cdot \\
& -1 \cdot (-1 \cdot -1 \cdot \sin(x) \cdot \cos(\sin(x)) + -1 \cdot \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \\
& \cdot \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \\
& \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) + -1 \\
& \cdot \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \\
& \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) + -1 \\
& \cdot \sin(x) \cdot (-1 \cdot \sin(x) \cdot -1 \cdot \sin(\sin(x)) + \cos(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x))) \\
& + \cos(x) \cdot (-1 \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \\
& \cdot \cos(\sin(x)) + -1 \cdot \sin(x) \cdot -1 \cdot \cos(x) \cdot \cos(\sin(x)) + \cos(x) \cdot -1 \\
& \cdot (-1 \cdot \sin(x) \cdot \cos(\sin(x)) + \cos(x) \cdot \cos(x) \cdot -1 \cdot \sin(\sin(x)))) \\
& \hspace{10em} (3.207)
\end{aligned}$$

[illegible]



Taylor Series:

$$T(\cos(\sin(x)) + x^3) = 1 - 0.5 \cdot x^2 + x^3 + 0.208333 \cdot x^4 \dots \quad (3.209)$$

Afterword

Dear readers, I hope you have been able to spare a moment of your attention for this textbook and to realize its incredible obviousness. You will now excel in your exam, and if not, good luck next year.

The author also expresses great gratitude for the help in preparing this textbook to the students and professors of MIPT, namely to DED, mentor Kolya, and co-mentor Artyom, for actively seeking out the cringe in the code, which undoubtedly improved the quality of the materials. For this important work, the author wholeheartedly thanks all the assistants.

Bibliography:

- Textbooks by G.I. Arkhipov, V.A. Sadovnichy, and V.N. Chubarikov
 - Textbook by J. Stewart
 - Textbook by an unknown author, "The Obviousness of Math"
 - Lectures by A.L. Lukashov on Bipkas
 - Lectures by D.A. Dagaev on the poetry of the Mechanics and Mathematics Faculty