

# Influence of galaxy rotation and outflows in the Lyman Alpha spectral line

Maria Camila Remolina Gutiérrez

Advisor:  
Dr. Jaime E. Forero-Romero

A monograph presented for the degree of  
Physicist



Departamento de Física  
Facultad de Ciencias  
Universidad de los Andes  
Colombia

November, 2015

## *Dedicado a*

a mi papá quien nunca dejará de ser mi héroe,

a mi mamá quien nunca dejará de cuidarme,

a Pipe por crecer conmigo,

a Sofi por ser mi mejor amiga,

a Juanda por ser mi eterno confidente,

a Jaime por ser mi guía y amigo,

a Lucas por cambiarme la vida,

...

a la Universidad de los Andes y la Beca Quiero Estudiar

por hacer mis sueños realidad.

# Influence of galaxy rotation and outflows in the Lyman Alpha spectral line

Maria Camila Remolina Gutiérrez

Submitted for the degree of Physicist

November, 2015

## Abstract

Young galaxies in the Universe have a strong Ly- $\alpha$  emission caused by the ionized Hydrogen atoms in their interstellar medium. When the spectrum of a galaxy has an intense peak around the Ly- $\alpha$  natural frequency ( $2.46 \times 10^{15}$  Hz) it is called a Lyman Alpha Emitter (LAE). Typical LAEs are very distant ( $z \gtrsim 2$ ). This makes that all the data astronomers can obtain from them is their spectra, and from there all the physical information of the galaxy must be derived. Trying to solve this task requires the creation of a simplified and solid model. In this monograph I propose to consider LAEs as a spherical distribution of Hydrogen atoms that undergoes a solid body rotation and a radial expansion. I simulate the effect of rotational velocity, outflow velocity and optical depth of the LAE in the outgoing spectra. The main conclusion is that this new model reproduces LAEs observed features in a clear way and with consistent physical parameters. However, proper observational fits are left for future work. This monograph accomplishes the objective of extracting as much information as possible from a LAE's Ly- $\alpha$  line.

# Influencia de rotación y outflows en la línea espectral Lyman Alpha

Maria Camila Remolina Gutiérrez

Presentado para el título de Física  
Noviembre, 2015

## Abstract

Las galaxias jóvenes en el Universo tienen una fuerte línea de emisión Ly- $\alpha$  causada por los átomos de Hidrógeno ionizados en su medio interestelar. Cuando el espectro de una galaxia tiene un pico intenso alrededor de la frecuencia natural Ly- $\alpha$  ( $2.46 \times 10^{15}$  Hz) se le llama una Lyman Alpha Emitter (LAE). Las LAEs típicas están muy distantes ( $z \gtrsim 2$ ). Esto hace que los datos que los astrónomos puedan obtener de ellas sean solo sus espectros, y es de estos que toda la información física de la galaxia debe ser derivada. Tratar de resolver esta tarea requiere crear un modelo sólido y simplificado. En esta monografía, yo propongo considerar las LAEs como una distribución esférica de átomos de Hidrógeno, que rota como un cuerpo rígido y se expande radialmente. Yo simulo los efectos de velocidad rotacional, velocidad de outflows y profundidad óptica de la LAE en el espectro de salida. La conclusión principal es que este nuevo modelo reproduce las características principales de LAEs observadas. Si embargo, fits observacionales se dejan para trabajo futuro. Esta monografía logra el objetivo de extraer la mayor información posible de la línea Ly- $\alpha$  de una LAE.

# Declaration

The work in this monograph is based on research carried out at the Physics Department of Universidad de los Andes, Colombia. No part of this work has been submitted elsewhere for any other degree and it is all my own work unless referenced to the contrary in the text.

**Copyright © 2015 by MARIA CAMILA REMOLINA GUTIÉRREZ.**

“The copyright of this monograph rests with the author. No quotations from it should be published without the author’s prior written consent and information derived from it should be acknowledged”.

# Acknowledgements

I acknowledge my advisor Jaime Forero-Romero for all his guidance and teachings along my time at the university. Also to Juan Nicolas Garavito-Camargo for becoming my academic older brother. Finally, to AstroAndes for all the wonderful group meetings.

I acknowledge the Scientific Computational Cluster of Universidad de los Andes because all the simulations were run in its processors.

Most of my code benefits from the work of the IPython and Matplotlib communities ([1, 2]).

# Contents

<b>Abstract</b>	<b>iii</b>
<b>Resumen</b>	<b>iv</b>
<b>Declaration</b>	<b>v</b>
<b>Acknowledgements</b>	<b>vi</b>
<b>1 Introduction</b>	<b>1</b>
1.1 The physics of Lyman Alpha Emitters . . . . .	1
<b>2 A new LAE model</b>	<b>5</b>
2.1 A Ly- $\alpha$ Photon's Path in CLARA . . . . .	6
2.2 Galaxy's Viewing Angle . . . . .	8
<b>3 Results</b>	<b>10</b>
3.1 Parameters' Initial Values . . . . .	10
3.2 Influence of the viewing angle $\theta$ . . . . .	14
3.3 Morphology of Ly- $\alpha$ line . . . . .	15
3.3.1 Standard Deviation . . . . .	16
3.3.2 Skewness . . . . .	16
3.3.3 Sigma of Asymmetry . . . . .	17
3.4 Influences of the free parameters . . . . .	18
<b>4 Comparison against observations</b>	<b>19</b>

<b>Contents</b>	<b>viii</b>
<b>5 Conclusions</b>	<b>21</b>
5.1 Future work . . . . .	22
<b>6 Bibliography</b>	<b>23</b>
Appendix	28
<b>A Additional Figures</b>	<b>28</b>



# Chapter 1

## Introduction

Galaxies are key to understand our Universe. So far, human kind has been able to study in detail the closest ones. Distant galaxies are still challenging to detect. They contain however information of millions of stars and gas that could unravel mysteries like for example, how Milky Way type of galaxies formed. It is an open challenge to obtain as much information as possible from these young galaxies. One way to accomplish this is to look at their spectra. Astronomers noted that a that relevant fraction of these galaxies emitted a really strong line at a particular wavelength and named them Lyman Alpha Emitters (LAEs). The purpose of this monograph is to model and simulate LAEs in order to advance our knowledge about this galaxy population.

### 1.1 The physics of Lyman Alpha Emitters

In the early Universe the most abundant elements were Hydrogen(H) and Helium(He), causing young galaxies to be rich in H atoms. Also, due to stellar activity in a galaxy, the surrounding gas can be ionized. These two facts leave the H atoms prone to excite and emit radiation at definite frequencies. One of these frequencies corresponds to the Ly- $\alpha$  line. [3]

When a Hydrogen atom is excited, the only electron it has moves to a higher

energy level. However after some time, it is no longer able to maintain itself in this state. For this, the electron goes back to the ground level. Finally, as the energy obtained from this de-excitement has to be conserved, it is emitted as a photon with a wavelength of  $1215.67 \text{ \AA}$ . This is the Lyman Alpha (  $\text{Ly-}\alpha$  ) transition, discovered by Theodore Lyman in 1906. [4]

$\text{Ly-}\alpha$  radiation is emitted from the H atoms in a galaxy. Due to the amount of atoms inside it, the whole body could become a strong  $\text{Ly-}\alpha$  radiator. Astronomers can take spectra of a galaxy to detect emission at that wavelength. If the galaxy shows a strong  $\text{Ly-}\alpha$  line, it classifies as a Lyman Alpha Emitter (LAE). ([5], [6], [7], [8], [9], [10], [11], [12], [10], [13], [14], [15], [16], [17], [18])





Figure 1.1: **Radiative Transfer Sketch:** A  $\text{Ly-}\alpha$  photon being absorbed and re-emitted by Hydrogen atoms.

When a  $\text{Ly-}\alpha$  photon is emitted inside the galaxy, it travels through its interstellar medium (ISM). During the photon's path it can be absorbed and re-emitted by an H atom. The new frequency of the photon is different that the initial one, in an observer frame of reference due to the atom's velocity. In LAEs, the state of the ISM gas, before and after a photon's re-emission is pretty much the same. This allows a random walk approximation and consider photon-atom encounters as scatterings. A photon's scattering can happen several times, as seen in Fig. 1.1.

This stops only until the photon is able to escape the galaxy.



**Figure A3.** Emergent line profile for the static and dustless sphere test. The photons are emitted at the centre of a sphere with different hydrogen optical depths. The analytical solution follows Dijkstra et al. (2006).

Figure 1.2: **Figure A3 of Forero-Romero et al. paper:** CLARA's view on the escape fraction of Lyman- $\alpha$  photons in high redshift galaxies [19]. Reproduced with the permission of the author.

In a static galaxy, this random walk process produces a spectrum with two equal and symmetric peaks around the natural Ly- $\alpha$  wavelength. This can be seen in Figure A3 of Forero-Romero et al. paper [19] (Fig. 1.2). If now the gas has a bulk velocity, the shape of the Ly- $\alpha$  profile changes. I explore these effects in this work by proposing a new model for a LAE. It consists of an spherical distribution of H atoms undergoing a solid body rotation and radial expansion(outflows). This new model can help observers to determine physical characteristics of a LAE by just looking at its Ly- $\alpha$  line.

The main goal is to measure the effect of the model's physical parameters on the outgoing Ly- $\alpha$  line. In this case, due to the resonant nature of the Ly- $\alpha$  line, analytical solutions can not be derived. It becomes necessary to run simulations to

explore and test the model.

In this monograph I use a radiative transfer code that exists in the literature called CLARA (Code for Lyman Alpha Radiation Analysis). CLARA was written by Forero-Romero et al. [19]. It can simulate the Ly- $\alpha$  line of a spherical rotating LAE depending on its mass and velocity. In this work I modify CLARA to include outflows and explore the resulting consequences on the Ly- $\alpha$  line.

# Chapter 2

## A new LAE model

Galaxies are ellipsoidal. However, for simplicity I model them as a sphere. This shape facilitates the interpretation of the results in our numerical experiments. Furthermore, this approximation is commonly used in the literature, as it explains a wide variety of observational features ([20], [21], [22]).

There are 3 parameters in my model that define a LAE: the rotational velocity ( $v_{\text{rot}}$ ), the outflow velocity ( $v_{\text{out}}$ ) and the optical depth ( $\tau_{\text{H}}$ ).  $v_{\text{out}}$  is due to material ejected from the galaxy, by supernovas([21], [23], [24], [25]), and  $\tau_{\text{H}}$  roughly corresponds to number of H atoms found by a Ly- $\alpha$  photon if one traces a line from the center of the galaxy, to its edge.

In this model, the LAE has a bulk velocity which is the superposition of the rotation and the outflow, as shown in Fig. 2.1. The velocity components are written in Eqs. (2.0.1) (2.0.2) (2.0.3). In these equations,  $R$  is the radius of the sphere;  $x$ ,  $y$  and  $z$  are the coordinates in a cartesian frame; and the  $\mp$  signs in  $v_x$  and  $v_y$  indicate the direction of rotation, respectively. This rotation is a solid body rotation and its direction goes according to the right hand rule applied to the  $\hat{k}$  unit vector

$$v_x = \frac{x}{R}v_{\text{out}} - \frac{y}{R}v_{\text{rot}} \quad (2.0.1)$$

$$v_y = \frac{y}{R}v_{\text{out}} + \frac{x}{R}v_{\text{rot}} \quad (2.0.2)$$

$$v_z = \frac{z}{R}v_{\text{out}} \quad (2.0.3)$$




Figure 2.1: **Model:** Spherical LAE with tangential and radial velocity due to rotation and outflows.

These 3 parameters,  $v_{\text{rot}}$ ,  $v_{\text{out}}$  and  $\tau_{\text{H}}$ , leave the idea of a model of LAE, that although simple, considers the main galaxy's dynamics. All of them have been previously proposed and used by different authors, but never combined together ([26], [27], [28], [22], [21], [29], [30], [31], [32], [33], [34], [35], [36]).

## 2.1 A Ly- $\alpha$ Photon's Path in CLARA

In several problems in science it is useful to define dimensionless variables. This new formulation has many advantages. The numerical equations do not depend on the unit choice. This tends to make the results' analysis straight forward. In the

Ly- $\alpha$  radiative transfer problem, a commonly used dimensionless variable  $x$  is used to describe a photon's frequency and is defined as:

$$x \equiv \frac{(\nu - \nu_\alpha)}{\Delta\nu_D}, \quad (2.1.4)$$

where  $\nu$  is the photon's frequency and  $\nu_\alpha = 2.46 \times 10^{15}$  Hz is the Ly- $\alpha$  natural frequency. The denominator  $\Delta\nu_D$  is defined in Eq. 2.1.5.

$$\Delta\nu_D \equiv \nu_\alpha \sqrt{\frac{2kT}{m_p c^2}} \equiv \nu_\alpha \frac{v_{\text{th}}}{c}, \quad (2.1.5)$$

where  $\Delta\nu_D$  is known as the Doppler broadening of the Ly- $\alpha$  line. It depends on the neutral gas temperature  $T$  or equivalently the thermal velocity  $v_{\text{th}}$  of the atoms. In the model the temperature is kept constant at  $T = 10^4$ K and the thermal velocity is  $v_{\text{th}} = 12.8 \text{ km s}^{-1}$ .

If  $x < 0$ , the final frequency  $\nu < \nu_\alpha$ . This translates to the final wavelength being larger than the Ly- $\alpha$  natural one. The phenomenon is called a redshift in frequency. If, on the contrary,  $x > 0$ , the photon suffers a blueshift in frequency.

However, in the plots shown in this monograph I use instead velocity,  $V$ , units. This eases the comparison against observational data. The velocity units are defined by

$$V = x v_{\text{th}}. \quad (2.1.6)$$

The units of  $V$  are usually  $\text{km s}^{-1}$ . In this case, the photon is redshifted in frequency when the velocity is greater than 0, and blueshifted in the opposite case.

The initial emission of photons is taken at the center of the sphere. From here, 100000 photons are emitted with the natural Ly- $\alpha$  wavelength. Then, they start to behave as described in section 1.1 (See Fig. 1.1). When each photon is re-emitted, its new wavelength depends on the H atom's velocity (both thermal and bulk). However its new direction of propagation is random in the rest frame of the atom.

The individual scattering of all the photons is tracked through the complete 3D Hydrogen distribution. Once each photon escapes the galaxy, its final values are stored: position  $\vec{r}$ , direction of propagation  $\hat{k}$ , dimensionless frequency  $x$ , and number of scatterings  $N$ . To build the observed spectrum I make a histogram of the escape frequencies  $x$ .

## 2.2 Galaxy's Viewing Angle

An observer located far away, only receives photons emitted along its line of sight. That means, only photons escaping in the direction of the observer must be counted in the spectrum. In the simulation I approximate this by taking into account only the photons with escaping direction angle  $\theta$  respect to the rotation axis within the range  $[\theta_{min} - \theta_{max}]$ . I illustrate this in Fig. 2.2.

Therefore, in principle the spectra depend on two new parameters, the azimuthal and the polar angle. However, the galaxy's motion is symmetrical respect to its rotation axis. This implies that the resulting spectrum is independent from the azimuthal angle. Taking into account this symmetry, I only select photons on their polar angle, regardless of their azimuthal angle. Regarding the polar angle, I build the spectra for observers located on 3 different positions, with  $\theta$  intervals uniformly distributed in  $\cos(\theta)$ .

To summarize, the parameters influencing the spectra are  $v_{rot}$ ,  $v_{out}$ ,  $\tau_H$  and  $\theta$ . In the next chapter I evaluate the impact of each parameter on the resulting spectra.






Figure 2.2: **Viewing Angle Sketch:** The galaxy cut is in the  $y-z$  plane perspective and the observer is located at an specific viewing angle of the sphere. Only photons with a direction that enters in this range of vision are taken into account to build the observed spectrum.

# Chapter 3

## Results

### 3.1 Parameters' Initial Values

In order to define the ranges of  $\tau_{\text{H}}$ ,  $v_{\text{rot}}$  and  $v_{\text{out}}$ , it is necessary to refer to observational constraints. The common values for typical LAEs are in Tab. 3.1. I run CLARA for all the permutations of these 3 parameters.

$\tau_{\text{H}}$	$v_{\text{rot}}$ (km s <sup>-1</sup> )	$v_{\text{out}}$ (km s <sup>-1</sup> )
10 <sup>5</sup> , 10 <sup>6</sup> , 10 <sup>7</sup>	50, 100	5, 10, 15, 20, 25, 50, 75

Table 3.1: **Parameters' Values:** All consistent with a LAE's typical properties

The resulting sets of spectra are in Figs. 3.1, 3.3 and 3.2 for  $\theta \simeq 90^\circ$ . I do not include the spectra for  $v_{\text{out}} = 50, 75$  km s<sup>-1</sup> because they are the same as  $v_{\text{out}} = 25$  km s<sup>-1</sup>. For this reason, I limit in all the plots  $v_{\text{out}}$  to a maximum value of 25 km s<sup>-1</sup>. Nevertheless, the figures for  $v_{\text{out}} = 50, 75$  km s<sup>-1</sup> are in the Appendix A in Figs. A.1, A.2 and A.3.




Figure 3.1: **Ly- $\alpha$  profile for  $\tau_H = 10^5$ :** With  $v_{rot}$  ranging 50, 100  $\text{km s}^{-1}$  and  $v_{out}$  ranging 5, 10, 15, 20, 25  $\text{km s}^{-1}$ . The intensity is in arbitrary units. The viewing angle  $\theta \simeq 90^\circ$ . The dashed vertical line indicates the Ly- $\alpha$  line's natural frequency.




Figure 3.2: **Ly- $\alpha$  profile for  $\tau_H = 10^6$ :** Same as Fig. 3.1 but for  $\tau_H = 10^6$ .




Figure 3.3: **Ly- $\alpha$  profile for  $\tau_H = 10^7$ :** Same as Fig. 3.1 but for  $\tau_H = 10^7$ .

These last 3 plots show a clear creation of two asymmetric peaks around  $V = 0$  km s<sup>-1</sup> with the tallest peak is always redshifted, with a strong dependence on the outflow velocity.

## 3.2 Influence of the viewing angle $\theta$

I take now into account the viewing angle of the galaxy to build the observed spectra. For all of the physical parameters' combinations, the effect of  $\theta$  in the Ly- $\alpha$  line is always the same.




Figure 3.4: **Ly- $\alpha$  profile for different  $\theta$ :** With  $\tau_H = 10^5$ ,  $v_{\text{rot}} = 50$  km s<sup>-1</sup> and  $v_{\text{out}} = 20$  km s<sup>-1</sup>. The intensity is in arbitrary units.

From Figs. 3.4, 3.5 and 3.6 is clear that the intensity of the valley between the two peaks increases along with  $\theta$ . This causes an intensity decrease in the rest of the frequencies, thus a broadening of the line. The asymmetry also changes with the viewing angle.

In the next subsection, I show only the results of the angle at which an observer sees the galaxy's angular momentum vector perpendicular to the line of sight. The purpose of this is only to decrease the number of plots in the document, as there is an analogous behavior for all viewing angles.




Figure 3.5: **Ly- $\alpha$  profile for different  $\theta$ :** With  $\tau_H = 10^6$ ,  $v_{\text{rot}} = 100 \text{ km s}^{-1}$  and  $v_{\text{out}} = 5 \text{ km s}^{-1}$ . The intensity is in arbitrary units.




Figure 3.6: **Ly- $\alpha$  profile for different  $\theta$ :** With  $\tau_H = 10^7$ ,  $v_{\text{rot}} = 100 \text{ km s}^{-1}$  and  $v_{\text{out}} = 15 \text{ km s}^{-1}$ . The intensity is in arbitrary units.

### 3.3 Morphology of Ly- $\alpha$ line

To quantify the morphology of the Ly- $\alpha$  profile, I use three statistics: the standard deviation, the skewness and a factor sigma of asymmetry ( $\sigma_A$ ).

### 3.3.1 Standard Deviation

The standard deviation measures the dispersion of frequencies in the Ly- $\alpha$  line. If it increases, the frequencies spread to wider ranges.




Figure 3.7: **Standard deviation plot for each  $\tau_H$**  . The viewing angle  $\theta \simeq 90^\circ$ .

Fig. 3.7 shows that the standard deviation is inversely proportional to the outflow velocity. This implies that the greater  $v_{out}$  is, the less disperse is the Ly- $\alpha$  frequency distribution. This signals that the peaks start merging to one another. Also, the higher the  $\tau_H$ , the more inclined the curves. The horizontal dotted line in the plot is the value of standard deviation for an observational LAE, in particular Q2206BX151 (at redshift  $z \approx 2.1974$ ). It is shown in Fig. 4.1 in the third column with second row.

### 3.3.2 Skewness

The skewness measures the asymmetry of the Ly- $\alpha$  line. If the value is greater than zero, there is more weight in the left tail of the line, if it is less than zero, there is more weight in the right one.

Fig. 3.8 show that the skewness is proportional to the outflow velocity, at a fixed rotational velocity. The greater  $v_{out}$  is, the more asymmetric is the Ly-






Figure 3.8: **Skewness plot for each  $\tau_H$**  . The viewing angle  $\theta \simeq 90^\circ$ .

$\alpha$  frequency distribution. The horizontal dotted line is the value of skewness for Q2206BX151. As seen, the line intersects with the skewness curves. This would imply that Q2206BX151, according to the model, has its parameters narrowed to:  $v_{rot} \in [50, 100]$  km s<sup>-1</sup>,  $v_{out} \in [15, 20]$  km s<sup>-1</sup> and  $\tau_H \in [10^6, 10^7]$ . However, these values are not completely consistent with the standard deviation.


### 3.3.3 Sigma of Asymmetry

Sigma of asymmetry  $\sigma_A$  is another measure for the asymmetry of the Ly- $\alpha$  line. It was defined by McLinden et al. [37] as follows. The two peaks, the red one and the blue one, are fitted with a Gaussian curve. Their standard deviations  $\sigma_{red}$  and  $\sigma_{blue}$  are obtained. Then the factor  $\sigma_A$  is:

$$\sigma_A = \frac{\sigma_{red}}{\sigma_{blue}} \quad (3.3.1)$$

If the value is less than one, the left peak is thinner than the right peak. If it is greater than one, there left peak is wider than the right peak.

Fig. 3.9 shows that this parameters has high fluctuations. I can not see a clear

Figure 3.9: **Sigma of asymmetry  $\sigma_A$ .**

behavior or tendency that applies to all the curves. The only thing is that the values of  $\sigma_A$  start close together and  $v_{\text{out}}$  makes them separate.

### 3.4 Influences of the free parameters

To summarize the influence of the 3 parameters on the Ly- $\alpha$  morphology is the following:

- $\tau_H$  induces a redshift. Increasing the optical depth separates the line of the zero velocity line.
- $v_{\text{out}}$  decreases the right peak's intensity. Higher  $v_{\text{out}}$  make the left peak smaller until it merges with the right one.
- $v_{\text{rot}}$  broadens the line and decreases the maximum intensity. Higher  $v_{\text{rot}}$  implies a flatter spectrum. This effect has not been deeply studied in literature. Only Garavito et al. [31] has simulated its effect. Our results are consistent with their conclusions.

# Chapter 4


## Comparison against observations

There are several observations of LAEs in the literature. Kulas et al. [12] observed distant LAEs ( $z \sim 2 - 3$ ) with high resolution. In the Figure 3 of their paper (Fig. 4.1) they show the 18 Ly- $\alpha$  profiles.

The observed spectra in Fig. 4.1 are modified by the redshift  $z$  of the galaxies. This is, how much distance light had to travel to reach their telescopes. That causes a broadening of the line.

Taking into account this effect, I notice similitudes between them and the simulated spectra. The Ly- $\alpha$  profiles are mostly 2 peaks with an asymmetry between them, just as in the simulation. Also, the small peak (if existing) is always smaller than the tall one.

Many authors that only include the outflows effect are able to fit observational spectra but with a very large magnitude of  $v_{\text{out}}$ , on the order of  $50 - 100 \text{ km s}^{-1}$ . However,  $v_{\text{out}}$  is caused mainly by ejection of material out the galaxy. It is then common that a galaxy rotates faster than it expands, not the opposite. Other authors that include the rotational effect are able to obtain the two peaks but not their asymmetry. Simulating LAEs without outflows does not replicate the majority of the cases.



**Figure 3.** Ly $\alpha$  velocity profiles for the 18 objects in our sample. The zero-point velocity as measured from the nebular emission (H $\alpha$  at  $z \sim 2$  or [O III] $\lambda$ 5007 at  $z \sim 3$ ) is indicated on each image with a dashed line. Considerable variety is evident among the line profile peaks and their ratios. From top to bottom and left to right, the panels proceed from Group I to Group V profiles.

Figure 4.1: **Figure 3 of Kulas et al. paper:** The Kinematics of Multiple-peaked Ly- $\alpha$  Emission in Star-forming Galaxies at  $z \sim 2 - 3$  [12]. This plot is reproduced with the authors permission.

In my model, the combination of rotation and outflows velocities not only makes a lot of sense but seems to be able to fit observations. This new model proposed could replicate Ly- $\alpha$  profiles with typical LAE's values. The main result of this monograph is that I presented a new LAE model roughly consistent with observations.

# Chapter 5

## Conclusions

In this monograph, the objective was to analyze and measure the influence of galaxy rotation and outflows on the Ly- $\alpha$  line. The motivation for this is to be able to obtain physical information of a LAE by just looking at its Ly- $\alpha$  profile. In order to accomplish this objective, I propose a new model of a LAE consisting of an sphere of Hydrogen atoms that expands radially and rotates as a solid body. The program CLARA [19] is used to set the conditions and emulate the transfer code.

The conclusions obtained from this work are:

- The outgoing spectra depend on the angle an external observer is viewing the galaxy from. The closer it is to the equator of the galaxy, the higher the central valley of the frequency distribution.
- The effects of  $v_{\text{rot}}$ ,  $v_{\text{out}}$  and  $\tau_{\text{H}}$  are consistent with the different authors that have used them.  $v_{\text{rot}}$  broadens the Ly- $\alpha$  line.  $v_{\text{out}}$  increases the peaks asymmetry and  $\tau_{\text{H}}$  induces a redshift around the zero velocity.
- The  $v_{\text{rot}}$  effect is much larger than the  $v_{\text{out}}$  one. Only a little fraction of  $v_{\text{rot}}$ , in the radial direction, is necessary to obtain 2 asymmetric peaks. If it  $v_{\text{out}}$  approaches the value of  $v_{\text{rot}}$ , both peaks start merging until only the outflows

effect is visible.

- The final spectra obtained are roughly consistent with LAEs observations.

## 5.1 Future work

Due to the long time CLARA takes to run, it was not possible to fit an observational LAE and predict its parameters. However, the next step is to use tools as MCMC (Monte Carlo Markov Chain) to obtain the galaxy's  $\tau_{\text{H}}$ ,  $v_{\text{rot}}$  and  $v_{\text{out}}$  that would agree with my model.

All of this work is available online and free to use to anyone. The data, source code and instructions to replicate my monograph's results are in the GitHub repository: [https://github.com/astroandes/CLARA\\_RotationOutflows](https://github.com/astroandes/CLARA_RotationOutflows).

# Chapter 6

## Bibliography

- [1] Fernando Pérez and Brian E. Granger. Ipython: a system for interactive scientific computing. *Computing in Science and Engineering*, 9(3):21–29, May 2007.
- [2] J. D. Hunter. Matplotlib: A 2d graphics environment. *Computing In Science & Engineering*, 9(3):90–95, 2007.
- [3] R. B. Partridge and P. J. E. Peebles. Are Young Galaxies Visible? *ApJ*, 147:868, March 1967.
- [4] P. W. Bridgman. Theodore Lyman: A Biographical Memoir. *National Academy of Sciences*, 1957.
- [5] S. Djorgovski and D. J. Thompson. Searches for Primeval Galaxies. In B. Barbuy and A. Renzini, editors, *The Stellar Populations of Galaxies*, volume 149 of *IAU Symposium*, page 337, 1992.
- [6] J. E. Rhoads, S. Malhotra, A. Dey, D. Stern, H. Spinrad, and B. T. Jannuzi. First Results from the Large-Area Lyman Alpha Survey. *ApJ*, 545:L85–L88, December 2000.
- [7] E. Gawiser, H. Francke, K. Lai, K. Schawinski, C. Gronwall, R. Ciardullo, R. Quadri, A. Orsi, L. F. Barrientos, G. A. Blanc, G. Fazio, and J. J. Feldmeier. Ly $\alpha$ -Emitting Galaxies at  $z = 3.1$ : L\* Progenitors Experiencing Rapid Star Formation. *ApJ*, 671:278–284, December 2007.

- [8] R. S. Koehler, P. Schuecker, and K. Gebhardt. Probing dark energy with baryonic acoustic oscillations at high redshifts. *A&A*, 462:7–20, January 2007.
- [9] M. Ouchi, K. Shimasaku, M. Akiyama, C. Simpson, T. Saito, Y. Ueda, H. Furusawa, K. Sekiguchi, T. Yamada, T. Kodama, N. Kashikawa, S. Okamura, M. Iye, T. Takata, M. Yoshida, and M. Yoshida. The Subaru/XMM-Newton Deep Survey (SXDS). IV. Evolution of Ly $\alpha$  Emitters from  $z=3.1$  to 5.7 in the 1 deg<sup>2</sup> Field: Luminosity Functions and AGN. *ApJs*, 176:301–330, June 2008.
- [10] T. Yamada, Y. Nakamura, Y. Matsuda, T. Hayashino, R. Yamauchi, N. Morimoto, K. Kousai, and M. Umemura. Panoramic Survey of Ly $\alpha$  Emitters at  $z = 3.1$ . *AJ*, 143:79, April 2012.
- [11] M. A. Schenker, D. P. Stark, R. S. Ellis, B. E. Robertson, J. S. Dunlop, R. J. McLure, J.-P. Kneib, and J. Richard. Keck Spectroscopy of Faint  $3 < z < 8$  Lyman Break Galaxies: Evidence for a Declining Fraction of Emission Line Sources in the Redshift Range  $6 < z < 8$ . *ApJ*, 744:179, January 2012.
- [12] K. R. Kulas, A. E. Shapley, J. A. Kollmeier, Z. Zheng, C. C. Steidel, and K. N. Hainline. The Kinematics of Multiple-peaked Ly $\alpha$  Emission in Star-forming Galaxies at  $z \sim 2-3$ . *ApJ*, 745:33, January 2012.
- [13] T. S. Chonis, G. A. Blanc, G. J. Hill, J. J. Adams, S. L. Finkelstein, K. Gebhardt, J. A. Kollmeier, R. Ciardullo, N. Drory, C. Gronwall, A. Hagen, R. A. Overzier, M. Song, and G. R. Zeimann. The Spectrally Resolved Ly $\alpha$  Emission of Three Ly $\alpha$ -selected Field Galaxies at  $z \sim 2.4$  from the HETDEX Pilot Survey. *ApJ*, 775:99, October 2013.
- [14] S. L. Finkelstein, C. Papovich, M. Dickinson, M. Song, V. Tilvi, a. M. Koeke-moer, K. D. Finkelstein, B. Mobasher, H. C. Ferguson, M. Giavalisco, N. Reddy, M. L. N. Ashby, a. Dekel, G. G. Fazio, a. Fontana, N. a. Grogin, J.-S. Huang, D. Kocevski, M. Rafelski, B. J. Weiner, and S. P. Willner. A galaxy rapidly forming stars 700 million years after the Big Bang at redshift 7.51. *Nature*, 502(7472):524–527, October 2013.



- [15] G. Östlin, M. Hayes, F. Duval, A. Sandberg, T. Rivera-Thorsen, T. Marquart, I. Orlitova, A. Adamo, J. Melinder, L. Guaita, H. Atek, J. M. Cannon, P. Gruyters, E. C. Herenz, D. Kunth, P. Laursen, J. M. Mas-Hesse, G. Micheva, H. O.-F. S. A. Pardy, M. M. Roth, D. Schaerer, and A. Verhamme. The Lyman-alpha Reference Sample: I. Survey outline and first results for Markarian 259. *ArXiv e-prints*, September 2014.
- [16] M. Hayes, G. Östlin, F. Duval, A. Sandberg, L. Guaita, J. Melinder, A. Adamo, D. Schaerer, A. Verhamme, I. Orlitová, J. M. Mas-Hesse, J. M. Cannon, H. Atek, D. Kunth, P. Laursen, H. Otí-Floranes, S. Pardy, T. Rivera-Thorsen, and E. C. Herenz. The Lyman Alpha Reference Sample. II. Hubble Space Telescope Imaging Results, Integrated Properties, and Trends. *ApJ*, 782:6, February 2014.
- [17] A. L. Faisst, P. Capak, C. M. Carollo, C. Scarlata, and N. Scoville. Spectroscopic Observation of Ly $\alpha$  Emitters at  $z \sim 7.7$  and Implications on Re-ionization. *ApJ*, 788:87, June 2014.
- [18] M. Fumagalli, J. M. O’Meara, J. X. Prochaska, M. Rafelski, and N. Kanekar. Directly imaging damped Ly  $\alpha$  galaxies at  $z \sim 2$  - III. The star formation rates of neutral gas reservoirs at  $z \sim 2.7$ . *MNRAS*, 446 : 3178 – –3198, *January*2015.
- [19] J. E. Forero-Romero, G. Yepes, S. Gottlöber, S. R. Knollmann, A. J. Cuesta, and F. Prada. CLARA’s view on the escape fraction of Lyman  $\alpha$  photons in high-redshift galaxies. *MNRAS*, 415:3666–3680, August 2011.
- [20] S.-H. Ahn, H.-W. Lee, and H. M. Lee. P Cygni type Ly $\alpha$  from starburst galaxies. *MNRAS*, 340:863–869, April 2003.
- [21] A. Verhamme, D. Schaerer, and A. Maselli. 3D Ly $\alpha$  radiation transfer. I. Understanding Ly $\alpha$  line profile morphologies. *A&A*, 460:397–413, December 2006.
- [22] M. Dijkstra, Z. Haiman, and M. Spaans. Ly $\alpha$  Radiation from Collapsing Protogalaxies. I. Characteristics of the Emergent Spectrum. *ApJ*, 649:14–36, September 2006.

- [23] A. Orsi, C. G. Lacey, and C. M. Baugh. Can galactic outflows explain the properties of Ly  $\alpha$  emitters? *MNRAS*, 425:87–115, September 2012.
- [24] T. Hashimoto, A. Verhamme, M. Ouchi, K. Shimasaku, D. Schaerer, K. Nakajima, T. Shibuya, M. Rauch, Y. Ono, and R. Goto. A Close Comparison between Observed and Modeled Ly $\alpha$  Lines for  $z \sim 2.2$  Ly $\alpha$  Emitters. *ApJ*, 812:157, October 2015.
- [25] M. Gronke, P. Bull, and M. Dijkstra. A Systematic Study of Lyman- $\alpha$  Transfer through Outflowing Shells: Model Parameter Estimation. *ApJ*, 812:123, October 2015.
- [26] T. F. Adams. The Escape of Resonance-Line Radiation from Extremely Opaque Media. *ApJ*, 174:439, June 1972.
- [27] J. P. Harrington. The scattering of resonance-line radiation in the limit of large optical depth. *MNRAS*, 162:43, 1973.
- [28] D. A. Neufeld. The transfer of resonance-line radiation in static astrophysical media. *ApJ*, 350:216–241, February 1990.
- [29] J. E. Forero-Romero, G. Yepes, S. Gottlöber, and F. Prada. Modelling the fraction of Lyman break galaxies with strong Lyman  $\alpha$  emission at  $5 \leq z \leq 7$ . *MNRAS*, 419:952–958, January 2012.
- [30] C. L. Martin, M. Dijkstra, A. Henry, K. T. Soto, C. W. Danforth, and J. Wong. The Ly $\alpha$  Line Profiles of Ultraluminous Infrared Galaxies: Fast Winds and Lyman Continuum Leakage. *ApJ*, 803:6, April 2015.
- [31] J. N. Garavito-Camargo, J. E. Forero-Romero, and M. Dijkstra. The Impact of Gas Bulk Rotation on the Ly $\alpha$  Line. *ApJ*, 795:120, November 2014.
- [32] D. A. Neufeld. The escape of Lyman-alpha radiation from a multiphase interstellar medium. *ApJ*, 370:L85–L88, April 1991.

- [33] P. Laursen, J. Sommer-Larsen, and A. C. Andersen. Ly $\alpha$  Radiative Transfer with Dust: Escape Fractions from Simulated High-Redshift Galaxies. *ApJ*, 704:1640–1656, October 2009.
- [34] L. A. Barnes, M. G. Haehnelt, E. Tescari, and M. Viel. Galactic winds and extended Ly $\alpha$  emission from the host galaxies of high column density quasi-stellar object absorption systems. *MNRAS*, 416:1723–1738, September 2011.
- [35] A. Verhamme, Y. Dubois, J. Blaizot, T. Garel, R. Bacon, J. Devriendt, B. Guiderdoni, and A. Slyz. Lyman- $\alpha$  emission properties of simulated galaxies: interstellar medium structure and inclination effects. *A&A*, 546:A111, October 2012.
- [36] H. Yajima, Y. Li, Q. Zhu, T. Abel, C. Gronwall, and R. Ciardullo. Were Progenitors of Local L\* Galaxies Ly $\alpha$  Emitters at High Redshift? *ApJ*, 754:118, August 2012.
- [37] E. M. McLinden, S. L. Finkelstein, J. E. Rhoads, S. Malhotra, P. Hibon, M. L. A. Richardson, G. Cresci, A. Quirrenbach, A. Pasquali, F. Bian, X. Fan, and C. E. Woodward. First Spectroscopic Measurements of [O III] Emission from Ly $\alpha$  Selected Field Galaxies at  $z \sim 3.1$ . *ApJ*, 730:136, April 2011.

# Appendix A

## Additional Figures

Additional figures of spectra with  $v_{\text{rot}} = 50, 100 \text{ km s}^{-1}$  and  $v_{\text{out}} = 50, 75 \text{ km s}^{-1}$ , for all the different  $\tau_{\text{H}}$  .




Figure A.1: **Ly- $\alpha$  profile for  $\tau_H = 10^5$ :** With  $v_{rot}$  ranging 50, 100  $\text{km s}^{-1}$  and  $v_{out}$  ranging 50, 75  $\text{km s}^{-1}$ .




Figure A.2: **Ly- $\alpha$  profile for  $\tau_H = 10^6$ :** With  $v_{rot}$  ranging 50, 100  $\text{km s}^{-1}$  and  $v_{out}$  ranging 50, 75  $\text{km s}^{-1}$ .




Figure A.3: **Ly- $\alpha$  profile for  $\tau_H = 10^7$ :** With  $v_{rot}$  ranging 50, 100  $\text{km s}^{-1}$  and  $v_{out}$  ranging 50, 75  $\text{km s}^{-1}$ .