



Corrigendum



Corrigendum to “Mediational analysis of severe retinal injury causation in children with acute closed head injury” [Child Abuse & Neglect (2025) Vol. 167, Page 107539, <https://doi.org/10.1016/j.chiabu.2025.107539>]

Stephen C. Boos^{a,*}, Fang Wang^b, Ming Wang^b, Kent P. Hymel^c

^a Department of Pediatrics, UMass Chan Medical School—Baystate, Springfield, MA, United States of America

^b Department of Population and Quantitative Health Sciences, Case Western Reserve School of Medicine, Cleveland, OH, United States of America

^c Department of Pediatrics, Pennsylvania State University School of Medicine, Hershey, PA, United States of America

The authors regret an error reported in Table 3 of our original report. Following private communications from an astute reader, we repeated all calculations from our study. A correction to coding resulted in changes to the values in column four of Table 3, as indicated below. This results in higher odds ratios overall and particularly for contact and inertial subdural hemorrhages and diffuse HIE. Importantly, multiple variables remained statistically significant predictors of severe retinal injury. The corrected analysis continues to support our assertion that studies of retinal injury causation need to consider multiple variables simultaneously and cannot rely on tests of single associations. The premise and results of our mediational analyses are reaffirmed. The authors would like to apologise for any inconvenience caused.

Table 3

Results of multivariable logistic regression for the full model and comparative univariable data from Sokoloff.²

Predictor	OR (95 % CI) Sokoloff ²	OR (95 % CI) Full model	OR (95 % CI) Select model ²
Intercept	NA	0.03 (0.01–0.07)**	0.04 (0.02–0.08)**
Abuse	4.51 (3.24–6.29)**	2.06 (1.36–3.12)**	
Age	OR not calculated $p = 0.01$	0.96 (0.94–0.99)*	0.96 (0.94–0.98)**
Respiratory compromise	5.17 (93.73–7.15)**	1.58 (0.94–2.65)	1.53 (0.92–2.54)
Circulatory compromise	4.30 (3.05–6.06)**	1.80 (1.10–2.97)*	1.90 (1.17–3.11)**
Seizure	3.54 (2.58–4.86)**	1.67 (1.10–2.54)*	1.77 (1.18–2.65)**
Altered consciousness	5.62 (3.94–8.03)**	1.65 (0.96–2.84)	1.73 (1.02–2.94)*
Contact subdural	0.28 (0.018–0.045)**	3.77 (1.57–9.77)**	3.98 (1.68–10.20)**
Inertial subdural	8.12 (5.44–12.12)**	8.70 (4.13–20.34)**	10.88 (5.28–25.05)**
Subarachnoid hemorrhage	1.49 (1.09–2.04)**	0.94 (0.60–1.46)	
Cortical brain injury	0.90 (0.57–1.44)	0.89 (0.49–1.63)	
Subcortical brain injury	1.80 (1.01–3.23)*	0.82 (0.39–1.73)	
Focal HIE	1.09 (0.58–2.05)	1.55 (0.70–3.44)	
Diffuse HIE	5.69 (4.07–7.95)**	2.51 (1.58–4.00)**	2.47 (1.61–3.79)**

* $p \leq 0.05$.

** $p \leq 0.01$.

DOI of original article: <https://doi.org/10.1016/j.chiabu.2025.107539>.

* Corresponding author at: UMass Chan Medical School – Baystate, United States of America.

E-mail address: Stephen.Boos@Baystatehealth.org (S.C. Boos).

<https://doi.org/10.1016/j.chiabu.2025.107625>

Available online 8 August 2025

0145-2134/© 2025 Elsevier Ltd. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

*** $p \leq 0.001$.

1. Boos SC, Wang F, Wang M, Hymel KP. Meditational analysis of severe retinal injury causation in children with acute closed head injury. *Child Abuse Negl.* 2025; 167: 107538. <https://doi.org/10.1016/j.chiabu.2025.107539>
2. Sokoloff M, Feldman KW, Levin AV, et al. Retinal hemorrhage variation in inertial versus contact head injuries. *Child Abuse Negl.* 2024; 149: 106606. <https://doi.org/10.1016/j.chiabu.2023.106606>