

## Nymph description 1 of 15



## Distribution



**Nymph Description:** Colour more commonly brown but can be green. From above the thorax has a similar 'X' mark to the adult. In early instars the hind part of the 'X' is more pronounced. The rear leg femur has distinct bands, seen as 3 light and 3 dark from above in early instars, or 3 light and 2 dark in later ones. From the side the bands are at 90 degrees across the femur. The rear tibia is dark except for a light band at the top. From the side the rest of the body has general mottling rather than distinct patterns.

**Confusion with other nymphs:** Austroicetes share some of the same features. *A. cruciata* can also form bands during outbreaks but it is found only in the southern and more arid parts of the plague locust distribution. Compare photo with nymphs of similar species: | [Austroicetes](#) | [Aiolopus](#) | [Caledia](#) | [Heteropternis](#) | [Oedaleus](#) | [Peakesia](#) | [Urnisa](#) |

**Unique features:** The combination of the 'X' mark on top of the thorax with general mottling, transverse bands on the femur and a dark tibia with a light band at the top distinguish them from most other nymphs. Forms into dense aggregations, called bands, when numbers are high.

**Further information on this species:** [Description of adult](#), [More images](#), [Distribution](#), [Biology](#)

Source: [https://www.agriculture.gov.au/biosecurity-trade/pests-diseases-weeds/locusts/about/id-guide/description\\_of\\_nymphs/apl-nymph](https://www.agriculture.gov.au/biosecurity-trade/pests-diseases-weeds/locusts/about/id-guide/description_of_nymphs/apl-nymph)