

FYS4150 Project 2:

Schrödinger's equation for two electrons in a 3-dimensional harmonic oscillator well

Marie Foss, Maria Hammerstrøm

Abstract

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

1 Introduction

The aim of this project is to solve Schrödinger's equation for two electrons in a three-dimensional harmonic oscillator well with and without a repulsive Coulomb interaction. We are first interested in the solution of the radial part of Schrödinger's equation for *one* electron. This equation reads

$$-\frac{\hbar^2}{2m} \left(\frac{1}{r^2} \frac{d}{dr} r^2 \frac{d}{dr} - \frac{l(l+1)}{r^2} \right) R(r) + V(r)R(r) = ER(r). \quad (1)$$

In our case $V(r)$ is the harmonic oscillator potential $(1/2)kr^2$ with $k = m\omega^2$ and E is the energy of the harmonic oscillator in three dimensions. The quantum number l is the orbital momentum of the electron. The oscillator frequency is ω and the energies are

$$E_{nl} = \hbar\omega \left(2n + l + \frac{3}{2} \right), \quad (2)$$

with $n = 0, 1, 2, \dots$ and $l = 0, 1, 2, \dots$. In this project we use $l = 0$.

We will solve Eq. (1) by reformulating it in a discretized form as an eigenvalue equation to be solved with Jacobi's method.

After some substitutions and introducing the dimensionless variable $\rho = (1/\alpha)r$ where α is of dimension length, we can rewrite Eq. (1) as

$$-\frac{d^2}{d\rho^2} u(\rho) + \rho^2 u(\rho) = \lambda u(\rho). \quad (3)$$

This is the first equation to solve numerically. In three dimensions the eigenvalues for $l = 0$ are $\lambda_0 = 3, \lambda_1 = 7, \lambda_2 = 11, \dots$

We use the by now standard expression for the second derivative of a function u

$$u'' = \frac{u(\rho + h) - 2u(\rho) + u(\rho - h)}{h^2} + O(h^2), \quad (4)$$

where h is our step length. For a given number of steps n_{step} , the step length is defined as

$$h = \frac{\rho_{\text{max}} - \rho_{\text{min}}}{n_{\text{step}}}. \quad (5)$$

Next we define minimum and maximum values for the variable ρ , $\rho_{\text{min}} = 0$ and ρ_{max} , respectively. Define an arbitrary value of ρ as

$$\rho_i = \rho_{\text{min}} + ih \quad i = 0, 1, 2, \dots, n_{\text{step}}. \quad (6)$$

Now we can write the Schrödinger equation in a compact way

$$-\frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} + \rho_i^2 u_i = -\frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} + V_i u_i = \lambda u_i, \quad (7)$$

where $V_i = \rho_i^2$ is the harmonic oscillator potential.

Define first the diagonal matrix element

$$d_i = \frac{2}{h^2} + V_i, \quad (8)$$

and the non-diagonal matrix element (which is a mere constant, and are all equal)

$$e_i = -\frac{1}{h^2}. \quad (9)$$

With these definitions the Schrödinger equation takes the following form

$$d_i u_i + e_{i-1} u_{i-1} + e_{i+1} u_{i+1} = \lambda u_i, \quad (10)$$

where u_i is unknown. We can write the latter equation as a matrix eigenvalue problem

tions

$$\begin{pmatrix} d_1 & e_1 & 0 & 0 & \dots & 0 & 0 \\ e_1 & d_2 & e_2 & 0 & \dots & 0 & 0 \\ 0 & e_2 & d_3 & e_3 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & \dots & \dots & d_{n_{\text{step}}-2} & e_{n_{\text{step}}-1} \\ 0 & \dots & \dots & \dots & \dots & e_{n_{\text{step}}-1} & d_{n_{\text{step}}-1} \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \\ \dots \\ \dots \\ \dots \\ u_{n_{\text{step}}-1} \end{pmatrix} = \lambda \begin{pmatrix} u_1 \\ u_2 \\ \dots \\ \dots \\ \dots \\ u_{n_{\text{step}}-1} \end{pmatrix} \quad \mathbf{S}^T \mathbf{A} \mathbf{S} = \mathbf{B} \quad (15)$$

where \mathbf{A} is the matrix in our given problem, and \mathbf{S} is an $(n \times n)$ orthogonal transformation matrix

Secondly, we will consider the Schrödinger equation for *two* electrons by studying two electrons in a harmonic oscillator well which also interact via a repulsive Coulomb interaction.

We start by writing the single-electron equation as

$$-\frac{\hbar^2}{2m} \frac{d^2}{dr^2} u(r) + \frac{1}{2} kr^2 u(r) = E^{(1)} u(r), \quad (12)$$

where $E^{(1)}$ stands for the energy with one electron only. For two electrons with *no* repulsive Coulomb interaction, we have the following Schrödinger equation:

$$\left(-\frac{\hbar^2}{2m} \frac{d^2}{dr_1^2} - \frac{\hbar^2}{2m} \frac{d^2}{dr_2^2} + \frac{1}{2} kr_1^2 + \frac{1}{2} kr_2^2 \right) u(r_1, r_2) = E^{(2)} u(r_1, r_2). \quad (13)$$

After a series of substitutions, this can be written as

$$-\frac{d^2}{d\rho^2} \psi(\rho) + \omega_r^2 \rho^2 \psi(\rho) + \frac{1}{\rho} = \lambda \psi(\rho) \quad (14)$$

2 Methods

2.1 Jacobi's method

In this project we will use Jacobi's method to find the eigenvalues, which consists of doing a number of similarity transforma-

$$\mathbf{S} = \begin{pmatrix} 1 & 0 & \dots & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & \dots & 0 & 0 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & 0 & \dots \\ 0 & 0 & \dots & \cos \theta & 0 & \dots & 0 & \sin \theta \\ 0 & 0 & \dots & 0 & 1 & \dots & 0 & 0 \end{pmatrix} \quad (16)$$

(MATRIX NOT FINISHED)

3 Results

...

4 Conclusions

...

5 List of codes

The codes developed for this project are:

...