

# FYS4150 Project 4:

## The Ising model in two dimensions

Marie Foss (# 56), Maria Hammerstrøm (# 59)

### Abstract

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

**Github:** <https://github.com/mariahammerstrom/Project4>

## 1 Introduction

The Ising model is a model to study phase transitions at finite temperature for magnetic systems. The model was invented in 1920<sup>1</sup> and the analytical solution to the two-dimensional case was found in 1944<sup>2</sup>. This project deals with the Ising model in two dimensions without an external magnetic field.

### 1.1 Theory

In the two-dimensional Ising model, the simplest form of the **energy** for a specific microstate  $i$  is expressed as

$$E_i = -J \sum_{\langle kl \rangle}^N s_k s_l, \quad (1)$$

with  $s_k = \pm 1$  where  $+1$  denotes spin up and  $-1$  denotes spin down,  $N$  is the total number of spins and  $J$  is a coupling constant expressing the strength of the interaction between neighboring spins. The symbol  $\langle kl \rangle$  indicates that the sum is over nearest neighbors only. We assume ferromagnetic ordering, meaning  $J > 0$  and will make use of periodic boundary conditions.

The probability of finding the system in a given microstate  $i$  is expressed by the **Boltzmann probability distribution**

$$P_i(\beta) = \frac{e^{-\beta E_i}}{Z}, \quad (2)$$

where  $\beta = 1/kT$  with  $k$  the Boltzmann constant and  $T$  the temperature, and with  $Z$  as the **partition function** for the canonical ensemble defined as

$$Z = \sum_{i=1}^{\infty} e^{-\beta E_i}, \quad (3)$$

summing over all micro states  $i$ .

The **magnetic moment** of a given microstate is

$$M_i = \sum_j s_j. \quad (4)$$

Some quantities of interest are the **expectation values** for the energy  $\langle E \rangle$  and magnetic moment  $\langle M \rangle$ :

$$\begin{aligned} \langle E \rangle &= \frac{1}{Z} \sum_i E_i e^{-\beta E_i} = kT^2 \frac{\partial \ln Z}{\partial T} = - \frac{\partial \ln Z}{\partial \beta} \\ \langle M \rangle &= \frac{1}{Z} \sum_i M_i e^{-\beta E_i}, \end{aligned} \quad (5)$$

as well as the variances for the energy  $\sigma_E^2$  and for the magnetic moment  $\sigma_M^2$ :

$$\begin{aligned} \sigma_E^2 &= \langle E^2 \rangle - \langle E \rangle^2 \\ \sigma_M^2 &= \langle M^2 \rangle - \langle M \rangle^2, \end{aligned} \quad (6)$$

These values can be used to calculate the **heat capacity** of a fixed volume given by

<sup>1</sup>Lenz, W. (1920). "Beitrag zum Verständnis der magnetischen Eigenschaften in festen Körpern". *Physikalische Zeitschrift* 21: 613-615.

<sup>2</sup>Onsager, Lars (1944). "Crystal statistics. I. A two-dimensional model with an order-disorder transition". *Physical Review*, Series II 65 (3-4): 117-149.

$$C_V = \frac{\sigma_E^2}{kT^2} = \frac{\partial \langle E \rangle}{\partial T}, \quad (7)$$

and the **susceptibility**, which describes whether a material is attracted into or repelled out of a magnetic field, given by

$$\chi = \frac{\sigma_M^2}{kT} = \frac{\langle M^2 \rangle - \langle M \rangle^2}{kT}. \quad (8)$$

The model we are considering here undergoes a **phase transition**. Below a critical temperature  $T_C$  there is spontaneous magnetization  $\langle M \rangle \neq 0$  (magnetic phase), while above this temperature the average magnetization is zero (paramagnetic phase). Near  $T_C$  we can characterize the behavior of many physical quantities by a power law behavior.

An important quantity is the **correlation length**, which is expected to be of the order of the lattice spacing for  $T \gg T_C$ . Because the spins become more and more correlated as  $T$  approaches  $T_C$ , the correlation length increases as we get closer to the critical temperature. The divergent behavior of  $\xi$  near  $T_C$  is

$$\xi(T) \sim |T_C - T|^{-\nu}. \quad (9)$$

A second-order phase transition is characterized by a correlation length which spans the whole system. Since we are always limited to a finite lattice,  $\xi$  will be proportional with the size of the lattice. Through so-called finite size scaling relations it is possible to relate the behavior at finite lattices with the results for an infinitely large lattice. The critical temperature then scales as

$$T_C(L) - T_C(L = \infty) = aL^{-1/\nu}, \quad (10)$$

with  $a$  a constant and  $\nu$  defined in Eq. (9).

## 1.2 Analytical solution

First we will assume that we only have two spins in each dimension, that is  $L = 2$ , where  $L$  is the lattice length. The situation looks like this:

$$\begin{array}{cc} \uparrow_{(1)} & \uparrow_{(2)} \\ \uparrow_{(3)} & \uparrow_{(4)} \end{array}$$

where an upward arrow denotes spin up. We will make use of **periodic boundary conditions**, which means that the neighbor to the right of a given spin  $s_N$  takes the value of  $s_1$ . Similarly, the neighbor to the left of  $s_1$  takes the value of  $s_N$ . This way of treating the boundaries are often used when approximating an infinite system that has a repeating structure. In our case, this mean we will treat our system as though it looked like this (where the original system is highlighted in magenta):

$$\begin{array}{cccc} \uparrow_{(4)} & \uparrow_{(3)} & \uparrow_{(4)} & \uparrow_{(3)} \\ \uparrow_{(2)} & \uparrow_{(1)} & \uparrow_{(2)} & \uparrow_{(1)} \\ \uparrow_{(4)} & \uparrow_{(3)} & \uparrow_{(4)} & \uparrow_{(3)} \\ \uparrow_{(2)} & \uparrow_{(1)} & \uparrow_{(2)} & \uparrow_{(1)} \end{array}$$

Closed form expression can be found for the partition function in Eq. (3) and the corresponding expectation values for  $E$ ,  $|M|$ , the

specific heat  $C_V$  and the susceptibility  $\chi$  as functions of  $T$  using periodic boundary conditions. In the case  $L = 2$  we can write Eq. (1) for the different components as, making use of the periodic boundary conditions, giving

$$\begin{aligned} E_1 &= -J(s_1 s_2 + s_1 s_3) = -2J, \\ E_2 &= -J(s_1 s_2 + s_2 s_4) = -2J, \\ E_3 &= -J(s_1 s_3 + s_3 s_4) = -2J, \\ E_4 &= -J(s_3 s_4 + s_2 s_4) = -2J. \end{aligned}$$

Thus the total energy for the system is

$$E = E_1 + E_2 + E_3 + E_4 = -8J.$$

If going through this exercise for different configurations of spin up and spin down, we find values for energies, degeneracies and magnetization for different configurations as shown in Table 1, where magnetization is calculated from Eq. (4).

These calculations can be used to find the closed-form expressions, starting with the partition function, which can be expressed as

$$\begin{aligned} Z(\beta) &= \sum_E \Omega(E) e^{-\beta E} = 2e^{8J\beta} + 2e^{-8J\beta} + 12 \\ &= 4 \cosh(8J\beta) + 12. \end{aligned} \quad (11)$$

Then the expectation value can be written, using the expression in Eq. (6), as

$$\langle E \rangle = -\frac{8J \sinh(8J\beta)}{\cosh(8J\beta) + 3} \approx -8J \tanh(8J\beta), \quad (12)$$

where we have used that  $\cosh(8J\beta) \gg 3$  in the approximation. Using this approximation gives

$$C_V(\beta) = k \left( \frac{8J\beta}{\cosh(8J\beta)} \right)^2. \quad (13)$$

Similarly, using the values for magnetization in Table 1 and Eq. (4), gives

$$\begin{aligned} \langle |M| \rangle &= \frac{1}{Z} (4e^{8J\beta} + 4 \cdot 2 + 4 \cdot |-2| + |-4| e^{8J\beta}) \\ &= \frac{8}{Z} (e^{8J\beta} + 2), \end{aligned} \quad (14)$$

and

$$\begin{aligned} \langle M \rangle &= \frac{1}{Z} (4e^{8J\beta} + 4 \cdot 2 - 4 \cdot 2 - 4e^{8J\beta}) = 0, \\ \langle M^2 \rangle &= \frac{1}{Z^2} (4^2 e^{8J\beta} + 4 \cdot 2^2 + 4 \cdot (-2)^2 + (-4)^2 e^{8J\beta}) \\ &= \frac{32}{Z^2} (e^{8J\beta} + 1). \end{aligned} \quad (15)$$

Thus

$$\chi(\beta) = \frac{\langle M^2 \rangle - \langle M \rangle^2}{kT} = \frac{32\beta}{Z^2} (e^{8J\beta} + 1). \quad (16)$$

Number of spins up	Energy $E$	Degeneracy $\Omega$	Magnetization $M$
4	$-8J$	1	4
3	0	4	2
2	0	4	0
2	$8J$	2	0
1	0	4	-2
0	$-8J$	1	-4

**Table 1:** Energy and magnetization for the two-dimensional Ising model with  $N = 2 \times 2$  spins with periodic boundary conditions.

## 2 Methods

We wrote a code for the Ising model which computes the mean energy  $E$ , mean magnetization  $|M|$ , the specific heat  $C_V$  and the susceptibility  $\chi$  as functions of  $T$  using periodic boundary conditions for in the  $x$  and  $y$  directions. Using the Ising model in two dimensions, the number of configurations is given by  $2^N$  with  $N = L \times L$  number of spins for a lattice of length  $L$ .

### 2.1 Algorithm

We will use the **Metropolis algorithm**. The algorithm goes as follows:

1. Generate a random configuration in the lattice to create an initial state with energy  $E_b$ .
2. Change the initial configuration by flipping for example one spin only. Compute the energy of this state  $E_t$ .
3. Calculate  $\Delta E = E_t - E_b$ .
4. *The Metropolis test:* If  $\Delta E \leq 0$  the new configuration is accepted, meaning the energy is lowered and we are moving towards the energy minimum at a given temperature. If  $\Delta E > 0$ , calculate  $w = e^{-\beta\Delta E}$ . If  $w < r$  where  $r$  is a random number, accept the new configuration. If else, keep the old configuration.
5. Compute the new energy  $E' = E_t + \Delta E$ . Calculate various expectation values using this energy.
6. Repeat steps 2 – 5 for the chosen number of Monte Carlo cycles, meaning the number of times we should sweep through the lattice and summed over all spins.
7. Compute the various expectation values by dividing by the total number of cycles (and possibly the number of spins).

### 2.2 Parallelization

We have parallelized our code using ...

## 3 Results

Our numerical results from computing the mean energy  $E$ , mean magnetization  $|M|$ , the specific heat  $C_V$  and the susceptibility  $\chi$  as functions of  $T$  using periodic boundary conditions, can be

compared with expressions given in Eq. (11) – (16) for a temperature  $T = 1.0$  (in units of  $kT/J$ ). This gives the following results:

Quantity	Closed form	Numerical
Mean energy $E$	-8	-
Mean magnetization $ M $	4	-
Specific heat $C_V$	$\sim 0$	-
Susceptibility $\chi$	$\sim 0$	-

COMMENT. The number of Monte Carlo cycles needed to achieve good agreement is XXX. COMMENT.

Next we looked at the case of  $L = 20$  spins in the  $x$  and  $y$  directions. In this case we wanted to perform a study of the time (or number of Monte Carlo cycles) we need before reaching an equilibrium situation and can start computing the various expectation values.

A rough and study is done by plotting the various expectation values as a function of number of Monte Carlo cycles, for both  $T = 1.0$  kT/J and  $T = 2.4$  kT/J, as shown in Fig. XXX. PLOT + COMMENTS!

We also study the total number of accepted configurations as a function of the total number of Monte Carlo cycles, shown in Fig. XXX. PLOT + COMMENTS!

...

By simply counting the number of times a given energy appears in our computation we can compute the probability  $P_i$ . We start the counting after the steady state situation has been reached. This result is compared with the compute variance in energy  $\sigma_E^2$ . DISCUSS.

In our study of the behavior of our model close to the critical temperature we look at the various expectation values as a function of  $T$  for  $L = 20, L = 40, L = 60$  and  $L = 80$  for  $T \in [2.0, 2.4]$ .

.... MORE.


## 4 Conclusions

...

## 5 List of codes

The codes developed and used for this project are:

`main.cpp` – main program (C++).



**Figure 1:** Visualization of the spin matrix for a  $80 \times 80$  lattice for different temperatures  $T$ . White = spin up, black = spin down.

`plotting.py` – plotting program which makes plots to study the total number of accepted configurations as a function of total number of Monte Carlo cycles and as a function of temperature  $T$  (Python).

`visualization.py` – provided program for running the Metropolis algorithm and plotting the spin matrix, as shown in Fig. 1 (Python).