

Week 3: Recording synaptic currents

Maria Kesa

January 13, 2016

Abstract

This week, I measure the synaptic currents associated with action potentials.

1 Recording synaptic currents

I begin by initializing two pyramidal cells and connecting them with a synapse.

```
load_file("nrngui.hoc")
load_file("LAcells_template.hoc")
load_file("interneuron_template.hoc")
```

```
number_of_cells=2
objectvar cell_exc[number_of_cells]
```

```
for i = 0, number_of_cells-1 {
    cell_exc[i] = new Cell_A()
}
```

```
objref syn, nc
cell_exc[0].dend syn = new pyrD2pyrD_STFD(0.9)
cell_exc[1].soma nc = new NetCon(&v(0.5), syn, 0, 2, 1)
```

I then introduce the vectors for recording the synaptic currents.

```
objref tvec, ivec
tvec = new Vector()
tvec.record(&t)
ivec = new Vector()
ivec.record(&syn.iampa) //ivec.record(&syn.inmda)
```

Then I introduce a current clamp and run the simulation.

```
objectvar stim
cell_exc[1].soma stim = new IClamp(0.5)
```

```
stim.del = 100
stim.dur = 200
stim.amp = 0.4
```

```
tstop = 400
```




Figure 1: The voltage traces of pre-synaptic(green) and post-synaptic(blue) cells, 400 pA

```
run()
```

Finally, I save the data to file.

```
objref savdata
savdata = new File()
savdata.wopen("current.dat")

savdata.printf("t SThcells[2].soma.v(0.5)\n")
savdata.printf("%d\n", tvec.size())

for i=0, tvec.size()-1 {
    savdata.printf("%g %g\n", tvec.x(i), ivec.x(i))
}

savdata.close()
```

2 Results

For clarity, the voltage traces of the pre-synaptic (green) and post-synaptic (blue) cells are plotted in Figure 1.

In Figure 2, we plot the AMPA current time course and in Figure 3 the NMDA current time course.




Figure 2: AMPA currents of the post-synaptic cell




Figure 3: NMDA currents of the post-synaptic cell

3 Conclusion

This week I recorded synaptic currents going through AMPA and NMDA synapses in pyramidal cells.