
ASSESSMENT 1

START OF SEMESTER 1

Why go quantum?

1. What is quantum computing?
 - (a) Simulating quantum mechanics on modern computers.
 - (b) Making use of the weirdness of quantum mechanics to solve problems that normal computers cannot.
 - (c) Injecting quantum mechanics into classical computers to increase their computing speed.
 - (d) Using computers to discover new laws of quantum physics.
 - (e) I don't know
2. Which of the following does NOT describe a qubit?
 - (a) The outcome of a quantum algorithm.
 - (b) The basic unit of quantum information.
 - (c) The quantum analog of a classical bit.
 - (d) A two-level/state quantum mechanical system.
 - (e) I don't know
3. Which famous quantum algorithm enables efficient prime factorization and has the potential to ruin modern (RSA) cryptography?
 - (a) Deutsch-Josza Algorithm
 - (b) Grover's Algorithm
 - (c) Bose-Einstein Algorithm
 - (d) Variational Quantum Eigensolver
 - (e) Shor's Algorithm
 - (f) I don't know
4. What does Moore's Law tell us?

- (a) The number of transistors on a microchip doubles every two years.
 - (b) Quantum computers are exponentially improved each year.
 - (c) Qubits double in speed each year.
 - (d) Classical computation power improves linearly each year.
 - (e) I don't know
5. What field are people looking to apply quantum computing to?
- (a) Chemistry
 - (b) Finance
 - (c) Biology
 - (d) Machine Learning
 - (e) All of the above
 - (f) None of the above
 - (g) I don't know

Classical Computation

6. What is 13_{10} in binary?

- (a) 00000000000000_2
- (b) 1111111111111_2
- (c) 1101_2
- (d) 1110_2
- (e) 10001_2
- (f) 10100101_2
- (g) I don't know


7. What is 110001_2 in decimal?

- (a) 6
- (b) 37
- (c) 49
- (d) 50
- (e) 71
- (f) I don't know

8. Which of the following runtimes is $O(n^2)$?

- (a) $O(n!)$
- (b) $O(e^n)$
- (c) $O(1)$
- (d) $O(25n^2 + n^3)$
- (e) $O(3 + n + 2n^{20})$
- (f) I don't know

9. What is the output of the following logic circuit?



- (a) 0
- (b) 1
- (c) 10
- (d) 01
- (e) I don't know

Mathematics for Quantum

10. What is the standard form of the complex number $e^{i\theta}$?
- (a) $1 + i\theta$
 - (b) $\cos \theta + i \sin \theta$
 - (c) $(e + i) \cos \theta$
 - (d) I don't know
11. Which of the following forms an orthonormal basis in 2 dimensions?
- (a) $\left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right\}$
 - (b) $\left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ -1 \end{pmatrix} \right\}$
 - (c) $\left\{ \begin{pmatrix} -1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right\}$
 - (d) I don't know
12. Which of the following expressions gives the expectation of a random variable X ?
- (a) $\langle X \rangle = xP(X = x)$
 - (b) $\langle X \rangle = \sum_x xP(X = x)$
 - (c) $\langle X \rangle = \prod_x xP(X = x)$
 - (d) I don't know

Quantum Gates

13. Which of the following gates maps a pure qubit state into a superposition?
- (a) Hadamard
 - (b) Pauli X
 - (c) Pauli Y
 - (d) Pauli Z
 - (e) CNOT
 - (f) I don't know
14. Which of the following gates acts as a quantum phase-flip/shift?
- (a) Hadamard
 - (b) Pauli X
 - (c) Pauli Y
 - (d) Pauli Z
 - (e) CNOT
 - (f) I don't know
15. Which of the following gates acts as a quantum bit-flip?
- (a) Hadamard
 - (b) Pauli X
 - (c) Pauli Y
 - (d) Pauli Z
 - (e) CNOT
 - (f) I don't know

Quantum Mechanics

16. The double slit experiment demonstrates which quantum property of electrons?

- (a) Quantized Charge
- (b) Wave-particle duality
- (c) Electron-light interaction
- (d) I don't know

17. Which of the following is a superposition state?

- (a) $|0\rangle$
- (b) $\frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$
- (c) $|0\rangle \otimes |1\rangle$
- (d) I don't know

18. Which of the following is an entangled state?

- (a) $|01\rangle$
- (b) $\frac{1}{\sqrt{2}}(|00\rangle - |11\rangle)$
- (c) $\frac{1}{\sqrt{2}}(|00\rangle + |10\rangle)$
- (d) I don't know

19. What is the normalization condition for the quantum state $|\psi\rangle$?

- (a) $\langle\psi|\psi\rangle = 1$
- (b) $\langle\psi|\psi\rangle = 0$
- (c) $\langle 1|\psi\rangle = 1$
- (d) I don't know

20. What is the vector form of the quantum state $|\psi\rangle = \alpha|0\rangle + \beta|1\rangle$?

- (a) $\begin{pmatrix} 0 \\ \beta \end{pmatrix}$
- (b) $\begin{pmatrix} \alpha \\ \beta \end{pmatrix}$
- (c) $\begin{pmatrix} \beta \\ \alpha \end{pmatrix}$
- (d) $\begin{pmatrix} \alpha \\ 0 \end{pmatrix}$
- (e) I don't know

21. The geometric interpretation of a qubit state is a vector on a




- (a) Bloch Sphere
- (b) Minkowski Hyperboloid
- (c) 4D Hypercube
- (d) I don't know

22. Given the quantum state $|\psi\rangle = \frac{1}{\sqrt{5}}|\alpha\rangle - \frac{2i}{\sqrt{5}}|\beta\rangle$, what is the probability of measuring state $|\beta\rangle$?

- (a) $\frac{1}{25}$
- (b) $\frac{16}{25}$
- (c) $\frac{1}{5}$
- (d) $\frac{4}{5}$
- (e) $-\frac{1}{5}$
- (f) $\frac{4i}{5}$
- (g) I don't know




Quantum Circuits

23. Which circuit produces an equal superposition state?




(d) I don't know

24. Which circuit produces an entangled state?



(d) I don't know

25. What is the matrix representation of the following circuit?



(a) $\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$

(b) $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$

(c) $\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}$

(d) I don't know

Quantum Computing/Mechanics 101

26. Which famous thought experiment illustrates the idea of superposition?
- (a) The “Glove in Box” Experiment
 - (b) Pavlov’s Dog
 - (c) Schrodinger’s Cat
 - (d) Fermi Paradox
 - (e) I don’t know
27. Which of the following weird quantum properties enables us to create pairs/groups of quantum states that cannot be described independently of one another (irrespective of distance)?
- (a) Superposition
 - (b) Entanglement
 - (c) Quantum Interference
 - (d) I don’t know
28. Which of the following weird quantum properties enables us to create a quantum state which is a linear combinations of several classical states simultaneously?
- (a) Superposition
 - (b) Entanglement
 - (c) Quantum Interference
 - (d) I don’t know
29. Which of the following weird quantum properties enables quantum states to constructively and destructively interact with one another?
- (a) Superposition
 - (b) Entanglement
 - (c) Quantum Interference
 - (d) I don’t know
30. Which of the following best describes a *quantum circuit*?
- (a) Quantum logic applied to classical bits.
 - (b) A sequence of qubits to which quantum gates are applied and a quantum measurement is performed.
 - (c) XOR, AND, and COPY gates applied to qubits.
 - (d) A classical circuit used to control a quantum computer.
 - (e) I don’t know

31. Which of the following best describes *quantum advantage*?
- (a) Quantum computers will be able to do everything classical computers can do, but faster and better.
 - (b) Quantum algorithms will be much easier to write and execute than classical algorithms.
 - (c) Quantum circuits will be much easier to design and run than classical circuits.
 - (d) Quantum computers will leverage strange properties of quantum mechanics to solve certain challenging problems faster than classical computers.
 - (e) I don't know
32. Which of the following best describes a *quantum algorithm*?
- (a) A classical algorithm designed to simulate a quantum system and solve hard problems.
 - (b) A quantum circuit specially designed to leverage entanglement, superposition, and quantum interference to solve computing problems more efficiently than classical computers.
 - (c) A classical circuit which mimics quantum logic gates to demonstrate strange quantum effects.
 - (d) A classical algorithm which is run on a quantum computer, making it faster than an algorithm run on a classical computer.
 - (e) I don't know
33. If I repeatedly create and measure a qubit (via the standard $\{|0\rangle, |1\rangle\}$ basis) in the superposition state $\frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$, what outcome do I expect?
- (a) $|0\rangle$ 100% of the time
 - (b) $|1\rangle$ 100% of the time
 - (c) $\frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$ 100% of the time
 - (d) $|0\rangle$ 50% of the time and $|1\rangle$ 50% of the time
 - (e) $|0\rangle$ 25% of the time and $|1\rangle$ 75% of the time
 - (f) $|0\rangle$ 75% of the time and $|1\rangle$ 25% of the time
 - (g) I don't know
34. Imagine I create a qubit in the superposition state $\frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$, and perform a quantum measurement (via the standard $\{|0\rangle, |1\rangle\}$ basis). Suppose the outcome of this measurement is the state $|0\rangle$. What outcome would I expect if I measure this qubit again in the same basis (assuming I do not apply anymore gates to the qubit)?
- (a) $|0\rangle$
 - (b) $|1\rangle$
 - (c) $\frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$
 - (d) I don't know

35. Imagine I create the two-qubit entangled state $\frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$. Now, I perform a measurement on the first qubit and find that it is in the state $|1\rangle$. What can I say about the state of the second qubit?
- (a) It is in the state $|0\rangle$
 - (b) It is in the state $|1\rangle$
 - (c) It is in the state $\frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$
 - (d) There is a 50% chance it is in the state $|0\rangle$ and a 50% chance it is in the state $|1\rangle$
 - (e) I can't say anything about the state of the second qubit without measuring it as well
 - (f) I don't know
36. Imagine I create the two-qubit entangled state $\frac{1}{\sqrt{2}}(|01\rangle + |10\rangle)$. Now, I perform a measurement on the first qubit and find that it is in the state $|1\rangle$. What can I say about the state of the second qubit?
- (a) It is in the state $|0\rangle$
 - (b) It is in the state $|1\rangle$
 - (c) It is in the state $\frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$
 - (d) There is a 50% chance it is in the state $|0\rangle$ and a 50% chance it is in the state $|1\rangle$
 - (e) I can't say anything about the state of the second qubit without measuring it as well
 - (f) I don't know