



INTRO TO QUANTUM
COMPUTING
LECTURE #14

The Qubit & Bloch Sphere

Amir Karamlou

02/07/2021



© 2020 The Coding School
All rights reserved

Use of this recording is for personal use only. Copying, reproducing, distributing, posting or sharing this recording in any manner with any third party are prohibited under the terms of this registration. All rights not specifically licensed under the registration are reserved.



ANNOUNCEMENTS

TODAY'S LECTURE

- Use the Bloch Sphere to conceptualize the state of a qubit
- View quantum gates as actions on the Bloch sphere



What is a qubit?

- Building block of quantum computers
- A two-level system
- Can be in a superposition of two values



Qubit Reviewed

Superposition: a qubit can be $|0\rangle$ and $|1\rangle$ at the same time!

This is how we show it: $|\psi\rangle = \alpha|0\rangle + \beta|1\rangle$

$$|0\rangle \equiv \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$|\psi\rangle = \alpha \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \beta \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} \alpha \\ \beta \end{pmatrix}$$

$$|1\rangle \equiv \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$\alpha, \beta \in \mathbb{C} \cup \text{complex}$

$$|\alpha|^2 + |\beta|^2 = 1$$



Complex Numbers Review

A complex number consists of both a **real** and **imaginary** component.


$$z = \underset{\text{real}}{a} + i \underset{\text{imaginary}}{b}$$

complex conjugate


$$z^* = \underset{\text{real}}{a} - i \underset{\text{imaginary}}{b}$$

$$|z|^2 = z \cdot z^* = z^* \cdot z$$

$$\begin{aligned}(a+ib)(a-ib) &= a^2 - i\cancel{ab} + i\cancel{ab} - (i)^2 b^2 \\ &= a^2 + b^2\end{aligned}$$



EULER'S FORMULA & COMPLEX EXPONENTIALS



Source: Wikipedia (CC)

Euler's formula:

$$e^{i\varphi} = \cos \varphi + i \sin \varphi$$

Polar representation of complex numbers!

$$z = x + iy = |z|(\cos \varphi + i \sin \varphi) = re^{i\varphi}$$

$$r = |z| = \sqrt{x^2 + y^2}$$

(vector radius)

$$\varphi = \tan^{-1} \left(\frac{y}{x} \right)$$

(vector angle)

What is phase?

$\varphi \equiv \phi$ (phi)

$$e^{i\varphi}$$

$$e^{i\phi} = \cos(\phi) + i \sin(\phi)$$

$$|e^{i\phi}| = \sqrt{\cos^2(\phi) + \sin^2(\phi)} = 1$$

$$\phi = 0 \rightarrow e^{i\phi} = e^0 = \cos(0) + i \sin(0) = 1$$

$$\phi = \pi/2 \rightarrow e^{i\pi/2} = \cos(\pi/2) + i \sin(\pi/2) = i$$

$$\phi = \pi \rightarrow e^{i\pi} = \cos(\pi) + i \sin(\pi) = -1$$

In quantum mechanics global phase has **NO *physical significance*** and **NO *effect*** therefore can be ignored!!

$$e^{i\varphi} |\psi\rangle \rightarrow |\psi\rangle$$


*

$$\Theta |0\rangle \rightarrow |0\rangle$$
$$\frac{|0\rangle - |1\rangle}{\sqrt{2}} \neq \frac{|0\rangle + |1\rangle}{\sqrt{2}}$$



Bloch Sphere

The state of a qubit can be represented as a point on the Bloch Sphere



Unit sphere

$$r=1$$

Qubit

Back to our qubit: $|\psi\rangle = \alpha|0\rangle + \beta|1\rangle$

$$\begin{aligned} |\psi\rangle &= |\alpha| e^{i\phi_\alpha} |0\rangle + |\beta| e^{i\phi_\beta} |1\rangle \\ &= \cancel{e^{i\phi_\alpha}} \left[|\alpha| |0\rangle + |\beta| e^{i(\phi_\beta - \phi_\alpha)} |1\rangle \right] \end{aligned}$$

↓
global phase ↓
relative phase

$$= |\alpha| |0\rangle + |\beta| e^{i\phi} |1\rangle$$

$$|\psi\rangle = \cos(\theta_{1/2}) |0\rangle + \sin(\theta_{1/2}) e^{i\phi} |1\rangle$$

$\alpha, \beta \in \mathbb{C}$ are complex

$$\alpha = |\alpha| e^{i\phi_\alpha}$$

$$\beta = |\beta| e^{i\phi_\beta}$$

$$\phi = \phi_\beta - \phi_\alpha$$

$$|\alpha|^2 + |\beta|^2 = 1 \quad |\alpha| = \cos(\theta_{1/2})$$

$$\cos^2(\theta_{1/2}) + \sin^2(\theta_{1/2}) = 1 \quad |\beta| = \sin(\theta_{1/2})$$




Bloch Sphere

$$|\psi\rangle = \cos(\theta/2)|0\rangle + \sin(\theta/2)e^{i\varphi}|1\rangle$$

North pole: $|0\rangle$

South pole: $|1\rangle$

Classical bits are only on the North pole
or the South pole



Bloch Sphere


$$|\psi\rangle = \cos(\theta/2)|0\rangle + \sin(\theta/2)e^{i\varphi}|1\rangle$$

What is the state of the qubit based on the position on the Bloch sphere?

$$\theta=0: |\psi\rangle = \cos(0)|0\rangle + \sin(0)e^{i\phi}|1\rangle = |0\rangle$$

$$\theta=\pi: |\psi\rangle = \cos(\pi/2)|0\rangle + \sin(\pi/2)e^{i\phi}|1\rangle$$

global phase




Bloch Sphere

$$|\psi\rangle = \cos(\theta/2)|0\rangle + \sin(\theta/2)e^{i\varphi}|1\rangle$$

More examples:

$$\begin{aligned}\theta = \pi/2, \varphi = 0: |\psi\rangle &= \cos(\pi/4)|0\rangle + \sin(\pi/4)e^{i0}|1\rangle \\ &= \frac{1}{\sqrt{2}}|0\rangle + \frac{1}{\sqrt{2}}|1\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}\end{aligned}$$

$$\begin{aligned}\theta = \pi/2, \varphi = \pi: |\psi\rangle &= \cos(\pi/4)|0\rangle + \sin(\pi/4)e^{i\pi}|1\rangle \\ &= \frac{1}{\sqrt{2}}|0\rangle - \frac{1}{\sqrt{2}}|1\rangle = \frac{|0\rangle - |1\rangle}{\sqrt{2}}\end{aligned}$$





QUBIT
X QUBIT



10 MIN BREAK!

$$\begin{matrix} \text{SU(2)} & \longrightarrow & \text{SO(3)} \\ \uparrow & & \uparrow \\ \text{2d complex} & & \text{3D real} \end{matrix}$$

Quantum gates

- Maps one the quantum state to another quantum state
- Moves a point on the Bloch Sphere to another point
- We can represent them with matrices



Important Quantum Gates

Pauli gates:

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{Pauli-X operator (X)}$$

$$\sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \text{Pauli-Y operator (Y)}$$

$$\sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad \text{Pauli-Z operator (Z)}$$





Important Quantum Gates

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{Pauli-X operator (X)}$$

$$\sigma_x |0\rangle = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix} = |1\rangle$$

$$\sigma_x |1\rangle = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} = |0\rangle$$

(quantum bit-flip)



Rotates the state around the **X** axis by 180 degrees


Important Quantum Gates

$$\sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad \text{Pauli-Z operator (Z)}$$

$$\sigma_z |0\rangle = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} = |0\rangle$$

$$\sigma_z |1\rangle = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ -1 \end{pmatrix} = -|1\rangle$$

(phase operator)



Rotates the state around the **Z** axis by 180 degrees

Important Quantum Gates

$$\sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \text{Pauli-Y operator (Y)}$$


$$\sigma_y |0\rangle = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ i \end{pmatrix} = i |1\rangle$$

$$\sigma_y |1\rangle = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \begin{pmatrix} 0 \\ i \end{pmatrix} = \begin{pmatrix} -i \\ 0 \end{pmatrix} = -i |0\rangle$$

(combination of bit-flip and phase-flip)

$$\sigma_y = i \sigma_x \sigma_z$$


Rotates the state around the Y axis by 180 degrees



Rotations on the Bloch Sphere

We're not just limited by 180 degree rotations

For example we can rotate around the x axis by 90 degrees



Rotations on the Bloch Sphere

We can rotate by an angle θ around x or y or z axis:

$$R_x(\theta) = e^{-i \frac{\theta}{2} \sigma_x}$$

Hermitian

$$R_y(\theta) = e^{-i \frac{\theta}{2} \sigma_y}$$

$$R_z(\theta) = e^{-i \frac{\theta}{2} \sigma_z}$$



Rotations on the Bloch Sphere

Let's take rotation by an angle θ around the x axis as an example:

$$R_x(\theta) = e^{-i \frac{\theta}{2} \sigma_x} = \cos \frac{\theta}{2} I - i \sin \frac{\theta}{2} \sigma_x$$

Matrix representation?

$$\begin{aligned} R_x(\theta) &= \cos(\theta/2) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - i \sin(\theta/2) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \\ &= \begin{pmatrix} \cos(\theta/2) & 0 \\ 0 & \cos(\theta/2) \end{pmatrix} + \begin{pmatrix} 0 & -i \sin(\theta/2) \\ -i \sin(\theta/2) & 0 \end{pmatrix} = \begin{pmatrix} \cos(\theta/2) & -i \sin(\theta/2) \\ -i \sin(\theta/2) & \cos(\theta/2) \end{pmatrix} \end{aligned}$$



Rotations on the Bloch Sphere

Let's take rotation by an angle θ around the x axis as an example:

$$R_z(\theta) = e^{-i \frac{\theta}{2} \sigma_z} = \cos \frac{\theta}{2} I - i \sin \frac{\theta}{2} \sigma_z$$

Matrix representation?

$$\begin{aligned} R_z(\theta) &= \cos(\theta/2) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - i \sin(\theta/2) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \\ &= \begin{pmatrix} \cos(\theta/2) - i \sin(\theta/2) & 0 \\ 0 & \cos(\theta/2) - i \sin(\theta/2) \end{pmatrix} \end{aligned}$$




Rotations on the Bloch Sphere

Examples:

$$\begin{aligned} R_x(\pi) |0\rangle &= (\cos(\pi/2) \mathbb{I} - i \sin(\pi/2) \sigma_x) |0\rangle \\ &\stackrel{\text{X } \pi\text{-pulse}}{\downarrow} \\ &= \cos(\pi/2) \cdot \mathbb{I} |0\rangle - i \sin(\pi/2) \sigma_x |0\rangle \\ &= -i \sigma_x |0\rangle = \cancel{|1\rangle} \end{aligned}$$


$$\begin{aligned} R_x(\pi/4) |0\rangle &= (\cos(\pi/4) \cdot \mathbb{I} - i \sin(\pi/4) \sigma_x) |0\rangle \\ &= \cos(\pi/4) \cdot \mathbb{I} |0\rangle - i \sin(\pi/4) \sigma_x |0\rangle \\ &= \frac{1}{\sqrt{2}} |0\rangle - i \frac{1}{\sqrt{2}} |1\rangle = \frac{|0\rangle - i |1\rangle}{\sqrt{2}} \end{aligned}$$



Rotations on the Bloch Sphere

Examples:

$$\begin{aligned} R_y(\pi/2)|0\rangle &= \cos(\pi/4) \cdot |0\rangle - i \sin(\pi/4) \sigma_y |0\rangle \\ &= \frac{1}{\sqrt{2}}|0\rangle + \frac{i}{\sqrt{2}}|1\rangle = \frac{|0\rangle + i|1\rangle}{\sqrt{2}} \end{aligned}$$





$$\begin{aligned} R_z(\pi/2)|+\rangle &= \cos(\pi/4) \cdot |+\rangle - i \sin(\pi/4) \sigma_z |+\rangle \\ &= \frac{1}{\sqrt{2}}|+\rangle - \frac{i}{\sqrt{2}}|-\rangle \quad (\sigma_z |+\rangle = |-\rangle) \end{aligned}$$

$$= \frac{1}{\sqrt{2}} \left[|0\rangle + |1\rangle - i(|0\rangle + i|1\rangle) \right] = \frac{|0\rangle - i|1\rangle}{\sqrt{2}}$$

Experimental Implementations

- All quantum gates are engineered using rotations on the Bloch sphere
- In our quantum hardware we can control each transition with some level of accuracy
- Example: Rabi oscillation:



Quantum Universality

Any quantum computation operation can be made by using a combination of:

{CNOT, H, S, T}



Quantum Universality

What is the S gate?

$$S = R_z(\pi/2) = \begin{pmatrix} 1 & 0 \\ 0 & i \end{pmatrix}$$


$$S^2 = Z$$

What is the T gate?

$$T = R_z(\pi/4) = \begin{pmatrix} 1 & 0 \\ 0 & e^{i\pi/4} \end{pmatrix}$$

$$T^2 = S$$

$$T^4 = Z$$





© 2020 The Coding School
All rights reserved

Use of this recording is for personal use only. Copying, reproducing, distributing, posting or sharing this recording in any manner with any third party are prohibited under the terms of this registration. All rights not specifically licensed under the registration are reserved.