

# Decomposing labelled proof theory for intuitionistic modal logic

Sonia Marin\*

IT-Universitetet i København  
Denmark

Marianela Morales

Universidad Nacional de Córdoba  
Argentina

Lutz Straßburger

Inria Saclay & LIX  
France

## ACM Reference Format:

Sonia Marin, Marianela Morales, and Lutz Straßburger. 2018. Decomposing labelled proof theory for intuitionistic modal logic. In *Proceedings of . ACM*, New York, NY, USA, 1 page. <https://doi.org/>

*Labelled deduction* has been proposed by Gabbay [7] in the 80s as a unifying framework throughout proof theory in order to provide proof systems for a wide range of logics. For modal logics it can take for example the form of labelled natural deduction and labelled sequent systems, as used by Simpson [3], Viganò [4] and Negri [2].

These formalisms make explicit use not only of labels, but also of relational atoms referring to the accessibility relation of a Kripke model. In this short note we propose a system that represents both the *accessibility relation* (for modal logics) and the *preorder relation* (for intuitionistic logic), using the full power of the bi-relational semantics for intuitionistic modal logics, and developing fully the idea of [1].

A *bi-relational frame* [5, 6]  $\mathcal{B}$  is a triple  $\langle W, R, \leq \rangle$  of a non-empty set of worlds  $W$  equipped with an accessibility relation  $R$  and a preorder  $\leq$ , satisfying:

- (F<sub>1</sub>) For all worlds  $x, y, z$ , if  $xRy$  and  $y \leq z$ , there exists a  $u$  such that  $x \leq u$  and  $uRz$ .
- (F<sub>2</sub>) For all worlds  $x, y, z$ , if  $xRy$  and  $x \leq z$ , there exists a  $u$  such that  $y \leq u$  and  $zRu$ .

Reflecting this definition, we define our two-sided intuitionistic labelled sequents to be of the form  $\mathcal{B}, \mathcal{L} \Rightarrow \mathcal{R}$  where  $\mathcal{B}$  denotes a set of relational atoms  $xRy$  and preorder atoms  $x \leq y$ , and  $\mathcal{L}$  and  $\mathcal{R}$  are multi-sets of labelled formulas  $x : A$  (for  $x$  and  $y$  taken from the set of labels and  $A$  an intuitionistic modal formula).

Furthermore, our system has to incorporate the two semantic conditions into deductive rules as follows:

$$F_1 \frac{\mathcal{B}, xRy, y \leq z, x \leq u, uRz, \mathcal{L} \Rightarrow \mathcal{R}}{\mathcal{B}, xRy, y \leq z, \mathcal{L} \Rightarrow \mathcal{R}} u \text{ fresh}$$

$$F_2 \frac{\mathcal{B}, xRy, x \leq z, y \leq u, zRu, \mathcal{L} \Rightarrow \mathcal{R}}{\mathcal{B}, xRy, x \leq z, \mathcal{L} \Rightarrow \mathcal{R}} u \text{ fresh}$$

In the intuitionistic setting, the validity of a modal formula has to be defined using both the  $R$  and the  $\leq$  relation as:  $x \Vdash \Box A$  iff for all  $y$  and  $z$  s.t.  $x \leq y$  and  $yRz$ ,  $z \Vdash A$ .

\*Funded by the Qatar National Research Council project MetaCLF nb. xxx.

Again, our system reflects exactly this definition in the rules introducing the  $\Box$ -operator:

$$\Box_L \frac{\mathcal{B}, x \leq y, yRz, \mathcal{L}, x : \Box A, z : A \Rightarrow \mathcal{R}}{\mathcal{B}, \mathcal{L}, x \leq y, yRz, x : \Box A \Rightarrow \mathcal{R}}$$

$$\Box_R \frac{\mathcal{B}, x \leq y, yRz, \mathcal{L} \Rightarrow \mathcal{R}, z : A}{\mathcal{B}, \mathcal{L} \Rightarrow \mathcal{R}, x : \Box A} y, z \text{ fresh}$$

By complementing these rules with the standard labelled rules for intuitionistic modal logic of [3], we get a system that is sound and complete wrt. the birelational semantics.

In [6], Plotkin and Stirling give a correspondence result for intuitionistic modal logic extended with a family of axioms wrt. some classes of bi-relational frames. For example, the frames that validate the axiom  $4_\Diamond : \Diamond \Diamond A \supset \Diamond A$  are exactly the ones satisfying the condition:

( $\Diamond_4$ ) if  $wRv$  and  $vRu$ , there exists a  $u'$  s.t.  $u \leq u'$  and  $wRu'$ .

Incorporating the preorder symbol into the syntax of our sequents allows us to also obtain a sound and complete proof system for the intuitionistic modal logic extended with axiom  $4_\Diamond$ , by designing the following rule:

$$\Diamond_4 \frac{\mathcal{B}, wRv, vRu, u \leq u', wRu', \mathcal{L} \Rightarrow \mathcal{R}}{\mathcal{B}, wRv, vRu, \mathcal{L} \Rightarrow \mathcal{R}} u' \text{ fresh}$$

Therefore, we decompose further the formalism of labelled sequents and extend the reach of labelled deduction to the logics studied in [6]. These systems enjoy cut-elimination via usual arguments, the generality of the result is subject of ongoing study.

## References

- [1] Paolo Maffezoli, Alberto Naibo, and Sara Negri. *The Church-Fitch knowability paradox in the light of structural proof theory*. Synthese, 190(14):2677-2716, 2013.
- [2] Sara Negri. *Proof analysis in modal logics*. Journal of Philosophical Logic, 34:507-544, 2005.
- [3] Alex Simpson. *The Proof Theory and Semantics of Intuitionistic Modal Logic*. PhD thesis, University of Edinburgh, 1994.
- [4] Luca Viganò. *Labelled Non-Classical Logic*. Kluwer Academic Publisher, 2000.
- [5] Gisèle Fischer-Servi. *Axiomatizations for some intuitionistic modal logics*. Rendiconti del Seminario Matematico della Università Politecnica di Torino, 42(3):179-194, 1984.
- [6] Gordon D. Plotkin and Colin P. Stirling. *A framework for intuitionistic modal logic*. In J. Y. Halpern, editor, 1st Conference on Theoretical Aspects of Reasoning About Knowledge. Morgan Kaufmann, 1986.
- [7] Dov M. Gabbay. *Labelled Deductive Systems*. Clarendon Press, 1996.