

Decomposing labelled proof theory for intuitionistic modal logic

Sonia Marin*
IT-Universitetet i København
Denmark

Marianela Morales
Universidad Nacional de Córdoba
Argentina

Lutz Straßburger
Inria Saclay & LIX
France

ACM Reference Format:

Sonia Marin, Marianela Morales, and Lutz Straßburger. 2018. Decomposing labelled proof theory for intuitionistic modal logic. In *Proceedings of . ACM*, New York, NY, USA, 1 page. <https://doi.org/>

Labelled deduction has been proposed by Gabbay [7] in the 80's as a unifying framework throughout proof theory in order to provide proof systems for a wide range of logics. For modal logics it can take for example the form of labelled natural deduction and labelled sequent systems, as used by Simpson [3], Vigano [4] and Negri [2].

These formalisms make explicit use not only of labels, but also of relational atoms referring to the accessibility relation of a Kripke model. In this short note we propose a system that represents both the *accessibility relation* (for modal logics) and the *preorder relation* (for intuitionistic logic), using the full power of the bi-relational semantics for intuitionistic modal logics, and developing fully the idea of [1].

A *bi-relational frame* [5] \mathcal{B} is a triple $\langle W, R, \leq \rangle$ of a non-empty set of worlds W equipped with an accessibility relation R and a preorder \leq , satisfying:

- (F₁) For all worlds x, y, z , if xRy and $y \leq z$, there exists a u such that $x \leq u$ and uRz .
- (F₂) For all worlds x, y, z , if xRy and $x \leq z$, there exists a u such that $y \leq u$ and zRu .

Echoing this construction, we define our two-sided intuitionistic labelled sequents to be of the form $\mathcal{B}, \mathcal{L} \Rightarrow \mathcal{R}$ where \mathcal{B} denotes a set of relational atoms xRy and preorder atoms $x \leq y$, and \mathcal{L} and \mathcal{R} are multiset of labelled formulas $x : A$ (for x and y taken from the set of labels and A an intuitionistic modal formula).

Furthermore, our system has to incorporate the two semantic conditions into deductive rules as follows:

$$F_1 \frac{\mathcal{B}, xRy, y \leq z, x \leq u, uRz, \mathcal{L} \Rightarrow \mathcal{R}}{\mathcal{B}, xRy, y \leq z, \mathcal{L} \Rightarrow \mathcal{R}} u \text{ fresh}$$

$$F_2 \frac{\mathcal{B}, xRy, x \leq z, y \leq u, zRu, \mathcal{L} \Rightarrow \mathcal{R}}{\mathcal{B}, xRy, x \leq z, \mathcal{L} \Rightarrow \mathcal{R}} u \text{ fresh}$$

In the intuitionistic setting, the validity of a modal formula has to be defined using both the R and the \leq relation as: $x \Vdash \Box A$ iff for all y and z s.t. $x \leq y$ and yRz , $z \Vdash A$.

Again, our system reflects exactly this definition in the rules introducing the \Box -operator:

$$\Box_L \frac{\mathcal{B}, x \leq y, yRz, \mathcal{L}, x : \Box A, z : A \Rightarrow \mathcal{R}}{\mathcal{B}, \mathcal{L}, x \leq y, yRz, x : \Box A \Rightarrow \mathcal{R}}$$

$$\Box_R \frac{\mathcal{B}, x \leq y, yRz, \mathcal{L} \Rightarrow \mathcal{R}, z : A}{\mathcal{B}, \mathcal{L} \Rightarrow \mathcal{R}, x : \Box A} y, z \text{ fresh}$$

$$\text{id} \frac{}{x \leq w, w \leq w', w'Rv, v \leq v', v'Ru, w' \leq t, tRv', w \leq t, tRu, w : \Box A, u : A \Rightarrow}$$

$$\Box_L \frac{}{x \leq w, w \leq w', w'Rv, v \leq v', v'Ru, w' \leq t, tRv', w \leq t, tRu, w : \Box A \Rightarrow u :}$$

$$\blacksquare_4 \frac{}{x \leq w, w \leq w', w'Rv, v \leq v', v'Ru, w' \leq t, tRv', w \leq tw : \Box A \Rightarrow u : A}$$

$$\text{trans} \frac{}{x \leq w, w \leq w', w'Rv, v \leq v', v'Ru, w' \leq t, tRv', w : \Box A \Rightarrow u : A}$$

$$F_1 \frac{}{x \leq w, w \leq w', w'Rv, v \leq v', v'Ru, w : \Box A \Rightarrow u : A}$$

$$\Box_R \frac{}{x \leq w, w : \Box A \Rightarrow w : \Box \Box A}$$

$$\supset_R \frac{}{x : \Box A \supset \Box \Box A}$$

$$\text{id} \frac{}{x \leq w, wRv, vRu, u \leq u', wRu'u : A \Rightarrow w : \Diamond A, u' : A}$$

$$\Diamond_R \frac{}{x \leq w, wRv, vRu, u \leq u', wRu'u : A \Rightarrow w : \Diamond A}$$

$$\blacklozenge_4 \frac{}{x \leq w, wRv, vRu, u : A \Rightarrow w : \Diamond A}$$

$$\Diamond_L \frac{}{x \leq w, w : \Diamond \Diamond A \Rightarrow w : \Diamond A}$$

$$\supset_R \frac{}{x : \Diamond \Diamond A \supset \Diamond A}$$

References

- [1] Paolo Maffezioli, Alberto Naibo, and Sara Negri. *The Church-Fitch knowability paradox in the light of structural proof theory*. Synthese, 190(14):2677-2716, 2013.
- [2] Sara Negri. *Proof analysis in modal logics*. Journal of Philosophical Logic, 34:507-544, 2005.
- [3] Alex Simpson. *The Proof Theory and Semantics of Intuitionistic Modal Logic*. PhD thesis, University of Edinburgh, 1994.
- [4] Luca Viganò. *Labelled Non-Classical Logic*. Kluwer Academic Publisher, 2000.
- [5] Gisèle Fischer-Servi. *Axiomatizations for some intuitionistic modal logics*. Rendiconti del Seminario Matematico della Università Politecnica di Torino, 42(3):179-194, 1984.
- [6] Gordon D. Plotkin and Colin P. Stirling. *A framework for intuitionistic modal logic*. In J. Y. Halpern, editor, 1st Conference on Theoretical Aspects of Reasoning About Knowledge. Morgan Kaufmann, 1986.
- [7] Dov M. Gabbay. *Labelled Deductive Systems*. Clarendon Press, 1996.

*Funded by the Qatar National Research Council project MetaCLF nb. xxx.