

Course on Proof Theory - Lecture 5

Beyond classical logic

Gianluca Curzi, Marianna Girlando

University of Birmingham

Midlands Graduate School
Nottingham, 10-14 April 2022

From classical logic to intuitionistic logic

Semantics of intuitionistic logic

Sequent calculus LJ

Meta-properties of LJ

Automated proof search

References

On classical reasoning

- ▶ Classical logic based on the notion of **truth**, where **not false** = **true**:

$$A \vee \neg A \quad (\textit{excluded middle}) \qquad \neg\neg A \rightarrow A \quad (\textit{involutivity})$$

On classical reasoning

- ▶ Classical logic based on the notion of **truth**, where **not false** = **true**:

$$A \vee \neg A \quad (\textit{excluded middle}) \qquad \neg\neg A \rightarrow A \quad (\textit{involutivity})$$

- ▶ Truth-value of a formula is “**absolute**” (either **true** or **false**), whether or not we know it, prove it, or verify it in any possible way. E.g.

There are seven 7's in a row somewhere in the decimal representation of the real number R

Intuitionistic logic



Figure: From left, L.E.J. Brouwer (1881-1966) and Arend Heyting (1898-1980).

- ▷ **Intuitionistic logic**: there is no absolute truth, there is only the knowledge and intuitive construction of the idealised mathematician. A logical judgement is only considered “true” if we can **verify its correctness**.
- ▷ Intuitionistic logic **rejects** the excluded middle and involutivity.

An example

Theorem. There exist irrational numbers x and y , such that x^y is rational.

First proof. If $\sqrt{2}^{\sqrt{2}}$ is a rational number then we take $x = y = \sqrt{2}$; otherwise, we take $x = \sqrt{2}^{\sqrt{2}}$ and $y = \sqrt{2}$.

An example

Theorem. There exist irrational numbers x and y , such that x^y is rational.

First proof. If $\sqrt{2}^{\sqrt{2}}$ is a rational number then we take $x = y = \sqrt{2}$; otherwise, we take $x = \sqrt{2}^{\sqrt{2}}$ and $y = \sqrt{2}$.

Drawback: which of the two possibilities actually holds?

An example

Theorem. There exist irrational numbers x and y , such that x^y is rational.

First proof. If $\sqrt{2}^{\sqrt{2}}$ is a rational number then we take $x = y = \sqrt{2}$; otherwise, we take $x = \sqrt{2}^{\sqrt{2}}$ and $y = \sqrt{2}$.

Drawback: which of the two possibilities actually holds?

Second proof. Take $x = \sqrt{2}$ and $y = 2 \log_2 3$, so that $x^y = 3$.

Benefit: Proof is **constructive**, it exhibits a “witness”.

From classical logic to intuitionistic logic

Semantics of intuitionistic logic

Sequent calculus LJ

Meta-properties of LJ

Automated proof search


References

Kripke semantics

(Intuitionistic) Kripke model. (W, \leq, \Vdash) where:

- ▷ W nonempty set of states (or worlds)
- ▷ \leq is partial order on W
- ▷ \Vdash called forcing relation is binary relation between states and propositional variables such that:

$$w \leq w' \quad \text{and} \quad w \Vdash p \quad \implies \quad w' \Vdash p$$




$$w \Vdash p$$

(monotonicity)

Idea:

- ▷ $w \in W$ represents a state of knowledge
- ▷ \leq represents gaining of knowledge
- ▷ \Vdash determines which facts are true at a given state.



Kripke semantics, continued

Given (W, \leq, \Vdash) , we extend \Vdash to any formula:

with

 $\omega \vdash A$
$$w \Vdash A \wedge B \quad \text{iff} \quad w \Vdash A \text{ and } w \Vdash \underline{B}$$
$$w \Vdash A \vee B \quad \text{iff} \quad w \Vdash A \text{ **or** } w \Vdash B$$
$$w \Vdash A \rightarrow B \quad \text{iff} \quad \text{for all } w' \geq w, \quad (w' \Vdash A \text{ implies } w' \Vdash B)$$
 $w \not\perp \perp$
$$\Gamma \Vdash A \text{ iff for every } (W, \leq, \Vdash) \text{ and every } w \in W, \text{ if } w \Vdash \Gamma \text{ then } w \Vdash A.$$


Kripke semantics, continued

Given (W, \leq, \Vdash) , we extend \Vdash to any formula:

$w \Vdash A \wedge B$ iff $w \Vdash A$ **and** $w \Vdash B$

$w \Vdash A \vee B$ iff $w \Vdash A$ **or** $w \Vdash B$

$w \Vdash A \rightarrow B$ iff **for all** $w' \geq w$, $w' \Vdash A$ **implies** $w' \Vdash B$

$w \nVdash \perp$

$\Gamma \Vdash A$ iff for every (W, \leq, \Vdash) and every $w \in W$, if $w \Vdash \Gamma$ then $w \Vdash A$.

▷ **Monotonicity** “lifts” to formulas: $w \leq w'$ and $w \Vdash A \implies w' \Vdash A$



Kripke semantics, continued

Given (W, \leq, \Vdash) , we extend \Vdash to any formula:

$w \Vdash A \wedge B$ iff $w \Vdash A$ **and** $w \Vdash B$

$w \Vdash A \vee B$ iff $w \Vdash A$ **or** $w \Vdash B$

$w \Vdash A \rightarrow B$ iff **for all** $w' \geq w$, $w' \Vdash A$ **implies** $w' \Vdash B$

$w \nVdash \perp$

$\Gamma \Vdash A$ iff for every (W, \leq, \Vdash) and every $w \in W$, if $w \Vdash \Gamma$ then $w \Vdash A$.

▷ **Monotonicity** “lifts” to formulas: $w \leq w'$ and $w \Vdash A \implies w' \Vdash A$

▷ Intuitionistic logic **breaks dualities** of classical logic:

▷ Negation is **not involutive**:

$w \Vdash \neg A := A \rightarrow \perp$ iff $w' \nVdash A$ for all $w' \geq w$

Kripke semantics, continued

Given (W, \leq, \Vdash) , we extend \Vdash to any formula:

$w \Vdash A \wedge B$ iff $w \Vdash A$ **and** $w \Vdash B$

$w \Vdash A \vee B$ iff $w \Vdash A$ **or** $w \Vdash B$

$w \Vdash A \rightarrow B$ iff **for all** $w' \geq w$, $w' \Vdash A$ **implies** $w' \Vdash B$

$w \nVdash \perp$

$\Gamma \Vdash A$ iff for every (W, \leq, \Vdash) and every $w \in W$, if $w \Vdash \Gamma$ then $w \Vdash A$.

▷ **Monotonicity** “lifts” to formulas: $w \leq w'$ and $w \Vdash A \implies w' \Vdash A$

▷ Intuitionistic logic **breaks dualities** of classical logic:

▷ Negation is **not involutive**:


$w \Vdash \neg A := A \rightarrow \perp$ iff $w' \nVdash A$ for all $w' \geq w$

▷ The equivalence $w \Vdash A \rightarrow B \iff w \Vdash \neg A \vee B$ **does not** hold.

An example

Example. Let (W, \leq, \Vdash) be such that:

- ▷ $W = \{w, w', w''\}$
- ▷ $w \leq w', w \leq w''$ with w', w'' incomparable.
- ▷ $w' \Vdash p, \quad w'' \Vdash q, \quad \boxed{w \nVdash p}, \quad \boxed{w \nVdash q}$



We have $w \Vdash \neg\neg(p \vee q)$ and $w \Vdash (p \rightarrow q) \rightarrow q$. Notice that $w \nVdash p \vee \neg p$.

From classical logic to intuitionistic logic

Semantics of intuitionistic logic

Sequent calculus LJ LK

Meta-properties of LJ

Automated proof search

References

Sequent calculus for intuitionistic logic, naive attempt

LJ = restriction of LK where all sequents have exactly **one formula** on the right-hand side of the sequent.

$$\begin{array}{c} \text{init} \frac{}{A, \Gamma \vdash A} \\ \wedge_L \frac{A_1, A_2, \Gamma \vdash C}{A_1 \wedge A_2, \Gamma \vdash C} \\ \vee_L \frac{A_1, \Gamma \vdash C \quad A_2, \Gamma \vdash C}{A_1 \vee A_2, \Gamma \vdash C} \\ \rightarrow_L \frac{\Gamma \vdash A \quad B, \Gamma \vdash C}{A \rightarrow B, \Gamma \vdash C} \\ \text{cut} \frac{\Gamma \vdash A \quad A, \Gamma \vdash C}{\Gamma \vdash C} \end{array} \quad \begin{array}{c} \perp_L \frac{}{\perp, \Gamma \vdash A} \\ \wedge_R \frac{\Gamma \vdash A \quad \Gamma \vdash B}{\Gamma \vdash A \wedge B} \\ \vee_R \frac{\Gamma \vdash A_i}{\Gamma \vdash A_1 \vee A_2} \\ \rightarrow_R \frac{A, \Gamma \vdash B}{\Gamma \vdash A \rightarrow B} \end{array}$$

Examples

Double negation law is not intuitionistically (cut-free) provable ([Exercise](#)).

Examples

Double negation law is not intuitionistically (cut-free) provable ([Exercise](#)).

Peirce's law is not intuitionistically (cut-free) provable:

$$\begin{array}{c} \text{init} \frac{}{A \vdash A} \\ \rightarrow_R \frac{}{\vdash A, A \rightarrow B} \quad \text{init} \frac{}{A \vdash A} \\ \rightarrow_L \frac{}{(A \rightarrow B) \rightarrow A \vdash A} \\ \rightarrow_R \frac{}{\vdash ((A \rightarrow B) \rightarrow A) \rightarrow A} \end{array}$$

A counterexample to completeness

$$\frac{\frac{\frac{\text{???}}{\vdash A \vee (A \rightarrow \perp)}}{\rightarrow_L \frac{(A \vee (A \rightarrow \perp)) \rightarrow \perp \vdash \perp}{\rightarrow_R \vdash ((A \vee (A \rightarrow \perp)) \rightarrow \perp) \rightarrow \perp}} \text{init } \frac{}{\perp \vdash \perp}}{\vdash ((A \vee (A \rightarrow \perp)) \rightarrow \perp) \rightarrow \perp}$$

$\underbrace{\hspace{10em}}_{\neg\neg(A \vee \neg A)}$

A counterexample to completeness

$$\begin{array}{c}
 \text{init} \frac{}{((A \vee (A \rightarrow \perp)) \rightarrow \perp), A \vdash A} \\
 \vee_R \frac{}{((A \vee (A \rightarrow \perp)) \rightarrow \perp), A \vdash A \vee (A \rightarrow \perp)} \quad \text{init} \frac{}{\perp \vdash \perp} \\
 \rightarrow_L \frac{}{((A \vee (A \rightarrow \perp)) \rightarrow \perp), A \vdash \perp} \\
 \rightarrow_R \frac{}{((A \vee (A \rightarrow \perp)) \rightarrow \perp) \vdash A \rightarrow \perp} \\
 \vee_R \frac{}{((A \vee (A \rightarrow \perp)) \rightarrow \perp) \vdash A \vee (A \rightarrow \perp)} \quad \text{init} \frac{}{\perp \vdash \perp} \\
 \rightarrow_L \frac{}{(A \vee (A \rightarrow \perp)) \vdash \perp} \\
 \Rightarrow \frac{}{(A \vee (A \rightarrow \perp)) \vdash \perp} \\
 \rightarrow_R \frac{}{\vdash ((A \vee (A \rightarrow \perp)) \rightarrow \perp) \rightarrow \perp} \\
 \underbrace{\vdash ((A \vee (A \rightarrow \perp)) \rightarrow \perp) \rightarrow \perp}_{\neg\neg(A \vee \neg A)}
 \end{array}$$

Idea: To prove intuitionistically valid formula $\neg\neg(A \vee \neg A)$ we need to apply twice \rightarrow_L to the formula $((A \vee (A \rightarrow \perp)) \rightarrow \perp)$, so we need to “save” a copy of it in our set of hypothesis.

Sequent calculus for intuitionistic logic

$$\begin{array}{c} \text{init} \frac{}{A, \Gamma \vdash A} \qquad \perp_{\text{L}} \frac{}{\perp, \Gamma \vdash A} \\[10pt] \wedge_{\text{L}} \frac{A_1, A_2, \Gamma \vdash C}{A_1 \wedge A_2, \Gamma \vdash C} \qquad \wedge_{\text{R}} \frac{\Gamma \vdash A \quad \Gamma \vdash B}{\Gamma \vdash A \wedge B} \\[10pt] \vee_{\text{L}} \frac{A_1, \Gamma \vdash C \quad A_2, \Gamma \vdash C}{A_1 \vee A_2, \Gamma \vdash C} \qquad \vee_{\text{R}} \frac{\Gamma \vdash A_i}{\Gamma \vdash A_1 \vee A_2} \\[10pt] \rightarrow_{\text{L}} \frac{\Gamma \vdash A \quad B, \Gamma \vdash C}{A \rightarrow B, \Gamma \vdash C} \qquad \rightarrow_{\text{R}} \frac{A, \Gamma \vdash B}{\Gamma \vdash A \rightarrow B} \\[10pt] \text{cut} \frac{\Gamma \vdash A \quad A, \Gamma \vdash C}{\Gamma \vdash C} \end{array}$$

Sequent calculus for intuitionistic logic

$$\frac{A[\frac{t}{x}], (\forall x. A) \Rightarrow}{\forall x. A \Rightarrow}$$

$$\begin{array}{c}
 \text{init} \frac{}{A, \Gamma \vdash A} \qquad \perp_L \frac{}{\perp, \Gamma \vdash A} \\
 \wedge_L \frac{A_1, A_2, \Gamma \vdash C}{A_1 \wedge A_2, \Gamma \vdash C} \qquad \wedge_R \frac{\Gamma \vdash A \quad \Gamma \vdash B}{\Gamma \vdash A \wedge B} \\
 \vee_L \frac{A_1, \Gamma \vdash C \quad A_2, \Gamma \vdash C}{A_1 \vee A_2, \Gamma \vdash C} \qquad \vee_R \frac{\Gamma \vdash A_i}{\Gamma \vdash A_1 \vee A_2} \\
 \rightarrow_L \frac{A \rightarrow B, \Gamma \vdash A \quad B, \Gamma \vdash C}{A \rightarrow B, \Gamma \vdash C} \qquad \rightarrow_R \frac{A, \Gamma \vdash B}{\Gamma \vdash A \rightarrow B} \\
 \text{cut} \frac{\Gamma \vdash A \quad A, \Gamma \vdash C}{\Gamma \vdash C}
 \end{array}$$

N.B. Recall that $A \rightarrow B$ is a **primitive** connective ($A \rightarrow B \neq \neg A \vee B$).

From classical logic to intuitionistic logic

Semantics of intuitionistic logic

Sequent calculus LJ

Meta-properties of LJ

Automated proof search

References

Soundness and completeness for LJ

Theorem. LJ is **sound and complete** for Kripke models.

$$\underbrace{\Gamma \vdash_{\text{LJ}} A}_{\text{LJ}} \iff \Gamma \Vdash A$$

Soundness and completeness for LJ

Theorem. LJ is **sound and complete** for Kripke models.

$$\Gamma \vdash_{\text{LJ}} A \quad \Longleftrightarrow \quad \Gamma \Vdash A$$

Moreover, **cut-elimination still holds!** **Corollaries:**

▷ **Disjunction property:** $\vdash_{\text{LJ}} A \vee B$ \implies $\vdash_{\text{LJ}} A$ or $\vdash_{\text{LJ}} B$

Soundness and completeness for LJ

Theorem. LJ is **sound and complete** for Kripke models.

$$\Gamma \vdash_{\text{LJ}} A \quad \Longleftrightarrow \quad \Gamma \Vdash A$$

Moreover, **cut-elimination still holds!** **Corollaries:**

▷ **Disjunction property:** $\vdash_{\text{LJ}} A \vee B \implies \vdash_{\text{LJ}} A \text{ or } \vdash_{\text{LJ}} B$

▷ **Existential property:** $\vdash_{\text{LJ}} \exists x.A \implies \vdash_{\text{LJ}} A[t/x]$, for some term t

Soundness and completeness for LJ

Theorem. LJ is **sound and complete** for Kripke models.

$$\Gamma \vdash_{\text{LJ}} A \quad \Longleftrightarrow \quad \Gamma \Vdash A$$

Moreover, **cut-elimination still holds!** **Corollaries:**

- ▷ **Disjunction property:** $\vdash_{\text{LJ}} A \vee B \implies \vdash_{\text{LJ}} A$ or $\vdash_{\text{LJ}} B$
- ▷ **Existential property:** $\vdash_{\text{LJ}} \exists x.A \implies \vdash_{\text{LJ}} A[t/x]$, for some term t

Example.

$$\vdash \exists x.(\underbrace{\quad}_{\text{triangle}} + x = \underline{5}) \quad \rightsquigarrow \quad \underbrace{\quad}_{\text{triangle}} \quad \frac{\vdash \underline{3} + \underline{2} = \underline{5}}{\exists_{\mathbf{R}} \vdash \underline{\mathbf{3}}x.(\underline{3} + \underline{\mathbf{x}} = \underline{5})}$$


LK vs LJ

- ▶ LJ proves less formulas than LK (e.g. $A \vee \neg A$) ...

LK vs LJ

- ▶ LJ proves less formulas than LK (e.g. $A \vee \neg A$) ... but we can “see” classical logic within intuitionistic logic by wearing special glasses, called **Glivenko's glasses**, which turn any formula A into $\neg\neg A$.

Glivenko's theorem (1929). $\vdash_{LK} A \iff \vdash_{LJ} \neg\neg A$.



- ▶ LJ proves less formulas than LK (e.g. $A \vee \neg A$) ... but we can “see” classical logic within intuitionistic logic by wearing special glasses, called **Glivenko’s glasses**, which turn any formula A into $\neg\neg A$.

Glivenko’s theorem (1929). $\vdash_{LK} A \iff \vdash_{LJ} \neg\neg A$.


- ▶ Gödel and Gentzen’s glasses, or **double negation translation**:

$$\begin{aligned}
 p^G &:= \neg\neg p & (-)^G \\
 (\neg A)^G &:= \neg A^G \\
 (A \wedge B)^G &:= A^G \wedge B^G \\
 (A \vee B)^G &:= \neg(\neg A^G \wedge \neg B^G) \\
 (A \rightarrow B)^G &:= A^G \rightarrow B^G
 \end{aligned}$$

Gödel & Gentzen’s theorem (1933). $\vdash_{LK} A \iff \vdash_{LJ} A^G$.

Intermediate logics

- ▶ Are there logics in-between LJ and LK?



Intermediate logics

- Are there logics in-between LJ and LK?

$\text{LJ} + \{ \dots \}$

$\text{LJ} \xrightarrow[\uparrow]{???} \text{LK}$

$B \vee \neg B$

$B := \neg A$

- Yes ... and they are called intermediate logics!

Weak excluded middle logic $:= \text{LJ} + \{ \neg A \vee \neg\neg A \}$

Gödel-Dummett logic $:= \text{LJ} + \{ (A \rightarrow B) \vee (B \rightarrow A) \}$

\vdots

$(A \rightarrow B) \vee (B \rightarrow A)$

Intermediate logics

- ▶ Are there logics in-between LJ and LK?

$$\text{LJ} \overset{???}{\dashrightarrow} \text{LK}$$

- ▶ Yes ... and they are called **intermediate logics**!

Weak excluded middle logic $:= \text{LJ} + \{\neg A \vee \neg\neg A\}$

Gödel-Dummett logic $:= \text{LJ} + \{(A \rightarrow B) \vee (B \rightarrow A)\}$

\vdots

- ▶ **Question for the audience:** How many intermediate logics are there?

From classical logic to intuitionistic logic

Semantics of intuitionistic logic

Sequent calculus LJ


Meta-properties of LJ

Automated proof search

References


Automated proof search

How to check if A is a theorem of classical or intuitionistic logic?




Automated proof search

How to check if A is a theorem of classical or intuitionistic logic?



Sequent calculus can be used to implement a **decision procedure**, that is, a terminating and effective procedure to check if A is a theorem in LK or LJ.



Desirable features for root-first proof search, I

Wish 1: Don't have to make a "guess" when going from the conclusion to the premiss of a rule.

Cut-free sequent calculus

$$\frac{\Gamma \Rightarrow \Delta, \bar{A} \quad \bar{A}, \Gamma \Rightarrow \Delta}{\Gamma \Rightarrow \Delta}$$

$$\begin{array}{ccc} \Gamma \vdash_S \Delta & \begin{array}{c} \text{soundness} \\ \Longleftrightarrow \\ \text{completeness} \end{array} & \Gamma \models \Delta \\ \Downarrow & & \\ \Gamma \vdash_{S^-} \Delta & & \end{array}$$

for $S = \{\text{LK}, \text{LJ}\}$ and S^- denoting $S \setminus \{\text{cut}\}$

$$\begin{array}{l} \text{Soundness} \quad \Gamma \vdash_S \Delta \implies \Gamma \models \Delta \\ \text{Completeness} \quad \Gamma \models \Delta \implies \Gamma \vdash_S \Delta \implies \Gamma \vdash_{S^-} \Delta \end{array}$$

Desirable features for root-first proof search, II

Wish 2: don't have to make a “choice” when going from the conclusion to the premiss of a rule

\leadsto choices introduce **backtrack points** in proof search \vee

Example

$$\left(\begin{array}{c} \text{v}_{R1} \frac{\Gamma \vdash \Delta, A}{\Gamma \vdash \Delta, A \vee B} \\ \text{v}_{R2} \frac{\Gamma \vdash \Delta, B}{\Gamma \vdash \Delta, A \vee B} \end{array} \right) \quad \text{v}_R \frac{\Gamma \vdash \Delta, A, B}{\Gamma \vdash \Delta, A \vee B}$$

$$\text{v}_{L2} \frac{a \vdash \overset{o}{b}}{a \vdash a \vee b}$$

Desirable features for root-first proof search, II

Wish 2: don't have to make a “choice” when going from the conclusion to the premiss of a rule

↪ choices introduce **backtrack points** in proof search

	multiplicative	additive
right	$\wedge_R \frac{\Gamma \vdash \Delta, A \quad \Gamma' \vdash \Delta', B}{\Gamma, \Gamma' \vdash \Delta, \Delta', A \wedge B}$	$\wedge_R \frac{\Gamma \vdash \Delta, A \quad \Gamma \vdash \Delta, B}{\Gamma \vdash \Delta, A \wedge B}$
left	$\wedge_L \frac{A, B, \Gamma \vdash \Delta}{A \wedge B, \Gamma \vdash \Delta}$	$\wedge_L \frac{\Gamma, A_i \vdash \Delta}{\Gamma, A_1 \wedge A_2 \vdash \Delta}$

When possible:

- ▷ Choose the multiplicative version of one premisses-rules
- ▷ Choose the additive version of the two-premisses rules

Classical first-order logic

LK^- meet our two desiderata. Is this enough to ensure termination of root-first proof search? **no**

$$\forall_L \frac{A[t/x], \forall x.A, \Gamma \vdash \Delta}{\forall x.A, \Gamma \vdash \Delta} \quad \forall_R \frac{\Gamma \vdash \Delta, A[y/x]}{\Gamma \vdash \Delta, \forall x.A} * \quad \exists_L \frac{A[y/x], \Gamma \vdash \Delta}{\exists x.A, \Gamma \vdash \Delta} * \quad \exists_R \frac{\Gamma \vdash \Delta, A[t/x], \exists x.A}{\Gamma \vdash \Delta, \exists x.A}$$

* y does not occur free in Γ, Δ, A

$$\begin{array}{c} \vdots \\ \forall_R \frac{\vdash \exists x \forall y (P(x, y)), P(z, k), P(x, z)}{\vdash \exists x \forall y (P(x, y)), \forall y (P(\underline{z}, y)), P(x, z)} \\ \exists_R \frac{\vdash \exists x \forall y (P(x, y)), \forall y (P(\underline{z}, y)), P(x, z)}{\vdash \exists x \forall y (P(x, y)), P(x, \underline{z})} \\ \forall_R \frac{\vdash \exists x \forall y (P(x, y)), P(x, \underline{z})}{\vdash \exists x \forall y (P(x, y)), \forall y (P(x, y))} \\ \exists_R \frac{\vdash \exists x \forall y (P(x, y)), \forall y (P(x, y))}{\vdash \exists x \forall y (P(x, y))} \end{array}$$

Classical first-order logic

LK⁻ meet our two desiderata. Is this enough to ensure termination of root-first proof search? **no**

$$\begin{array}{c}
 \frac{A[t/x], \forall x.A, \Gamma \vdash \Delta}{\forall x.A, \Gamma \vdash \Delta} \quad \forall_L \qquad \frac{\Gamma \vdash \Delta, A[y/x]}{\Gamma \vdash \Delta, \forall x.A} \quad \forall_R \qquad \frac{A[y/x], \Gamma \vdash \Delta}{\exists x.A, \Gamma \vdash \Delta} \quad \exists_L \qquad \frac{\Gamma \vdash \Delta, A[t/x], \exists x.A}{\Gamma \vdash \Delta, \exists x.A} \quad \exists_R \\
 * \quad y \text{ does not occur free in } \Gamma, \Delta, A \\
 \\
 \vdots \\
 \frac{\vdash \exists x \forall y (P(x, y)), P(z, k), P(x, z)}{\vdash \exists x \forall y (P(x, y)), \forall y (P(z, y)), P(x, z)} \quad \forall_R \\
 \frac{\vdash \exists x \forall y (P(x, y)), \forall y (P(z, y)), P(x, z)}{\vdash \exists x \forall y (P(x, y)), P(x, z)} \quad \exists_R \\
 \frac{\vdash \exists x \forall y (P(x, y)), P(x, z)}{\vdash \exists x \forall y (P(x, y)), \forall y (P(x, y))} \quad \forall_R \\
 \frac{\vdash \exists x \forall y (P(x, y)), \forall y (P(x, y))}{\vdash \exists x \forall y (P(x, y))} \quad \exists_R
 \end{array}$$

First-order logic is **semi-decidable**: for every formula A ,

- ▷ If A is a theorem, the algorithm produces a proof;
- ▷ Otherwise, the algorithm either produces a failed proof or does not terminate.

Classical propositional logic

Propositional LK^- meets our two desiderata. Is this enough to ensure termination of root-first proof search? yes

$$\begin{array}{c} \text{init} \frac{}{p, \Gamma \vdash \Delta, p} \quad \neg^L \frac{\Gamma \vdash \Delta, A}{\neg A, \Gamma \vdash \Delta} \quad \neg^R \frac{A, \Gamma \vdash \Delta}{\Gamma \vdash \Delta, \neg A} \\[10pt] \wedge^L \frac{A, B, \Gamma \vdash \Delta}{A \wedge B, \Gamma \vdash \Delta} \quad \wedge^R \frac{\Gamma \vdash \Delta, A \quad \Gamma \vdash \Delta, B}{\Gamma \vdash \Delta, A \wedge B} \quad \vee^L \frac{A, \Gamma \vdash \Delta \quad B, \Gamma \vdash \Delta}{A \vee B, \Gamma \vdash \Delta} \quad \vee^R \frac{\Gamma \vdash \Delta, A, B}{\Gamma \vdash \Delta, A \vee B} \\[10pt] \rightarrow^L \frac{\Gamma \vdash \Delta, A \quad B, \Gamma \vdash \Delta}{A \rightarrow B, \Gamma \vdash \Delta} \quad \rightarrow^R \frac{A, \Gamma \vdash \Delta, B}{\Gamma \vdash \Delta, A \rightarrow B} \end{array}$$

Proof search strategy

- ▶ Apply the rules of LK^- in whatever order
- ▶ The calculus has the subformula property: all formulas get decomposed into smaller ones
- ▶ Proof search comes to an end in a finite number of steps.

Classical propositional logic is **decidable**.

Intuitionistic propositional logic

Propositional LJ^- is cut-free, but we have to make choices on some rules.

Is this enough to ensure termination of root-first proof search? **no**

$$\begin{array}{c} \text{init} \frac{}{p, \Gamma \vdash p} \quad \perp_{\text{L}} \frac{}{\perp, \Gamma \vdash A} \\ \wedge_{\text{L}} \frac{A, B, \Gamma \vdash C}{A \wedge B, \Gamma \vdash C} \quad \wedge_{\text{R}} \frac{\Gamma \vdash A \quad \Gamma \vdash B}{\Gamma \vdash A \wedge B} \quad \vee_{\text{L}} \frac{A, \Gamma \vdash C \quad B, \Gamma \vdash C}{A \vee B, \Gamma \vdash C} \quad \vee_{\text{R}}^i \frac{\Gamma \vdash A_i}{\Gamma \vdash A_1 \vee A_2} \quad i \in \{1, 2\} \\ \rightarrow_{\text{L}} \frac{A \rightarrow B, \Gamma \vdash A \quad B, \Gamma \vdash C}{A \rightarrow B, \Gamma \vdash C} \quad \rightarrow_{\text{R}} \frac{A, \Gamma \vdash B}{\Gamma \vdash A \rightarrow B} \end{array}$$

Intuitionistic propositional logic

Propositional LJ^- is cut-free, but we have to make choices on some rules.

Is this enough to ensure termination of root-first proof search? **no**

$$\begin{array}{c}
 \text{init} \frac{}{p, \Gamma \vdash p} \quad \perp_L \frac{}{\perp, \Gamma \vdash A} \\
 \wedge_L \frac{A, B, \Gamma \vdash C}{A \wedge B, \Gamma \vdash C} \quad \wedge_R \frac{\Gamma \vdash A \quad \Gamma \vdash B}{\Gamma \vdash A \wedge B} \quad \vee_L \frac{A, \Gamma \vdash C \quad B, \Gamma \vdash C}{A \vee B, \Gamma \vdash C} \quad \vee_R i \frac{\Gamma \vdash A_i}{\Gamma \vdash A_1 \vee A_2} \quad i \in \{1, 2\} \\
 \rightarrow_L \frac{A \rightarrow B, \Gamma \vdash A \quad B, \Gamma \vdash C}{A \rightarrow B, \Gamma \vdash C} \quad \rightarrow_R \frac{A, \Gamma \vdash B}{\Gamma \vdash A \rightarrow B}
 \end{array}$$

$$\begin{array}{c}
 \vdots \\
 \frac{a \rightarrow \perp \vdash a \quad \perp \vdash a}{\rightarrow_L} \\
 \frac{\rightarrow_L \quad a \rightarrow \perp \vdash a \quad \perp \vdash a}{\rightarrow_L} \\
 \frac{\rightarrow_L \quad a \rightarrow \perp \vdash a \quad \perp \vdash \perp}{\rightarrow_L} \\
 \frac{\rightarrow_L \quad a \rightarrow \perp \vdash \perp}{\rightarrow_R} \vdash (a \rightarrow \perp) \rightarrow \perp
 \end{array}$$

Solution

Proof search strategy [Troelstra, Schwichtenberg, Basic proof theory]

- ▷ Apply rule \rightarrow_L after all the other rules.
- ▷ Before applying rule \rightarrow_L to a sequent $\Gamma \vdash \Delta$, check if there is a sequent in the branch containing the same formulas as $\Gamma \vdash \Delta$.
- ▷ The calculus has the subformula property: all formulas get decomposed into smaller ones.
- ▷ Proof search comes to an end in a finite number of steps.
- ▷ If a proof has not been found, backtrack on all choice points.

Intuitionistic propositional logic is **decidable**.

Many other solutions exist: [Dyckhoff, Intuitionistic decision procedures since Gentzen, 2015]

11

From classical logic to intuitionistic logic

Semantics of intuitionistic logic

Sequent calculus LJ

Meta-properties of LJ

Automated proof search

References

References

- ▷ Dyckhoff, R. (2015). Intuitionistic decision procedures since Gentzen. In: Advances in proof theory.
- ▷ Buss, S. R., editor (1998). Handbook of Proof Theory, volume 137 of Studies in Logic and the Foundations of Mathematics. Elsevier.
- ▷ Negri, S. and von Plato, J. (2001). Structural proof theory. Cambridge University Press.
- ▷ Smullyan, R. M. (1968). First-Order Logic. Springer-Verlag.
- ▷ Troelstra, A. S. and Schwichtenberg, H. (1996). Basic Proof Theory. Cambridge University Press, New York, NY, USA.