

# Investigating unconventional superconductivity in the 2D Hubbard-Kanamori model using Functional Renormalization Group (FRG)

210003218

February-May 2025

## Contents

<b>1</b>	<b>Abstract</b>	<b>2</b>
<b>2</b>	<b>Introduction</b>	<b>2</b>
<b>3</b>	<b>Theoretical Background</b>	<b>2</b>
3.1	Unconventional superconductivity . . . . .	2
3.1.1	Spin-fluctuation mediated superconductivity . . . . .	2
3.2	Hubbard-Kanamori Model . . . . .	2
3.2.1	Tight Binding Models . . . . .	2
3.2.2	Hubbard Model . . . . .	3
3.2.3	Hubbard-Kanamori Model . . . . .	3
3.3	Theoretical Background in FRG . . . . .	4
3.3.1	Flow equation . . . . .	4
3.3.2	Truncation scheme . . . . .	6
3.3.3	Decoupling of flow equation . . . . .	6
3.3.4	Instability calculation . . . . .	6
<b>4</b>	<b>Computational Methods</b>	<b>6</b>
4.1	Diverge . . . . .	6
4.2	Convergence of Calculation . . . . .	6
4.2.1	Form factor convergence . . . . .	6
4.2.2	Number of k points convergence . . . . .	6
4.3	Calculation of Susceptibilities . . . . .	6
4.3.1	Nesting vectors . . . . .	6
4.3.2	Superconducting order parameters . . . . .	6

<b>5</b>	<b>Results and discussion</b>	<b>6</b>
5.1	1NN Model . . . . .	6
5.1.1	Superconductivity in the 1NN Model . . . . .	9
5.1.2	Magnetic stripes in the 1NN Model . . . . .	9
5.2	Effect of next-nearest neighbour hopping (1NNN model) . . . . .	9
5.2.1	Superconductivity in the 1NNN Model . . . . .	12
5.2.2	Magnetic stripes in the 1NNN model . . . . .	12
5.2.3	Continuous variation of next-nearest neighbour hopping . . . . .	12
5.3	Effect of bi-orbital system (1NN2 model) . . . . .	12
5.3.1	Superconductivity in the 1NN2 model . . . . .	12
5.3.2	Effect of orbital hybridisation in the 1NN2 model . . . . .	12
<b>6</b>	<b>Conclusion and Outlook</b>	<b>12</b>

# 1 Abstract

# 2 Introduction

# 3 Theoretical Background

## 3.1 Unconventional superconductivity

### 3.1.1 Spin-fluctuation mediated superconductivity

## 3.2 Hubbard-Kanamori Model

### 3.2.1 Tight Binding Models

The Tight Binding Model is a central element of condensed matter physics (1). In this model, electrons are bound in orbitals (called sites) around the lattice ions. Due to the overlap between the quantum mechanical wavefunctions that describe these sites, electrons are allowed to 'hop' to neighbouring sites. The probability that this hopping process will occur is given by a tunnelling amplitude, which can be calculated using a hopping integral.

This work is carried out in the tight-binding model framework, where the magnitude of the tunnelling amplitudes are at first treated as free parameters. Here, we build from a simple 2D nearest neighbour hopping Hubbard model (Section. 3.2.2) and investigate the effect of introducing and varying the strength of the next-nearest neighbour hopping amplitude (See Fig. 1). We extend these models further by considering the two-orbital per site case.

$$\hat{H}_{TB}(\underline{\mathbf{R}}) = \sum_{ij\sigma} t_{ij}(\hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma} + h.c) \quad (1)$$




Figure 1: **Two-Dimensional Tight-Binding Models:** Three pannels showing the 1NN, 1NNN and 1NN2 models discussed in Section 5.1, 5.2 and 5.3 respectively. Fig a) shows the Nearest-neighbour hopping case, where  $t$  depicts the hopping amplitude between the neighbouring sites. Fig b) shows the inclusion of the next-nearest neighbour hopping, the magnitude given by  $t'$ . Fig c) Shows the extension to the two-orbital case, depicting the same orbital ( $t_{orb1}$ ,  $t_{orb2}$ ) and different orbital ( $t_{orb1,2}$ ) nearest neighbour hopping. Note that this is just a pictorial representation of the orbitals, and that it does not correspond to a particular choice of orbitals or their real space projection.

### 3.2.2 Hubbard Model

The tight binding model as defined above fails to account for any interactions between neighbouring electrons. This motivates the extension of this model to the Hubbard model (2), which includes the (onsite) Coulomb repulsion between electrons. Despite its simple form, this model can describe very rich physical phenomena. In particular, it becomes very interesting to study when  $U$  and  $t$  are of comparable order, since it highlights the competing phenomena that takes place in correlated systems. The 2D Hubbard model remains unsolved to date, but is able to predict all sorts of correlated phases: it describes metals, insulators, superconductors and other exotic phases (REFERENCE). This model has been widely studied since it resembles the structure of the cuprate high-temperature superconductors (REFERENCE). While it has shown to accurately capture the magnetic and superconducting behaviour that is expected of this family of superconductors, recent findings suggest that the superconducting groundstate could indeed be an artefact of some numerical approximation (REFERENCE-ABSENCE...). A large part of this thesis focuses on exploring this toy model and the effect of the nearest-neighbour hopping parameter.

$$\hat{H} = \sum_{ij\sigma} -t_{ij}(\hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma} + h.c) + U \sum_i \hat{n}_{i\uparrow} \hat{n}_{i\downarrow} \quad (2)$$

### 3.2.3 Hubbard-Kanamori Model

In the case of materials with a multi-band and/or multi-orbital nature, the Hubbard model is not sufficient to capture all of the physical phenomena. This motivates the extension of the Hubbard Model to the Hubbard- Kanamori model by including a Hund's coupling term. Solving this Hamiltonian is rather challenging, which is why we resort to numerical techniques such as FRG to do so.

$$H_{int} = U \sum_{is} n_{i,s\uparrow} n_{i,s\downarrow} + \frac{V}{2} \sum_{i,s,t \neq s} n_{is} n_{it} - \frac{J}{2} \sum_{i,s,t \neq s} \vec{S}_{is} \cdot \vec{S}_{it} + \frac{J'}{2} \sum_{i,s,t \neq s} \sum_{\sigma} c_{is\sigma}^{\dagger} c_{is\bar{\sigma}}^{\dagger} c_{it\bar{\sigma}} c_{it\sigma} \quad (3)$$

Here,  $U$  and  $V$  represent the intraorbital interaction of electrons in the same and different orbitals respectively. For generality, the intraorbital exchange  $J$  and the 'pair hopping' term  $J'$  following from Hund's rule coupling have been separated. Note that this Hamiltonian is relevant for the later section of this project, where the model is extended to a two-orbital, two-dimensional Hubbard Model.

### 3.3 Theoretical Background in FRG

In this section the theoretical framework in which FRG calculations are performed is outlined. The central element of FRG is a flow equation that describes the evolution of the system's action with respect to a scalar parameter Lambda  $\Lambda$  (*See section 3.3.1*). The flow equation can be solved exactly for a limited number of systems (REFERENCE), so for most scenarios an approximation has to be made in order to reach a solution in a reasonable computational time. More details of how this approximation is performed and the limitations it presents can be found in Section 3.3.2. For the systems explored in this project only two-particle interactions are considered and any higher order terms are neglected. This allows for the (effective) action of the system to be separated into three channels, which correspond to the three physical channels: Superconductivity, Spin-Density and Charge-Density Waves. The calculation is then performed to determine the "winning channel" which will correspond to the respective physical phase that the material exhibits. This rather "hand-wavy" overview of FRG can be viewed as a flowchart in Fig.2.

#### 3.3.1 Flow equation

The central elements of statistical physics are the partition function, the canonical potential and its Legendre transformations. One can derive all physical observables from these. In quantum many body physics, this partition function is replaced by a partition functional, defined as follows:

$$\mathcal{Z}[\bar{\eta}, \eta] = \int \mathcal{D}\bar{\psi} \mathcal{D}\psi e^{\mathcal{S}[\bar{\psi}, \psi]_e(\bar{\eta}, \psi) + (\eta, \bar{\psi})} \quad (4)$$

In the case of fermionic systems, the action in the exponent of Eq.4 takes the form shown below.

$$\mathcal{S}[\psi, \bar{\psi}] = -(\bar{\psi}, G_0^{-1} \psi) + V[\psi, \bar{\psi}] \quad (5)$$

Here,  $V[\psi, \bar{\psi}]$  is an arbitrary many-body interaction and  $G_0$  represents the propagator of the noninteracting system. This equation contains the shorthand notation (...), which represents the sum  $\sum_x \bar{\psi}(x)(G_0^{-1}\psi)(x)$ ,  $(G_0^{-1}\psi)(x) = \sum_{x'} G_0^{-1}(x, x')\psi(x')$ . The Grassman field index  $x$  represents all the quantum numbers of the single-particle basis and imaginary time. Note that in the limiting case where  $V = 0$ , the path integral is exactly solvable. The main idea behind FRG is to introduce a cutoff in the non interacting Green's function ( $G_0$ ). The cutoff is then interpolated




Figure 2: **FRG Flowchart:** Schematic diagram outlining the TU<sup>2</sup>FRG calculation steps. Starting from the flow equation and applying the truncation scheme in order to decouple the action into three "physical" terms. The flow equation can then be solved by calculating each channel separately and taking the dominating phase to be the channel that diverges.

between the solveable initial state and the full path integral solution by subsequently including electronic correlations.

For convenience, the equations are then formulated in terms of the effective action (the Legendre transform of  $\mathcal{G}[\eta, \bar{\eta}]$ ):

$$\mathcal{T}[\psi, \bar{\psi}] = (\bar{\eta}, \psi) + (\bar{\psi}, \eta) \mathcal{G}[\eta, \bar{\eta}] \quad (6)$$

where  $\mathcal{G}[\eta, \bar{\eta}]$  is given by:

$$\mathcal{G}[\eta, \bar{\eta}] = -\ln \int \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-\mathcal{S}[\psi, \bar{\psi}]} e^{(\bar{\eta}, \psi) + (\bar{\psi}, \eta)} \quad (7)$$

The next step is to introduce a scalar flow parameter  $\lambda$  into the generating functionals defined above. This has to be performed such that the generators recover their original structure at  $\lambda = 0$ . After a series of algebraic manipulations, which are omitted here but can be found in (REFERENCE), one arrives at the exact functional flow equation for the effective action:

$$\frac{d}{d\lambda} \mathcal{T}^\Lambda[\psi, \bar{\psi}] = (\bar{\psi}, \dot{Q}_0^\Lambda \psi) - \frac{1}{2} \text{tr}(\dot{Q}_0^\Lambda (\Gamma^{(2)\Lambda}[\psi, \bar{\psi}])^{-1}). \quad (8)$$

Where  $\Gamma^{(2)\lambda}[\psi, \bar{\psi}]$  and  $\underline{Q}_0^\Lambda$  are given by equations (9) and (10) respectively.

$$\Gamma^{(2)\lambda}[\psi, \bar{\psi}] = \begin{bmatrix} \bar{\delta} \delta \Gamma[\psi, \bar{\psi}](x', x) & \bar{\delta} \bar{\delta} \Gamma[\psi, \bar{\psi}](x', x) \\ \delta \delta \Gamma[\psi, \bar{\psi}](x', x) & \delta \bar{\delta} \Gamma[\psi, \bar{\psi}](x', x) \end{bmatrix} \quad (9)$$

$$\underline{Q}_0^\Lambda = \begin{bmatrix} Q_0^\Lambda & 0 \\ 0 & -Q_0^{\Lambda t} \end{bmatrix} = \text{diag}(Q_0^\Lambda, -Q_0^{\Lambda t}), \quad (10)$$

### 3.3.2 Truncation scheme

### 3.3.3 Decoupling of flow equation

### 3.3.4 Instability calculation

## 4 Computational Methods

### 4.1 Diverge

### 4.2 Convergence of Calculation

#### 4.2.1 Form factor convergence

#### 4.2.2 Number of k points convergence

### 4.3 Calculation of Susceptibilities

#### 4.3.1 Nesting vectors

#### 4.3.2 Superconducting order parameters

## 5 Results and discussion

### 5.1 1NN Model

We define the 1NN model as the 2D Hubbard model with a single orbital per site, allowing hopping between nearest-neighbour sites (*a diagrammatic representation can be found in Fig.1.a*). As outlined in Section 3.2.2, solving this model near half-filling becomes a hard endeavour. This section explores the solution to the 1NN Model Hamiltonian using two-particle-interaction-truncated FRG. Here, a phase diagram in terms of the On-site Coulomb repulsion  $U$  and chemical potential  $\mu$  is presented and discussed in detail. The calculations were carried out using a  $20 \times 5 \ n_k \times n_{k_f}$  grid and a form factor of  $4\text{\AA}$ , with a nearest-neighbour hopping parameter of  $1\text{eV}$ . The choice of convergence parameters was determined by the methods discussed in Section 4.2. Results are presented for Coulomb repulsion values ranging from  $1\text{-}20\text{ eV}$  and chemical potential values spanning the entire energetic bandwidth of the model (from  $-4\text{eV}$  to  $4\text{eV}$ ). Whilst there exists previous work on the 2D Hubbard Model using FRG (REFERENCE), most of it focuses on small areas of the phase diagram and none of it explores the effect of varying the onsite Coulomb repulsion. Therefore, the work presented here covers a much wider range of parameters than what has previously been explored.

In real materials, the chemical potential is an easily tuneable parameter due to its strong connection to the electronic doping of the system. However, controlling the magnitude of the Coulomb repulsion between electrons is not that straightforward. Therefore, the analysis performed in this

project does not necessarily provide a route to enhance the superconductivity in real materials that resemble the models investigated. Instead, the main aim is to understand how varying these parameters affects the correlated phases observed in the model. It is also important to note, as discussed in Section 3.3.2, that any analysis on the transition temperature of the superconducting regions must be treated qualitatively and that this work does not claim to have found Superconducting regions at temperatures of the order of thousands of Kelvin in any of the models discussed.

The complete phase diagram for the values discussed above is shown in Fig.3. This particle-hole symmetric phase diagram shows a magnetic dome sandwiched between two superconducting regions. Moreover, narrow magnetic stripes appear for even integer chemical potential values.

Paragraph comparing results to the Cuprates!




Figure 3: **Phase diagram for the 1NN model:** ( $t = 1\text{eV}$ ,  $n_k \times n_{kf} = 20 \times 5$ ,  $ff = 4\text{\AA}$ ) as a function of On-site Coulomb Repulsion  $U$  and chemical potential  $\mu$ . Figure shows the four phases observed in the 1NN model: SC (Superconductivity), SDW (Spin-Density Wave), CDW (Charge Density Wave) and FL (Fermi-Liquid). Calculated points in the phase diagram are shown by the 'x' markers and a lighter-coloured background is used to depict interpolated regions between these points.



### 5.1.1 Superconductivity in the 1NN Model




Figure 4: **Superconductivity in the 1NN model:** Fig a) Superconducting order parameter projected on Fermi Surface at  $\mu = 1.00$  eV, for  $U = 10.00$  eV. The order parameter is antisymmetric about a 90 degree rotation and hence it exhibits d-wave symmetry. Fig b) Transition temperature ( $T_c$ ) as a function of chemical potential ( $\mu$ ) for  $U = 10.00$  eV.  $T_c$  is enhanced closest to the magnetic instability.

### 5.1.2 Magnetic stripes in the 1NN Model

## 5.2 Effect of next-nearest neighbour hopping (1NNN model)




Figure 5: **Magnetic stripes in the 1NN model:** Fig a) Transition temperature as a function of Coulomb repulsion ( $U$ ) along the Magnetic stripe at  $\mu = 1\text{eV}$ . Fig b). Magnetic susceptibility along high symmetry path for  $U = 2.00$  eV. Fig c) Magnetic susceptibility along high symmetry path for  $U = 18.00\text{eV}$ . Plots showing the susceptibility as a function of  $q$  for both magnetic regions. The Ferromagnetic SDW is suppressed by a superconducting phase at  $U \approx 6.00\text{eV}$ . At larger values of  $U$ , the SDW phase is recovered but with an Anti-Ferromagnetic ordering instead.




Figure 6: **1NNN model:** a) Phase diagram for the 1NNN model as a function of Coulomb repulsion  $U$  and chemical potential  $\mu$  ( $t = 1\text{eV}$ ,  $n_{k_x n_{k_y}} = 20 \times 5$ ,  $f_f = 4\text{\AA}$ ) for  $t' = 0.00, 0.25, 0.50, 0.75$  eV. b) Transition temperature for the superconducting region. c) Single plot of the band structure along high-symmetry path for corresponding values of  $t'$ . d) Single plot of the Fermi surface for all values of  $t'$  considered.

### 5.2.1 Superconductivity in the 1NNN Model

### 5.2.2 Magnetic stripes in the 1NNN model

$t'$ (eV)	$\mu$ (eV)	Competing with other phase?	Nesting vector	Magnetic ordering
0.00	1.00	Yes (SC)	(0,0)-( $\pi$ , $\pi$ )	Changing from FM-AFM
0.00	2.00	No	(0,0)-(0, $\pi$ )	Changing from FM to Commensurate
0.25	2.40	No	(0,0)	FM
0.25	1.40	Yes (CDW)	(0,0)	FM
0.25	-2.00	No	(0,0)-( $\pi$ , $\pi$ )	FM - AFM
0.25	0.60	No	(0,0)	FM

Table 1: **Survey of stripes in the 1NNN model.** Table summarising the location and magnetic ordering of the Stripes in the 1NNN model .

### 5.2.3 Continuous variation of next-nearest neighbour hopping

## 5.3 Effect of bi-orbital system (1NN2 model)

Model	$t_{3z^2-r^2}^{[1,0,0]}$	$t_{3z^2-r^2}^{[0,1,0]}$	$t_{x^2-y^2}$	$t_{x^2-y^2-3z^2-r^2}^{[1,0,0]}$	$t_{x^2-y^2-3z^2-r^2}^{[0,1,0]}$	$t_{\perp}$
	-0.781	-0.719	-0.375	-0.402	-0.310	-2.50
1NN2MN	on	on	on	off	off	off
1NN2MY	on	on	on	on	on	off

Table 2: **Nearest neighbour hopping parameters for 2D two-orbital Hubbard models.** First row shows the calculated hopping parameters, note that  $t_{\perp}$  corresponds to the intralayer hopping between the  $d_{3z^2-r^2}$  orbitals. Other rows show which hopping parameters were included in each of the two models.

### 5.3.1 Superconductivity in the 1NN2 model

### 5.3.2 Effect of orbital hybridisation in the 1NN2 model

## 6 Conclusion and Outlook

## References