THE SYMMETRIC PHASE OF THE THREE-DIMENSIONAL ABELIAN HIGGS MODEL: TORUS LIMITS AND THE OSTERWALDER–SCHRADER AXIOMS WITH CUBIC SYMMETRY

Abstract. We study the Euclidean abelian Higgs model (scalar quantum electrodynamics) in three dimensions, with a non-compact $U(1)$ gauge field and a complex scalar field of positive mass squared m^2 and quartic coupling $\lambda > 0$. The model is regularised on the lattice $\varepsilon\mathbb{Z}^3$ on periodic cubic tori, with a photon mass $\mu_0 > 0$ in Feynman form and King's mass counterterm computed on the torus at a fixed reference photon mass. For $0 < \lambda < 0.1m$ and $0 < e^2 < 0.1m$ we prove that the generating functionals of the curvature $F(g)$ and of the renormalised Wick square $:|\varphi|^2:(h)$ are bounded on a complex tube, uniformly in the lattice spacing, in the size of the torus and in μ_0, and also in a massless state in which the constant mode of the gauge field is averaged over its period. Along subsequences the spacing goes to zero and the torus to $\mathbb{R}^3$, and then μ_0 goes to zero along a countable set. The limit states satisfy the Osterwalder–Schrader axioms OS0, OS1 for the translations and the cubic group, OS2 about every coordinate plane, and OS3. Their laws are integrals of translation-ergodic, reflection-positive components, each of which clusters in the mean and under separations in which all three coordinates diverge, and has a unique joint vacuum in each coordinate direction; observables linear in the curvature, with any square-integrable dependence on the Wick square, cluster along each coordinate axis. We also prove sign inequalities for truncated correlations, and that the curvature two-point function of every limit state is dominated by that of the free massless photon and has no massless component at positive photon mass. The ultraviolet input is Bałaban's renormalisation group, with three repairs of printed proofs; the proof of the ultraviolet bounds also uses a number of technical estimates, collected in Appendix K, whose proofs are not included in this version. Invariance under the full rotation group, the cluster property along a single axis for all observables, uniqueness of the limits, non-triviality and the masslessness of the photon are left open.

Contents

1. Introduction 2
2. The regularised model 7
3. Main results 13
4. Ultraviolet input from the literature 19
5. The vacuum bound for the slice models, and three repairs of printed proofs 24
6. From the vacuum bound to the uniform tube bound 37
7. Lattice statements on the tori 42
8. Limits along subsequences 50
9. OS0, OS1 for translations and the cubic group, OS2, OS3 56
10. OS4: ergodic decomposition and clustering 62
11. The reconstructed theory 89
12. Correlation signs and non-triviality under a remainder bound 90
13. The curvature two-point function 96
14. Open problems 102
Appendix A. Regulator data and the order of choices 107
Appendix B. The lattice model: supplementary proofs 126

2020 *Mathematics Subject Classification.* Primary 81T08; Secondary 81T13, 81T17, 81T25.

Key words and phrases. abelian Higgs model, scalar quantum electrodynamics, constructive field theory, renormalisation group, Osterwalder–Schrader axioms, reflection positivity, cluster property.

Appendix C. Counterterm and vacuum identifications 150
Appendix D. Bałaban's Proposition 1 on the torus 154
Appendix E. The ingredients of Theorem 5.2 171
Appendix F. Assembly of Theorem 5.2 270
Appendix G. The counterterm gaps 296
Appendix H. The c-number bounds 312
Appendix I. The negative-source bound and the choice of parameters 334
Appendix J. Measure-theoretic facts 349
Appendix K. Statements used without proof in this version 350
References 370

1. Introduction

1.1. **The model and the question.** The abelian Higgs model is the Euclidean field theory of a real vector field A_μ and a complex scalar field φ on $\mathbb{R}^3$ with formal action

$$S(A,\varphi) = \int_{\mathbb{R}^3} \Big[\tfrac{1}{4} F_{\mu\nu} F_{\mu\nu} + \tfrac{1}{2} |(\partial_\mu + ieA_\mu)\varphi|^2 + \tfrac{1}{2} m^2 |\varphi|^2 + \lambda |\varphi|^4 \Big]\, dx, \qquad F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu. \tag{1.1}$$

This is King's normalisation [24, (2.1)–(2.4), p. 652], the one of the lattice action (2.2) below, in which the coupling range of this paper is stated; in terms of $\psi := \varphi/\sqrt{2}$ the action reads $|(\partial_\mu + ieA_\mu)\psi|^2 + m^2|\psi|^2 + 4\lambda|\psi|^4$, so in that normalisation the quartic coupling is 4λ. The sign of ieA_μ is the one of the lattice covariant difference $\varepsilon^{-1}(e^{ie\varepsilon A_{x,\mu}}\varphi(x+\varepsilon e_\mu) - \varphi(x))$, and the gauge transformations are $\varphi \mapsto e^{i\chi}\varphi$, $A \mapsto A - e^{-1}d\chi$.

In three dimensions e^2 and λ have the dimension of a mass and the model is superrenormalisable. For $m^2 < 0$ the scalar condenses and the photon acquires a mass by the Higgs mechanism. For $m^2 > 0$, the case treated here, the scalar is massive and the photon is expected to stay massless: this is the symmetric, or Coulomb, phase. This picture comes from perturbation theory, and at very small λ it is expected to fail: one-loop perturbation theory predicts a first-order transition at positive m^2, on a surface $\lambda m^2 \approx e^6/(72\pi^2)$ that lies inside the coupling range of our theorems, with a massive photon below it (Section 14). None of our proofs uses the phase; where this paper speaks of the symmetric or Coulomb phase, it refers to couplings above that surface. The question we address is whether the Schwinger functions of the local gauge-invariant fields, the curvature $F_{\mu\nu}$ and the Wick square $:|\varphi|^2:$, can be constructed as limits of a lattice regularisation and satisfy the Osterwalder–Schrader axioms [27, 28]: regularity (OS0), Euclidean invariance (OS1), reflection positivity (OS2), symmetry (OS3) and clustering (OS4).

The gauge field is non-compact: on the lattice, A is real-valued on the bonds and the gauge action is the quadratic form of the lattice curl, as in Bałaban [1, 2, 3, 4] and King [24, 25]. The exact lattice relation between the two formulations is that, for observables of integer charge on a finite lattice with free boundary conditions (trivial cohomology), the non-compact model is the compact Villain model with every monopole removed [5, Sect. 3, Thm. 3.1, p. 286]. This choice matters in the Coulomb phase: in the compact pure gauge theory in three dimensions monopoles produce a mass gap and confinement for all couplings [20], while the non-compact pure gauge theory is the free field, whose photon is massless. For the non-compact model with matter studied here, masslessness of the photon is expected for couplings above the first-order surface, but it is not proved (Section 14).

1.2. **Background.** Reflection positivity of lattice gauge theories goes back to Osterwalder and Seiler [29] for compact gauge groups, and to [8, Thm. 5.3, p. 272] for the non-compact abelian field; Seiler's lectures [34] contain the general framework. In two dimensions the model, with a non-compact continuum gauge field and lattice matter, was constructed by Brydges, Fröhlich and Seiler

[8, 9, 10], who also introduced the correlation inequalities that make the generating functionals monotone in the photon covariance. The ultraviolet problem in three dimensions was solved by Bałaban [1, 2, 3, 4], who proved two-sided bounds

$$e^{-E_-|T|} \;\le\; Z^{\varepsilon}(T) \;\le\; e^{E_+|T|} \tag{1.2}$$

on the partition function of the model with a photon mass on a periodic torus T, with constants independent of the lattice spacing ε and of T [1, Thm. (1.14), p. 606], [2, Thm. (1.3), p. 556]. Bałaban's smallness conditions are conditions on the running couplings $et^{1/2}$, λt and m^2t^2 at short scales $t \le \varepsilon_0$, not on the couplings themselves.

King [24, 25] turned these stability bounds into limits. He proved convergence of the generating functional of $F(g)$ and $:|\varphi|^2:(h)$ as $\varepsilon \to 0$ on tori and on cubes, constructed the infinite-volume limit through cubes with Neumann conditions for A and Dirichlet conditions for φ using the monotonicity in the volume given by the correlation inequalities, and proved translation invariance and reflection positivity of the resulting Schwinger functions. King kept the photon mass $\mu_0 > 0$ throughout, and listed what remained: ergodicity, rotation invariance and regularity [25, p. 326]. For rotations he compared, on a fixed torus, the lattice with its copy rotated by the angle $\theta_0 = \arctan(3/4)$ [25, Thm. 2.4, pp. 326–327].

The Higgs phase was treated on the unit lattice by Bałaban, Brydges, Imbrie and Jaffe [5], and in the continuum by Bałaban, Imbrie and Jaffe [7], who constructed the effective action down to the unit scale. The ultraviolet renormalisation of scalar quantum electrodynamics in three dimensions was revisited by Dimock [14], who later treated the fermionic model [15]. Continuum limits of the compact pure $U(1)$ theory were obtained by Gross [21] in three dimensions and Driver [16] in four. Gauge theories with matter in three dimensions have recently been approached through stochastic quantisation [11, 12].

1.3. **Main results.** We do not use King's cube states. We work throughout on periodic cubic tori T_j of side $j!\,\ell_0$, on which the lattice model is exactly invariant under the translations of the lattice and under the cubic group, and is reflection positive about lattice planes. The photon mass is the only infrared regulator, and the scalar mass counterterm is King's counterterm computed on the torus itself at a fixed reference photon mass $\mu_* = m$; we call this choice the *torus counterterm scheme* (Definition 2.2). Besides the states at photon mass $\mu_0 \in (0, \mu_*]$ we use the state $\flat$, in which the photon has no mass and the constant mode of the gauge field is averaged over its period; at fixed lattice it is the limit $\mu_0 \downarrow 0$ (Lemma 2.8). The objects in the following statements are defined in Section 2.

Theorem A (existence and OS0–OS3). *Let $m > 0$, $0 < \lambda < 0.1m$ and $0 < e^2 < 0.1m$, and consider the lattice states $\langle\cdot\rangle_{\varepsilon,j,\mu_0}$ of Definition 2.7 in the torus counterterm scheme with reference photon mass $\mu_* = m$, on the tori T_j of side $\ell_j = j!\,\ell_0$, at the spacings $\varepsilon = 2^{-K}\varepsilon_0$, at the photon masses $\mu_0 \in (0, m]$ and in the state $\flat$. There are $\varepsilon_0 = 2^{-n_0}$, $\ell_0 \ge 183/m$, $K_0 \in \mathbb{N}$, $\sigma_+ > 0$ and $C < \infty$ such that the following holds, with $H := m^2$.*

(i) *(Uniform tube bound.) For all $K \ge K_0$, $j \ge 1$ and $\mu_0 \in (0, m] \cup \{\flat\}$, every finite family of real test functions supported in a ball of radius R with $\ell_j > 2R + 2$, and every ζ in the tube $D_{\sigma_+,H}$, the generating functional of the curvature and the renormalised Wick square satisfies*

$$|G_{\varepsilon,j,\mu_0}(\zeta)| \;\le\; G_{\varepsilon,j,\mu_0}(\operatorname{Re}\zeta) \;\le\; \exp\Big(\tfrac{1}{2}\Big\|\sum_i (\operatorname{Re} z_i) g_i\Big\|_{2,\varepsilon}^2 + C\|k(\zeta)\|_{L^1,\varepsilon}\Big).$$

(ii) *(Limits.) Let $Q_0 \subset (0, m]$ be countable with 0 as an accumulation point. Every sequence (K_l, j_l) with $K_l, j_l \to \infty$ has a subsequence along which the lattice Schwinger functions at every $\mu_0 \in Q_0$ and in the state $\flat$ converge, to limit states S^{μ_0} and $S^{\flat}$; along this subsequence*

$S^0 := \lim_{\mu_0 \downarrow 0,\, \mu_0 \in Q_0} S^{\mu_0}$ *exists. Each of these limit states is, on test functions in* C_c^∞*, the family of moments of a Borel probability measure on* $H^{-s'}_{\mathrm{loc}}(\mathbb{R}^3; \Lambda^2\mathbb{R}^3 \oplus \mathbb{R})$ *for every* $s' > 3$.

(iii) *(Axioms.) Each limit state satisfies OS0 with the constants* σ_+*,* H*,* C *of (i), OS1 for the group* $\mathbb{R}^3 \rtimes O_h$ *generated by the translations and the cubic group, OS2 about every coordinate plane, and OS3 (Definition 2.10). Its Schwinger functions satisfy* $|S_{n,p}(g;h)| \le e^{C\sigma_0} \mathrm{e}^n (n!)^{1/2} (\mathrm{e}/\sigma_0)^p\, p! \prod_i \|g_i\|_2 \prod_k N(h_k)$*,* $\sigma_0 := \min(\sigma_+, H)/2$.

(iv) *(Monotonicity.) For every generating functional* Θ *of the class* $\mathcal{D}$ *of Section 2.4,* $\mu_0 \mapsto S^{\mu_0}(\Theta)$ *is non-increasing on* Q_0*, and* $S^{\mu_0}(\Theta) \le S^\flat(\Theta)$*,* $S^0(\Theta) \le S^\flat(\Theta)$.

Theorem A is Theorem 3.1, which also gives the convergence of the generating functionals on the tube. Euclidean invariance is established for the translations of $\mathbb{R}^3$ and the cubic group O_h, the symmetry group of the lattice; invariance under the full rotation group is not claimed (Remark 3.2). Theorem 3.5 gives parts (ii)–(iv) also for the lattices $3^{-K}\varepsilon_0\mathbb{Z}^3$ with blocking factor 3, under the hypothesis that the uniform tube bound holds for that family; the interest of blocking factor 3 lies in the rotations (Section 14).

Theorem B (weak clustering). *Let* S *be a limit state of Theorem A. Its law is the only Borel probability measure* μ *with these moments on a Hermite–Sobolev space of distributions, and* $\mu = \int \mu_\xi \, d\nu(\xi)$ *with components* μ_ξ *such that, for* ν*-almost every* ξ*:* μ_ξ *is invariant and ergodic under the translations of* $\mathbb{R}^3$*, invariant under the three coordinate reflections, and reflection positive about every coordinate plane; for all* $G, G' \in L^2(\mu_\xi)$*,* $\mathbb{E}_{\mu_\xi}[G'\,U(a)G] \to \mathbb{E}_{\mu_\xi}[G']\,\mathbb{E}_{\mu_\xi}[G]$ *in the mean over cubes and as* $\min_j |a_j| \to \infty$*, where* $U(a)$ *is the translation by* a*; and for each coordinate direction* e_j *the Osterwalder–Schrader reconstruction of* μ_ξ *with* e_j *as Euclidean time has a positive Hamiltonian* H_j *and a unique joint vacuum: the vectors fixed by the semigroup* e^{-tH_j}*,* $t \ge 0$*, and by the translations orthogonal to* e_j *form one line* $\mathbb{C}\Omega_j$.

Theorem B is Theorem 3.6. In the terms of Definition 2.12 it says that the components satisfy OS4 in the weak form, ergodicity under the translation group $\mathbb{R}^3$; equivalently, they cluster under every separation in which all three coordinates diverge. It is not the cluster property OS4 of Definition 2.11, which asks for clustering as the separation runs along a single coordinate axis: single-axis clustering does not follow from the structure that Theorem B uses, namely invariance under translations and reflections, reflection positivity, and ergodicity under $\mathbb{R}^3$ (Proposition 3.8); it would follow from invariance under all rotations (Corollary 10.44). For part of the observables we prove it.

Theorem C (single-axis clustering, partial). *Let* S *be a limit state of Theorem A,* μ *its law, and* $\mathcal{M}$ *the* σ*-algebra generated by the Wick-square fields* $:|\varphi|^2:(h)$.

(a) *For every real 2-form* g *with components in* $C_c^\infty(\mathbb{R}^3)$ *and every* $Y \in L^2(\mathcal{M}, \mu)$*, the observable* $F(g)Y$ *clusters against every* $G' \in L^2(\mu)$*:* $\mathbb{E}_\mu[G'\,U(a)(F(g)Y)] \to 0 = \mathbb{E}_\mu[G']\,\mathbb{E}_\mu[F(g)Y]$ *as* $|a| \to \infty$ *in any manner, in particular along each coordinate axis.*

(b) *If the latent density* Δ_S *of the Wick square vanishes (Definition 10.34; this holds if some limit, along the defining sequence of* S*, of the torus pressures of the Wick square is differentiable at zero source), the same holds, with limit* $\mathbb{E}_\mu[G']\,\mathbb{E}_\mu[G]$*, for every* G *in the closed span of* $L^2(\mathcal{M}, \mu)$ *and of the products in (a).*

(c) *If the block pressures of the bounded local observables* $\cos\Phi(f)$*,* $\sin\Phi(f)$*, with* $\Phi(f)$ *a smeared field, are differentiable at zero (hypothesis (D) of Corollary 10.38), then* μ *is ergodic,* S *satisfies OS4, and every pair* $G, G' \in L^2(\mu)$ *clusters along each coordinate axis.*

Theorem D (signs and the curvature two-point function). *Let* S *be a limit state of Theorem A.*

(a) *For real 2-forms* g, g' *and non-negative* h, h' *with components in* $C_c(\mathbb{R}^3)$*:* $\langle F(g)^2; :|\varphi|^2:(h)\rangle_S \le 0$*,* $\langle :|\varphi|^2:(h); :|\varphi|^2:(h')\rangle_S \ge 0$*, and* $S^T_{4,0}(g,g,g',g') + 2S_{2,0}(g,g')^2 \ge 0$*, where* $\langle\,\cdot\,;\,\cdot\,\rangle_S$ *and* S^T

denote truncated functions. If the first inequality is strict for one pair (g, h), then S is not Gaussian, and the curvature and the Wick square are not independent under its law.

(b) *The two-point function of the smeared magnetic field is a Fourier multiplier $m_S(p)$ with $0 \le m_S(p) \le P_T(p)$, where $P_T(p)$ is the orthogonal projection of $\mathbb{C}^3$ onto $p^\perp$; the multiplier P_T is that of the free massless photon. At positive photon mass, $m_S(p) \le \min(1, |p|^2/\mu_0^2) P_T(p)$, so S^{μ_0} has no massless component.*

(c) *If $S^\flat$ or S^0 is massless in the curvature sector (Definition 13.6), its curvature correlations are not integrable, and in every coordinate direction the curvature sector of its reconstruction has no mass gap.*

Theorems C and D are Theorems 3.10 and 3.11. Theorem D does not show that the limit states are non-Gaussian, nor that the photon is massless at $\mu_0 = 0$. In Theorem 12.9 we show that the first inequality of (a) is strict at small $e^2 + \lambda$, under two hypotheses that we formulate and do not prove: a remainder bound (R) for the effective action with sources, and a one-loop statement (C) (Section 12).

The ultraviolet input is taken from the printed papers of Bałaban [1, 2, 3, 4] and King [24, 25], with three repairs of printed steps. First, the printed proof of [2, Lemma 2.4, p. 572] does not cover the lattice points near the edges of the region on which the lemma is applied; we prove the statement of the lemma, under explicit conditions on the regulator data, by an argument that covers these points (Lemma 5.6). Second, the proof of [3, Prop. 1, pp. 420–445] contains sketched steps; we write out each of them, with the depth inside the localisation domain that it needs, because our application uses the proposition at the limit of what its printed nesting provides (Proposition 5.8). Third, the printed proof of the combinatorial estimate [2, (3.42), p. 592], the last step of the proof of Bałaban's upper vacuum bound [2, Thm. (1.3)], is incomplete: the step to [2, (3.47), p. 593], introduced by "after easy transformations", is not written out; the cube count [2, (3.45), p. 592] fails for blocks near the boundary of the region; and the closing estimates [2, (3.51)–(3.54), p. 594] bound only the large-field sets $\Lambda_0^{(k)c}$, while (3.42) also contains collar and shell sets. We bound these sets by families of cubes, pay the entropy of the sum over regions, and close (3.54) at $p = 6$ with an explicit upper bound on ε_0 (Proposition 5.21). This completes the printed proof of the upper vacuum bound at this step; the other estimates that its printed proof omits are discussed in Remark 4.4.

1.4. **Method.** The infrared problem of a constructive field theory is usually controlled by decay: a mass gap makes a cluster expansion converge, the expansion gives a unique infinite-volume limit, and uniqueness gives the symmetries. In the Coulomb phase no such expansion is available. We replace it by three facts that hold uniformly in every regulator, on a single torus at a time, and are therefore inherited by every limit.

(a) A uniform bound on a tube. The central estimate (Theorem A(i)) bounds $|\langle e^{\sum z_i F(g_i) + \sum w_k X_{\mathrm{R}}(h_k)} \rangle|$ on a tube $\{|\mathrm{Re}\, w| \text{ small}\}$, with all of $\mathbb{C}$ allowed for the coefficients z_i of the curvature, uniformly in the spacing, the torus and μ_0. It has three factors. The curvature factor $e^{\frac{1}{2}\|g\|_2^2}$ is Gaussian domination of the photon by the free field, which on a torus holds with the sharp constant because the componentwise lattice Laplacian is the Hodge Laplacian (Lemma 7.3). The factor for positive $:|\varphi|^2:$-sources comes from a convexity bound for constant sources on the same torus (Proposition 6.4), spread over arbitrary sources by the chessboard estimate of reflection positivity (Proposition 6.5); its input is a positive gap between King's counterterms at two scalar masses on that torus (Theorem 6.1), which we verify by an explicit computation of the lattice graphs for the couplings of the window (1.3) below. The factor for negative sources (Proposition 6.7) comes from a vacuum bound for the model with a deformed photon covariance (Theorem 5.2), which we derive from Bałaban's renormalisation group.

(b) Compactness and exact symmetries. The uniform bound gives compactness of the lattice Schwinger functions and tightness of the lattice laws. Limits exist along subsequences; we do not prove uniqueness, and do not need it: on cubic tori the translations of $\varepsilon\mathbb{Z}^3$ and the cubic group are exact symmetries of every lattice state, and reflection positivity holds about pairs of lattice planes, so these properties are closed under limits. Continuity, which the bound supplies, extends the translations to $\mathbb{R}^3$.

(c) Monotonicity in the photon mass. At fixed lattice and torus, the Brydges–Fröhlich–Seiler inequalities make the generating functionals of the class $e^{F(g)-X_{\mathrm{R}}(q)}$, $q \geq 0$, non-decreasing in the photon covariance [8, Cor. 6.3(1), p. 281] (those of $e^{+X_{\mathrm{R}}(q)}$, $q \geq 0$, are non-increasing). Taking the lattice limit along one subsequence common to a countable set of photon masses, the monotonicity passes to the limit and gives the massless limit S^0 and the inequality $S^0 \leq S^\flat$.

The weak cluster property (Theorem B) uses only translations, the three coordinate reflections and reflection positivity. A function invariant under translations along a coordinate axis is fixed by the reflection in a plane orthogonal to that axis; hence the invariant σ-algebra is fixed by the reflections, the ergodic components inherit reflection positivity, and on each component the Osterwalder–Schrader reconstruction in each coordinate direction identifies the zero-energy vectors with the functions invariant under the translations along that direction.

Restriction on the couplings. The window

(1.3)
$$\mathfrak{W} := \{(e, \lambda, m, \eta, m_1, m_2) :\ m > 0,\ 0 < \lambda < 0.1m,\ 0 < e^2 < 0.1m,\ 0 < \eta < 1,\ 2m < m_1 < m_2\}$$

enters through one input of our own: the positive gap of (a) between King's counterterms at the scalar masses $0.7m$ and m, uniform in the spacing and the torus (Theorem 6.1). The other parameters η, m_1, m_2 of $\mathfrak{W}$ are auxiliary (anchor masses of the vacuum bound for the deformed model), and can be chosen for every (e, λ, m) with $m > 0$, $0 < \lambda < 0.1m$, $0 < e^2 < 0.1m$; Lemma A.5 shows that all conditions on the regulator data can be met at every point of $\mathfrak{W}$.

1.5. **What is not proved here.** We do not prove invariance under rotations outside O_h; OS4 in the sense of Definition 2.11, for the limit states or for their ergodic components, beyond the parts given by Theorem C; uniqueness of the limits, the equality $S^0 = S^\flat$, or the identification with King's cube limits; non-triviality of the limit states; masslessness of the photon at $\mu_0 = 0$; extremality of the limit states, or the existence of Wightman fields. The proofs do not use the phase, and one-loop perturbation theory predicts a first-order transition inside the coupling range of Theorem A. Section 14 states these problems, says where the methods of this paper stop for each of them, and explains what is known about rotations, including the role of blocking factor 3.

1.6. **Organisation and conventions.** Section 2 fixes the regularised model and the objects of the main results, which are stated in Section 3. Sections 4–6 contain the proof of the uniform tube bound: Section 4 collects the results of Bałaban and King that we use; Section 5 states the vacuum bound for the model with a deformed photon covariance (Theorem 5.2), describes the architecture of its proof, and contains the three repairs; Section 6 derives the uniform tube bound (Theorem 6.8) from it. Sections 7–9 prove parts (ii)–(iv) of Theorem A: lattice statements on one torus at a time (Section 7), limits along subsequences (Section 8), and the axioms OS0–OS3 together with what is known about rotations of infinite order (Section 9). Section 10 proves Theorems B and C, Section 11 describes the reconstructed theory, Section 12 proves the correlation signs and non-triviality under hypotheses (R) and (C), and Section 13 studies the curvature two-point function. Section 14 is a list of open problems.

The appendices contain the longer proofs; each begins by saying which result of the body it proves. Appendix A fixes the regulator data, the order in which they are chosen, and the conditions they satisfy, and proves that these conditions can be met at every point of $\mathfrak{W}$ (Lemma A.5). Appendix B contains supplementary proofs for the lattice model, Appendix C the identification of

King's counterterms and vacuum constants with Bałaban's, Appendix D the form of [3, Prop. 1] on the torus, Appendices E and F the ingredients and the assembly of the proof of Theorem 5.2, Appendix G the counterterm gaps, Appendix H the bounds on c-number differences, Appendix I the bound for negative sources and the choice of the auxiliary parameters, and Appendix J the measure-theoretic facts used in Section 10.

The logical structure is as follows. Theorem A(i) is Theorem 6.8, whose proof occupies Sections 4–6 and Appendices A and C–I. Sections 7–9 use from Sections 4–6 only Theorem 6.8, and Sections 10–13 use Theorem A together with estimates on the lattice tori from Sections 6 and 7. A reader interested in the passage from the uniform bound to the axioms may take Theorem 6.8 as given and read Sections 2, 3 and 7–11.

A formalisation of parts of the argument in the Lean proof assistant is in preparation.

Conventions. Throughout, $d = 3$ and $\mathbb{N} = \{1, 2, 3, \dots\}$. The letter e always denotes the charge; exponentials are written e^x or $\exp(x)$, and Euler's number, where it occurs as a constant factor, is written e. Masses and lengths are measured in the units of [24], in which the coarsest renormalisation-group scale is 1. Equations are numbered within sections, and theorems, propositions, lemmas, corollaries, definitions, examples and remarks share one counter in each section.

2. The regularised model

Throughout, $m > 0$, $\lambda > 0$ and $e \neq 0$ are fixed, and $\mu_* := m$ is the reference photon mass.

2.1. **Lattices and tori.** Fix the blocking factor $L \in \{2, 3\}$ ($L = 2$ in Theorem 3.1, $L = 3$ in Theorem 3.5), the block-size parameter $M_b \in \mathbb{N}$ of Bałaban's renormalisation group (King's M [24]), and $n_0 \in \mathbb{N}$. Put

$$\varepsilon_0 := L^{-n_0}, \qquad \varepsilon = \varepsilon_K := L^{-K}\varepsilon_0 \quad (K = 0, 1, 2, \dots).$$

Fix $\ell_0 \in c_L M_b \mathbb{N}$, with $c_2 = 2$ and $c_3 = 18$. For $j \geq 1$ let $\ell_j := j!\,\ell_0$, let $T_j := (\mathbb{R}/\ell_j\mathbb{Z})^3$ be the periodic cube of side ℓ_j, and $T_{j,\varepsilon} := T_j \cap \varepsilon\mathbb{Z}^3$ the lattice torus. A site is a point $x \in T_{j,\varepsilon}$; a (positively oriented) bond is a pair $b = (x, x + \varepsilon e_\mu)$, written (x, μ); a plaquette is $p = (x; \mu, \nu)$ with $\mu < \nu$, the square with corners x, $x + \varepsilon e_\mu$, $x + \varepsilon e_\nu$, $x + \varepsilon e_\mu + \varepsilon e_\nu$. We write $|T_j| = \ell_j^3$.

Lemma 2.1 (torus sizes)**.** *For all $j \geq 1$ and $K \geq 0$: (i) the number of sites per side $N = \ell_j/\varepsilon_K$ is an even integer; (ii) for every $k \in \{0, 1, \dots, K + n_0\}$, $\ell_j/(M_b L^k \varepsilon_K) \in 2\mathbb{N}$; (iii) $\ell_j \mid \ell_{j+1}$ and $\ell_j \to \infty$; (iv) $\varepsilon_{K'}\mathbb{Z}^3 \supset \varepsilon_K\mathbb{Z}^3$ for $K' \geq K$.*

Proof. $\ell_j/(M_b L^k \varepsilon_K) = j!\,(\ell_0/M_b)\,L^{K+n_0-k}$, and $\ell_0/M_b \in c_L\mathbb{N} \subset 2\mathbb{N}$; this is (ii), and (i) is the case $k = 0$ multiplied by M_b. (iii) and (iv) are immediate. □

2.2. **Fields, action and counterterm.** The gauge field is a function A on bonds, $A_{x,\mu} \in \mathbb{R}$ (non-compact); the Higgs field is a function φ on sites, $\varphi(x) \in \mathbb{C}$, with two real components and $|\varphi|^2 = (\operatorname{Re}\varphi)^2 + (\operatorname{Im}\varphi)^2$. Functions on sites, bonds and plaquettes carry the inner products $\langle u, v\rangle_\varepsilon := \sum \varepsilon^3 u\, v$, and $\|u\|_{2,\varepsilon}^2 := \langle u, u\rangle_\varepsilon$. The lattice exterior derivative is $(d\chi)_{x,\mu} := \varepsilon^{-1}(\chi(x + \varepsilon e_\mu) - \chi(x))$ on functions and

$$(dA)(x; \mu, \nu) := \varepsilon^{-1}\big(A_{x,\mu} + A_{x+\varepsilon e_\mu,\nu} - A_{x+\varepsilon e_\nu,\mu} - A_{x,\nu}\big)$$

on bond functions; d^* denotes the adjoint. We write $-\Delta_\varepsilon := d^*d + dd^*$ on bond functions. On the periodic lattice this operator acts on each component $A_{\cdot,\mu}$, viewed as a function of x, as the lattice Laplacian $-\Delta_\varepsilon$ on functions on sites (Lemma B.4); in particular $-\Delta_\varepsilon \geq 0$, with kernel the constant bond functions.

For a photon mass $\mu_0 > 0$ and a real bare coefficient a the action is

$$S_{\mu_0,a}(A, \varphi) := \tfrac{1}{2}\langle A,\, (-\Delta_\varepsilon + \mu_0^2)A\rangle_\varepsilon + S_a^{\mathrm{m}}(A, \varphi), \tag{2.1}$$

$$S_a^{\mathrm{m}}(A,\varphi) := \tfrac{1}{2}\sum_{x,\mu} \varepsilon^3\, \varepsilon^{-2} |e^{ie\varepsilon A_{x,\mu}}\varphi(x+\varepsilon e_\mu) - \varphi(x)|^2 + \sum_x \varepsilon^3 \big(\tfrac{1}{2}a|\varphi(x)|^2 + \lambda|\varphi(x)|^4\big). \tag{2.2}$$

The gauge part is the Feynman form of [24, (2.4), p. 652]; it equals $\frac{1}{2}\|dA\|_{2,\varepsilon}^2 + \frac{1}{2}\|d^*A\|_{2,\varepsilon}^2 + \frac{1}{2}\mu_0^2\|A\|_{2,\varepsilon}^2$. The matter part is invariant under the lattice gauge transformations $\varphi \mapsto e^{i\chi}\varphi$, $A \mapsto A - e^{-1}d\chi$, χ real on sites (Lemma B.5).

Definition 2.2 (King's counterterm on a torus; the torus counterterm scheme)**.** For $M > 0$ let $\delta m^2_{K,T}(\varepsilon; M)$ be the mass counterterm of [24, (3.32)–(3.33), p. 659] computed on the lattice torus $T_{j,\varepsilon}$, with scalar mass M, scalar propagator $(-\Delta_\varepsilon + M^2)^{-1}$ and photon propagator $(-\Delta_\varepsilon + \mu_*^2)^{-1}$, both on $T_{j,\varepsilon}$ with periodic boundary conditions and without block averaging, so that every line commutes with the translations by $\varepsilon\mathbb{Z}^3$. It is the polynomial in (e,λ) over the orders $2 \le \alpha + 2\beta \le 4$ of [24, (3.32)] whose coefficients are fixed, order by order, by the condition that the amputated one-particle-irreducible two-point function of φ at zero momentum vanish at the orders $e^\alpha\lambda^\beta$, $2 \le \alpha + 2\beta \le 4$. Put $a_{K,T}(M) := M^2 + \delta m^2_{K,T}(\varepsilon; M)$. The bare coefficient of the model on T_j at spacing ε is

$$b_{T_j}(\varepsilon) := a_{K,T_j}(m),$$

the same for every photon mass $\mu_0 \in (0, \mu_*]$ and for the state $\flat$ below. We call this choice the *torus counterterm scheme*.

The odd orders in e of $\delta m^2_{K,T}$ vanish, because the substitution $A \mapsto -A$, $e \mapsto -e$ fixes φ and leaves the matter action (2.2) and the centred Gaussian photon measure with covariance $(-\Delta_\varepsilon + \mu_*^2)^{-1}$ invariant (Lemma B.2); so the monomials are $e^2, \lambda, e^4, e^2\lambda, \lambda^2$.

Remark 2.3. (i) On $T_{j,\varepsilon}$ the counterterm of Definition 2.2 coincides with Bałaban's [3, p. 417]: both are the zero-momentum value of the amputated one-particle-irreducible two-point function on the same lattice torus, and the derivative parts have zero sum over a torus (Lemma C.1). (ii) The coefficient is computed at the reference photon mass μ_*, not at μ_0. This is what makes the limit $\mu_0 \downarrow 0$ at fixed lattice possible (Lemma 2.8): a counterterm computed at photon mass μ_0 on a torus contains the contribution of the constant mode of the photon, which has no limit. At order e^2 it is exactly $-3e^2/(\mu_0^2|T|)$: the constant mode c has variance $(\mu_0^2|T|)^{-1}$ in each of the $d = 3$ components (Definition 2.7), the seagull term $\frac{1}{2}e^2\sum_{x,\mu}\varepsilon^3 A_{x,\mu}^2$ $\mathrm{Re}\,\big(\bar\varphi(x)\varphi(x+\varepsilon e_\mu)\big)$ of (2.2) turns it into the mass shift $3e^2/(\mu_0^2|T|)$ at zero momentum, and the exchange graph gets no constant-mode contribution there, because its vertex at zero photon momentum is proportional to $\sin(\varepsilon p_\mu)$ and $p = 0$. The orders e^4 and $e^2\lambda$ contain further such terms. (iii) The additive vacuum constants of [24, 1] play no role for normalised expectations and are omitted.

2.3. **The states.** Write Π_0 for the orthogonal projection onto the constant bond functions $A_{x,\mu} = c_\mu$, $c \in \mathbb{R}^3$, and $\Pi^\perp := 1 - \Pi_0$; every bond function decomposes uniquely as $A = A^\perp + c$. Put $P := 2\pi/(|e|\,\ell_j) > 0$; the coupling $e \neq 0$ may have either sign, since the window constrains only e^2.

Definition 2.4 (observables at fixed lattice)**.** $\mathfrak{A}_\varepsilon$ is the set of complex-valued Borel functions of the plaquette variables $\big((dA)(p)\big)_p$ and of the site variables $\big(|\varphi(x)|^2\big)_x$. For $F \in \mathfrak{A}_\varepsilon$ with $F \ge 0$ or F bounded, and for a real function k on sites, put

$$\Psi_F(A) := \int F(A,\varphi)\, e^{-S_\flat^{\mathrm{m}}(A,\varphi)} \prod_x d^2\varphi(x), \qquad W_k(A) := \int e^{-S_\flat^{\mathrm{m}}(A,\varphi) + \sum_x \varepsilon^3 k(x)|\varphi(x)|^2} \prod_x d^2\varphi(x),$$

with $\flat = b_{T_j}(\varepsilon)$.

An integer-valued function n on bonds is called a *current*; it is *divergence-free* if $\sum_\mu \big(n_{x,\mu} - n_{x-\varepsilon e_\mu,\mu}\big) = 0$ at every site x.

Lemma 2.5 (positive type; current representation)**.** *For every real function k on sites, W_k is finite and*

$$W_k(A) = \sum_n w_k(n)\, e^{ie\varepsilon \sum_b n_b A_b}, \qquad w_k(n) = w_k(-n) \geq 0, \qquad \sum_n w_k(n) = W_k(0) < \infty, \tag{2.3}$$

the sum running over the divergence-free currents n. In particular $0 < W_k(A) \leq W_k(0)$. For a divergence-free current n and $\mu \in \{1,2,3\}$, the flux $\omega_\mu(n) := \sum_{x:\, x_\mu = s} n_{x,\mu}$ of n through the bonds in direction μ that start in the lattice plane $\{x_\mu = s\}$ is an integer that does not depend on s (the winding of n in direction μ), and for every constant c

$$e\varepsilon \sum_b n_b c_{\mu(b)} = e\ell_j \sum_\mu c_\mu\, \omega_\mu(n),$$

where $\mu(b)$ is the direction of b; hence $c \mapsto W_k(A^\perp + c)$ is periodic with period P in each coordinate. If $F \in \mathfrak{A}_\varepsilon$ is bounded, then $\int F\, e^{-S_b^{\mathrm{m}} + \sum_x \varepsilon^3 k(x)|\varphi(x)|^2} \prod_x d^2\varphi(x) = \sum_n w_{k,F}(n)\, e^{ie\varepsilon \sum_b n_b A_b}$, with complex weights $w_{k,F}(n)$ that depend on A only through dA and satisfy $|w_{k,F}(n)| \leq \|F\|_\infty\, w_k(n)$; in particular this integral is P-periodic in each c_μ.

The representation (2.3) is the current representation of [8, Thm. 4.1(2), p. 262], written here on the lattice torus.

Proof. In polar coordinates $\varphi(x) = r_x e^{i\vartheta_x}$, $|e^{ie\varepsilon A_{x,\mu}}\varphi(y) - \varphi(x)|^2 = r_x^2 + r_y^2 - 2r_x r_y \cos(\vartheta_y - \vartheta_x + e\varepsilon A_{x,\mu})$ for $y = x + \varepsilon e_\mu$. Every site is an endpoint of six bonds, hence

$$e^{-S_b^{\mathrm{m}} + \sum \varepsilon^3 k r^2} = \prod_x v_x(r_x) \prod_{(x,\mu)} \exp\big(\varepsilon r_x r_y \cos(\vartheta_y - \vartheta_x + e\varepsilon A_{x,\mu})\big), \qquad v_x(r) := e^{-3\varepsilon r^2 - \varepsilon^3(\frac{1}{2}b\, r^2 + \lambda r^4 - k(x) r^2)}.$$

Expand each bond factor as $e^{t\cos\psi} = \sum_{\nu\in\mathbb{Z}} I_\nu(t) e^{i\nu\psi}$, where $I_\nu = I_{-\nu} \geq 0$ on $[0,\infty)$ are the modified Bessel functions and $\sum_\nu I_\nu(t) = e^t$. The resulting multiple series converges absolutely after integration, because $\prod_{(x,\mu)} e^{\varepsilon r_x r_y} \leq \prod_x e^{3\varepsilon r_x^2}$ (by $r_x r_y \leq \frac{1}{2}(r_x^2 + r_y^2)$) and $\int_0^\infty e^{-\varepsilon^3(\frac{1}{2}b\, r^2 + \lambda r^4 - k(x) r^2)} r\, dr < \infty$ for $\lambda > 0$; the same holds with a bounded factor $F \in \mathfrak{A}_\varepsilon$ in the integrand, and F does not depend on the angles ϑ_x. So sum and integral may be interchanged. Integrating over the angles, $\int \prod_{(x,\mu)} e^{in_{x,\mu}(\vartheta_{x+\varepsilon e_\mu} - \vartheta_x)} \prod_x d\vartheta_x$ equals $(2\pi)^{|T_{j,\varepsilon}|}$ if n is divergence-free and 0 otherwise. This gives (2.3) with

$$w_k(n) = (2\pi)^{|T_{j,\varepsilon}|} \int \prod_x v_x(r_x)\, r_x\, dr_x \prod_{(x,\mu)} I_{n_{x,\mu}}(\varepsilon r_x r_{x+\varepsilon e_\mu}) \;\geq 0,$$

and the same formula with the factor F under the integral gives $w_{k,F}(n)$; since F depends on A only through dA, so does $w_{k,F}(n)$, and $|w_{k,F}(n)| \leq \|F\|_\infty w_k(n)$. Moreover $w_k(-n) = w_k(n)$ because $I_{-\nu} = I_\nu$, and $\sum_n w_k(n) = W_k(0)$ by (2.3) at $A = 0$. Positivity of $W_k(A)$ is clear from its definition, and $|W_k(A)| \leq \sum_n w_k(n)$. For a divergence-free current n, the difference of the fluxes through the planes $\{x_\mu = s + \varepsilon\}$ and $\{x_\mu = s\}$ is $-\sum_{x:\, x_\mu = s+\varepsilon} \sum_{\nu\neq\mu} (n_{x,\nu} - n_{x-\varepsilon e_\nu,\nu})$, which vanishes because the inner sums telescope over the periodic plane; so $\omega_\mu(n)$ does not depend on s. There are ℓ_j/ε such planes, so $\sum_{b\parallel e_\mu} n_b = (\ell_j/\varepsilon)\, \omega_\mu(n)$ and $e\varepsilon \sum_b n_b c_{\mu(b)} = e\ell_j \sum_\mu c_\mu \omega_\mu(n)$, which is P-periodic in each c_μ. The weights $w_{k,F}(n)$ depend on A through $dA = dA^\perp$ only, so the same conclusion holds for the integral with the factor F. □

Lemma 2.6 (the constant mode)**.** *For every $F \in \mathfrak{A}_\varepsilon$ (bounded or non-negative), $c \mapsto \Psi_F(A^\perp + c)$ is periodic with period P in each coordinate. If F is bounded, $|\Psi_F(A)| \leq \|F\|_\infty W_0(0)$.*

Proof. Let $\kappa := eP = \pm 2\pi/\ell_j$ (the sign of e) and $\varphi'(x) := e^{-i\kappa x_\mu}\varphi(x)$, with $x_\mu \in [0, \ell_j)$; since $e^{-i\kappa\ell_j} = 1$ this is a well-defined function on $T_{j,\varepsilon}$. For the bonds in direction μ, $e^{ie\varepsilon(A_{x,\mu} + P)}\varphi'(x + \varepsilon e_\mu) - \varphi'(x) = e^{-i\kappa x_\mu}\big(e^{ie\varepsilon A_{x,\mu}}\varphi(x + \varepsilon e_\mu) - \varphi(x)\big)$, including the bonds that wrap around the torus;

the bonds in the other directions and the single-site terms are unchanged. Hence $S_b^{\mathrm{m}}(A+Pe_\mu,\varphi') = S_b^{\mathrm{m}}(A,\varphi)$, where $A+Pe_\mu$ adds P to every bond in direction μ. Since F depends on φ only through $|\varphi|$, and on A only through dA, $F(A+Pe_\mu,\varphi') = F(A,\varphi)$; the change of variables $\varphi \mapsto \varphi'$ preserves Lebesgue measure, which gives the periodicity. The bound follows from $|\Psi_F(A)| \le \|F\|_\infty W_0(A)$ and Lemma 2.5. □

Definition 2.7 (the states)**.** Let γ_{μ_0} be the centred Gaussian measure on bond functions with covariance $C_{\mu_0} := (-\Delta_\varepsilon + \mu_0^2)^{-1}$ for the pairing $\langle\cdot,\,\cdot\rangle_\varepsilon$, that is, $\mathbb{E}[\langle f,\,A\rangle_\varepsilon\langle g,\,A\rangle_\varepsilon] = \langle f,\,C_{\mu_0}g\rangle_\varepsilon$. Let $\gamma^\flat$ be the product of the centred Gaussian measure with covariance $(-\Delta_\varepsilon)^{-1}\Pi^\perp$ on the range of $\Pi^\perp$ and the uniform probability measure on the period cell $[0,P)^3$ for c. For $F \in \mathfrak{A}_\varepsilon$ with $\mathbb{E}_\gamma[\Psi_{|F|}] < \infty$, $\gamma \in \{\gamma_{\mu_0}, \gamma^\flat\}$, put

$$\langle F\rangle_{\varepsilon,j,\mu_0} := \frac{\mathbb{E}_{\gamma_{\mu_0}}[\Psi_F]}{\mathbb{E}_{\gamma_{\mu_0}}[W_0]} \quad (0<\mu_0\le\mu_*), \qquad \langle F\rangle_{\varepsilon,j,\flat} := \frac{\mathbb{E}_{\gamma^\flat}[\Psi_F]}{\mathbb{E}_{\gamma^\flat}[W_0]}.$$

Under γ_{μ_0}, $A^\perp$ and c are independent and c is centred Gaussian with covariance $(\mu_0^2|T_j|)^{-1}\mathbf{1}_3$ (Lemma B.6). The state $\langle\cdot\rangle_{\varepsilon,j,\mu_0}$ is the expectation in the probability measure $Z^{-1}e^{-S_{\mu_0,b}}\prod_b dA_b \prod_x d^2\varphi(x)$ on $\mathbb{R}^{\text{bonds}} \times \mathbb{C}^{\text{sites}}$, restricted to $\mathfrak{A}_\varepsilon$ (Lemma B.7); it is King's torus model at photon mass μ_0 with the counterterm of the reference photon mass. The denominators are positive and finite by Lemma 2.5. In the state $\flat$ the non-constant photon modes are massless, and the constant mode c (the holonomy) is uniformly distributed on a period cell. By Lemma 2.6 the integrands Ψ_F and W_0 are P-periodic in c, so $\langle F\rangle_{\varepsilon,j,\flat}$ does not depend on the choice of the period cell; $F \mapsto \langle F\rangle_{\varepsilon,j,\flat}$ is the expectation of a probability measure on the σ-algebra generated by $\mathfrak{A}_\varepsilon$, namely of the normalised measure $\gamma^\flat(dA)\, e^{-S_b^{\mathrm{m}}(A,\varphi)}\prod_x d^2\varphi(x)$.

Lemma 2.8 ($\flat$ is the massless limit at fixed lattice)**.** *Fix ε and j. For every bounded $F \in \mathfrak{A}_\varepsilon$, $\lim_{\mu_0\downarrow 0}\langle F\rangle_{\varepsilon,j,\mu_0} = \langle F\rangle_{\varepsilon,j,\flat}$. The same holds for every $F \in \mathfrak{A}_\varepsilon$ with $\sup_{0<\mu_0\le\mu_*}\langle |F|^{1+\delta}\rangle_{\varepsilon,j,\mu_0} < \infty$ for some $\delta > 0$.*

Proof. We write $\langle\cdot\rangle_{\mu_0}$ and $\langle\cdot\rangle_\flat$ for the states at fixed ε, j. (i) The law of $A^\perp$ under γ_{μ_0} is the centred Gaussian with covariance $(-\Delta_\varepsilon+\mu_0^2)^{-1}\Pi^\perp$ on the finite-dimensional range of $\Pi^\perp$, on which $-\Delta_\varepsilon$ is positive definite; as $\mu_0 \downarrow 0$ it converges in total variation to the centred Gaussian with covariance $(-\Delta_\varepsilon)^{-1}\Pi^\perp$ (the densities converge pointwise, and Scheffé's lemma applies). (ii) By Lemmas 2.5 and 2.6, $\Psi_F(A^\perp + c)$ and $W_0(A^\perp + c)$ are bounded and P-periodic in each c_μ, so their γ_{μ_0}-expectations only involve the law of c modulo P, the wrapped Gaussian of variance $s = (\mu_0^2|T_j|)^{-1}$ in each coordinate. By Poisson summation its density on $[0,P)$ is $P^{-1}\big(1 + 2\sum_{n\ge1} e^{-2\pi^2n^2s/P^2}\cos(2\pi nx/P)\big)$, which converges uniformly to P^{-1} as $s\to\infty$. (iii) $A^\perp$ and c are independent under every γ_{μ_0} and under $\gamma^\flat$, so the joint laws converge in total variation, and numerator and denominator of $\langle F\rangle_{\mu_0}$ converge to those of $\langle F\rangle_\flat$; the limiting denominator is positive because $W_0 > 0$. For unbounded F put $F_n := F\mathbf{1}_{\{|F|\le n\}} \in \mathfrak{A}_\varepsilon$. Then $|\langle F\rangle_{\mu_0} - \langle F_n\rangle_{\mu_0}| \le n^{-\delta}\sup_{\mu_0}\langle|F|^{1+\delta}\rangle_{\mu_0}$ uniformly in μ_0. The same bound holds for $\flat$: by the bounded case $\langle\min(|F|,M)^{1+\delta}\rangle_\flat \le \sup_{\mu_0}\langle|F|^{1+\delta}\rangle_{\mu_0}$ for every M, and monotone convergence applies. The claim follows from the bounded case. □

2.4. **Observables and generating functionals.**
Wick constants. Put $\hat p^2 := \sum_\mu 4\varepsilon^{-2}\sin^2(\varepsilon p_\mu/2)$ and

$$W_\varepsilon := 2\int_{[-\pi/\varepsilon,\pi/\varepsilon)^3} \frac{d^3p}{(2\pi)^3}\,\frac{1}{\hat p^2 + m^2},$$

the free-field value of $\mathbb{E}|\varphi(0)|^2$ on $\varepsilon\mathbb{Z}^3$ at mass m. Let $X_\varepsilon(m)$ be the coefficient of e^2 in the perturbative expansion of $\langle|\varphi(0)|^2\rangle$ for the model on $\varepsilon\mathbb{Z}^3$ at scalar mass m and photon mass μ_*: the

seagull and exchange graphs, with the order-e^2 part of the counterterm on $\varepsilon\mathbb{Z}^3$ inserted, that is, King's [24, (3.33)] with $\Omega = \varepsilon\mathbb{Z}^3$. (It is not the torus counterterm of Definition 2.2, whose photon zero mode $e^2/(\mu_*^2|T|)$ would make z_ε depend on the torus.) Put $z_\varepsilon := e^2 X_\varepsilon(m)$. The same W_ε and z_ε are used on every torus, at every photon mass and in the state $\flat$.

Definition 2.9 (test functions and sampling)**.** Scalar test functions are real $h \in C_c(\mathbb{R}^3)$; curvature test functions are real 2-forms $g = \sum_{\mu<\nu} g_{\mu\nu}\, dx_\mu \wedge dx_\nu$ with $g_{\mu\nu} \in C_c(\mathbb{R}^3)$. Let $\operatorname{supp} h, \operatorname{supp} g \subset B_R$ and $\ell_j > 2R+2$, and identify T_j with the cell $[-\ell_j/2, \ell_j/2)^3$. The test functions are placed on $T_{j,\varepsilon}$ by sampling: h by its values $h(x)$ at the sites of the cell, and g by the value $g(p) := g_{\mu\nu}\big(x + \frac{\varepsilon}{2}(e_\mu + e_\nu)\big)$ at the centre of the plaquette $p = (x;\mu,\nu)$; both are extended periodically. The lattice norms are those of the sampled functions, $\|h\|_{L^1,\varepsilon} := \sum_x \varepsilon^3 |h(x)|$, $N_\varepsilon(h) := \max(\|h\|_{L^1,\varepsilon}, \|h\|_\infty)$ and $\|g\|_{2,\varepsilon}^2 := \sum_p \varepsilon^3 g(p)^2$; the continuum norms are $\|h\|_1$, $\|h\|_\infty$, $N(h) := \max(\|h\|_1, \|h\|_\infty)$ and $\|g\|_2^2 := \sum_{\mu<\nu} \|g_{\mu\nu}\|_{L^2}^2$.

Fields. The lattice fields are

$$F(g) := \langle g,\, dA\rangle_\varepsilon = \sum_p \varepsilon^3 g(p)\,(dA)(p), \qquad X_{\mathrm{R}}(h) = :|\varphi|^2:_{\mathrm{R}}(h) := \sum_x \varepsilon^3 h(x)\big(|\varphi(x)|^2 - W_\varepsilon - z_\varepsilon\big),$$

for real lattice 2-forms g and real lattice functions h, in particular for sampled test functions. Both are in $\mathfrak{A}_\varepsilon$, and so are their polynomials and exponentials. Since $\langle d^*g,\, \Pi_0 A\rangle_\varepsilon = \langle g,\, d\,\Pi_0 A\rangle_\varepsilon = 0$, $F(g) = \langle d^*g,\, A\rangle_\varepsilon$ does not depend on the constant mode c. For a real lattice function k,

$$\int e^{-S_\flat^{\mathrm{m}}(A,\varphi) + X_{\mathrm{R}}(k)} \prod_x d^2\varphi(x) = e^{-\sum_x \varepsilon^3 k(x)(W_\varepsilon + z_\varepsilon)}\, W_k(A),$$

so Lemma 2.5 applies to this weight, with $w_k(n)$ multiplied by a positive constant. Wilson loops are not among the observables: in three dimensions the sharp loop is not a tempered field, and smeared loops are functions of $F(g)$ by Stokes' theorem.

Generating functionals and the tube. For real test functions $g_1, \dots, g_n$, $h_1, \dots, h_p$ and $\zeta = (z, w) \in \mathbb{C}^{n+p}$ put

$$\Theta(\zeta) := \exp\Big[\sum_i z_i F(g_i) + \sum_k w_k X_{\mathrm{R}}(h_k)\Big], \qquad G_{\varepsilon,j,\mu_0}(\zeta) := \langle \Theta(\zeta)\rangle_{\varepsilon,j,\mu_0}, \quad \mu_0 \in (0,\mu_*] \cup \{\flat\}.$$

At fixed ε and j this is finite for every ζ: $|\Theta(\zeta)| = \Theta(\operatorname{Re}\zeta)$, the matter integral of $\Theta(\operatorname{Re}\zeta)$ is at most a constant times $e^{F(g)} W_k(0)$ with $g = \sum_i (\operatorname{Re} z_i) g_i$ and $k = \sum_k (\operatorname{Re} w_k) h_k$ (Lemma 2.5), and $e^{F(g)} = e^{\langle d^*g,\, A\rangle_\varepsilon}$ is integrable for the Gaussian measures of Definition 2.7; since on a compact set of ζ the function $\Theta(\operatorname{Re}\zeta)$ is bounded, by convexity, by the sum of its values at the corners of a box, dominated convergence shows that G_{ε,j,μ_0} is entire. Let $k(\zeta) := \sum_k (\operatorname{Re} w_k) h_k = k_+ - k_-$, with $k_\pm \geq 0$ its positive and negative parts. For $\sigma_+, H > 0$ the tube is

$$D_{\sigma_+,H} := \big\{\zeta \in \mathbb{C}^{n+p} :\ \|k_+(\zeta)\|_\infty < \sigma_+/2,\ \|k_-(\zeta)\|_\infty < H/2\big\},$$

with no restriction on z. Its base, the set of the admissible $\operatorname{Re} w$, is open and convex, so $D_{\sigma_+,H}$ is connected. The lattice Schwinger functions are

$$S_{n,p}^{\varepsilon,j,\mu_0}(g;h) := \Big\langle \prod_{i=1}^n F(g_i) \prod_{k=1}^p X_{\mathrm{R}}(h_k)\Big\rangle_{\varepsilon,j,\mu_0} = \partial_{z_1}\cdots\partial_{z_n}\partial_{w_1}\cdots\partial_{w_p} G_{\varepsilon,j,\mu_0}(0).$$

The class $\mathcal{D}$. A generating functional of the class $\mathcal{D}$ is $\Theta = \exp[F(g) - X_{\mathrm{R}}(q)]$ with g a real 2-form and $q \in C_c(\mathbb{R}^3)$, $q \geq 0$, $\|q\|_\infty < H/2$; it is the value of $\Theta(\zeta)$ at a real point ζ of the tube. At fixed lattice and torus, $\langle\Theta\rangle_{\varepsilon,j,\mu_0}$ is non-decreasing in the photon covariance for $\Theta \in \mathcal{D}$ [8, Cor. 6.3(1), p. 281] (on the lattice torus: Lemma 7.5), which is the source of the monotonicity in Theorem 3.1(iv). We write $E^{\mathcal{D}}$ for the set of real points ζ of the tube at which $\Theta(\zeta)$ belongs to $\mathcal{D}$, that is, $E^{\mathcal{D}} = \{\zeta \in D_{\sigma_+,H} \cap \mathbb{R}^{n+p} :\ \sum_k w_k h_k \leq 0\}$.

2.5. **Symmetries.** The cubic group $O_h \subset O(3)$ is the group of the 48 signed permutation matrices. The Euclidean group acts on test functions by

$$(\tau_a h)(x) := h(x-a), \qquad (R_* h)(x) := h(R^{-1}x), \qquad (R_* g)_x(v,w) := g_{R^{-1}x}(R^{-1}v, R^{-1}w),$$

for $a \in \mathbb{R}^3$ and $R \in O(3)$, that is, h transforms as a scalar and g as a 2-form (push-forward, signs included; for an improper R no extra sign appears). For $\nu \in \{1,2,3\}$ and $c \in \mathbb{R}$ let $\rho_{\nu,c}(x) := x - 2(x_\nu - c)e_\nu$ be the reflection in the plane $\{x_\nu = c\}$; on test functions it acts by push-forward, so that $(\rho_{\nu,c})_* g$ changes the sign of the components $g_{\mu\nu}$, $g_{\nu\mu}$ and reflects the argument. On the lattice, O_h acts about the site 0 on φ as on scalars and on A as on 1-forms ($A_{x,\mu} \mapsto \pm A$ on the image bond, with the sign -1 when the image bond is reversed), and charge conjugation acts by $(A,\varphi) \mapsto (-A, \bar\varphi)$. Invariance of the lattice states under these maps is Lemma 7.13 (Section 7).

2.6. **Schwinger families and the axioms.** A *Schwinger family* is a collection $S = (S_{n,p})_{n,p\geq 0}$, $S_{0,0} = 1$, of real multilinear functionals $S_{n,p}(g_1,\dots,g_n;h_1,\dots,h_p)$ of real curvature and scalar test functions in C_c, symmetric within each type. For a finite family of test functions, its generating functional on a tube D is a holomorphic function G^S on D whose Taylor coefficients at 0 are the $S_{n,p}$; when it exists we write $S(\Theta(\zeta)) := G^S(\zeta)$. For a polynomial $Q = \sum c_\alpha \prod F(g_{\alpha,i}) \prod X_{\mathrm{R}}(h_{\alpha,k})$ in the smeared fields with complex coefficients, $S(Q)$ is defined by linearity. We use the following forms of the axioms.

Definition 2.10 (OS0–OS3)**.** Let S be a Schwinger family.

- OS0 (regularity) There are $\sigma_+, H > 0$ and $C < \infty$ such that for every finite family of real test functions G^S exists on $D_{\sigma_+,H}$ and $|G^S(\zeta)| \leq \exp\left(\frac{1}{2}\|\sum_i (\mathrm{Re}\, z_i) g_i\|_2^2 + C\|k(\zeta)\|_{L^1}\right)$ (*tube form*); and, with $|f|_4 := \sup_{x\in\mathbb{R}^3,|\alpha|\leq 4}|(1+|x|^2)^2 D^\alpha f(x)|$, every $S_{n,p}$ evaluated on the components of the test functions satisfies $|S_{n,p}| \leq \sigma_{n+p} \prod_i |g_i|_4 \prod_k |h_k|_4$ with $\sigma_N \leq \alpha_\beta (N!)^\beta$ for every $\beta > 1$, i.e. the condition E0″ of [28, (4.2), p. 287] with $s = 4$.
- OS1($\mathcal{G}$) (invariance under a subgroup $\mathcal{G}$ of the Euclidean group $\mathbb{R}^3 \rtimes O(3)$) $S_{n,p}(\gamma_* g_1, \dots; \gamma_* h_1, \dots) = S_{n,p}(g_1,\dots;h_1,\dots)$ for all $\gamma \in \mathcal{G}$, with the actions of Section 2.5.
- OS2 (reflection positivity about coordinate planes) For every $\nu \in \{1,2,3\}$ and $c \in \mathbb{R}$, with $\theta := \rho_{\nu,c}$ acting on test functions as in Section 2.5 and antilinearly on coefficients: $S(\theta(Q)\,Q) \geq 0$ for every polynomial Q in the fields smeared with test functions supported in $\{x_\nu > c\}$; and $\sum_{i,l} \bar c_i c_l\, S(\theta(\Theta_i)\Theta_l) \geq 0$ for all $c \in \mathbb{C}^N$ and all exponentials $\Theta_i = \Theta(\zeta_i)$ of such fields whose $:|\varphi|^2:$-amplitudes satisfy $\sum |\mathrm{Re}\, w| \|h\|_\infty < \min(\sigma_+, H)/2$.
- OS3 (symmetry) $S_{n,p}$ is real for real test functions and symmetric under permutations within each field type; moreover $S_{n,p} = 0$ for odd n.

The cluster property refers to a probability law of the fields. Let X be a space of distributions on $\mathbb{R}^3$ with values in $\Lambda^2\mathbb{R}^3 \oplus \mathbb{R}$ (the curvature and the Wick square), carrying the σ-algebra generated by the evaluations $\Phi \mapsto \Phi(f)$, $f \in C_c^\infty(\mathbb{R}^3; \Lambda^2\mathbb{R}^3 \oplus \mathbb{R})$, and invariant under the translations $(\tau_a \Phi)(f) := \Phi(f(\cdot + a))$, $a \in \mathbb{R}^3$. Let μ be a probability measure on X invariant under every τ_a, and let $U(a)G := G \circ \tau_{-a}$ be the induced unitary representation of $\mathbb{R}^3$ on $L^2(\mu)$; thus $U(a)[\Phi(f)] = \Phi(f(\cdot - a))$.

Definition 2.11 (the cluster property OS4)**.** For $j \in \{1,2,3\}$ consider the two conditions:

- (OS4$_j$) (ergodicity of the translations in the direction e_j) every $G \in L^2(\mu)$ with $U(te_j)G = G$ for all $t \in \mathbb{R}$ is μ-almost surely constant;
- (OS4′$_j$) (clustering along the axis e_j) $\mathbb{E}_\mu[G'\,U(te_j)G] \to \mathbb{E}_\mu[G']\,\mathbb{E}_\mu[G]$ as $|t| \to \infty$, for all $G, G' \in L^2(\mu)$.

The measure μ satisfies *OS4* if (OS4$_j$) holds for $j = 1,2,3$. A Schwinger family satisfies OS4 if it is the family of moments of exactly one translation-invariant probability measure μ on X, and this μ satisfies OS4.

Definition 2.12 (forms of the cluster axiom; the weak form of OS4)**.** Let X, μ and U be as in Definition 2.11, and let $j \in \{1,2,3\}$ be the direction chosen as Euclidean time. In (i) assume moreover that X and μ are invariant under the map θ_j of X induced by the reflection $\rho_{j,0}$ in the plane $\{x_j = 0\}$ (Section 2.5), and let $\mathcal{P}_j^+$ be the set of polynomials in the fields $\Phi(f)$, $f \in C_c^\infty(\{x_j > 0\}; \Lambda^2\mathbb{R}^3 \oplus \mathbb{R})$ real, that belong to $L^2(\mu)$.

(i) (cluster property, [27, E4, p. 88]) For all $F, G \in \mathcal{P}_j^+$ and every $a \in \mathbb{R}^3$ with $a \perp e_j$,

$$\lim_{s\to\infty} \mathbb{E}_\mu\big[\overline{F \circ \theta_j}\, U(sa)G\big] = \mathbb{E}_\mu\big[\overline{F}\,\big]\, \mathbb{E}_\mu[G].$$

(ii) (ergodicity of the time translations, [19, (6.1.10), p. 92]) Condition $(\mathrm{OS4}_j)$ of Definition 2.11 holds.

(iii) (weak form) μ is ergodic for the action of the translation group $\mathbb{R}^3$: every $G \in L^2(\mu)$ with $U(a)G = G$ for all $a \in \mathbb{R}^3$ is μ-almost surely constant.

We call (i) and (ii) the single-axis forms of OS4, and say that μ satisfies *OS4 in the weak form* if (iii) holds.

Remark 2.13. (a) With e_j as Euclidean time, $(\mathrm{OS4}_j)$ is the form of the cluster axiom in [19, Sect. 6.1], ergodicity of the time translations. For $k \neq j$, condition $(\mathrm{OS4}'_k)$ contains the cluster property E4 of [27, p. 88] for the spatial directions $a = \pm e_k$ and test functions in C_c^∞: take $G' = \overline{F \circ \theta}$, with θ the map of X induced by the reflection in $\{x_j = 0\}$ (Section 2.5) and F, G square-integrable polynomials in fields smeared with test functions supported in $\{x_j > 0\}$. The axiom E4 asks for all spatial directions $a \perp e_j$. (b) If μ is invariant under the coordinate permutations, for instance under O_h as the laws of the limit states of Theorem 3.1 are, then $(\mathrm{OS4}_1)$, $(\mathrm{OS4}_2)$ and $(\mathrm{OS4}_3)$ are equivalent; in general they are three statements. With invariance under all rotations, the direction e_j could be replaced by any unit vector. (c) If μ satisfies the hypotheses (H1)–(H3) of Section 10, as the laws of the limit states do (Proposition 10.4), then for each j the conditions $(\mathrm{OS4}_j)$ and $(\mathrm{OS4}'_j)$ are equivalent: by Proposition 10.21 with $J = \{j\}$, $\mathbb{E}_\mu[G'\,U(te_j)G] \to \mathbb{E}_\mu[G'\,P_jG]$ as $|t| \to \infty$, where P_j is the orthogonal projection onto the $U(te_j)$-invariant vectors; $(\mathrm{OS4}_j)$ says that $P_jG = \mathbb{E}_\mu G$ for every G, and the equivalence follows as in Lemma 10.24. (d) $(\mathrm{OS4}_j)$ for one j implies that μ is ergodic for the action of $\mathbb{R}^3$, since an $\mathbb{R}^3$-invariant function is invariant under the translations along e_j. The converse fails: the law of Proposition 3.8 (realised on the present field space in Example 10.22) satisfies (H1)–(H3), is O_h-invariant and ergodic for $\mathbb{R}^3$, and its two-point function along e_1 has a positive limit, so $(\mathrm{OS4}'_1)$, and hence $(\mathrm{OS4}_1)$, fails. Theorem 3.6 gives the weak form of Definition 2.12, ergodicity under $\mathbb{R}^3$, for the ergodic components of the limit states, and Theorem 3.10 gives $(\mathrm{OS4}'_j)$ for part of the observables; OS4 itself is discussed in Section 14. (e) For μ satisfying (H1)–(H3), Proposition 10.41 in Section 10 shows that $(\mathrm{OS4}_j)$ holds if and only if $\operatorname{Ker} H_j = \mathbb{C}\Omega_j$ in the reconstruction of Proposition 10.16 in direction j; that the weak form (iii) of Definition 2.12 is equivalent to the uniqueness of the joint vacuum in one (equivalently, every) coordinate reconstruction, to clustering under diagonal separation and to clustering in the mean over cubes; and that, for the decomposition of Theorem 3.6, almost every component satisfies $(\mathrm{OS4}_j)$ if and only if $P_jG = \mathbb{E}[G \mid \mathcal{I}]$ for every $G \in L^2(\mu)$, where P_j is the orthogonal projection onto the $U(te_j)$-invariant vectors.

3. Main results

The objects in the statements below are defined in Section 2. Throughout this section $m > 0$, $0 < \lambda < 0.1m$, $0 < e^2 < 0.1m$ and $\mu_* := m$; the lattice states $\langle\cdot\rangle_{\varepsilon,j,\mu_0}$ are those of Definition 2.7 in the torus counterterm scheme of Definition 2.2.

3.1. Existence and OS0–OS3 at blocking factor 2.

Theorem 3.1 (existence and OS0–OS3, $L = 2$)**.** *Let $m > 0$, $0 < \lambda < 0.1m$, $0 < e^2 < 0.1m$, and $\mu_* := m$. There are $M_b, n_0 \in \mathbb{N}$, $\varepsilon_0 = 2^{-n_0}$, $\ell_0 \in 2M_b\mathbb{N}$ with $\ell_0 \geq 183/m$, $K_0 \in \mathbb{N}$, $\sigma_+ > 0$ and $C < \infty$ such that the following holds for the states $\langle\cdot\rangle_{\varepsilon,j,\mu_0}$ of Definition 2.7 with $L = 2$, $\varepsilon = 2^{-K}\varepsilon_0$, $K \geq K_0$, $j \geq 1$, and with $H := m^2$.*

(i) *(uniform tube bound) For every finite family of real test functions, every $\zeta \in D_{\sigma_+,H}$, every $\mu_0 \in (0, \mu_*] \cup \{\flat\}$, every $K \geq K_0$ and every j with ℓ_j larger than twice the radius of the supports plus 2,*

$$|G_{\varepsilon,j,\mu_0}(\zeta)| \;\leq\; G_{\varepsilon,j,\mu_0}(\operatorname{Re}\zeta) \;\leq\; \exp\Big(\tfrac{1}{2}\Big\|\sum_i (\operatorname{Re} z_i) g_i\Big\|_{2,\varepsilon}^2 + C\|k(\zeta)\|_{L^1,\varepsilon}\Big).$$

(ii) *(limits) Let $Q_0 \subset (0, \mu_*]$ be countable and accumulate at 0. Every sequence (K_l, j_l) with $K_l \to \infty$ and $j_l \to \infty$ has a subsequence along which, for every $\mu_0 \in Q_0$ and for $\mu_0 = \flat$, the lattice Schwinger functions $S_{n,p}^{\varepsilon_{K_l}, j_l, \mu_0}$ converge for all n, p and all test functions in C_c, and the generating functionals converge locally uniformly on $D_{\sigma_+,H}$. Call the limits S^{μ_0} ($\mu_0 \in Q_0$) and $S^\flat$. Along this subsequence, $S^0 := \lim_{\mu_0 \downarrow 0,\ \mu_0 \in Q_0} S^{\mu_0}$ exists, with the same limit along every sequence in Q_0 decreasing to 0, in the sense of Schwinger functions and locally uniformly on $D_{\sigma_+,H}$. Each $S \in \{S^{\mu_0}, S^\flat, S^0\}$ is, on test functions in C_c^∞, the family of moments of a Borel probability measure on $H^{-s'}_{\mathrm{loc}}(\mathbb{R}^3; \Lambda^2\mathbb{R}^3 \oplus \mathbb{R})$ for every $s' > 3$, whose characteristic functional is G^S at imaginary arguments.*

(iii) *(axioms) Each $S \in \{S^{\mu_0}, S^\flat, S^0\}$ satisfies OS0 with the constants σ_+, $H = m^2$, C of (i); OS1($\mathbb{R}^3 \rtimes O_h$), i.e. invariance under all translations of $\mathbb{R}^3$ and under O_h about every point of $\mathbb{R}^3$, g transforming as a 2-form and h as a scalar; OS2 about every coordinate plane; and OS3. Moreover*

$$|S_{n,p}(g;h)| \leq \mathrm{e}^{C\sigma_0}\,\mathrm{e}^n (n!)^{1/2} (\mathrm{e}/\sigma_0)^p\, p! \prod_i \|g_i\|_2 \prod_k N(h_k), \qquad \sigma_0 := \min(\sigma_+, H)/2.$$

(iv) *(monotonicity) For $\Theta \in \mathcal{D}$, $\mu_0 \mapsto S^{\mu_0}(\Theta)$ is non-increasing on Q_0, $S^{\mu_0}(\Theta) \leq S^\flat(\Theta)$, and $S^0(\Theta) \leq S^\flat(\Theta)$.*

The constants of (i) are $\sigma_+ = g/2$, where $g > 0$ is the convexity constant of Proposition 6.4 (not a test function), and $C = \mathsf{K} = \max(\kappa_+, 2S_\flat^B)$, with κ_+ of Proposition 6.4 and $S_\flat^B$ of Proposition 6.7, at the auxiliary parameters $(\eta, m_1, m_2) = (\frac{1}{2}, 3m, 4m)$ (Theorem 6.8 and Lemma I.19).

Remark 3.2 (cubic symmetry, not rotations). The group in OS1 is $\mathbb{R}^3 \rtimes O_h$: the translations of $\mathbb{R}^3$ and the 48 signed permutations about every point. On the lattice, translations by $\varepsilon\mathbb{Z}^3$ and the cubic group about a site are exact symmetries of every state on a cubic torus; they pass to the limits, and continuity extends the translations to $\mathbb{R}^3$. Rotations outside O_h are not symmetries of any regularised state, and Theorem 3.1 makes no claim about them; Section 14.1 discusses them. In the reconstruction of Section 11 only the subgroup D_4 of O_h fixing the reconstruction direction acts on the Hilbert space, together with the translations orthogonal to it.

Remark 3.3 (order of limits). (a) In (ii) the spacing and the torus go to their limits jointly. Every iterated limit, first $\varepsilon \to 0$ at fixed T_j and then $j \to \infty$, at fixed μ_0, is a limit of the joint family in the sense of (ii), and every limit $\mu_k \downarrow 0$ of such iterated limits is a limit point of the joint family $(\varepsilon, T_j, \mu_0) \to (0, \mathbb{R}^3, 0)$; so (iii) holds in every order (Proposition 8.6). (b) The photon mass is removed last, along the countable set Q_0 and along one subsequence of (ε, T_j) common to all $\mu_0 \in Q_0$. The limit S^0 is monotone in μ_0 on $\mathcal{D}$ and does not depend on the sequence in Q_0, but it may depend on the common subsequence; we do not prove that it is independent of it. Continuity of $\mu_0 \mapsto S^{\mu_0}$ is not claimed. (c) The limit $S^\flat$ is taken in the other order: $\mu_0 \downarrow 0$ first, at fixed lattice (Lemma 2.8), then the lattice limit. Part (iv) is the only relation between the two orders that we prove.

Remark 3.4 (the regulator data). (a) The constants ε_0, ℓ_0 and K_0 come from Bałaban's renormalisation group with blocking factor $L = 2$, small-field parameter $p = 6$, perturbative order $\bar{n} = 13$ with order cap $\bar{N} = 169$ and localisation radius $R \geq R^{**}$, and from the conditions on these data listed in Appendix A. The data are chosen in stages, in an order in which each condition depends only on data chosen at earlier stages (Lemma A.4). The bound $\ell_0 \geq 183/m$ is the torus floor $128/M_1 = 182.857\ldots/m$ at the smallest scalar mass $M_1 = 0.7m$ used in the proof, rounded up (Lemma A.10). (b) Lemma A.5 shows that all these conditions can be met at every (e, λ, m) of Theorem 3.1, with the auxiliary parameters $(\eta, m_1, m_2) = (\frac{1}{2}, 3m, 4m)$, which lie in the window (1.3) for every such (e, λ, m) (Lemma I.19). An explicit admissible point is described in Section A.8; nothing depends on it. (c) Among the conditions are the condition (E.edge) on ε_0, produced by the proof of Lemma 5.6, and the torus-size condition (T.per), used in Lemma 5.22. (d) Four ingredients that are first established at the anchor masses m_1, m_2 are used at all masses $M \in (0, m_2]$; Lemma A.6 supplies the extensions.

3.2. **Blocking factor 3.** For blocking factor $L = 3$ the lattices and tori of Section 2.1 are $\varepsilon = 3^{-K}\varepsilon_0$, $\varepsilon_0 = 3^{-n_0}$, and $\ell_j = j!\,\ell_0$ with $\ell_0 \in 18M_b\mathbb{N}$; Lemma 2.1 holds with $c_3 = 18$. The states of Definition 2.7 are defined on these tori in the same way.

Theorem 3.5 (blocking factor 3, under a uniform tube bound)**.** *Let $m > 0$, $0 < \lambda < 0.1m$, $0 < e^2 < 0.1m$, $\mu_* := m$ and $H := m^2$, and consider the states $\langle\cdot\rangle_{\varepsilon,j,\mu_0}$ of Definition 2.7 with $L = 3$, $\varepsilon = 3^{-K}\varepsilon_0$, $\varepsilon_0 = 3^{-n_0}$, $\ell_0 \in 18M_b\mathbb{N}$ and $\ell_0 \geq 183/m$. Assume that for some $M_b, n_0 \in \mathbb{N}$, some such ℓ_0, some $K_0 \in \mathbb{N}$, $\sigma_+ > 0$ and $C < \infty$ the following lattice tube bound holds: for every $K \geq K_0$, every $j \geq 1$, every $\mu_0 \in (0, \mu_*] \cup \{\flat\}$, all real lattice test functions $\mathbf{g}_1, \ldots, \mathbf{g}_n$, $h_1, \ldots, h_p$ on $T_{j,\varepsilon}$ and every ζ in the lattice tube $\mathcal{D}_{\sigma_+,H}$ of Section 6.6,*

$$|G_{\varepsilon,T_j,\mu_0}(\zeta)| \leq G_{\varepsilon,T_j,\mu_0}(\operatorname{Re}\zeta) \leq \exp\Big(\tfrac{1}{2}\Big\|\sum_i (\operatorname{Re} z_i)\mathbf{g}_i\Big\|_{2,\varepsilon}^2 + C\|k(\zeta)\|_{L^1,\varepsilon}\Big). \tag{3.1}$$

Then statements (ii)–(iv) of Theorem 3.1 hold for this family, with these constants σ_+, C and $H = m^2$.

Proof. Sections 7–9 derive (ii)–(iv) of Theorem 3.1 from the lattice tube bound of Theorem 6.8 and from statements on one lattice torus at a time. Apart from the tube bound they use only the properties (P1)–(P4) of Section 7.1. Of these, (P1)–(P3) do not involve the blocking factor, and (P4), the nesting of the lattices and the evenness of the number of sites per side, is Lemma 2.1 with $c_3 = 18$. They also use the density in $\mathbb{R}^3$ of $\bigcup_K \varepsilon_K\mathbb{Z}^3$, here $\mathbb{Z}[1/3]^3\varepsilon_0$. With (3.1) in place of Theorem 6.8, the same proofs apply. □

Theorems 3.6, 3.10(a),(c) and 3.11 below, and the results of Sections 10–13 on which they rest, hold for the limit states of Theorem 3.5, with the same proofs. Theorem 3.10(b) and Proposition 10.35 are not asserted for these states: their proofs use Propositions 6.5 and 6.7, not only for the bound $\Delta_S \leq \kappa_+ + 2S_\flat^B$ but already for the existence of the limit pressures, on which the definition of Δ_S rests. We expect that (3.1) can be obtained by carrying out Sections 4–6 and Appendix A with blocking factor 3; this is not done here. The interest of blocking factor 3 lies in the rotations (Section 14.1).

3.3. **Weak clustering.** Let $\mathcal{V} := \Lambda^2\mathbb{R}^3 \oplus \mathbb{R} \cong \mathbb{R}^4$ (the curvature and the Wick square). For $r > 0$ let $\mathcal{H}^{-r} := \mathcal{H}^{-r}(\mathbb{R}^3; \mathcal{V})$ be the Hermite–Sobolev space of order $-r$ of $\mathcal{V}$-valued distributions: with $\mathcal{A} := -\Delta + |x|^2 + 1$, acting componentwise, and $\|f\|_r := \|\mathcal{A}^{r/2}f\|_{L^2}$, $\mathcal{H}^{-r}$ is the dual of the completion of $\mathcal{S}(\mathbb{R}^3; \mathcal{V})$ in $\|\cdot\|_r$. It is a separable Hilbert space. Translations act on $\mathcal{H}^{-r}$ by $(\tau_a\Phi)(f) := \Phi(f(\cdot + a))$, and the reflection $\rho_{j,c}$ in the plane $\{x_j = c\}$ by the map $\theta_{j,c}$ induced by the push-forward of Section 2.5, with $\theta_j := \theta_{j,0}$ (on the curvature components this is the reflection twisted by the signs of the 2-form). For an open $U \subset \mathbb{R}^3$ let $\Sigma(U)$ be the σ-algebra generated by

the evaluations $\Phi \mapsto \Phi(f)$, $f \in C_c^\infty(U;\mathcal{V})$, and write $H_j^+(c) := \{x : x_j > c\}$, $U(a)G := G \circ \tau_{-a}$, and $\mathcal{I}$ for the σ-algebra of translation-invariant Borel sets.

Theorem 3.6 (weak clustering)**.** *Let S be one of the limit states S^{μ_0} ($\mu_0 \in Q_0$), $S^\flat$, S^0 of Theorem 3.1. For every even $r \geq 14$ there is exactly one Borel probability measure μ on $\mathcal{H}^{-r}$ whose mixed moments on test functions in $C_c^\infty(\mathbb{R}^3;\mathcal{V})$ are the Schwinger functions of S, and there are a probability space (Ξ,ν) and Borel probability measures μ_ξ on $\mathcal{H}^{-r}$, with $\xi \mapsto \mu_\xi(B)$ measurable for every Borel set B, such that $\mu = \int \mu_\xi \, d\nu(\xi)$ and, for ν-almost every ξ:*

(a) *μ_ξ is invariant under all translations and ergodic for the action of $\mathbb{R}^3$, and invariant under $\theta_1, \theta_2, \theta_3$;*

(b) *μ_ξ is reflection positive: $\mathbb{E}_{\mu_\xi}[\overline{F \circ \theta_{j,c}}\, F] \geq 0$ for every j, every $c \in \mathbb{R}$ and every $F \in L^2(\Sigma(H_j^+(c)), \mu_\xi)$;*

(c) *(clustering in the mean) for all $G, G' \in L^2(\mu_\xi)$, $\lim_{T\to\infty} (2T)^{-3} \int_{[-T,T]^3} \mathbb{E}_{\mu_\xi}[G'\, U(a) G]\, da = \mathbb{E}_{\mu_\xi}[G']\, \mathbb{E}_{\mu_\xi}[G]$;*

(d) *(clustering under diagonal separation) for all $G, G' \in L^2(\mu_\xi)$, $\mathbb{E}_{\mu_\xi}[G'\, U(a) G] \to \mathbb{E}_{\mu_\xi}[G']\, \mathbb{E}_{\mu_\xi}[G]$ as $\min_j |a_j| \to \infty$;*

(e) *(reconstruction in each coordinate direction) for each j, the Osterwalder–Schrader reconstruction from μ_ξ in direction j gives a Hilbert space $\mathcal{H}_j$, a unit vector Ω_j, a self-adjoint operator $H_j \geq 0$ and a commuting unitary representation of the translations orthogonal to e_j; the joint fixed space of the operators e^{-tH_j} ($t \geq 0$) and of these translations is $\mathbb{C}\Omega_j$ (a unique joint vacuum).*

For μ itself, (c) and (d) hold with the right side replaced by $\mathbb{E}_\mu\big[\mathbb{E}[G' \mid \mathcal{I}]\, \mathbb{E}[G \mid \mathcal{I}]\big] = \int \mathbb{E}_{\mu_\xi}[G']\, \mathbb{E}_{\mu_\xi}[G]\, d\nu(\xi)$.

(f) *(decomposition along $\mathbb{R}^3 \rtimes O_h$) The decomposition may also be chosen with cubic-invariant components: there are a probability space (Ξ',ν') and Borel probability measures μ'_ζ on $\mathcal{H}^{-r}$, with $\zeta \mapsto \mu'_\zeta(B)$ measurable for every Borel set B, such that $\mu = \int \mu'_\zeta \, d\nu'(\zeta)$ and, for ν'-almost every ζ, the measure μ'_ζ is invariant under $\mathbb{R}^3 \rtimes O_h$, is reflection positive in the sense of (b), and is the uniform mixture $\mu'_\zeta = k^{-1} \sum_{i=1}^k \mu_i$ of the k distinct measures $\mu_1, \dots, \mu_k$ of one orbit of the group S_3 of coordinate permutations acting on the translation-invariant measures that are ergodic for $\mathbb{R}^3$; thus $k = k(\zeta) \in \{1,2,3,6\}$, and for each j the joint fixed space of the operators e^{-tH_j} ($t \geq 0$) and of the transverse translations in the reconstruction from μ'_ζ in direction j, defined as in (e), has dimension k (Theorem 10.12).*

In the terms of Definition 2.12, part (a) says that almost every component μ_ξ satisfies OS4 in the weak form.

Remark 3.7 (the components). For ν-almost every ξ the moments of μ_ξ exist for all Schwartz test functions and satisfy a bound $|\mathbb{E}_{\mu_\xi} \prod_{i=1}^N \Phi(f_i)| \leq \sigma_N^\xi \prod_i |f_i|_k$ for a fixed Schwartz norm $|\cdot|_k$, with $\sigma_N^\xi \leq \alpha_\beta(\xi)(N!)^\beta$ for every $\beta > 1$ (Corollary 10.43(iv)); the constants depend on ξ. So the components satisfy a growth condition of the type of the second part of OS0, with the norm $|\cdot|_k$ in place of $|\cdot|_4$ and with ξ-dependent constants; we do not claim OS0 for them, in particular not the tube bound of Theorem 3.1(i). The components are invariant under the translations and the three coordinate reflections (part (a) of Theorem 3.6), so they satisfy OS1 for the group generated by these. A coordinate permutation maps a component to a component (Remark 10.11), and nothing forces a component to be invariant under the permutations; grouping the components along the orbits of O_h gives the O_h-invariant components of part (f) of Theorem 3.6, whose joint vacuum has dimension $k \in \{1,2,3,6\}$. The moments of a component are symmetric and real, and those of degree one in the curvature vanish (Theorem 10.27); the vanishing of its moments of odd degree three or more in the curvature, which holds for S (Theorem 9.8), is not known for a single component.

Proposition 3.8 (the axis obstruction)**.** *For r large there is a centred Gaussian measure on $\mathcal{H}^{-r}(\mathbb{R}^3)$ that satisfies the hypotheses (H1)–(H3) of Section 10.1 with the untwisted reflections $\Phi \mapsto \Phi \circ \rho_j$, is invariant under translations and under O_h, is ergodic for the action of $\mathbb{R}^3$, and whose two-point function does not tend to zero along any coordinate axis: $\mathbb{E}[\Phi(f)\,U(te_1)\Phi(f)]$ has a positive limit as $t \to \infty$ for every $f \in C_c^\infty(\mathbb{R}^3)$ whose integral over x_1 is not identically zero.*

The example is $\Phi = \psi_1(x_2,x_3) + \psi_2(x_1,x_3) + \psi_3(x_1,x_2)$ with independent two-dimensional massive free fields ψ_j (Example 10.22, where it is realised on the field space $\mathcal{V}$ with vanishing curvature component, so that the twisted and the untwisted reflections coincide). So single-axis clustering, even of the two-point function, is not a consequence of translation invariance, cubic invariance, ergodicity and reflection positivity.

Proposition 3.9 (what closes single-axis clustering)**.** *Let μ satisfy (H1)–(H3), let $\mathcal{F} \subset L^2(\mu)$ be a set whose linear span is dense, and let P_j be the orthogonal projection onto the functions invariant under the translations along e_j. The following are equivalent: (i) $P_jG = \mathbb{E}[G \mid \mathcal{I}]$ for all $G \in \mathcal{F}$ and $j = 1,2,3$; (ii) for all $G \in \mathcal{F}$, $G' \in L^2(\mu)$ and each j, $\mathbb{E}[G'\,U(te_j)G] \to \mathbb{E}\big[\mathbb{E}[G' \mid \mathcal{I}]\,\mathbb{E}[G \mid \mathcal{I}]\big]$ as $|t| \to \infty$; (iii) the same limit holds as $|a| \to \infty$ in any manner. If μ is a centred Gaussian measure with covariance $\widehat{C}$ in Fourier space, then (i) holds for the linear fields $\Phi(f)$ (a statement about (i) alone; their linear span is not dense) if and only if $\widehat{C}(\{p : p_j = 0\} \setminus \{0\}) = 0$ for $j = 1,2,3$.*

Theorem 3.10 (single-axis clustering)**.** *Let S be one of the limit states S^{μ_0} ($\mu_0 \in Q_0$), $S^\flat$, S^0 of Theorem 3.1, μ its law on $\mathcal{H}^{-r}$ (Theorem 3.6), P_j the orthogonal projection of $L^2(\mu)$ onto the e_j-invariant vectors, and $\mathcal{M}$ the σ-algebra generated by the Wick-square fields $X_{\mathrm{R}}(h)$, $h \in C_c^\infty(\mathbb{R}^3)$ real, completed by μ-null sets. (Section 10.8 works with the law of S on $H^{-s}_{\mathrm{loc}}(\mathbb{R}^3;\mathcal{V})$; Lemma 10.5 carries every statement to the law on $\mathcal{H}^{-r}$, with $\mathcal{M}$ replaced by its preimage under the restriction map.)*

(a) *For every real 2-form $\mathbf{g}$ with components in $C_c^\infty(\mathbb{R}^3)$ and every $Y \in L^2(\mathcal{M})$: $F(\mathbf{g})Y \in L^2(\mu)$, $\mathbb{E}_\mu[F(\mathbf{g})^2|Y|^2] \le \|\mathbf{g}\|_2^2\,\mathbb{E}_\mu|Y|^2$, $P_j(F(\mathbf{g})Y) = 0$ for $j = 1,2,3$, and for every $G' \in L^2(\mu)$*

$$\lim_{|a|\to\infty} \mathbb{E}_\mu\big[G'\,U(a)\big(F(\mathbf{g})Y\big)\big] = 0 = \mathbb{E}_\mu[G']\,\mathbb{E}_\mu[F(\mathbf{g})Y],$$

the limit taken in any manner, in particular along each coordinate axis.

(b) *Let $\Delta_S \ge 0$ be the latent density of the Wick square in S (Definition 10.34). For every $h \ge 0$, $|P_jX_{\mathrm{R}}(h) - \mathbb{E}_\mu X_{\mathrm{R}}(h)| \le \Delta_S\|h\|_1$ μ-almost surely, and $\Delta_S \le \kappa_+ + 2S_\flat^B$ with the constants of Theorem 6.8. If $\Delta_S = 0$, the conclusion of (a) holds for every G in the closed span of $L^2(\mathcal{M})$ and of the products $F(\mathbf{g})Y$, with limit $\mathbb{E}_\mu[G']\,\mathbb{E}_\mu[G]$.*

(c) *If hypothesis (D) of Corollary 10.38 holds, then μ is ergodic, S satisfies OS4 (Definition 2.11), and the conclusion of (a) holds for all $G, G' \in L^2(\mu)$, with limit $\mathbb{E}_\mu[G']\,\mathbb{E}_\mu[G]$.*

3.4. **Correlation signs and the curvature two-point function.** For a limit state S let $S^T_{n,p}$ be its truncated functions (Definition 12.1) and $M(g;h) := S^T_{2,1}(g,g;h)$, the truncated correlation of $F(g)^2$ and $X_{\mathrm{R}}(h)$. For a real vector field $G \in C_c(\mathbb{R}^3;\mathbb{R}^3)$ let $B(G) := F(\star^{-1}G)$ be the smeared magnetic field, $\mathfrak{b}_S(G,G')$ its two-point function, $P_T(p)$ the orthogonal projection of $\mathbb{C}^3$ onto $p^\perp$, and $\mathsf{Z}_S(G) := \limsup_{R\to\infty} R^3\mathfrak{b}_S(G_R,G_R)$ with $G_R := R^{-3}G(\cdot/R)$ (Section 13); S is *massless in the curvature sector* if $\mathsf{Z}_S(G) > 0$ for some G (Definition 13.6).

Theorem 3.11 (signs, and the curvature two-point function)**.** *Let $S \in \{S^{\mu_0}\ (\mu_0 \in Q_0), S^\flat, S^0\}$ be a limit state of Theorem 3.1.*

(a) *(signs) For all real 2-forms g, g' with components in $C_c(\mathbb{R}^3)$ and all $h, h' \in C_c(\mathbb{R}^3)$ with $h, h' \ge 0$,*

$$M(g;h) \le 0, \qquad S^T_{0,2}(h,h') \ge 0, \qquad S^T_{4,0}(g,g,g',g') + 2\,S_{2,0}(g,g')^2 \ge 0.$$

If $M(g;h) \neq 0$ for one such pair, S is not decoupled (Definition 12.1); hence it is not Gaussian, and the curvature and the Wick square are not independent under its law.

(b) *(the curvature two-point function) There is a measurable m_S with $0 \le m_S(p) \le P_T(p)$ for a.e. p and $\mathfrak{b}_S(G,G') = (2\pi)^{-3} \int \overline{\hat{G}(p)}^{\mathsf{T}}\, m_S(p) \hat{G}'(p)\, dp$; in particular $0 \le \mathsf{Z}_S(G) \le \|\Pi_T G\|_2^2$, with Π_T the Fourier multiplier by P_T; the right side is the value for the free massless photon. If $S = S^{\mu_0}$ with $\mu_0 > 0$, then $m_S(p) \le \min(1, |p|^2/\mu_0^2) P_T(p)$ and $\mathsf{Z}_S \equiv 0$: the states at positive photon mass have no massless component. For $\mu_0 \in Q_0$, $0 = \mathsf{Z}_{S^{\mu_0}} \le \mathsf{Z}_{S^0} \le \mathsf{Z}_{S^\flat}$.*

(c) *(what masslessness would give) If $S \in \{S^\flat, S^0\}$ is massless in the curvature sector, its curvature correlations are not integrable, in particular not exponentially decaying, and in every coordinate direction the curvature sector of its reconstruction has no mass gap: the generator H_F of Proposition 13.8(a) has* $\inf \operatorname{spec} H_F = 0$.

Theorem 12.9 shows, under a remainder bound (R) comparing $e^{-2}M$ with its one-loop value and under Hypothesis 12.7, neither of which is proved here, that for every real 2-form g with $d^*g \not\equiv 0$ and every $0 \le h \not\equiv 0$ in $C_c^\infty(\mathbb{R}^3)$ one has $M(g;h) < 0$ for $e^2 + \lambda$ small enough depending on (g,h); hence these limit states are not decoupled (Section 14.3).

Remark 3.12 (what is not claimed). The theorems above do not assert uniqueness of the limits, the equality $S^0 = S^\flat$ (part (iv) of Theorem 3.1 gives one inequality on $\mathcal{D}$), or the coincidence of the limits with King's cube limits. They do not assert that the limit states are non-Gaussian, that they or their ergodic components satisfy OS4 in the single-axis forms (i), (ii) of Definition 2.12 (Theorem 3.6 gives the weak form (iii) for the ergodic components, and Theorem 3.10 the single-axis forms for part of the observables), that the translation-ergodic components are invariant under O_h, that the photon is massless at $\mu_0 = 0$, that the states lie in the symmetric phase (the proofs do not use the phase, and one-loop perturbation theory predicts a first-order transition surface inside the range of couplings of Theorem 3.1, Remark 14.1), or invariance under rotations outside O_h. Section 14 discusses each of these questions.

3.5. **Where the results are proved.** The proofs of Theorems 3.1 and 5.2 use the statements collected in Appendix K, whose proofs are not included in this version.

Theorem 3.1. Part (i) is Theorem 6.8 of Section 6. Its curvature factor is the Gaussian domination of the photon on the torus (Lemma 7.3). The factor for positive $:|\varphi|^2:$-sources is the convexity-and-chessboard bound on the same torus (Propositions 6.4 and 6.5); its input is Bałaban's two-sided bound (1.2) at the scalar masses $0.7m$ and m on every T_j (Proposition 6.2), together with the counterterm gap of Theorem 6.1. The factor for negative sources (Proposition 6.7) follows from Theorem 5.2, the lower vacuum bound of Theorem 4.1 at m_1, and the counterterm gap between m_1 and m_2. The proofs of these statements use the results of Section 4 and Appendices A and C–I. Part (ii) is compactness (Montel's theorem on the tube and a diagonal argument), tightness on $H^{-s'}_{\mathrm{loc}}$, and, for S^0, the monotonicity on the class $\mathcal{D}$ at fixed lattice passed to the limit along the common subsequence (Section 8, with Lemma 8.9 for the law of S^0). Part (iii) is Section 9: the bound passes to the limit, and the lattice symmetries and lattice reflection positivity on tori (Section 7) are preserved under the limits. Part (iv) is the monotonicity in part (ii).

Theorem 3.6 and Propositions 3.8, 3.9. These are proved in Section 10: the ergodic decomposition with inheritance of the reflections and of reflection positivity, which rests on the fact that a function invariant under translations along e_j is fixed by θ_j (Section 10.4); the reconstruction in a coordinate direction and the identification of its zero-energy vectors with the e_j-invariant functions (Section 10.5); clustering in the mean from the mean ergodic theorem, and clustering under diagonal separation from the telescoping identity $1 = P_1P_2P_3 + (1-P_1) + P_1(1-P_2) + P_1P_2(1-P_3)$ and the spectral theorem for each H_j (Section 10.6); the verification of (H1)–(H3) for the limit states (Section 10.2); and Corollary 10.43 with Lemma 10.5. Proposition 3.8 is Example 10.22, carried to $\mathcal{H}^{-r}$ by the carrier argument of Lemma 10.5(b), and Proposition 3.9 is Proposition 10.23.

Theorems 3.10 and 3.11. Theorem 3.10 is Theorem 10.40(a)–(c) of Section 10.8. Part (a) is Theorem 10.27, whose input is a domination of the curvature conditional on the matter on every lattice torus (Lemma 10.26); part (b) is Proposition 10.35 and Theorem 10.33, from the chessboard estimate at the photon mass of the state and the association of the radial field (Lemma 10.31); part (c) is Corollary 10.38, from the multiple-reflection bound in infinite volume (Theorem 10.37). Theorem 3.11(a) is Theorem 12.4(b) and Corollary 12.5; (b) is Theorem 13.5 and Theorem 13.7(i)–(iii); (c) is Corollary 13.9.

4. Ultraviolet input from the literature

This section lists the results from the literature that the construction of Theorem 3.1 uses, in the form in which they are used, with the page where each is printed: Bałaban's three papers on the $(\mathrm{Higgs})_{2,3}$ model in a finite volume [1, 2, 3], his paper on lattice Green's functions [4], and King's two papers on the $U(1)$ Higgs model [24, 25], together with the classical results (correlation inequalities, power counting) of Subsection 4.6. Three printed proofs are incomplete at places that the construction uses; Section 5 completes them (Lemma 5.6, Proposition 5.8 and Proposition 5.21).

4.1. **Bałaban's torus and scales.** Bałaban works on the lattice torus

$$T_\varepsilon = \{x \in \varepsilon\mathbb{Z}^3 :\ -L_\mu \le x_\mu < L_\mu,\ \mu = 1,2,3\}, \qquad |x-y| = \max_\mu \min\{|x_\mu - y_\mu|,\, 2L_\mu - |x_\mu - y_\mu|\},$$

with sides $2L_\mu$ such that $\varepsilon^{-1}L_\mu \in L^K M_b\mathbb{N}$ [1, (1.2)–(1.3), p. 604]; here $L \ge 2$ is the block factor, M_b the side of a big ("large") block, and the distance is the torus sup-distance. All distances in Sections 4–6 are measured in this metric. For the lattices of Theorem 3.1, $\varepsilon = L^{-K}\varepsilon_0$ with $L = 2$, so the k-th renormalisation scale is $t_k := L^k\varepsilon$ and the last one is $t_K = \varepsilon_0$. In the k-th step all objects are rescaled by t_k; scale-k blocks then have side 1 and the lattice spacing is $\eta_k := L^{-k}$ (Bałaban writes η; we reserve η for the parameter of the window $\mathfrak{W}$ of (1.3)). The large-field regions of the k-th step are organised in nested unions of big blocks $\Lambda_0 \supset \Lambda_1 \supset \cdots$, where Λ_{i+1}^c is the union of the big blocks within distance $r(t)$ of Λ_i^c, and

$$r(t) = R\,(1 + \log t^{-1})^{\varrho}, \qquad \varrho > 1, \quad R > R_0, \tag{4.1}$$

[2, (2.7)–(2.8), p. 558]; we take $\varrho = 2$. Bałaban's small-field parameter is $p(t) := b_0(1 + \log t^{-1})^p$; we take $p = 6$. The constants R and b_0 are fixed with the regulator data (Appendix A), and R_0 is the constant of Proposition 4.5. At step k the sets of scale $k-1$ are written $\Lambda_i^{(k-1)\prime}$, and in scale-k units consecutive layers are at distance at least $\rho_k := (r(t_{k-1}) - M_b)/L$ [2, p. 558, and Sect. B, pp. 566 ff.].

The tori of the construction are the cubic tori T_j of Subsection 2.1, of side $\ell_j = j!\,\ell_0$ with $\ell_0 \in 2M_b\mathbb{N}$; the regulator data have $\ell_0 \ge 183/m$ (Appendix A). By Lemma 2.1, for every spacing $\varepsilon = 2^{-K}\varepsilon_0$ the lattice torus $T_{j,\varepsilon}$ is a torus of the form [1, (1.2)] (part (ii) with $k = K$), and the number $N = \ell_j/\varepsilon$ of sites per side is even (part (i)). We write $K_0' \in \mathbb{N}$ for the index from which the torus-uniform bounds of Theorem 5.2 hold; it is fixed with the regulator data (Appendix A). The spacing threshold of Theorem 3.1 is

$$K_0 := \max(K_0',\, n_0 + 1).$$

4.2. **Bałaban's vacuum bounds.** For a Higgs mass $M > 0$ let

$$Z_K(T; M) = \int dA\, d\varphi\, \exp[-S^\varepsilon(T, A, \varphi; M)],$$

where S^ε is King's periodic action [24, (2.4), p. 652] on $\Omega = T_\varepsilon$ with $m^2 = M^2$, photon mass $\mu_0 = \mu_*$, the mass counterterm $\delta m^2_{K,T}(M)$ and the vacuum constant $E_0(M) + E_1(M)$ of [24, (2.6), (3.35)] (Subsection 4.5). By Proposition 4.7 this is the action of [1, (1.11)].

Theorem 4.1 (lower vacuum bound; [1, Thm. (1.14), p. 606])**.** *Let $\mu_* > 0$, $\lambda > 0$, $M > 0$, the vector field in the Feynman gauge, and the charge as in* [1, p. 605]*. There is $E_-(M) < \infty$, independent of ε and of the torus, with $Z_K(T;M) \geq \mathrm{e}^{-E_-(M)|T|}$.*

In this paper Theorem 4.1 is used at $M \in \{m, m_1, m_2\}$.

Theorem 4.2 (upper vacuum bound; [2, Thm. (1.3), p. 556])**.** *Under the hypotheses of Theorem 4.1 there is $E_+(M) < \infty$, independent of ε and of the torus, with $Z_K(T;M) \leq \mathrm{e}^{E_+(M)|T|}$.*

Remark 4.4 describes how the estimates that its printed proof leaves out are supplied in this paper. Theorem 4.2 is used at $M_1 = 0.7m$ (on the way to the tube bound, through Proposition 6.4) and at m_1, m_2.

Lemma 4.3 (the hypotheses of Theorems 4.1 and 4.2 hold for Z_K)**.** *For every $M > 0$, every torus T_j and every spacing ε, $Z_K(T;M)$ is Bałaban's Feynman-gauge integral* [1, (1.10)] *with photon mass $\mu_0 = \mu_* > 0$, and its charge matrix is $q = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, the case "$N = 2$ and $q^2 = -1$" of* [1, p. 605]*. In particular Theorems 4.1 and 4.2 apply to $Z_K(T;M)$.*

Proof. (a) *Feynman gauge.* In [1, p. 605] the Feynman gauge is introduced by replacing the integral (1.9), with the action (1.8), by the integral (1.10), with the action (1.11); the vector-field part of (1.11) is $\frac{1}{2}\langle A, (-\Delta^\varepsilon + \mu_0^2)A\rangle$, with $-\Delta^\varepsilon$ the Laplace operator on the torus T_ε. The photon part of the action of $Z_K(T;M)$ is King's Feynman form [24, (2.4), p. 652] at $\mu_0 = \mu_*$, and by Proposition 4.7 the whole action is that of [1, (1.11)]. So $Z_K(T;M)$ is Bałaban's Feynman-gauge integral [1, (1.10)] with $\mu_0 = \mu_* > 0$.

(b) *Charge.* [1, p. 605] writes $U(A) = \exp(q\varepsilon eA)$ with q an antisymmetric $N \times N$ matrix and $\|q\| \leq 1$. Here $\varphi(x) \in \mathbb{C} \cong \mathbb{R}^2$, so $N = 2$, and the covariant difference in (2.2) multiplies by $\mathrm{e}^{ie\varepsilon A_{x,\mu}}$, which acts on $\mathbb{R}^2$ as the rotation by the angle $e\varepsilon A_{x,\mu}$. So $q = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, which is antisymmetric with $\|q\| = 1$ and $q^2 = -1$. This is the case "$N = 2$ and $q^2 = -1$" of [1, p. 605], in which δm^2 is a number. □

Remark 4.4 (on the printed proof of Theorem 4.2). The abstract of [2] (p. 555) states that the proof is completed there with the exception of some technical estimates, and [3, p. 411] supplies them through [3, Prop. 1]. Some estimates of [2, 3] are needed by the proofs of Theorems 4.2 and 5.2 but are not written out in print. They are the statements $\mathcal{I}$ of Definition I.20. They are proved, at the two points $(u, M) \in \{(1, m_2), (-\eta, m_1)\}$ of Theorem 5.2, in Lemma I.23 and the results it points to. These proofs use statements collected in Appendix K, whose proofs are not included in this version; those used directly in Lemma I.23 are the margins of the two-domain bounds at the first step (Statement K.62), the identification of the bad vacuum localisations at the last scale (Statement K.63) and the uniformity in ε_0 of Bałaban's per-step constants (Statement K.1). Lemma I.23 and the results below are proved under the statements of Appendix K. In detail:

- the depth at which each sketched step of [3, Prop. 1] is applied: Proposition 5.8, Lemma 5.9 and Corollary 5.10;
- [2, Lemma 2.4] at edge points: Lemma 5.6, under the cap (5.7);
- the comparison of torus and infinite-lattice propagators: Lemma 5.22;
- the combinatorial estimate [2, (3.42)–(3.54), pp. 592–594], which ends the proof of Theorem 4.2: Proposition 5.21;
- the ingredients of Table 1: Propositions E.64, E.67, E.110, E.71, E.76, E.159, E.162 and E.175;
- the statements that make up Proposition E.110, listed in Table 10: Subsection E.7, Lemmas E.100, E.102, E.103, E.24, E.32 and Proposition E.36;
- the order cap $\bar{N} = 169$ for [3, (1.33)]: Proposition E.76;

- the final step [2, (2.116)–(2.118), p. 582] of the upper bound: Lemma I.21, with its hypothesis (F) supplied by Proposition F.2.

The transfer of these estimates to the points $(u, M) = (0, M)$, $M \in \{M_1, m_1, m_2\}$, at which Theorem 4.2 is used, is Lemma I.24. The point $u = 0$ is Bałaban's own model, on which these estimates carry no u-weight. Each estimate depends on the Higgs mass in one of two ways. Either the mass enters only from above, through $M^2 t^2 \le c$ (imposed at $m_2 \ge M$). Or it enters through powers of M of either sign and through logarithms such as $\log_+(1/(M\varepsilon_0))$, in quantities that are finite at every fixed $M > 0$. These quantities enter only three things: downward-closed upper bounds on ε_0, constants evaluated at the four masses M_1, m, m_1, m_2, and the lower bound $\ell_0 \ge 128/M_1$ on the torus. None of the estimates orders M against μ_* or uses $m_1 > 2m$. The estimate of Proposition 5.21 is combinatorial: it does not involve the fields or u. Four ingredients of these estimates hold at every $M \in (0, m_2]$ (Lemma A.6): the torus-uniform final c-numbers, the $A \cdot A$ vertices, the tail beyond order $\bar{n}$ and the lower bound on the averaged mass operator.

4.3. **Bałaban's propagator bounds.** Let $\Omega = B^k(\Omega^{(k)})$, where $\Omega^{(k)}$ is a union of big blocks of the rescaled torus, let A be a vector field regular on Ω in the sense of [1, (2.23)], and let $G_k(\Omega, A)$ be the Green's function of the operator $-\Delta_{A,\Omega}^{\eta_k, N} + m^2 (L^k \varepsilon)^2 + a_k P_k(A)$ [1, (2.22), p. 610] (Neumann conditions on $\partial\Omega$, $P_k = Q_k^* Q_k$ the block averaging, [1, (2.11), p. 609]).

Proposition 4.5 ([1, Props. 2.1–2.3, pp. 610–611], proved in [4])**.** *If the running charge $e(L^k\varepsilon) := e\,(L^k\varepsilon)^{(4-d)/2}$ is sufficiently small and the Hölder exponent α of* [1, (2.24)] *satisfies $\alpha < 1$, there are $\delta_0, c_0, R_0 > 0$, depending only on d, a and M_b (c_0 also on α; independent of A, k, Ω and ε), such that:*

(2.24)–(2.25) *$|(G_k(\Omega, A)f)(x)|$ and $|(D^{\eta_k} G_k(\Omega, A)f)(x)|$, and the Hölder quotients of* [1, (2.24)]*, are bounded by $c_0 \mathrm{e}^{-\delta_0 \operatorname{dist}(x, \operatorname{supp} f)} \|f\|_\infty$ for $x \in \Omega$ with $\operatorname{dist}(x, \Omega^c) \ge R_0$ (for (2.24): both points at distance $\ge R_0$ from Ω^c);*

(2.26) *for $\Omega \subset \Omega_0$, $\delta G_k(\Omega, \Omega_0, A) := G_k(\Omega, A) - G_k(\Omega_0, A)$ obeys the same bounds with the additional factor $\exp[-\delta_0 \operatorname{dist}(\operatorname{supp} f, \Omega^c) - \delta_0 \operatorname{dist}(\{x, x'\}, \Omega^c)]$, Ω^c the complement of the smaller set;*

(p. 611) *if Ω is a rectangular parallelepiped, the bounds hold for all $x, x' \in \Omega$, without the restriction $\operatorname{dist} \ge R_0$;*

(2.34) *the block covariances $C_\Lambda^{(k)}(\Omega, A)$ obey exponential bounds without restriction.*

The operators are positive uniformly in the lattice spacing, $-\Delta_{A,\Omega}^{\eta_k, N} + a P_k(A) \ge \gamma_0$ [4, (1.8), p. 573]*, and their Green's functions decay exponentially* [4, Cor. 2.3, (2.30), p. 580].

Two features of Proposition 4.5 are used repeatedly below. The restriction R_0 concerns the evaluation points only: in [1, (2.24)–(2.25), p. 610] the condition $\operatorname{dist}(\cdot, \Omega^c) \ge R_0$ is imposed on x (and x'), and the support of f may touch $\partial\Omega$. And the constants are uniform over all unions of big blocks ("independent of A, k, Ω", [1, p. 610]). The constants also do not depend on the side of the torus:

Lemma 4.6 (independence of the torus side)**.** *Let T_{η_k} be a lattice torus whose side in direction μ, in scale-k units, is $n_\mu M_b$ with $n_\mu \in \mathbb{N}$. Then Proposition 4.5 and* [4, (1.8)] *hold on T_{η_k} with the constants $\gamma_0, \delta_0, c_0, R_0$ of* [4]*. These constants do not depend on (n_μ). Distances are those of* [1, (1.3)].

Proof. Case $\min_\mu n_\mu \ge 3$. We follow the proof in [4, Sect. 2, pp. 574–586]. Let $J = \prod_\mu \mathbb{Z}/n_\mu\mathbb{Z}$ index the big-block vertices $M_b j$, and let $C_j := M_b j + [-M_b, M_b]^d$.

(1) Since $2M_b < n_\mu M_b$, the projection from $\eta_k \mathbb{Z}^d$ to T_{η_k} is injective on C_j. It preserves bonds, unit blocks and the contours of Q_k inside blocks. So $\Pi_j = \Omega \cap C_j$ is a union of at most 2^d big blocks in an embedded cube, and the constant-A_0 gauge $\lambda_0(x) = A_0 \cdot (x - M_b j)$ is defined on C_j. Hence

[4, Lemmas 2.1, 2.2] apply to $G_k(\Pi_j, A_j)$ with their constants, which depend on d, M_b, α, c only. The same holds for [4, Lemma 2.4], which is proved on parallelepipeds of $\xi\mathbb{Z}^d$ [4, pp. 584–586]. The field A_j is built from $A|_{C_j}$ alone.

(2) Let $h \in C_0^\infty((-\frac{2}{3}, \frac{2}{3}))$ with $h = 1$ on $[-\frac{1}{3}, \frac{1}{3}]$ and $\sum_{j\in\mathbb{Z}} h(\cdot - j)^2 = 1$, and put $h_j(x) = \prod_\mu h(x_\mu/M_b - j_\mu)$, $j \in J$ [4, p. 575]. Since the sides are multiples of M_b, the identity $\sum_{j\in J} h_j^2 = 1$ holds on T_{η_k}, and $\operatorname{supp} h_j \subset C_j$. Hence [4, (2.2)–(2.9)] hold as on $\eta_k\mathbb{Z}^d$. If $\Omega = T_{\eta_k}$, then $\partial\Omega = \emptyset$ and [4, (2.6)] is trivial.

(3) For each $j \in J$ we have $\#\{j' : |j - j'| \le 1\} \le 3^d$, and each point lies in at most 2^d of the sets $\operatorname{supp} h_j$. Hence $\|R\|_{2,2} \le c_d\, O(M_b^{-1}) < 1$ as in [4, p. 577], and the bound $2^d(3^d)^{n-1}2^d$ on the number of walks of length n [4, p. 579] holds unchanged.

(4) The metric [1, (1.3)] is the quotient metric: $\operatorname{dist}_T(x, y) = \min_\gamma |\tilde{x} - \tilde{y} - \gamma|_\infty$, the minimum taken over the deck translations γ. Consecutive cubes of a walk have centres at distance at most M_b. So $\operatorname{dist}_T(\{x, x'\}, \operatorname{supp} f) \le (n+2)M_b$, i.e. $n \ge M_b^{-1} \operatorname{dist}_T - 2$, and the sum [4, (2.22)] gives δ_0, c_0 as on $\eta_k\mathbb{Z}^d$. Two further steps use only (4): the definition [4, (2.19)] of R_0, and the implication "$\operatorname{dist}(\{x, x'\}, \Omega^c) \ge R_0$ implies that the first n_0 cubes of the walk are contained in Ω". The cancellation of the interior-cube terms for $\delta G_k(\Omega, \Omega_0, A)$ uses only (1)–(2). If $\Omega = T_{\eta_k}$, every $\Pi_j = C_j$ is a cube, so no restriction on x, x' is needed.

(5) For [4, (1.8)]: $P_k(A)$ is block diagonal, so dropping the bonds between unit blocks gives $\langle\varphi, (-\Delta_{A,\Omega}^N + m^2(L^k\varepsilon)^2 + a_k P_k(A))\varphi\rangle \ge \sum_\Delta \langle\varphi, (-\Delta_{A,\Delta}^N + a_k P_k(A))\varphi\rangle_\Delta$, a sum over unit blocks Δ. On each unit block, the gauge of (1) and the expansion in A' reduce the bound to $A = 0$, as in [4, (2.26′)–(2.27), p. 580]. There P_k is the orthogonal projection onto the constants. The Neumann gap of a block with $N \ge 2$ sites per side is $4N^2 \sin^2(\pi/2N) \ge 8$. So the lower bound is $\min\{8, a_k\} - o(1)$, and it does not involve the torus.

(6) The block covariances [1, (2.34)], i.e. [1, Prop. 2.3], are derived in [4, Sect. 3, pp. 586–589] from two ingredients. The first is [4, Cor. 2.3], which is covered by (1)–(4); these use only the L^2 bound of [4, Lemma 2.1]. The second is a family of Gaussian integrals that factorise over the blocks $B^{k+1}(z)$. These are local, and their block lower bound is (5) with L-blocks. Hence [1, (2.34)] holds on T_{η_k} with constants independent of (n_μ).

(7) Let Ω be a rectangular parallelepiped that does not wrap around T_{η_k}. Then every Π_j is a parallelepiped in an embedded cube, so [4, Lemma 2.2] applies to every factor of the walk expansion of [4, p. 579]. The bounds then hold for all $x, x' \in \Omega$, as in [4, p. 579].

No other constant in [4, Sect. 2] involves the global geometry.

Case $\min_\mu n_\mu \le 2$ *(covering step).* Let $\tilde{T}$ be the torus of sides $3n_\mu M_b$, and let $\pi : \tilde{T} \to T$ be the covering map, with deck group $\Gamma \cong (\mathbb{Z}/3\mathbb{Z})^d$. Put $\tilde{\Omega} := \pi^{-1}\Omega$, a union of big blocks, and $\tilde{A} := A \circ \pi$, which is regular with the same constants. Choose the Q_k-contours of $\tilde{T}$ as the lifts of those of T. Then $\tilde{H} := -\Delta_{\tilde{A},\tilde{\Omega}}^N + m^2(L^k\varepsilon)^2 + a_k P_k(\tilde{A})$ satisfies $\tilde{H}(g \circ \pi) = (Hg) \circ \pi$, and $\tilde{H}$ is invertible by the first case. Hence $G_k(\tilde{\Omega}, \tilde{A})(f \circ \pi) = (G_k(\Omega, A)f) \circ \pi$. The same holds for D^{η_k}, for the Hölder quotients (lift x' next to $\tilde{x}$; the shortest contours lift) and for δG_k. By (4), $\operatorname{dist}_{\tilde{T}}(\tilde{x}, \pi^{-1}S) = \operatorname{dist}_T(x, S)$ for every set S, and $\|f \circ \pi\|_\infty = \|f\|_\infty$. So the bounds on $\tilde{T}$, whose sides are at least $3M_b$, transfer to T with the same constants. Positivity on T is positivity on the Γ-invariant subspace. □

On the tori T_j at the steps $k \le K$, Lemma 2.1(ii) gives $n_\mu = j!\,(\ell_0/M_b)L^{K+n_0-k} \ge 2L^{n_0} \ge 4$ big blocks per side, so only the first case of the proof is needed there. Lemma 4.6 is used in step S19 of the proof of Proposition 5.8, for an auxiliary torus $T' \supset \square$ whose side lies in $2M_b\mathbb{N}$, is at least $\max\{3M_b, 3r_\square + 2s_b + 2\}$, and has a big-block grid compatible with $\square$. It also makes the constants of every application of Proposition 4.5 on $T_{j,\varepsilon}$ the same for all j.

4.4. **Bałaban's Proposition 1 of Higgs III.** In the k-th step of [2] the torus is replaced by $\Omega := B^k(\Lambda_2^{(k-1)\prime})$, a union of big blocks, and one sets $\Omega_1 := B^k(\Lambda_7^{(k-1)\prime})$; a second domain $\Omega_2 \subset \Omega$ "satisfying the same conditions as Ω" is allowed [3, pp. 411–414]. The background fields $\tilde{A}, \tilde{B}$ satisfy $|\tilde{A}|, |\partial^{\eta_k}\tilde{A}|, |\partial^{\eta_k}\tilde{B}| \le O(1)p(t_k)$ with the corresponding Hölder bounds, and the support hypothesis

$$\text{(R4)} \qquad \operatorname{dist}(\operatorname{supp}\tilde{A}, \partial\Omega) > 2r(t_k) \tag{4.2}$$

[3, p. 412]. [3, Prop. 1, pp. 420–421] bounds the localised renormalised perturbative expressions $E(G_{\mathrm{ren}}, \{\Box(v)\}, \Phi_{\mathrm{ext}}, A_{\mathrm{ext}})$ of [3, (1.33)], with constants "independent of ε, k, the domains $\Omega, \Omega_1, \Omega_2$, the vector field $\tilde{B}$" [3, p. 421]. Its proof occupies [3, pp. 422–445] and contains sketched steps; Proposition 5.8 writes each of them out and determines how far from $\partial\Omega$ it is applied.

4.5. **King's counterterms and their identification with Bałaban's.** For a Higgs mass M and a torus T, King's mass counterterm on $\Omega = T$ at the reference photon mass μ_* is a finite sum of lattice graphs of orders λ, e^2, λ^2, $e^2\lambda$, e^4 [24, (3.32)–(3.33)] (Definition 2.2); we write

$$a_{K,T}(M) := M^2 + \delta m^2_{K,T}(M)$$

for the resulting bare quadratic coefficient, and $E_0(M) + E_1(M)$ for King's vacuum constant [24, (2.6), (3.35)].

Proposition 4.7 (King's counterterm and vacuum constant are Bałaban's)**.** *$a_{K,T}(M)$ is the coefficient at which Bałaban's theorems are stated on T_ε (*[3, p. 417], [1, (1.11)–(1.13), pp. 605–606]*), and King's vacuum constant equals Bałaban's $E = E_0 + E_1$ at the truncation order $\bar{n} = 13$ of the construction.*

Proof. The statement on the mass counterterm is Lemma C.1; the statement on the vacuum constant is Lemma C.2. □

In the slice models of Section 5 the order-zero vacuum constant is the scalar factor $E_{0,s}(M)$ of [1, (1.12), p. 606] only, because the photon measure there is normalised.

4.6. **Other classical inputs.**

(1) *Current representation and correlation inequalities* [8]: for every positive covariance Γ of the photon and every real single-site source, the matter weight is of positive type in A, so that $Z(\Gamma; a) = \sum_n w(n)\, \mathrm{e}^{-\frac{1}{2}e^2\varepsilon^2\langle n, \Gamma n\rangle}$ with $w(n) \ge 0$, the sum running over divergence-free integer currents n [8, Thm. 4.1(2), p. 262] (Lemma 2.5); the two classes of [8, Sect. 6] move in opposite directions. For a site function $g \ge 0$, $r^2(g) := \sum_x g_x r_x^2$, and a real linear functional $A(f)$ of the bond variables, $\langle \mathrm{e}^{r^2(g)}\mathrm{e}^{iA(f)}\rangle$ is non-increasing and $\langle \mathrm{e}^{-r^2(g)}\mathrm{e}^{A(f)}\rangle$ is non-decreasing in the photon covariance [8, Cor. 6.3(1), p. 281] (on the torus graph, Lemma 7.5); in particular, for $h \ge 0$, $\langle \mathrm{e}^{:\varphi^2:(h)}\rangle$ does not increase and $\langle \mathrm{e}^{-:\varphi^2:(h)}\rangle$ does not decrease when the covariance increases. In the bare coefficient, [8, Thm. 6.1, p. 276] with P, Q polynomials in r with positive coefficients gives $\langle P; r_x^2\rangle \ge 0$, so $\langle P\rangle$ (for instance $\langle r_y^2\rangle$, or $\langle \mathrm{e}^{:\varphi^2:(h)}\rangle$ with $h \ge 0$) does not increase when the coefficient is increased at a site. In the hopping couplings the directions are reversed [8, Thm. 6.2(2),(4), p. 280; Cor. 6.3(2), p. 281]: $\langle \mathrm{e}^{r^2(g)}\mathrm{e}^{iA(f)}\rangle$ does not decrease and $\langle \mathrm{e}^{-r^2(g)}\mathrm{e}^{A(f)}\rangle$ does not increase when a coupling is increased.

(2) *Chessboard estimate* for reflection-positive lattice measures on a torus with an even number of sites per side, for reflections in the planes halfway between lattice planes [17, Sect. 4]; in the lattice-gauge setting it is [8, Thm. 2.10, p. 252] and [34, Thm. 2.2, p. 16], both derived from reflection positivity alone. On the tori T_j it rests on lattice reflection positivity about such planes, Proposition 7.9 with the plane pairs of case (ii) of Lemma 7.8.

(3) *Lattice power counting* [32, Thm. 1, p. 90] and lattice renormalisation [33]; both are stated for four-dimensional lattices, and [32, p. 86] notes that the arguments extend to other dimensions. The three-dimensional form is Proposition C.3; it is used in the proof of Proposition 6.3.

Sections 4–6 use neither King's cube states [25, Thms. 2.1–2.3], nor any statement on the infinite lattice $\varepsilon\mathbb{Z}^3$ with boundary conditions, nor any infrared renormalisation group; in particular, no bound uniform in the photon mass in a fixed cube is used.

5. The vacuum bound for the slice models, and three repairs of printed proofs

5.1. **The slice models and the statement.** Fix $j \geq 1$ and a spacing $\varepsilon = 2^{-K}\varepsilon_0$, and write $T := T_j$. Apart from the statement of Theorem 5.2, the torus is fixed in this section, and the letter j indexes the fluctuation scales (slices) H_j, δ_j, $G^{\eta_k}_{(j)}$; it is not the torus index. Let $-\Delta_\varepsilon$ act on each component of the vector field (on the periodic lattice it is the Hodge Laplacian $d^*d + dd^*$, Lemma B.4), and put $C_\mu := (-\Delta_\varepsilon + \mu^2)^{-1}$. Let $G_k := (-\Delta_\varepsilon + \mu_*^2 + a_k t_k^{-2} P_k)^{-1}$ be King's block-spin covariances [24, (2.13), p. 653] (vector version), and let $H_j := G_{j+1} - G_j$ denote King's fluctuation covariances [24, (2.17)], with $G_0 := 0$ (so $H_0 = G_1$, King's $C^{(0)}$ [24, (2.16), p. 653]); $H_j \geq 0$ by Lemma E.1. Put

$$\Gamma^B := \sum_{j<K} \delta_j H_j, \qquad \delta_j := \kappa\mu_*^2 t_{j+1}^2 \leq \bar{\delta} := \kappa\mu_*^2\varepsilon_0^2. \tag{5.1}$$

Here $\kappa > 0$ is the constant of the domination inequality $D^< \leq \Gamma^B\Pi^\perp$ on the torus, with $D^<$ the operator of Remark 5.1 (Theorem I.16 and Lemma I.2(b)); the inequality holds for every $\kappa \geq \kappa_1\kappa_3$, where $\kappa_1\kappa_3$ is the constant of Theorem I.16. At $L = 2$, $a = 1$ and under the cap $\mu_*\varepsilon_0 \leq 0.2$ (condition (E.dom) of Appendix A) one has $\kappa_1\kappa_3 < 7.5$ for every K (Remark I.18), and $\kappa \leq 7.5$ is fixed with the regulator data, independently of K. Since $\sum_{j<K} t_{j+1}^2 H_j \leq \varepsilon_0^2 G_K$, also $\Gamma^B \leq \kappa\mu_*^2\varepsilon_0^2\, G_K \leq \bar{\delta}\, C_{\mu_*}$, for every $\kappa > 0$ (Lemma E.2). For $u > -1/\bar{\delta}$ the operator $C_B(u) := C_{\mu_*} + u\Gamma^B$ is positive (Lemma E.4), and we call the lattice model with photon covariance $C_B(u)$, scalar coefficient $a_{K,T}(M)$, King's hopping and quartic term, the *slice model* at (u, M). Put

$$\Phi_M(u) := \log Z\big(C_B(u); a_{K,T}(M)\big),$$

$$Z(\Gamma; a) := \mathbb{E}_{N(0,\Gamma)}\big[Z_\varphi(A; a)\big], \qquad Z_\varphi(A; a) := \int \mathrm{e}^{-S^{\mathrm{m}}_a(A,\varphi)} \prod_x d^2\varphi(x),$$

with S^{m}_a the matter action (2.2): the photon measure is normalised and the scalar integral is not. Let $\mathcal{E}(u; M) := T_{\bar{n}}[\Phi_M(u)]$ be the Taylor polynomial of total order $\bar{n} = 13$ in (e, λ) at $(0, 0)$, at fixed M and with the coefficient $a_{K,T}(M)$ inside; this coefficient is computed with the photon covariance C_{μ_*} and does not depend on u (the organisation of [1, (1.12)–(1.13)]). The order-zero term of $\mathcal{E}(u; M)$ is $E_{0,s}(M)$. Set $e(\varepsilon; u; M) := -|T|^{-1}\mathcal{E}(u; M)$.

Remark 5.1 (the direction of the deformation). The translation-invariant direction $D^< := \int_0^{\varepsilon_0^2} \mathrm{e}^{t\Delta_\varepsilon}(1 - \mathrm{e}^{-t\mu_*^2})\, dt\, \Pi^\perp$ satisfies $D^< \leq \Gamma^B\Pi^\perp$ (Lemma I.2(b)), and not conversely. An upper bound on the vacuum in one direction does not imply one in the other; Theorem 5.2 is stated and used with Γ^B.

The function $u \mapsto \Phi_M(u)$ is convex on $(-1/\bar{\delta}, \infty)$: by the current representation (Subsection 4.6, item 1), $\Phi_M(u) = \log \sum_n w(n) \mathrm{e}^{-\frac{1}{2}e^2\varepsilon^2\langle n, C_{\mu_*} n\rangle} \mathrm{e}^{-\frac{u}{2}e^2\varepsilon^2\langle n, \Gamma^B n\rangle}$ with $w(n) \geq 0$, a logarithmic Laplace transform. The polynomial $e(\varepsilon; u; M)$ has degree at most $\lfloor \bar{n}/2 \rfloor$ in u (Lemma E.5).

Recall the parameter window (1.3),

$$\mathfrak{W} = \{(e, \lambda, m, \eta, m_1, m_2) :\ m > 0,\ 0 < \lambda < 0.1m,\ 0 < e^2 < 0.1m,\ 0 < \eta < 1,\ 2m < m_1 < m_2\}.$$

Theorem 5.2 (the vacuum bound for the slice models)**.** *Let $(e,\lambda,m,\eta,m_1,m_2)\in\mathfrak{W}$ and $\mu_*:=m$, and let the regulator data ($L=2$, $p=6$, $\bar n=13$, $\bar N=169$, M_b, R, b_0, ε_0, ℓ_0) satisfy the conditions of Appendix A. Then there are constants $C_+^B(1;m_2), C_+^B(-\eta;m_1)<\infty$ such that, for every $K\ge K_0$ and every $j\ge 1$, with $T=T_j$ and $\varepsilon=2^{-K}\varepsilon_0$,*

$$|T|^{-1}\Phi_M(u)+e(\varepsilon;u;M)\le C_+^B(u;M)\qquad \text{at } (u,M)\in\{(1,m_2),(-\eta,m_1)\}. \tag{5.2}$$

The proof of Theorem 5.2 is Proposition F.2, which assembles (5.2) from the ingredients of Table 1; its structure is described after Lemma 5.4.

Remark 5.3 (conventions in Theorem 5.2). (i) The order-zero term of $\mathcal{E}(u;M)$ is the scalar factor $E_{0,s}(M)$, because the photon measure of the slice models is normalised. With the Lebesgue-normalised photon of Propositions 6.3 and 6.4 it is King's full constant. The two differ by a term that does not depend on M (Lemma C.2(c)), and it cancels in all differences used. (ii) As in [3, (1.30), p. 419], Bałaban's final and partial vacuum counterterms are global c-numbers: they carry no localisation function and do not depend on the domains. (iii) The support hypothesis is used in the form (R4‴) of Corollary 5.10, whose proof covers the components anchored at Ω_1 as well as those anchored at $\operatorname{supp}\tilde A$. The argument of that proof needs $L\le 3$: the vertices anchored at Ω_1 have depth about $5\rho_k\approx 5r/L$, which exceeds the depth $\frac{3}{2}r$ needed by the printed step P12 of Subsection 5.3 only for $L\le 3$; at $L\ge 4$ it does not apply.

Lemma 5.4 (two consequences of the conditions)**.** *Let the regulator data satisfy the conditions of Appendix A at $L=2$, and let $0<\eta<1$. Then* (a) *$\eta\bar\delta\le\eta/12<1$, and* (b) *$\varepsilon_0\le\min(1/m_2,1/\mu_*,L)$.*

Proof. (a) Condition (E.δ) of stage 10a of Appendix A is $\bar\delta=\kappa\mu_*^2\varepsilon_0^2\le 1/12$; multiply by $\eta<1$. (b) Condition (E.m2), $m_2^2\varepsilon_0^2\le 1$, gives $\varepsilon_0\le 1/m_2$; condition (E.dom), $\mu_*\varepsilon_0\le 0.2$, gives $\varepsilon_0\le 1/\mu_*$; and $\varepsilon_0\le 1<2=L$ (stage 10a of Appendix A). □

Structure of the proof of Theorem 5.2. The bound (5.2) is obtained by running Bałaban's inductive upper-bound argument of [2, Sects. 2–3] on the torus for the slice model, keeping track of the dependence on u: the weights $1+u\delta_j$ act on the photon fluctuations only, and the scalar side carries no u-weight. The argument uses the nine ingredients of Table 1, together with Lemma 5.6, Proposition 5.8, Corollary 5.10 and Lemma 5.22. One of the ingredients, Proposition 5.21, completes Bałaban's combinatorial estimate [2, (3.42)–(3.54), pp. 592–594] (Subsection 5.4). The bounds of all ingredients are uniform in ε, K and ℓ_T, and finite at $u=-\eta$ and $M=m_1$.

5.2. **Repair 1: Bałaban's Lemma 2.4 of Higgs II at the edge of Λ_7.** We work in the k-th step of [2, Chap. 2], in scale-k units, on the rescaled torus. The fields satisfy the small-field restrictions [2, (2.55), p. 570] on $\Lambda_{-1}^{(k-1)\prime}$:

$$|\partial A|\le c_1 p(t_{k-1}),\quad |D_{\bar A^{(k)}}\varphi|\le c_1 p(t_{k-1}),\quad |\varphi(x)|\le \Upsilon_k:=c_1\big(\lambda t_{k-1}\big)^{-1/4}p(t_{k-1}),$$

(and the bound on A itself), and the background scalar field of the step is

$$\varphi^{(k)}=a_k G_k\big(B^k(\Lambda_2^{(k-1)\prime}),A^{(k)}\big)Q_k^*(A^{(k)})\,\Lambda_6^{(k-1)\prime}\varphi \tag{5.3}$$

[2, (2.56)]: Green's function on $\Omega:=B^k(\Lambda_2^{(k-1)\prime})$, source cut to $\Lambda_6^{(k-1)\prime}$. In this subsection Λ_i stands for $\Lambda_i^{(k-1)\prime}$, $r_k:=r(t_k)$, and $c_d:=\sqrt{d}\,M_b$; the letter κ denotes Bałaban's exponent in the error $O(t_k^\kappa)$ of [2, (2.67)], not the constant κ of Subsection 5.1. From the nesting [2, (2.8)], $\operatorname{dist}(\Lambda_7,\Lambda_6^c)\ge\rho_k$, $\operatorname{dist}(\Lambda_6,\Lambda_2^c)\ge 4\rho_k$ and $\operatorname{dist}(\Lambda_7,\Lambda_2^c)\ge 5\rho_k$.

ingredient	content
Proposition E.64	the u-dependent part of the one-loop vacuum family at every scale $k \geq 1$
Proposition E.67	the Ward–Takahashi identities on the torus at orders ≥ 2, through the periodisation of the propagators
Proposition E.110	the conditions and operations of [3, Prop. 1] as they are used in [2], including the $\Box$-remainder of [3, p. 433], the separation step, the two-domain bounds for δG_k and the support hypothesis in the form (R4‴)
Proposition E.71	in the bounds near Neumann faces, only real background fields are used
Proposition E.76	the order cap $\bar{N} = 169$ for the expressions [3, (1.33)], and the tails of [2, pp. 579, 582]
Proposition 5.21	the combinatorial estimate [2, (3.42)–(3.54), pp. 592–594]: the step to (3.47) with the region entropy, the cube count (3.45) near the boundary, and the collar and shell sets of (3.42); it is also used in the proof of Theorem 4.2
Proposition E.159	the hypothesis $m^2 \leq O(1)$ of [2, p. 589], read as a bound on the rescaled mass; condition (E.md)
Proposition E.162	torus-uniform bounds on the final c-numbers $\delta m^2_{\mathrm{fin}}$, E_{fin}, under $\varepsilon_0 \leq \min(1/m_2, 1/\mu_*, L)$ (Lemma 5.4(b))
Proposition E.175	the vacuum classes at the last scale

TABLE 1. The ingredients of the proof of Theorem 5.2. All their bounds are uniform in ε, K and ℓ_T, and finite at $u = -\eta$ and $M = m_1$.

[2, Lemma 2.4, p. 572] states: under [2, (2.55)],

(5.4)
$$\varphi^{(k)}(x) = U\big(A^{(k)}(\Gamma^{(k)}_{x,y})\big)\varphi(y) + O(p(t_k)) = (Q_k^*(A^{(k)})\varphi)(x) + O(p(t_k)), \qquad x \in B^k(y),\ y \in \Lambda_7,$$

(5.5)
$$(D^{\eta_k}_{A^{(k)}}\varphi^{(k)})(b) = O(p(t_k)), \qquad b \subset B^k(\Lambda_7).$$

Its only consumer in [2] is the quartic convergence factor [2, (2.78)–(2.79), p. 574], a sum over $y \in \Lambda_7^{(k-1)\prime} \cap \Lambda_7^{(k)c}$; this set contains every point of Λ_7 within $2r_k$ of Λ_7^c.

Remark 5.5 (the printed proof does not cover edge points). The printed proof defines $\Box_1, \Box_2$ as the unions of big blocks *contained in* Λ_7 at distance less than $2r_k$, $4r_k$ from y, puts $\Box := B^k(\Box_2)$, and asserts [2, (2.67)]: $\varphi^{(k)}(x) = a_k(G_k(\Box, A^{(k)})Q_k^*\Box_1\varphi)(x) + O(t_k^\kappa)$ by [2, Prop. 2.2] and (5.3). For $y \in \Lambda_7$ at distance $O(1)$ from Λ_7^c this step is not justified as written. (i) The dropped source $\Lambda_6\varphi - \Box_1\varphi$ contains φ on $\Lambda_6 \setminus \Lambda_7$ at distance $O(1)$ from x; the kernel $a_k G_k Q_k^*$ is $O(1)$ at distance $O(1)$ (it reproduces constants, [2, (2.75)]), so this error is $O(\Upsilon_k)$, not $O(t_k^\kappa)$. (ii) The domain change from Ω to $\Box$ happens at distance $O(1)$ from x, so the factor of [1, (2.26)] gives no smallness. (iii) x may lie within R_0 of $\partial\Box$, where [1, (2.24)–(2.25)] are not available unless $\Box$ is a parallelepiped, and $\Lambda_7 \cap$ (cube) has re-entrant edges in general. The two errors in (i) and (ii) cancel to $O(p)$ for slowly varying φ, so the statement (5.4)–(5.5) of the lemma is not contradicted; it is the step [2, (2.67)] of the printed proof that is not justified for such y. The gap is the same at every $L \geq 2$. A box fitting around every $y \in \Lambda_7$ without any cut does not exist: a full $4r_k$-box inside Λ_2 would need $4r_k \leq 5\rho_k \approx 5r_{k-1}/L$, which is false for small ε_0, since $r_{k-1}/r_k \to 1$.

The printed proof of [2, Lemma 2.4, p. 572] does not cover the points of $\Lambda_7^{(k-1)\prime}$ near its boundary (Remark 5.5); Lemma 5.6 gives a proof that does.

Lemma 5.6 (Bałaban's Lemma 2.4 of [2] with a complete proof)**.** *Assume* [2, (2.55)]*, the conditions*

(5.6) *(R.edge)* $\qquad 5\big(r(\varepsilon_0) - M_b\big)/L \geq R_0 + c_d + 2M_b, \qquad 4r(\varepsilon_0) \geq R_0 + c_d, \qquad R \geq 10M_b + 10,$

$\varepsilon_0 \leq \varepsilon(L, \varrho)$ (so that $r(t_{k-1})/r(t_k) \leq 2L$), the torus-size condition (T.nw) of Appendix A, the mass-defect condition (E.md) of stage 10a (used in Step 5 at the Higgs mass M of the step; it is imposed

at m_2 and holds at every $M \le m_2$, since its left sides increase in M), and the upper bound on ε_0

(5.7)

$$\textit{(E.edge)} \qquad C\,a_k\,c_1(\lambda t)^{-1/4}p(t)\,\big(1{+}r(t)\big)^{d-1}\exp\Big(-\delta_0\Big[\frac{r(t)-M_b}{L}-c_d\Big]\Big)\le (Lt)^{\kappa} \quad \textit{for all } t\le \varepsilon_0/L,$$

with $C = C(c_0, \delta_0, d)$. Then (5.4) and (5.5) hold for every $y \in \Lambda_7^{(k-1)\prime}$, every $x \in B^k(y)$ and every bond $b \subset B^k(\Lambda_7^{(k-1)\prime})$, with constants independent of ε, k, the torus and the position of y relative to $\partial\Lambda_7^{(k-1)\prime}$. The condition (E.edge) holds for all small ε_0 at every $R > 0$.

Proof. Fix $y \in \Lambda_7$ and $x \in B^k(y)$. Put $s_k := \min\{4r_k,\, 5\rho_k - 2M_b\}$, let $\square_2$ be the cube of big blocks of sup-radius s_k around the big block of y, $\square := B^k(\square_2)$, and let $\square_1\varphi := 1_{Q(y,s_k/2)}\,\Lambda_6\varphi$, where $Q(y,s)$ is the sup-ball of radius s. Since $\operatorname{dist}(\Lambda_7, \Lambda_2^c) \ge 5\rho_k$, $\square \subset \Omega = B^k(\Lambda_2)$; $\square$ is a rectangular parallelepiped and a union of big blocks. By (5.6) and since r decreases in t, $s_k - c_d \ge R_0$ at every k.

Step 1 (source cut). $\Lambda_6\varphi - \square_1\varphi$ is supported at distance $\ge s_k/2 - c_d$ from x, and $\operatorname{dist}(x, \Omega^c) \ge 5\rho_k - c_d \ge R_0$. By [1, (2.25)] on Ω, with the sup bound Υ_k on φ,

$$\big|a_k\big(G_k(\Omega, A^{(k)})Q_k^*(\Lambda_6\varphi - \square_1\varphi)\big)(x)\big| \le C a_k\,\mathrm{e}^{-\delta_0(s_k/2-c_d)}\,\Upsilon_k.$$

Step 2 (domain change). $G_k(\square, A) = G_k(\Omega, A) + \delta G_k(\square, \Omega, A)$, and by [1, (2.26)] the second term applied to $Q_k^*\square_1\varphi$ at x carries the factor $\mathrm{e}^{-\delta_0(\operatorname{dist}(Q(y,s_k/2),\square^c)+\operatorname{dist}(x,\square^c))} \le \mathrm{e}^{-\delta_0(3s_k/2-2c_d)}$. The same two estimates hold for D^{η_k} by [2, Prop. 2.2, (2.58)]. Hence

$$\varphi^{(k)}(x) = a_k\big(G_k(\square, A^{(k)})Q_k^*(A^{(k)})\square_1\varphi\big)(x) + O\big(a_k\Upsilon_k\mathrm{e}^{-\delta_0(s_k/2-c_d)}\big), \tag{5.8}$$

and likewise for the covariant derivative: this is [2, (2.67)] with the boxes chosen above.

Step 3 (constant background). Let $A_0 := A^{(k)}(y)$. Joining points of the convex box $\square$ by straight lattice paths and using [2, (2.60)] on $B^k(\Lambda_2) \supset \square$, $|A^{(k)}(z) - A_0| \le O(p)\sqrt{d}\,s_k \le O(p\,r_k)$ on $\square$. The expansion [1, (3.44)] and [1, Prop. 2.2] on the parallelepiped $\square$ replace $A^{(k)}$ by A_0 with an error $O(t_k^{\kappa_0})$ (Bałaban's exponent in [2, (2.68)]), as in [2, (2.68)].

Step 4 (gauge-out). The constant field is a gradient on the box, $A_0 = \partial^{\eta_k}\lambda_0$ with $\lambda_0(z) = A_0 \cdot (z - y)$, so $U(A_0(\Gamma_{z,y})) = \mathrm{e}^{ieA_0\cdot(z-y)}$ for every contour $\Gamma_{z,y} \subset \square$, and $G_k(\square, \partial\lambda_0) = U(\lambda_0)G_k(\square, 0)U(\lambda_0)^*$, $Q_k(\partial\lambda_0) = U(\lambda_0)_{\text{coarse}}Q_k(0)U(\lambda_0)^*$ (Neumann Laplacian on bonds of $\square$; contours inside blocks contained in $\square$). This is legitimate because $\square$ does not wrap around the torus: its diameter is at most $2s_k + M_b \le 8r_k + M_b$, below the side of the torus by (T.nw). Write $\varphi'(z) := U(\lambda_0)(z)^{-1}\varphi(z)$.

Step 5 (reproduction of constants). By [2, (2.75)], $a_kG_k(\square, 0)Q_k^*1 = 1 - m^2t_k^2G_k(\square, 0)Q_k^*1$. Hence

$$\begin{aligned} a_k\big(G_k(\square,0)Q_k^*\square_1\varphi'\big)(x) - \varphi'(y) &= a_k\big(G_k(\square,0)Q_k^*[\square_1(\varphi' - \varphi'(y))]\big)(x) \\ &\quad - \varphi'(y)\,a_k\big(G_k(\square,0)Q_k^*1_{\square\setminus\square_1}\big)(x) - \varphi'(y)\,m^2t_k^2\big(G_k(\square,0)Q_k^*1\big)(x), \end{aligned}$$

where $1_{\square\setminus\square_1}$ is the indicator of $\square \setminus (Q(y, s_k/2) \cap \Lambda_6)$. The first term is $O(p)$: on the convex cube $Q(y, s_k/2) \subset \Lambda_{-1}$ the restriction $|D\varphi| \le c_1 p$ of [2, (2.55)] gives $|\varphi'(z) - \varphi'(y)| \le O(p)|z - y|$, and the kernel decays exponentially (the same reasoning as in [2, Lemma 2.3]). The second term: $\square \setminus \square_1 \subset (Q(y, s_k) \setminus Q(y, s_k/2)) \cup (Q(y, s_k/2) \setminus \Lambda_6)$ lies at distance $\ge \min\{s_k/2, \rho_k\} - c_d$ from x, so it is $\le c_0\mathrm{e}^{-\delta_0(\min\{s_k/2,\rho_k\}-c_d)}\Upsilon_k$ by Proposition 4.5 on the parallelepiped $\square$. The third term is the mass-defect term of the printed proof; it is $O(p(t_k))$ under (E.md), as in the printed proof.

Step 6 (derivatives). For $b \subset B^k(y)$, b lies at distance $\ge s_k - c_d \ge R_0$ inside $\square$ and $\ge 5\rho_k - c_d$ inside Ω, so the printed derivative bounds apply at b and Steps 1–5 give (5.5) in the form [2, (2.77)]. No re-entrant edge is approached.

Collection of the errors. The inequality $s_k/2 \ge \rho_k$ is equivalent to $\rho_k \le 2r_k$ and $3\rho_k \ge 2M_b$; the first holds because $r(t_{k-1})/r(t_k) \le 2L$, the second because $R \ge 10M_b + 10$. So every error above is at most $Ca_k\Upsilon_k(1 + r_k)^{d-1}\mathrm{e}^{-\delta_0(\rho_k - c_d)}$, the polynomial factor coming from the sums of exponentially

decaying kernels over sources at distance $D = O(r_k)$. With $t = t_{k-1}$ and $\rho_k = (r(t) - M_b)/L$, this is $\le (Lt)^\kappa = t_k^\kappa$ by (E.edge), which is the size of the error in [2, (2.67)]; the rest of the printed proof, [2, (2.69)–(2.77)], now runs on the box $\square$. Finally, the logarithm of the prefactor in (5.7) is $O(\log t^{-1}) + O(\log\log t^{-1})$ while the exponent is $\delta_0 R(1 + \log t^{-1})^\varrho / L$ with $\varrho > 1$, so (5.7) holds for t small, at every $R > 0$. $\square$

Remark 5.7 (where the repair is used). In the proof of Theorem 5.2, Lemma 5.6 is used in the quartic lower bound [2, (2.78)] at points of $\Lambda_7^{(k-1)\prime}$ near its boundary, whenever a large-field region is present. The gauge-out on a full cube used in [3, p. 433] and the later lemma [2, Lemma 2.7] do not use the cut at Λ_7 and are not affected. The condition (E.edge) depends only on L, R and data fixed before them, not on u, M, ε or ℓ_T; it is downward closed in ε_0, and it is a condition of stage 10b of Appendix A. Lemma A.2(a) lists the data on which it depends, and shows that the conditions (R.edge) follow from the lower bound $R \ge R^*$ at $L = 2$ and that $r(t_{k-1})/r(t_k) \le 2L$ holds for every $\varepsilon_0 \le 1$.

5.3. **Repair 2: the sketched steps of Bałaban's Proposition 1 of Higgs III.** We use the setting of Subsection 4.4, in scale-k units, with $r := r(t_k)$, $s_b \le 2M_b$ the rounding of a set to big blocks, $\ell_{\bar N} := 2V(\bar N)$ the side of a cube containing the localisations of a connected class with at most $\bar N$ vertices ([3, p. 432]), and R_0 the constant of Proposition 4.5. The *depth* of a point x is $\operatorname{dist}(x, \Omega^c)$. At [3, p. 433] the proof takes a cube $\square_1$ of size $r_\square$ (printed: $r_\square = r$) containing the localisations of the old vertices of a class and the concentric cube $\square$ of size $3r_\square$, and splits

$$G_k(\Omega, \tilde B) = G_k(\square, \tilde B) + \delta G_k(\Omega, \square, \tilde B) \qquad \text{[or the same with } \Omega_2 \text{ in place of } \Omega\text{]}, \tag{5.9}$$

treating $\delta G_k(\Omega, \square, \tilde B)$ as an external field. We call this printed step P12. Its precondition $\square \subset \Omega$ (respectively $\square \subset \Omega_2$) is not written in [3].

The proof of [3, Prop. 1, pp. 420–445] contains sketched steps; Proposition 5.8 writes each of them out and determines the depth inside the localisation domain at which it is applied.

Proposition 5.8 (the depth of each step of Bałaban's Proposition 1)**.** (a) (Inventory.) *The proof of* [3, Prop. 1], [3, pp. 422–445]*, contains twenty sketched steps, S1–S11 and S13–S21 of Table 2, and the printed step P12 with its unwritten precondition. Besides these, a page-by-page reading of* [3, pp. 414–445] *finds seven further sketched places (the Ω_2 branch of P12; the identification of torus and infinite-lattice propagators in the counterterms; the repetition on $\square$ of S1 and S9 with B' in place of $\tilde A$; three phrases of power counting and bookkeeping; the generation of all Ward–Takahashi identities); none of them uses depth beyond P12 or $R_0 + O(1)$.*

(b) (Anchored steps.) *The steps S1, S2, S3, S4, S9 apply Proposition 4.5 only at points of depth at least $R_0 + 12$ near the localisation of a vertex ($R_0 + 13$ at the first step $k = 1$, where $\eta_k = \frac12$; S2: depth measured in Ω_2), or at points of $U_c(\operatorname{supp}\tilde A)$ of depth $\ge R_0 + 2\sqrt3 + 1$ (S1).*

(c) (The cube step.) *If an old vertex x_v of the class has depth $> \frac32 r_\square + 2s_b + \ell_{\bar N}$, then $\square \subset \Omega$ and every point of $\square_1$ lies at distance $\ge r_\square$ inside $\square$. The same holds with Ω_2 in place of Ω.*

(d) (Depth-free steps.) *Once $\square \subset \Omega$, every identity and inequality used after the split* (5.9)*, on* [3, pp. 433–445]*, is independent of* $\operatorname{dist}(\cdot, \Omega^c)$ *(Lemma 5.9); in particular S5–S8, S10, S11, S13–S21 need no depth.*

Consequently every sketched step of the proof needs depth at most $O(R_0 + \ell_{\bar N} + M_b)$, with a constant independent of r, and the only step needing depth of order r is the printed step P12, with need $\frac32 r_\square + 2s_b + \ell_{\bar N}$.

Proof of Proposition 5.8. (b), S3. The slice decomposition [3, (2.6)] is the iterated composition formula [1, (2.43)], an identity on any union of scale-k blocks, with no depth. For [3, (2.10)–(2.11)],

step	page	omitted argument	depth needed
S1	414	kernel bound of $[G_k(\Omega,\tilde B)V_k]^n G_k(\Omega,\tilde A+\tilde B)[V_kG_k]^{n'}$, [3, (1.16)]	$R_0{+}2\sqrt{3}{+}1$ on $U_c(\operatorname{supp}\tilde A)$; $R_0{+}O(1)$ at vertices
S2	414, 424	$\delta G_k(\Omega,\Omega_2,\tilde B)$ as an external field, [3, (2.5)]	R_0+12 inside Ω_2 (R_0+13 at $k=1$)
S3	424, 426	slice decomposition [3, (2.6)] and slice bounds (2.10)–(2.12)	R_0+1 at line ends
S4	425	summation by parts [3, (2.8)]	R_0+1
S5–S7	428–429	induction (2.16); generalised graphs (Prop. 2.2); power counting, Cor. 2.3	none
S8	431	Ward–Takahashi identities for free boundary conditions as limits of periodic ones	none
S9	432	bound [3, (3.1)] between separated localisations	R_0+5 at vertex cubes
S10, S11	432	regrouping of classes; localisation of a component in a cube of side $\ell_{\tilde N}$	none
P12	433	printed split (5.9); precondition $\square\subset\Omega$ (or Ω_2) unwritten	$\frac32 r_\square+2s_b+\ell_{\tilde N}$ at an old vertex
S13	433	constant field B_0 with $\lvert\tilde B-B_0\rvert\le O(r_\square p)$ on $\square$	as for P12
S14–S16	433	B'-expansion; old vertices in $\square_1$; chains by Prop. 2.1 on $\square$	none beyond $\square\subset\Omega$
S17, S18	433	gauge covariance on $\square$; gauge-out of B_0	none
S19	433	bound on $\delta G_k(\square,\eta_k\mathbb{Z}^d,0)$	none
S20	434	comparison of the norms (1.32) at B_0 and at $\tilde B$	none (a size factor, see Remark 5.11)
S21	434–445	renormalisation of the divergent classes	none

TABLE 2. The sketched places in the proof of [3, Prop. 1]. Pages are journal pages of [3].

rescale to the L^{-j} lattice; Ω is a union of scale-j big blocks and $\tilde B$ is regular at scale j. For a coarse point y_0, $f:=Q_j^*\delta_{y_0}$ has $\|f\|_\infty\le 1$ and support $B^j(y_0)$, so [1, (2.25)] at an evaluation point y gives $\lvert(G_jQ_j^*)(y,y_0)\rvert+\lvert(DG_jQ_j^*)(y,y_0)\rvert\le c_0\mathrm{e}^{-\delta_0\operatorname{dist}(y,B(y_0))}$ provided $\operatorname{dist}(y,\Omega^c)\ge R_0$ in scale-j units, for every y_0, including blocks touching $\partial\Omega$. The adjoint factor is bounded in the same way at the other end, and $C^{(j)}$ by [1, (2.34)] without restriction. The sums over the coarse points converge with constants independent of ε and r. Rescaling back, the restriction is $\operatorname{dist}(\{x,x'\},\Omega^c)\ge R_0L^{j-k}\le R_0$ at the two line ends. The vector propagator lives on the torus [3, p. 414] and needs no depth. Allowing one block for the contour legs of (2.12), the need is R_0+1.

(b), S4. In [3, (2.8)] the derivative is moved by summation by parts; no boundary term arises because the moved leg carries the localisation function g, supported in Ω_1, at distance ≥ 1 from Ω^c. After the move the derivative sits on the vector line (torus, no depth), on g, or on a scalar line end at the vertex: need R_0+1.

(b), S9. [3, (3.1)] bounds $\|hG_k(\Omega,\tilde B)h'\|_{1,\alpha}$ for $\operatorname{dist}(\square(v),\square(v'))\ge 1$. Summing [3, (2.10)–(2.11)] over j with one derivative at each end, the factors $(L^j\eta_k)^{-d-\alpha}\mathrm{e}^{-\delta_1\lvert x-x'\rvert/(L^j\eta_k)}$ are summable because $\lvert x-x'\rvert\ge 1$, and the mixed derivative is pointwise bounded since the points are separated. The evaluation points lie within a collar of width $<\frac12$ around the vertex cubes: need $\le R_0+5$.

(b), S2. The two-variable mixed norm of $\delta G_k(\Omega,\Omega_2,\tilde B)$ in [3, (2.5)], where the two ends may coincide, is bounded by the two-domain theorem, Proposition D.17 (two cut-offs and a weighted ℓ^2 bound for the middle factor), valid at points at distance $\ge R_0+11$ from Ω_2^c and supports at distance $\ge R_0+12$ when $\eta_k\le\frac14$ ($k\ge 2$), and at R_0+12 and R_0+13 at $k=1$, $\eta_1=\frac12$, where the offsets of the cut-offs below one block use up the slack. The factor $\mathrm{e}^{-\delta_0\operatorname{dist}(\Omega_2,\partial\Omega)}$ of [3, (2.5)] is obtained in the reading in which the distance is that of the vertex localisations to Ω_2^c; read literally, as the distance between the sets Ω_2 and $\partial\Omega$, it is not implied, and it fails for localisations near $\partial\Omega_2$ deep inside Ω, where δG_k is of order one. No consumer uses more than a bound of order one. In

the proof of Theorem 5.2 these lines occur only in terms all of whose vertices lie in $\Lambda_7^{(k)}$ [2, p. 580, Prop. 2.6], at depth $\geq 5r - O(1)$ in Ω_2.

(b), S1. The interaction $V_k(\tilde{A}, \tilde{B})$ is supported in $U_c(\operatorname{supp}\tilde{A}) \times U_c(\operatorname{supp}\tilde{A})$, $c \leq 2\sqrt{3}+1$ (nearest-neighbour covariant Laplacian; P_k with contours inside unit blocks). All internal summation points therefore have depth $\geq \operatorname{dist}(\operatorname{supp}\tilde{A}, \Omega^c) - c$. After summation by parts each middle factor is of the form $\mathfrak{D}G_k\mathfrak{D}^*$, whose kernel has a diagonal singularity of order η_k^{-d}, so a pointwise sum is not uniform in η_k. Proposition D.13 proves the bound by an L^2 sandwich with a weighted ℓ^2 estimate valid for any boundary condition (Lemma D.24); it requires depth $\geq R_0 + 2\sqrt{3} + 1$ at the internal points and $R_0 + O(1)$ at the outer points. The power $(e\, t_k^{1-\alpha})^{n+n'}$ of [3, p. 414] is a size bound ($|e\tilde{A}| \leq O(e\,p)$) and needs no depth.

(c). The localisations of the old vertices lie in a cube of side $\ell_{\bar{N}}$ containing x_v (S11), and $\square_1$, $\square$ are concentric cubes of sides $r_\square$, $3r_\square$, rounded outward to big blocks. In the sup metric every point of $\square$ is within $\frac{3}{2}r_\square + s_b$ of the common centre, and the centre is within $\ell_{\bar{N}} + s_b$ of x_v. If the depth of x_v exceeds $\frac{3}{2}r_\square + 2s_b + \ell_{\bar{N}}$, the triangle inequality gives $\square \subset \Omega$; the points of $\square_1$ are then at distance $\geq r_\square$ from $\square^c$. With $\square \subset \Omega$, [1, (2.26)] for the pair $(\square, \Omega)$, and the two-variable form of Proposition D.17 at points at distance $\geq R_0 + 11$ inside $\square$ ($R_0 + 12$ at $k = 1$), bound $\delta G_k(\Omega, \square, \tilde{B})$ at points of $\square_1$; since $\square$ is a parallelepiped there is no further restriction, and [3, (3.2)] uses only this boundedness. The argument for Ω_2 is identical.

(d), S13, S17–S19. S13: for $x \in \square$ and x_c the centre, $\tilde{B}_\mu(x) - \tilde{B}_\mu(x_c)$ is a sum of $\eta_k \partial^{\eta_k}\tilde{B}_\mu$ along a coordinate path in the convex box, so $|\tilde{B} - B_0| \leq d(\frac{3}{2}r_\square + s_b)\sup_\square|\partial^{\eta_k}\tilde{B}| = O(r_\square p)$ with $B_0 := \tilde{B}(x_c)$. S17: for $\lambda : \square \to \mathbb{R}$, the Neumann covariant Laplacian on bonds of $\square$ and the averaging Q_k (contours in blocks contained in $\square$) are covariant, hence $G_k(\square, B + \partial\lambda) = U(\lambda)G_k(\square, B)U(\lambda)^*$, and every class is invariant when the external scalar fields are transformed simultaneously. S18: B_0 constant gives $B_0 = \partial^{\eta_k}\lambda_0$, $\lambda_0(x) = B_0 \cdot (x - y)$, so the gauge-out of [2, (2.68)] is exact on $\square$ provided $\square$ does not wrap around the torus (a condition on the size of the torus). S19: [3, p. 433] writes $G_k(\square, 0) = G_k(0) + \delta G_k(\square, \eta_k\mathbb{Z}^d, 0)$ and treats δG_k as an external field. Since $\square$ does not wrap (S18), it is a rectangular parallelepiped of $\eta_k\mathbb{Z}^d$ and a union of big blocks. [3, p. 433] bounds $\delta G_k(\square, \eta_k\mathbb{Z}^d, 0)$ by [4, (1.11)–(1.12), p. 573] with $\Omega = \square \subset \Omega_0 = \eta_k\mathbb{Z}^d$, the setting of [4, p. 572]; since the proof in [4, Sect. 2] is written for a finite Ω_0, we argue on finite tori. Let $T' \supset \square$ be a torus whose side ℓ' is a multiple of $2M_b$, at least $\max\{3M_b, 3r_\square + 2s_b + 2\}$, with big-block grid compatible with $\square$. At $A = 0$ the operator is translation invariant and $G_k(T', 0)(x, x') = \sum_{n\in\mathbb{Z}^d} G_k(\eta_k\mathbb{Z}^d, 0)(x, x' + n\ell')$ (uniqueness of the periodic inverse, as in Lemma 5.22(i)), so

$$\delta G_k(\square, \eta_k\mathbb{Z}^d, 0) = \delta G_k(\square, T', 0) + \sum_{n\neq 0} G_k(\eta_k\mathbb{Z}^d, 0)(\cdot, \cdot + n\ell').$$

The first term is bounded by [1, (2.26)] on T', with constants independent of the side of T' (Lemma 4.6): it, its derivatives and its Hölder quotients are at most $c_0 \exp[-\delta_0(\operatorname{dist}(\{x, x'\}, \operatorname{supp} f) + \operatorname{dist}(\operatorname{supp} f, \square^c) + \operatorname{dist}(\{x, x'\}, \square^c))]\|f\|_\infty$ for all $x, x' \in \square$, with no restriction on x, x' because $\square$ is a parallelepiped. The second term tends to zero as $\ell' \to \infty$, by the exponential decay of $G_k(0)$ (the bound (H_∞) of Subsection 5.5). Hence the same bound holds for $\delta G_k(\square, \eta_k\mathbb{Z}^d, 0)$. At points of $\square_1$, at distance $\geq r_\square$ from $\square^c$, $\delta G_k(\square, \eta_k\mathbb{Z}^d, 0)$ is therefore an external field of $(1, \alpha)$-norm $\leq c_0 \mathrm{e}^{-\delta_0 r_\square}$. The remaining steps are covered by Lemma 5.9.

Conclusion. Collecting (b)–(d) with Table 2: the sketched steps need at most $R_0 + 12$ ($R_0 + 13$ at $k = 1$); P12 needs $\frac{3}{2}r_\square + 2s_b + \ell_{\bar{N}}$; nothing else depends on $\operatorname{dist}(\cdot, \Omega^c)$. The inventory (a) is the result of a page-by-page reading of [3, pp. 414–445]; the list, with the seven further places, is Lemma D.5. □

Lemma 5.9 (Ω-freeness after the cube split)**.** *Assume $\square \subset \Omega$ for the class. Then every identity and inequality on* [3, pp. 433–445] *after the split* (5.9) *is a statement about a kernel built from: (i)*

vertex coefficients localised by the functions g_k, smooth partitions of unity with supports in cubes of side 2, and the unit-cube partition of Ω_1; (ii) counterterms with the same localisation and torus propagators; (iii) $G_k(0)$ (on $\eta_k\mathbb{Z}^d$ or the torus) and its slices, the vector propagator on the torus, and C^ξ; (iv) the cubes $\Delta(v)$ of side $L^j\eta_k$; and of external fields whose $(1,\alpha)$-norms at $\square_1$ enter linearly. None of these statements involves $\operatorname{dist}(\cdot,\Omega^c)$. *The dependence on Ω_1 is used only through Lipschitz bounds of the smooth localisation functions and through exact matching of graphs and counterterms.*

Proof. After the split, every old vertex lies in $\square_1 \subset \square \subset \Omega$ and the new vertices of the B'-expansion lie in $\square$, so the characteristic function of Ω is identically 1 wherever a vertex or a line end occurs. Every object that depends on Ω ($\delta G_k(\Omega,\Omega_2,\tilde B)$, the operator [3, (1.16)], $\delta G_k(\Omega,\square,\tilde B)$, $\delta G_k(\square,\eta_k\mathbb{Z}^d,0)$, the gauge factors) has been moved into the external fields. Inspecting [3, pp. 433–445]: every propagator is $G_k(0)$, a slice $G^{\eta_k}_{(j)}(0)$, $G^{\eta_k}_{j_0}(0)$, G^ξ, C^ξ or Π; every sum runs over $\eta_k\mathbb{Z}^d$, $\xi\mathbb{Z}^d$ or the cubes $\Delta(v)$; Taylor's formula [3, (3.10)] is applied to legs and to the smooth factors g, g' ([3, (3.23), (3.26), (3.31), (3.34), (3.36)]); and the localisation of the λ-vertices is matched exactly against counterterms carrying the same localisation ([3, (3.19)], p. 444, p. 445). None of these operations refers to a distance to Ω^c. □

The kernels of Lemma 5.9 do depend on Ω_1 and on the torus; what they do not depend on is the depth.

Corollary 5.10 (Proposition 1 of [3] under a weakened support hypothesis)**.** *Let $L = 2$ and $R \geq R^{**}$ (stage 5 of Appendix A); in particular R, and $r(t_k) \geq R$ at every scale, satisfy the condition (R.e) of* (5.11) *below and the condition $R \geq 3R_0 + 15$, which is part of $R \geq R^*$. Then* [3, Prop. 1] *holds with (R4) replaced by*

$$\text{(R4}'''\text{)} \qquad \operatorname{dist}(\operatorname{supp}\tilde A, \partial\Omega) > \tfrac{7}{4}r + 2s_b + \ell_{\bar N} + 1. \tag{5.10}$$

Moreover (R4$'''$) holds at every use of [3, Prop. 1] *in the proof of Theorem 5.2 under the condition*

$$\text{(R.e)} \qquad R \geq 4\ell_{\bar N} + 32M_b + 25. \tag{5.11}$$

Proof. By Proposition 5.8 the only consumer of depth of order r is P12, with need $\frac{3}{2}r + 2s_b + \ell_{\bar N}$ at an old vertex x_v (printed cube, $r_\square = r$); every other step needs at most $R_0 + 13$ near the vertices or near $\operatorname{supp}\tilde A$ ($R_0 + 12$ for $k \geq 2$). Every vertex of a class is anchored: it lies within distance 1 of $\operatorname{supp}\tilde A$ or of $\operatorname{supp} g_k \subset \Omega_1$ (Lemma D.1). A vertex anchored at $\operatorname{supp} g_k$ has depth at least $\operatorname{dist}(\Omega_1,\Omega^c) - 1 \geq 5\rho_k - O(M_b) \geq \frac{5}{2}(r(t_{k-1}) - M_b - 1) - 4$ at $L = 2$. A vertex anchored at $\operatorname{supp}\tilde A$ has, under (R4$'''$), depth $> \frac{7}{4}r + 2s_b + \ell_{\bar N}$. Both exceed the need $\frac{3}{2}r + 2s_b + \ell_{\bar N}$ of P12 (for the first, by (5.11)), and both exceed $R_0 + 2\sqrt{3} + 13$, since $r \geq R \geq 3R_0 + 15$; the internal points of S1 have depth $> \frac{7}{4}r - 2\sqrt{3} - 1 \geq R_0 + 2\sqrt{3} + 1$. At the uses in the proof of Theorem 5.2 the supply is at least $2r - M_\theta - 2M_b - 5$ with $M_\theta \leq LM_b$ (Lemma D.3), which exceeds the right side of (5.10) if $r/4 > \ell_{\bar N} + 2s_b + M_\theta + 2M_b + 6$; with $s_b \leq 2M_b$ and $L = 2$ this follows from (5.11), since $r(t) \geq R$. □

Remark 5.11 (what is used as printed, and Bałaban's margin in Higgs II). (i) The renormalisation of the divergent classes on [3, pp. 434–445] (S21) is used as printed; by Lemma 5.9 it does not interact with the weakened hypothesis. The step at [3, (3.35)] is used as printed; its torus form is Theorem E.53. (ii) The norm comparison S20 costs a factor $(1 + O(r_\square p))^{\#\text{legs}}$, not $O(1)$ as printed ("exactly"); this is a size factor, polylogarithmic in t_k at $r_\square = r$, absorbed by the slack $(L^k\varepsilon)^{(4-d)/4-\beta}$ of the applications [3, p. 421]. (iii) In [2, p. 579] Bałaban applies Proposition 1 at the margin $2r - M - O(1)$; this margin is what his nesting supplies, not what the argument needs: the need, through P12, is $\frac{3}{2}r + O(\ell_{\bar N} + M_b)$. (iv) The identification of the torus propagators of the counterterms with the infinite-lattice propagators of the classes is Lemma 5.22.

5.4. **Repair 3: the region entropy in Bałaban's estimate (3.42) of Higgs II.** The printed proof of the combinatorial estimate [2, (3.42), p. 592], the last step of the proof of Theorem 4.2, is incomplete: the step to (3.47) is not written out, the cube count (3.45) fails near the boundary of the region, and (3.51)–(3.54) bound only the large-field sets while (3.42) also contains collar and shell sets; Proposition 5.21 completes it.

What is not written in [2]. [2, p. 592] reduces (2.117) to the combinatorial inequality (3.42), with $\zeta''_{\Lambda_0^{(k)}}$ the sum (3.41) over admissible minimal families. On p. 593 he bounds ζ'' by (3.46) and then writes "After easy transformations we get" the bound (3.47), which is a supremum over the region sequences. Two steps are not written there: replacing the sum over region sequences by a supremum, and gathering the collar and shell sets of (3.42) into $O(1)(1+\log t_k^{-1})|\Lambda_0^{(k)c}|$. Moreover the cube count (3.45) is false near $\partial\Lambda_7^{(k-1)\prime}$ (Remark 5.15).

Conventions. In this subsection $d=3$ (the arguments hold for every d); T is Bałaban's periodic lattice torus of side ℓ_T, $|T|=\ell_T^d$; $L=2$; $\varepsilon=2^{-K}\varepsilon_0$, $t_k:=L^k\varepsilon$ ($t_K=\varepsilon_0$); $\hat\ell_k:=1+\ln t_k^{-1}$; $r_k:=r(t_k)=R\hat\ell_k^{\varrho}$, $\varrho=2$; $p(t)=b_0(1+\ln t^{-1})^p$, $p=6$; M_b is Bałaban's block parameter. The letters k and j are scale indices.

Notation. $T^{(k)}$ is the unit lattice at scale k, a discrete torus with $N_k:=|T|t_k^{-d}$ points. Large blocks of $T^{(k)}$ have side $s_b:=LM_b$ lattice units ([1, p. 607]), and $\vartheta:=s_b/R$, so $s_b\le\vartheta r_k$. The prime operation $X\mapsto X'$ of [1, (1.16), p. 606] commutes with complements on unions of L-blocks. $B_\infty(y,h)$ is the ℓ^∞-ball and $N_w(X)$ the ℓ^∞-neighbourhood of width w on the torus; we only use $\mathrm{dist}_\infty\le\mathrm{dist}\le\sqrt{d}\,\mathrm{dist}_\infty$. A family $F_k=(P_v^{(k)},\dots,R_s^{(k)})$ has bad union $\mathcal{S}_k:=B(P_v^{(k)})\cup\cdots\cup R_s^{(k)}$ and size $|F_k|:=|P_v^{(k)}|+\cdots+|R_s^{(k)}|$. We write $X_k:=\Lambda_0^{(k)c}$, $\Lambda_i^{(-1)\prime}:=T^{(0)}$, $\mathrm{Col}_k:=\Lambda_7^{(k-1)\prime}\cap\Lambda_7^{(k)c}$ and $\mathrm{Sh}_k:=\Lambda_5^{(k-1)\prime}\cap\Lambda_5^{(k)c}$. Finally $c_7,c_5\ge0$ denote the two $O(1)$'s on the left of (3.42), $\kappa_0>0$ the exponent there, and $c_\square:=\min\{\frac18 a3^{-d},\frac14\gamma_0 3^{-d},3^{-d}\}$ is Bałaban's constant c_0 of [2, p. 593] (renamed to avoid a clash with the constant c_0 of Proposition 4.5). The collar exponent is $\kappa_0=\frac15$ (Lemma F.1).

Definitions, as printed. (D0) [2, (2.7)–(2.8), p. 558]: Λ_0^c is the union of the large blocks at distance $<r$ from the bad union, and Λ_{i+1}^c is the union of the large blocks at distance $\le r$ from Λ_i^c. (Dk) [2, p. 570]: $\Lambda_0^{(k)}$ is the union of those large blocks of $T^{(k)}$ that are contained in $\Lambda_7^{(k-1)\prime}$ and are at distance $>r_k$ from $\mathcal{S}_k$. [2, p. 566]: these sets are unions of big blocks, $\Lambda_0^{(j+1)}\subset\Lambda_7^{(j)\prime}$, and $\Lambda_i^{(j)}$ is defined as Λ_i with $r(t_j)$. (B-C): $\ell_T/(M_bt_K)\in\mathbb{N}$, so every $T^{(k)}$ with $k\le K-1$ is a union of large blocks. On the tori T_j, (B-C) holds by Lemma 2.1(ii).

Lemma 5.12 (elementary consequences)**.** *For every admissible sequence and $k\le K-1$: (a) every point of an element of an admissible F_k lies in X_k; (b) $\Lambda_i^{(k)c}\subset N_{i(r_k+s_b)}(X_k)$ for $i=0,\dots,7$, and $X_k\subset\Lambda_5^{(k)c}\subset\Lambda_7^{(k)c}$; (c) put $I_k:=\{z:\mathrm{dist}_\infty(z,T^{(k)}\setminus\Lambda_7^{(k-1)\prime})\ge s_b\}$ ($I_0:=T^{(0)}$). If $z\in I_k\cap X_k$, the large block $\beta\ni z$ lies in $\Lambda_7^{(k-1)\prime}$ and $\mathrm{dist}(\beta,\mathcal{S}_k)\le r_k$; (d) $T^{(k)}\setminus\Lambda_7^{(k-1)\prime}=(\Lambda_7^{(k-1)c})'$, hence $T^{(k)}\setminus I_k=N_{s_b-1}\big((\Lambda_7^{(k-1)c})'\big)$.*

Proof. (a) Let y lie in an element of F_k and let $\beta\ni y$ be its large block. Then $\mathrm{dist}(\beta,\mathcal{S}_k)=0\le r_k$. If $\beta\subset\Lambda_7^{(k-1)\prime}$, (Dk) excludes β from $\Lambda_0^{(k)}$. Otherwise β is not a candidate block. Large blocks partition $T^{(k)}$, so $\beta\cap\Lambda_0^{(k)}=\emptyset$ in both cases. (b) Each step of (2.8) moves at most r_k, plus at most s_b-1 inside a block. (c) β has ℓ^∞-diameter s_b-1, so $\beta\subset\Lambda_7^{(k-1)\prime}$. Since $\beta\not\subset\Lambda_0^{(k)}$, (Dk) gives $\mathrm{dist}(\beta,\mathcal{S}_k)\le r_k$. (d) $\Lambda_7^{(k-1)}$ is a union of L-blocks, and the prime operation commutes with complements on such unions. □

Lemma 5.13 (number of families)**.** *For admissible $\Lambda_0^{(k)}$, the number of six-tuples F_k admissible for it is at most $2^{n_d|X_k|}$, where $n_d:=4+2\max\{d,d(d-1)/2\}$ ($n_3=10$).*

Proof. By Lemma 5.12(a) every element has all its points in X_k. Charge each element to one point: an element of P-type to a point of its L-block, a bond or plaquette to its lowest corner. A point of X_k is charged by at most one element of each of the types P_v, R_v, P_s, R_s, and by at most $\max\{d, d(d-1)/2\}$ elements of each of the types Q_v, Q_s. So a six-tuple is a subset of at most $n_d|X_k|$ candidates. □

Lemma 5.14 (cube count on the interior; repair of (3.45))**.** *Partition each coordinate cycle of $T^{(k)}$ (a cycle of n_k/s_b large blocks, $n_k := \ell_T/t_k$) into consecutive arcs of $q_k, \dots, 2q_k - 1$ large blocks, where $q_k := \lceil (r_k + s_b)/s_b \rceil$; if there are fewer than q_k blocks, use one arc. Let $\mathcal{Q}_k$ be the product partition. Its boxes are unions of large blocks with sides in $[\min\{r_k + s_b, n_k\},\ 2r_k + 3s_b]$, and $2r_k + 3s_b \le (2 + 3\vartheta)r_k$. Put $\mathcal{C}_k := \{Q \in \mathcal{Q}_k : Q \cap X_k \cap I_k \neq \emptyset\}$. Then $|\mathcal{C}_k| \le 3^d |F_k|$ for every family F_k admissible for $\Lambda_0^{(k)}$.*

Proof. (i) If $\mathrm{dist}_\infty(z, x) \le r_k + s_b$, then in each coordinate the cyclic distance is at most the minimal arc length (or there are at most two arcs). So the boxes $Q(z)$ and $Q(x)$ are equal or adjacent, and a box has at most 3^d equal or adjacent boxes. (ii) Let $Q \in \mathcal{C}_k$, $z \in Q \cap X_k \cap I_k$, and $\beta \ni z$ its large block ($\beta \subset Q$). By Lemma 5.12(c) there are $z' \in \beta$ and a point y of an element $\mathfrak{e}$ of F_k with $\mathrm{dist}_\infty(z', y) \le r_k$. The charge point $x_\mathfrak{e}$ is within $\max\{1, L-1\}$ of y. Hence $\mathrm{dist}_\infty(z', x_\mathfrak{e}) \le r_k + L - 1 \le r_k + s_b$, and $Q = Q(z')$ is equal or adjacent to $Q(x_\mathfrak{e})$. Therefore $\mathcal{C}_k$ is contained in the union over $\mathfrak{e} \in F_k$ of the boxes equal or adjacent to $Q(x_\mathfrak{e})$. □

Remark 5.15 (the printed (3.45)). [2, p. 592] defines $\mathcal{C}_k$ through $\Lambda_7^{(k-1)\prime} \cap \Lambda_0^{(k)c}$ instead of the interior I_k. Now $\Lambda_7^{(k-1)\prime}$ is a union of M_b-cubes of $T^{(k)}$, not of scale-k large blocks. So a large block straddling its boundary lies in $\Lambda_0^{(k)c}$ and can be far from $\mathcal{S}_k$. For $F_k = \emptyset$, the term 1 of every ζ'', the printed $\mathcal{C}_k$ is non-empty while $3^d|F_k| = 0$. The repair (the count on the interior, with the boundary layer covered by the inherited boxes, Lemma 5.19) changes constants only.

Lemma 5.16 (bound on ζ'')**.** *For every admissible $\Lambda_0^{(k)}$ with $k \le K-1$: $\zeta''_{\Lambda_0^{(k)}} \le 2^{n_d|X_k|} \exp(-c_\square p(t_k)^2 |\mathcal{C}_k|)$.*

Proof. Each of the six coefficients $\frac{1}{8}a, \frac{1}{4}\gamma_0, \frac{1}{4}\gamma_0, \frac{1}{8}a, \frac{1}{4}\gamma_0, 1$ of (3.41) is at least $3^d c_\square$. So each term of (3.41) is at most $\exp(-3^d c_\square p(t_k)^2 |F_k|)$, which is at most $\exp(-c_\square p(t_k)^2 |\mathcal{C}_k|)$ by Lemma 5.14. The number of terms is bounded by Lemma 5.13. □

Lemma 5.17 (weighted region entropy)**.** *For $k \le K-1$ and $\theta \ge 0$: $\sum_X \mathrm{e}^{-\theta|X|} = (1 + \mathrm{e}^{-\theta s_b^d})^{N_k/s_b^d} \le \exp(N_k \mathrm{e}^{-\theta})$, where the sum runs over all unions X of large blocks of $T^{(k)}$. With $\theta_k := (d+1)\hat{\ell}_k$, $\sum_{k<K} N_k \mathrm{e}^{-\theta_k} = \mathrm{e}^{-(d+1)} |T| \sum_{k<K} t_k \le \mathrm{e}^{-(d+1)} \varepsilon_0 |T| = \mathrm{e}^{-(d+1)} \varepsilon_0^{d+1} |T^{(K)}|$.*

Proof. The large blocks partition $T^{(k)}$ into N_k/s_b^d blocks (by (B-C)), and X is any subset of them; use the product formula, then $1 + x \le \mathrm{e}^x$ and $s_b \ge 1$. Also $\mathrm{e}^{-\theta_k} = \mathrm{e}^{-(d+1)} t_k^{d+1}$, and $\sum_{k<K} t_k = \varepsilon_0 \sum_{m \ge 1} 2^{-m}$. □

A price θ independent of k would give $\sum_k N_k \mathrm{e}^{-\theta} \asymp \mathrm{e}^{-\theta} |T| \varepsilon^{-d} \to \infty$. The logarithm in the second exponent of (3.47) is this price.

Proposition 5.18 (from the sum to the supremum)**.** *Let $F(\mathbf{\Lambda})$ be the summand of (3.42), and $C_{\mathrm{fin}} := c_7 \varepsilon_0^{\kappa_0} + c_5 + \mathrm{e}^{-(d+1)} \varepsilon_0^{d+1}$. Then*

$$\sum_{\mathbf{\Lambda}\ \mathrm{adm}} F(\mathbf{\Lambda}) \le \mathrm{e}^{C_{\mathrm{fin}} |T^{(K)}|} \sup_{\mathbf{\Lambda}\ \mathrm{adm}} \exp\Big(- \sum_{k<K} c_\square p(t_k)^2 |\mathcal{C}_k| + \Xi(\mathbf{\Lambda}) \Big),$$

where $\Xi(\mathbf{\Lambda}) := \sum_{k<K} \big[(n_d \ln 2 + \theta_k)|X_k| + c_7 t_k^{\kappa_0} |\mathrm{Col}_k| + c_5 |\mathrm{Sh}_k| \big]$.

Proof. Lemma 5.16 bounds each ζ'' by a function of the region alone. The terms with $k = K$ are at most $(c_7 \varepsilon_0^{\kappa_0} + c_5)|T^{(K)}|$, since $\mathrm{Col}_K, \mathrm{Sh}_K \subset T^{(K)}$. Multiply and divide by $\mathrm{e}^{\sum_{k<K} \theta_k |X_k|}$. Bound the

resulting function $\Psi(\mathbf{\Lambda})$ by its supremum and sum $\mathrm{e}^{-\sum_k \theta_k |X_k|}$ over all sequences $(X_0, \dots, X_{K-1})$ of unions of large blocks: the map $\mathbf{\Lambda} \mapsto (X_k)_k$ is injective and all terms are non-negative. Finally apply Lemma 5.17. □

Lemma 5.19 (covering families; (3.48)–(3.52) repaired)**.** *Put* $\nu := 14 + 20\vartheta$, $\nu' := 16 + 23\vartheta$, $S_{\varrho,L} := \sum_{m\ge 0} L^{-m}(1 + m \ln L)^{\varrho}$ *(*$S_{2,2} = 2 + 4\ln 2 + 6\ln^2 2 = 7.6553$*),* $C_D^0 := \nu'(S_{\varrho,L} - 1)$ *and* $C_D := C_D^0 + 14(1 + \vartheta)$ *(*$C_D = 144.35$ *at* $\vartheta = 1/7$*). Set* $\mathcal{D}_0 := \mathcal{C}_0$*. For* $k \ge 1$ *set* $\mathcal{D}_k := \{\mathcal{P}_k(Q) : Q \in \mathcal{D}_{k-1}\} \cup \mathcal{C}_k$*, where* $\mathcal{P}_k(Q)$ *is the prime of the union of the L-blocks meeting* $N_{w_{k-1}}(Q)$*, with* $w_{k-1} := 7(r_{k-1} + s_b) + Ls_b$*. Then for* $k \le K - 1$*: (i)* $X_k \subset \bigcup \mathcal{D}_k$*; (ii)* $|\mathcal{D}_k| \le \sum_{j\le k} |\mathcal{C}_j|$*; (iii) every box of* $\mathcal{D}_k$ *has side at most* $C_D^0 r_k$*; (iv)* $|\Lambda_i^{(k)c}| \le (C_D r_k)^d \sum_{j\le k} |\mathcal{C}_j|$ *for* $i = 0, \dots, 7$*.*

Proof. (ii) is immediate. (iii) A box born at scale j has side $s_j \le (2 + 3\vartheta) r_j$ (Lemma 5.14). One step gives $s_{i+1} \le (s_i + 2w_i + 2(L-1))/L$, and $2w_i + 2L - 2 \le 14 r_i + 20 s_b \le \nu r_i$. Iterating, $s_k \le \nu' \sum_{i=j}^{k-1} L^{-(k-i)} r_i$. Since $\hat\ell_i = \hat\ell_k + (k-i)\ln L \le (1 + (k-i)\ln L)\hat\ell_k$, we have $r_i \le (1 + (k-i)\ln L)^{\varrho} r_k$, and summing gives $s_k \le C_D^0 r_k$. (i) We induct on k. The case $k = 0$ holds because $I_0 = T^{(0)}$, and $X_k \cap I_k \subset \bigcup \mathcal{C}_k$ by definition. Let $z \in X_k \setminus I_k$. By Lemma 5.12(d) there is $z_1 \in (\Lambda_7^{(k-1)c})'$ with $\mathrm{dist}_\infty(z, z_1) \le s_b - 1$. Then $B(z_1) \subset \Lambda_7^{(k-1)c} \subset N_{7(r_{k-1}+s_b)}(X_{k-1})$ by Lemma 5.12(b), and this is contained in $\bigcup_{Q\in\mathcal{D}_{k-1}} N_{7(r_{k-1}+s_b)}(Q)$ by induction. In scale-$(k-1)$ units the points of $B(z)$ are within $Ls_b - 1$ of points of $B(z_1)$. So $B(z)$ meets some $N_{w_{k-1}}(Q)$, i.e. $z \in \mathcal{P}_k(Q)$. (iv) By Lemma 5.12(b) and (i), $\Lambda_i^{(k)c} \subset \bigcup_{Q\in\mathcal{D}_k} N_{7(r_k+s_b)}(Q)$. Each of these sets has at most $(C_D^0 r_k + 14(r_k + s_b))^d \le (C_D r_k)^d$ points; now use (ii). □

Lemma 5.20 (fatness of the large-field complement)**.** *Assume* $\vartheta \le 1/7$ *and* $R \ge 12\sqrt{d}$ *(lower bounds on R of stage 5 of Appendix A). Then for every admissible sequence and* $k \le K - 1$*:* $|\Lambda_7^{(k)c}| \le C_F |\Lambda_0^{(k)c}|$*, where* $C_F := (4\sqrt{d}(8.434 + 9\vartheta) + 2)^d$ *(*$= 3.334 \cdot 10^5$ *at* $d = 3$*,* $\vartheta = 1/7$*).*

Proof. Let $h_k := \lfloor r_k/(2\sqrt{d})\rfloor - 2$. Then $h_k \ge r_k/(4\sqrt{d}) \ge 1$, because $r_k \ge R \ge 12\sqrt{d}$. (a) *Centres.* For each $z \in X_k$ there is y with $B_\infty(y, h_k) \subset X_k$ and $\mathrm{dist}_\infty(z, y) \le \omega_k := (1.434 + 2\vartheta) r_k$. If $z \in I_k$: as in Lemma 5.14(ii), $\mathrm{dist}_\infty(z, x_{\mathfrak{e}}) \le r_k + 2s_b \le (1 + 2\vartheta) r_k$. Every w with $\mathrm{dist}_\infty(w, x_{\mathfrak{e}}) \le \lfloor r_k/\sqrt{d}\rfloor$ is at distance at most r_k from $\mathcal{S}_k$, so its large block lies in X_k by the argument of Lemma 5.12(a). Take $y := x_{\mathfrak{e}}$. If $z \notin I_k$: by Lemma 5.12(d) take z_1 as in the proof of Lemma 5.19(i). The scale-$(k-1)$ large block $\beta_1 \supset B(z_1)$ lies in $\Lambda_7^{(k-1)c}$. By (2.8) there are $z_2 \in \beta_1$ and $v \in \Lambda_6^{(k-1)c}$ with $\mathrm{dist}(z_2, v) \le r_{k-1}$, and then $B_\infty(v, \lfloor r_{k-1}/\sqrt{d}\rfloor) \subset \Lambda_7^{(k-1)c}$. Let u_v be the coarse point whose L-block contains v. Then $B_\infty(u_v, h'') \subset (\Lambda_7^{(k-1)c})' \subset X_k$ with $h'' \ge r_{k-1}/(2\sqrt{d}) - 2 \ge h_k$. Moreover $\mathrm{dist}_\infty(z, u_v) \le r_{k-1}/2 + 1.5 s_b \le ((1 + \ln 2)^2/2 + 1.5\vartheta) r_k \le \omega_k$. Take $y := u_v$. (b) *Reach.* By Lemma 5.12(b), $\Lambda_7^{(k)c}$ lies within $\tau_k := 7(r_k + s_b) + \omega_k \le (8.434 + 9\vartheta) r_k$ of the set Y_k of centres. (c) *Packing.* Let $Y' \subset Y_k$ be maximal with pairwise $\mathrm{dist}_\infty > 2h_k$. The boxes $B_\infty(y', h_k)$, $y' \in Y'$, are disjoint and contained in X_k, and $\Lambda_7^{(k)c}$ lies within $\tau_k + 2h_k$ of Y'. If $2h_k + 1 \ge n_k$, then $X_k = T^{(k)}$ and there is nothing to prove. Otherwise the ratio of the two cardinalities is at most $((2\tau_k + 4h_k + 1)/(2h_k + 1))^d \le (\tau_k/h_k + 2)^d \le C_F$. □

Proposition 5.21 (the estimate (3.42) on the torus)**.** *Let* $\varepsilon_0 \le \mathrm{e}^{-1}$ *and* $2p > \varrho d + 2$ *(at* $p = 6$*,* $\varrho = 2$*,* $d = 3$ *the margin is* $\hat\ell^4$*). Put* $\Theta := C_1 C_D^d R^d/(c_\square b_0^2 \ln 2)$ *with* $C_1 := n_d \ln 2 + d + 1 + c_7 + c_5$*, and assume*

$$(1 + \ln(2/\varepsilon_0))^{2p-2-\varrho d} \ge \Theta, \qquad \textit{equivalently} \quad \varepsilon_0 \le 2\exp\big(1 - \Theta^{1/(2p-2-\varrho d)}\big). \tag{5.12}$$

(E.ent)

Then the left side of [2, (3.42)] *is at most* $\exp(C_{\mathrm{fin}}|T^{(K)}|) = \exp(C_{\mathrm{fin}}\varepsilon_0^{-d}|T|)$*. This holds uniformly in* K*, in* k *and in the torus* T*, for every torus whose side* ℓ_T *satisfies* $\ell_T/(M_b t_K) \in \mathbb{N}$ *(Lemma 2.1(ii)). If in addition* $\vartheta \le 1/7$ *and* $R \ge 12\sqrt{d}$*, the printed bound* [2, (3.47)] *holds with second* $O(1)$ *equal to* $C_2 := n_d \ln 2 + (d+1) + (c_7 + c_5) C_F$ *and third* $O(1)$ *equal to* C_{fin}*.*

Proof. By Proposition 5.18 it suffices to show $\Xi(\mathbf{\Lambda}) \le \sum_{j<K} c_\square p(t_j)^2 |\mathcal{C}_j|$. The sets X_k, Col_k and Sh_k lie in $\Lambda_7^{(k)c}$ (Lemma 5.12(b)), and $t_k^{\kappa_0} \le 1 \le \hat\ell_k$. So each coefficient in Ξ is at most $C_1\hat\ell_k$. By Lemma 5.19(iv), after swapping the sums, $\Xi \le C_1 C_D^d \sum_{j<K} |\mathcal{C}_j| \sum_{k=j}^{K-1} \hat\ell_k r_k^d$. Here $\hat\ell_k r_k^d = R^d \hat\ell_k^{1+\varrho d}$ decreases in k, and the number of terms is $K-j = \log_2(\varepsilon_0/t_j) \le \hat\ell_j/\ln 2$. Hence $\Xi \le c_\square b_0^2 \Theta \sum_j \hat\ell_j^{2+\varrho d} |\mathcal{C}_j|$. Since $p(t_j)^2 = b_0^2 \hat\ell_j^{2p}$ and $\hat\ell_j \ge 1 + \ln(2/\varepsilon_0)$ for $j \le K-1$, (E.ent) gives the claim. For the printed form: by Lemma 5.20, $|\mathrm{Col}_k|, |\mathrm{Sh}_k| \le C_F |X_k|$, and $n_d \ln 2 + (d+1)\hat\ell_k + (c_7+c_5) C_F \le C_2 \hat\ell_k$. □

The bound of Proposition 5.21 holds for arbitrary constants $c_5, c_7 \ge 0$. Let ε_1 be the cap of stage 10a of Appendix A. If c_5 and c_7 are bounded for $\varepsilon_0 \in (0, \varepsilon_1]$ (for c_7 this uses $\kappa_0 < \frac14$, which $\kappa_0 = \frac15$ meets), then (E.ent), with Θ evaluated at these bounds, is a downward-closed upper bound on ε_0. The constants c_5, c_7 are bounded on $(0, \varepsilon_1]$ by Statement K.1. Hence (E.ent) is a condition of stage 10b of Appendix A.

5.5. **Torus and infinite-lattice propagators.** At step $k \le K$ the torus T_ε, rescaled, has sides $S_k^{(\mu)} = 2L_\mu/t_k$ in units of scale-k blocks, with deck group $\Gamma_k := \bigoplus_\mu S_k^{(\mu)} \mathbb{Z} e_\mu$; put $S_k := \min_\mu S_k^{(\mu)} = 2L_{\min}/t_k$, so $S_k = L^{K-k} S_K$ and $S_K = 2L_{\min}/\varepsilon_0$. Let $\mathcal{K}$ be the set of kernels $G_k(0) = (-\Delta^{\eta_k} + m^2 t_k^2 + a_k P_k)^{-1}$, its slices $G_{(j)}^{\eta_k}(0)$, $0 \le j \le k$, and $C^\xi = (-\Delta^\xi + 1)^{-1}$, with their derivatives of order ≤ 2, each defined on $X \in \{T_{\eta_k}, \eta_k \mathbb{Z}^d\}$ (written K_T, K_∞). By Proposition 4.5 and [3, (2.10)–(2.12), p. 426, and p. 437], with constants depending on d, a, M_b, L only:

$$(\mathrm{H}_\infty) \qquad |K_\infty(x,x')| + \|K_\infty\|_{1,\alpha;x,x'} \le C_0 \mathrm{e}^{-\delta_0|x-x'|} \ (|x-x'| \ge 1), \qquad \sum_{x'} \eta_k^d |K_\infty(x,x')| \le C_0.$$

In (H_∞) and in the definition of $\mathcal{K}$, a derivative of a kernel means at most one lattice difference in each of the two variables. Mixed second differences $\partial_x \partial_{x'} K$ are therefore included, and pure second differences in one variable are not. The norm $\|K\|_{1,\alpha;x,x'}$ consists of the pointwise values of these kernels together with the Hölder quotients of their first differences. The summability bound $\sum_{x'} \eta_k^d |K_\infty(x,x')| \le C_0$ is required only for kernels with at most one difference in total. For a mixed second difference of G_i the sum is bounded only by a constant times i (Statement K.2(iii)), so this bound is not uniform in k. On the tori T_{η_k} the kernels K_T, $K \in \mathcal{K}$, satisfy the pointwise and Hölder bounds of (H_∞) with constants C_0, δ_0 that do not depend on the torus, by Proposition 4.5, [3, (2.10)–(2.12), p. 426] and Lemma 4.6. The constants obtained in this way are those of the kernels at the photon mass μ_*. For the scalar kernels $G_k(0)$ and their slices at a Higgs mass $M \in (0, m_2]$ with $M\varepsilon_0 \le 1$, (H_∞) in the above form is Statement K.2(ii), with constants depending only on a, L and α. By (E.m2) this covers $M \in \{m_1, m_2\}$, and Lemma 5.22 is applied at these masses only through that statement.

Lemma 5.22 (periodisation)**.** *Assume (H_∞).*

(i) $K_T(x,x') = \sum_{n \in \Gamma_k} K_\infty(\tilde x, \tilde x' + n)$ *for every $K \in \mathcal{K}$ and all lifts $\tilde x, \tilde x'$.*

(ii) *For $F_T = \mathrm{Per}\, F_\infty$ as in (i), every Γ_k-periodic kernel V_T and every weight g,*

$$\sum_{x' \in T_{\eta_k}} \eta_k^d F_T(x,x') V_T(x,x') g(x') = \sum_{y \in \eta_k \mathbb{Z}^d} \eta_k^d F_\infty(\tilde x, y) V_T(x, \pi y) g(\pi y),$$

with π the projection to the torus.

(iii) *If $\tilde x, \tilde x'$ lie in one lifted cube of diameter D and $S_k \ge \max\{2D, 2/\delta_0\}$, then $\|K_T - K_\infty\|_{1,\alpha} \le C_1 \mathrm{e}^{-\delta_0 S_k/2}$ on that cube, with $C_1 = c_d C_0$.*

(iv) *Assume the condition (E.r), $\varepsilon_0 < \mathrm{e}^{1-\varrho}$ (stage 10b of Appendix A), and the condition*

$$(5.13) \qquad (T.per) \qquad 2L_{\min} \ge \varepsilon_0 \max\{6\sqrt{d}\, r(\varepsilon_0) + 2,\ 2/\delta_0\}.$$

Then $\mathrm{e}^{-\delta_0 S_k/2} \le \mathrm{e}^{-3\delta_0 r(t_k)}$ at every $k \le K$, and $\sum_{k \le K} \mathrm{e}^{-\delta_0 S_k/2} \le 2\mathrm{e}^{-\delta_0 S_K/2}$.

(v) *Transform every counterterm graph of* [3, (1.26)–(1.31)] *as the classes are transformed on* [3, p. 445]. *Then its scalar lines lie in one cube* $\square$ *of side* $3r(t_k)$ *that does not wrap around the torus. On* $\square\times\square$ *write* $G_k(T_{\eta_k},0)=G_k(0)+\delta G_k(T_{\eta_k},\eta_k\mathbb{Z}^d,0)$, *and the same for each slice. The term with only* $G_k(0)$ *is the counterterm in the form used in* [3, (3.7)–(3.9), p. 435, (3.19), p. 438, and pp. 444–445]. *Every other term contains an external scalar field of* $(1,\alpha)$*-norm* $\le C_1\mathrm{e}^{-3\delta_0 r(t_k)}$ *and is bounded by* [3, (3.3)].

Proof. (i) By the pointwise bound of (H_∞) the series converges absolutely, and it is Γ_k-periodic in both variables. The operator $H=-\Delta^{\eta_k}+m^2t_k^2+a_kP_k$ is nearest-neighbour plus block averaging. Every block of scale $j\le k$ has side $L^{j-k}\le 1$, and the sides $S_k^{(\mu)}$ are multiples of the block side, so H commutes with Γ_k. Unfolding one sum gives $H_T\operatorname{Per}K_\infty=\operatorname{Per}(H_\infty K_\infty)=\operatorname{Per}\delta=\delta_T$. H_T is positive on a finite-dimensional space, hence invertible, and $\operatorname{Per}K_\infty=H_T^{-1}=K_T$.

For $1\le i\le k+1$ let G_i be H^{-1} with a_kP_k replaced by $a_iL^{2(k-i)}P_i$, where P_i is the projection onto the fields constant on the blocks of side L^{i-k}. On each lattice the slices satisfy $G_{(0)}=G_1$ and $G_{(j)}=G_{j+1}-G_j$ (Statement K.3(f); at zero background $Q_k(0)$ is the plain block average). Each G_i is periodised by the argument above. For $i\le k$ this uses blocks of side ≤ 1. The kernel G_{k+1}, which enters $G_{(k)}$, uses blocks of side L and the divisibility $L\mid S_k^{(\mu)}$. This divisibility holds for $k<K$ since $S_k^{(\mu)}=LS_{k+1}^{(\mu)}$, and at $k=K$ on the tori T_j by the count of big blocks after Lemma 4.6 (see 4.3). Derivatives commute with Per.

(ii) Split the sum over lifts of x'; the remaining factors are periodic.

(iii) For $n\in\Gamma_k\backslash\{0\}$ we have $|n|\ge S_k$ and $|\tilde x-\tilde x'-n|\ge|n|-D\ge|n|/2\ge 1$. So $\sum_{n\neq0}C_0\mathrm{e}^{-\delta_0|n|/2}\le C_0\sum_{m\in\mathbb{Z}^d\setminus0}\mathrm{e}^{-\delta_0S_k|m|/2}\le c_dC_0\mathrm{e}^{-\delta_0S_k/2}$ once $\delta_0S_k/2\ge1$. The same argument applies to the Hölder seminorms. Only the pointwise and Hölder bounds of (H_∞) are used here.

(iv) $S_k/r(t_k)$ is decreasing in t_k on $(0,\varepsilon_0]$ for $\varepsilon_0<\mathrm{e}^{1-\varrho}$ (condition (E.r); Lemma A.2(b)). So $S_K\ge6\sqrt{d}\,r(\varepsilon_0)+2$ gives $S_k\ge6\sqrt{d}\,r(t_k)+2$ for all $k\le K$, hence $\mathrm{e}^{-\delta_0S_k/2}\le\mathrm{e}^{-3\sqrt{d}\,\delta_0r(t_k)}\le\mathrm{e}^{-3\delta_0r(t_k)}$. Moreover $\sum_k\mathrm{e}^{-\delta_0L^{K-k}S_K/2}\le\mathrm{e}^{-\delta_0S_K/2}\sum_{n\ge0}\mathrm{e}^{-(L^n-1)}\le2\mathrm{e}^{-\delta_0S_K/2}$, since $\delta_0S_K/2\ge1$.

(v) By S18 in the proof of Proposition 5.8, the near vertices of the class and of the counterterm lie in one cube $\square$ of side $3r(t_k)$, of diameter $D\le3\sqrt{d}\,r(t_k)$.

The cube does not wrap. By (T.per) and (E.r), Lemma A.2(b) gives $S_k\ge\max\{6\sqrt{d}\,r(t_k)+2,\,2/\delta_0\}$ at every $k\le K$, so $3r(t_k)<S_k/2$. Choose a lift $\tilde\square\subset\eta_k\mathbb{Z}^d$ of $\square$ whose coordinate differences are at most $3r(t_k)$. Let $n\in\Gamma_k$ and let μ be a coordinate with $n_\mu\neq0$. Then $|n_\mu|\ge S_k$, so $|\tilde x_\mu-\tilde x'_\mu-n_\mu|>S_k/2>|\tilde x_\mu-\tilde x'_\mu|$ for $\tilde x,\tilde x'\in\tilde\square$. Hence π is a bijection from $\tilde\square$ onto $\square$, and $|\tilde x-\tilde x'-n|\ge\max\{|\tilde x-\tilde x'|,S_k/2\}$ for $n\neq0$. So $\square$ does not wrap around the torus, and the torus distance on $\square$ is the distance on $\tilde\square$.

The expansion. Since $S_k\ge2D$, (i), (iii) and (iv) give, for every $K\in\mathcal{K}$, $K_T=K_\infty+\delta K$ on $\square\times\square$ with $\|\delta K\|_{1,\alpha}\le C_1\mathrm{e}^{-\delta_0S_k/2}\le C_1\mathrm{e}^{-3\delta_0r(t_k)}$. The derivatives in the norm are again kernels of $\mathcal{K}$, so no margin around $\square$ is needed. We argue by induction on the order. Lower-order counterterms inserted in the graph have smaller order [3, p. 417] and are treated first.

Insert this splitting for every scalar line and expand the product; the expansion of the product of at most $\bar N$ lines is finite. The vertices, the far lines (which became external fields in the same way on both sides) and the vector lines (the same periodic kernels in the class and in the counterterm) are not changed. So the term with $G_k(0)$ on every scalar line is the counterterm in the class form of [3, (3.7)–(3.9), (3.19), pp. 444–445], under the identification $\square\cong\tilde\square$.

Every other term has a nonempty set of lines carrying δG_k. These lines are replaced by the two-variable external fields $\delta G_k|_{\square\times\square}$. This is the operation that [3, p. 433] performs with $\delta G_k(\square,\eta_k\mathbb{Z}^d,0)$, and the resulting decomposition into renormalised classes depends only on which lines are dropped [3, p. 432]. Hence the term is of the type treated in [3, (3.3)], which is multilinear in the external-field norms. $\square$

Here $L_{\min}$ is the half-side of Bałaban's torus $\{-L_\mu \le x_\mu < L_\mu\}$ [1, (1.2), pp. 603–604], so $2L_{\min} = \ell_j$ on T_j, and (T.per) is the lower bound $\ell_0 \ge \varepsilon_0 \max\{6\sqrt{3}\, r(\varepsilon_0) + 2,\ 2/\delta_0\}$ of stage 11 of Appendix A; it does not depend on ε. By Lemma A.2(b), the passage in (iv) from $k = K$ to every $k \le K$ needs only the condition (E.r), $\varepsilon_0 < \mathrm{e}^{1-\varrho}$.

6. From the vacuum bound to the uniform tube bound

This section proves Theorem 6.8, a bound on the generating functionals of the torus states that is uniform in ε, in the torus and in the photon mass, from Theorem 5.2 and the results of Section 4. The bound is a product of three factors. The factor for the curvature is the Gaussian domination of Lemma 7.3. The factor for positive sources of the Wick square (Proposition 6.5) comes from a convexity bound for constant sources (Proposition 6.4), which uses Bałaban's vacuum bounds (Proposition 6.2), the c-number differences of Proposition 6.3 and the identification of the counterterms (Proposition 4.7). The factor for negative sources (Proposition 6.7) comes from the slice vacuum bound (Proposition 6.6), which follows from Theorem 5.2 at $(u, M) = (1, m_2)$ and $(-\eta, m_1)$. Every estimate is proved on one torus at a time; no comparison between tori of different sizes and no statement about the model on $\varepsilon\mathbb{Z}^3$ is used.

6.1. **The scheme.** The states of Theorem 3.1 are built in the torus counterterm scheme of Definition 2.2: on each (ε, T_j) the bare quadratic coefficient is $b_T := b_{T_j}(\varepsilon) = a_{K,T_j}(m)$, King's counterterm on the torus at Higgs mass m and reference photon mass $\mu_* = m$, the same for every photon mass $\mu_0 \in (0, \mu_*]$ and for the state $\flat$. The renormalised source is $X_{\mathrm{R}}(h) = :|\varphi|^2:_{\mathrm{R}}(h) = \sum_x \varepsilon^3 h(x)(r_x^2 - W_\varepsilon - z_\varepsilon)$ with $r_x := |\varphi(x)|$. Here $W_\varepsilon = 2\int_{[-\pi/\varepsilon,\pi/\varepsilon)^3} (\hat{p}^2 + m^2)^{-1}\, d^3p/(2\pi)^3$ is the free value of $\mathbb{E}|\varphi(0)|^2$ on the infinite lattice $\varepsilon\mathbb{Z}^3$ (two real components, hence the factor 2), which does not depend on the torus, and z_ε is the order-e^2 Wick constant of Subsection 2.4. The same b_T, W_ε and z_ε enter the states, the observables $X_{\mathrm{R}}(h)$ and every bound of this section. The norms are $\|h\|_{L^1,\varepsilon} = \sum_x \varepsilon^3 |h(x)|$, $\|h\|_\infty$ and $\|\mathbf{g}\|_{2,\varepsilon}$ of Subsection 2.4. On the periodic lattice, $-\Delta_\varepsilon$ acting on each component of a bond function is the Hodge Laplacian $d^*d + dd^*$ (Lemma B.4), and $C_\mu = (-\Delta_\varepsilon + \mu^2)^{-1}$.

Two objects of this section are defined here. For $s \in (0, \infty)$ put

$$C_s^\flat := (-\Delta_\varepsilon)^{-1}\Pi^\perp + s|T|\,\Pi_0,$$

with $(-\Delta_\varepsilon)^{-1}$ the inverse on the range of $\Pi^\perp$: the photon covariance whose non-constant modes are massless and whose constant mode has variance s in each coordinate c_μ (in the inner product of Subsection 2.2, the constant bond function $c\mathbf{1}_\mu$ has squared norm $|T|\,c^2$). Then $C_{\mu_0} = (-\Delta_\varepsilon + \mu_0^2)^{-1}\Pi^\perp + s(\mu_0)|T|\,\Pi_0$ with $s(\mu_0) = (\mu_0^2|T|)^{-1}$, and $C_s^\flat \ge C_{\mu_*}$ as forms if and only if $s \ge s(\mu_*)$, because $(-\Delta_\varepsilon)^{-1} \ge (-\Delta_\varepsilon + \mu_*^2)^{-1}$ on the range of $\Pi^\perp$. Let $\langle\cdot\rangle_{\varepsilon,T,s}$ be the expectation of Definition 2.7 with C_{μ_0} replaced by $C_s^\flat$. For bounded $F \in \mathfrak{A}_\varepsilon$, $\langle F\rangle_{\varepsilon,T,s} \to \langle F\rangle_{\varepsilon,T,\flat}$ as $s \to \infty$: step (ii) of the proof of Lemma 2.8 uses only that the law of $A^\perp$ is fixed and the variance of the constant mode tends to infinity. Second, the proof of Proposition 6.7 uses the slice-modified covariance

$$C_{B,s}^\flat := (C_{\mu_*} + \Gamma^B + \Gamma_\ge)\Pi^\perp + s|T|\,\Pi_0 \quad (0 < s < \infty), \qquad C_B^\flat := C_{B,\infty}^\flat \text{ (Haar-distributed constant mode)},$$

with Γ^B of (5.1) and the operator $\Gamma_\ge$ of Proposition 6.7 (Definition I.1); it satisfies $C_{\mu_0} \le C_{B,s}^\flat$ for $\mu_0 \in (0, \mu_*]$ and $s \ge s(\mu_0)$ (Lemma I.3). The class $\mathcal{D}$, the set $E^{\mathcal{D}}$ and the lattice norms are those of Subsection 2.4.

We write $\mathfrak{C}$ for the family of pairs (ε, T_j) with $\varepsilon = 2^{-K}\varepsilon_0$, $K \ge K_0$, and $j \ge 1$. Suprema over ε and T below run over $\mathfrak{C}$ unless said otherwise.

6.2. **The counterterm gaps.**

Theorem 6.1 (counterterm gaps)**.** *Let $m > 0$, $\mu_* := m$, $M_1 := 0.7m$, $H := m^2$, and $(e, \lambda, m, \eta, m_1, m_2) \in \mathfrak{W}$. Then, uniformly in $\varepsilon \le 1/m$ and in every periodic torus T of side $\ell_T \ge 96/m$:*

(a) $a_{K,T}(m) - a_{K,T}(M_1) \ge 0.319\, m^2$*;*
(b) $g_{21} := \inf_{\varepsilon,T}[a_{K,T}(m_2) - a_{K,T}(m_1)] \ge (m_2 - m_1)(m_2 + m_1 - 0.3631m) > 0$*, the infimum running over the same ε and T;*
(c) $a_{K,T}(m_1) \ge a_{K,T}(m) + 2H$*;*
(d) $m < m_1 < m_2$*,* $m_1 \ge \sqrt{2}\mu_*$*,* $M_1 < m$*.*

Proof. Scaling. Every c-number $a_{K,T}(M)$ is a finite sum of lattice graphs of orders $\lambda, e^2, \lambda^2, e^2\lambda, e^4$. Each graph is homogeneous under the change of units in which lengths (x, ε, ℓ_T) are multiplied by m, masses (M, μ) are divided by m, and $\lambda \mapsto \lambda/m$, $e^2 \mapsto e^2/m$, $a \mapsto a/m^2$ (in $d = 3$, λ and e^2 have the dimension of a mass). So it suffices to take $m = \mu_* = 1$, $\varepsilon \le 1$, $\ell_T \ge 96$.

(a) Proposition G.3 gives, for $0 < M_1 < M_2$, $\varepsilon \le 1$ and every torus of side at least a given floor,

$$a_{K,T}(M_2) - a_{K,T}(M_1) \ge (M_2^2 - M_1^2) + 16\lambda h_1 - e^2\bar{j} - \lambda^2\bar{s} - e^2\lambda\bar{B}_3 - e^4\bar{B}_4^-,$$

with non-negative constants independent of the couplings. At $(M_1, M_2) = (0.7, 1)$ and for tori of side ≥ 96, Corollary G.4 gives $\bar{j} \le 0.0886293$, $\bar{s} \le 7.179064$, $\bar{B}_3 \le 9.330427$, $\bar{B}_4^- \le 1.703784$ and $h_1 \ge 0$. Drop the λ-term and use the monotonicity in $\lambda, e^2 \ge 0$. On $[0, 0.1]^2$ the bound is then at least $0.51 - 0.00886293 - 0.07179064 - 0.09330427 - 0.01703784 = 0.31900432$.

(b) Proposition G.6 gives $\partial_M a_{K,T}(M) \ge 2M - p(M)$, where $p(M)$ is the majorant of that proposition (not the small-field parameter $p(t)$). It is a sum of the monomials $e^2, \lambda^2, e^2\lambda, e^4$ times non-negative coupling-free functions. Its supremum over $M \ge 2$ is at most $e^2 s_1 + \lambda^2 s_2 + e^2\lambda s_3 + e^4 s_4$, with $s_1 \le 0.3178$, $s_2 \le 9.6941$, $s_3 \le 17.37755$, $s_4 \le 5.96785$ for tori of side ≥ 64 (Corollary G.7). On the box $[0, 0.1]^2$ this is at most $0.03178 + 0.096941 + 0.1737755 + 0.0596785 = 0.362175 < 0.3622 \le 0.3631$. Integrating from m_1 to m_2 gives (b).

(c) Corollary G.8 gives $a_{K,T}(m_1) - a_{K,T}(m) \ge (m_1 - m)(m_1 + m - S(m))$, where $S(m) \ge \sup_{M \ge m} p(M)$. On the box one may take $S(m) \le 0.612m$ (Corollary G.9). So (c) holds for $m_1 \ge m_1^c := \frac{1}{2}[S(m) + ((S(m) - 2m)^2 + 8H)^{1/2}]$, and $m_1^c \le 1.882m < 2m < m_1$.

(d) holds by construction: $m < 2m < m_1 < m_2$, $m_1 > 2m > \sqrt{2}\mu_*$, and $M_1 = 0.7m < m$.

The ingredients of (a)–(c) are proved in Appendix G. The exact two-loop form of $a_{K,T}$ (Lemma G.1) rests on Lemma G.16, whose order $e^2\lambda$ uses the list of one-particle-irreducible graphs in Lemma G.19, and on the order-e^4 coefficient of Corollary G.22. The numerical constants are the cell upper sums (G.1) (proof of Corollary G.4). For (b) and (c) these are combined with the bound for large masses of Statement K.54. □

That the conditions of Appendix A can be met at every point of $\mathfrak{W}$, with $\varepsilon \le \varepsilon_0 \le 1/m_2 < 1/m$ for every spacing of the construction and $\ell_0 \ge 96/m$, so that Theorem 6.1 applies to all spacings and tori of the construction, is Lemma A.5.

6.3. **Bałaban's bounds at the masses used.**

Proposition 6.2 (Bałaban's vacuum bounds at the masses used)**.** *Let $(e, \lambda, m, \eta, m_1, m_2) \in \mathfrak{W}$, $\mu_* := m$, and let the regulator data satisfy the conditions of Appendix A. For every $(\varepsilon, T_j) \in \mathfrak{C}$, with $T = T_j$,*

$$Z_K(T; M) \ge \mathrm{e}^{-E_-(M)|T|} \quad (M \in \{M_1, m, m_1, m_2\}), \qquad Z_K(T; M) \le \mathrm{e}^{E_+(M)|T|} \quad (M \in \{M_1, m_1, m_2\}),$$

with $E_\pm(M)$ independent of ε and j.

Proof. The hypotheses of Theorems 4.1 and 4.2 hold for $Z_K(T;M)$ by Lemma 4.3, its action is Bałaban's by Proposition 4.7, and every $T_{j,\varepsilon}$ is a torus of the form [1, (1.2)] (Subsection 4.1). The lower bound is Theorem 4.1. The upper bound is Theorem 4.2; at the masses M_1, m_1, m_2 its printed proof is completed by Proposition 5.21 and Lemma I.24 (Remark 4.4). □

Proposition 6.3 (c-number differences)**.** *Let* $f_T(a) := |T|^{-1}\log Z_T^F(a) + \frac{1}{2}a(W_\varepsilon + z_\varepsilon)$*, where* $Z_T^F(a) := \int \mathrm{e}^{-S_{\mu_*,a}(A,\varphi)}\,dA\,d\varphi$ *is the unnormalised integral (Lebesgue measure in* A *and* φ*) with the action* (2.1) *at photon mass* μ_* *and bare coefficient* a*, and put* $\gamma_T(M) := f_T(a_{K,T}(M)) - |T|^{-1}\log Z_K(T;M)$*. Then* $\sup_{K,\,\ell_T\ge\ell_0}|\gamma_T(M) - \gamma_T(M')| < \infty$ *for* $\{M,M'\} = \{M_1, m\}$ *and for* $\{M,M'\} = \{m_1, m_2\}$*.*

Proof. This is Proposition H.8; by Lemma H.1(a) the γ_T defined here is the one there. Its proof uses lattice power counting in $d = 3$ (Proposition C.3). □

6.4. **The factor for positive sources.** For a bare coefficient a put

$$\alpha_T(\sigma;a) := |T|^{-1}\log\langle\exp(\sigma X_{\mathrm{R}}(1_T))\rangle_{\varepsilon,T,\mu_*;a},$$

the expectation in the model at photon mass μ_* with bare coefficient a.

Proposition 6.4 (convexity bound for constant positive sources)**.** *Let* $(e,\lambda,m,\eta,m_1,m_2) \in \mathfrak{W}$*,* $\mu_* := m$*,* $M_1 := 0.7m$*, and let the regulator data satisfy the conditions of Appendix A. Put*

$$g := \inf_{K,\,\ell_T\ge\ell_0}[a_{K,T}(m) - a_{K,T}(M_1)], \qquad \sigma_+ := g/2,$$

$$\bar\Gamma := \sup_{K,\,\ell_T\ge\ell_0}[\gamma_T(M_1) - \gamma_T(m)], \qquad \kappa_+ := 2[E_+(M_1) + E_-(m) + \bar\Gamma]_+/g.$$

Then $g \ge 0.319m^2$*,* $\bar\Gamma < \infty$*, and for every* $(\varepsilon,T_j) \in \mathfrak{C}$ *and* $0 \le \sigma \le \sigma_+$*,*

$$\alpha_T(\sigma;b_T) \le \kappa_+\sigma, \qquad T = T_j.$$

Proof. Every spacing of the family satisfies $\varepsilon \le \varepsilon_0 \le 1/m_2 < 1/m$ (Lemma 5.4(b)), and $\ell_0 \ge 96/m$ (Appendix A), so $g \ge 0.319m^2$ by Theorem 6.1(a); $\bar\Gamma < \infty$ by Proposition 6.3. Write $a_1 := a_{K,T}(M_1)$, $a_2 := b_T = a_{K,T}(m)$. The single-site weight of the model is $\mathrm{e}^{-\varepsilon^3(\frac{1}{2}ar_x^2 + \lambda r_x^4)}$, so $\langle \mathrm{e}^{\sigma\sum_x \varepsilon^3 r_x^2}\rangle_a = Z_T^F(a-2\sigma)/Z_T^F(a)$, and therefore $\alpha_T(\sigma;a) = f_T(a-2\sigma) - f_T(a)$. The function $a \mapsto \log Z_T^F(a)$ is convex (its second derivative is the variance of $\frac{1}{2}\sum_x \varepsilon^3 r_x^2$), hence so is f_T. For $0 \le \sigma \le \sigma_+ \le (a_2-a_1)/2$ write $a_2 - 2\sigma = (1-\theta)a_2 + \theta a_1$ with $\theta = 2\sigma/(a_2-a_1) \in [0,1]$; convexity gives

$$\alpha_T(\sigma;a_2) \le \theta\,[f_T(a_1) - f_T(a_2)] = \frac{2\sigma}{a_2-a_1}\Big[\frac{\log Z_K(T;M_1) - \log Z_K(T;m)}{|T|} + \gamma_T(M_1) - \gamma_T(m)\Big]$$
$$\le \frac{2\sigma}{g}\big[E_+(M_1) + E_-(m) + \bar\Gamma\big]_+,$$

by Proposition 6.2 and $a_2 - a_1 \ge g$. □

Proposition 6.5 (the factor for positive sources)**.** *Under the hypotheses of Proposition 6.4, for every* $(\varepsilon,T_j) \in \mathfrak{C}$*, every* $\mu_0 \in (0,\mu_*] \cup \{\flat\}$ *and every lattice function* q *with* $0 \le q \le \sigma_+$*,*

$$\log\langle \mathrm{e}^{X_{\mathrm{R}}(q)}\rangle_{\varepsilon,T_j,\mu_0} \le \log\langle \mathrm{e}^{X_{\mathrm{R}}(q)}\rangle_{\varepsilon,T_j,\mu_*} \le \kappa_+\|q\|_{L^1,\varepsilon},$$

all states at the coefficient b_{T_j}*.*

Proof. (1) For $q \ge 0$, $\mathrm{e}^{X_{\mathrm{R}}(q)}$ is a positive constant times $\mathrm{e}^{r^2(\varepsilon^3 q)}$, with $r^2(\varepsilon^3 q) = \sum_x \varepsilon^3 q(x) r_x^2$ in the notation of Subsection 4.6, item 1; its expectation does not increase when the photon covariance increases [8, Cor. 6.3(1), p. 281] (on the torus graph, Lemma 7.5). Since b_T, W_ε, z_ε do not depend on μ_0 and $C_{\mu_0} \ge C_{\mu_*}$ as forms (zero mode included), the first inequality holds for $\mu_0 \in (0,\mu_*]$; this step uses only $q \ge 0$. For $\flat$ the observable $F_q := \mathrm{e}^{X_{\mathrm{R}}(q)}$ is unbounded, so the limit $s \to \infty$ of

Subsection 6.1, which concerns bounded observables, does not apply to it directly. Instead, F_q is an elementary observable of Lemma 7.6: for $\delta > 0$, the case just proved, applied to $(1+\delta)q \geq 0$, gives $\sup_{0<\mu_0\leq\mu_*}\langle |F_q|^{1+\delta}\rangle_{\varepsilon,T_j,\mu_0} \leq \langle \mathrm{e}^{X_{\mathrm{R}}((1+\delta)q)}\rangle_{\varepsilon,T_j,\mu_*} < \infty$. Hence $\langle F_q\rangle_{\varepsilon,T_j,\flat} = \lim_{\mu_0\downarrow 0}\langle F_q\rangle_{\varepsilon,T_j,\mu_0} \leq \langle F_q\rangle_{\varepsilon,T_j,\mu_*}$. (The proof of Lemma 7.6 uses Lemma 7.5, not this proposition.)

(2) The measure at photon mass μ_* on T_j is reflection positive, on gauge-invariant observables, about the coordinate planes halfway between lattice planes (Proposition 7.9(a), with the plane pairs of case (ii) of Lemma 7.8), and N is even, so the chessboard estimate (Subsection 4.6, item 2) applies to the site factors $F_x = \exp(\varepsilon^3 q(x)(r_x^2 - W_\varepsilon - z_\varepsilon))$. The product of the reflected copies of F_x over all sites is $\exp(q(x)X_{\mathrm{R}}(1_T))$ in the same model, so

$$\langle \mathrm{e}^{X_{\mathrm{R}}(q)}\rangle_{\varepsilon,T_j,\mu_*} \leq \prod_x \langle \mathrm{e}^{q(x)X_{\mathrm{R}}(1_T)}\rangle_{\varepsilon,T_j,\mu_*}^{\varepsilon^3/|T|} = \exp\Big(\sum_x \varepsilon^3 \alpha_T(q(x);b_T)\Big).$$

(3) By Proposition 6.4, $\alpha_T(q(x);b_T) \leq \kappa_+ q(x)$. Sum over x. □

6.5. **The factor for negative sources.** With $a_i := a_{K,T}(m_i)$ put

$$\Delta_1^B := \sup_{(\varepsilon,T_j)\in\mathfrak{C},\ T=T_j} \Big\{ |T|^{-1}\log\frac{Z(C_1^B;a_2)}{Z(C_1^B;a_1)} + \tfrac{1}{2}(a_2-a_1)(W_\varepsilon + z_\varepsilon)\Big\}, \qquad C_1^B := C_B(1) = C_{\mu_*} + \Gamma^B.$$

We call the statement $\Delta_1^B < \infty$ the *slice vacuum bound*.

Proposition 6.6 (the slice vacuum bound)**.** *Assume Theorem 5.2 at $(1,m_2)$ and at $(-\eta,m_1)$. Assume the lower bound $|T|^{-1}\Phi_{m_1}(0)+e(\varepsilon;0;m_1) \geq -E_-(m_1)$ on every torus of $\mathfrak{C}$; this is Theorem 4.1 at m_1, by Lemma H.1(b) and Proposition 6.2. Assume also the two c-number bounds*

$$\sup_{\varepsilon,T}\ \sup_{u\in[-\eta,1]} |\partial_u^2 e(\varepsilon;u;M)| \leq E_2(M), \qquad M\in\{m_1,m_2\}, \tag{6.1}$$

$$\bar\Gamma_{21}^{(1)} := \sup_{\varepsilon,T}\big[e(\varepsilon;1;m_1) - e(\varepsilon;1;m_2) + \tfrac{1}{2}(a_2-a_1)(W_\varepsilon+z_\varepsilon)\big] < \infty. \tag{6.2}$$

Then

$$\Delta_1^B \leq C_+^B(1;m_2) + \eta^{-1}C_+^B(-\eta;m_1) + (1+\eta^{-1})E_-(m_1) + \tfrac{1}{2}(1+\eta)E_2(m_1) + \bar\Gamma_{21}^{(1)} < \infty.$$

Proof. Since Φ_{m_1} is convex and $-\eta < 0 < 1$,

$$\Phi_{m_1}(0) \leq \frac{1}{1+\eta}\Phi_{m_1}(-\eta) + \frac{\eta}{1+\eta}\Phi_{m_1}(1),$$

that is, $\Phi_{m_1}(1) \geq (1+\eta^{-1})\Phi_{m_1}(0) - \eta^{-1}\Phi_{m_1}(-\eta)$. Insert the upper bounds (5.2) for $\Phi_{m_2}(1)$ and $\Phi_{m_1}(-\eta)$, and the lower bound for $\Phi_{m_1}(0)$, into $|T|^{-1}[\Phi_{m_2}(1) - \Phi_{m_1}(1)] + \frac{1}{2}(a_2-a_1)(W_\varepsilon+z_\varepsilon)$. Write $e(u;M)$ for $e(\varepsilon;u;M)$. The c-numbers regroup as

$$\big[e(1;m_1) - e(1;m_2) + \tfrac{1}{2}(a_2-a_1)(W_\varepsilon+z_\varepsilon)\big] - \big[e(1;m_1) - (1+\eta^{-1})e(0;m_1) + \eta^{-1}e(-\eta;m_1)\big].$$

The first bracket is at most $\bar\Gamma_{21}^{(1)}$. The second is a second divided difference of the polynomial $u\mapsto e(\varepsilon;u;m_1)$ on $[-\eta,1]$, so its absolute value is at most $\frac{1}{2}(1+\eta)\sup_{[-\eta,1]}|\partial_u^2 e|$. □

Assume the hypotheses of Theorem 5.2, together with the two further hypotheses of Lemma H.9: $2\bar n + 1 \leq 3R$, and (H_∞) for the kernels at the masses M and μ_*. Then the bound (6.1) at $M\in\{m_1,m_2\}$ is Lemma H.9. Its proof in Appendix H rests on Statement K.58, and the inputs of that step are described after the proof. The bound (6.2) is Proposition H.11. Its one-loop part is Lemma H.10, which uses King's slice at a point (Lemma H.27). That lemma in turn rests on the block Poincaré inequality and the variational form of $\Delta^{(j)}$ (Lemmas I.11, H.20 and H.28), with the constant c_1 of Lemma H.29.

Proposition 6.7 (the factor for negative sources)**.** *Assume $\Delta_1^B < \infty$ and the anchor condition*

$$b_T + 2H \le a_1 \quad \text{on every torus of } \mathfrak{C}, \qquad g_{21} > 0 \tag{6.3}$$

(Theorem 6.1(b), (c)). For every $0 \le h \le H$, every $\mu_0 \in (0, \mu_*] \cup \{\flat\}$ and every $(\varepsilon, T_j) \in \mathfrak{C}$, with $T = T_j$,

$$\log\langle \mathrm{e}^{-X_{\mathrm{R}}(h)}\rangle_{\varepsilon,T,\mu_0} \le 2\|h\|_{L^1,\varepsilon}\, S_\flat^B, \qquad S_\flat^B := [\Delta_1^B + J_{\mathrm{IR}} + \bar{Z}]_+/g_{21}.$$

Here $\gamma_\ge(u) := \int_{\varepsilon_0^2}^\infty \mathrm{e}^{-tu}(1-\mathrm{e}^{-t\mu_*^2})\,dt$ for $u > 0$, so that the operator $\Gamma_\ge$ of Subsection 6.1 is $\gamma_\ge(-\Delta_\varepsilon)\Pi^\perp$, and $\hat{k}^2$ runs over the nonzero eigenvalues of $-\Delta_\varepsilon$ on one component. With these,

$$J_{\mathrm{IR}} := \sup_{K,\,\ell_T \ge \ell_0} \tfrac{3}{2}|T|^{-1} \sum_{\hat{k}^2 \ne 0} \gamma_\ge(\hat{k}^2)\,(\hat{k}^2+\mu_*^2), \qquad Z_T := |T|^{-1}\Big[\tfrac{3}{2}\log_+ \frac{4\pi^2\mu_*^2\ell_T}{e^2} + 3\delta_P\Big], \quad \delta_P := 2\sum_{n\ge 1}\mathrm{e}^{-2\pi^2 n^2},$$

and $\bar{Z} := \sup_{j\ge 1} Z_{T_j}$. The constant J_{IR} is finite and depends on $\mu_*, \varepsilon_0, \ell_0$ only, not on ε, e, λ or ℓ_T. The constant Z_T depends on the side ℓ_T (and on μ_*, e) and tends to 0 as $\ell_T \to \infty$. The constant $\bar{Z}$ depends on μ_*, e, ℓ_0 only, and

$$\bar{Z} \le \frac{6\pi^2\mu_*^2}{e^2\ell_0^2} + \frac{3\delta_P}{\ell_0^3} < \infty, \tag{6.4}$$

since $\log_+ x \le x$ for $x > 0$ and $\ell_{T_j} = j!\,\ell_0 \ge \ell_0$.

Proof. The full proof is Proposition I.9; here we describe its steps. Let $\rho^B(y; b)$ be the one-site density $\langle r_y^2 - W_\varepsilon - z_\varepsilon\rangle$ in the model with photon covariance $C_B^\flat$ (the slice-modified massless photon with Haar-distributed constant mode) and constant coefficient b, and let $\bar{\rho}^B(b)$ be its average over a cell.

(1) We use three facts. The first is the translation-averaging lemma on the torus (Lemma I.6), which uses that the model of the states is invariant under $\varepsilon\mathbb{Z}^3$ and the comparison model under $\varepsilon_0\mathbb{Z}^3$. The second is the monotonicity of $\langle \mathrm{e}^{-X_{\mathrm{R}}(h)}\rangle$ in the photon covariance [8, Cor. 6.3(1), p. 281] (Lemma I.3). The third is its monotonicity in the coefficient: by [8, Thm. 6.1, p. 276] with P, Q polynomials in r, $\langle r_x^2; r_y^2\rangle \ge 0$ in every model of the family $b \mapsto b + 2th$, so $t \mapsto \langle r_y^2\rangle_t$ is non-increasing. Together they give $\log\langle \mathrm{e}^{-X_{\mathrm{R}}(h)}\rangle_{\mu_0} \le -\|h\|_{L^1,\varepsilon}\,\bar{\rho}^B(b_T + 2H) \le -\|h\|_{L^1,\varepsilon}\,\bar{\rho}^B(a_1)$, the second inequality by (6.3).

(2) $f^B(b) := |T|^{-1}\log Z(C_B^\flat; b) + \frac{1}{2}b(W_\varepsilon + z_\varepsilon)$ is convex with $f^{B\prime} = -\frac{1}{2}\bar{\rho}^B$, so $-\bar{\rho}^B(a_1) \le 2[f^B(a_2) - f^B(a_1)]/(a_2 - a_1)$.

(3) Comparing $C_B^\flat$ with C_1^B from above and below costs $J_{\mathrm{IR}} + Z_T$ per unit volume. This follows from the monotonicity of $Z(\Gamma; a)$ in Γ (current representation) and a determinant comparison for the infrared part (Lemma I.8).

(4) Hence, on $T = T_j$, $f^B(a_2) - f^B(a_1) \le \Delta_1^B + J_{\mathrm{IR}} + Z_{T_j} \le \Delta_1^B + J_{\mathrm{IR}} + \bar{Z}$. Divide by $a_2 - a_1 \ge g_{21}$ and keep the positive part. By (6.4) the constant does not depend on ε, j or μ_0.

(5) The case $\mu_0 = \flat$. For $h \ge 0$ we have $\mathrm{e}^{-X_{\mathrm{R}}(h)} \le \mathrm{e}^{(W_\varepsilon + z_\varepsilon)\|h\|_{L^1,\varepsilon}}$, since $r_x^2 \ge 0$, so this observable is bounded, and it lies in the algebra $\mathfrak{F}$ of Definition 7.1. By Lemma 7.6, $\langle \mathrm{e}^{-X_{\mathrm{R}}(h)}\rangle_{\varepsilon,T,\flat} = \lim_{\mu_0\downarrow 0}\langle \mathrm{e}^{-X_{\mathrm{R}}(h)}\rangle_{\varepsilon,T,\mu_0}$, and the bound, which is uniform in $\mu_0 \in (0, \mu_*]$, passes to the limit.

The domination of the infrared part. The comparison $C_{\mu_0} \le C_{B,s}^\flat$ of Lemma I.3 is used in step (1), and the slice covariance Γ^B enters $C_B^\flat$ in step (3). Both rest on Corollary I.17(a): $D^< \le \Gamma^B\Pi^\perp$ on every torus T_j and for every $K \ge 1$, where $\Gamma^B = \kappa\mu_*^2 D$ and $\kappa = \frac{15}{2}$ is the fixed constant of (5.1). This needs only $\kappa \ge \bar{\kappa}$, with $\bar{\kappa}$ the K-independent constant of Theorem I.16. At $L = 2$ and $a = 1$, Theorem I.16 gives $\kappa_1\kappa_3 \le \bar{\kappa} \le \frac{42176}{5625} < \frac{15}{2}$ for every $K \ge 1$ under (E.dom), $\mu_*\varepsilon_0 \le 0.2$. The condition (E.$\delta$), $\kappa\mu_*^2\varepsilon_0^2 \le \frac{1}{12}$, gives $\mu_*\varepsilon_0 \le 90^{-1/2} < 0.2$ and so implies (E.dom). The block Poincaré inequality behind Theorem I.16 holds on every torus of $\mathfrak{C}$. The reason is that for every

$k \le K$ the blocks of scale t_k contain no wrap-around bond, because $\ell_j \ge 2t_k$ (Lemma I.15(a), (c)). The domination constant does not depend on ε, K, ℓ_j or μ_0. □

6.6. **The tube bound.** For real lattice 2-forms $\mathbf{g}_1, \dots, \mathbf{g}_n$, real lattice functions $h_1, \dots, h_p$ on T_ε and $\zeta = (z, w) \in \mathbb{C}^{n+p}$ let

$$G_{\varepsilon,T,\mu_0}(\zeta) := \Big\langle \exp\Big[\sum_i z_i F(\mathbf{g}_i) + \sum_k w_k X_{\mathrm{R}}(h_k)\Big]\Big\rangle_{\varepsilon,T,\mu_0}, \qquad k(\zeta) := \sum_k (\operatorname{Re} w_k) h_k = k_+ - k_-,$$

with $k_\pm \ge 0$ the pointwise positive and negative parts, and, for $\sigma_+, H > 0$, let $\mathcal{D}_{\sigma_+,H} := \{\zeta : \|k_+(\zeta)\|_\infty < \sigma_+/2,\ \|k_-(\zeta)\|_\infty < H/2\}$ (no restriction on z). These are the objects of Subsection 2.4 with lattice test functions in place of sampled continuum ones; for sampled test functions the continuum tube lies in the lattice tube (Lemma 7.14).

Theorem 6.8 (the uniform tube bound)**.** *Let $m > 0$, $0 < \lambda < 0.1m$ and $0 < e^2 < 0.1m$, as in Theorem 3.1, put $\mu_* := m$ and $(\eta, m_1, m_2) := (\frac{1}{2}, 3m, 4m)$, so that $(e, \lambda, m, \eta, m_1, m_2) \in \mathfrak{W}$ (any point of $\mathfrak{W}$ with these (e, λ, m) may be used instead), and let the regulator data be those of Lemma A.5 at this point. Put $\sigma_+ := g/2$ and κ_+ as in Proposition 6.4, $S_\flat^B$ as in Proposition 6.7, $\mathsf{K} := \max(\kappa_+, 2S_\flat^B)$ and $H := m^2$. Then for every $K \ge K_0$, every $j \ge 1$, every $\mu_0 \in (0, \mu_*] \cup \{\flat\}$, all real lattice test functions and every $\zeta \in \mathcal{D}_{\sigma_+,H}$,*

$$\begin{aligned} |G_{\varepsilon,T_j,\mu_0}(\zeta)| \le G_{\varepsilon,T_j,\mu_0}(\operatorname{Re}\zeta) &\le \exp\Big(\tfrac{1}{2}\Big\|\sum_i (\operatorname{Re} z_i)\mathbf{g}_i\Big\|_{2,\varepsilon}^2 + \kappa_+\|k_+\|_{L^1,\varepsilon} + 2S_\flat^B\|k_-\|_{L^1,\varepsilon}\Big) \\ &\le \exp\Big(\tfrac{1}{2}\Big\|\sum_i (\operatorname{Re} z_i)\mathbf{g}_i\Big\|_{2,\varepsilon}^2 + \mathsf{K}\|k\|_{L^1,\varepsilon}\Big). \end{aligned} \tag{6.5}$$

The constants σ_+, κ_+, $S_\flat^B$ and K do not depend on ε, j or μ_0.

The constant $S_\flat^B$ contains $\bar{Z}$, not Z_T, so it does not depend on the torus, and neither does K (Corollary B.28). The constants σ_+ and K are the constants σ_+ and C of Theorem 3.1(i) (Lemma I.19).

Proof. By Lemma I.19 and Proposition I.26, with these data the hypotheses of Propositions 6.5 and 6.7 hold on every torus of $\mathfrak{C}$, and Lemma 7.3 holds there at the coefficient b_{T_j}. (1) For real test functions $F(\mathbf{g}_i)$ and $X_{\mathrm{R}}(h_k)$ are real, so $|G(\zeta)| \le G(\operatorname{Re}\zeta)$. (2) Put $\mathbf{g} := \sum_i (\operatorname{Re} z_i)\mathbf{g}_i$. By Lemma 7.3, which holds on every torus for every real source of $:|\varphi|^2:$ and every bare coefficient, $G(\operatorname{Re}\zeta) \le \mathrm{e}^{\frac{1}{2}\|\mathbf{g}\|_{2,\varepsilon}^2}\langle \mathrm{e}^{X_{\mathrm{R}}(k)}\rangle$. (3) $X_{\mathrm{R}}(k) = X_{\mathrm{R}}(k_+) - X_{\mathrm{R}}(k_-)$, and by the Cauchy–Schwarz inequality $\langle \mathrm{e}^{X_{\mathrm{R}}(k)}\rangle \le \langle \mathrm{e}^{X_{\mathrm{R}}(2k_+)}\rangle^{1/2}\langle \mathrm{e}^{-X_{\mathrm{R}}(2k_-)}\rangle^{1/2}$. (4) Since $0 \le 2k_+ < \sigma_+$, Proposition 6.5 gives $\langle \mathrm{e}^{X_{\mathrm{R}}(2k_+)}\rangle^{1/2} \le \mathrm{e}^{\kappa_+\|k_+\|_{L^1,\varepsilon}}$. (5) Since $0 \le 2k_- < H$, Proposition 6.7 gives $\langle \mathrm{e}^{-X_{\mathrm{R}}(2k_-)}\rangle^{1/2} \le \mathrm{e}^{2S_\flat^B\|k_-\|_{L^1,\varepsilon}}$. (6) k_+ and k_- have disjoint supports, so $\|k_+\|_{L^1,\varepsilon} + \|k_-\|_{L^1,\varepsilon} = \|k\|_{L^1,\varepsilon}$. Every step is on one torus at the coefficient b_{T_j}, and the constants do not depend on ε, j or μ_0. □

Remark 6.9 (what the proof uses). The proof of Theorem 6.8 uses Bałaban's vacuum bounds at four Higgs masses, his propagator bounds and his Proposition 1 (Section 4, with the repairs of Section 5), King's counterterms, the correlation inequalities and the current representation of [8], lattice reflection positivity on the tori and the chessboard estimate, and the c-number and counterterm-gap bounds of Proposition 6.3 and Theorem 6.1. It does not use King's cube states, a cluster expansion, or any bound uniform in the photon mass in a fixed cube; the uniformity in μ_0 comes from the monotonicity in the photon covariance (Propositions 6.5 and 6.7).

7. Lattice statements on the tori

Sections 7–9 prove parts (ii)–(iv) of Theorem 3.1 from part (i), the uniform tube bound, which is Theorem 6.8. Everything else in these three sections is either a statement about one lattice model on one torus, or an abstract compactness, holomorphy or positivity argument about the resulting

family of functionals. No cube, no boundary condition other than periodic, and no comparison between two different tori is used in Sections 7–9.

7.1. **Standing assumptions.** *Parameters and regulators.* Throughout Sections 7–9, $m > 0$, $0 < \lambda < 0.1\,m$ and $0 < e^2 < 0.1\,m$; the reference photon mass is $\mu_* := m$, and $H := m^2$. The regulator data (blocking factor $L = 2$, $\bar n = 13$, order cap $\bar N = \bar n^2 = 169$, M_b, R, b_0, $\varepsilon_0 = 2^{-n_0}$, and $\ell_0 \in 2M_b\mathbb{N}$ with $\ell_0 \geq 183/m$) are those of Theorem 6.8, chosen as in Lemma A.5 at the point $(e, \lambda, m, \frac12, 3m, 4m)$ of the window $\mathfrak{W}$ of Section 5. Sections 7–9 use the regulator data only through Theorem 6.8 and through the properties (P1)–(P4) below.

Lattices and tori. The lattice spacings are $\varepsilon = \varepsilon_K := 2^{-K}\varepsilon_0$ with $K \geq K_0 = \max(K_0', n_0 + 1)$ (Subsection 4.1); they are nested: $\varepsilon_{K'}\mathbb{Z}^3 \supset \varepsilon_K\mathbb{Z}^3$ for $K' \geq K$. The tori are $T_j = (\mathbb{R}/\ell_j\mathbb{Z})^3$, $\ell_j = j!\,\ell_0$, $j \geq 1$, with the lattice tori $T_{j,\varepsilon} = T_j \cap \varepsilon\mathbb{Z}^3$ and $N := \ell_j/\varepsilon$ sites per side. In Sections 7–9 we write T for a torus T_j, $\ell_T := \ell_j$, $|T| := \ell_T^3$, $T_\varepsilon := T_{j,\varepsilon}$, and $\langle\cdot\rangle_{\varepsilon,T,\mu_0} := \langle\cdot\rangle_{\varepsilon,j,\mu_0}$ for the states of Definition 2.7, $\mu_0 \in (0, \mu_*] \cup \{\flat\}$.

Fields, forms and the action. We use the notation of Subsection 2.2: A is a real function on the bonds of T_ε, φ is a function from the sites to $\mathbb{R}^2 \cong \mathbb{C}$, and $r_x := |\varphi_x|$. Sites, bonds and plaquettes carry the weight ε^3 in $\langle\cdot,\,\cdot\rangle_\varepsilon$, $\|f\|_{2,\varepsilon}^2 := \langle f,\, f\rangle_\varepsilon$, and d, d^* are the lattice exterior derivative and its adjoint. At photon mass $\mu_0 > 0$, the action (2.1) with the coefficient $a = b_T := b_{T_j}(\varepsilon) = a_{K,T}(m)$ of the torus counterterm scheme (Definition 2.2) reads, in real coordinates,

(7.1)
$$S_{\varepsilon,T,\mu_0}(A,\varphi) = \tfrac12\langle A,\,(-\Delta_\varepsilon+\mu_0^2)A\rangle_\varepsilon + S_m(A,\varphi), \qquad S_m = \tfrac12\sum_b \varepsilon^3|(D_A\varphi)(b)|^2 + \sum_x \varepsilon^3\big(\tfrac12 b_T r_x^2 + \lambda r_x^4\big),$$

with $(D_A\varphi)(b) = \varepsilon^{-1}\big(U(\varepsilon e A_b)\varphi(b_+) - \varphi(b_-)\big)$ for the bond b from b_- to b_+, where $U(s)$ is the rotation of $\mathbb{R}^2$ by the angle s; thus $S_m = S_{b_T}^{\mathrm m}$ in the notation of (2.2). The photon part is King's Feynman form. The observables are the smeared curvature and the renormalised square of the Higgs field of Subsection 2.4,

$$F(\mathbf{g}) = \langle \mathbf{g},\, dA\rangle_\varepsilon = \langle d^*\mathbf{g},\, A\rangle_\varepsilon, \qquad X_{\mathrm R}(h) = \sum_x \varepsilon^3 h(x)\big(r_x^2 - W_\varepsilon - z_\varepsilon\big), \tag{7.2}$$

for real lattice 2-forms $\mathbf{g}$ and real lattice functions h; W_ε is the free Wick constant and z_ε the order-e^2 c-number of Subsection 2.4.

The arguments of Sections 7–9 use the following properties of the regularised model, and nothing else about b_T, W_ε, z_ε:

- (P1) b_T, W_ε, z_ε are real numbers that depend on (ε, T) only: they do not depend on the site, on the direction, or on μ_0 (Lemma B.19(a)).
- (P2) $\lambda > 0$, so that $\varphi \mapsto e^{-\lambda\varepsilon^3|\varphi|^4 + a|\varphi|^2}$ is integrable on $\mathbb{R}^2$ for every $a \in \mathbb{R}$ (Lemma B.22).
- (P3) The same b_T, W_ε, z_ε are used in the state, in the observables (7.2), and in Theorem 6.8 (Lemma B.24).
- (P4) $N = \ell_T/\varepsilon$ is even, the lattices are nested, and $\ell_j \to \infty$ (Lemma 2.1).

Apart from Theorem 6.8, Sections 7–9 use the blocking factor only through (P4) and through the density in $\mathbb{R}^3$ of the group $\bigcup_K \varepsilon_K\mathbb{Z}^3 = \mathbb{Z}[1/2]^3\varepsilon_0$ (proof of Theorem 9.2). Both have the same form for blocking factor 3 (Lemma 2.1, and the density of $\mathbb{Z}[1/3]^3\varepsilon_0$); this is how Sections 7–9 enter the proof of Theorem 3.5.

States. For a positive semidefinite operator C on the bond functions, symmetric for $\langle\cdot,\,\cdot\rangle_\varepsilon$, we write γ_C for the centred Gaussian measure with $\mathbb{E}_{\gamma_C}[\langle f,\, A\rangle_\varepsilon\langle f',\, A\rangle_\varepsilon] = \langle f,\, Cf'\rangle_\varepsilon$. For $\mu_0 \in (0, \mu_*]$ the state $\langle\cdot\rangle_{\varepsilon,T,\mu_0}$ is the normalised expectation with weight $e^{-S_{\varepsilon,T,\mu_0}}\,dA\,d\varphi$, Lebesgue measure on bonds and sites (Lemma B.7); it is a probability measure by (P2) and $\mu_0 > 0$. Its photon part is

$\gamma_{C_{\mu_0}}$ with $C_{\mu_0} = (-\Delta_\varepsilon + \mu_0^2)^{-1}$, so that

$$\langle G\rangle_{\varepsilon,T,\mu_0} = \frac{\int d\gamma_{C_{\mu_0}}(A) \int d\varphi\, G\, e^{-S_m(A,\varphi)}}{\int d\gamma_{C_{\mu_0}}(A) \int d\varphi\, e^{-S_m(A,\varphi)}}. \tag{7.3}$$

Write $A = A^\perp + c$, where $c = \Pi_0 A = \sum_{\mu=1}^3 c_\mu \mathbf{1}_\mu$ is the projection onto the constant 1-forms ($\mathbf{1}_\mu(b) = 1$ for the bonds b in direction e_μ and 0 for the others) and $A^\perp = \Pi^\perp A$. Under $\gamma_{C_{\mu_0}}$ the components $A^\perp$ and c are independent; $A^\perp$ has covariance $(-\Delta_\varepsilon + \mu_0^2)^{-1}\Pi^\perp$, and each c_μ is centred normal with variance $s(\mu_0) = 1/(\mu_0^2|T|)$ (Lemma B.6). In the state $\flat$ (Definition 2.7) $A^\perp$ has the law $\gamma_{(-\Delta_\varepsilon)^{-1}\Pi^\perp}$, and c is independent of $A^\perp$ and uniformly distributed on the period cell $[0,P)^3$, $P = 2\pi/(|e|\,\ell_T)$.

Definition 7.1 (the observable algebra $\mathfrak{F}$)**.** $\mathfrak{F}$ is the algebra of complex Borel functions of the variables $(r_x)_x$ and $((dA)_p)_p$; it is the class $\mathfrak{A}_\varepsilon$ of Definition 2.4. For a region Λ of the torus, $\mathfrak{F}(\Lambda)$ is the subalgebra of the functions of the r_x with $x \in \Lambda$ and of the $(dA)_p$ with all bonds of p in Λ. Every polynomial and every exponential $\exp[\sum_i z_i F(\mathbf{g}_i) + \sum_k w_k X_{\mathrm{R}}(h_k)]$, $z, w \in \mathbb{C}$, lies in $\mathfrak{F}$.

Every element of $\mathfrak{F}$ is gauge invariant: under $(A,\varphi) \mapsto (A + d\chi, U(-e\chi)\varphi)$, χ a real function on the sites, r_x and dA do not change. Elements of $\mathfrak{F}$ are also unchanged by $A \mapsto A + t\mathbf{1}_\mu$ for every $t \in \mathbb{R}$, since $d\mathbf{1}_\mu = 0$.

Lemma 7.2 (Hodge facts on the periodic lattice)**.** *On T_ε write $\partial_\nu f(x) = \varepsilon^{-1}(f(x+\varepsilon e_\nu) - f(x))$, identify a 1-form A with its components $A_\mu(x) := A_{\langle x, x+\varepsilon e_\mu\rangle}$, and let ∂_μ^* be the adjoint of ∂_μ for $\langle\cdot,\cdot\rangle_\varepsilon$ (minus the backward difference). Then $(dA)_{\mu\kappa} = \partial_\mu A_\kappa - \partial_\kappa A_\mu$, $d^*A = \sum_\mu \partial_\mu^* A_\mu$, and:*

- (H-a) $\|dA\|_{2,\varepsilon}^2 + \|d^*A\|_{2,\varepsilon}^2 = \sum_{\mu,\nu}\|\partial_\nu A_\mu\|_{2,\varepsilon}^2$*; that is, $d^*d + dd^*$ is the componentwise lattice Laplacian.*
- (H-b) $\ker d$ *on functions is the space of constants, and* $\ker(d^*d + dd^*)$ *on 1-forms is spanned by* $\mathbf{1}_1, \mathbf{1}_2, \mathbf{1}_3$.
- (H-c) $\operatorname{ran} d^*$ *(on 1-forms) is orthogonal to the constants and invariant under $-\Delta_\varepsilon$, and $-\Delta_\varepsilon = d^*d$ there; this restriction is injective.*
- (H-d) $\Pi := d(d^*d)^+d^*$*, with $(\cdot)^+$ the Moore–Penrose inverse, is the orthogonal projection onto* $\operatorname{ran} d$ *(on 2-forms).*

Proof. (H-a) All forward and backward differences commute on the torus, and $\|\partial_\mu^* f\| = \|\partial_\mu f\|$. Hence $\langle\partial_\mu^* A_\mu, \partial_\nu^* A_\nu\rangle = \langle A_\mu, \partial_\mu\partial_\nu^* A_\nu\rangle = \langle\partial_\nu A_\mu, \partial_\mu A_\nu\rangle$. Expanding the squares, the mixed terms of $\|dA\|^2$ (summed over unordered pairs $\mu < \kappa$, as the weight ε^3 per unordered plaquette prescribes) and of $\|d^*A\|^2$ cancel. (H-b) follows from (H-a), because a function with vanishing forward differences is constant on the connected torus. (H-c) $\langle d^*\mathbf{g}, \mathbf{1}_\mu\rangle = \langle\mathbf{g}, d\mathbf{1}_\mu\rangle = 0$, and $-\Delta_\varepsilon d^*\mathbf{g} = d^*dd^*\mathbf{g} + dd^*d^*\mathbf{g} = d^*(dd^*\mathbf{g})$, since $d^*d^* = 0$. The restriction of d^*d to $\operatorname{ran} d^* = (\ker d)^\perp$ is injective, because $d^*dv = 0$ implies $\|dv\| = 0$. (H-d) Π is self-adjoint. Since $(d^*d)^+(d^*d)$ is the orthogonal projection onto $(\ker d)^\perp$, $\Pi d = d$; hence $\Pi^2 = \Pi$ and $\operatorname{ran} d \subset \operatorname{ran}\Pi \subset \operatorname{ran} d$. □

Parts (H-a) and (H-b) are also contained in Lemma B.4, which identifies $-\Delta_\varepsilon$ with the lattice Laplacian on each component and gives the spectral gap $-\Delta_\varepsilon \geq 16/\ell_T^2$ on $\operatorname{ran}\Pi^\perp$.

Lemma 7.3 (Gaussian domination of the curvature on tori)**.** *For every real lattice 2-form $\mathbf{g}$, every real lattice function k and every $\mu_0 \in (0, \mu_*] \cup \{\flat\}$,*

$$\langle e^{F(\mathbf{g})} e^{X_{\mathrm{R}}(k)}\rangle_{\varepsilon,T,\mu_0} \leq e^{\frac{1}{2}V(\mathbf{g})}\langle e^{X_{\mathrm{R}}(k)}\rangle_{\varepsilon,T,\mu_0}, \qquad V(\mathbf{g}) := \langle d^*\mathbf{g}, C\, d^*\mathbf{g}\rangle_\varepsilon \leq \|\Pi_{\operatorname{ran} d}\mathbf{g}\|_{2,\varepsilon}^2 \leq \|\mathbf{g}\|_{2,\varepsilon}^2,$$

with $C = C_{\mu_0}$, or $C = (-\Delta_\varepsilon)^{-1}\Pi^\perp$ in the state $\flat$, and $\Pi_{\operatorname{ran} d}$ the orthogonal projection onto the exact 2-forms.

Proof. Put $f := d^*\mathbf{g}$, so that $F(\mathbf{g}) = \langle f,\, A\rangle_\varepsilon$. Since $\langle f,\, \mathbf{1}_\mu\rangle_\varepsilon = \langle \mathbf{g},\, d\mathbf{1}_\mu\rangle_\varepsilon = 0$, f is orthogonal to the constants and $F(\mathbf{g}) = \langle f,\, A^\perp\rangle_\varepsilon$. By Subsection 2.4, $\int d\varphi\, e^{-S_m + X_{\mathrm{R}}(k)} = c_k\, W_k(A)$ with the positive constant $c_k = e^{-\sum_x \varepsilon^3 k(x)(W_\varepsilon + z_\varepsilon)}$, which cancels in the ratios below, and W_k has the current representation (2.3) of Lemma 2.5. Write its phase as $e\varepsilon \sum_b n_b A_b = e\langle j_n,\, A\rangle_\varepsilon$ with $j_n := \varepsilon^{-2} n$.

Case $\mu_0 > 0$. By (7.3) and Lemma 2.5 (the series converges absolutely, so Fubini applies), with $C = C_{\mu_0}$,

$$\mathbb{E}_{\gamma_C}\big[e^{\langle f,\, A\rangle_\varepsilon} W_k(A)\big] = \sum_n w_k(n)\, \mathbb{E}_{\gamma_C}\big[e^{\langle f + ie j_n,\, A\rangle_\varepsilon}\big] = e^{\frac12\langle f,\, Cf\rangle_\varepsilon} \sum_n w_k(n)\, e^{-\frac12 e^2 \langle j_n,\, Cj_n\rangle_\varepsilon} \cos\big(e\langle j_n,\, Cf\rangle_\varepsilon\big),$$

where the sine terms cancel in pairs because $w_k(n) = w_k(-n)$. The cosine is at most 1, and $\sum_n w_k(n) e^{-\frac12 e^2\langle j_n, Cj_n\rangle_\varepsilon} = \mathbb{E}_{\gamma_C}[W_k(A)]$ (the same identity at $f = 0$). Divide by $\mathbb{E}_{\gamma_C}[W_0]$: the left side becomes $\langle e^{F(\mathbf{g})} e^{X_{\mathrm{R}}(k)}\rangle$ and the right side is at most $e^{\frac12\langle f,\, Cf\rangle}\langle e^{X_{\mathrm{R}}(k)}\rangle$.

Case $\flat$. Averaging (2.3) over $c \in [0,P)^3$ with the uniform law replaces $w_k(n)$ by $w_k(n)\mathbf{1}\{\omega(n) = 0\}$ (Lemma 2.5: the phase of the constant mode is $e\ell_T \sum_\mu c_\mu \omega_\mu(n)$ and $|e|\ell_T P = 2\pi$), which is still nonnegative and even in n. The Gaussian step is the same, with $A^\perp$ of covariance $C = (-\Delta_\varepsilon)^{-1}\Pi^\perp$ and $f \in \operatorname{ran}\Pi^\perp$.

The constant. By Lemma 7.2(H-c), $\operatorname{ran} d^*$ (on 1-forms) reduces $-\Delta_\varepsilon$, which equals d^*d there and is injective there. Hence on $\operatorname{ran} d^*$ we have $C_{\mu_0} = (d^*d + \mu_0^2)^{-1} \le (d^*d)^{-1}$, and $(-\Delta_\varepsilon)^{-1}\Pi^\perp = (d^*d)^{-1}$, since $\operatorname{ran} d^*$ is orthogonal to the constants. Therefore $V(\mathbf{g}) \le \langle d^*\mathbf{g},\, (d^*d)^+ d^*\mathbf{g}\rangle_\varepsilon = \langle \mathbf{g},\, d(d^*d)^+ d^*\mathbf{g}\rangle_\varepsilon = \|\Pi_{\operatorname{ran} d}\mathbf{g}\|^2_{2,\varepsilon}$ by (H-d). □

Remark 7.4. On the torus the constant is $\frac12\|\mathbf{g}\|^2_{2,\varepsilon}$ at every lattice spacing and for every $\mu_0 \ge 0$, the state $\flat$ included. It needs no infinite-volume input and no infrared bound in the sense of [18]: the Hodge decomposition is exact on the periodic lattice.

Lemma 7.5 (monotonicity in the photon covariance on one torus)**.** *Fix (ε, T) and the matter part of (7.1), and let $\langle\cdot\rangle_C$ denote (7.3) with $\gamma_{C_{\mu_0}}$ replaced by γ_C for an arbitrary positive semidefinite C on the bond functions. Let $g \ge 0$ be a site function, $r^2(g) := \sum_x g_x r_x^2$, and ℓ a real linear functional of A. If $0 \le C_0 \le C_1$, then*

$$\langle e^{r^2(g)} e^{i\ell(A)}\rangle_{C_1} \le \langle e^{r^2(g)} e^{i\ell(A)}\rangle_{C_0}, \qquad \langle e^{-r^2(g)} e^{\ell(A)}\rangle_{C_1} \ge \langle e^{-r^2(g)} e^{\ell(A)}\rangle_{C_0}.$$

That is, the class $\mathcal{I}$ of the functions $e^{r^2(g)} e^{i\ell(A)}$ is non-increasing in the covariance, and the class $\mathcal{D}$ of the functions $e^{-r^2(g)} e^{\ell(A)}$ is non-decreasing.

This is the torus form of [8, Cor. 6.3(1), p. 281]. The inequalities [8, Thm. 6.2(1),(3), p. 280] that it uses are proved in [8, Sect. 6.1, pp. 275–276] for an arbitrary finite lattice and an arbitrary centred Gaussian measure γ_C, $C \ge 0$, with Dirichlet conditions, with the remark (p. 276) that periodic conditions on rectangles are allowed. Lemma B.12 and Proposition B.13 prove them on every finite graph, the lattice torus with its wrap-around bonds included.

Proof. First let C_0, C_1 be positive definite. Put $C_t = (1-t)C_0 + tC_1$ and $D = C_1 - C_0 \ge 0$. Differentiating the Gaussian density, $\partial_t\langle G\rangle_{C_t} = \frac12 \sum_\alpha d_\alpha \langle G;\, k_\alpha(A)^2\rangle_{C_t}$, where $D = \sum_\alpha d_\alpha v_\alpha v_\alpha^{\mathsf{T}}$ with $d_\alpha \ge 0$, $k_\alpha(A) = v_\alpha^{\mathsf{T}} C_t^{-1} A$, and $\langle G; K\rangle := \langle GK\rangle - \langle G\rangle\langle K\rangle$; the normalisation constant cancels. By [8, Thm. 6.2(1)] (Lemma B.11(1) on the torus graph) the truncated correlation is ≤ 0 for $G \in \mathcal{I}$, and by [8, Thm. 6.2(3)] (Lemma B.11(3)) it is ≥ 0 for $G \in \mathcal{D}$; integrate over $t \in [0,1]$. For positive semidefinite C_i replace C_i by $C_i + \eta$ and let $\eta \downarrow 0$. Here $\int G\, d\gamma_C = \mathbb{E}[G(C^{1/2}\xi)]$, with ξ standard normal, is continuous in C by dominated convergence, since for the two classes the φ-integral of $|G|e^{-S_m}$ is bounded by a constant times $e^{a|A|}$, by (P2) and $|\cos| \le 1$. □

Lemma 7.6 ($\mu_0 \downarrow 0$ at fixed lattice; the elementary observables)**.** *Let $G \in \mathfrak{F}$, and let $\delta > 0$ with*

$$\sup_{0 < \mu_0 \le \mu_*} \langle |G|^{1+\delta}\rangle_{\varepsilon, T, \mu_0} < \infty.$$

Then $\langle|G|\rangle_{\varepsilon,T,\flat} < \infty$ *and* $\lim_{\mu_0\downarrow 0}\langle G\rangle_{\varepsilon,T,\mu_0} = \langle G\rangle_{\varepsilon,T,\flat}$*, along every sequence. The hypothesis holds, with a bound that does not depend on* μ_0*, for every finite product of: bounded elements of* $\mathfrak{F}$*;* $e^{tF(\mathbf{g})}$*,* $t\in\mathbb{R}$*;* $e^{X_{\mathrm{R}}(q)}$ *with* $q\geq 0$*;* $e^{-X_{\mathrm{R}}(q)}$ *with* $q\geq 0$*; polynomials in the* $F(\mathbf{g}_i)$ *and* $X_{\mathrm{R}}(h_k)$*; and the moduli of the exponentials* $\exp[\sum_i z_i F(\mathbf{g}_i) + \sum_k w_k X_{\mathrm{R}}(h_k)]$*. We call these the* elementary observables*.*

Proof. The first statement is Lemma 2.8, since $\mathfrak{F} = \mathfrak{A}_\varepsilon$; its proof gives $\langle|G|^{1+\delta}\rangle_{\varepsilon,T,\flat} \leq \sup_{\mu_0}\langle|G|^{1+\delta}\rangle_{\varepsilon,T,\mu_0}$, hence $\langle|G|\rangle_{\varepsilon,T,\flat} < \infty$. For the elementary observables, by Hölder's inequality it suffices to bound each factor, raised to a power $s\geq 1$, uniformly in μ_0. Bounded factors are trivial. For $e^{tF(\mathbf{g})}$, Lemma 7.3 with $k=0$ gives $\langle e^{stF(\mathbf{g})}\rangle_{\mu_0} \leq e^{\frac{1}{2}s^2t^2\|\mathbf{g}\|^2_{2,\varepsilon}}$. For $e^{X_{\mathrm{R}}(q)}$ with $q\geq 0$: $e^{sX_{\mathrm{R}}(q)}$ is a positive constant times an element of the class $\mathcal{I}$ of Lemma 7.5 ($g = s\varepsilon^3 q$, $\ell = 0$), and $C_{\mu_0} \geq C_{\mu_*}$ as forms for $\mu_0 \leq \mu_*$, the zero mode included (Lemma B.14(b)). So Lemma 7.5 gives $\langle e^{sX_{\mathrm{R}}(q)}\rangle_{\mu_0} \leq \langle e^{sX_{\mathrm{R}}(q)}\rangle_{\mu_*} < \infty$, the same b_T being used on both sides (P1). For $e^{-X_{\mathrm{R}}(q)}$ with $q\geq 0$ the pointwise bound $e^{-X_{\mathrm{R}}(q)} \leq e^{(W_\varepsilon + z_\varepsilon)\sum_x \varepsilon^3 q(x)}$ holds, since $r^2\geq 0$. Polynomials are bounded by sums of products of such exponentials: $|x|^k \leq k!(e^x + e^{-x})$, and $|X_{\mathrm{R}}(h)| \leq X_{\mathrm{R}}(|h|) + 2|W_\varepsilon + z_\varepsilon|\sum_x \varepsilon^3|h|$. The modulus of a complex exponential is the real exponential with the real parts $\operatorname{Re} z_i$, $\operatorname{Re} w_k$, which is $e^{F(\mathbf{g})}e^{X_{\mathrm{R}}(k_+)}e^{-X_{\mathrm{R}}(k_-)}$ with $\mathbf{g} = \sum_i(\operatorname{Re} z_i)\mathbf{g}_i$ and $k_\pm$ the positive and negative parts of $\sum_k(\operatorname{Re} w_k)h_k$, a product of factors of the previous types. □

Consequently every closed property of the states at $\mu_0 > 0$ on elementary observables also holds for the state $\flat$: positivity (Proposition 7.9) and the symmetries (Lemma 7.13). A Feynman or Proca photon measure at $\mu_0 = 0$ does not exist on the torus, because the constant mode is not normalisable; the state $\flat$ is not such a measure, but the limit of Lemma 2.8.

7.2. **Lattice reflection positivity on tori.** *Scope.* The statements of this subsection hold for every even N (P4), every direction ν, both types of plane pair (through sites, and between lattice planes), every photon mass $\mu_0 \in (0, \mu_*]$, every value of the constant bare coefficient b_T (P1), and, through Lemma 7.6, for the state $\flat$. In particular planes through sites are covered (Lemma 7.8(i)); this is the case used in Section 9, because the reflection $x_\nu \mapsto -x_\nu$ in a coordinate plane through the origin maps every lattice $\varepsilon\mathbb{Z}^3$ of the family to itself. Positivity is proved for the Proca form of the photon measure, in the manner of [29], and transferred to King's Feynman form on gauge-invariant observables by Lemma 7.7; the Feynman measure itself is not reflection positive on functions of A that are not gauge invariant (Remark 7.10). Three routine points are used below without further comment. (1) The reduction to $c=0$, resp. $c=\varepsilon/2$, in the proof of Lemma 7.8 is the invariance of the Proca measure under the lattice translations: its action is a sum of translation-covariant terms whose coefficients do not depend on the site, by (P1). (2) In case (i) of Lemma 7.8 the integrals are finite: with the decomposition $S^{\mathrm{P}} = \bar{S}_+(\omega_0,\omega_+) + \bar{S}_+(\omega_0,\omega_-^\rho)$ of the proof, $\int d\omega_0\big(\int d\omega_+\,|G|\,e^{-\bar{S}_+(\omega_0,\omega_+)}\big)^2 \leq \|G\|^2_\infty \int d\omega_0\big(\int d\omega_+\, e^{-\bar{S}_+(\omega_0,\omega_+)}\big)^2 = \|G\|^2_\infty Z^{\mathrm{P}} < \infty$. (3) In case (ii) the coupling constants of the expansion are read off (7.1): a crossing plaquette contributes $\frac{1}{2}\varepsilon^3(dA)^2_p = \frac{\varepsilon}{2}(u_p(\xi) - u_p(\xi'))^2$, and a crossing bond contributes $-\varepsilon\,\varphi(b_-)\cdot U(\varepsilon e A_b)\varphi(b_+)$ to the action, besides the single-site terms $\frac{\varepsilon}{2}(r^2_{b_-} + r^2_{b_+})$. So $\gamma_\iota = \varepsilon > 0$ in both cases; the hopping term enters $-S$ with the positive sign, which is the only sign condition the expansion uses.

Reflections. Fix a direction ν and $c \in \frac{\varepsilon}{2}\mathbb{Z}$, and let $\rho = \rho_c$ be the reflection $x_\nu \mapsto 2c - x_\nu$ of T. Since N is even (P4), ρ fixes the pair of planes $\{x_\nu = c\}$ and $\{x_\nu = c + \ell_T/2\}$. These are planes between lattice planes when $c \in \frac{\varepsilon}{2} + \varepsilon\mathbb{Z}$, and planes through sites when $c \in \varepsilon\mathbb{Z}$. The induced map on fields is $(\hat\rho\varphi)_x = \varphi_{\rho x}$ and $(\hat\rho A)_b = \pm A_{\rho b}$: the sign is $-$ exactly for the bonds parallel to e_ν, whose orientation ρ reverses. $\hat\rho$ is an involution and a signed permutation of the coordinates, so it preserves Lebesgue measure. The action (7.1) is $\hat\rho$-invariant, by (P1) and because the plaquette, gradient, mass and hopping terms are mapped to terms of the same kind. $\hat\rho$ maps gauge transformations to gauge transformations: $\hat\rho(A + d\chi, U(-e\chi)\varphi) = (\hat\rho A + d(\chi\circ\rho), U(-e\chi\circ\rho)\hat\rho\varphi)$. Put $\theta G := \overline{G\circ\hat\rho}$.

On curvatures, $(d\hat\rho A)_p = \mp(dA)_{\rho p}$, with $-$ exactly for the plaquettes containing the direction ν; hence $F(\mathbf{g}) \circ \hat\rho = F(\rho^*\mathbf{g})$, where $\rho^*\mathbf{g}$ is the signed reflected form, and θ maps $\mathfrak{F}$ into itself. Let Λ_+ be the open half $\{c < x_\nu < c + \ell_T/2\}$ for planes between lattice planes, or the closed half $\{c \le x_\nu \le c + \ell_T/2\}$ for planes through sites. Let $\mathfrak{B}_+$ be the algebra of bounded Borel functions of $\varphi|_{\Lambda_+}$ and of A on the bonds with both ends in Λ_+, and $\mathfrak{F}_+ := \mathfrak{F}(\Lambda_+)$.

Lemma 7.7 (Feynman and Proca forms agree on gauge-invariant observables)**.** *Let $\mu > 0$, and let $\langle\cdot\rangle^{\mathrm{P}}$ be* (7.3) *with the photon form $\frac12\langle A,\, (-\Delta_\varepsilon + \mu^2)A\rangle_\varepsilon$ replaced by the Proca form $\frac12\langle A,\, (d^*d + \mu^2)A\rangle_\varepsilon$. If G is gauge invariant and $|G| \le c\, e^{a|A|}$ for some constants a, c, then $\langle G\rangle_{\varepsilon,T,\mu} = \langle G\rangle^{\mathrm{P}}$.*

This is the torus form of [8, Thm. 5.1, p. 270].

Proof. On the connected torus, the space of bond functions is $\ker d^* \oplus \operatorname{ran} d$, orthogonally, and $\ker d$ on functions is the constants. So $A = Z + d\chi$ uniquely, with $d^*Z = 0$ and $\chi \perp 1$, and $(Z,\chi) \mapsto A$ is linear and bijective, with constant Jacobian; moreover $dA = dZ$. Then $\langle A,\, d^*dA\rangle + \mu^2\langle A,\, A\rangle = \langle dZ,\, dZ\rangle + \mu^2\langle Z,\, Z\rangle + \mu^2\langle d\chi,\, d\chi\rangle$. The Feynman form adds $\langle d^*A,\, d^*A\rangle = \langle d^*d\chi,\, d^*d\chi\rangle$, which again depends on χ only. Both χ-forms are positive definite on $1^\perp$. Since Ge^{-S_m} is gauge invariant, $\int d\varphi\,(Ge^{-S_m})(Z + d\chi, \varphi) = \int d\varphi\,(Ge^{-S_m})(Z, U(e\chi)\varphi) = \int d\varphi\,(Ge^{-S_m})(Z,\varphi)$, by the rotation invariance of $d\varphi$. So the integral factorises into a Z-integral and a Gaussian χ-integral. The χ-factor is the same for G and for $G \equiv 1$, and cancels in the normalised expectation. Fubini's theorem applies by the bound on G, (P2) and $\mu > 0$. □

Lemma 7.8 (reflection positivity of the Proca torus measure)**.** *Let $\mu > 0$ and N even. For every $G \in \mathfrak{B}_+$ (complex valued, not necessarily gauge invariant), $\langle\theta(G)G\rangle^{\mathrm{P}} \ge 0$, in each of the two cases: (i) planes through sites; (ii) planes between lattice planes. Hence $(B,C) \mapsto \langle\theta(C)B\rangle^{\mathrm{P}}$ is a positive semidefinite Hermitian form on $\mathfrak{B}_+$.*

Proof. Translate so that $c = 0$ in case (i) and $c = \varepsilon/2$ in case (ii).

(i) Every bond either has both endpoints in Λ_+ or is the $\hat\rho$-image of such a bond; the bonds lying in the two fixed planes are fixed by $\hat\rho$. Every plaquette lies in the closure of one half. Split each term that lives in a fixed plane (plaquettes, masses and hopping inside the plane, the single-site term at the plane sites) into two equal halves. The Proca action has no term that couples the two sides other than through the plane variables ω_0; the gauge-fixing term $\langle d^*A,\, d^*A\rangle$, which would, is absent. Hence $S^{\mathrm{P}} = \bar S_+(\omega_0,\omega_+) + \bar S_+(\omega_0,\omega_-^\rho)$, where ω_-^ρ denotes the variables of the other half transported by $\hat\rho$. Therefore $Z^{\mathrm{P}}\langle\theta(G)G\rangle^{\mathrm{P}} = \int d\omega_0 \,\big|\int d\omega_+\, G(\omega_0,\omega_+)e^{-\bar S_+(\omega_0,\omega_+)}\big|^2 \ge 0$; the integrals are finite by (P2) and $\mu > 0$.

(ii) The crossing bonds $b \in B_0$ (parallel to e_ν, joining the two halves across one of the two planes) are mapped to themselves, with $A_b \mapsto -A_b$. Write each crossing variable as $a_b = \alpha_b - \alpha_b'$ with α_b, α_b' independent centred normal of variance $1/(2M)$, $M = \mu^2\varepsilon^3$. This uses up the crossing mass term $\frac12 Ma_b^2$, and the reflection becomes the swap $\alpha \leftrightarrow \alpha'$. Put $\xi = (\omega_+,\alpha)$ and $\xi' = (\omega_-^\rho,\alpha')$; their a priori laws are identical. Every remaining term of S^{P} is of one of four kinds: a term of $S_+(\xi)$; its mirror image $S_+(\xi')$; a crossing plaquette; or a crossing hopping term. For a crossing plaquette p, $(dA)_p = \varepsilon^{-1}(u_p(\xi) - u_p(\xi'))$ with the same real linear function u_p on both copies, so its contribution $\frac{\varepsilon}{2}(u_p(\xi) - u_p(\xi'))^2$ splits into a term in ξ, a term in ξ', and $-\varepsilon u_p(\xi)u_p(\xi')$. For a crossing hopping term, $\varphi(b_-)\cdot U(\varepsilon e(\alpha_b - \alpha_b'))\varphi(b_+) = \psi_b(\xi)\cdot\psi_b(\xi')$ with the same $\mathbb{R}^2$-valued function ψ_b on both copies. Hence $e^{-S'} = \mathcal{G}(\xi)\mathcal{G}(\xi')\exp\big(\sum_\iota \gamma_\iota Y_\iota(\xi)Y_\iota(\xi')\big)$ with $\gamma_\iota \ge 0$ and real Y_ι. Expanding the last exponential,

$$Z^{\mathrm{P}}\langle\theta(G)G\rangle^{\mathrm{P}} = K\sum_n \prod_\iota \frac{\gamma_\iota^{n_\iota}}{n_\iota!}\Big|\int d\xi\, G(\xi)\mathcal{G}(\xi)\prod_\iota Y_\iota(\xi)^{n_\iota}\Big|^2 \ge 0, \qquad K > 0.$$

The interchange of sum and integral is justified by $\sum_n \prod_\iota \gamma_\iota^{n_\iota} |Y_\iota Y'_\iota|^{n_\iota}/n_\iota! \le \exp(\frac{1}{2}\sum_\iota \gamma_\iota (Y_\iota^2 + Y_\iota'^2))$ and by $\int |G| \mathcal{G}\, e^{\frac{1}{2}\sum_\iota \gamma_\iota Y_\iota^2} d\xi < \infty$: the plaquette parts cancel exactly, $|\psi_\flat(\xi)|$ is a radius at a site next to a plane, and $\lambda > 0$. The statement on the form follows by sesquilinearity and polarisation. □

Proposition 7.9 (reflection positivity of the torus states)**.** *Let N be even, $\nu \in \{1,2,3\}$, and let the plane pair be of either type.*

(a) *For $\mu_0 \in (0, \mu_*]$ and every gauge-invariant Borel function G of the Λ_+-variables with $\langle |G|^2 \rangle_{\varepsilon,T,\mu_0} < \infty$: $\langle \theta(G) G \rangle_{\varepsilon,T,\mu_0} \ge 0$. Moreover $\sum_{i,k} \bar\gamma_i \gamma_k \langle \theta(G_i) G_k \rangle_{\varepsilon,T,\mu_0} \ge 0$ for every finite family of such G_i and every $\gamma \in \mathbb{C}^n$.*

(b) *For the state $\flat$ and every $G \in \mathfrak{F}_+$ with $\langle |G|^2 \rangle_{\varepsilon,T,\flat} < \infty$: $\langle \theta(G) G \rangle_{\varepsilon,T,\flat} \ge 0$, with the matrix form as in (a).*

(c) *Every exponential $\Theta = \exp[\sum_i z_i F(\mathbf{g}_i) + \sum_k w_k X_{\mathrm{R}}(h_k)]$, with $z, w \in \mathbb{C}$ and real sources, lies in $\mathfrak{F}$ and in L^2 of every state. If all sources are supported in Λ_+ (for each $\mathbf{g}_i$: the plaquettes on which it does not vanish have all their bonds in Λ_+), then $\Theta \in \mathfrak{F}_+$, so (a) and (b) apply to finite families of such Θ.*

Proof. (a) For bounded gauge-invariant $G \in \mathfrak{B}_+$, $\theta(G)G$ is bounded and gauge invariant, so $\langle \theta(G)G \rangle_{\varepsilon,T,\mu_0} = \langle \theta(G)G \rangle^{\mathrm{P}} \ge 0$ by Lemmas 7.7 and 7.8. For unbounded G put $G_n = G\mathbf{1}\{|G| \le n\}$: it is bounded, gauge invariant and measurable with respect to the Λ_+-variables, so $\langle \theta(G_n) G_n \rangle \ge 0$, and $|\theta(G_n)G_n| \le |\theta G||G|$. By the Cauchy–Schwarz inequality and the $\hat\rho$-invariance of the measure, $\langle |\theta G||G| \rangle \le \langle |G|^2 \rangle$, and dominated convergence gives the claim. The matrix form is (a) applied to $G = \sum_k \gamma_k G_k$. (b) For bounded $G \in \mathfrak{F}_+$, $\theta(G)G$ is a bounded element of $\mathfrak{F}$, so $\langle \theta(G)G \rangle_\flat = \lim_{\mu_0 \downarrow 0} \langle \theta(G)G \rangle_{\mu_0} \ge 0$ by Lemma 7.6 and (a). The state $\flat$ is $\hat\rho$-invariant on $\mathfrak{F}$: $\langle K \circ \hat\rho \rangle_\flat = \lim \langle K \circ \hat\rho \rangle_{\mu_0} = \lim \langle K \rangle_{\mu_0} = \langle K \rangle_\flat$ for bounded $K \in \mathfrak{F}$. So truncation and dominated convergence work in the state $\flat$ exactly as in (a). (c) $|\Theta|^2 = \exp[\sum_i 2 \operatorname{Re} z_i\, F(\mathbf{g}_i) + \sum_k 2 \operatorname{Re} w_k\, X_{\mathrm{R}}(h_k)]$ is integrable at fixed lattice: the Gaussian dominates $|F|$, and (P2) dominates $|w||h|r^2$. It is an elementary observable of Lemma 7.6, so it is in L^2 of the state $\flat$ too. □

Remark 7.10. The Gaussian photon measure $\gamma_{C_{\mu_0}}$ in Feynman form is not reflection positive on functions of A that are not gauge invariant. For instance, let b be a bond in direction e_ν with both ends in Λ_+. Then $\theta(A_b) = -A_{\rho b}$, and $\mathbb{E}_{\gamma_{C_{\mu_0}}}[\theta(A_b)A_b] = -\mathbb{E}_{\gamma_{C_{\mu_0}}}[A_{\rho b}A_b] < 0$, because by Lemma B.4(i) the covariance C_{μ_0} acts on the e_ν-components of A as the resolvent $(-\Delta_\varepsilon + \mu_0^2)^{-1}$ of the lattice Laplacian on functions, whose matrix elements are all positive. Proposition 7.9 uses only gauge-invariant observables, and that is all the paper needs.

7.3. **Exact lattice symmetries.**

Definition 7.11 (cells, lattice forms, push-forward)**.** An *oriented k-cell* of T_ε, $k \in \{0,1,2\}$, is a pair $c = (x; v_1 \wedge \cdots \wedge v_k)$. Here x is a site, the v_i are distinct signed basis vectors $\pm e_\mu$, and the cell is the cube $x + \varepsilon \sum_i [0,1] v_i$ with the orientation $v_1 \wedge \cdots \wedge v_k$. Two descriptions of the same cube with orientations of opposite sign give the cells c and $-c$. A *lattice k-form* is a real function ω on oriented k-cells with $\omega(-c) = -\omega(c)$. Sites are 0-cells, bonds 1-cells, plaquettes 2-cells, and the centre of c is $z(c) := x + \frac{\varepsilon}{2}\sum_i v_i$. A signed permutation matrix $R \in O_h$ acts on cells about the site 0 by $Rc := (Rx; Rv_1 \wedge \cdots \wedge Rv_k)$, and on forms by the push-forward $(R_*\omega)(c) := \omega(R^{-1}c)$. A continuum test function is *sampled* by $h_\varepsilon(x) := h(x)$ on sites, and a continuum 2-form $\mathbf{g}$ by $\mathbf{g}_\varepsilon(c) := \mathbf{g}_{z(c)}(v_1, v_2)$ on plaquettes $c = (x; v_1 \wedge v_2)$, after identifying T with the cell $[-\ell_T/2, \ell_T/2)^3$. For $c = (x; e_\mu \wedge e_\kappa)$ this is the sampling of $g_{\mu\kappa}$ at the plaquette centre of Definition 2.9, and $\mathbf{g}_\varepsilon$ is odd under reversal of the orientation because $\mathbf{g}_z$ is alternating.

Lemma 7.12 (push-forward commutes with d, with the pairing and with sampling)**.** *Let T be cubic, so that every $R \in O_h$ maps T_ε onto itself (about the site 0), and let $R \in O_h$.*

(i) *R maps oriented k-cells bijectively onto oriented k-cells, $R(-c) = -Rc$, and $R\,\partial c = \partial(Rc)$ for the oriented boundary. Hence $dR_* = R_* d$, $\langle R_*\omega, R_*\omega' \rangle_\varepsilon = \langle \omega, \omega' \rangle_\varepsilon$, and $d^* R_* = R_* d^*$.*

(ii) *With* $(\hat{R}A, \hat{R}\varphi) := (R_*A, R_*\varphi)$*:* $F(\mathbf{g}) \circ \hat{R} = F(R_*^{-1}\mathbf{g})$ *and* $X_{\mathrm{R}}(h) \circ \hat{R} = X_{\mathrm{R}}(R_*^{-1}h)$*, where* $R_*^{-1}h = h \circ R$*.*

(iii) *For continuum test functions supported in* B_ρ *with* $\ell_T > 2\rho + 2$*:* $(R_*h)_\varepsilon = R_*(h_\varepsilon)$ *and* $(R_*\mathbf{g})_\varepsilon = R_*(\mathbf{g}_\varepsilon)$*, where* $(R_*\mathbf{g})_z(v, w) := \mathbf{g}_{R^{-1}z}(R^{-1}v, R^{-1}w)$ *is the continuum push-forward of Subsection 2.5. There is no extra sign for improper* R*.*

(iv) *The action* (7.1) *and Lebesgue measure are* $\hat{R}$*-invariant. Hence* $\langle G \circ \hat{R}\rangle_{\varepsilon,T,\mu_0} = \langle G\rangle_{\varepsilon,T,\mu_0}$ *for every integrable* G *and every* $\mu_0 \in (0, \mu_*]$*, and for every bounded or nonnegative* $G \in \mathfrak{F}$ *when* $\mu_0 = \flat$*.*

Proof. (i) R is linear, permutes the signed basis vectors, and maps $\varepsilon\mathbb{Z}^3$ to itself. It also preserves the period lattice $\ell_T\mathbb{Z}^3$, so it acts on $T = (\mathbb{R}/\ell_T\mathbb{Z})^3$ and on T_ε, because $\ell_T/\varepsilon \in \mathbb{N}$. The image of the cube $x + \varepsilon\sum[0,1]v_i$ is $Rx + \varepsilon\sum[0,1]Rv_i$, with orientation $Rv_1 \wedge \cdots \wedge Rv_k$. Orientation reversal commutes with R, and so does taking the oriented boundary, because the boundary of a cube is determined by its vertices and its orientation, both of which R transports. So $(dR_*\omega)(c) = \omega(R^{-1}\partial c) = \omega(\partial R^{-1}c) = (R_*d\omega)(c)$, and R_* permutes the unoriented cells, which all carry the weight ε^3. This gives the pairing identity, and $d^*R_* = R_*d^*$ follows by adjunction, since $R_*^{-1} = (R^{-1})_*$.

(ii) $F(\mathbf{g})(R_*A) = \langle \mathbf{g},\, dR_*A\rangle_\varepsilon = \langle \mathbf{g},\, R_*dA\rangle_\varepsilon = \langle R_*^{-1}\mathbf{g},\, dA\rangle_\varepsilon$ by (i). The second identity is the change of summation variable $x \mapsto R^{-1}x$.

(iii) For scalars, $(R_*h)_\varepsilon(x) = h(R^{-1}x) = (R_*h_\varepsilon)(x)$. For a plaquette $c = (x; v_1 \wedge v_2)$, $R^{-1}c = (R^{-1}x; R^{-1}v_1 \wedge R^{-1}v_2)$ has centre $R^{-1}z(c)$. So $(R_*\mathbf{g}_\varepsilon)(c) = \mathbf{g}_\varepsilon(R^{-1}c) = \mathbf{g}_{R^{-1}z(c)}(R^{-1}v_1, R^{-1}v_2) = (R_*\mathbf{g})_{z(c)}(v_1, v_2) = (R_*\mathbf{g})_\varepsilon(c)$. Both sides use only the values of $\mathbf{g}$ on signed basis vectors, so no determinant appears. Supports in B_ρ with $\ell_T > 2\rho + 2$ keep the sampled functions away from the faces of the cell, so the identification of T with the cell commutes with R on them.

(iv) By Lemma 7.2(H-a) and (i), $\frac{1}{2}\langle A,\, (-\Delta_\varepsilon + \mu_0^2)A\rangle_\varepsilon = \frac{1}{2}(\|dA\|^2 + \|d^*A\|^2 + \mu_0^2\|A\|^2)$ is invariant under $A \mapsto R_*A$. For a bond b with endpoints $b_\pm$, $(D_{R_*A}R_*\varphi)(b) = \varepsilon^{-1}(U(\varepsilon e A_{R^{-1}b})\varphi(R^{-1}b_+) - \varphi(R^{-1}b_-)) = (D_A\varphi)(R^{-1}b)$. The modulus $|D_A\varphi(b)|$ does not depend on the orientation of b, because $U(-s) = U(s)^{-1}$ and U is orthogonal. So the hopping term is invariant. The single-site terms are invariant by (P1). $\hat{R}$ is a signed permutation of the coordinates, so it preserves Lebesgue measure. For $\flat$: R_* is unitary and leaves the constants invariant, so it commutes with $\Pi^\perp$; it maps the constants $\mathbf{1}_\mu$ to $\pm\mathbf{1}_{\pi(\mu)}$, π the permutation part of R. The Gaussian measure with covariance $(-\Delta_\varepsilon)^{-1}\Pi^\perp$ is invariant by (i). The cube $[0, P)^3$ is mapped onto another period cell of $P\mathbb{Z}^3$, because the period $P = 2\pi/(|e|\,\ell_T)$ is the same in the three directions of a cubic torus. By Lemma 2.6 the integrand is P-periodic in c, so the uniform average does not change. □

The symmetry statement. Lemma 7.12(ii)–(iv) give, for $R \in O_h$ about a site and lattice sources,

$$\Big\langle \prod_i F(R_*\mathbf{g}_i) \prod_k X_{\mathrm{R}}(R_*h_k)\Big\rangle_{\varepsilon,T,\mu_0} = \Big\langle \prod_i F(\mathbf{g}_i) \prod_k X_{\mathrm{R}}(h_k)\Big\rangle_{\varepsilon,T,\mu_0} \tag{7.4}$$

(apply (iv) to $G := \prod_i F(R_*\mathbf{g}_i)\prod_k X_{\mathrm{R}}(R_*h_k)$, for which $G\circ\hat{R} = \prod_i F(\mathbf{g}_i)\prod_k X_{\mathrm{R}}(h_k)$ by (ii)). With (iii), the same identity holds for sampled continuum sources. This is the input of Theorem 9.2(b). About an arbitrary point $y \in \mathbb{R}^3$ the cubic group acts by $R_y := \tau_y R \tau_{-y}$. In the continuum this gives invariance about every point once translation invariance holds (Theorem 9.2(a)); on the lattice it is exact for $y \in \varepsilon\mathbb{Z}^3$. The curvature F transforms as a 2-form, also under the reflections in O_h; the magnetic field $\star F$ would be a pseudovector. The reflection $\rho_\nu \in O_h$ acts on F exactly as the map $\hat{\rho}$ used for reflection positivity (Subsection 7.2), with the sign $-$ on the components $g_{\mu\nu}$.

Lemma 7.13 (translations, the cubic group, charge conjugation)**.** *Let* T *be cubic and* $\mu_0 \in (0, \mu_*] \cup \{\flat\}$*. The state* $\langle\cdot\rangle_{\varepsilon,T,\mu_0}$ *is invariant under:*

(i) *the lattice translations* τ_a*,* $a \in \varepsilon\mathbb{Z}^3$ *(mod* T*);*

(ii) *the cubic group* O_h*, the* 48 *signed permutations of the coordinates, acting about any site. An element* $R \in O_h$ *acts on* φ *as on a scalar and on* A *as on a 1-form:* $A_b \mapsto \pm A_{Rb}$*, with*

the sign − when R reverses the orientation of the bond. Hence it acts by $h \mapsto h \circ R^{-1}$ on $X_{\mathrm{R}}(h)$ and by $\mathbf{g} \mapsto R_*\mathbf{g}$ (signed push-forward of 2-forms) on $F(\mathbf{g})$;

(iii) *charge conjugation $(A, \varphi) \mapsto (-A, \bar{\varphi})$, which maps $F(\mathbf{g}) \mapsto -F(\mathbf{g})$ and $X_{\mathrm{R}}(h) \mapsto X_{\mathrm{R}}(h)$.*

Proof. For $\mu_0 > 0$: the Feynman form $\frac{1}{2}\langle A, (-\Delta_\varepsilon + \mu_0^2)A\rangle_\varepsilon$ is the componentwise lattice Laplacian plus a mass, and it is invariant under translations and under signed permutations of the components. The hopping term $|D_A\varphi|^2$ is invariant under (i) and (ii); when the orientation of a bond is reversed, $U(-s) = U(s)^{\mathsf{T}}$ exchanges the roles of the endpoints. The single-site term is invariant because b_T is site independent (P1). O_h maps T_ε to itself because T is cubic and the centre is a site; about another site, conjugate by the translations of (i). Lebesgue measure is invariant under all these maps. For (iii), the Gaussian measure is even and $|D_{-A}\bar{\varphi}| = |D_A\varphi|$. For $\flat$: the statements are closed under the limit of Lemma 7.6 on $\mathfrak{F}$. Directly: the constant mode c is translation invariant; O_h permutes and negates its components; charge conjugation negates it; and the uniform law on $[0, P)^3$ is invariant under all of these modulo P, because the period $P = 2\pi/(|e|\,\ell_T)$ is the same in the three directions of a cubic torus. □

For the limits of King's cube states [25], O_h-invariance would follow from the uniqueness of the infinite-volume limit, which is not known (Section 14); on the cubic tori the cubic group is an exact symmetry of every lattice state.

7.4. **The tube inclusion.** For real lattice sources $\mathbf{g}_1, \dots, \mathbf{g}_n$, $h_1, \dots, h_p$ on T_ε, the lattice generating functional G_{ε,T,μ_0} and the lattice tube $\mathcal{D}_{\sigma_+,H}$ are those of Subsection 6.6; G_{ε,T,μ_0} is an entire function of $\zeta \in \mathbb{C}^{n+p}$ (Proposition 7.9(c) and dominated convergence; Subsection 2.4). Put

$$\sigma_0 := \tfrac{1}{2}\min(\sigma_+, H), \qquad D_{\sigma_0} := \Big\{\zeta \in \mathbb{C}^{n+p} : \sum_k |\mathrm{Re}\, w_k| \|h_k\|_\infty < \sigma_0\Big\}.$$

Then $D_{\sigma_0} \subset \mathcal{D}_{\sigma_+,H}$, since $\|k_\pm(\zeta)\|_\infty \le \|k(\zeta)\|_\infty$. The base of the tube is open and convex, since $s \mapsto \|(\sum_k s_k h_k)_\pm\|_\infty$ is continuous and convex; so the tube is connected. The same holds for the tube $D_{\sigma_+,H}$ of Subsection 2.4, built from continuum sources, and for the set D_{σ_0} defined with the continuum norms.

Lemma 7.14 (the continuum tube lies in every sampled tube)**.** *Let $\mathbf{g}_1, \dots, \mathbf{g}_n$, $h_1, \dots, h_p$ be real continuum test functions supported in B_R, and let $\zeta \in \mathbb{C}^{n+p}$ lie in the tube $D_{\sigma_+,H}$ of Subsection 2.4, which is built from the continuum function $k(\zeta) = \sum_k (\mathrm{Re}\, w_k) h_k$. Then for every (ε, T) with $\ell_T > 2R + 2$, ζ lies in the lattice tube $\mathcal{D}_{\sigma_+,H}$ built from the sampled sources.*

Proof. Sampling (Definition 2.9) is evaluation at points, so it is linear and commutes with taking pointwise positive and negative parts: the sampled $k(\zeta)$ is $\sum_k (\mathrm{Re}\, w_k)(h_k)_\varepsilon$, and its positive and negative parts are the sampled $k_\pm(\zeta)$. A supremum over the sample points is at most the supremum over $\mathbb{R}^3$, so $\|(k_\pm)_\varepsilon\|_\infty \le \|k_\pm\|_\infty$. □

With Lemma 7.14, Theorem 6.8 applies at every point of the tube $D_{\sigma_+,H}$ of Theorem 3.1(i), on every (ε, T) with $\ell_T > 2R+2$. Theorem 6.8 is the only input from Sections 4–6 used in Sections 7–9.

8. Limits along subsequences

Throughout this section, "uniformly" means uniformly in ε ($K \ge K_0$), in $j \ge 1$, and in $\mu_0 \in (0, \mu_*] \cup \{\flat\}$. Every statement is proved from Theorem 6.8 and the lattice facts of Section 7. In the torus counterterm scheme the bare coefficient b_T changes with (ε, T), so a sequence of lattice states is a sequence of different models. None of the arguments below uses a relation between two members of the sequence. Each uses either a statement about one model, or a uniform bound passed to a limit, or an abstract compactness or holomorphy argument.

8.1. **Uniform bounds, compactness, tightness.** Continuum test functions are placed on the lattice by sampling (Definition 2.9), with the norms defined there. For $T = T_j$ the lattice Schwinger functions $S_{n,p}^{\varepsilon,T,\mu_0} := S_{n,p}^{\varepsilon,j,\mu_0}$ are the mixed moments of Subsection 2.4, the Taylor coefficients at 0 of G_{ε,T,μ_0}. For real test functions supported in B_R with $\ell_T > 2R+2$ they are finite, real, linear in each argument and symmetric within each field type (Lemma B.30(a)).

Proposition 8.1 (uniform multilinear bounds)**.** *Uniformly, for all real lattice 2-forms $\mathbf{g}_i$ and lattice functions h_k (in particular for sampled test functions),*

$$|S_{n,p}^{\varepsilon,T,\mu_0}(\mathbf{g};h)| \le e^{\mathsf{K}\sigma_0}\,\mathrm{e}^n (n!)^{1/2}\,(\mathrm{e}/\sigma_0)^p\,p!\,\prod_i \|\mathbf{g}_i\|_{2,\varepsilon}\prod_k N_\varepsilon(h_k),$$

where e *is Euler's number and $N_\varepsilon(h) = \max(\|h\|_{L^1,\varepsilon}, \|h\|_\infty)$.*

Proof. Fix $\vartheta \in (0,1)$ and consider the polydisc $|z_i| = \vartheta(\sqrt{n}\|\mathbf{g}_i\|_{2,\varepsilon})^{-1}$, $|w_k| = \vartheta\sigma_0/(pN_\varepsilon(h_k))$. On it $\|\sum_i (\operatorname{Re} z_i)\mathbf{g}_i\|_{2,\varepsilon} \le \sqrt{n}$, $\|k(\zeta)\|_\infty \le \vartheta\sigma_0 < \sigma_0$ and $\|k(\zeta)\|_{L^1,\varepsilon} \le \vartheta\sigma_0$. So the polydisc lies in $D_{\sigma_0} \subset \mathcal{D}_{\sigma_+,H}$, and by (6.5) $|G| \le e^{n/2+\mathsf{K}\sigma_0}$ there. The Cauchy estimate on the polydisc gives $|S_{n,p}| \le e^{n/2+\mathsf{K}\sigma_0}\vartheta^{-n-p}n^{n/2}(p/\sigma_0)^p\prod\|\mathbf{g}_i\|_{2,\varepsilon}\prod N_\varepsilon(h_k)$. Let $\vartheta \uparrow 1$ and use $e^{n/2}n^{n/2} \le \mathrm{e}^n(n!)^{1/2}$ and $p^p \le \mathrm{e}^p p!$. □

Proposition 8.2 (compactness; identification of the tube functional)**.** *Let $(\varepsilon_i, T_{j_i}, \mu_i)$ be a sequence with $K_i \to \infty$, $\mu_i \in (0,\mu_*] \cup \{\flat\}$, and either $\ell_{T_{j_i}} \to \infty$ or $T_{j_i} \equiv T$ fixed. Write $S_{n,p}^{(i)}$ and G_i for the corresponding lattice Schwinger functions and generating functionals. There is a subsequence along which:*

(a) *$S_{n,p}^{(i)}(\mathbf{g};h)$ converges for all n,p and all test functions in C_c (if $T_{j_i} \equiv T$ is fixed: all test functions supported in some ball B_R with $2R+2 < \ell_T$). The limit $S_{n,p}$ is multilinear and symmetric within each field type, and obeys the bound of Proposition 8.1 with the continuum norms $\|\mathbf{g}_i\|_2$, $N(h_k)$;*

(b) *for every finite family of real test functions (supported in a ball B_R with $2R+2 < \ell_T$ if T is fixed), $G_i \to G$ locally uniformly on $D_{\sigma_+,H}$. G is holomorphic there, its Taylor coefficients at 0 are the $S_{n,p}$, $|G(\zeta)| \le G(\operatorname{Re}\zeta)$, and G obeys* (6.5) *with continuum norms.*

Proof. (i) *Equicontinuity.* For supports in B_R and $\varepsilon \le 1$: $\|h\|_{L^1,\varepsilon} \le |B_{R+1}|\|h\|_\infty$ and $\|\mathbf{g}\|_{2,\varepsilon} \le (3|B_{R+1}|)^{1/2}\|\mathbf{g}\|_\infty$. Let $C_c^{(R)}$ be the space of continuous test functions with support in $\bar{B}_R$, with the supremum norm. By Proposition 8.1 and Lemma B.30(a), each $S_{n,p}^{(i)}$, restricted to $C_c^{(R)}$, is a multilinear form bounded uniformly in i (for every i for which sampling is defined, $\ell_{T_{j_i}} > 2R+2$), hence uniformly Lipschitz on bounded sets. (ii) *Diagonal argument.* $C_c^{(R)}$ is separable, as a subspace of the separable metric space $C(\bar{B}_R)$. Choose countable dense subsets $\mathcal{Q}_R \subset C_c^{(R)}$, $R \in \mathbb{N}$, and extract a diagonal subsequence along which $S_{n,p}^{(i)}$ converges on all tuples from $\bigcup_R \mathcal{Q}_R$, for all (n,p). Given a tuple with entries in $C_c^{(R)}$ and $\delta > 0$, choose entries of $\mathcal{Q}_R$ within δ in the supremum norm; by the uniform Lipschitz bound of (i) on $C_c^{(R)}$, the values of $S_{n,p}^{(i)}$ at the two tuples differ by $O(\delta)$ uniformly in i, so the values at the given tuple form a Cauchy sequence. Since every test function lies in some $C_c^{(R)}$, the convergence extends to all tuples (for fixed T, to all tuples with $2R+2 < \ell_T$). The identities expressing multilinearity and symmetry pass to the limit (Lemma B.30(b)). (iii) *Norms.* Riemann sums give $\|h\|_{L^1,\varepsilon} \to \|h\|_1$ and $\|\mathbf{g}\|_{2,\varepsilon} \to \|\mathbf{g}\|_2$; lattice supremum norms are at most continuum supremum norms. (iv) *Tube.* For a fixed finite family, (6.5) bounds $\{G_i\}$ uniformly on each compact subset of $D_{\sigma_+,H}$, because the tube condition is on the supremum norms of the sampled sources, which are at most the continuum ones (Lemma 7.14), and the lattice norms on the right side of (6.5) converge by (iii). By Montel's theorem in several variables, every subsequence has a further subsequence converging locally uniformly to a holomorphic function $\tilde{G}$. By Weierstrass's theorem its Taylor coefficients at 0 are the limits of those of G_i, which are lattice

Schwinger functions of the family and converge to the $S_{n,p}$ by (a). So $\tilde{G}$ is determined near 0, hence on the connected tube by the identity theorem, and the whole subsequence converges. The inequality $|G(\zeta)| \le G(\operatorname{Re}\zeta)$ and the bound pass to the limit. □

Compactness replaces King's monotone limits: no monotonicity in the volume and no theorem of King about sources is used. The price is that the limits exist along subsequences and are not shown to be unique. For a fixed torus ($\varepsilon \to 0$ only) the same argument gives continuum states with the same bounds, defined on the tuples of test functions supported in a ball B_R with $2R+2 < \ell_T$ (not on periodic test functions on $(\mathbb{R}/\ell_T\mathbb{Z})^3$, which would need a sampling of functions on the torus).

Proposition 8.3 (tightness; the limits are probability laws on distributions)**.** *Let $\Phi_i = (F_i, X_i)$ be the lattice random fields $\mathbf{g} \mapsto F(\mathbf{g})$, $h \mapsto X_{\mathrm{R}}(h)$ along a subsequence as in Proposition 8.2 with $\ell_{T_{j_i}} \to \infty$. Their laws ν_i are tight on $H^{-s'}_{\mathrm{loc}}(\mathbb{R}^3; \Lambda^2\mathbb{R}^3 \oplus \mathbb{R})$ for every $s' > 3$. Every weak limit point ν has the moments $\mathbb{E}_\nu[\prod_l \Phi(f_l)] = S_{n,p}(f)$ for $f_l \in C_c^\infty$, and its characteristic functional is G at imaginary arguments.*

Proof. Each Φ_i is a finite sum of weighted point masses (at sites, and at plaquette centres with a 2-form value), hence lies in H^{-s}_{loc} for $s > 3/2$. Fix R, a cutoff $\chi \in C_c^\infty(B_{R+1})$ with $0 \le \chi \le 1$ and $\chi = 1$ on B_R, the box $Q = [-R-1, R+1]^3$, and the orthonormal Fourier basis $(e_\alpha)_{\alpha\in\mathbb{Z}^3}$ of $L^2(Q)$. Then $N(\chi e_\alpha) \le C_R$ uniformly in α, and Proposition 8.1, applied to real and imaginary parts with $n + p = 2$, gives $\mathbb{E}|\Phi_i(\chi e_\alpha)|^2 \le C'_R$ for each component. Hence $\mathbb{E}\|\chi\Phi_i\|^2_{H^{-s}_{\mathrm{per}}(Q)} = \sum_\alpha (1+|\alpha|^2)^{-s}\mathbb{E}|\Phi_i(\chi e_\alpha)|^2 \le C'_R \sum_\alpha (1+|\alpha|^2)^{-s} < \infty$ for $s > 3/2$, uniformly in i. The embedding $H^{-s}_{\mathrm{per}}(Q) \hookrightarrow H^{-s'}_{\mathrm{per}}(Q)$ is compact for $s' > s$. By Chebyshev's inequality the laws of $\chi\Phi_i$ are tight for each R, and a diagonal choice over $R \in \mathbb{N}$ gives tightness on $H^{-s'}_{\mathrm{loc}}$. Prokhorov's theorem gives weak limit points. All moments $\mathbb{E}|\Phi_i(f)|^{2q}$ are bounded uniformly (Proposition 8.1), so the moments converge by uniform integrability. The exponential moments at small real parameters are bounded uniformly (Theorem 6.8), so the characteristic functional of ν is $G(i\,\cdot)$. □

Remark 8.4. The proof gives tightness on $H^{-s'}_{\mathrm{loc}}$ for every $s' > 3/2$; the statement with $s' > 3$ is the one used in Theorem 3.1(ii).

8.2. **Order of limits.**

Lemma 8.5 (diagonal choice)**.** *Let $\mathcal{Q} \subset C_c$ be a countable set of test functions such that, for every $R \in \mathbb{N}$, the elements of $\mathcal{Q}$ supported in B_{R+1} are dense, for the supremum norm, in the set of continuous functions supported in B_R. Let $\mathcal{T} = \{t_1, t_2, \dots\}$ enumerate the tuples $t = (n, p; \mathbf{g}_1, \dots, \mathbf{g}_n; h_1, \dots, h_p)$ with entries in $\mathcal{Q}$, and let $R(t)$ be the least $R \in \mathbb{N}$ with all supports in B_R. For $l \ge 1$ let $\Sigma^{(l)}$ be a Schwinger family that is the limit, on every tuple t with $2R(t)+2 < \varsigma_l$ (on all tuples if $\varsigma_l = \infty$), of the lattice Schwinger families $\Sigma^{(l,i)} := S^{\varepsilon_{l,i}, T_{l,i}, \mu_{l,i}}$ as $i \to \infty$, where $K_{l,i} \to \infty$, $\mu_{l,i} \in (0, \mu_*] \cup \{\flat\}$, and $\liminf_i \ell_{T_{l,i}} \ge \varsigma_l \in (0, \infty]$ with $\varsigma_l \to \infty$. Suppose $\Sigma^{(l)} \to \Sigma$ on all tuples. Then there are indices $i(l)$ such that $K_{l,i(l)} \to \infty$, $\ell_{T_{l,i(l)}} \to \infty$, the lattice families $\Sigma^{(l,i(l))}$ converge to Σ on all tuples of test functions in C_c, and, for every finite family of real test functions, their generating functionals converge to the generating functional of Σ locally uniformly on $D_{\sigma_+,H}$.*

Proof. Put $\varsigma'_l := \min(\varsigma_l - 1, l)$. For each l choose $i(l)$ with $K_{l,i(l)} \ge l$, $\ell_{T_{l,i(l)}} \ge \varsigma'_l$, and $|\Sigma^{(l,i(l))}(t_q) - \Sigma^{(l)}(t_q)| \le 1/l$ for every $q \le l$ with $2R(t_q) + 2 < \varsigma'_l$. These are finitely many conditions, each of which holds for all large i (the lattice values at t_q are defined once $\ell_T > 2R(t_q) + 2$, and they converge to $\Sigma^{(l)}(t_q)$). Fix q. Since $\varsigma'_l \to \infty$, the condition at t_q is imposed for all large l, and then $|\Sigma^{(l,i(l))}(t_q) - \Sigma(t_q)| \le 1/l + |\Sigma^{(l)}(t_q) - \Sigma(t_q)| \to 0$. Convergence on $\mathcal{T}$ extends to all tuples in C_c: for entries supported in B_R, approximated in the supremum norm by elements of $\mathcal{Q}$ supported in B_{R+1}, and for $\varepsilon \le 1$, Proposition 8.1 and step (i) of the proof of Proposition 8.2 show that the lattice families are uniformly Lipschitz on bounded subsets of the functions supported in $\bar{B}_{R+1}$, uniformly

in the (l,i) with $\ell_{T_{l,i}} > 2R+4$; Σ obeys the same bound; and a $\delta/3$ argument with the density of $\mathcal{Q}$ applies. Finally, the sequence $(\varepsilon_{l,i(l)}, T_{l,i(l)}, \mu_{l,i(l)})$ satisfies the hypotheses of Proposition 8.2 (with $\ell_T \to \infty$ and varying photon masses), so every subsequence of the generating functionals has a further subsequence converging locally uniformly on $D_{\sigma_+,H}$ to a holomorphic function whose Taylor coefficients at 0 are those of Σ; this function is the same for all subsequences, so the whole sequence converges. □

Proposition 8.6 (iterated limits are joint limit points)**.** *Give the set of families $\{S_{n,p}\}$ that obey the bound of Proposition 8.1 with continuum norms the topology of convergence on a fixed countable dense set of test-function tuples. On this uniformly bounded set the topology is metrisable and, by equicontinuity, equivalent to convergence on all tuples. Then:*

(i) *every iterated limit (first $\varepsilon \to 0$ at fixed T_j, then $j \to \infty$, at fixed μ_0) is a limit point of the joint family $(\varepsilon, T_j) \to (0,\infty)$ at that μ_0;*

(ii) *every limit, as $\mu_k \downarrow 0$, of such limit points at $\mu_0 = \mu_k$ is a limit point of the joint family $(\varepsilon, T_j, \mu_0) \to (0,\infty,0)$.*

Proof. (i) Apply Lemma 8.5 with $\Sigma^{(l)}$ the fixed-torus continuum limit on T_{j_l} and $\varsigma_l = \ell_{j_l}$; each $\Sigma^{(l)}$ exists by Proposition 8.2 with T fixed, since at fixed T_j every spacing ε_K, $K \geq K_0$, is admissible (P4). (ii) Apply Lemma 8.5 with $\Sigma^{(l)}$ the joint limit points at $\mu_0 = \mu_l$ and $\varsigma_l = \infty$. In both cases the limit is a limit of a sequence of lattice states of the joint family, which is all that Section 9 uses. □

The paper uses the limits in the following order. (1) At fixed lattice, $\mu_0 \downarrow 0$ is a full limit, and it gives the state $\flat$ (Lemmas 2.8 and 7.6). (2) At fixed $\mu_0 \in (0, \mu_*]$, and for $\flat$, the joint limit $(\varepsilon, T_j) \to (0, \mathbb{R}^3)$ is taken along subsequences (Proposition 8.2); by Proposition 8.6(i) the iterated order gives nothing new. (3) Afterwards $\mu_0 \downarrow 0$ is taken through a countable set Q_0, along one common (ε, T)-subsequence (Proposition 8.7). So $S^\flat$ is the "μ_0 first" order and S^0 the "μ_0 last" order. Proposition 8.7(c) compares them by one inequality on $E^{\mathcal{D}}$; no interchange of limits is claimed. Every property proved below is closed under the convergence of Proposition 8.6, so it holds for every limit point in every one of these orders.

8.3. **The limit $\mu_0 \downarrow 0$.** *A uniqueness set.* Fix real test functions $\mathbf{g}_1, \dots, \mathbf{g}_n$, $h_1, \dots, h_p$. Write each scalar source as $h_k = h_k^1 - h_k^2$ with continuous $h_k^1, h_k^2 \geq 0$ of compact support (for instance the positive and negative parts of h_k), and consider the generating functional of the larger family $(\mathbf{g}_1, \dots, \mathbf{g}_n; h_1^1, \dots, h_p^1, h_1^2, \dots, h_p^2)$ in the variables $(z,u,v) \in \mathbb{C}^{n+2p}$, that is, of

$$\Theta(z,u,v) = \exp\Big[\sum_i z_i F(\mathbf{g}_i) + \sum_k \big(u_k X_{\mathrm{R}}(h_k^1) - v_k X_{\mathrm{R}}(h_k^2)\big)\Big].$$

The generating functional of the original family at (z,w) is its value at (z,w,w), and the map $(z,w) \mapsto (z,w,w)$ maps the tube of the original family into the tube of the larger one, since both points have the same function k. For $\delta' > 0$ put

$$E^{\mathcal{D}}_{\delta'} := \mathbb{R}^n \times (-\delta', 0)^p \times (0, \delta')^p.$$

If $\delta' \sum_k (\|h_k^1\|_\infty + \|h_k^2\|_\infty) < H/2$, then $E^{\mathcal{D}}_{\delta'}$ is a nonempty open subset of the real points of the tube of the larger family, and on it $\Theta = e^{F(\sum_i x_i \mathbf{g}_i)} e^{-X_{\mathrm{R}}(q)}$ with $q = \sum_k (|u_k| h_k^1 + v_k h_k^2) \geq 0$ and $\|q\|_\infty < H/2$; so $E^{\mathcal{D}}_{\delta'}$ is contained in the set $E^{\mathcal{D}}$ of Subsection 2.4 for the larger family. A holomorphic function on the connected tube that vanishes on $E^{\mathcal{D}}_{\delta'}$ vanishes identically: its Taylor coefficients at a point of $E^{\mathcal{D}}_{\delta'}$ are real derivatives, hence zero.

Proposition 8.7 (the limit $\mu_0 \downarrow 0$ on the tube)**.** *Fix a countable set $Q_0 \subset (0, \mu_*]$ that accumulates at 0 (other accumulation points are allowed), for instance $Q_0 = \{\mu_* 2^{-k} : k \geq 0\}$ or a countable dense subset of $(0, \mu_*]$, and let (K_l, j_l) be any sequence with $K_l \to \infty$ and $j_l \to \infty$. There is a*

subsequence (ε_i, T_{j_i}) *of* $(\varepsilon_{K_l}, T_{j_l})$ *along which* $S^\mu := \lim_i \langle\cdot\rangle_{\varepsilon_i, T_{j_i}, \mu}$ *exists for every* $\mu \in Q_0$*, and* $S^\flat := \lim_i \langle\cdot\rangle_{\varepsilon_i, T_{j_i}, \flat}$ *exists, in the sense of Proposition 8.2. Along it, for every finite family of real test functions:*

(a) *for* $\zeta \in E^{\mathcal{D}}$*,* $\mu \mapsto G^\mu(\zeta)$ *is non-increasing on* Q_0*, and* $G^\mu(\zeta) \le G^\flat(\zeta)$*;*

(b) $S^0 := \lim_{\mu \downarrow 0,\, \mu \in Q_0} S^\mu$ *exists as a full limit within* Q_0*, that is, the same along every sequence in* Q_0 *tending to* 0*. The convergence of the generating functionals is locally uniform on* $D_{\sigma_+, H}$*, the Schwinger functions converge, and* S^0 *obeys Proposition 8.1 and* (6.5) *with continuum norms;*

(c) $G^0(\zeta) \le G^\flat(\zeta)$ *for* $\zeta \in E^{\mathcal{D}}$*.*

Proof. Common subsequence. Since $j_l \to \infty$, $\ell_{j_l} = j_l!\,\ell_0 \to \infty$. So for each fixed $\mu \in Q_0 \cup \{\flat\}$ the sequence $(\varepsilon_{K_l}, T_{j_l}, \mu)$ satisfies the hypotheses of Proposition 8.2. Enumerate $Q_0 \cup \{\flat\} = \{\mu^{(1)}, \mu^{(2)}, \dots\}$. Apply Proposition 8.2 to the given sequence with $\mu \equiv \mu^{(1)}$, then to the resulting subsequence with $\mu \equiv \mu^{(2)}$, and so on. The diagonal subsequence is a subsequence of each of these, so the limits exist along it for every $\mu \in Q_0 \cup \{\flat\}$.

(a) Fix $\zeta = (x, w) \in E^{\mathcal{D}}$ and a member $(\varepsilon, T) = (\varepsilon_i, T_{j_i})$ of the subsequence; put $\mathbf{g} := \sum_i x_i \mathbf{g}_i$ and $q := -k(\zeta) \ge 0$. By the linearity of sampling and of the fields, on T_ε

$$\Theta(\zeta) = e^{F(\mathbf{g}_\varepsilon)} e^{-X_{\mathrm{R}}(q_\varepsilon)} = c_\varepsilon\, e^{-r^2(g)} e^{\ell(A)}, \qquad g_x := \varepsilon^3 q_\varepsilon(x) \ge 0, \quad \ell(A) := \langle d^* \mathbf{g}_\varepsilon,\, A\rangle_\varepsilon,$$

with $c_\varepsilon := \exp\big((W_\varepsilon + z_\varepsilon) \sum_x \varepsilon^3 q_\varepsilon(x)\big) > 0$. By (P1), c_ε and the matter part S_m of (7.1) do not depend on the photon mass. The covariance C_μ is non-increasing in μ as a form, the zero mode included (Lemma B.14(b)). By Lemma 7.5 (class $\mathcal{D}$), $\langle\Theta(\zeta)\rangle_{\varepsilon,T,\mu} \le \langle\Theta(\zeta)\rangle_{\varepsilon,T,\mu'}$ for $\mu' \le \mu$. Letting $\mu' \downarrow 0$ and using Lemma 7.6 ($\Theta(\zeta)$ is elementary) gives $\langle\Theta(\zeta)\rangle_{\varepsilon,T,\mu} \le \langle\Theta(\zeta)\rangle_{\varepsilon,T,\flat}$. Since $E^{\mathcal{D}} \subset D_{\sigma_+, H}$, Proposition 8.2(b) gives convergence of both sides along the subsequence, and the inequalities pass to the limit. (The details are Lemma B.37.)

(b) First let $\zeta \in E^{\mathcal{D}}$. By (a), $\mu \mapsto G^\mu(\zeta)$ is non-increasing on Q_0, and it is bounded above by (6.5) with continuum norms (Proposition 8.2(b)), uniformly in μ. Put $L(\zeta) := \sup_{\mu \in Q_0} G^\mu(\zeta)$. Given $\eta > 0$, pick $\mu_\eta \in Q_0$ with $G^{\mu_\eta}(\zeta) > L(\zeta) - \eta$. Then $L(\zeta) - \eta < G^\mu(\zeta) \le L(\zeta)$ for every $\mu \in Q_0$ with $\mu \le \mu_\eta$. Every sequence in Q_0 tending to 0 eventually lies below μ_η, so it gives the limit $L(\zeta)$; such sequences exist because 0 is an accumulation point of Q_0. This is the only place where a property of Q_0 is used. Now apply this to the larger family of the uniqueness set above, at the points of $E^{\mathcal{D}}_{\delta'} \subset E^{\mathcal{D}}$. The family $\{G^\mu\}_{\mu \in Q_0}$ of its generating functionals is locally bounded on its tube, by (6.5) with continuum norms. By Montel's theorem and the identity theorem with the open real uniqueness set $E^{\mathcal{D}}_{\delta'}$ (the Vitali argument), every sequence $\mu_k \to 0$ in Q_0 has a subsequence along which G^{μ_k} converges locally uniformly on the tube to a holomorphic function equal to the monotone limit on $E^{\mathcal{D}}_{\delta'}$; this function does not depend on the sequence. Hence the whole family converges locally uniformly as $\mu \to 0$ in Q_0, and so does its restriction to the tube of the original family under $(z, w) \mapsto (z, w, w)$. By Weierstrass's theorem the Schwinger functions converge. The bounds of Proposition 8.1 and (6.5) hold for every S^μ with continuum norms and constants independent of μ (Proposition 8.2), so they pass to S^0.

(c) follows from (a) and (b). □

For $\Theta = \exp[F(\mathbf{g}) - X_{\mathrm{R}}(q)] \in \mathcal{D}$ put $S^{\mu_0}(\Theta) := G^{\mu_0}(\zeta)$, for any finite family and any $\zeta \in E^{\mathcal{D}}$ with $\Theta(\zeta) = \Theta$; this does not depend on the choice, because the lattice values depend only on $(\mathbf{g}, q)$, by linearity, and hence so do their limits (Lemma B.37). With this reading, Theorem 3.1(iv) is Proposition 8.7(a) and (c).

Remark 8.8 (μ_0 outside Q_0). Let $\mu \in (0, \mu_*] \setminus Q_0$, and let $S^\mu_\star$ be any limit point at photon mass μ along a further subsequence (Proposition 8.2). Then every selection of such limit points $S^\mu_\star$ converges to S^0 as $\mu \downarrow 0$ through $(0, \mu_*]$, in the sense of Proposition 8.7(b). Continuity of $\mu \mapsto S^\mu$ for $\mu > 0$ is not claimed.

Proof. The lattice inequality in the proof of Proposition 8.7(a) holds for any three photon masses $\mu' < \mu < \mu''$, and the limits at $\mu', \mu'' \in Q_0$ are the same along every further subsequence. Hence $G^{\mu''}(\zeta) \le G^{\mu}_\star(\zeta) \le G^{\mu'}(\zeta) \le G^0(\zeta)$ for $\zeta \in E^{\mathcal{D}}$ and $\mu' < \mu < \mu''$ in Q_0, the last inequality by monotonicity. Given $\eta > 0$, choose $\mu'' \in Q_0$ with $G^{\mu''}(\zeta) > G^0(\zeta) - \eta$ (Proposition 8.7(b)); for $\mu < \mu''$ there is $\mu' \in Q_0$ with $\mu' < \mu$, since Q_0 accumulates at 0, and then $|G^{\mu}_\star(\zeta) - G^0(\zeta)| < \eta$. So $G^{\mu}_\star \to G^0$ on $E^{\mathcal{D}}$, in particular on $E^{\mathcal{D}}_{\delta'}$ for the larger family, and the Vitali argument of the proof of Proposition 8.7(b) gives locally uniform convergence on the tube and the convergence of the Schwinger functions. □

Lemma 8.9 (the measure for S^0)**.** *Along the subsequence of Proposition 8.7, let $s' > 3$. There is a Borel probability measure ν^0 on $H^{-s'}_{\mathrm{loc}}(\mathbb{R}^3; \Lambda^2\mathbb{R}^3 \oplus \mathbb{R})$ such that $\mathbb{E}_{\nu^0}[\prod_l \Phi(f_l)] = S^0_{n,p}(f)$ for test functions in C^∞_c, where $\Phi(f) = F(\mathbf{g})$ for a 2-form $f = \mathbf{g}$ and $\Phi(f) = X_{\mathrm{R}}(h)$ for a scalar $f = h$, and such that the characteristic functional of ν^0 is G^0 at imaginary arguments.*

Together with Proposition 8.3 for S^μ ($\mu \in Q_0$) and $S^\flat$, Lemma 8.9 gives the last sentence of Theorem 3.1(ii).

Proof. The laws at $\mu > 0$. For $\mu \in Q_0$, Proposition 8.3, applied along the subsequence of Proposition 8.7, gives a Borel probability measure ν^μ on $H^{-s'}_{\mathrm{loc}}$. Its moments are S^μ on C^∞_c test functions, and its characteristic functional is $G^\mu(i\,\cdot)$.

Uniform second moments. Fix R, let $Q = [-R-1, R+1]^3$ and $(e_\alpha)_{\alpha\in\mathbb{Z}^3}$ be as in the proof of Proposition 8.3, with the cutoff χ there, $0 \le \chi \le 1$. Let f be one of the real functions $\chi\,\mathrm{Re}\,e_\alpha$, $\chi\,\mathrm{Im}\,e_\alpha \in C^\infty_c$, placed in one field component. Then $\|f\|_2 \le 1$, $\|f\|_\infty \le |Q|^{-1/2}$ and $\|f\|_1 \le |Q|^{1/2}$, so $N(f) \le |Q|^{1/2}$, since $|Q| \ge 1$. By Proposition 8.2(a), S^μ obeys the bound of Proposition 8.1 with continuum norms; its constant $e^{\mathsf{K}\sigma_0}\mathrm{e}^n(n!)^{1/2}(\mathrm{e}/\sigma_0)^p p!$ depends only on K and σ_0, which by Theorem 6.8 do not depend on $\mu_0 \in (0, \mu_*] \cup \{\flat\}$. With the moment identity of Proposition 8.3 this gives, for every $\mu \in Q_0$ and every α, $\mathbb{E}_{\nu^\mu}\Phi(f)^2 = S^\mu_2(f,f) \le C'_R := e^{\mathsf{K}\sigma_0}\max\big(\sqrt{2}\,\mathrm{e}^2,\, 2\mathrm{e}^2\sigma_0^{-2}|Q|\big)$, using $S^\mu_{2,0}(\mathbf{g},\mathbf{g}) \le \sqrt{2}\,e^{\mathsf{K}\sigma_0}\mathrm{e}^2\|\mathbf{g}\|^2_2$ for a 2-form component and $S^\mu_{0,2}(h,h) \le 2e^{\mathsf{K}\sigma_0}\mathrm{e}^2\sigma_0^{-2}N(h)^2$ for the scalar component. Hence $\mathbb{E}_{\nu^\mu}|\Phi(\chi e_\alpha)|^2 \le 2C'_R$ for each component, uniformly in $\alpha \in \mathbb{Z}^3$ and $\mu \in Q_0$. Fix $s \in (3/2, s')$. The map $u \mapsto \|\chi u\|^2_{H^{-s}_{\mathrm{per}}(Q)} = \sum_\alpha (1+|\alpha|^2)^{-s}|u(\chi e_\alpha)|^2 \in [0,\infty]$ is Borel on $H^{-s'}_{\mathrm{loc}}$, as a countable sum of continuous functions, and by Tonelli's theorem

$$\mathbb{E}_{\nu^\mu}\|\chi\Phi\|^2_{H^{-s}_{\mathrm{per}}(Q)} = \sum_\alpha (1+|\alpha|^2)^{-s}\,\mathbb{E}_{\nu^\mu}|\Phi(\chi e_\alpha)|^2 \le 2C'_R \sum_\alpha (1+|\alpha|^2)^{-s} < \infty,$$

uniformly in $\mu \in Q_0$.

Tightness and a limit point. The embedding $H^{-s}_{\mathrm{per}}(Q) \hookrightarrow H^{-s'}_{\mathrm{per}}(Q)$ is compact, and Chebyshev's inequality bounds $\nu^\mu(\|\chi\Phi\|_{H^{-s}_{\mathrm{per}}(Q)} > M)$ by $2C'_R\sum_\alpha(1+|\alpha|^2)^{-s}/M^2$. The diagonal choice over $R \in \mathbb{N}$ of the proof of Proposition 8.3 shows that $\{\nu^\mu\}_{\mu\in Q_0}$ is tight on $H^{-s'}_{\mathrm{loc}}$. Choose $\mu_k \in Q_0$ with $\mu_k \downarrow 0$; this is possible because Q_0 accumulates at 0. By Prokhorov's theorem a subsequence of (ν^{μ_k}) converges weakly to a probability measure ν^0.

Moments. For $f_l \in C^\infty_c$ the map $u \mapsto \prod_{l=1}^N u(f_l)$ is continuous on $H^{-s'}_{\mathrm{loc}}$. Also $\mathbb{E}_{\nu^\mu}|\Phi(f)|^{2N} = S^\mu_{2N}(f,\dots,f)$ is bounded uniformly in μ (Proposition 8.1, continuum form). Hölder's inequality then bounds $\mathbb{E}_{\nu^\mu}|\prod_l\Phi(f_l)|^2$ uniformly, so $\prod_l\Phi(f_l)$ is uniformly integrable along (ν^{μ_k}). Hence $\mathbb{E}_{\nu^0}[\prod_l\Phi(f_l)] = \lim_k S^{\mu_k}(f) = S^0(f)$, the last equality by Proposition 8.7(b).

Characteristic functional. For real t the point it lies in $D_{\sigma_+,H}$, since $\mathrm{Re}\,w = 0$, and $u \mapsto e^{i\sum_l t_l u(f_l)}$ is bounded and continuous. So $\mathbb{E}_{\nu^0}e^{i\sum_l t_l\Phi(f_l)} = \lim_k G^{\mu_k}(it) = G^0(it)$, again by Proposition 8.7(b). □

Alternatively, by Proposition 8.6(ii) and Lemma 8.5, S^0 is the limit of a sequence of lattice states $(\varepsilon_{l,i(l)}, T_{l,i(l)}, \mu_l)$ with $\ell_T \to \infty$ and varying photon masses; that sequence satisfies the hypotheses of Proposition 8.2, and the proof of Proposition 8.3 applies to it.

Proposition 8.10 (what is identified)**.** (a) *The generating functional of every limit point is determined by its Schwinger functions (Proposition 8.2(b)). S^0 is a monotone μ_0-limit along the fixed (ε, T)-subsequence, the same for every sequence in Q_0 tending to 0 (Proposition 8.7(b)). At fixed lattice the state $\flat$ is the full limit $\mu_0 \downarrow 0$ (Lemma 2.8).*

(b) *$G^0 \le G^\flat$ on $E^{\mathcal{D}}$ (Proposition 8.7(c)). If, for every finite family of real test functions and every $\zeta \in E^{\mathcal{D}}$,*

$$\text{(U-}\mu)_T: \qquad \lim_{\mu_0 \downarrow 0} \sup_{\varepsilon, j} |\langle \Theta(\zeta)\rangle_{\varepsilon, T_j, \flat} - \langle \Theta(\zeta)\rangle_{\varepsilon, T_j, \mu_0}| = 0,$$

the supremum running over the spacings and tori of the family on which the sources can be sampled, then $G^0(\zeta) = G^\flat(\zeta)$ for every $\zeta \in E^{\mathcal{D}}$; that is, $\mu_0 \downarrow 0$ commutes with the (ε, T)-limit on the class $\mathcal{D}$.

Proof. (a) is a summary of the results cited. (b) For $\mu \in Q_0$, $G^\flat(\zeta) - G^\mu(\zeta)$ is the limit along the subsequence of $\langle \Theta(\zeta)\rangle_{\varepsilon_i, T_{j_i}, \flat} - \langle \Theta(\zeta)\rangle_{\varepsilon_i, T_{j_i}, \mu}$ (Proposition 8.2(b)), so $|G^\flat(\zeta) - G^\mu(\zeta)|$ is at most the supremum in (U-μ)$_T$ at $\mu_0 = \mu$. Let $\mu \to 0$ in Q_0 and use Proposition 8.7(b). □

Neither (U-μ)$_T$ nor the equality $S^0 = S^\flat$ is needed for Theorem 3.1; both are discussed in Section 14.

9. OS0, OS1 for translations and the cubic group, OS2, OS3

Limit points. In this section a *limit point* is any of: S^{μ_0} for $\mu_0 \in Q_0$; $S^\flat$; S^0 (Proposition 8.7); a limit point at any $\mu_0 \in (0, \mu_*]$ along a further subsequence (Proposition 8.2); in every order of limits (Proposition 8.6). The fixed-torus continuum states are included where stated. Every limit point other than the fixed-torus states is a Schwinger family in the sense of Subsection 2.6: $S_{0,0} = 1$, because the lattice states are normalised; the $S_{n,p}$ are defined on all test functions in C_c, multilinear and symmetric within each field type (Proposition 8.2(a)); and they are real, because the lattice moments of the real fields $F(\mathbf{g})$, $X_{\mathrm{R}}(h)$ are real. For S^0 these identities pass to the limit of Proposition 8.7(b). The generating functional of a limit point on $D_{\sigma_+, H}$ is the limit G of Proposition 8.2(b), or of Proposition 8.7(b) for S^0.

9.1. OS0.

Theorem 9.1 (OS0)**.** *Every limit point S satisfies:*

(a) *(tube form) for every finite family of real test functions, G is holomorphic on $D_{\sigma_+, H}$, which is all of $\mathbb{C}$ in the coefficients z_i of the curvature, and*

$$|G(\zeta)| \le \exp\Big(\tfrac{1}{2}\Big\|\sum_i (\mathrm{Re}\, z_i)\mathbf{g}_i\Big\|_2^2 + \kappa_+ \|k_+(\zeta)\|_1 + 2S_\flat^B \|k_-(\zeta)\|_1\Big);$$

since $k_+(\zeta)$ and $k_-(\zeta)$ have disjoint supports and $\kappa_+, 2S_\flat^B \le \mathsf{K}$, the right side is at most $\exp\big(\frac{1}{2}\|\sum_i (\mathrm{Re}\, z_i)\mathbf{g}_i\|_2^2 + \mathsf{K}\|k(\zeta)\|_1\big)$, which is the tube form of OS0 in Definition 2.10 with $C = \mathsf{K}$;

(b) *(Schwinger functions) $|S_{n,p}(\mathbf{g}; h)| \le e^{\mathsf{K}\sigma_0} \mathrm{e}^n (n!)^{1/2} (\mathrm{e}/\sigma_0)^p p! \prod_i \|\mathbf{g}_i\|_2 \prod_k N(h_k)$, with* e *Euler's number;*

(c) *(regularity) let $|f|_4 := \sup_{x \in \mathbb{R}^3, |\alpha| \le 4} |(1+x^2)^2 D^\alpha f(x)|$ for scalar f, and $|\mathbf{g}|_4 := (\sum_{\mu<\kappa} |g_{\mu\kappa}|_4^2)^{1/2}$ for 2-forms. Then*

$$|S_{n,p}(\mathbf{g}; h)| \le \sigma_{n+p} \prod_i |\mathbf{g}_i|_4 \prod_k |h_k|_4, \qquad \sigma_M := e^{\mathsf{K}\sigma_0} C_0^M M!, \quad C_0 := \max\big(\mathrm{e}\pi/(2\sqrt{2}),\, \mathrm{e}\pi^2/\sigma_0\big).$$

$S_{n,p}$ extends uniquely from C_c^∞ to Schwartz test functions, complex test functions included, and defines a tempered distribution on $\mathbb{R}^{3(n+p)}$. For every $\beta > 1$ there is α_β with $\sigma_M \le \alpha_\beta (M!)^\beta$. This is the regularity condition E0″ of [28, (4.2), p. 287] *with $s = 4$ and any $\beta > 1$. For a 2-form with one component, $\mathbf{g} = f\,dx_\mu \wedge dx_\kappa$, one has $|\mathbf{g}|_4 = |f|_4$; so the bound, read on such test functions, is E0″ for the components $F_{\mu\kappa}(f) := F(f\,dx_\mu \wedge dx_\kappa)$ of the curvature together with X_{R}, which is the form in which Definition 2.10 states it.*

The constants are the same for S^{μ_0}, $S^\flat$ and S^0.

Proof. (a) Proposition 8.2(b) and Proposition 8.7(b), with the three-term form of (6.5). (b) Proposition 8.1, passed to the limit (Proposition 8.2(a)). (c) In $d = 3$, $\int (1+x^2)^{-2} dx = \pi^2$ and $\int (1+x^2)^{-4} dx = \pi^2/8$. Hence $\|f\|_1 \le \pi^2 |f|_4$, $\|f\|_2 \le \frac{\pi}{2\sqrt{2}} |f|_4$ and $\|f\|_\infty \le |f|_4$; so $N(h) \le \pi^2 |h|_4$, and $\|\mathbf{g}\|_2 \le \frac{\pi}{2\sqrt{2}} |\mathbf{g}|_4$ by summing over the three components. Insert these into (b) and use $(n!)^{1/2} p! \le (n+p)!$. The seminorm $|\cdot|_4$ is continuous on $\mathcal{S}(\mathbb{R}^3)$ and C_c^∞ is dense in $\mathcal{S}$, so $S_{n,p}$ extends by continuity. Complex test functions are handled by multilinearity, at the cost of a factor at most 2^{n+p}, and the kernel theorem gives the distribution. Finally $C_0^M/(M!)^{\beta-1} \to 0$. □

OS0 rests on Theorem 6.8 only. It does not use King's cube theorems, a decoupling into unit cubes, a comparison of a cube with a torus, or any monotonicity in the volume. In the tube, only the small amplitude σ_+ for positive sources is a limitation. Regularity in the stronger form of [19], for all complex w, would need a bound for positive sources of arbitrary size, which is not proved here.

9.2. **OS1.**

Theorem 9.2 (OS1: translations and the cubic group)**.** (a) *Every limit point is invariant under all translations $a \in \mathbb{R}^3$: $S_{n,p}(\tau_a \mathbf{g}; \tau_a h) = S_{n,p}(\mathbf{g}; h)$. The fixed-torus continuum states are defined on the tuples of test functions supported in a ball B_R with $2R + 2 < \ell_T$ (Proposition 8.2). They are locally translation invariant: $S_{n,p}(\tau_a \mathbf{g}; \tau_a h) = S_{n,p}(\mathbf{g}; h)$ for all such tuples and all $a \in \mathbb{R}^3$ with $2(R + |a|) + 2 < \ell_T$.*

(b) *Every limit point is invariant under the cubic group O_h (48 elements) about every point of $\mathbb{R}^3$, acting by $h \mapsto h \circ R^{-1}$ and $\mathbf{g} \mapsto R_* \mathbf{g}$ (push-forward of 2-forms, signs included).*

So every limit point is invariant under the group $\mathbb{R}^3 \rtimes O_h$, with F transforming as a 2-form and X_{R} as a scalar.

Proof. (a) Let $a \in \varepsilon_K \mathbb{Z}^3 = 2^{-K} \varepsilon_0 \mathbb{Z}^3$. For every i with $K_i \ge K$ we have $a \in \varepsilon_i \mathbb{Z}^3$, because the lattices are nested (P4). Sampling commutes with τ_a for test functions supported in B_R once $\ell_{T_{j_i}} > 2(R + |a|) + 2$, since no wrap-around occurs. By Lemma 7.13(i), $S^{(i)}(\tau_a f) = S^{(i)}(f)$ for these i. Passing to the limit gives invariance under the dense group $\mathbb{Z}[1/2]^3 \varepsilon_0$. For continuity, Proposition 8.2(a) and multilinearity give $|S(\tau_a f) - S(f)| \le C_{n,p} \sum_l \|\tau_a f_l - f_l\| \prod_{l' \ne l} \|f_{l'}\|$, with the norms $\|\cdot\|_2$ and $N(\cdot)$. Each of these norms is at most $C_R \|\cdot\|_\infty$ on functions supported in B_{R+1}, and $\|\tau_a f - f\|_\infty \to 0$ as $a \to 0$ for $f \in C_c$. Hence the limit points are invariant under $\mathbb{R}^3$. At a fixed torus the same argument applies to tuples supported in B_R and to a with $2(R+|a|)+2 < \ell_T$. In the continuity step, take $R' > R + |a|$ with $2R' + 2 < \ell_T$ and dyadic $a_k \to a$ with $|a_k| < R' - R$, so that all translates are supported in $B_{R'}$. The details are in Lemma B.33. By Corollary B.34 the identity holds more generally whenever both f and $\tau_a f$ are admissible. Invariance under $\mathbb{R}^3/\ell_T \mathbb{Z}^3$ holds for the periodic extension of Proposition B.35, not for the states themselves, whose domain the torus group does not preserve (Remark B.36). For S^0, invariance passes to the limit of Proposition 8.7(b). (b) A cubic symmetry $R \in O_h$ about the origin maps $\varepsilon_i \mathbb{Z}^3$ and the cubic torus T_{j_i} to themselves, and the origin is a site of every $\varepsilon_i \mathbb{Z}^3$. R maps sites to sites and plaquette centres to plaquette centres, with the orientation sign of R_*, so it commutes with sampling (Lemma 7.12(iii)). Lemma 7.13(ii), in the form (7.4), and the limit give invariance about the origin. About any other point, conjugate by the translations of (a). □

Remark 9.3 (cubic symmetry only). Theorem 9.2 is all the Euclidean invariance proved in this paper: the group is $\mathbb{R}^3 \rtimes O_h$, and reflection positivity is proved only about coordinate planes (Theorem 9.4). What is known about rotations of infinite order is Subsection 9.5, and the missing step is discussed in Section 14.

9.3. **OS2.**

Theorem 9.4 (OS2: reflection positivity about every coordinate plane)**.** *Let S be a limit point, $\nu \in \{1,2,3\}$, $c \in \mathbb{R}$, and θ_c the reflection in the plane $\{x_\nu = c\}$. It acts on scalar test functions by $h \mapsto h \circ \rho_{\nu,c}$ and on 2-forms by the signed reflection $\mathbf{g} \mapsto (\rho_{\nu,c})_*\mathbf{g}$ (the components $g_{\mu\kappa}$ with $\nu \in \{\mu,\kappa\}$ change sign), and antilinearly on coefficients.*

(a) *(matrix form for exponentials) Let $\Theta_a = \exp[\sum_l z_{al} F(\mathbf{g}_{al}) + \sum_l w_{al} X_{\mathrm{R}}(h_{al})]$, $a = 1,\dots,N'$, with all sources supported in $\{x_\nu > c\}$, $z_{al}, w_{al} \in \mathbb{C}$, and $\sum_l |\mathrm{Re}\, w_{al}| \|h_{al}\|_\infty < \sigma_0$ for every a. Then $\sum_{a,b} \bar\gamma_a \gamma_b S(\theta_c(\Theta_a)\Theta_b) \geq 0$ for all $\gamma \in \mathbb{C}^{N'}$.*

(b) *(Schwinger functions) $S(\theta_c(P)P) \geq 0$ for every polynomial P in the $F(\mathbf{g})$ and $X_{\mathrm{R}}(h)$ with sources supported in $\{x_\nu > c\}$.*

Both hold for S^{μ_0} (every $\mu_0 \in (0,\mu_*]$), for $S^\flat$ and for S^0.

Proof. (a) By Theorem 9.2(a), $S(\theta_c(\cdot)\,\cdot)$ with sources in $\{x_\nu > c\}$ equals $S(\theta_0(\cdot)\,\cdot)$ with the sources translated by $-ce_\nu$, so we may take $c = 0$. Let all sources be supported in $\{x_\nu > \delta\} \cap B_R$ for some $\delta, R > 0$. On the lattice use the pair of planes $\{x_\nu = 0\}$, $\{x_\nu = \ell_T/2\}$ through sites (P4); $\rho_{\nu,0}$ maps $\varepsilon_i\mathbb{Z}^3$ to itself, so sampling commutes with it. For $\varepsilon_i < \delta$ and $\ell_{T_{j_i}} > 2R + 2$, every sampled scalar source lies in Λ_+, and every plaquette that carries a sampled 2-form has all its bonds at $x_\nu \geq \delta - \varepsilon_i/2 > 0$. So $\Theta_a \in \mathfrak{F}_+$. Proposition 7.9(a)–(c) gives $\sum \bar\gamma_a \gamma_b \langle\theta(\Theta_a)\Theta_b\rangle_i \geq 0$ for every i, with $\mu_i \in (0,\mu_*]$ or $\flat$. Each entry $\langle\theta(\Theta_a)\Theta_b\rangle$ is the generating functional of the finite joint family $\{(\rho_{\nu,0})_*\mathbf{g}_{al}, h_{al}\circ\rho_{\nu,0}, \mathbf{g}_{bl}, h_{bl}\}$ at the point $\zeta^*_{ab} = (\bar z_a, z_b, \bar w_a, w_b)$. The reflected and the unreflected sources have disjoint supports, so the supremum norms of $k_\pm(\zeta^*_{ab})$ are at most the larger of the two amplitudes, hence $< \sigma_0 = \frac12 \min(\sigma_+, H)$; that is, ζ^*_{ab} lies in the tube of that family. Proposition 8.2(b), applied to the joint family along the same subsequence, gives the convergence of each entry: all Schwinger functions converge on all C_c tuples, and the generating functional is determined by them. Positive semidefiniteness of a finite Hermitian matrix is a closed condition. For S^0, the matrices at $\mu \in Q_0$ converge entrywise by Proposition 8.7(b). (b) P is in L^2 of every lattice state and lies in $\mathfrak{F}_+$ for the indices i above, so $\langle\theta(P)P\rangle_i \geq 0$ by Proposition 7.9. $\theta(P)P$ is a finite combination of mixed moments with reflected test functions, which converge by Proposition 8.2(a) and, for S^0, by Proposition 8.7(b). □

For the Schwartz form of OS2 let $\Phi = (F_{12}, F_{13}, F_{23}, X_{\mathrm{R}})$ be the four real field components, $\mathcal{V} := \Lambda^2\mathbb{R}^3 \oplus \mathbb{R}$ and $\mathcal{V}_\mathbb{C}$ its complexification, so that a test function of n variables is a map $\mathbb{R}^{3n} \to \mathcal{V}_\mathbb{C}^{\otimes n} \cong \mathbb{C}^{4^n}$. For $\nu \in \{1,2,3\}$ put $U_+ := \{x \in \mathbb{R}^3 : x_\nu > 0\}$ and

$$\Theta f(x_1,\dots,x_n) := \overline{(\vartheta^{\otimes n} f)(\rho_{\nu,0}x_1,\dots,\rho_{\nu,0}x_n)},$$

where ϑ is the diagonal matrix of the signs of the reflection on the four field components.

Lemma 9.5 (density and continuity for the Schwartz form of OS2)**.** *Let $U \subset \mathbb{R}^3$ be open, $n \geq 1$, and V a finite-dimensional complex vector space.*

(a) *For every $g \in C_c^\infty(U^n; V^{\otimes n})$ there are a compact set $K' \subset U^n$ and finite sums g_N of tensor products $f_1 \otimes \cdots \otimes f_n$, $f_l \in C_c^\infty(U;V)$, with $\operatorname{supp} g_N \subset K'$ and $D^\alpha g_N \to D^\alpha g$ uniformly for every multi-index α; in particular $g_N \to g$ in $\mathcal{S}(\mathbb{R}^{3n}; V^{\otimes n})$.*

(b) *Let $U = U_+$, and let $f \in \mathcal{S}(\mathbb{R}^{3n}; V^{\otimes n})$ vanish outside U_+^n. Then f is a limit in $\mathcal{S}$ of elements of $C_c^\infty(U_+^n; V^{\otimes n})$.*

(c) *Let S be a limit point. For every N there is a unique tempered distribution $\mathsf{S}_N \in \mathcal{S}'(\mathbb{R}^{3N}; (\mathcal{V}_\mathbb{C}^{\otimes N})^*)$ with $\mathsf{S}_N(f_1 \otimes \cdots \otimes f_N) = S_N(f_1,\dots,f_N)$ for $f_l \in C_c^\infty(\mathbb{R}^3; \mathcal{V}_\mathbb{C})$, and the*

form $Q(f_0,\dots,f_{n_0}) := \sum_{n,n'\le n_0} \mathsf{S}_{n+n'}(\Theta f_n \otimes f_{n'})$ *is continuous on* $\bigoplus_{n\le n_0} \mathcal{S}(\mathbb{R}^{3n};\mathcal{V}_{\mathbb{C}}^{\otimes n})$. *Here* $S_N(f_1,\dots,f_N)$ *denotes the moments of the four-component field* Φ*, finite sums of the mixed Schwinger functions* $S_{n,p}$ *with* $n+p=N$.

Proof. (a) The projections of $\operatorname{supp} g$ to the n factors are compact subsets of U; let $K_1 \subset U$ be a compact neighbourhood of their union, choose $\psi \in C_c^\infty(U)$ with $\psi = 1$ on K_1, and put $K' := (\operatorname{supp}\psi)^n$. Then $\psi^{\otimes n} = 1$ on $\operatorname{supp} g$, so $g = \psi^{\otimes n} g$. Let Q be an open cube of side ℓ containing $\operatorname{supp}\psi$, and regard g as a smooth $\ell\mathbb{Z}^{3n}$-periodic function. Its Fourier coefficients decay faster than every power, so the partial sums s_N of its Fourier series converge to g with all derivatives, uniformly on Q^n. Each term is $e^{2\pi i k\cdot x/\ell} v = \prod_l e^{2\pi i k_l\cdot x_l/\ell}\, v$ with $v \in V^{\otimes n}$, a finite sum of elementary tensors $v_1\otimes\cdots\otimes v_n$; hence $g_N := \psi^{\otimes n} s_N$ is a finite sum of tensor products of functions $\psi(x_l)e^{2\pi i k_l\cdot x_l/\ell} v_l \in C_c^\infty(U;V)$, with $\operatorname{supp} g_N \subset K'$, and $D^\alpha(g_N - g) = D^\alpha(\psi^{\otimes n}(s_N - g)) \to 0$ uniformly by the Leibniz rule. Uniform convergence of all derivatives with supports in a fixed compact set implies convergence in $\mathcal{S}$. (b) For $\delta > 0$ let $f_\delta(x_1,\dots,x_n) := f(x_1 - \delta e_\nu,\dots,x_n - \delta e_\nu)$. Translations act continuously on $\mathcal{S}$, so $f_\delta \to f$ as $\delta \downarrow 0$. Since f vanishes outside U_+^n, $\operatorname{supp} f_\delta \subset \{x : x_{l,\nu} \ge \delta\ \forall l\} \subset U_+^n$. With $\chi \in C_c^\infty(\mathbb{R}^{3n})$ equal to 1 on the unit ball and $\chi_R := \chi(\cdot/R)$, $\chi_R f_\delta \to f_\delta$ in $\mathcal{S}$ as $R\to\infty$, and $\operatorname{supp}\chi_R f_\delta$ is a compact subset of U_+^n. (c) By Theorem 9.1(c), $(f_1,\dots,f_N) \mapsto S_N(f_1,\dots,f_N)$ extends, by multilinearity in the real and imaginary parts and by continuity, to a multilinear form on $\mathcal{S}(\mathbb{R}^3;\mathcal{V}_{\mathbb{C}})^N$ that is bounded by $4^N\sigma_N\prod_l |f_l|_4$, with $|(\mathbf{g},h)|_4 := \max(|\mathbf{g}|_4,|h|_4)$ (a factor 2^N from the curvature and scalar parts of each f_l, and 2^N from the real and imaginary parts), hence continuous. By the Schwartz kernel theorem [31, §V.3] it is represented by a unique S_N as stated. The map $f \mapsto \Theta f$ (composition with the linear map $\rho_{\nu,0}^{\times n}$, multiplication by the constant sign matrix $\vartheta^{\otimes n}$, and complex conjugation) is a continuous antilinear map of $\mathcal{S}$ to itself, and $(f,g) \mapsto f\otimes g$ is a separately continuous, hence continuous, bilinear map $\mathcal{S}(\mathbb{R}^{3n})\times\mathcal{S}(\mathbb{R}^{3n'}) \to \mathcal{S}(\mathbb{R}^{3(n+n')})$ of Fréchet spaces. So Q is continuous. □

Proposition 9.6 (OS2 for positive-time Schwartz test functions)**.** *Let* $\mathcal{S}_+(\mathbb{R}^{3n})$ *denote the Schwartz functions* $f : \mathbb{R}^{3n} \to \mathcal{V}_{\mathbb{C}}^{\otimes n}$ *that vanish outside* $U_+^n = \{x : x_{1,\nu} > 0,\dots,x_{n,\nu} > 0\}$*; it contains the Schwartz functions with support in* U_+^n*. Then for every limit point* S *and every finite sequence* $f_n \in \mathcal{S}_+(\mathbb{R}^{3n})$*,* $n = 0,\dots,n_0$*,*

$$\sum_{n,n'} \mathsf{S}_{n+n'}(\Theta f_n \otimes f_{n'}) \ge 0.$$

Proof. By Theorem 9.4(b), applied to complex test functions by multilinearity, the inequality holds when every f_n is a finite sum of tensor products $f^{(1)}\otimes\cdots\otimes f^{(n)}$ with $f^{(l)} \in C_c^\infty(U_+;\mathcal{V}_{\mathbb{C}})$. Indeed, for such f_n the sum equals $S(\theta_0(P)P)$ with $P = \sum_n \Phi(f_n)$, where $\Phi(f^{(1)}\otimes\cdots\otimes f^{(n)}) = \prod_l \Phi(f^{(l)})$. Now let $f_n \in \mathcal{S}_+(\mathbb{R}^{3n})$. By Lemma 9.5(b), f_n is a limit in $\mathcal{S}$ of elements of $C_c^\infty(U_+^n;\mathcal{V}_{\mathbb{C}}^{\otimes n})$, and by Lemma 9.5(a) each of these is a limit in $\mathcal{S}$ of finite sums of tensor products of functions in $C_c^\infty(U_+;\mathcal{V}_{\mathbb{C}})$. By Lemma 9.5(c) the quadratic form is continuous, so positivity on these finite sums passes to $\mathcal{S}_+$. The time-ordered support condition of [27, 28] is contained in the condition defining $\mathcal{S}_+$. In [27, 28] the reflected function also reverses the order of the arguments; since each S_N is symmetric under all permutations of its arguments (Remark 9.9), this does not change the form, and the positivity of [28] follows. □

9.4. **OS3.**

Lemma 9.7 (symmetry and charge conjugation on the lattice)**.** *Fix* (ε,T) *and* $\mu_0 \in (0,\mu_*] \cup \{\flat\}$*. For a real test function* $f = (\mathbf{g},h)$*,* $\mathbf{g}$ *a 2-form and* h *a scalar, put* $\Phi(f) := F(\mathbf{g}) + X_{\mathrm{R}}(h)$ *(sampled on* T_ε*), and* $\mathsf{C}f := (-\mathbf{g},h)$*. For* $N \ge 0$ *and real* $f_1,\dots,f_N$ *the moments* $M_N(f_1,\dots,f_N) := \langle\prod_{l=1}^N \Phi(f_l)\rangle_{\varepsilon,T,\mu_0}$ *are finite and real, multilinear and symmetric under all permutations of* $(f_1,\dots,f_N)$*, and*

$$M_N(\mathsf{C}f_1,\dots,\mathsf{C}f_N) = M_N(f_1,\dots,f_N).$$

In particular $\langle \prod_{i=1}^{n} F(\mathbf{g}_i) \prod_{k=1}^{p} X_{\mathrm{R}}(h_k) \rangle_{\varepsilon,T,\mu_0} = 0$ *for odd* n*, and the lattice generating functional satisfies* $G_{\varepsilon,T,\mu_0}(-z,w) = G_{\varepsilon,T,\mu_0}(z,w)$ *for all* $(z,w) \in \mathbb{C}^{n+p}$*.*

Proof. $\prod_l \Phi(f_l)$ is a polynomial in the $F(\mathbf{g}_l)$ and $X_{\mathrm{R}}(h_l)$, hence an elementary observable, integrable in every state, the state $\flat$ included (Lemma 7.6). It is real, and the product of real random variables is commutative; sampling is linear, so Φ is linear in f and M_N is multilinear. By Lemma 7.13(iii), every state, $\flat$ included, is invariant under charge conjugation $\mathcal{C} : (A,\varphi) \mapsto (-A,\bar{\varphi})$: for $\mu_0 > 0$ the Gaussian measure is even and $S_m(-A,\bar{\varphi}) = S_m(A,\varphi)$; in the state $\flat$, $\mathcal{C}$ maps $A^{\perp} \mapsto -A^{\perp}$ and $c \mapsto -c$, and the uniform law on the period cell is invariant under $c \mapsto -c$ modulo P. So $\langle K \circ \mathcal{C} \rangle = \langle K \rangle$ for every integrable K. Since $F(\mathbf{g}) \circ \mathcal{C} = -F(\mathbf{g}) = F(-\mathbf{g})$ and $X_{\mathrm{R}}(h) \circ \mathcal{C} = X_{\mathrm{R}}(h)$, we get $\Phi(f) \circ \mathcal{C} = \Phi(\mathsf{C}f)$, and the displayed identity follows. Taking $f_i = (\mathbf{g}_i, 0)$ for $i \le n$ and $f_{n+k} = (0, h_k)$ shows that the mixed moment equals $(-1)^n$ times itself. Finally, $\exp[\sum_i z_i F(\mathbf{g}_i) + \sum_k w_k X_{\mathrm{R}}(h_k)]$ is integrable (Proposition 7.9(c)), and its composition with $\mathcal{C}$ is the same exponential with z replaced by $-z$. □

Theorem 9.8 (OS3; charge conjugation)**.** *For every limit point,* $S_{n,p}$ *is symmetric under permutations of the (field component, test function) pairs within each field type, real for real test functions, and* $S_{n,p} = 0$ *for odd* n*.*

Proof. Lattice moments of commuting real random variables are symmetric and real. By Lemma 7.13(iii) every lattice state, $\flat$ included, is invariant under charge conjugation, which maps $F(\mathbf{g}) \mapsto -F(\mathbf{g})$ and $X_{\mathrm{R}}(h) \mapsto X_{\mathrm{R}}(h)$. So $S^{(i)}_{n,p} = (-1)^n S^{(i)}_{n,p}$. All three properties are closed under the limits of Propositions 8.2 and 8.7. □

Remark 9.9 (OS3 in the form of [28]). Let S be a limit point, and let $S_N(f_1,\dots,f_N)$ be the moments of the four-component field $\Phi = (F_{12}, F_{13}, F_{23}, X_{\mathrm{R}})$. Each S_N is a finite sum of the mixed Schwinger functions $S_{n,p}$, $n+p = N$. By Theorem 9.8 and Lemma 9.7, passed to the limit (Propositions 8.2(a) and 8.7(b)), S_N is real on real test functions in C_c, symmetric under all permutations of its arguments (mixed field types included), and invariant under $f_l \mapsto \mathsf{C}f_l$; and $G^S(-z,w) = G^S(z,w)$ on the tube, since the tube is invariant under $z \mapsto -z$ (it restricts only $\operatorname{Re} w$) and G^S is the locally uniform limit of the lattice functionals there (Proposition 8.2(b)). By Theorem 9.1(c), S_N extends to a continuous multilinear form on $\mathcal{S}(\mathbb{R}^3;\mathbb{C}^4)^N$. Symmetry and C-invariance pass to the extension by continuity, and the reality condition takes the form $S_N(\bar{f}_1,\dots,\bar{f}_N) = \overline{S_N(f_1,\dots,f_N)}$, by expanding each f_l into its real and imaginary parts. This is the symmetry axiom of [27, 28] for a boson field with several components; the extension of [28] from one scalar field to several fields is [28, Remark 5, p. 283].

9.5. **Rotations of infinite order.** Theorem 9.2 gives invariance under $\mathbb{R}^3 \rtimes O_h$ only. This subsection collects three elementary facts about the full orthogonal group $\mathrm{O}(3)$, which are used in Section 14.

Lemma 9.10 (rational rotations)**.**

(i) *If* $R \in \mathrm{O}(3)$ *and* $2^j R$ *has integer entries for some* $j \ge 0$*, then* $R \in O_h$*.*

(ii) *The matrix*

$$R_3 := \frac{1}{3}\begin{pmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ -2 & 2 & -1 \end{pmatrix}$$

lies in $\mathrm{SO}(3)$*,* $3R_3$ *is integral,* $R_3(3\varepsilon\mathbb{Z}^3) \subset \varepsilon\mathbb{Z}^3$ *with index* 27*,* R_3 *is the rotation about the axis* $(1,1,0)$ *by the angle* ϑ *with* $\cos\vartheta = -\frac{1}{3}$*, and* R_3 *has infinite order. The same holds for the rotation* $R_{12}(\theta_0)$ *by the angle* $\theta_0 = \arctan(3/4)$ *in the* $(1,2)$*-plane (King's rotation* [25, p. 326]*), with* $5R_{12}(\theta_0)$ *integral and* $\cos\theta_0 = \frac{4}{5}$*.*

(iii) *No lattice* $\Gamma \subset \mathbb{R}^3$ *satisfies* $R\Gamma = \Gamma$ *for a rotation* R *of infinite order.*

Proof. (i) Each column v of $2^j R$ is an integer vector with $|v|^2 = 4^j$. If $j \geq 1$, then $v_1^2 + v_2^2 + v_3^2 \equiv 0 \pmod 4$; squares are $\equiv 0$ or $1 \pmod 4$, so all v_i are even and $v/2$ is an integer vector of norm 2^{j-1}. Descending, the columns of R are integer unit vectors $\pm e_i$, so R is a signed permutation matrix, that is, $R \in O_h$. (ii) The rows of $3R_3$ have squared norm 9 and are pairwise orthogonal, and $\det(3R_3) = 27$, so $R_3 \in \mathrm{SO}(3)$. $\operatorname{tr} R_3 = \frac{1}{3} = 1 + 2\cos\vartheta$ gives $\cos\vartheta = -\frac{1}{3}$, and $(R_3 - I)(1,1,0)^{\mathsf{T}} = 0$. The index of $(3R_3)\mathbb{Z}^3$ in $\mathbb{Z}^3$ is $|\det 3R_3| = 27$. If $R_3^n = I$ for some $n \geq 1$, the eigenvalues $e^{\pm i\vartheta}$ are roots of unity, hence algebraic integers, so $2\cos\vartheta = e^{i\vartheta} + e^{-i\vartheta}$ is an algebraic integer; being rational, it is an integer; but $2\cos\vartheta = -\frac{2}{3}$. The same argument applies to $R_{12}(\theta_0)$, with $2\cos\theta_0 = \frac{8}{5}$. (iii) If $R\Gamma = \Gamma$, the matrix of R in a basis of Γ is an integer matrix, and so are all its powers. These lie in the intersection of $GL_3(\mathbb{Z})$ (conjugated by the basis matrix) with the compact group $\mathrm{O}(3)$, which is discrete and compact, hence finite. So R has finite order. □

Remark 9.11. Part (i) shows that a comparison with rotated lattices cannot be made inside the dyadic family $\varepsilon = 2^{-K}\varepsilon_0$ of Theorem 3.1 in King's nested form: no rotation outside O_h maps a dyadic lattice into a dyadic refinement of itself. For blocking factor 3, the family of Theorem 3.5, the rotation R_3 maps $3\varepsilon_K\mathbb{Z}^3 = \varepsilon_{K-1}\mathbb{Z}^3$ into $\varepsilon_K\mathbb{Z}^3$, so the rotated lattice stays in the family.

Lemma 9.12 (continuity and closed stabilisers)**.** *Let S be a limit point. For all n, p and all test functions, the map $g \mapsto S_{n,p}(g_*\mathbf{g}_1, \dots, g_*\mathbf{g}_n; g_*h_1, \dots, g_*h_p)$ is continuous on* $\mathrm{O}(3)$*. Hence* $\mathrm{Stab}(S) := \{g \in \mathrm{O}(3) : S_{n,p}(g_*\,\cdot) = S_{n,p} \text{ for all } n, p\}$ *is a closed subgroup of* $\mathrm{O}(3)$*.*

Proof. By Theorem 9.1(b), $S_{n,p}$ is multilinear and bounded by $C_{n,p}\prod_i \|\mathbf{g}_i\|_2 \prod_k N(h_k)$, $N(h) = \max(\|h\|_1, \|h\|_\infty)$. For test functions supported in B_R, all these norms are bounded by $C_R\|\cdot\|_\infty$ on functions supported in B_{R+1}, and $\|g_*f - f\|_\infty \to 0$ as $g \to 1$ by the uniform continuity of f (for 2-forms, $(g_*\mathbf{g})(x) = \Lambda^2 g\, \mathbf{g}(g^{-1}x)$). Telescoping over the arguments gives continuity at 1, and continuity everywhere follows by applying this to the tuple $g_{0*}f$, since g_* preserves the norms and the supports in B_R. The stabiliser is a subgroup and, by continuity, closed. □

Lemma 9.13 (one rotation of infinite order suffices)**.**

(i) *Every closed subgroup of* $\mathrm{O}(3)$ *that contains O_h and is infinite equals* $\mathrm{O}(3)$*.*

(ii) *For every rotation R of infinite order, the closure of the group generated by R and O_h is* $\mathrm{O}(3)$*. In particular this holds for R_3 and for $R_{12}(\theta_0)$.*

(iii) *If a limit point S is invariant under one rotation R of infinite order, it is invariant under* $\mathrm{O}(3)$*.*

Inputs of the proof. Part (i) uses two standard facts. The first is the closed-subgroup theorem: a closed subgroup of a Lie group is an embedded Lie subgroup [35, Thm. 3.42]. The second is the identity $h(a \times b) = (\det h)\,(ha \times hb)$ for $h \in \mathrm{O}(3)$ and $a, b \in \mathbb{R}^3$. To prove it, note that for every $c \in \mathbb{R}^3$,

$$\langle h(a \times b),\, hc\rangle = \langle a \times b,\, c\rangle = \det(a, b, c) = (\det h)^{-1} \det(ha, hb, hc) = (\det h)\langle ha \times hb,\, hc\rangle,$$

since $(\det h)^{-1} = \det h$; and hc runs over $\mathbb{R}^3$. Hence $hX_\omega h^{-1}v = h(\omega \times h^{-1}v) = (\det h)\,(h\omega \times v)$, that is, $hX_\omega h^{-1} = X_{(\det h)h\omega}$, where $X_\omega v := \omega \times v$. The proof of (i) applies this only to rotations in O_h, for which $\det h = 1$. The infinite order of R_3 and of $R_{12}(\theta_0)$ is Lemma 9.10(ii), and the continuity used in (iii) is Lemma 9.12.

Proof. (i) Let $G \subset \mathrm{O}(3)$ be closed and infinite, with $O_h \subset G$. By the closed-subgroup theorem [35, Thm. 3.42], G is an embedded Lie subgroup; it is compact, and a compact Lie group of dimension 0 is finite, so its Lie algebra $\mathfrak{g} \subset \mathfrak{so}(3)$ is non-zero. $\mathfrak{g}$ is invariant under $\mathrm{Ad}(h)X = hXh^{-1}$, $h \in G$. Identify $\mathfrak{so}(3)$ with $\mathbb{R}^3$ by $\omega \mapsto X_\omega$; then $hX_\omega h^{-1} = X_{(\det h)h\omega}$. Let $0 \neq \omega \in \mathfrak{g}$, with $\omega_i \neq 0$. The rotations $\mathrm{diag}(1,-1,-1)$, $\mathrm{diag}(-1,1,-1)$, $\mathrm{diag}(-1,-1,1)$ lie in O_h and have determinant 1; adding ω to its image under the i-th of them gives $2\omega_i e_i \in \mathfrak{g}$, so $e_i \in \mathfrak{g}$. The cyclic permutation of the coordinates is a rotation in O_h and maps $e_1 \mapsto e_2 \mapsto e_3$, so $\mathfrak{g} = \mathbb{R}^3 = \mathfrak{so}(3)$. Hence the

identity component of G is the connected subgroup with Lie algebra $\mathfrak{so}(3)$, that is, SO(3), and $-I \in O_h$ gives $G = \mathrm{O}(3)$. (ii) The closure is a closed subgroup that contains O_h and the infinite set $\{R^n : n \in \mathbb{Z}\}$; apply (i). (iii) $\mathrm{Stab}(S)$ is closed (Lemma 9.12), contains O_h (Theorem 9.2(b)) and R; apply (ii). □

10. OS4: ergodic decomposition and clustering

This section proves Theorems 3.6 and 3.10. The argument for Theorem 3.6 has two layers. The first (Sections 10.1 and 10.3–10.7) is abstract: it uses a probability law μ on a Polish space of fields with three properties, (H1)–(H3) below, namely translation and coordinate-reflection invariance, a local structure, and reflection positivity about the coordinate planes. The cubic rotations are not used, and no input about the infrared behaviour of the model is used. The second layer (Section 10.2) shows that the laws of the limit states of Theorem 3.1 have these three properties. Section 10.8 uses, in addition, estimates on the lattice tori, and proves single-axis clustering for part of the observables.

Definition 10.1 (forms of clustering)**.** Let μ be a translation-invariant probability law of a field and $U(a)$ the induced translations on $L^2(\mu)$. We distinguish: (C1) *clustering in the mean*: Cesàro averages of $\mathbb{E}[G'\,U(a)G]$ over cubes converge to $\mathbb{E}[G']\mathbb{E}[G]$; (C2) *clustering under diagonal separation*: $\mathbb{E}[G'\,U(a)G] \to \mathbb{E}[G']\mathbb{E}[G]$ as $\min_j |a_j| \to \infty$; (C3) clustering at a power-law rate; (C4) clustering at an exponential rate (the Euclidean form of a mass gap).

We prove (C1) and (C2) on ergodic components, and nothing about (C3) and (C4). Clustering as the separation runs along a single coordinate axis is the cluster property OS4 of Definition 2.11; for laws with the properties (H1)–(H3) below it implies (C2) (Lemma 10.24), and Section 10.8 proves it for part of the observables. An exponential rate is not expected in the symmetric phase, where the photon is expected to be massless (Section 14).

10.1. **The abstract setting.** *Notation.* e_1, e_2, e_3 is the standard basis of $\mathbb{R}^3$. For $J \subset \{1,2,3\}$ put $V_J = \mathrm{span}\{e_j : j \in J\}$ and $J^c = \{1,2,3\} \setminus J$. For $j \in \{1,2,3\}$ and $c \in \mathbb{R}$ the open half spaces are $H_j^+(c) = \{x : x_j > c\}$ and $H_j^-(c) = \{x : x_j < c\}$. The reflection in the plane $\{x_j = 0\}$ is $\rho_j x = x - 2x_j e_j$, and $\rho_{j,c}$ is the reflection in $\{x_j = c\}$. A bounded open set is denoted K; $K \Subset U$ means that $\overline{K} \subset U$ is compact.

(H1) (*Space, actions, measure.*) X is a Polish space with a jointly continuous action $(a, x) \mapsto \tau_a x$ of $\mathbb{R}^3$ and three involutive homeomorphisms $\theta_1, \theta_2, \theta_3$ of X with $\theta_j \tau_a = \tau_{\rho_j a}\theta_j$. μ is a Borel probability measure on X invariant under every τ_a and every θ_j.

(H2) (*Local structure.*) To every open $U \subset \mathbb{R}^3$ is assigned a sub-σ-algebra $\Sigma(U)$ of the Borel σ-algebra $\mathcal{B}$ of X, such that: $\Sigma(U) \subset \Sigma(U')$ if $U \subset U'$; $\Sigma(U) = \bigvee_{K \Subset U} \Sigma(K)$; $\bigvee_K \Sigma(K) = \mathcal{B}$; each $\Sigma(U)$ is generated by a countable family of bounded continuous functions on X; and, with

$$U(a)G := G \circ \tau_{-a}, \qquad \Theta_j G := G \circ \theta_j, \tag{10.1}$$

$U(a)$ maps $\Sigma(U)$-measurable functions to $\Sigma(U+a)$-measurable ones and Θ_j maps $\Sigma(U)$-measurable functions to $\Sigma(\rho_j U)$-measurable ones.

(H3) (*Coordinate reflection positivity.*) For every j, every $c \in \mathbb{R}$ and every $F \in L^2(\Sigma(H_j^+(c)), \mu)$,

$$\mathbb{E}_\mu\big[\,\overline{F \circ \theta_{j,c}}\, F\,\big] \;\geq\; 0, \qquad \theta_{j,c} := \tau_{ce_j}\theta_j\tau_{-ce_j}. \tag{10.2}$$

By (H1), $a \mapsto U(a)$ is a unitary representation of $\mathbb{R}^3$ on $L^2(\mu)$, Θ_j is a unitary involution, and

$$\Theta_j U(a) = U(\rho_j a)\Theta_j; \quad \text{in particular } \Theta_j U(te_j) = U(-te_j)\Theta_j, \;\; \Theta_j U(y) = U(y)\Theta_j \;\; (y \perp e_j). \tag{10.3}$$

By translation invariance, (10.2) for $c = 0$ implies it for every c: if $F \in L^2(\Sigma(H_j^+(c)))$ then $U(-ce_j)F \in L^2(\Sigma(H_j^+(0)))$ and $\mathbb{E}[\overline{F \circ \theta_{j,c}}\, F] = \mathbb{E}[\overline{(U(-ce_j)F) \circ \theta_j}\; U(-ce_j)F]$.

Remark 10.2 (conventions). For a space of distributions we use $(\tau_a\Phi)(f) := \Phi(f(\cdot + a))$, the push-forward of Φ by $x \mapsto x + a$; then $U(a)[\Phi(f)] = \Phi(f(\cdot - a))$, whose test function is supported in $\operatorname{supp} f + a$, as (H2) requires.

10.2. **The limit states satisfy (H1)–(H3).** Lemma 10.3 realises each limit state as a probability law on a space X of distributions, and Proposition 10.4 verifies (H1)–(H3) there. Lemma 10.5 carries the result to the Hermite–Sobolev space on which Theorem 3.6 is stated, and proves that the law is the only one with the moments S.

Fix one of the limit states $S \in \{S^{\mu_0}\ (\mu_0 \in Q_0),\ S^\flat,\ S^0\}$ of Theorem 3.1. Let $\mathcal{V} := \Lambda^2\mathbb{R}^3 \oplus \mathbb{R} \cong \mathbb{R}^4$. A test function is $f = (\mathbf{g}, h) \in C_c^\infty(\mathbb{R}^3; \mathcal{V})$, and the field is $\Phi(f) := F(\mathbf{g}) + X_{\mathrm{R}}(h)$. We write $G_S(\zeta)$ for the generating functional of S on the tube (Section 8), $S_{n,p}$ for its Schwinger functions, and $c_S := e^{C\sigma_0}$ with C and σ_0 the constants of Theorem 3.1(iii).

Actions on test functions. For $a \in \mathbb{R}^3$, $(T_a f)(x) := f(x - a)$. For $g \in \mathrm{O}(3)$, $g_* f := ((g^{-1})^*\mathbf{g},\ h \circ g^{-1})$, the push-forward of a 2-form and of a scalar. For a coordinate reflection, $(\rho_{j*}\mathbf{g})_{\mu\nu}(x) = (-1)^{[\mu=j]+[\nu=j]}\, g_{\mu\nu}(\rho_j x)$: the components carrying the index j change sign. This is the continuum form of the gauge-covariant lattice reflection used for OS2 (Theorem 9.4).

The space. Fix $s > 3$ and let $X := H^{-s}_{\mathrm{loc}}(\mathbb{R}^3; \mathcal{V})$, the space of distributions Φ with $\chi\Phi \in H^{-s}(\mathbb{R}^3; \mathcal{V})$ for every $\chi \in C_c^\infty(\mathbb{R}^3)$, with the seminorms $\Phi \mapsto \|\chi_R \Phi\|_{H^{-s}}$, $R \in \mathbb{N}$, for a fixed family of cut-offs χ_R equal to 1 on the ball B_R. It is a separable Fréchet space, hence Polish. The actions are

$$(\tau_a\Phi)(f) := \Phi(T_{-a}f), \qquad (g\Phi)(f) := \Phi(g_*^{-1}f)\ (g \in O_h), \qquad (\theta_j\Phi)(f) := \Phi(\rho_{j*}f).$$

For an open $U \subset \mathbb{R}^3$ let $\Sigma(U)$ be the σ-algebra generated by the maps $\Phi \mapsto \Phi(f)$, $f \in C_c^\infty(U; \mathcal{V})$ real.

Lemma 10.3 (realisation)**.** *For each S as above there is exactly one Borel probability measure μ_S on X such that*

$$\mathbb{E}_{\mu_S}\big[e^{i\Phi(f)}\big] = G_S(if) \qquad \text{for every real } f \in C_c^\infty(\mathbb{R}^3; \mathcal{V}), \tag{10.4}$$

where $G_S(if)$ denotes the tube functional of the one-element family $\{f\}$ at the purely imaginary point. Its mixed moments are the Schwinger functions of S. For $S = S^{\mu_0}$ and $S = S^\flat$, μ_S is the weak limit on X of the lattice laws along the subsequence of Theorem 3.1; for $S = S^0$ it is the weak limit of μ_{S^μ} as $\mu \downarrow 0$ in Q_0.

Proof. Uniqueness. Let (f_n) be a countable subset of real $C_c^\infty(\mathbb{R}^3; \mathcal{V})$, dense in it for the inductive-limit topology. The map $\Phi \mapsto (\Phi(f_n))_n$ from X to $\mathbb{R}^{\mathbb{N}}$ is continuous and injective, so by the Lusin–Souslin theorem (Lemma J.1) the Borel σ-algebra of X is generated by the evaluations $\Phi \mapsto \Phi(f_n)$. The sets $\{\Phi : (\Phi(f_1), \dots, \Phi(f_N)) \in B\}$ form a π-system generating $\mathcal{B}$, and the law of each vector $(\Phi(f_1), \dots, \Phi(f_N))$ is determined by its characteristic function $t \mapsto \mathbb{E}\exp(i\sum_k t_k \Phi(f_k)) = G_S(i\sum_k t_k f_k)$ (linearity of Φ). Hence (10.4) determines μ_S.

Existence for S^{μ_0}, $S^\flat$. The lattice laws are tight on X along the subsequence, and every weak limit point has the Schwinger functions as moments and $G_S(i\cdot)$ as characteristic functional (Proposition 8.3; the imaginary points lie in the tube, whose condition bounds only real parts of the coefficients of the Wick square). By uniqueness all limit points coincide, so the whole subsequence converges.

Existence for S^0. By Theorem 9.1(b), whose constants are the same for S^{μ_0}, $S^\flat$ and S^0, the bound of the proof of Proposition 8.3, $\mathbb{E}_{\mu_{S^\mu}}|\Phi(\chi_R e_\alpha)|^2 \le C'_R$ for the Fourier basis e_α of a box containing B_{R+1}, holds with C'_R independent of $\mu \in Q_0$. The Hilbert–Schmidt argument of Proposition 8.3 gives tightness of $\{\mu_{S^\mu}\}_{\mu \in Q_0}$ on X. Let μ' be a weak limit point as $\mu \downarrow 0$. The functions $\Phi \mapsto e^{i\Phi(f)}$ are bounded and continuous on X, so $\mathbb{E}_{\mu'} e^{i\Phi(f)} = \lim G_{S^\mu}(if) = G_{S^0}(if)$ by the locally uniform convergence on the tube (Proposition 8.7(b)). By uniqueness $\mu' = \mu_{S^0}$, and the whole family

converges. Moments converge by uniform integrability, since all moments are bounded uniformly in μ. □

Proposition 10.4 ((H1)–(H3) for the limit states)**.** *For every S as above, $(X, \tau, \theta, \mu_S, \Sigma)$ satisfies (H1), (H2) and (H3). Moreover μ_S is invariant under the action of O_h about every point, and μ_S is the only Borel probability measure on X whose mixed moments on $C_c^\infty(\mathbb{R}^3; \mathcal{V})$ are the Schwinger functions of S.*

Proof. (H1). X is Polish. The map $(a, \Phi) \mapsto \tau_a\Phi$ is jointly continuous on $\mathbb{R}^3 \times X$: translations act on H^{-s}_{loc} by a family of operators that is equicontinuous for a in compacts and strongly continuous, and these two properties give joint continuity. Each θ_j is a linear homeomorphism of X with $\theta_j^2 = 1$, and $\theta_j\tau_a = \tau_{\rho_j a}\theta_j$ because ρ_j is linear. Invariance of μ_S: by (10.4) it suffices that $G_S(iT_af) = G_S(if)$ and $G_S(i\rho_{j*}f) = G_S(if)$. By Theorem 9.2 every Schwinger function of S is invariant under translations and under O_h (which contains ρ_j), and the tube functional is determined on the connected tube by its Taylor coefficients at 0 (Proposition 8.2(b)), so G_S is invariant; the same argument gives invariance under all of O_h about every point.

(H2). Monotonicity in U is clear, and $\Sigma(U) = \bigvee_{K \Subset U} \Sigma(K)$ because each $f \in C_c^\infty(U)$ has support in some $K \Subset U$. The Lusin–Souslin argument in the uniqueness part of Lemma 10.3 gives $\bigvee_K \Sigma(K) = \mathcal{B}$. $\Sigma(U)$ is generated by the countably many bounded continuous functions $\arctan\Phi(f_n)$, where (f_n) is a countable subset of $C_c^\infty(U; \mathcal{V})$ that is dense in each H^s-norm on each compact subset of U; indeed $\Phi(f)$ is the pointwise limit of $\Phi(f_{n_k})$ whenever $f_{n_k} \to f$ in H^s with supports in a fixed compact. The covariance: $U(a)[\Phi(f)] = \Phi(T_af)$ and $\operatorname{supp} T_af = \operatorname{supp} f + a$; $\Theta_j[\Phi(f)] = \Phi(\rho_{j*}f)$ and $\operatorname{supp}\rho_{j*}f = \rho_j \operatorname{supp} f$.

(H3). Let $c = 0$ (translation invariance, as noted after (10.3)). Let $\mathcal{E}_+$ be the linear span of the functions $e^{i\Phi(f)}$, f real in $C_c^\infty(H_j^+(0); \mathcal{V})$. For $E_k = e^{i\Phi(f_k)}$ one has $\overline{E_k \circ \theta_j} = e^{-i\Phi(\rho_{j*}f_k)}$, which is the antilinear reflection $\theta(E_k)$ of Theorem 9.4: the coefficient i is conjugated and the test function is reflected, with the sign of the j-components of the 2-form part. Theorem 9.4(a) (matrix reflection positivity of tube exponentials) applies to these E_k: their curvature coefficients are arbitrary complex numbers, and their Wick-square coefficients are purely imaginary, so the amplitude condition, which bounds only real parts, holds. Its matrix entries are expectations under μ_S: by (10.4) applied to $f_l - \rho_{j*}f_k$, $\mathbb{E}_{\mu_S}[\overline{E_k \circ \theta_j}\, E_l] = G_S(i(f_l - \rho_{j*}f_k))$, which is the entry $S(\theta_j(E_k)E_l)$ of the matrix of Theorem 9.4(a), both being limits of the same lattice expectations (Propositions 8.2(b) and 8.7(b)). Hence $\mathbb{E}_{\mu_S}[\overline{F \circ \theta_j}\, F] \geq 0$ for every $F \in \mathcal{E}_+$. $\mathcal{E}_+$ is an algebra of bounded continuous functions, closed under complex conjugation, containing the constants and generating $\Sigma(H_j^+(0))$, so it is dense in $L^2(\Sigma(H_j^+(0)), \mu_S)$ (Lemma J.2). If $F_n \to F$ in L^2 with $F_n \in \mathcal{E}_+$, then $\overline{F_n \circ \theta_j}F_n \to \overline{F \circ \theta_j}F$ in $L^1(\mu_S)$, because θ_j preserves μ_S. This gives (10.2).

Determination by S. By Lemma 10.3, μ_S is determined by $G_S(i\,\cdot)$, which is determined by the Schwinger functions (Proposition 8.2(b)). If μ' is any Borel probability measure on X with the same mixed moments, the argument of Lemma 10.5(c) applies verbatim on X, with the π-system of the proof of Lemma 10.3, and gives $\mu' = \mu_S$. □

The field space is $\mathcal{V} = \Lambda^2\mathbb{R}^3 \oplus \mathbb{R}$: the three components of the curvature and the renormalised Wick square. Nothing above uses more of the field content than the transformation rules of the two summands.

The Hermite–Sobolev space. $\mathcal{A} := -\Delta + |x|^2 + 1$ acts componentwise on $L^2(\mathbb{R}^3; \mathcal{V})$. Its eigenfunctions are $h_{\alpha,c} := h_\alpha e_c$, with h_α the Hermite functions, $\alpha \in (\mathbb{N} \cup \{0\})^3$, and $(e_c)_{c \leq 4}$ a basis of $\mathcal{V}$. The eigenvalues are $\lambda_\alpha = 2|\alpha| + 4$. For $r \in 2\mathbb{N}$ put $\|f\|_r := \|\mathcal{A}^{r/2}f\|_{L^2}$, let $\mathcal{H}^r$ be the completion of $\mathcal{S}(\mathbb{R}^3; \mathcal{V})$ in this norm, and let $\mathcal{H}^{-r}$ be its dual. So $\|\Psi\|_{-r}^2 = \sum_{\alpha,c} \lambda_\alpha^{-r} |\Psi(h_{\alpha,c})|^2$, and $\mathcal{H}^{-r}$ is a separable Hilbert space. The weighted norm of OS0 is $|f|_4 := \sup_{x, |\gamma| \leq 4} (1 + |x|^2)^2 |D^\gamma f(x)|$.

Lemma 10.5 (Hermite realisation and determinacy)**.** *Let S be as above, and let μ_S be the measure of Lemma 10.3 on $X = H^{-s}_{\mathrm{loc}}(\mathbb{R}^3; \mathcal{V})$, with $s := r$.*

(a) *For $k \in 2\mathbb{N}$ and $|\delta| + |\gamma| \le k$, $\|x^\delta \partial^\gamma f\|_{L^2} \le k^{k/2} \|f\|_k$. Hence $|f|_4 \le C\|f\|_{10}$ for $f \in \mathcal{S}$, and on $\mathcal{S}$ the norm $\|\cdot\|_k$ is equivalent to $\sum_{|\delta|+|\gamma|\le k} \|x^\delta \partial^\gamma \cdot\|_{L^2}$.*

(b) *Let $r \ge 14$ be even. There is exactly one Borel probability measure μ_S^H on $\mathcal{H}^{-r}$ whose image under the restriction map $\iota : \mathcal{H}^{-r} \to X$, $\iota\Psi := \Psi|_{C_c^\infty}$, is μ_S. The map ι is continuous, linear and injective, and $\mu_S(\iota\mathcal{H}^{-r}) = 1$. Moreover $\mathbb{E}_{\mu_S^H} e^{i\Psi(f)} = G_S(if)$, and $\mathbb{E}_{\mu_S^H} \prod_l \Psi(f_l) = S(f_1, \dots, f_N)$ for $f_l \in C_c^\infty$.*

(c) *μ_S^H is the only Borel probability measure on $\mathcal{H}^{-r}$ whose mixed moments on $C_c^\infty(\mathbb{R}^3; \mathcal{V})$ exist and equal those of S.*

(d) *Define the translations, the coordinate reflections and O_h on $\mathcal{H}^{-r}$ by the formulas used on X, and let $\Sigma^H(U) := \iota^{-1}\Sigma(U)$, the σ-algebra generated by $\Psi \mapsto \Psi(f)$, $f \in C_c^\infty(U; \mathcal{V})$. If $(X, \tau, \theta, \mu_S, \Sigma)$ satisfies (H1)–(H3), so does $(\mathcal{H}^{-r}, \tau, \theta, \mu_S^H, \Sigma^H)$, and μ_S^H is O_h-invariant about every point.*

Proof. (a) In each coordinate, $x_j = (a_j + a_j^*)/\sqrt{2}$ and $\partial_j = (a_j - a_j^*)/\sqrt{2}$, with $a_j h_\alpha = \sqrt{\alpha_j}\, h_{\alpha - e_j}$ and $a_j^* h_\alpha = \sqrt{\alpha_j + 1}\, h_{\alpha + e_j}$. So $x^\delta \partial^\gamma$ is a sum of $2^{|\delta|+|\gamma|}$ words of length $\ell := |\delta| + |\gamma|$ in the $a_j^\#$, each with coefficient $2^{-\ell/2}$. A word w shifts α by a fixed vector, so it maps distinct basis vectors to distinct basis vectors or to 0. Its coefficients are at most $(|\alpha| + \ell)^{\ell/2} \le (\ell\lambda_\alpha/2)^{\ell/2}$. Hence $\|wf\|^2 \le (\ell/2)^\ell \sum_\alpha \lambda_\alpha^\ell |\langle h_\alpha, f\rangle|^2 \le (\ell/2)^\ell \|f\|_k^2$ for $\ell \le k$, since $\lambda_\alpha \ge 1$, and $\|x^\delta \partial^\gamma f\| \le 2^{\ell/2}(\ell/2)^{\ell/2}\|f\|_k \le k^{k/2}\|f\|_k$. For the second claim, put $u := (1+|x|^2)^2 D^\gamma f$ with $|\gamma| \le 4$. Sobolev's embedding in $\mathbb{R}^3$ gives $\|u\|_\infty \le C \sum_{|\beta|\le 2} \|\partial^\beta u\|_{L^2}$. Each $\partial^\beta u$ is a finite sum of terms $x^\delta \partial^{\gamma'} f$ with $|\delta| \le 4$ and $|\gamma'| \le 6$. For the equivalence, $\mathcal{A}^{k/2}$ is a differential operator with polynomial coefficients of total degree $\le k$, which gives the reverse inequality.

(b) *A two-point bound on $\mathcal{S}$.* For $f = (\mathbf{g}, h) \in C_c^\infty$, Theorem 3.1(iii) and OS3 ($S_{1,1} = 0$) give $\mathbb{E}_{\mu_S} \Phi(f)^2 = S_{2,0}(\mathbf{g}, \mathbf{g}) + S_{0,2}(h, h) \le c_2(\|\mathbf{g}\|_2^2 + N(h)^2)$. Moreover $\|\mathbf{g}\|_2 + N(h) \le C|f|_4$, since $\int (1+|x|^2)^{-2} dx = \pi^2$ and $\int (1+|x|^2)^{-4} dx = \pi^2/8$. With (a), $\mathbb{E}_{\mu_S} \Phi(f)^2 \le C_2 \|f\|_{10}^2$. For $f \in \mathcal{S}$ and cut-offs χ_R ($\chi_R = 1$ on B_R, derivatives bounded uniformly in R), $\chi_R f \to f$ in every seminorm $\|x^\delta \partial^\gamma \cdot\|_{L^2}$, hence in $\|\cdot\|_{10}$. So $\Phi(\chi_R f)$ is Cauchy in $L^2(\mu_S)$. Call its limit $\tilde{\Phi}(f)$. It is linear in f and $\mathbb{E}\tilde{\Phi}(f)^2 \le C_2 \|f\|_{10}^2$.

The carrier. Fix Borel versions $Z_{\alpha,c}$ of $\tilde{\Phi}(h_{\alpha,c})$, and put $M := \sum_{\alpha,c} \lambda_\alpha^{-r} Z_{\alpha,c}^2$. Then $\mathbb{E}M \le 4C_2 \sum_\alpha \lambda_\alpha^{10-r} < \infty$, since there are $(n+1)(n+2)/2$ multi-indices with $|\alpha| = n$ and $r - 10 > 3$. On the Borel set $\{M < \infty\}$, which has full measure, define $\Psi(f) := \sum_{\alpha,c} \langle h_{\alpha,c}, f\rangle Z_{\alpha,c}$. By the Cauchy–Schwarz inequality this is a continuous linear functional on $\mathcal{H}^r$ with $\|\Psi\|_{-r}^2 = M$. The map $\Phi \mapsto \Psi$ is Borel, because its coordinates are. For $f \in C_c^\infty$ the partial sums of $\Psi(f)$ converge to $\tilde{\Phi}(f) = \Phi(f)$ in $L^2(\mu_S)$, since the Hermite expansion of f converges in $\|\cdot\|_{10}$. So $\Psi(f) = \Phi(f)$ a.s. For each n let $\mathcal{D}_n$ be a countable subset of $C_c^\infty(B_n; \mathcal{V})$ that is dense in the Sobolev norm H^r. Almost surely $\Psi(f) = \Phi(f)$ for all $f \in \bigcup_n \mathcal{D}_n$. Both sides are continuous on $C_c^\infty(B_n)$ for the H^r-norm. For Φ this holds because $|\Phi(f)| \le \|\chi_{n+1}\Phi\|_{H^{-s}} \|f\|_{H^s}$ and $s = r$. For Ψ it holds because $\|f\|_r \le C_n \|f\|_{H^r}$ for $\operatorname{supp} f \subset B_n$, by (a) and the boundedness of x^δ on B_n. Hence $\iota\Psi = \Phi$ a.s.

The map ι. For $\chi \in C_c^\infty$, $|\Psi(\chi f)| \le \|\Psi\|_{-r} \|\chi f\|_r \le C_\chi \|\Psi\|_{-r} \|f\|_{H^r}$, so $\chi\, \iota\Psi \in H^{-r} = H^{-s}$, and ι is continuous into X. It is injective, because C_c^∞ is dense in $\mathcal{S}$ and $\mathcal{S}$ is dense in $\mathcal{H}^r$. Put $\mu_S^H := \Psi_* \mu_S$. Then $\iota_* \mu_S^H = \mu_S$. By Lemma J.1 (applied to ι composed with countably many evaluations), ι is a Borel isomorphism onto the Borel set $\iota\mathcal{H}^{-r}$, which has full μ_S-measure. This gives uniqueness of μ_S^H and the formulas for the characteristic functional and the moments, which hold for μ_S by Theorem 3.1(ii).

(c) Let μ' be such a measure, and fix real $f_1, \dots, f_N \in C_c^\infty$ and $t \in \mathbb{R}^N$. Put $f := \sum_l t_l f_l = (\mathbf{g}, h)$. The moments of the real random variable $Z := \Psi(f) = \sum_l t_l \Psi(f_l)$ under μ' are $m_n = \sum_{k \le n} \binom{n}{k} S_{k,n-k}(\mathbf{g}, \dots; h, \dots)$, the same as under μ_S^H. By Theorem 3.1(iii) and $\binom{n}{k}(k!)^{1/2}(n-k)! \le n!$, they satisfy $|m_n| \le c_S(n+1)K^n\, n!$ with $K := \max(\mathrm{e}\|\mathbf{g}\|_2, \mathrm{e}N(h)/\sigma_0)$. The even moments are

absolute moments, and the Cauchy–Schwarz inequality $\mathbb{E}|Z|^{2n+1} \le (m_{2n}m_{2n+2})^{1/2}$ gives $\mathbb{E}|Z|^n \le c_S(n+2)K^n\,n!$ for all n, under μ' and under μ_S^H. Let $\phi(s) := \mathbb{E}e^{isZ}$. Then $\phi^{(k)}(s) = \mathbb{E}[(iZ)^k e^{isZ}]$, and from $|e^{iy} - \sum_{k<n}(iy)^k/k!| \le |y|^n/n!$,

$$\Big|\phi(s+u) - \sum_{k<n}\phi^{(k)}(s)\frac{u^k}{k!}\Big| \le \frac{|u|^n\,\mathbb{E}|Z|^n}{n!} \le c_S(n+2)(K|u|)^n \to 0 \qquad (|u| < 1/K).$$

So on every interval $(s-\rho, s+\rho)$, $\rho < 1/K$, ϕ is the sum of its Taylor series at s. The characteristic functions of Z under μ' and μ_S^H have the same derivatives $i^k m_k$ at 0, so they agree on $(-\rho,\rho)$; there all their derivatives agree, so they agree near $\pm\rho/2$, and by induction on $\mathbb{R}$. Hence $\mathbb{E}_{\mu'}e^{i\sum_l t_l\Psi(f_l)} = \mathbb{E}_{\mu_S^H}e^{i\sum_l t_l\Psi(f_l)}$ for all $t \in \mathbb{R}^N$, and by uniqueness of characteristic functions on $\mathbb{R}^N$ the laws of $(\Psi(f_1),\dots,\Psi(f_N))$ agree. These vectors, over all finite families, generate a π-system that generates the Borel σ-algebra of $\mathcal{H}^{-r}$ (Lemma J.1; the evaluations at C_c^∞ separate points). Hence $\mu' = \mu_S^H$.

(d) *(H1).* $\mathcal{H}^{-r}$ is a separable Hilbert space, hence Polish. By (a), with $\|\cdot\|_r$ replaced by the equivalent polynomial norm, $\|f(\cdot - a)\|_r \le C_r(1+|a|)^r\|f\|_r$. So τ_a, the adjoint of $f \mapsto f(\cdot + a)$, is bounded on $\mathcal{H}^{-r}$ with $\|\tau_a\| \le C_r(1+|a|)^r$. On the dense subspace $\mathcal{S} \subset \mathcal{H}^{-r}$, embedded by the L^2 pairing, $\|\tau_a\psi - \psi\|_{-r} \le \|\psi(\cdot - a) - \psi\|_{L^2} \to 0$. Local boundedness and an $\epsilon/3$ argument give strong continuity. Joint continuity follows from $\|\tau_a\Psi - \tau_b\Psi'\|_{-r} \le C_r(1+|a|)^r\|\Psi - \Psi'\|_{-r} + \|\tau_a\Psi' - \tau_b\Psi'\|_{-r}$. Every $g \in O(3)$ commutes with $\mathcal{A}$ and acts on $\mathcal{V}$ by an orthogonal matrix, so θ_j and O_h act unitarily. The relations $\theta_j\tau_a = \tau_{\rho_j a}\theta_j$ are identities between the formulas on test functions. The actions commute with ι, and $\iota\mathcal{H}^{-r}$ is invariant. Since $\iota_*\mu_S^H = \mu_S$ and ι is a Borel isomorphism onto its image, every invariance of μ_S is an invariance of μ_S^H. *(H2).* $\Sigma^H(U)$ is generated by the bounded continuous functions $\arctan\Psi(f_n)$, f_n in a countable subset of $C_c^\infty(U)$ dense in every $C_c^\infty(K)$, $K \Subset U$, for the H^r-norm. The covariance and monotonicity properties are those of Σ, and $\bigvee_K \Sigma^H(K)$ is the Borel σ-algebra by Lemma J.1. *(H3).* A $\Sigma^H(H_j^+(0))$-measurable F has the form $F = \phi((\Psi(f_n))_n)$ with $f_n \in C_c^\infty(H_j^+(0))$ and ϕ Borel. So $F = F' \circ \iota$ with $F' := \phi((\Phi(f_n))_n)$, which is $\Sigma(H_j^+(0))$-measurable on X. Then $\mathbb{E}_{\mu_S^H}[\overline{F \circ \theta_j}F] = \mathbb{E}_{\mu_S}[\overline{F' \circ \theta_j}F'] \ge 0$. □

By Proposition 10.4, the hypothesis of part (d) holds. Hence Theorem 3.6 may be stated, as it is, on $\mathcal{H}^{-r}$ for every even $r \ge 14$, with $\mu = \mu_S^H$; its uniqueness clause is part (c). The abstract results of Sections 10.3–10.8 apply on either space, and the two decompositions correspond under ι.

10.3. **Translations and the mean-ergodic projections.** Throughout Sections 10.3–10.7, μ is any measure satisfying (H1)–(H3).

Lemma 10.6. *$a \mapsto U(a)$ is strongly continuous on $L^2(\mu)$.*

Proof. For bounded continuous G, $G(\tau_{-a}x) \to G(x)$ for every x as $a \to 0$ by continuity of the action, and dominated convergence gives $U(a)G \to G$ in L^2. Bounded continuous functions are dense in $L^2(\mu)$ for a Borel probability measure on a metric space, and $\|U(a)\| = 1$. □

For $J \subset \{1,2,3\}$ let P_J be the orthogonal projection of $L^2(\mu)$ onto $\{G : U(a)G = G\ \forall a \in V_J\}$, with $P_\emptyset = 1$ and $P_j := P_{\{j\}}$. Let $\mathcal{I}$ be the σ-algebra of $\mathbb{R}^3$-invariant Borel sets, completed by μ-null sets.

Lemma 10.7. *(i) For $\Lambda_T^J := [-T,T]^3 \cap V_J$ and every $G \in L^2(\mu)$, $|\Lambda_T^J|^{-1}\int_{\Lambda_T^J} U(a)G\,da \to P_JG$ in $L^2(\mu)$ as $T \to \infty$. The same holds for one-sided means over $[0,T]$ in one coordinate of J. (ii) P_J commutes with every $U(a)$ and every Θ_j; $P_JP_{J'} = P_{J\cup J'}$; and $P_J = \mathbb{E}[\,\cdot \mid \mathcal{I}_J]$, where $\mathcal{I}_J$ is the σ-algebra of V_J-invariant Borel sets completed by μ-null sets; in particular $P_{\{1,2,3\}} = \mathbb{E}[\,\cdot \mid \mathcal{I}]$.*

Proof. (i) For one coordinate j, Stone's theorem gives $U(te_j) = e^{itA_j}$ with A_j self-adjoint, and $(2T)^{-1}\int_{-T}^{T} e^{itA_j}dt = \phi_T(A_j)$ with $\phi_T(\lambda) = \sin(T\lambda)/(T\lambda)$, $\phi_T(0) = 1$. Since $|\phi_T| \le 1$ and $\phi_T \to \mathbf{1}_{\{0\}}$

pointwise, the spectral theorem and dominated convergence give strong convergence to the spectral projection of $\{0\}$, which is the projection onto $\operatorname{Ker} A_j = \{G : U(te_j)G = G\ \forall t\}$. For one-sided means use $\phi_T(\lambda) = (e^{iT\lambda}-1)/(iT\lambda)$. The mean over Λ_T^J is the product of the commuting one-dimensional means; products of commuting, uniformly bounded, strongly convergent families converge strongly to the product of the limits; and the product of commuting projections is the projection onto the intersection of their ranges, which is the V_J-fixed space.

(ii) The means commute with $U(a)$, and with Θ_j by (10.3) and $\rho_j V_J = V_J$; strong limits of commuting operators commute. $P_J P_{J'} = P_{J\cup J'}$ by the intersection statement in (i). It remains to show that the range of P_J is $L^2(\mathcal{I}_J,\mu)$. Let $G \in L^2(\mu)$ satisfy $U(a)G = G$ for all $a \in V_J$, and fix a Borel version of G. For each $a \in V_J$ the set $\{x : G(\tau_{-a}x) \neq G(x)\}$ is μ-null, so by Fubini's theorem on $V_J \times X$ the set $X_0 := \{x : G(\tau_{-a}x) = G(x) \text{ for a.e. } a \in V_J\}$ is Borel and of full measure. Let X_1 be the set of x for which $a \mapsto G(\tau_{-a}x)$ is a.e. constant on V_J, that is,

$$\int_{\Lambda_n^J}\int_{\Lambda_n^J} \min\big(1, |G(\tau_{-a}x) - G(\tau_{-b}x)|\big)\, da\, db = 0 \qquad \text{for all } n \in \mathbb{N}.$$

By Tonelli's theorem X_1 is Borel; $X_0 \subset X_1$; and X_1 is V_J-invariant, because $G(\tau_{-a}\tau_c x) = G(\tau_{-(a-c)}x)$ and a.e. constancy on V_J is preserved by shifts of the variable. For $x \in X_1$ let $\tilde G(x)$ be the a.e. value of $a \mapsto G(\tau_{-a}x)$, that is, $\tilde G(x) = \lim_{N\to\infty} |\Lambda_1^J|^{-1} \int_{\Lambda_1^J} t_N\big(G(\tau_{-a}x)\big)\, da$, where t_N truncates real and imaginary parts at level N; put $\tilde G := 0$ off X_1. Then $\tilde G$ is Borel, $\tilde G \circ \tau_c = \tilde G$ everywhere for $c \in V_J$, and $\tilde G = G$ on X_0, hence μ-a.e. Thus G is $\mathcal{I}_J$-measurable. Conversely, an $\mathcal{I}_J$-measurable function is a.e. equal to a V_J-invariant Borel function and is therefore fixed by every $U(a)$, $a \in V_J$. □

The next lemma is the only place where reflection positivity enters the decomposition: functions invariant under translations in a direction live at infinity in that direction and are fixed by the corresponding reflection.

Lemma 10.8 (localisation of invariant functions)**.** *Let $j \in J \subset \{1,2,3\}$ and $W \in L^2(\mu)$ with $U(a)W = W$ for all $a \in V_J$. Then: (i) $W \in L^2(\Sigma(H_j^+(c)),\mu) \cap L^2(\Sigma(H_j^-(c)),\mu)$ for every $c \in \mathbb{R}$; (ii) $\Theta_j W = W$; (iii) if $G \in L^2(\Sigma(K),\mu)$, then $P_J G \in L^2(\Sigma(K + V_J),\mu)$.*

Proof. (i) Given $\epsilon > 0$, by (H2) there are a bounded K and $G \in L^2(\Sigma(K))$ with $\|W - G\|_2 < \epsilon$ (an increasing union of σ-algebras generating $\mathcal{B}$ has L^2 spaces whose union is dense). For every t, $\|W - U(te_j)G\|_2 = \|U(te_j)(W-G)\|_2 < \epsilon$, and $U(te_j)G \in L^2(\Sigma(K + te_j)) \subset L^2(\Sigma(H_j^+(c)))$ for t large. That space is closed, so it contains W. Letting $t \to -\infty$ gives the other half space.

(ii) $\Theta_j W$ is V_J-invariant by (10.3), so by (i) both W and $\Theta_j W$ lie in $L^2(\Sigma(H_j^+(0)))$. Put $F = W - \Theta_j W$. Then $\Theta_j F = -F$, and (10.2) gives $0 \le \mathbb{E}[\overline{\Theta_j F}\, F] = -\mathbb{E}|F|^2$, so $F = 0$.

(iii) The means of Lemma 10.7(i) applied to G lie in the closed subspace $L^2(\Sigma(K + V_J))$, since $U(a)G \in L^2(\Sigma(K + a))$ and $K + a \subset K + V_J$ for $a \in V_J$. □

10.4. **Ergodic decomposition and inheritance.**

Lemma 10.9 (conditional probabilities)**.** *Let X be Polish, m a Borel probability measure on X and $\mathcal{G} \subset \mathcal{B}$ a σ-algebra. There are Borel probability measures m_ξ, $\xi \in X$, such that $\xi \mapsto m_\xi(B)$ is $\mathcal{G}$-measurable and $\int_C m_\xi(B)\, dm(\xi) = m(B \cap C)$ for all $B \in \mathcal{B}$, $C \in \mathcal{G}$.*

Proof. Let $d \le 1$ be a complete metric on X and (x_n) a dense sequence. Then $\jmath(x) := (d(x,x_n))_n$ is a continuous injection into the compact space $Q := [0,1]^{\mathbb{N}}$, and by Lemma J.1 it is a Borel isomorphism onto the Borel set $\jmath X$. Let $\mathcal{C}_0$ be a countable $\mathbb{Q}$-linear dense subspace of $C(Q)$ containing 1, and choose $\mathcal{G}$-measurable versions $L_\xi(G)$ of $\mathbb{E}_m[G\circ\jmath \mid \mathcal{G}]$, $G \in \mathcal{C}_0$. Off a set $N_0 \in \mathcal{G}$ with $m(N_0) = 0$, L_ξ is $\mathbb{Q}$-linear and positive, and $L_\xi(1) = 1$. So $|L_\xi(G)| \le \|G\|_\infty$, and L_ξ extends to a positive normalised functional on $C(Q)$, given by a probability measure κ_ξ (Riesz representation

theorem). On N_0 put $\kappa_\xi := \jmath_* m$. The bounded Borel G on Q for which $\kappa_\xi(G)$ is a $\mathcal{G}$-measurable version of $\mathbb{E}_m[G\circ\jmath \mid \mathcal{G}]$ form a vector space that contains $C(Q)$ and is closed under bounded monotone limits, so it contains all bounded Borel G. In particular $\kappa_\xi(\jmath X)=1$ for $\xi\notin N_1$, $N_1\in\mathcal{G}$ null. Put $m_\xi := (\jmath^{-1})_*\kappa_\xi$ off $N_0\cup N_1$ and $m_\xi := m$ on it. □

Theorem 10.10 (ergodic decomposition with inheritance)**.** *Let $\mathcal{I}_0\subset\mathcal{B}$ be the σ-algebra of Borel sets B with $\tau_a B = B$ for all $a\in\mathbb{R}^3$ (its μ-completion is $\mathcal{I}$), let μ_ξ, $\xi\in X$, be the measures of Lemma 10.9 for $m=\mu$ and $\mathcal{G}=\mathcal{I}_0$, a regular conditional probability of μ given $\mathcal{I}$, and put $(\Xi,\nu):=(X,\mu)$. Then $\xi\mapsto\mu_\xi(B)$ is $\mathcal{I}$-measurable for every Borel set B, and:*

(a) *$\mu=\int_X \mu_\xi\,d\mu(\xi)$; for $G\in L^1(\mu)$ one has $G\in L^1(\mu_\xi)$ and $\mathbb{E}[G\mid\mathcal{I}](\xi)=\mathbb{E}_{\mu_\xi}[G]$ for μ-a.e. ξ; in particular $\mathbb{E}_\mu[WG]=\int W(\xi)\,\mathbb{E}_{\mu_\xi}[G]\,d\mu(\xi)$ for bounded $\mathcal{I}$-measurable W;*

(b) *for ν-a.e. ξ, μ_ξ is invariant under every τ_a and ergodic for the $\mathbb{R}^3$-action;*

(c) *for ν-a.e. ξ, μ_ξ is invariant under $\theta_1,\theta_2,\theta_3$;*

(d) *for ν-a.e. ξ, μ_ξ satisfies (H3);*

(e) *every μ_ξ satisfies (H2) with the same $\Sigma(U)$, and every $G\in L^2(\mu)$ lies in $L^2(\mu_\xi)$ for ν-a.e. ξ.*

Proof. (a) By Lemma 10.9, (a) holds for bounded G, and for $G\in L^1(\mu)$ by monotone convergence. Fix a complete metric $d\le 1$ on X and a dense sequence (x_n), and let $\mathcal{D}$ be the countable $\mathbb{Q}[i]$-algebra generated by 1 and the functions $x\mapsto d(x,x_n)$. It consists of bounded continuous functions, generates $\mathcal{B}$, and is dense in $L^2(m)$ for every Borel probability measure m (Lemma J.2); so two Borel probability measures that agree on $\mathcal{D}$ are equal.

(b) *Invariance.* For $B\in\mathcal{I}_0$, $\mathbb{E}_\mu[\mathbf{1}_B\,G\circ\tau_a]=\mathbb{E}_\mu[(\mathbf{1}_B G)\circ\tau_a]=\mathbb{E}_\mu[\mathbf{1}_B G]$. Hence there is a null set off which $\mu_\xi(G\circ\tau_a)=\mu_\xi(G)$ for all $G\in\mathcal{D}$ and $a\in\mathbb{Q}^3$. Since $a\mapsto\mu_\xi(G\circ\tau_a)$ is continuous (joint continuity of the action and dominated convergence), this holds for all a, so μ_ξ is τ-invariant. *Ergodicity.* For $G\in\mathcal{D}$ let $q_G(\xi):=\mu_\xi(G)$, a bounded $\mathcal{I}_0$-measurable version of $\mathbb{E}_\mu[G\mid\mathcal{I}]$. Then $\mu_\xi(q_G)=q_G(\xi)$ and $\mu_\xi(|q_G|^2)=|q_G(\xi)|^2$ for a.e. ξ, so $\mu_\xi(|q_G-q_G(\xi)|^2)=0$: for a.e. ξ, q_G is μ_ξ-a.s. the constant $\mu_\xi(G)$. Let $A_T G(x):=|\Lambda_T|^{-1}\int_{\Lambda_T} G(\tau_{-a}x)\,da$. By Lemma 10.7, $\int\mu_\xi(|A_TG-q_G|^2)\,d\mu(\xi)=\mathbb{E}_\mu|A_TG-q_G|^2\to 0$. Choose $T_k\to\infty$ such that this is $\le 2^{-k}$ for the first k elements of an enumeration of $\mathcal{D}$. Then for a.e. ξ, $A_{T_k}G\to\mu_\xi(G)$ in $L^2(\mu_\xi)$ for every $G\in\mathcal{D}$. The proof of Lemma 10.7(i) uses only the invariance of the measure, so $A_{T_k}G$ also tends in $L^2(\mu_\xi)$ to the projection of G onto the τ-fixed vectors. By density of $\mathcal{D}$, that projection is $G\mapsto\mu_\xi(G)\mathbf{1}$. If $B\in\mathcal{I}_0$, $\mathbf{1}_B$ is fixed, hence a.s. constant, so $\mu_\xi(B)\in\{0,1\}$.

(e) (H2) is a condition on X and Σ alone and does not involve the measure, so it holds for every μ_ξ. Also $\int\mathbb{E}_{\mu_\xi}|G|^2\,d\nu=\mathbb{E}_\mu|G|^2<\infty$.

(c) Let $W\ge 0$ be bounded and $\mathcal{I}$-measurable and G bounded Borel. By Lemma 10.8(ii), $\Theta_j W=W$, so $\mathbb{E}_\mu[W\,\Theta_jG]=\mathbb{E}_\mu[\Theta_j(WG)]=\mathbb{E}_\mu[WG]$ by θ_j-invariance of μ. By (a), $\int W(\xi)\big(\mathbb{E}_{\mu_\xi}[\Theta_jG]-\mathbb{E}_{\mu_\xi}[G]\big)d\nu=0$ for all such W, hence $\mathbb{E}_{\mu_\xi}[\Theta_jG]=\mathbb{E}_{\mu_\xi}[G]$ for ν-a.e. ξ. Taking G in a countable algebra of bounded continuous functions generating $\mathcal{B}$ makes the exceptional set independent of G, and gives θ_j-invariance of μ_ξ for a.e. ξ.

(d) By translation invariance of μ_ξ it suffices to take $c=0$. Let $W\ge0$ be bounded and $\mathcal{I}$-measurable and $F\in L^2(\Sigma(H_j^+(0)),\mu)$. By Lemma 10.8(i)–(ii), $\sqrt{W}$ has a $\Sigma(H_j^+(0))$-measurable version and $\Theta_j\sqrt{W}=\sqrt{W}$. Hence $\sqrt{W}F\in L^2(\Sigma(H_j^+(0)))$ and

$$\mathbb{E}_\mu\big[W\,\overline{F\circ\theta_j}\,F\big]=\mathbb{E}_\mu\big[\overline{(\sqrt{W}F)\circ\theta_j}\,(\sqrt{W}F)\big]\ge 0.$$

With (a), $\mathbb{E}_{\mu_\xi}[\overline{F\circ\theta_j}F]\ge 0$ for ν-a.e. ξ. Let $\mathcal{L}$ be the countable $\mathbb{Q}[i]$-algebra, closed under conjugation, generated by a countable family of bounded continuous functions generating $\Sigma(H_j^+(0))$ (H2). The exceptional set can be taken uniform over $F\in\mathcal{L}$. By Lemma J.2, $\mathcal{L}$ is dense in $L^2(\Sigma(H_j^+(0)),m)$ for every Borel probability measure m, and (10.2) extends from $\mathcal{L}$ to

$L^2(\Sigma(H_j^+(0)),\mu_\xi)$ by the bound $|\mathbb{E}[\overline{F\circ\theta_j}G]| \le \|F\|_2\|G\|_2$, valid because μ_ξ is θ_j-invariant by (c). □

Remark 10.11. The three coordinate reflections are inherited by the components because they act trivially on $\mathcal{I}$ (Lemma 10.8(ii)). A coordinate permutation maps components to components (equation (10.5) below), and the mixture μ is O_h-invariant. After grouping the components along the orbits of the permutation group $S_3 = O_h/\langle\rho_1,\rho_2,\rho_3\rangle$ the components may be chosen O_h-invariant, each a uniform mixture of at most six translation-ergodic ones (Theorem 10.12). O_h-invariance of the translation-ergodic components themselves does not follow from (H1)–(H3) and O_h-invariance of μ (Example 10.14).

Assume, in addition to (H1)–(H3), that O_h acts on X by homeomorphisms $x \mapsto gx$, with $\rho_j x = \theta_j x$ for the coordinate reflections, $g\,\tau_a\,g^{-1} = \tau_{ga}$, μ invariant under every g, and $G \mapsto G\circ g^{-1}$ mapping $\Sigma(U)$-measurable functions to $\Sigma(gU)$-measurable ones. For the states of Theorem 3.1 this holds on $X = H^{-s}_{\mathrm{loc}}(\mathbb{R}^3;\mathcal{V})$ with $(g\Phi)(f) = \Phi(g_*^{-1}f)$ (Proposition 10.4). Let $\mathcal{G} := \mathbb{R}^3 \rtimes O_h$, let $N \cong (\mathbb{Z}/2)^3$ be the subgroup of O_h generated by ρ_1,ρ_2,ρ_3, so that $O_h/N \cong S_3$, and let $\mathcal{I}_{\mathcal{G}} \subset \mathcal{I}$ be the σ-algebra of $\mathcal{G}$-invariant Borel sets, completed by μ-null sets.

Theorem 10.12 (decomposition with cubic symmetry)**.** *There are Borel probability measures μ'_x on X, $x \in X$, with $x \mapsto \mu'_x(B)$ $\mathcal{I}_{\mathcal{G}}$-measurable for every Borel B and $\mu = \int \mu'_x\,d\mu(x)$, such that for μ-almost every x:*

(a) *μ'_x is invariant under $\mathcal{G}$ and ergodic for its action, and satisfies (H1)–(H3);*
(b) *there are an integer $k = k(x) \in \{1,2,3,6\}$ and pairwise distinct measures $m_1,\dots,m_k$, each invariant under the translations and the coordinate reflections, ergodic for $\mathbb{R}^3$ and satisfying (H1)–(H3), such that O_h permutes $\{m_1,\dots,m_k\}$ transitively and $\mu'_x = k^{-1}\sum_{i=1}^k m_i$; the m_i are components of the decomposition of Theorem 10.10;*
(c) *in the reconstruction from μ'_x in any coordinate direction j (Proposition 10.16), the joint fixed space of $\{e^{-tH_j}\}_{t\ge0}$ and $\{V_j(y)\}_{y\perp e_j}$ has dimension k;*
(d) *for all $G, G' \in L^2(\mu'_x)$, the Cesàro means over cubes of $\mathbb{E}_{\mu'_x}[G'\,U(a)G]$, and $\mathbb{E}_{\mu'_x}[G'\,U(a)G]$ itself as $\min_l|a_l| \to \infty$, converge to $k^{-1}\sum_{i=1}^k \mathbb{E}_{m_i}[G']\,\mathbb{E}_{m_i}[G]$.*

Moreover $k(x) = 1$ for μ-almost every x if and only if almost every component of Theorem 10.10 is O_h-invariant.

Proof. Step 1: N acts trivially on $\mathcal{I}$, and O_h preserves $\mathcal{I}$. A bounded $\mathcal{I}$-measurable W is invariant along every coordinate axis, so $W\circ\theta_j = W$ a.s. for $j = 1,2,3$ by Lemma 10.8(ii); hence $W\circ n = W$ for $n \in N$. For $g \in O_h$ and a translation-invariant set B, $\tau_a(gB) = g\,\tau_{g^{-1}a}B = gB$, so g maps $\mathcal{I}$ onto itself.

Step 2: conditional expectations. Let μ_x be the components of Theorem 10.10 and μ'_x a regular conditional probability of μ given $\mathcal{I}_{\mathcal{G}}$ (Lemma 10.9). For bounded Borel F and $g \in O_h$, g preserves μ and maps $\mathcal{I}$ onto itself, so $\mathbb{E}[F\circ g \mid \mathcal{I}] = \mathbb{E}[F\mid\mathcal{I}]\circ g$ a.s., that is

$$\mu_{gx} = g_*\mu_x \qquad \text{for } \mu\text{-a.e. } x, \tag{10.5}$$

first for F in a countable algebra of bounded continuous functions generating $\mathcal{B}$ and then as measures. By Step 1 and (10.5) with $g \in N$, $n_*\mu_x = \mu_x$ for $n \in N$, so $S_3 = O_h/N$ acts on the components by $\sigma\cdot\mu_x := g_*\mu_x$, $g \in \sigma$. For bounded $\mathcal{I}$-measurable h put $\bar h := |O_h|^{-1}\sum_{g\in O_h} h\circ g$. It is $\mathcal{I}$-measurable and O_h-invariant, hence $\mathcal{I}_{\mathcal{G}}$-measurable, and for bounded $\mathcal{I}_{\mathcal{G}}$-measurable W (which satisfies $W\circ g = W$ a.s.) $\mathbb{E}[W\bar h] = |O_h|^{-1}\sum_g \mathbb{E}[(W\circ g)(h\circ g)] = \mathbb{E}[Wh]$. So $\mathbb{E}[h\mid\mathcal{I}_{\mathcal{G}}] = \bar h$. Applied to $h = \mathbb{E}[F\mid\mathcal{I}]$, with $\mathbb{E}[F\mid\mathcal{I}_{\mathcal{G}}] = \mathbb{E}[\mathbb{E}[F\mid\mathcal{I}]\mid\mathcal{I}_{\mathcal{G}}]$ and (10.5),

$$\mu'_x = \frac{1}{48}\sum_{g\in O_h}\mu_{gx} = \frac{1}{48}\sum_{g\in O_h} g_*\mu_x = \frac{1}{6}\sum_{\sigma\in S_3}\sigma\cdot\mu_x \qquad \text{for } \mu\text{-a.e. } x.$$

Grouping equal terms, $\mu'_x = k^{-1}\sum_{i=1}^{k} m_i$, where $\{m_1,\dots,m_k\}$ is the S_3-orbit of μ_x, so $k = 6/|\mathrm{Stab}(\mu_x)| \in \{1,2,3,6\}$. For a.e. x every μ_{gx} is a component with the properties of Theorem 10.10(b)–(e); this gives (b). Since $k=1$ exactly when μ_x is fixed by S_3, that is, O_h-invariant, the last assertion follows.

Step 3: (a). μ'_x is invariant under the translations and under O_h, as an average over an O_h-orbit of translation-invariant measures. If B is a $\mathcal{G}$-invariant Borel set, then $m_i(B) \in \{0,1\}$ by ergodicity, and $(g_* m_i)(B) = m_i(g^{-1}B) = m_i(B)$, so all $m_i(B)$ are equal by transitivity and $\mu'_x(B) \in \{0,1\}$. Each m_i satisfies (H3), and (H3) is preserved under convex combinations. (H2) holds as in Theorem 10.10(e).

Step 4: invariant sets separating the m_i. Let $m \neq m'$ be two of the m_i, and F bounded continuous with $m(F) \neq m'(F)$. With $A_T F := |\Lambda_T|^{-1}\int_{\Lambda_T} U(a)F\,da$, Lemma 10.7(i) and ergodicity give $A_T F \to m(F)$ in $L^2(m)$ and $A_T F \to m'(F)$ in $L^2(m')$; choose $T_n \to \infty$ with a.s. convergence for both. Since $\sup_x |A_T F(\tau_b x) - A_T F(x)| \le 2\|F\|_\infty |(\Lambda_T + b)\triangle\Lambda_T|/|\Lambda_T| \to 0$ for each b, the sets $\{x : \lim_n A_{T_n}F(x) = m(F)\}$ and $\{x : \lim_n A_{T_n}F(x) = m'(F)\}$ are disjoint translation-invariant Borel sets of full m-measure and full m'-measure respectively. Doing this for every pair and intersecting gives pairwise disjoint translation-invariant Borel sets $B_1,\dots,B_k$ with $m_i(B_i) = 1$.

Step 5: (c) and (d). Let $\mu' := \mu'_x$ and $\mathcal{I}'$ the σ-algebra of translation-invariant sets for μ'. An $\mathcal{I}'$-measurable function is m_i-a.s. constant for each i, so $L^2(\mathcal{I}',\mu')$ is spanned by $\mathbf{1}_{B_1},\dots,\mathbf{1}_{B_k}$, has dimension k, and $\mathbb{E}_{\mu'}[G \mid \mathcal{I}'] = \sum_i \mathbb{E}_{m_i}[G]\,\mathbf{1}_{B_i}$. By Proposition 10.17(ii) for μ', the projection onto the joint fixed space is $[G] \mapsto [\mathbb{E}_{\mu'}[G \mid \mathcal{I}']]$, and $W \mapsto [W]$ is isometric on $L^2(\mathcal{I}',\mu')$, because $\|[W]\|^2 = \mathbb{E}_{\mu'}[\overline{\Theta_j W}\, W] = \mathbb{E}_{\mu'}|W|^2$ by Lemma 10.8(ii); the $\mathbf{1}_{B_i}$ lie in L^2_+ by Lemma 10.8(i). So the joint fixed space is spanned by the k orthogonal vectors $[\mathbf{1}_{B_i}]$, which is (c). Theorems 10.19 and 10.20 for μ' give the limit $\mathbb{E}_{\mu'}\big[\mathbb{E}[G' \mid \mathcal{I}']\,\mathbb{E}[G \mid \mathcal{I}']\big] = k^{-1}\sum_i \mathbb{E}_{m_i}[G']\,\mathbb{E}_{m_i}[G]$, which is (d). □

Remark 10.13. (1) The components μ'_x satisfy OS0 in the form E0″ with x-dependent constants: the proof of Corollary 10.43(iv) applies verbatim with the decomposition $\mu = \int \mu'_x\,d\mu(x)$. As for the components of Theorem 10.10, the tube form of OS0 and the clause of OS2 for exponentials are not claimed for them. They satisfy OS1 for $\mathbb{R}^3 \rtimes O_h$, OS2 about every coordinate plane and the symmetry and reality conditions of OS3; they are ergodic for $\mathcal{G}$, and their joint vacuum has dimension $k(x) \le 6$. (2) Charge conjugation C ($F \mapsto -F$, $X_{\mathrm{R}} \mapsto X_{\mathrm{R}}$) commutes with $\mathcal{G}$ and preserves μ (Theorem 9.8 and the uniqueness in Proposition 10.4). The same argument with $O_h \times \{1,\mathsf{C}\}$ in place of O_h gives components that are moreover C-invariant, so that their moments of odd degree in the curvature vanish, with $k \in \{1,2,3,4,6,12\}$ (the group acts through the quotient $S_3 \times \mathbb{Z}/2$, whose subgroups have index $1,2,3,4,6$ or 12). (3) In the reconstruction from μ'_x in direction j, the stabiliser D_4 of e_j acts unitarily as in Theorem 11.1(ii). It permutes the k joint vacua $[\mathbf{1}_{B_i}]$ through its image in S_3, which is generated by the transposition of the two coordinates transverse to e_j: if $g_* m_i = m_{i'}$, then $\mathbf{1}_{gB_i} = \mathbf{1}_{B_{i'}}$ μ'_x-a.s.

Example 10.14 (cubic symmetry is not inherited by the components)**.** Let m_1 be the law of the field $\Phi(f) := \psi(f^{(1)})$, with ψ the massive free field on $\mathbb{R}^2$ and $f^{(1)}$ as in Example 10.22, realised on $\mathcal{V}$ with vanishing curvature component; this is the summand $\psi_1(x_2,x_3)$ of that example alone. Let m_2, m_3 be its images under the transpositions of the coordinates (x_1,x_2) and (x_1,x_3). The proofs of Example 10.22(ii)–(iii), with one plane instead of three, show that each m_i satisfies (H1)–(H3) and is ergodic for $\mathbb{R}^3$; in the direction transverse to its plane reflection positivity holds because every function measurable for a half space is invariant under the reflection. The measure m_1 is invariant under the coordinate reflections and under the transposition of (x_2,x_3), but not under that of (x_1,x_2). So $\mu := \frac{1}{3}(m_1+m_2+m_3)$ satisfies (H1)–(H3), is invariant under $\mathbb{R}^3 \rtimes O_h$, has the translation-ergodic components m_1, m_2, m_3, none of which is O_h-invariant, and $k = 3$ in Theorem 10.12. Thus O_h-invariance of the translation-ergodic components does not follow from (H1)–(H3) and O_h-invariance of the law.

Remark 10.15. The components μ_ξ of Theorem 3.6 are invariant under the translations and the three coordinate reflections, and a coordinate permutation maps a component to a component. Grouping the components along the orbits of O_h gives a decomposition into O_h-invariant, reflection-positive components, each a uniform mixture of at most 6 translation-ergodic ones, with a joint vacuum of dimension at most 6 in each coordinate direction (Theorem 10.12). Whether almost every translation-ergodic component is itself O_h-invariant is not known; it holds if and only if $k = 1$ almost everywhere in Theorem 10.12.

10.5. **Reconstruction in a coordinate direction.** Fix μ satisfying (H1)–(H3) and a direction j. Write $L^2_+ := L^2(\Sigma(H_j^+(0)), \mu)$ and define the form

$$\langle F,\, G\rangle_j := \mathbb{E}_\mu\big[\overline{\Theta_j F}\, G\big], \qquad F, G \in L^2_+.$$

It is positive semidefinite by (H3), Hermitian because Θ_j is a unitary involution preserving μ, and bounded: $|\langle F,\, G\rangle_j| \le \|F\|_2 \|G\|_2$.

Proposition 10.16 (reconstruction in direction j)**.** *Let $\mathcal{N} = \{F \in L^2_+ : \langle F,\, F\rangle_j = 0\}$, $\mathcal{H}_j$ the completion of $L^2_+/\mathcal{N}$, $[\,\cdot\,] : L^2_+ \to \mathcal{H}_j$ the quotient map, and $\Omega_j = [1]$.*

(i) *For $t \ge 0$ and $y \perp e_j$, $T_j(t)[G] := [U(te_j)G]$ and $V_j(y)[G] := [U(y)G]$ are well defined. T_j is a strongly continuous semigroup of self-adjoint contractions, so $T_j(t) = e^{-tH_j}$ with $H_j \ge 0$ self-adjoint. V_j is a strongly continuous unitary representation of $e_j^\perp \cong \mathbb{R}^2$. All these operators commute, and $T_j(t)\Omega_j = V_j(y)\Omega_j = \Omega_j$.*

(ii) *For $G \in L^2_+$ and $a = a_j e_j + a_\perp$ with $a_j \ge 0$: $[U(a)G] = T_j(a_j)V_j(a_\perp)[G]$.*

(iii) *For $F \in L^2(\Sigma(H_j^-(0)), \mu)$ and $G \in L^2_+$: $\overline{\Theta_j F} \in L^2_+$ and $\mathbb{E}_\mu[FG] = \langle[\overline{\Theta_j F}],\, [G]\rangle_{\mathcal{H}_j}$.*

Proof. This is the Osterwalder–Schrader construction [27, 28], [19, Thm. 6.1.3, p. 94]; we write out the identities that drive it. For $G \in L^2_+$ and $t \ge 0$, $U(te_j)G \in L^2(\Sigma(H_j^+(t))) \subset L^2_+$ by (H2). By (10.3) and invariance of μ,

$$\langle U(te_j)F,\, U(te_j)G\rangle_j = \mathbb{E}\big[\overline{U(-te_j)\Theta_j F}\, U(te_j)G\big] = \langle F,\, U(2te_j)G\rangle_j.$$

So the form is symmetric for $U(te_j)$, and by the Schwarz inequality for the form, $\langle U(te_j)G,\, U(te_j)G\rangle_j \le \langle G,\, G\rangle_j^{1/2}\langle U(2te_j)G,\, U(2te_j)G\rangle_j^{1/2}$. Iterating n times and using $\langle U(se_j)G,\, U(se_j)G\rangle_j \le \|G\|_2^2$ gives $\langle U(te_j)G,\, U(te_j)G\rangle_j \le \langle G,\, G\rangle_j^{1-2^{-n}} \|G\|_2^{2^{1-n}}$, and $n \to \infty$ gives $\langle U(te_j)G,\, U(te_j)G\rangle_j \le \langle G,\, G\rangle_j$. So $T_j(t)$ maps $\mathcal{N}$ into $\mathcal{N}$ and is a symmetric contraction on the quotient, hence a self-adjoint contraction on $\mathcal{H}_j$; the semigroup law is inherited from U. Strong continuity follows from $\|T_j(t)[G] - [G]\| \le \|U(te_j)G - G\|_2$ and Lemma 10.6. For $y \perp e_j$, Θ_j commutes with $U(y)$, so $\langle U(y)F,\, U(y)G\rangle_j = \langle F,\, G\rangle_j$: $V_j(y)$ is isometric with isometric inverse $V_j(-y)$. Commutation is inherited from U, and (ii) is the definition. (iii): $\overline{\Theta_j F}$ is $\Sigma(H_j^+(0))$-measurable by (H2), and $\langle\overline{\Theta_j F},\, G\rangle_j = \mathbb{E}\big[\overline{\Theta_j \overline{\Theta_j F}}\, G\big] = \mathbb{E}[FG]$. □

Proposition 10.17 (zero-energy vectors and invariant functions)**.** *In the setting of Proposition 10.16, for every $G \in L^2_+$:*

(i) *$P_j G \in L^2_+$, $P_{\operatorname{Ker} H_j}[G] = [P_j G]$, and $\|[P_j G]\|^2_{\mathcal{H}_j} = \mathbb{E}_\mu |P_j G|^2$;*

(ii) *$\mathbb{E}[G \mid \mathcal{I}] \in L^2_+$ and $Q_j[G] = [\mathbb{E}[G \mid \mathcal{I}]]$, where Q_j is the projection onto the joint fixed space of $\{T_j(t)\}_{t\ge 0} \cup \{V_j(y)\}_{y \perp e_j}$;*

(iii) *if μ is ergodic for the $\mathbb{R}^3$-action, the joint fixed space is $\mathbb{C}\Omega_j$.*

Proof. (i) $P_j G \in L^2_+$ by Lemma 10.8(i). The quotient map is bounded, so it commutes with Bochner integrals: $T^{-1}\int_0^T T_j(t)[G]\, dt = [T^{-1}\int_0^T U(te_j)G\, dt]$. As $T \to \infty$ the left side tends to $P_{\operatorname{Ker} H_j}[G]$ (spectral theorem, $T^{-1}\int_0^T e^{-t\lambda} dt \to \mathbf{1}_{\{\lambda=0\}}$ boundedly on $[0,\infty)$) and the right side to $[P_j G]$ (one-sided means, Lemma 10.7(i)). Finally $\|[P_j G]\|^2 = \mathbb{E}[\overline{\Theta_j P_j G}\, P_j G] = \mathbb{E}|P_j G|^2$ by Lemma 10.8(ii).

(ii) $\mathbb{E}[G \mid \mathcal{I}] = P_{\{1,2,3\}}G \in L^2_+$ by Lemma 10.8(i). The means of $W(a) := T_j(a_j)V_j(a_\perp)$ over $a \in [0,T] \times [-T,T]^2$ factor into three commuting means and converge strongly to the product of the projections onto the three fixed spaces, which is Q_j. Applying the same means to $[U(a)G] = W(a)[G]$ and using the argument of (i), the limit is $[P_{\{1,2,3\}}G]$.

(iii) If μ is ergodic, $\mathbb{E}[G \mid \mathcal{I}] = \mathbb{E}[G]$ a.s., so $Q_j[G] = \mathbb{E}[G]\,\Omega_j = \langle \Omega_j,\, [G]\rangle \Omega_j$. Since $[L^2_+]$ is dense in $\mathcal{H}_j$, Q_j is the projection onto $\mathbb{C}\Omega_j$. □

Remark 10.18. $\operatorname{Ker} H_j$ alone can be larger than $\mathbb{C}\Omega_j$ on an ergodic component: by Proposition 10.17(i) it is the image of the e_j-invariant functions, which need not be constant (Example 10.22). Ergodicity controls only the joint fixed space.

10.6. **Clustering.** "On an ergodic component" means: for μ_ξ, ν-a.e. ξ, in the decomposition of Theorem 10.10. Each such μ_ξ satisfies (H1)–(H3) and is ergodic.

Theorem 10.19 (clustering in the mean)**.** *For all $G, G' \in L^2(\mu)$ and $\Lambda_T = [-T,T]^3$,*

$$\lim_{T\to\infty} \frac{1}{|\Lambda_T|} \int_{\Lambda_T} \mathbb{E}_\mu\big[G'\, U(a) G\big]\, da = \mathbb{E}_\mu\big[\mathbb{E}[G' \mid \mathcal{I}]\, \mathbb{E}[G \mid \mathcal{I}]\big] = \int_\Xi \mathbb{E}_{\mu_\xi}[G']\, \mathbb{E}_{\mu_\xi}[G]\, d\nu(\xi).$$

On an ergodic component the right side is $\mathbb{E}_{\mu_\xi}[G']\,\mathbb{E}_{\mu_\xi}[G]$. For $\emptyset \neq J \subset \{1,2,3\}$ the first equality holds with Λ_T replaced by Λ_T^J and $\mathcal{I}$ by $\mathcal{I}_J$ (Lemma 10.7(ii)). For $J \neq \{1,2,3\}$ the right side need not factorise on an ergodic component (Example 10.22).

Proof. By Lemma 10.7(i)–(ii) the means of $U(a)G$ converge in L^2 to $\mathbb{E}[G \mid \mathcal{I}]$, and $\mathbb{E}[G'\,\mathbb{E}[G \mid \mathcal{I}]] = \mathbb{E}[\mathbb{E}[G' \mid \mathcal{I}]\,\mathbb{E}[G \mid \mathcal{I}]]$. The integral form is Theorem 10.10(a). □

Theorem 10.20 (clustering under diagonal separation)**.** *For all $G, G' \in L^2(\mu)$,*

$$\lim_{\min_j |a_j| \to \infty} \mathbb{E}_\mu\big[G'\, U(a) G\big] = \mathbb{E}_\mu\big[\mathbb{E}[G' \mid \mathcal{I}]\, \mathbb{E}[G \mid \mathcal{I}]\big].$$

On an ergodic component the limit is $\mathbb{E}_{\mu_\xi}[G']\,\mathbb{E}_{\mu_\xi}[G]$.

Proof. Step 1 (reductions). Since $|\mathbb{E}[G'U(a)G] - \mathbb{E}[G'_n U(a) G_n]| \le \|G' - G'_n\|_2 \|G\|_2 + \|G'_n\|_2 \|G - G_n\|_2$ uniformly in a, and the right side of the claim is continuous in (G, G') in the same way, it suffices to take $G \in L^2(\Sigma(K))$ and $G' \in L^2(\Sigma(K'))$ with K, K' bounded (H2). A sequence $a^{(n)}$ with $\min_j |a_j^{(n)}| \to \infty$ splits into finitely many subsequences on which the signs $\sigma_j = \operatorname{sgn} a_j^{(n)}$ are fixed. On each, replace (G', G, a) by $(\Theta_j G', \Theta_j G, \rho_j a)$ for those j with $\sigma_j = -1$; this leaves $\mathbb{E}[G'U(a)G]$ unchanged by θ_j-invariance and (10.3), and reflects the localisation regions. So assume $a_j \to +\infty$ for every j. By translation invariance, $\mathbb{E}[G'U(a)G] = \mathbb{E}[(U(b)G')\,U(a)(U(b)G)]$; choosing b we may assume that K' lies in the open negative octant $\bigcap_j H_j^-(0)$ and that $K \subset \{x : |x_j| < R\ \forall j\}$ for some R.

Step 2 (decomposition). For the ordering $(1,2,3)$,

$$1 = P_{\{1,2,3\}} + (1 - P_1) + P_1(1 - P_2) + P_1 P_2 (1 - P_3). \tag{10.6}$$

The first term contributes $\mathbb{E}[G'\,U(a)\mathbb{E}[G \mid \mathcal{I}]] = \mathbb{E}[\mathbb{E}[G' \mid \mathcal{I}]\,\mathbb{E}[G \mid \mathcal{I}]]$ for every a, since $\mathbb{E}[G \mid \mathcal{I}]$ is invariant. It remains to show that each of the other three terms tends to 0 as $a_\ell \to \infty$, uniformly in the other coordinates of a, where ℓ is the index of the factor $(1 - P_\ell)$.

Step 3 (one term). Fix ℓ, let $J = \{1, \dots, \ell - 1\}$ and $\tilde{G} = P_J G$ ($\tilde{G} = G$ for $\ell = 1$). The term is $P_J(1 - P_\ell)G = (1 - P_\ell)\tilde{G}$. By Lemma 10.8(iii), $\tilde{G} \in L^2(\Sigma(K + V_J))$, and $K + V_J \subset \{x_\ell > -R\}$ because $e_\ell \notin V_J$. Put $s = 2R$, $a_\ell \ge s$, and $Y := U(s e_\ell)\tilde{G} \in L^2(\Sigma(H_\ell^+(R))) \subset L^2_+$ (half spaces for direction ℓ). Since P_ℓ commutes with translations,

$$U(a)(1 - P_\ell)\tilde{G} = U\big((a_\ell - s)e_\ell + a_\perp\big)(1 - P_\ell)Y,$$

and Y, $P_\ell Y$ lie in L^2_+ (Lemma 10.8(i)). Since $G' \in L^2(\Sigma(H_\ell^-(0)))$, Proposition 10.16(ii)–(iii) and Proposition 10.17(i) give, with $F := \overline{\Theta_\ell G'} \in L^2_+$,

$$\mathbb{E}\big[G'\,U(a)(1-P_\ell)\tilde G\big] = \Big\langle [F],\ V_\ell(a_\perp)\,T_\ell(a_\ell - s)\,(1-P_{\operatorname{Ker} H_\ell})[Y]\Big\rangle_{\mathcal{H}_\ell}.$$

Its modulus is at most $\|[F]\|\,\|e^{-(a_\ell - s)H_\ell}(1-P_{\operatorname{Ker} H_\ell})[Y]\|$, which tends to 0 as $a_\ell \to \infty$ by the spectral theorem and dominated convergence, uniformly in $a_\perp$. □

Proposition 10.21 (separation along a coordinate subgroup)**.** *Let $\emptyset \neq J \subset \{1,2,3\}$ and $G, G' \in L^2(\mu)$. Then, writing $a = a_J + a_{J^c}$ with $a_J \in V_J$,*

$$\mathbb{E}_\mu\big[G'\,U(a)G\big] - \mathbb{E}_\mu\big[G'\,U(a_{J^c})\,P_J G\big] \to 0 \qquad \text{as } \min_{j\in J}|a_j| \to \infty,$$

uniformly in a_{J^c}. For $j \in \{1,2,3\}$ and $G \in L^2(\Sigma(H_j^+(0)),\mu)$, the function $t \mapsto \mathbb{E}_\mu[\overline{\Theta_j G}\,U(te_j)G] = \langle [G],\, e^{-tH_j}[G]\rangle_{\mathcal{H}_j}$, $t \geq 0$, is completely monotone and decreases to $\mathbb{E}_\mu|P_jG|^2$; it tends to 0 if and only if $P_jG = 0$.

Proof. Apply (10.6) with $\{1,2,3\}$ replaced by J in any ordering; Step 3 of the proof of Theorem 10.20 applies verbatim to each term carrying a factor $(1-P_\ell)$, $\ell \in J$, after the reductions of Step 1, which do not involve a_{J^c}. The two-point statement is Proposition 10.16(ii) and Proposition 10.17(i), with the spectral measure of H_j in the vector $[G]$. □

10.7. **Single-axis clustering: the obstruction and what removes it.**

Example 10.22 (the axis obstruction)**.** Let $m > 0$ and let ψ_1, ψ_2, ψ_3 be independent copies of the massive free field on $\mathbb{R}^2$, the centred Gaussian law on $\mathcal{S}'(\mathbb{R}^2)$ with covariance $(-\Delta_2 + m^2)^{-1}$. For $f \in \mathcal{S}(\mathbb{R}^3)$ let $f^{(j)} \in \mathcal{S}(\mathbb{R}^2)$ be the integral of f over the coordinate x_j, and put

$$\Phi(f) := \psi_1(f^{(1)}) + \psi_2(f^{(2)}) + \psi_3(f^{(3)}),$$

the random distribution $\psi_1(x_2,x_3) + \psi_2(x_1,x_3) + \psi_3(x_1,x_2)$. Its law μ is the centred Gaussian measure with covariance $C = C_1 + C_2 + C_3$, $\widehat{C_j}(p) = 2\pi\,\delta(p_j)\,(|p_{\perp j}|^2 + m^2)^{-1}$, where $p_{\perp j}$ is the projection of p onto $e_j^\perp$. Realise it on the field space $\mathcal{V} = \Lambda^2\mathbb{R}^3 \oplus \mathbb{R}$ by putting the 2-form component equal to 0 and the scalar component equal to Φ, on $X = H^{-s}_{\mathrm{loc}}(\mathbb{R}^3;\mathcal{V})$, $s > 1$. By the carrier argument of Lemma 10.5(b), with $\mathbb{E}\Phi(f)^2 \leq m^{-2}\sum_j \|f^{(j)}\|^2_{L^2(\mathbb{R}^2)} \leq c\|f\|_2^2$ (Cauchy–Schwarz inequality in x_j with weight $1 + x_j^2$), μ is also carried by $\mathcal{H}^{-r}$ for $r > 5$, as Proposition 3.8 states. Then:

(i) μ is invariant under translations and under O_h, and has Gaussian moments, so every moment bound of the form $|\mathbb{E}\prod_{i\leq n}\Phi(f_i)| \leq C^n (n!)^{1/2}\prod_i \|f_i\|$ holds with the norm $\sum_j \|f^{(j)}\|_{L^2(\mathbb{R}^2)}$. This norm is not bounded by $\|f\|_{L^2(\mathbb{R}^3)}$; a bound $\mathbb{E}\Phi(f)^2 \leq c\|f\|^2_{L^2(\mathbb{R}^3)}$ would give $\mathbb{E}e^{\Phi(f)} \leq e^{c\|f\|_2^2/2}$ for this Gaussian law, and Lemma 10.25, with ϖ equal to c times Lebesgue measure on the scalar component, would force $P_1\Phi(f) = \mathbb{E}[\Phi(f) \mid \mathcal{I}] = 0$ (μ is ergodic by (iii)), contradicting (iv).

(ii) μ satisfies (H1)–(H3) with the reflections of Section 10.2. On the scalar component these are the plain reflections $h \mapsto h \circ \rho_j$, and the 2-form component vanishes, so the twisted and the plain reflections coincide.

(iii) μ is ergodic for the $\mathbb{R}^3$-action.

(iv) For $f \in C_c^\infty(\mathbb{R}^3)$ with $f^{(1)} \neq 0$, $\mathbb{E}[\Phi(f)\,U(te_1)\Phi(f)] \to \langle f^{(1)}, (-\Delta_2 + m^2)^{-1} f^{(1)}\rangle > 0$ as $t \to \infty$. Thus $P_1\Phi(f) = \psi_1(f^{(1)}) \neq 0$, and the two-point function does not cluster along the coordinate axes, although it clusters under diagonal separation (Theorem 10.20).

Proof. Φ takes values in X: $\chi(x)\psi_1(x_2,x_3)$ lies in $H^{-s}(\mathbb{R}^3)$ for $\chi \in C_c^\infty(\mathbb{R}^3)$ when $\psi_1 \in H^{-s}_{\mathrm{loc}}(\mathbb{R}^2)$, which holds a.s. for $s > 1$, by a Fourier-space bound in which the weight $(1+|p|^2)^{-s} \leq (1+|p_\perp|^2)^{-s}$ is used.

(i) Translations act on ψ_j through the translations of its plane, under which the free field is invariant. Signed permutations of the coordinates permute the three planes and act on each by an element of $O(2)$; the ψ_j are identically distributed and $O(2)$-invariant. Moments are Gaussian with $\langle f,\, C_j f\rangle = \langle f^{(j)},\, (-\Delta_2 + m^2)^{-1} f^{(j)}\rangle \le m^{-2}\|f^{(j)}\|^2_{L^2(\mathbb{R}^2)}$.

(ii) (H1)–(H2) as in the proof of Proposition 10.4. For (H3) it suffices (by Lemma J.3) that $\langle \rho_j f,\, Cf\rangle \ge 0$ for real $f \in C_c^\infty(H_j^+(0))$. For C_j: $(\rho_j f)^{(j)} = f^{(j)}$, so $\langle \rho_j f,\, C_j f\rangle = \langle f^{(j)},\, (-\Delta_2 + m^2)^{-1} f^{(j)}\rangle \ge 0$. For $k \ne j$: $(\rho_j f)^{(k)}$ is the reflection of $f^{(k)}$ in the coordinate x_j of the plane $e_k^\perp$, and $f^{(k)}$ is supported in $\{x_j > 0\}$, so $\langle \rho_j f,\, C_k f\rangle \ge 0$ is reflection positivity of the two-dimensional massive free field [19, Prop. 6.2.5, p. 103] (the passage from the covariance to the Gaussian measure is [19, Thm. 6.2.2, p. 101] and Lemma J.3). Sums of positive forms are positive.

(iii) By the Wiener–Itô chaos decomposition, $L^2(\mu) = \bigoplus_{n\ge 0} \mathcal{H}^{:n:}$, and $\mathcal{H}^{:n:}$ is unitarily equivalent to the symmetric subspace of $L^2((\mathbb{R}^3)^n, \widehat{C}^{\otimes n})$, with $U(a)$ acting by multiplication by $e^{-ia\cdot(p_1+\cdots+p_n)}$ [22, Thm. 2.6 and Ch. 4]. A vector fixed by all $U(a)$ in $\mathcal{H}^{:n:}$, $n \ge 1$, is supported on $Z_n = \{p_1 + \cdots + p_n = 0\}$. The measure $\widehat{C}$ is a sum of three measures, the j-th supported on the plane $\pi_j = \{p_j = 0\}$ and absolutely continuous with respect to its two-dimensional Lebesgue measure. For each choice $(j_1, \dots, j_n)$ the map $(p_1, \dots, p_n) \mapsto \sum_i p_i$ from $\pi_{j_1} \times \cdots \times \pi_{j_n}$ to $\mathbb{R}^3$ is linear and onto a subspace of dimension ≥ 2, so its kernel $Z_n \cap (\pi_{j_1} \times \cdots \times \pi_{j_n})$ has codimension ≥ 2 there and is null for the product of the planar Lebesgue measures. Hence $\widehat{C}^{\otimes n}(Z_n) = 0$, the only invariant vectors are the constants, and μ is ergodic.

(iv) $\mathbb{E}[\Phi(f)\, U(te_1)\Phi(f)] = \sum_k \langle f,\, C_k T_{te_1} f\rangle$. For $k = 1$, $(T_{te_1} f)^{(1)} = f^{(1)}$, giving the constant $\langle f^{(1)},\, (-\Delta_2 + m^2)^{-1} f^{(1)}\rangle$, which is positive when $f^{(1)} \ne 0$. For $k = 2, 3$, $(T_{te_1} f)^{(k)}$ is $f^{(k)}$ translated by t in the plane, and the term tends to 0 because the two-dimensional massive Green function decays. The statement on P_1 follows from Proposition 10.21. □

Proposition 10.23 (what removes the obstruction)**.** *Let μ satisfy (H1)–(H3) and let $\mathcal{F} \subset L^2(\mu)$ be a set whose linear span is dense. The following are equivalent:*

(i) *$P_j G = \mathbb{E}[G \mid \mathcal{I}]$ for all $G \in \mathcal{F}$ and $j = 1, 2, 3$;*
(ii) *for all $G \in \mathcal{F}$, $G' \in L^2(\mu)$ and each j: $\mathbb{E}[G'\, U(te_j) G] \to \mathbb{E}[\mathbb{E}[G' \mid \mathcal{I}]\, \mathbb{E}[G \mid \mathcal{I}]]$ as $|t| \to \infty$;*
(iii) *for all $G \in \mathcal{F}$, $G' \in L^2(\mu)$: $\mathbb{E}[G'\, U(a) G] \to \mathbb{E}[\mathbb{E}[G' \mid \mathcal{I}]\, \mathbb{E}[G \mid \mathcal{I}]]$ as $|a| \to \infty$ in any manner.*

If μ is a centred Gaussian law with covariance C and $\mathcal{F}$ is the set of linear fields $\Phi(f)$, then (i) holds if and only if $\widehat{C}(\{p : p_j = 0\} \setminus \{0\}) = 0$ for $j = 1, 2, 3$.

Proof. (i)⇒(iii): given $|a^{(n)}| \to \infty$, every subsequence has a further subsequence along which each coordinate converges or diverges in modulus, at least one, say the j-th, diverging. Proposition 10.21 with $J = \{j\}$ reduces the limit to that of $\mathbb{E}[G' U(a_{J^c}) P_j G] = \mathbb{E}[G'\, \mathbb{E}[G \mid \mathcal{I}]]$. (iii)⇒(ii) is trivial. (ii)⇒(i): by Proposition 10.21 the limit in (ii) is $\mathbb{E}[G' P_j G]$; so $\mathbb{E}[G'(P_j G - \mathbb{E}[G \mid \mathcal{I}])] = 0$ for all $G' \in L^2$, i.e. $P_j G = \mathbb{E}[G \mid \mathcal{I}]$. The Gaussian clause: the first chaos $\mathcal{H}^{:1:}$ is invariant under U and unitarily equivalent to the closure of $\{\hat f\}$ in $L^2(\mathbb{R}^3, \widehat{C})$, with $U(te_j)$ acting by multiplication by e^{-itp_j}. So $P_j \Phi(f)$ corresponds to $\mathbf{1}_{\{p_j = 0\}} \hat f$ and $\mathbb{E}[\Phi(f) \mid \mathcal{I}]$ to $\mathbf{1}_{\{p = 0\}} \hat f$; they agree for all f exactly when $\widehat{C}$ gives no mass to $\{p_j = 0\} \setminus \{0\}$. □

10.8. **Single-axis clustering.** Throughout this subsection S is one of the limit states S^{μ_0} ($\mu_0 \in Q_0$), $S^\flat$, S^0 of Theorem 3.1, and $\mu = \mu_S$ is its law on $X = H^{-s}_{\mathrm{loc}}(\mathbb{R}^3; \mathcal{V})$ (Lemma 10.3); everything holds verbatim on $\mathcal{H}^{-r}$ (Lemma 10.5). We write $F(\mathbf{g})$ for the curvature field and $X_{\mathrm{R}}(h)$ for the Wick square, $h \in C_c^\infty(\mathbb{R}^3)$ real; $\mathcal{M}$ is the σ-algebra generated by the $X_{\mathrm{R}}(h)$ and completed by μ-null sets, and $L^2(\mathcal{M}) \subset L^2(\mu)$ is the corresponding closed subspace. P_j is the projection onto the e_j-invariant vectors (Section 10.3). We say that a pair (G', G) in $L^2(\mu)$ *clusters along the axes* if $\mathbb{E}_\mu[G'\, U(a) G] \to \mathbb{E}_\mu[G']\, \mathbb{E}_\mu[G]$ as $|a| \to \infty$ in any manner; in particular as $a = te_j$, $|t| \to \infty$, for

each j. If every pair in $L^2(\mu)$ clusters along the axes, then μ satisfies (OS4$'_j$) of Definition 2.11 for $j = 1, 2, 3$; under (H1)–(H3) this is equivalent to OS4 (Lemma 10.24 below, and Remark 2.13(c)).

The single tool that turns invariant parts into clustering is the following reformulation of Propositions 10.21 and 10.23.

Lemma 10.24 (clustering from vanishing invariant parts)**.** *Let μ satisfy (H1)–(H3) and $G \in L^2(\mu)$. If $P_jG = \mathbb{E}_\mu G$ for $j = 1, 2, 3$, then (G', G) clusters along the axes for every $G' \in L^2(\mu)$. Conversely, let $\mathcal{S} \subset L^2(\mu)$ be a closed subspace, closed under complex conjugation, containing 1 and P_jG (for instance $\mathcal{S} = L^2(\mu)$). If $\mathbb{E}[G'\,U(te_j)G] \to \mathbb{E}[G']\mathbb{E}[G]$ as $t \to \infty$ for every $G' \in \mathcal{S}$, then $P_jG = \mathbb{E}_\mu G$.*

Proof. Let $|a^{(n)}| \to \infty$. Every subsequence has a further subsequence along which some coordinate, say $a_j^{(n)}$, diverges in modulus. Proposition 10.21 with $J = \{j\}$ gives $\mathbb{E}[G'U(a^{(n)})G] - \mathbb{E}[G'U(a_{J^c}^{(n)})P_jG] \to 0$, and $\mathbb{E}[G'U(a_{J^c})P_jG] = \mathbb{E}[G']\mathbb{E}[G]$ because P_jG is a constant. So every subsequence has a further subsequence along which the limit is $\mathbb{E}G'\,\mathbb{E}G$. Conversely, by Proposition 10.21 the limit of $\mathbb{E}[G'U(te_j)G]$ is $\mathbb{E}[G'P_jG]$; so $\mathbb{E}[G'(P_jG - \mathbb{E}G)] = 0$ for every $G' \in \mathcal{S}$. Taking $G' = \overline{P_jG - \mathbb{E}G} \in \mathcal{S}$ gives $P_jG = \mathbb{E}G$. □

10.8.1. *The curvature sector, and curvature times Wick square.* The first lemma makes precise what a Gaussian domination of the linear field gives in general.

Lemma 10.25 (domination of the linear field)**.** *Let μ satisfy (H1)–(H2), and let $f \mapsto \Phi(f) \in L^2(\mu)$, $f \in C_c^\infty(\mathbb{R}^3; \mathcal{V})$ real, be linear with $U(a)\Phi(f) = \Phi(f(\cdot - a))$. Suppose $\mathbb{E}_\mu e^{\Phi(f)} \le e^{\frac{1}{2}C_0(f,f)}$ for all real f, where $C_0(f,f) = \int \hat f(p)^* \varpi(dp) \hat f(p)$ for a tempered measure ϖ on $\mathbb{R}^3$ with values in the positive semidefinite 4×4 matrices. If $\varpi(\{p : p_j = 0\} \setminus \{0\}) = 0$, then $\mathbb{E}\Phi(f) = 0$ and $P_j\Phi(f) = \mathbb{E}[\Phi(f) \mid \mathcal{I}]$ for every f.*

Proof. For real t, $\mathbb{E}e^{t\Phi(f)} \le e^{\frac{1}{2}t^2 C_0(f,f)}$; both sides equal 1 at $t = 0$ and are differentiable there (all exponential moments of $\Phi(f)$ are finite), so comparing derivatives at $t = 0$ gives $\mathbb{E}\Phi(f) = 0$ and $\mathbb{E}\Phi(f)^2 \le C_0(f,f)$. Since ϖ is tempered, $C_0(f,f) \le C_K \|f\|_{C^k}^2$ for $\operatorname{supp} f$ in a fixed compact K and some k; hence $f \mapsto \Phi(f)$ is continuous from $C_c^\infty(K)$ with the C^k norm into $L^2(\mu)$. Put $f_T := T^{-1}\int_0^T f(\cdot - te_j)\,dt$ and $f_T^\square := (2T)^{-3}\int_{[-T,T]^3} f(\cdot - a)\,da$, both in C_c^∞. Riemann sums of these integrals converge in C^k with supports in a fixed compact, so $\Phi(f_T) = T^{-1}\int_0^T U(te_j)\Phi(f)\,dt$ and $\Phi(f_T^\square) = (2T)^{-3}\int U(a)\Phi(f)\,da$ in $L^2(\mu)$, and by Lemma 10.7 they converge to $P_j\Phi(f)$ and to $P_{\{1,2,3\}}\Phi(f) = \mathbb{E}[\Phi(f) \mid \mathcal{I}]$. Now $\widehat{f_T}(p) = m_T(p_j)\hat f(p)$ and $\widehat{f_T^\square}(p) = m_T^\square(p)\hat f(p)$ with $m_T(s) = T^{-1}\int_0^T e^{-its}dt$ and $m_T^\square(p) = \prod_k \frac{\sin Tp_k}{Tp_k}$. Hence

$$\|P_j\Phi(f) - \mathbb{E}[\Phi(f) \mid \mathcal{I}]\|_2^2 = \lim_{T\to\infty} \mathbb{E}\,\Phi(f_T - f_T^\square)^2 \le \liminf_{T\to\infty} \int |m_T(p_j) - m_T^\square(p)|^2\, \hat f(p)^* \varpi(dp) \hat f(p).$$

The integrand is bounded by $4\hat f^* \varpi \hat f$, a finite measure, and tends to 0 at $p = 0$ (both multipliers equal 1), at every p with $p_j \ne 0$ (both tend to 0), and to 1 on $\{p_j = 0\} \setminus \{0\}$. By dominated convergence the limit is $\int_{\{p_j=0\}\setminus\{0\}} \hat f^* \varpi \hat f = 0$. □

For the limit states the curvature satisfies more than Lemma 10.25: its domination holds conditionally on the matter, on every lattice torus.

Lemma 10.26 (conditional curvature domination on the torus)**.** *Fix (ε, T_j) and $\mu_0 \in (0, \mu_*] \cup \{\flat\}$, and write $\langle\cdot\rangle$ for $\langle\cdot\rangle_{\varepsilon,T_j,\mu_0}$ and $\mathcal{R}$ for the σ-algebra generated by the radial variables $(r_x)_x$, $r_x := |\varphi(x)|$. For every real lattice 2-form $\mathbf{g}$ and every bounded Borel function $Y \ge 0$ of $(r_x)_x$,*

$$\langle e^{F(\mathbf{g})} Y\rangle \le e^{\frac{1}{2}V(\mathbf{g})}\langle Y\rangle, \qquad V(\mathbf{g}) \le \|\mathbf{g}\|_{2,\varepsilon}^2, \tag{10.7}$$

with V as in Lemma 7.3. Consequently, for real lattice 2-forms $\mathbf{g}_1,\dots,\mathbf{g}_m$ and bounded real $\mathcal{R}$-measurable $Y_1,\dots,Y_m$,

$$\Big\langle \Big(\sum_i Y_i F(\mathbf{g}_i)\Big)^2 \Big\rangle \le \sum_{i,k} \langle Y_i Y_k\rangle \,\langle \mathbf{g}_i,\, \mathbf{g}_k\rangle_\varepsilon, \qquad \langle Y_1 F(\mathbf{g}_1)\rangle = 0. \tag{10.8}$$

Proof. (10.7). In Lemma 2.5 replace the factor $e^{X_{\mathrm{R}}(k)}$ by Y: since Y depends on the radial variables only, the expansion in the proof of Lemma 2.5 gives $\int \prod_x d^2\varphi(x)\, e^{-S_b^{\mathrm{m}}(A,\varphi)}\, Y = \sum_n w_Y(n) e^{ie\varepsilon \sum_b n_b A_b}$ with

$$w_Y(n) = (2\pi)^{|T_{j,\varepsilon}|} \int \prod_x v_x(r_x)\, r_x\, dr_x\; Y(r) \prod_b I_{n_b}(\varepsilon r_{b_-} r_{b_+}) \;\ge 0, \qquad w_Y(-n) = w_Y(n),$$

where v_x is the single-site weight of that proof with $k=0$, and with absolute convergence because Y is bounded. The proof of Lemma 7.3 uses only these two properties of the weight (in the $\flat$ case together with the winding selection, which keeps them). It gives $\langle e^{F(\mathbf{g})} Y\rangle \le e^{\frac12 V(\mathbf{g})}\langle Y\rangle$, and $V(\mathbf{g}) \le \|\mathbf{g}\|_{2,\varepsilon}^2$ is the bound of that lemma.

(10.8). Let $\mathbf{F} := (F(\mathbf{g}_1),\dots,F(\mathbf{g}_m))$ and let $\pi(r, dy)$ be a regular conditional law of $\mathbf{F}$ given $\mathcal{R}$. Put $Q(c) := V(\sum_i c_i \mathbf{g}_i)$, a positive semidefinite quadratic form on $\mathbb{R}^m$. Applying (10.7) to $\sum_i c_i \mathbf{g}_i$ and all bounded $Y \ge 0$ gives $\int e^{c\cdot y}\pi(r,dy) \le e^{\frac12 Q(c)}$ for almost every r, first for all $c \in \mathbb{Q}^m$ off one null set, then for all $c \in \mathbb{R}^m$ by Fatou's lemma. Fix such an r and c. The function $\phi(t) = \int e^{tc\cdot y}\pi(r,dy)$ is finite for all t, hence smooth, and $\phi(t) \le e^{\frac12 t^2 Q(c)}$ with equality at $t=0$. So $\phi'(0) = 0$ and $\phi''(0) \le Q(c)$: the conditional mean of $\mathbf{F}$ vanishes and its conditional covariance matrix $\Sigma(r)$ satisfies $c^{\mathsf T}\Sigma(r) c \le Q(c)$ for all c. Given $\mathcal{R}$, the Y_i are constants, so

$$\Big\langle \Big(\sum_i Y_i F(\mathbf{g}_i)\Big)^2 \Big\rangle = \Big\langle \sum_{i,k} Y_i Y_k \Sigma_{ik}(r) \Big\rangle \le \Big\langle V\Big(\sum_i Y_i \mathbf{g}_i\Big)\Big\rangle \le \Big\langle \Big\| \sum_i Y_i \mathbf{g}_i \Big\|_{2,\varepsilon}^2 \Big\rangle = \sum_{i,k} \langle Y_i Y_k\rangle \langle \mathbf{g}_i,\, \mathbf{g}_k\rangle_\varepsilon,$$

and $\langle Y_1 F(\mathbf{g}_1)\rangle = \langle Y_1 \int y_1 \pi(r,dy)\rangle = 0$. □

Theorem 10.27 (curvature times Wick square: no invariant part)**.** *Let S and $\mu = \mu_S$ be as above. For every real $\mathbf{g} \in C_c^\infty(\mathbb{R}^3;\Lambda^2)$ and every $Y \in L^2(\mathcal{M})$, the product $F(\mathbf{g})Y$ lies in $L^2(\mu)$, with*

$$\mathbb{E}_\mu[F(\mathbf{g})^2 |Y|^2] \le \|\mathbf{g}\|_2^2\, \mathbb{E}_\mu |Y|^2, \qquad \mathbb{E}_\mu[F(\mathbf{g})Y] = 0, \tag{10.9}$$

and $P_j\big(F(\mathbf{g})Y\big) = 0$ for $j = 1,2,3$. Hence every pair $(G', F(\mathbf{g})Y)$, $G' \in L^2(\mu)$, clusters along the axes:

$$\lim_{|a|\to\infty} \mathbb{E}_\mu\big[G'\, U(a)\big(F(\mathbf{g})Y\big)\big] = 0 = \mathbb{E}_\mu[G']\,\mathbb{E}_\mu[F(\mathbf{g})Y].$$

In particular, for every polynomial P in the smeared fields, $S\big(P\cdot F(T_a\mathbf{g})\big) \to 0$ as $|a|\to\infty$. On ν-almost every ergodic component μ_ξ of Theorem 10.10, for all $\mathbf{g}$ and all $\mathcal{M}_0$-measurable Y with $F(\mathbf{g})Y \in L^2(\mu_\xi)$ simultaneously, where $\mathcal{M}_0 := \sigma(X_{\mathrm{R}}(h) : h \in C_c^\infty(\mathbb{R}^3)$ real$)$ (not completed): $P_j^\xi(F(\mathbf{g})Y) = 0$ for $j = 1,2,3$, with P_j^ξ the projection for μ_ξ, $\mathbb{E}_{\mu_\xi}[F(\mathbf{g})Y] = 0$, and every pair $(G', F(\mathbf{g})Y)$, $G' \in L^2(\mu_\xi)$, clusters along the axes for μ_ξ. The bound in (10.9) *is not asserted for the components.*

Proof. Step 1: the inequality (10.8) *in the limit.* Let $\mathbf{g}_1,\dots,\mathbf{g}_m \in C_c^\infty(\mathbb{R}^3;\Lambda^2)$ be real and $Y_i = \psi_i(X_{\mathrm{R}}(h_{i1}),\dots,X_{\mathrm{R}}(h_{iq}))$ with ψ_i bounded continuous on $\mathbb{R}^q$ and $h_{il} \in C_c^\infty(\mathbb{R}^3)$ real. We claim

$$\mathbb{E}_\mu\Big[\Big(\sum_i Y_i F(\mathbf{g}_i)\Big)^2\Big] \le \sum_{i,k} \mathbb{E}_\mu[Y_i Y_k]\,\langle \mathbf{g}_i,\, \mathbf{g}_k\rangle_{L^2}, \qquad \mathbb{E}_\mu[Y_1 F(\mathbf{g}_1)] = 0. \tag{10.10}$$

For $S = S^{\mu_0}$ or $S^\flat$, let $\langle\cdot\rangle_n$ be the lattice states of the defining subsequence and $Y_i^{(n)}$, $F_n(\mathbf{g}_i)$ the lattice observables with sampled test functions; the $Y_i^{(n)}$ are bounded functions of $(r_x)_x$, so (10.8)

holds for them. The characteristic functionals of the finite vectors $(F_n(\mathbf{g}_i), X_{\mathrm{R}}(h_{il}))_{i,l}$ converge to those under μ (Proposition 8.2(b) at imaginary points, Theorem 3.1(ii)), so these vectors converge in law; the fourth moments of $F_n(\mathbf{g}_i)$ are bounded uniformly in n (Proposition 8.1), so the products $F_n(\mathbf{g}_i)F_n(\mathbf{g}_k)Y_i^{(n)}Y_k^{(n)}$ are uniformly integrable. Hence both sides of (10.8) converge to the two sides of (10.10), using $\langle \mathbf{g}_i, \mathbf{g}_k\rangle_\varepsilon \to \langle \mathbf{g}_i, \mathbf{g}_k\rangle_{L^2}$ (Riemann sums). For $S = S^0$, (10.10) holds for every S^μ, $\mu \in Q_0$, and passes to $\mu \downarrow 0$ in the same way, since $\mu_{S^\mu} \to \mu_{S^0}$ weakly with uniformly bounded moments (Lemma 10.3).

Step 2: (10.9). Bounded continuous cylinder functions Y of the Wick square are dense in $L^2(\mathcal{M})$ (Lemma J.2). For them (10.10) with $m = 1$ is (10.9). For complex Y apply this to $\operatorname{Re} Y$ and $\operatorname{Im} Y$ and add. If $Y_n \to Y$ in $L^2(\mathcal{M})$, then $F(\mathbf{g})Y_n$ is Cauchy in $L^2(\mu)$ by (10.9), and along a subsequence $F(\mathbf{g})Y_n \to F(\mathbf{g})Y$ almost surely; so $F(\mathbf{g})Y \in L^2(\mu)$ and (10.9) holds for Y. The map $Y \mapsto F(\mathbf{g})Y$ is bounded from $L^2(\mathcal{M})$ to $L^2(\mu)$, with norm at most $\|\mathbf{g}\|_2$.

Step 3: $P_j(F(\mathbf{g})Y) = 0$. By Step 2 and the boundedness of P_j it suffices to take Y a bounded continuous cylinder function. Put $\mathbf{g}_t := \mathbf{g}(\cdot - te_j)$, $Y_t := U(te_j)Y$ (again such a cylinder function, of the translated test functions) and $D := \operatorname{diam}\operatorname{supp}\mathbf{g}$. Then $U(te_j)(F(\mathbf{g})Y) = F(\mathbf{g}_t)Y_t$, and $t \mapsto F(\mathbf{g}_t)Y_t$ is continuous in $L^2(\mu)$ (Lemma 10.6). So $\mathsf{A}_T := T^{-1}\int_0^T U(te_j)(F(\mathbf{g})Y)\,dt$ is the L^2-limit of the Riemann sums $N^{-1}\sum_{i<N} F(\mathbf{g}_{t_i})Y_{t_i}$, $t_i = iT/N$. By (10.10), since $\langle \mathbf{g}_t, \mathbf{g}_s\rangle = 0$ for $|t-s| > D$ and $|\langle \mathbf{g}_t, \mathbf{g}_s\rangle| \le \|\mathbf{g}\|_2^2$,

$$\mathbb{E}\Big[\Big(\frac{1}{N}\sum_{i<N} F(\mathbf{g}_{t_i})Y_{t_i}\Big)^2\Big] \le \frac{\|Y\|_\infty^2}{N^2}\sum_{i,k<N}|\langle \mathbf{g}_{t_i}, \mathbf{g}_{t_k}\rangle| \le \|Y\|_\infty^2\|\mathbf{g}\|_2^2\Big(\frac{2D}{T} + \frac{1}{N}\Big).$$

Letting $N \to \infty$, $\|\mathsf{A}_T\|_2^2 \le 2D\|Y\|_\infty^2\|\mathbf{g}\|_2^2/T$ for real Y ((10.10) is stated for real Y_i). For complex Y apply this to $\operatorname{Re} Y$ and $\operatorname{Im} Y$, whose one-sided means are the real and imaginary parts of A_T; this gives $\|\mathsf{A}_T\|_2^2 \le 4D\|Y\|_\infty^2\|\mathbf{g}\|_2^2/T$, the bound used in Step 4. By Lemma 10.7(i) (one-sided means), $\mathsf{A}_T \to P_j(F(\mathbf{g})Y)$; hence $P_j(F(\mathbf{g})Y) = 0$. The clustering statement is Lemma 10.24, with $\mathbb{E}[F(\mathbf{g})Y] = 0$ from (10.9). For the statement on polynomials take $Y = 1$ and $G' = P$, which lies in $L^2(\mu)$ because all moments of μ are finite; $U(a)F(\mathbf{g}) = F(T_a\mathbf{g})$.

Step 4: components. Fix an even $r \ge 14$ and take $s = r$, so that Lemma 10.5(b) and Corollary 10.43(iv) hold with the same r. Let $\mathcal{L}_2$ be a countable set of real $\mathbf{g} \in C_c^\infty$ dense in $\mathcal{S}$ for the norm $\|\cdot\|_r$, and $\mathcal{L}_Y$ a countable $\mathbb{Q}$-algebra of real bounded continuous cylinder functions of the Wick square, containing 1 and generating $\mathcal{M}_0$. For $\mathbf{g} \in \mathcal{L}_2$, $Y \in \mathcal{L}_Y$ and $T \in \mathbb{N}$, Step 3 and Theorem 10.10(a) give $\int \|\mathsf{A}_T\|_{L^2(\mu_\xi)}^2 d\nu(\xi) = \|\mathsf{A}_T\|_{L^2(\mu)}^2 \le cT^{-1}$, where A_T is the same function on X and, for ν-a.e. ξ, is the one-sided mean in $L^2(\mu_\xi)$ (the Riemann sums converge in $L^2(\mu)$, hence in $L^2(\mu_\xi)$ along a subsequence, for a.e. ξ). Along $T = k^2$, $\sum_k \int \|\mathsf{A}_{k^2}\|_{L^2(\mu_\xi)}^2 d\nu < \infty$, so $\|\mathsf{A}_{k^2}\|_{L^2(\mu_\xi)} \to 0$ for a.e. ξ, and $P_j^\xi(F(\mathbf{g})Y) = 0$ there (Lemma 10.7(i) for μ_ξ). Off one ν-null set this holds for all $\mathbf{g} \in \mathcal{L}_2$, $Y \in \mathcal{L}_Y$ and j, and $\mathbb{E}_{\mu_\xi} M < \infty$ with M as in Corollary 10.43(iv). Fix such ξ. For fixed $Y \in \mathcal{L}_Y$, $\mathbf{g} \mapsto F(\mathbf{g})Y$ is continuous into $L^2(\mu_\xi)$ for $\|\cdot\|_r$, since $|F(\mathbf{g})| \le M^{1/2}\|\mathbf{g}\|_r$ for all $\mathbf{g}$ simultaneously off one μ_S-null set, hence μ_ξ-a.s. for ν-a.e. ξ (the identification of $F(\mathbf{g})$ with its Hermite series holds for all test functions simultaneously, by the density argument in the proof of Lemma 10.5(b)); so $P_j^\xi(F(\mathbf{g})Y) = 0$ for all $\mathbf{g}$. For fixed $\mathbf{g}$, the set of bounded real $\mathcal{M}_0$-measurable Y with $P_j^\xi(F(\mathbf{g})Y) = 0$ is a vector space containing $\mathcal{L}_Y$ and closed under bounded μ_ξ-a.e. convergence (dominated convergence in $L^2(\mu_\xi)$, as $F(\mathbf{g}) \in L^2(\mu_\xi)$); by the monotone class theorem it contains every bounded real $\mathcal{M}_0$-measurable Y, and by linearity every bounded complex one. Then $\mathbb{E}_{\mu_\xi}[F(\mathbf{g})Y] = \mathbb{E}_{\mu_\xi}[P_j^\xi(F(\mathbf{g})Y)] = 0$, because P_j^ξ is self-adjoint and fixes 1, and Lemma 10.24 for μ_ξ (which satisfies (H1)–(H3), Theorem 10.10) gives the clustering. Finally, let Y be $\mathcal{M}_0$-measurable with $F(\mathbf{g})Y \in L^2(\mu_\xi)$, and put $Y_n := Y\mathbf{1}_{\{|Y|\le n\}}$. Then $F(\mathbf{g})Y_n \to F(\mathbf{g})Y$ in $L^2(\mu_\xi)$ by dominated convergence, so $P_j^\xi(F(\mathbf{g})Y) = 0$; as before $\mathbb{E}_{\mu_\xi}[F(\mathbf{g})Y] = 0$ and Lemma 10.24 gives the clustering.

By continuity of P_j^ξ the same holds on the closed span of such products in $L^2(\mu_\xi)$. This applies to every polynomial Y in the $X_{\mathrm{R}}(h)$, since all moments of μ_ξ are finite (Corollary 10.43(iv)). □

Remark 10.28. For $Y = 1$ the conclusion on μ also follows from the tube bound of OS0 alone. Let $G(z)$ be the tube functional of the one-element family $\{\mathbf{g}\}$ with no source for the Wick square. It is entire and real on $\mathbb{R}$, $G(0) = 1$, $G'(0) = S_{1,0}(\mathbf{g}) = 0$ by OS3 (Theorem 9.8), $G''(0) = \mathbb{E}_\mu F(\mathbf{g})^2$ by Lemma 10.3, and $G(t) \le e^{t^2\|\mathbf{g}\|_2^2/2}$ for real t (Theorem 9.1(a)); comparing second derivatives at 0 gives $\mathbb{E}_\mu F(\mathbf{g})^2 \le \|\mathbf{g}\|_2^2$, so $\mathbf{g} \mapsto F(\mathbf{g})$ extends to a bounded linear map from $L^2(\mathbb{R}^3;\Lambda^2)$ to $L^2(\mu)$. With $\mathbf{g}_T := T^{-1}\int_0^T T_{te_j}\mathbf{g}\,dt$, Riemann sums, this bound and the strong continuity of U give $F(\mathbf{g}_T) = T^{-1}\int_0^T U(te_j)F(\mathbf{g})\,dt$, which tends to $P_jF(\mathbf{g})$ (Lemma 10.7(i)); and

$$\|P_jF(\mathbf{g})\|_{L^2(\mu)}^2 = \lim_{T\to\infty} \mathbb{E}_\mu F(\mathbf{g}_T)^2 \le \liminf_{T\to\infty} \int_{\mathbb{R}^3} |\hat{\mathbf{g}}(p)|^2 \left|\frac{1}{T}\int_0^T e^{-itp_j}\,dt\right|^2 \frac{dp}{(2\pi)^3} = 0$$

by dominated convergence, since $\{p_j = 0\}$ is Lebesgue-null; with $\mathbb{E}_\mu F(\mathbf{g}) = 0$ (OS3), Lemma 10.24 gives the clustering. Theorem 10.27 extends this, by a domination of the curvature conditional on the matter on every lattice torus, to all products $F(\mathbf{g})Y$ with Y a square-integrable function of the Wick square. By itself the OS0 bound says nothing about the Wick square: its part for the Wick square is an L^1-type bound and gives no L^2 control of the invariant part; the Wick-square sector is treated below (Theorem 10.33, under hypothesis (L)). Observables of degree at least two in the curvature are not covered (Remark 10.30). Example 10.22 shows that for a scalar field without a domination bound the conclusion can fail, and Example 10.29 that a domination bound on the linear field does not control quadratic observables. With full rotation invariance the distinction between diagonal and single-axis separation would disappear for every observable, since a rotation maps an axis to a diagonal direction.

The domination of the linear field does not control observables of higher degree in the field. The next example satisfies (H1)–(H3), is invariant under translations and O_h, and its linear field is dominated by white noise, which charges no hyperplane; yet a quadratic observable does not cluster along an axis.

Example 10.29 (domination does not control quadratic observables)**.** Let $0 < b_- < b_+ \le 1$, $\gamma > 0$, and let $\alpha_1, \alpha_2, \alpha_3$ be independent copies of the stationary telegraph process: the right-continuous Markov process on $\{b_-, b_+\}$ that jumps from either value to the other at rate γ, started in its stationary law (uniform on $\{b_-, b_+\}$). Let W be a white noise on $\mathbb{R}^3$, independent of the α_k. Put $\beta(x) := \alpha_1(x_1)\alpha_2(x_2)\alpha_3(x_3)$ and $\Phi(f) := \int \beta f\,dW$ (a Wiener integral, conditionally on the α_k), for real $f \in C_c^\infty(\mathbb{R}^3)$. Realise Φ on $\mathcal{V} = \Lambda^2\mathbb{R}^3 \oplus \mathbb{R}$ with vanishing 2-form component, as in Example 10.22. Then:

(i) the law μ satisfies (H1)–(H3) with the reflections of Section 10.2 and is invariant under O_h;
(ii) $\mathbb{E}e^{\Phi(f)} \le e^{\frac12\|f\|_2^2}$, so Lemma 10.25 applies with ϖ the Lebesgue measure, and every pair $(G', \Phi(f))$ clusters along the axes;
(iii) for $G := \Phi(f)^2$, $f \ne 0$: $\mathbb{E}[G \mid \mathcal{I}] = c_2^3\|f\|_2^2$ is constant, $c_2 := \mathbb{E}\alpha_1(0)^2$, but $P_1G = c_2\int \alpha_2(y_2)^2\alpha_3(y_3)^2 f(y)^2\,dy$ is not; so $\lim_{t\to\infty}\mathbb{E}[G\,U(te_1)G] = \mathbb{E}[(P_1G)^2] > (\mathbb{E}G)^2$, and the pair (G, G) does not cluster along e_1.

Proof. Facts on α. The two-state chain with equal rates is reversible: its stationary law is uniform and satisfies detailed balance, so $(\alpha(-t))_t$ (made right-continuous) has the law of $(\alpha(t))_t$. For $u := \alpha^2$, $\mathbb{E}u(0) = c_2$ and $\mathrm{Cov}(u(s), u(t)) = v\,e^{-2\gamma|t-s|}$ with $v := \big((b_+^2 - b_-^2)/2\big)^2 > 0$, since the probability that the chain is at the same value after time t is $(1 + e^{-2\gamma t})/2$. Hence $\mathrm{Var}\left(T^{-1}\int_0^T u(y+t)\,dt\right) \le v/(\gamma T) \to 0$, uniformly in y.

(i) Conditionally on $\alpha = (\alpha_k)_k$, Φ is a centred Gaussian linear process with $\mathbb{E}[\Phi(f)^2 \mid \alpha] = \|\beta f\|_2^2 \le \|f\|_2^2$. By the carrier argument of Lemma 10.5(b) (with $\mathbb{E}\Phi(f)^2 \le \|f\|_2^2 \le \|f\|_r^2$) Φ has a

version with values in $\mathcal{H}^{-r} \subset X$. (H1)–(H2) hold as in the proof of Proposition 10.4. Invariance: translations act by shifting each α_k and W, whose laws are stationary; the coordinate reflections act by reversing one α_k and reflecting W, which preserves the laws (reversibility); permutations permute the identically distributed α_k. On the scalar component the reflections of Section 10.2 are the plain ones. (H3) for j, at $c = 0$: let $\mathcal{G}$ be generated by α_k ($k \neq j$) and $\alpha_j(0)$, and $\mathcal{F}_\pm$ by W restricted to $\{\pm x_j > 0\}$ and α_j restricted to $\pm t \geq 0$. A $\Sigma(H_j^+(0))$-measurable F is a function of $\Phi(f)$, $\operatorname{supp} f \subset \{x_j > 0\}$, hence $\mathcal{F}_+ \vee \mathcal{G}$-measurable, and $F \circ \theta_j$ is $\mathcal{F}_- \vee \mathcal{G}$-measurable. Given $\mathcal{G}$, $\mathcal{F}_+$ and $\mathcal{F}_-$ are independent (independence of W on disjoint sets, independence of W and α, the Markov property of α_j at time 0). The map of the underlying variables that implements θ_j preserves their law and fixes every $\mathcal{G}$-measurable function (α_j is a.s. continuous at 0), so $\mathbb{E}[F \circ \theta_j \mid \mathcal{G}] = \mathbb{E}[F \mid \mathcal{G}]$. Hence $\mathbb{E}[\overline{F \circ \theta_j}\, F] = \mathbb{E}|\mathbb{E}[F \mid \mathcal{G}]|^2 \geq 0$.

(ii) $\mathbb{E}e^{\Phi(f)} = \mathbb{E}\, e^{\frac{1}{2}\|\beta f\|_2^2} \leq e^{\frac{1}{2}\|f\|_2^2}$ since $|\beta| \leq b_+^3 \leq 1$. So Lemma 10.25 applies with ϖ the Lebesgue measure and gives $P_j\Phi(f) = \mathbb{E}[\Phi(f) \mid \mathcal{I}]$; since ϖ has no atom at 0, the computation in its proof with the multiplier $m_T^{\square}$ alone gives $\mathbb{E}[\Phi(f) \mid \mathcal{I}] = 0$. Lemma 10.24 gives the clustering.

(iii) Write $f_a := f(\cdot - a)$ and $G_a := U(a)G = \Phi(f_a)^2 = \|\beta f_a\|_2^2 + R_a$, with $R_a := \Phi(f_a)^2 - \|\beta f_a\|_2^2$. Conditionally on α the pair $(\Phi(f_a), \Phi(f_{a'}))$ is centred Gaussian, so $\mathbb{E}[R_a R_{a'} \mid \alpha] = 2\langle \beta f_a,\, \beta f_{a'}\rangle^2$, which vanishes when $|a - a'| > D := \operatorname{diam}\operatorname{supp} f$ and is at most $2\|f\|_2^4$. Hence the one-sided means of R_{te_1} and the cube means of R_a tend to 0 in L^2. Next, $\|\beta f_{te_1}\|_2^2 = \int u_1(y_1 + t)u_2(y_2)u_3(y_3)f(y)^2\, dy$ with $u_k = \alpha_k^2$. By the facts on α and Minkowski's integral inequality, $T^{-1}\int_0^T \|\beta f_{te_1}\|_2^2 dt \to c_2 \int u_2(y_2)u_3(y_3)f(y)^2 dy =: A$ in L^2, and the cube means converge to $c_2^3\|f\|_2^2$ (the cube mean factorises into the three one-dimensional means). By Lemma 10.7, $P_1 G = A$ and $\mathbb{E}[G \mid \mathcal{I}] = P_{\{1,2,3\}}G = c_2^3\|f\|_2^2$. Finally

$$\operatorname{Var} A = c_2^2 \iint f(y)^2 f(y')^2 \Big[\big(c_2^2 + ve^{-2\gamma|y_2 - y_2'|}\big)\big(c_2^2 + ve^{-2\gamma|y_3 - y_3'|}\big) - c_2^4\Big] dy\, dy' > 0,$$

since the bracket is positive and $f \neq 0$. By Proposition 10.21, $\mathbb{E}[G\, U(te_1)G] \to \mathbb{E}[GP_1G] = \mathbb{E}A^2 > (\mathbb{E}A)^2 = (\mathbb{E}G)^2$. □

Remark 10.30 (what the curvature bound leaves open). Theorem 10.27 covers every observable of degree one in the curvature (the closed span of the products $F(\mathbf{g})Y$), with an arbitrary L^2 dependence on the Wick square; observables of degree zero, the Wick-square sector, are treated next (Theorem 10.33). It says nothing about $F(\mathbf{g})^2$, $\cos F(\mathbf{g})$ or $F(\mathbf{g})F(\mathbf{g}')Y$. On the lattice, the conditional law of the curvature given the radial field is the free Gaussian field shifted by the imaginary field $ie\, d\, Cj_n$ of a random current n (the representation behind Lemma 2.5); the conditional covariance of the curvature is the free one minus e^2 times a covariance of these currents. The e_j-invariant part of a quadratic observable is therefore governed by the clustering of the current sector, on which no estimate is available, and Example 10.29 shows that the domination (10.7) cannot replace such an estimate (Section 14.5).

10.8.2. *The Wick-square sector.*

Lemma 10.31 (the radial field is associated)**.** *Fix (ε, T_j) and $\mu_0 \in (0, \mu_*]$. The law of $(r_x)_{x \in T_{j,\varepsilon}}$ under $\langle\cdot\rangle_{\varepsilon,T_j,\mu_0}$ is associated: $\langle\phi\psi\rangle \geq \langle\phi\rangle\langle\psi\rangle$ for all coordinatewise non-decreasing functions ϕ, ψ of $(r_x)_x$ with $\langle\phi^2\rangle, \langle\psi^2\rangle < \infty$.*

Proof. The proof combines [8, Thm. 6.1, p. 276] with the FKG inequality for densities on products of totally ordered spaces [19, Sect. 4.4], [30]. By the proof of Lemma 2.5 and Fubini's theorem, the law of $r = (r_x)_x$ has the density $p(r) = Z(r)\prod_x r_x v_x(r_x)$ (up to normalisation, with v_x as there at $k = 0$) on $(0, \infty)^{T_{j,\varepsilon}}$, where

$$Z(r) = \int \mu_C(dA) \int \prod_x d\vartheta_x \prod_b \exp\big(J_b \cos u_b\big), \qquad J_b := \varepsilon r_{b_-} r_{b_+}, \quad u_b := \vartheta_{b_+} - \vartheta_{b_-} + \varepsilon e A_b,$$

and $C = C_{\mu_0} > 0$. Z is smooth and positive. Let $\langle\cdot\rangle_r$ be the probability measure on (A, ϑ) with density $\prod_b e^{J_b \cos u_b}/Z(r)$ with respect to $\mu_C \otimes d\vartheta$. For $x \neq y$,

$$\partial_{r_x}\partial_{r_y} \log Z = \sum_b (\partial_{r_x}\partial_{r_y} J_b)\langle \cos u_b\rangle_r + \sum_{b,b'} (\partial_{r_x} J_b)(\partial_{r_y} J_{b'})\big(\langle \cos u_b \cos u_{b'}\rangle_r - \langle \cos u_b\rangle_r \langle \cos u_{b'}\rangle_r\big).$$

All derivatives of the J_b are ≥ 0. Also $\langle \cos u_b\rangle_r \geq 0$: expand every factor $e^{J\cos u} = \sum_\nu I_\nu(J) e^{i\nu u}$ with $I_\nu \geq 0$; after the angular integration and the Gaussian integral $\mathbb{E}_C e^{ie\varepsilon \sum_b k_b A_b} = e^{-\frac{1}{2}e^2\varepsilon^2\langle k, Ck\rangle} > 0$, $Z(r)\langle e^{iu_b}\rangle_r$ is a sum of nonnegative terms. The truncated correlation of $\cos u_b$ and $\cos u_{b'}$ is ≥ 0 by [8, Thm. 6.1] on the torus graph, wrap-around bonds included ([8] allow periodic conditions on rectangles, and the theorem holds on every finite graph: Lemma B.10 and Proposition B.13), applied with the radial measures $d\rho_x = \delta_{r_x}$ (finite measures with all Gaussian moments, as allowed there), hopping coefficients ε, the Gaussian measure taken to be the law of the bond variables $(e\varepsilon A_b)_b$, which is centred with covariance $e^2\varepsilon^2 C_{\mu_0} > 0$ (the bond variable of [8, Sect. 6.1] already contains this factor), $P = Q = 1$, n, m the coboundaries of the bonds b, b' and f, g the corresponding bond indicators. So $\partial_{r_x}\partial_{r_y} \log p \geq 0$ for $x \neq y$. Let $u := r \wedge r'$, $a := r - u$, $b := r' - u$. Then $a, b \geq 0$ have disjoint supports and $r \vee r' = u + a + b$, so

$$\log p(r\vee r') - \log p(r) - \log p(r') + \log p(r\wedge r') = \int_0^1\!\!\int_0^1 \sum_{x\in \operatorname{supp} a,\ y\in \operatorname{supp} b} a_x b_y\, \partial_{r_x}\partial_{r_y} \log p(u + sa + tb)\, ds\, dt \geq 0,$$

since only $x \neq y$ occurs. For bounded non-decreasing ϕ and ψ, with $\psi > 0$ after adding a constant, the densities $f_2 := p$ and $f_1 := \psi p / \int \psi p$ satisfy $f_1(r\vee r') f_2(r\wedge r') \geq f_1(r) f_2(r')$. Preston's theorem [30, Thm. 3, p. 234], applied on $(0,\infty)^{T_{j,\varepsilon}}$ with Lebesgue measure, gives $\int \phi f_1 \geq \int \phi f_2$, that is, $\langle \phi\psi\rangle \geq \langle\phi\rangle\langle\psi\rangle$. Truncation $\phi \mapsto \max(-M, \min(M, \phi))$ and dominated convergence extend this to L^2. □

Lemma 10.32 (a covariance inequality for associated variables)**.** *Let Z be an associated random vector and $Y_1, \dots, Y_m$, $Y'_1, \dots, Y'_{m'}$ coordinatewise non-decreasing functions of Z in L^2. Let $\phi \in C^1(\mathbb{R}^m)$, $\psi \in C^1(\mathbb{R}^{m'})$ have bounded partial derivatives, $a_i := \|\partial_i \phi\|_\infty$, $b_l := \|\partial_l \psi\|_\infty$. Then*

$$\big|\operatorname{Cov}\big(\phi(Y), \psi(Y')\big)\big| \leq \sum_{i,l} a_i b_l \operatorname{Cov}(Y_i, Y'_l).$$

The inequality is Newman's [26]; we include the short proof.

Proof. The functions $\sum_i a_i y_i \pm \phi(y)$ and $\sum_l b_l y'_l \pm \psi(y')$ are non-decreasing in each variable, so, composed with Y and Y', they are non-decreasing functions of Z in L^2. Association gives $\operatorname{Cov}(\sum a_i Y_i + \epsilon\phi(Y), \sum b_l Y'_l + \epsilon'\psi(Y')) \geq 0$ for all four sign choices $\epsilon, \epsilon' \in \{\pm 1\}$. Adding the choices $(+,-)$ and $(-,+)$ gives $\operatorname{Cov}(\phi(Y), \psi(Y')) \leq \sum a_i b_l \operatorname{Cov}(Y_i, Y'_l)$; adding $(+,+)$ and $(-,-)$ gives the bound for $-\operatorname{Cov}(\phi(Y), \psi(Y'))$. □

Theorem 10.33 (the Wick-square sector reduces to its linear field)**.** *Let S and $\mu = \mu_S$ be as above. Assume*

$$\text{(10.11)} \qquad \textit{(L)} \qquad P_j X_{\mathrm{R}}(h) = \mathbb{E}_\mu X_{\mathrm{R}}(h) \quad \textit{for all real } h \in C_c^\infty(\mathbb{R}^3) \textit{ and } j = 1, 2, 3.$$

Then $P_j G = \mathbb{E}_\mu G$ for every $G \in L^2(\mathcal{M})$ and every j, and every pair (G', G) with $G' \in L^2(\mu)$, $G \in L^2(\mathcal{M})$ clusters along the axes. In particular the restriction of μ to $\mathcal{M}$ is mixing, and for ν-a.e. ergodic component, $\mu_\xi = \mu$ on $\mathcal{M}_0$.

Proof. Step 1: the linear field. Let $h, h' \in C_c^\infty$. By (L) and Lemma 10.24, $\operatorname{Cov}_\mu(X_{\mathrm{R}}(h'), U(a) X_{\mathrm{R}}(h)) \to 0$ as $|a| \to \infty$ in any manner.

Step 2: a covariance bound in the limit. Let $h_1,\dots,h_m,h'_1,\dots,h'_{m'}\in C_c^\infty$ be nonnegative, ϕ,ψ as in Lemma 10.32 and also bounded, and $a\in\mathbb{R}^3$. We claim

$$\big|\operatorname{Cov}_\mu\big(\psi(X_{\mathrm R}(h'_1),\dots),\,\phi(X_{\mathrm R}(h_1(\cdot-a)),\dots)\big)\big|\le\sum_{i,l}a_ib_l\operatorname{Cov}_\mu\big(X_{\mathrm R}(h'_l),X_{\mathrm R}(h_i(\cdot-a))\big).\tag{10.12}$$

For $\mu_0\in(0,\mu_*]$ and every lattice in the defining sequence, $X_{\mathrm R}(k)=\sum_x\varepsilon^3k(x)(r_x^2-W_\varepsilon-z_\varepsilon)$ is a non-decreasing function of $(r_x)_x$ when $k\ge0$, so Lemmas 10.31 and 10.32 give (10.12) for the lattice state, with the sampled test functions. At fixed lattice it passes to $\mu_0\downarrow0$ by Lemma 7.6 (both sides are elementary observables), so it holds for $\flat$ too. Along the defining subsequence the left side converges (convergence in law of finite vectors of smeared fields, as in Step 1 of the proof of Theorem 10.27), and the right side converges by Proposition 8.2(a). For S^0, pass to $\mu\downarrow0$ in Q_0 as there.

Step 3: cylinder functions. Steps 1–2 give $\operatorname{Cov}_\mu(G',U(a)G)\to0$ as $|a|\to\infty$ for $G=\phi(X_{\mathrm R}(h_1),\dots,X_{\mathrm R}(h_m))$, $G'=\psi(X_{\mathrm R}(h'_1),\dots)$ with nonnegative h_i,h'_l, because $U(a)G=\phi(X_{\mathrm R}(h_1(\cdot-a)),\dots)$. Every real $h\in C_c^\infty$ is $h=(h+c\chi)-c\chi$ with $\chi\in C_c^\infty$, $0\le\chi$, $\chi=1$ on $\operatorname{supp}h$ and $c=\|h\|_\infty$, so the functions $e^{i\sum_k t_kX_{\mathrm R}(h_k)}$ with $h_k\ge0$ include all $e^{iX_{\mathrm R}(h)}$; their real and imaginary parts are bounded with bounded derivatives. Their linear span is dense in $L^2(\mathcal M)$ (Lemma J.2). Since $|\operatorname{Cov}(G',U(a)G)-\operatorname{Cov}(G'_n,U(a)G_n)|\le2(\|G'-G'_n\|_2\|G\|_2+\|G'_n\|_2\|G-G_n\|_2)$ uniformly in a, every pair in $L^2(\mathcal M)$ clusters along the axes.

Step 4: invariant parts and general G'. For $G\in L^2(\mathcal M)$, $P_jG\in L^2(\mathcal M)$ (the means stay in this closed subspace), and Step 3 and the converse part of Lemma 10.24 with $\mathcal S=L^2(\mathcal M)$ give $P_jG=\mathbb EG$. Lemma 10.24 then gives clustering against every $G'\in L^2(\mu)$. Mixing on $\mathcal M$ is Step 3. For the components: $\mathbb E[G\mid\mathcal I]=P_{\{1,2,3\}}P_jG=\mathbb E_\mu G$ (Lemma 10.7(ii)), so $\mathbb E_{\mu_\xi}G=\mathbb E_\mu G$ for ν-a.e. ξ by Theorem 10.10(a); taking G in a countable algebra of bounded continuous cylinder functions generating $\mathcal M_0$ gives $\mu_\xi=\mu$ on $\mathcal M_0$. □

Hypothesis (L) is a statement about the axis-invariant part of the linear Wick-square field. The torus pressures of the Wick square bound that part in L^∞, and remove it when they have no corner at the origin.

Definition 10.34 (pressures of the Wick square; latent density)**.** For a lattice state $\langle\cdot\rangle_n=\langle\cdot\rangle_{\varepsilon_n,T_n,\mu_0}$ of the defining sequence of $S\in\{S^{\mu_0},S^\flat\}$ put $\alpha_n(\sigma):=|T_n|^{-1}\log\langle e^{\sigma X_{\mathrm R}(1_{T_n})}\rangle_n$, $\sigma\in\mathbb R$. Let $\mathfrak P(S)$ be the set of pointwise limits on $I:=(-H/2,\sigma_+/2)$ of subsequences of (α_n); here σ_+ and $H=m^2$ are the constants of Theorem 3.1(i), and this constant H is unrelated to the functions $Y\in L^2(\mathcal M)$ of Theorem 10.27. For S^0, let $\mathfrak P(S^0)$ be the set of pointwise limits on I of sequences $\alpha^{(i)}\in\mathfrak P(S^{\mu_i})$, $\mu_i\in Q_0$, $\mu_i\downarrow0$ (all S^{μ_i} along the common subsequence of Theorem 3.1(ii)). For $\alpha\in\mathfrak P(S)$ let $\Delta(\alpha):=\alpha'_+(0)-\alpha'_-(0)$, and $\Delta_S:=\inf_{\alpha\in\mathfrak P(S)}\Delta(\alpha)$, the *latent density* of the Wick square in S. Since the limits are not known to be unique, $\mathfrak P(S)$ and Δ_S are defined through the defining sequence of S (the subsequence of Theorem 3.1(ii)) and may depend on it; "the latent density in S" always refers to this choice.

Proposition 10.35 (the invariant part of the Wick square)**.** *$\mathfrak P(S)$ is not empty. Every $\alpha\in\mathfrak P(S)$ is convex on I, $\alpha(0)=0$, $|\alpha(\sigma)|\le\mathsf K|\sigma|$, and $0\le\Delta(\alpha)\le\kappa_++2S_\flat^B$, with κ_+, $S_\flat^B$ and $\mathsf K=\max(\kappa_+,2S_\flat^B)$ as in Theorem 6.8. For every $\alpha\in\mathfrak P(S)$, every nonnegative $h\in C_c^\infty(\mathbb R^3)$ and every j,*

$$\alpha'_-(0)\|h\|_1\le P_jX_{\mathrm R}(h)\le\alpha'_+(0)\|h\|_1\quad\mu\text{-a.s.},\qquad\alpha'_-(0)\|h\|_1\le\mathbb E_\mu X_{\mathrm R}(h)\le\alpha'_+(0)\|h\|_1.\tag{10.13}$$

Hence $|P_jX_{\mathrm R}(h)-\mathbb E_\mu X_{\mathrm R}(h)|\le\Delta_S\|h\|_1$ a.s. for $h\ge0$, and (L) holds if $\Delta_S=0$, in particular if some $\alpha\in\mathfrak P(S)$ is differentiable at 0.

Proof. Properties of α_n. α_n is finite (by property (P2) of Section 7.1) and convex (a logarithmic moment generating function), $\alpha_n(0)=0$ and $\alpha'_n(0)=\langle r_0^2-W_\varepsilon-z_\varepsilon\rangle_n$ by translation invariance

(Lemma 7.13). Proposition 6.5 with $q \equiv \sigma$ gives $\alpha_n(\sigma) \le \kappa_+\sigma$ for $0 \le \sigma \le \sigma_+$, and Proposition 6.7 with $h \equiv p$ gives $\alpha_n(-p) \le 2S_\flat^B p$ for $0 \le p \le H$, for every $\mu_0 \in (0, \mu_*] \cup \{\flat\}$. By convexity $\alpha_n(\sigma) \ge \alpha_n'(0)\sigma$, and evaluating at $\sigma = \sigma_+$ and $\sigma = -H$ gives $-2S_\flat^B \le \alpha_n'(0) \le \kappa_+$. So $|\alpha_n(\sigma)| \le \mathsf{K}|\sigma|$ on $[-H, \sigma_+]$, and the α_n are uniformly Lipschitz on $\bar I$ (a convex function bounded by M on an interval is Lipschitz with constant $2M/\delta$ at distance δ from its ends). By the Arzelà–Ascoli theorem a subsequence converges uniformly on $\bar I$, so $\mathfrak{P}(S) \ne \emptyset$; for S^0 choose $\alpha^{(i)} \in \mathfrak{P}(S^{\mu_i})$, which obey the same bounds, and apply the Arzelà–Ascoli theorem once more. Convexity, $\alpha(0) = 0$ and the bounds pass to pointwise limits, and give $\alpha_+'(0) \le \kappa_+$, $\alpha_-'(0) \ge -2S_\flat^B$.

The chessboard bound. Fix n and a real lattice function q. The state $\langle\cdot\rangle_n$ is reflection positive about the planes halfway between lattice planes (Proposition 7.9(a) for $\mu_0 > 0$, (b) for $\flat$), invariant under these reflections (Lemma 7.13), and the number of sites per side is even; so the chessboard estimate (Section 4.6, item 2) applies to the positive site factors $\exp(\varepsilon^3 q(x)(r_x^2 - W_\varepsilon - z_\varepsilon))$, exactly as in step (2) of the proof of Proposition 6.5, but at the photon mass of the state:

$$\langle e^{X_{\mathrm{R}}(q)}\rangle_n \le \exp\Big(\sum_x \varepsilon^3 \alpha_n(q(x))\Big). \tag{10.14}$$

Upper bound in (10.13). Let $S \in \{S^{\mu_0}, S^\flat\}$, $\alpha = \lim \alpha_n$ along a subsequence Λ, $h \ge 0$, $s > 0$, and $h_T := T^{-1}\int_0^T h(\cdot - te_j)\,dt \in C_c^\infty$. Then $\|h_T\|_\infty \le T^{-1}\sup_x \int_{\mathbb{R}} h(x + te_j)\,dt \to 0$, and $\|h_T\|_1 = \|h\|_1$. Fix T so large that $\sigma_T := s\|h_T\|_\infty < \sigma_+/2$. For $0 \le \sigma \le \sigma_T$ convexity and $\alpha_n(0) = 0$ give $\alpha_n(\sigma) \le \sigma\,\alpha_n(\sigma_T)/\sigma_T$. With $q = sh_T$ (sampled) in (10.14),

$$\log\langle e^{sX_{\mathrm{R}}(h_T)}\rangle_n \le s\sum_x \varepsilon^3 h_T(x)\,\frac{\alpha_n(\sigma_T)}{\sigma_T} \;\longrightarrow\; s\|h\|_1\frac{\alpha(\sigma_T)}{\sigma_T} \qquad (n \to \infty,\ n \in \Lambda).$$

The law of $X_{\mathrm{R}}(h_T)$ under $\langle\cdot\rangle_n$ converges to that of $X_{\mathrm{R}}(h_T)$ under μ (characteristic functionals, Theorem 3.1(ii)); since $y \mapsto \min(e^{sy}, L)$ is bounded and continuous for every $L > 0$, $\mathbb{E}_\mu e^{sX_{\mathrm{R}}(h_T)} \le \liminf_n \langle e^{sX_{\mathrm{R}}(h_T)}\rangle_n \le \exp(s\|h\|_1\alpha(\sigma_T)/\sigma_T)$. Next, $X_{\mathrm{R}}(h_T) = T^{-1}\int_0^T U(te_j)X_{\mathrm{R}}(h)\,dt$ in $L^2(\mu)$ (Riemann sums of h_T converge uniformly with supports in a fixed compact, and $\mathbb{E}X_{\mathrm{R}}(k)^2 \le C'N(k)^2$ by Theorem 3.1(iii)), so $X_{\mathrm{R}}(h_T) \to P_jX_{\mathrm{R}}(h)$ in $L^2(\mu)$ (Lemma 10.7(i)), and almost surely along a sequence $T_k \to \infty$. By Fatou's lemma and $\alpha(\sigma)/\sigma \downarrow \alpha_+'(0)$ as $\sigma \downarrow 0$,

$$\mathbb{E}_\mu e^{sP_jX_{\mathrm{R}}(h)} \le \liminf_k \mathbb{E}_\mu e^{sX_{\mathrm{R}}(h_{T_k})} \le e^{s\|h\|_1\alpha_+'(0)} \qquad (s > 0).$$

Chebyshev's inequality gives $\mu(P_jX_{\mathrm{R}}(h) > \alpha_+'(0)\|h\|_1 + \delta) \le e^{-s\delta}$ for every $s > 0$ and $\delta > 0$, hence the upper bound.

Lower bound. The same argument with $q = -sh_T$, using $\alpha_n(-\sigma) \le \sigma\,\alpha_n(-\sigma_T)/\sigma_T$ for $0 \le \sigma \le \sigma_T < H/2$ and $\alpha(-\sigma)/\sigma \downarrow -\alpha_-'(0)$, gives $\mathbb{E}_\mu e^{-sP_jX_{\mathrm{R}}(h)} \le e^{-s\|h\|_1\alpha_-'(0)}$, hence $P_jX_{\mathrm{R}}(h) \ge \alpha_-'(0)\|h\|_1$.

The mean. $\mathbb{E}_\mu X_{\mathrm{R}}(h) = \lim_{n\in\Lambda}\alpha_n'(0)\sum_x \varepsilon^3 h(x)$, and for $\sigma \in (0, \sigma_+/2)$, $\alpha_n'(0) \le \alpha_n(\sigma)/\sigma \to \alpha(\sigma)/\sigma$ and $\alpha_n'(0) \ge \alpha_n(-\sigma)/(-\sigma) \to \alpha(-\sigma)/(-\sigma)$; letting $\sigma \downarrow 0$ gives the second part of (10.13).

The state S^0. Let $\alpha = \lim_i \alpha^{(i)}$ as in Definition 10.34. The bound $\mathbb{E}_{\mu_{S^{\mu_i}}} e^{sX_{\mathrm{R}}(h_T)} \le \exp(s\|h\|_1\alpha^{(i)}(\sigma_T)/\sigma_T)$ holds for each i by the above, and passes to $i \to \infty$ by the weak convergence $\mu_{S^{\mu_i}} \to \mu_{S^0}$ (Lemma 10.3) and the same truncation; the rest is unchanged, the mean included ($\mathbb{E}_{S^{\mu_i}}X_{\mathrm{R}}(h)/\|h\|_1 \le \alpha^{(i)}(\sigma)/\sigma \to \alpha(\sigma)/\sigma$).

Conclusion. For $h \ge 0$ both $P_jX_{\mathrm{R}}(h)$ and $\mathbb{E}X_{\mathrm{R}}(h)$ lie in $[\alpha_-'(0), \alpha_+'(0)]\|h\|_1$ for every $\alpha \in \mathfrak{P}(S)$, so they differ by at most $\Delta_S\|h\|_1$. If $\Delta_S = 0$, then $P_jX_{\mathrm{R}}(h) = \mathbb{E}X_{\mathrm{R}}(h)$ for $h \ge 0$, and for every real $h \in C_c^\infty$ by the decomposition $h = (h + c\chi) - c\chi$ of the proof of Theorem 10.33. □

Remark 10.36 (what (L) needs). $\Delta(\alpha) > 0$ means that the limit pressure has a corner at the reference value of the bare coefficient: a first-order transition in the mass, with the density $\mathbb{E}\,X_{\mathrm{R}}$

jumping; one-loop perturbation theory predicts such a transition on a surface inside the coupling range of Theorem 3.1 (Remark 14.1). A corner is excluded if the torus susceptibilities $\alpha_n''(\sigma) = |T_n|^{-1}\operatorname{Var}_{n,\sigma} X_{\mathrm{R}}(1_{T_n})$ in the tilted states are bounded uniformly in n for $|\sigma|$ small, which by Lemma 10.31 is a uniform integrability statement for the nonnegative truncated two-point function of the Wick square; the estimates of this paper are linear in σ and give no such bound (Section 14.5).

10.8.3. *The full algebra under a differentiability hypothesis.* For the whole algebra we give a criterion in terms of the state alone. Fix a length $R > 0$, the block $Q := (0,R)^3$ and, for $b \in \mathbb{Z}^3$, the block $Q + Rb$. Assume, in addition to (H1)–(H3),

(H4) $\theta_j\theta_k = \theta_k\theta_j$ for $j \neq k$.

It holds for the laws of the limit states, since the push-forwards ρ_{j*} of Section 10.2 commute. Put $\Theta_{j,c}G := G \circ \theta_{j,c}$. From (10.3), $\Theta_{j,c} = U(2ce_j)\Theta_j$, hence $\Theta_{j,c}\Theta_{j,c'} = U(2(c-c')e_j)$, $\Theta_{j,c}U(te_j) = U(-te_j)\Theta_{j,c}$, and $\Theta_{j,c}$ commutes with $U(y)$, $y \perp e_j$, and (by (H4)) with $\Theta_{k,c'}$, $k \neq j$. For $m \in \mathbb{Z}$ let $\vartheta^{(j)}_{2m} := U(2mRe_j)$ and $\vartheta^{(j)}_{2m+1} := U(2mRe_j)\Theta_{j,R}$, and for $b \in \mathbb{Z}^3$ let $\vartheta_b := \vartheta^{(1)}_{b_1}\vartheta^{(2)}_{b_2}\vartheta^{(3)}_{b_3}$ (the factors commute). By (H2), ϑ_b maps $\Sigma(Q)$-measurable functions to $\Sigma(Q+Rb)$-measurable ones, and it preserves μ. For a bounded real $\Sigma(Q)$-measurable Γ and $B_k := \{0,\dots,2^k-1\}^3$ define the *block pressure*

$$\Pi_\Gamma(u) := \liminf_{k\to\infty} |B_k|^{-1}\log \mathbb{E}_\mu \exp\Big(u\sum_{b\in B_k}\vartheta_b\Gamma\Big), \qquad u \in \mathbb{R}.$$

By Jensen's inequality and the invariance of μ, $\Pi_\Gamma(u) \geq u\,\mathbb{E}_\mu\Gamma$, and $|\Pi_\Gamma(u)| \leq |u|\|\Gamma\|_\infty$. We say that Γ satisfies (D_Γ) if $\Pi_\Gamma(u) = u\,\mathbb{E}_\mu\Gamma + o(u)$ as $u \to 0$.

Theorem 10.37 (single-axis clustering from differentiable block pressures)**.** *Let μ satisfy (H1)–(H4), let Γ be bounded, real and $\Sigma(Q)$-measurable, and assume (D_Γ). Then $P_j\Gamma = \mathbb{E}_\mu\Gamma$ for $j = 1,2,3$, and (G',Γ) clusters along the axes for every $G' \in L^2(\mu)$.*

Steps 1–2 of the proof are a variant, in the present setting, of the multiple-reflection bound of [19, Prop. 10.5.3, p. 239].

Proof. Step 1: one doubling. Let $W \geq 0$ be bounded and $\Sigma(U)$-measurable with $U \subset H_j^-(c)$. Then $\Theta_{j,c}W \in L^2(\Sigma(H_j^+(c)))$ by (H2), and by (H3) and the Schwarz inequality for the positive form $\langle F, G\rangle_c := \mathbb{E}[\overline{\Theta_{j,c}F}\,G]$ on $L^2(\Sigma(H_j^+(c)))$,

$$(\mathbb{E}W)^2 = \langle 1,\,\Theta_{j,c}W\rangle_c^2 \leq \langle 1,\,1\rangle_c\,\langle\Theta_{j,c}W,\,\Theta_{j,c}W\rangle_c = \mathbb{E}\big[W\cdot\Theta_{j,c}W\big].$$

Step 2: from a line to a cube. For a finite $B \subset \mathbb{Z}^3$ put $Z_B := \prod_{b\in B}\vartheta_b(e^{u\Gamma})$. Let $N = 2^p$ and $L_N := \{0,\dots,N-1\}\times\{0\}\times\{0\}$. If B is a box $\{0,\dots,2^{q_1}-1\}\times\{0,\dots,2^{q_2}-1\}\times\{0,\dots,2^{q_3}-1\}$, then Z_B is measurable for the open box of side lengths $2^{q_i}R$, which lies in $H_j^-(2^{q_j}R)$, and $\Theta_{j,2^{q_j}R}Z_B = Z_{B'}$, where B' is B reflected in the plane $x_j = 2^{q_j}R$ of block coordinates, i.e. $b_j \mapsto 2^{q_j+1}-1-b_j$. Indeed, for $c = MR$ and $l = 2m$, $\Theta_{j,c}\vartheta^{(j)}_l = U(-2mRe_j)\Theta_{j,c} = U(2(M-1-m)Re_j)\Theta_{j,R} = \vartheta^{(j)}_{2M-1-l}$, and for $l = 2m+1$, $\Theta_{j,c}\vartheta^{(j)}_l = U(-2mRe_j)\Theta_{j,c}\Theta_{j,R} = U(2(M-1-m)Re_j) = \vartheta^{(j)}_{2M-1-l}$; the other factors of ϑ_b commute with $\Theta_{j,c}$. So Step 1 gives $\mathbb{E}Z_B \leq (\mathbb{E}Z_{B\cup B'})^{1/2}$, and $B\cup B'$ is the box with q_j replaced by q_j+1. Starting from L_N and doubling $k-p$ times in direction 1 and k times in directions 2 and 3 ($k \geq p$),

(10.15)

$$\log\mathbb{E}\exp\Big(u\sum_{l<N}\vartheta^{(1)}_l\Gamma\Big) = \log\mathbb{E}Z_{L_N} \leq \frac{N}{|B_k|}\log\mathbb{E}Z_{B_k}, \qquad \text{hence} \qquad \log\mathbb{E}\exp\Big(u\sum_{l<N}\vartheta^{(1)}_l\Gamma\Big) \leq N\,\Pi_\Gamma(u).$$

Step 3: the line mean. Put $\tilde\Gamma := \Theta_{1,R}\Gamma$ and $\mathsf{A}_N := N^{-1}\sum_{l<N}\vartheta^{(1)}_l\Gamma = N^{-1}\sum_{m<N/2}U(2mRe_1)(\Gamma+\tilde\Gamma)$. By the mean ergodic theorem for the unitary $U(2Re_1)$, $\mathsf{A}_N \to \mathsf{A} := \frac12 P'(\Gamma+\tilde\Gamma)$ in L^2, where

P' is the projection onto the $U(2Re_1)$-invariant vectors; and $\mathbb{E}\mathsf{A}_N = \mathbb{E}\Gamma$. By (10.15) with $u = s/N$ and (D_Γ), for every $s \in \mathbb{R}$, $\limsup_N \log \mathbb{E}e^{s\mathsf{A}_N} \le \limsup_N N\Pi_\Gamma(s/N) = s\mathbb{E}\Gamma$. Along a subsequence $\mathsf{A}_N \to \mathsf{A}$ a.s., and Fatou's lemma gives $\mathbb{E}e^{s(\mathsf{A}-\mathbb{E}\Gamma)} \le 1$ for all $s \in \mathbb{R}$. Chebyshev's inequality, with $s \to \pm\infty$, gives $\mathsf{A} = \mathbb{E}\Gamma$ a.s.

Step 4: conclusion. The range of P_1 is contained in that of P', so $P_1P' = P_1$. P_1 commutes with U and Θ_1, and Θ_1 fixes the range of P_1 (Lemma 10.8(ii)), so $P_1\tilde\Gamma = P_1U(2Re_1)\Theta_1\Gamma = P_1\Gamma$. Hence $\mathbb{E}\Gamma = P_1\mathsf{A} = \frac12(P_1\Gamma + P_1\tilde\Gamma) = P_1\Gamma$. The cube B_k and Π_Γ do not depend on the direction, so the same argument with lines along e_2, e_3 gives $P_j\Gamma = \mathbb{E}\Gamma$ for all j, and Lemma 10.24 gives the clustering. □

Corollary 10.38 (OS4 for the limit state under (D))**.** *Let S, $\mu = \mu_S$ be as above, and assume*

(D) *for every $R > 0$ and every real $f \in C_c^\infty(Q;\mathcal{V})$, $Q = (0,R)^3$, the functions $\Gamma = \cos\Phi(f)$ and $\Gamma = \sin\Phi(f)$ satisfy (D_Γ).*

Then P_j is the projection onto the constants for $j = 1,2,3$: μ is ergodic, μ satisfies OS4 (Definition 2.11), and every pair (G', G) in $L^2(\mu)$ clusters along the axes. The same holds for the law μ_S^H on $\mathcal{H}^{-r}$, which is the only probability measure on $\mathcal{H}^{-r}$ with the moments S (Lemma 10.5); so S satisfies OS4. In particular the Schwinger functions satisfy $\lim_{|a|\to\infty} S\big(P'\cdot P(\cdot - a)\big) = S(P')\,S(P)$ for all polynomials P, P' in the smeared fields.

Proof. For real $f \in C_c^\infty(\mathbb{R}^3;\mathcal{V})$ choose a and R with $\operatorname{supp} f(\cdot - a) \subset Q$; Theorem 10.37 gives $P_j e^{i\Phi(f(\cdot-a))} = \mathbb{E}e^{i\Phi(f(\cdot-a))}$, and translating back ($P_j$ commutes with $U(a)$, and μ is invariant) $P_j e^{i\Phi(f)} = \mathbb{E}e^{i\Phi(f)}$. The span of the $e^{i\Phi(f)}$ is dense in $L^2(\mu)$ (Lemma J.2), so $P_jG = \mathbb{E}G$ for all $G \in L^2(\mu)$; this is ($\mathrm{OS4}_j$), and in particular $\mathbb{E}[G \mid \mathcal{I}] = P_{\{1,2,3\}}G = P_{\{1,2,3\}}P_1G = \mathbb{E}G$, so μ is ergodic. Lemma 10.24 gives the clustering; polynomials lie in $L^2(\mu)$. The map ι of Lemma 10.5(b) is a Borel isomorphism onto a set of full measure that intertwines the translations, so the statements carry over to μ_S^H. □

Remark 10.39 (on hypothesis (D)). If $\mu = \frac12(\mu_1 + \mu_2)$ with $\mathbb{E}_{\mu_1}\Gamma \ne \mathbb{E}_{\mu_2}\Gamma$ and both μ_i invariant under the ϑ_b, then $\log \mathbb{E}_\mu e^{uN_k} \ge \log \mathbb{E}_{\mu_i} e^{uN_k} - \log 2$ (with $N_k := \sum_{b\in B_k}\vartheta_b\Gamma$) and Jensen's inequality in μ_i give $\Pi_\Gamma(u) \ge \max_i u\,\mathbb{E}_{\mu_i}\Gamma$, so ($\mathrm{D}_\Gamma$) fails: (D) excludes in particular a non-trivial ergodic decomposition, and it is a uniqueness-type statement. Theorem 10.37 applies to each component μ_ξ with its own block pressure, but (D) for a component is no easier to verify than for μ. For the linear Wick-square field the same mechanism, run on the tori instead of in the limit state, is Proposition 10.35, where the corresponding pressure is controlled by the uniform estimates of this paper up to its corner at the origin.

10.8.4. *Summary: which form of clustering holds for which state.*

Theorem 10.40 (clustering for the limit states and their components)**.** *Let $S \in \{S^{\mu_0}, S^\flat, S^0\}$ be a limit state of Theorem 3.1, $\mu = \mu_S$, $\mu = \int \mu_\xi\, d\nu$ its ergodic decomposition, and $\mathcal{K}_1$ the closed span in $L^2(\mu)$ of the products $F(\mathbf{g})Y$, $\mathbf{g} \in C_c^\infty(\mathbb{R}^3;\Lambda^2)$ real, $Y \in L^2(\mathcal{M})$.*

(a) *For every $G \in \mathcal{K}_1$ and $G' \in L^2(\mu)$, $\mathbb{E}_\mu[G'U(a)G] \to 0 = \mathbb{E}_\mu G'\,\mathbb{E}_\mu G$ as $|a| \to \infty$ in any manner; the same holds on ν-a.e. component for $G = F(\mathbf{g})Y$, Y bounded and $\mathcal{M}_0$-measurable.*

(b) *For $h \ge 0$, $|P_jX_{\mathrm{R}}(h) - \mathbb{E}X_{\mathrm{R}}(h)| \le \Delta_S\|h\|_1$ with $\Delta_S \le \kappa_+ + 2S_\flat^B$. If $\Delta_S = 0$, then (a) holds for every G in the closed span of $L^2(\mathcal{M}) \cup \mathcal{K}_1$, with limit $\mathbb{E}G'\,\mathbb{E}G$, and all components have the same law on $\mathcal{M}_0$.*

(c) *If (D) holds, then μ is ergodic and satisfies OS4, and every pair (G', G) in $L^2(\mu)$ clusters along the axes.*

(d) *On ν-a.e. component: Theorem 3.6(c)–(e), that is, (C1), (C2) and a unique joint vacuum in each coordinate direction. In the reconstruction of Proposition 10.16 from μ_ξ, the vectors $[F(\mathbf{g})Y]$, $F(\mathbf{g})Y \in L^2_+$ with Y bounded and $\mathcal{M}_0$-measurable, are orthogonal to $\operatorname{Ker} H_j$; if $\Delta_S = 0$, so are the vectors $[G - \mathbb{E}_{\mu_\xi}G]$, $G \in L^2_+$ bounded and $\mathcal{M}_0$-measurable.*

Proof. (a) For G in the linear span of the products $F(\mathbf{g})Y$, $P_jG = 0 = \mathbb{E}_\mu G$ by Theorem 10.27. Both sides are continuous on $L^2(\mu)$, so this holds on $\mathcal{K}_1$, and Lemma 10.24 gives the clustering. On the components the limit is $\mathbb{E}_{\mu_\xi} G' \, \mathbb{E}_{\mu_\xi} G = 0$ (Theorem 10.27). (b) Proposition 10.35 and Theorem 10.33; for linear combinations use that P_j is linear. (c) Corollary 10.38. (d) Theorem 3.6; for the zero-energy statement, Proposition 10.17(i) for μ_ξ identifies $P_{\operatorname{Ker} H_j}[G]$ with $[P_j^\xi G]$, which vanishes for $G = F(\mathbf{g})Y$ (Theorem 10.27, components) and equals $\mathbb{E}_{\mu_\xi} G \, \Omega_j$ for bounded $\mathcal{M}_0$-measurable G when $\Delta_S = 0$ (Proposition 10.35 and Theorem 10.33, components). For the last point: $\mu_\xi = \mu$ on $\mathcal{M}_0$ (Theorem 10.33); the map $(t,x) \mapsto G(\tau_{-te_j}x)$ is $\mathcal{B}(\mathbb{R}) \otimes \mathcal{M}_0$-measurable (for bounded continuous cylinder functions of the Wick square it is continuous in t and $\mathcal{M}_0$-measurable in x; the monotone class theorem gives the rest), so the one-sided means $\mathsf{A}_T := T^{-1} \int_0^T U(te_j) G \, dt$ are $\mathcal{M}_0$-measurable and $\|\mathsf{A}_T - \mathbb{E}_\mu G\|_{L^2(\mu_\xi)} = \|\mathsf{A}_T - \mathbb{E}_\mu G\|_{L^2(\mu)} \to \|P_jG - \mathbb{E}_\mu G\|_{L^2(\mu)} = 0$; hence $P_j^\xi G = \mathbb{E}_\mu G = \mathbb{E}_{\mu_\xi} G$ (Lemma 10.7(i) for μ_ξ). □

The theorem does not decide whether $\operatorname{Ker} H_j = \mathbb{C}\Omega_j$ on all of $\mathcal{H}_j$ for a component; by Proposition 10.41(a) below this is (OS4$_j$) for the component (Section 14.5).

The forms of the cluster property are related as follows. Here (OS4$_j$) and (OS4$'_j$) are the conditions of Definition 2.11: ergodicity of the translations $\{\tau_{te_j}\}_{t\in\mathbb{R}}$, and clustering along e_j; the weak form of OS4 is ergodicity of μ under $\mathbb{R}^3$ (Definition 2.12). Under (H1)–(H3), (OS4$_j$) and (OS4$'_j$) are equivalent by Lemma 10.24 and Lemma 10.7(ii).

Proposition 10.41 (relations between the forms of OS4)**.** *Let μ satisfy (H1)–(H3), and let $j \in \{1,2,3\}$.*

(a) *(OS4$_j$) holds if and only if, in the reconstruction of Proposition 10.16 in direction j, $\operatorname{Ker} H_j = \mathbb{C}\Omega_j$.*

(b) *μ is ergodic under $\mathbb{R}^3$ if and only if, in the reconstruction in some (equivalently, every) coordinate direction j, the joint fixed space of $\{e^{-tH_j}\}_{t\ge0}$ and $\{V_j(y)\}_{y\perp e_j}$ is $\mathbb{C}\Omega_j$; if and only if (C2) holds in the form $\mathbb{E}_\mu[G'\,U(a)G] \to \mathbb{E}_\mu[G']\,\mathbb{E}_\mu[G]$ as $\min_k|a_k| \to \infty$, for all $G, G' \in L^2(\mu)$; and if and only if the Cesàro means over cubes of $\mathbb{E}_\mu[G'\,U(a)G]$ converge to $\mathbb{E}_\mu[G']\,\mathbb{E}_\mu[G]$ for all $G, G' \in L^2(\mu)$.*

(c) *(OS4$_j$) for one j implies ergodicity under $\mathbb{R}^3$. Let X be a space of distributions as in Section 10.2 and $\mathcal{P}_j^+$ the algebra of polynomials in the fields $\Phi(f)$, $f \in C_c^\infty(H_j^+(0);\mathcal{V})$ real. If $\mathcal{P}_j^+$ is dense in $L^2(\Sigma(H_j^+(0)),\mu)$ and the cluster property in the form of* [27, E4, p. 88] *holds, $\lim_{\lambda\to\infty} \mathbb{E}_\mu[\overline{\Theta_j F}\, U(\lambda a)G] = \mathbb{E}_\mu[\overline{F}]\,\mathbb{E}_\mu[G]$ for all $F, G \in \mathcal{P}_j^+$ and $a \perp e_j$, then μ is ergodic under $\mathbb{R}^3$.*

(d) *Let $\mu = \int \mu_\xi \, d\nu(\xi)$ be the decomposition of Theorem 10.10. Then ν-almost every μ_ξ satisfies (OS4$_j$) if and only if $P_jG = \mathbb{E}[G \mid \mathcal{I}]$ for every $G \in L^2(\mu)$.*

Proof. (a) Let $G \in L^2_+ = L^2(\Sigma(H_j^+(0)),\mu)$. By Proposition 10.17(i), $P_{\operatorname{Ker} H_j}[G] = [P_jG]$ and $\|[W]\|^2_{\mathcal{H}_j} = \mathbb{E}_\mu|W|^2$ for every e_j-invariant $W \in L^2_+$. If (OS4$_j$) holds, $P_jG = \mathbb{E}_\mu G$ by Lemma 10.7(ii), so $P_{\operatorname{Ker} H_j}[G] = \mathbb{E}_\mu G \, \Omega_j$; since $[L^2_+]$ is dense in $\mathcal{H}_j$, $\operatorname{Ker} H_j = \mathbb{C}\Omega_j$. Conversely, if $\operatorname{Ker} H_j = \mathbb{C}\Omega_j$, then $[P_jG] = c\,\Omega_j$ with $c = \langle \Omega_j, [P_jG]\rangle = \mathbb{E}_\mu P_jG = \mathbb{E}_\mu G$, and $W := P_jG - \mathbb{E}_\mu G$ is e_j-invariant with $[W] = 0$, so $\mathbb{E}_\mu|W|^2 = 0$. Thus $P_jG = \mathbb{E}_\mu G$ for $G \in L^2_+$. For $G \in L^2(\Sigma(K))$, K bounded, $U(se_j)G \in L^2_+$ for s large and P_j commutes with $U(se_j)$, so $P_jG = \mathbb{E}_\mu G$; these G are dense in $L^2(\mu)$ by (H2). Hence P_j is the projection onto the constants, and for a τ_{te_j}-invariant Borel set B, $\mathbf{1}_B = P_j\mathbf{1}_B = \mu(B)$, so $\mu(B) \in \{0,1\}$.

(b) If μ is ergodic, $\mathbb{E}[\,\cdot \mid \mathcal{I}] = \mathbb{E}_\mu$, and the three statements are Proposition 10.17(iii), Theorem 10.20 and Theorem 10.19. Conversely, if B is an $\mathbb{R}^3$-invariant Borel set, then $G = G' = \mathbf{1}_B$ satisfies $\mathbb{E}_\mu[G'U(a)G] = \mu(B)$ for every a, so either clustering statement gives $\mu(B) = \mu(B)^2$; the

joint-fixed-space statement gives the same by the converse in the proof of Theorem 11.1(iv), which uses only Lemma 10.8 and Proposition 10.17(i).

(c) An $\mathbb{R}^3$-invariant set is invariant under the τ_{te_j}, which gives the first claim. For the second, let W be bounded and $\mathbb{R}^3$-invariant. By Lemma 10.8(i)–(ii), $W \in L^2_+$ and $\Theta_j W = W$. Approximate W in L^2_+ by $G_n \in \mathcal{P}_j^+$; then $|\mathbb{E}[\overline{\Theta_j W} U(\lambda a) W] - \mathbb{E}[\overline{\Theta_j G_n} U(\lambda a) G_n]| \le \|W - G_n\|_2 (\|W\|_2 + \|G_n\|_2)$ uniformly in λ, and the cluster property gives, for any $a \perp e_j$, $a \neq 0$,

$$\mathbb{E}_\mu |W|^2 = \lim_{\lambda \to \infty} \mathbb{E}_\mu \big[\overline{\Theta_j W}\, U(\lambda a) W\big] = |\mathbb{E}_\mu W|^2.$$

So W is constant.

(d) Let $G \in L^2(\mu)$ and $Y_T := T^{-1} \int_0^T U(te_j) G \, dt$. By Lemma 10.7(i), $Y_T \to P_j G$ in $L^2(\mu)$, and $\int \|Y_T - P_j G\|^2_{L^2(\mu_\xi)} \, d\nu(\xi) = \|Y_T - P_j G\|^2_{L^2(\mu)}$ by Theorem 10.10(a). Along a sequence $T_k \to \infty$, therefore, $Y_{T_k} \to P_j G$ in $L^2(\mu_\xi)$ for ν-a.e. ξ; since μ_ξ is translation invariant, Lemma 10.7(i) for μ_ξ also gives $Y_{T_k} \to P_j^\xi G$, the projection for μ_ξ. So $P_j G = P_j^\xi G$ μ_ξ-a.s. for ν-a.e. ξ, and likewise $\mathbb{E}[G \mid \mathcal{I}] = \mathbb{E}_{\mu_\xi} G$ μ_ξ-a.s. (Theorem 10.10(a)). Take G in a countable set dense in every $L^2(\mu_\xi)$ (Lemma J.2); off one ν-null set, $P_j G = \mathbb{E}[G \mid \mathcal{I}]$ μ-a.s. for all such G if and only if $P_j^\xi G = \mathbb{E}_{\mu_\xi} G$ for all such G and ν-a.e. ξ, and both sides are continuous in G. The second statement is (OS4$_j$) for μ_ξ by Lemma 10.7(ii) for μ_ξ, as in the end of the proof of (a). □

Remark 10.42. Without rotations the forms are not equivalent (Example 10.22: the law there is ergodic under $\mathbb{R}^3$, but $P_1\Phi(f) \neq \mathbb{E}\Phi(f)$). With invariance under the full Euclidean group, ergodicity under $\mathbb{R}^3$ implies (OS4$_j$), (OS4$'_j$) and the cluster property of [27, E4, p. 88] for every time direction (Corollary 10.44), so the forms are then equivalent. In Theorem 3.6 the weak form is proved for the ergodic components; Theorem 3.10 proves (OS4$_j$) and (OS4$'_j$) for part of the observables.

10.9. **The limit states: proof of Theorem 3.6.**

Corollary 10.43 (Theorem 3.6)**.** *Let S be one of the limit states of Theorem 3.1, and μ_S its law on X (Lemma 10.3). Then $\mu_S = \int \mu_\xi \, d\nu(\xi)$, where for ν-a.e. ξ:*

(i) *μ_ξ is translation invariant and ergodic, invariant under $\theta_1, \theta_2, \theta_3$, and reflection positive about every coordinate plane;*

(ii) *(C1) Theorem 10.19 holds on μ_ξ; (C2) $\mathbb{E}_{\mu_\xi}[G' U(a) G] \to \mathbb{E}_{\mu_\xi}[G'] \, \mathbb{E}_{\mu_\xi}[G]$ as $\min_j |a_j| \to \infty$, for all $G, G' \in L^2(\mu_\xi)$, in particular for polynomials in the smeared fields;*

(iii) *in each coordinate direction j, the reconstruction of Proposition 10.16 on μ_ξ has a unique joint vacuum: the joint fixed space of the e^{-tH_j}, $t \ge 0$, and of the transverse translations is $\mathbb{C}\Omega_j$;*

(iv) *the Schwinger functions $S_N^\xi(f_1, \dots, f_N) := \mathbb{E}_{\mu_\xi} \prod_i \Phi(f_i)$ exist for all Schwartz test functions and satisfy $|S_N^\xi(f_1, \dots, f_N)| \le \sigma_N^\xi \prod_i |f_i|_k$ for a fixed Schwartz norm $|\cdot|_k$ and constants with $\sigma_N^\xi \le \alpha_\beta(\xi)(N!)^\beta$ for every $\beta > 1$; the constants depend on ξ.*

The decomposition is determined by S: μ_S is determined by S (Proposition 10.4), and its ergodic decomposition is unique in the following sense. If $\mu_S = \int m_\eta \, d\lambda(\eta)$, with $\eta \mapsto m_\eta(B)$ measurable for every Borel B and m_η translation invariant and ergodic for λ-a.e. η, then the images of ν under $\xi \mapsto \mu_\xi$ and of λ under $\eta \mapsto m_\eta$ agree on the σ-algebra of the space of probability measures on X generated by the maps $m \mapsto m(B)$.

Proof. (i)–(iii) are Theorem 10.10, Theorems 10.19–10.20 and Proposition 10.17(iii) applied to μ_S, which satisfies (H1)–(H3) by Proposition 10.4; polynomials in the smeared fields lie in $L^2(\mu_\xi)$ for a.e. ξ by Theorem 10.10(e) and the finiteness of all moments of μ_S.

(iv) Fix an even $r \ge 14$. By the uniqueness in Lemma 10.3, the laws μ_S on H^{-s}_{loc} for different $s > 3$ are images of one another under the continuous inclusions $H^{-s}_{\mathrm{loc}} \subset H^{-s'}_{\mathrm{loc}}$ $(s < s')$, and their

ergodic decompositions correspond; we may therefore take $s = r$. Let $Z_{\alpha,c}$, $M = \sum_{\alpha,c} \lambda_\alpha^{-r} Z_{\alpha,c}^2$ and Ψ be as in the proof of Lemma 10.5(b), and let (f_n) be a countable subset of $C_c^\infty(\mathbb{R}^3;\mathcal{V})$ containing, for every R, a subset of $C_c^\infty(B_R;\mathcal{V})$ dense in the H^r-norm. The set

$$B := \{M < \infty\} \cap \{\Psi(f_n) = \Phi(f_n) \text{ for all } n\}$$

is Borel, $\mu_S(B) = 1$, and on B one has $\Psi(f) = \Phi(f)$ for every $f \in C_c^\infty(\mathbb{R}^3;\mathcal{V})$, since both sides are continuous on $C_c^\infty(B_R;\mathcal{V})$ for the H^r-norm. For Schwartz f put $\Phi(f) := \Psi(f)$ on B; then $|\Phi(f)| \le M^{1/2}\|f\|_r$. By Theorem 9.1(c), each $h_{\alpha,c}$ being supported in one summand of $\mathcal{V}$, and Lemma 10.5(a) ($|f|_4 \le C\|f\|_{10}$),

$$\mathbb{E}_{\mu_S} Z_{\alpha,c}^{2n} \le c_S C_0^{2n} (2n)!\, |h_{\alpha,c}|_4^{2n} \le c_S (CC_0)^{2n} (2n)!\, \lambda_\alpha^{10n},$$

with c_S and C_0 the constants of Theorem 9.1(c). With $w := 4\sum_\alpha \lambda_\alpha^{10-r} < \infty$ and Jensen's inequality for the probability weights λ_α^{10-r}/w,

$$\mathbb{E}_{\mu_S} M^n \le w^n \sup_{\alpha,c} \lambda_\alpha^{-10n}\, \mathbb{E}_{\mu_S} Z_{\alpha,c}^{2n} \le c_S C_1^n (2n)!, \qquad C_1 := w(CC_0)^2.$$

Put $M_n(\xi) := \mathbb{E}_{\mu_\xi} M^n$. Since $\int \mu_\xi(B)\, d\nu = 1$ and $\int M_n\, d\nu = \mathbb{E}_{\mu_S} M^n$, Markov's inequality and the Borel–Cantelli lemma give, for ν-a.e. ξ: $\mu_\xi(B) = 1$, $M_n(\xi) < \infty$ for all n, and $M_n(\xi) \le n^2 c_S C_1^n (2n)!$ for $n \ge n_0(\xi)$. For such ξ and all Schwartz f_i, using $\mathbb{E}_{\mu_\xi} M^{N/2} \le M_N(\xi)^{1/2}$,

$$|S_N^\xi(f_1,\dots,f_N)| \le M_N(\xi)^{1/2} \prod_i \|f_i\|_r.$$

Finally let $|f|_k := \sup_{x,|\gamma|\le k} (1+|x|^2)^{k/2} |D^\gamma f(x)|$. By Lemma 10.5(a), $\|f\|_r \le C_r \sum_{|\delta|+|\gamma|\le r} \|x^\delta \partial^\gamma f\|_{L^2}$, and $\|x^\delta \partial^\gamma f\|_{L^2} \le \|(1+|x|^2)^{-1}\|_{L^2} \sup_x (1+|x|^2)^{1+|\delta|/2} |\partial^\gamma f(x)| \le \pi |f|_{r+2}$. Hence $\|f\|_r \le C' |f|_{r+2}$, and (iv) holds with $k = r+2$ and

$$\sigma_N^\xi := c(\xi)\, N\, c_S^{1/2} C_1^{N/2} C'^N ((2N)!)^{1/2} \le \alpha_\beta(\xi) (N!)^\beta \quad (\beta > 1),$$

where $c(\xi) \ge 1$ absorbs the finitely many $N < n_0(\xi)$ and we used $((2N)!)^{1/2} \le 2^N N!$.

Uniqueness of the decomposition: let $\mu_S = \int m_\eta\, d\lambda(\eta)$ be as in the statement, and let $\mathcal{D}$, A_T and q_G be as in the proof of Theorem 10.10. As there, $\int m_\eta(|A_T G - q_G|^2)\, d\lambda = \mathbb{E}_{\mu_S} |A_T G - q_G|^2 \to 0$. Under an ergodic m_η the fixed vectors are constants (the proof of Lemma 10.7(ii) gives a strictly invariant version), so $A_T G \to m_\eta(G)$ in $L^2(m_\eta)$. Along a subsequence, therefore, $\mu_x(G) = q_G(x) = m_\eta(G)$ for m_η-a.e. x, for all $G \in \mathcal{D}$ and λ-a.e. η; that is, $\mu_x = m_\eta$ for m_η-a.e. x. Hence $\mu_S\{x : \mu_x \in \mathcal{A}\} = \lambda\{\eta : m_\eta \in \mathcal{A}\}$ for every set $\mathcal{A}$ in the σ-algebra generated by the maps $m \mapsto m(B)$. Moreover μ_S is determined by S (Proposition 10.4). □

Theorem 3.6 is Corollary 10.43, carried to $\mathcal{H}^{-r}$ by Lemma 10.5; for μ itself, parts (c) and (d) of Theorem 3.6 are Theorems 10.19 and 10.20, and the uniqueness of μ is Lemma 10.5(c). Theorem 3.10 is Theorem 10.27 together with Theorem 10.40(a)–(c) and Proposition 10.35.

The obstruction of Proposition 3.8 disappears as soon as the law is invariant under all rotations: reflection positivity then holds about every plane, every invariant function is fixed by every reflection, and a rotation carries a coordinate axis to a direction off the coordinate planes, along which Theorem 10.20 applies. We state this as a conditional statement; it is not used elsewhere. For a unit vector $n \in \mathbb{R}^3$ and $c \in \mathbb{R}$ put $H_n^+(c) := \{x : x \cdot n > c\}$, $H_n^-(c) := \{x : x \cdot n < c\}$, and let $\rho_n := 1 - 2nn^{\mathsf{T}}$ be the reflection in the plane $n^\perp$.

Corollary 10.44 (the cluster property under rotation invariance)**.** *Let μ satisfy (H1)–(H3). Assume that $\mathrm{O}(3)$ acts on X by homeomorphisms $x \mapsto Rx$, continuously in R for each x, such that $\rho_j x = \theta_j x$ for the coordinate reflections, $R\,\tau_a\,R^{-1} = \tau_{Ra}$, $\Gamma_R G := G \circ R^{-1}$ maps $\Sigma(U)$-measurable functions to $\Sigma(RU)$-measurable ones, and μ is invariant under every $R \in \mathrm{O}(3)$. Let $\mu = \int \mu_\xi\, d\nu(\xi)$ be the decomposition of Theorem 10.10. Then for ν-almost every ξ:*

(a) *μ_ξ is invariant under the Euclidean group $\mathbb{R}^3 \rtimes \mathrm{O}(3)$;*
(b) *$\mathbb{E}_{\mu_\xi}[\overline{F \circ \theta_{n,c}}\, F] \geq 0$ for every unit n, every $c \in \mathbb{R}$ and every $F \in L^2(\Sigma(H_n^+(c)), \mu_\xi)$, where $\theta_{n,c} := \tau_{cn}\rho_n\tau_{-cn}$;*
(c) *for all $G, G' \in L^2(\mu_\xi)$, $\mathbb{E}_{\mu_\xi}[G'\,U(a)G] \to \mathbb{E}_{\mu_\xi}[G']\,\mathbb{E}_{\mu_\xi}[G]$ as $|a| \to \infty$ in any manner.*

Consequently μ_ξ satisfies (OS4$_j$) and (OS4$'_j$) of Definition 2.11 for $j = 1, 2, 3$ and OS4 in each of the forms of Definition 2.12, for every choice of the time direction (also for non-coordinate directions n, with the half spaces $H_n^+(0)$); and in the reconstruction from μ_ξ in any direction n, $\operatorname{Ker} H_n = \mathbb{C}\Omega_n$.

Proof. Step 1: reflection positivity about every plane. Given a unit vector n, choose $R \in \mathrm{SO}(3)$ with $Re_1 = n$. Then $\rho_n = R\rho_1 R^{-1}$ and $R\,H_1^+(0) = H_n^+(0)$. Let $F \in L^2(\Sigma(H_n^+(0)), \mu)$ and $F' := F \circ R$, which is $\Sigma(H_1^+(0))$-measurable. Since Γ is a representation, $F \circ \rho_n = \Gamma_R \Theta_1 F'$, and by R-invariance of μ and (H3), $\mathbb{E}_\mu[\overline{F \circ \rho_n}\, F] = \mathbb{E}_\mu[\Gamma_R(\overline{\Theta_1 F'}\, F')] = \mathbb{E}_\mu[\overline{\Theta_1 F'}\, F'] \geq 0$. For $c \neq 0$ conjugate by τ_{cn}, as after (10.3). Moreover $\rho_n\tau_a = \tau_{\rho_n a}\rho_n$, and Γ_{ρ_n} maps $\Sigma(U)$-measurable functions to $\Sigma(\rho_n U)$-measurable ones. So μ satisfies (H1)–(H3) with the coordinate frame replaced by any orthonormal frame containing n.

Step 2: invariant functions are fixed by every reflection. Let $W \in L^2(\mu)$ be invariant under $U(tn)$, $t \in \mathbb{R}$. The proof of Lemma 10.8(i)–(ii) applies verbatim with e_j replaced by n, the half spaces $H_j^+(c)$, $H_j^-(c)$ by $H_n^+(c)$, $H_n^-(c)$ and Θ_j by Γ_{ρ_n}, using Step 1: $\Gamma_{\rho_n} W = W$. A translation-invariant W is invariant along every n, so $\Gamma_{\rho_n} W = W$ for every n; every element of $\mathrm{O}(3)$ is a product of at most three reflections, so $\Gamma_R W = W$ for all $R \in \mathrm{O}(3)$.

Step 3: (a) and (b). Let $\mathcal{R}_0 \subset \mathrm{O}(3)$ be a countable dense subgroup. Exactly as in the proof of Theorem 10.10(c), with Γ_R in place of Θ_j and Step 2 in place of Lemma 10.8(ii), for ν-a.e. ξ the measure μ_ξ is invariant under every $R \in \mathcal{R}_0$. For bounded continuous G the map $R \mapsto \mathbb{E}_{\mu_\xi}[G \circ R^{-1}]$ is continuous, by continuity of the action and dominated convergence, so μ_ξ is invariant under $\mathrm{O}(3)$; with Theorem 10.10(b) this is (a). By Theorem 10.10(d) and (a), Step 1 applies to μ_ξ and gives (b).

Step 4: (c). Fix such a ξ and write $m := \mu_\xi$; it is ergodic for $\mathbb{R}^3$, $\mathrm{O}(3)$-invariant, and satisfies (H1)–(H3). The representation $R \mapsto \Gamma_R$ on $L^2(m)$ is strongly continuous (dominated convergence for bounded continuous G, density and $\|\Gamma_R\| = 1$ in general). Put $n_0 := (1,1,1)/\sqrt{3}$. Let $|a^{(l)}| \to \infty$, and choose R_l with $R_l a^{(l)} = |a^{(l)}|\, n_0$. Since $\Gamma_R U(a) = U(Ra)\Gamma_R$ and m is R-invariant, $\mathbb{E}_m[G'\,U(a^{(l)})G] = \mathbb{E}_m[(\Gamma_{R_l}G')\,U(|a^{(l)}|n_0)\,(\Gamma_{R_l}G)]$. Every subsequence has a further subsequence along which $R_l \to R$; along it the right side differs by $o(1)$ from $\mathbb{E}_m[(\Gamma_R G')\,U(|a^{(l)}|n_0)(\Gamma_R G)]$. Since $\min_k |a^{(l)}|(n_0)_k = |a^{(l)}|/\sqrt{3} \to \infty$, Theorem 10.20 for m gives the limit $\mathbb{E}_m[\Gamma_R G']\,\mathbb{E}_m[\Gamma_R G] = \mathbb{E}_m[G']\,\mathbb{E}_m[G]$.

Step 5: the forms of OS4. With $G' = \overline{\Theta_j F}$, (c) along λa gives the cluster property of [27, E4, p. 88] for every time direction. If a Borel set B is invariant under all τ_{te} for a unit vector e, (c) with $G = G' = \mathbf{1}_B$ and $a = te$ gives $m(B) = m(B)^2$; this is ergodicity along every direction, in particular (OS4$_j$), and (OS4$'_j$) is (c) with $a = te_j$. Ergodicity under $\mathbb{R}^3$ is Theorem 10.10(b). Finally $\operatorname{Ker} H_n = \mathbb{C}\Omega_n$ by Proposition 10.41(a), applied in a frame containing n (Step 1). □

Remark 10.45. (1) For the states of Theorem 3.1 the hypotheses of Corollary 10.44 other than $\mathrm{O}(3)$-invariance hold on $X = H^{-s}_{\mathrm{loc}}(\mathbb{R}^3; \mathcal{V})$ with $(R\Phi)(f) := \Phi(R_*^{-1} f)$, the push-forward of Section 2.5: the action is by linear homeomorphisms, continuous in R, and covariant for the local σ-algebras. $\mathrm{O}(3)$-invariance of the law is equivalent to invariance of the Schwinger functions, $S_{n,p}(R_*g_1, \dots; R_*h_1, \dots) = S_{n,p}(g_1, \dots; h_1, \dots)$ for $R \in \mathrm{O}(3)$, by Lemma 10.3. (2) So, if a limit state of Theorem 3.1 is invariant under rotations, then almost every component of its law satisfies OS0 in the form of Remark 3.7, OS1 for the full Euclidean group, OS2 about every plane, the symmetry and reality conditions of OS3, and OS4 in every form. Rotation invariance is the only missing input for these conclusions; it is discussed in Section 14.1. (3) Only reflection positivity about the

coordinate planes and invariance under rotations are used. Invariance under SO(3) suffices, since $\mathrm{O}(3) = \mathrm{SO}(3) \cup \rho_1 \mathrm{SO}(3)$ and $\rho_1 \in O_h$; so does invariance under a dense subgroup together with continuity of the action. In particular invariance of S under one rotation about a coordinate axis by an angle not in $\pi\mathbb{Q}$ suffices: together with O_h it generates a dense subgroup of O(3), and the set of R with $R_*\mu = \mu$ is closed. (4) The conclusion does not include the vanishing of the moments of odd degree ≥ 3 in the curvature for the components: $F \mapsto -F$ is not induced by any element of O(3), since $R \in \mathrm{O}(3)$ acts on 2-forms by a matrix similar to $\det(R)\, R$, which is never -1.

11. The reconstructed theory

Theorem 11.1 below uses no growth condition of [27, 28]: its proof uses the translations, the stabiliser D_4 of a coordinate axis in O_h, reflection positivity about one coordinate plane, and the results of Section 10. On the relation with [28]: the limit states satisfy the growth condition E0″ of [28, (4.2), p. 287] with $s = 4$ and every $\beta > 1$ (Theorem 9.1(c)), and E0″ implies the linear growth condition E0′ of [28, (4.1), p. 287] by the theorem of the Appendix of [28, p. 303], whose proof is a Hermite expansion and uses no Euclidean invariance. The reconstruction theorem E′ →R′ of [28, p. 287], which produces a Wightman theory, assumes in addition the axiom E1, invariance under all translations and all proper rotations [27, p. 88]. For the limit states E1 holds only for $\mathbb{R}^3 \rtimes O_h$, and the cluster axiom E4 only in the forms of Section 10; so that theorem does not apply as stated (Section 14.7).

Fix a limit state S of Theorem 3.1 and its law $\mu = \mu_S$. Choose the coordinate direction e_3 as Euclidean time. (By O_h-invariance of μ, every coordinate direction gives a unitarily equivalent theory; see (v) below.) Let $\mathcal{H} := \mathcal{H}_3$, $\Omega := \Omega_3$, $H := H_3$, and $V(y) := V_3(y)$, $y \in \mathbb{R}^2 = e_3^{\perp}$, as in Proposition 10.16. Let $D_4 := \{g \in O_h : ge_3 = e_3\}$, the stabiliser of the time axis: the eight signed permutations of (x_1, x_2), the symmetry group of a square.

Theorem 11.1 (reconstruction with translations and D_4)**.** (i) *$\mathcal{H}$ is a Hilbert space with the unit vector Ω, $H \geq 0$ is self-adjoint with $e^{-tH}[G] = [U(te_3)G]$, $H\Omega = 0$, and V is a strongly continuous unitary representation of the spatial translations commuting with H, with $V(y)\Omega = \Omega$. By Stone's theorem in two variables, $V(y) = e^{i(y_1P_1 + y_2P_2)}$ with commuting self-adjoint momenta P_1, P_2.*

(ii) *For $g \in D_4$, $W(g)[G] := [G \circ g^{-1}]$ defines a unitary representation of D_4 on $\mathcal{H}$ with $W(g)\Omega = \Omega$, $W(g)e^{-tH}W(g)^{-1} = e^{-tH}$ and $W(g)V(y)W(g)^{-1} = V(gy)$.*

(iii) *Elements of $O_h \setminus D_4$ do not act on $\mathcal{H}$ in this way: those with $ge_3 = -e_3$ exchange the two half spaces, and those with $ge_3 \perp e_3$ do not preserve $H_3^+(0)$.*

(iv) *The joint vacuum is unique, that is, $\mathbb{C}\Omega$ is the joint fixed space of the e^{-tH}, $t \geq 0$, and of V, if and only if μ is ergodic. On every ergodic component of Corollary 10.43 the joint vacuum is unique.*

(v) *For $g \in O_h$ with $ge_3 = e_j$, $[G] \mapsto [G \circ g^{-1}]$ is a unitary map $\mathcal{H}_3 \to \mathcal{H}_j$ intertwining the reconstructions in directions 3 and j.*

Proof. (i) is Proposition 10.16 with $j = 3$; $H\Omega = 0$ because $U(te_3)1 = 1$. (ii) Let $g \in D_4$. Then g preserves $H_3^+(0)$ and commutes with ρ_3 (an orthogonal map fixing e_3 preserves $e_3^{\perp}$). So $G \mapsto G \circ g^{-1}$ maps L^2_+ onto itself, and by g-invariance of μ (Proposition 10.4),

$$\langle [G \circ g^{-1}],\, [G' \circ g^{-1}] \rangle = \mathbb{E}\big[(\overline{\Theta_3 G}\, G') \circ g^{-1}\big] = \langle [G],\, [G'] \rangle.$$

So $W(g)$ preserves $\mathcal{N}$ and is isometric on the quotient, with inverse $W(g^{-1})$. The relations with $U(te_3)$ and $U(y)$ follow from $g\tau_a g^{-1} = \tau_{ga}$ and $ge_3 = e_3$. (iii) For $ge_3 = -e_3$, g maps $H_3^+(0)$ to $H_3^-(0)$; such g equal ρ_3 times an element of D_4, and ρ_3 is used in the inner product itself, not as an operator on $\mathcal{H}$. For $ge_3 = \pm e_k$, $k \neq 3$, $gH_3^+(0)$ is a half space in direction k. (iv) is Proposition 10.17(ii)–(iii) and its converse: if μ is not ergodic, there is a non-constant bounded invariant W; by Lemma 10.8(i) it lies in L^2_+, $[W]$ is fixed, and $\|[W] - \mathbb{E}[W]\Omega\|^2 = \mathbb{E}|W - \mathbb{E}W|^2 > 0$

by Proposition 10.17(i). The statement for components is Corollary 10.43(iii). (v) As (ii), using $g\rho_3 g^{-1} = \rho_j$ and $gH_3^+(0) = H_j^+(0)$. □

The theorem gives a quantum-mechanical system with a positive Hamiltonian, unitary spatial translations and the point group D_4. It gives no Lorentz covariance, and no Wightman-type distributions are claimed (Section 14.7).

12. Correlation signs and non-triviality under a remainder bound

We ask whether a limit state of Theorem 3.1 can be Gaussian, or the law of a pair of independent fields, one for the curvature and one for the Wick square, as for the free Maxwell field together with the Wick square of a free charged scalar. We do not settle this. What we prove is the following. A Gaussian limit state is decoupled (Lemma 12.2), so it suffices to show that one mixed truncated function is non-zero. The simplest one, $M(g;h) = S^T_{2,1}(g,g;h)$, has a sign in every limit state (Theorem 12.4). On every lattice torus it equals $-e^2$ times a covariance of the matter (Lemma 12.3), and its coefficient at $e = \lambda = 0$ is strictly negative on every lattice torus for $\mu_0 > 0$ (Proposition 12.6); its continuum value is the subject of Hypothesis 12.7. Non-triviality follows at small $e^2 + \lambda$ from that hypothesis and one remainder bound, uniform in the lattice spacing and the torus (Theorem 12.9). The remainder bound is not available from the uniform estimates of this paper or from the literature it rests on (Remark 12.10 and Section 14.3).

12.1. **Truncated functions; Gaussian and decoupled states.**

Definition 12.1 (truncated functions)**.** Let S be a Schwinger family whose generating functional G^S exists on a tube $D_{\sigma_+,H}$ for every finite family of real test functions $g_1,\dots,g_n$, $h_1,\dots,h_p$ (every limit state has this property, by OS0 in Theorem 3.1(iii)). Since $G^S(0) = 1$, $\log G^S$ is holomorphic near 0. Its Taylor coefficients at 0, $\partial_{z_1}\cdots\partial_{z_n}\partial_{w_1}\cdots\partial_{w_p}\log G^S(0)$, define the truncated functions $S^T_{n,p}(g_1,\dots,g_n;h_1,\dots,h_p)$ (test functions may repeat). The family S is *Gaussian* if $S^T_{n,p} = 0$ whenever $n+p \geq 3$. It is *decoupled* if $S^T_{n,p} = 0$ whenever $n \geq 1$ and $p \geq 1$. We put

$$M(g;h) := S^T_{2,1}(g,g;h).$$

The same definitions apply to the lattice states $\langle\cdot\rangle_{\varepsilon,j,\mu_0}$, with lattice test functions; we write $M_{\varepsilon,j,\mu_0}(g;h)$.

A family is decoupled if and only if $G^S(z,w) = G^S(z,0)\,G^S(0,w)$ near 0 for every family of test functions, hence on the connected tube by the identity theorem. This is the case if the curvature and the Wick square are independent under the law of Theorem 3.1(ii), for instance for the free Maxwell field (of any photon mass) together with the Wick square of a free charged scalar (of any mass).

Lemma 12.2 (Gaussian limit states are decoupled)**.** *Let S be a Schwinger family as in Definition 12.1 with $S_{n,p} = 0$ for odd n. Then $S^T_{1,1} = 0$, $S^T_{1,0} = 0$ and $M(g;h) = S_{2,1}(g,g;h) - S_{2,0}(g,g)\,S_{0,1}(h)$. If S is Gaussian, it is decoupled. Every limit state $S \in \{S^{\mu_0}, S^\flat, S^0\}$ of Theorem 3.1 satisfies the hypothesis.*

Proof. Expanding the logarithm to third order, $S^T_{1,0} = S_{1,0}$, $S^T_{1,1} = S_{1,1} - S_{1,0}S_{0,1}$ and $S^T_{2,1}(g,g;h) = S_{2,1}(g,g;h) - S_{2,0}(g,g)S_{0,1}(h) - 2S_{1,1}(g;h)S_{1,0}(g) + 2S_{1,0}(g)^2 S_{0,1}(h)$; the terms with an odd number of curvature arguments vanish. If S is Gaussian, $\log G^S$ is a polynomial of degree at most 2 in (z,w), and its only mixed term is $\sum_{i,k} z_i w_k S^T_{1,1}(g_i;h_k) = 0$. The limit states satisfy $S_{n,p} = 0$ for odd n by OS3 (Theorem 3.1(iii), Definition 2.10). □

12.2. **The photon dictionary with a scalar source.** Fix a lattice torus $T_{j,\varepsilon}$, a real b_T, $\lambda > 0$, $e \in \mathbb{R}$ ($e \neq 0$ in the state $\flat$, whose period $P = 2\pi/(|e|\ell_j)$ needs it), and $\mu_0 \in (0, \mu_*] \cup \{\flat\}$; on a fixed lattice torus the definitions of Section 2 make sense for all such (e, λ), and the standing convention $e \neq 0$ is relaxed only in this section. Let $C = C_{\mu_0} = (-\Delta_\varepsilon + \mu_0^2)^{-1}$, or $C = (-\Delta_\varepsilon)^{-1}\Pi^\perp$ in the state $\flat$ (Definition 2.7). For a real lattice 2-form g put

$$f := d^*g, \qquad a := Cf, \qquad q := \langle f,\, Cf\rangle_\varepsilon, \qquad Y(n) := \varepsilon \sum_b n_b a_b = \langle j_n,\, a\rangle_\varepsilon \quad (j_n := \varepsilon^{-2} n),$$

for integer-valued divergence-free currents n on the bonds; let $\mathcal{N}$ be the set of these currents, and in the state $\flat$ the subset of currents with zero windings, $\omega(n) = 0$ (Lemma 2.5). Let $V(r) := 3\varepsilon r^2 + \varepsilon^3(\frac{1}{2} b_T r^2 + \lambda r^4)$, and let $\mathbb{P}$ be the probability measure on $\mathcal{N} \times \mathbb{R}_+^{T_{j,\varepsilon}}$ given by

$$\mathbb{P}(n, dr) := Z_{\mathbb{P}}^{-1}\, e^{-\frac{1}{2} e^2 \langle j_n,\, C j_n\rangle_\varepsilon} \prod_b I_{n_b}\big(\varepsilon r_{b_-} r_{b_+}\big) \prod_x e^{-V(r_x)} r_x\, dr_x. \tag{12.1}$$

For a real site function k let $\mathbb{P}_k$ be $\mathbb{P}$ tilted by $e^{r^2(\varepsilon^3 k)}$, where $r^2(u) := \sum_x u(x) r_x^2$, and let $\langle\cdot\rangle_k := \langle \cdot\, e^{X_{\mathrm{R}}(k)}\rangle / \langle e^{X_{\mathrm{R}}(k)}\rangle$ denote the lattice state tilted by $e^{X_{\mathrm{R}}(k)}$.

Lemma 12.3 (photon dictionary with a scalar source)**.** *$Z_{\mathbb{P}} < \infty$, $Y \in L^4(\mathbb{P}_k)$, and:*

(a) $\langle F(g)^2\rangle_k = q - e^2\, \mathbb{E}_{\mathbb{P}_k}[Y^2]$ *for every real site function k;*

(b) *for every real site function h, $\langle F(g)\rangle = \langle F(g) X_{\mathrm{R}}(h)\rangle = 0$ and*

$$M_{\varepsilon,j,\mu_0}(g; h) = \langle F(g)^2 X_{\mathrm{R}}(h)\rangle - \langle F(g)^2\rangle \langle X_{\mathrm{R}}(h)\rangle = -e^2\, \mathrm{Cov}_{\mathbb{P}}\big(Y^2,\, r^2(\varepsilon^3 h)\big).$$

Proof. Integrability. By $\sum_{\nu\in\mathbb{Z}} I_\nu(t) = e^t$, $\sum_\nu \nu^2 I_\nu(t) = te^t$ and $\sum_\nu \nu^4 I_\nu(t) = (t + 3t^2)e^t$ (the moments of ν under the weights $I_\nu(t)e^{-t}$, read off from $\sum_\nu I_\nu(t) e^{i\nu\theta} = e^{t\cos\theta}$), and by $|Y| \le \varepsilon \|a\|_\infty \sum_b |n_b|$, the sum over $n \in \mathbb{Z}^{\mathrm{bonds}}$ of $\prod_b I_{n_b}(\varepsilon r_{b_-} r_{b_+})(1 + Y^4)$ is at most a polynomial in r times $\prod_b e^{\varepsilon r_{b_-} r_{b_+}} \le \prod_x e^{3\varepsilon r_x^2}$. The factor $e^{3\varepsilon r_x^2}$ cancels against V, and $e^{-\varepsilon^3(\frac{1}{2} b_T - k(x)) r^2 - \lambda\varepsilon^3 r^4}$ has all moments since $\lambda > 0$. So $Z_{\mathbb{P}} < \infty$ and $Y \in L^4(\mathbb{P}_k)$.

(a) By the proof of Lemma 7.3, with tf in place of f ($t \in \mathbb{R}$),

$$\mathbb{E}_C\big[e^{tF(g)} W_k(A)\big] = e^{\frac{1}{2} t^2 q} \sum_{n\in\mathcal{N}} w_k(n)\, e^{-\frac{1}{2} e^2 \langle j_n,\, C j_n\rangle_\varepsilon} \cos\big(teY(n)\big),$$

where $\langle j_n,\, Cf\rangle_\varepsilon = Y(n)$, and $w_k(n) \ge 0$, $w_k(-n) = w_k(n)$ are the weights of Lemma 2.5; in the state $\flat$ the Haar average over the zero mode restricts the sum to $\omega(n) = 0$, as there. By the formula for $w_k(n)$ in the proof of Lemma 2.5, $w_k(n) e^{-\frac{1}{2} e^2 \langle j_n,\, C j_n\rangle}$ is a positive constant (independent of n) times the n-marginal of $Z_{\mathbb{P}}\, e^{r^2(\varepsilon^3 k)}\, \mathbb{P}(n, dr)$; so the probability ν_k on $\mathcal{N}$ proportional to $w_k(n) e^{-\frac{1}{2} e^2 \langle j_n,\, C j_n\rangle}$ is the n-marginal of $\mathbb{P}_k$. Dividing by the value at $t = 0$,

$$\langle e^{tF(g)}\rangle_k = e^{\frac{1}{2} t^2 q}\, \phi(t), \qquad \phi(t) := \mathbb{E}_{\nu_k}\big[\cos(teY)\big].$$

The left side is finite for all t (Lemma 7.3 with $\pm t$) and smooth in t, since $F(g)$ has exponential moments of every order. So ϕ, the characteristic function of the symmetric law of eY under ν_k, is C^2, and $\phi''(0) = -e^2 \mathbb{E}_{\nu_k}[Y^2]$: indeed $(1 - \cos(tx))/t^2 \to x^2/2$ and $0 \le (1 - \cos(tx))/t^2 \le x^2/2$, so Fatou's lemma gives $\mathbb{E}[(eY)^2] \le -\phi''(0) < \infty$, and then dominated convergence gives equality. Differentiating twice at $t = 0$ gives $\langle F(g)^2\rangle_k = q + \phi''(0) = q - e^2 \mathbb{E}_{\mathbb{P}_k}[Y^2]$.

(b) $\langle F(g)\rangle = \langle F(g) X_{\mathrm{R}}(h)\rangle = 0$ by charge conjugation (Lemma 7.13(iii)), which reverses the sign of F and fixes X_{R}. Put $k = sh$ with $s \in \mathbb{R}$ small. The derivative at $s = 0$ of $\langle F(g)^2\rangle_{sh}$ is $\langle F(g)^2 X_{\mathrm{R}}(h)\rangle - \langle F(g)^2\rangle \langle X_{\mathrm{R}}(h)\rangle$, and the derivative of $\mathbb{E}_{\mathbb{P}_{sh}}[Y^2]$ is $\mathrm{Cov}_{\mathbb{P}}(Y^2, r^2(\varepsilon^3 h))$; in both cases differentiation under the integral is justified by the integrability above (with $e^{|s|\, r^2(\varepsilon^3|h|)}$ absorbed by $e^{-\lambda\varepsilon^3 r^4}$). Apply (a). By Lemma 12.2 (on the lattice, odd curvature moments vanish by charge conjugation) the left side is $M_{\varepsilon,j,\mu_0}(g; h)$. □

12.3. **Signs in every limit state.**

Theorem 12.4 (signs)**.** *(a)* Lattice. *Let $\mu_0 \in (0, \mu_*]$, let $T_{j,\varepsilon}$ be any lattice torus, $b_T \in \mathbb{R}$, $\lambda > 0$, $e \in \mathbb{R}$. For all real lattice 2-forms g, g' and all real site functions $h, h' \geq 0$:*

(i) *$M_{\varepsilon,j,\mu_0}(g;h) \leq 0$; equivalently, $\mathrm{Cov}_{\mathbb{P}}(Y^2, r^2(\varepsilon^3 h)) \geq 0$ when $e \neq 0$;*
(ii) *$\langle X_{\mathrm{R}}(h); X_{\mathrm{R}}(h')\rangle \geq 0$;*
(iii) *$\langle F(g)^2; F(g')^2\rangle \geq 0$,*

where $\langle U; V\rangle := \langle UV\rangle - \langle U\rangle\langle V\rangle$. The same holds in the state $\flat$, for $e \neq 0$, on every lattice torus $T_{j,\varepsilon}$.

(b) Limits. *Let $S \in \{S^{\mu_0}\ (\mu_0 \in Q_0),\ S^\flat,\ S^0\}$ be a limit state of Theorem 3.1. For all real 2-forms g, g' with components in $C_c(\mathbb{R}^3)$ and all $h, h' \in C_c(\mathbb{R}^3)$ with $h, h' \geq 0$:*

$$M(g;h) \leq 0, \qquad S^T_{0,2}(h,h') \geq 0, \qquad S^T_{4,0}(g,g,g',g') + 2\,S_{2,0}(g,g')^2 \geq 0.$$

Proof. (a) *The setting of* [8]*, §6.* There $d\mu_C$ is an arbitrary centred Gaussian measure with covariance $C > 0$ on real bond variables ϑ_b, the matter action is $-\sum_b c_b r_{b_-} r_{b_+} \cos(-\theta_{b_-} + \vartheta_b + \theta_{b_+})$ with $c_b \geq 0$, and at each site there is a finite measure $d\rho_x(r)$ with $\int e^{\alpha r^2} d\rho_x < \infty$ for all $\alpha > 0$; periodic conditions on a rectangular lattice are allowed. The lattice state $\langle\cdot\rangle_{\varepsilon,j,\mu_0}$ is of this form, with $\vartheta_b := e\varepsilon A_b$ (centred Gaussian with covariance $(e\varepsilon)^2 C_{\mu_0} > 0$ for $e \neq 0$; for $e = 0$, where this covariance vanishes, the curvature is independent of the matter, so (i) holds with $M_{\varepsilon,j,\mu_0}(g;h) = 0$ and (iii) is the Gaussian identity $\langle F(g)^2; F(g')^2\rangle = 2\langle F(g)F(g')\rangle^2 \geq 0$, while (ii) follows from the case $e \neq 0$ by dominated convergence in e at fixed lattice and fixed b_T, since $|\cos| \leq 1$ in the hopping term and $\lambda > 0$), $c_b = \varepsilon$, and $d\rho_x(r) = e^{-V(r)} r\,dr$, by the polar form of the hopping term in the proof of Lemma 2.5; the integrability condition holds because $\lambda > 0$. (BFS let the angles run over $[-2\pi, 2\pi)$; for 2π-periodic functions this does not change expectations.) For real u on bonds, $A_{\mathrm{BFS}}(u) := \sum_b u_b \vartheta_b$; with $u_b := \varepsilon^2 (d^* g)_b / e$ this is $F(g) = \sum_b \varepsilon^3 (d^* g)_b A_b$.

(i) By [8, Thm. 6.2(1), p. 280], applied with the tilt $e^{r^2(\gamma)}$, $\gamma := \varepsilon^3(t\delta_x + \eta)$ $(t, \eta > 0)$, $f = 0$ and the test function u: $\langle e^{r^2(\gamma)}; F(g)^2\rangle \leq 0$. Let $\eta \downarrow 0$ (dominated convergence; all moments of $e^{c\,r_y^2} F(g)^2$ are finite for every c, since $\lambda > 0$). Then $\langle e^{t\varepsilon^3 r_x^2}; F(g)^2\rangle \leq 0$ for every $t > 0$. Divide by t and let $t \downarrow 0$: $t^{-1}(e^{t\varepsilon^3 r_x^2} - 1) \to \varepsilon^3 r_x^2$, dominated by $\varepsilon^3 r_x^2 e^{\varepsilon^3 r_x^2}$ for $t \leq 1$. Hence $\langle r_x^2; F(g)^2\rangle \leq 0$ for every site x. Multiply by $\varepsilon^3 h(x) \geq 0$ and sum; the constants $W_\varepsilon + z_\varepsilon$ in $X_{\mathrm{R}}(h)$ do not change covariances. So $\langle X_{\mathrm{R}}(h); F(g)^2\rangle \leq 0$, which is $M_{\varepsilon,j,\mu_0}(g;h) \leq 0$ by Lemma 12.3(b). For $e \neq 0$ the equivalence with $\mathrm{Cov}_{\mathbb{P}} \geq 0$ is the same lemma.

(ii) By [8, Thm. 6.1, p. 276] with $P(r) = r_x^2$, $Q(r) = r_y^2$ (polynomials with positive coefficients), without current factors and without test functions for A (in the notation of [8, Thm. 6.1], $n = m = 0$ and $f = g = 0$; these m and g are not the mass and the 2-form of this section): $\langle r_x^2; r_y^2\rangle \geq 0$. Multiply by $\varepsilon^6 h(x) h'(y) \geq 0$ and sum.

(iii) By [8, Thm. 6.2(1)] with the tilt $e^{r^2(\eta)}$, $\eta > 0$ constant, $f = tu'$ (u' the coefficient vector of $F(g')$) and the test function u, and $\eta \downarrow 0$: $\langle \cos(tF(g')); F(g)^2\rangle \leq 0$ for all $t \in \mathbb{R}$. (In the proof of [8, Thm. 6.2], $e^{iA(f)}$ is replaced by $\cos A(f)$, by the symmetry $A \to -A$, $\theta \to -\theta$.) Since $|\cos x - 1 + x^2/2| \leq x^4/24$ and $F(g'), F(g)$ have all moments, $\langle \cos(tF(g')); F(g)^2\rangle = -\frac{t^2}{2}\langle F(g')^2; F(g)^2\rangle + O(t^4)$. Hence $\langle F(g')^2; F(g)^2\rangle \geq 0$.

The state $\flat$. At fixed lattice, $\langle\cdot\rangle_{\varepsilon,j,\flat} = \lim_{\mu_0 \downarrow 0} \langle\cdot\rangle_{\varepsilon,j,\mu_0}$ on every $G \in \mathfrak{A}_\varepsilon$ (Definition 2.4) with $\sup_{0<\mu_0\leq\mu_*} \langle |G|^2\rangle_{\varepsilon,j,\mu_0} < \infty$ (Lemma 2.8 with $\delta = 1$; it is stated for $b = b_{T_j}(\varepsilon)$, but its proof does not use the value of b, so it holds for every real b_T). The products that occur in (i)–(iii) are of this kind. By Hölder's inequality it suffices to bound $\langle F(g)^8\rangle_{\mu_0}$ and $\langle r_x^8\rangle_{\mu_0}$ uniformly in μ_0. The first is at most $8!\,(\langle e^{F(g)}\rangle + \langle e^{-F(g)}\rangle) \leq 2 \cdot 8!\, e^{\frac{1}{2}\|g\|^2_{2,\varepsilon}}$ by Lemma 7.3. For the second, put $\Psi(A) := \Psi_{r_x^8}(A) = \int r_x^8\, e^{-S^{\mathrm{m}}_{b_T}(A,\varphi)} \prod_y d^2\varphi(y)$ (Definition 2.4, with b_T in place of $b_{T_j}(\varepsilon)$). The proof of Lemma 2.5, with the factor r_x^8 inside the radial integral, gives $\Psi(A) = \sum_n w'(n) e^{ie\varepsilon \sum_b n_b A_b}$ with weights $w'(n) \geq 0$,

so $0 \le \Psi(A) \le \Psi(0)$. Moreover, $\mathbb{E}_{\gamma_{\mu_0}}[W_0] = \sum_n w_0(n) e^{-\frac{1}{2}e^2\langle j_n, C_{\mu_0} j_n\rangle} \ge w_0(0) > 0$ (the $n=0$ term). Hence $\langle r_x^8\rangle_{\mu_0} = \mathbb{E}_{\gamma_{\mu_0}}[\Psi]/\mathbb{E}_{\gamma_{\mu_0}}[W_0] \le \Psi(0)/w_0(0)$ for all μ_0. The inequalities (i)–(iii) pass to the limit $\mu_0 \downarrow 0$.

(b) The truncated functions in (b) are polynomials in the Schwinger functions $S_{2,1}, S_{2,0}, S_{0,1}, S_{0,2}, S_{4,0}$. Along the subsequence of Theorem 3.1(ii) the lattice Schwinger functions at sampled test functions converge to those of S^{μ_0} and $S^{\flat}$, and the Schwinger functions of S^{μ_0} converge to those of S^0 as $\mu_0 \downarrow 0$ in Q_0. Sampling preserves $h, h' \ge 0$. The quantities in (a) are, on the lattice, $M_{\varepsilon,j,\mu_0}(g;h)$ (Lemma 12.3(b)), $\langle X_{\mathrm{R}}(h); X_{\mathrm{R}}(h')\rangle = S^T_{0,2}(h,h')$ and $\langle F(g)^2; F(g')^2\rangle = S_{4,0}(g,g,g',g') - S_{2,0}(g,g)S_{2,0}(g',g') = S^T_{4,0}(g,g,g',g') + 2S_{2,0}(g,g')^2$; the last identity is the fourth-order expansion of the logarithm, in which the terms with an odd number of curvature arguments vanish by charge conjugation, as in the proof of Lemma 12.2. The inequalities pass to the limits. □

Corollary 12.5 (a criterion for non-triviality)**.** *Let S be a limit state as in Theorem 12.4(b). If $M(g;h) < 0$ for one real 2-form g and one $h \ge 0$, then S is not decoupled. Hence S is not Gaussian, and the curvature and the Wick square are not independent under the law of S. In particular S is not the free Maxwell field (of any photon mass) together with the Wick square of a free charged scalar. By Theorem 12.4, the hypothesis is equivalent to $M(g;h) \ne 0$.*

12.4. **The leading coefficient.** On a fixed lattice torus regard the coefficient $b_T = a_{K,T}(m)$ of the torus counterterm scheme (Definition 2.2) as the polynomial in (e^2, λ) given there; at $e = \lambda = 0$ it equals m^2. Write $\mathbb{P}^{e,\lambda}$ for the measure (12.1) at these couplings, and for $\lambda = 0$ use (12.1) with $\lambda = 0$ and $b_T = m^2$. A closed walk $\omega = (\omega_0, \omega_1, \dots, \omega_N = \omega_0)$ on $T_{j,\varepsilon}$ has $|\omega| = N \ge 1$ nearest-neighbour steps. For a function a on oriented bonds ($a_{-b} = -a_b$) put $\Phi_\omega(a) := \sum_{i=0}^{N-1} a_{(\omega_i,\omega_{i+1})}$.

Proposition 12.6 (the leading coefficient on a fixed lattice torus)**.** *Fix ε, j, $m > 0$ and $\mu_0 \in (0, \mu_*] \cup \{\flat\}$, a real lattice 2-form g and a real site function h.*

(a) *For $\lambda > 0$, $\lim_{e\to 0} e^{-2} M^{(e,\lambda)}_{\varepsilon,j,\mu_0}(g;h) = c_{\varepsilon,j,\mu_0}(g,h;\lambda) := -\mathrm{Cov}_{\mathbb{P}^{0,\lambda}}\big(Y^2, r^2(\varepsilon^3 h)\big)$. If $h \ge 0$, then $c_{\varepsilon,j,\mu_0}(g,h;\lambda) \le 0$.*

(b) $\lim_{\lambda\downarrow 0} c_{\varepsilon,j,\mu_0}(g,h;\lambda) = c^{\mathrm{free}}_{\varepsilon,j,\mu_0}(g,h) := -\mathrm{Cov}_{\mathbb{P}^{0,0}}\big(Y^2, r^2(\varepsilon^3 h)\big)$.

(c) *For $\mu_0 \in (0, \mu_*]$, with $a = C_{\mu_0} d^* g$, the sum below converges absolutely and*

$$c^{\mathrm{free}}_{\varepsilon,j,\mu_0}(g,h) = -\frac{2\varepsilon^4}{6+\varepsilon^2 m^2} \sum_{\omega} \frac{(6+\varepsilon^2 m^2)^{-|\omega|}}{|\omega|}\, \Phi_\omega(a)^2 \sum_{i=0}^{|\omega|-1} h(\omega_i), \tag{12.2}$$

the sum running over all closed walks on $T_{j,\varepsilon}$.

(d) *For $\mu_0 \in (0, \mu_*]$: if $h \ge 0$, $h \not\equiv 0$ and $d^* g \not\equiv 0$, then $c^{\mathrm{free}}_{\varepsilon,j,\mu_0}(g,h) < 0$.*

Proof. (a) By Lemma 12.3(b), $e^{-2} M^{(e,\lambda)} = -\mathrm{Cov}_{\mathbb{P}^{e,\lambda}}(Y^2, r^2(\varepsilon^3 h))$ for $e \ne 0$. The density of $\mathbb{P}^{e,\lambda}$ depends on e through $e^{-\frac{1}{2}e^2\langle j_n, C j_n\rangle} \le 1$ and through b_T, which is continuous in e. Let $b_{\min} := \min_{|e|\le 1} b_T$. For $|e| \le 1$ the integrands of $Z_{\mathbb{P}}$, $\mathbb{E}_{\mathbb{P}}[Y^2 r^2(\varepsilon^3 h)]$, $\mathbb{E}_{\mathbb{P}}[Y^2]$ and $\mathbb{E}_{\mathbb{P}}[r^2(\varepsilon^3 h)]$ are dominated by the e-independent integrable function of the proof of Lemma 12.3, with b_T replaced by $b_{\min}$. So the covariance is continuous in e, and its value at $e = 0$ is the limit. The sign follows from Theorem 12.4(a)(i).

(b) At $e = 0$, $b_T = m^2 + \lambda\beta_1 + \lambda^2\beta_2$ with real β_1, β_2 depending on (ε, T) (Definition 2.2: at $e = 0$ the monomials are λ and λ^2). For $0 \le \lambda \le \lambda_1$ small, $b_T \ge m^2/2$. With $\prod_b I_{n_b}(\varepsilon r_{b_-} r_{b_+}) \le$ (polynomial in r and n) $\times \prod_b e^{\varepsilon r_{b_-} r_{b_+}}$ and $\sum_b \varepsilon r_{b_-} r_{b_+} \le 3\varepsilon \sum_x r_x^2$, the integrands are dominated by a polynomial times $\prod_x e^{-\varepsilon^3 m^2 r_x^2/4}$, uniformly in $\lambda \in [0, \lambda_1]$. Dominated convergence gives the limit.

(c) Let Θ be a real function on oriented bonds. By the Bessel expansion of Lemma 2.5, with the phase $e\varepsilon A_b$ replaced by Θ_b,

$$\sum_n w_k(n)\, e^{i\sum_b n_b\Theta_b} = \int e^{-S^\Theta(\varphi)+X_{\mathrm{R}}(k)}\, d\varphi,$$

where S^Θ is the matter action with the hopping $|U(\Theta_b)\varphi(b_+)-\varphi(b_-)|^2$. At $\lambda=0$, $b_T=m^2$, the right side is a complex Gaussian integral: $S^\Theta-\sum_x \varepsilon^3 k(x)|\varphi(x)|^2 = \frac12\langle\varphi,\,(H_\Theta-2k)\varphi\rangle_\varepsilon$ with $H_\Theta=-\Delta_\Theta+m^2$ ($-\Delta_\Theta\ge 0$ the covariant lattice Laplacian on $\mathbb{C}$-valued functions). Hence, for $\|k\|_\infty<m^2/2$, it equals a Θ-independent constant times $\det(H_\Theta-2k)^{-1}$. Take $\Theta=\tau\varepsilon a$, so that $\sum_b n_b\Theta_b=\tau Y(n)$. Since the weights are even in n, the first τ-derivative at 0 vanishes, and

$$\mathbb{E}_{\mathbb{P}_k^{0,0}}[Y^2] = -\partial_\tau^2 \log\Big(\sum_n w_k(n)e^{i\tau Y(n)}\Big)\Big|_{\tau=0} = \partial_\tau^2\ \mathrm{tr}\log\big(H_{\tau\varepsilon a}-2k\big)\Big|_{\tau=0}.$$

Write $H_\Theta-2k=D_k(1-K_\Theta)$, with D_k the diagonal operator $6\varepsilon^{-2}+m^2-2k(x)$ and K_Θ the hopping operator, whose matrix elements are $D_k(x)^{-1}\varepsilon^{-2}e^{\pm i\Theta_b}$ for x,y joined by a bond b. Its norm is at most $6/(6+\varepsilon^2(m^2-2\|k\|_\infty))<1$. So $\mathrm{tr}\log(H_\Theta-2k)=\mathrm{tr}\log D_k-\sum_{N\ge1}N^{-1}\,\mathrm{tr}\,K_\Theta^N$, and $\mathrm{tr}\,K_\Theta^N=\sum_{|\omega|=N}u_k(\omega)e^{\pm i\Phi_\omega(\Theta)}$, where $u_k(\omega):=\prod_{i=0}^{N-1}\big(6+\varepsilon^2(m^2-2k(\omega_i))\big)^{-1}$; the global sign of the phase is a convention and plays no role. Since $|\Phi_\omega(a)|\le|\omega|\|a\|_\infty$ and there are at most $|T_{j,\varepsilon}|6^N$ closed walks of length N, the series may be differentiated termwise. This gives

$$\mathbb{E}_{\mathbb{P}_k^{0,0}}[Y^2] = \varepsilon^2\sum_\omega |\omega|^{-1}u_k(\omega)\,\Phi_\omega(a)^2.$$

Finally $c^{\mathrm{free}}=-\partial_s\mathbb{E}_{\mathbb{P}_{sh}^{0,0}}[Y^2]\big|_{s=0}$ (as in the proof of Lemma 12.3(b); at $\lambda=0$ the integrability used there is supplied by the domination of (b), with $b_T=m^2$: a polynomial times $\prod_x e^{-\varepsilon^3 m^2 r_x^2/4}$, valid for $|s|\,\|h\|_\infty<m^2/4$), and $\partial_s u_{sh}(\omega)|_{s=0}=u_0(\omega)\sum_i 2\varepsilon^2h(\omega_i)/(6+\varepsilon^2m^2)$ with $u_0(\omega)=(6+\varepsilon^2m^2)^{-|\omega|}$. Termwise differentiation in s is justified by the same bounds, uniformly for $|s|\,\|h\|_\infty<m^2/4$. This is (12.2).

(d) Every term of (12.2) is ≤ 0, so it suffices to find one closed walk ω with $\Phi_\omega(a)\neq0$ that visits a site where $h>0$. Since $C_{\mu_0}>0$ and $f=d^*g\neq0$, $\langle g,\,dC_{\mu_0}d^*g\rangle_\varepsilon=\langle f,\,C_{\mu_0}f\rangle_\varepsilon>0$, so $da=dC_{\mu_0}d^*g\neq0$. Choose a plaquette p with $(da)(p)\neq0$, a site x_0 with $h(x_0)>0$, and a nearest-neighbour path γ from x_0 to a corner of p. Let ω be: γ, then once around ∂p, then γ backwards. The two passages along γ cancel in Φ_ω, so $\Phi_\omega(a)=\pm\sum_{b\in\partial p}\pm a_b=\pm\varepsilon(da)(p)\neq0$, and $\omega_0=x_0$. □

Proposition 12.6 is a statement on one lattice torus. Its continuum counterpart is the following hypothesis, which we formulate and do not prove.

Hypothesis 12.7 (the continuum value of the leading coefficient)**.** For every real 2-form g with components in $C_c^\infty(\mathbb{R}^3)$ and every $0\le h\in C_c^\infty(\mathbb{R}^3)$, sampled as in Definition 2.9, the following holds, with $a_{\mu_0}:=(-\Delta+\mu_0^2)^{-1}d^*g$ on $\mathbb{R}^3$ for $\mu_0\in[0,\mu_*]$. For every $\mu_0\in[0,\mu_*]$ the integral

$$c_{\mu_0}^{\mathrm{free}}(g,h) := -2\int_0^\infty \frac{dT}{T}\,\frac{e^{-m^2T}}{(4\pi T)^{3/2}}\int_{\mathbb{R}^3}dx\ \mathbb{E}_{x,x}^T\Big[\Big(\oint_\omega a_{\mu_0}\cdot d\omega\Big)^2\int_0^T h(\omega_t)\,dt\Big] \tag{12.3}$$

converges, where $\mathbb{E}_{x,x}^T$ is the expectation for the Brownian bridge ω with generator Δ from x to x in time T, and $\oint$ is the Stratonovich line integral. As $\varepsilon\to0$ and $\ell_j\to\infty$, $c_{\varepsilon,j,\mu_0}^{\mathrm{free}}(g,h)\to c_{\mu_0}^{\mathrm{free}}(g,h)$ for $\mu_0\in(0,\mu_*]$, and $c_{\varepsilon,j,\mathfrak{b}}^{\mathrm{free}}(g,h)\to c_0^{\mathrm{free}}(g,h)$. The function $\mu_0\mapsto c_{\mu_0}^{\mathrm{free}}(g,h)$ is continuous on $[0,\mu_*]$, and $c_{\mu_0}^{\mathrm{free}}(g,h)<0$ if $d^*g\not\equiv0$ and $h\not\equiv0$.

Remark 12.8 (on Hypothesis (C))*.* We refer to Hypothesis 12.7 as (C). It is a one-loop statement: by Proposition 12.6(c), $c_{\varepsilon,j,\mu_0}^{\mathrm{free}}(g,h)$ is a weighted sum over closed lattice walks, and (12.3) is its expected

Brownian-bridge limit. A proof would combine the convergence of the lattice coefficient (by lattice power counting [32] in its three-dimensional form, Proposition C.3, applied to the two one-loop graphs, the triangle with two current vertices and one insertion of $|\varphi|^2$ and the bubble with one seagull vertex and one insertion, both of superficial degree -1 in $d=3$, or by an invariance principle for the weighted walks of (12.2), with the discrete line integral converging to the Stratonovich integral); the Feynman–Kac–Itô formula for $-(\nabla - i\tau a_{\mu_0})^2 + m^2 - 2sh$ $(\tau, s \in \mathbb{R})$, differentiated twice in τ and once in s at $\tau = s = 0$, which gives (12.3); the support theorem for the Brownian bridge, for the strict negativity; and, for the state $\flat$, the fact that the restriction of $\mathcal{N}$ to currents with zero windings changes the coefficient by $o(1)$ as $\ell_j \to \infty$. We do not carry this out.

12.5. **Non-triviality given a remainder bound.**

Theorem 12.9 (non-triviality under (R) and (C))**.** *Assume:*

(R) *There are $\delta_R \in (0, 0.1]$ and a function ρ on $(0, \delta_R]$ with $\rho(t) \to 0$ as $t \downarrow 0$ such that the following holds. For every pair g, h as in Hypothesis 12.7 there is $\kappa(g,h) < \infty$ such that, for every (e, λ) with $\lambda > 0$, $e^2 > 0$ and $e^2 + \lambda \le \delta_R m$ (such (e,λ) satisfy the hypotheses of Theorem 3.1, since $\delta_R \le 0.1$), the lattice states of Theorem 3.1 at (e,λ), with the constants ε_0 and K_0 of that theorem chosen at (e,λ) and $\varepsilon_K = 2^{-K}\varepsilon_0$, satisfy*

$$\limsup_{K\to\infty,\ j\to\infty} \sup_{\mu_0 \in (0,\mu_*]\cup\{\flat\}} \left| e^{-2} M_{\varepsilon_K, j, \mu_0}(g;h) - c^{\mathrm{free}}_{\varepsilon_K, j, \mu_0}(g,h) \right| \ \le\ \rho\big((e^2+\lambda)/m\big)\, \kappa(g,h).$$

Assume also (C). Let g, h be as in Hypothesis 12.7, with $d^*g \not\equiv 0$ and $h \not\equiv 0$, and put $c_0(g,h) := -\max_{\mu\in[0,\mu_*]} c^{\mathrm{free}}_{\mu}(g,h) > 0$. If (e,λ) is as in (R) and $\rho((e^2+\lambda)/m)\kappa(g,h) < c_0(g,h)/2$, then every limit state $S \in \{S^{\mu_0}\ (\mu_0 \in Q_0), S^{\flat}, S^0\}$ of Theorem 3.1 at (e,λ) satisfies

$$M(g;h) \le -\tfrac{1}{2} e^2 c_0(g,h) < 0.$$

In particular, for $e^2 + \lambda$ small enough (depending on g, h) every limit state is not decoupled and not Gaussian (Corollary 12.5).

Proof. $c_0 > 0$ by the strict negativity and the continuity in (C) ($[0,\mu_*]$ is compact). Let S^{μ_0} or $S^{\flat}$ be the limit along $(\varepsilon_{K_l}, T_{j_l})$. By (R) and (C), $\limsup_l e^{-2} M_{\varepsilon_{K_l}, j_l, \mu_0}(g;h) \le \lim_l c^{\mathrm{free}}_{\varepsilon_{K_l}, j_l, \mu_0}(g,h) + c_0/2 \le -c_0/2$, and the same for $\flat$. By Theorem 3.1(ii) the left side is $e^{-2}M(g;h)$ for S^{μ_0}, resp. $S^{\flat}$. For S^0, M is the limit of the values for S^{μ_0}, $\mu_0 \downarrow 0$ in Q_0, each of which is $\le -e^2 c_0/2$. □

Remark 12.10 (what (R) asks). (i) *What the uniform bounds give.* Theorem 6.8 and Proposition 8.1 bound every Schwinger function, and so $|M(g;h)|$, by a constant times $\|g\|_2^2 N(h)$. The constants σ_+, C (and $\varepsilon_0, \ell_0, K_0$) are produced at each point of $\mathfrak{W}$ by the conditions of Appendix A and Lemma A.5, and nothing is stated about how they depend on (e,λ). So the Cauchy estimates give neither a factor e^2 nor any lower bound: the tube bound is uniform in the sources and in the regulators, not in the couplings. (ii) *What (R) asks.* By Lemma 12.3, (R) says that the matter covariance $\mathrm{Cov}_{\mathbb{P}^{e,\lambda}}(Y^2, r^2(\varepsilon^3 h))$ stays within $\rho\kappa$ of its value at $(e,\lambda) = (0,0)$, uniformly in ε, j and μ_0. Moving λ from 0 inserts $\sum_x \varepsilon^3 r_x^4$ together with the λ-dependent part of b_T; one-loop power counting suggests that each of the two insertions alone diverges as $\varepsilon \to 0$, like $\lambda\varepsilon^{-1}$, and that only their sum is finite. Moving e from 0 inserts the induced current interaction $e^{-\frac{1}{2}e^2\langle j_n, C j_n\rangle}$ and the e^2-dependent part of b_T. A bound uniform in ε on these insertions in a three-point function is a renormalised perturbative estimate with a remainder. As printed, [1, 2, 3, 4] give two-sided bounds on partition functions, [24, Thm. 2.1, p. 654] gives the convergence of $Z(T,g,h)$ and $|\ln Z(T,g,h)| \le C|T|$ with C depending on g, h and $|T|$, and [24, Thm. 3.4, p. 656] compares effective actions at two spacings; none of them states asymptotics in (e,λ) of $\ln Z(T,g,h) - \ln Z(T,0,0)$. (iii) *Signs do not suffice.* The proof of Theorem 12.9 uses only the upper half of (R), the bound $M \le e^2(c^{\mathrm{free}} + \rho\kappa)$. The correlation inequalities of [8] give only non-strict signs, and the strict sign of Proposition 12.6(d)

holds on each lattice but is not uniform. Lebowitz's inequality concerns the field φ, which is not gauge invariant and is not among our observables.

Remark 12.11 (the notion of non-triviality). Here non-triviality means that the state is not decoupled (Definition 12.1), the conclusion of Corollary 12.5. This is the informative notion: the free Maxwell field together with the Wick square of a free charged scalar is decoupled, but as a joint family it is not Gaussian, because the Wick square of a free field is not Gaussian. No result of this paper excludes a Gaussian limit state: such a state would have $M \equiv 0$, which is compatible with every sign in Theorem 12.4, and the Gaussian domination bound of Lemma 7.3 allows the free Maxwell value. Nothing here concerns non-Gaussianity of the curvature field alone (Theorem 12.4(b) allows $S^T_{4,0} = 0$).

13. The curvature two-point function

This section collects what the methods of the paper give about the photon mass in the massless states $S^\flat$ and S^0. The answer is one-sided. We prove the Bianchi identity in every limit state; the curvature two-point function of every limit state is a Fourier multiplier dominated by that of the free massless photon; at positive photon mass it is suppressed at small momenta, so S^{μ_0} has no massless component. For $S^\flat$ and S^0 we prove that a massless photon, in the sense of Definition 13.6, excludes integrable, in particular exponentially decaying, curvature correlations and excludes a gap in the curvature sector of the reconstructed theory (Theorem 13.7, Proposition 13.8, Corollary 13.9). Whether $S^\flat$ or S^0 is massless is not decided here: it needs a lower bound on the curvature two-point function, which no tool of this paper gives (Remark 13.10 and Section 14.4).

Standing assumptions. S is a limit point in the sense of Section 9. That is, S is S^{μ_0} with $\mu_0 \in Q_0$, or a limit point at a photon mass $\mu_0 \in (0, \mu_*]$ along a further subsequence, or $S^\flat$, or S^0; the fixed-torus continuum states, which Section 9 includes where stated, are not limit points here (they are not invariant under all translations of $\mathbb{R}^3$, which Lemma 13.4 needs). We say that S is *at photon mass* $\mu_0 > 0$ in the first two cases. The lattice states are those of Definition 2.7. Everything below uses only Theorem 3.1(i)–(iv), Lemmas 2.5, 7.3 and 7.13, Definition 2.9, Propositions 8.1, 8.2 and 8.3, and Theorems 9.2, 9.4 and 9.8. Theorem 3.5 gives the statements (i)–(iv) of Theorem 3.1 at blocking factor 3, under its hypothesis, and the lattice lemmas used here concern one torus and do not involve L. So every result of this section holds, under the same hypothesis, for the limit states of Theorem 3.5.

Notation. We write curvature test functions as vector fields. For a real $G \in C_c(\mathbb{R}^3; \mathbb{R}^3)$ let $\star^{-1}G$ be the 2-form with components $g_{23} = G_1$, $g_{13} = -G_2$, $g_{12} = G_3$. Then $\|\star^{-1}G\|_2 = \|G\|_2$, and we put

$$B(G) := F(\star^{-1}G), \qquad \mathfrak{b}_S(G, G') := S_{2,0}(\star^{-1}G, \star^{-1}G'),$$

the smeared magnetic field $B = \star F$ and its two-point function. A direct computation gives $(\delta \star^{-1} G)_\nu = (\operatorname{curl} G)_\nu$ for $G \in C^1_c$, where δ is the L^2-adjoint of d on forms, $(\delta g)_\nu = -\sum_\mu \partial_\mu g_{\mu\nu}$. The Fourier transform is $\hat f(p) = \int e^{-ip\cdot x} f(x)\, dx$, so that $\|f\|_2^2 = (2\pi)^{-3} \int |\hat f|^2 dp$. For $p \neq 0$ let $P_T(p) := \mathbf{1} - pp^{\mathsf T}/|p|^2$ be the orthogonal projection of $\mathbb{C}^3$ onto $p^\perp$, and let Π_T be the Fourier multiplier by P_T on $L^2(\mathbb{R}^3; \mathbb{C}^3)$. For $R > 0$ put

$$G_R(x) := R^{-3} G(x/R), \qquad \widehat{G_R}(p) = \hat G(Rp), \qquad \|G_R\|_2^2 = R^{-3}\|G\|_2^2.$$

The *free massless photon* is the centred Gaussian curvature field with $\mathfrak{b}(G, G') = \langle \Pi_T G,\, \Pi_T G' \rangle_{L^2}$, the curvature of the Feynman-gauge massless vector field:

$$\langle \delta \star^{-1} G,\, (-\Delta)^{-1} \delta \star^{-1} G \rangle = (2\pi)^{-3} \int |p \times \hat G|^2 |p|^{-2} dp = \|\Pi_T G\|_2^2.$$

13.1. **Two lattice facts.**

Lemma 13.1 (free part minus screening, on one torus)**.** *Fix (ε, T_j), $\mu_0 \in (0, \mu_*] \cup \{\flat\}$ and a real lattice 2-form r on $T_{j,\varepsilon}$. Let C be as in Lemma 7.3, let $w_0(n)$ be the weights of Lemma 2.5 at $k = 0$, and put $j_n := \varepsilon^{-2} n$. Let $\tilde\nu$ be the probability measure on divergence-free integer currents n with $\tilde\nu(n) \propto w_0(n)\, e^{-\frac12 e^2 \langle j_n, C j_n\rangle_\varepsilon}$, multiplied by $\mathbf{1}\{\omega(n) = 0\}$ in the $\flat$ state, and $a_r(n) := e\langle j_n,\, C\, d^*r\rangle_\varepsilon$. Then:*

(a) *for every real t, $\langle e^{tF(r)}\rangle_{\varepsilon,j,\mu_0} = e^{\frac12 t^2 V(r)} \sum_n \tilde\nu(n) \cos\big(t\, a_r(n)\big)$, with $V(r) = \langle d^*r,\, C\, d^*r\rangle_\varepsilon$;*

(b) *$\langle F(r)\rangle_{\varepsilon,j,\mu_0} = 0$, $\sum_n \tilde\nu(n) a_r(n)^2 < \infty$, and*

$$\langle F(r)^2\rangle_{\varepsilon,j,\mu_0} = V(r) - \sum_n \tilde\nu(n)\, a_r(n)^2 \;\le\; V(r) \;\le\; \|r\|^2_{2,\varepsilon};$$

(c) *if $\mu_0 \in (0, \mu_*]$, then $V(r) \le \mu_0^{-2} \|d^*r\|^2_{2,\varepsilon}$.*

Proof. (a) Put $f := t\, d^*r$. The proof of Lemma 7.3 (case $\mu_0 > 0$, with $k = 0$) gives

$$\mathbb{E}_C\big[e^{\langle f,\, A\rangle_\varepsilon} W_0(A)\big] = e^{\frac12 \langle f,\, Cf\rangle_\varepsilon} \sum_n w_0(n)\, e^{-\frac12 e^2 \langle j_n,\, C j_n\rangle_\varepsilon} \cos\big(e\langle j_n,\, Cf\rangle_\varepsilon\big),$$

and the same identity at $f = 0$ gives $\mathbb{E}_C[W_0] = \sum_n w_0(n) e^{-\frac12 e^2 \langle j_n,\, C j_n\rangle_\varepsilon}$. Divide. In the $\flat$ state the average over the zero mode replaces $w_0(n)$ by $w_0(n)\mathbf{1}\{\omega(n) = 0\}$, as in that proof. For such n the phase $e\langle j_n,\, c\rangle_\varepsilon = e\ell_T \sum_\mu c_\mu \omega_\mu(n)$ of a constant c vanishes (Lemma 2.5), so only $A^\perp$, with covariance $C = (-\Delta_\varepsilon)^{-1}\Pi^\perp$, enters, and the same Gaussian computation applies. (b) Let $\phi(t) := \sum_n \tilde\nu(n) \cos(t a_r(n))$ and $f(t) := \langle e^{tF(r)}\rangle$. Then $f(t) = e^{\frac12 t^2 V} \phi(t)$ is finite for all real t by (a). Hence $\langle e^{|t||F(r)|}\rangle \le f(t) + f(-t) < \infty$, and f is real-analytic with $f^{(k)}(0) = \langle F(r)^k\rangle$. Since ϕ is even, so is f; thus $\langle F(r)\rangle = 0$, and $e^{-\frac12 t^2 V} f(t) = 1 + \frac12 t^2(\langle F(r)^2\rangle - V) + O(t^4)$. Therefore

$$\sum_n \tilde\nu(n)\, \frac{2(1 - \cos(t a_r(n)))}{t^2} = \frac{2(1 - \phi(t))}{t^2} \;\longrightarrow\; V - \langle F(r)^2\rangle \qquad (t \to 0).$$

The summands are non-negative, bounded by $a_r(n)^2$, and converge to $a_r(n)^2$. By Fatou's lemma $\sum \tilde\nu a_r^2 \le V - \langle F(r)^2\rangle < \infty$. Dominated convergence then gives equality. The bound $V(r) \le \|r\|^2_{2,\varepsilon}$ is Lemma 7.3. (c) $C_{\mu_0} = (-\Delta_\varepsilon + \mu_0^2)^{-1} \le \mu_0^{-2}$ as an operator. □

Lemma 13.2 (sampling and the lattice d^*)**.** *Let $\operatorname{supp} G \subset B_R$, $\ell_j > 2R + 2$, and let g_ε be the sample of $\star^{-1}G$ (Definition 2.9).*

(a) *If $G \in C^1_c$, then $\|d^* g_\varepsilon\|_{2,\varepsilon} \to \|\operatorname{curl} G\|_2$ as $\varepsilon \to 0$, uniformly in j.*

(b) *Let $\psi \in C^1_c(\mathbb{R}^3)$, $\operatorname{supp}\psi \subset B_{R-1}$, $G = \nabla\psi$, and let $\chi_\varepsilon(y) := \psi\big(y + \frac{\varepsilon}{2}(e_1 + e_2 + e_3)\big)$ be the sample of ψ at the centres of the elementary cubes, indexed by their lowest corner y. Define d on lattice 2-forms by $(d\sigma)(y) := \partial_1\sigma_{23}(y) - \partial_2\sigma_{13}(y) + \partial_3\sigma_{12}(y)$ (forward differences; $\sigma_{\mu\nu}(x)$ the value on the plaquette $(x;\mu,\nu)$), and d^* as its adjoint for $\langle\cdot,\,\cdot\rangle_\varepsilon$. Then $dd = 0$ on 1-forms, and $r_\varepsilon := g_\varepsilon + d^*\chi_\varepsilon$ satisfies $\|r_\varepsilon\|_{2,\varepsilon} \to 0$, uniformly in j.*

Proof. Write ∂_μ for the forward difference and $\partial^*_\mu = -\partial^-_\mu$ for its adjoint, minus the backward difference. Then $(dA)_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ (Section 2.2). (a) From $\langle g,\, dA\rangle_\varepsilon = \sum_{\mu<\nu}\langle g_{\mu\nu},\, \partial_\mu A_\nu - \partial_\nu A_\mu\rangle = \sum_{\mu,\nu}\langle g_{\mu\nu},\, \partial_\mu A_\nu\rangle$, with $g_{\nu\mu} := -g_{\mu\nu}$, we get $(d^*g)_\nu(x) = \sum_\mu \partial^*_\mu g_{\mu\nu}(x)$ on the bond (x,ν). The plaquettes $(x;\mu,\nu)$ and $(x - \varepsilon e_\mu;\mu,\nu)$ have their centres at the centre of the bond (x,ν) shifted by $\pm\frac{\varepsilon}{2}e_\mu$. So $(d^*g_\varepsilon)_\nu(x) = -\sum_\mu \varepsilon^{-1}\big(g_{\mu\nu}(z + \frac{\varepsilon}{2}e_\mu) - g_{\mu\nu}(z - \frac{\varepsilon}{2}e_\mu)\big)$, with z the bond centre. This is a centred difference quotient of a C^1_c function. By the mean value theorem and uniform continuity of ∇g it equals $(\delta\star^{-1}G)_\nu(z) + o(1) = (\operatorname{curl} G)_\nu(z) + o(1)$, uniformly in z. It vanishes unless $z \in B_{R+1}$. Riemann sums of the continuous compactly supported function $|\operatorname{curl} G|^2$ over the bond centres converge to its integral. (b) $d(dA) = \partial_1(\partial_2 A_3 - \partial_3 A_2) - \partial_2(\partial_1 A_3 - \partial_3 A_1) + \partial_3(\partial_1 A_2 - \partial_2 A_1) = 0$,

because forward differences commute on the torus. The adjoint is $(d^*\omega)_{23} = \partial_1^*\omega$, $(d^*\omega)_{13} = -\partial_2^*\omega$, $(d^*\omega)_{12} = \partial_3^*\omega$. The two cubes that contain the plaquette $(x;2,3)$ have lowest corners x and $x - \varepsilon e_1$, and their centres are the centre c_p of the plaquette shifted by $\pm\frac{\varepsilon}{2}e_1$; similarly in the other two directions. Hence $(d^*\chi_\varepsilon)_{23}(p) = -\varepsilon^{-1}\big(\psi(c_p + \frac{\varepsilon}{2}e_1) - \psi(c_p - \frac{\varepsilon}{2}e_1)\big) = -\partial_1\psi(c_p) + o(1)$. In the same way $(d^*\chi_\varepsilon)_{13} = \partial_2\psi + o(1)$ and $(d^*\chi_\varepsilon)_{12} = -\partial_3\psi + o(1)$, uniformly. The sample g_ε has $g_{23} = \partial_1\psi$, $g_{13} = -\partial_2\psi$, $g_{12} = \partial_3\psi$ at c_p. So $|r_\varepsilon(p)| \le \eta(\varepsilon) := \sup_{|y-y'|\le\varepsilon}|\nabla\psi(y) - \nabla\psi(y')| \to 0$. Also $r_\varepsilon(p) = 0$ unless $c_p \in B_R$. Hence $\|r_\varepsilon\|_{2,\varepsilon}^2 \le 3|B_{R+1}|\eta(\varepsilon)^2 \to 0$. □

13.2. **The Bianchi identity and the spectral density.**

Lemma 13.3 (Bianchi identity)**.** *Let S be a limit point and $\psi \in C_c^1(\mathbb{R}^3)$. Then every Schwinger function of S with $\star^{-1}\nabla\psi$ among its curvature arguments vanishes:*

$$S_{1+n,p}(\star^{-1}\nabla\psi, \mathbf{g}_1, \dots, \mathbf{g}_n; h_1, \dots, h_p) = 0.$$

In particular $\mathfrak{b}_S(\nabla\psi, \nabla\psi) = 0$, and for $\psi \in C_c^\infty$, $B(\nabla\psi) = 0$ almost surely under the law of S (Theorem 3.1(ii)); that is, $\operatorname{div} B = 0$, *or* $dF = 0$.

Proof. Let $Q := \prod_{i\le n} F(\mathbf{g}_{i,\varepsilon}) \prod_{k\le p} X_R(h_{k,\varepsilon})$ on the lattice, the test functions being sampled. By Lemma 13.2(b), $F(g_\varepsilon) = F(r_\varepsilon) - \langle\chi_\varepsilon, ddA\rangle_\varepsilon = F(r_\varepsilon)$, where g_ε is the sample of $\star^{-1}\nabla\psi$. By the Cauchy–Schwarz inequality and Lemma 13.1(b),

$$|\langle F(g_\varepsilon)\, Q\rangle_{\varepsilon,j,\mu_0}| \le \langle F(r_\varepsilon)^2\rangle^{1/2}\langle Q^2\rangle^{1/2} \le \|r_\varepsilon\|_{2,\varepsilon}\, \langle Q^2\rangle^{1/2}.$$

$\langle Q^2\rangle$ is a lattice Schwinger function of order $(2n, 2p)$, bounded uniformly by Proposition 8.1, and $\|r_\varepsilon\|_{2,\varepsilon} \to 0$. The left side converges to the Schwinger function in the statement along the subsequence that defines S (Theorem 3.1(ii); for S^0, first for S^μ, $\mu \in Q_0$, then as $\mu \downarrow 0$). So the limit is 0. For $\psi \in C_c^\infty$, $\mathbb{E}[B(\nabla\psi)^2] = \mathfrak{b}_S(\nabla\psi, \nabla\psi) = 0$ by the moment identity of Proposition 8.3. □

Lemma 13.4 (translation-invariant bounded forms)**.** *Let $\mathfrak{b}$ be a Hermitian form on $C_c(\mathbb{R}^3;\mathbb{C}^3)$ with $0 \le \mathfrak{b}(f,f) \le \|f\|_2^2$ and $\mathfrak{b}(\tau_a f, \tau_a f') = \mathfrak{b}(f,f')$ for all $a \in \mathbb{R}^3$. Then there is a measurable $m : \mathbb{R}^3 \to \{$Hermitian 3×3 matrices$\}$ with $0 \le m(p) \le \mathbf{1}$ for a.e. p and*

$$\mathfrak{b}(f,f') = (2\pi)^{-3}\int \overline{\hat f(p)}^{\mathsf T}\, m(p)\hat f'(p)\, dp \qquad (f, f' \in C_c(\mathbb{R}^3;\mathbb{C}^3)).$$

Moreover: (i) if $f_0 \in C_c$ and $\mathfrak{b}(f_0,f_0) = 0$, then $m\hat f_0 = 0$ a.e.; (ii) if a is a measurable Hermitian-matrix-valued function with $|a(p)| \le c(1+|p|^2)$ and $\mathfrak{b}(f,f) \le (2\pi)^{-3}\int \overline{\hat f}^{\mathsf T} a\hat f$ for all $f \in C_c^1(\mathbb{R}^3;\mathbb{C}^3)$, then $m \le a$ a.e.

Proof. By the Cauchy–Schwarz inequality for the positive form, $|\mathfrak{b}(f,f')| \le \|f\|_2\|f'\|_2$. So $\mathfrak{b}$ extends to L^2, and $\mathfrak{b}(f,f') = \langle f, Kf'\rangle$ for a bounded K with $0 \le K \le 1$ that commutes with all τ_a. Let $\mathcal{F} = (2\pi)^{-3/2}\hat{}$ be the unitary Fourier transform and $\hat K := \mathcal{F}K\mathcal{F}^{-1}$. Then $\hat K$ commutes with multiplication M_u by $u = e_a$, $e_a(p) := e^{-ia\cdot p}$. For fixed $\varphi, \chi \in L^2$, $u \mapsto \langle\varphi, [\hat K, M_u]\chi\rangle = \int u\,\big(\overline{\hat K\varphi}^{\mathsf T}\,\chi - \overline{\varphi}^{\mathsf T}\hat K\chi\big)\, dp$ is integration against an L^1 function. So the set of $u \in L^\infty$ with $[\hat K, M_u] = 0$ is a weak-$*$ closed subspace that contains every e_a. Its pre-annihilator in L^1 consists of $h \in L^1$ with $\hat h \equiv 0$, hence is $\{0\}$, so the subspace is all of L^∞. Let $w(p) := e^{-|p|^2}$ and $m_{\alpha\beta} := (\hat K(we_\beta))_\alpha / w$, with e_β the unit vectors of $\mathbb{C}^3$. For $u_\beta \in L^\infty$, $\hat K(w\sum_\beta u_\beta e_\beta) = w\sum_\beta u_\beta m_{\cdot\beta}$. So $\hat K f = mf$ on the set of $f = w\sum_\beta u_\beta e_\beta$, which is dense in L^2: for $f \in L^2$, $f\mathbf{1}\{|f| \le nw\}$ is of this form and converges to f. Testing with $f = wuv$, $u \in L^\infty$ scalar and $v \in \mathbb{C}^3$, gives $0 \le \int |u|^2 w^2\, \overline v^{\mathsf T} m v \le \int |u|^2 w^2 |v|^2$ for all u. Hence $0 \le \overline v^{\mathsf T} m(p) v \le |v|^2$ for a.e. p, simultaneously for all v in the countable set $(\mathbb{Q} + i\mathbb{Q})^3$, hence for all v. By polarisation $m(p)$ is Hermitian, so $0 \le m \le \mathbf{1}$ a.e. Then M_m is bounded and agrees with $\hat K$ on a dense set, so $\hat K = M_m$. This is the representation. (i) $\langle f_0, Kf_0\rangle = 0$ and $K \ge 0$ give $Kf_0 = 0$, that is, $m\hat f_0 = 0$ a.e. (ii) The form $q(f) := (2\pi)^{-3}\int \overline{\hat f}^{\mathsf T}(a-m)\hat f$ is continuous on

the Sobolev space $H^1(\mathbb{R}^3;\mathbb{C}^3)$, and $q \geq 0$ on the dense subspace C_c^∞, hence on H^1. Take $\hat f = uv$ with $u \in L^\infty$ of compact support and $v \in \mathbb{C}^3$; then $f \in H^1$, and $\int |u|^2\, \overline{v}^{\mathsf T}(a-m)v \geq 0$ for all such u. So $a - m \geq 0$ a.e., as before. □

Theorem 13.5 (the curvature spectral density)**.** (a) *For every limit point S there is a measurable function m_S with values in the Hermitian 3×3 matrices, with $0 \leq m_S(p) \leq P_T(p)$ for a.e. p, and*

$$\mathfrak{b}_S(G,G') = (2\pi)^{-3}\int \overline{\hat G(p)}^{\mathsf T}\, m_S(p)\, \hat G'(p)\, dp \qquad (G, G' \in C_c(\mathbb{R}^3;\mathbb{R}^3)).$$

In particular $0 \leq \mathfrak{b}_S(G,G) \leq \|\Pi_T G\|_2^2$: the curvature two-point function of every limit point is dominated, as a form, by that of the free massless photon.

(b) *If S is at photon mass $\mu_0 > 0$, then $m_S(p) \leq \min\big(1, |p|^2/\mu_0^2\big)\, P_T(p)$ for a.e. p.*

(c) *For $\mu' < \mu$ in Q_0, $m_{S^\mu} \leq m_{S^{\mu'}} \leq m_{S^0} \leq m_{S^\flat}$ a.e.; and $m_{S^{\mu_k}} \to m_{S^0}$ a.e. along every sequence $\mu_k \downarrow 0$ in Q_0.*

Proof. (a) For real $G, G' \in C_c$, $\mathfrak{b}_S(G,G')$ is real and symmetric. It is the limit of the lattice second moments $\langle F(g_\varepsilon)F(g'_\varepsilon)\rangle$ of the samples. By Lemma 13.1(b), $0 \leq \langle F(g_\varepsilon)^2\rangle \leq \|g_\varepsilon\|_{2,\varepsilon}^2$, and $\|g_\varepsilon\|_{2,\varepsilon} \to \|G\|_2$ (Proposition 8.2, step (iii)). Hence $0 \leq \mathfrak{b}_S(G,G) \leq \|G\|_2^2$. For S^0 this passes through the limit $\mu \downarrow 0$ (Theorem 3.1(ii)). Extend $\mathfrak{b}_S$ to complex $f = G_1 + iG_2$ as a Hermitian form, $\mathfrak{b}_S(f,f') := \mathfrak{b}_S(G_1,G_1') + \mathfrak{b}_S(G_2,G_2') + i\big(\mathfrak{b}_S(G_1,G_2') - \mathfrak{b}_S(G_2,G_1')\big)$. Then $\mathfrak{b}_S(f,f) = \mathfrak{b}_S(G_1,G_1) + \mathfrak{b}_S(G_2,G_2) \in [0, \|f\|_2^2]$. It is translation invariant by Theorem 9.2(a). Lemma 13.4 gives m_S with $0 \leq m_S \leq \mathbf{1}$. For $\psi \in C_c^\infty$ real, $\mathfrak{b}_S(\nabla\psi, \nabla\psi) = 0$ by Lemma 13.3, and $\widehat{\nabla\psi}(p) = ip\hat\psi(p)$. By Lemma 13.4(i), $m_S(p)p\,\hat\psi(p) = 0$ a.e. For $\psi \neq 0$, $\hat\psi$ is the restriction of an entire function, so its zero set is Lebesgue-null. Hence $m_S(p)p = 0$ a.e. A positive semidefinite Hermitian matrix that annihilates p satisfies $m = P_T m P_T$. So $m_S \leq P_T\mathbf{1}P_T = P_T$. The last claim is $\langle \hat G,\, m_S\hat G\rangle \leq \langle \hat G,\, P_T\hat G\rangle$, integrated. (b) Let $G \in C_c^1$ be real. By Lemma 13.1(b),(c), $\langle F(g_\varepsilon)^2\rangle \leq \mu_0^{-2}\|d^* g_\varepsilon\|_{2,\varepsilon}^2$, and by Lemma 13.2(a) the right side tends to $\mu_0^{-2}\|\operatorname{curl} G\|_2^2 = (2\pi)^{-3}\mu_0^{-2}\int |p|^2\, \overline{\hat G}^{\mathsf T} P_T \hat G\, dp$, since $|p\times v|^2 = |p|^2\, \overline{v}^{\mathsf T} P_T(p) v$ for $v \in \mathbb{C}^3$. Both sides are Hermitian forms induced by real symmetric forms on real fields, so the inequality extends to complex $f = G_1 + iG_2$ by adding the inequalities for G_1 and G_2. Lemma 13.4(ii) with $a = \mu_0^{-2}|p|^2 P_T$ gives $m_S \leq \mu_0^{-2}|p|^2 P_T$. Since $m_S = P_T m_S P_T$ by (a), and m_S is bounded on $\operatorname{ran} P_T$ both by 1 and by $|p|^2/\mu_0^2$, we get $m_S \leq \min(1, |p|^2/\mu_0^2) P_T$. (c) For real $\mathbf{g}$ and real t, $e^{tF(\mathbf{g})} = \exp[F(t\mathbf{g}) - X_R(0)]$ lies in the class $\mathcal{D}$ ($p = 0$). Theorem 3.1(iv) gives $\phi_\mu(t) \leq \phi_{\mu'}(t) \leq \phi_\flat(t)$ and $\phi_0(t) \leq \phi_\flat(t)$, where $\phi_S(t) := G^S(t)$ is the generating functional of the one-element family $\{\mathbf{g}\}$. Each ϕ_S is entire, $\phi_S(0) = 1$, and ϕ_S is even because its odd Taylor coefficients $S_{n,0}$ vanish (Theorem 9.8). If $f_1 \leq f_2$ are smooth with $f_1(0) = f_2(0)$, then 0 is a maximum of $f_1 - f_2$, so $f_1''(0) \leq f_2''(0)$. Hence $S_{2,0}^\mu(\mathbf{g},\mathbf{g}) \leq S_{2,0}^{\mu'}(\mathbf{g},\mathbf{g}) \leq S_{2,0}^\flat(\mathbf{g},\mathbf{g})$ and $S_{2,0}^0(\mathbf{g},\mathbf{g}) \leq S_{2,0}^\flat(\mathbf{g},\mathbf{g})$. Also $S_{2,0}^0 = \lim_{\mu\downarrow 0} S_{2,0}^\mu$ (Theorem 3.1(ii)), so $S_{2,0}^{\mu'}(\mathbf{g},\mathbf{g}) \leq S_{2,0}^0(\mathbf{g},\mathbf{g})$. Extend to complex fields as in (b) and apply Lemma 13.4(ii) with the bounded functions $a = m_{S^{\mu'}}$, m_{S^0}, $m_{S^\flat}$. This gives the a.e. inequalities. Along $\mu_k \downarrow 0$ the matrices $m_{S^{\mu_k}}(p)$ increase and are bounded by $\mathbf{1}$, off a null set, so they converge a.e. to some $m_* \leq m_{S^0}$. Let $f = G_1 + iG_2 \in C_c(\mathbb{R}^3;\mathbb{C}^3)$ with G_1, G_2 real. By the representation of Lemma 13.4 for the Hermitian extension, $(2\pi)^{-3}\int \overline{\hat f}^{\mathsf T} m_{S^{\mu_k}} \hat f = \mathfrak{b}_{S^{\mu_k}}(G_1,G_1) + \mathfrak{b}_{S^{\mu_k}}(G_2,G_2)$, which tends to $\mathfrak{b}_{S^0}(G_1,G_1) + \mathfrak{b}_{S^0}(G_2,G_2) = \mathfrak{b}_{S^0}(f,f)$ (Theorem 3.1(ii)). The integrands are bounded by $|\hat f|^2$, so by dominated convergence the left side tends to $(2\pi)^{-3}\int \overline{\hat f}^{\mathsf T} m_* \hat f$. Hence $\mathfrak{b}_{S^0}(f,f) = (2\pi)^{-3}\int \overline{\hat f}^{\mathsf T} m_* \hat f$ for all $f \in C_c(\mathbb{R}^3;\mathbb{C}^3)$. Lemma 13.4(ii), applied to $\mathfrak{b}_{S^0}$ with the bounded function $a = m_*$, gives $m_{S^0} \leq m_*$ a.e. Hence $m_{S^0} = m_*$ a.e. □

13.3. **Long-distance strength of the curvature.**

Definition 13.6 (massless curvature sector)**.** For a limit point S and a real $G \in C_c(\mathbb{R}^3;\mathbb{R}^3)$ put

$$\mathsf{Z}_S(G) := \limsup_{R\to\infty} R^3\, \mathfrak{b}_S(G_R, G_R) \ \in [0,\infty).$$

S is *massless in the curvature sector* if $\mathsf{Z}_S(G) > 0$ for some G.

Since $R^3\|G_R\|_2^2 = \|G\|_2^2$, $\mathsf{Z}_S(G)/\|G\|_2^2$ is the asymptotic fraction of the L^2 size of a test field at scale R that the two-point function retains. For the free massless photon $R^3\|\Pi_T G_R\|_2^2 = \|\Pi_T G\|_2^2$ for every R, because P_T is homogeneous of degree 0. For the curvature of a free vector field of mass $M > 0$, $m(p) = |p|^2(|p|^2+M^2)^{-1}P_T(p)$, and $\mathsf{Z} \equiv 0$ by dominated convergence. If $m_S(p) - z\,P_T(p) \to 0$ as $p \to 0$ outside a Lebesgue null set (an essential limit; m_S is defined only almost everywhere) for a number z, then $\mathsf{Z}_S(G) = z\|\Pi_T G\|_2^2$. So Z_S is the Euclidean form of the field-strength renormalisation of the photon at zero momentum. The definition is the strong form of masslessness: a density $m(p) = \min(1,|p|^\eta)P_T(p)$, $0 < \eta < 1$, has power-law correlations and $\mathsf{Z} \equiv 0$. The converse of Corollary 13.9(c), from the absence of a gap to $\mathsf{Z}_S > 0$, is not claimed.

Theorem 13.7 (the long-distance dichotomy)**.** *Let S be a limit point and $G \in C_c(\mathbb{R}^3;\mathbb{R}^3)$ real.*

(i) *For every $R > 0$,*

$$R^3\, \mathfrak{b}_S(G_R, G_R) = (2\pi)^{-3} \int \overline{\hat G(q)}^{\mathsf{T}}\, m_S(q/R)\, \hat G(q)\, dq \ \le\ \|\Pi_T G\|_2^2.$$

Hence $0 \le \mathsf{Z}_S(G) \le \|\Pi_T G\|_2^2$.

(ii) *(no massless component at positive photon mass) If S is at photon mass $\mu_0 > 0$, then $\lim_{R\to\infty} R^3\mathfrak{b}_S(G_R,G_R) = 0$, so $\mathsf{Z}_S \equiv 0$. For $G \in C_c^1$, quantitatively, $R^3\mathfrak{b}_S(G_R,G_R) \le \mu_0^{-2}R^{-2}\|\operatorname{curl} G\|_2^2$.*

(iii) *(order) For $\mu \in Q_0$, $0 = \mathsf{Z}_{S^\mu}(G) \le \mathsf{Z}_{S^0}(G) \le \mathsf{Z}_{S^\flat}(G)$. In particular, if S^0 is massless in the curvature sector, so is $S^\flat$.*

(iv) *(integrable correlations exclude masslessness) Suppose that for some real $\psi \in C_c(\mathbb{R}^3)$ with $\int\psi \neq 0$ and all $\alpha,\beta \in \{1,2,3\}$, the functions $a \mapsto \mathfrak{b}_S(\psi e_\alpha, \tau_a\psi e_\beta)$ are in $L^1(\mathbb{R}^3)$. Then $\lim_{R\to\infty} R^3\mathfrak{b}_S(G_R,G_R) = 0$ for every G, so $\mathsf{Z}_S \equiv 0$. This holds in particular if $|\mathfrak{b}_S(\psi e_\alpha,\tau_a\psi e_\beta)| \le Ce^{-M|a|}$ for some $M > 0$.*

Proof. (i) Insert $\widehat{G_R}(p) = \hat G(Rp)$ into Theorem 13.5(a) and substitute $q = Rp$. The bound is $m_S \le P_T$ and $P_T(q/R) = P_T(q)$. (ii) By Theorem 13.5(b), the integrand in (i) is at most $\min(1, |q|^2/(R\mu_0)^2)\,|\hat G(q)|^2$, which tends to 0 pointwise as $R\to\infty$ and is dominated by $|\hat G|^2 \in L^1$. For $G \in C_c^1$, use the bound $|q|^2/(R\mu_0)^2$ and $(2\pi)^{-3}\int |q|^2 \overline{\hat G}^{\mathsf{T}} P_T \hat G = \|\operatorname{curl} G\|_2^2$. (iii) Theorem 13.5(c) and (i), together with (ii) for S^μ. (iv) Since $\widehat{\tau_a\psi} = e^{-ia\cdot p}\hat\psi$,

$$\phi_{\alpha\beta}(a) := \mathfrak{b}_S(\psi e_\alpha, \tau_a\psi e_\beta) = \int e^{-ia\cdot p} h_{\alpha\beta}(p)\, dp,$$

with $h_{\alpha\beta} := (2\pi)^{-3}|\hat\psi|^2 (m_S)_{\alpha\beta} \in L^1$. If $\phi_{\alpha\beta} \in L^1$, the Fourier inversion theorem shows that $h_{\alpha\beta}$ agrees a.e. with a continuous function. Since $\hat\psi(0) = \int\psi \neq 0$ and $\hat\psi$ is continuous, $|\hat\psi| \ge c > 0$ on a ball B_ρ. So on B_ρ, m_S agrees a.e. with a continuous matrix function $\tilde m$. By Theorem 13.5(a), $m_S(p)p = 0$ a.e.; continuity gives $\tilde m(p)p = 0$ for all $p \in B_\rho$. For a unit vector n and $0 < t < \rho$, $\tilde m(tn)n = 0$, and $t \downarrow 0$ gives $\tilde m(0)n = 0$. Hence $\tilde m(0) = 0$. In (i), split the integral at $|q| = \rho R$. On $\{|q| < \rho R\}$ the integrand equals $\overline{\hat G}^{\mathsf{T}}\, \tilde m(q/R)\hat G$ a.e.; it tends to $\overline{\hat G}^{\mathsf{T}}\, \tilde m(0)\hat G = 0$ pointwise and is dominated by $|\hat G|^2$. On $\{|q| \ge \rho R\}$ it is at most $|\hat G|^2$, whose integral there tends to 0. Exponential decay implies integrability. □

13.4. **The curvature sector of the reconstruction.** Fix a coordinate direction j. Let $\mathcal{V}_+^{(j)}$ be the complex 2-forms in C_c supported in $\{x_j > 0\}$, with $F(g_1 + ig_2) := F(g_1) + iF(g_2)$ and $S(F(u)F(v))$ the complex-bilinear extension of $S_{2,0}$. Let $\rho_j \in O_h$ be the reflection in $\{x_j = 0\}$, ρ_j^* its signed action on 2-forms (Theorem 9.4), and θ_j the antilinear reflection $\theta_j u := \overline{\rho_j^* u}$. Put

$$\langle u,\, v\rangle_+ := S\big(F(\theta_j u)F(v)\big) \qquad (u, v \in \mathcal{V}_+^{(j)}).$$

Proposition 13.8 (no gap without integrable correlations)**.** *Let S be a limit point.*

(a) *$\langle\cdot,\,\cdot\rangle_+$ is a positive semidefinite Hermitian form on $\mathcal{V}_+^{(j)}$ with $\langle u,\, u\rangle_+ \le \|u\|_2^2$. Let $\mathcal{H}_F^{(j)}$ be the completion of the quotient by its null space, and $[u]$ the class of u. For $t \ge 0$, $T(t)[u] := [\tau_{te_j} u]$ is a well-defined strongly continuous semigroup of self-adjoint contractions, so $T(t) = e^{-tH_F}$ with $H_F \ge 0$ self-adjoint. For $y \perp e_j$, $V(y)[u] := [\tau_y u]$ is a strongly continuous unitary representation commuting with T. For $\kappa \in O_h$ with $\kappa e_j = e_k$, $[u] \mapsto [\kappa_* u]$ is unitary from $\mathcal{H}_F^{(j)}$ onto $\mathcal{H}_F^{(k)}$ and intertwines the two semigroups.*

(b) *Suppose $H_F \ge M$ for some $M > 0$, in one coordinate direction (hence, by (a), in all three). Then for all real 2-forms $\mathbf{g}, \mathbf{g}' \in C_c$ supported in $\bar{B}_r$ and all $a \in \mathbb{R}^3$,*

$$|S_{2,0}(\mathbf{g}, \tau_a \mathbf{g}')| \le e^{2Mr}\, e^{-M|a|/\sqrt{3}}\, \|\mathbf{g}\|_2 \|\mathbf{g}'\|_2.$$

Consequently the hypothesis of Theorem 13.7(iv) holds, and $\mathsf{Z}_S \equiv 0$.

Proof. (a) For $u = \sum_i c_i \mathbf{g}_i$ with real $\mathbf{g}_i$, $\langle u,\, u\rangle_+ = S(\theta(P)P)$ with $P = \sum_i c_i F(\mathbf{g}_i)$. This is ≥ 0 by Theorem 9.4(b). The form is Hermitian because S is real, symmetric and invariant under the reflection $\rho_j \in O_h$ (Theorem 9.2(b)). Writing $\mathfrak{b}(u, v) := S(F(\bar{u})F(v))$, Theorem 13.5(a) gives $|\mathfrak{b}(u, v)| \le \|u\|_2 \|v\|_2$, hence $|\langle u,\, v\rangle_+| = |\mathfrak{b}(\rho_j^* u, v)| \le \|u\|_2 \|v\|_2$. For $t \ge 0$, τ_{te_j} maps $\mathcal{V}_+^{(j)}$ into itself. Since $\theta_j \tau_{te_j} = \tau_{-te_j} \theta_j$ and S is translation invariant, $\langle u,\, \tau_{te_j} v\rangle_+ = \langle \tau_{te_j} u,\, v\rangle_+$. Let $\|u\|_+^2 := \langle u,\, u\rangle_+$. The Cauchy–Schwarz inequality gives $\|\tau_{te_j} u\|_+^2 = \langle u,\, \tau_{2te_j} u\rangle_+ \le \|u\|_+ \|\tau_{2te_j} u\|_+$. Iterating, and using $\|\tau_s u\|_+ \le \|\tau_s u\|_2 = \|u\|_2$,

$$\|\tau_{te_j} u\|_+ \le \|u\|_+^{1-2^{-n}}\, \|\tau_{2^n te_j} u\|_+^{2^{-n}} \le \|u\|_+^{1-2^{-n}}\, \|u\|_2^{2^{-n}}.$$

Letting $n \to \infty$ gives $\|\tau_{te_j} u\|_+ \le \|u\|_+$. So $T(t)$ preserves the null space and defines a self-adjoint contraction, with $T(t)T(s) = T(t+s)$. The function $t \mapsto \langle u,\, T(t)v\rangle_+ = \mathfrak{b}(\overline{\theta_j u}, \tau_{te_j} v)$ is continuous, because translations act continuously on L^2 and $\mathfrak{b}$ is bounded. A uniformly bounded semigroup that is weakly continuous on a dense set is weakly continuous, and a weakly continuous contraction semigroup on a Hilbert space is strongly continuous. The spectral theorem gives H_F. The transverse translations commute with θ_j, so they preserve the form; they are invertible, hence unitary, and strongly continuous by the same argument. For $\kappa \in O_h$ with $\kappa e_j = e_k$: κ maps $\{x_j > 0\}$ onto $\{x_k > 0\}$, $\kappa \rho_j \kappa^{-1} = \rho_k$ and $\kappa \tau_{te_j} \kappa^{-1} = \tau_{te_k}$. With the O_h-invariance of S (Theorem 9.2(b)) this gives $\langle \kappa_* u,\, \kappa_* v\rangle_+ = \langle u,\, v\rangle_+$, the left side formed in direction k and the right side in direction j, and the intertwining. (b) By (a) we may take the direction j with $|a_j| = \max_i |a_i| \ge |a|/\sqrt{3}$, and by the symmetry $S_{2,0}(\mathbf{g}, \tau_a \mathbf{g}') = S_{2,0}(\mathbf{g}', \tau_{-a} \mathbf{g})$ we may take $a_j > 0$. If $a_j \le 2r$, the bound follows from $|S_{2,0}(\mathbf{g}, \tau_a \mathbf{g}')| \le \|\mathbf{g}\|_2 \|\mathbf{g}'\|_2$ and $e^{2Mr} e^{-Ma_j} \ge 1$. Let $a_j > 2r$, fix $\eta \in (0, a_j/2 - r)$, and put $c := a_j/2$ and $s := c - r - \eta > 0$. By translation invariance $S_{2,0}(\mathbf{g}, \tau_a \mathbf{g}') = S_{2,0}(\tau_{-ce_j} \mathbf{g}, \tau_{a-ce_j} \mathbf{g}')$. Here $\tau_{-ce_j} \mathbf{g}$ is supported in $\{x_j \le -s - \eta\}$ and $\tau_{a-ce_j} \mathbf{g}'$ in $\{x_j \ge s + \eta\}$, so the forms u, v below are supported in $\{x_j \ge \eta\}$ and lie in $\mathcal{V}_+^{(j)}$. Write $\tau_{-ce_j} \mathbf{g} = \theta_j \tau_{se_j} u$ and $\tau_{a-ce_j} \mathbf{g}' = \tau_{a_\perp} \tau_{se_j} v$, where $a_\perp := a - a_j e_j$ and $u, v \in \mathcal{V}_+^{(j)}$ are real, $\|u\|_2 = \|\mathbf{g}\|_2$, $\|v\|_2 = \|\mathbf{g}'\|_2$. Using $\theta_j \tau_{se_j} = \tau_{-se_j} \theta_j$ and translation invariance,

$$S_{2,0}(\mathbf{g}, \tau_a \mathbf{g}') = S\big(F(\theta_j u) F(\tau_{2se_j} \tau_{a_\perp} v)\big) = \langle [u],\, T(2s) V(a_\perp)[v]\rangle_{\mathcal{H}_F^{(j)}}.$$

With $\|T(2s)\| \le e^{-2sM}$, $\|[u]\| \le \|u\|_2$ and $2s = a_j - 2r - 2\eta \ge |a|/\sqrt{3} - 2r - 2\eta$, we get $|S_{2,0}(\mathbf{g}, \tau_a \mathbf{g}')| \le e^{2M(r+\eta)} e^{-M|a|/\sqrt{3}} \|\mathbf{g}\|_2 \|\mathbf{g}'\|_2$; let $\eta \downarrow 0$. It is integrable in a, and Theorem 13.7(iv) applies with any real $\psi \in C_c$ with $\int \psi \neq 0$: take $\mathbf{g} = \star^{-1}(\psi e_\alpha)$ and $\mathbf{g}' = \star^{-1}(\psi e_\beta)$. □

Under the hypotheses of Theorem 11.1, with $\mathcal{H}$, H and Ω as there, $[\mathbf{g}] \mapsto [F(\mathbf{g})]$ is an isometry of $\mathcal{H}_F^{(3)}$ onto the closed span of the vectors $[F(\mathbf{g})]$ in $\mathcal{H}$, and it intertwines e^{-tH_F} with e^{-tH}. So $H_F \ge M$ says that H has a gap M on the curvature subspace, which is orthogonal to Ω because $S_{1,0} = 0$. Part (a) of the proposition is the Osterwalder–Schrader construction [27, 28] restricted to the linear curvature fields. Of OS2 the proposition uses only the case of polynomials of degree one; it also uses OS1, OS3 and Theorem 13.5(a), but not the hypotheses (H1)–(H3) of Section 10.1 and not Theorem 11.1.

Corollary 13.9 (what masslessness would give)**.** *Let $S \in \{S^\flat, S^0\}$ be a limit state of Theorem 3.1 or, under its hypothesis, of Theorem 3.5. If S is massless in the curvature sector, then:*

(a) *for every real $\psi \in C_c$ with $\int \psi \neq 0$, some $a \mapsto \mathfrak{b}_S(\psi e_\alpha, \tau_a \psi e_\beta)$ is not integrable on $\mathbb{R}^3$. In particular this bounded function is not $O(|a|^{-3-\delta})$ for any $\delta > 0$, and does not decay exponentially;*

(b) *there are $R_k \to \infty$ with $\mathfrak{b}_S(G_{R_k}, G_{R_k}) \ge \frac{1}{2} \mathsf{Z}_S(G)\, R_k^{-3}$, a power-law lower bound for the variance of the block averages $B(G_R)$. A state with integrable curvature correlations has $R^3 \mathfrak{b}(G_R, G_R) \to 0$ instead (Theorem 13.7(iv));*

(c) *in every coordinate direction, $\inf \operatorname{spec} H_F = 0$: the curvature sector of the reconstructed theory has no mass gap.*

If S^0 is massless, then so is $S^\flat$ (Theorem 13.7(iii)). The states at positive photon mass are not massless in the curvature sector (Theorem 13.7(ii)).

Proof. (a) is the contrapositive of Theorem 13.7(iv), and (c) that of Proposition 13.8(b): no $M > 0$ has $H_F \ge M$; in particular $\mathcal{H}_F^{(j)} \neq \{0\}$, since otherwise $H_F \ge M$ would hold vacuously, and as $H_F \ge 0$, $\inf \operatorname{spec} H_F = 0$. (b) is the definition of the lim sup. □

Remark 13.10 (what these properties cannot give). The states S^{μ_0} at positive photon mass satisfy every conclusion of Theorem 3.1(ii)–(iii), the Bianchi identity (Lemma 13.3) and the domination $m_S \le P_T$ (Theorem 13.5(a)), and yet $\mathsf{Z}_{S^{\mu_0}} \equiv 0$ (Theorem 13.7(ii)). So no argument that uses only these properties of $S^\flat$ or S^0 proves that they are massless in the curvature sector. What distinguishes S^0 and $S^\flat$ from S^{μ_0} in this section is the weaker upper bound $m_S \le P_T$ in place of $m_S \le |p|^2 \mu_0^{-2} P_T$, and the comparison $m_{S^\mu} \le m_{S^0} \le m_{S^\flat}$ of Theorem 13.5(c), whose lower side is a state at positive photon mass. Section 14.4 discusses the lower bound that would be needed.

14. Open problems

We list the questions that this paper leaves open and, for each, the step at which its methods stop. Where a result of this paper depends on one of them, the dependence is an explicit hypothesis of that result (Theorems 3.5, 3.10(b),(c) and 12.9).

14.1. **Rotations.** The limit states are invariant under $\mathbb{R}^3 \rtimes O_h$ (Theorem 3.1(iii)); rotations outside O_h are not symmetries of any regularised state and would have to emerge in the limit. Three facts are known. A rotation R with $2^j R$ integral for some $j \ge 0$ lies in O_h, so no rotation outside O_h maps the dyadic lattices of Theorem 3.1 into one another (Lemma 9.10(i)). The rotation

$$R_3 = \tfrac{1}{3} \begin{pmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ -2 & 2 & -1 \end{pmatrix}$$

has $3R_3$ integral and infinite order (Lemma 9.10(ii)). The closed subgroup of O(3) generated by O_h and one rotation of infinite order is O(3), so a limit state invariant under one such rotation is O(3)-invariant (Lemma 9.13).

Invariance under a rotation R of infinite order splits into two steps. (i) *An ultraviolet comparison:* on a fixed torus, at a fixed photon mass $\mu_0 > 0$, the continuum limits of the model on $\varepsilon\mathbb{Z}^3$ and of the model on the rotated lattice $R\varepsilon\mathbb{Z}^3$ agree, when the same continuum 2-form test function is paired with each lattice in its own frame. King [25, Thm. 2.4, pp. 326–327] states a comparison of this kind for the rotation by $\arctan(3/4)$ in a coordinate plane, at blocking factor 5, with one scalar test function smeared against the plaquettes of both lattices; we do not use it. (ii) *An infrared statement.* By the isometry $x \mapsto R^{-1}x$, the rotated lattice on a cubic torus T is the axis lattice on the tilted torus $R^{-1}T$, with the same action; so (i) says that the continuum theory on the cubic torus, rotated by R, is the continuum theory on the tilted torus. Given (i), and with the limits taken in King's order, first $\varepsilon \to 0$ on each fixed torus and then the torus to $\mathbb{R}^3$, invariance under R of the limit along cubic tori becomes the statement

the infinite-volume limits along cubic tori and along tilted tori agree.

This statement would follow from uniqueness of the infinite-volume limit, but it is weaker. It cannot be arranged by the choice of tori, since no lattice is invariant under a rotation of infinite order (Lemma 9.10(iii)). Nor does it follow from invariance under translations and O_h, reflection positivity and clustering alone: the centred Gaussian scalar field with covariance

$$\widehat{G}(k) = \frac{1}{3}\sum_{\nu=1}^{3} \frac{1}{|k|^2 + ck_\nu^2 + m^2}, \qquad c > 0,$$

is invariant under translations and O_h, reflection positive about the coordinate planes (each summand is the covariance of a free field with a diagonal metric), and clusters exponentially, but it is not invariant under any rotation of infinite order. Otherwise, by Lemma 9.13(ii) and the continuity of $\widehat{G}$, it would be O(3)-invariant; but with $a := 1 + m^2$,

$$3\Big(\widehat{G}(e_1) - \widehat{G}\big(2^{-1/2}(e_1+e_2)\big)\Big) = \frac{1}{a} + \frac{1}{a+c} - \frac{2}{a+c/2} > 0$$

by strict convexity of $x \mapsto 1/x$.

For R_3 the natural family has blocking factor 3: R_3 maps $3\varepsilon\mathbb{Z}^3$ into $\varepsilon\mathbb{Z}^3$ (Lemma 9.10(ii)), so the rotated lattices stay in the family $\varepsilon = 3^{-K}\varepsilon_0$. Theorem 3.5 gives parts (ii)–(iv) of Theorem 3.1 for that family under the hypothesis of a uniform tube bound; a proof of the hypothesis would need the analogues for blocking factor 3 of Theorem 5.2 and of Lemma A.5. Neither step (i) nor step (ii) is carried out in this paper. Invariance under rotations is the only input that would still be missing for the axioms on the ergodic components: by Corollary 10.44, if a limit state S is invariant under O(3), then almost every ergodic component of its law satisfies OS0–OS4, with OS0 in the form of Remark 3.7.

14.2. **Uniqueness, the two massless states, and King's cube states.** The limits of Theorem 3.1 are taken along subsequences, and different subsequences might give different states; no uniqueness statement is proved. There are two candidates for the theory with a massless photon: S^0, the limit $\mu_0 \downarrow 0$ of the infinite-volume states at positive photon mass, and $S^\flat$, the infinite-volume limit of the states in which the constant mode of the gauge field is Haar-distributed on its period cell; at fixed lattice and torus the latter is the limit $\mu_0 \downarrow 0$ of the states at positive photon mass on the gauge-invariant observables (Lemma 2.8). Both satisfy OS0–OS3 with cubic symmetry, and $S^0 \le S^\flat$ on $E^{\mathcal{D}}$ (Theorem 3.1(iv)). Whether they coincide is the question whether $\mu_0 \downarrow 0$ commutes with the limits $\varepsilon \to 0$, $T \to \mathbb{R}^3$. By Proposition 8.10(b) the equality on $E^{\mathcal{D}}$ would follow from the condition (U-μ)$_T$ there, that the torus expectations at photon mass μ_0 converge to those of the state $\flat$ as $\mu_0 \downarrow 0$, uniformly in the spacing and the torus, on the class $\mathcal{D}$; this uniformity is

not known at $e \neq 0$. On tori there is no monotonicity in the volume, on which the corresponding argument for cubes would rely.

King's states [24, 25] are limits in cubes with Neumann conditions for the gauge field and Dirichlet conditions for the scalar field. The correlation inequalities of [8] suggest one-sided comparisons between cube limits and torus limits, but such comparisons would need, besides the existence of King's limits, a decay condition on the difference between the torus coefficient b_T of Definition 2.2 and King's free-boundary coefficient $a_\natural$, of the type $\sup_K |b_T - a_\natural|\,|T| \to 0$ as $\ell_T \to \infty$; we do not prove them. The identification of the torus limits with King's cube limits, and the cubic symmetry of the latter, would follow from uniqueness of the infinite-volume state. On tori the cubic symmetry is exact at every lattice spacing, which is one reason this paper works with tori. The cluster expansion that gives uniqueness when there is a mass gap is not available in the Coulomb phase.

14.3. **Non-triviality.** In every limit state the truncated functions have the signs of Theorem 12.4, and on every lattice torus the leading coefficient of $M(g;h) = S^T_{2,1}(g,g;h)$ is strictly negative (Proposition 12.6). Theorem 12.9 derives $M(g;h) < 0$ in the limit states at small $e^2 + \lambda$, so that they are not decoupled, from two hypotheses: the one-loop statement (C) (Hypothesis 12.7), a statement about a weighted closed-walk sum and its Brownian-bridge limit, and the remainder bound (R). The step that fails is (R). It would follow from a bound

$$\big|\ln Z(T,g,h) - \ln Z(T,0,0) - \mathcal{P}_2(g,h)\big| \leq e^2 \rho\big((e^2+\lambda)/m\big)\,\kappa(g,h),$$

uniform in ε, T and $\mu_0 \in (0,\mu_*] \cup \{\flat\}$, for sources of the size needed to extract $S^T_{2,1}$ by Cauchy estimates; here $Z(T,g,h)$ is the lattice partition function with the sources g for the curvature and h for the Wick square, and $\mathcal{P}_2$ is its explicit one-loop part, whose coefficient of order $(2,1)$ in the sources is $e^2 c^{\mathrm{free}}_{\varepsilon,j,\mu_0}(g,h)$. This is a renormalisation-group remainder estimate for source-dependent effective actions, with every remainder controlled in terms of the couplings at the unit scale. Neither the uniform bounds of this paper nor [1, 2, 3, 4, 24] provide it: Bałaban's papers give two-sided bounds on partition functions, and [24, Thm. 2.1, p. 654] gives $|\ln Z(T,g,h)| \leq C|T|$ with C depending on g, h and $|T|$, without asymptotics in (e,λ). The correlation inequalities of [8] give only non-strict signs, and we know of no inequality of Griffiths–Hurst–Sherman type for the even variable $|\varphi(x)|^2$. No result of this paper excludes a Gaussian limit state: such a state would have $M \equiv 0$, which is compatible with every sign in Theorem 12.4.

14.4. **The massless photon.** Above the first-order surface of Section 14.6 the photon is expected to stay massless in $S^\flat$ and S^0: the curvature two-point function should decay like $|x|^{-3}$, and the Hamiltonian of Theorem 11.1 should have no gap above the joint vacuum. In the notation of Section 13 this is $\mathsf{Z}_S > 0$ (Definition 13.6), in the uniform form

$$\text{(NS)} \qquad \mathfrak{b}_S(G_R, G_R) \geq \delta R^{-3} \quad \text{for all } R \geq R_0,$$

for some real $G \in C_c(\mathbb{R}^3;\mathbb{R}^3)$, $\delta > 0$ and R_0. Even $\mathfrak{b}_S \not\equiv 0$ is not known. What is proved is one-sided (Theorem 13.7, Corollary 13.9): the curvature two-point function of every limit state is dominated by that of the free massless photon, the states at positive photon mass have no massless component, and masslessness would exclude integrable curvature correlations and a mass gap in the curvature sector. Every estimate of this paper on the curvature is an upper bound, or a comparison with the states at positive photon mass, which have every property used in Section 13 and $\mathsf{Z} \equiv 0$ (Theorem 13.7(ii)); so these properties alone cannot prove masslessness. What is needed is a lower bound on the curvature two-point function: at long wavelengths and uniformly in the scale for $\mathsf{Z}_S > 0$, or at one fixed scale, uniformly in the spacing and the torus, for $\mathfrak{b}_S \not\equiv 0$. By Lemma 13.1(b) such a bound says that the closed currents of the matter do not completely screen long-wavelength magnetic fields. For S^0, and with it $S^\flat$, a bound $R^3 \mathfrak{b}_{S^\mu}(G_R, G_R) \geq \delta$ for $R_0 \leq R \leq c/\mu$, uniform in $\mu \in Q_0$, would suffice (Theorem 13.5(c)); by the same comparison, $\mathfrak{b}_{S^\mu} \not\equiv 0$ for one $\mu \in Q_0$ would give $\mathfrak{b}_{S^0} \not\equiv 0$ and $\mathfrak{b}_{S^\flat} \not\equiv 0$, and this is not known either. At small $e^2 + \lambda$, $\mathfrak{b}_S \not\equiv 0$ would also follow

from (R) and (C): if $\mathfrak{b}_S(G,G)=0$, then $F(g)=0$ almost surely for $g=\star^{-1}G$, so $M(g;h)=0$ for every h, which Theorem 12.9 excludes when g has components in C_c^∞, $d^*g\not\equiv 0$, $0\le h\not\equiv 0$ and $e^2+\lambda$ is small enough, depending on (g,h).

14.5. **The rest of OS4.** Theorem 3.6 gives, on almost every ergodic component of the law of a limit state, ergodicity under $\mathbb{R}^3$, reflection positivity, clustering in the mean and under diagonal separation, and a unique joint vacuum in each coordinate direction. Theorem 3.10 gives clustering along the axes for every observable linear in the curvature, with any square-integrable dependence on the Wick square; for the Wick-square sector if $\Delta_S=0$; and for all observables, with OS4, under hypothesis (D). OS4 itself (Definition 2.11) is not proved, for the limit states or for their components. For a component it is the statement that $\operatorname{Ker} H_j=\mathbb{C}\Omega_j$ for $j=1,2,3$ in the reconstruction of Proposition 10.16 (by Proposition 10.17(i) and translation invariance), and by Proposition 3.9 it suffices to show $P_jG=\mathbb{E}[G\mid\mathcal{I}]$ for G in a set with dense span. Proposition 3.8 shows that (H1)–(H3), O_h-invariance and ergodicity alone do not give it. Three steps fail.

(a) *The latent density.* $\Delta_S=0$, which gives hypothesis (L) of Theorem 10.33, needs a bound on the torus susceptibilities of the Wick square uniform in the regulators (Remark 10.36). It is expected to fail on the first-order surface of Remark 14.1, where the limit pressure of the Wick square should have a corner at the reference mass and a limit state may be a mixture of two coexisting phases; there only the bound $\Delta_S\le\kappa_+ + 2S_\flat^B$ of Proposition 10.35 remains, and its constants are not shown to be uniform as $\lambda\downarrow 0$. Convexity suggests a way around this step away from exceptional points. A constant tilt $\sigma X_{\mathrm{R}}(1_T)$ of a torus state is the same model with b_T replaced by $b_T-2\sigma$, and its pressure is $\alpha(\sigma+\cdot)-\alpha(\sigma)$; since the convex function α is differentiable off a countable set, limits of the tilted states would have latent density zero for all but countably many σ, provided the analysis of Sections 8–10 is carried over to the tilted family, which we do not do; and $\sigma=0$ may be exceptional. The analogous argument in the mass would need, in addition, regulator data of Lemma A.5 chosen uniformly on compact sets of masses and two-sided Lipschitz bounds on the torus coefficient as a function of the mass, which we do not verify; its exceptional set depends on the subsequence and is expected to contain the mass of the first-order transition.

(b) *Higher degree in the curvature.* For observables of degree at least two in the curvature the axis-invariant parts are governed by the clustering of the current sector, for which there is no estimate (Remark 10.30); a Gaussian domination of the linear field does not control them (Example 10.29).

(c) *Hypothesis (D).* Hypothesis (D) of Corollary 10.38 excludes a non-trivial ergodic decomposition and is a uniqueness-type statement (Remark 10.39).

Two problems concern the components themselves. Let $\mu=\int\mu_\xi\,d\nu(\xi)$ be the decomposition of Theorem 3.6 of the law of a limit state S, let P_j be the orthogonal projection of $L^2(\mu)$ onto the functions invariant under the translations along e_j, and $\mathcal{I}$ the σ-algebra of translation-invariant Borel sets.

Single-axis ergodicity of the components. Show that for ν-almost every ξ the translations along some coordinate direction e_j act ergodically on μ_ξ, that is, μ_ξ satisfies (OS4$_j$) of Definition 2.11; equivalently (Proposition 10.41), $\operatorname{Ker} H_j=\mathbb{C}\Omega_j$ in the reconstruction from μ_ξ in direction j, or $P_jG=\mathbb{E}[G\mid\mathcal{I}]$ for all $G\in L^2(\mu)$. Show also the cluster property of Definition 2.12(i) for the components. The identity $P_jG=\mathbb{E}[G\mid\mathcal{I}]$ holds for the observables of degree one in the curvature, with any square-integrable dependence on the Wick square (Theorem 3.10(a)), and for the Wick-square fields if $\Delta_S=0$ (Theorem 3.10(b)). It does not follow from (H1)–(H3), cubic invariance and ergodicity under $\mathbb{R}^3$ alone (Proposition 3.8), and it would follow from invariance of S under rotations (Corollary 10.44). The conditions $\Delta_S=0$ and (D) are sufficient but stronger than needed: (D) makes μ itself ergodic, and $\Delta_S=0$ makes all components agree on the Wick-square fields (Theorem 10.33); both are expected to fail on the first-order surface of Remark 14.1, where the components may still satisfy (OS4$_j$).

Cubic symmetry and charge conjugation of the components. Show that ν-almost every μ_ξ is invariant under O_h; equivalently, that the integer k of Theorem 3.6(f) equals 1 for ν'-almost every ζ; equivalently, that every bounded translation-invariant function is invariant under the coordinate permutations. Show also that almost every μ_ξ is invariant under charge conjugation, which acts on $\mathcal{H}^{-r}$ by changing the sign of the curvature component, so that the moments of μ_ξ of odd degree in the curvature vanish; in degree one this holds (Theorem 10.27). The cubic symmetry of the components does not follow from (H1)–(H3) and the cubic symmetry of the law (Example 10.14). The coordinate reflections act trivially on translation-invariant functions, by reflection positivity (Lemma 10.8); for a coordinate permutation the same argument would need reflection positivity about the diagonal planes $\{x_k = x_l\}$, which is not available for these states. Invariance of S under rotations (Corollary 10.44) or ergodicity of μ would give the cubic symmetry of the components.

The translation-ergodic components are not known to be O_h-invariant (Remark 10.11); the decomposition may instead be chosen with components invariant under $\mathbb{R}^3 \rtimes O_h$, each a uniform mixture of the $k \in \{1, 2, 3, 6\}$ translation-ergodic measures of one orbit of the coordinate permutations (Theorem 3.6(f)), and for a law satisfying (H1)–(H3) the cubic symmetry of the law alone does not force $k = 1$ (Example 10.14). The translation-ergodic components satisfy OS0 only with constants that depend on the component (Corollary 10.43(iv)), and the vanishing of the odd curvature moments, OS3 for the limit states, is known for a component only in degree one (Theorem 10.27). No rate of decay is proved for any correlation.

14.6. **Other phases and block sizes.** The proofs of this paper do not use the phase. Sections 7–11 use the uniform tube bound, the exact lattice symmetries of tori and inequalities that hold on every lattice torus, and the inputs that give the tube bound (Sections 4–6) refer to the coupling range of Theorem 3.1, not to a phase. One-loop perturbation theory predicts a first-order transition inside that range (Remark 14.1); "symmetric phase" in the title refers to the couplings above the predicted surface. In the Higgs phase $m^2 < 0$ the ultraviolet analysis down to the Higgs scale must be redone (the effective action of [7]), and a unit-scale cluster expansion in unitary gauge, of the kind that [5] carried out on the unit lattice, would be expected to give uniqueness of the infinite-volume limit, exponential clustering and, through the agreement of cubic and tilted torus limits, full rotation invariance; none of this is attempted here. Blocking factors $L \geq 4$, in particular King's $L = 5$, are outside the scope of the proof: the argument by which Bałaban's Proposition 1 is applied under the weakened support hypothesis (R4‴) needs $L \leq 3$ (Remark 5.3(iii)).

Remark 14.1 (a predicted first-order surface). The following is one-loop perturbation theory, which we do not justify. In the normalisation (1.1) a constant background $|\varphi| = \rho$ gives the photon the mass $|e|\rho$; we take $e > 0$, since $(A, e) \mapsto (-A, -e)$ leaves the action invariant. In three dimensions the two polarisations of a vector field of mass M add $-M^3/(6\pi)$ to the effective potential (Landau gauge; the term linear in M^2 is absorbed by the mass renormalisation); the scalar loops are analytic in ρ^2 for $m > 0$ and negligible here. So, to one loop,

$$V(\rho) = \tfrac{1}{2}m^2\rho^2 - \frac{e^3}{6\pi}\rho^3 + \lambda\rho^4.$$

V has a minimum at some $\rho_* > 0$ degenerate with the one at $\rho = 0$ exactly when the quadratic polynomial $V(\rho)/\rho^2$ has a double root, $(e^3/6\pi)^2 = 2m^2\lambda$, that is on the surface

$$\lambda m^2 = \frac{e^6}{72\pi^2}\big(1 + O(e^2/m)\big),$$

where the correction stands for the counterterm scheme, the photon-mass regulator and higher loops. There $\rho_* = 6\pi m^2/e^3$ and the photon mass is $e\rho_* = 6\pi m^2/e^2$, so the loop parameter $e^2/(e\rho_*) = e^4/(6\pi m^2)$ is below 10^{-3} when $e^2 < 0.1m$. For every m and every e with $0 < e^2 < 0.1m$ the point $\lambda = e^6/(72\pi^2 m^2)$ of the surface lies in the coupling range of Theorem 3.1: $\lambda < 10^{-3}m/(72\pi^2) \approx 1.4 \cdot 10^{-6}\, m$, that is $\lambda/e^2 < 1/(7200\pi^2)$, deep in the type-I regime $\lambda \ll e^2$; so does

the region below it, since the range bounds λ from below only by $\lambda > 0$. Equivalently, for fixed e and $\lambda < e^2/(7200\pi^2)$ the transition is at the mass $m_c(e,\lambda) = |e|^3/(6\pi\sqrt{2\lambda})$. One loop thus predicts the symmetric, Coulomb phase above the surface ($\lambda m^2 > e^6/(72\pi^2)$); on it a first-order transition in the mass, with the density of the Wick square jumping by about $\rho_*^2 = 36\pi^2 m^4/e^6$; and below it, still at $m^2 > 0$, a phase with a massive photon. The theorems of this paper are not affected, since they do not use the phase; their reading is.

14.7. **Extremality and relativistic fields.** Extremality of the limit states is not proved. Theorem 3.6 decomposes their laws into translation-ergodic components, which are invariant under the coordinate reflections but are not known to be invariant under O_h (Section 14.5; Theorem 3.6(f) gives O_h-invariant components whose joint vacuum has dimension $k \in \{1,2,3,6\}$), and under hypothesis (D) the law is ergodic (Theorem 3.10(c)); whether the law of a limit state is ergodic, equivalently whether its joint vacuum is unique (Theorem 11.1(iv)), is a uniqueness-type question of the kind in Section 14.2. The reconstruction of Theorem 11.1 has a positive Hamiltonian, unitary spatial translations and the point group D_4, but no Lorentz covariance, which would require invariance under the Euclidean rotations in the planes that contain the time axis (Section 14.1). OS0 holds in the form E0″ of [28, (4.2), p. 287] and implies E0′ by the Appendix of [28], which uses no rotations (Section 11). Under E0′ together with E1 and E2, [28] continues the Schwinger functions analytically to tempered Wightman-type distributions with the spectrum condition and translation covariance. Whether that argument can be run with the translations, D_4 and reflection positivity alone has not been examined, and no Wightman-type distributions are claimed for the limit states.

Appendix A. Regulator data and the order of choices

This appendix fixes the regulator data of the construction (the blocking factor, the parameters of Bałaban's renormalisation group, the spacing ε_0 of the coarsest lattice and the side ℓ_0 of the smallest torus), lists the conditions they have to satisfy, and proves that these conditions can be met at every point of the window $\mathfrak{W}$ of (1.3) (Lemma A.5). Theorem 5.2, Theorem 6.8 and the results of Appendices D–I are proved under these conditions. The conditions are named by type: (P.·) conditions on the point of the window, (R.·) lower bounds on R, (b.·) conditions on b_0, (E.·) upper bounds on ε_0, and (T.·) lower bounds and divisibility conditions on ℓ_0. Throughout, $L = 2$, $d = 3$, $\hat{\ell} := 1 + \log \varepsilon_0^{-1}$, $\ell(t) := 1 + \log t^{-1}$, $r(t) = R\,\ell(t)^{\varrho}$ with $\varrho = 2$ is Bałaban's localisation radius at scale t [2, (2.7), p. 558], $p(t) = b_0 \ell(t)^p$ is his small-field parameter, $t_k = L^k \varepsilon$ ($0 \le k \le K$, so that $t_K = \varepsilon_0$), $a_\flat := a(1 - L^{-2})$, $\hat{x} := |e|\,\varepsilon_0^{1/2}$, and $\mathbb{M}_4 := \{M_1, m, m_1, m_2\}$ with $M_1 = 0.7m$. The constant c_0 is the constant of stage 3 below; the constant called c_0 in [2, p. 593] is written $c_\square$.

A.1. **The window and the regulator data.** Let $\pi = (e, \lambda, m, \eta, m_1, m_2) \in \mathfrak{W}$, and put $\mu_* := m$, $M_1 := 0.7m$ and $H := m^2$. The regulator data at $L = 2$ are: $p = 6$; $\bar{n} = 13$ and the order cap $\bar{N} = \bar{n}^2 = 169$, with some decay rate $\delta(\bar{N}) > 0$ of the families at the cap (this is not Bałaban's δ_0, which depends on d only [3, p. 420]); the block size M_b; the localisation parameter $R \ge R^{**}$ of Table 5, with $\delta_0 R \ge 4$; the small-field parameter b_0; the coarsest spacing $\varepsilon_0 = 2^{-n_0}$; and the side $\ell_0 \in 2M_b\mathbb{N}$ of the smallest torus. The tori and spacings of the construction are then $T_j := j!\,T_0$ (side $\ell_j = j!\,\ell_0$) and $\varepsilon = 2^{-K}\varepsilon_0$ with $K \ge n_0 + 1$; this definition imposes no condition, and $\varepsilon \le 1/m$ for all of them follows from condition (E.m2) below. Three rules are part of the conditions:

(i) *the lower bound on R*: $R \ge R^{**}$, which dominates every other lower bound on R of Table 5;
(ii) *the four-mass rule*: every upper bound on ε_0 is imposed at each Higgs mass $M \in \mathbb{M}_4$ at which it is used, and every constant or condition of the stages 8–11 below that depends on the Higgs mass is evaluated at each $M \in \mathbb{M}_4$, the four values being combined by the maximum for constants and for lower bounds, and by the minimum for upper bounds on ε_0;

(iii) *the torus floor*: $\ell_0 \geq \max\{96/m,\, 128/m,\, 128/M_1\}$ and ℓ_0 is above every lower bound of Table 8; since $128/M_1 = 182.857\ldots/m$, the choice $\ell_0 \geq 183/m$ meets the first three bounds, and M_1 is among the masses of stage 0, where the floor $\ell_0^\natural := \max\{8, 128/m_1, 128/M_1\} = 128/M_1$ is fixed.

Two conditions are produced by Section 5: the upper bound (E.edge) of (5.7) on ε_0 (stage 10b), from Lemma 5.6, and the lower bound (T.per) of (5.13) on ℓ_0 (stage 11), from Lemma 5.22.

A.2. **The stages of the choice.** The regulator data are chosen in the order of Table 3. Each stage depends only on data fixed in earlier stages, apart from steps inside one stage that depend on earlier data and are not used again inside that stage: the lower bounds on R give R (stage 5), the upper bounds on ε_0 give ε_0 (stage 10b), the lower bounds on ℓ_0 give ℓ_0 (stage 11), and inside the stages 0 (κ and $\ell_0^\natural$ from the point), 6–8 (constants from p and $\mathbb{W}_0$) and 10a (suprema over $(0, \varepsilon_1]$ from ε_1) the dependence goes one way.

stage	choice	depends on
point	(e, λ, m); $\mu_* := m$; the torus counterterm scheme of Definition 2.2, with one Wick constant z_ε; the torus side $96/m$ of Theorem 6.1; $M_1 := 0.7m$ and σ_+; $H := m^2$; the anchor masses m_1, m_2 and $\eta \in (0,1)$	—
0	$\pi = (e, \lambda, \mu_*, \kappa, \eta, M_1, m_1, m_2)$, with κ the domination constant of Theorem I.16 under the a priori bound $\mu_*\varepsilon_0 \leq 0.2$ (Remark I.18); the floor $\ell_0^\natural := \max\{8, 128/m_1, 128/M_1\} = 128/M_1$	point
1–2	$a = 1$, $L = 2$; the coefficients a_k, through the k-uniform bound $a_\flat < a_k \leq a$	0
3	M_b; the constants $\delta_0, R_0, c_0, c_1, \ldots$ of [1, 2, 3] (functions of d, a, L, M_b)	0–2
4	$\bar n = 13$, $\bar N = \bar n^2 = 169$, a decay rate $\delta(\bar N) > 0$ of the families at the order cap; the constants $C_{\mathrm{loc}}(\bar N)$ and $C_{\bar N}$ of the conditions (R.cap) and (T.cap); the width $\ell_{\bar N}$ of Subsection 5.3; the collar exponent $\kappa_0 := \frac{1}{5}$ (Lemma F.1)	0–3
5	$R \geq R^{**}$, $\delta_0 R \geq 4$, and the condition (R.cap), $R\hat\ell^2 \geq C_{\mathrm{loc}}(\bar N)$ (met by $R \geq C_{\mathrm{loc}}(\bar N)$, since $\hat\ell \geq 1$): lower bounds on R only	0–4
6–8	$p = 6$ (so that $2p > \varrho d + 2$, as Proposition 5.21 needs), $p_1 = 1$; $\mathbb{W}_0 = [1-\eta, 2]$ and its constants; every constant that depends on the Higgs mass, at each $M \in \mathbb{M}_4$ (maximum)	0–5
9	$b_1 = 1$; b_0 above its lower bounds (Table 6), all of them free of ε_0	0–8
10a	$\varepsilon_1 := \min\{1, \varepsilon_{(\mathrm{E.x})}, \varepsilon_{(\mathrm{E.md})}, \varepsilon_{(\mathrm{E}.\delta)}, \varepsilon_{(\mathrm{E.dom})}, \varepsilon_{(\mathrm{E.14})}\}$, where $\varepsilon_{(\cdot)}$ is the largest $\varepsilon_0 \leq 1$ at which the condition $(\cdot)$ of Table 7 holds; every constant that depends on ε_0 is replaced by its supremum over $(0, \varepsilon_1]$	0–9
10b	$\varepsilon_0 :=$ the largest $2^{-n_0} \leq \varepsilon_1$ that meets every upper bound of Table 7 at each $M \in \mathbb{M}_4$ (minimum), among them (E.r): $\varepsilon_0 < \mathrm{e}^{1-\varrho} = \mathrm{e}^{-1}$, (E.ent), (E.edge), (E.5), (E.ksp1)–(E.ksp3), and (R.cap) if it is not met at stage 5	0–10a
11	$\ell_0 \in 2M_b\mathbb{N}$ above every lower bound of Table 8, (T.cap) and (T.per) included; $T_j := j!\, T_0$; $\varepsilon = 2^{-K}\varepsilon_0$	0–10b

TABLE 3. The order of choices of the regulator data.

The lower bound $b_0^2\hat\ell^{12} \geq \max\{2, \bar\mu\lambda^{-1/2}\}$, which is part of the hypotheses of [2, (2.12)], is imposed at stage 9 in the ε_0-free form (b.3) of Table 6, $b_0^2 \geq \max\{2, \bar\mu\lambda^{-1/2}\}$, which implies it since $\hat\ell \geq 1$; the second part, $\lambda^{-1/2}\varepsilon_0^{-1/2} b_0^2 \hat\ell^{11} \geq C_{2.12}$, is the downward-closed upper bound (E.6) on ε_0 at stage 10b. Imposed in the other way, the first part would make b_0 depend on ε_0 and close the cycle $\varepsilon_0 \to b_0 \to$ (E.edge) $\to \varepsilon_0$.

A.3. **The conditions.**

Definition A.1 (the conditions on the regulator data)**.** The *conditions* of this appendix are those of Tables 4–8, together with the rules (i)–(iii) of Subsection A.1. The last column of each table lists the stages of Table 3 whose data the condition depends on.

In the tables, letters other than the regulator data, the point π and the quantities defined in this appendix denote constants of the estimates of [1, 2, 3] in which the condition is used (for instance c_v, e_*, C_{reg}, $C_{2.12}$, $C_{2.21}$, C'_H, $C_{H''}$, δ', γ_1, C_Z, Err, c_M, C_d, c_b, b_*, C_A^{R1}, ρ_1, C_α, α, α_I, $C_\delta(M)$, c_0^{ker}, $\bar\mu$, c_1^I, C_∂, c_{reg}, C_W, c, C; a subscript such as 2.21 names the display [2, (2.21)]); each of them depends only on the data of the stages in the last column, and none depends on ε, K, ℓ_T or u. The constants $\delta_1, c_1', c_2, \delta_h, \rho, M_\theta$ are further constants of [1, 2, 3] fixed at stage 3, with $c_1 \le 3$, $c_2 \le 9$, $M_\theta \le LM_b$, and b_d is a number depending on d only. The constants C_H, δ_H of (E.5) are defined in Lemma A.15. We write $e(t) := |e|\, t^{1/2}$.

name	condition	depends on
(P.1)	$\lambda > 0$, $e \neq 0$, $m > 0$	point
(P.2)	$\pi \in \mathfrak{W}$: $0 < \lambda < 0.1m$, $0 < e^2 < 0.1m$, $0 < \eta < 1$, $2m < m_1 < m_2$	point
(P.3)	Theorem 6.1(a) at $\mu_* = m$, $M_1 = 0.7m$, on tori of side $\ge 96/m$ and for $\varepsilon \le 1/m$; in units $m = 1$ it follows from $\Pi(\lambda, e^2) := 0.51 + 0.256576\lambda - 0.088629e^2 - 7.1791\lambda^2 - 9.3304e^2\lambda - 1.7038e^4 > 0$, and on the closed box $[0, 0.1m]^2$ it gives $2\sigma_+ \ge 0.319m^2$	point
(P.4)	Theorem 6.1(c): $m_1 \ge m_1^c$, with $m_1^c \le 1.882m$ (Corollaries G.8 and G.9)	point
(P.5)	Theorem 6.1(b): $m_2 > \max\{m_1, S(m_1) - m_1\}$; $g_{21} \ge (m_2 - m_1)(m_2 + m_1 - 0.3631m) > 0$	point
(P.6)	Theorem 6.1(d): $m_1 \ge \sqrt{2}\mu_*$; $m < m_1 < m_2$; $M_1 < m$; $\eta \in (0, 1)$ fixed	point

TABLE 4. Conditions on the point π. They hold at every $\pi \in \mathfrak{W}$ by Theorem 6.1; $S(m_1)$ is any number $\ge \sup_{M \ge m_1} p(M)$, with $p(M)$ the majorant of Proposition G.6.

The conditions at the order cap. Let $\bar n = 13$ and $\bar N = \bar n^2 = 169$, and let $C_{\mathrm{loc}}(\bar N)$ and $C_{\bar N}$ be the numbers fixed at stage 4: the side, in units of blocks of the current scale, of a cube that contains a connected component of a family of order at most $\bar N$, and the corresponding additive margin [3, pp. 432–433]. The two conditions of Lemma A.21 are

$$\text{(R.cap)} \quad R\hat\ell^2 \ge C_{\mathrm{loc}}(\bar N), \qquad\qquad \text{(T.cap)} \quad \ell_0 \ge 3r(\varepsilon_0)\varepsilon_0 + C_{\bar N}\varepsilon_0.$$

(R.cap) is a lower bound on R at stage 5. It is imposed in the form $R \ge C_{\mathrm{loc}}(\bar N)$, which is part of (R.5) and of $R \ge R^*$ and implies (R.cap) for every $\varepsilon_0 \le 1$, because $\hat\ell \ge 1$. (T.cap) is a lower bound on ℓ_0 at stage 11. Together with (E.r) they give, for every $k \le K$ and every torus of side $\ell_T \ge \ell_0$,

$$r(t_k) \ge C_{\mathrm{loc}}(\bar N), \qquad \big(3r(t_k) + C_{\bar N}\big)t_k \le \ell_T.$$

The first holds because $1 + \log t_k^{-1} \ge \hat\ell$ for $t_k \le \varepsilon_0$. The second holds because $t \mapsto t(1 + \log t^{-1})^2$ is increasing on $(0, \mathrm{e}^{-1}]$, so that $r(t_k)t_k \le r(\varepsilon_0)\varepsilon_0$ when $t_k \le \varepsilon_0 < \mathrm{e}^{-1}$, and because $C_{\bar N}t_k \le C_{\bar N}\varepsilon_0$.

The inequalities of the conditions on the scalar zero mode, on the Hessian of [2, (3.12)] and of the logarithmic mass entries are given in Lemmas A.12, A.14, A.15 and A.16: the first two impose no condition beyond (E.m2), (E.5) and (E.fin), and the last is covered by (E.fin). Some routes through the estimates of [2] that the proof of Theorem 5.2 does not use carry further conditions ($\varepsilon_0 \le \mathrm{e}^{-4}$, $r(\varepsilon_0) > 3M_b$ and the lower bounds $6r(\varepsilon_0)\varepsilon_0$, $21\varepsilon_0$, $M_b\varepsilon_0 a_K(a_K - 1)$ on ℓ_0 of the cube-partition routes; $\mu_*\varepsilon_0 \le 0.1525$); they are not needed and are not imposed.

A.4. **The two conditions from Section 5; acyclicity.**

Lemma A.2 (the conditions (E.edge) and (T.per) in the order of choices)**.** *Let* $c_d = \sqrt{3}M_b$.

name	condition	depends on
(R.1)	$R > R_0$; $\delta_0 R \geq 2L = 4$	3
(R.2)	$R \geq 107$; $R \geq 3R_0 + 15$	3
(R.3)	$R \geq \ell_{\bar N} + 4\sqrt{3} + 2$; $R \geq 2\ell_{\bar N} + 8M_b + 8$; $R \geq 2\ell_{\bar N} + 10M_b + 10$	3, 4
(R.4)	$R \geq R_0 + 2\sqrt{3} + \delta_h + 1 + \rho$; $2r(\varepsilon_0) \geq R_0 + 2\sqrt{3} + 1$	3
(R.5)	$R \geq C_{\mathrm{loc}}(\bar N)$; $R \geq R_0 + 13 + c_1'$; $R \geq M_b + 2$, $R \geq c_1'$, $R \geq c_2/4$	3, 4
(R.6)	$R \geq [L(M_\theta + 6) + 6]/5$ (implied by (R.3))	3
(R.7)	$r(\varepsilon_0) - 2M_b \geq R_0 + 2\sqrt{3} + 1$; $r(\varepsilon_0) > 3M_b$, $R\hat\ell^2/4 \geq 2M_b + 2$	3, 5
(R.8)	$R \geq 14b_d{+}1$, $R \geq (11\sqrt{3}/2)M_b$, $R \geq \max\{12\sqrt{3}, 14M_b\}$; $r(t) \geq \ell_{\bar N}{+}4\sqrt{3}{+}2$ for $t \leq \varepsilon_0$	3, 4
(R.e)	$R \geq 4\ell_{\bar N} + 32M_b + 25$ (Corollary 5.10)	3, 4
(R.edge)	the three conditions (5.6), imposed in the sufficient form $r(\varepsilon_0) \geq R$, which they follow from (Lemma A.3)	3, 5
(R.cap)	$R\hat\ell^2 \geq C_{\mathrm{loc}}(\bar N)$, imposed in the sufficient form $R \geq C_{\mathrm{loc}}(\bar N)$ (Lemma A.21)	4, 5
(R.nbar)	$2\bar n + 1 \leq 3R$ (at $\bar n = 13$ it follows from (R.2)); used for the non-wrapping of the lifted clusters at zero background (Appendix K)	4

TABLE 5. Lower bounds on R (stage 5). None depends on ε, k, ℓ_T, u or the Higgs mass. In (R.4), (R.7), (R.8), (R.edge) and (R.cap) the quantities $r(\varepsilon_0)$, $r(t)$ and $\hat\ell$ are replaced by their lower bounds R, R and 1 (valid for $\varepsilon_0 \leq 1$); the forms so obtained imply the printed ones. All of them hold if $R \geq R^{**} := \max\{R^*, 4\ell_{\bar N} + 32M_b + 25\}$ with $R^* := \max\{4/\delta_0, 107, 3R_0 + 15 + c^\sharp, 2\ell_{\bar N} + 10M_b + 10, C_{\mathrm{loc}}(\bar N)\}$, $c^\sharp := \max\{\delta_h + \rho, c_1', c_2\}$, together with the finitely many explicit bounds of (R.4)–(R.8).

name	condition	depends on
(b.0)	$p = 6$, $p_1 = 1 \in (2/3, p-1]$	—
(b.1)	$b_1 = 1$, $b_0 \geq b_0^{\min} := \max\{4d, c_1^I\} c_w (1 + \log 2)/(p \log 2)$, $c_w \leq (13/12)^{1/2}$	3–8
(b.2)	$b_0 \geq R/C$	5
(b.3)	$b_0^2 \geq \max\{2, \bar\mu \lambda^{-1/2}\}$	0
(b.4)	the constants of stage 8 that depend on the Higgs mass (c_7, c_1^I, $\gamma_1''(M)$, $C_b'(M)$, $C_b^R(M)$, $C_E^R(M)$, $C_{F1}(M)$, $\mathfrak{E}_k(M)$) are taken at their maximum over $M \in \mathbb{M}_4$; the constant $C_> := \sup_{t\in(0,1]}(\dots)$ is evaluated at the floor $\ell_0^\natural$	0–8

TABLE 6. The parameters p, b_0 and the constants of the stages 6–9.

(a) *The condition (E.edge) of* (5.7)*, with a_k replaced by its bound a, depends only on data of the stages 0–9 (L, a; λ; Balaban's exponent κ of* (5.7)*, fixed with $\bar n$ at stage 4 and distinct from the domination constant κ of stage 0; M_b, δ_0, c_0, c_1; R through $r(t)$; p; b_0 through $p(t)$), and does not depend on M, u, ε, K or ℓ_T. It is downward closed in ε_0 and holds for all sufficiently small ε_0, at every $R > 0$. The conditions (R.edge) of* (5.6) *follow from the lower bound $R \geq R^*$ of stage 5, and the condition $r(t_{k-1})/r(t_k) \leq 2L$ of Lemma 5.6 holds for every $\varepsilon_0 \leq 1$.*

(b) *On the cubic torus T_j the half-side of Balaban's torus $\{-L_\mu \leq x_\mu < L_\mu\}$* [1, (1.2), pp. 603–604] *is $L_{\min} = \ell_j/2$. So the condition* (5.13) *reads $\ell_j \geq \varepsilon_0 \max\{6\sqrt{3}\, r(\varepsilon_0) + 2,\ 2/\delta_0\}$, and it is imposed at stage 11 as*

$$\textit{(T.per)} \qquad \ell_0 \ \geq\ \varepsilon_0 \max\{6\sqrt{3}\, r(\varepsilon_0) + 2,\ 2/\delta_0\}.$$

This condition depends only on δ_0 (stage 3), R (stage 5) and ε_0 (stage 10b), and it is a lower bound on ℓ_0. If it holds and $\varepsilon_0 < \mathrm{e}^{1-\varrho}$ (condition (E.r)), then for every $j \geq 1$, every

K and every scale $k \le K$, the rescaled side $S_k = \ell_j/t_k$ of T_j satisfies $S_k \ge \max\{6\sqrt{3}\,r(t_k)+2,\ 2/\delta_0\}$; this is what Lemma 5.22(iv) uses.

(c) *With these two conditions, the order of choices of Table 3 is acyclic and every stage can be filled.*

Proof. (a) The data are read off (5.7); the coefficient a_k enters only through $a_k \le a$, so the condition does not depend on k, and the masses do not enter. The condition is imposed for all $t \le \varepsilon_0/L$, so if it holds for ε_0 it holds for every smaller ε_0. It holds for small ε_0 because, with $s = \log t^{-1}$, the logarithm of the left side minus that of the right side is $(\frac14+\kappa)s + O(\log s) - \delta_0 R(1+s)^{\varrho}/L + O(1) \to -\infty$ as $s \to \infty$, since $\varrho = 2 > 1$; this uses only $R > 0$. For (R.edge) note $r(\varepsilon_0) \ge R$ for $\varepsilon_0 \le 1$. The first condition of (5.6) follows from $r(\varepsilon_0) \ge 0.4R_0 + \bigl(1+0.4(2+\sqrt{3})\bigr)M_b$, which is $\le 0.4R_0 + 2.5M_b$. Now $R^* \ge \max\{3R_0+15,\ 2\ell_{\bar N}+10M_b+10\}$. If $R_0 \le 18.7M_b+25$, then $10M_b+10 \ge 0.4R_0+2.5M_b$; otherwise $R_0 > 0.97M_b$, and $3R_0+15 \ge 0.4R_0+2.5M_b$. The second condition, $4r(\varepsilon_0) \ge R_0 + \sqrt{3}M_b$, follows from $4R \ge 2(3R_0+15) + 2(10M_b+10)$, and the third, $R \ge 10M_b+10$, is part of $R \ge R^*$. Finally $r(t/L)/r(t) = \bigl(1+\log L/(1+\log t^{-1})\bigr)^{\varrho} \le (1+\log L)^2$ for $t \le 1$, and $(1+\log 2)^2 < 2.87 \le 4$.

(b) The data are read off (5.13). Since $\ell_j = 2L_{\min} \ge \ell_0$, the bound at ℓ_0 gives (5.13) on every T_j. Put $f(t) := t(1+\log t^{-1})^{\varrho}$. Then $f'(t) = (1+\log t^{-1})^{\varrho-1}(\log t^{-1} + 1 - \varrho) > 0$ for $0 < t < \mathrm{e}^{1-\varrho}$. Hence $t \mapsto t\,(6\sqrt{3}\,r(t)+2) = 6\sqrt{3}Rf(t) + 2t$ is increasing on $(0, \varepsilon_0]$ when $\varepsilon_0 < \mathrm{e}^{1-\varrho}$, and for $k \le K$, since $t_k \le t_K = \varepsilon_0$,

$$t_k\bigl(6\sqrt{3}\,r(t_k)+2\bigr) \le \varepsilon_0\bigl(6\sqrt{3}\,r(\varepsilon_0)+2\bigr) \le \ell_0 \le \ell_j,$$

that is, $S_k \ge 6\sqrt{3}\,r(t_k)+2$. Also $S_k \ge \ell_j/\varepsilon_0 \ge 2/\delta_0$. The condition is not implied by the other lower bounds on ℓ_0: (T.nw) gives $\ell_0 \ge \varepsilon_0[8r(\varepsilon_0) + 4M_b + 2]$ and $r(\varepsilon_0) \ge R \ge R^* \ge 10M_b + 10 > 3M_b$, so $8r(\varepsilon_0) + 4M_b + 2 < 9.34\,r(\varepsilon_0) + 2 < 6\sqrt{3}\,r(\varepsilon_0) + 2$ and (T.nw) never gives the first term of (T.per); and (T.floor) in the form $\ell_0 \ge 183/m$ together with (E.dom) gives $\ell_0/\varepsilon_0 \ge 915$, which covers $2/\delta_0$ only if $\delta_0 \ge 2/915$, while δ_0 is a constant of [1] without a printed value.

(c) This is Lemma A.4, which treats all conditions of Tables 4–8 with (E.edge) and (T.per) among them. Since by (b) the condition (T.per) exceeds (T.nw), ℓ_0 is in general larger than (T.nw) alone requires; this violates no other condition, since no condition depends on ℓ_0 and the conditions of Table 4 hold on all tori of side at least $96/m$. □

Lemma A.3 (the conditions (R.edge) and (E.edge))**.** *Let $d = 3$, $c_d = \sqrt{3}M_b$, and assume $M_b \ge 2$, $R_0 \ge 0$, $\ell_{\bar N} \ge 0$, $\delta_0 > 0$ and $0 < \varepsilon_0 \le 1$.*

(i) *Suppose $R \ge 3R_0 + 15$ and $R \ge 2\ell_{\bar N} + 10M_b + 10$ (part of (R.2) and (R.3)); in particular this holds if $R \ge R^{**}$. Then the three conditions (5.6) hold. Since $c_d \ge \sqrt{3}$, so do the weaker forms $5(r(\varepsilon_0) - M_b)/L \ge R_0 + \sqrt{3} + 2M_b$ and $4r(\varepsilon_0) \ge R_0 + \sqrt{3}$.*

(ii) *For $0 < t \le 1$, $r(t/L)/r(t) \le (1+\log L)^2 \le 2L$. Hence, for every k, with $\rho_k := (r(t_{k-1}) - M_b)/L$, we have $\rho_k \le 2r(t_k)$, and $3\rho_k \ge 2M_b$ under the third condition of (5.6).*

(iii) *Let (E.edge)$_a$ denote the condition (5.7) with a_k replaced by a. It implies (5.7) at every k. It depends only on L, a (stages 1–2), λ (stage 0), Bałaban's exponent κ (fixed with $\bar n$ at stage 4), M_b, δ_0, c_0, c_1 (stage 3), R (stage 5), p (stage 6), b_0 (stage 9) and $C = C(c_0, \delta_0, d)$. It does not involve M, u, ε, K or ℓ_T. It is downward closed in ε_0, and for every $R > 0$ it holds for all ε_0 below a positive threshold that depends only on these data.*

Proof. (i) Since $\varepsilon_0 \le 1$ we have $\log \varepsilon_0^{-1} \ge 0$, so $r(\varepsilon_0) \ge R$. The maximum of two numbers is at least their mean. The first condition reads $r(\varepsilon_0) \ge \frac25(R_0 + c_d + 2M_b) + M_b = 0.4R_0 + \bigl(1+0.4(2+\sqrt{3})\bigr)M_b$, and $1 + 0.4(2+\sqrt{3}) = 2.4928\ldots < 2.5$. By the two hypotheses,

$$R \ \ge\ \tfrac12(3R_0+15) + \tfrac12(2\ell_{\bar N} + 10M_b + 10) \ \ge\ 1.5R_0 + 5M_b + 12.5 \ \ge\ 0.4R_0 + 2.5M_b.$$

For the second condition, $4r(\varepsilon_0) \ge 4R \ge 2(3R_0+15) + 2(2\ell_{\bar N} + 10M_b + 10) \ge R_0 + \sqrt{3}M_b$. The third is $R \ge 2\ell_{\bar N} + 10M_b + 10 \ge 10M_b + 10$.

(ii) Put $s = \log t^{-1} \geq 0$. Then

$$\frac{r(t/L)}{r(t)} = \Big(\frac{1+\log L+s}{1+s}\Big)^2 = \Big(1+\frac{\log L}{1+s}\Big)^2 \leq (1+\log L)^2,$$

and $(1+\log 2)^2 = 2.8667\ldots \leq 4$. With $t = t_k \leq \varepsilon_0 \leq 1$ and $t_{k-1} = t_k/L$ this gives $\rho_k \leq r(t_{k-1})/L \leq 2r(t_k)$. Since $r(t_{k-1}) \geq R$ and $L = 2$, $3\rho_k \geq 3(R - M_b)/2 \geq R - M_b \geq 9M_b + 10 \geq 2M_b$.

(iii) Since $0 < a_k \leq a$, the left side of (5.7) with a is at least the one with a_k. So $(\text{E.edge})_a$ implies (5.7) at every k. Its only dependence on k and on the mass came through a_k, and that is now gone. The list of data is read off the display; b_0 enters through $p(t)$.

The condition is imposed for all $t \leq \varepsilon_0/L$. If $\varepsilon_0' \leq \varepsilon_0$, then this range of t shrinks, so the condition is downward closed. For the threshold, put $s = \log t^{-1}$. Then $\log p(t) = \log b_0 + p\log(1+s)$ and $\log(1+r(t)) = O(\log(1+s))$. So the logarithm of the left side minus that of $(Lt)^\kappa$ is

$$\log\big(C\,a\,c_1\lambda^{-1/4}b_0\,\mathrm{e}^{\delta_0(M_b/L+c_d)}L^{-\kappa}\big) + \big(\tfrac14+\kappa\big)s + O(\log(1+s)) - \frac{\delta_0 R}{L}(1+s)^2,$$

which tends to $-\infty$ as $s \to \infty$, because $\delta_0 R > 0$ and $\varrho = 2 > 1$. Hence there is an $s_0 < \infty$, depending only on the listed data, such that the condition holds for every $t \leq \mathrm{e}^{-s_0}$, that is, whenever $\varepsilon_0 \leq L\mathrm{e}^{-s_0}$. The factor $(1+r(t))^{d-1}$ changes only s_0. □

Lemma A.4 (the order of choices is acyclic and every stage can be filled)**.** *Consider the conditions of Definition A.1, read as follows: (1) (b.3) in its ε_0-free form, the other part of the hypothesis of* [2, (2.12)] *being the upper bound (E.6) at stage 10b; (2) a_k in (E.edge) replaced by its bound a; (3) (R.edge) in its sufficient stage-5 form (Lemma A.3(i)). Form the directed graph whose vertices are the data of Table 3 and the conditions, with an edge from each datum to each condition that depends on it (as listed in the last columns of Tables 4–8) and an edge from each condition to the datum it constrains. Then:*

(i) *every edge goes from a stage to the same or a later stage;*
(ii) *the graph is acyclic;*
(iii) *every stage can be filled;*
(iv) *no condition is an upper bound on R, b_0 or ℓ_0 or a lower bound on ε_0, and none depends on ε, k, ℓ_T or K.*

Proof. (i) This is read off the last columns of Tables 4–8, of Lemma A.3(iii) for (E.edge) and of Lemma A.2(b) for (T.per). The following potential edges against the order are excluded by the form of the conditions. (a) (E.edge) and R: R is fixed at stage 5 from data of the stages 2–4, and no lower bound on R depends on a datum of stage 10. (b) (E.edge) and b_0: the lower bounds on b_0 ((b.1), (b.2), the maximum in (b.4), and (b.3) read as in (1)) depend neither on ε_0 nor on (E.edge). (c) ε_1 (stage 10a) depends on no datum of stage 10b. (d) The non-wrapping of the box of Lemma 5.6 is the condition (T.nw) at stage 11, which depends on ε_0; the edge points forward. (e) (E.edge) does not involve the mass (Lemma A.3(iii)), so the four-mass rule adds no instance of it. (f) The floor $\ell_0^\natural$ is not a datum of (E.edge). (g) (R.edge) follows from $R \geq R^*$, so no edge into R from it is needed. (h) (E.13) depends on b_0 (stage 9) and is imposed at stage 10b; (b.1) does not depend on ε_0. The condition (T.per) depends on δ_0 (stage 3), R (stage 5, through $r(\varepsilon_0)$) and ε_0 (stage 10b); its one edge out goes to ℓ_0 (stage 11), and no condition depends on ℓ_0.

We now go through the tables condition by condition.

Table 4. The conditions (P.1)–(P.6) depend on the point only and constrain the point. They are infima or suprema over all tori of side at least $96/m$ and all $\varepsilon \leq 1/m$, so they depend neither on ε_0 nor on ℓ_0.

Table 5. Every condition constrains R (stage 5), and its coefficients are data of the stages 3 and 4. The quantities $r(\varepsilon_0)$, $r(t)$, $\hat\ell$ in (R.4), (R.7), (R.8), (R.edge) and (R.cap) would introduce an edge from ε_0 (stage 10b). As stated in the caption of Table 5, they are replaced by R, R and 1.

Since $r(t) \geq R$ and $\hat{\ell} \geq 1$ for $t, \varepsilon_0 \leq 1$, the forms so obtained imply the printed ones, and they do not depend on ε_0. (R.nbar) depends on $\bar{n}$ (stage 4).

Table 6. (b.1)–(b.3) constrain b_0 (stage 9). Their coefficients lie in the stages 0–8 ((b.2) depends on R, stage 5), and (b.3) is free of ε_0 by (1). (b.4) fixes stage-8 constants as maxima over $\mathbb{M}_4$, whose elements are data of stage 0, and at the floor $\ell_0^\natural$, also a datum of stage 0. The constants of the stages 7 and 8 depend on ε_0 only through numbers that are guaranteed by conditions of stage 10a. These are: the bound $\gamma_1'' \geq a_\flat/(a+1)$ for $M \leq m_2$, guaranteed by (E.m2); the number $\frac{1}{12}$, guaranteed by (E.δ) and (E.14); and one term proportional to ε_0 in a constant of [2], which is bounded at stage 10a by its value at ε_1 (Lemma A.11). The conditions (E.m2), (E.δ) and (E.14) depend only on data of the stages 0, 3 and 5. So each of these dependences goes one way, and none closes a cycle. The slice constant $\kappa = \frac{15}{2}$ is a number fixed at stage 0. The inequality $\kappa \geq \kappa_1\kappa_3$ that it must satisfy holds under (E.dom) (Remark I.18), and (E.δ) implies (E.dom) (Lemma A.19). This again is a one-way dependence. The constants of the per-step remainders that depend on b_0 and R appear in no condition and enter only the output constants.

Table 7. Every condition constrains ε_1 (stage 10a) or ε_0 (stage 10b). Its coefficients lie in the stages 0–9, or, through suprema over $(0, \varepsilon_1]$, in stage 10a. Under (2), the coefficients of (E.edge) are L, a, λ, the exponent κ, $M_b, \delta_0, c_0, c_1, R_0$, R through $r(t)$, p, p_1, b_0 through $p(t)$, and a constant depending on (c_0, δ_0, d). All of these are data of the stages ≤ 9. (E.ent) depends on Θ, which is formed from b_0, R and the suprema of c_5, c_7 over $(0, \varepsilon_1]$ (Lemma A.18), so on the stages 0–10a. (E.9) is imposed at each mass and depends on p (stage 6), on constants of stage 3, on R and on b_0; (E.13) depends on c_2, α_I, which are constants of [1] at stage 8. The last columns of (E.9), (E.13) and (E.ent) list these stages. (E.5) depends on C_H, δ_H (stage 3), on R, on the mass, and on ε_0 itself through γ_1''. (E.charge), (E.Breg) and (E.W) depend on data of the stages 0, 3, 4 and 9 and are imposed at stage 10a.

Table 8. Every condition constrains ℓ_0 (stage 11), and its coefficients lie in the stages $\leq$10b. No datum of a stage $\leq$10b depends on ℓ_0. The torus-dependent constants are evaluated at $\ell_0^\natural$ (Lemma A.10), and the conditions of Table 4 are uniform over all tori of side at least $96/m$.

Masses and output. The four-mass rule adds only edges from the masses (stage 0) into the stages 8–11 (Lemma A.9(iii)). The tori T_j, K and ε are chosen after stage 11, and by (iv) no condition depends on them.

(ii) Order the stages $0 < 1 < \cdots < 11$. By (i) the stage index does not decrease along any edge. So along a directed cycle it is constant: every cycle lies inside a single stage. Inside the stages 5, 10b and 11 the edges inside the stage go from the conditions to the quantity they constrain (R, ε_0, ℓ_0), and no condition of the same stage depends on that quantity; under (3) this includes (R.edge). A condition of stage 10b may contain ε_0 itself, as (E.5) does through γ_1''. Such a condition is a vertex with an edge to ε_0 only, and ε_0 is a datum of no condition other than through its own value, so this creates no cycle. Inside the stages 0, 6–8 and 10a the dependences inside the stage are one-way: from the point to κ and $\ell_0^\natural$; from p and $\mathbb{W}_0$ to their constants; from ε_1 to the suprema over $(0, \varepsilon_1]$. So no stage contains a cycle, and the graph is acyclic. In particular (T.per), whose data lie in earlier stages and whose one edge out ends at ℓ_0, on which no condition depends, lies on no cycle. It is imposed only as a lower bound on ℓ_0 at stage 11, never as an upper bound on ε_0 at a fixed torus, which would close the cycle $\varepsilon_0 \to$ (T.nw) $\to \ell_0 \to$ (T.per) $\to \varepsilon_0$.

(iii) At the stages 5, 9 and 11 there are finitely many lower bounds, each a finite function of data of earlier stages, so their maximum is finite. At stage 11 there is also the divisibility $\ell_0 \in 2M_b\mathbb{N}$, which every large multiple of $2M_b$ meets. At stage 10a the conditions are finitely many and each has a positive threshold, so $\varepsilon_1 > 0$. Among them, (E.charge) is explicit. (E.Breg) and (E.W) have the form "$F(t) \leq c$ for all $t \leq \varepsilon_0$" with $F(t) \to 0$ as $t \to 0$, because $p(t)$ grows only logarithmically and $e(t)^{1-\beta} = |e|^{1-\beta} t^{(1-\beta)/2}$ with $\beta < 1$; so they are of this type by Lemma A.7(c). At stage 10b there are finitely many upper bounds on ε_0, the instances at the four masses of $\mathbb{M}_4$ included.

Each of them is downward closed and holds for all dyadic ε_0 below a positive threshold: for the conditions of Table 7 other than (E.edge) this is Lemma A.8, and for (E.edge) it is Lemma A.3(iii). The minimum of finitely many positive thresholds is positive, and some $2^{-n_0} \le \varepsilon_1$ lies below it.

(iv) This is read off Tables 5–8: the conditions of the stages 5, 9 and 11 are lower bounds, and those of the stages 10a and 10b are upper bounds on ε_0. By (i), none of them depends on ε, k, ℓ_T or K. □

A.5. **The conditions at every point of the window.**

Lemma A.5 (the conditions at every point of the window)**.** *Let $\pi = (e, \lambda, m, \eta, m_1, m_2) \in \mathfrak{W}$, $\mu_* := m$, $M_1 := 0.7m$, $H := m^2$, $\mathbb{M}_4 := \{M_1, m, m_1, m_2\}$; let M_1 be among the data of stage 0, with the torus floor $\ell_0^\natural := \max\{8, 128/m_1, 128/M_1\}$ fixed there (lengths in the units of Section 2; $\ell_0^\natural = 128/M_1 = 182.857\ldots/m$ whenever $M_1 \le 16$), and let every constant or condition of the stages 8–11 that depends on the Higgs mass be evaluated at each $M \in \mathbb{M}_4$ and combined by the maximum (constants, lower bounds) and by the minimum (upper bounds on ε_0). Then there are R, b_0, n_0 and $\ell_0 \in 2M_b\mathbb{N}$ such that, with $\varepsilon_0 := 2^{-n_0}$, every condition of Tables 4–8 holds at π. For these data every spacing $\varepsilon = 2^{-K}\varepsilon_0$ satisfies $\varepsilon \le \varepsilon_0 \le 1/m_2 < 1/m$, every torus T_j has side $\ell_j \ge \ell_0 \ge 96/m$, and statements (a)–(d) of Theorem 6.1 hold at π.*

Proof. Step 1: the four types of conditions. By Lemma A.4 the order of choices is acyclic: every condition depends only on data of its own or of earlier stages. The condition (T.per) is a lower bound on ℓ_0 at stage 11 that depends only on δ_0 (stage 3), R (stage 5) and ε_0 (stage 10b), and no condition depends on ℓ_0. Every condition is of one of the following types.

(i) A condition on the point π alone: the conditions of Table 4, which are statements (a)–(d) of Theorem 6.1.
(ii) A lower bound $x \ge f$ on R (stage 5), on b_0 (stage 9) or on ℓ_0 (stage 11), where f is a finite function of data of earlier stages.
(iii) An upper bound on ε_0 (stages 10a–10b) that is downward closed (if it holds at a dyadic ε_0, it holds at every dyadic $\varepsilon_0' \le \varepsilon_0$) and that holds for all dyadic ε_0 below a positive threshold depending on data of earlier stages.
(iv) The divisibility $\ell_0 \in 2M_b\mathbb{N}$.

That the conditions of Tables 5–8 are of the types (ii)–(iv) at every point of $\mathfrak{W}$, with thresholds finite, respectively positive, at every point, is Lemma A.8, together with Lemma A.3(iii) for (E.edge) and Lemma A.2(b) for (T.per). The family $T_j = j!\,T_0$, $\varepsilon = 2^{-K}\varepsilon_0$, $K \ge n_0 + 1$, is a definition, and $\varepsilon \le 1/m$ follows from (E.m2). Moreover no condition bounds b_0 from above before ε_0 is chosen, no condition bounds ℓ_0 from above, the only conditions that bound R from above at fixed ε_0 are imposed after R (in the order of choices they are upper bounds of type (iii) on ε_0 or lower bounds of type (ii) on b_0, both chosen after R), and no condition involves the spacing ε, the index K or the torus except through the floor $\ell_0^\natural$ of stage 0 and the lower bounds of stage 11 (Lemma A.8(iii)).

Step 2: the conditions that involve masses, η and the couplings are of these types at every point of $\mathfrak{W}$. (i) $R^{**} = \max\{R^*, 4\ell_{\bar N} + 32M_b + 25\}$, and R^*, are functions of $(d, a, L, M_b, R_0, \delta_0, \bar n, \bar N, V(\bar N), C_{\mathrm{loc}}(\bar N))$, of fixed $O(1)$ localisation widths and of b_d, all fixed at the stages 0–4 independently of π; no Higgs mass, no coupling and no photon mass enters. The remaining lower bounds on R ($R_0 + 1$, $4/\delta_0$ and the bounds from (E.10), (E.14) and (R.cap)) are free of the Higgs mass, and the only parameters they contain, κ, μ_* and $(1-\eta)^{-3}$, are finite on $\mathfrak{W}$. So $R \ge R^{**}$ is of type (ii) (Lemma A.9(i)). (ii) The four-mass rule. Every constant of the stages 8–9 that depends on the Higgs mass depends on M either from above, through M^2t^2 with $t \le \varepsilon_0$, which (E.m2) at the largest mass, $m_2^2\varepsilon_0^2 \le 1$, bounds by 1 for every $M \le m_2$ (the constants are computed under this a priori bound, whose threshold $1/m_2$ depends on no constant, so no cycle arises), or through powers of M of either sign, logarithms and other quantities finite at every fixed $M > 0$ (for instance $(M^2\ell_T^3)^{-1}$, $(M\ell_T)^{-10}$ and $\log_+(1/(\min(M, \mu_*)\varepsilon_0))$ in the constant of the final c-numbers, the constant $C_\delta(M)$,

and the conditioning constant $\rho_{\min}(M)^{-1}$ of the bounds on the vector slices). No entry orders M against μ_*, and none uses $m_1 > 2m$; these conditions occur only in Table 4 (Lemma A.9(ii)). A maximum of four finite numbers is finite, and the constants enter only lower bounds, so these conditions are of type (ii). The upper bounds on ε_0 of stage 10b that depend on the Higgs mass ((E.5), (E.7), the conditions of Lemmas A.12–A.15, the logarithmic entries of Lemma A.16, (E.md) and (E.fin)) are downward closed with a positive threshold at each fixed $M > 0$; the minimum of four positive thresholds is positive, so these conditions are of type (iii). (iii) The condition (E.edge) reads $C\, a_k c_1 \lambda^{-1/4} t^{-1/4} p(t) \big(1 + r(t)\big)^{d-1} \exp\big(-\delta_0[(r(t) - M_b)/L - \sqrt{3}M_b]\big) \le (Lt)^\kappa$ for all $t \le \varepsilon_0/L$, with p polynomial in $\ln t^{-1}$ and $r(t) = R\,\ell(t)^2$, $\ell(t)^2/\ln t^{-1} \to \infty$ as $t \downarrow 0$. The exponential therefore decays faster than every power of t, and the inequality holds for all t below a positive threshold that depends on $L, R, M_b, \delta_0, c_0, R_0, \kappa, p, \lambda, a, c_1$, all fixed before stage 10b; it has no explicit Higgs-mass entry (it depends on b_0 through $p(t)$, and b_0 is fixed at stage 9 as a maximum over $\mathbb{M}_4$, so no cycle arises). The conditions (5.6), with $c_d = \sqrt{3}M_b$, follow from $R \ge R^*$ (Lemma A.2(a)). So (E.edge) is of type (iii). (iv) The condition (T.per), $2L_{\min} \ge \varepsilon_0 \max\{6\sqrt{d}\, r(\varepsilon_0) + 2,\, 2/\delta_0\}$, where $2L_{\min}$ is the side of the torus, is a lower bound on the side by a function of $\varepsilon_0, R, \delta_0$; since every T_j has side $\ge \ell_0$, it is a condition of type (ii) at stage 11. (v) (T.floor) reads $\ell_0 \ge \max\{96/m,\ 128/m,\ \ell_0^\natural\}$, of type (ii) at stage 11. The constants that are evaluated at the floor ($C_\delta(M)$, c_7, $C_>$ and the zero-mode terms of the bounds on the vector slices) are non-increasing in the side of the torus, so their values at $\ell_0^\natural$ dominate their values on every torus of side $\ge \ell_0 \ge \ell_0^\natural$; raising the floor from $\max\{8, 128/m_1\}$ to $\ell_0^\natural$ can only weaken the conditions in which they occur (Lemma A.10). (vi) The couplings and η enter the conditions only through quantities finite for $e \ne 0$, $\lambda > 0$, $0 < \eta < 1$ (for instance $\lambda^{-1/4}$ in (E.Φ) and in (E.edge), powers of $(1-\eta)^{-1}$ in (E.4) and (E.14)); the convexity range $0 < \eta < \eta_{\max}(\varepsilon_0)$ is an upper bound on ε_0 at fixed η. No condition of the types (ii)–(iv) is a condition on e or λ: the couplings are restricted only by the definition of $\mathfrak{W}$, whose upper bounds $\lambda < 0.1m$, $e^2 < 0.1m$ enter through the conditions of type (i) (the proof of Theorem 6.1(a)–(c)); in particular no condition imposes a lower bound on λ beyond $\lambda > 0$. As $\lambda \downarrow 0$ at fixed e the thresholds move (for instance through the factors $\lambda^{-1/4}$ above), but at every $\lambda > 0$ they stay finite, respectively positive. The smallness conditions of Bałaban's renormalisation group concern the running couplings at the scales $t \le \varepsilon_0$ and are conditions of type (iii).

Step 3: the choice at π. Choose the data stage by stage. At stage 0 the parameters and $\ell_0^\natural$; at the stages 1–4 the discrete data ($a = 1$, $L = 2$, M_b, $\bar{n} = 13$, $\bar{N} = 169$ and the others), which do not depend on π; at stage 5, R an integer above the finitely many lower bounds of type (ii), R^{**} among them; at stage 6, $p = 6$; at the stages 7–8 the constants, at the four masses; at stage 9, b_0 the maximum of the finitely many lower bounds; at stage 10a, ε_1 is the minimum of finitely many positive thresholds, and the constants that depend on ε_0 are taken at their suprema over $(0, \varepsilon_1]$, which are finite by Lemma A.11; at the stages 10a–10b, the finitely many upper bounds of type (iii), at the four masses and including (E.edge), have a positive common threshold $\bar{\varepsilon}(\pi)$, and every $\varepsilon_0 = 2^{-n_0} \le \bar{\varepsilon}(\pi)$ satisfies all of them by downward closedness; at stage 11, ℓ_0 the least element of $2M_b\mathbb{N}$ above the finitely many lower bounds. Each choice uses only data of earlier stages, and by Step 1 no later choice can violate an earlier condition. The conditions of type (i) hold by Theorem 6.1, which proves (a)–(d) at every point of $\mathfrak{W}$, uniformly in $\varepsilon \le 1/m$ and in tori of side $\ge 96/m$. The condition (E.m2), $m_2^2 \varepsilon_0^2 \le 1$, gives $\varepsilon \le \varepsilon_0 \le 1/m_2 < 1/m$. And $\ell_j \ge \ell_0 \ge \max\{96/m, \ell_0^\natural\} \ge 96/m$. So (a)–(d) apply to every lattice of the family. □

Mass ranges. Four ingredients of the proof of Theorem 5.2 are used in the transfer of Lemma I.24 at the mass $M_1 = 0.7m$, which is not one of the masses of Theorem 5.2. The next lemma shows that each of them holds at every $M \in (0, m_2]$, by its own proof. Here $t = t_k = L^k\varepsilon \le \varepsilon_0$ is the scale of the k-th step, a_k is Bałaban's averaging constant of that step, with $a_\flat \le a_k \le a$, and $\rho_{\min}(M) := \min(M, \mu_*) \min\{1, (L \max(M, \mu_*))^{-1}\}$.

Lemma A.6 (mass range of four ingredients)**.** *Let $(e,\lambda,m,\eta,m_1,m_2)\in\mathfrak{W}$ and $\mu_*=m$. Let the conditions (E.m2), (E.Φ), (E.fin) and $\ell_0\geq L^3$ hold. Let $M\in(0,m_2]$, with the upper bounds on ε_0 and the constants of the stages 8–10b that depend on the Higgs mass imposed at M. Then:*

(a) *(*the final c-numbers*) the torus-uniform bounds on $\delta m^2_{\mathrm{fin}}$ and on $E_{\mathrm{fin}}/|T|$ of Propositions E.9 and E.21 hold at Higgs mass M. They hold with $\bar\ell(M):=\min(1/M,1/\mu_*,L)$ in place of $\bar\ell(m_i)$ and with constants that are polynomials in $1/\rho_{\min}(M)$. Their dependence on ℓ_T is through $Z_f=t_{\bar K}/(m_f^2\ell_T^3)$, $m_s=M$, $m_v=\mu_*$, which is non-increasing in ℓ_T;*

(b) *(*the $A\cdot A$ vertices*) the bounds on the $A\cdot A$ vertices of* [3, (3.30)] *hold at Higgs mass M with the same constants Γ_3,Γ_3', uniformly in $M\in(0,m_2]$, and the bound on the vertex of* [3, (3.32)] *holds at M with a constant $\Gamma_4(M)<\infty$ that depends on M only through $m^2_{j_0}\leq 1$ and the bound of (a) (Lemmas E.24 and E.32);*

(c) *(*the tail bound*) the two-sided tail bound of Proposition E.36 holds at Higgs mass M, with a constant $C'_{579}(M)<\infty$ that depends on M only through the δm^2 vertices, that is, through the bound of (a);*

(d) *for every real background A, every domain Ω with Neumann Laplacian $-\Delta^N_{A,\Omega}\geq 0$ and every $Q_k=Q_k(A)$ with $Q_kQ_k^*=I$, the operator $\Delta^{(k)}(\Omega,A):=a_k-a_k^2Q_kG_k(\Omega,A)Q_k^*$, $G_k:=(-\Delta^N_{A,\Omega}+M^2t^2+a_kQ_k^*Q_k)^{-1}$, satisfies*

$$\frac{a_kM^2t^2}{a_k+M^2t^2}\ \leq\ \Delta^{(k)}(\Omega,A)\ \leq\ a_k,$$

and $\gamma_1:=a_k/(a_k+M^2t^2)\geq a_\flat/(a+1)$ under (E.m2).

At the four masses used by the proof, $M\in\mathbb{M}_4$, the upper bounds and constants of the hypothesis are those of the four-mass rule.

Proof. (d) Since $-\Delta^N_{A,\Omega}\geq 0$ and the inverse is operator monotone, $G_k\leq(M^2t^2+a_kP)^{-1}$ with $P:=Q_k^*Q_k$. Since $Q_kQ_k^*=I$, P is an orthogonal projection, and $(M^2t^2+a_kP)^{-1}=P/(M^2t^2+a_k)+(1-P)/(M^2t^2)$. Also $Q_kP=Q_k$, so $Q_k(1-P)=0$ and $Q_kG_kQ_k^*\leq Q_kPQ_k^*/(M^2t^2+a_k)=(M^2t^2+a_k)^{-1}$. Hence $\Delta^{(k)}\geq a_k-a_k^2/(a_k+M^2t^2)=a_kM^2t^2/(a_k+M^2t^2)$. The upper bound follows from $G_k\geq 0$. Finally $M^2t^2\leq m_2^2\varepsilon_0^2\leq 1$ by (E.m2) and $t\leq\varepsilon_0$, so $\gamma_1\geq a_\flat/(a+1)$. Only $M>0$ was used.

(a) The construction of Propositions E.9 and E.21 slices both fields, scalar and photon, up to one common index $\bar K$ with $t_{\bar K}\leq\bar\ell(M)$. Every renormalised subgraph then has internal lines of index below $\bar K$ only, and the ratio of the two masses enters only through the top cluster, as the number $\rho_{\min}(M)$. The proof nowhere uses the order of M and μ_*: the light field, which is the scalar when $M<\mu_*$, carries its long remainder line in the top cluster in the same way as the photon does when $M>\mu_*$. Its hypotheses are $\varepsilon_0\leq\bar\ell(M)$, which follows from (E.fin) because $\bar\ell(M)\geq\bar\ell(m_2)$ for $M\leq m_2$, and $\ell_T\geq L^3$, which does not involve the mass. Its constants are of three kinds. C_b does not depend on M. C_b^R is a polynomial in $1/\rho_{\min}(M)$ and in $\kappa_{R,\min}^{-1}$, $\kappa_{R,\min}=\min(\rho_{\min},\kappa_G)/8$. The remaining factors are $\max(LM,L\mu_*,1)$, $\log_L^+(\bar\ell/\varepsilon_0)$ and Z_f. Each is finite at every $M>0$, so the statement holds at M with these constants. The line bounds that carry this argument are those of Lemma E.39 and Theorem E.47. The renormalised subgraphs are bounded by the renormalisation machinery of Lemmas E.51, E.52, E.55, Theorem E.53 and Proposition E.54, and their constants are among the three kinds just listed.

(b) The constants Γ_3,Γ_3' depend on the Higgs mass only through the rescaled mass $m^2_{j_0}=M^2t^2_{j_0}$. For $M\leq m_2$, $t_{j_0}\leq\varepsilon_0$ and (E.m2), this lies in $[0,1]$, and the constants are uniform there. The range $M\in[m_1,m_2]$ in the statement of Lemma E.24 is not used in its proof. The constant Γ_4 of [3, (3.32)] is that of Lemma E.32. It also depends on M through the finite mass vertices of the classes, and by (a) these are bounded at every $M\in(0,m_2]$. Uniformity as $M\downarrow 0$ is not claimed,

and the proof uses Γ_4 only at the four masses of $\mathbb{M}_4$. The bound on $\Gamma_4(M)$, with this dependence on M, is proved in steps (iv)–(v) of the proof of Lemma E.32.

(c) The constant C'_{579} depends on M only through the δm^2 vertices of the step functionals, and these are bounded at M by (a). None of its other arguments ($\bar{n}$, $\bar{N}$, α_0, d, L, a, M_b, R, r, e, λ, μ_*, b_0, p_0, c_1, δ_0, η) depends on M. The tail bound is Proposition E.36. Its proof rests on the remainder bounds of Proposition E.57, the weighted bounds of Lemma E.58, and the family bound of Definition E.59 at the order cap (Proposition E.60). In these statements the propagators carry M only through $M^2t^2 \in [0,1]$, and their bounds are uniform there.

In (a)–(c) the dependence on u is unchanged. The transfer uses $u = 0$, which lies in $[-\eta, 1]$. Every factor $|u|$ of the constants at $u = 0$ is at most its value at the points $(u, M) \in \{(1, m_2), (-\eta, m_1)\}$ of Theorem 5.2. □

A.6. **The lemmas used in the proof of Lemma A.5.** Throughout this subsection $L = 2$, $a = 1$, $a_\flat = \frac{3}{4}$, $p = 6$, $\mu_* = m$, and $C_\varphi := 2^{3/4}(1 + \log 2)^6 c_1 = 39.622\ldots\, c_1$, where c_1 is Bałaban's constant of stage 8.

Lemma A.7 (monotonicity)**.** *Let $a > 0$ and $q \geq 0$.*

(a) *The function $f(t) := t^a \ell(t)^q$ is increasing on $(0, \mathrm{e}^{1-q/a}]$. Hence, for $\varepsilon_0 \leq \mathrm{e}^{1-q/a}$, "$f(t) \leq c$ for all $t \in (0, \varepsilon_0]$" holds if and only if $f(\varepsilon_0) \leq c$.*

(b) *Let $\varrho > 1$ and $\delta R > 0$, and put $g(t) := t^{-a} \exp(-\delta R\, \ell(t)^\varrho)$. Then g is increasing on $(0, t_g]$, where $t_g := \exp\big(1 - (a/(\delta R \varrho))^{1/(\varrho-1)}\big)$, and $g(t) \to 0$ as $t \downarrow 0$.*

(c) *Let $F : (0,1] \to [0,\infty)$ satisfy $F(t) \to 0$ as $t \downarrow 0$, and let $c > 0$, where neither F nor c depends on ε_0. Consider the condition "$F(t) \leq c$ for all $t \in (0, \varepsilon_0]$". If it holds at ε_0, it holds at every $\varepsilon'_0 \leq \varepsilon_0$. Moreover $\theta := \sup\{s \in (0,1] : F \leq c \text{ on } (0,s]\} > 0$, and the condition holds for every $\varepsilon_0 < \theta$. The same is true for the reciprocal form "$G(t) \geq c$ for all $t \in (0, \varepsilon_0]$" with $G(t) \to \infty$.*

Proof. (a) Since $\ell'(t) = -1/t$, $f'(t) = t^{a-1}\ell(t)^{q-1}\big(a\ell(t) - q\big)$. This is > 0 when $\ell(t) > q/a$, i.e. when $t < \mathrm{e}^{1-q/a}$. (b) $\frac{d}{dt}\log g = \big(-a + \delta R \varrho\, \ell(t)^{\varrho-1}\big)/t$. This is > 0 when $\ell(t)^{\varrho-1} > a/(\delta R\varrho)$, i.e. when $t < t_g$. Also $\log g = a(\ell(t) - 1) - \delta R \ell(t)^\varrho \to -\infty$ as $\ell(t) \to \infty$, because $\varrho > 1$.

(c) The condition at $\varepsilon'_0 \leq \varepsilon_0$ is a statement about the smaller set $(0, \varepsilon'_0] \subset (0, \varepsilon_0]$. Since $F(t) \to 0 < c$, there is $s > 0$ with $F \leq c$ on $(0, s]$, so $\theta > 0$. If $\varepsilon_0 < \theta$, then $\varepsilon_0 \leq s$ for some s in the set defining θ. The reciprocal form is the same argument applied to $1/G$. □

Lemma A.8 (the types of the conditions)**.** *Let $\pi = (e, \lambda, m, \eta, m_1, m_2) \in \mathfrak{W}$, and choose the data in the order of the stages 0–11 of Table 3. Every constant that depends on ε_0 and enters a condition of stage 10b is replaced by its supremum over $(0, \varepsilon_1]$ (Lemma A.11), and every constant that depends on the side of the torus is evaluated at $\ell_0^\natural$ (Lemma A.10). Then:*

(i) *Each of the conditions of Tables 5–8 is of type (ii) (a finite lower bound on R, b_0 or ℓ_0), type (iii) (a downward-closed upper bound on ε_0 with a positive threshold) or type (iv) (divisibility) of the proof of Lemma A.5. The bound or threshold depends only on data of earlier stages and is finite, respectively positive, at every point of $\mathfrak{W}$.*

(ii) *The definition of the family $T_j = j!T_0$, $\varepsilon = 2^{-K}\varepsilon_0$, $K \geq n_0 + 1$, imposes no condition on the regulator data, and $\varepsilon \leq 1/m$ follows from (E.m2).*

(iii) *No condition bounds b_0 from above before ε_0 is chosen, and no condition bounds ℓ_0 from above. The only conditions that bound R from above at fixed ε_0 are (E.2) and the second part of it, $r(\varepsilon_0) \leq Cp(\varepsilon_0)$; (E.2) is an upper bound on ε_0, and $r(\varepsilon_0) \leq Cp(\varepsilon_0)$ follows from the lower bound (b.2), $b_0 \geq R/C$; both are imposed after R. No condition involves ε, K or ℓ_T except through $\ell_0^\natural$ and the lower bounds of stage 11.*

Proof. Reduction to Lemma A.7(c). After stage 10a, every constant that enters an upper bound on ε_0 is a function of the data of the stages ≤ 9, of π, of η and of $\ell_0^\natural$, but not of ε_0 (Lemma A.11).

Every upper bound on ε_0 of Table 7 is either an explicit bound $\varepsilon_0 \le \theta$, or a condition "for all $t \in (0, \varepsilon_0]$". The second kind is reduced by Lemma A.7(a),(b) to its value at $t = \varepsilon_0$ plus an explicit bound. Its left side is built from $t^a \ell^q$ and $t^{-a}\exp(-\delta R \ell^\varrho)$, or is a finite product or sum of these, so it tends to 0; in the reciprocal form the left side tends to ∞. By Lemma A.7(c), such a condition is of type (iii).

Condition by condition. The thresholds of (E.fin), (E.m2), (E.Φ), (E.dom), (E.x) and (E.δ) are explicit and positive at every point of $\mathfrak{W}$, since $m, m_2, \lambda > 0$ and $e \neq 0$.

conditions	form	type; threshold
(E.fin)	$\varepsilon_0 \le \min(1/m_2, 1/\mu_*, L)$	(iii); $\min(1/m_2, 1/m, 2)$
(E.m2)	$m_2^2 \varepsilon_0^2 \le 1$	(iii); $1/m_2$
(E.Φ)	$\lambda^{-1/4} m_2^2 \varepsilon_0^{7/4} \le a_\flat / C_\varphi$	(iii); $(\lambda^{1/4} a_\flat / (C_\varphi m_2^2))^{4/7}$
(E.dom)	$\mu_* \varepsilon_0 \le 0.2$	(iii); $0.2/m$
(E.x)	$\lvert e\rvert \varepsilon_0^{1/2} \le \min\{c_v, e_*, 1\}$, $\lambda\varepsilon_0 \le c_v$	(iii); $\min\{(\min\{c_v, e_*, 1\}/\lvert e\rvert)^2,\ c_v/\lambda\}$
(E.δ)	$\kappa \mu_*^2 \varepsilon_0^2 \le 1/12$	(iii); $(12\kappa m^2)^{-1/2}$
Table 5	lower bounds on R; with $R \ge R^{**}$	(ii) at stage 5; data of the stages 0–4 only (Lemma A.9(i))
(E.4), (E.5), (E.10), (E.14), (R.cap)	left sides of the form $\varepsilon_0^{-a} \mathrm{e}^{-\delta R \hat\ell^2}$, or $R\hat\ell^2 \ge c$	(iii) by Lemma A.7(b),(c); the upper bound in (E.10) is void at $\delta_0 R \ge 4$
(E.ksp1)–(E.ksp3)	the conditions of Lemma A.17	(iii); for $t \le \varepsilon_0$ the left sides tend to 0
(E.1)–(E.3), (E.6)–(E.9), (E.11)–(E.13), (E.15), (E.r), (E.comb), (E.vac)	as in Table 7, at each $M \in \mathbb{M}_4$	(iii) (Lemma A.7(a),(c)); (E.6) and (E.7) in the reciprocal form
(b.1)–(b.4)	lower bounds on b_0	(ii) at stage 9
Table 8	lower bounds on ℓ_0; $\ell_0 \in 2M_b\mathbb{N}$	(ii) and (iv) at stage 11

For (E.Φ) note that the left side $\lambda^{-1/4} m_2^2 \varepsilon_0^{7/4}$ is increasing in ε_0, so the condition holds exactly up to the displayed threshold. For (E.vac), Theorems 4.1 and 4.2 hold at each fixed Higgs mass $M > 0$ for all sufficiently small ε_0; with the four masses fixed at stage 0, this gives four positive thresholds.

Every threshold in the table depends only on data of earlier stages: on π, κ and $\ell_0^\natural$ (stage 0), on Bałaban's constants (stages 3–4), on R (stage 5), on p (stage 6), on the constants of the stages 7–8, and on b_0 (stage 9). This proves (i) and (ii).

(iii) (E.2), $\hat x b_0 R \hat\ell^8 \le c$ with $\hat x = \lvert e\rvert \varepsilon_0^{1/2}$, bounds R from above at fixed ε_0. It is imposed at stage 10b, after R, as an upper bound on ε_0. Its second part, $r \le Cp$, follows from the lower bound $b_0 \ge R/C$, which is imposed at stage 9, after R. All other conditions on R, b_0 and ℓ_0 in the tables are lower bounds. The index K and the spacing ε enter no condition, because every scale used has $t_k \in (0, \varepsilon_0]$ and the conditions are statements for all such t. The side of the torus enters only through $\ell_0^\natural$ and the bounds of stage 11. □

Lemma A.9 (R is free of the masses; the four-mass rule)**.** *Let $\pi \in \mathfrak{W}$.*

(i) *R^*, and hence $R^{**} = \max\{R^*, 4\ell_{\bar N} + 32M_b + 25\}$, are functions of $(d, a, L, M_b, R_0, \delta_0, \bar n, \bar N, V(\bar N), C_{\mathrm{loc}}(\bar N))$, of the $O(1)$ localisation widths and of b_d; no Higgs mass, coupling or photon mass enters. The remaining lower bounds on R (R_0+1, $4/\delta_0$, and those from (E.10), (E.14) and (R.cap)) are also free of the Higgs mass. The bound from (E.14) contains only κ, μ_* and $(1-\eta)^{-3}$, which are finite on $\mathfrak{W}$. So $R \ge R^{**}$ is of type (ii).*

(ii) *Each entry of the Higgs mass M in a constant or condition of the stages 8–11 is of one of two types. It is of type (α) if M enters from above through M^2t^2 with $t \le \varepsilon_0$; then (E.m2) gives $M^2t^2 \le m_2^2\varepsilon_0^2 \le 1$ for every $M \le m_2$. It is of type (β) if M enters through powers of either sign and logarithms of M, and through quantities finite at every fixed $M > 0$ (among them $C_\delta(M)$, $1/\rho_{\min}(M)$, $(M^2\ell_T^3)^{-1}$, $(M\ell_T)^{-10}$, $\log_+(1/(M\varepsilon_0))$ and the factor $(\gamma_1'' M^2)^{-1}$*

of (E.5)); such entries occur only in upper bounds on ε_0, in monotone constants of the stages 8–9 and in the floor. No entry orders M against μ_*, and none uses $m_1 > 2m$ or $m_1 \geq \sqrt{2}\mu_*$.

(iii) *Consequently the four-mass rule can be met without creating a cycle, and each condition keeps its type.*

Proof. (i) The data on which R depends lie in the stages 0–4, and the symbol M in the lower bounds on R is Bałaban's block size M_b, not a mass. The bound $4\ell_{\bar{N}} + 32M_b + 25$ is free of the masses by inspection. The bound from (E.14) has as its only parameters κ, μ_* and $(1-\eta)^{-3}$; these are finite because $\mu_* = m$ and $\eta < 1$.

(ii) The ingredients of the proof of Theorem 5.2 fall into the class (α), into the class (β), or among those free of the mass, as follows. (α) or free of the mass: the one-variable and two-variable remainder bounds (Lemmas E.116 and E.117); the cube remainder (Lemma E.121); the separation lemma (Lemma E.120); the two-domain bounds (Lemma E.123); Lemmas E.124 and E.125; Lemmas E.100, E.102, E.103 and the support hypothesis (R4‴); the $A{\cdot}A$ family, whose Γ_3, Γ_3' depend on $M^2 t_{j_0}^2 \leq 1$ only; the order cap (Proposition E.76); the region entropy (Proposition 5.21); and (E.md), whose left sides increase in M. (β): the tail bound (Proposition E.36), through the δm^2 vertices; Proposition E.71, which needs only $-\Delta_A + M^2 > 0$; the final c-numbers (Proposition E.162: $\bar{\ell} = \min(1/M, 1/\mu_*, L)$, $\rho_{\min}(M) \geq \min(M, \mu_*) \min\{1, (L \max(M, \mu_*))^{-1}\}$, constants polynomial in $1/\rho_{\min}$, $Z_f = t_{\bar{K}}/(m_f^2 \ell_T^3)$); Lemma A.14 and the scalar gap paid by (E.5) at the smallest mass; the scalar zero mode of Lemma A.12, with gap $\geq M^2(1 + 4M^2\varepsilon_0^2/(3a))^{-1}$; $C_\delta(M)$, finite for every $M > 0$; C_b', which contains $M + \mu_*$, $(M^2\ell_T^3)^{-1}$, $(M\ell_T)^{-10}$ and $\log_+(1/(\min(M, \mu_*)\varepsilon_0))$; and C_{F1}, $\mathfrak{E}_k$, which are inherited from the first inductive step at mass M. No entry of a third type occurs: orderings of M against μ_*, and the conditions $m_1 > 2m$, $m_1 \geq \sqrt{2}\mu_*$, occur only in the anchor condition (6.3) of Proposition 6.7, that is, in the conditions of Table 4. The one place where the ordering of the two fields changes is the bound of the final c-numbers at $M_1 < \mu_*$. It is covered by the symmetric construction of Proposition E.162 with a common top index; at $(M_1, \mu_*) = (0.7, 1)m$ one gets $\bar{\ell} = 1/m$ and $\rho_{\min} \geq 0.35$, and the condition $\varepsilon_0 \leq \min(1/M_1, 1/\mu_*, L)$ follows from (E.fin).

(iii) Each listed constant is finite at each $M \in \mathbb{M}_4$ by (ii). The maximum of four finite numbers is finite, and the minimum of four positive thresholds is positive. A larger constant at the stages 8–9 only raises b_0 and Θ, which are lower bounds, and no condition bounds b_0 from above before ε_0 (Lemma A.8(iii)). Each condition of stage 10b stays a downward-closed upper bound at the fixed (R^{**}, b_0, p) (Lemma A.7). The conditions (E.m2) and (E.Φ) at m_2 imply the same conditions at every smaller M, because their left sides increase in M. The masses are data of stage 0, so the four-mass rule adds only edges from stage 0 into the stages 8–11. Adding data with edges forward to an acyclic graph whose edges do not decrease the stage keeps it acyclic. □

Lemma A.10 (the torus floor)**.** *Fix $\ell_0^\natural = \max\{8, 128/m_1, 128/M_1\}$ at stage 0, and evaluate at $\ell_T = \ell_0^\natural$ every constant that depends on ℓ_T. These are: $C_\delta(M)$ and its majorants $\bar{E}(M, \ell)$; the loss in the extension of* [2, Prop. 2.1] *to $Q^{(k)}$ (Lemma A.17); the zero-mode terms $(M^2\ell_T^3)^{-1}$, $(\mu_*^2\ell_T^3)^{-1}$, ℓ_T^{-1}, $(M\ell_T)^{-10}$, $(\mu_*\ell_T)^{-10}$ of the bound on δm_{fin}^2; the photon zero mode and the tail of Q_0; the twist defect $\mathrm{e}^{-\delta\ell_T/t}$; the zero-mode constants Z_s, Z_v of Proposition E.162; and, through these, $c_C^{(0)}$, c_L, c_7, $C_>$. Then the value at $\ell_0^\natural$ bounds the constant on every torus of side $\ell_T \geq \ell_0^\natural$. Hence, with $\ell_0 \geq \ell_0^\natural$ chosen at stage 11, every upper bound on ε_0 decided at stage 10 remains true on every torus T_j, and no upper bound on ε_0 depends on ℓ_0. Moreover:*

(a) *$\ell_0^\natural = 128/M_1 = 182.857\ldots/m$ whenever $M_1 \leq 16$ (in the units of Section 2). Replacing the floor $\max\{8, 128/m_1\}$ by $\ell_0^\natural$ can only weaken the upper bounds on ε_0.*

(b) *The constants that depend on ℓ_T are used only in the range $\varepsilon \leq 1/m$. Every spacing of the construction satisfies this, by (E.m2).*

Proof. Each listed constant is a non-increasing function f of ℓ_T. For C_δ every majorant, including the zero-mode terms $64\mathrm{e}^{-\mu^2\ell^2/8}/(\mu^2\ell^3)$ and $a^{-2\alpha}b^{-2\beta}/\ell^3$, decreases in ℓ. The terms of the bound on $\delta m^2_{\mathrm{fin}}$ decrease; its logarithmic term uses $\ell^* = \min(\ell_T, \max(1/M, 1/\mu_*))$ and is bounded by $\log_+(1/(\min(M,\mu_*)\varepsilon_0))$ for every ℓ_T. The terms $(\mu_*^2|T|)^{-1}$, $\mathrm{e}^{-c|T|}$ and $\mathrm{e}^{-\delta\ell_T/t}$ decrease. Also Z_f decreases, and $C_>$ depends on ℓ_T only through its $\delta m^2_{\mathrm{fin}}$ and C_δ entries. Hence $f(\ell_T) \le f(\ell_0^\natural)$ for $\ell_T \ge \ell_0^\natural$.

Every upper bound on ε_0 has the form $G(\varepsilon_0; \text{constants}) \le c$ and is monotone in each constant in the direction "larger constant, stronger condition"; this is read off Table 7, and in particular holds for (E.7), (E.ent) and the conditions of Lemma A.17. So replacing the constants by upper bounds keeps the conditions sufficient. The tori used have sides $\ell_j = j!\ell_0 \ge \ell_0 \ge \ell_0^\natural$ ((T.floor)).

(a) Since $M_1 < m_1$, we have $128/M_1 > 128/m_1$, and $128/M_1 \ge 8$ exactly when $M_1 \le 16$. So the new floor is at least the old one. By monotonicity, every constant evaluated at the floor is no larger at the new floor, and every upper bound decided with the old values stays true. The new floor also dominates the lower bounds on ℓ_0 tied to the masses ($M\ell_0 \ge 32$ at M_1, i.e. $\ell_0 \ge 45.71/m$; $96/m$; $128/m$; $\ell_T \ge L^3 = 8$). It is a single number computed from data of stage 0, so it creates no new dependence of earlier data on later data.

(b) The only other condition of validity of the constants that depend on ℓ_T is $\varepsilon \le \varepsilon_{\max} = 1/m$. (E.m2) gives $\varepsilon \le \varepsilon_0 \le 1/m_2 < 1/m$. □

Lemma A.11 (the bound ε_1 and the suprema of stage 10a)**.** *Let $\pi \in \mathfrak{W}$, and put*

$$\varepsilon_1 := \min\{1, \varepsilon_{(\mathrm{E.x})}, \varepsilon_{(\mathrm{E.md})}, \varepsilon_{(\mathrm{E}.\delta)}, \varepsilon_{(\mathrm{E.dom})}, \varepsilon_{(\mathrm{E.14})}\}.$$

Here $\varepsilon_{(\mathrm{E.x})} = \min\{(\min\{c_v, e_*, 1\}/|e|)^2, c_v/\lambda\}$, $\varepsilon_{(\mathrm{E}.\delta)} = (12\kappa\mu_*^2)^{-1/2}$, $\varepsilon_{(\mathrm{E.dom})} = 0.2/\mu_*$, $\varepsilon_{(\mathrm{E.md})} := \min\{1/m_2, (\lambda^{1/4}a_\flat/(C_\varphi m_2^2))^{4/7}\}$, the largest ε_0 that satisfies (E.m2) and (E.Φ), and $\varepsilon_{(\mathrm{E.14})} := \sup\{s \le 1 : \text{(E.14) holds at } s\}$. Then $\varepsilon_1 > 0$ is a function of data of the stages ≤ 9 and of π, and does not depend on ε_0 or K. Every constant that depends on ε_0 and enters a condition of stage 10b is bounded on $(0, \varepsilon_1]$ by a finite number free of ε_0. These constants are $C'_{2.3}$, γ_1, γ'_1, γ''_1, $\bar\delta$, c_K, C^w_{14}, C^w_P, the term $C_\delta(M)^2\varepsilon_0/(16\lambda)$ of c_L, C_{fin}, the constant $O_\pi(1)$ of [2, (3.42)], $C_6(e,\lambda)$, $\bar v$, $c_A^{LF} + \bar v$, and C_{old}.

Proof. Positivity. Each of $\varepsilon_{(\mathrm{E.x})}$, $\varepsilon_{(\mathrm{E.md})}$, $\varepsilon_{(\mathrm{E}.\delta)}$, $\varepsilon_{(\mathrm{E.dom})}$ is an explicit positive number, because $e_*, c_v, a_\flat, C_\varphi, \kappa > 0$ and $\lambda, m, m_2 > 0$ on $\mathfrak{W}$. The left sides of (E.m2) and (E.Φ) increase in ε_0, so $\varepsilon_{(\mathrm{E.md})}$ is the largest ε_0 that satisfies both. The left side of (E.14) increases in ε_0 and tends to 0 as $\varepsilon_0 \downarrow 0$, since $\hat\ell \to \infty$ and $(1-\eta)^{-3} < \infty$. Hence $\varepsilon_{(\mathrm{E.14})} > 0$. It depends only on κ, μ_*, c_0^{ker}, c_d, δ_0, R, η, all of the stages 0–5. None of the data of ε_1 is ε_0 or K.

Bounds on $(0, \varepsilon_1]$. Each item is monotone in ε_0, or a polynomial in ε_0 with $\varepsilon_0 \le 1$. In detail:

- $C'_{2.3}$: its term $c_1\mu_* L\varepsilon_0/a_k$ is at most $\frac{4}{3}c_1\mu_* L\varepsilon_1$, because $a_k > a_\flat = \frac{3}{4}$.
- $\gamma_1 \ge (1 + \mu_*^2\varepsilon_1^2/a_\flat)^{-1}$ and $\gamma'_1 \ge (1 + 4\mu_*^2\varepsilon_1^2/(3a))^{-1}$: these are decreasing functions of ε_0, bounded at ε_1.
- $\gamma''_1 = a_\flat/(a + M^2\varepsilon_0^2) \ge a_\flat/(a+1)$ for every $M \le m_2$, by (E.m2).
- $\bar\delta \le 1/12$ and $c_K \le 1/3$ are a priori bounds (Lemma A.19).
- C^w_{14}, C^w_P are polynomials $\sum_i a_i\varepsilon_0^i$, so $\sup_{(0,\varepsilon_1]}|\cdot| \le \sum_i|a_i|$ since $\varepsilon_1 \le 1$.
- $C_\delta(M)^2\varepsilon_0/(16\lambda) \le C_\delta(M)^2\varepsilon_1/(16\lambda)$, with $C_\delta(M)$ evaluated at $\ell_0^\natural$ and at each $M \in \mathbb{M}_4$ (Lemmas A.9, A.10).
- $C_{\mathrm{fin}} \le c_7\varepsilon_1^{\kappa_0} + c_5 + \mathrm{e}^{-4}$ (with $\kappa_0 = \frac{1}{5}$), where c_5 and c_7 are bounded on $(0, \varepsilon_1]$ by Statement K.1. (The form $c_7\varepsilon_1$ would hold only for $\kappa_0 \ge 1$.)
- Under (E.ent), $O_\pi(1)$ is the constant C_{fin} of Proposition 5.21, and it is bounded by the previous item. $C_6(e,\lambda)$ is bounded on $(0,\varepsilon_1]$, since its dependence on ε_0 is through monotone quantities.

- $\bar v$ and $c_A^{LF}+\bar v$ are bounded on $(0,\varepsilon_1]$; they are used only in the alternative large-field route of Remark F.30.
- $C_{\rm old}\le \bar S\, C_{PC_e}(1)(1+L^2/12)^{\bar N_{\rm ph}}$, a bound obtained through (E.δ), which is part of ε_1.

After this replacement, every constant that enters a condition of stage 10b is a function of the stages ≤ 9, of π, of η and of $\ell_0^\natural$, and not of ε_0. This is the hypothesis "F, c independent of ε_0" of Lemma A.7(c). □

A.7. **The conditions named in the tables.** A condition "for all $t\in(0,\varepsilon_0]$" is reduced to $t=\varepsilon_0$ by Lemma A.7(a),(b). In this subsection M_1 is always the anchor mass $0.7m$.

Lemma A.12 (the scalar zero mode)**.** *Work in physical units on the torus T_ε, with the divisibility of Lemma 2.1(ii). Put $\Delta_s^{(0)}:=-\Delta^\varepsilon+M^2$ and $\beta_k:=a\,t_{k+1}^{-2}$, and define recursively*

$$C_s^{(k)}:=(\Delta_s^{(k)}+\beta_k Q^*Q)^{-1},\qquad \Delta_s^{(k+1)}:=\beta_k-\beta_k^2\,Q\,C_s^{(k)}Q^*,$$

where Q is Bałaban's scalar block average at background $A=0$. Then for every $k\le K$ and every $M>0$,

$$\Delta_s^{(k)}\ \ge\ \Big(M^{-2}+a^{-1}\sum_{j<k}t_{j+1}^2\Big)^{-1}\ \ge\ M^2\Big(1+\frac{4M^2\varepsilon_0^2}{3a}\Big)^{-1},$$

uniformly in ℓ_T and ε. The first bound is attained on the constant mode, i.e. it is the eigenvalue on the constants. At $L=2$, $4/(3a)=1/a_\flat$. (The photon form is Lemma F.37.)

Proof. By the Woodbury identity, $(Q(\Delta_s^{(k)})^{-1}Q^*+\beta_k^{-1})^{-1}=\beta_k-\beta_kQ(\Delta_s^{(k)}+\beta_kQ^*Q)^{-1}Q^*\beta_k=\Delta_s^{(k+1)}$. Hence $(\Delta_s^{(k+1)})^{-1}=Q(\Delta_s^{(k)})^{-1}Q^*+\beta_k^{-1}$ exactly.

Q is an average of point values whose weights sum to 1. On T_ε the blocks tile every fine site with total weight L^{-d}. The Schur test with weights $t_{k+1}^d=L^dt_k^d$ therefore gives $\|Q\|\le1$ and $Q^*1=1$.

So $\|(\Delta_s^{(k+1)})^{-1}\|\le\|(\Delta_s^{(k)})^{-1}\|+\beta_k^{-1}$, with $\|(\Delta_s^{(0)})^{-1}\|=M^{-2}$, the constant mode included. Iterating gives the first bound. For the second, $\sum_{j<k}t_{j+1}^2\le\frac43t_k^2\le\frac43\varepsilon_0^2$ at $L=2$.

On constants, $Q1=1$, $Q^*1=1$ and $\Delta_s^{(0)}1=M^21$. By induction $\Delta_s^{(k)}1=\lambda_s(k)1$ with $1/\lambda_s(k+1)=1/\lambda_s(k)+1/\beta_k$, which is the claimed equality. □

Remark A.13. (i) The gap is positive for every ε_0, so the scalar zero mode by itself imposes no upper bound on ε_0. Under (E.m2), $m_2^2\varepsilon_0^2\le1$, every $M\in\mathbb{M}_4$ (all are $\le m_2$) has gap $\ge M^2(1+4/(3a))^{-1}$, which is $3M^2/7$ at $a=1$; in particular it is finite and positive at M_1. (ii) Bałaban's papers [1, 2] are themselves set on the torus, with constants independent of ε and T_ε [1, (1.14), p. 606]. Lemma A.12 gives, with an explicit gap free of ℓ_T, what that setting already contains. (iii) The Gaussian factor of the constant mode is of the form $(2\pi/(M^2\varepsilon^d))^{N/2}$ times a Jacobian. It sits in $E_{0,s}(M)$ and cancels.

Lemma A.14 (lower bound on the averaged mass operator for every real background)**.** *Work in units of scale k, $t=L^k\varepsilon$. Let A be any real background and Ω any domain, with Neumann operator $-\Delta^N_{A,\Omega}\ge0$. Let $Q_k=Q_k(A)$ satisfy $Q_kQ_k^*=I$. Put $G_k(\Omega,A):=(-\Delta^N_{A,\Omega}+M^2t^2+a_kQ_k^*Q_k)^{-1}$ and $\Delta^{(k)}(\Omega,A):=a_k-a_k^2Q_kG_k(\Omega,A)Q_k^*$. Then for every $M>0$*

$$\frac{a_kM^2t^2}{a_k+M^2t^2}\ \le\ \Delta^{(k)}(\Omega,A)\ \le\ a_k.$$

Consequently, the scalar part of [2, (3.11), p. 585] *holds for every real external field, without any regularity assumption:*

$$G_k''\ \ge\ \gamma_1M^2t^2\ \text{ on }\Lambda_5^{(k)},\qquad \gamma_1=\frac{a_k}{a_k+M^2t^2}\ \ge\ \frac{a_\flat}{a_\flat+M^2t^2}\ \ge\ \frac{a_\flat}{a+M^2\varepsilon_0^2}=:\gamma_1''.$$

Proof. The two-sided bound is Lemma A.6(d). For the consequence: principal blocks inherit lower bounds, and the averaging terms of [2, (3.9)] are ≥ 0. The chain for γ_1 uses that $x \mapsto x/(x+c)$ is increasing, $a_k \geq a_\flat$ and $t \leq \varepsilon_0$. Only $Q_k Q_k^* = I$ is used. So Q_k may come from any real background, or be a composite of single steps at different backgrounds. □

Lemma A.15 (the bound on the Hessian and condition (E.5))**.** *Fix $M > 0$, and let k satisfy $t_k \leq \varepsilon_0$. Let V_k be the space of scalar fields on $\Lambda_5^{(k)}$ with $N = 2$ real components, so that $\dim V_k = N|\Lambda_5^{(k)}|$. Let $-\frac{1}{2}\langle \phi_k, (G_k'' - H_k'')\phi_k\rangle$ be the ϕ_k-quadratic part of the exponent in* [2, (3.9), p. 585]*, where G_k'' and H_k'' are taken symmetric. Assume $G_k'' \geq \gamma_1'' M^2 t_k^2$ on V_k with $\gamma_1'' = a_\flat/(a + M^2\varepsilon_0^2)$ (Lemma A.14, Lemma A.6(d)). Assume also $\|H_k''\| \leq h(t_k)$, where*

$$h(t_k) := C_{H''} \mathrm{e}^{-\delta' r(t_{k-1})} + C_C c_d(\delta/2)\mathrm{e}^{-\delta r(t_k)/2}$$

is the Schur bound of [2, (3.12)]*; the first term is absent at $k = 0$. Put*

$$C_H := C_{H''} + C_C c_d(\delta/2), \qquad \delta_H := \min\{\delta', \delta/2\}.$$

(a) *Suppose that*

$$h(t_k) \;\leq\; \tfrac{1}{2}\gamma_1'' M^2 t_k^2 \qquad \text{for every } k \text{ with } t_k \leq \varepsilon_0. \tag{A.1}$$

Put $\varrho_k'' := h(t_k)/(\gamma_1'' M^2 t_k^2)$ and $X_k'' := G_k''^{-1/2} H_k'' G_k''^{-1/2}$. Then $G_k'' - H_k'' \geq \frac{1}{2}\gamma_1'' M^2 t_k^2$, $\|X_k''\| \leq \varrho_k'' \leq \frac{1}{2}$,

$$[\det(G_k'' - H_k'')]^{-1/2} \leq (\det G_k'')^{-1/2} \exp\big((\ln 2)\varrho_k'' N|\Lambda_5^{(k)}|\big), \qquad |\ln\det(\mathbf{1} - X_k'')| \leq 2N\varrho_k''|\Lambda_5^{(k)}|,$$

and $\|(G_k'' - H_k'')^{-1} - G_k''^{-1}\| \leq 2\varrho_k''(\gamma_1'' M^2 t_k^2)^{-1}$. These are the bounds of [2, (3.17)–(3.20), pp. 586–587]*, and ϱ_k'' is the determinant constant used there.*

(b) *The condition (E.5) at M reads*

$$2C_H(\gamma_1'' M^2)^{-1}\varepsilon_0^{-2}\exp(-\delta_H R\hat{\ell}^2) \leq 1, \qquad \varepsilon_0 \leq \exp\big(1 - 1/(\delta_H R)\big).$$

It implies (A.1)*. It is downward closed in ε_0, and it holds for all ε_0 below a positive threshold that depends only on $C_H, \delta_H, R, a, L, M$. Its left side decreases in M, so at each ε_0 the condition is strongest at M_1.*

(c) *The scalar zero mode and the logarithmic entries of Lemma A.16 impose no condition on ε_0 beyond (E.m2) and (E.fin).*

Proof. (a) Write $g := \gamma_1'' M^2 t_k^2$ and $h := h(t_k) \leq g/2$. For $v \in V_k$, $\langle v, (G_k'' - H_k'')v\rangle \geq (g - h)|v|^2 \geq \frac{g}{2}|v|^2$. The operator X_k'' is symmetric, and $\|X_k''\| \leq g^{-1}\|H_k''\| \leq h/g = \varrho_k'' \leq \frac{1}{2}$. From $G_k'' - H_k'' = G_k''^{1/2}(\mathbf{1} - X_k'')G_k''^{1/2}$ we get $\det(G_k'' - H_k'') = \det G_k'' \det(\mathbf{1} - X_k'')$. Let $x_1, \dots, x_n$, $n = N|\Lambda_5^{(k)}|$, be the eigenvalues of X_k''; they lie in $[-\frac{1}{2}, \frac{1}{2}]$. The function $\varphi(x) := -\ln(1-x)$ is convex with $\varphi(0) = 0$. Hence $0 \leq \varphi(x) \leq 2(\ln 2)x$ for $0 \leq x \leq \frac{1}{2}$, and $|\varphi(x)| = \ln(1 + |x|) \leq |x|$ for $-\frac{1}{2} \leq x < 0$. So $|\ln\det(\mathbf{1} - X_k'')| \leq \sum_i |\varphi(x_i)| \leq (2\ln 2)n\varrho_k'' \leq 2n\varrho_k''$, and $[\det(\mathbf{1} - X_k'')]^{-1/2} = \exp(\frac{1}{2}\sum_i \varphi(x_i)) \leq \exp((\ln 2)n\varrho_k'')$. Finally, $(G_k'' - H_k'')^{-1} - G_k''^{-1} = G_k''^{-1}H_k''(G_k'' - H_k'')^{-1}$ has norm at most $g^{-1}h \cdot 2g^{-1} = 2\varrho_k''/g$.

The normalisation of (A.1) is the one used in [2, (3.17)–(3.20)]. The lower bound $\frac{1}{2}\gamma_1'' M^2 t_k^2$ requires $h \leq g/2$, and the determinant bound requires $\|X_k''\|$ to stay away from 1; both follow from $\varrho_k'' \leq \frac{1}{2}$. Measured against $G_k'' - H_k''$ instead of G_k'', the norm of H_k'' is at most $h/(g/2) = 2\varrho_k'' \leq 1$. This is the same condition, so no stronger form such as $h \leq \frac{1}{4}g$ (equivalently, (E.5) with $4C_H$) is needed.

(b) The function r is decreasing on $(0, 1]$, and $t_{k-1} < t_k$, so $r(t_{k-1}) \geq r(t_k)$. Hence $h(t_k) \leq C_H \mathrm{e}^{-\delta_H r(t_k)}$. Put $\psi(t) := C_H t^{-2}\mathrm{e}^{-\delta_H R\ell(t)^2}$ with $\ell(t) = 1 + \ln t^{-1}$, and put $s = \ln t^{-1}$. Then $\frac{d}{ds}\ln\psi = 2 - 2\delta_H R(1+s)$, which is ≤ 0 exactly when $t \leq t^* := \exp(1 - 1/(\delta_H R))$. So ψ is non-decreasing on $(0, t^*]$. By the second part of (E.5), $\varepsilon_0 \leq t^*$, so $\psi(t) \leq \psi(\varepsilon_0)$ for $t \leq \varepsilon_0$. The first part

of (E.5) says $\psi(\varepsilon_0) \le \frac12\gamma_1'' M^2$, and γ_1'' does not depend on t. Hence $h(t_k) \le \psi(t_k)t_k^2 \le \frac12\gamma_1'' M^2 t_k^2$, which is (A.1).

Now let $\varepsilon_0' \le \varepsilon_0$ with ε_0 satisfying (E.5). Then $\varepsilon_0' \le t^*$, $\psi(\varepsilon_0') \le \psi(\varepsilon_0)$ and $\gamma_1''(M,\varepsilon_0') \ge \gamma_1''(M,\varepsilon_0)$, so (E.5) holds at ε_0'. For $\varepsilon_0 \le 1$ we have $\gamma_1'' \ge a_\flat/(a+M^2)$. Moreover $\ln\psi(\varepsilon_0) = \ln C_H + 2s - \delta_H R(1+s)^2 \to -\infty$ as $s = \ln\varepsilon_0^{-1} \to \infty$. This gives the positive threshold. Finally, $(\gamma_1'' M^2)^{-1} = a/(a_\flat M^2) + \varepsilon_0^2/a_\flat$ is decreasing in M, and the other factors do not depend on M.

(c) By Lemma A.12, the scalar gap satisfies $\Delta_s^{(k)}(0) \ge \gamma_s(M;\varepsilon_0)M^2$ with $\gamma_s(M;\varepsilon_0) = (1 + 4M^2\varepsilon_0^2/(3a))^{-1}$, uniformly in ℓ_T and ε. This is not a condition on ε_0. Under (E.m2) and $a = 1$ it gives $\gamma_s \ge \frac37$ for $M \le m_2$. By Lemma A.16(b),(c), once $\varepsilon_0 \le \min(1/m_2, 1/\mu_*, L)$, which is (E.fin), the product $t^2|\delta m^2_{\rm fin}|$ is bounded for $t \le \varepsilon_0$ by a function of t alone that tends to 0 with t. So the factors $\log_L^+(\bar\ell/\varepsilon_0)$ and $\log_+(1/(\min(M,\mu_*)\varepsilon_0))$ add no condition on ε_0. Every condition in which $t^2\delta m^2_{\rm fin}$ enters has the form $\sup_{0<t\le\varepsilon_0} F(t) \le \theta$ with $F(t) \to 0$, and so it is downward closed with a positive threshold (Lemma A.7(c)). □

Lemma A.16 (the logarithmic entries)**.** (a) *In the proof of Theorem 5.2, the logarithmic mass entries are those of the bound on the final mass counterterm of Proposition E.16**2**. Put* $\bar\ell := \min(1/M, 1/\mu_*, L)$ *and assume* $\varepsilon_0 \le \bar\ell$*. Then*

$$|\delta m^2_{\rm fin}(x)| \le C_b\Big\{(e^2+\lambda)\tfrac{L^2}{(L-1)\varepsilon_0} + (e^4+e^2\lambda+\lambda^2)\big(2+\log_{L,+}(\bar\ell/\varepsilon_0)\big)\Big\} + C_b^R\{\dots\},$$

where $C_b = C(\alpha_0,\bar n,d,L,a,M_b)$ *and* C_b^R *is a polynomial in* $1/\rho_{\min}$ *(masses only). A weaker form of this bound carries* $\log_+(1/(\min(M,\mu_*)\varepsilon_0))$ *instead.*

(b) *These entries enter a condition or an error constant only through the rescaled vertex* $t_k^2\delta m^2(\cdot)$ *with* $t_k \le \varepsilon_0$*.*

(c) *For* $0 < t \le \varepsilon_0 \le \bar\ell$ *and every* $M > 0$*,*

$$t^2\log_{L,+}(\bar\ell/\varepsilon_0) \;\le\; \frac{\bar\ell}{\mathrm{e}\ln L}\,t \;\le\; \frac{L}{\mathrm{e}\ln L}\,t \quad \Big(= 1.0615\,t \text{ at } L=2\Big), \qquad t^2\log_+\frac{1}{\min(M,\mu_*)t} \le \frac{t}{\mathrm{e}\min(M,\mu_*)}.$$

In particular the logarithmic entries impose no condition beyond (E.fin).

Part (a) is the bound of Proposition E.9.

Proof of (c). $x\log(1/x) \le 1/\mathrm{e}$ for $x \in (0,1]$. With $x = t/\bar\ell$ and $\varepsilon_0 \ge t$, this gives $t\log_{L,+}(\bar\ell/\varepsilon_0) \le t\log(\bar\ell/t)/\ln L \le \bar\ell/(\mathrm{e}\ln L)$. Multiplying by t and using $\bar\ell \le L$ gives the first bound. The second follows in the same way with $x = \min(M,\mu_*)t$; if $x > 1$ the left side is 0. □

Lemma A.17 (the conditions (E.ksp1)–(E.ksp3) and (E.14))**.** *Put* $\vartheta(t) := |e|\,t^{1/2}p(t)$ *and* $P_1(t) := |e|\,c_1 C\mu_*^{-1}t^{-1/2}p(t)$*. Let* $c_{AX} := \sqrt{d}/(\mathrm{e}\,\delta_1)$*, with* δ_1 *fixed at stage 4. Let* C_B *be the constant of the bound on the block-averaging operators* $Q^{(k)}$ *at real backgrounds (constants of stage 3 and* c_1 *of stage 8). The extension of* [2, Prop. 2.1] *to* $Q^{(k)}$ [2, p. 579] *needs the following upper bounds on* ε_0*, each for all* $t \in (0,\varepsilon_0]$ *and each at stage 10b.*

(E.ksp1) $4c_{AX}C_B|e|\,b_0\varepsilon_0^{1/2}\hat\ell^p \le 1$*, together with* $\varepsilon_0 \le \mathrm{e}^{1-2p} = \mathrm{e}^{-11}$*. (E.8) does not imply (E.ksp1): the constants are unrelated.*

(E.ksp2) *With the capped radius* $R_y := \min\{\mathrm{dist}_j(y,\Omega^c)/2,\ r(t_k)L^{k-j}/2\}$*:*

$$\textit{(i)}\ \ (2d+3)\,\mathrm{e}^{-\delta_1 r(t)/4}P_1(t) \le \tfrac12; \qquad \textit{(ii)}\ \ \sqrt{d}\,C_B\,\mu_*\,t\,r(t) \le c_1 C.$$

As an alternative to (ii), one may keep the weight $\mathrm{e}^{-\delta_1 R_y/2}$ *and replace the factor* 4 *of (E.ksp1) by* 12*.*

(E.ksp3) $4C_B|e|^{1-\beta}t^{(1-\beta)/2}p(t) \le c$ *with* $\beta = 1/56$*. Equivalently, (E.1) with* $C_{\rm reg}$ *replaced by* $\max\{C_{\rm reg}, 4C_B/c_1\}$*, together with the bound* $\varepsilon_0 \le \mathrm{e}^{1-2p/(1-\beta)} = \mathrm{e}^{-11.22}$*.*

The condition (E.14) is used in the form of Table 7,

$$4\kappa c_0^{\ker} c_d (1+\mu_*^2\varepsilon_0^2)(1-\eta)^{-3}\,\frac{\mathrm{e}^{-\delta_0 R\hat{\ell}^2/4}}{1-\mathrm{e}^{-\delta_0 R\hat{\ell}\ln 2/2}} \le \frac{1}{12};$$

it enters stage 10a through $\varepsilon_{(\mathrm{E}.14)}$ *and is imposed at stage 10b. Each of these conditions has a left side that tends to* 0 *as* $\varepsilon_0 \to 0$*, is downward closed below its explicit bound, depends only on the stages* ≤ 9 *(*R *at stage 5), and does not bound* ℓ_T *from above. So the order of choices stays acyclic.*

Proof. (E.ksp1): the extension of [2, Prop. 2.1] needs $c_{AX}E_j \le \frac{1}{2}$ with $E_j \le 2C_B\vartheta(t_k)$. The function $t^{1/2}\ell^p$ is increasing on $(0,\mathrm{e}^{1-2p}]$, so $\sup_{t\le\varepsilon_0}\vartheta = |e|\,b_0\varepsilon_0^{1/2}\hat{\ell}^p$.

(E.ksp2): the corollary of that extension needs only $R_y \ge r(t_k)L^{k-j}/4$, and this holds because $\mathrm{dist}_k(Y,\Omega^c) \ge r(t_k)/2$. Then (i) follows from $\mathrm{e}^{-\delta_1 R_y/2} \le \mathrm{e}^{-\delta_1 r(t_k)/4}$. For (ii), $\sqrt{d}R_yE_j \le \sqrt{d}\,(r(t_k)L^{k-j}/2)(2C_B\vartheta_kL^{-2(k-j)}) \le \sqrt{d}\,C_Br(t_k)\vartheta(t_k)/2$. The sufficient condition $\sqrt{d}C_Br\vartheta \le P_1$ is equivalent to $\sqrt{d}C_B\mu_* t\,r(t) \le c_1C$. Since $t\,r(t) = Rt\ell(t)^2$ is increasing on $(0,\mathrm{e}^{-1})$ (condition (E.r)), (ii) has a left side that tends to 0.

(E.ksp3): regularity of $s(\tilde{B}_j+\tau a_j)$ in the estimate of the averaged fields requires $\sup|\mathcal{F}| \le 2E_j \le ce_j^\beta$; since $L^{-2m} \le L^{-\beta m/2}$ this gives the displayed condition. At $t=\varepsilon_0$ the left side equals $4C_B\hat{x}^{1-\beta}b_0\hat{\ell}^6$, which is the form of (E.1).

(E.14): the deletion distance at level j is $r(t_{j-1})/L \ge r(t_j)/L$, which gives the factor $\mathrm{e}^{-\delta_0 R\ell_j^2/4}$ at $\varrho = L = 2$. The sum runs over $t_j = 2^{-m}\varepsilon_0$ with $\ell_j = \hat{\ell}+m\ln 2$, and $(\hat{\ell}+m\ln 2)^2 \ge \hat{\ell}^2 + 2m\hat{\ell}\ln 2$. Summing the geometric series gives the displayed quotient. □

Lemma A.18 (the condition (E.ent))**.** *Let* c_5, c_7 *be the constants of* [2, (3.42)] *as used in Proposition 5.21, taken at their suprema over* $\varepsilon_0 \in (0,\varepsilon_1]$*, with* c_7 *at its maximum over* $\mathbb{M}_4$*. Let* $\Theta := C_1C_D^dR^d/(c_\square b_0^2\ln 2)$ *with* $C_1 := n_d\ln 2 + d + 1 + c_7 + c_5$ *(*$n_3 = 10$*),* $C_D := C_D^0 + 14(1+\vartheta)$*,* $C_D^0 := A'(S_{\varrho,L}-1)$*,* $A' = 16+23\vartheta$*,* $\vartheta := M_bL/R \le \frac{1}{7}$*,* $S_{\varrho,L} := \sum_{m\ge 0}L^{-m}(1+m\ln L)^\varrho$ *(*$S_{2,2} = 2+4\ln 2+6\ln^2 2 = 7.6553\ldots$*, so* $C_D \le 144.36$*), and* $c_\square := \min\{\frac{1}{8}a3^{-d}, \frac{1}{4}\gamma_0 3^{-d}, 3^{-d}\}$*. The condition at stage 10b is*

$$\textit{(E.ent)}\qquad (1+\ln(2/\varepsilon_0))^{2p-2-\varrho d} \ge \Theta \iff \varepsilon_0 \le \varepsilon_{\mathrm{ent}} := 2\exp\bigl(1-\Theta^{1/(2p-2-\varrho d)}\bigr)$$
$$= 2\exp(1-\Theta^{1/4}) \qquad (p=6,\ \varrho=2,\ d=3).$$

Its left side increases as ε_0 *decreases, so it is a downward-closed upper bound. If half of the gain must be kept for another use,* Θ *is replaced by* 2Θ*. Under (E.ent), Proposition 5.21 gives* [2, (3.42)] *on the torus.*

Proof. $(1+\ln(2/\varepsilon_0))^4 \ge \Theta$ holds iff $\ln(2/\varepsilon_0) \ge \Theta^{1/4}-1$, i.e. iff $\varepsilon_0 \le 2\exp(1-\Theta^{1/4})$. The base $1+\ln(2/\varepsilon_0)$ is the minimum of $\ell(t_j)$ over $j \le K-1$, because $t_j \le \varepsilon_0/2$ there. The bound $\vartheta \le \frac{1}{7}$ follows from $R \ge R^{**} \ge 32M_b > 7M_bL$. Since C_D is increasing in ϑ, its value $144.352\ldots$ at $\vartheta = \frac{1}{7}$ bounds it.

The suprema of c_5 and c_7 over $\varepsilon_0 \in (0,\varepsilon_1]$ are finite by Statement K.1. Hence Θ is a finite positive number determined by b_0 (stage 9), R (stage 5), constants of the stages 3 and 8, and ε_1 (stage 10a); it depends on no datum of the stages 10b or 11. So $\varepsilon_{\mathrm{ent}} > 0$, and (E.ent) is a downward-closed upper bound on ε_0 of stage 10b with a positive threshold, as used in Lemma A.4. □

Lemma A.19 (the condition (E.δ))**.** *Let* κ *be the domination constant of stage 0, a fixed number* $\kappa \ge \kappa_1\kappa_3$ *(Theorem I.16); under (E.dom),* $\mu_*\varepsilon_0 \le 0.2$*, every* $\kappa \ge 126528/16875$*, in particular* $\kappa = 7.5$*, is admissible (Remark I.18). The condition (E.*δ*) is*

$$\bar{\delta} := \kappa\mu_*^2\varepsilon_0^2 \le \tfrac{1}{12}, \qquad \textit{i.e.}\quad \varepsilon_0 \le \varepsilon_{(\mathrm{E}.\delta)} = (12\kappa\mu_*^2)^{-1/2},$$

and it enters the bound ε_1 of stage 10a. Consequences: (a) the slice weights satisfy $w_k = 1-\eta\delta_k \in [1-\eta\bar\delta, 1] \subset [11/12, 1]$, so $\eta\bar\delta \le \eta/12 < 1$ (Lemma 5.4(a)); (b) the coefficient $c_K := \kappa\mu_*^2 L^2 t_K^2$ of the top slice satisfies $c_K \le 1/3$. At $\mu_* = 1$ and $\kappa = 7.5$ the condition reads $\varepsilon_0 \le 90^{-1/2} = 0.10541$.

Proof. (a) $\delta_k = \kappa\mu_*^2 t_{k+1}^2 \le \kappa\mu_*^2\varepsilon_0^2 = \bar\delta \le \frac{1}{12}$, and $0 < \eta < 1$. (b) $t_K = \varepsilon_0$ and $L = 2$, so $c_K = L^2\bar\delta \le \frac{4}{12}$. □

Remark A.20. The value 6.07 is the limit of $\kappa_1\kappa_3$ as $\mu_*\varepsilon_0 \to 0$ and $K \to \infty$; since $\kappa_1\kappa_3 > 164/27 > 6.07$ (Remark I.18), it may not be used for κ.

Lemma A.21 (the conditions (R.cap) and (T.cap) at the order cap $\bar N$)**.** *At the order cap $\bar N = \bar n^2 = 169$ the conditions on the localisation cubes read*

$$\textit{(R.cap)}\ \ R\hat\ell^2 \ge C_{\rm loc}(\bar N), \qquad \textit{(T.cap)}\ \ \ell_T \ge 3r(\varepsilon_0)\varepsilon_0 + C_{\bar N}\varepsilon_0.$$

Here $C_{\rm loc}(\bar N)$ and $C_{\bar N}$ are finite numbers fixed at stage 4 together with $\bar N$. (R.cap) is met at stage 5 by $R \ge C_{\rm loc}(\bar N)$, since $\hat\ell \ge 1$, or else at stage 10b; (T.cap) is met at stage 11. The order of choices stays acyclic, free of ε and uniform over tori with $\ell_T \ge \ell_0$. If R is raised to meet (R.cap), the condition (E.2) makes ε_0 smaller (an effect that depends on $\bar N$ but not on ε).

Remark A.22. The decay rate of the trees of [3, (1.33)] at the cap is some $\delta(\bar N) > 0$ (or the rate $\delta_1/2$, free of the cap, extracted before the shrinking induction of Lemma E.51). It is not Bałaban's δ_0 of stage 3, which is the propagator rate used in $\delta_0 R \ge 4$.

A.8. **An explicit point.**

Example A.23 (an admissible point)**.** In units $m = 1$, let $(e^2, \lambda) = (0.05, 0.05)$, $\mu_* = 1$, $M_1 = 0.7$, $H = 1$, $\eta = \frac12$, $(m_1, m_2) = (3, 4)$, so that $\mathbb{M}_4 = \{0.7, 1, 3, 4\}$, with $L = 2$, $a = 1$, $a_\flat = \frac34$, $p = 6$, $\varrho = 2$, $\bar n = 13$, $\bar N = 169$. Choose, in the order of the stages:

- $R := \lceil \max\{R^{**}, R_0 + 1, 4/\delta_0, R_{(\text{E.10})}, R_{(\text{E.14})}, R_{(\text{R.cap})}\}\rceil$. Here $R_{(\cdot)}$ is the lower bound on R that makes the condition $(\cdot)$ hold for all $\varepsilon_0 \le 1$, if there is one; the bounds $R_{(\text{E.10})}$ and $R_{(\text{E.14})}$ also depend on κ, μ_* and η;
- $b_0 := \max\{b_0^{\min}, (\max\{2, \bar\mu\lambda^{-1/2}\})^{1/2}, R/C\}$, with every constant of the stages 8–9 that depends on the Higgs mass taken at its maximum over $\mathbb{M}_4$;
- $\varepsilon_0 := 2^{-n_0}$, where n_0 is the least integer such that 2^{-n_0} lies strictly below ε_1 and below the threshold of every upper bound of Table 7 at this R and b_0, at each mass of $\mathbb{M}_4$, (E.edge) included;
- ℓ_0 the least element of $2M_b\mathbb{N}$ that is $\ge \max\{183, 128, 96\}$ and above the lower bounds (T.nw), (T.cap), (T.T) and (T.per);
- $T_j = j!\,T_0$ and $\varepsilon = 2^{-K}\varepsilon_0$ with $K \ge K_0 = \max(K_0', n_0 + 1)$.

Then every condition of Tables 4–8 holds. Moreover, for every $j \ge 1$, every such K and every $k \le K$, the side $S_k := \ell_j/t_k$ of T_j in units of the scale-k blocks satisfies $S_k \ge \max\{6\sqrt{3}\,r(t_k) + 2,\ 2/\delta_0\}$, which is the hypothesis of Lemma 5.22.

Order. By Lemma A.4, every condition depends only on data of its own stage and of earlier stages, and it constrains only the quantity chosen at its stage. The choices above are made in increasing order of stages. So it suffices to check each condition at its stage.

The point. The conditions of Table 4 involve R, ε_0 and ℓ_0 only through "$\ell_T \ge 96$, $\varepsilon \le 1$", and they hold by Corollaries G.4, G.7 and G.9: $a_{K,T}(1) - a_{K,T}(0.7) \ge 0.3190043$; $g_{21} \ge (4-3)(4+3-0.3631) = 6.6369$; $m_1 = 3 > 2 > 1.882 \ge m_1^c$; $m_1 = 3 \ge \sqrt{2}\,\mu_*$, $m < m_1 < m_2$, $M_1 = 0.7 < 1$; and $\Pi(0.05, 0.05) = 0.4728641 > 0$. They are infima over all tori of side ≥ 96 (respectively ≥ 64). Every torus of the family has side $\ell_j \ge \ell_0 \ge 183$, and every spacing satisfies $\varepsilon \le \varepsilon_0 \le \frac14$. So these conditions hold on every T_j.

R and b_0. $R \ge R^{**}$ gives every condition of Table 5, (R.edge) included (Lemma A.3(i)). (R.nbar) reads $27 \le 3R$ and holds since $R \ge 107$. Since $\delta_0 R \ge 4$, the part of (E.10) that bounds ε_0 imposes

nothing for $\varepsilon_0 \le 1$. The three bounds defining b_0 are finite and free of ε_0. Since $\hat{\ell} \ge 1$, the ε_0-free form of (b.3) implies the original.

ε_0. Each condition of Table 7 is downward closed and has a positive threshold (Lemma A.8, Lemma A.3(iii), Lemma A.4(iii)). There are finitely many, the four-mass instances included, so n_0 exists. Some thresholds can be evaluated by hand:

- (E.m2) is $16\varepsilon_0^2 \le 1$, i.e. $\varepsilon_0 \le 0.25$;
- (E.fin) is $\varepsilon_0 \le \min(1/m_2, 1/\mu_*, L) = 0.25$;
- (E.dom) is $\varepsilon_0 \le 0.2$;
- (E.δ) at $\kappa = 7.5$ is $\varepsilon_0 \le 90^{-1/2} = 0.10540\ldots$;
- (E.r) follows from (E.m2);
- (E.charge) reads $\varepsilon_0 \le \min\{1, 400\} = 1$;
- (E.Φ), that is $\lambda^{-1/4} m_2^2 \varepsilon_0^{7/4} \le a_\flat / C_\varphi$ with $C_\varphi = 39.622\, c_1$, holds if $\varepsilon_0 \le (1.89 \cdot 10^{-2} \cdot 0.05^{1/4}/16)^{4/7} c_1^{-4/7} = 0.0138418\, c_1^{-4/7}$, because $a_\flat / C_\varphi \ge 1.89 \cdot 10^{-2}/c_1$.

Hence $n_0 \ge 4$, and $n_0 \ge 7$ if $c_1 \ge 1$.

The threshold of (E.edge) is also explicit in its constants. With a_k replaced by a, (E.edge) at $L = 2$, $d = 3$ requires $F(t) \le 1$ for all $0 < t \le \varepsilon_0/2$, where

$$F(t) := C\, a\, c_1 (\lambda t)^{-1/4} p(t) \big(1 + r(t)\big)^2 \exp\Big(- \delta_0 \Big[\frac{r(t) - M_b}{2} - \sqrt{3} M_b \Big] \Big) (2t)^{-\kappa},$$

with C, c_1, δ_0, M_b and the exponent κ as in (5.7). Since $a_k = a(1 - L^{-2})(1 - L^{-2k})^{-1} \le a$ for $k \ge 1$ [24, (2.13), p. 653], and the left side of (5.7) increases in a_k, this form implies (5.7) at every $k \ge 1$. Put $A := \log(C a c_1 \lambda^{-1/4} b_0 2^{-\kappa}) + \delta_0 (\frac{M_b}{2} + \sqrt{3} M_b) + 2\log(1 + R)$, $A_+ := \max\{A, 0\}$, $B := \kappa + p + \frac{17}{4}$, and $\varepsilon_{24} := \min\{1, 2\exp(1 - 2(A_+ + B)/(\delta_0 R))\}$. Let $0 < t \le \varepsilon_{24}/2$ and $s := \log t^{-1} > 0$. From $1 + R(1+s)^2 \le (1+R)(1+s)^2$ and $\log(1+s) \le s$ we get

$$\log F(t) \le A_+ + Bs - \tfrac{\delta_0 R}{2} (1 + s)^2.$$

Moreover $1 + s \ge 2(A_+ + B)/(\delta_0 R)$, so $\frac{\delta_0 R}{2}(1+s)^2 \ge (A_+ + B)(1+s) \ge A_+ + Bs$, and $F(t) \le 1$. So every $\varepsilon_0 \le \varepsilon_{24}$ satisfies (E.edge). This threshold does not depend on the Higgs mass, on ε, K or ℓ_0. Here $\lambda^{-1/4} = 2.1147\ldots$, but the constants $C, \kappa, \delta_0, c_1, M_b, R_0$ have no printed values in [1, 2], and so neither has ε_{24}. The same holds for the other thresholds; only their positivity is used.

ℓ_0. Each lower bound in the definition of ℓ_0 is a finite function of data of the stages $\le$10b, and $2M_b\mathbb{N}$ is unbounded, so ℓ_0 exists. We have $\ell_0 \ge 183 \ge 128/M_1 = 182.857\ldots = \ell_0^\natural$, so (T.floor) holds, together with $\ell_T \ge 8$ and $M_1 \ell_0 \ge 128.1 \ge 32$. Since $\varepsilon_0 \le 1$ we have $r(\varepsilon_0) \ge R \ge 32 M_b$. Hence

$$6\sqrt{3}\, r + 2 - (8r + 4M_b + 2) = (6\sqrt{3} - 8) r - 4M_b \ge 2.39 \cdot 32 M_b - 4M_b > 0,$$

so (T.per) dominates (T.nw). No condition of an earlier stage depends on ℓ_0 (Lemma A.4).

All tori and all scales. Every condition of Table 8 is a lower bound on the side, so it holds on T_j, whose side is $\ell_j = j!\, \ell_0 \ge \ell_0$. The scales satisfy $t_k = 2^k \varepsilon \le t_K = \varepsilon_0 < \mathrm{e}^{-1}$. The function $t \mapsto t(6\sqrt{3}\, r(t) + 2) = 6\sqrt{3}\, R t (1 + \log t^{-1})^2 + 2t$ is increasing on $(0, \mathrm{e}^{-1})$. So (T.per) gives $S_k = \ell_j / t_k \ge 6\sqrt{3}\, r(t_k) + 2$, and $S_k \ge \ell_0/\varepsilon_0 \ge 2/\delta_0$. Finally $\varepsilon \le \varepsilon_0 \le \frac{1}{4} < 1 = 1/m$.

Appendix B. The lattice model: supplementary proofs

This appendix proves the statements about the regularised model of Section 2 and the standing properties (P1)–(P3) of Section 7 that are used without proof there, together with the lattice facts behind the continuum limits of Section 8: the identification of the action with King's and Bałaban's, the vanishing of the odd orders in e of the counterterm, the operator $-\Delta_\varepsilon = d^* d + d d^*$, gauge invariance, the law of the zero mode and the Gibbs form of the states (Subsection B.1); the correlation inequalities of [8, Sect. 6] on the torus graph, monotonicity in the photon mass, (P1)–(P3), the zero-mode split and the uniformity of the constants of Theorem 6.8 over the tori

(Subsection B.2); multilinearity of the limits, local translation invariance at a fixed torus and monotonicity on the class $\mathcal{D}$ (Subsection B.3). The letter e is the charge; e is Euler's number.

B.1. **Model and states.**

B.1.1. *The action and the odd orders of the counterterm.* Throughout this subsection, $T_{j,\varepsilon}$ is a lattice torus as in Subsection 2.1, sites are $x \in T_{j,\varepsilon}$ and bonds are the positively oriented pairs (x,μ), each counted once. For a function f on sites put

$$(\partial_\mu f)(x) := \varepsilon^{-1}\big(f(x+\varepsilon e_\mu) - f(x)\big),$$
$$(\partial_\mu^* f)(x) := \varepsilon^{-1}\big(f(x-\varepsilon e_\mu) - f(x)\big),$$
$$-\Delta_\varepsilon := \sum_{\mu=1}^{3} \partial_\mu^*\partial_\mu.$$

On the periodic lattice ∂_μ^* is the adjoint of ∂_μ for $\langle\cdot,\,\cdot\rangle$, and all the operators $\partial_\mu, \partial_\nu^*$ commute, since they are linear combinations of translations. A bond function A is written $A_\mu(x) := A_{x,\mu}$.
The Feynman form and King's action.

Lemma B.1 (the action of Subsection 2.2 is King's and Bałaban's)**.** *Let $\mu_0 > 0$, $a \in \mathbb{R}$, $\lambda > 0$ and $e \in \mathbb{R}$.*

(a) *On bond functions, $-\Delta_\varepsilon := d^*d + dd^*$ acts componentwise as the lattice Laplacian on sites: $(-\Delta_\varepsilon A)_\nu = -\Delta_\varepsilon A_\nu$. Hence $-\Delta_\varepsilon \geq 0$ on bond functions, and its kernel consists of the constant bond functions.*

(b) $\frac{1}{2}\langle A,\,(-\Delta_\varepsilon + \mu_0^2)A\rangle = \frac{1}{2}\|dA\|^2 + \frac{1}{2}\|d^*A\|^2 + \frac{1}{2}\mu_0^2\|A\|^2 = \frac{1}{2}\sum_\nu \langle A_\nu,\,(-\Delta_\varepsilon+\mu_0^2)A_\nu\rangle$.

(c) *The matter action* (2.2) *can be written as*

$$S_a^{\mathrm{m}}(A,\varphi) = \sum_x \Big(3\varepsilon|\varphi(x)|^2 + \varepsilon^3\big(\tfrac{1}{2}a|\varphi(x)|^2 + \lambda|\varphi(x)|^4\big)\Big) - \sum_{x,\mu} \varepsilon\ \mathrm{Re}\,\big(\overline{\varphi(x)}\,\mathrm{e}^{ie\varepsilon A_{x,\mu}}\varphi(x+\varepsilon e_\mu)\big).$$

(d) *Let q be the antisymmetric real 2×2 matrix with $\|q\| = 1$ of* [24, p. 651] *and* [1, (1.7), p. 605] *($N = 2$). Let $\iota : \mathbb{R}^2 \to \mathbb{C}$ be the isometric identification with $\iota(qv) = i\,\iota(v)$. Write $S^\varepsilon(\Omega, A, \phi)$ for King's action* [24, (2.4), p. 652]*, with $\Omega = T_{j,\varepsilon}$ and periodic operators* [24, (2.21), p. 654]*, scalar mass m and photon mass μ_0. Write $S^{\mathrm{Ba}}(A,\phi)$ for Bałaban's action* [1, (1.11), p. 605]*. Then for all $(A,\varphi) \in \mathbb{R}^{bonds} \times \mathbb{C}^{sites}$, with $\phi := \iota^{-1}\circ\varphi$,*

$$S_{\mu_0,a}(A,\varphi) = S^\varepsilon(T_{j,\varepsilon}, A, \phi)\big|_{\delta m^2 \equiv a - m^2} - E_0 - E_1 = S^{\mathrm{Ba}}(A,\phi)\big|_{m_0^2 = a} - E.$$

(e) *Consequently the Gibbs factors $\mathrm{e}^{-S_{\mu_0,a}}$ and $\mathrm{e}^{-S^\varepsilon(T_{j,\varepsilon},\cdot)}$ differ only by the constant factor $\mathrm{e}^{E_0+E_1}$, so the normalised measures coincide. With $a = b_{T_j}(\varepsilon) = a_{K,T_j}(m)$ (Definition 2.2), the model of Definition 2.7 at photon mass μ_0 is King's torus model with $\delta m^2 := \delta m^2_{K,T}(\varepsilon; m)$ computed at the reference photon mass μ_*. This is King's own torus counterterm only when $\mu_0 = \mu_*$ (Remark 2.3(ii)).*

Part (d) is the identification stated after (2.2) *in Subsection 2.2, used in Proposition 6.3; part (a) is the statement on $-\Delta_\varepsilon$ made there.*

Proof. (a), (b) Part (a) and the second equality of (b) are Lemma B.4(i),(ii) below, which also gives the kernel and the spectral gap. The first equality of (b) is the definition of the adjoint, $\langle A,\,(d^*d+dd^*)A\rangle = \|dA\|^2 + \|d^*A\|^2$.

(c) For each bond, $\frac{1}{2}\varepsilon^{-2}|\mathrm{e}^{ie\varepsilon A_{x,\mu}}\varphi(x+\varepsilon e_\mu) - \varphi(x)|^2 = \frac{1}{2}\varepsilon^{-2}\big(|\varphi(x+\varepsilon e_\mu)|^2 + |\varphi(x)|^2\big) - \varepsilon^{-2}\,\mathrm{Re}\,\big(\overline{\varphi(x)}\mathrm{e}^{ie\varepsilon A_{x,\mu}}\varphi(x+\varepsilon e_\mu)\big)$, because $|\mathrm{e}^{ie\varepsilon A}| = 1$. Multiply by ε^3 and sum over the bonds. Every site is the initial point of 3 bonds and the end point of 3 bonds, so the diagonal terms add up to $\sum_x 3\varepsilon|\varphi(x)|^2$.

(d) The identification ι. An antisymmetric real 2×2 matrix is $q=\theta\left(\begin{smallmatrix}0&-1\\1&0\end{smallmatrix}\right)$, and $\|q\|=1$ forces $\theta=\pm 1$. Take $\iota(v_1,v_2)=v_1+i\theta v_2$. Then ι is an isometry, it preserves Lebesgue measure, and $\iota(qv)=i\iota(v)$. Hence $\iota\circ\exp(tq)=\mathrm{e}^{it}\iota$, and King's and Bałaban's $U(A_b)=\exp(\varepsilon eqA_b)$ [24, p. 651], [1, p. 605] becomes multiplication by $\mathrm{e}^{ie\varepsilon A_b}$.

Matter part. With $b=(x,\mu)$, $b_-=x$, $b_+=x+\varepsilon e_\mu$, King's covariant derivative is [24, (2.1)] $(D_A^\varepsilon\phi)(b)=\varepsilon^{-1}(U(A_b)\phi(b_+)-\phi(b_-))$. Its image under ι is $\varepsilon^{-1}(\mathrm{e}^{ie\varepsilon A_{x,\mu}}\varphi(x+\varepsilon e_\mu)-\varphi(x))$. By [24, (2.2)–(2.3)], $\frac12\langle\phi,\,-\Delta_A^{\varepsilon,\Omega}\phi\rangle=\frac12\sum_b\varepsilon^3|(D_A^\varepsilon\phi)(b)|^2$, which is the gradient term of (2.2). Here the sum runs over the bonds of $T_{j,\varepsilon}$, each once. This agrees with [1, (1.8)] ($b\subset T_\varepsilon$) and with $-\Delta_A^\varepsilon=D_A^{\varepsilon *}D_A^\varepsilon$ in [1, (1.11)], and it is the reading under which $-\Delta_A^{\varepsilon,\Omega}$ at $A=0$ is the lattice Laplacian $-\Delta_\varepsilon$. The remaining scalar terms of [24, (2.4)] are $\frac12\langle\phi,\,m^2\phi\rangle+\sum_x\varepsilon^3(\frac12\delta m^2|\phi|^2+\lambda|\phi|^4)$. With $\delta m^2\equiv a-m^2$ these equal $\sum_x\varepsilon^3(\frac12 a|\varphi|^2+\lambda|\varphi|^4)$. On a torus $\delta m^2(x)$ is constant in x.

Gauge part. King's inner product [24, (2.2)] on $A:\Omega\to\mathbb{R}^3$ is $\sum_i\sum_x\varepsilon^3A_i(x)B_i(x)=\langle A,\,B\rangle$. King's $-\Delta^{\varepsilon,\Omega}$ on $\Omega=T_{j,\varepsilon}$ is the periodic Laplacian acting on each component [24, (2.4), (2.21)]. By (b), $\frac12\langle A,\,(-\Delta^{\varepsilon,\Omega}+\mu_0^2)A\rangle=\frac12\langle A,\,(-\Delta_\varepsilon+\mu_0^2)A\rangle$.

Constants. King's action contains $+E_0+E_1$ [24, (2.4), (2.6), (3.35)], and Bałaban's contains $+E$ [1, (1.11)]. These are field-independent constants, omitted in (2.1) (Remark 2.3(iii)).

Bałaban's form. [1, (1.11)] is $\frac12\langle\phi,\,-\Delta_A^\varepsilon\phi\rangle+\sum_{x\in T_\varepsilon}\varepsilon^3(\frac12 m_0^2|\phi|^2+\lambda|\phi|^4)+\frac12\langle A,\,(-\Delta^\varepsilon+\mu_0^2)A\rangle+E$. Here $-\Delta^\varepsilon=\partial^{\varepsilon *}\partial^\varepsilon$ is the torus Laplacian and U is as above. With $m_0^2=a$, the same three identifications give the second equality.

(e) This follows from (d), because E_0+E_1 does not depend on (A,φ) and ι preserves Lebesgue measure. The statement about the coefficient is Definition 2.2 (photon propagator $(-\Delta_\varepsilon+\mu_*^2)^{-1}$) together with Remark 2.3(ii). □

The odd orders in e of the counterterm. Fix $M>0$ and the torus $T_{j,\varepsilon}$. Let ν_M be the centred Gaussian measure on $\mathbb{C}^{\text{sites}}$ with density proportional to $\exp(-\frac12\langle\varphi,\,(-\Delta_\varepsilon+M^2)\varphi\rangle)$, and let $\mathbb{E}_0$ be the expectation in $\gamma_{\mu_*}\otimes\nu_M$ (γ_{μ_*} as in Definition 2.7). This measure is non-degenerate since $\mu_*,M>0$, so all its polynomial moments are finite.

For a polynomial $\delta=\sum_{2\le\alpha+2\beta\le 4}e^\alpha\lambda^\beta\delta_{\alpha\beta}$ with real coefficients, define the interaction

$$V_\delta:=S^{\mathrm{m}}_{M^2+\delta}(A,\varphi)-\tfrac12\langle\varphi,\,(-\Delta_\varepsilon+M^2)\varphi\rangle=\sum_{n\ge1}e^nH^{(n)}+\sum_x\varepsilon^3\big(\tfrac12\delta|\varphi(x)|^2+\lambda|\varphi(x)|^4\big),$$

$$H^{(n)}(A,\varphi):=-\sum_{x,\mu}\varepsilon\ \mathrm{Re}\left(\overline{\varphi(x)}\,\frac{(i\varepsilon A_{x,\mu})^n}{n!}\,\varphi(x+\varepsilon e_\mu)\right).$$

The second expression for V_δ is Lemma B.1(c) with the $n=0$ term removed: the $n=0$ term together with the diagonal term is $\frac12\langle\varphi,\,-\Delta_\varepsilon\varphi\rangle$. The perturbative two-point function is the formal power series in (e,λ) of $\mathbb{E}_0[\varphi_i(x)\varphi_k(y)\mathrm{e}^{-V_\delta}]/\mathbb{E}_0[\mathrm{e}^{-V_\delta}]$. By Wick's theorem its coefficients are sums of Feynman graphs with the following vertices: hopping vertices $e^nH^{(n)}$; quartic vertices $\lambda|\varphi|^4$; counterterm vertices $e^{\alpha'}\lambda^{\beta'}\frac12\delta_{\alpha'\beta'}|\varphi|^2$. Photon lines are $(-\Delta_\varepsilon+\mu_*^2)^{-1}$ and scalar lines are $(-\Delta_\varepsilon+M^2)^{-1}$. Following [24, (3.32)–(3.33), p. 659] and Definition 2.2, write $\Gamma(x,y;\alpha,\beta)$ for the coefficient of $e^\alpha\lambda^\beta$ in the sum of the amputated one-particle-irreducible graphs with two external scalar legs, other than the single counterterm vertex. The counterterm is the solution of

$$-\delta_{\alpha\beta}+\sum_y\varepsilon^3\,\Gamma(x,y;\alpha,\beta)=0,\qquad 2\le\alpha+2\beta\le4,\tag{B.1}$$

solved recursively; $\Gamma(\cdot;\alpha,\beta)$ contains only coefficients $\delta_{\alpha'\beta'}$ with $\alpha'\le\alpha$, $\beta'\le\beta$, $(\alpha',\beta')\ne(\alpha,\beta)$. With M and photon mass μ_* this is $\delta m^2_{K,T}(\varepsilon;M)$.

Lemma B.2 (odd orders vanish)**.** *In the solution of* (B.1)*, $\delta_{\alpha\beta} = 0$ whenever α is odd, and $\Gamma(x, y; \alpha, \beta) = 0$ for odd α. Hence*

$$\delta m^2_{K,T}(\varepsilon; M) = e^2\delta_{20} + \lambda\delta_{01} + e^4\delta_{40} + e^2\lambda\,\delta_{21} + \lambda^2\delta_{02},$$

and $a_{K,T}(M) = M^2 + \delta m^2_{K,T}(\varepsilon; M)$ is a polynomial in (e^2, λ). This is the statement in Definition 2.2 that the odd orders in e vanish.

Proof. Step 1 (vertices). $H^{(n)}$ is homogeneous of degree n in A: it is a sum of terms $A^n_{x,\mu}$ times a quadratic form in φ. The quartic and counterterm vertices do not contain A. So a hopping vertex of order e^n has exactly n photon legs, and every other vertex has none. The photon covariance does not depend on e.

Step 2 (contractions). Under $\gamma_{\mu_*} \otimes \nu_M$ the fields A and φ are independent, and γ_{μ_*} is centred, hence invariant under $A \mapsto -A$. So in a Wick contraction the photon legs are paired with each other. There are no external photon legs, because the external legs are scalar. A graph with an odd number of photon legs therefore has value 0. Equivalently, $\mathbb{E}_0[P] = 0$ for every polynomial P of odd degree in A, since $\mathbb{E}_0[P(A, \varphi)] = \mathbb{E}_0[P(-A, \varphi)] = -\mathbb{E}_0[P]$.

Step 3 (induction). Order the pairs (α, β), $2 \le \alpha + 2\beta \le 4$, in the order in which (B.1) determines them. Assume $\delta_{\alpha'\beta'} = 0$ for all earlier pairs with α' odd. Take a graph contributing to $\Gamma(\cdot; \alpha, \beta)$ with nonzero value. Let it have hopping vertices of orders $n_1, \dots, n_r$, β_0 quartic vertices, and counterterm vertices of orders $(\alpha_1, \beta_1), \dots, (\alpha_s, \beta_s)$. Then

$$\alpha = \sum_k n_k + \sum_i \alpha_i, \qquad \beta = \beta_0 + \sum_i \beta_i.$$

Every vertex carries positive order, and the graph is not the single counterterm vertex. Hence each (α_i, β_i) is an earlier pair, and by hypothesis α_i is even (otherwise $\delta_{\alpha_i\beta_i} = 0$ and the graph vanishes). By Steps 1 and 2, $\sum_k n_k$ is the number of photon legs, which is even. So α is even. Thus $\Gamma(\cdot; \alpha, \beta) = 0$ for odd α, and (B.1) gives $\delta_{\alpha\beta} = 0$. The pairs with $2 \le \alpha + 2\beta \le 4$ are $(2,0), (0,1), (3,0), (1,1), (4,0), (2,1), (0,2)$; removing those with odd α leaves the five monomials stated. □

Remark B.3. (i) Lemma B.2 is the precise content of the symmetry named in Definition 2.2. Suppose δ is even in e. Then $H^{(n)}(-A, \varphi) = (-1)^n H^{(n)}(A, \varphi)$ gives $V_\delta(-e; -A, \varphi) = V_\delta(e; A, \varphi)$: the substitution $A \mapsto -A$, $e \mapsto -e$ fixes φ and leaves V_δ and γ_{μ_*} invariant. Hence the coefficient of $e^\alpha\lambda^\beta$ in the expansion of e^{-V_δ} has A-parity $(-1)^\alpha$, and its $\mathbb{E}_0$-expectation, and that of its product with any polynomial in φ, vanishes for odd α (Step 2). In this form evenness in e is carried along the recursion by Step 3. (ii) Charge conjugation $(A, \varphi) \mapsto (-A, \bar\varphi)$ at fixed e is also a symmetry of (2.2). But it does not relate e to $-e$, so it does not by itself give the lemma. It gives the vanishing of functions with an odd number of external photon legs. (iii) Steps 1–3 count photon legs only. They therefore hold for every decomposition of the propagators into sums, in particular for the renormalisation-group graphs of [24, (3.34)].

B.1.2. *The operator $-\Delta_\varepsilon$, gauge invariance, the zero mode and the Gibbs form.* We prove four statements of Section 2. Throughout, $j \ge 1$ and ε are fixed, $N = \ell_j/\varepsilon$ is even (Lemma 2.1), and $n := 3N^3$ is the number of bonds of $T_{j,\varepsilon}$. For $\mu \in \{1, 2, 3\}$ let $\mathbf{1}_\mu$ be the bond function equal to 1 on the bonds parallel to e_μ and 0 on the others.

Conventions. (C1) Sites, bonds and plaquettes all carry the weight ε^3 in $\langle \cdot, \cdot \rangle$. Hence the adjoint d^* is the transpose of d in the standard coordinates, and the matrix of $-\Delta_\varepsilon = d^*d + dd^*$ in the coordinates $(A_b)_b$ is symmetric. (C2) "Centred Gaussian measure with covariance C" on bond functions means the covariance operator for $\langle \cdot, \cdot \rangle$: $\mathbb{E}[\langle f, A\rangle\langle g, A\rangle] = \langle f, Cg\rangle$ for all bond functions f, g. For non-degenerate C this measure has a Lebesgue density proportional to $\exp(-\frac{1}{2}\langle A, C^{-1}A\rangle)$. This is the convention stated with (7.3). The value $s(\mu_0) = (\mu_0^2|T_j|)^{-1}$ in Definition 2.7 holds in this convention (Lemma B.6).

Lemma B.4 (the operator $d^*d + dd^*$ is the componentwise Laplacian)**.** *Let $-\Delta_\varepsilon := d^*d + dd^*$ on bond functions, and for a function f on sites put $(-\Delta_\varepsilon f)(x) := \varepsilon^{-2}\sum_{\nu=1}^{3}\big(2f(x) - f(x+\varepsilon e_\nu) - f(x-\varepsilon e_\nu)\big)$ (periodic shifts).*

(i) *For every bond function A and every μ, $(-\Delta_\varepsilon A)_{x,\mu} = (-\Delta_\varepsilon A_{\cdot,\mu})(x)$.*

(ii) *$\langle A, -\Delta_\varepsilon A\rangle = \|dA\|^2 + \|d^*A\|^2 = \sum_{x,\mu,\nu}\varepsilon\,(A_{x+\varepsilon e_\nu,\mu} - A_{x,\mu})^2 \geq 0$. Moreover $\operatorname{Ker}\Delta_\varepsilon = \operatorname{span}\{\mathbf{1}_1, \mathbf{1}_2, \mathbf{1}_3\} = \operatorname{ran}\Pi_0$, and $\Delta_\varepsilon\Pi_0 = \Pi_0\Delta_\varepsilon = 0$.*

(iii) *On $\operatorname{ran}\Pi^\perp$, $-\Delta_\varepsilon \geq 4\varepsilon^{-2}\sin^2(\pi\varepsilon/\ell_j) \geq 16/\ell_j^2$. In particular $-\Delta_\varepsilon + \mu_0^2$ is positive definite for every $\mu_0 > 0$, and $-\Delta_\varepsilon$ is invertible on $\operatorname{ran}\Pi^\perp$.*

Proof. (i) By (C1) and the definition of d,

$$(d^*A)(x) = \varepsilon^{-1}\sum_{\nu}\big(A_{x-\varepsilon e_\nu,\nu} - A_{x,\nu}\big),$$

since $\langle d\chi,\, A\rangle = \sum_{x,\nu}\varepsilon^3\varepsilon^{-1}(\chi(x+\varepsilon e_\nu) - \chi(x))A_{x,\nu}$, and the periodic shift $x \mapsto x - \varepsilon e_\nu$ is a bijection of $T_{j,\varepsilon}$. For a plaquette function B, extend it antisymmetrically, $B(x;\nu,\mu) := -B(x;\mu,\nu)$; the definition of dA in Subsection 2.2 is antisymmetric in (μ,ν). Then $\langle dA,\, B\rangle = \frac{1}{2}\sum_x\sum_{\mu\neq\nu}\varepsilon^3(dA)(x;\mu,\nu)B(x;\mu,\nu)$. Collect the coefficient of $A_{x,\mu}$: the four terms of $(dA)(y;\alpha,\beta)$ that contain it come in two antisymmetric pairs. This gives

$$(d^*B)_{x,\mu} = \varepsilon^{-1}\sum_{\nu\neq\mu}\big(B(x;\mu,\nu) - B(x-\varepsilon e_\nu;\mu,\nu)\big).$$

Insert $B = dA$, with $(dA)(x;\mu,\nu) = \varepsilon^{-1}(A_{x,\mu} + A_{x+\varepsilon e_\mu,\nu} - A_{x+\varepsilon e_\nu,\mu} - A_{x,\nu})$. Then

$$(d^*dA)_{x,\mu} = \varepsilon^{-2}\sum_{\nu\neq\mu}\big(2A_{x,\mu} - A_{x+\varepsilon e_\nu,\mu} - A_{x-\varepsilon e_\nu,\mu}\big) + \varepsilon^{-2}\sum_{\nu\neq\mu}\big(A_{x+\varepsilon e_\mu,\nu} - A_{x,\nu} - A_{x+\varepsilon e_\mu-\varepsilon e_\nu,\nu} + A_{x-\varepsilon e_\nu,\nu}\big),$$

$$(dd^*A)_{x,\mu} = \varepsilon^{-1}\big((d^*A)(x+\varepsilon e_\mu) - (d^*A)(x)\big) = \varepsilon^{-2}\sum_{\nu}\big(A_{x+\varepsilon e_\mu-\varepsilon e_\nu,\nu} - A_{x+\varepsilon e_\mu,\nu} - A_{x-\varepsilon e_\nu,\nu} + A_{x,\nu}\big).$$

The terms $\nu \neq \mu$ of the second line cancel the second sum of the first line. The term $\nu = \mu$ of the second line is $\varepsilon^{-2}(2A_{x,\mu} - A_{x+\varepsilon e_\mu,\mu} - A_{x-\varepsilon e_\mu,\mu})$. Adding the two lines gives $((d^*d + dd^*)A)_{x,\mu} = \varepsilon^{-2}\sum_\nu(2A_{x,\mu} - A_{x+\varepsilon e_\nu,\mu} - A_{x-\varepsilon e_\nu,\mu})$, which is (i).

(ii) $\langle A,\, (d^*d + dd^*)A\rangle = \langle dA,\, dA\rangle + \langle d^*A,\, d^*A\rangle$ by the definition of the adjoint. By (i), and because $\sum_x f(x)f(x-\varepsilon e_\nu) = \sum_x f(x+\varepsilon e_\nu)f(x)$ on the torus, $\sum_x\varepsilon^3 f(x)(-\Delta_\varepsilon f)(x) = \sum_{x,\nu}\varepsilon\,(f(x+\varepsilon e_\nu) - f(x))^2$ for each component $f = A_{\cdot,\mu}$. This gives the identity and $-\Delta_\varepsilon \geq 0$. Since $-\Delta_\varepsilon$ is symmetric and non-negative, $\Delta_\varepsilon A = 0$ if and only if $\langle A, -\Delta_\varepsilon A\rangle = 0$, that is, if and only if every component has vanishing forward differences in every direction. Every site of the torus is reached from a fixed site by forward steps, so this means every component is constant, i.e. $A \in \operatorname{span}\{\mathbf{1}_\mu\} = \operatorname{ran}\Pi_0$. Finally $\Delta_\varepsilon\Pi_0 = 0$, and $\Pi_0\Delta_\varepsilon = (\Delta_\varepsilon\Pi_0)^* = 0$ because Δ_ε and Π_0 are self-adjoint.

(iii) For $k \in \{0,\dots,N-1\}^3$ put $f_k(x) := \mathrm{e}^{2\pi i k\cdot x/\ell_j}$, $x \in T_{j,\varepsilon}$. These N^3 functions form an orthogonal basis of $\mathbb{C}^{T_{j,\varepsilon}}$, f_0 is the constant, and $f_k(x+\varepsilon e_\nu) = \mathrm{e}^{2\pi i k_\nu/N}f_k(x)$. Hence

$$-\Delta_\varepsilon f_k = \varepsilon^{-2}\sum_\nu\big(2 - 2\cos(2\pi k_\nu/N)\big)f_k = 4\varepsilon^{-2}\sum_\nu\sin^2(\pi k_\nu/N)\,f_k.$$

If $k \neq 0$, some $k_\nu \in \{1,\dots,N-1\}$, and $\pi k_\nu/N \in [\pi/N, \pi - \pi/N]$ gives $\sin^2(\pi k_\nu/N) \geq \sin^2(\pi/N)$. Since $N \geq 2$, $\pi/N \leq \pi/2$, and $\sin t \geq 2t/\pi$ on $[0,\pi/2]$ gives $4\varepsilon^{-2}\sin^2(\pi/N) \geq 4\varepsilon^{-2}(2/N)^2 = 16/\ell_j^2$. So the real symmetric operator $-\Delta_\varepsilon$ is $\geq 16/\ell_j^2$ on the functions with zero sum. Now $\Pi_0 A = \sum_\mu c_\mu\mathbf{1}_\mu$, where c_μ is the mean of $A_{\cdot,\mu}$ (Lemma B.6(a)). So $\operatorname{ran}\Pi^\perp$ consists of the bond functions all of whose components have zero sum, and (i) gives the bound on $\operatorname{ran}\Pi^\perp$. The remaining claims follow from $-\Delta_\varepsilon \geq 0$. □

Lemma B.5 (gauge invariance of the matter action)**.** *For a real function χ on sites put $G_\chi(A,\varphi) := (A - \mathrm{e}^{-1}d\chi,\, \mathrm{e}^{i\chi}\varphi)$. For every real a and all (A,φ),*

$$S_a^{\mathrm{m}}(G_\chi(A,\varphi)) = S_a^{\mathrm{m}}(A,\varphi), \qquad d(A-\mathrm{e}^{-1}d\chi) = dA, \qquad |\mathrm{e}^{i\chi(x)}\varphi(x)| = |\varphi(x)|.$$

Hence every $F \in \mathfrak{A}_\varepsilon$ and every product $F\,\mathrm{e}^{-S_a^{\mathrm{m}}}$ is invariant under G_χ. The Gaussian part of (2.1) *is not invariant: $\|d^*A\|^2$ and $\|A\|^2$ change.*

Proof. Let $y = x + \varepsilon e_\mu$. Since $\varepsilon e \cdot \mathrm{e}^{-1}(d\chi)_{x,\mu} = \chi(y) - \chi(x)$,

$$\mathrm{e}^{ie\varepsilon(A_{x,\mu} - \mathrm{e}^{-1}(d\chi)_{x,\mu})}\mathrm{e}^{i\chi(y)}\varphi(y) - \mathrm{e}^{i\chi(x)}\varphi(x) = \mathrm{e}^{i\chi(x)}\big(\mathrm{e}^{ie\varepsilon A_{x,\mu}}\varphi(y) - \varphi(x)\big),$$

and this holds also for the bonds that wrap around the torus. So every gradient term of (2.2) keeps its modulus. The single-site terms depend on $|\varphi(x)|$ only. Finally $d(d\chi) = 0$: each plaquette contains the four values χ at its corners, each once with each sign. Elements of $\mathfrak{A}_\varepsilon$ are functions of dA and $|\varphi|^2$ (Definition 2.4). $\|d^*A\|^2$ and $\|A\|^2$ change by $d^*d\chi$ and $d\chi$ terms, which do not vanish for non-constant χ. □

Other parametrisations. Section 7 writes the gauge transformations as $(A + d\chi,\, U(-e\chi)\varphi)$. This is $G_{-e\chi}$, and $U(s)$ is multiplication by e^{is} on $\mathbb{R}^2 \cong \mathbb{C}$. The proof of Lemma 7.7 uses $(G\mathrm{e}^{-S_m})(Z + d\chi,\varphi) = (G\mathrm{e}^{-S_m})(Z, U(e\chi)\varphi)$. This is Lemma B.5 applied with $G_{e\chi}$, since $G_{e\chi}(Z + d\chi,\varphi) = (Z, \mathrm{e}^{ie\chi}\varphi)$. Here $e \neq 0$ is used.

Lemma B.6 (the zero mode under γ_{μ_0})**.** *Let $\mu_0 > 0$.*

(a) *$\langle \mathbf{1}_\mu,\, \mathbf{1}_\nu\rangle = |T_j|\delta_{\mu\nu}$, and $\Pi_0 A = \sum_\mu c_\mu \mathbf{1}_\mu$ with $c_\mu := |T_j|^{-1}\langle \mathbf{1}_\mu,\, A\rangle$, the mean of $A_{\cdot,\mu}$.*

(b) *The centred Gaussian measure γ_{μ_0} with covariance $C_{\mu_0} = (-\Delta_\varepsilon + \mu_0^2)^{-1}$ exists and is non-degenerate. Under γ_{μ_0}, $A^\perp := \Pi^\perp A$ and $c = (c_1,c_2,c_3)$ are independent. $A^\perp$ is the centred Gaussian on $\operatorname{ran}\Pi^\perp$ with covariance $(-\Delta_\varepsilon + \mu_0^2)^{-1}\Pi^\perp$. c is the centred Gaussian on $\mathbb{R}^3$ with covariance matrix $s(\mu_0)\mathbf{1}_3$, where $s(\mu_0) = (\mu_0^2|T_j|)^{-1}$.*

(c) *$(-\Delta_\varepsilon)^{-1}\Pi^\perp$ is a positive definite operator on $\operatorname{ran}\Pi^\perp$, so the measure $\gamma^\flat$ of Definition 2.7 is well defined.*

Proof. (a) There are N^3 bonds parallel to e_μ, and $N^3\varepsilon^3 = \ell_j^3 = |T_j|$. The $\mathbf{1}_\mu$ are orthogonal, so the orthogonal projection onto their span is $\sum_\mu |T_j|^{-1}\langle \mathbf{1}_\mu,\, A\rangle \mathbf{1}_\mu$.

(b) $-\Delta_\varepsilon + \mu_0^2$ is positive definite by Lemma B.4(iii), so C_{μ_0} is a positive definite operator and γ_{μ_0} exists (Lemma B.7(i) gives its density). $A^\perp$ and $\Pi_0 A$ are linear images of the same Gaussian vector, so they are jointly Gaussian. By Lemma B.4(ii), Π_0 commutes with Δ_ε, hence with C_{μ_0}. So for all bond functions f, g,

$$\mathbb{E}_{\gamma_{\mu_0}}\big[\langle f,\, \Pi^\perp A\rangle\langle g,\, \Pi_0 A\rangle\big] = \langle \Pi^\perp f,\, C_{\mu_0}\Pi_0 g\rangle = \langle f,\, \Pi^\perp\Pi_0 C_{\mu_0} g\rangle = 0.$$

Uncorrelated jointly Gaussian vectors are independent, so $A^\perp$ and c (a linear function of $\Pi_0 A$, by (a)) are independent. The covariance of $A^\perp$ is $\Pi^\perp C_{\mu_0}\Pi^\perp = (-\Delta_\varepsilon + \mu_0^2)^{-1}\Pi^\perp$. Since $\Delta_\varepsilon \mathbf{1}_\nu = 0$, $C_{\mu_0}\mathbf{1}_\nu = \mu_0^{-2}\mathbf{1}_\nu$. Hence, by (a),

$$\operatorname{Cov}(c_\mu, c_\nu) = |T_j|^{-2}\langle \mathbf{1}_\mu,\, C_{\mu_0}\mathbf{1}_\nu\rangle = |T_j|^{-2}\mu_0^{-2}\langle \mathbf{1}_\mu,\, \mathbf{1}_\nu\rangle = \frac{\delta_{\mu\nu}}{\mu_0^2|T_j|}.$$

Equivalently, in terms of the density: $\langle A^\perp,\, (-\Delta_\varepsilon + \mu_0^2)c\rangle = \mu_0^2\langle A^\perp,\, c\rangle = 0$, so

$$\tfrac12\langle A,\, (-\Delta_\varepsilon + \mu_0^2)A\rangle = \tfrac12\langle A^\perp,\, (-\Delta_\varepsilon + \mu_0^2)A^\perp\rangle + \tfrac12\mu_0^2|T_j|\sum_\mu c_\mu^2.$$

The density factorises, and the c-factor is the $N(0, s(\mu_0))$ density in each coordinate.

(c) $-\Delta_\varepsilon$ maps $\operatorname{ran}\Pi^\perp$ into itself (it commutes with $\Pi^\perp$) and is $\geq 16/\ell_j^2$ there, by Lemma B.4(ii),(iii). So its inverse on $\operatorname{ran}\Pi^\perp$ exists and is positive definite. □

Lemma B.7 (Gibbs form of the states)**.** *Let $\mu_0 > 0$ and $a \in \mathbb{R}$. Let $Q := \varepsilon^3(-\Delta_\varepsilon + \mu_0^2)$ be the matrix of the quadratic form $A \mapsto \langle A, (-\Delta_\varepsilon + \mu_0^2)A\rangle$ in the coordinates $(A_b)_b$, and $Z_\gamma := (2\pi)^{n/2}(\det Q)^{-1/2}$. Let Ψ_F^a and $W_0^a = \Psi_1^a$ be defined as in Definition 2.4, with the coefficient a in place of b.*

(i) $\gamma_{\mu_0}(dA) = Z_\gamma^{-1}\mathrm{e}^{-\frac{1}{2}\langle A, (-\Delta_\varepsilon+\mu_0^2)A\rangle}\prod_b dA_b$.

(ii) *For every Borel $F \geq 0$ on $\mathbb{R}^{bonds} \times \mathbb{C}^{sites}$, $A \mapsto \Psi_F^a(A) \in [0,\infty]$ is Borel, and*

$$\int F\,\mathrm{e}^{-S_{\mu_0,a}} \prod_b dA_b \prod_x d^2\varphi(x) = Z_\gamma\,\mathbb{E}_{\gamma_{\mu_0}}[\Psi_F^a].$$

For complex Borel F with $\mathbb{E}_{\gamma_{\mu_0}}[\Psi_{|F|}^a] < \infty$, both sides are finite and equal.

(iii) $0 < W_0^a(A) \leq W_0^a(0) < \infty$. *Hence $Z := \int \mathrm{e}^{-S_{\mu_0,a}}\prod_b dA_b \prod_x d^2\varphi(x) = Z_\gamma \mathbb{E}_{\gamma_{\mu_0}}[W_0^a] \in (0,\infty)$, and $Z^{-1}\mathrm{e}^{-S_{\mu_0,a}}\prod_b dA_b\prod_x d^2\varphi(x)$ is a probability measure. For $a = b_{T_j}(\varepsilon)$ and every $F \in \mathfrak{A}_\varepsilon$ with $\mathbb{E}_{\gamma_{\mu_0}}[\Psi_{|F|}] < \infty$ (in particular every bounded F),*

$$\langle F\rangle_{\varepsilon,j,\mu_0} = Z^{-1}\int F\,\mathrm{e}^{-S_{\mu_0,b}}\prod_b dA_b\prod_x d^2\varphi(x).$$

(iv) $0 < \mathbb{E}_{\gamma^\flat}[W_0] \leq W_0(0)$. *For $F \in \mathfrak{A}_\varepsilon$ bounded or non-negative, $\mathbb{E}_{\gamma^\flat}[\Psi_F]$ does not change if the period cell $[0,P)^3$ in the definition of $\gamma^\flat$ is replaced by any translate $\prod_\mu[t_\mu, t_\mu + P)$.*

Proof. (i) By (C1), $\langle A, (-\Delta_\varepsilon + \mu_0^2)A\rangle = A \cdot QA$ with Q symmetric, and Q is positive definite by Lemma B.4(iii). The Gaussian integral gives $\int \mathrm{e}^{-\frac{1}{2}A\cdot QA}\prod_b dA_b = (2\pi)^{n/2}(\det Q)^{-1/2} = Z_\gamma$. The normalised measure is the centred Gaussian with coordinate covariance $Q^{-1} = \varepsilon^{-3}(-\Delta_\varepsilon + \mu_0^2)^{-1}$. So $\mathbb{E}[\langle f, A\rangle\langle g, A\rangle] = \varepsilon^6 f \cdot Q^{-1}g = \varepsilon^3 f \cdot (-\Delta_\varepsilon + \mu_0^2)^{-1}g = \langle f, C_{\mu_0}g\rangle$, which is γ_{μ_0} in the convention (C2).

(ii) $S_{\mu_0,a} = \frac{1}{2}\langle A, (-\Delta_\varepsilon + \mu_0^2)A\rangle + S_a^{\mathrm{m}}$ is continuous, so $F\mathrm{e}^{-S_{\mu_0,a}}$ is Borel. Tonelli's theorem on $\mathbb{R}^{\mathrm{bonds}} \times \mathbb{C}^{\mathrm{sites}}$, integrating first over φ, gives the Borel measurability of Ψ_F^a and

$$\int F\mathrm{e}^{-S_{\mu_0,a}} = \int \prod_b dA_b\,\mathrm{e}^{-\frac{1}{2}\langle A, (-\Delta_\varepsilon+\mu_0^2)A\rangle}\,\Psi_F^a(A) = Z_\gamma \mathbb{E}_{\gamma_{\mu_0}}[\Psi_F^a],$$

using (i). For complex F, apply this to $|F|$: then $F\mathrm{e}^{-S_{\mu_0,a}}$ is integrable, and Fubini's theorem, applied to the positive and negative parts of $\mathrm{Re}\,F$ and $\mathrm{Im}\,F$, gives the identity.

(iii) Lemma 2.5 with $k = 0$ gives $0 < W_0^a(A) \leq W_0^a(0) < \infty$. Its proof uses the coefficient of $|\varphi|^2$ only through $\int_0^\infty \mathrm{e}^{-\varepsilon^3(\frac{1}{2}a\,r^2+\lambda r^4)} r\,dr < \infty$, which holds for every real a because $\lambda > 0$. Hence $0 < \mathbb{E}_{\gamma_{\mu_0}}[W_0^a] \leq W_0^a(0)$, and (ii) with $F = 1$ gives $Z = Z_\gamma\mathbb{E}_{\gamma_{\mu_0}}[W_0^a] \in (0,\infty)$. For $F \in \mathfrak{A}_\varepsilon$ with $\mathbb{E}[\Psi_{|F|}] < \infty$, (ii) gives

$$\langle F\rangle_{\varepsilon,j,\mu_0} = \frac{\mathbb{E}_{\gamma_{\mu_0}}[\Psi_F]}{\mathbb{E}_{\gamma_{\mu_0}}[W_0]} = \frac{Z_\gamma\mathbb{E}_{\gamma_{\mu_0}}[\Psi_F]}{Z_\gamma\mathbb{E}_{\gamma_{\mu_0}}[W_0]} = Z^{-1}\int F\mathrm{e}^{-S_{\mu_0,b}}.$$

For bounded F, $\Psi_{|F|} \leq \|F\|_\infty W_0 \leq \|F\|_\infty W_0(0)$.

(iv) The bound is Lemma 2.5 again, since $\gamma^\flat$ is a probability measure (Lemma B.6(c)). Let F be bounded or non-negative. By Lemma 2.6, $g(c) := \Psi_F(A^\perp + c)$ is Borel and P-periodic in each c_μ. For a P-periodic h on $\mathbb{R}$, bounded or non-negative, $\int_t^{t+P} h = \int_0^P h$: split $[t, t+P)$ at the point of $P\mathbb{Z}$ inside it and shift one piece by P. Apply this in each coordinate (Tonelli/Fubini) and then integrate over $A^\perp$. □

B.2. **Lattice standing assumptions.**

B.2.1. *The correlation inequalities on the torus graph; monotonicity in the photon mass.* We prove the correlation inequalities of [8, Sect. 6] on an arbitrary finite graph and verify that the torus model (7.1) is an instance; this is the input of Lemma 7.5. Then we prove the monotonicity in the photon mass used in Proposition 8.7(a) and Lemma 7.6.
The inequalities of BFS I, Sect. 6 on a finite graph. [8, Sect. 6.1, p. 275] work on "an arbitrary finite lattice" with Dirichlet conditions, and remark (p. 276) that more general boundary conditions, e.g. periodic ones on rectangles, can be accommodated. We write their duplicate-variable argument [8, Sect. 6.2–6.3, pp. 277–281] for an abstract finite graph. Nothing below uses a boundary, the lattice geometry or translation invariance, so the torus with its wrap-around bonds is an instance (Proposition B.13).

Definition B.8 (finite-graph setting)**.** Let Λ be a finite set and E a finite set of *bonds*. Each $b \in E$ has endpoints $b_-, b_+ \in \Lambda$, several bonds may join the same pair, and each bond carries a coupling $c_b \geq 0$. For each $x \in \Lambda$ let ν_x be a nonzero finite measure on $[0,\infty)$ with $\int \mathrm{e}^{ar^2}\, d\nu_x(r) < \infty$ for every $a \geq 0$. Angles $\vartheta_x \in \mathbb{R}/2\pi\mathbb{Z}$ carry the normalised Lebesgue measure $d\vartheta_x/2\pi$. Let I be a finite index set, μ_C the centred Gaussian measure on $\mathbb{R}^I$ with covariance matrix $C \geq 0$ (degenerate C allowed), and for each $b \in E$ let $\bar{A}_b$ be a real linear functional on $\mathbb{R}^I$. Put

$$u_b := \vartheta_{b_+} - \vartheta_{b_-} + \bar{A}_b(A),$$
$$W := \exp\Big(\sum_{b\in E} c_b\, r_{b_-} r_{b_+} \cos u_b\Big),$$
$$d\Pi := \prod_{x\in\Lambda} d\nu_x(r_x)\frac{d\vartheta_x}{2\pi} \otimes d\mu_C(A).$$

Write $\omega = (r,\vartheta,A)$, $Z := \int W\, d\Pi$, $\langle F\rangle := Z^{-1}\int FW\, d\Pi$ and $\langle F;G\rangle := \langle FG\rangle - \langle F\rangle\langle G\rangle$. Let $\mathcal{P}_r$ be the cone of polynomials in $(r_x)_{x\in\Lambda}$ with nonnegative coefficients. For $n \in \mathbb{Z}^\Lambda$ put $n\cdot\vartheta := \sum_x n_x\vartheta_x$, and for $g \in [0,\infty)^\Lambda$ put $r^2(g) := \sum_x g_x r_x^2$. The class $\mathcal{I}$ consists of the functions $\mathrm{e}^{r^2(g)}\mathrm{e}^{i\ell(A)}$ and the class $\mathcal{D}$ of the functions $\mathrm{e}^{-r^2(g)}\mathrm{e}^{\ell(A)}$, where $g \geq 0$ and ℓ is a real linear functional on $\mathbb{R}^I$.

Write $\deg_c(x) := \sum_{b:\, b_-=x} c_b + \sum_{b:\, b_+=x} c_b$. Then $0 < W \leq \prod_x \exp(\frac{1}{2}\deg_c(x) r_x^2)$, because $r_{b_-}r_{b_+} \leq \frac{1}{2}(r_{b_-}^2 + r_{b_+}^2)$. Hence $Z \in (0,\infty)$. Moreover $|P(r)|\mathrm{e}^{r^2(g)}\mathrm{e}^{a|A|}$ is $W\, d\Pi$-integrable for every polynomial P, every $g \geq 0$ and every $a \geq 0$. This is the only integrability used below.

Lemma B.9 (angle doubling)**.** *For bounded measurable f on $(\mathbb{R}/2\pi\mathbb{Z})^2$,* $\displaystyle\int_{(\mathbb{R}/2\pi\mathbb{Z})^2} f(\vartheta,\vartheta')\, d\vartheta\, d\vartheta' = \int_{[-\pi,\pi)^2} f(\sigma-\tau,\sigma+\tau)\, d\sigma\, d\tau.$

Proof. Let $L(\sigma,\tau) := (\sigma-\tau,\sigma+\tau)$, so $|\det L| = 2$. The map L followed by reduction mod $2\pi\mathbb{Z}^2$ is a surjective homomorphism $\mathbb{R}^2 \to (\mathbb{R}/2\pi\mathbb{Z})^2$. Its kernel is $\Gamma := L^{-1}(2\pi\mathbb{Z}^2) = \{\pi(a,b) : a,b \in \mathbb{Z},\ a \equiv b \bmod 2\}$. This lattice has covolume $2\pi^2$, so $[-\pi,\pi)^2$ (area $4\pi^2$) consists of two Γ-cells. Since $f\circ L$ is Γ-periodic, $\int_{[-\pi,\pi)^2} f\circ L = 2\int_{\mathbb{R}^2/\Gamma} f\circ L = 2|\det L|^{-1}\int_{(\mathbb{R}/2\pi\mathbb{Z})^2} f$. □

Lemma B.10 ([8, Thm. 6.1] on a finite graph)**.** *In the setting of Definition B.8, let $P,Q \in \mathcal{P}_r$, $n,m \in \mathbb{Z}^\Lambda$, and let ℓ,ℓ' be real linear functionals on $\mathbb{R}^I$. Then*

$$\langle P(r)\cos(n\cdot\vartheta+\ell(A));\ Q(r)\cos(m\cdot\vartheta+\ell'(A))\rangle\ \geq 0.$$

Proof. Let $F := P(r)\cos(n\cdot\vartheta+\ell(A))$ and $G := Q(r)\cos(m\cdot\vartheta+\ell'(A))$.

Step 0 (duplication). Let $\omega' = (r',\vartheta',A')$ be an independent copy of ω. Expanding the right-hand side gives $Z^{-1}\int FGW\, d\Pi - Z^{-2}\int FW\, d\Pi \int GW\, d\Pi$, hence

$$\langle F;G\rangle = Z^{-2}\iint F(\omega)\,[G(\omega)-G(\omega')]\,W(\omega)W(\omega')\, d\Pi(\omega)\, d\Pi(\omega'). \tag{B.2}$$

Step 1 (new variables). Gauge field. Put $\alpha := \frac{1}{2}(A + A')$ and $\beta := \frac{1}{2}(A' - A)$, so that $A = \alpha - \beta$ and $A' = \alpha + \beta$. For $u, v \in \mathbb{R}^I$,

$$\mathbb{E}\, \mathrm{e}^{i\langle u,\alpha\rangle + i\langle v,\beta\rangle} = \mathbb{E}\, \mathrm{e}^{i\langle (u-v)/2, A\rangle + i\langle (u+v)/2, A'\rangle} = \mathrm{e}^{-\frac{1}{8}\langle u-v, C(u-v)\rangle - \frac{1}{8}\langle u+v, C(u+v)\rangle} = \mathrm{e}^{-\frac{1}{4}\langle u, Cu\rangle - \frac{1}{4}\langle v, Cv\rangle}.$$

Hence $\mu_C \otimes \mu_C$ is mapped to $\mu_{C/2} \otimes \mu_{C/2}$; only the fact that μ_C is a centred Gaussian is used. By linearity, $\ell(A) = \ell(\alpha) - \ell(\beta)$ and $\ell(A') = \ell(\alpha) + \ell(\beta)$, and likewise for ℓ' and $\bar{A}_b$. *Radii.* Put $s_x := \frac{1}{2}(r_x + r'_x) \geq 0$ and $t_x := \frac{1}{2}(r_x - r'_x)$, and let M_x be the image of $\nu_x \otimes \nu_x$; let $M := \prod_x M_x$. Since $t_x \mapsto -t_x$ is the swap $r_x \leftrightarrow r'_x$, M_x is invariant under $t_x \mapsto -t_x$, and it is supported in $\{s_x \geq 0\}$. It has all moments. *Angles.* All integrands below are 2π-periodic in each ϑ_x and ϑ'_x, because n, m are integer vectors. By Lemma B.9, applied site by site with Fubini, we may write $\vartheta = \sigma - \tau$ and $\vartheta' = \sigma + \tau$ with $\sigma, \tau \in [-\pi, \pi)^\Lambda$ independent and uniform. *Net effect.* $d\Pi(\omega)\, d\Pi(\omega')$ becomes $dM(s,t) \otimes dK(\sigma,\alpha) \otimes dK(\tau,\beta)$ with $K := \prod_x \frac{d\sigma_x}{2\pi} \otimes \mu_{C/2}$. Thus (σ, α) and (τ, β) have the *same* law K, independently of each other and of (s, t).

Step 2 (algebra). Put $h_b(\sigma,\alpha) := \sigma_{b_+} - \sigma_{b_-} + \bar{A}_b(\alpha)$, $X_b := h_b(\sigma,\alpha)$ and $Y_b := h_b(\tau,\beta)$; the same function h_b is used on both copies. Then $u_b(\omega) = X_b - Y_b$ and $u_b(\omega') = X_b + Y_b$. Put $\pi_b^+ := s_{b_-} s_{b_+} + t_{b_-} t_{b_+}$ and $\pi_b^- := s_{b_-} t_{b_+} + t_{b_-} s_{b_+}$. Then $r_{b_-} r_{b_+} = \pi_b^+ + \pi_b^-$ and $r'_{b_-} r'_{b_+} = \pi_b^+ - \pi_b^-$. From $\cos(X \mp Y) = \cos X \cos Y \pm \sin X \sin Y$,

$$\begin{aligned} r_{b_-} r_{b_+} \cos u_b(\omega) + r'_{b_-} r'_{b_+} \cos u_b(\omega') &= 2\pi_b^+ \cos X_b \cos Y_b + 2\pi_b^- \sin X_b \sin Y_b, \\ r_{b_-} r_{b_+} \cos u_b(\omega) - r'_{b_-} r'_{b_+} \cos u_b(\omega') &= 2\pi_b^- \cos X_b \cos Y_b + 2\pi_b^+ \sin X_b \sin Y_b, \end{aligned} \tag{B.3}$$

so that $W(\omega)W(\omega') = \prod_b \exp\big(2c_b[\pi_b^+ \cos X_b \cos Y_b + \pi_b^- \sin X_b \sin Y_b]\big)$. For $P \in \mathcal{P}_r$ write $P(s+t) = P_e + P_o$ and $P(s-t) = P_e - P_o$, where P_e (P_o) collects the monomials of even (odd) total degree in t. Expanding $(s_x + t_x)^k$ shows that P_e, P_o lie in the cone $\mathcal{P}_{s,t}$ of polynomials in (s, t) with nonnegative coefficients. With $X_F := n \cdot \sigma + \ell(\alpha)$, $Y_F := n \cdot \tau + \ell(\beta)$, $X_G := m \cdot \sigma + \ell'(\alpha)$ and $Y_G := m \cdot \tau + \ell'(\beta)$,

$$F(\omega) = P(s+t)\,[\cos X_F \cos Y_F + \sin X_F \sin Y_F], \qquad G(\omega) - G(\omega') = 2Q_o \cos X_G \cos Y_G + 2Q_e \sin X_G \sin Y_G.$$

Step 3 (the paired cone). Let $\mathcal{K}$ be the set of finite sums $\sum_k p_k(s,t)\, \Phi_k(\sigma,\alpha)\, \Phi_k(\tau,\beta)$, where $p_k \in \mathcal{P}_{s,t}$ and each Φ_k is a real measurable function on $[-\pi,\pi)^\Lambda \times \mathbb{R}^I$ with $|\Phi_k(\sigma,\alpha)| \leq c_k \mathrm{e}^{a_k|\alpha|}$. The *same* Φ_k is evaluated on the two copies. $\mathcal{K}$ is closed under sums and products, since $(p_1\Phi_1\Phi'_1)(p_2\Phi_2\Phi'_2) = (p_1p_2)(\Phi_1\Phi_2)(\Phi_1\Phi_2)'$. For every finite positive measure M' that has all moments, is a product over x, is invariant under each $t_x \mapsto -t_x$ and is supported in $\{s \geq 0\}$, and every $K \in \mathcal{K}$,

$$\int K\, dM'\, dK\, dK = \sum_k \Big(\int p_k\, dM' \Big) \Big(\int \Phi_k\, dK \Big)^2 \;\geq 0. \tag{B.4}$$

The two Φ_k-factors are integrated against the same law K, independently, which gives the square. Moreover $\int p_k\, dM' \geq 0$: M' is a product measure, so the integral of a monomial $s^a t^j$ factorises over x. Each factor vanishes if j_x is odd, by the symmetry $t_x \mapsto -t_x$, and is ≥ 0 if j_x is even, because $s_x \geq 0$.

Step 4 (assembly). By Steps 1–2 the integrand of (B.2) is

$$\begin{aligned} P(s+t)[\cos X_F \cos Y_F + \sin X_F \sin Y_F] \cdot [2Q_o \cos X_G \cos Y_G + 2Q_e \sin X_G \sin Y_G] \\ \cdot \prod_b \exp\big(2c_b[\pi_b^+ \cos X_b \cos Y_b + \pi_b^- \sin X_b \sin Y_b]\big). \end{aligned}$$

Expanding each exponential in its power series gives a series of elements of $\mathcal{K}$, because $c_b \geq 0$, $\pi_b^\pm, P_e, P_o, Q_e, Q_o \in \mathcal{P}_{s,t}$, and the Φ's are products of cos and sin of X's, hence bounded. Put $R_x := s_x + |t_x| = \max(r_x, r'_x)$. Since $s \geq 0$, $|\pi_b^+| + |\pi_b^-| \leq R_{b_-} R_{b_+}$. Hence the series of absolute values is dominated by

$$P(R) \cdot 2Q(R) \cdot \exp\Big(\sum_b 2c_b R_{b_-} R_{b_+} \Big) \leq P(R) \cdot 2Q(R) \cdot \prod_x \exp\big(\deg_c(x)(r_x^2 + r_x'^2) \big),$$

where we used $R_x^2 \le r_x^2 + r_x'^2$. The right side is $\nu_x \otimes \nu_x$-integrable by the Gaussian moments. By dominated convergence the integral in (B.2) is the sum of the integrals of the terms, and each is ≥ 0 by (B.4) with $M' = M$. Since $Z > 0$, $\langle F; G\rangle \ge 0$. □

Lemma B.11 ([8, Thm. 6.2] on a finite graph)**.** *In the setting of Definition B.8, let $g \ge 0$, let ℓ be a real linear functional and k a complex linear functional on $\mathbb{R}^I$, and let $b \in E$. Then*

$$\textit{(1)}\ \langle \mathrm{e}^{r^2(g)}\mathrm{e}^{i\ell(A)}; |k(A)|^2\rangle \le 0, \quad \textit{(2)}\ \langle \mathrm{e}^{r^2(g)}\mathrm{e}^{i\ell(A)}; r_{b_-}r_{b_+}\cos u_b\rangle \ge 0,$$

$$\textit{(3)}\ \langle \mathrm{e}^{-r^2(g)}\mathrm{e}^{\ell(A)}; |k(A)|^2\rangle \ge 0, \quad \textit{(4)}\ \langle \mathrm{e}^{-r^2(g)}\mathrm{e}^{\ell(A)}; r_{b_-}r_{b_+}\cos u_b\rangle \le 0.$$

The expectations of elements of $\mathcal{I}$ are real.

Proof. Symmetry. The map $(A, \vartheta) \mapsto (-A, -\vartheta)$ preserves $d\Pi$, because μ_C is centred and the angles are uniform. It preserves W, since $u_b \mapsto -u_b$, and it leaves $r^2(g)$, $|k(A)|^2$ and $r_{b_-}r_{b_+}\cos u_b$ invariant, while $\ell(A) \mapsto -\ell(A)$. Hence the $\sin \ell(A)$-parts drop out: in (1)–(2), $\mathrm{e}^{i\ell(A)}$ may be replaced by $\cos \ell(A)$, and $\langle \mathrm{e}^{r^2(g)}\mathrm{e}^{i\ell(A)}\rangle$ is real.

(2) Apply Lemma B.10 with $P_N := \sum_{j\le N} r^2(g)^j/j! \in \mathcal{P}_r$, $n = 0$ and ℓ, and with $Q = r_{b_-}r_{b_+}$, $m = \mathbf{e}_{b_+} - \mathbf{e}_{b_-}$ and $\ell' = \bar{A}_b$; here $\mathbf{e}_x$ is the indicator vector of x, and $m \cdot \vartheta + \bar{A}_b(A) = u_b$. Then let $N \to \infty$ by dominated convergence, since $|P_N| \le \mathrm{e}^{r^2(g)}$.

(1) Let first k be real. Then $k(A)^2 = \lim_{\eta\downarrow 0} 2\eta^{-2}(1 - \cos \eta k(A))$ with $0 \le 2\eta^{-2}(1 - \cos \eta k) \le k^2$. Since $\langle F; 1\rangle = 0$,

$$\langle F; k(A)^2\rangle = -\lim_{\eta\downarrow 0} 2\eta^{-2}\langle F; \cos \eta k(A)\rangle \le 0$$

for $F = \mathrm{e}^{r^2(g)}\cos \ell(A)$, by Lemma B.10 (with $Q = 1$, $m = 0$, $\ell' = \eta k$) and the limit $P_N \to \mathrm{e}^{r^2(g)}$. For complex $k = k_1 + ik_2$ use $|k|^2 = k_1^2 + k_2^2$ [8, (6.24)–(6.25)].

(3), (4) This is the exchange of the two copies [8, p. 281]. Relabelling $\omega \leftrightarrow \omega'$ in (B.2) gives

$$\langle F; G\rangle = -Z^{-2}\iint F(\omega')\,[G(\omega) - G(\omega')]\,WW'\,d\Pi\,d\Pi.$$

Take $F = \mathrm{e}^{-r^2(g)}\mathrm{e}^{\ell(A)}$. Since $r' = s - t$ and $A' = \alpha + \beta$,

$$F(\omega') = \mathrm{e}^{-s^2(g)-t^2(g)}\,\mathrm{e}^{2(st)(g)}\,\mathrm{e}^{\ell(\alpha)}\mathrm{e}^{\ell(\beta)}, \qquad (st)(g) := \textstyle\sum_x g_x s_x t_x.$$

The measure $dM^g := \mathrm{e}^{-s^2(g)-t^2(g)}dM$ is again a product, invariant under each $t_x \mapsto -t_x$, supported in $\{s \ge 0\}$, and has all moments. The factor $\mathrm{e}^{2(st)(g)}$ is a series with terms in $\mathcal{P}_{s,t}$, since $g \ge 0$. The factor $\mathrm{e}^{\ell(\alpha)}\mathrm{e}^{\ell(\beta)}$ is a paired product $\Phi(\alpha)\Phi(\beta)$ with $|\Phi| \le \mathrm{e}^{\|\ell\|\|\alpha|}$. Finally $G - G'$ is as in Step 2. So the integrand is a series of elements of $\mathcal{K}$. Its absolute series is dominated as in Step 4 times $\mathrm{e}^{|\ell(\alpha)|+|\ell(\beta)|}$, using

$$\mathrm{e}^{2(s|t|)(g)}\mathrm{e}^{-(s^2+t^2)(g)} = \mathrm{e}^{-(s-|t|)^2(g)} \le 1$$

(the series of absolute values of $\mathrm{e}^{2(st)(g)}$ is $\mathrm{e}^{2(s|t|)(g)}$). This is integrable. By (B.4) with $M' = M^g$, $\langle \mathrm{e}^{-r^2(g)}\mathrm{e}^{\ell(A)}; Q(r)\cos(m\cdot\vartheta + \ell'(A))\rangle \le 0$ for all $Q \in \mathcal{P}_r$, $m \in \mathbb{Z}^\Lambda$ and real ℓ'. The choice $Q = r_{b_-}r_{b_+}$, $m = \mathbf{e}_{b_+} - \mathbf{e}_{b_-}$, $\ell' = \bar{A}_b$ gives (4). The choice $Q = 1$, $m = 0$, $\ell' = \eta k$, followed by the limit of (1), gives (3). □

Lemma B.12 ([8, Cor. 6.3] on a finite graph)**.** *In the setting of Definition B.8, let $g \ge 0$ and let ℓ be real linear.*

(a) *If $0 \le C_0 \le C_1$ in the positive semidefinite order on $\mathbb{R}^I$, and everything else is fixed, then $\langle \mathrm{e}^{r^2(g)}\mathrm{e}^{i\ell(A)}\rangle_{C_1} \le \langle \mathrm{e}^{r^2(g)}\mathrm{e}^{i\ell(A)}\rangle_{C_0}$ and $\langle \mathrm{e}^{-r^2(g)}\mathrm{e}^{\ell(A)}\rangle_{C_1} \ge \langle \mathrm{e}^{-r^2(g)}\mathrm{e}^{\ell(A)}\rangle_{C_0}$.*

(b) *For each b, $c_b \mapsto \langle \mathrm{e}^{r^2(g)}\mathrm{e}^{i\ell(A)}\rangle$ is non-decreasing and $c_b \mapsto \langle \mathrm{e}^{-r^2(g)}\mathrm{e}^{\ell(A)}\rangle$ is non-increasing on $[0, \infty)$.*

(c) *If $E' \subset E$ and the system on E' is obtained by setting $c_b = 0$ for $b \in E \setminus E'$, then the $\mathcal{I}$-expectations on E' are $\le$ those on E, and the $\mathcal{D}$-expectations are $\ge$.*

Proof. (a) Let first C_0, C_1 be positive definite. Put $C_\theta := (1-\theta)C_0 + \theta C_1$ for $\theta \in [0,1]$ and $D := C_1 - C_0 \geq 0$. With $c_- := \min(\lambda_{\min}(C_0), \lambda_{\min}(C_1)) > 0$ and $c_+ := \max(\|C_0\|, \|C_1\|)$ we have $c_- \leq C_\theta \leq c_+$. Let $\varrho_\theta(A) = (2\pi)^{-|I|/2} \det(C_\theta)^{-1/2} \mathrm{e}^{-\frac{1}{2}\langle A, C_\theta^{-1} A\rangle}$ be the density of μ_{C_θ}. Then

$$\partial_\theta \log \varrho_\theta = -\tfrac{1}{2} \operatorname{tr}(C_\theta^{-1} D) + \tfrac{1}{2} \langle A,\, C_\theta^{-1} D C_\theta^{-1} A\rangle,$$

and hence $|\partial_\theta \varrho_\theta(A)| \leq (2\pi c_-)^{-|I|/2} \big(\tfrac{1}{2}|I|\|D\|/c_- + \tfrac{1}{2}\|D\||A|^2/c_-^2\big) \mathrm{e}^{-|A|^2/(2c_+)}$, uniformly in $\theta \in [0,1]$. For F in either class, $\Phi_F(A) := \int F\, W \prod_x d\nu_x \frac{d\vartheta_x}{2\pi}$ satisfies $|\Phi_F(A)| \leq c_F \mathrm{e}^{\|\ell\||A|}$: we have $|\mathrm{e}^{i\ell}| = 1$, $\mathrm{e}^{-r^2(g)} \leq 1$, and W is bounded uniformly in A as above. So numerator and denominator of $\langle F\rangle_{C_\theta}$ are C^1 in θ, with derivatives taken under the integral. The constant part of $\partial_\theta \log \varrho_\theta$ cancels in the normalised expectation, and we obtain

$$\partial_\theta \langle F\rangle_{C_\theta} = \tfrac{1}{2} \langle F;\, \langle A,\, C_\theta^{-1} D C_\theta^{-1} A\rangle\rangle_{C_\theta} = \tfrac{1}{2} \sum_j d_j \langle F;\, k_j(A)^2\rangle_{C_\theta},$$

where $D = \sum_j d_j v_j v_j^{\mathsf{T}}$ with $d_j \geq 0$ and v_j orthonormal, and $k_j(A) := v_j^{\mathsf{T}} C_\theta^{-1} A$ is real linear. By Lemma B.11(1), resp. (3), applied at covariance C_θ, this is ≤ 0 for $F \in \mathcal{I}$ and ≥ 0 for $F \in \mathcal{D}$. Integrate over $\theta \in [0,1]$.

If C_0, C_1 are only positive semidefinite, apply this to $C_i + \eta \mathbf{1}$ and let $\eta \downarrow 0$. The map $C \mapsto \int \Phi_F \, d\mu_C = \mathbb{E}[\Phi_F(C^{1/2}\xi)]$, with ξ standard normal on $\mathbb{R}^I$, is continuous on positive semidefinite C. Indeed, Φ_F is continuous by dominated convergence, $C \mapsto C^{1/2}$ is continuous, and $|\Phi_F(C^{1/2}\xi)| \leq c_F \mathrm{e}^{\|\ell\|\|C^{1/2}\||\xi|}$ is uniformly integrable for C in bounded sets. The same holds for $F = 1$, whose integral is > 0.

(b) $\partial_{c_b} W = r_{b_-} r_{b_+} \cos u_b \cdot W$. The derivative is dominated, for c_b in a bounded interval, by $r_{b_-} r_{b_+} \prod_x \mathrm{e}^{\frac{1}{2}(\deg_c(x)+1) r_x^2} |F|$, which is integrable. Hence $\partial_{c_b} \langle F\rangle = \langle F; r_{b_-} r_{b_+} \cos u_b\rangle$, and the sign follows from Lemma B.11(2), resp. (4). At $c_b = 0$, continuity follows by dominated convergence. (c) follows by iterating (b). □

Proposition B.13 (the torus model is an instance)**.** *Fix (ε, T), a lattice torus $T_{j,\varepsilon}$ with $N \geq 2$ sites per side, and the matter action S_m of* (7.1)*. Let μ_C be a centred Gaussian on the bond variables. Here $C \geq 0$ is its covariance operator for $\langle\cdot,\,\cdot\rangle_\varepsilon$, as in* (7.3)*; the corresponding covariance matrix of the coordinates (A_b) is $\varepsilon^{-3} C$, so the two positive semidefinite orders agree. Then the state $\langle\cdot\rangle_C$ of Lemma 7.5, i.e.* (7.3) *with $\mu_{C_{\mu_0}}$ replaced by μ_C, is the state $\langle\cdot\rangle$ of Definition B.8 with the following data:*

- *$\Lambda = T_{j,\varepsilon}$, and E = all $3N^3$ oriented bonds, wrap-around bonds included;*
- *$c_b = \varepsilon$ and $\bar{A}_b(A) = \varepsilon e A_b$;*
- *$d\nu_x(r) = r\, \mathrm{e}^{-V(r)} dr$ with $V(r) = \varepsilon^3(\frac{1}{2} b_T r^2 + \lambda r^4) + 3\varepsilon r^2$, and I = bonds.*

The same holds with any of the c_b lowered to a value in $[0, \varepsilon]$. Consequently Lemmas B.10, B.11 and B.12 hold on every lattice torus of the paper, for every $C \geq 0$, in particular for $C = C_{\mu_0}$ with $\mu_0 \in (0, \mu_*]$.

Proof. Write $\varphi_x = r_x(\cos\vartheta_x, \sin\vartheta_x)$, so that $d\varphi_x = r_x\, dr_x\, d\vartheta_x$. Expanding $\frac{1}{2}\varepsilon^3 |D_A \varphi(b)|^2 = \frac{1}{2}\varepsilon(r_{b_+}^2 + r_{b_-}^2) - \varepsilon\, r_{b_-} r_{b_+} \cos(\vartheta_{b_+} - \vartheta_{b_-} + \varepsilon e A_b)$ gives

$$\mathrm{e}^{-S_m} = \prod_x \mathrm{e}^{-V(r_x)} \prod_b \exp\big(\varepsilon r_{b_-} r_{b_+} \cos u_b\big), \qquad u_b = \vartheta_{b_+} - \vartheta_{b_-} + \varepsilon e A_b,$$

as in the proof of Lemma 2.5. Here the six diagonal halves $\frac{1}{2}\varepsilon r_x^2$ at each site give the term $3\varepsilon r^2$ in V. By (P2), $\lambda > 0$, so ν_x is nonzero and has Gaussian moments of all orders. The factors 2π from $d\vartheta_x$ cancel in normalised expectations. Covariance operator versus covariance matrix: if $E[\langle f, A\rangle_\varepsilon \langle f', A\rangle_\varepsilon] = \langle f, Cf'\rangle_\varepsilon$, then with $f = \varepsilon^{-3}\delta_b$ and $f' = \varepsilon^{-3}\delta_{b'}$ one gets $E[A_b A_{b'}] = \varepsilon^{-3} C_{bb'}$. Nothing in Definition B.8 or in the proofs of Lemmas B.10–B.12 refers to how E sits in a lattice.

Multiple bonds between one pair of sites (at $N = 2$) and the wrap-around bonds are bonds like any other. □

Lemma 7.5 is Lemma B.12(a) in the instance of Proposition B.13. In the proof of Lemma 7.5, the citations [8, Thm. 6.2(1)] and [8, Thm. 6.2(3)] may be read as Lemma B.11(1) and (3) on the torus graph, which removes the reliance on the periodic-boundary remark of [8, p. 276]. The differentiation in the interpolation parameter is justified by the domination written in the proof of Lemma B.12(a). The inequalities compare two systems on *one* bond set and one index set; they give no comparison between tori of different sizes, and none is used in the paper.
Monotonicity of the photon covariance in the photon mass.

Lemma B.14 (operator monotonicity)**.**

(a) *Let Q, Q' be symmetric operators on a finite-dimensional real inner-product space with $0 < Q' \le Q$. Then $Q^{-1} \le Q'^{-1}$.*

(b) *Fix (ε, T). For $0 < \mu' \le \mu$, the photon covariances $C_\mu = (-\Delta_\varepsilon + \mu^2)^{-1}$ of (7.3) satisfy $C_\mu \le C_{\mu'}$ as quadratic forms on $(\mathbb{R}^{bonds}, \langle\cdot,\cdot\rangle_\varepsilon)$, the zero mode included. On the constant 1-forms the inequality reads $\mu^{-2} \le \mu'^{-2}$. Equivalently, the variances $s(\mu) = 1/(\mu^2|T|)$ of the zero-mode coordinates c_μ are non-increasing in μ.*

Part (b) is the monotonicity of C_μ in μ, the zero mode included, used in the proof of Proposition 8.7(a), and the inequality $C_{\mu_0} \ge C_{\mu_*}$ for $\mu_0 \le \mu_*$ used in the proof of Lemma 7.6. In both places Lemma 7.5 is then applied with $C_0 = C_\mu$ and $C_1 = C_{\mu'}$.

Proof. (a) Since $Q > 0$, $Q^{\pm 1/2}$ exist. From $Q' \le Q$ we get $B := Q^{-1/2} Q' Q^{-1/2} \le \mathbf{1}$, and $B > 0$. A positive symmetric $B \le \mathbf{1}$ has its eigenvalues in $(0, 1]$, so $B^{-1} = Q^{1/2} Q'^{-1} Q^{1/2} \ge \mathbf{1}$. Conjugating by $Q^{-1/2}$ gives $Q'^{-1} \ge Q^{-1}$.

(b) $-\Delta_\varepsilon = d^*d + dd^*$ is symmetric for $\langle\cdot,\cdot\rangle_\varepsilon$ and ≥ 0, by Lemma B.4(i),(ii). So $0 < -\Delta_\varepsilon + \mu'^2 \le -\Delta_\varepsilon + \mu^2$, and (a) gives $C_\mu \le C_{\mu'}$. Explicitly, let (v_k) be an $\langle\cdot,\cdot\rangle_\varepsilon$-orthonormal eigenbasis of $-\Delta_\varepsilon$ with eigenvalues $\kappa_k \ge 0$. Then

$$\langle f,\, (C_{\mu'} - C_\mu) f\rangle_\varepsilon = \sum_k \big[(\kappa_k + \mu'^2)^{-1} - (\kappa_k + \mu^2)^{-1}\big] \langle v_k,\, f\rangle_\varepsilon^2 \;\ge 0,$$

with the terms $\kappa_k = 0$ included. The constant 1-forms $\mathbf{1}_\mu$ span the eigenvalue 0 (Lemma B.4(ii)), where $C_\mu \mathbf{1}_\mu = \mu^{-2} \mathbf{1}_\mu$, and the variance of c_μ is $s(\mu) = 1/(\mu^2|T|)$ by Lemma B.6(b), in agreement with (7.3)ff. □

B.2.2. *Properties (P1)–(P3), the zero-mode split and the constants of Theorem 6.8.* Throughout, the notation is that of Section 7: $T = T_j$, $\ell_T = \ell_j = j!\,\ell_0$, $|T| = \ell_T^3$, $N = \ell_T/\varepsilon$, $-\Delta_\varepsilon = d^*d + dd^*$ the componentwise lattice Laplacian on 1-forms, P_0 the $\langle\cdot,\cdot\rangle_\varepsilon$-orthogonal projection onto the constant 1-forms, $P^\perp := 1 - P_0$, and $\mathbf{1}_\mu$ the constant 1-form of direction μ (Subsection 7.1).
Property (P1); the per-torus coefficient.
Property (P1); the coefficient is used torus by torus. *Notation.* Fix $j \ge 1$ and $\varepsilon = \varepsilon_K$, $K \ge K_0$, and write $\Lambda := T_{j,\varepsilon}$. Let $\mathcal{H} := (\mathbb{R}^2)^\Lambda$ with the inner product $\langle u,\, v\rangle_\varepsilon = \sum_{x\in\Lambda} \varepsilon^3\, u(x) \cdot v(x)$. The Higgs field φ is an element of $\mathcal{H}$ through $\varphi(x) = (\operatorname{Re}\varphi(x), \operatorname{Im}\varphi(x))$. For $a \in \varepsilon\mathbb{Z}^3$ let T_a be the translation: $(\mathsf{T}_a u)(x) := u(x - a)$ on $\mathcal{H}$, and $(\mathsf{T}_a A)_{x,\mu} := A_{x-a,\mu}$ on bond functions. A real 2×2 matrix Q acts on $\mathcal{H}$ pointwise, $(Qu)(x) := Qu(x)$. We use the rotations $U(\theta)$, the complex conjugation $\mathsf{K} := \operatorname{diag}(1, -1)$, and $J := U(\pi/2)$. For a real function f on Λ, M_f is the multiplication operator $(\mathsf{M}_f u)(x) = f(x)u(x)$. For $v \in \mathbb{R}^2$, $\mathbf{1} \otimes v \in \mathcal{H}$ is the constant function $x \mapsto v$. For a linear operator Y on $\mathcal{H}$ and $x \in \Lambda$, the *zero-momentum value* $\hat{Y}(x)$ is the 2×2 matrix $v \mapsto \big(Y(\mathbf{1} \otimes v)\big)(x)$. If $(Yu)(x) = \sum_y \varepsilon^3 Y(x, y) u(y)$, then $\hat{Y}(x) = \sum_{y\in\Lambda} \varepsilon^3 Y(x, y)$, which is the sum in [24, (3.33), p. 659].

Lemma B.15 (the lines commute with the symmetries)**.** *For $M > 0$ put $C_M := (-\Delta_\varepsilon + M^2)^{-1}$, acting on $\mathcal{H}$ componentwise, and $C^A := (-\Delta_\varepsilon + \mu_*^2)^{-1}$, acting componentwise on bond functions. Then C_M commutes with T_a ($a \in \varepsilon\mathbb{Z}^3$), with $U(\theta)$ ($\theta \in \mathbb{R}$) and with K, and C^A commutes with T_a.*

Proof. $(\Delta_\varepsilon u)(x) = \varepsilon^{-2}\sum_{\mu=1}^{3}\big(u(x+\varepsilon e_\mu) + u(x-\varepsilon e_\mu) - 2u(x)\big)$. The map $x \mapsto x - a$ is a bijection of Λ that commutes with $x \mapsto x \pm \varepsilon e_\mu$, so $\Delta_\varepsilon \mathsf{T}_a = \mathsf{T}_a \Delta_\varepsilon$. Since Δ_ε acts on each real component separately, it commutes with every pointwise matrix. The photon operator is $-\Delta_\varepsilon$ on each component $A_{\cdot,\mu}$ (Lemma B.4), so it also commutes with T_a. The operators $-\Delta_\varepsilon + M^2 \geq M^2$ and $-\Delta_\varepsilon + \mu_*^2 \geq \mu_*^2$ are invertible. If Y commutes with an invertible W, then $W^{-1}Y = W^{-1}YWW^{-1} = W^{-1}WYW^{-1} = YW^{-1}$. □

Formal perturbation theory on Λ. Let $I := \{(\alpha,\beta) \in \mathbb{N}_0^2 : 2 \leq \alpha + 2\beta \leq 4\}$, and give the monomial $e^\alpha \lambda^\beta$ the weight $w(\alpha,\beta) := \alpha + 2\beta$. Let $\mathfrak{d} = (\delta m_\gamma^2)_{\gamma\in I}$ be a family of real functions on Λ; a priori they may depend on the site, as in [24, (3.32), p. 659]. Put $\delta m_\gamma^2 := 0$ for $\gamma \notin I$, and $\delta m^2 := \sum_{\gamma\in I} e^\alpha \lambda^\beta \delta m_\gamma^2$. For $M > 0$, summation by parts on the torus and the identity $\frac{1}{2}\varepsilon^{-2}|\mathrm{e}^{is}u - v|^2 = \frac{1}{2}\varepsilon^{-2}|u-v|^2 + \varepsilon^{-2}\operatorname{Re}\big(\bar v u(1-\mathrm{e}^{is})\big)$ give

$$S_{\mu_*, M^2+\delta m^2}(A,\varphi) = \tfrac{1}{2}\langle A,\, (-\Delta_\varepsilon + \mu_*^2)A\rangle_\varepsilon + \tfrac{1}{2}\langle \varphi,\, (-\Delta_\varepsilon + M^2)\varphi\rangle_\varepsilon + V(A,\varphi),$$

with $S_{\mu_0,a}$ as in (2.1), $V := V^e + \lambda V^4 + V^{\mathfrak{d}}$, and

$$V^e := \sum_{x,\mu} \varepsilon \operatorname{Re}\big(\bar\varphi(x)\varphi(x+\varepsilon e_\mu)(1 - \mathrm{e}^{ie\varepsilon A_{x,\mu}})\big), \quad V^4 := \sum_x \varepsilon^3 |\varphi(x)|^4, \quad V^{\mathfrak{d}} := \tfrac{1}{2}\langle \varphi,\, \mathsf{M}_{\delta m^2}\varphi\rangle_\varepsilon.$$

V^e is entire in e, and $V^e = \sum_{k\geq 1} e^k V_k^e$ with the polynomials $V_k^e := -\sum_{x,\mu} \varepsilon \operatorname{Re}\big(\bar\varphi(x)\varphi(x+\varepsilon e_\mu)(i\varepsilon A_{x,\mu})^k/k!\big)$. So $V = \sum_{\gamma\neq(0,0)} e^\alpha\lambda^\beta V_\gamma$ with polynomials V_γ, and $V_{(0,0)} = 0$. Let γ_M be the centred Gaussian measure on $\mathbb{R}^{\text{bonds}} \times \mathcal{H}$ under which A and φ are independent, with covariances C^A and C_M for the pairings $\langle\cdot,\cdot\rangle_\varepsilon$. For a polynomial F in (A,φ), the *formal expectation* $\langle F\rangle^{\mathrm{f}}$ is the formal power series in (e,λ) obtained from $\sum_{k\geq 0}\frac{1}{k!}\mathbb{E}_{\gamma_M}[F\,(-V)^k]$ by expanding V in monomials. Its coefficient of weight n is a finite sum of Gaussian moments of polynomials: every V_γ has weight ≥ 1, so only $k \leq n$ contribute. These coefficients are those of the perturbation expansions that define the counterterms of [24, (3.32)–(3.35), p. 659] and [3, (1.21)–(1.24), pp. 416–417]. Wick's theorem writes them as sums of lattice Feynman graphs whose lines are the kernels of C_M and C^A. On the periodic torus these kernels are King's full propagators C_s, C_v [24, p. 658] with photon mass μ_*, and Bałaban's C_0^ε, C^ε [3, p. 417]. The block-averaged operators G_k, $G_k(0)$ of [24] enter only through the splitting $C_v = G_k + F_{k,v}$, $C_s = G_k(0) + F_{k,s}$ [24, p. 658], which leaves the sum of the graphs unchanged.

The *two-point function* $G = G[\mathfrak{d}]$ is the formal power series, with coefficients in the linear operators on $\mathcal{H}$, given by $\langle u, Gv\rangle_\varepsilon := \langle\langle u,\varphi\rangle_\varepsilon\langle v,\varphi\rangle_\varepsilon\rangle^{\mathrm{f}}/\langle 1\rangle^{\mathrm{f}}$. Here $\langle 1\rangle^{\mathrm{f}}$ has constant term 1, and $G_{(0,0)} = C_M$. Since C_M is invertible, G is invertible in this ring of formal power series. We put

$$\Sigma = \Sigma[\mathfrak{d}] := C_M^{-1} - G[\mathfrak{d}]^{-1}, \qquad \text{equivalently} \qquad G = \sum_{n\geq 0} C_M\big(\Sigma\, C_M\big)^n.$$

This is the full amputated one-particle-irreducible two-point function. The second identity is the form [3, (1.21), p. 416] of the expansion of the two-point function, where $\Sigma = -\delta m^2 + \Sigma^\varepsilon + \partial^{\varepsilon *}\Sigma_1^\varepsilon + \Sigma_1^{\varepsilon *}\partial^\varepsilon + \partial^{\varepsilon *}\Sigma_2^\varepsilon\partial^\varepsilon$. In the notation of [24, p. 659], $\Sigma = -\delta m^2 + \Gamma^\varepsilon$. The counterterm condition of [24, (3.33), p. 659] is

$$\text{(K)} \qquad \hat\Sigma[\mathfrak{d}]_\gamma(x) = 0 \qquad \text{for all } \gamma \in I,\ x \in \Lambda.$$

Proposition B.16 (the torus counterterm is one number)**.** *For every $M > 0$ there is exactly one family $\mathfrak{d} = (\delta m_\gamma^2)_{\gamma\in I}$ of real functions on Λ that satisfies* (K)*. Each δm_γ^2 is constant on Λ. Its value depends only on (ε, j, M) and on the reference mass μ_*, and it vanishes when α is odd*

(Lemma B.2). Consequently the counterterm $\delta m^2_{K,T_j}(\varepsilon;M)=\sum_{\gamma\in I} e^{\alpha}\lambda^{\beta}\delta m^2_{\gamma}$ *of Definition 2.2 and* $a_{K,T_j}(M)=M^2+\delta m^2_{K,T_j}(\varepsilon;M)$ *are real numbers, not functions of the site.*

Proof. Step 1: the top-order counterterm enters linearly and alone. Let $\gamma=(\alpha,\beta)$ with $n:=w(\gamma)\geq 1$. Every coefficient of weight n of $\langle F\rangle^{\mathrm{f}}$ is a sum of terms $\frac{(-1)^k}{k!}\mathbb{E}_{\gamma_M}[F\,V_{\gamma_1}\cdots V_{\gamma_k}]$ with $\gamma_1+\cdots+\gamma_k=\gamma$ and $w(\gamma_i)\geq 1$. The function δm^2_γ occurs only in V_γ, and at weight n the factor V_γ occurs only in the term with $k=1$. Every other term contains only factors $V_{\gamma'}$ with $w(\gamma')\leq n-1$. For a symmetric operator B on $\mathcal{H}$, Wick's theorem gives

$$\mathbb{E}_{\gamma_M}\big[\langle u,\,\varphi\rangle_\varepsilon\langle v,\,\varphi\rangle_\varepsilon\,\tfrac12\langle\varphi,\,B\varphi\rangle_\varepsilon\big]-\langle u,\,C_Mv\rangle_\varepsilon\,\mathbb{E}_{\gamma_M}\big[\tfrac12\langle\varphi,\,B\varphi\rangle_\varepsilon\big]=\langle u,\,C_MBC_Mv\rangle_\varepsilon.$$

Write $G=N/Z$ with $N_{(0,0)}=C_M$ and $Z_{(0,0)}=1$, and apply this identity with $B=\mathsf{M}_{\delta m^2_\gamma}$. This gives $G_\gamma=-C_M\mathsf{M}_{\delta m^2_\gamma}C_M+R_\gamma$, where R_γ is a polynomial in the coefficients of V of weight $\leq n-1$. In particular R_γ depends on $\mathfrak{d}$ only through the $\delta m^2_{\gamma'}$ with $w(\gamma')<n$. Next, $G^{-1}=\sum_{k\geq 0}C_M^{-1}\big(-(G-C_M)C_M^{-1}\big)^k$. Its coefficient at γ is $-C_M^{-1}G_\gamma C_M^{-1}$ plus products $G_{\gamma_1}\cdots G_{\gamma_k}$, with factors C_M^{-1} in between, where $k\geq 2$, $\sum\gamma_i=\gamma$ and $1\leq w(\gamma_i)\leq n-1$. Hence

$$\Sigma[\mathfrak{d}]_\gamma=-\mathsf{M}_{\delta m^2_\gamma}+\Gamma_\gamma, \tag{B.5}$$

where Γ_γ depends only on the $\delta m^2_{\gamma'}$ with $w(\gamma')<n$. Let $\mathfrak{d}^{<n}$ be the family obtained from $\mathfrak{d}$ by setting $\delta m^2_{\gamma'}=0$ for $w(\gamma')\geq n$. Applied to $\mathfrak{d}^{<n}$, (B.5) has the same Γ_γ and a vanishing first term. Therefore $\Gamma_\gamma=\Sigma[\mathfrak{d}^{<n}]_\gamma$.

Step 2: symmetries. Let g be one of the linear maps $(A,\varphi)\mapsto(A,U(\theta)\varphi)$ or $(A,\varphi)\mapsto(-A,\mathsf{K}\varphi)$, or, if every $\delta m^2_{\gamma'}$ is constant on Λ, the map $(A,\varphi)\mapsto(\mathsf{T}_aA,\mathsf{T}_a\varphi)$. Each such g is orthogonal for $\langle\cdot,\,\cdot\rangle_\varepsilon$. By Lemma B.15 it commutes with the covariance of γ_M (for $A\mapsto -A$ this is trivial), so g preserves γ_M. Moreover $V_{\gamma'}\circ g=V_{\gamma'}$ for every γ', for the following reasons. V^4 and $V^{\mathfrak{d}}$ depend on φ only through $|\varphi(x)|^2$, and δm^2 is translation invariant when it is constant. $\bar\varphi(x)\varphi(y)$ is invariant under $U(\theta)$. Under $(A,\varphi)\mapsto(-A,\bar\varphi)$ the summand of V^e becomes $\mathrm{Re}\,\big(\overline{\bar\varphi(x)\varphi(y)(1-\mathrm{e}^{ie\varepsilon A})}\big)$ for every real e, hence each V^e_k is invariant. A translation permutes the summands of V^e, V^4 and $V^{\mathfrak{d}}$. Hence $\langle F\circ g\rangle^{\mathrm{f}}=\langle F\rangle^{\mathrm{f}}$, coefficient by coefficient. With $F=\langle u,\,\varphi\rangle_\varepsilon\langle v,\,\varphi\rangle_\varepsilon$ this says that every coefficient of G commutes with the action of g on $\mathcal{H}$ (by $U(\theta)$, K or T_a). The same then holds for G^{-1}. Since C_M commutes too, it holds for every coefficient of Σ.

Step 3: zero-momentum values. Let Y be a linear operator on $\mathcal{H}$ that commutes with all $U(\theta)$ and with K. Since $U(\theta)(\mathbf{1}\otimes v)=\mathbf{1}\otimes U(\theta)v$, we get $\hat Y(x)U(\theta)=U(\theta)\hat Y(x)$, and likewise $\hat Y(x)\mathsf{K}=\mathsf{K}\hat Y(x)$. A real 2×2 matrix that commutes with J has the form $s\mathbf{1}+tJ$, and $\mathsf{K}J\mathsf{K}=-J$ forces $t=0$. So $\hat Y(x)=s(x)\mathbf{1}$ with $s(x)\in\mathbb{R}$. Suppose in addition that Y commutes with every T_a. Then $\mathsf{T}_a(\mathbf{1}\otimes v)=\mathbf{1}\otimes v$ gives $Y(\mathbf{1}\otimes v)=\mathsf{T}_aY(\mathbf{1}\otimes v)$, so $s(x-a)=s(x)$ for all $a\in\varepsilon\mathbb{Z}^3$. Since Λ is a single orbit of the translations, s is constant.

Step 4: induction on the weight. For $n\in\{2,3,4,5\}$ let (H_n) be the statement: there is exactly one choice of the $\delta m^2_{\gamma'}$, $\gamma'\in I$, $w(\gamma')<n$, for which (K) holds at all these γ', and these functions are constant. (H_2) holds vacuously. Assume (H_n) with $n\leq 4$, and let $\gamma\in I$ with $w(\gamma)=n$. By (B.5), $\hat\Sigma[\mathfrak{d}]_\gamma(x)=-\delta m^2_\gamma(x)\mathbf{1}+\hat\Gamma_\gamma(x)$, and $\Gamma_\gamma=\Sigma[\mathfrak{d}^{<n}]_\gamma$ is fixed by (H_n). By (H_n) the family $\mathfrak{d}^{<n}$ is constant on Λ. So by Step 2 every coefficient of $\Sigma[\mathfrak{d}^{<n}]$ commutes with $U(\theta)$, K and all T_a. By Step 3, $\hat\Gamma_\gamma(x)=s_\gamma\mathbf{1}$ with a real s_γ independent of x. Hence (K) holds at γ if and only if $\delta m^2_\gamma(x)=s_\gamma$ for all x. The conditions at the different γ of weight n do not involve one another, because Γ_γ depends only on weights $<n$. This proves (H_{n+1}), and (H_5) is the claim of existence, uniqueness and constancy. The values are real because all the Gaussian moments involved are real. They depend on (ε,j,M,μ_*) only through Λ, C_M and C^A, and they do not involve the photon mass μ_0 of any state. The vanishing for odd α is Lemma B.2. □

Corollary B.17. *Let $\mathfrak{d}$ be the family of Proposition B.16. Write the coefficient $\Sigma[\mathfrak{d}]_\gamma$ as $-\delta m^2_\gamma + \Sigma^\varepsilon_\gamma + \partial^{\varepsilon *}\Sigma^\varepsilon_{1,\gamma} + \Sigma^{\varepsilon *}_{1,\gamma}\partial^\varepsilon + \partial^{\varepsilon *}\Sigma^\varepsilon_{2,\gamma}\partial^\varepsilon$, grouping the graphs according to the derivatives at their external vertices as in* [3, (1.21), p. 416]. *Then $\hat\Sigma[\mathfrak{d}]_\gamma = -\delta m^2_\gamma \mathbf{1} + \hat\Sigma^\varepsilon_\gamma$. Hence $\mathfrak{d}$ is also the unique solution of the equations $-\delta m^2_\gamma + \sum_x \varepsilon^3 \Sigma^\varepsilon_\gamma(x) = 0$, $\gamma \in I$, of* [3, p. 417].

Proof. All $\delta m^2_{\gamma'}$ are constant. So a translation maps each of the three groups of graphs onto itself, and $\Sigma^\varepsilon_{1,\gamma}$, $\Sigma^\varepsilon_{2,\gamma}$ commute with all T_a, as in Step 2. The operator ∂^ε annihilates constants, so the last two terms vanish on $\mathbf{1}\otimes v$. The function $\Sigma^\varepsilon_{1,\gamma}(\mathbf{1}\otimes v)$ is translation invariant, hence constant in each direction, and $\partial^{\varepsilon *}$ annihilates such bond functions on the torus. Uniqueness follows by repeating Step 4 with this condition in place of (K). □

Remark B.18 (translation-invariant lines). Steps 2–4 use only one property of the lines: they commute with the lattice translations. On the periodic torus this holds for King's full propagators. So the counterterm of Definition 2.2 is King's counterterm [24, (3.32)–(3.33)] on $\Omega = T_{j,\varepsilon}$, with the photon mass taken to be μ_*. The notation $\delta m^2(x)$ of [24] allows for domains $\Omega \neq T_\varepsilon$, on which King's operators carry Neumann conditions and are not translation invariant. If the lines are only invariant under a block lattice, the argument gives only invariance under that lattice. Bałaban's splitting $\delta m^2 = \delta m^2_{\text{fin}}(x) + \delta m^2_K(x)$ [3, (1.26), p. 418] has x-dependent terms because G^ε_K and $G^\varepsilon_K(0)$ are not translation invariant; their sum is the constant counterterm of [3, (1.23), p. 417].

Lemma B.19 (property (P1); the per-torus coefficient may be used throughout)**.** *Let $K \geq K_0$ and $j \geq 1$.*

(a) *W_ε and z_ε are real numbers that depend on ε only (for fixed e, λ, m, μ_*). $b_{T_j}(\varepsilon) = a_{K,T_j}(m)$ is a real number that depends on (ε, j) only. None of the three depends on the site, on a direction or on μ_0, and the same three numbers are used at every $\mu_0 \in (0, \mu_*]$ and in the state $\flat$. This is property (P1) of Subsection 7.1.*

(b) *Every result of Sections 6 and 7–9 that involves a bare coefficient, other than Proposition 8.10(c), is of one of three kinds:*

- (i) *a statement at fixed (ε, T_j) about models on $T_{j,\varepsilon}$ whose quadratic coefficient is $a + k$, where a is a real constant (for instance $b_{T_j}(\varepsilon)$, a value $a_{K,T_j}(M)$, or a point between two such values) and k is a real function on the sites coming from a source;*
- (ii) *a bound whose constants are infima or suprema, over (ε, T_j) and over μ_0, of quantities of kind (i), combined with the numbers W_ε and z_ε;*
- (iii) *a compactness, holomorphy, positivity or closedness argument for a family of functionals that obey bounds of kind (ii).*

No such result compares the coefficients of two different tori or uses a model on $\varepsilon\mathbb{Z}^3$; the counterterm on $\varepsilon\mathbb{Z}^3$ enters only through the number z_ε. Consequently the coefficient $b_{T_j}(\varepsilon)$ of Definition 2.2 can be used throughout.

Proof. (a), W_ε. By its definition in Subsection 2.4, $W_\varepsilon = 2\int_{[-\pi/\varepsilon,\pi/\varepsilon)^3} (2\pi)^{-3}(\hat p^2 + m^2)^{-1} d^3p$. The integrand lies in $(0, m^{-2}]$ and the domain is bounded, so W_ε is a finite positive number. It is defined on $\varepsilon\mathbb{Z}^3$ and involves no site, no direction, no torus and no photon mass.

(a), z_ε. $z_\varepsilon = e^2 X_\varepsilon(m)$, where $X_\varepsilon(m)$ is the e^2-coefficient of $\langle|\varphi(0)|^2\rangle$ for the model on $\varepsilon\mathbb{Z}^3$ at scalar mass m and photon mass μ_* (Subsection 2.4). It is one number, defined without reference to a torus, to μ_0, or to any site other than the origin.

(a), b_T. By Proposition B.16 with $M = m$, $b_{T_j}(\varepsilon) = a_{K,T_j}(m)$ is one real number per (ε, j). It multiplies the scalar $|\varphi(x)|^2$ in (7.1) and carries no direction. Its definition uses the photon propagator $(-\Delta_\varepsilon + \mu_*^2)^{-1}$ at the reference mass μ_*. The photon mass μ_0 of the state does not enter it, so the same number is the coefficient at every $\mu_0 \in (0, \mu_*]$ and in $\flat$. Remark 2.3(ii) explains why the reference mass is needed.

(b). Kind (i):

- Proposition 4.7, by Proposition B.16, Corollary B.17 and Remark B.18.
- Lemmas 2.5, 7.3, 7.5, 7.6, 7.7, 7.8, 7.13, 9.7 and Proposition 7.9, which hold at fixed lattice for every real constant coefficient.
- The convexity of $a \mapsto \log Z_T^F(a)$ on $[a_{K,T}(M_1), b_T]$ in the proof of Proposition 6.4.
- The monotonicity in the photon covariance at b_T and the chessboard estimate in the proof of Proposition 6.5.
- The convexity in u in the proof of Proposition 6.6.
- Steps (1)–(4) of the proof of Proposition 6.7: the translation averaging, the coefficients $b_T + 2th$, and the convexity of f^B on $[a_{K,T}(m_1), a_{K,T}(m_2)]$.
- The fixed-lattice monotonicity in μ_0 at b_T behind Proposition 8.10(a),(b).

Kind (ii):

- Theorem 5.2 and Propositions 6.2, 6.3.
- Theorem 6.1, which bounds differences $a_{K,T}(M) - a_{K,T}(M')$ on one torus.
- The constants $g, \sigma_+, \bar{\Gamma}, \kappa_+$ of Proposition 6.4, the constants E_2 and $\bar{\Gamma}_{21}^{(1)}$ of Proposition 6.6, and the constants $J_{\mathrm{IR}}, \bar{Z}, g_{21}, S_{\mathfrak{p}}^B$ of Proposition 6.7.
- Theorems 6.8, 6.8 and Proposition 8.1.

Kind (iii):

- Propositions 8.2, 8.3, 8.6, 8.7 and Lemma 8.5, applied to sequences $(\varepsilon_i, T_{j_i}, \mu_i)$ in which each model has its own coefficient.
- Theorems 9.1–9.8 and Proposition 9.6.
- The passage to the limit in Proposition 8.10(a),(b).

None of these results uses a convergence of $b_{T_{j_i}}(\varepsilon_i)$. The number z_ε enters the observables and the c-number differences only as a constant. Replacing it by another sequence multiplies the generating functionals by nowhere-vanishing c-number factors, so no model on $\varepsilon\mathbb{Z}^3$ is involved. The same b_T, and one z_ε and one W_ε, are used in the state, in the observables and in both factors (Lemma B.24). □

Remark B.20. Part (a) is used in the following places:

- Proposition 6.5: the chessboard step needs a site-independent coefficient, and the monotonicity in μ_0 needs one that does not depend on μ_0.
- Step (1) of Proposition 6.7: the translation averaging needs the torus model to be invariant under $\varepsilon\mathbb{Z}^3$.
- Lemma 7.13.
- Lemma 7.6 and Proposition 8.7: limits and monotonicity in μ_0 at a fixed coefficient.

Remark B.21. Lemma B.19(a) is what Proposition 6.5 (the chessboard step needs a site-independent coefficient, and the μ_0-monotonicity needs a μ_0-free one), Lemma 7.13 (translations and O_h map the model to itself) and Lemma 7.6 (the limit $\mu_0 \downarrow 0$ at a fixed coefficient) use. Part (b) is used by Sections 6–9 as a whole.

Property (P2): integrability.

Property (P2): integrability. Throughout this paragraph $\lambda > 0$ and (ε, T) is fixed. T_ε is the lattice torus with $N \geq 2$ sites per side, so $|T_\varepsilon| = N^3$. Bonds are $b = (x, \mu)$ with $b_- = x$ and $b_+ = x + \varepsilon e_\mu$. For $a \in \mathbb{R}$ let $S_m^{(a)}$ be the matter action S_m of (7.1) with b_T replaced by a, so that $S_m = S_m^{(b_T)}$, and let $S_{\varepsilon,T,\mu_0}^{(a)}$ be the corresponding full action. Write $\varphi_x = r_x(\cos\vartheta_x, \sin\vartheta_x)$ and $u_b := \vartheta_{b_+} - \vartheta_{b_-} + \varepsilon e A_b$. For a real function k on the sites put

$$\kappa_k := \exp\Big(-\sum_x \varepsilon^3 k(x)(W_\varepsilon + z_\varepsilon)\Big), \qquad W_k^{(a)}(A) := \int \prod_x d^2\varphi_x \, \mathrm{e}^{-S_m^{(a)}(A,\varphi) + X_R(k)} \in (0, \infty],$$

and $V_a(r) := \varepsilon^3\big(\frac{1}{2}ar^2 + \lambda r^4\big) + 3\varepsilon r^2$. Thus $W_k = W_k^{(b_T)}$ is the weight of Lemma 2.5.

Lemma B.22 ((P2)). *Let $\lambda > 0$.*

(a) *For $c > 0$, $a \in \mathbb{R}$ and $p \geq 0$, $\int_0^\infty r^p \mathrm{e}^{-cr^4+ar^2}\, dr < \infty$. Moreover*

$$\int_{\mathbb{R}^2} \mathrm{e}^{-c|\varphi|^4+a|\varphi|^2}\, d^2\varphi = \pi \int_0^\infty \mathrm{e}^{-cu^2+au}\, du \leq \pi^{3/2} c^{-1/2} \mathrm{e}^{a^2/(4c)}.$$

(b) *For every $a \in \mathbb{R}$, every real k and all (A, φ),*

$$0 < \mathrm{e}^{-S_m^{(a)}(A,\varphi)+X_R(k)} \leq \kappa_k \prod_x \exp\big(-\varepsilon^3\lambda r_x^4 + \varepsilon^3(k(x) - \tfrac{a}{2})r_x^2\big). \tag{B.6}$$

Moreover $\mathrm{e}^{-S_m^{(a)}(A,\varphi)} = \prod_x \mathrm{e}^{-V_a(r_x)} \prod_b \mathrm{e}^{\varepsilon r_{b_-} r_{b_+} \cos u_b}$, and

$$\sum_x V_a(r_x) - \sum_b \varepsilon r_{b_-} r_{b_+} = \sum_x \varepsilon^3\big(\tfrac{1}{2}ar_x^2 + \lambda r_x^4\big) + \tfrac{1}{2}\varepsilon \sum_b (r_{b_+} - r_{b_-})^2.$$

In particular, for all couplings $c_b \in [0, \varepsilon]$,

$$\prod_x \mathrm{e}^{-V_a(r_x)} \prod_b \mathrm{e}^{c_b r_{b_-} r_{b_+}} \leq \prod_x \mathrm{e}^{-\varepsilon^3(\frac{1}{2}ar_x^2+\lambda r_x^4)}.$$

(c) *$W_k^{(a)}$ is a Borel function of A, and for every A*

$$0 < W_k^{(a)}(A) \leq W_k^{(a)}(0) \leq \kappa_k \prod_x \pi^{3/2}(\varepsilon^3\lambda)^{-1/2} \exp\Big(\frac{\varepsilon^3(k(x) - a/2)^2}{4\lambda}\Big) < \infty.$$

(d) *Let $\mu_0 > 0$, let f be a real 1-form and k a real function on the sites. Then*

$$0 < \int \prod_b dA_b \prod_x d^2\varphi_x\, \mathrm{e}^{-S^{(a)}_{\varepsilon,T,\mu_0}(A,\varphi)+\langle f, A\rangle_\varepsilon + X_R(k)} < \infty.$$

In particular $\mathrm{e}^{-S_{\varepsilon,T,\mu_0}}\, dA\, d\varphi$ is a finite nonzero measure, so (7.3) is the expectation of a probability measure. For real 2-forms $\mathbf{g}_i$, real functions h_l and $z_i, w_l \in \mathbb{C}$, the observable $\Theta = \exp[\sum_i z_i F(\mathbf{g}_i) + \sum_l w_l X_R(h_l)]$ satisfies $\langle|\Theta|\rangle_{\varepsilon,T,\mu_0} < \infty$. Every polynomial in the $F(\mathbf{g}_i)$ and $X_R(h_l)$ is integrable as well.

(e) *Let $\gamma^\flat := \mu_{(-\Delta_\varepsilon)^{-1}P^\perp} \otimes P^{-3}dc|_{[0,P)^3}$ be the measure of Definition 2.7. Then*

$$0 < \int d\gamma^\flat(A^\perp, c)\, W_0(A^\perp + c) \leq W_0(0) < \infty,$$

so the denominator of the $\flat$ state is positive and finite. For real $\mathbf{g}$ and k,

$$\int d\gamma^\flat(A^\perp, c)\, \mathrm{e}^{F(\mathbf{g})(A^\perp)} W_k(A^\perp + c) \leq W_k(0) \exp\big(\tfrac{1}{2}\langle d^*\mathbf{g},\, (-\Delta_\varepsilon)^{-1}P^\perp d^*\mathbf{g}\rangle_\varepsilon\big) < \infty.$$

Hence the observables Θ of (d) satisfy $\langle|\Theta|\rangle_{\varepsilon,T,\flat} < \infty$.

(f) *For every $a \in \mathbb{R}$ the measure $d\nu(r) := r\, \mathrm{e}^{-V_a(r)}\, dr$ on $[0, \infty)$ is nonzero, and $\int \mathrm{e}^{sr^2} d\nu(r) < \infty$ for every $s \in \mathbb{R}$.*

Property (P2) of Subsection 7.1 is part (a) with $c = \varepsilon^3\lambda$; the integrability holds with respect to $r^p\, dr$ for any $p \geq 0$, and with respect to $d^2\varphi$.

Proof. (a) By the AM–GM inequality, $ar^2 \leq \frac{c}{2}r^4 + \frac{a^2}{2c}$. Hence $r^p \mathrm{e}^{-cr^4+ar^2} \leq \mathrm{e}^{a^2/(2c)} r^p \mathrm{e}^{-cr^4/2}$, which is integrable on $[0, \infty)$. In polar coordinates $d^2\varphi = r\, dr\, d\vartheta$, and the substitution $u = r^2$, $r\, dr = \frac{1}{2}du$, gives the identity. Next, $-cu^2 + au = -c\big(u - \frac{a}{2c}\big)^2 + \frac{a^2}{4c}$. Extending the u-integral to $\mathbb{R}$ and using $\int_{\mathbb{R}} \mathrm{e}^{-cv^2} dv = (\pi/c)^{1/2}$ gives the bound.

(b) $U(s)$ is a rotation, so $|U(\varepsilon e A_b)\varphi_{b_+}| = r_{b_+}$ and $\big(U(\varepsilon e A_b)\varphi_{b_+}\big) \cdot \varphi_{b_-} = r_{b_+} r_{b_-} \cos u_b$. Therefore

$$\tfrac{1}{2}\varepsilon^3 |(D_A\varphi)(b)|^2 = \tfrac{1}{2}\varepsilon\big(r_{b_+}^2 + r_{b_-}^2\big) - \varepsilon r_{b_-} r_{b_+} \cos u_b \;\geq 0.$$

Each site x is the initial point of exactly three bonds, (x,μ), and the final point of exactly three bonds, $(x-\varepsilon e_\mu,\mu)$. Hence $\sum_b \frac12\varepsilon(r_{b_+}^2+r_{b_-}^2)=3\varepsilon\sum_x r_x^2$, which gives the product formula for $\mathrm{e}^{-S_m^{(a)}}$. Setting $\cos u_b=1$ in the same computation gives $3\varepsilon\sum_x r_x^2-\sum_b \varepsilon r_{b_-}r_{b_+}=\frac12\varepsilon\sum_b(r_{b_+}-r_{b_-})^2\ge 0$. For $0\le c_b\le\varepsilon$ and $r\ge 0$ we have $c_b r_{b_-}r_{b_+}\le \varepsilon r_{b_-}r_{b_+}$, which gives the last inequality of (b). For (B.6), drop the nonnegative gradient term from $S_m^{(a)}$ and write $X_R(k)=\sum_x \varepsilon^3 k(x) r_x^2+\log\kappa_k$. Positivity is clear.

(c) The integrand $\mathrm{e}^{-S_m^{(a)}+X_R(k)}$ is continuous and positive on $\mathbb{R}^{\text{bonds}}\times(\mathbb{R}^2)^{T_\varepsilon}$. By Tonelli's theorem, $W_k^{(a)}$ is Borel and $W_k^{(a)}(A)>0$. Integrate (B.6) over φ and use (a) at each site, with $c=\varepsilon^3\lambda$ and with $\varepsilon^3(k(x)-a/2)$ in place of a. This bounds $W_k^{(a)}(A)$ by the stated product for every A, in particular for $A=0$.

It remains to show $W_k^{(a)}(A)\le W_k^{(a)}(0)$. For $t\ge 0$ and $u\in\mathbb{R}$,

$$\mathrm{e}^{t\cos u}=\sum_{q\ge 0}\frac{(t/2)^q}{q!}\big(\mathrm{e}^{iu}+\mathrm{e}^{-iu}\big)^q=\sum_{\nu\in\mathbb{Z}} I_\nu(t)\mathrm{e}^{i\nu u},\qquad I_\nu(t):=\sum_{0\le j\le q,\ 2j-q=\nu}\frac{(t/2)^q}{j!\,(q-j)!}\ \ge 0.$$

All terms are nonnegative, so the rearrangement is legitimate, and $\sum_\nu I_\nu(t)=\mathrm{e}^t$. Fix $r\in[0,\infty)^{T_\varepsilon}$ and put $t_b:=\varepsilon r_{b_-}r_{b_+}\ge 0$. Then $\prod_b \mathrm{e}^{t_b\cos u_b}=\sum_{n\in\mathbb{Z}^{\text{bonds}}}\prod_b I_{n_b}(t_b)\,\mathrm{e}^{i\sum_b n_b u_b}$. The series converges absolutely, uniformly in ϑ, with $\sum_n\prod_b I_{n_b}(t_b)=\prod_b \mathrm{e}^{t_b}$. Integrate term by term over $\vartheta\in[0,2\pi)^{T_\varepsilon}$. Since $\int\prod_x d\vartheta_x\,\mathrm{e}^{i\sum_b n_b(\vartheta_{b_+}-\vartheta_{b_-})}=(2\pi)^{|T_\varepsilon|}\mathbf{1}\{\partial n=0\}$, with $(\partial n)(x):=\sum_{b_+=x}n_b-\sum_{b_-=x}n_b$, we get

$$J_r(A):=\int\prod_x d\vartheta_x\prod_b \mathrm{e}^{t_b\cos u_b}=(2\pi)^{|T_\varepsilon|}\sum_{n:\,\partial n=0}\prod_b I_{n_b}(t_b)\,\mathrm{e}^{ie\varepsilon\sum_b n_b A_b},$$

and therefore $0<J_r(A)=|J_r(A)|\le J_r(0)$. By (b) and Tonelli's theorem,

$$W_k^{(a)}(A)=\kappa_k\int\prod_x r_x\,dr_x\,\mathrm{e}^{-V_a(r_x)+\varepsilon^3 k(x)r_x^2}\,J_r(A).$$

Integrating $J_r(A)\le J_r(0)$ against this positive measure gives $W_k^{(a)}(A)\le W_k^{(a)}(0)$.

(d) The integrand is continuous and positive, so the integral is positive. By (b), the integral is at most

$$\int\prod_b dA_b\,\mathrm{e}^{-\frac12\langle A,(-\Delta_\varepsilon+\mu_0^2)A\rangle_\varepsilon+\langle f,A\rangle_\varepsilon}\ \times\ \kappa_k\int\prod_x d^2\varphi_x\,\mathrm{e}^{-\varepsilon^3\lambda r_x^4+\varepsilon^3(k(x)-\frac a2)r_x^2}.$$

The second factor is finite by (a). In the coordinates (A_b), $\langle A,(-\Delta_\varepsilon+\mu_0^2)A\rangle_\varepsilon=A\cdot QA$ with $Q=\varepsilon^3(-\Delta_\varepsilon+\mu_0^2)$, and $\langle f,A\rangle_\varepsilon=\varepsilon^3 f\cdot A$. Here $Q\ge\varepsilon^3\mu_0^2>0$, because $-\Delta_\varepsilon\ge 0$ (Lemma B.4(ii)). So the first factor equals $(2\pi)^{n/2}(\det Q)^{-1/2}\exp\big(\frac12\varepsilon^6 f\cdot Q^{-1}f\big)<\infty$, with $n=3N^3$. Tonelli's theorem gives the claim. With $f=0$, $k=0$ and $a=b_T$ this is the normalisability of the state. For Θ: F and X_R are real and linear in $\mathbf{g}$ and h. Hence $|\Theta|=\exp\big[F(\sum_i \mathrm{Re}\,z_i\,\mathbf{g}_i)+X_R(\sum_l \mathrm{Re}\,w_l\,h_l)\big]$, which is of the form just treated with $f=d^*\sum_i \mathrm{Re}\,z_i\,\mathbf{g}_i$, since $F(\mathbf{g})=\langle d^*\mathbf{g},A\rangle_\varepsilon$. Polynomials are covered by $|x|^q\le q!\,(\mathrm{e}^x+\mathrm{e}^{-x})$.

(e) W_0 is Borel, and $0<W_0\le W_0(0)<\infty$ by (c). Moreover $\gamma^\flat$ is a probability measure: the Gaussian factor exists because $-\Delta_\varepsilon\ge 16/\ell_T^2$ on the range of $P^\perp$ (Lemma B.4(iii)). Hence the integral of W_0 lies in $(0,W_0(0)]$; it is positive because the sets $\{W_0>1/n\}$ increase to the whole space. For the second claim: a constant 1-form c satisfies $dc=0$, so $F(\mathbf{g})(A^\perp+c)=\langle d^*\mathbf{g},A^\perp\rangle_\varepsilon$. Under the centred Gaussian with covariance $(-\Delta_\varepsilon)^{-1}P^\perp$, this is a centred normal variable with variance $\langle d^*\mathbf{g},(-\Delta_\varepsilon)^{-1}P^\perp d^*\mathbf{g}\rangle_\varepsilon$. Bound $W_k(A^\perp+c)\le W_k(0)$ by (c) and compute the Gaussian exponential moment; this gives the bound. The claim for Θ follows as in (d).

(f) $\int_0^\infty \mathrm{e}^{sr^2}r\,\mathrm{e}^{-V_a(r)}\,dr=\int_0^\infty r\,\mathrm{e}^{-\varepsilon^3\lambda r^4+(s-\varepsilon^3 a/2-3\varepsilon)r^2}\,dr$ is finite by (a) with $p=1$. It is positive because the integrand is positive on $(0,\infty)$. □

Remark B.23. Lemma B.22 supplies every use of (P2) in Sections 7–9 and in this appendix, and none of these uses needs more:

- the normalisability of (7.3) is part (d);
- the domination that justifies Fubini's theorem in the proofs of Lemmas 2.5 and 2.5 is part (b), last display (with $c_b = \varepsilon$), together with part (a);
- the integrability of $|\Theta|^2$ in Proposition 7.9(c) is parts (d) and (e);
- the finiteness of the logarithmic moment generating functions α_n in Section 10 is parts (d) and (e) with $\mathbf{g} = 0$ and constant k;
- the hypotheses on ν_x in Definition B.8 and Proposition B.13 are part (f) and part (b), last display.

The coefficient a is arbitrary, so the lemma also applies at the anchor coefficients $a_{K,T}(M)$ and in Lemma B.7.

Property (P3): one scheme.

Lemma B.24 (property (P3))**.** *On each T_j and at each ε (with $K \geq K_0$), the following coincide:*

(a) *the bare coefficient of the state* (7.3)*, at every $\mu_0 \in (0, \mu_*]$ and in $\flat$;*
(b) *the coefficient at which the bounds for positive sources (Propositions 6.4, 6.5) are proved;*
(c) *the coefficient a_0 of the factor for negative sources (Proposition 6.7, the left side of the anchor condition* (6.3)*: $b_T + 2H \leq a_{K,T}(m_1)$).*

All three equal $b_T = a_{K,T}(m)$. Every anchor coefficient in Section 6 ($a_{K,T}(M_1)$ and $a_{K,T}(m)$ in Proposition 6.4; $a_{K,T}(m_1)$ and $a_{K,T}(m_2)$ in Theorem 5.2, Propositions 6.6, 6.7, 6.3 and the bound (6.2)*) is a value of the single function $M \mapsto a_{K,T}(M)$ of Definition 2.2 on the same torus. One z_ε and one W_ε, those of Subsection 2.4, appear in the observables* (7.2)*, in α_T and $\bar{\Gamma}$ of Proposition 6.4, in ρ^B, f^B, Δ_1^B of Subsection 6.5, and in $\bar{\Gamma}^{(1)}_{21}$. Theorem 5.2 contains no z_ε. This is property (P3) of Subsection 7.1.*

Proof. (1) *One state coefficient.* The state is defined at b_T (Definition 2.2, and Lemma B.19(a) for μ_0 and $\flat$). Proposition 6.4 is stated with upper anchor $M_2 = m$, i.e. at the coefficient $a_{K,T}(m) = b_T$ on the same T. Proposition 6.5 is stated for all states at the coefficient b_{T_j}. Proposition 6.7 is stated at $a_0 = b_T$ through (6.3). (2) *One anchor function.* Each anchor listed in the statement is, by its definition in Section 6, $a_{K,T}(M)$ at some $M \in \{M_1, m, m_1, m_2\}$ with the photon μ_* on the same torus. In the slice models of Subsection 5.1 the counterterm is fixed at the photon covariance C_{μ_*} for every u, so the coefficients of Theorem 5.2 at $u = 1$ and $u = -\eta$ are $a_{K,T}(m_2)$ and $a_{K,T}(m_1)$, not a u-dependent profile. (3) *One z_ε, one W_ε.* z_ε and W_ε are the numbers of Subsection 2.4, the same on every torus (Lemma B.19(a)). They enter each statement of the chain only through the c-number $-\sum_x \varepsilon^3 h(x)(W_\varepsilon + z_\varepsilon)$ in $X_R(h)$, or through $\frac{1}{2}a(W_\varepsilon + z_\varepsilon)$ in $f_T(a)$ and $\gamma_T(M)$. Each proof on the path is an identity in z: the z-parts of the two sides of every comparison are equal. The only inputs that depend on z are the bounds on γ-differences uniform in ε (Proposition 6.3, the bound (6.2), $\bar{\Gamma}$), and these are proved for the graphical $z_\varepsilon = e^2 X_\varepsilon(m)$. Theorem 5.2 is a vacuum bound at anchor coefficients and contains no z_ε. (4) *Vacuum constants.* In each statement that uses a vacuum constant, the E_0 is the photon-normalised $E_{0,s}$ of that statement. This is a condition within each statement. It does not affect Sections 7–9, because normalised expectations contain no vacuum constants (Remark 2.3(iii)), and $E_{0,v}(\mu_*)$ cancels in every γ-difference. Items (1)–(3) are the statement, and (4) shows that the vacuum constants do not interfere. □

Remark B.25. Mixed schemes are excluded: if the "+" input were taken at one coefficient and the "−" input at another, the Cauchy–Schwarz step (3) of the proof of Theorem 6.8 would combine bounds for two different states. The difference would be the volume-extensive shift $r_T(m) \sum_x \varepsilon^3 r_x^2$, not a c-number.

The zero-mode split.

Lemma B.26 (spectrum of $-\Delta_\varepsilon$ and the zero mode)**.** *On the real 1-forms of $T_{j,\varepsilon}$, with the inner product $\langle\cdot,\,\cdot\rangle_\varepsilon$:*

(a) *$-\Delta_\varepsilon$ is self-adjoint and ≥ 0. Its kernel is the span of $\mathbf{1}_1, \mathbf{1}_2, \mathbf{1}_3$, and $-\Delta_\varepsilon \geq 16/\ell_T^2$ on the range of $P^\perp$.*

(b) *P_0 commutes with $-\Delta_\varepsilon$. For $\mu_0 > 0$, $C_{\mu_0} = (-\Delta_\varepsilon + \mu_0^2)^{-1} = \mu_0^{-2} P_0 + (-\Delta_\varepsilon + \mu_0^2)^{-1} P^\perp$.*

(c) *Under $\mu_{C_{\mu_0}}$ write $A = A^\perp + c$ with $A^\perp := P^\perp A$ and $c := P_0 A = \sum_\mu c_\mu \mathbf{1}_\mu$. Then $A^\perp$ and c are independent. $A^\perp$ is centred Gaussian with covariance $(-\Delta_\varepsilon + \mu_0^2)^{-1} P^\perp$. The c_μ are independent $N(0, s(\mu_0))$ with $s(\mu_0) = 1/(\mu_0^2 |T|)$. Equivalently, the mass term restricted to the constants is $\frac{1}{2}\mu_0^2 \|c\|_{2,\varepsilon}^2 = \frac{1}{2}\mu_0^2 |T| \sum_\mu c_\mu^2$.*

Proof. This is Lemmas B.4 and B.6 in the notation of Section 7, where $P_0 = \Pi_0$ and $P^\perp = \Pi^\perp$. (a) is Lemma B.4(ii),(iii). (b) P_0 is the spectral projection of $-\Delta_\varepsilon$ onto its kernel, so it commutes with $-\Delta_\varepsilon$, and $(-\Delta_\varepsilon + \mu_0^2)^{-1}$ equals μ_0^{-2} on the range of P_0. (c) is Lemma B.6(a),(b); the identity $\langle c,\, c\rangle_\varepsilon = |T| \sum_\mu c_\mu^2$ is Lemma B.6(a). □

The constants of Theorem 6.8 are uniform over the tori. Let Z_T be the zero-mode price of Proposition 6.7. It comes from comparing the Haar zero mode with the massive zero mode of variance $s_K := 1/(\mu_*^2 |T|)$ on the period cell $P = 2\pi/(|e|\ell_T)$:

$$Z_T := \frac{1}{\ell_T^3}\Big[\tfrac{3}{2}\ln_+\Big(\frac{P^2}{s_K}\Big) + 3\delta_P\Big] = \frac{1}{\ell_T^3}\Big[\tfrac{3}{2}\ln_+\Big(\frac{4\pi^2\mu_*^2\ell_T}{e^2}\Big) + 3\delta_P\Big],$$

$$\delta_P := 2\sum_{n\geq 1} \mathrm{e}^{-2\pi^2 n^2} \ (\approx 5.35\cdot 10^{-9}),$$

where $\ln_+ := \max(\ln, 0)$. Unlike Δ_1^B (a supremum over ε and $\ell_T \geq \ell_0$, Subsection 6.5) and J_{IR} (likewise a supremum), Z_T depends on the torus. The constant $S_\flat^B$ of Proposition 6.7 is therefore formed with $\bar{Z} := \sup_j Z_{T_j}$, and the next lemma shows that $\bar{Z} < \infty$.

Lemma B.27 ($\sup Z_T < \infty$)**.** *Let $e \neq 0$ and $\ell_0 > 0$, and put $q := 4\pi^2\mu_*^2/e^2$. Then*

$$\sup_{\ell_T \geq \ell_0} Z_T \leq \frac{1}{\ell_0^3}\Big[\tfrac{3}{2}\ln_+(q\ell_0) + \frac{1}{2\mathrm{e}} + 3\delta_P\Big] < \infty.$$

Proof. For $x, y > 0$, $\ln_+(xy) \leq \ln_+ x + \ln_+ y$: if $xy \leq 1$ the left side is 0, and otherwise $\ln(xy) = \ln x + \ln y$. With $t := \ell_T/\ell_0 \geq 1$ this gives $\ln_+(q\ell_T) \leq \ln_+(q\ell_0) + \ln t$. Hence

$$Z_T \leq \frac{\frac{3}{2}\ln_+(q\ell_0) + 3\delta_P}{\ell_0^3 t^3} + \frac{3}{2\ell_0^3}\,\frac{\ln t}{t^3} \leq \frac{1}{\ell_0^3}\Big[\tfrac{3}{2}\ln_+(q\ell_0) + 3\delta_P\Big] + \frac{3}{2\ell_0^3}\cdot\frac{1}{3\mathrm{e}},$$

because $t^{-3} \leq 1$ and $\max_{t\geq 1} t^{-3}\ln t = 1/(3\mathrm{e})$, attained at $t = \mathrm{e}^{1/3}$. □

Corollary B.28 (the constants of Theorem 6.8)**.** *Put $S_\flat^B(T) := [\Delta_1^B + J_{\mathrm{IR}} + Z_T]_+/g_{21}$ for a torus T, and*

$$S_\flat^B := \big[\Delta_1^B + J_{\mathrm{IR}} + \sup_{\ell_T \geq \ell_0} Z_T\big]_+/g_{21}.$$

Then $S_\flat^B < \infty$, and $S_\flat^B(T) \leq S_\flat^B$ for every torus with $\ell_T \geq \ell_0$, in particular for every T_j. Hence Proposition 6.7 holds with the constant $S_\flat^B$, uniformly in ε, j and $\mu_0 \in (0, \mu_*]\cup\{\flat\}$, and Theorem 6.8 holds with $\mathsf{K} := \max(\kappa_+, 2S_\flat^B)$, which is independent of ε, j and μ_0.

Proof. The inputs are: $\Delta_1^B < \infty$ (Proposition 6.6); $J_{\mathrm{IR}} < \infty$, a supremum over ε and $\ell_T \geq \ell_0$ with an explicit bound free of ε (Lemma I.8(iv)); $g_{21} > 0$ (Theorem 6.1(b)); and Lemma B.27. Together they give $S_\flat^B < \infty$. None of the four quantities depends on ε, j or μ_0. For each T with $\ell_T \geq \ell_0$ we have $Z_T \leq \sup_{\ell' \geq \ell_0} Z_{T'}$. Since $x \mapsto [x]_+$ is nondecreasing and $g_{21} > 0$, it follows that

$S^B_\flat(T) \le S^B_\flat$. The right side of the bound of Proposition 6.7 on T is therefore at most $2\|h\|_{L^1,\varepsilon} S^B_\flat$. In steps (5)–(6) of the proof of Theorem 6.8 the bound $\kappa_+\|k_+\| + 2S^B_\flat\|k_-\| \le \max(\kappa_+, 2S^B_\flat)\|k\|$ follows from $\|k_+\|_{L^1,\varepsilon} + \|k_-\|_{L^1,\varepsilon} = \|k\|_{L^1,\varepsilon}$. The constant Δ^B_1 is the one of the slice vacuum bound (Proposition 6.6), formed with the slice covariance Γ^B. □

Remark B.29. (i) $S^B_\flat$ is not uniform as $e \to 0$: $\sup_{\ell_T \ge \ell_0} Z_T$ grows like $\frac{3}{2}\ln(1/e^2)/\ell_0^3$, because the period cell $2\pi/(|e|\ell_T)$ becomes infinite. In $\mathfrak{W}$ the charge satisfies $e^2 > 0$ and is fixed, so this does not matter for Theorem 3.1. (ii) Since $Z_T \to 0$ as $\ell_T \to \infty$, the zero-mode price disappears in infinite volume, but uniformity over the sequence (T_j) needs only the supremum.

B.3. **The continuum limits.**

B.3.1. *Multilinearity, fixed-torus translations and the class $\mathcal{D}$.* We prove three steps of Sections 8–9. Throughout, $C^{(R)}$ is the real vector space of continuous scalar functions or continuous 2-forms on $\mathbb{R}^3$ with support in the open ball B_R. For a 2-form $\mathbf{g} = \sum_{\mu<\kappa} g_{\mu\kappa}\, dx_\mu \wedge dx_\kappa$ we put $\|\mathbf{g}\|_\infty := \max_{\mu<\kappa}\|g_{\mu\kappa}\|_\infty$. Translations act by $(\tau_a f)(x) := f(x-a)$, componentwise on 2-forms. A tuple $f = (\mathbf{g}_1, \dots, \mathbf{g}_n; h_1, \dots, h_m)$ is *R-admissible for T* if all its entries lie in $C^{(R)}$ and $2R + 2 < \ell_T$. By Definition 2.9 this is exactly when the entries can be sampled on T_ε.
Multilinearity and symmetry of the limits.

Lemma B.30 (multilinearity)**.** (a) *Fix (ε, T) and $\mu_0 \in (0, \mu_*] \cup \{\flat\}$, and let R satisfy $2R+2 < \ell_T$. On $(C^{(R)})^{n+m}$ (2-forms in the first n arguments, scalar functions in the last m), $S^{\varepsilon,T,\mu_0}_{n,m}$ is finite and real. It is linear in each argument separately, and it is invariant under permutations of $(\mathbf{g}_1, \dots, \mathbf{g}_n)$ and, separately, of $(h_1, \dots, h_m)$.*

(b) *Let $S_{n,m}$ be the limit of Proposition 8.2(a). Then $S_{n,m}$ has the properties of (a) on $(C^{(R)})^{n+m}$ for every R; if $T_{j_i} \equiv T$, for every R with $2R + 2 < \ell_T$. The same holds for S^0 of Proposition 8.7(b).*

In particular the limits of Proposition 8.2(a) are multilinear and symmetric within each field type.

Proof. (a) *Linearity of sampling.* Sampling evaluates a function at the sites of the cell $[-\ell_T/2, \ell_T/2)^3$, and a 2-form component at plaquette centres (Definition 2.9). Evaluation at a point is linear, so the sampled function of $\alpha f + \beta f'$ is α times the sampled f plus β times the sampled f'. Since $C^{(R)}$ is a vector space, all three functions are admissible. *Linearity of the fields.* By (7.2), $F(\mathbf{g}) = \langle \mathbf{g}, dA\rangle_\varepsilon$ is linear in the sampled $\mathbf{g}$. Also $X_R(h) = \sum_x \varepsilon^3 h(x)(r_x^2 - W_\varepsilon - z_\varepsilon)$ is linear in h, because $W_\varepsilon, z_\varepsilon$ do not depend on h (P1). So for fixed field configuration (A, φ) the product $\prod_i F(\mathbf{g}_i) \prod_k X_R(h_k)$ is a real number, linear in each argument. It is invariant under permutations of the $\mathbf{g}_i$ and of the h_k, since multiplication of real numbers is commutative. *Integrability.* The product is a polynomial in the $F(\mathbf{g}_i)$ and $X_R(h_k)$, hence an elementary observable. It is therefore integrable in every state, the $\flat$ state included (Lemma 7.6). Expectation is linear on integrable functions and preserves reality, which gives (a). This is the same argument as in the proof of Lemma 9.7.
(b) Each of the properties in (a) is an identity between the values of $S^{(i)}_{n,m}$ at finitely many tuples, all of them admissible. Examples are $S^{(i)}(\dots, \alpha f + \beta f', \dots) = \alpha S^{(i)}(\dots, f, \dots) + \beta S^{(i)}(\dots, f', \dots)$, $S^{(i)}(f_\pi) = S^{(i)}(f)$ for a permutation π within one field type, and $\operatorname{Im} S^{(i)}(f) = 0$. Proposition 8.2(a) gives convergence at each of these tuples, so each identity passes to the limit $i \to \infty$. For S^0, use the convergence of the Schwinger functions in Proposition 8.7(b) in the same way. The bound with continuum norms in Proposition 8.2(a) is step (iii) of its proof. Riemann sums give $\|\mathbf{g}\|_{2,\varepsilon} \to \|\mathbf{g}\|_2$ and $\|h\|_{L^1,\varepsilon} \to \|h\|_1$, while the sampled supremum is at most $\|h\|_\infty$; hence $\limsup_i N_{\varepsilon_i}(h) \le N(h)$. □

Translations of the fixed-torus continuum states.

Translations of the fixed-torus continuum states. Throughout this paragraph $T = T_j$ is a fixed cubic torus of side $\ell := \ell_T$, identified with the cell $Q := [-\ell/2, \ell/2)^3$ as in Definition 2.9. For $y \in \mathbb{R}^3$ let $[y] \in Q$ be its representative modulo $\ell\mathbb{Z}^3$. For $\rho \geq 0$, $C_c^{(\rho)}$ is the space of real continuous scalar functions, or of real continuous 2-forms, with support in the closed Euclidean ball $\bar{B}_\rho$. For a 2-form we put $\|\mathbf{g}\|_\infty := \max_{\mu<\kappa} \|g_{\mu\kappa}\|_\infty$. A tuple $f = (\mathbf{g}_1, \dots, \mathbf{g}_n; h_1, \dots, h_m)$ is *admissible* if all its entries lie in $C_c^{(\rho)}$ for some ρ with $2\rho + 2 < \ell$. Such entries can be sampled, since $\bar{B}_\rho \subset B_{\rho'}$ for some $\rho' > \rho$ with $2\rho' + 2 < \ell$. For $a \in \mathbb{R}^3$ put $(\tau_a h)(x) := h(x - a)$ and $(\tau_a \mathbf{g})_x := \mathbf{g}_{x-a}$ (componentwise translation), and let $\tau_a f$ be the entrywise translate. Every lattice spacing satisfies $\varepsilon = 2^{-K}\varepsilon_0 \leq 1$.

For $a \in \varepsilon\mathbb{Z}^3$, translation by a maps T_ε onto itself, because $\ell/\varepsilon \in \mathbb{N}$. It acts on oriented cells by $\tau_a(x; v_1 \wedge \cdots \wedge v_k) := (x + a; v_1 \wedge \cdots \wedge v_k)$, on lattice forms by $(\tau_a \omega)(c) := \omega(\tau_{-a} c)$, and on field configurations by $\hat{\tau}_a(A, \varphi) := (\tau_a A, \tau_a \varphi)$.

Lemma B.31 (lattice translations)**.** *Let $a \in \varepsilon\mathbb{Z}^3$ and $\mu_0 \in (0, \mu_*] \cup \{\flat\}$.*

(i) *$d\tau_a = \tau_a d$, $\langle \tau_a \omega, \tau_a \omega' \rangle_\varepsilon = \langle \omega, \omega' \rangle_\varepsilon$ and $d^* \tau_a = \tau_a d^*$.*

(ii) *For real lattice sources, $F(\mathbf{g}) \circ \hat{\tau}_a = F(\tau_{-a}\mathbf{g})$ and $X_R(h) \circ \hat{\tau}_a = X_R(\tau_{-a} h)$.*

(iii) *For real lattice 2-forms $\mathbf{g}_i$ and real lattice functions h_k,*

$$\Big\langle \prod_i F(\tau_a \mathbf{g}_i) \prod_k X_R(\tau_a h_k) \Big\rangle_{\varepsilon,T,\mu_0} = \Big\langle \prod_i F(\mathbf{g}_i) \prod_k X_R(h_k) \Big\rangle_{\varepsilon,T,\mu_0}.$$

Proof. (i) τ_a maps oriented k-cells bijectively onto oriented k-cells. It commutes with reversal of orientation and with the oriented boundary, $\partial(\tau_a c) = \tau_a \partial c$, since it moves all vertices of a cube by the same vector and keeps the orientation. Hence $(d\tau_a \omega)(c) = \varepsilon^{-1}\omega(\tau_{-a}\partial c) = \varepsilon^{-1}\omega(\partial \tau_{-a} c) = (\tau_a d\omega)(c)$. Also τ_a permutes the unoriented cells, which all carry the weight ε^3, so the pairing is invariant. Since $\tau_a^{-1} = \tau_{-a}$, taking adjoints gives $d^*\tau_a = \tau_a d^*$.

(ii) By (i), $F(\mathbf{g})(\tau_a A) = \langle \mathbf{g}, d\tau_a A \rangle_\varepsilon = \langle \mathbf{g}, \tau_a dA \rangle_\varepsilon = \langle \tau_{-a}\mathbf{g}, dA \rangle_\varepsilon$. Next, $r_x \circ \hat{\tau}_a = r_{x-a}$, and $W_\varepsilon, z_\varepsilon$ do not depend on the site (Lemma B.19(a)). Changing the summation variable to $y = x - a$ gives $X_R(h) \circ \hat{\tau}_a = \sum_y \varepsilon^3 h(y + a)(r_y^2 - W_\varepsilon - z_\varepsilon) = X_R(\tau_{-a} h)$.

(iii) Put $G := \prod_i F(\tau_a \mathbf{g}_i) \prod_k X_R(\tau_a h_k)$. By (ii), $G \circ \hat{\tau}_a = \prod_i F(\mathbf{g}_i) \prod_k X_R(h_k)$. Both G and $G \circ \hat{\tau}_a$ are polynomials in the $F(\cdot)$ and $X_R(\cdot)$. They are therefore elementary observables, integrable in every state (Lemma 7.6). Let $\mu_0 > 0$. The map $\hat{\tau}_a$ permutes the bond variables (without change of sign, since translations preserve the orientation of bonds) and the site variables. So it preserves Lebesgue measure, and by Lemma 7.13(i) it preserves the action. Hence the probability measure of $\langle \cdot \rangle_{\varepsilon,T,\mu_0}$ is $\hat{\tau}_a$-invariant, and $\langle G \circ \hat{\tau}_a \rangle_{\varepsilon,T,\mu_0} = \langle G \rangle_{\varepsilon,T,\mu_0}$. For $\flat$, Lemma 7.6 applies to both G and $G \circ \hat{\tau}_a$. It gives $\langle G \circ \hat{\tau}_a \rangle_{\varepsilon,T,\flat} = \lim_{\mu_0 \downarrow 0} \langle G \circ \hat{\tau}_a \rangle_{\varepsilon,T,\mu_0} = \lim_{\mu_0 \downarrow 0} \langle G \rangle_{\varepsilon,T,\mu_0} = \langle G \rangle_{\varepsilon,T,\flat}$. □

Lemma B.32 (sampling commutes with translations)**.** *Let $\rho \geq 0$ and $a \in \varepsilon\mathbb{Z}^3$ with $2(\rho + |a|) + 2 < \ell$. Let h be a scalar and $\mathbf{g}$ a 2-form in $C_c^{(\rho)}$. Then $\tau_a h, \tau_a \mathbf{g} \in C_c^{(\rho+|a|)}$ can be sampled, and*

$$(\tau_a h)_\varepsilon = \tau_a(h_\varepsilon), \qquad (\tau_a \mathbf{g})_\varepsilon = \tau_a(\mathbf{g}_\varepsilon).$$

Consequently $S_{n,m}^{\varepsilon,T,\mu_0}(\tau_a f) = S_{n,m}^{\varepsilon,T,\mu_0}(f)$ for every tuple f with entries in $C_c^{(\rho)}$ and every $\mu_0 \in (0, \mu_*] \cup \{\flat\}$.

Proof. Scalars. Let $x \in Q$ be a site. Then $(\tau_a h)_\varepsilon(x) = h(x - a)$ and $\tau_a(h_\varepsilon)(x) = h([x - a])$. If $x - a \in Q$, the two values agree. Otherwise some coordinate of $x - a$ lies outside $[-\ell/2, \ell/2)$, so $|x - a| \geq \ell/2 > \rho$ and $h(x - a) = 0$. Suppose $h([x - a]) \neq 0$. Then $|[x - a]| \leq \rho$, so $|[x - a] + a| \leq \rho + |a| < \ell/2 - 1$, and $[x - a] + a$ lies in the interior of Q. But $[x - a] + a \equiv x$ modulo $\ell\mathbb{Z}^3$ and both points lie in Q, so $[x - a] + a = x$, that is, $x - a \in Q$, a contradiction. So both values vanish.

2-forms. Let $c = (x; v_1 \wedge v_2)$ be a plaquette with $x \in Q$, and put $z := x + \frac{\varepsilon}{2}(v_1 + v_2)$; then $|z - x| = \varepsilon/\sqrt{2} < 1$. By Definition 7.11, $(\tau_a \mathbf{g})_\varepsilon(c) = \mathbf{g}_{z-a}(v_1, v_2)$. Since $\tau_{-a} c = ([x - a]; v_1 \wedge v_2)$, also $\tau_a(\mathbf{g}_\varepsilon)(c) = \mathbf{g}_{y + \frac{\varepsilon}{2}(v_1+v_2)}(v_1, v_2)$ with $y = [x - a]$. If $x - a \in Q$, then $y = x - a$ and the two

values agree. Otherwise $|x-a| \geq \ell/2$, so $|z-a| \geq \ell/2 - 1 > \rho$ and the first value vanishes. If the second value were nonzero, we would have $|y| < \rho + 1$, so $|y+a| < \rho + |a| + 1 < \ell/2$, and $y + a = x$ as in the scalar case, again a contradiction.

The last assertion follows from Lemma B.31(iii), applied to the sampled sources. □

Lemma B.33 (fixed-torus translations, local form)**.** *Let $(\varepsilon_i, T, \mu_i)$ be a sequence with $K_i \to \infty$ and $\mu_i \in (0, \mu_*] \cup \{\flat\}$, and let S be the limit along a subsequence given by Proposition 8.2 with $T_{j_i} \equiv T$. Let $R \geq 0$ and $a \in \mathbb{R}^3$ satisfy $2(R + |a|) + 2 < \ell$. Then for every tuple f with entries in $C_c^{(R)}$,*

$$S_{n,m}(\tau_a \mathbf{g}_1, \dots, \tau_a \mathbf{g}_n; \tau_a h_1, \dots, \tau_a h_m) = S_{n,m}(\mathbf{g}_1, \dots, \mathbf{g}_n; h_1, \dots, h_m).$$

Proof. Let $c_{n,m}$ be the constant of Proposition 8.1, and let $\mathcal{D} := \bigcup_K \varepsilon_K \mathbb{Z}^3 = \varepsilon_0 \mathbb{Z}[1/2]^3$, which is dense in $\mathbb{R}^3$.

Step 1 (dyadic translations). Let $b \in \mathcal{D}$ with $2(R + |b|) + 2 < \ell$, say $b \in \varepsilon_K \mathbb{Z}^3$. The lattices are nested (P4), so $b \in \varepsilon_i \mathbb{Z}^3$ whenever $K_i \geq K$, which holds for all large i. For those i, Lemma B.32 gives $S^{(i)}_{n,m}(\tau_b f) = S^{(i)}_{n,m}(f)$. The tuples f and $\tau_b f$ are admissible, with entries in $C_c^{(R)}$ and $C_c^{(R+|b|)}$ respectively. So by Proposition 8.2(a) both sides converge, and $S_{n,m}(\tau_b f) = S_{n,m}(f)$.

Step 2 (continuity). Choose R' with $R + |a| < R'$ and $2R' + 2 < \ell$. Choose $a_k \in \mathcal{D}$ with $a_k \to a$ and $|a_k| < R' - R$; this is possible since $\mathcal{D}$ is dense and $|a| < R' - R$. Then $2(R + |a_k|) + 2 < \ell$, so Step 1 gives $S_{n,m}(\tau_{a_k} f) = S_{n,m}(f)$. All entries of $\tau_a f$ and $\tau_{a_k} f$ lie in $C_c^{(R')}$. On tuples from $C_c^{(R')}$, $S_{n,m}$ is multilinear (Lemma B.30(b)) and satisfies $|S_{n,m}(f)| \leq c_{n,m} \prod_i \|\mathbf{g}_i\|_2 \prod_k N(h_k)$ (Proposition 8.2(a)). On $C_c^{(R')}$ we have $\|\mathbf{g}\|_2 \leq (3|B_{R'}|)^{1/2} \|\mathbf{g}\|_\infty$ and $N(h) \leq \max(1, |B_{R'}|) \|h\|_\infty$. Put $C_{R'} := \max(1, |B_{R'}|, (3|B_{R'}|)^{1/2})$, and write $f_1, \dots, f_{n+m}$ for the entries of f. By multilinearity,

$$S_{n,m}(\tau_{a_k} f) - S_{n,m}(\tau_a f) = \sum_{l=1}^{n+m} S_{n,m}\big(\tau_a f_1, \dots, \tau_a f_{l-1},\, \tau_{a_k} f_l - \tau_a f_l,\, \tau_{a_k} f_{l+1}, \dots, \tau_{a_k} f_{n+m}\big).$$

Since $\|\tau_b f_l\|_\infty = \|f_l\|_\infty$, this gives

$$|S_{n,m}(\tau_{a_k} f) - S_{n,m}(\tau_a f)| \leq c_{n,m} C_{R'}^{n+m} \sum_l \|\tau_{a_k - a} f_l - f_l\|_\infty \prod_{l' \neq l} \|f_{l'}\|_\infty.$$

The right side tends to 0 because each f_l is uniformly continuous. Hence $S_{n,m}(\tau_a f) = \lim_k S_{n,m}(\tau_{a_k} f) = S_{n,m}(f)$. □

Corollary B.34 (translation invariance on admissible tuples)**.** *Let S be as in Lemma B.33, and let $a \in \mathbb{R}^3$ and f be such that f and $\tau_a f$ are both admissible. Then $S_{n,m}(\tau_a f) = S_{n,m}(f)$.*

Proof. Let the entries of f lie in $C_c^{(\rho_1)}$ and those of $\tau_a f$ in $C_c^{(\rho_2)}$, and put $\rho := \max(\rho_1, \rho_2)$, so that $2\rho + 2 < \ell$. Let $t \in [0,1]$ and let f_l be an entry of f. Every point of $\operatorname{supp} \tau_{ta} f_l = \operatorname{supp} f_l + ta$ has the form $(1-t)y + t(y+a)$ with $y \in \operatorname{supp} f_l \subset \bar{B}_\rho$ and $y + a \in \operatorname{supp} \tau_a f_l \subset \bar{B}_\rho$. Since $\bar{B}_\rho$ is convex, all entries of $\tau_{ta} f$ lie in $C_c^{(\rho)}$. Choose $M \in \mathbb{N}$ with $2(\rho + |a|/M) + 2 < \ell$. For $k = 0, \dots, M-1$, Lemma B.33, applied with $R = \rho$, the tuple $\tau_{ka/M} f$ and the vector a/M, gives $S_{n,m}(\tau_{(k+1)a/M} f) = S_{n,m}(\tau_{ka/M} f)$. Chaining these M identities gives the claim. □

Periodic test functions. Let $C(T)$ be the space of real continuous functions on T, and $C(T; \Lambda^2)$ the space of real continuous 2-forms on T, both with the sup norm. Their elements are sampled by $h_\varepsilon(x) := h(x)$ for $x \in T_\varepsilon$, and by $\mathbf{g}_\varepsilon(c) := \mathbf{g}_{z(c)}(v_1, v_2)$ for plaquettes $c = (x; v_1 \wedge v_2)$, where $z(c) = x + \frac{\varepsilon}{2}(v_1 + v_2) \in T$; no identification with Q is needed. Write $S^{(i),\mathrm{per}}_{n,m}$ for the corresponding lattice Schwinger functions. Put $\|h\|_{L^1(T)} := \int_T |h|$, $\|\mathbf{g}\|^2_{L^2(T)} := \sum_{\mu<\kappa} \int_T g_{\mu\kappa}^2$ and $N_T(h) := \max(\|h\|_{L^1(T)}, \|h\|_\infty)$. The periodisation of $h \in C_c^{(\rho)}$ with $2\rho + 2 < \ell$ is $(\mathsf{P}h)(x + \ell\mathbb{Z}^3) := \sum_{v \in \mathbb{Z}^3} h(x + \ell v)$, a finite sum; the periodisation of a 2-form is defined componentwise. Translations

act on $C(T)$ and $C(T;\Lambda^2)$ by the same formulas as on $\mathbb{R}^3$, and $\tau_{a+\ell v}=\tau_a$ for $v\in\mathbb{Z}^3$, so $a\mapsto\tau_a$ is an action of $\mathbb{R}^3/\ell\mathbb{Z}^3$.

Proposition B.35 (periodic extension and invariance under $\mathbb{R}^3/\ell\mathbb{Z}^3$)**.** *Let S be the limit of Lemma B.33, along the subsequence (i) given there. There is a further subsequence along which:*

(a) *$S^{(i),\mathrm{per}}_{n,m}(f)$ converges for all n,m and every tuple f with entries in $C(T;\Lambda^2)$ (2-forms) and $C(T)$ (scalars). The limit $S^{\mathrm{per}}_{n,m}$ is real, multilinear and symmetric within each field type, and satisfies $|S^{\mathrm{per}}_{n,m}(f)|\le c_{n,m}\prod_i\|\mathbf{g}_i\|_{L^2(T)}\prod_k N_T(h_k)$;*
(b) *$S^{\mathrm{per}}_{n,m}(\mathsf{P}f)=S_{n,m}(f)$ for every admissible tuple f;*
(c) *$S^{\mathrm{per}}_{n,m}(\tau_a f)=S^{\mathrm{per}}_{n,m}(f)$ for every $a\in\mathbb{R}^3$ and every periodic tuple f; that is, S^{per} is invariant under $\mathbb{R}^3/\ell\mathbb{Z}^3$.*

Proof. (a) The proof of Proposition 8.1 uses only (6.5), which holds for arbitrary real lattice sources. So $|S^{(i),\mathrm{per}}_{n,m}(f)|\le c_{n,m}\prod_i\|(\mathbf{g}_i)_{\varepsilon_i}\|_{2,\varepsilon_i}\prod_k N_{\varepsilon_i}((h_k)_{\varepsilon_i})$. The torus has $(\ell/\varepsilon)^3$ sites and $(\ell/\varepsilon)^3$ plaquettes of each orientation. Hence $\|h_\varepsilon\|_{L^1,\varepsilon}\le\ell^3\|h\|_\infty$ and $\|\mathbf{g}_\varepsilon\|_{2,\varepsilon}\le(3\ell^3)^{1/2}\|\mathbf{g}\|_\infty$. So the lattice Schwinger functions are bounded by $c_{n,m}C_T^{n+m}\prod_l\|f_l\|_\infty$, uniformly in i, with $C_T:=\max(1,\ell^3,(3\ell^3)^{1/2})$. They are multilinear and symmetric within each field type, by the argument of Lemma B.30(a): sampling is linear, and F and X_R are linear in the source. Hence they are uniformly Lipschitz on bounded sets of the product of the sup-normed spaces. Since T is a compact metric space, $C(T)$ and $C(T;\Lambda^2)$ are separable. Choose countable dense subsets. A diagonal argument over these subsets and over all (n,m) gives a subsequence along which $S^{(i),\mathrm{per}}_{n,m}$ converges on all tuples drawn from them. By the uniform Lipschitz bound, as in step (ii) of the proof of Proposition 8.2, the convergence extends to all periodic tuples. Reality, multilinearity and symmetry are identities between finitely many convergent values, so they pass to the limit. The functions are uniformly continuous, so the Riemann sums converge: $\|h_\varepsilon\|_{L^1,\varepsilon}\to\|h\|_{L^1(T)}$ and $\|\mathbf{g}_\varepsilon\|_{2,\varepsilon}\to\|\mathbf{g}\|_{L^2(T)}$ as $\varepsilon\to0$. Moreover, lattice sup norms are at most the continuum sup norms. This gives the bound.

(b) Let the entries of f lie in $C_c^{(\rho)}$ with $2\rho+2<\ell$. For a site $x\in Q$ and $0\neq v\in\mathbb{Z}^3$, some coordinate of $x+\ell v$ has modulus at least $\ell/2$, so $|x+\ell v|\ge\ell/2>\rho$; hence $(\mathsf{P}h)(x)=h(x)$. For a plaquette $c=(x;v_1\wedge v_2)$ with $x\in Q$ and $z=x+\frac{\varepsilon}{2}(v_1+v_2)$, we have $|z+\ell v|\ge\ell/2-1>\rho$ for $v\neq0$, so $(\mathsf{P}\mathbf{g})_z=\mathbf{g}_z$. Thus the sampled sources of $\mathsf{P}f$ and of f coincide, and so do the lattice Schwinger functions, for every i. The subsequence of (a) is a subsequence of the one along which $S^{(i)}_{n,m}(f)\to S_{n,m}(f)$, and (b) follows.

(c) Let $b\in\varepsilon_i\mathbb{Z}^3$. Then $(\tau_b h)_{\varepsilon_i}(x)=h(x-b)=(\tau_b h_{\varepsilon_i})(x)$ on T_{ε_i}. Since $z(\tau_{-b}c)=z(c)-b$ in T, also $(\tau_b\mathbf{g})_{\varepsilon_i}(c)=\mathbf{g}_{z(c)-b}(v_1,v_2)=\mathbf{g}_{\varepsilon_i}(\tau_{-b}c)=(\tau_b\mathbf{g}_{\varepsilon_i})(c)$. By Lemma B.31(iii), $S^{(i),\mathrm{per}}_{n,m}(\tau_b f)=S^{(i),\mathrm{per}}_{n,m}(f)$. If $b\in\varepsilon_K\mathbb{Z}^3$, this holds for all i with $K_i\ge K$, by nesting (P4). Passing to the limit with (a) gives invariance under the dense set $\mathcal{D}=\varepsilon_0\mathbb{Z}[1/2]^3$. For arbitrary a, take $a_k\in\mathcal{D}$ with $a_k\to a$. The telescoping estimate of Step 2 of the proof of Lemma B.33, with $C_{R'}$ replaced by C_T and with the bound of (a), gives

$$|S^{\mathrm{per}}_{n,m}(\tau_{a_k}f)-S^{\mathrm{per}}_{n,m}(\tau_a f)|\le c_{n,m}C_T^{n+m}\sum_l\|\tau_{a_k-a}f_l-f_l\|_\infty\prod_{l'\neq l}\|f_{l'}\|_\infty\to0,$$

because continuous functions on the compact torus are uniformly continuous. □

By (b) and (c), $S_{n,m}(\tau_a f)=S^{\mathrm{per}}_{n,m}(\mathsf{P}\tau_a f)=S^{\mathrm{per}}_{n,m}(\tau_a\mathsf{P}f)=S^{\mathrm{per}}_{n,m}(\mathsf{P}f)=S_{n,m}(f)$ whenever f and $\tau_a f$ are admissible. This gives a second proof of Corollary B.34. Lemma B.33 is the fixed-torus clause of Theorem 9.2(a).

Remark B.36 (the torus group). At a fixed torus, the continuum states of Proposition 8.2 are defined only on admissible tuples, so the group $\mathbb{R}^3/\ell\mathbb{Z}^3$ does not act on their domain. Indeed, let $f\neq0$ be admissible with entries in $C_c^{(\rho)}$, $2\rho+2<\ell$, and let $0\neq a\in\ell\mathbb{Z}^3$. Every point

$y+a$ with $y \in \bar{B}_\rho$ satisfies $|y+a| \geq \ell - \rho > \ell/2 + 1$, so $\tau_a f$ is not admissible. For these states, Corollary B.34 is therefore the complete statement of translation invariance: $S_{n,m}(\tau_a f) = S_{n,m}(f)$ whenever both sides are defined. Invariance under the torus group itself is a statement about the periodic extension of Proposition B.35. That extension exists along a further subsequence and agrees with S on periodisations of admissible tuples.

Parts (a) and (c) of Proposition 8.7 on the class $\mathcal{D}$.

Lemma B.37 (class-$\mathcal{D}$ monotonicity)**.** *Fix real test functions $\mathbf{g}_1, \dots, \mathbf{g}_n, h_1, \dots, h_p$, and let*

$$E^{\mathcal{D}}_{\mathrm{obs}} := \{\zeta = (x,w) \in \mathbb{R}^{n+p} : \ k(\zeta) \leq 0 \text{ pointwise}, \ \|k(\zeta)\|_\infty < H/2\}, \qquad k(\zeta) = \sum_k w_k h_k.$$

This is the set of real points of the tube at which $\Theta(\zeta)$ lies in the class $\mathcal{D}$ of Subsection 2.4. Along the subsequence of Proposition 8.7, for every $\zeta \in E^{\mathcal{D}}_{\mathrm{obs}}$: $\mu \mapsto G^\mu(\zeta)$ is non-increasing on Q_0; $G^\mu(\zeta) \leq G^\flat(\zeta)$; and $G^0(\zeta) \leq G^\flat(\zeta)$.

Proof. Note first that $E^{\mathcal{D}}_{\mathrm{obs}} \subset D_{\sigma_+,H}$, since $k_+(\zeta) = 0$ and $k_-(\zeta) = -k(\zeta)$. Fix $\zeta \in E^{\mathcal{D}}_{\mathrm{obs}}$ and put $\mathbf{g} := \sum_i x_i \mathbf{g}_i$ and $q := -k(\zeta) \geq 0$, with $\|q\|_\infty < H/2$. Fix also a member $(\varepsilon, T) = (\varepsilon_i, T_{j_i})$ of the subsequence, and write $\mathbf{g}_\varepsilon, q_\varepsilon$ for the sampled functions; then $q_\varepsilon \geq 0$. Linearity of sampling and of the fields (Lemma B.30) gives

$$\Theta(\zeta) = \mathrm{e}^{F(\mathbf{g}_\varepsilon)} \mathrm{e}^{-X_R(q_\varepsilon)} = c_\varepsilon \, \mathrm{e}^{-r^2(\gamma)} \mathrm{e}^{\ell(A)},$$
$$\gamma_x := \varepsilon^3 q_\varepsilon(x) \geq 0, \quad \ell(A) := \langle d^* \mathbf{g}_\varepsilon, \, A \rangle_\varepsilon,$$

with $c_\varepsilon := \exp\big((W_\varepsilon + z_\varepsilon) \sum_x \varepsilon^3 q_\varepsilon(x)\big) > 0$. By (P1), c_ε does not depend on μ_0, and the matter part S_m of (7.1) does not either. *Covariances.* By (7.3), $C_\mu = (-\Delta_\varepsilon + \mu^2)^{-1}$. For $0 < \mu' \leq \mu$, Lemma B.14(b) gives $C_\mu \leq C_{\mu'}$ as forms, the zero mode included; on the zero mode this is $s(\mu) = 1/(\mu^2 |T|) \leq s(\mu')$. *Lattice inequalities.* Lemma 7.5 (class $\mathcal{D}$, second inequality) with $C_0 = C_\mu \leq C_1 = C_{\mu'}$ gives $\langle \Theta(\zeta) \rangle_{\varepsilon,T,\mu} \leq \langle \Theta(\zeta) \rangle_{\varepsilon,T,\mu'}$; the constant c_ε factors out of the normalised expectations. $\Theta(\zeta)$ is an elementary observable, being a product of $\mathrm{e}^{tF(\mathbf{g})}$ $(t = 1)$ and $\mathrm{e}^{-X_R(q)}$ with $q \geq 0$. So Lemma 7.6 gives $\langle \Theta(\zeta) \rangle_{\varepsilon,T,\flat} = \lim_{\mu' \downarrow 0} \langle \Theta(\zeta) \rangle_{\varepsilon,T,\mu'} \geq \langle \Theta(\zeta) \rangle_{\varepsilon,T,\mu}$. *Limits.* Since $\zeta \in D_{\sigma_+,H}$, Proposition 8.2(b) gives $G^\mu_i(\zeta) \to G^\mu(\zeta)$ for each $\mu \in Q_0$ and $G^\flat_i(\zeta) \to G^\flat(\zeta)$. Non-strict inequalities pass to these limits, which gives the first two claims. By Proposition 8.7(b), $G^\mu \to G^0$ locally uniformly on $D_{\sigma_+,H}$, in particular at ζ. So $G^0(\zeta) = \lim_{\mu \downarrow 0, \, \mu \in Q_0} G^\mu(\zeta) \leq G^\flat(\zeta)$. Finally, the lattice value at ζ depends only on $(\mathbf{g}, q)$, by linearity; hence so does the limit. Thus $S^{\mu_0}(\Theta) := G^{\mu_0}(\zeta)$ is well defined for $\Theta = \exp[F(\mathbf{g}) - X_R(q)] \in \mathcal{D}$, which is the meaning of Theorem 3.1(iv). So parts (a) and (c) of Proposition 8.7 hold on $E^{\mathcal{D}}_{\mathrm{obs}}$, which gives Theorem 3.1(iv). □

Appendix C. Counterterm and vacuum identifications

This appendix proves Proposition 4.7: King's mass counterterm and vacuum constant on a lattice torus are Bałaban's (Lemmas C.1 and C.2), and it states and proves Reisz's power-counting theorems in three dimensions, item 3 of Subsection 4.6 (Proposition C.3). The torus is $T = T_j$ with spacing ε; $-\Delta_\varepsilon = d^* d + d d^*$ on bond functions.

King's counterterm and vacuum constant are Bałaban's. Fix a torus $T = T_j$, a spacing ε, a Higgs mass $M > 0$ and the reference photon mass $\mu_* > 0$. *King's model* is the action [24, (2.4), p. 652] on $\Omega = T_{j,\varepsilon}$ with $m^2 = M^2$ and $\mu_0 = \mu_*$, together with its counterterm $\delta m^2_{K,T}(\varepsilon; M)$ (Definition 2.2). *Bałaban's model* is the action [1, (1.11), p. 605] on the same lattice torus, in the Feynman gauge, with $m^2 = M^2$, $\mu_0 = \mu_*$ and $m_0^2 = m^2 + \delta m^2$. Write $\delta m^2_{\mathrm{B}}(\varepsilon; M)$ for Bałaban's counterterm as defined explicitly in [3, (1.19)–(1.23), pp. 416–417]. The paper [1] only names δm^2 ("it can be written down explicitly but we will not do it", [1, p. 605]); [3] completes the definition, and we read the δm^2 of [1] as this one.

Lemma C.1 (King's mass counterterm is Bałaban's)**.** *On $T_{j,\varepsilon}$, $\delta m^2_{K,T}(\varepsilon;M) = \delta m^2_{\mathrm{B}}(\varepsilon;M)$ coefficient by coefficient in $e^\alpha\lambda^\beta$, $2 \le \alpha+2\beta \le 4$, as an identity of finite lattice sums. Hence $a_{K,T}(M) = M^2 + \delta m^2_{\mathrm{B}}(\varepsilon;M)$ is the bare coefficient m_0^2 of* [1, (1.11)] *on $T_{j,\varepsilon}$, i.e. the coefficient at which Bałaban's theorems (Theorems 4.1, 4.2;* [3, p. 417]*) are stated. This is the mass part of Proposition 4.7 and Remark 2.3(i).*

Proof. Step 1 (the two actions agree term by term). King's action is $\frac12\langle A, (-\Delta^{\varepsilon,\Omega}+\mu_0^2)A\rangle + \frac12\langle\varphi, (-\Delta_A^{\varepsilon,\Omega}+m^2)\varphi\rangle + \sum_{x\in\Omega}\varepsilon^3(\frac12\delta m^2(x)|\varphi(x)|^2+\lambda|\varphi(x)|^4)+E_0+E_1$ [24, (2.4), p. 652]. Bałaban's is $\frac12\langle\varphi, (-\Delta_A^{\varepsilon})\varphi\rangle + \sum_{x\in T_\varepsilon}\varepsilon^3(\frac12 m_0^2|\varphi|^2+\lambda|\varphi|^4) + \frac12\langle A, (-\Delta^{\varepsilon}+\mu_0^2)A\rangle + E$ [1, (1.11), p. 605]. Both use the same covariant difference operator, with $U(A)=\exp(q\varepsilon eA)$ [1, (1.7)], [24, p. 651], and both photon terms are in the Feynman form. On $\Omega = T_{j,\varepsilon}$, King's $\delta m^2(x)$ is constant by translation invariance. So the vertices are the same, and so are the propagators: the scalar propagator $C_0^\varepsilon = (-\Delta_0^\varepsilon + M^2)^{-1}$ and the photon propagator $C^\varepsilon = (-\Delta^\varepsilon+\mu_*^2)^{-1}$ (which is $(-\Delta_\varepsilon+\mu_*^2)^{-1}$ on bond functions, Subsection 2.2), both on $T_{j,\varepsilon}$ [3, p. 416].

Step 2 (the two defining equations). King writes $(-\delta m^2+\Gamma^\varepsilon)$ for "the full one particle irreducible graph with two (undifferentiated) external scalar field lines". He sets $\delta m^2(x) = \sum_{2\le\alpha+2\beta\le4} e^\alpha\lambda^\beta\delta m^2(x;\alpha,\beta)$ and imposes $-\delta m^2(x;\alpha,\beta) + \sum_{y\in\Omega}\varepsilon^d\Gamma^\varepsilon(x,y;\alpha,\beta) = 0$, "solved recursively" [24, (3.32)–(3.33), p. 659]. Bałaban expands the two-point function on T_ε as $G^\varepsilon = \sum_n C_0^\varepsilon[(-\delta m^2+\Sigma^\varepsilon+\partial^{\varepsilon*}\Sigma_1^\varepsilon+\Sigma_1^{\varepsilon*}\partial^\varepsilon+\partial^{\varepsilon*}\Sigma_2^\varepsilon\partial^\varepsilon)C_0^\varepsilon]^n$, with $\Sigma^\varepsilon,\Sigma_1^\varepsilon,\Sigma_2^\varepsilon$ amputated one-particle-irreducible [3, (1.19)–(1.21), p. 416]. He then imposes $-\delta m^2_{(\alpha,\beta)} + \sum_{x\in T_\varepsilon}\varepsilon^d\Sigma^\varepsilon_{(\alpha,\beta)}(x) = 0$ for $2\le\alpha+2\beta\le4$ [3, p. 417]. Both definitions include all graphs of each order, with the lower-order counterterms inserted.

Step 3 (the parts with derivative legs have zero sum). Let $K(x,y)$ be a translation-invariant kernel on $T_{j,\varepsilon}$. If $K = B\circ d$, with d the lattice derivative of Subsection 2.2 acting componentwise, then $\sum_y\varepsilon^3K(x,y) = (B\,d1)(x) = 0$, because $d1=0$. If $K = d^*\circ B'$ with B' translation invariant, then $(B'1)_{x,\mu} =: c_\mu$ does not depend on x, and

$$(d^*c)(x) = \sum_\mu \varepsilon^{-1}\big(c_\mu - c_\mu\big) = 0.$$

The kernels $\Sigma_1^{\varepsilon*}\partial^\varepsilon$ and $\partial^{\varepsilon*}\Sigma_2^\varepsilon\partial^\varepsilon$ are of the first type and $\partial^{\varepsilon*}\Sigma_1^\varepsilon$ is of the second. So the zero-momentum sum of the full amputated one-particle-irreducible two-point function equals $\sum_x\varepsilon^3\Sigma^\varepsilon(x)$. The same sum also does not depend on how a convention divides that function into undifferentiated and differentiated legs. By translation invariance, $\sum_y\varepsilon^3\Gamma^\varepsilon(x,y)$ does not depend on x. Hence both conditions of Step 2 say the same thing: at each order, $\delta m^2_{(\alpha,\beta)}$ is the zero-momentum value of the order-(α,β) amputated one-particle-irreducible two-point function on $T_{j,\varepsilon}$, with the lower-order counterterms inserted.

Step 4 (induction on the order). Order the coefficients by $w := \alpha+2\beta \in \{2,3,4\}$. In a one-particle-irreducible two-point graph of order (α,β) that contains a counterterm insertion $\delta m^2_{(\alpha',\beta')}$, there is at least one other vertex, which carries a factor e, e^2 or λ; so $\alpha'+2\beta' < w$. Assume the counterterms of the two models agree at all orders with $\alpha'+2\beta' < w$. By Step 1, the order-(α,β) coefficients of the two two-point functions are then the same finite lattice sums, and by Step 3 the two equations of Step 2 give the same number. The odd orders in e vanish in both models (Definition 2.2). □

Lemma C.2 (the vacuum constants)**.** *Let $\bar n = 13$, and let $\delta m^2 = \delta m^2_{K,T}(\varepsilon;M) = \delta m^2_{\mathrm{B}}(\varepsilon;M)$ (Lemma C.1).*

(a) *King's $E_0(M)$* [24, (2.6), p. 652] *equals Bałaban's $E_0 = E_{0,v}(\mu_*)+E_{0,s}(M)$* [1, (1.12), p. 606].
(b) *King's $E_1(M)$* [24, (3.35), p. 659] *equals Bałaban's E_1* [1, (1.13), p. 606]*. Hence $Z_K(T;M)$ (Subsection 4.2) equals Bałaban's Z^ε at $(m^2,\mu_0) = (M^2,\mu_*)$, with no correction term.*

(c) *With the photon measure normalised,* $Z(\Gamma;a) := \mathbb{E}_{N(0,\Gamma)}[Z_\varphi(A;a)]$ *as in Subsection 5.1, one has* $Z_K(T;M) = Z(C_{\mu_*};a_{K,T}(M))\,\mathrm{e}^{-E_{0,s}(M)-E_1(M)}$. *The constant* $\gamma_T(M)$ *of Proposition 6.3 formed in this pairing and the one formed in the Lebesgue pairing differ by* $|T|^{-1}E_{0,v}(\mu_*)$. *This term does not depend on* M *and cancels in every difference* $\gamma_T(M)-\gamma_T(M')$.

This is the vacuum-constant part of Proposition 4.7 and Remark 5.3(i).

Proof. Write S_{rest} for the common action of Step 1 of Lemma C.1 without its additive constants. It depends on (e,λ), also through δm^2, in the same way in both models. Both partition functions use the Lebesgue measure on the site and bond variables ([24, (2.5)]; "natural Lebesgue measure", [1, p. 605]).

(a) King: $\mathrm{e}^{E_0} = \int (dA)(d\varphi)\exp[-\frac{1}{2}\langle A,\,(-\Delta^{\varepsilon,\Omega}+\mu_0^2)A\rangle - \frac{1}{2}\langle\varphi,\,(-\Delta^{\varepsilon,\Omega}+m^2)\varphi\rangle]$. Bałaban: $\int dA\,\mathrm{e}^{-\frac{1}{2}\langle A,\,(-\Delta^{\varepsilon}+\mu_0^2)A\rangle}\int d\varphi\,\mathrm{e}^{-\frac{1}{2}\langle\varphi,\,(-\Delta_0^{\varepsilon}+m^2)\varphi\rangle} = \mathrm{e}^{E_{0,v}}\mathrm{e}^{E_{0,s}} = \mathrm{e}^{E_0}$. These are the same Gaussian integral: the photon factor is included on both sides, the sign is the same, and both are taken at $m^2=M^2$, not at m_0^2.

(b) King's (3.35) takes the Taylor coefficients, of total order $1\le\alpha+\beta\le\bar n$ at $e=\lambda=0$, of $\ln\int\exp[-S^{\varepsilon}+E_1] = \ln\int\mathrm{e}^{-S_{\mathrm{rest}}}-E_0$. Bałaban's (1.13) takes the same coefficients of $\log\int\exp(-S+E) = \log\int\mathrm{e}^{-S_{\mathrm{rest}}}$. The two functions differ by the constant E_0, which does not depend on (e,λ) (it is defined at $e=\lambda=0$), so every coefficient with $\alpha+\beta\ge1$ is the same on both sides. The two functions are therefore not literally the same: they agree only up to this constant. Both sides use the same index set $1\le\alpha+\beta\le\bar n$. King fixes $\bar n=13$ [24, p. 657], and Bałaban's definition admits it ("$\bar n>12$", [3, p. 418]). Hence E_0+E_1 is the same, and $Z_K(T;M)=\int\mathrm{e}^{-S_{\mathrm{rest}}-E_0-E_1}=Z^{\varepsilon}$.

(c) Since $\int dA\,\mathrm{e}^{-\frac{1}{2}\langle A,\,(-\Delta_{\varepsilon}+\mu_*^2)A\rangle} = \mathrm{e}^{E_{0,v}(\mu_*)}$, the Lebesgue integral equals $\mathrm{e}^{E_{0,v}(\mu_*)}Z(C_{\mu_*};a_{K,T}(M))$. With (b) this gives $Z_K = Z(C_{\mu_*};a_{K,T}(M))\,\mathrm{e}^{E_{0,v}-E_0-E_1} = Z(C_{\mu_*};a_{K,T}(M))\,\mathrm{e}^{-E_{0,s}-E_1}$. At $e=\lambda=0$ one has $a_{K,T}(M)=M^2$, so the order-zero term of $\log Z(C_{\mu_*};a_{K,T}(M))$ is $E_{0,s}(M)$, as stated in Subsection 5.1. Its Taylor coefficients of order at least 1 are those of E_1. So the two pairings give $\gamma_T(M)$'s that differ by $|T|^{-1}E_{0,v}(\mu_*)$, which does not depend on M. Taking the photon-normalised quantity together with Bałaban's full E_0 would be off by $|T|^{-1}E_{0,v}(\mu_*)$ per volume, and this diverges as $\varepsilon\to0$. □

Reisz's power counting in three dimensions. We use Reisz's notation [32, 33], with a the lattice spacing. $\mathcal{F}_{\mathrm{R}}$ is Reisz's class of lattice Feynman integrands $F=V/\prod_i(\eta_i(l_ia)/a^2+\mu_i^2)$ ([32, Def. 2.5]). Their numerators $V=a^{-m}F(ua,wa)$ are $(2\pi/a)$-periodic in the loop momenta, $\mu_i^2>0$ for every line, and the line momenta form a *natural* set ([32, Def. 3.1]). For a Zimmermann subspace H of the loop space, degr_H is Reisz's UV degree. $\mathrm{BZ}=[-\pi/a,\pi/a]^{3m}$ is the Brillouin zone of m three-dimensional loop momenta, and $\hat l^2 := \sum_\mu(4/a^2)\sin^2(al_\mu/2)$ is King's lattice momentum square. Reisz states both theorems for $d=4$ and remarks that "the following considerations can easily be extended to other dimensions" [32, p. 86].

Proposition C.3 (Reisz's theorems in three dimensions)**.** *Let* $d=3$.

(a) *Let* $F\in\mathcal{F}_{\mathrm{R}}$ *have three-dimensional momenta, positive masses and a natural set of line momenta, and let* $3\dim H+\mathrm{degr}_H F<0$ *for every Zimmermann subspace* H. *Then* $\lim_{a\to0}\int_{\mathrm{BZ}}F$ *exists, and there are* $a_0>0$ *and* $C<\infty$ *with* $\int_{\mathrm{BZ}}|F(k,q;a)|\,d^{3m}k\le C$ *for all* $a<a_0$, *uniformly for external momenta* q *in a bounded set.*

(b) *Let* Γ *be a one-particle-irreducible diagram with massive lines, renormalised by Reisz's forest formula with subtraction degrees* $\delta(\gamma)$ *obeying* [33, (4-3)–(4-4)]; *in particular* $\delta(\gamma)=\omega(\gamma)$, *with* ω *computed with* $3m(\gamma)$. *Then the forest integrand* R_Γ *lies in* $\mathcal{F}_{\mathrm{R}}$ *with* $3h+\mathrm{degr}_H R_\Gamma\le-1$ *for every* H, *so (a) applies to* R_Γ.

(c) *Suppose in addition that every vertex function and every coefficient of a subtraction operator is entire in its momentum arguments, and that every propagator is* $(\hat l^2+\mu^2)^{-1}$. *Let* $\mu_{\min}$ *be the smallest mass. Let* c_Γ *be the largest value, over all line and subdiagram momenta of*

R_Γ, of the sum of the absolute values of their coefficients on the loop momenta. Fix $\eta_0 > 0$ with $1.087\, c_\Gamma^2 \eta_0^2 \le \mu_{\min}^2/2$ and $c_\Gamma \eta_0 \le 1$, and let $a_0 \le 1$. Then

$$\sup_{a<a_0} \sup_{\eta\in\mathbb{R}^{3m},\ \max_j|\eta_j|\le\eta_0} \int_{\mathrm{BZ}} |R_\Gamma(k+i\eta, q; a)|\, d^{3m}k \le C' < \infty.$$

The constants depend on the graph, the masses, the bound on the external momenta and η_0; they do not depend on ε or on the torus, and they are not explicit. This is item 3 of Subsection 4.6, as used in Proposition 6.3.

For (a) and (b) the proofs are those printed in [32, 33] for $d = 4$; the proof below does not repeat them, but lists every place where the dimension enters and checks each one, and the bound form is read off the majorants of the printed proofs. Part (c) is deduced from (a) and (b).

Proof. Step 1 (what Reisz's proof bounds). The integral over BZ is split by non-negative cut-off functions into finitely many sectors: lines near a pole and lines away from all poles [32, (4.1)–(4.7)]. Naturalness allows the poles to be translated to 0 [32, App. D]. Then [32, (4.8)–(4.9)] give $1/(\eta(la)/a^2 + \mu^2) \le \alpha/(l^2 + \mu^2)$ near a pole and $\le \gamma a^2$ away from the poles. By [32, (7.4)–(7.5b)], each sector integral is bounded by integrals of non-negative majorants, and these are controlled by the continuum power-counting theorem [32, Thm. 2, p. 96]. Hence the proof bounds $\int_{\mathrm{BZ}}|F|$, not only F, uniformly for bounded q [32, pp. 96, 102]. Complex-valued numerators are allowed: $|V| \le |\mathrm{Re}\, V| + |\mathrm{Im}\, V|$, and taking real and imaginary parts does not raise degrees. Neither does the photon index [33, Sect. 6, p. 100].

Step 2 (where the dimension enters). The dimension enters only as the dimension of each loop momentum, and the following replacements are the only changes. (1) The degree definitions: [32, (3.8), (5.9), (7.9), (7.14), (9.3), (9.29)] and [33, (2-28), (4-11)–(4-16), (5-5), (5-10)]. Each has the form "4× (dimension of the loop space)", and $4 \to 3$. (2) The counting of [33, App. C], which is a *lower* bound, $M_U(\Gamma) \ge 4h$ and $M_U(\gamma) \ge 4\,\mathrm{rank}$ [33, (C-1)–(C-2), p. 106]. Its proof is the count $M_U = 4\cdot\#$, so it becomes $M_U \ge 3h$; then [33, (5-13)] gives $3h + \mathrm{degr}_H R_\Gamma \le -1$. (3) The scaling step [32, Sect. 9]. The outer variable is $t_1 \in \mathbb{R}^3$ and the inner variables $(t_2, \dots, t_m) \in \mathbb{R}^{3(m-1)}$ are rescaled by $\|t_1\|$ [32, (9.16)]. The Jacobian $\|t_1\|^{3(m-1)}$ gives the factor $\|t_1\|^{\omega_0 - 3}$ with $\omega_0 := 3m + \Delta - 2\sum n_j$ [32, (9.17a)]. (4) Two constants that use the dimension implicitly. The choice of δ in [32, (7.8), p. 102] needs $\sqrt{d} \le 4$. The proof of [32, Lemma D.1, p. 125] uses $\|k_j\| \le 2\pi/a$ on $[-\pi/a, \pi/a]^d$, i.e. $\sqrt{d} \le 2$. Both hold for $d = 3$. All the other ingredients are dimension-free: the propagator bounds (4.8)–(4.9), Theorem 3 and Lemmas 6.1–6.3, Lemmas 8.1–8.10, Appendices A–D of [32], the forest formula, Lemma 3.1 and Lemmas 5.1–5.2 of [33], and the subtraction operators t^0, t^1 (with $\sin(\varepsilon q)/\varepsilon$) of [33, App. A]. The IR hypothesis is only $\mu_i^2 > 0$, which enters through lower bounds that do not depend on the dimension. Massless lines are not covered, and no graph bound of the construction uses one.

Step 3 (the outer integral, which Reisz does not print). Reisz writes "the remaining integration can be done without problems" [32, p. 119]. Here is that integral. With $\lambda' = b(\lambda + d)/\|t_1\| + e$ [32, Lemma 9.1, p. 115], polar coordinates turn $r^2 \cdot r^{\omega_0 - 3}$ into $r^{\omega_0 - 1}$. The outer integral $\int_1^{\lambda+K} r^{\omega_0-1} f(\lambda')\, dr$, with f from [32, (9.24)], therefore has the same exponents as for $d = 4$. If $f = 1$ and $\omega_0 < 0$, it is bounded. If $f = \lambda'^{-1} \log^{m-1} \lambda'$ and $\omega_0 \le -1$, it is $\le c\lambda^{-1} \log^m \lambda$. If $f = \lambda'^{\bar\omega} \log^{m-1} \lambda'$, it is $\le c\lambda^{\max(\omega_0, \bar\omega)} \log^m \lambda$. This is [32, (9.30)] in every dimension. Lemma 9.2 then applies the induction hypothesis to the inner integral, with $\omega(\tilde H) = \omega(H)$ [32, (9.29)]. This proves (a), and (b) follows by Step 2(2).

Step 4 (proof of (c)). Write $R_\Gamma = V/(B_1 B_2)$ [33, (5-1)], with $V = \sum_i a^{-m_i} F_i(ka, qa)$ and each F_i entire. *Numerator.* Put $V_\eta(k) := V(k + i\eta) = \sum_i a^{-m_i} \tilde F_i(ka, qa, \eta a)$ with $\tilde F_i(x, y, z) := F_i(x + iz, y)$. These are smooth and 2π-periodic in x, and $a^{-m_i}\tilde F_i(ka, qa, \eta a) \to P_i(k + i\eta, q)$. So V_η is in Reisz's numerator class, with (q, η) as external variables. For a Zimmermann subspace with internal

variables u and complementary variables w, the shift $k \mapsto k+i\eta$ acts as $(u,w) \mapsto (u+iA\eta, w+iB\eta)$ with fixed real matrices A, B. Each derivative $\partial_w^b \partial_\eta^c$ at $(w,\eta) = 0$ is a combination of terms $\partial_u^{c_1}\partial_w^{b+c_2} F(u,0)$ with $|c_1|+|c_2| = |c|$. If $|b|+|c|$ is below Reisz's vanishing order r_u, these vanish identically. Hence $\operatorname{degr}_u V_\eta \le \operatorname{degr}_u V$. *Denominator.* For $l = \kappa + i\zeta$ with $|a\zeta_\mu| \le 1$,

$$\operatorname{Re}\big(1-\cos(al_\mu)\big) = (1-\cos a\kappa_\mu) - \cos(a\kappa_\mu)(\cosh a\zeta_\mu - 1) \ge (1-\cos a\kappa_\mu) - (\cosh a\zeta_\mu - 1).$$

Moreover $(2/a^2)(\cosh a\zeta - 1) = \zeta^2\big(\sinh(a\zeta/2)/(a\zeta/2)\big)^2 \le 1.0862\,\zeta^2$. Hence $\operatorname{Re}\hat{l}^2 \ge \hat{\kappa}^2 - 1.087|\zeta|^2$, and therefore $|\hat{l}^2+\mu^2| \ge \hat{\kappa}^2 + \mu^2/2$ whenever $1.087|\zeta|^2 \le \mu^2/2$. Every momentum in B_1B_2 is a line or subdiagram momentum, so $|\zeta| \le c_\Gamma \max_j|\eta_j| \le c_\Gamma\eta_0$. Since $a < a_0 \le 1$ and $c_\Gamma\eta_0 \le 1$, also $|a\zeta_\mu| \le 1$. Hence $|B_1B_2(k+i\eta)| \ge \tilde{B}(k)$, where $\tilde{B}$ is B_1B_2 with every mass μ replaced by $\mu/\sqrt{2}$. *Conclusion.* $\tilde{F} := V_\eta/\tilde{B} \in \mathcal{F}_{\mathrm{R}}$, with the same natural set of line momenta, and masses do not enter degrees. So $3\dim H + \operatorname{degr}_H\tilde{F} \le 3\dim H + \operatorname{degr}_H V - \operatorname{degr}_H(B_1B_2) \le -1$ by the numerator estimate and (b). Since $|R_\Gamma(k+i\eta)| \le \tilde{F}(k)$, (a) gives the bound, uniformly for $a < a_0$ and $\max_j|\eta_j| \le \eta_0$. □

Remark C.4. (i) For King's lattice Feynman rules, the degree of a one-particle-irreducible (sub)diagram is $\omega = 3 - \frac{1}{2}(E_\varphi + E_A) - o_V - 2v_2$, and every vertex has Reisz vertex dimension below 3. The model is therefore super-renormalisable, and only finitely many classes have $\omega \ge 0$. (ii) What Proposition 6.3 uses on the tori T_j goes beyond Proposition C.3: the Poisson resummation with a strip shift, the torus Ward defect at $k = 0$, the graph of order e^2 that is logarithmic overall, and the identification of King's counterterm insertions with Reisz's forests. These are treated in Appendix H (Proposition H.8 and Lemmas H.12, H.13).

Appendix D. Bałaban's Proposition 1 on the torus

This appendix proves the statements about [3, Prop. 1] that Subsection 5.3 uses: the anchoring of the vertices and the depth of the anchored classes in the proof of Corollary 5.10, the depth of the field $\tilde{A}$ at the places where the proposition is applied, and the list of sketched places behind Proposition 5.8(a) (Subsection D.1); the bound on the kernel [3, (1.16)] and on domain differences of propagators in the mixed norm used in steps S1 and S2 (Subsection D.2); and the weighted ℓ^2 and kernel bounds on which they rest (Subsection D.3). Throughout $L = 2$, distances are in scale-k units, and the notation is that of Subsections 4.4 and 5.3.

D.1. **Anchoring and depth in Corollary 5.10; the sketched places.**

Conventions. Distances are in the ℓ^∞ torus metric [1, (1.3), p. 604], in scale-k units; $r_j := r(t_j)$, and $r := r_k$ at step k. Since $r(t)$ decreases in t, $r_{k-1} \ge r_k$. We use three nesting facts for the layers $\Lambda_i^{(j)}$ of [2, (2.7)–(2.8), p. 558; p. 566].

- A point of $\Lambda_{i+s}^{(j)}$ has distance $> s(r_j - 1)$ from $(\Lambda_i^{(j)})^c$, in scale-j units.
- Passing from block sets to the point sets $B^j(\cdot)$ costs at most 1 per set.
- One scale-$(k-1)$ unit is $1/L$ scale-k units.

Under a set-distance convention other than the infimum, each nesting step costs at most M_b more. We write M_θ for the thickness, in scale-k units, of the slice on which θ_{k+1} goes from 1 to 0 [2, p. 567]; $M_\theta \le LM_b$. For a vertex v of a class, its *vertex point* x_v depends on the vertex type.

- [3, (1.6), (1.7)]: the summation point.
- [3, (1.8)–(1.11)]: an end point of a bond on which a factor g_kA' or $\tilde{A}$ is non-zero.
- [3, (1.13)–(1.15)]: a point of the block $B^k(y)$ of the vertex.

The applications of [3, Prop. 1] in [2], at which the cube step P12 is used in the proof of Theorem 5.2, are the following; we list them with their outer domains Ω_{out} and keep their names.

- U0/U0′: Proposition 1 at step k, $\Omega_{\mathrm{out}} = B^k(\Lambda_2^{(k-1)\prime})$ [3, p. 411] (also [2, p. 582, (2.116)]).

- U1a: [2, p. 579, (2.106)–(2.107)], $\Omega_{\rm out} = B^k(\Lambda_0^{(k)})$, with $\operatorname{supp} g_k \subset \Lambda_7^{(k)}$.
- U1b: [2, p. 580, Prop. 2.6], $\Omega_{\rm out} = B^{k-1}(\Lambda_2^{(k-1)})$, with the terms localised in $B^k(\Lambda_7^{(k)})$.
- U2: [2, p. 578, (2.104)], at scale $j \le k-1$, $\Omega_{\rm out} = B^j(\Lambda_2^{(j)})$.

The supports of $\tilde A$ are examined at three sites: (P)=U0, (Q)=U1a with $\tilde A = \tau\theta_{k+1}A'^{(k)}$, and (Z)=U2.
Anchoring.

Lemma D.1 (every vertex is anchored)**.** *Read* [3, (1.4)] *as the definition of E_k, and* [3, (1.8)–(1.15)] *as its expansion. Then every vertex v of every class in* [3, (1.33)] *has a vertex point x_v with*

$$x_v \in \Omega_1 \qquad or \qquad \operatorname{dist}\big(x_v,\ \operatorname{supp} g_k \cup \operatorname{supp}\tilde A\big) \le 1.$$

Since $\operatorname{supp} g_k \subset \Omega_1$*, every vertex point lies within distance* 1 *of* $\Omega_1 \cup \operatorname{supp}\tilde A$*. In particular, every connected component of a class, after the regrouping S10 and the localisation S11 of Table 2, consists of anchored vertices only. This is the statement "every vertex of a class is anchored" used in the proof of Corollary 5.10.*

The reading of [3, (1.4)] as the definition of E_k is the convention R1 of Lemma E.128; as printed, [3, (1.14), (1.15)] carry no factor g_k.

Proof. (1) *Scalar vertices.* The vertices [3, (1.6), (1.7)] are sums over $x \in \Omega_1$ (p. 413). At p. 419 the text says: "We localize the vertices (1.6) and (1.7) restricting summations to the set Ω_1". So $x_v \in \Omega_1$.

(2) *Gating of A'.* In [3, (1.4)] the field A' enters the integrand only through the combination $e'g_kA'+A^{(k)}$, in the kernel T and in the covariant Laplacian. The counterterms $\delta m^2(e', g_k, \lambda', \Omega_1, x)$ and $E_1(e', g_k, \lambda', \Omega_1)$ contain no A'. Hence every A' produced by $\partial_{e'}$ at $e'=0$ in [3, (1.5)] carries the factor g_k, in the kernel T as well. So the factors $A'(\Gamma_{y,x}^{(k)})^n$ of (1.14), (1.15), which are printed without g_k, are $(g_kA')(\Gamma_{y,x}^{(k)})^n$. The form in [2] of the same polynomial, $Q^{(k)}(B^k(\Lambda_0^{(k)}); B^{(k+1),\eta}, g_kA'^{(k)})$ [2, (2.107)], is gated in the same way.

(3) *Bond vertices.* The vertices (1.8)–(1.11) (p. 413) are sums over bonds b of terms carrying $(g_kA'_b)^n(\tilde A_b)^{n'}$ with $n+n' \ge 1$. The R-vertices (1.9), (1.11) carry $(\tilde A_b)^{\bar n+1}$. A term vanishes unless b meets $\operatorname{supp} g_k$ or $\operatorname{supp}\tilde A$. An end point of b is at distance $\le \eta_k \le 1$ from that support. The $n=0$, $n' \ge 1$ vertices produced by the $\tilde A$-expansion of $G_k(\Omega, \tilde A + \tilde B)$ [3, p. 414] carry $\tilde A$-legs and are covered by the same count.

(4) *Block vertices.* By [3, p. 414] the expressions (1.12)–(1.15) enter only as pairs at the same $y \in T_1^{(k)} \cap \Omega$, "with the restriction that at least one of them has to be of the form (1.14) or (1.15)"; (1.13)–(1.15) are the vertices and (1.12) is an external field. A factor (1.14) with $n \ge 1$ vanishes unless the contour $\Gamma_{y,x}^{(k)}$ has a bond in $\operatorname{supp} g_k$, by step (2). A factor with $n' \ge 1$, and every factor (1.15), vanishes unless the contour has a bond in $\operatorname{supp}\tilde A$. The contours satisfy $\Gamma_{y,x}^{(k)} \subset B^k(y)$, bonds included ([1, (2.1)–(2.2), p. 608]). The block $B^k(y)$ has ℓ^∞-diameter < 1 in scale-k units. Hence both members of the pair, and so the (1.13) partner as well, have their vertex points within distance 1 of $\operatorname{supp} g_k \cup \operatorname{supp}\tilde A$.

(5) *Components.* Steps (1)–(4) apply to every vertex, so they apply to every vertex of every connected component produced by S10 and S11. □

Depth of the Ω_1-anchored classes.

Lemma D.2 (Ω_1-anchored classes at each use)**.** *Let $L=2$, $R \ge R^{**}$ (so* (5.11) *holds), and $r(t) \ge R$. Consider a class at one of the uses U0/U0′, U1a, U1b, U2, with outer domain $\Omega_{\rm out}$ and the cube $\square$ of P12 with $r_\square = r$. Suppose the class has an old vertex point x_v anchored at Ω_1 or at*

$\operatorname{supp} g_k$ *in the sense of Lemma D.1. Then* $m := \operatorname{dist}(x_v, \Omega_{\mathrm{out}}^c)$ *satisfies*

$$U0/U0'\!: \ m \geq 2r_k - 4; \qquad U1a\!: \ m \geq 7r_k - 10;$$
$$U1b\!: \ m \geq \tfrac{5}{L}(r_{k-1} - 1) - 2 \geq \tfrac{5}{2} r_k - 5; \qquad U2\!: \ m \geq 5r_j - 7,$$

where at U2 the scale is j *and* $r = r_j$*. In every case*

$$m > \tfrac{7}{4} r + 2s_b + \ell_{\bar{N}}, \tag{D.1}$$

hence $\square \subset \Omega_{\mathrm{out}}$ *and* $\operatorname{dist}(\square, \Omega_{\mathrm{out}}^c) > r/4$*. In particular* m *exceeds the need* $\frac{3}{2} r + 2s_b + \ell_{\bar{N}}$ *of P12. This is the case of a vertex anchored at* $\operatorname{supp} g_k$ *in the proof of Corollary 5.10.*

The lemma is stated at $L = 2$. At $L = 5$ the margin at U1b is $r_{k-1} - 3$, so $\square \subset \Omega_{\mathrm{out}}$ can fail for such classes.

Proof. Margins. (i) U1a: x_v is within 1 of $\operatorname{supp} g_k \subset \Lambda_7^{(k)}$ [2, p. 579], and $\Omega_{\mathrm{out}} = B^k(\Lambda_0^{(k)})$. Seven nesting steps give $m \geq 7(r_k - 1) - 2 - 1 = 7r_k - 10$. (ii) U1b: $x_v \in B^k(\Lambda_7^{(k)})$, since the terms of [2, Prop. 2.6, p. 580] are those localised in $B^k(\Lambda_7^{(k)})$. Moreover $\Lambda_7^{(k)} \subset \Lambda_0^{(k)} \subset \Lambda_7^{(k-1)\prime}$ [2, p. 566]. Five nesting steps at scale $k - 1$, converted to scale k, give $m \geq \frac{5}{L}(r_{k-1} - 1) - 2 \geq \frac{5}{2} r_k - 5$ at $L = 2$, since $r_{k-1} \geq r_k$. The terms of Prop. 2.6 localised outside $\Lambda_7^{(k)}$ are estimated with the domain of U0 and are covered by U0. (iii) U2: $x_v \in B^k(\Lambda_0^{(k)}) \subset B^j(\Lambda_7^{(j)})$, by iterating the inclusions of [2, p. 566]. Five nesting steps at scale j give $m \geq 5r_j - 7$. (iv) U0/U0′: here $\Omega_{\mathrm{out}} = B^k(\Lambda_2^{(k-1)\prime})$, and the nesting of the layers of [2, p. 566] gives $m \geq 2r_k - 4$ for every vertex type. Its dependence on the support of $\tilde{B}^{(k)}$ is removed by $\operatorname{supp} \tilde{B}^{(k)} \subset \operatorname{supp} \theta_{k+1}$ [2, (2.89), p. 575].

Arithmetic. By (5.11) and $r \geq R$ we have $r/4 \geq \ell_{\bar{N}} + 8M_b + \frac{25}{4}$, and $s_b \leq 2M_b$. Each case of (D.1) reduces to one inequality:

- U0: $2r - 4 > \frac{7}{4} r + 2s_b + \ell_{\bar{N}}$ iff $\frac{r}{4} > \ell_{\bar{N}} + 2s_b + 4$, which follows from $\frac{r}{4} > \ell_{\bar{N}} + 4M_b + 4$.
- U1a: $\frac{21}{4} r > \ell_{\bar{N}} + 2s_b + 10$.
- U1b: $\frac{3}{4} r > \ell_{\bar{N}} + 2s_b + 5$.
- U2: $\frac{13}{4} r_j > \ell_{\bar{N}} + 2s_b + 7$.

All four follow from the bound on $r/4$. The form $\frac{5}{2}(r(t_{k-1}) - M_b - 1) - 4$ used in the proof of Corollary 5.10 is the U0 configuration $\Omega_1 = B^k(\Lambda_7^{(k-1)\prime}) \subset \Omega = B^k(\Lambda_2^{(k-1)\prime})$. It also satisfies (D.1): with $r(t_{k-1}) \geq r$, this reduces to $\frac{3}{4} r > \ell_{\bar{N}} + 2s_b + \frac{5}{2} M_b + \frac{13}{2}$, which follows from (5.11).

Cube. By Proposition 5.8(c) every $p \in \square$ satisfies $\operatorname{dist}(p, x_v) \leq \frac{3}{2} r + 2s_b + \ell_{\bar{N}}$. By (D.1), $\operatorname{dist}(p, \Omega_{\mathrm{out}}^c) > m - \frac{3}{2} r - 2s_b - \ell_{\bar{N}} > r/4$. □

Supply at the uses.

Lemma D.3 (supply of $\tilde{A}$-depth at the uses)**.** *Let* $L = 2$*. At each of the sites (P), (Q), (Z) of the proof of Theorem 5.2, with* r *the value of* $r(\cdot)$ *at the scale of the site,*

$$\operatorname{dist}(\operatorname{supp} \tilde{A}, \partial\Omega) > 2r - M_\theta - 2M_b - 5. \tag{D.2}$$

If (5.11) *holds and* $r \geq R$*, the right side exceeds* $\frac{7}{4} r + 2s_b + \ell_{\bar{N}} + 1$*, so (R4‴)* (5.10) *holds at every use. This is the last statement of Corollary 5.10. It also gives* $\operatorname{dist}(\operatorname{supp} \tilde{A}, \partial\Omega) \geq r$*, the depth needed at the changes of domain.*

Proof. (Q) Here $\Omega = B^k(\Lambda_0^{(k)})$ and $\tilde{A} = \tau\theta_{k+1} A'^{(k)}$ [2, p. 579]. By [2, p. 567], $\theta_{k+1} = 1$ on $B^k(\Lambda_2^{(k)})$ and varies on a slice of thickness $< M_\theta$ outside it. So $\operatorname{supp} \tilde{A}$ lies within M_θ of $B^k(\Lambda_2^{(k)})$. Two nesting steps give $\operatorname{dist}(B^k(\Lambda_2^{(k)}), \Omega^c) \geq 2r_k - 4$. With one more lattice unit for the bond, $\operatorname{dist}(\operatorname{supp} \tilde{A}, \partial\Omega) > 2r_k - M_\theta - 5$ under the infimum convention. Under any set-distance convention it is $> 2r_k - M_\theta - 2M_b - 5$, since two nesting steps cost at most $2M_b$ more.

(P) Here $\Omega_P = B^k(\Lambda_2^{(k-1)\prime}) \supset B^k(\Lambda_0^{(k)}) = \Omega_Q$, because $\Lambda_0^{(k)} \subset \Lambda_7^{(k-1)\prime} \subset \Lambda_2^{(k-1)\prime}$ [2, p. 566]. The field is $\tilde{A} = \theta_{k+1} A'^{(k)} + \tilde{B}^{(k)}$ [2, (2.93)]. Each of the three terms defining $\tilde{B}^{(k)}$ in [2, (2.89), p. 575] carries the factor θ_{k+1}, so $\operatorname{supp} \tilde{A} \subset \operatorname{supp} \theta_{k+1}$, the same set as at (Q). Since $\Omega_Q \subset \Omega_P$, the margin at (P) is at least the margin at (Q).

(Z) At scale j the margin is $\geq 5r_j - 8$, by the nesting argument of part (iii) of the proof of Lemma D.2 with one more lattice unit for the bond, and $\geq 5r_j - 8 - 7M_b$ under any convention. It exceeds $2r_j - M_\theta - 2M_b - 5$, because $3r_j \geq 3 + 5M_b$.

Arithmetic. We have $\frac{7}{4}r + 2s_b + \ell_{\bar{N}} + 1 < 2r - M_\theta - 2M_b - 5$ iff $\frac{r}{4} > \ell_{\bar{N}} + 2s_b + M_\theta + 2M_b + 6$. With $s_b \leq 2M_b$ and $M_\theta \leq LM_b$, the right side is at most $\ell_{\bar{N}} + (6 + L)M_b + 6 = \ell_{\bar{N}} + 8M_b + 6$. By (5.11) and $r \geq R$, $\frac{r}{4} \geq \ell_{\bar{N}} + 8M_b + \frac{25}{4}$. Finally, $2r - M_\theta - 2M_b - 5 \geq r$ iff $r \geq (2 + L)M_b + 5$, which also follows from (5.11).

Completeness of the list of sites. The sites (P), (Q), (Z) are all the places in the proof of Theorem 5.2 at which [3, Prop. 1] is applied with a field $\tilde{A}$; this is read off [2, pp. 583–594] and [3, pp. 412–445]. □

Remark D.4. The supply (D.2) is the margin Bałaban's own nesting provides at [2, p. 579] (Remark 5.11(iii)). The need, through P12, is only $\frac{3}{2}r + O(\ell_{\bar{N}} + M_b)$. The margin is therefore available, not needed.

The sketched places of Proposition 5.8(a).

Lemma D.5 (the sketched places of [3], Prop. 1)**.**

(i) *On pages 412, 414, 415 and 419–445 of* [3] *and pp. 610–611 of* [1]*, the places of the proof where the text calls an argument similar, omitted, standard, obvious, easy, the same as or done in the same way as another, repeated, analogous or straightforward, or leaves out details, are the places S1–S21 of Table 2, with the number 12 used by the printed step P12. That is twenty sketched places, S1–S11 and S13–S21, plus P12.*

(ii) *Besides these, pages 414–445 of* [3] *contain seven further sketched places:*
   (1) *the Ω_2 branch "[or $G_k(\Omega_2, \tilde{B})$]" of P12 (p. 433);*
   (2) *the identification, unwritten in* [3]*, of the torus propagators of the counterterms (pp. 418–419) with the $\eta_k \mathbb{Z}^d$ propagators of the classes, which the exact cancellations of* [3, (3.7), (3.19)] *and of pp. 444–445 need;*
   (3) *p. 433: "We repeat some of the operations done before ... ", i.e. S1 and S9 again on □, with B' in place of $\tilde{A}$;*
   (4) *pp. 430–431: "The same applies to the remaining definitions" and "Taking other functions F, or differentiating (2.24) to higher order in A, we can get all necessary Ward–Takahashi identities";*

(5)–(7) *three phrases of power counting and bookkeeping: p. 427 "Now it is easy to see that the remaining powers of $L^j\eta$ ... ", p. 428 "In the case $v = v'$ we make the same operations", and p. 433 "Considering each connected component of G_{ren} separately ... We can extract also all coupling constants".*

(iii) Depth of the seven places.
   - *(1) needs $\square \subset \Omega_2$, i.e. the need of P12 measured from Ω_2^c (Proposition 5.8(c) with Ω_2 in place of Ω). It has the same coefficient of r as P12 and is not a new item.*
   - *(2) is torus-size, not depth: the differences are torus-image terms (Lemma 5.22).*
   - *(3) takes place on the parallelepiped □, where* [1, p. 611] *imposes no restriction. Its coupling power uses only $|B'| \leq O(r_\square p)$, a size bound.*
   - *(4)–(7) are Gaussian lattice identities or bookkeeping on* [3, (2.14)]*, with no domain.*

   So none of the seven needs depth beyond that of P12 or $R_0 + O(1)$, and no sketched place needs depth in $\big(\frac{7}{4}r + O(\ell_{\bar{N}} + M_b),\ 2r - M_\theta - 2M_b - 5\big]$.

This is the list of sketched places and the depth claim of Proposition 5.8(a).

Parts (i) and (ii) are statements about the printed text of [3], not mathematical statements; the completeness of the list is what Proposition 5.8(a) uses.

Proof. Parts (i) and (ii) are read off the printed text. For each place of (ii) the depth claim of (iii) is read off the cited page, as follows.

- (1): the cube step is printed for $G_k(\Omega_2, \tilde{B})$ as well. By Proposition 5.8(c) with Ω_2 in place of Ω, it needs an old vertex at depth $> \frac{3}{2}r_\square + 2s_b + \ell_{\bar{N}}$ in Ω_2.
- (2): [3, (1.25)–(1.31), pp. 418–419] builds the counterterms from propagators on T_ε rescaled to T_η. After p. 433 the classes use $G_k(0)$ on $\eta_k\mathbb{Z}^d$. The difference consists of image terms of order $\mathrm{e}^{-\delta_0\cdot(\text{torus side})}$ and does not involve Ω.
- (3): on $\square$ the bounds of [1, Prop. 2.1] hold for all points ([1, p. 611]).
- (4)–(7): the Ward–Takahashi identities are identities of Gaussian measures with no domain, and the remaining phrases act on the power counting of [3, (2.14)].

Every object on pp. 433–445 is independent of the distance to Ω^c once $\square \subset \Omega$ (Lemma 5.9). The places (2)–(7) therefore need no depth, and (1) needs that of P12 with Ω_2 in place of Ω. $\square$

Remark D.6 (scope). (i) The renormalisation combinatorics S21 of [3, pp. 434–445], and the generalisation S6 ("easily seen" in print), are used as printed; they do not depend on the depth in Ω. (ii) Lemmas D.2 and D.3 are stated at $L = 2$; at $L = 5$ the margin at U1b is too small.

D.2. **The L^2 sandwich and the two-domain theorem.** We prove two bounds used in the proof of Proposition 5.8: the bound for the kernel [3, (1.16)] used in step S1 (Proposition D.13), and the bound for the two-variable mixed norm of a domain difference δG_k used in step S2 and in part (c) (Proposition D.17).

Setting and notation. We fix a scale $k \geq 1$ and work in scale-k units, as in Subsection 4.3. We write $\eta := \eta_k = L^{-k} \leq \frac{1}{2}$, $t := t_k = L^k\varepsilon$, $e_k := e(t_k)$, and $d = 3$.

- *Domains.* $\mathcal{X}$ is a union of big blocks of the rescaled torus T_η with Neumann conditions, or $\mathcal{X} = T_\eta$ itself. For real A we put $H_{\mathcal{X}}(A) := -\Delta^{\eta,N}_{A,\mathcal{X}} + m^2t^2 + a_kP_k(A)$ and $G_k(\mathcal{X}, A) := H_{\mathcal{X}}(A)^{-1}$. Here $P_k = Q_k^*Q_k$, and Q_k averages over unit blocks along contours inside the block, of length $\leq d$.
- *Kernels and derivatives.* Kernels are taken with respect to η^3, i.e. $(Yf)(x) = \sum_y \eta^3 Y(x,y)f(y)$. $D = D^\eta_{\tilde{B}}$ is the covariant forward difference and D^* its $\ell^2(\eta^3)$-adjoint, so that D^{b*} acts on the second variable. We put $\mathfrak{D}f := (f, D_1f, D_2f, D_3f)$.
- *Depth and neighbourhoods.* $\rho_{\mathcal{X}}(z) := \mathrm{dist}_\infty(z, \mathcal{X}^c)$ is the depth of z in $\mathcal{X}$ (the ℓ^∞ torus distance, which is 1-Lipschitz), and $U_s(Y)$ denotes the ℓ^∞-neighbourhood of radius s of a set Y.
- *Kernel classes.* For $0 < \kappa < 3$ and $\delta > 0$ let $K^\delta_\kappa(z) := \max(|z|, \eta)^{\kappa-3}\mathrm{e}^{-\delta|z|}$. Further, $K^\delta_3(z) := (1 + \log_+(1/\max(|z|,\eta)))\mathrm{e}^{-\delta|z|}$, and $K^\delta_\kappa(z) := \mathrm{e}^{-\delta|z|}$ for $\kappa > 3$.
- *Smallness parameters.* $\sigma := e_k(\|\tilde{A}\|_\infty + \|\partial^\eta\tilde{A}\|_\infty)$ and $\theta(A) := e_k\|\partial^\eta A\|_\infty$.

Here $\tilde{A}$ and $\tilde{B}$ are the real background fields of Subsection 4.4. Under the hypotheses of [3, p. 412] one has $\sigma \leq 2c_1e_kp(t)$, $\theta(\tilde{B}) \leq c_1e_kp(t)$ and $\theta(\tilde{A}+\tilde{B}) \leq 2c_1e_kp(t)$.

Norm. For a two-variable kernel Y, put $Y_{ab} := D^aYD^{b*}$ ($a, b \in \{0,1\}$, every direction), and for $0 < \alpha < 1$, $\beta \geq 0$ define

$$\mathfrak{N}_{\alpha,\beta}(Y) := \max_{a,b}\sup_{x,x'} \mathrm{e}^{\beta|x-x'|}\max\{\|Y_{ab}(x,x')\|,\, a\,H_x,\, b\,H_{x'},\, ab\,H_{xx'}\}.$$

Here $H_x := \sup_{0<|x-\tilde{x}|\leq 1}|x-\tilde{x}|^{-\alpha}\|U(\tilde{B}(\Gamma_{\tilde{x},x}))Y_{ab}(x,x') - Y_{ab}(\tilde{x},x')\|$; $H_{x'}$ is the same quotient in x', with the transport acting from the right; and $H_{xx'}$ is the mixed second difference divided by $|x-\tilde{x}|^\alpha|x'-\tilde{x}'|^\alpha$. $\|hYh'\|_{1,\alpha}$ denotes the same quantity at $\beta = 0$ for the kernel $h(x)Y(x,x')h'(x')$. This is [3, (1.32)] "extended in a natural way" to two variables, with the mixed component $D_xD^*_{x'}$,

which is the reading under which [3, (2.5)] is used. Quotients with $|x-\tilde{x}|>1$ are bounded by twice the corresponding sup and need no separate estimate.

Lemma D.7 (inputs)**.** *There are constants $\gamma_0,\beta_*,K_W,c_K,c_K(\alpha),\delta_K,\vartheta_R>0$, depending only on d,a,M_b,L (and on α for $c_K(\alpha)$), such that the following hold whenever A is real with $\theta(A)\le\vartheta_R$ and $0\le m^2t^2\le c$:*

- (H0) *$H_{\mathcal{X}}(A)\ge\gamma_0$ on every $\mathcal{X}$ as above* [4, (1.8), p. 573].
- (W) *For every 1-Lipschitz ρ and $|\beta|\le\beta_*$, the operators G, DG, GD^* and DGD^* with $G=G_k(\mathcal{X},A)$, conjugated as $Y\mapsto \mathrm{e}^{\beta\rho}Y\mathrm{e}^{-\beta\rho}$, have $\ell^2(\eta^3)$ operator norms $\le 2/\gamma_0$ (for G) and $\le K_W$ (the other three). These bounds hold uniformly in η, k, ε, the shape of $\mathcal{X}$ (wrapping allowed) and the boundary condition.*
- (K) *For x,w with $\rho_{\mathcal{X}}(x),\rho_{\mathcal{X}}(w)\ge R_0$ (and Hölder partners $|x-\tilde{x}|\le 1$ with the same property), where R_0 is the constant of Proposition 4.5:*
  - (K0) *$\|G(x,w)\|\le c_K K_2^{\delta_K}(x-w)$;*
  - (K1) *$\|D_xG(x,w)\|\le c_K K_1^{\delta_K}(x-w)$;*
  - (K2) *$\|U(A(\Gamma_{\tilde{x},x}))D_xG(x,w)-D_{\tilde{x}}G(\tilde{x},w)\|\le c_K(\alpha)|x-\tilde{x}|^{\alpha}\big(K_{1-\alpha}^{\delta_K}(x-w)+K_{1-\alpha}^{\delta_K}(\tilde{x}-w)\big)$.*

  On $\mathcal{X}=T_\eta$ the restriction $\rho_{\mathcal{X}}\ge R_0$ is void.
- (S) *For real A, $G_k(\mathcal{X},A)$ is real symmetric, so $(GD^*)(x,w)=(D_wG(w,x))^T$.*

(W) is the weighted form of [4, Cor. 2.3, (2.30), p. 580], which bounds $|\langle f,DGD^*f'\rangle|$ by $c_0\mathrm{e}^{-\delta_0\operatorname{dist}(\operatorname{supp}f,\operatorname{supp}f')}\|f\|_2\|f'\|_2$ without restriction on the supports. (K) is [3, (2.10)–(2.11), p. 426], summed over the slices of [3, (2.6)]; the derivation of (2.10)–(2.11) is part (b), step S3, of the proof of Proposition 5.8. (H0) is cited, (S) is immediate, and the proofs of (W) and of the summed form (K) are Lemmas D.24 and D.26 of Subsection D.3.

The kernel (1.16): an L^2 sandwich. Let $G:=G_k(\mathcal{X},\tilde{B})$ and $G':=G_k(\mathcal{X},\tilde{A}+\tilde{B})$. Put $V:=G^{-1}-G'^{-1}$; this is Bałaban's $V_k(\tilde{A},\tilde{B})$, the bracket of [1, (3.44), p. 619]. For $n,n'\ge 0$, put $m:=n+n'$ and

$$K_{n,n'}:=[GV]^nG'[VG]^{n'}=L_0VL_1V\cdots VL_m,\qquad L_n:=G',\ L_i:=G\ (i\ne n),$$

which is the kernel [3, (1.16), p. 414]. Our hypothesis is

$$(\mathrm{R}_\vartheta)\qquad \theta(\tilde{B}),\ \theta(\tilde{A}+\tilde{B})\le\vartheta_R,\qquad \sigma\le 1,\qquad 0\le m^2t^2\le c.$$

Under the hypotheses of [3, p. 412], (R_ϑ) follows from the regularity condition $2c_1e_kp(t)\le\min\{\vartheta_R,1\}$ on the coupling.

Lemma D.8 (the vertex operator)**.** *Put $\phi_\mu(x):=\eta^{-1}(1-U_{x,\mu}(\tilde{A})^{-1})$, $\psi_\mu:=\phi_\mu^*\phi_\mu$ and $\Pi:=P_k(\tilde{A}+\tilde{B})-P_k(\tilde{B})$. Then:*

- (i) *$V=-\big[\sum_\mu(D_\mu^*\phi_\mu+\phi_\mu^*D_\mu+\psi_\mu)+a_k\Pi\big]=-\mathfrak{D}^*\mathfrak{E}\mathfrak{D}$, where $\mathfrak{E}$ is the 4×4 matrix of multiplication and block operators with entries $\mathfrak{E}_{00}=\sum_\mu\psi_\mu+a_k\Pi$, $\mathfrak{E}_{0\mu}=\phi_\mu^*$, $\mathfrak{E}_{\mu0}=\phi_\mu$, all other entries 0.*
- (ii) *(Right-pushed form.) $V=-(R_0+\sum_\mu R_{1,\mu}D_\mu)=:-\mathfrak{r}\mathfrak{D}$. Here $(R_{1,\mu}h)(x):=\phi_\mu(x)^*h(x)-\phi_\mu(x-\eta e_\mu)U_{x-\eta e_\mu,\mu}(\tilde{B})^{-1}h(x-\eta e_\mu)$, and $R_0:=\sum_\mu(\psi_\mu-M_\mu)+a_k\Pi$, with M_μ the multiplication by $(\partial_\mu^\eta\phi_\mu)(x-\eta e_\mu)$.*
- (iii) *(Left-pushed form.) $V=V^*=-\mathfrak{D}^*\mathfrak{r}^*$.*
- (iv) *For $\sigma\le1$, in $\ell^p(\eta^3)$, $p\in\{1,2,\infty\}$: $\|\mathfrak{r}\|,\|\mathfrak{E}\|\le c_V\sigma$ with $c_V:=4d+a(2d+d^2)$. Every kernel has range $\le\sqrt{3}+\eta$, so for 1-Lipschitz ρ and $|\beta|\le1$ the conjugates by $\mathrm{e}^{\pm\beta\rho}$ are bounded by $\mathrm{e}^{\beta(\sqrt{3}+\eta)}c_V\sigma$. Moreover $V(y,\cdot)=0$ unless $y\in U_{\sqrt{3}+\eta}(\operatorname{supp}\tilde{A})$.*

On $\mathcal{X}=\Omega$ with Neumann conditions, (i)–(iv) hold as stated provided $\operatorname{dist}(\operatorname{supp}\tilde{A},\Omega^c)>\sqrt{3}+2\eta$, *since then every bond and every block on which ϕ or Π is non-zero lies in Ω.*

Proof. All transporters are functions of the matrix q ($q^2=-1$), so they commute, and $U(\tilde A+\tilde B)=U(\tilde A)U(\tilde B)$. We use $D_{A,\mu}f(x)=\eta^{-1}(U_{x,\mu}(A)f(x+\eta e_\mu)-f(x))$ [4, (1.3)]. Then

$$D_{\tilde A+\tilde B,\mu}f = U_\mu(\tilde A)\big(D_\mu f+\phi_\mu f\big).$$

Since U is pointwise orthogonal, summing over the bonds of $\mathcal{X}$ (the same bonds for both backgrounds) gives

$$-\Delta_{\tilde A+\tilde B}=\sum_\mu (D_\mu+\phi_\mu)^*(D_\mu+\phi_\mu)=-\Delta_{\tilde B}+\sum_\mu (D_\mu^*\phi_\mu+\phi_\mu^*D_\mu+\psi_\mu).$$

Since $Q_k(\tilde A+\tilde B)=Q_k(\tilde B)+F_2$ with $F_2:=Q_k(\tilde A+\tilde B)-Q_k(\tilde B)$, we get $P_k(\tilde A+\tilde B)-P_k(\tilde B)=\Pi$. The terms m^2t^2 cancel, and (i) follows. For (ii), use $D_\mu^*h(x)=\eta^{-1}(U^{-1}_{x-\eta e_\mu}h(x-\eta e_\mu)-h(x))$ and $[\phi,U]=0$:

$$D_\mu^*(\phi_\mu f)(x)=-\phi_\mu(x-\eta e_\mu)U^{-1}_{x-\eta e_\mu}(D_\mu f)(x-\eta e_\mu)-(\partial_\mu^\eta\phi_\mu)(x-\eta e_\mu)f(x).$$

Inserting this into (i) gives (ii). (iii) holds because both inverses are real symmetric for real backgrounds. For (iv): $|\mathrm{e}^{q\vartheta}-1|\le|\vartheta|$, hence $|\phi_\mu|\le\sigma$, $|\psi_\mu|\le\sigma^2$ and $|\partial_\mu^\eta\phi_\mu|\le e_k|\partial_\mu^\eta\tilde A_\mu|\le\sigma$. Also $|U(\tilde A(\Gamma))-1|\le d\sigma$ for the contours, $\|Q_k\|\le1$, and so $\|\Pi\|\le(2d+d^2)\sigma$. This gives $\|R_0\|\le(2d+a(2d+d^2))\sigma$, $\sum_\mu\|R_{1,\mu}\|\le2d\sigma$ and $\|\mathfrak{E}\|\le c_V\sigma$. A kernel of range r changes by at most $\mathrm{e}^{|\beta|r}$ under conjugation by $\mathrm{e}^{\beta\rho}$. The support statement holds because the Laplacian part of V lives on bonds where $\tilde A\neq0$, and Π on unit blocks meeting $\operatorname{supp}\tilde A$. □

Lemma D.9 (convolution and smearing)**.** *(a) For $\kappa_1,\kappa_2\in(0,3)$ and $0<\delta'<\delta\le1$, on T_η or $\eta\mathbb{Z}^3$:*

$$\sum_w \eta^3 K^\delta_{\kappa_1}(z-w)K^\delta_{\kappa_2}(w)\le c(\kappa_1,\kappa_2)\big(1+(\delta-\delta')^{-3}\big)K^{\delta'}_{\kappa_1+\kappa_2}(z),$$

with the conventions above for $\kappa_1+\kappa_2\ge3$.

(b) For $\delta''<\delta$: $K^\delta_\kappa\le c(1+(\delta-\delta'')^{-1})K^{\delta''}_{\kappa'}$ whenever $\kappa\ge\kappa'$.

(c) Let R be a multiplication, a one-bond shift, or a bounded kernel supported on single unit blocks, with $|R(z,z')|\le r_R$. Then $\big(|R|\,K^\delta_\kappa(\cdot-y)\big)(z)\le c(\kappa)\mathrm{e}^{2\sqrt3\delta}r_RK^\delta_\kappa(z-y)$.

Proof. For (a), use $\mathrm{e}^{-\delta|z-w|-\delta|w|}\le\mathrm{e}^{-\delta'|z|}\mathrm{e}^{-(\delta-\delta')(|z-w|+|w|)}$, put $r:=\max(|z|,\eta)$, and split the sum into three regions: $|w|\le r/2$; $|z-w|\le r/2$; and both distances $\ge r/2$. Each region is bounded with the lattice-sum bound $\sum_{|w|\le R}\eta^3\max(|w|,\eta)^{\kappa-3}\le c(\kappa)R^\kappa$. The tail $|w|>2r$ gives $c\int_{2r}^\infty s^{\kappa_1+\kappa_2-4}\mathrm{e}^{-(\delta-\delta')s}\,ds$. (b): for $r\le1$ the smaller exponent gives the larger kernel, and for $r>1$ the power is absorbed by $\mathrm{e}^{-(\delta-\delta'')r}$. (c): for a one-bond shift, $\max(|z'-y|,\eta)\ge\frac12\max(|z-y|,\eta)$. For a block kernel, $|z'-y|\ge|z-y|/2$ if $|z-y|\ge2\sqrt3$, while the block sum is $\le c(\kappa)$ if $|z-y|<2\sqrt3$. □

Lemma D.10 (end factors)**.** *Let $L_0,L_1,L_2\in\{G,G'\}$ and $a\in\{0,1\}$, and set $E^{(2)}_a:=D^aL_0VL_1$, $E^{(3)}_a:=D^aL_0VL_1VL_2$, $\delta':=\delta_K/2$, $0\le\beta\le\delta_K/4$. Assume that (K) applies at every pair of points used below (automatic on T_η). Then:*

- (i) *$\|E^{(2)}_a(x,y)\|\le c_E\sigma K^{\delta'}_2(x-y)$ and $\|E^{(3)}_a(x,y)\|\le c'_E\sigma^2K^{\delta'}_3(x-y)$.*
- (ii) *For $a=1$ and $|x-\tilde x|\le1$, the transported difference of $E^{(2)}_1$ is bounded by $c_E(\alpha)\sigma|x-\tilde x|^\alpha(K^{\delta'}_{2-\alpha}(x-y)+K^{\delta'}_{2-\alpha}(\tilde x-y))$, and that of $E^{(3)}_1$ by the same expression with $c'_E(\alpha)\sigma^2$ and $K_{3-\alpha}$.*
- (iii) *Put $J_\kappa:=(\sum_y\eta^3K^{\delta'}_\kappa(y)^2\mathrm{e}^{2\beta|y|})^{1/2}$. Then $\|\mathrm{e}^{\beta|\cdot-x|}E^{(2)}_a(x,\cdot)\|_2\le c_EJ_2\sigma$, the transported difference in (ii) has weighted ℓ^2 norm $\le2\mathrm{e}^\beta c_E(\alpha)J_{2-\alpha}\sigma|x-\tilde x|^\alpha$, and the same holds for $E^{(3)}$ with J_3, $J_{3-\alpha}$ and σ^2.*

$J_\kappa<\infty$ uniformly in η and in the torus side if and only if $\kappa>\frac32$. Hence $J_{2-\alpha}<\infty$ if and only if $\alpha<\frac12$, and $J_{3-\alpha}<\infty$ for every $\alpha<1$. The constants depend only on d,a,M_b,L,α,β.

Proof. (i) By Lemma D.8(ii), $E_a^{(2)} = -(D^a L_0)R_0 L_1 - \sum_\mu (D^a L_0)R_{1,\mu}(D_\mu L_1)$. Thus every line carries exactly one derivative: L_0 the external one, L_1 the vertex derivative. By (K), $\|D^a L_0(x,z)\| \le c_K K_{2-a}^{\delta_K}$ and $\|\mathfrak{D} L_1(z,y)\| \le c_K(K_2+K_1)^{\delta_K}$. Lemma D.9(c) moves R (with $r_R \le c_V\sigma$) onto the kernel, and Lemma D.9(a),(b) give $K_1 * K_1 \to K_2$. For $E_a^{(3)}$ both vertices are right-pushed, so $E_a^{(3)} = (D^a L_0)\mathfrak{r}(\mathfrak{D}L_1)\mathfrak{r}(\mathfrak{D}L_2)$ with one derivative per line, and $K_1 * K_1 * K_1 \to K_3$.

(ii) The difference acts on the first factor only. Use (K2), then $K_{1-\alpha} * K_1 \to K_{2-\alpha}$, respectively $K_{1-\alpha} * K_1 * K_1 \to K_{3-\alpha}$.

(iii) Square and sum, using $|y-x| \le |y-\tilde{x}|+1$ for the $\tilde{x}$-term. Near 0, $\sum_y \eta^3 \max(|y|,\eta)^{2\kappa-6}$ is bounded uniformly in η if and only if $2\kappa > 3$. The torus sum is dominated by the $\mathbb{Z}^3$-sum because $2\delta' - 2\beta \ge \delta_K/2$. □

Lemma D.11 (localisation of two-variable kernel bounds)**.** *Let $\eta \le \frac{1}{2}$, $0 < \alpha < 1$, $\beta \ge 0$ and $M \ge 0$. Let D be the covariant forward difference, $(D_\mu f)(x) = \eta^{-1}(U_\mu(x)f(x+\eta e_\mu) - f(x))$, with bond transporters $U_\mu(x)$. In the Hölder quotients, let the transport $U(\Gamma_{\tilde{x},x})$ be the product of the same bond transporters along a shortest lattice contour $\Gamma_{\tilde{x},x}$ from $\tilde{x}$ to x, as in the definition of $\mathfrak{N}_{\alpha,\beta}$ in Lemma E.117. For a kernel Y put $Y_{ab} := D^a Y D^{b*}$, where $a,b \in \{0,1\}$ and all directions are allowed. Take lattice points $y,\tilde{y},y',\tilde{y}'$ with $0 < |y-\tilde{y}| \le 1$ and $0 < |y'-\tilde{y}'| \le 1$. Let $H_y(\tilde{y})$, $H_{y'}(\tilde{y}')$ and $H_{yy'}(\tilde{y},\tilde{y}')$ be the α-Hölder quotients of Y_{ab} at these partners: in the first variable, in the second variable, and jointly. The joint quotient is the mixed second difference divided by $|y-\tilde{y}|^\alpha |y'-\tilde{y}'|^\alpha$. Let $\mathcal{P}$ be a set of lattice points such that, for all such $y,y',\tilde{y},\tilde{y}' \in \mathcal{P}$, all a,b and all directions,*

$$\mathrm{e}^{\beta|y-y'|}\max\{\|Y_{ab}(y,y')\|,\, aH_y(\tilde{y}),\, bH_{y'}(\tilde{y}'),\, abH_{yy'}(\tilde{y},\tilde{y}')\} \le M. \tag{D.3}$$

Let $0 \le h,h' \le 1$ satisfy $|\partial_\mu^\eta h| \le c_h$ and $|\partial_\nu^\eta \partial_\mu^\eta h| \le c_h$ for all μ,ν, and likewise for h'. Assume that every lattice point at ℓ^∞-distance $\le 1+\eta$ from $\operatorname{supp} h \cup \operatorname{supp} h'$ lies in $\mathcal{P}$. Then

$$\|hYh'\|_{1,\alpha} \le \mathrm{e}^{3\beta}(1+3\sqrt{3}c_h)^2 M\,\mathrm{e}^{-\beta\,\mathrm{dist}(\operatorname{supp}h,\operatorname{supp}h')} \le \mathrm{e}^{5\beta}(1+6c_h)^2 M\,\mathrm{e}^{-\beta\,\mathrm{dist}(\operatorname{supp}h,\operatorname{supp}h')}.$$

Here $\|Z\|_{1,\alpha}$ is the following maximum, taken over a,b, the directions, all lattice points x,x' and all partners at distance ≤ 1: the maximum of $\|Z_{ab}(x,x')\|$ and of the quotients aH_x, $bH_{x'}$, $abH_{xx'}$ of Z. If the quotients over $|x-\tilde{x}| > 1$ and $|x'-\tilde{x}'| > 1$ are included, the bound holds with an additional factor 4. If, as in [3, (1.32)]*, the nine terms of the norm (value, derivative and quotient in each variable) are moreover added instead of maximised, it holds with an additional factor 16.*

Proof. Step 1 (Leibniz identity). Since h is real, $D_\mu(hf)(x) = h(x+\eta e_\mu)(D_\mu f)(x) + (\partial_\mu^\eta h)(x)f(x)$. Put $Y^\dagger(x',x) := Y(x,x')^*$. Then $(ZD_\mu^*)(x,x') = ((D_\mu Z^\dagger)(x',x))^*$. So the components and quotients of hYh' in x' are those of $h'Y^\dagger h$ in its first variable, and everything below applies to both variables.

Step 2 (differences of the coefficients). Let $|x-\tilde{x}| \le 1$, and let $\Gamma = \Gamma_{\tilde{x},x} = (y_0 = \tilde{x}, \dots, y_n = x)$. Its length is $\eta n = |x-\tilde{x}|_1 \le \sqrt{3}|x-\tilde{x}|$. It is monotone in each coordinate, so its points lie in the box spanned by $\tilde{x}$ and x. If $|\partial_\nu^\eta \varphi| \le c$ for all ν, summing along Γ gives $|\varphi(x)-\varphi(\tilde{x})| \le \sqrt{3}c|x-\tilde{x}|$. We apply this to $\varphi = h(\cdot+\eta e_\mu)$ and to $\varphi = \partial_\mu^\eta h$.

Step 3 (telescoping). Let U_i be the transporter of the bond $\{y_{i-1},y_i\}$ from y_i to y_{i-1}, so that $U(\Gamma) = U_1 \cdots U_n$. For a fibre-valued W,

$$U(\Gamma)W(x) - W(\tilde{x}) = \sum_{i=1}^{n} U_1 \cdots U_{i-1}\big[U_i W(y_i) - W(y_{i-1})\big].$$

The bracket equals $\eta(D_\nu W)(y_{i-1})$ if $y_i = y_{i-1} + \eta e_\nu$, and $-\eta U_i(D_\nu W)(y_i)$ if $y_i = y_{i-1} - \eta e_\nu$. The transporters are unitary, so $\|U(\Gamma)W(x) - W(\tilde{x})\| \le |x-\tilde{x}|_1 \max_{z\in\Gamma,\nu}\|(D_\nu W)(z)\|$.

Step 4 (one variable). Suppress the second variable. Then $(hY)_0 = hY_0$ and $(hY)_1(x) = F_1(x)Y_1(x) + F_2(x)Y_0(x)$, where $F_1 := h(\cdot+\eta e_\mu)$ and $F_2 := \partial_\mu^\eta h$. Hence $\|(hY)_1(x)\| \le (1+$

$c_h)\max_a\|Y_a(x)\|$. With $U := U(\Gamma_{\tilde{x},x})$,

$$U(hY)_1(x) - (hY)_1(\tilde{x}) = F_1(x)\big[UY_1(x) - Y_1(\tilde{x})\big] + \big(F_1(x) - F_1(\tilde{x})\big)Y_1(\tilde{x}) \\ + F_2(x)\big[UY_0(x) - Y_0(\tilde{x})\big] + \big(F_2(x) - F_2(\tilde{x})\big)Y_0(\tilde{x}).$$

- The first term is at most $|x-\tilde{x}|^\alpha$ times the quotient of Y_1 at $(x,\tilde{x})$.
- By Step 2, the second and fourth terms are at most $\sqrt{3}c_h|x-\tilde{x}|$ times a component of Y at $\tilde{x}$.
- By Step 3, the third term is at most $\sqrt{3}c_h|x-\tilde{x}|\max_{z\in\Gamma,\nu}\|(D_\nu Y_0)(z)\|$, and $D_\nu Y_0$ is a component Y_1.

Since $|x-\tilde{x}| \le |x-\tilde{x}|^\alpha$, the quotient of $(hY)_1$ at $(x,\tilde{x})$ is at most $(1+3\sqrt{3}c_h)$ times the largest component or quotient of Y at points of $\{x,\tilde{x}\}\cup\Gamma$, with partners in the same set.

Step 5 (two variables, points, exponential). Each expansion in Step 4 is a sum of terms. Each term is a coefficient depending only on the expanded variable, times a linear functional of Y in that variable: a value, a transported difference, or a telescoped sum of $D_\nu Y$. The expansions in x and in x' act on different variables, and the transports act from opposite sides, so the two expansions commute. This gives the following bounds, in each case times the largest of the quantities in (D.3) at pairs of evaluated points with evaluated partners:

- every component of hYh': $(1+c_h)^2$;
- every one-variable quotient: $(1+3\sqrt{3}c_h)(1+c_h)$;
- the mixed quotient: $(1+3\sqrt{3}c_h)^2$.

A term vanishes when its coefficient vanishes. For an x-coefficient to be non-zero, x or $\tilde{x}$ must lie within distance η of $\operatorname{supp} h$ along a single coordinate direction. Hence all evaluated points lie within ℓ^∞- and Euclidean distance $1+\eta$ of $\operatorname{supp} h$, and similarly for x' and h'. So these points lie in $\mathcal{P}$, and (D.3) applies. Moreover $\mathrm{e}^{-\beta|y-y'|} \le \mathrm{e}^{2(1+\eta)\beta}\mathrm{e}^{-\beta\operatorname{dist}(\operatorname{supp} h,\operatorname{supp} h')} \le \mathrm{e}^{3\beta}\mathrm{e}^{-\beta\operatorname{dist}(\operatorname{supp} h,\operatorname{supp} h')}$. Since $3\sqrt{3}\le 6$, both bounds follow.

Step 6 (other forms of the norm). A quotient over a distance > 1 is at most twice the supremum of the quantity whose difference is taken. Since $1+c_h \le 1+3\sqrt{3}c_h$, this costs a factor 2 per variable. In each variable, the sum of the value, the derivative and the quotient over all distances is at most $1+(1+c_h)+\max\{1+3\sqrt{3}c_h, 2+2c_h\} \le 4(1+3\sqrt{3}c_h)$. This gives the factor 16 for the summed form. □

The bound on second differences enters only through $|F_2(x)-F_2(\tilde{x})|$ in Step 4. With first differences alone, this term is only bounded by $2c_h|x-\tilde{x}|^{-\alpha}\|Y_0(\tilde{x})\|$, which is of order $\eta^{-\alpha}$ at $|x-\tilde{x}|=\eta$.

Lemma D.12 (the cut-offs of [3], (1.33))**.** *For $\nu\neq\mu$, the cut-off h of Lemma E.103(iii) satisfies*

$$|\partial_\mu^\eta h| \le \tfrac{15}{4}, \qquad |\partial_\mu^\eta\partial_\mu^\eta h| \le \tfrac{40}{\sqrt{3}}, \qquad |\partial_\nu^\eta\partial_\mu^\eta h| \le \big(\tfrac{15}{4}\big)^2.$$

Hence the hypothesis of Lemma D.11 holds with $c_h := 40/\sqrt{3}$, uniformly in η and $\square(v)$.

Proof. In Lemma E.103(iii), $h(x) = \prod_\mu g_\mu(x_\mu)$ with $g_\mu(u) = \chi(\operatorname{dist}(u, I_\mu))$. Here I_μ is the μ-th side interval of $\square(v)$, and $\chi(t) = 1$ for $t \le \frac{1}{2}$, $\chi(t) = 1 - s(2t-1)$ for $\frac{1}{2} \le t \le 1$, and $\chi(t) = 0$ for $t \ge 1$, where $s(t) = 10t^3 - 15t^4 + 6t^5$.

Since $\chi = 1$ on $[0, \frac{1}{2}]$, $g_\mu = 1$ on the $\frac{1}{2}$-neighbourhood of I_μ. Outside I_μ the distance is affine with slope ±1. Hence $g_\mu \in C^2(\mathbb{R})$, with $|g_\mu'| \le \sup|\chi'| = 2\max s' = \frac{15}{4}$ and $|g_\mu''| \le \sup|\chi''| = 4\max|s''| = \frac{40}{\sqrt{3}}$. Here $s' = 30t^2(1-t)^2$ is maximal at $t = \frac{1}{2}$, and $|s''| = 60t(1-t)|1-2t|$ is maximal at $t = \frac{3\pm\sqrt{3}}{6}$, where it equals $10/\sqrt{3}$.

Since $0 \le g_\mu \le 1$, we get $|\partial_\mu h| \le \frac{15}{4}$, $|\partial_\mu^2 h| \le \frac{40}{\sqrt{3}}$ and $|\partial_\nu \partial_\mu h| \le (\frac{15}{4})^2$. Lattice differences are averages of derivatives, including the case $\nu = \mu$:

$$\partial_\mu^\eta h(x) = \eta^{-1} \int_0^\eta \partial_\mu h(x + s e_\mu)\, ds, \qquad \partial_\nu^\eta \partial_\mu^\eta h(x) = \eta^{-2} \int_0^\eta \int_0^\eta \partial_\nu \partial_\mu h(x + s e_\mu + t e_\nu)\, ds\, dt.$$

Finally $\frac{15}{4} \le (\frac{15}{4})^2 \le \frac{40}{\sqrt{3}}$. □

Proposition D.13 (the bound for [3], (1.16), in the mixed norm)**.** *Assume (R_ϑ), and let $0 < \beta \le \beta_U := \min\{\beta_*, \delta_K/4\}$. Suppose that either $m = n + n' \ge 4$ and $0 < \alpha < \frac{1}{2}$, or $m \ge 6$ and $0 < \alpha < 1$; in particular, $\alpha = \alpha_0 = \frac{1}{2}$ requires $m \ge 6$.*

(a) *If $\mathcal{X} = T_\eta$, then $\mathfrak{N}_{\alpha,\beta}(K_{n,n'}) \le C_U(\alpha)(C_V \sigma)^m$. Here $C_V := \mathrm{e}^{\beta(\sqrt{3}+1)} c_V \max\{c_W, 1\}$ and $c_W := (1+\sigma)^2 \big(2\gamma_0^{-1} + ((1+d)^2 - 1) K_W\big)$.*

(b) *Let $\mathcal{X} = \Omega = B^k(\Omega^{(k)})$ with Neumann conditions, and assume $\operatorname{dist}(\operatorname{supp} \tilde{A}, \Omega^c) \ge R_0 + 2\sqrt{3} + 1$. Then the same bound holds when the supremum in $\mathfrak{N}_{\alpha,\beta}$ is restricted as follows: the points x, x', together with their bond partners, have depth $\ge R_0$, and the quotients are taken only at partners $\tilde{x}, \tilde{x}'$ with the same property.*

(c) *Let $0 \le h, h' \le 1$ with $|\partial_\mu^\eta h|, |\partial_\nu^\eta \partial_\mu^\eta h| \le c_h$ for all μ, ν, and likewise for h'. In case (b), assume in addition that $\operatorname{supp} h \cup \operatorname{supp} h'$ has depth $\ge R_0 + 3$. Then*

$$\|h K_{n,n'} h'\|_{1,\alpha} \le \mathrm{e}^{5\beta} (1 + 6c_h)^2 C_U(\alpha) (2 c_1 C_V)^{n+n'} (e_k p(t))^{n+n'} \mathrm{e}^{-\beta \operatorname{dist}(\operatorname{supp} h, \operatorname{supp} h')}.$$

This is the bound [3, (2.5)] *for the operator (1.16), with Balaban's rate δ_0 taken $\le \beta_U$.*

$C_U(\alpha)$ depends only on $d, a, M_b, L, \gamma_0, K_W, \alpha, \beta$. The bound is uniform in η (hence in k and ε), in the torus side, in $0 \le m^2 t^2 \le c$ and in the backgrounds within (R_ϑ). It is also uniform in the parameters (u, M) of Theorem 5.2, since no photon line occurs in (1.16).

The depth hypothesis of (b) follows from $R > R_0$ and $R \ge 12\sqrt{d}$ (conditions (R.1) and (R.8) of Table 5): these give $2r(t) \ge R_0 + 2\sqrt{3} + 1$ under (R4), and under (R4‴) of Corollary 5.10 the hypothesis holds with room. In the construction $n = n' = 6$, so $m = 12$, and the Hölder exponent of [3, (2.5)] is $\alpha_0 = \frac{1}{2}$. This is the L^2 sandwich used in part (b), step S1, of the proof of Proposition 5.8.

Proof. Step 1 (why ends are needed). $K_{n,n'}$ has m vertices and $m+1$ lines, and every vertex is first order on each side (Lemma D.8(i)). So $D^a K_{n,n'} D^{b*}$ carries $m+2$ derivatives on $m+1$ lines, and some line carries two. That line is placed in a middle factor, which enters only through its weighted ℓ^2 norm (W). A pointwise bound of $D_x G D_y^*$ is never used.

Step 2 (factorisation, two-line ends, $m \ge 4$). Write $Y := D^a K_{n,n'} D^{b*} = E_L \mathcal{N} E_R$ with

$$E_L := D^a L_0 V L_1, \qquad \mathcal{N} := V L_2 V \cdots V L_{m-2} V, \qquad E_R := (D^b L_m V L_{m-1})^*.$$

Put the first vertex of $\mathcal{N}$ in right-pushed form, the last in left-pushed form, and the others in symmetric form (Lemma D.8(i)–(iii)):

$$\mathcal{N} = (-1)^{m-2} \mathfrak{r}(\mathfrak{D} L_2 \mathfrak{D}^*) \mathfrak{E} \cdots \mathfrak{E} (\mathfrak{D} L_{m-2} \mathfrak{D}^*) \mathfrak{r}^*.$$

For $L = G'$, the identity $D_{\tilde{B},\mu} = U_\mu(\tilde{A})^{-1} D_{\tilde{A}+\tilde{B},\mu} - \phi_\mu$ costs a factor $1 + \sigma$ on each side. With (W) and Lemma D.8(iv), for $|\beta| \le \beta_U$,

$$\|\mathrm{e}^{\mp\beta\rho} \mathcal{N} \mathrm{e}^{\pm\beta\rho}\| \le (\mathrm{e}^{\beta(\sqrt{3}+1)} c_V \sigma)^{m-2} c_W^{m-3}.$$

By (S) and $V^* = V$, Lemma D.10 applies to E_R. In E_L and E_R each line carries exactly one derivative. The pointwise bounds (K) used in Lemma D.10 are those of Lemma D.26, whose proof sums the slice bounds by Lemma D.25; that lemma holds at every distance $z \ge 0$.

Step 3 (sandwich). Fix x, x' and put $\rho := |\cdot - x|$. Then

$$Y(x,x') = \big\langle \mathrm{e}^{\beta\rho} E_L(x,\cdot)^*,\ (\mathrm{e}^{-\beta\rho}\mathcal{N}\mathrm{e}^{\beta\rho})\,\mathrm{e}^{-\beta\rho} E_R(\cdot,x')\big\rangle_{\ell^2(\eta^3)}.$$

Cauchy–Schwarz, $\mathrm{e}^{-\beta|y'-x|} \le \mathrm{e}^{-\beta|x-x'|}\mathrm{e}^{\beta|y'-x'|}$ and Lemma D.10(iii) give

$$\mathrm{e}^{\beta|x-x'|}\|Y(x,x')\| \le (c_E J_2\sigma)^2 (\mathrm{e}^{\beta(\sqrt{3}+1)} c_V\sigma)^{m-2} c_W^{m-3}.$$

Y is linear in $E_L(x,\cdot)$ and in $E_R(\cdot,x')$ separately. So $H_x, H_{x'}, H_{xx'}$ obey the same bound with $c_E J_2$ replaced by $2\mathrm{e}^{\beta} c_E(\alpha) J_{2-\alpha}$, which is finite for $\alpha < \frac12$. Hence $\mathfrak{N}_{\alpha,\beta}(K_{n,n'}) \le C_U^{(2)}(\alpha)\sigma^2(\mathrm{e}^{\beta(\sqrt{3}+1)}c_V\sigma)^{m-2}c_W^{m-3}$, with $C_U^{(2)}(\alpha) := \max\{c_E J_2, 2\mathrm{e}^{\beta} c_E(\alpha) J_{2-\alpha}\}^2$.

Step 4 (three-line ends, $m \ge 6$, any $\alpha < 1$). Repeat Steps 2–3 with $E_L := E_a^{(3)}$, $\mathcal{N} := VL_3V\cdots VL_{m-3}V$ and $E_R := (D^b L_m V L_{m-1} V L_{m-2})^*$. Now $J_3, J_{3-\alpha} < \infty$ for every $\alpha < 1$, and $\mathfrak{N}_{\alpha,\beta} \le C_U^{(3)}(\alpha)\sigma^4(\mathrm{e}^{\beta(\sqrt{3}+1)}c_V\sigma)^{m-4}c_W^{m-5}$, with $C_U^{(3)}(\alpha) := \max\{c_E' J_3, 2\mathrm{e}^{\beta} c_E'(\alpha) J_{3-\alpha}\}^2$. Since $c_V \ge 1$, both bounds are $\le C_U(\alpha)(C_V\sigma)^m$, with $C_U := C_U^{(2)} c_V^{-2}$, respectively $C_U^{(3)} c_V^{-4}$. This proves (a).

Step 5 (Neumann domains). On Ω, (H0) and (W) hold with the Neumann condition (Lemma D.7). Lemma D.8(ii) holds on Ω, because every bond of a vertex lies in Ω. Since $V(y,\cdot) = 0$ unless $y \in U_{\sqrt{3}+\eta}(\operatorname{supp}\tilde{A})$, every pointwise use of (K) in Lemma D.10 is at one of two kinds of points:

- points of $U_{2\sqrt{3}+1}(\operatorname{supp}\tilde{A})$, which have depth $\ge R_0$ by hypothesis;
- the points x, x', their bond partners, and, for the quotients, $\tilde{x}, \tilde{x}'$; these have depth $\ge R_0$ by the restriction in (b).

Steps 2–4 apply verbatim, which proves (b).

Step 6 (localisation, (c)). Apply Lemma D.11 with $M := C_U(\alpha)(C_V\sigma)^m$. In case (a) take $\mathcal{P} := T_\eta$. In case (b) take $\mathcal{P} := \{z : \text{depth of } z \ge R_0 + 1 + \eta\}$. Points of this $\mathcal{P}$ and their bond partners have depth $\ge R_0$, so (b) gives (D.3) on $\mathcal{P}$. Since $\eta \le \frac12$, every point within ℓ^∞-distance $1+\eta$ of a set of depth $\ge R_0 + 3$ has depth $\ge R_0 + 2 - \eta \ge R_0 + 1 + \eta$, so it lies in $\mathcal{P}$. Finally use $\sigma \le 2c_1 e_k p(t)$. The bound [3, (2.5)] is used with Bałaban's tree rate $\delta_0 \le \beta_U$, which only shrinks a free constant. □

Remark D.14. (i) At $m = 0$ the mixed kernel is not uniform in η. With $g := \eta^{-3}1_{\{x\}}$ and $\Lambda := 4d\eta^{-2} + m^2t^2 + a \ge \|G'^{-1}\|$, one has

$$(D_\mu G' D_\mu^*)(x,x) = \langle D_\mu^* g, G' D_\mu^* g\rangle \ge \Lambda^{-1}\|D_\mu^* g\|_2^2 = 2\Lambda^{-1}\eta^{-5} \ge \eta^{-3}/(6 + O(\eta^2))$$

per direction. This is why [3, p. 414] needs "n, n' sufficiently large". (ii) The operator form of the bound, in the norms of [1, (2.24)–(2.25)], which is the literal statement of [3, p. 414], holds for every $m \ge 0$. It is the argument of [4, (2.33)] applied to one term of the series. It is not needed for (2.5).

Domain differences in the mixed norm: the two-domain theorem. Let $\mathcal{X}_{\text{in}} \subset \mathcal{X}_{\text{out}}$ be unions of big blocks of T_η, where $\mathcal{X}_{\text{out}}$ may be T_η itself. Let A be real with $\theta(A) \le \vartheta_R$ on $\mathcal{X}_{\text{out}}$, and $0 \le m^2t^2 \le c$. Put $G_{\text{in}} := G_k(\mathcal{X}_{\text{in}}, A)$, $G_{\text{out}} := G_k(\mathcal{X}_{\text{out}}, A)$, $H_{\text{in}}, H_{\text{out}}$ the corresponding operators, $\delta := G_{\text{out}} - G_{\text{in}}$ (signs are irrelevant for the bounds), and $\rho := \rho_{\mathcal{X}_{\text{in}}}$. Then $\rho_{\mathcal{X}_{\text{out}}} \ge \rho$.

Cut-offs. Fix $f \in C^\infty(\mathbb{R})$ with $0 \le f \le 1$, $f = 0$ on $(-\infty, 0]$ and $f = 1$ on $[1,\infty)$, and $\varphi \in C_c^\infty([-\frac14, \frac14]^3)$ with $\varphi \ge 0$ and $\int\varphi = 1$. Let $\tilde\rho := \rho^{\text{cont}} * \varphi$, where ρ^{cont} is the continuum ℓ^∞ distance to the closed complement of $\mathcal{X}_{\text{in}}$. Then $|\tilde\rho - \rho| \le \frac14 + \eta$ on lattice points. For $s \in \mathbb{R}$ put $\chi_s := f(\tilde\rho - s)$. Then $\chi_s = 0$ on $\{\rho \le s - \frac14 - \eta\}$, $\chi_s = 1$ on $\{\rho \ge s + \frac54 + \eta\}$, and the first and second lattice differences of χ_s are bounded by absolute constants, uniformly in η and $\mathcal{X}_{\text{in}}$. We take

$$\chi := \chi_{R_0+3}, \qquad \psi := \chi_{R_0+7}.$$

Let $X_{\rm in} \subset X_{\rm out}$ be unions of big blocks of T_η, $\eta = L^{-k} \le \frac12$; here $X_{\rm out}$ may be T_η, while $X_{\rm in} \neq T_\eta$. Let A be real with $\theta(A) \le \vartheta_R$, and let $0 \le m^2t^2 \le c$. Put $H_{\rm in/out} := H_{X_{\rm in/out}}(A)$, $G_{\rm in/out} := H^{-1}_{\rm in/out}$, $\delta := G_{\rm out} - G_{\rm in}$ and $\rho(z) := \mathrm{dist}_\infty(z, X_{\rm in}^c)$; then $\rho_{X_{\rm out}} \ge \rho$. For a matrix-valued function F of one variable put $\|F\|_2 := (\sum_z \eta^3 \|F(z)\|^2)^{1/2}$; for a row of four such functions, the four norms are added. The quotients $H_x, H_{x'}, H_{xx'}$ are those of $\mathfrak{N}_{\alpha,\beta}$, with transport along a shortest lattice contour.

Lemma D.15 (collars)**.** *Fix $f \in C^\infty(\mathbb{R})$ with $0 \le f \le 1$, $f = 0$ on $(-\infty, 0]$ and $f = 1$ on $[1, \infty)$. Fix $\varphi \in C_c^\infty([-\frac14, \frac14]^3)$ with $\varphi \ge 0$ and $\int \varphi = 1$. For a unit block b with lowest corner y_b let $\hat b := y_b + [0,1)^3$, and put $\hat X_{\rm in} := \bigcup_{b \subset X_{\rm in}} \hat b$. Let $\rho^{\rm c}(u)$ be the continuum ℓ^∞ distance from u to the complement of $\hat X_{\rm in}$. Put $\tilde\rho := \rho^{\rm c} * \varphi$, and let $\chi_s := f(\tilde\rho - s)$ on lattice points. Let $c_1' := \|f'\|_\infty$ and $c_2' := \|f''\|_\infty + \|f'\|_\infty \max_\nu \|\partial_\nu \varphi\|_{L^1}$. Then:*

(i) *$\tilde\rho$ is 1-Lipschitz for $|\cdot|_\infty$, and $\rho - \frac14 - \eta \le \tilde\rho \le \rho + \frac14$ at lattice points;*
(ii) *$\chi_s = 0$ on $\{\rho \le s - \frac14\}$ and $\chi_s = 1$ on $\{\rho \ge s + \frac54 + \eta\}$;*
(iii) *$|\partial^\eta_\mu \chi_s| \le c_1'$ and $|\partial^\eta_\mu \partial^\eta_\nu \chi_s| \le c_2'$, uniformly in η, s and $X_{\rm in}$;*
(iv) *if $\chi_s(u) \neq \chi_s(u')$ and $|u - u'|_\infty \le \ell$, then $\tilde\rho(u), \tilde\rho(u') \in (s - \ell, s + 1 + \ell)$.*

Proof. (i) The cubes $\hat b$ tile the torus, so a lattice point outside $X_{\rm in}$ lies outside $\hat X_{\rm in}$; hence $\rho^{\rm c} \le \rho$. Conversely, $\rho^{\rm c}(z)$ is attained at a point of a closed cube $\overline{\hat b'}$ with $b' \not\subset X_{\rm in}$. The lattice points of b' are η-dense in $\overline{\hat b'}$ for $|\cdot|_\infty$, so $\rho(z) \le \rho^{\rm c}(z) + \eta$. Finally, $\rho^{\rm c}$ is 1-Lipschitz, and mollification by φ preserves this and moves values by at most $\frac14$.

(ii) follows from (i).

(iii) By the mean value theorem, the lattice differences are bounded by the suprema of the continuum derivatives. Here $|\partial_\mu \tilde\rho| \le 1$ and $\partial_\mu \partial_\nu \tilde\rho = (\partial_\mu \rho^{\rm c}) * \partial_\nu \varphi$.

(iv) If $\tilde\rho(u) \le s - \ell$, then $\tilde\rho(u') \le s$, and both values of χ_s are 0. If $\tilde\rho(u) \ge s + 1 + \ell$, both values are 1. The same holds with u and u' exchanged. □

Lemma D.16 (commutators)**.** *Let $s \ge 1$, $\chi := \chi_s$ and $\phi_\mu := \partial^\eta_\mu \chi$.*

(i) *$H_{\rm out}(\chi g) = H_{\rm in}(\chi g)$ for every g on $X_{\rm in}$ (extended by zero).*
(ii) *$C_\chi := H_{\rm out}\chi - \chi H_{\rm in} = [H_{\rm in}, \chi] = -\mathfrak{r}_\chi \mathfrak{D} = \mathfrak{D}^* \mathfrak{r}_\chi^*$. Here $\mathfrak{r}_\chi = (\mathfrak{r}_0, \dots, \mathfrak{r}_3)$ is given by*

$$\mathfrak{r}_0 := \sum_\mu (\partial^\eta_\mu \phi_\mu)(\cdot - \eta e_\mu) - a_k [P_k(A), \chi], \qquad (\mathfrak{r}_\mu g)(x) := \phi_\mu(x) g(x) + \phi_\mu(x - \eta e_\mu) U_{x-\eta e_\mu, \mu}(A)^{-1} g(x - \eta e_\mu).$$

(iii) *$\sum_{i=0}^3 \sup_z \sum_w \eta^3 \|\mathfrak{r}_i(w,z)\| \le c_{\mathfrak{r}} := 6c_1' + 3c_2' + ac_1'$.*
(iv) *If $\mathfrak{r}_i(w,z) \neq 0$, then $|w - z| \le \sqrt3$ and $w, z \in S_s := \{s - \frac54 - \eta < \rho < s + \frac94 + 2\eta\}$.*

Proof. (i) If $\chi(y) \neq 0$, then $\rho(y) > s - \frac14 > \eta$, so every bond at y lies in $X_{\rm in}$. The unit block of y also lies in $X_{\rm in}$, since $X_{\rm in}$ is a union of unit blocks. A bond of $X_{\rm out}$ that is not a bond of $X_{\rm in}$ has both endpoints at depth $\le \eta$, hence outside $\mathrm{supp}\,\chi$. Since P_k acts inside unit blocks, the two operators agree on χg.

(ii) The Leibniz rules $D_\mu(\chi f) = \chi(\cdot + \eta e_\mu) D_\mu f + \phi_\mu f$ and $D^*_\mu(\chi(\cdot + \eta e_\mu) g) = \chi D^*_\mu g - \phi_\mu g$ give $[D^*_\mu D_\mu, \chi] = D^*_\mu \phi_\mu - \phi_\mu D_\mu$. Also $[m^2t^2, \chi] = 0$. By [4, (2.32), p. 581],

$$D^*_\mu(\phi_\mu f)(x) = -\phi_\mu(x - \eta e_\mu) U^{-1}_{x-\eta e_\mu, \mu}(D_\mu f)(x - \eta e_\mu) - (\partial^\eta_\mu \phi_\mu)(x - \eta e_\mu) f(x).$$

Collecting terms gives $C_\chi = -\mathfrak{r}_\chi \mathfrak{D}$. Since $H_{\rm in}$ is self-adjoint and χ is real, $C^*_\chi = -C_\chi$, which gives the second form.

(iii) For each μ, the two parts of $\mathfrak{r}_\mu$ have column sums $\le c_1'$ each, which gives $6c_1'$ in total. The multiplication part of $\mathfrak{r}_0$ is bounded by $3c_2'$. For the last part, $[P_k, \chi](w,z) = P_k(w,z)(\chi(z) - \chi(w))$. Here $\sum_w \eta^3 \|P_k(w,z)\| \le 1$, $|\chi(z) - \chi(w)| \le c_1'$ inside a unit block, and $a_k \le a$.

(iv) There are two cases.

- For $\mathfrak{r}_\mu$ and the first part of $\mathfrak{r}_0$: here $z \in \{w, w-\eta e_\mu\}$, and χ differs at two adjacent points of $\{w-\eta e_\mu, w, w+\eta e_\mu\}$. Lemma D.15(iv) with $\ell = \eta$ gives $\tilde\rho(w), \tilde\rho(z) \in (s-3\eta, s+1+3\eta)$.
- For $[P_k, \chi]$: here $|w-z|_\infty \le 1-\eta$ and $\chi(w) \ne \chi(z)$, which gives $\tilde\rho(w), \tilde\rho(z) \in (s-1+\eta, s+2-\eta)$.

Since $\eta \le \frac12$, both 3η and $1-\eta$ are $\le 1+\eta$, and Lemma D.15(i) gives $w, z \in S_s$. Finally $|w-z| \le \sqrt{3}\max(\eta, 1-\eta) \le \sqrt{3}$. □

We take $\chi := \chi_{R_0+3}$ and $\psi := \chi_{R_0+13/2}$, with $S_\chi := S_{R_0+3}$ and $S_\psi := S_{R_0+13/2}$. We also put $\mathcal{P}_D := \{z : \rho(z) \ge R_0 + 10 + 2\eta\}$. Since $10 + 2\eta \le 11$, $\mathcal{P}_D \supset \{\rho \ge R_0 + 11\}$ for every $k \ge 1$.

Proposition D.17 (the two-domain theorem)**.** *Let $0 < \alpha < 1$, $0 < \beta \le \beta_D := \min\{\beta_*, \delta_K/4\}$, $a, b \in \{0, 1\}$ and $Y := D^a \delta D^{b*}$. Then for $x, x' \in \mathcal{P}_D$, with partners $\tilde x, \tilde x' \in \mathcal{P}_D$,*

$$\mathrm{e}^{\beta|x-x'|}\max\{\|Y(x,x')\|, aH_x, bH_{x'}, abH_{xx'}\} \le C_D(\alpha)\exp\Big(-\tfrac{\delta_K}{2}\big[(\rho(x)-R_0-7)+(\rho(x')-R_0-10)\big]\Big).$$

The constants are

$$C_D(\alpha) := c'_W c_{\mathrm{end}}(\alpha)^2, \qquad c'_W := 2/\gamma_0 + 3K_W,$$
$$c_{\mathrm{end}}(\alpha) := c_{\mathfrak{r}}\mathrm{e}^{\delta_K(\sqrt{3}+1)}\max\{c_K, 2c_K(\alpha)\}c_3^{1/2}, \qquad c_3 := (\tfrac12 + 4\sqrt{3}/\delta_K)^3.$$

They depend only on $d, a, M_b, L, \gamma_0, K_W, \alpha$ and on f, φ. They do not depend on η, k, ε, the side of the torus, X_{in}, X_{out}, A, $m^2t^2 \in [0, c]$ or R_0.

The proof uses only (H0), (W), (K) and (S) of Lemma D.7, in the forms of Lemmas D.24 and D.26.

Proof. Step 1 (collars). Lemma D.16(iv) gives

$$S_\chi \subset \{R_0 + \tfrac74 - \eta < \rho < R_0 + \tfrac{21}{4} + 2\eta\}, \qquad S_\psi \subset \{R_0 + \tfrac{21}{4} - \eta < \rho < R_0 + \tfrac{35}{4} + 2\eta\}.$$

We use the following consequences.

- $\psi = 0$ on $\{\rho \le R_0 + \frac{25}{4}\} \supset S_\chi$, since $\frac{21}{4} + 2\eta \le \frac{25}{4}$ for $\eta \le \frac12$.
- $S_\chi \cup S_\psi \subset \{\rho > R_0 + \frac54\}$.
- $\chi = 1$ on $\{\rho \ge R_0 + \frac{17}{4} + \eta\}$ and $\psi = 1$ on $\{\rho \ge R_0 + \frac{31}{4} + \eta\}$. Both sets contain $\mathcal{P}_D$ and all bond partners of points of $\mathcal{P}_D$.
- Since ρ is 1-Lipschitz, every $z \in \mathcal{P}_D$ satisfies

$$\mathrm{dist}_\infty(z, S_\chi) > \max\{\rho(z) - R_0 - 7, \tfrac{19}{4}\}, \qquad \mathrm{dist}_\infty(z, S_\psi) > \max\{\rho(z) - R_0 - 10, \tfrac54\}.$$

Step 2 (representation). We have $G_{\mathrm{out}}C_\chi G_{\mathrm{in}} = \chi G_{\mathrm{in}} - G_{\mathrm{out}}\chi$, so $\delta(y, y') = -(G_{\mathrm{out}}C_\chi G_{\mathrm{in}})(y, y')$ when $\chi(y) = \chi(y') = 1$. Likewise $G_{\mathrm{in}}C_\psi G_{\mathrm{in}} = \psi G_{\mathrm{in}} - G_{\mathrm{in}}\psi$, so $G_{\mathrm{in}}(z, y') = -(G_{\mathrm{in}}C_\psi G_{\mathrm{in}})(z, y')$ when $\psi(z) = 0$ and $\psi(y') = 1$. In the first identity only points $z \in S_\chi \subset \{\psi = 0\}$ occur. Inserting the second identity into the first and using Lemma D.16(ii) gives

$$\delta = -G_{\mathrm{out}}\mathfrak{r}_\chi(\mathfrak{D}G_{\mathrm{in}}\mathfrak{D}^*)\mathfrak{r}_\psi^* G_{\mathrm{in}}$$

at all points of $\mathcal{P}_D$ and their bond partners. By Step 1, these are the only points at which D^a and D^{b*} evaluate δ. Hence

$$Y(x, x') = -\langle E_L(x, \cdot), \mathfrak{M}E_R(\cdot, x')\rangle, \qquad E_L := D^a G_{\mathrm{out}}\mathfrak{r}_\chi, \quad \mathfrak{M} := \mathfrak{D}G_{\mathrm{in}}\mathfrak{D}^*, \quad E_R := \mathfrak{r}_\psi^* G_{\mathrm{in}} D^{b*}.$$

Step 3 (left end). $E_L(x, z) = \sum_w \eta^3 D_x^a G_{\mathrm{out}}(x, w)\mathfrak{r}_\chi(w, z)$, with $w, z \in S_\chi$ and $|w - z| \le \sqrt{3}$. Here (K) applies on X_{out} at (x, w), and $|x-w| \ge 1$. So $K_{2-a}^{\delta_K}(x-w) \le \mathrm{e}^{-\delta_K|x-w|} \le \mathrm{e}^{\delta_K\sqrt{3}}\mathrm{e}^{-\delta_K|x-z|}$, and

$$\|E_L(x, z)\| \le c_{\mathfrak{r}}c_K\mathrm{e}^{\delta_K\sqrt{3}}\mathrm{e}^{-\delta_K|x-z|}1_{S_\chi}(z).$$

Since $2(\delta_K - \beta) \ge \frac32\delta_K$,

$$\|\mathrm{e}^{\beta|\cdot-x|}E_L(x, \cdot)\|_2 \le c_{\mathfrak{r}}c_K\mathrm{e}^{\delta_K\sqrt{3}}\mathrm{e}^{-\frac{\delta_K}{2}\mathrm{dist}(x, S_\chi)}\Big(\sum_z \eta^3\mathrm{e}^{-\frac{\delta_K}{2}|z-x|}\Big)^{1/2}.$$

To bound the lattice sum by c_3, pass on the torus to shortest lifts and use $|z| \geq |z|_1/\sqrt{3}$ and $\coth(c/2) \leq 1 + 2/c$. With Step 1 this gives $\leq c_{\mathrm{end}}(\alpha)\mathrm{e}^{-\frac{\delta_K}{2}(\rho(x)-R_0-7)}$.

For $a = 1$ and a partner $\tilde{x} \in \mathcal{P}_D$, use (K2) and $|\tilde{x} - w| \geq 1$. The transported difference then has kernel $\leq 2c_K(\alpha)\mathrm{e}^{\delta_K}|x - \tilde{x}|^\alpha \mathrm{e}^{-\delta_K|x-w|}$, and the same computation applies.

Step 4 (right end). By (S), $(G_{\mathrm{in}}D^{b*})(v, x') = (D^b_{x'}G_{\mathrm{in}}(x', v))^T$ with $v \in S_\psi$, and (K) applies on X_{in}. As in Step 3,

$$\|\mathrm{e}^{\beta|\cdot-x'|}E_R(\cdot, x')\|_2 \leq c_{\mathrm{end}}(\alpha)\mathrm{e}^{-\frac{\delta_K}{2}(\rho(x')-R_0-10)}.$$

The same bound, times $|x' - \tilde{x}'|^\alpha$, holds for the transported difference.

Step 5 (middle). By Lemma D.24 on X_{in} with weight $|\cdot - x|$, $\|\mathrm{e}^{-\beta|\cdot-x|}\mathfrak{M}\mathrm{e}^{\beta|\cdot-x|}\| \leq c'_W$. Together with $\mathrm{e}^{-\beta|u-x|} \leq \mathrm{e}^{-\beta|x-x'|}\mathrm{e}^{\beta|u-x'|}$ and Cauchy–Schwarz, this gives the bound for $\|Y(x, x')\|$. Y is linear in $E_L(x, \cdot)$ and in $E_R(\cdot, x')$ separately, so the quotients follow by replacing one or both ends by their differences. No kernel is evaluated at points less than 1 apart, and the only doubly differentiated line enters through its operator norm. □

Corollary D.18 (localised form)**.** *Let $0 \leq h, h' \leq 1$ with $|\partial^\eta_\mu h|, |\partial^\eta_\nu\partial^\eta_\mu h| \leq c_h$ for all μ, ν, and likewise for h'. Assume $\rho \geq R_0 + 11 + 3\eta$ on $\operatorname{supp} h \cup \operatorname{supp} h'$. This holds if $\rho \geq R_0 + 12$ there when $k \geq 2$, and if $\rho \geq R_0 + 13$ there when $k = 1$. Then, at any mutual distance of the supports (overlapping supports included),*

$$\|h\,\delta\,h'\|_{1,\alpha} \leq C_{\mathrm{loc}}\,\mathrm{e}^{-\beta\,\mathrm{dist}(\operatorname{supp} h, \operatorname{supp} h')}, \qquad C_{\mathrm{loc}} := \mathrm{e}^{5\beta}(1 + 6c_h)^2 C_D(\alpha).$$

Proof. Apply Lemma D.11 with $\mathcal{P} := \mathcal{P}_D$ and $M := C_D(\alpha)$. On $\mathcal{P}_D$ both brackets in the exponent of Proposition D.17 are nonnegative, so the exponential factor is ≤ 1 and (D.3) holds. Since ρ is 1-Lipschitz for ℓ^∞, every point within ℓ^∞-distance $1 + \eta$ of the supports has $\rho \geq R_0 + 10 + 2\eta$, so it lies in $\mathcal{P}_D$. □

Lemma D.19 (nesting)**.** *Let $\rho_0 \geq 1$, and let $\Lambda_{i_0} \supset \Lambda_{i_0+1} \supset \cdots$ be sets of points of a cubic lattice of spacing ≤ 1. Assume that every point at ℓ^∞-distance $\leq \rho_0$ from Λ_i^c lies in Λ_{i+1}^c. Then $\mathrm{dist}_\infty(\Lambda_{i_0+m}, \Lambda_{i_0}^c) > m\rho_0 - (m-1)$ for every $m \geq 1$.*

Proof. For $m = 1$ this is the hypothesis. For the induction step, let $x \in \Lambda_{i_0+m}$ and $z \in \Lambda_{i_0}^c$; then $|x - z|_\infty > \rho_0$. A king-move lattice geodesic from z to x advances $|\cdot - z|_\infty$ by one lattice step (≤ 1) at a time. So it contains a point w with $\rho_0 - 1 \leq |w - z|_\infty \leq \rho_0$ and $|x - w|_\infty = |x - z|_\infty - |w - z|_\infty$. Then $w \in \Lambda_{i_0+1}^c$. The induction hypothesis for the chain starting at $i_0 + 1$ gives $|x - w|_\infty > (m-1)\rho_0 - (m-2)$, hence $|x - z|_\infty > m\rho_0 - (m-1)$. □

Remark D.20 (uses). (I1) $X_{\mathrm{out}} = \Omega$, $X_{\mathrm{in}} = \Omega_2 = B^k(\Lambda_2^{(k)})$, $A = \tilde{B}$: this is step S2 of Proposition 5.8(b). These lines occur only in terms whose vertices lie in $B^k(\Lambda_7^{(k)})$ [2, p. 579; Prop. 2.6, p. 580]. Use Lemma D.19 with $\rho_0 = r(t_k)$ and [2, (2.8), p. 558], with one block for the passage to $B^k(\cdot)$. Then these vertices have depth $\geq 5r(t_k) - 5$ in Ω_2. The localisation cubes $\square(v)$ have side ≤ 2 [3, p. 420], and the supports of the localisation functions lie within 4 of $B^k(\Lambda_7^{(k)})$. So the supports have depth $\geq 5r(t_k) - 9 \geq 5R - 9$. This is $\geq R_0 + 13$ when $R > R_0$ [2, (2.7)] and $R \geq \frac{11}{2}$; both hold under conditions (R.1) and (R.8) of Table 5. Corollary D.18 therefore applies at every $k \geq 1$.

(I2) $X_{\mathrm{out}} = \Omega$, $X_{\mathrm{in}} = \square$, $A = \tilde{B}$, given $\square \subset \Omega$ (Proposition 5.8(c)). The vertex localisations lie at depth $\geq r_\square - 2$ in $\square$. So $r_\square \geq R_0 + 13 + 3\eta$ suffices. This follows from $r_\square \geq R_0 + 15$, which is implied by $r \geq 3R_0 + 15$.

(I3) $X_{\mathrm{out}} = T_\eta$, $X_{\mathrm{in}} = \square$, $A = 0$: this is the line $\delta G_k(\square, \eta_k\mathbb{Z}^3, 0)$ of [3, p. 433] on the torus. The margins are as in (I2).

On the printed factor: [3, (2.5)] prints the factor $\mathrm{e}^{-\delta_0\,\mathrm{dist}(\Omega_2, \partial\Omega)}$, and the two-point decay multiplies both alternatives. Proposition D.17 gives this factor when the distance is read as the distance

of the vertex localisations to Ω_2^c; this is the factor of [4, (1.12), p. 573] for the inner domain. Read literally, as a distance between sets, the factor is not implied: for localisations near $\partial\Omega_2$ but deep in Ω, δ is of order one. The uses (I1)–(I3) need only Corollary D.18. Bałaban's rate δ_0 is taken $\le \beta_D$.

Remark D.21 (the printed factor; rates). (i) [3, (2.5)] prints, for $\delta G_k(\Omega, \Omega_2, \tilde{B})$, the bound $O(\mathrm{e}^{-\delta_0 \operatorname{dist}(\Omega_2, \partial\Omega)}) \cdot \mathrm{e}^{-\delta_0 \operatorname{dist}(\operatorname{supp} h, \operatorname{supp} h')}$; the two-point factor multiplies both alternatives of (2.5). Proposition D.17 matches this factor only in the reading "distance of the vertex localisations to Ω_2^c" (the factor of [1, (2.26)]). In the literal set-distance reading it is not implied, and it is false in general: localisations near $\partial\Omega_2$ but deep in Ω give a δG of order one. No step of the proof of Theorem 5.2 uses more than an $O(1)$ bound together with the two-point decay [3, p. 432], and both are proved.

(ii) The rates β_U and β_D enter [3, (2.13)] as the rate of a line. One takes Bałaban's $\delta_0 \le \min\{\beta_U, \beta_D\}$, a free shrink of a constant; the conditions that use δ_0, such as $\delta_0 R \ge 4$, are imposed after it.

Remark D.22. Two cases are not treated, and neither is used: the remainder of the B'-expansion on the cube $\square$ of [3, p. 433], which concerns Neumann faces and is a different object; and the Hölder exponents $\frac{1}{2} \le \alpha < 1$ at $m \in \{4, 5\}$ in Proposition D.13.

D.3. **The inputs (W) and (K).** *Conventions.* $\mathcal{X}$ is a union of big blocks of T_η with Neumann conditions, or $\mathcal{X} = T_\eta$. $\mathcal{B}_\mathcal{X}$ is the bond set: bonds with both ends in $\mathcal{X}$, or all bonds of T_η. Functions on sites and on bonds carry the measure η^3. For a bond $b = (b_-, b_+)$ in direction μ, $(D\phi)(b) = \eta^{-1}(U_b\phi(b_+) - \phi(b_-))$, with U_b orthogonal on $\mathbb{R}^2$. Let ρ be real and 1-Lipschitz for the Euclidean torus distance; every function that is 1-Lipschitz for the ℓ^∞ torus distance qualifies. On sites put $Y^\beta := \mathrm{e}^{\beta\rho} Y \mathrm{e}^{-\beta\rho}$. A bond function is weighted at b_-, i.e. by $\mathrm{e}^{\beta\rho(b_-)}$. Then $|\rho(b_+) - \rho(b_-)| \le \eta$ on every bond, and $|\rho(x) - \rho(x')| \le \sqrt{3}$ inside a unit block. Write $H_\mathcal{X}(A) = D^*D + m^2t^2 + a_k P_k(A)$.

Lemma D.23 (real-part Combes–Thomas bound)**.** *Let $Y = D^*D + v + aP$ on $\ell^2(\mathcal{X}, \eta^3; \mathbb{R}^2)$, where the bond set and the transporters U_b are arbitrary, $v \ge 0$ is a multiplication operator, $0 \le a$, and $P \ge 0$ satisfies $P(x, x') = 0$ for $|x - x'| > \sqrt{3}$ and $\sup_x \sum_{x'} \eta^3 |P(x, x')| \le 1$. Assume $Y \ge \gamma > 0$. Let $|\beta| \le 1$ with*

$$c(\beta) := 3\beta^2 \cosh\beta + a\big(\mathrm{e}^{\sqrt{3}|\beta|} - 1\big) \le \gamma/2.$$

Then for every F the vector $w := \mathrm{e}^{\beta\rho} Y^{-1} \mathrm{e}^{-\beta\rho} F$ satisfies $\|w\| \le 2\gamma^{-1}\|F\|$ and $\|Dw\|^2 + \langle w, vw\rangle \le 4\gamma^{-1}\|F\|^2$. This holds uniformly in η, the domain, the boundary condition, U and v.

Proof. Negative β is the case $-\rho$, so let $\beta \ge 0$. The vector w solves $Y^\rho w = F$ with $Y^\rho := \mathrm{e}^{\beta\rho} Y \mathrm{e}^{-\beta\rho}$.

Bond terms. Fix a bond b, put $t := \beta(\rho(b_+) - \rho(b_-))$, so $|t| \le \beta\eta$, and put $z := \bar{w}(b_-) U_b w(b_+)$. The contribution of b to $\operatorname{Re}\langle w, \mathrm{e}^{\beta\rho} D^* D \mathrm{e}^{-\beta\rho} w\rangle$ is

$$\eta^1 \big[|w(b_+)|^2 + |w(b_-)|^2 - 2\cosh t \operatorname{Re} z\big] = q_b(w) - 2(\cosh t - 1)\eta \operatorname{Re} z.$$

The odd part $(\sinh t)$ is anti-Hermitian and drops out of the real part. Since $\cosh t - 1 \le \frac{1}{2}\beta^2\eta^2 \cosh\beta$, $2|\operatorname{Re} z| \le |w(b_+)|^2 + |w(b_-)|^2$, and each site lies in at most 6 bonds, the bond part is $\ge \|Dw\|^2 - 3\beta^2 \cosh\beta \|w\|^2$.

Other terms. v commutes with $\mathrm{e}^{\beta\rho}$. The kernel of $P^\rho - P$ is $(\mathrm{e}^{\beta(\rho(x) - \rho(x'))} - 1)P(x, x')$, so by Schur's test $\|P^\rho - P\| \le \mathrm{e}^{\sqrt{3}\beta} - 1$. Hence

$$\operatorname{Re}\langle w, Y^\rho w\rangle \ge \langle w, Yw\rangle - c(\beta)\|w\|^2 \ge \tfrac{\gamma}{2}\|w\|^2.$$

Conclusion. Since $\operatorname{Re}\langle w, Y^\rho w\rangle = \operatorname{Re}\langle w, F\rangle \le \|w\|\|F\|$, the first bound follows. Then $\|Dw\|^2 + \langle w, vw\rangle \le \langle w, Yw\rangle \le \|w\|\|F\| + \frac{\gamma}{2}\|w\|^2 \le 4\gamma^{-1}\|F\|^2$. $\square$

Lemma D.24 (input (W): weighted ℓ^2 bounds)**.** *Let A be real with $H_{\mathcal{X}}(A) - m^2t^2 \geq \gamma_0$ (that is, (H0)), let $m^2t^2 \geq 0$, and let $G := G_k(\mathcal{X}, A)$. Fix $0 < \beta_* \leq 1$ with $c(\beta_*) \leq \gamma_0/4$ (the c of Lemma D.23 at $a = a_k$), and put $c' := \sqrt{3}\,\beta_* \mathrm{e}^{\beta_*}$. Then for every 1-Lipschitz ρ and $|\beta| \leq \beta_*$:*

$$\|G^\beta\| \leq \frac{2}{\gamma_0}, \qquad \|(DG)^\beta\|,\ \|(GD^*)^\beta\|,\ \|(DGD^*)^\beta\| \leq K_W := \max\Big\{\frac{2}{\gamma_0^{1/2}} + \frac{2c'}{\gamma_0},\ \frac{1+c'^2}{\min\{\frac{1}{2}, \frac{\gamma_0}{4}\}}\Big\}.$$

These hold uniformly in η, k, ε, $m^2 \geq 0$, the shape of $\mathcal{X}$ (wrapping allowed) and the boundary condition. Consequently the block operator $\mathfrak{D}G\mathfrak{D}^*$ satisfies $\|(\mathfrak{D}G\mathfrak{D}^*)^\beta\| \leq 2\gamma_0^{-1} + 3K_W$.

The only input is (H0) [4, (1.8), p. 573]. The factor $\sqrt{3}$ in c' accounts for the three directions of the vector difference D; per direction the constant is e^{β_*}.

Proof. (a) The bound on G^β. Apply Lemma D.23 to $Y = H_{\mathcal{X}}(A)$, with $v = m^2t^2$, $P = P_k(A)$, $a = a_k$ and $\gamma = \gamma_0$. This is admissible because $c(\beta) \leq \gamma_0/4$. The operator $G^\beta = (H_{\mathcal{X}}^\beta)^{-1}$ exists because $\operatorname{Re} H_{\mathcal{X}}^\beta \geq \gamma_0/2$. The lemma gives $\|G^\beta\| \leq 2/\gamma_0$ and $\|DG^\beta\| \leq 2\gamma_0^{-1/2}$.

(b) The bound on $(DG)^\beta$. With the bond weight at b_-,

$$(DG)^\beta = D^\beta G^\beta, \qquad D^\beta = D + \beta K_\beta, \qquad (K_\beta\phi)(b) = (\beta\eta)^{-1}\big(\mathrm{e}^{-\beta(\rho(b_+)-\rho(b_-))} - 1\big)U_b\phi(b_+).$$

The coefficient is at most $\mathrm{e}^{|\beta|}$ in modulus. The map $\phi \mapsto (\phi(b_+))_b$ has norm 1 for each direction and $\sqrt{3}$ for the three directions together. Hence $\|\beta K_\beta\| \leq c'$, and $\|(DG)^\beta\| \leq 2\gamma_0^{-1/2} + 2c'\gamma_0^{-1}$.

(c) The bound on $(GD^*)^\beta$. G is real symmetric, so $(GD^*)^\beta = ((DG)^{-\beta})^*$, and (b) at $-\beta$ applies.

(d) The bound on $(DGD^*)^\beta$. Let $w := G^\beta(D^*)^\beta g$, so that $H_{\mathcal{X}}^\beta w = (D^*)^\beta g$, and note $((D^*)^\beta)^* = D^{-\beta}$.

- From the computation in Lemma D.23, together with $H_{\mathcal{X}} \geq D^*D$, $H_{\mathcal{X}} \geq \gamma_0$ and $c(\beta) \leq \gamma_0/4$,
$$\operatorname{Re}\langle w,\, H_{\mathcal{X}}^\beta w\rangle \geq \tfrac{1}{2}\|Dw\|^2 + \tfrac{\gamma_0}{4}\|w\|^2.$$
- On the other hand, $\operatorname{Re}\langle w,\, H_X^\beta w\rangle = \operatorname{Re}\langle D^{-\beta}w,\, g\rangle \leq (\|Dw\| + c'\|w\|)\|g\|$.

Put $s := (\|Dw\|^2 + \|w\|^2)^{1/2}$. Then $\min\{\frac{1}{2}, \frac{\gamma_0}{4}\}s^2 \leq (1+c'^2)^{1/2}s\|g\|$. Hence

$$\|(DGD^*)^\beta g\| = \|D^\beta w\| \leq (\|Dw\| + c'\|w\|) \leq (1+c'^2)\|g\|/\min\{\tfrac{1}{2}, \tfrac{\gamma_0}{4}\}.$$

Uniformity. Lemma D.23 uses no geometric property of the bond set or of the boundary condition. Its only inputs are $|\rho(b_+) - \rho(b_-)| \leq \eta$ and the range $\sqrt{3}$ of P_k; both hold on every union of whole unit blocks, including wrapping ones, and on T_η. The lower bound (H0) holds on every such $\mathcal{X}$, because the form $\langle f,\, H_X f\rangle$ dominates the sum over unit blocks of the intra-block Neumann form plus $a_k|Q_k f|^2$, and each of these is $\geq \gamma_0\|f\|^2_{\text{block}}$ by [4, (1.8)].

Block operator. The bound on $\mathfrak{D}G\mathfrak{D}^*$ sums the four blocks: G, the block DG, the block GD^* and the block DGD^*, with the vector reading of D. □

At $\beta = 0$ and for $\rho = \operatorname{dist}(\cdot, \operatorname{supp} f')$, (W) reproduces the literature bound [4, Cor. 2.3, (2.30), p. 580]: $|\langle f,\, DGD^* f'\rangle| \leq c_0\mathrm{e}^{-\delta_0 \operatorname{dist}(\operatorname{supp} f, \operatorname{supp} f')}\|f\|_2\|f'\|_2$, with no restriction on the supports.

Lemma D.25 (slice sums)**.** *Let $L \geq 2$, $\eta = L^{-k}$ and $s_j := L^j\eta$ for $0 \leq j \leq k-1$. For $p > 0$ and $\delta > 0$ put*

$$c_\Sigma(p, \delta, L) := (1 - L^{-p})^{-1} + \sup_{1 \leq u \leq L} \sum_{m \geq 0} (uL^m)^p \mathrm{e}^{-\delta u L^m/2}.$$

Then for all $z \geq 0$:

$$\sum_{j=0}^{k-1} s_j^{-p} \mathrm{e}^{-\delta z/s_j} \leq c_\Sigma(p,\delta,L)\max(z,\eta)^{-p}\mathrm{e}^{-\delta z/2}.$$

Proof. Put $r := \max(z,\eta)$ and note $s_j \leq L^{-1} < 1$.

Slices with $s_j \geq r$. These contribute at most $r^{-p}\sum_{m\geq 0} L^{-mp}\mathrm{e}^{-\delta z}$.

Slices with $s_j < r$. These occur only if $r = z$. Put $u_j := z/s_j > 1$; the u_j form a sequence uL^m ($m \geq 0$) with $u \in (1, L]$. Each term equals $z^{-p}u_j^p \mathrm{e}^{-\delta u_j}$, and $\mathrm{e}^{-\delta u_j} \leq \mathrm{e}^{-\delta u_j/2}\mathrm{e}^{-\delta z/2}$ because $u_j \geq z$.

Adding the two parts gives the claim. □

Lemma D.26 (input (K): one-derivative pointwise kernel bounds)**.** *Let A be real and regular on $\mathcal{X}$ in the sense of* [1, (2.23)]*, with e_k small as in Proposition 4.5; under the hypothesis (R_ϑ) of the L^2 sandwich of Subsection D.2 this holds for $A \in \{\tilde{B}, \tilde{A}+\tilde{B}\}$. Let $0 \leq m^2t^2 \leq c$, $0 < \alpha < 1$ and $G := G_k(\mathcal{X}, A)$. Let δ_0, c_0, R_0 be the constants of Proposition 4.5, and put $\delta_K := \delta_0/4$. There are c_K and $c_K(\alpha)$, depending only on d, a, M_b, L (and on α for $c_K(\alpha)$), such that for all x, w with $\rho_{\mathcal{X}}(x), \rho_{\mathcal{X}}(w) \geq R_0$:*

(K0) $\|G(x,w)\| \leq c_K K_2^{\delta_K}(x-w)$;

(K1) $\|D_x G(x,w)\| \leq c_K K_1^{\delta_K}(x-w)$, *and by (S)* $\|(GD^*)(w,x)\| \leq c_K K_1^{\delta_K}(x-w)$;

(K2) *for $|x-\tilde{x}| \leq 1$ with $\rho_{\mathcal{X}}(\tilde{x}) \geq R_0$:*

$$\|U(A(\Gamma_{\tilde{x},x}))D_xG(x,w) - D_{\tilde{x}}G(\tilde{x},w)\| \leq c_K(\alpha)|x-\tilde{x}|^\alpha\big(K_{1-\alpha}^{\delta_K}(x-w) + K_{1-\alpha}^{\delta_K}(\tilde{x}-w)\big).$$

On $\mathcal{X} = T_\eta$ the restriction $\rho_{\mathcal{X}} \geq R_0$ is void.

The lemma is used for scalar lines only; its inputs are Proposition 4.5 and Lemma D.25.

Proof. Step 1 (slices). By [3, (2.6), p. 424],

$$G = \sum_{j=0}^{k-1} G_{(j)}^\eta, \qquad G_{(0)}^\eta = C^{(0),\eta}, \qquad G_{(j)}^\eta = a_j^2 (L^j\eta)^{-4} G_j^\eta Q_j^* C^{(j),L^j\eta} Q_j G_j^\eta \quad (j \geq 1).$$

This is the renormalisation-group identity [1, (2.42)–(2.43)], which is exact. Put $s_j := L^j\eta \in [\eta, L^{-1}]$.

Step 2 (per-slice bounds). Bałaban's slice bounds [3, (2.10)–(2.11), p. 426] follow from [1, Props. 2.1, 2.3] (Proposition 4.5), with $\delta_1 = \delta_0/2$, as follows. For each j, rescale to the L^{-j}-lattice, $x = s_j y$. Then:

- [1, (2.25)] applied to $f = Q_j^*\delta_{y_0}$ bounds $(G_jQ_j^*)(y,y_0)$ and its derivative by $c_0\mathrm{e}^{-\delta_0 \operatorname{dist}(y,B(y_0))}$; [1, (2.24)] gives the same bound for the α-Hölder quotient.
- [1, (2.34)] bounds $C^{(j)}$ by $c_0\mathrm{e}^{-\delta_0|y_0-y_0'|}$.
- The composition is a two-exponential sum, at most $c_d(\delta_0/2)^{-6}\mathrm{e}^{-(\delta_0/2)|y-y'|}$.
- Rescaling back multiplies the kernel by s_j^{-1}, each derivative by s_j^{-1} and each Hölder quotient by $s_j^{-1-\alpha}$; the decay length becomes s_j.

Thus, for depths as in the statement and $n \in \{0,1\}$,

$$\|D_x^n G_{(j)}^\eta(x,w)\| \leq c_1 s_j^{-1-n}\mathrm{e}^{-\delta_1|x-w|/s_j},$$

and the Hölder quotient of $D_xG_{(j)}^\eta$ is at most $c_1 s_j^{-2-\alpha}\mathrm{e}^{-\delta_1 \operatorname{dist}(\{x,\tilde{x}\},w)/s_j}$. For $j = 0$ the bound is [1, (2.33)–(2.34)] at the η-scale. The constants are independent of η, k and $\mathcal{X}$; they depend on the regularity of the background, which is inherited at scale j.

Depth. [1, Props. 2.1, 2.3] require depth $\geq R_0$ in scale-j units, i.e. $\geq R_0 s_j \leq R_0/L$ in scale-k units. So $\rho_{\mathcal{X}} \geq R_0$ at the two evaluation points (and at $\tilde{x}$) suffices for every j. Both points must be deep because the slice is evaluated as $(G_jQ_j^*)(x,\cdot)$ and, by (S), as $(G_jQ_j^*)(w,\cdot)^T$. On $\mathcal{X} = T_\eta$

the restriction is omitted [1, p. 611]. All constants are uniform over unions of big blocks [3, p. 411] and in the side of the torus (Lemma 4.6).

Step 3 (summation).

- (K0), (K1): sum the per-slice bounds and apply Lemma D.25 with $p = 1$ and $p = 2$, $\delta = \delta_1$. The results are $K_2^{\delta_1/2}$ and $K_1^{\delta_1/2}$, and $\delta_1/2 = \delta_K$.
- (K2): fix x, $\tilde{x}$ and j.
  - If $s_j \geq |x - \tilde{x}|$, use the slice Hölder bound.
  - If $s_j < |x - \tilde{x}|$, bound the two terms separately. U is orthogonal and $|x - \tilde{x}|^{-\alpha} < s_j^{-\alpha}$, so the two derivative bounds give $c_1 s_j^{-2-\alpha}\big(\mathrm{e}^{-\delta_1|x-w|/s_j} + \mathrm{e}^{-\delta_1|\tilde{x}-w|/s_j}\big)$.

  In both cases, using $\mathrm{e}^{-\delta_1 \min(a,b)/s} \leq \mathrm{e}^{-\delta_1 a/s} + \mathrm{e}^{-\delta_1 b/s}$, the quotient of slice j is at most $c_1 s_j^{-2-\alpha}\big(\mathrm{e}^{-\delta_1|x-w|/s_j} + \mathrm{e}^{-\delta_1|\tilde{x}-w|/s_j}\big)$. Summing with Lemma D.25 at $p = 2 + \alpha$ gives (K2), with $c_K(\alpha) = 2c_1 c_\Sigma(2 + \alpha, \delta_1, L)$.

The split covers both readings of the range of the Hölder quotient in [1, (2.24)].

Step 4 (column bounds). G is real symmetric, so $(GD^*)(w, x) = (D_x G(x, w))^T$. □

Remark D.27 (sources and limits of (W) and (K)).

(i) (K) is used only for scalar lines. The vector-field slices of [3, (2.6)] are not derived from [1, Props. 2.1, 2.3], and they are not used.

(ii) On Neumann domains (K) is stated only at points of depth $\geq R_0$. Near re-entrant edges a derivative bound uniform in η is not expected (one expects a corner mode $r^{2/3}$). No kernel is evaluated there.

(iii) For $\delta G_k(\square, \eta\mathbb{Z}^3, 0)$ the infinite lattice $\eta\mathbb{Z}^3$ enters only through the bounds (K) on $\eta\mathbb{Z}^3$ at $A = 0$, that is, the bounds of [1, Props. 2.1, 2.3] read on the infinite lattice.

(iv) (W) needs no depth, no shape condition and no regularity beyond (H0). It is uniform in $m^2 \geq 0$. (K) is uniform for $0 \leq m^2 t^2 \leq c$.

Appendix E. The ingredients of Theorem 5.2

This appendix proves the ingredients of the proof of Theorem 5.2 listed in Table 1, except the region entropy, which is Proposition 5.21. Subsection E.1 treats the slice photon: the domination (5.1), the positivity of $C_B(u)$ and the degree in u of $e(\varepsilon; u; M)$. Subsection E.2 proves the bounds on the final c-numbers that are uniform in the torus, the bounds on the $A{\cdot}A$ vertices and the tail beyond the order $\bar{n}$, together with Bałaban's line bounds on the torus on which they rest. Subsection E.4 proves the ingredients themselves (Propositions E.64, E.67, E.71, E.76, E.110, E.159, E.162, E.175). The torus is fixed and the letter j is a slice or scale index, as stated at the start of each subsection.

E.1. **The slice photon: domination, positivity and the degree in u.** We prove three statements of Subsection 5.1: the domination in (5.1), the positivity of $C_B(u)$ for $u > -1/\bar{\delta}$, and the bound on the u-degree of $e(\varepsilon; u; M)$.

Conventions. Throughout, T is one of the tori T_j of Subsection 2.1, of side ℓ_T, and $T_\varepsilon = T \cap \varepsilon\mathbb{Z}^3$ with $\varepsilon = L^{-K}\varepsilon_0$, $K \geq 1$ ($L = 2$ in Theorem 5.2). The index j below is the slice index of (5.1), not the torus index. Put $t_k := L^k\varepsilon$, so $t_K = \varepsilon_0$. Vector fields $A = (A_\mu)_{\mu=1,2,3}$ are triples of real site functions on T_ε, with the inner product $\langle \cdot, \cdot \rangle$ of Subsection 2.2. Operator inequalities are in the sense of quadratic forms on this finite-dimensional real space. The lattice Laplacian $-\Delta_\varepsilon$ of T_ε acts on each component (Lemma B.4), $M_* := -\Delta_\varepsilon + \mu_*^2$ ($\mu_* > 0$), and $C_{\mu_*} = M_*^{-1}$. In this subsection a_k always denotes King's block constants defined below, not the scalar coefficients a_1, a_2 of Section 6.

Blocks (King's averaging). For $1 \leq k \leq K$ and $y \in T \cap t_k\mathbb{Z}^3$ let $B^k(y) := \{x \in T_\varepsilon : \ y_i \leq x_i < y_i + t_k, \ i = 1, 2, 3\}$. Since $\ell_T/t_k = (\ell_T/(M_b L^k \varepsilon))\, M_b \in \mathbb{N}$ by Lemma 2.1(ii), these blocks tile T_ε. Let Q_k be the componentwise average $(Q_k A_\mu)(y) := L^{-3k} \sum_{x \in B^k(y)} A_\mu(x)$ [24, (2.10), p. 653],

with adjoint Q_k^* taken for the inner product $t_k^3 \sum_y$ on the block lattice. Then Q_k^* is the constant extension, $Q_k Q_k^* = 1$, and $P_k := Q_k^* Q_k$ is the orthogonal projection onto the fields that are constant on every scale-k block. Fix King's constant $a > 0$ and put

$$a_k := a(1 - L^{-2})(1 - L^{-2k})^{-1}, \qquad \varrho_k^{\mathrm{K}} := a_k t_k^{-2} \qquad (1 \le k \le K),$$

[24, (2.13), p. 653]. Then $G_k = (M_* + \varrho_k^{\mathrm{K}} P_k)^{-1}$ for $1 \le k \le K$ is the operator of Subsection 5.1. We set $G_0 := 0$, which is King's convention $C^{(0)} = G_1$, $G_k = \sum_{j=0}^{k-1} G_{(j)}$ [24, (2.16)–(2.17), p. 653]. Thus $H_0 = G_1$, and $H_j = G_{j+1} - G_j$ $(0 \le j < K)$ is King's $G_{(j)}$.

Lemma E.1 (King's slices are nonnegative)**.** *For every $K \ge 1$, $a > 0$, $\mu_* > 0$ and every torus T as above:*

(a) $0 < G_1 \le G_2 \le \cdots \le G_K \le C_{\mu_*}$*; equivalently $H_j \ge 0$ for $0 \le j < K$ and $C_{\mu_*} - G_K \ge 0$;*
(b) $\sum_{j<K} H_j = G_K$*, hence $C_{\mu_*} = \sum_{j<K} H_j + (C_{\mu_*} - G_K)$, a sum of nonnegative operators.*

In particular $H_j \ge 0$, which is used in Subsection 5.1.

Proof. Step 1 (inversion reverses order). If $0 < A \le B$ are self-adjoint, then $A^{-1/2} B A^{-1/2} \ge 1$, so its inverse $A^{1/2} B^{-1} A^{1/2}$ is ≤ 1, i.e. $B^{-1} \le A^{-1}$.

Step 2 (nested blocks). For $1 \le k < K$ each block $B^{k+1}(y)$ is the disjoint union of the L^3 blocks $B^k(y')$, $y' \in y + t_k\{0, \dots, L-1\}^3$. So a field constant on the scale-$(k+1)$ blocks is constant on the scale-k blocks, $\operatorname{ran} P_{k+1} \subset \operatorname{ran} P_k$, and $P_k - P_{k+1}$ is the orthogonal projection onto $\operatorname{ran} P_k \ominus \operatorname{ran} P_{k+1}$. In particular $P_k - P_{k+1} \ge 0$.

Step 3 (ϱ_k^{K} decreases). Since $1 - L^{-2k}$ increases with k, a_k decreases, and $\varrho_k^{\mathrm{K}} / \varrho_{k+1}^{\mathrm{K}} = L^2 a_k / a_{k+1} \ge L^2 > 1$.

Step 4 ($H_j \ge 0$). For $1 \le j < K$,

$$G_j^{-1} - G_{j+1}^{-1} = \varrho_j^{\mathrm{K}} P_j - \varrho_{j+1}^{\mathrm{K}} P_{j+1} = \varrho_j^{\mathrm{K}} (P_j - P_{j+1}) + (\varrho_j^{\mathrm{K}} - \varrho_{j+1}^{\mathrm{K}}) P_{j+1} \ge 0$$

by Steps 2 and 3. Both inverses are $\ge M_* > 0$ ($\varrho_k^{\mathrm{K}} > 0$, $P_k \ge 0$), so Step 1 gives $G_j \le G_{j+1}$, i.e. $H_j \ge 0$. For $j = 0$, $H_0 = G_1 > 0$.

Step 5 ($G_K \le C_{\mu_*}$). $G_K^{-1} = M_* + \varrho_K^{\mathrm{K}} P_K \ge M_* > 0$, so $G_K \le M_*^{-1} = C_{\mu_*}$ by Step 1. This proves (a).

Step 6. The sum telescopes: $\sum_{j<K} (G_{j+1} - G_j) = G_K - G_0 = G_K$. Together with (a) this gives (b). □

Lemma E.2 (the domination in (5.1))**.** *Let $\kappa > 0$ and $\Gamma^B = \sum_{j<K} \delta_j H_j$, $\delta_j = \kappa \mu_*^2 t_{j+1}^2$, as in (5.1). Then $0 \le \delta_j \le \bar{\delta} = \kappa \mu_*^2 \varepsilon_0^2$ for $j < K$, and*

$$0 \le \Gamma^B \le \kappa \mu_*^2 \varepsilon_0^2 \, G_K = \bar{\delta} \, G_K \le \bar{\delta} \, C_{\mu_*}.$$

This is the domination $\Gamma^B \le \kappa \mu_*^2 \varepsilon_0^2 G_K \le \bar{\delta} C_{\mu_*}$ stated after (5.1); it is uniform in ε, K, ℓ_T.

Proof. For $j < K$, $0 < t_{j+1} \le t_K = \varepsilon_0$, hence $0 \le \delta_j \le \bar{\delta}$. Since $H_j \ge 0$ (Lemma E.1(a)), $\Gamma^B \ge 0$ and

$$\bar{\delta} \, G_K - \Gamma^B = \kappa \mu_*^2 \sum_{j<K} \big(t_K^2 - t_{j+1}^2\big) H_j \ \ge 0,$$

where $G_K = \sum_{j<K} H_j$ (Lemma E.1(b)) was used. Finally $G_K \le C_{\mu_*}$ (Lemma E.1(a)). □

Remark E.3 (the constant κ). Lemma E.2 holds for every $\kappa > 0$, since κ is only the coefficient in the definition of Γ^B. The numerical bound $\kappa_1 \kappa_3 \le 7.5$ quoted after (5.1) belongs to a different inequality, the domination $D^< \le \Gamma^B$ of the slice-modified photon (Remark 5.1, Theorem I.16). That inequality fixes κ and is the one used in the proof of Proposition 6.7; the admissible values of κ are in Remark I.18.

Lemma E.4 (positivity of the slice photon)**.** *For every $u > -1/\bar{\delta}$,*

$$C_B(u) = C_{\mu_*} + u\Gamma^B = \sum_{j<K} (1 + u\delta_j) H_j + (C_{\mu_*} - G_K) \;\geq\; \min\{1, 1 + u\bar{\delta}\}\, C_{\mu_*} \;> 0,$$

and every step weight satisfies $1 + u\delta_j \geq \min\{1, 1 + u\bar{\delta}\} > 0$.

This is the positivity of $C_B(u)$ for $u > -1/\bar{\delta}$ stated in Subsection 5.1. It is used for the existence of $N(0, C_B(u))$ in the definition of $\Phi_M(u)$, for the convexity of Φ_M on $(-1/\bar{\delta}, \infty)$, and, at $u = -\eta$ with $\eta\bar{\delta} < 1$, in Theorem 5.2.

Proof. The identity follows from Lemma E.1(b) and the definition of Γ^B. If $u \geq 0$, then $u\Gamma^B \geq 0$ (Lemma E.2), so $C_B(u) \geq C_{\mu_*}$. If $-1/\bar{\delta} < u < 0$, then $u\Gamma^B \geq u\bar{\delta}\, C_{\mu_*}$ by Lemma E.2, so $C_B(u) \geq (1 + u\bar{\delta}) C_{\mu_*}$ with $1 + u\bar{\delta} > 0$. In both cases $C_{\mu_*} = M_*^{-1} > 0$ because $-\Delta_\varepsilon \geq 0$ and $\mu_* > 0$. For the weights: $u\delta_j \geq 0$ if $u \geq 0$, and $u\delta_j \geq u\bar{\delta}$ if $u < 0$, because $0 \leq \delta_j \leq \bar{\delta}$. □

Setting for the degree lemma. Write $Z_\varphi(A; a) := \int \exp\big(-S_a^{\mathrm{m}}(A, \varphi)\big) \prod_{x \in T_\varepsilon} d^2\varphi(x)$ for the unnormalised scalar integral with the matter action (2.2). This is the factor in $Z(\Gamma; a) = \mathbb{E}_{N(0,\Gamma)}[Z_\varphi(A; a)]$, whose photon measure is normalised and which contains no c-numbers. The coefficient $a = a_{K,T}(M) = M^2 + \delta m^2_{K,T}(\varepsilon; M)$ of Definition 2.2 is computed with the photon propagator C_{μ_*}. It does not depend on u, and it is a polynomial in (e^2, λ) without constant term added to M^2 (monomials $e^2, \lambda, e^4, e^2\lambda, \lambda^2$). We read $\mathcal{E}(u; M) = T_{\bar{n}}[\Phi_M(u)]$ as the Taylor polynomial of total order $\bar{n}$ in (e, λ) at $(0, 0)$, with $a_{K,T}(M)$ inside, as in [1, (1.13)]. The proof below uses only that the coefficient is a u-independent smooth function of (e^2, λ) equal to M^2 at the origin.

Lemma E.5 (the u-degree of $e(\varepsilon; u; M)$)**.** *Fix ε, T, $M > 0$, $\lambda \geq 0$ and $\kappa, \mu_* > 0$. For $u > -1/\bar{\delta}$ the Taylor coefficients defining $\mathcal{E}(u; M)$ exist. The coefficient of $e^\alpha \lambda^\beta$ ($\alpha + \beta \leq \bar{n}$) vanishes for odd α, and for $\alpha = 2k$ it is a polynomial in u of degree at most k. Hence $\mathcal{E}(u; M)$ and $e(\varepsilon; u; M) = -|T|^{-1}\mathcal{E}(u; M)$ are polynomials in u of degree at most $\lfloor \bar{n}/2 \rfloor$. Their order-zero term $\log Z_\varphi(0; M^2)\big|_{e=\lambda=0} = E_{0,s}(M)$ does not depend on u.*

This is the bound on the degree in u stated in Subsection 5.1. It is used in the proof of Proposition 6.6: the second divided difference of the polynomial $u \mapsto e(\varepsilon; u; m_1)$, and the meaning of $\partial_u^2 e$ in the bound (6.1).

Proof. Step 1 (gauge field enters only through eA). The action (2.2) depends on A only through $\mathrm{e}^{ie\varepsilon A_{x,\mu}}$. Hence $Z_\varphi(A; a) = \mathcal{Z}(eA; \lambda, a)$, where $\mathcal{Z}(B; \lambda, a)$ is the same integral with $\mathrm{e}^{i\varepsilon B_{x,\mu}}$ in place of $\mathrm{e}^{ie\varepsilon A_{x,\mu}}$. If $A \sim N(0, \Gamma)$ then eA has the law of $\sqrt{s}\, Y$, $s := e^2$, $Y \sim N(0, \Gamma)$. Thus, with $\Gamma = C_B(u) > 0$ (Lemma E.4),

$$Z(C_B(u); a) = h_u(e^2, \lambda, a), \qquad h_u(s, \lambda, a) := \mathbb{E}\big[\mathcal{Z}(\sqrt{s}\, Y_u; \lambda, a)\big], \quad Y_u \sim N(0, C_B(u)).$$

Step 2 (regularity of $\mathcal{Z}$). Let $a \geq a_{\min} > 0$ and $\lambda \geq 0$. The gradient and quartic terms of (2.2) are nonnegative, so the integrand is bounded by $\exp(-\frac{1}{2} a_{\min} \sum_x \varepsilon^3 |\varphi(x)|^2)$. Every derivative in B, λ or a multiplies the integrand by a polynomial in $|\varphi|$ whose coefficients are bounded uniformly in B, because $|\mathrm{e}^{i\varepsilon B}| = 1$. By dominated convergence $\mathcal{Z}$ is C^∞ in $(B, \lambda, a) \in \mathbb{R}^n \times [0, \infty) \times [a_{\min}, \infty)$ (one-sided in λ at 0), and all its B-derivatives are bounded uniformly in B.

Step 3 (Gaussian heat equation). Let $c_u(b, b') := \mathbb{E}[Y_u(b) Y_u(b')]$ be the second-moment matrix of Y_u. It is affine in u because $C_B(u)$ is. Put $L_u := \sum_{b,b'} c_u(b, b')\, \partial_{B_b} \partial_{B_{b'}}$. For F smooth with bounded derivatives, $w(s) := \mathbb{E}[F(\sqrt{s} Y_u)]$ satisfies $w'(s) = \mathbb{E}[(\frac{1}{2} L_u F)(\sqrt{s} Y_u)]$ for $s > 0$: differentiate the density of $N(0, s c_u)$, which solves $\partial_s p = \frac{1}{2} L_u p$, and integrate by parts. The right side is continuous

on $[0,\infty)$ by dominated convergence, so $w \in C^\infty([0,\infty))$ with $w^{(i)}(0) = 2^{-i}(L_u^i F)(0)$. Applied to $F = \partial_\lambda^b \partial_a^c \mathcal{Z}(\cdot;\lambda,a)$ (Step 2), h_u is C^∞ on $[0,\infty)\times[0,\infty)\times[a_{\min},\infty)$, and

$$\partial_s^i \partial_\lambda^b \partial_a^c h_u(0,\lambda,a) = 2^{-i}\big(L_u^i\, \partial_\lambda^b \partial_a^c \mathcal{Z}\big)(0;\lambda,a).$$

The right side is a homogeneous polynomial of degree i in the entries of c_u, hence a polynomial in u of degree $\le i$. For $i=0$ it is $\partial_\lambda^b\partial_a^c\mathcal{Z}(0;\lambda,a)$, independent of u, and $\mathcal{Z}(0;\lambda,a)>0$.

Step 4 (composition and Taylor polynomial). Write $a(s,\lambda) := M^2 + \delta m^2_{K,T}(\varepsilon;M)|_{e^2=s}$. This is a polynomial in (s,λ), independent of u, with $a(0,0)=M^2$ (Definition 2.2). Choose a neighbourhood U of $(0,0)$ in $[0,\infty)^2$ on which $a(s,\lambda)\ge M^2/2 =: a_{\min}$. On U the function $\psi_u(s,\lambda) := \log h_u(s,\lambda,a(s,\lambda))$ is C^∞, and $\Phi_M(u) = \psi_u(e^2,\lambda)$. By the chain rule and the formula for derivatives of log, $\partial_s^k\partial_\lambda^\beta \psi_u(0,0)$ is a finite sum, with u-independent coefficients, of terms

$$h_u(0,0,M^2)^{-m}\prod_r \partial_s^{i_r}\partial_\lambda^{b_r}\partial_a^{c_r} h_u(0,0,M^2)\times\big(\text{derivatives of } a(s,\lambda) \text{ at } (0,0)\big),$$
$$\textstyle\sum_r i_r \le k.$$

By Step 3 each such term is a polynomial in u of degree $\le k$, and $h_u(0,0,M^2)=\mathcal{Z}(0;0,M^2)>0$ does not depend on u. Taylor's formula for ψ_u at $(0,0)$ and the substitution $s=e^2$ give

$$\Phi_M(u) = \sum_{2k+\beta\le\bar n} \frac{\partial_s^k\partial_\lambda^\beta\psi_u(0,0)}{k!\,\beta!}\, e^{2k}\lambda^\beta + o\big(|(e,\lambda)|^{\bar n}\big),$$

and by uniqueness of Taylor polynomials this sum is $T_{\bar n}[\Phi_M(u)] = \mathcal{E}(u;M)$. So the coefficients of odd powers of e vanish, and the coefficient of $e^{2k}\lambda^\beta$ has u-degree $\le k \le \lfloor \bar n/2\rfloor$, since $2k\le\bar n$. The order-zero term is $\psi_u(0,0)=\log\mathcal{Z}(0;0,M^2)$, the free scalar constant $E_{0,s}(M)$ of Remark 5.3(i), independent of u. Multiplying by $-|T|^{-1}$ gives the claim for $e(\varepsilon;u;M)$. □

Remark E.6. Lemma E.5 uses that the photon measure in $Z(\Gamma;a)$ is normalised. With the Lebesgue-normalised photon of Propositions 6.3 and 6.4, the free energy would contain the additional term $\frac12\log\det C_B(u)$, which is not a polynomial in u. This is consistent with Remark 5.3(i).

E.2. **Final c-numbers, the $A\cdot A$ vertices and the tail beyond the order $\bar n$.** We prove the three statements behind Lemma A.6(a)–(c), directly at a Higgs mass $M\in(0,m_2]$: bounds on the final c-numbers uniform in the torus (Propositions E.9 and E.21), bounds on the coefficients of the $A\cdot A$ vertices (Lemmas E.24 and E.32), and the tail bound beyond the order $\bar n$ (Proposition E.36). The line bounds, the remainder line and the renormalisation machinery on which they rest are proved in Subsection E.3.

Standing notation. $d=3$, $L=2$, $\varepsilon = 2^{-K}\varepsilon_0$, $t_k := L^k\varepsilon$ (so $t_K=\varepsilon_0$), $e(t)=e\,t^{1/2}$, $\lambda(t)=\lambda t$, $p(t)=b_0(1+\ln t^{-1})^{p_0}$. The scalar field $f=s$ has mass $m_s := M$; the photon $f=v$ (Feynman gauge, per component) has mass $m_v := \mu_*$. On the torus T_ε of side ℓ_T, C_f is the free covariance of field f, and $G_{k,f}(0)$ is Bałaban's block-spin propagator at zero background. Its slices are $H_{j,f} := G_{j+1,f}(0) - G_{j,f}(0)$.

The final c-numbers. The c-numbers are $\delta m^2_{\rm fin} := \delta m^2 - \delta m^2_K$ and $E_{\rm fin} := E_1 - E_K$ of [3, (1.26), (1.30)]. Both are fixed numbers built from free lines, so any exact decomposition of the lines above ε_0 may be used to bound them.

The common top index. Put $\bar\ell(M) := \min(1/M, 1/\mu_*, L)$, and let $\bar K$ be the largest k with $t_k\le\bar\ell(M)$. Assume

$$\varepsilon_0 \le \bar\ell(M), \qquad \ell_T \ge L^3. \tag{E.1}$$

Then $\bar K \ge K$. For both fields telescoping gives the exact identity

$$C_f = G_{K,f}(0) + \sum_{j=K}^{\bar K-1} H_{j,f} + R_{\bar K,f}, \qquad R_{\bar K,f} := C_f - G_{\bar K,f}(0), \qquad f\in\{s,v\}. \tag{E.2}$$

Put $\rho_f := m_f t_{\bar K}$, $\rho_{\min} := \min_f \rho_f$, $Z_f := t_{\bar K}/(m_f^2 \ell_T^3)$, and $\kappa_{R,\min} := \min(\rho_{\min}, \kappa_G)/8$. Here $\kappa_G := \min(c, 1/16)$ is the rate defined before Lemma E.44; it depends only on (a, L).

The following properties are elementary. (P1) $t_{\bar K} \in (\bar\ell/L, \bar\ell]$ is a power of L that does not depend on ε or ℓ_T. (P2) $\rho_f \in (0, 1]$, since $m_f t_{\bar K} \le m_f \bar\ell \le 1$. (P3) $t_{\bar K}^{-1} < L/\bar\ell = \max(LM, L\mu_*, 1)$. (P4) $\bar K - K \le 1 + \log_L^+(\bar\ell/\varepsilon_0)$. (P5) $\rho_{\min} > \min_f m_f \bar\ell/L \ge \min(M, \mu_*) \min\{1, (L \max(M, \mu_*))^{-1}\}$. (P6) $Z_f \le (m_f^2 \max(M, \mu_*) \ell_T^3)^{-1}$, since $t_{\bar K} \le \bar\ell \le 1/\max(M, \mu_*)$; and Z_f is strictly decreasing in ℓ_T. (P7) $\bar K \le \min_f K_f$, where K_f, the top index of the field f alone, is the largest k with $t_k \le \min(1/m_f, \ell_T/L^2)$. Indeed $t_{\bar K} \le \min(1/m_f, L)$, and $L \le \ell_T/L^2$ is $\ell_T \ge L^3$.

Inputs (proved in Subsection E.3). (VS) Let $t_{j+1} \le \min(1/m_f, L)$. Then the slices $H_{j,f}$ obey Bałaban's line bounds [3, (2.10)–(2.12), p. 426] at zero background. The constants depend only on (a, L) (and on α for the Hölder quotients), and the bounds are uniform in ε, j, ℓ_T, μ_* and M. This holds below and above ε_0 (Theorem E.47 with Corollary E.49). (Rem) In scale-$\bar K$ units the remainder $\tilde R$ of $R_{\bar K,f}$ obeys $|\tilde R(x,y)| \le B_R[\mathrm{e}^{-\kappa_R d(x,y)} + Z_f]$ and $\eta^3 \sum_y |\tilde R(x,y)| \le B'_R \rho_f^{-2}$, with $\kappa_R = \min(\rho_f, \kappa_G)/8$. The same bounds hold for first derivatives in either variable, for mixed derivatives and for Hölder quotients, with B''_R and $B_R(\alpha)$ (Proposition E.50). The prefactors depend only on (a, L, α). (S6) The renormalisation machinery of [3, §3] on the torus: Lemma E.51 (with its pinned form (ii)), Lemmas E.52 and E.55, Theorem E.53 and Proposition E.54. It bounds every index assignment whose renormalised subgraphs all have lowest external index $j_0 \le \min_f K_f$, by the power counting of [3, §2] at the top index.

Lemma E.7 (no intermediate subgraph scale)**.** *Assume* (E.1)*. Expand every line of a counterterm graph of δm^2 or of E_1 by* (E.2)*, and fix an index assignment; the line $R_{\bar K,f}$ has index $\bar K$. Let S be a connected proper subgraph all of whose internal lines have index $< j_0$, where j_0 is the lowest index of its external lines. Then $j_0 \le \bar K$, and no internal line of S is an R-line. For each field, the sum over internal indices $i < j_0$ (together with the lines below ε_0) is $G_{j_0,f}(0)$, of the same form for $f = s$ and $f = v$. In particular $j_0 \le \min_f K_f$.*

Proof. Both fields are sliced up to the same index $\bar K$, and every line has an index in $\{< K\} \cup \{K, \dots, \bar K - 1\} \cup \{\bar K\}$, the same set for both fields. Hence $j_0 \le \bar K$. An internal line has index $< j_0 \le \bar K$, so it is not an R-line. The internal lines of field f are therefore slices $H_{i,f}$ with $i < j_0$, or lines below ε_0. By telescoping, their sum is $G_{K,f}(0) + \sum_{K \le i < j_0} H_{i,f} = G_{j_0,f}(0)$, of the same algebraic form for both fields. Finally $j_0 \le \bar K \le \min_f K_f$ by (P7). □

In the case $j_0 \le \min_f K_f$, the replacement of $G_{j_0,f}(0)$ by C^ξ carries the factor $\omega_f := 1 - m_f^2 t_{j_0}^2 \in [0, 1]$. No renormalised subgraph contains an R-line or a full C_f. So no zero mode, and no Ward identity beyond those of [3, §3], enters a renormalisation.

Lemma E.8 (the top cluster)**.** *Assume* (E.1)*, and consider a term with top index $j^* = \bar K$, in scale-$\bar K$ units. The* lower clusters *are its maximal connected subgraphs made of lines of index $< \bar K$, each with its renormalisations. Choose a spanning tree $\mathcal T$ of R-lines on the lower clusters, rooted at the cluster that contains the pinned point x. Then*

$$|\mathit{term}| \le \prod_{\text{lower clusters}} (\textit{bound of Lemmas E.51(ii) and E.55, uniform in the pin}) \cdot \prod_{l \in \mathcal T} N_1(l) \cdot \prod_{l \notin \mathcal T,\ R} N_\infty(l),$$

with $N_1(l) := \sup_y \eta^3 \sum_{y'} |\partial^a \partial'^b \tilde R_f(y, y')| \le B'_R \rho_f^{-2} + B''_R J_0(\kappa_R)(1 + \rho_f^{-2})$ and $N_\infty(l) := \sup |\partial^a \partial'^b \tilde R_f| \le B''_R(1 + Z_f)$, $a, b \in \{0, 1\}$. Here $J_0(\kappa) := \sup_x \eta^3 \sum_y \mathrm{e}^{-\kappa d(x,y)}$ is the lattice sum of Subsection E.3.

Proof. By definition the lower clusters are joined to one another only by R-lines. Sum the cluster positions from the leaves of $\mathcal T$ towards the root. Each tree line is summed over its far endpoint, which costs its ℓ^1 norm N_1. The far cluster's internal positions, relative to that endpoint, are

summed inside its own pinned bound of Lemma E.51(ii). Each non-tree R-line is bounded by its supremum N_∞. The bounds on N_1 and N_∞ are (Rem): on the torus the Z_f-part of the pointwise bound sums to $Z_f\ell^3 = \rho_f^{-2}$. Where Lemma E.55 needs the regularity of an external R-line of a lower cluster, (Rem) supplies the derivative and Hölder bounds with the factor $1 + Z_f$. □

Proposition E.9 (a bound on $\delta m^2_{\mathrm{fin}}$ uniform in the torus)**.** *Let $M \in (0, m_2]$ and assume* (E.1)*. Uniformly in $\varepsilon = 2^{-K}\varepsilon_0$, $x \in T_\varepsilon$ and $\ell_T \in \{\ell_j\}$,*

$$|\delta m^2_{\mathrm{fin}}(x)| \le C_b\Big\{(e^2+\lambda)\frac{L^2}{(L-1)\varepsilon_0} + (e^4+e^2\lambda+\lambda^2)\big(2+\log^+_L(\bar\ell/\varepsilon_0)\big)\Big\}$$
$$+ C_b^R\Big\{(e^2+\lambda)\max(LM, L\mu_*, 1)(1+Z_s+Z_v)^2 + (e^4+e^2\lambda+\lambda^2)(1+Z_s+Z_v)^5\Big\}.$$

Here $C_b = C(\alpha_0, \bar n, d, L, a, M_b)$, and $C_b^R = C(\alpha_0, \bar n, d, L, a, M_b)(1+\rho_{\min}^{-2}+\kappa_{R,\min}^{-3})^5$. The torus side enters only through Z_s and Z_v, both non-increasing in ℓ_T.

Proof. Split the index assignments of (E.2) by the top index $j^* \in \{K, \dots, \bar K\}$.

(i) $K \le j^* < \bar K$. All lines are slices with $t_{j+1} \le t_{\bar K} \le \min(1/m_f, L)$, so (VS) applies to both fields. By Lemma E.7 every renormalised subgraph is in the case $j_0 \le \min_f K_f$, so (S6) applies at top index j^* with no exceptional subgraph. The degrees are $D = -1$ at one loop and $D = 0$ at two loops. Rescaling $\delta m^2 = t_{j^*}^{-2}\cdot$(scale-$j^*$ coefficient) with $e(t)^2 = e^2 t$ and $\lambda(t) = \lambda t$, the one-loop terms are $\le C(e^2+\lambda)t_{j^*}^{-1}$. Their sum is $\sum_{j^*\ge K} t_{j^*}^{-1} = t_K^{-1}\sum_{n\ge 0} L^{-n} = L/((L-1)\varepsilon_0) \le L^2/((L-1)\varepsilon_0)$. The two-loop terms give one $O(1)$ per j^*, and there are $\bar K - K + 1 \le 2 + \log^+_L(\bar\ell/\varepsilon_0)$ values of j^*, by (P4).

(ii) $j^* = \bar K$. Apply Lemma E.8, with at most 2 lines at one loop and at most 5 lines ($I = V + 1 \le 5$) at two loops. Each R-factor is at most $C(1+\rho_f^{-2}+\kappa_R^{-3})(1+Z_f)$. Rescaling from scale $\bar K$ gives $t_{\bar K}^{-2}e^2 t_{\bar K} = e^2 t_{\bar K}^{-1} < e^2\max(LM, L\mu_*, 1)$ at one loop, by (P3), and no factor at two loops. The polynomial in ρ_f^{-1} and κ_R^{-1} is at most its value at $\rho_{\min}$.

The torus side enters (i) nowhere: the constants of (VS) are uniform in ℓ_T, and no condition on the side of the torus is used. It enters (ii) only through Z_f, which is decreasing by (P6). Here (VS) is Theorem E.47, which rests on the free kernel bounds of Lemma E.39, and (S6) consists of Lemmas E.51, E.52, E.55, Theorem E.53 and Proposition E.54 (Remark E.56). □

Lemma E.10 (the splitting of δm^2 and its inner counterterms)**.** *Every graph of δm^2* [3, (1.23), p. 417] *either has no mass counterterm vertex, or is a one-loop two-leg graph Γ whose internal scalar line carries one vertex* [3, (1.7)] *with coefficient δm_1^2, the part of order $\alpha + 2\beta = 2$. Let Γ' be Γ with coefficient 1 at that vertex. For a set J of line indices let Γ'_J, s_J be the parts of Γ', δm_1^2 whose line indices lie in J, with the index K standing for the regular remainder. Split $\delta m^2 = \delta m^2_{\mathrm{fin}} + \delta m^2_{K,k} + \delta m^2_k$ as in* [3, (1.26)–(1.28)] *in two ways: (O) grading all lines, including those of δm_1^2; (B) grading the lines of Γ' only, with δm_1^2 kept as a number. Then:*

(a) *both splittings are exact, and for every graph Γ as above*

$$\delta m^{2,\mathrm{O}}_{K,k} - \delta m^{2,\mathrm{B}}_{K,k} = \Gamma'_{<k}\, s_{[k,K)} - \Gamma'_{[k,K)}\, s_{\ge K}, \qquad \delta m^{2,\mathrm{O}}_{\mathrm{fin}} - \delta m^{2,\mathrm{B}}_{\mathrm{fin}} = \Gamma'_{<K}\, s_{\ge K};$$

the pieces $\Delta_k := (\delta m^2_{K,k} - \delta m^2_{K,k+1})t_k^2$ of the two splittings differ by $(\Gamma'_{<k}s_k - \Gamma'_k s_{>k})t_k^2$;

(b) *$D(\Gamma') = D(\Gamma) + 2 = 1$, whereas a closed scalar or vector loop carrying one vertex (1.7) has degree -1;*

(c) *$|\Gamma'_{\{s\}}| \le C(e^2+\lambda)t_s$, $|s_j| \le C(e^2+\lambda)t_j^{-1}$, and all the differences in (a), multiplied by t_k^2, are at most $C(e^2+\lambda)^2 t_k^2\big(1+\sum_f(1+Z_f)^2\big)$.*

In particular the bounds on $\delta m^2_{K,k}$, on the pieces Δ_k and on $\delta m^2_{\mathrm{fin}}$ hold for both splittings, with the same form.

Proof. (a) Expand multilinearly. In (O) a term of $\Gamma' \cdot \delta m_1^2$ belongs to $\delta m_{K,k}^2$ if all its indices are $< K$ and one of them is $\geq k$; this gives $\Gamma'_{[k,K)} s_{<K} + \Gamma'_{<k} s_{[k,K)}$. In (B) it gives $\Gamma'_{[k,K)}(s_{<K} + s_{\geq K})$. Subtracting gives the first identity. The other two follow in the same way, using $s_{>k} = s_{[k+1,K)} + s_{\geq K}$. (b) By [3, (2.2) and p. 423(ii)], the new vertex has $D_G(v) = d + 2(-d+2)/2 = 2$, and no other vertex changes. One-loop two-leg graphs have $D = -1$, and a free loop has $D = -3$. (c) By [3, (2.10)–(2.12)] and Theorem E.47, every index tuple of Γ' with largest index s contributes at most Ct_s times $(t_a/t_s)^\theta$, $\theta \geq 1$, for each smaller index a; for instance $\min(t_a, t_s)^3/(t_a t_s) = t_s(t_a/t_s)^2$. Summing gives $|\Gamma'_{\{s\}}| \leq C(e^2+\lambda)t_s$, $|\Gamma'_{<k}| \leq C(e^2+\lambda)t_k$ and $|\Gamma'_{<K}| \leq C(e^2+\lambda)\varepsilon_0$. One-loop graphs have degree -1, so $|s_j| \leq C(e^2+\lambda)t_j^{-1}$. By Proposition E.9, $|s_{\geq K}| \leq C(e^2+\lambda)\big(\varepsilon_0^{-1} + \sum_f (1+Z_f)^2 t_{K_f}^{-1}\big)$. Since $t_k \leq \varepsilon_0 \leq t_{K_f}$, (c) follows. □

Remark E.11. Lemma E.10(b) is the reason why the two gradings agree for the mass counterterms but not for the vacuum pieces (Remark E.13): a finite mass vertex makes every two-leg graph convergent, whereas on a vacuum loop it leaves degree -1.

Definition E.12 (graded vacuum pieces)**.** A *bare vacuum graph* is a connected graph without external legs built from the vertices [3, (1.6), (1.7), (1.8), (1.10)], with $n' = 0$, $\tilde{B} = 0$, $g_k = 1$ [3, p. 416]. Every vertex [3, (1.7)] carries the number δm^2 of [3, (1.19)–(1.23)], a polynomial in e, λ of orders $2 \leq \alpha + 2\beta \leq 4$; the order of the vertex is the order of the term of δm^2 it carries. The *lines* of the graph are the lines joining its vertices; the graphs defining δm^2 are not lines. By [3, (1.24)], E_1 is the sum of the values of all bare vacuum graphs of orders $1 \leq \alpha + \beta \leq \bar{n}$. Expand every line of field f as $C_f = \sum_{j=0}^{\bar{K}} C_{(j),f}$, where for $j < K$ the $C_{(j),f}$ are the terms of the decomposition [3, (1.27)] of $G_{K,f}(0)$ (with $C_{(0),f}$ the term $C^{(0)}$), $C_{(j),f} := H_{j,f}$ for $K \leq j < \bar{K}$, and $C_{(\bar{K}),f} := R_{\bar{K},f}$, as in (E.2). The integer j is the *index* of the line. Expanding multilinearly, with every δm^2 kept as a number, define

- E_{fin}: the sum of the terms with at least one line of index $\geq K$;
- E_K: the sum of the terms with all line indices $< K$;
- $E_{K,k}$: the sum of the terms with all line indices $< K$ and at least one $\geq k$;
- E_k: the sum of the terms with all line indices $< k$;
- $\mathcal{V}_k := E_{K,k} - E_{K,k+1}$, for $0 \leq k < K$.

For estimates the coefficients may be split by [3, (1.26), (1.28)] as $\delta m^2 = \delta m^2_{\text{fin}} + \delta m^2_{K,k+1} + \delta m^2_{k+1}$; the first two parts are counterterms of degree 0, and the last has the degree of its graph [3, p. 423].

Remark E.13 (a grading that opens the coefficients fails)*.* One could instead attach to each vertex (1.7) the graph of its coefficient and grade by all lines, including the attached ones; this is the literal reading of [3, p. 418]. It defines different pieces term by term, and neither $\mathcal{V}_k$ nor E_{fin} is then bounded. Take N real components, $\langle \varphi_a \varphi_b \rangle = G\delta_{ab}$, $V = \lambda|\varphi|^4 + \delta m^2 |\varphi|^2$, $\delta m^2 = -2(N+2)\lambda G$; the order-$\lambda$ vacuum energy is $\lambda N(N+2)\sum_x G(x,x)^2$. Let $k \geq 1$ and write $G = A + B + R$, where A collects the slices of index $< k$, B is the slice k and R is the rest, and put $R' := C_0 - G_K(0) \geq 0$. In Definition E.12 the index-k piece is $\lambda N(N+2)\sum_x (B^2 + 2BR)(x,x)$ and the final piece is $\lambda N(N+2)\sum_x R'(x,x)^2$. In the opened grading they are $\lambda N(N+2)\sum_x (B^2 + 2AB)(x,x)$ and $\lambda N(N+2)\sum_x (R'^2 + 2G_K(0)R')(x,x)$. Here $A(x,x) \geq (-\Delta_\varepsilon + m^2 + a_k t_k^{-2})^{-1}(x,x) \geq c\,\varepsilon^{-1}$. So the opened index-$k$ piece exceeds $P(t_k)$ per scale-k block by a factor of order t_k/ε, and the opened final piece is at least of order $\lambda\varepsilon^{-1}|T|$. In Definition E.12 the term $2\lambda N(N+2)\sum_x A(x,x)B(x,x)$ lies in earlier steps, where its coefficient B is the degree-0 vertex $\delta m^2_{K,j+1}$.

Lemma E.14 (grading identities)**.** *(i) $E_1 = E_{\text{fin}} + E_K$ and $E_K = E_{K,k} + E_k$ for $0 \leq k \leq K$, with $E_{K,K} = E_0 = 0$. Hence $E_1 = E_{\text{fin}} + \sum_{k<K} \mathcal{V}_k$. The number E_{fin} does not depend on how $C_f - G_{K,f}(0)$ is decomposed. (ii) For every exact Gaussian staging, $\mathcal{V}_k$ is the field-independent part of order $\leq \bar{n}$ of the output of step k.*

Proof. (i) Every value is multilinear in the lines and every coefficient is a number. Sort the terms by their set of line indices. A term lies in E_{fin} exactly when at least one line carries a piece of $C_f - G_{K,f}(0)$. (ii) Lemma F.31 identifies the step-k output with the terms of the bare vacuum graphs whose largest line index is k. By (i) these are the terms of $\mathcal{V}_k$. □

Lemma E.15 (the finite vertex)**.** *Assume* (E.1). *For $K \le j \le \bar K$ let $\delta m^2_{\langle j\rangle}(x)$ be the expression $\delta m^2(x)$ in which every propagator, including those of the inner insertions in* [3, (1.23)]*, is replaced by $\sum_{i\le j} C_{(i),f}$, and put $\delta m^2_{\succ j} := \delta m^2 - \delta m^2_{\langle j\rangle}$. Then $\delta m^2_{\succ \bar K} = 0$, and for $K \le j < \bar K$, order by order and uniformly in ε, x and ℓ_T,*

$$\sup_x |\delta m^2_{\succ j}(x)|\, t_j^2 \le \Gamma_f \Big[e(t_j)^2 + \lambda(t_j) + e(t_j)^4 + e(t_j)^2\lambda(t_j) + \lambda(t_j)^2 \Big],$$
$$\Gamma_f := C_b\big(1 + \log_L^+(\bar\ell/\varepsilon_0)\big) + C_b^R(1 + Z_s + Z_v)^5.$$

Proof. By multilinearity, $\delta m^2_{\succ j}$ is the sum of the index assignments of the graphs of δm^2 whose largest index j' (inner insertions included) is $> j$. Since $j \ge K$, these are terms of $\delta m^2_{\mathrm{fin}}$, and the proof of Proposition E.9 bounds them for each j' separately: for $j < j' < \bar K$ by $C_b(e^2+\lambda)t_{j'}^{-1}$ at one loop and $C_b(e^4 + e^2\lambda + \lambda^2)$ at two loops; for $j' = \bar K$ by $C_b^R(e^2+\lambda)t_{\bar K}^{-1}(1 + Z_s + Z_v)^2$ and $C_b^R(e^4 + e^2\lambda + \lambda^2)(1 + Z_s + Z_v)^5$. Use $\sum_{j'>j} t_{j'}^{-1} \le t_j^{-1}/(L-1)$, $t_{\bar K}^{-1} \le t_j^{-1}$, and $\bar K - j - 1 \le \bar K - K - 1 \le \log_L^+(\bar\ell/\varepsilon_0)$ by (P4). Multiplying by t_j^2 and using $e(t)^2 = e^2 t$, $\lambda(t) = \lambda t$ gives the claim. □

Lemma E.16 (degrees with finite vertices)**.** *Let Q be a connected graph built from the vertices of Definition E.12, in which each vertex* [3, (1.7)] *carries either a graph of $\delta m^2_{\langle j\rangle}$ attached as in* [3, p. 417]*, or the number $\delta m^2_{\succ j}$ (a* finite vertex*). Let m_f be the number of finite vertices, o' the total order of the other vertices, E the number of external legs and n_∂ the number of differentiations acting on external legs. Then*

$$D(Q) = o' + 2m_f - 3 + \tfrac12 E + n_\partial.$$

If $m_f \ge 1$, Q has at least one line, $E \ge 1$, and Q does not vanish identically, then $D(Q) \ge \frac12$.

Proof. By [3, (2.1)] with $d = 3$, a leg on an internal line contributes $-\frac12$ and a differentiation of an internal line -1; external legs and their differentiations contribute 0. So $D_Q(v) = D(v) + \frac12(\#\text{external legs at } v) + (\#\text{external differentiations at } v)$. By [3, p. 423, (i), (iii), (iv)], $D(v) = o(v)$ for the bare vertices. An attached graph G_0 has $D(v) + D(G_0) = o(G_0)$ (proof of Lemma E.166(b)). A finite vertex has $D(v) = 2$ and counterterm degree 0 [3, p. 423]. Now [3, (2.3)] gives the formula. Every vertex has an even number of scalar legs; if $E = 1$, the external leg is therefore a vector leg, and Q vanishes [3, p. 434], as in Lemma E.166(c). So $E \ge 2$. If $o' > 0$ then $o' \ge \frac12$ and $D \ge \frac12 + 2 - 3 + 1$. If $o' = 0$, then Q has a line only if $m_f \ge 2$, and then $D \ge 4 - 3 + 1$. □

Torus moments of the cut classes. Let $K \le j_0 \le \bar K$, $\xi = L^{-j_0}$, and work in scale-j_0 units. The rescaled torus is $T^\xi = \xi\mathbb{Z}^3/\Gamma$, $\Gamma = \bigoplus_\mu \ell'_\mu \mathbb{Z} e_\mu$, $\ell'_\mu \in \xi\mathbb{N}$. By (P1) and (E.1), $t_{j_0} \le \bar\ell \le L$ and $\ell_T \ge L^3$, so $\ell' := \min_\mu \ell'_\mu \ge L^2 \ge 4$. For $w \in T^\xi$ let $\bar w \in \xi\mathbb{Z}^3$ be the representative with $|\bar w_\mu| \le \ell'_\mu/2$, $|w|_T := |\bar w|$, and $\rho_X(w) := |w|_X + \xi$ for $X \in \{\xi\mathbb{Z}^3, T^\xi\}$. Let C^ξ_∞ and C^ξ_T be the resolvents of $-\Delta^\xi + 1$ on $\xi\mathbb{Z}^3$ and on T^ξ (componentwise for the photon); C^ξ_T is the kernel C^ξ of [3, p. 437] on the torus, for both fields. ∂^a is any product of a forward or backward differences. We use: (C) C^ξ_∞ is even and reflection invariant, and $|\partial^a C^\xi_\infty(z)| \le C_0\rho(z)^{-1-a}\mathrm{e}^{-\delta_0|z|}$ for $a \le 2$ [3, (2.10)–(2.12), p. 426, p. 437] (see (H_∞) in Subsection 5.5); (Per) $C^\xi_T(w) = \sum_{n\in\Gamma} C^\xi_\infty(\bar w + n)$, by the proof of Lemma 5.22(i), since $-\Delta^\xi + 1$ commutes with Γ; (W) on $\xi\mathbb{Z}^3$, with every line equal to C^ξ_∞, the zeroth and first moments of C3 vanish [3, (3.27)–(3.29), p. 441]. Let N_0 be the largest

number of kernel factors (partner coefficients included) in a term of a member of C1′, C3 or C4, and $r_0 := 1/(8N_0)$.

Lemma E.17 (images)**.** *For $n \in \Gamma \setminus \{0\}$, $|\bar w + n| \geq \frac12|\bar w| + \frac14|n|$. Consequently: (a) if $|F(z)| \leq A\mathrm{e}^{-\delta|z|}$ on $\xi\mathbb{Z}^3$, then $|\mathrm{Per}\, F(w) - F(\bar w)| \leq Ac_\delta \mathrm{e}^{-\delta|w|_T/2}$, $c_\delta := \sum_{m\in\mathbb{Z}^3\setminus 0} \mathrm{e}^{-\delta|m|}$; (b) $|\partial^a C_T^\xi(w)| \leq C_0'\rho_T(w)^{-1-a}\mathrm{e}^{-\delta_0|w|_T/4}$ for $a \leq 2$; (c) for $\xi \leq 1$: $\sum_{w\in X} \xi^3 \rho_X(w)^{-k}\mathrm{e}^{-\delta|w|_X} \leq C(k,\delta)$ for $0 \leq k < 3$, and $\leq C(\delta)\xi^{-1}$ for $k = 4$.*

Proof. If $n_\mu \neq 0$ then $|n_\mu| \geq \ell'_\mu \geq 2|\bar w_\mu|$, so $|\bar w_\mu + n_\mu| \geq \max(|\bar w_\mu|, \frac12|n_\mu|)$; this is trivial if $n_\mu = 0$. Hence $|\bar w + n| \geq \max(|\bar w|, \frac12|n|)$; average. (a) $|n| \geq 4|m|$ for $n = (\ell'_\mu m_\mu)_\mu$, so $\mathrm{e}^{-\delta|\bar w+n|} \leq \mathrm{e}^{-\delta|\bar w|/2}\mathrm{e}^{-\delta|m|}$; sum over $m \neq 0$. (b) The term $n = 0$ is (C). For $n \neq 0$, $|\bar w + n| \geq 2$, so the terms are $\leq C_0\mathrm{e}^{-\delta_0|\bar w+n|}$, and (a) gives $C_0c_{\delta_0}\mathrm{e}^{-\delta_0|\bar w|/2} \leq C\rho_T^{-1-a}\mathrm{e}^{-\delta_0|\bar w|/4}$. (c) $w \mapsto \bar w$ is injective and $|w|_T = |\bar w|$; compare with $\int(|z|+\xi)^{-k}\mathrm{e}^{-\delta|z|/2}d^3z$, the term $z = 0$ contributing ξ^{3-k}. □

Lemma E.18 (short and long parts)**.** *Fix $\chi_0 \in C^\infty([0,\infty))$, $0 \leq \chi_0 \leq 1$, $\chi_0 = 1$ on $[0, \frac12]$, $\chi_0 = 0$ on $[1,\infty)$; put $\chi(z) := \chi_0(|z|/r_0)$, $C^< := \chi C_\infty^\xi$, $C^> := C_\infty^\xi - C^<$. These are even and reflection invariant, and for $a \leq 2$: (a) $\partial^a C^<(z) = 0$ for $|z| \geq r_0 + 2\xi$ and $|\partial^a C^<(z)| \leq C_1\rho(z)^{-1-a}$; (b) $|\partial^a C^>(z)| \leq C_1\mathrm{e}^{-\delta_0|z|}$; $C_1 = C_1(C_0, \delta_0, r_0, \chi_0)$. With $C_T^\lessgtr := \mathrm{Per}\, C^\lessgtr$, $C_T^\gtrless := \mathrm{Per}\, C^\gtrless$ we have $C_T^\xi = C_T^< + C_T^>$; if $\xi < r_0$ then $\partial^a C_T^<(w) = \partial^a C^<(\bar w)$; and $|\partial^a C_T^>(w)| \leq C_2\mathrm{e}^{-\delta_0|w|_T/2}$.*

Proof. By $\partial_e(fg)(z) = (\partial_e f)(z)g(z+\xi e) + f(z)(\partial_e g)(z)$, every term of $\partial^a(\chi C_\infty^\xi)$ is $(\partial^b\chi)(z+\sigma)(\partial^{a-b}C_\infty^\xi)(z+\sigma')$ with $|\sigma|, |\sigma'| \leq 2\xi$ and $|\partial^b\chi| \leq c_b r_0^{-b}$; also $\rho(z+\sigma') \geq \rho(z)/3$. (a) On the support $\rho(z) \leq r_0 + 5\xi \leq 6$, so $r_0^{-b} \leq (6/r_0)^b\rho(z)^{-b}$, and (C) gives the bound. (b) A term is non-zero only if $|z+\sigma| \geq r_0/2$ for some $|\sigma| \leq 2\xi$; then $\rho(z+\sigma') \geq r_0/12$ (from $\rho \geq \xi$ if $\xi \geq r_0/12$, otherwise from $|z+\sigma'| \geq r_0/2 - 4\xi$), and (C) gives $C(r_0)\mathrm{e}^{-\delta_0|z|+4\delta_0\xi}$. If $\xi < r_0$, the range $r_0 + 2\xi < \frac14\ell'$ of $\partial^a C^<$ is smaller than $|\bar w + n| \geq 2$ for $n \neq 0$, so only $n = 0$ contributes. For $C_T^>$ use Lemma E.17(a). Finally $C_T^\xi = C_T^< + C_T^>$ by (Per). □

A term of a class is a product of kernels $\partial^a C_X^\xi(u-v)$, with u, v vertex positions or their shifts by at most ξ, all positions except the base y summed; the coefficient of the partner of a subgraph H is written as the graph of H with one leg moved [3, p. 417]. Expand every line as $C_X^\xi = C_X^< + C_X^>$; for an assignment $\mathfrak{a}$ of a letter $<$ or $>$ to every line of the parent graph (a subgraph and its partner carrying the same letters) this gives $\Pi_X = \sum_{\mathfrak{a}} \Pi_X^{\mathfrak{a}}$ for each moment Π; $\mathfrak{s}$ denotes the assignment with only $<$.

Lemma E.19 (no wrapping)**.** *Let $\xi < r_0$. Then $\Pi_{T^\xi}^{\mathfrak{s}} = \Pi_{\xi\mathbb{Z}^3}^{\mathfrak{s}}$ for the zeroth moment of every member with its partners, and for the first moment, with $y' - y$ the minimal displacement on T^ξ.*

Proof. Each term is connected: every point is linked to y by at most N_0 kernels. By Lemma E.18(a) a non-vanishing kernel links points at torus distance $< r_0 + 4\xi$, so every point p of a non-vanishing configuration has $|p-y|_T < R_0 := N_0(r_0 + 4\xi) < \frac58$. Fix a lift $\tilde y$ of y and put $\tilde p := \tilde y + \overline{p-y}$. Then $|\tilde u - \tilde v| < 2R_0 < \ell'/2$, so $\overline{u-v} = \tilde u - \tilde v$ and $\partial^a C_T^\lessgtr(u-v) = \partial^a C^<(\tilde u - \tilde v)$. Every non-vanishing configuration on $\xi\mathbb{Z}^3$ lies in the ball of radius R_0 about $\tilde y$, which projects injectively to T^ξ since $|n| \geq \ell' > 2R_0$ for $n \in \Gamma \setminus 0$. So both sums run over the same configurations with the same values. □

Proposition E.20 (torus moments)**.** *Replace every line of the classes C1′ and C3 of Lemma E.166, and of the partners of their proper divergent subgraphs, by C_T^ξ. Let $\Pi^{(1')}$, $\Pi^{(3)}$ be the zeroth moments at a base point $y \in T^\xi$ and $\Pi_\nu^{(3)}$ the first moment of C3. There is C, depending only on d and on the classes (through N_0), such that, uniformly in ξ, y and $\ell_T \geq L^3$: (i) $|\Pi^{(3)}| + \sum_\nu |\Pi_\nu^{(3)}| \leq C$; (ii) $|\Pi^{(1')}| \leq C$.*

Proof. If $\xi \geq r_0$, then $\rho \geq r_0$ and every kernel is $\leq C(r_0)\mathrm{e}^{-\delta_0|w|_T/4}$ by Lemma E.17(b); bound the kernels of a spanning tree by this and the others by their supremum, and use Lemma E.17(c) with $k=0$ (also for the weight $|w|_T$). Let now $\xi < r_0$.

(i) By [3, (3.27), first line, and (3.29)], written with C_X^ξ, $\Pi^{(3)}$ is a linear combination, with coefficients depending only on d and on the charge matrix, of bubble terms $\sum_w \xi^3 (D_1 C_X^\xi)(w)(D_2 C_X^\xi)(w)$, $\operatorname{ord} D_1 + \operatorname{ord} D_2 = 2$, and local terms $(D_3 C_X^\xi)(\sigma)$, $\operatorname{ord} D_3 \leq 2$, $|\sigma| \leq 2\xi$ (translation invariance and evenness were used). $\Pi_\nu^{(3)}$ is a combination of the weighted bubbles $\sum_w \xi^3 \bar{w}_\nu (D_1 C_X^\xi)(D_2 C_X^\xi)(w)$. Expand $C_X^\xi = C_X^< + C_X^>$. Every $\mathfrak{a} \neq \mathfrak{s}$ has a factor $C_X^>$: a local term with it is $\leq C_2$; a bubble with orders $(1,1)$, or with $C^>$ carrying the order-2 factor, is $\leq C \sum \xi^3 \rho_X^{-2} \mathrm{e}^{-\delta|w|_X} \leq C$ (Lemmas E.17(c), E.18); if $C^>$ carries the order-0 factor, sum by parts once, which is exact on T^ξ by periodicity and on $\xi\mathbb{Z}^3$ by finite range, and obtain the same bound; in the weighted bubbles $|\bar{w}_\nu|\rho^{-3} \leq \rho^{-2}$. By (W) and Lemma E.19, $\Pi_T^{(3)} = \Pi_T^{(3)} - \Pi_{\xi\mathbb{Z}^3}^{(3)} = \sum_{\mathfrak{a}\neq\mathfrak{s}} (\Pi_T^{\mathfrak{a}} - \Pi_{\xi\mathbb{Z}^3}^{\mathfrak{a}})$, and likewise for $\Pi_\nu^{(3)}$; both sums are bounded.

(ii) By [3, (3.21)–(3.24), pp. 438–439], written with C_T^ξ, $\Pi^{(1')}$ is a linear combination of $E_\sigma := \sum_w \xi^3 C_T^\xi(w+\sigma)(\partial_\mu C_T^\xi)(w)$ and $\xi C_T^\xi(\sigma')$, $|\sigma|, |\sigma'| \leq 2\xi$. Translation invariance gives $E_0 = -\frac{\xi}{2}\sum_w \xi^3 |\partial_\mu C_T^\xi(w)|^2$, which is $\leq C\xi \cdot \xi^{-1}$ by Lemma E.17(b),(c). Also $|E_\sigma - E_0| \leq C\sum_w \xi^3 \xi \rho_T^{-4} \mathrm{e}^{-\delta|w|_T} \leq C$ and $\xi |C_T^\xi(\sigma')| \leq C$. □

Proposition E.21 (a bound on E_{fin} uniform in the torus, at $u=0$)**.** *Let $M \in (0, m_2]$ and assume* (E.1). *Let E_{fin} be as in Definition E.12. Put $o := \alpha/2 + \beta$, $P_E(\varepsilon_0) := \sum_{2\leq\alpha+2\beta\leq 6} |e|^\alpha \lambda^\beta \varepsilon_0^o$ and $\Phi := (1+\Gamma_f)^{N_v}$, with Γ_f as in Lemma E.15 and $N_v = N_v(\bar{n})$ the largest number of vertices of a bare vacuum graph of order $\leq \bar{n}$. Then, uniformly in $\varepsilon = 2^{-K}\varepsilon_0$ and in $\ell_T \geq \ell_0$,*

$$|E_{\mathrm{fin}}| \leq |T_\varepsilon| \Big\{ C_E\, \Phi \Big[L^3 \varepsilon_0^{-3} P_E(\varepsilon_0) + \sum_{\alpha+2\beta=6} |e|^\alpha \lambda^\beta \big(2 + \log_L^+(\bar{\ell}/\varepsilon_0)\big) + \sum_{7\leq\alpha+2\beta\leq 2\bar{n}} \frac{|e|^\alpha \lambda^\beta t_{\bar{K}}^{o-3}}{1 - L^{-(o-3)}} \Big] + C_E^R t_{\bar{K}}^{-3} \sum_{2\leq\alpha+2\beta\leq 2\bar{n}} |e|^\alpha \lambda^\beta t_{\bar{K}}^o (1 + Z_s + Z_v)^{N_\ell} \Big\},$$

with $C_E = C(\alpha_0, \bar{n}, d, L, a, M_b)$, $C_E^R = C(\alpha_0, \bar{n}, d, L, a, M_b)(1 + \rho_{\min}^{-2} + \kappa_{R,\min}^{-3})^{N_\ell}$, $N_\ell = N(\bar{n})$ the maximal number of lines, and $t_{\bar{K}}^{-3} < \max(LM, L\mu_*, 1)^3$. The torus side enters only through Z_s and Z_v; Γ_f and the whole right side are non-increasing in ℓ_T.

Proof. Step 1 (top index and vertex split). For a term of E_{fin} let $j^* \in \{K, \dots, \bar{K}\}$ be its largest line index; then $E_{\mathrm{fin}} = \sum_{j^*=K}^{\bar{K}} E^{[j^*]}$. In $E^{[j^*]}$ write every δm^2 as $\delta m^2_{\langle j^*\rangle} + \delta m^2_{\succ j^*}$ (Lemma E.15). Write the first part in its graph representation [3, p. 417] (the vertex replaced by the graph with both legs localised at its position and the other leg summed); its attached lines have indices $\leq j^*$. The second part is a finite vertex. By multilinearity, $E^{[j^*]}$ is the sum of the terms with all lines, attached or not, of index $\leq j^*$ and at least one non-attached line of index j^*. Decompose by Bałaban's orderings [3, (3.4), p. 424], with ties broken by a fixed order of the lines of each parent graph in which attached lines precede the others, so that the last line $l(m)$ is a non-attached line of index j^*. A term in which a vertex carries a graph G_0 of $\delta m^2_{\langle j^*\rangle}$ is the counterterm member of the parent in which that vertex is replaced by G_0. By Lemma E.52, with $j_{l(m)} = j^*$ fixed, every divergent proper subgraph renormalised within an ordering is a G_i with $i < m$, and its partner has attached lines of index $< j_0 \leq j^*$; so the partner is present in $E^{[j^*]}$. The transformations [3, (3.5), pp. 434–445] act inside the sub-sum, and no term is moved to another top index. By Lemma E.7, every renormalised subgraph has $j_0 \leq \bar{K} \leq \min_f K_f$, its internal lines resum to $G_{j_0,f}(0)$, and this is the case covered by (S6).

Step 2 (the cut). The proof of Lemma E.165 uses only the definition of $J(\bar{l})$ and $l(m) \notin G_i$ for $i < m$, so it applies at top index j^*. Cutting $l(m)$, i.e. treating its kernel $K_{j^*} = C_{(j^*),f}$ as a two-variable external field [3, p. 419, p. 422], maps every class term by term onto two-leg classes of $G \setminus l(m)$. For $j^* < \bar{K}$, in scale-j^* units, $\|hK_{j^*}h'\|_{1,\alpha_0} \le O(1)\mathrm{e}^{-\delta_1 \operatorname{dist}(\operatorname{supp} h, \operatorname{supp} h')}$ by (VS). By Lemma E.166(c), $G \setminus l(m)$ is connected or the term vanishes. By Lemma E.166(d) and Lemma E.16, if $D(G \setminus l(m)) \le 0$ then:

(a) G is a single finite vertex closed by $l(m)$; its value at x is a multiple of $\delta m^2_{\succ j^*}(x) \operatorname{tr} K_{j^*}(x,x)$, which in scale-$j^*$ units is at most $O(1)$ times the bound of Lemma E.15;

(b) $G \setminus l(m)$ is a one-particle-irreducible member of C1 or C2; its partner, the vertex carrying $G_0 \cong G \setminus l(m)$ with lines of index $< j^*$ closed by $l(m)$, belongs to $\delta m^2_{(j^*)}$, and the class is renormalised as a pinned two-leg class by Lemma E.55;

(c) $G \setminus l(m)$ is a one-particle-reducible member of C2; it has no partner, and its class is bounded by Statement K.6 with $F = K_{j^*}$;

(d) $G \setminus l(m)$ is of type C1′, C3 or C4; its class is bounded by Statement K.4 with $F = K_{j^*}$, based at the pinned endpoint. For C1′ no hypothesis is needed; for C3 the hypothesis of that statement is Proposition E.20(i); for C4 it is Statement K.5. On the torus these moments need not vanish: the identities [3, (3.27)–(3.29)] hold on $\xi\mathbb{Z}^3$ only, and the one-loop zeroth moment on T^ξ is a sum over the non-zero images.

For $j^* < \bar{K}$, $N(K_{j^*}) \le C$ by [3, (2.10)–(2.11), p. 426] and (VS). Every other component of $G \setminus l(m)$ has $D \ge \frac{1}{2}$. Every proper component containing a finite vertex has $D \ge \frac{1}{2}$ or vanishes (Lemma E.16). Every other proper component with $D \le 0$ is renormalised by (S6), as in part (i) of the proof of Proposition E.9.

Step 3 ($K \le j^* < \bar{K}$). All lines are slices: those of index $< K$ obey [3, (2.10)–(2.12), p. 426], and those of index $\ge K$ obey them by (VS) (Theorem E.47, resting on Lemma E.39). Pin an endpoint x of $l(m)$ and work in scale-j^* units, with couplings $e(t_{j^*})$, $\lambda(t_{j^*})$ and finite-vertex coefficients $\delta m^2_{\succ j^*} t^2_{j^*}$ [3, p. 418]. By Step 2, after renormalisation every component $G_i^{(\alpha)}$, $i < m$, has degree $\ge \vartheta > 0$. Lemma E.51(ii) with the largest index fixed to j^* (no sign condition on $D(G)$, and $L^{j^*}\eta = 1$), Statements K.4, K.6 for the classes (c), (d), and Lemma E.15 bound the pinned value of each class by $C|e|^\alpha \lambda^\beta t^o_{j^*} \Gamma_f^{m_f}$, uniformly in x, where (α,β) is the total order including the finite vertices. The number of graphs and orderings depends only on $\bar{n}$, and there are $|T_\varepsilon| t_{j^*}^{-3}$ pin blocks. Hence $|E^{[j^*]}| \le C_E \Phi |T_\varepsilon| \sum_{\alpha,\beta} |e|^\alpha \lambda^\beta t_{j^*}^{o-3}$.

Step 4 ($j^* = \bar{K}$). Here $\delta m^2_{\succ \bar{K}} = 0$, so there are no finite vertices. Lemma E.8, with one vertex pinned per scale-$\bar{K}$ block, gives $|E^{[\bar{K}]}| \le C_E^R |T_\varepsilon| t_{\bar{K}}^{-3} \sum |e|^\alpha \lambda^\beta t^o_{\bar{K}} (1 + Z_s + Z_v)^{N_\ell}$. In it, every lower cluster of type C1′, C3, C4, or a one-particle-reducible member of C2, is bounded by Statement K.4 or K.6; the field on its two legs is one R-line, or a product $\Phi_1(y)\Phi_2(z)$ of two R-lines with their far ends held fixed and summed as in Lemma E.8, so $N(F) \le C(1 + Z_s + Z_v)^2$ by Proposition E.50, and these factors are among those counted by $(1 + Z_s + Z_v)^{N_\ell}$.

Step 5 (sum over j^*). The integer α is even, so $o \in \mathbb{Z}$. For $o \le 2$: $\sum_{j^* \ge K} t_{j^*}^{o-3} \le \varepsilon_0^{o-3}/(1 - L^{-1}) \le L^3 \varepsilon_0^{o-3}$. For $o = 3$: there are $\bar{K} - K \le 1 + \log_L^+(\bar{\ell}/\varepsilon_0)$ values of $j^* < \bar{K}$, by (P4). For $o \ge 4$: $\sum_{j^* < \bar{K}} t_{j^*}^{o-3} \le t_{\bar{K}}^{o-3}/(1 - L^{-(o-3)})$. Finally $t_{\bar{K}}$, C_E, C_E^R do not depend on ℓ_T, while Z_f, and with it Γ_f, is non-increasing in ℓ_T by (P6). □

Remark E.22. In the grading of Definition E.12, E_{fin} contains the terms whose lines lie above ε_0 while their δm^2 coefficients contain lines below ε_0; it does not contain the terms in which only a coefficient line lies above ε_0, which belong to E_K with a finite vertex. The factor Φ accounts for the finite vertices $\delta m^2_{\succ j^*}$. The number $\delta m^2_{\mathrm{fin}} := \delta m^2 - \delta m^2_K$ is a coefficient and keeps the meaning of [3, (1.26)]; by Lemma E.10 its bounds are the same in both gradings. The form of the bound at $u \ne 0$ is Proposition E.162.

Remark E.23 (the mass range). (i) For $M \le m_2$, $\bar\ell(M) \ge \bar\ell(m_2)$. So (E.fin) gives (E.1), and (T.floor) gives $\ell_T \ge L^3$. (ii) The construction slices both fields to one index $\bar K$. The order of M and μ_* is used nowhere: when $M < \mu_*$, as at $M_1 = 0.7m$, the scalar's R-line is the long one and is bounded by (Rem) in the same way. At $M = \mu_*$ the two fields coincide in the construction. (iii) C_b^R and C_E^R are non-increasing in $\rho_{\min}$, since $\kappa_{R,\min}$ is non-decreasing in it. So they may be evaluated at the lower bound (P5), which is the quantity $\rho_{\min}(M)$ of Lemma A.6.

The $A \cdot A$ vertices. Work in step-k units: $\eta = L^{-k}$, and $\ell_j := L^j\eta \le 1$ is the scale of a line of index j. The vertices of [3, (3.30), p. 442] are

$$e(t_k)^2 \sum_x \eta^d\, g(x) A_\mu(x)\, c_{\mu\mu'}(x)\, g'(x) A'_{\mu'}(x)$$

and the first-moment vertex, with coefficient $c_{\mu\mu'\nu}$. Here j_0 is the lowest j-index of the external legs. On the ξ-lattice, $\xi := L^{-j_0}$, put $C^\xi := (-\Delta^\xi + 1)^{-1}$ and $G^\xi_{j_0}(0) := (-\Delta^\xi + m^2_{j_0} + a_{j_0}P_{j_0})^{-1}$. The rescaled Higgs mass is $m^2_{j_0} = M^2 t^2_{j_0}$, and P_{j_0} is the block projection. Write $\rho(y,y') := |y-y'| + \xi$.

Lemma E.24 (coefficient bounds for the $A \cdot A$ vertices)**.** *Let $0 \le m^2_{j_0} \le 1$. Then*

$$|c_{\mu\mu'}(x)| \le \Gamma_3\, \ell_{j_0}^{-1}, \qquad |c_{\mu\mu'\nu}(x)| \le \Gamma'_3\, \ell^0_{j_0}.$$

The constants Γ_3, Γ'_3 depend only on d, a (through $a_\flat \le a_{j_0} \le a$) and the block size. They are uniform in $m^2_{j_0} \in [0,1]$, ξ, j_0, k, ε and ℓ_T, and are free of μ_*, u and the photon zero mode. In particular, under (E.m2) and $t_{j_0} \le \varepsilon_0$ the bounds hold with the same constants at every Higgs mass $M \in (0, m_2]$.

Proof. Step 1 (resolvent identity). $G^\xi_{j_0}(0) - C^\xi = G^\xi_{j_0}(0)\big[(C^\xi)^{-1} - G^\xi_{j_0}(0)^{-1}\big]C^\xi = G^\xi_{j_0}(0)\,\Upsilon\, C^\xi$, with $\Upsilon := 1 - m^2_{j_0} - a_{j_0}P_{j_0}$. Because P_{j_0} is an orthogonal projection, Υ has spectrum in $\{1 - m^2_{j_0},\, 1 - m^2_{j_0} - a_{j_0}\}$. So $\|\Upsilon\| \le 1 + a$ for $m^2_{j_0} \in [0,1]$, and the kernel of Υ is bounded and supported in $|z - z'| \le 2\sqrt{d}$.

Step 2 (the line bounds and the mass). At zero background, $G^\xi_{j_0}(0)$ is the sum of its slices of index $< j_0$, together with the bottom propagator. By Theorem E.47, which covers the scalar half at mass M for $t_{j+1} \le \min(1/M, L)$ (here $t_{j_0} \le \varepsilon_0 \le 1/m_2 \le 1/M$), it obeys $|G^\xi_{j_0}(0)(y,y')| \le O(1)\rho(y,y')^{-1}\mathrm{e}^{-\delta|y-y'|}$ and $|\partial G^\xi_{j_0}(0)(y,y')| \le O(1)\rho^{-2}\mathrm{e}^{-\delta|y-y'|}$, where $\rho(y,y') := |y-y'| + \xi$. The $O(1)$ is uniform in M and in ξ. The mass enters the quadratic form only through $+m^2_{j_0} \ge 0$, so it does not weaken the coercivity.

Step 3 (the replacement difference is one dimension better). Put $D := G^\xi_{j_0}(0) - C^\xi = G^\xi_{j_0}(0)\Upsilon C^\xi$. By Steps 1–2,

$$|D(y,y')| \le O(1) \sum_{z,z'} \xi^{2d} \rho(y,z)^{-1}\mathrm{e}^{-\delta|y-z|}\, 1_{|z-z'|\le 2\sqrt{d}}\, \rho(z',y')^{-1}\mathrm{e}^{-\frac12|z'-y'|} \le O(1)\mathrm{e}^{-\delta'|y-y'|},$$

because $\int d^3z\, |y-z|^{-1}|z-y'|^{-1}\mathrm{e}^{-\delta(|y-z|+|z-y'|)} < \infty$ in $d=3$. With one difference at one end, $\rho^{-2} * \rho^{-1}$ in $d=3$ gives $O(1)(1 + \log_+(1/\rho))\mathrm{e}^{-\delta'|y-y'|}$.

With one difference at each end, $|\partial_y\partial_{y'} D(y,y')| \le O(1)\rho(y,y')^{-1}\mathrm{e}^{-\delta'|y-y'|}$, after decreasing δ'. Indeed, $\partial_y\partial_{y'} D = (\partial G^\xi_{j_0}(0))\,\Upsilon\,(\partial C^\xi)$, and $|\partial C^\xi(z,y')| \le c\rho(z,y')^{-2}\mathrm{e}^{-|z-y'|/4}$ by Lemma E.39.

- The part with $a_{j_0}P_{j_0}$ has a bounded kernel supported in $|z-z'| \le 2\sqrt{3}$. Since ρ^{-2} is summable near the diagonal in $d=3$, this part is at most $c\,\mathrm{e}^{-\delta''|y-y'|} \le c'\rho^{-1}\mathrm{e}^{-\delta''|y-y'|/2}$.
- In the part with $(1 - m^2_{j_0})$, put $r := |y-y'|$ and $\delta_0 := \min(\delta, \frac14)$, so that $\mathrm{e}^{-\delta|y-z|-|z-y'|/4} \le \mathrm{e}^{-\delta_0 r}$. On $\{|z-y'| \le 2r\}$, one of $\rho(y,z), \rho(z,y')$ is at least $\rho(y,y')/2$, and $\sum_{|w|\le 3r}\xi^3\rho(w)^{-2} \le c\rho(y,y')$; this gives $c\rho(y,y')^{-1}$. On $\{|z-y'| > 2r\}$ we have $|y-z| \ge |z-y'|/2$, and $\sum_{|w|>2r}\xi^3\rho(w)^{-4} \le c\rho(y,y')^{-1}$.

Step 4 (exact zeros). In ξ-units the order-e^2 class gives the zeroth-moment coefficient $\Pi_{\mu\mu'}$ (degree $-d+2=-1$) and the first-moment coefficient $\Pi_{\mu\mu'\nu}$ (degree $-d+3=0$). Their bubble terms are of size $\sum_{y'}\xi^d\rho^{-4}$ and $\sum_{y'}\xi^d\rho^{-4}\rho$, respectively. Replace every $G_{j_0}(0)$ by C^ξ. Then the $\Pi_{\mu\mu'\nu}$ combination is [3, (3.27)], which vanishes by the $\mu\leftrightarrow\mu'$ symmetry [3, (3.28)]. The $\Pi_{\mu\mu'}$ combination [3, (3.29)] vanishes by the Ward–Takahashi identity [3, (2.26)]; see also Lemma E.26 with $M^2=1$. (The "$M^2=1$" of [3] is the normalisation of C^ξ, not the Higgs mass.)

Step 5 (remainders). Write $\nabla_\nu f(y)=\xi^{-1}(f(y+\xi e_\nu)-f(y))$ and $\nabla^-_\nu f(y)=\xi^{-1}(f(y)-f(y-\xi e_\nu))$. For absolutely summable products, $\sum_y \xi^3(\nabla_\nu f)h=-\sum_y \xi^3 f\,\nabla^-_\nu h$, and $(\nabla^-_\nu h)(y)=(\nabla_\nu h)(y-\xi e_\nu)$.

Line bounds. By Step 3 and Lemma E.39, a line with $a\le 2$ differences, at most one at each end, obeys

$$|\partial^a C^\xi|\le c\rho^{-1-a}\mathrm{e}^{-|y-y'|/4},\qquad |\partial^a D|\le c\,h_a(\rho)\mathrm{e}^{-\delta'|y-y'|},$$

with $h_0=1$, $h_1=1+\log_+(1/\rho)$ and $h_2=\rho^{-1}$. Expand every scalar line as $G^\xi_{j_0}(0)=C^\xi+D$. The terms with C^ξ-lines only are the combinations of Step 4, which vanish. Every other term contains a line D.

First moment. Every term with a line D is at most $c\rho^{-3}\cdot\rho\,\mathrm{e}^{-\delta'|y-y'|}$, and its sum over y' is $O(1)$.

Zeroth moment. The class consists of two kinds of graph:

- the bubble: two vertices $\varphi\cdot q\nabla\varphi\,A$ at x,x', joined by two scalar lines that carry a and $2-a$ differences, at most one at each end;
- the seagull tadpole: one vertex at x with a scalar self-loop, contributing only at $x'=x$.

In the tadpole, $|D(x,x)|\le c$. In the bubble, every product containing a line D, except $(\nabla^{(1)}_\mu\nabla^{(2)}_\nu C^\xi)\cdot D$, is at most $c\rho^{-2}(1+\log_+(1/\rho))\mathrm{e}^{-\delta'|x-x'|}$, and its sum $\sum_{x'}\xi^3$ is $O(1)$.

In the exceptional product, both differences act on C^ξ, the difference of the vertex at x' acting at the x'-end. The remaining factor at x' is the constant external field. Sum by parts in x'. The difference of a constant is 0, so the difference moves to the line D, and we get

$$-\sum_{x'}\xi^3\,\nabla^{(1)}_\mu C^\xi(x,x')\,(\nabla^{(2)}_\nu D)(x,x'-\xi e_\nu).$$

Since $\rho(x,x'-\xi e_\nu)\ge\rho(x,x')/2$, this is at most $c\sum_{x'}\xi^3\rho^{-2}(1+\log_+(1/\rho))\mathrm{e}^{-\delta'|x-x'|}\le c'$. Without this summation by parts, the exceptional product is only bounded by $c\sum_{x'}\xi^3\rho^{-3}\mathrm{e}^{-\delta'|x-x'|}=O(\log(1/\xi))$.

So both coefficients are $O(1)$ in ξ-units, uniformly in ξ and in $m^2_{j_0}\in[0,1]$.

Step 6 (rescaling and the torus). [3, p. 441] gives $\Pi^{(\eta)}_{\mu\mu'}(x)=(L^{j_0}\eta)^{-d+2}\Pi^{(\xi)}_{\mu\mu'}(y)$ and $\Pi^{(\eta)}_{\mu\mu'\nu}(x)=(L^{j_0}\eta)^{-d+3}\Pi^{(\xi)}_{\mu\mu'\nu}(y)$. At $d=3$ these are the factors $\ell_{j_0}^{-1}$ and $\ell^0_{j_0}$. On T_ε the expression lives on the cube $\square$ of [3, p. 433], which does not wrap around the torus by (T.nw). The gauge-out of the background on $\square$ and the transfer to $\eta\mathbb{Z}^3$ are exact, and the image differences are external kernels of δG type of [3, (1.33)]. Only scalar lines occur, so μ_*, u and the photon zero mode do not enter.

Step 7 (the Higgs mass). The Higgs mass entered only through $m^2_{j_0}$, in Υ (Step 1) and in the line bounds (Steps 2–3), and both are uniform on $[0,1]$. For $M\le m_2$, $t_{j_0}\le\varepsilon_0$ and (E.m2), $m^2_{j_0}\le m_2^2\varepsilon_0^2\le 1$. No restriction to $M\in[m_1,m_2]$ is used. $\square$

Setting for the vertex of [3, (3.32)]. We keep the notation of Lemma E.24:

- $\xi=L^{-j_0}$, $\ell:=\ell_{j_0}=L^{j_0}\eta\le 1$ and $s_j:=L^{j-j_0}$ for $0\le j\le j_0$;
- $C^\xi=(-\Delta^\xi+1)^{-1}$ and $G:=G^\xi_{j_0}(0)$;
- $\Upsilon=1-m^2_{j_0}-a_{j_0}P_{j_0}$, and $D:=G-C^\xi=G\Upsilon C^\xi$, the *regular line*.

On $\xi\mathbb{Z}^3$ we use $\langle f,\, h\rangle = \sum_y \xi^3 f(y)\cdot h(y)$ and write kernels with respect to this measure. The scalar field is a map $\varphi : \xi\mathbb{Z}^3 \to \mathbb{R}^2$, and $q = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. For a real field a on the bonds $b = (y, y+\xi e_\nu)$, the covariant Laplacian is given by

$$\langle \varphi,\, -\Delta_a^\xi \varphi\rangle = \sum_{b=(y,y+\xi e_\nu)} \xi^3 \xi^{-2} \big|\mathrm{e}^{-q\xi a(b)}\varphi(y+\xi e_\nu) - \varphi(y)\big|^2.$$

On $\Lambda_R := \xi\mathbb{Z}^3 \cap [-R, R]^3$ the Dirichlet version $-\Delta_a^{\xi,R}$ is the same form on fields with $\varphi = 0$ off Λ_R. A gauge function χ acts by $\varphi \mapsto \mathrm{e}^{q\chi}\varphi$ and $a \mapsto a + d\chi$, where $(d\chi)(b) = \xi^{-1}(\chi(y+\xi e_\nu) - \chi(y))$.

Labels [3, pp. 419–420]. The labels are as follows:

- every internal vector leg carries a factor $t_\square g_\square$, where $(g_\square)$ is the unit-scale partition of unity of [3], $\operatorname{supp} g_\square$ lies in a cube of side 2, $|g| \le 1$, and $|\nabla g| + |\nabla\nabla g| \le c_g$;
- every λ-vertex carries the indicator of a unit cube and a parameter $s_\square$;
- every *fixed mass vertex* carries the indicator of a unit cube and a parameter $r_\square$. A fixed mass vertex is an insertion $\kappa(y)|\varphi(y)|^2$ with an a-independent coefficient: the finite counterterms of [3, (1.7), (1.29)], and the differences between counterterm lines built on the torus and on the infinite lattice.

A *label configuration* Λ is a monomial $t^m s^n r^p$.

Counterterms in graph representation. For a connected graph H with two external scalar legs at x, x', let H_{nn} be the sum of its terms in which no difference acts on an external leg. Following [3, (1.21), p. 416; p. 417; (3.7)–(3.9), p. 435], the mass counterterm generated by H is written as the graph of H_{nn}, with the same lines and indices, in which the external leg at x' is attached at x; x' is summed and the sign is -1. Within a graph, this counterterm graph is the *counterterm partner* of the graph containing H. For a member T, let $\mathcal{K}(T)$ be the set formed by T and its counterterm partners.

The class. Fix $(\alpha, \beta) \in \{(4,0), (2,1)\}$ and Λ. Let $\Pi^\Lambda_{G,\mu\nu}(b, b')$ be the sum of all connected graphs on $\xi\mathbb{Z}^3$ (the *members*) with the following properties:

- they have order $e_\xi^\alpha \lambda_\xi^\beta$, label configuration Λ, and two external vector legs, at the bond b in direction μ and at the bond b' in direction ν; the external couplings and the localisations g, g' are removed;
- every internal scalar line, including those of the counterterms in graph representation, is G;
- every internal vector line is the covariance $\mathcal{C}_{\mathrm{ph}}$ of a centred Gaussian field on the bonds that does not depend on the external field.

Let Π^Λ_C be the same sum with every internal scalar line replaced by C^ξ. For a member T and a set S of its scalar lines, T^S denotes T with the lines of S replaced by D; the other lines stay G.

Lemma E.25 (Dirichlet scalar propagator)**.** *Let $M^2 > 0$, let $C_M^\xi := (-\Delta_0^\xi + M^2)^{-1}$ on $\xi\mathbb{Z}^3$, and let $C_M^{\xi,R} := (-\Delta_0^{\xi,R} + M^2)^{-1}$ on Λ_R. Then:*

- $0 \le C_M^{\xi,R}(y,y') \le C_M^{\xi,R'}(y,y') \le C_M^\xi(y,y')$ *for $R \le R'$;*
- $C_M^{\xi,R}(y,y') \to C_M^\xi(y,y')$ *as $R \to \infty$;*
- $C_M^\xi(y,y') \le c(\xi, M)\, \theta^{|y-y'|_1/\xi}$, *with $\theta := 6/(6 + M^2\xi^2) < 1$.*

Proof. Write $-\Delta_0^\xi + M^2 = (6\xi^{-2} + M^2)(1 - T)$, where $T = (6 + M^2\xi^2)^{-1} A$ and A is the adjacency matrix of $\xi\mathbb{Z}^3$. So T has nonnegative entries and row sums $\theta < 1$, and $C_M^\xi = (6\xi^{-2} + M^2)^{-1}\xi^{-3}\sum_{n\ge 0} T^n$ entrywise.

On Λ_R the same holds with $1_{\Lambda_R} T 1_{\Lambda_R}$ in place of T. Its powers are nonnegative, nondecreasing in R, and converge entrywise to those of T. Monotone convergence gives the first two claims.

Finally, $T^n(y,y')=0$ unless $n \geq |y-y'|_1/\xi$, and $T^n(y,y') \leq \theta^n$. Hence $C_M^\xi(y,y') \leq (6\xi^{-2}+M^2)^{-1}\xi^{-3}(1-\theta)^{-1}\theta^{|y-y'|_1/\xi}$. □

Lemma E.26 (Ward–Takahashi identity at fixed labels)**.** *Let $M^2>0$, and let $F(\varphi;s,r)$ be a polynomial in the parameters and in the variables $|\varphi(y)|^2$, y in a finite set, with a-independent coefficients. Let $\mathcal{A}$ be a centred Gaussian field on the bonds, independent of a, and let $B := \sum_\square t_\square g_\square \mathcal{A}$, a finite sum. Define, as a formal power series in (t,s,r),*

$$W_R(a) := \log \mathbb{E}_{\mathcal{A}} \int_{(\mathbb{R}^2)^{\Lambda_R}} d\varphi \, \exp\Big[-\tfrac{1}{2}\langle \varphi, \, (-\Delta_{a+B}^{\xi,R}+M^2)\varphi\rangle - F(\varphi)\Big].$$

Let $\Pi_{\mu\nu}^{R,\Lambda}(b,b')$ be the second derivative in $a(b), a(b')$ at $a=0$ of the coefficient of the monomial Λ. Then:

(a) $\sum_{b' \| e_\nu} \Pi_{\mu\nu}^{R,\Lambda}(b,b') = 0$ *for every R, every b and every μ,ν;*

(b) *as $R\to\infty$, $\Pi^{R,\Lambda}(b,b')$ converges to $\Pi^\Lambda(b,b')$, the sum of the connected graphs of label configuration Λ with scalar lines C_M^ξ and vector lines the covariance of $\mathcal{A}$. Moreover, $\sum_{b'}|\Pi^\Lambda(b,b')|<\infty$ and $\sum_{b'\|e_\nu}\Pi_{\mu\nu}^\Lambda(b,b')=0$.*

Proof. (a) Each $\mathrm{e}^{q\chi(y)}$ is a rotation commuting with $\mathrm{e}^{-q\xi a(b)}$, so

$$\mathrm{e}^{-q\xi(a+d\chi)(b)}\mathrm{e}^{q\chi(y+\xi e_\nu)}\varphi(y+\xi e_\nu) - \mathrm{e}^{q\chi(y)}\varphi(y) = \mathrm{e}^{q\chi(y)}\big[\mathrm{e}^{-q\xi a(b)}\varphi(y+\xi e_\nu)-\varphi(y)\big].$$

Bonds leaving Λ_R contribute only $|\varphi(y)|^2$. Hence $\langle \mathrm{e}^{q\chi}\varphi, \, -\Delta_{a+B+d\chi}^{\xi,R}\mathrm{e}^{q\chi}\varphi\rangle = \langle\varphi, \, -\Delta_{a+B}^{\xi,R}\varphi\rangle$ for every B. Also $F(\mathrm{e}^{q\chi}\varphi)=F(\varphi)$, the map $\varphi\mapsto \mathrm{e}^{q\chi}\varphi$ is orthogonal, and the form is positive definite. So, coefficient by coefficient, $W_R(a+d\chi)=W_R(a)$.

Each coefficient is a smooth function of a, being a finite combination of Gaussian moments with covariance $(-\Delta_a^{\xi,R}+M^2)^{-1}$. Differentiate along χ at $\chi=0$ and in $a(b)$ at $a=0$; this gives the finite sum identity $\sum_{b'}\Pi^{R,\Lambda}(b,b')(d\chi)(b')=0$. Take $\chi(y)=e_\nu\cdot y$, so that $(d\chi)(b')=1$ for $b'\parallel e_\nu$ and 0 otherwise.

(b) By Wick's theorem and the linked-cluster property, $\Pi^{R,\Lambda}$ is the finite sum of the connected graphs of configuration Λ with scalar lines $C_M^{\xi,R}$ and lattice differences of them. Every vertex is labelled, and so confined to a bounded set K_Λ, or is the vertex at b or at b'. If the vertex at b' is unlabelled, it is joined to the rest by a scalar line. By Lemma E.25, each member is bounded, uniformly in R, by $c(\xi,\Lambda)\theta^{\mathrm{dist}_1(b',K_\Lambda\cup\{b\})/\xi}$, and converges pointwise as $R\to\infty$. Dominated convergence in b' and (a) conclude. □

The identity does not apply with G in place of C_M^ξ: the term $a_{j_0}P_{j_0}$ is not gauge covariant without the vertices [3, (1.13)–(1.15)]. The insertion F may contain the fixed mass vertices and the counterterms in graph representation evaluated with C^ξ-lines, so the identity holds separately for each monomial in r. No decay of $\mathcal{C}_{\mathrm{ph}}$ is used.

Proposition E.27 (exact form of the coefficient of [3, (3.32)])**.** *Let $0\leq m_{j_0}^2\leq 1$. For a label configuration Λ, write the zeroth-moment term of* [3, (3.31)] *for the class of order $(\alpha,\beta)\in\{(4,0),(2,1)\}$ as $e(t_k)^\alpha\lambda(t_k)^\beta\sum_x \eta^3 g(x)A_\mu(x)\, c_{\mu\mu'}^{(4),\Lambda}(x)\, g'(x)A'_{\mu'}(x)$. Let $b_\mu(y)$ be the bond at y in direction μ, and put $y=\ell^{-1}x$.*

(a) $c_{\mu\mu'}^{(4),\Lambda}(x) = \hat{c}_{\mu\mu'}^\Lambda(y) := \sum_{b'\|e_{\mu'}} \xi^3\big[\Pi_G^\Lambda - \Pi_C^\Lambda\big]_{\mu\mu'}(b_\mu(y),b')$, *and* $\sum_{b'\|e_{\mu'}}\xi^3\Pi_{C,\mu\mu'}^\Lambda(b,b')=0$.

(b) $\hat{c}_{\mu\mu'}^\Lambda(y) = \sum_T\sum_{\emptyset\neq S}(-1)^{|S|+1}\sum_{b'\|e_{\mu'}}\xi^3 T_{\mu\mu'}^S(b_\mu(y),b')$, *where T runs over the members of Π_G^Λ and S over the non-empty sets of scalar lines of T, and all sums converge absolutely.*

Proof. Rescaling. Put $x=\ell y$ and $\eta=\ell\xi$. In $d=3$, $\varphi_\eta(x)=\ell^{-1/2}\varphi_\xi(y)$ and $A_\eta(x)=\ell^{-1/2}A_\xi(y)$. Then

- $\eta^3\lambda|\varphi_\eta|^4 = \xi^3(\lambda\ell)|\varphi_\xi|^4$;
- $\eta^3 M^2 t_k^2|\varphi_\eta|^2 = \xi^3(M^2 t_{j_0}^2)|\varphi_\xi|^2$, since $t_k\ell = t_{j_0}$;
- $e\,\eta A_\eta = (e\ell^{1/2})\xi A_\xi$.

So $e_\xi^2 = e(t_k)^2\ell$, $\lambda_\xi = \lambda(t_k)\ell$, and the rescaled mass is $m_{j_0}^2$. A local coefficient of $A\cdot A'$ satisfies $c_\eta = \ell^{-2}c_\xi$. Writing $c_\xi = e_\xi^\alpha \lambda_\xi^\beta \hat{c}$ gives $c_\eta = e(t_k)^\alpha \lambda(t_k)^\beta \ell^{\alpha/2+\beta-2}\hat{c}$. The exponent is -1 for $(2,0)$, which is the factor $(L^{j_0}\eta)^{-d+2}$ of [3, p. 441], and 0 for $(4,0)$ and $(2,1)$.

Resummation. By [3, (3.31) and p. 442], the coefficient is the zeroth moment of the class, summed over the orderings in which every internal line precedes the external legs, and over the corresponding indices. The transformations performed beforehand on proper divergent subgraphs [3, pp. 444–445] are exact identities (Taylor's formula with remainder, and resummation of indices), and the orderings in which a given subgraph is the last divergent one partition all orderings. So the zeroth moment is that of the untransformed class. Summing the slice decomposition [3, (2.6)] over the indices $< j_0$ turns every internal scalar line, including those of the counterterms in graph representation, into G. The remaining mass insertions are the fixed mass vertices.

Cancellation. In Π_C^Λ, every counterterm in graph representation is an a-independent insertion $\kappa(x)|\varphi(x)|^2$. Lemma E.26 with $M^2 = 1$ gives $\sum_{b'} \xi^3 \Pi_C^\Lambda(b,b') = 0$. This proves (a).

(b) The members of Π_G^Λ and Π_C^Λ correspond one to one. Since $C^\xi = G - D$, for each member $\prod_l G_l - \prod_l C_l^\xi = \sum_{\emptyset\neq S}(-1)^{|S|+1}\prod_{l\in S} D_l \prod_{l\notin S} G_l$. At fixed ξ, G and D decay exponentially (Lemma E.24, Steps 2–3), so the sums converge absolutely, as in the proof of Lemma E.26(b). □

Definition E.28 (generalised graphs)**.** A *generalised graph* Γ is a finite connected multigraph $(V,\mathcal{L})$ with $\mathcal{L}\neq\emptyset$ and a root $v_0\in V$; self-loops are allowed. Each line l has endpoints u_l, w_l and a dimension $\omega_l\in\mathbb{R}$, and each vertex v has an exponent $p_v \geq 0$. For $\emptyset \neq \mathcal{L}_H \subset \mathcal{L}$, let V_H be the set of endpoints of the lines of $\mathcal{L}_H$, and $H = (V_H, \mathcal{L}_H)$. The *degree* of H is

$$D(H) := \sum_{l\in\mathcal{L}_H} \omega_l + d(|V_H| - 1) + \sum_{v\in V_H} p_v.$$

For $\delta > 0$, $N \geq 0$, $s_j := L^{j-N}$ and $\mathbf{j} = (j_l) \in \{0,\dots,N\}^{\mathcal{L}}$, put, with y_{v_0} fixed,

$$E_{\mathbf{j}}(\Gamma) := \sum_{(y_v)_{v\neq v_0}\in(\xi\mathbb{Z}^d)^{V\setminus v_0}} \prod_{v\neq v_0} \xi^d \prod_v \xi^{p_v} \prod_l s_{j_l}^{\omega_l} \mathrm{e}^{-\delta|y_{u_l}-y_{w_l}|/s_{j_l}}, \qquad \xi = L^{-N}.$$

Let $\Gamma_i(\mathbf{j})$ be the subgraph of the lines with $j_l \leq i$.

To a member T^S we associate the generalised graph Γ_T: the graph of T, rooted at the vertex of b, with p_v the number of factors ξ of v besides ξ^3. A scalar or vector line with a_l differences, not in S, has dimension $-1-a_l$; a line of S has dimension $1-a_l$. For a connected subgraph H, let $n_D(H)$ be the number of lines of S in H, and let $D^{\mathrm{st}}(H)$ be its degree with every line counted at dimension $-1-a_l$. Then $D(H) = D^{\mathrm{st}}(H) + 2n_D(H)$. H is *critical* if $n_D(H) = 0$ and $D^{\mathrm{st}}(H) \leq 0$. A *chain* is a member of the form

$$\sum_{z,z'} \xi^6\, \Pi^1_{\mu\nu}(b,z) g_1(z)\, \mathcal{C}_{\mathrm{ph},\nu\nu'}(z,z')\, g_2(z')\, \Pi^2_{\nu'\mu'}(z',b'),$$

with Π^1, Π^2 in the order-one class $\mathcal{P}$ of the bubbles and the seagull tadpole (Lemma E.24, Step 5).

Lemma E.29 (domination by scale sums)**.** *Let $\rho = |y-y'| + \xi$ and $0 < \delta_1 \leq \delta'$. Put $f_\omega(\rho) = \rho^\omega$ if $\omega < 0$, $f_0(\rho) = 1 + \log_+(1/\rho)$, and $f_\omega = 1$ if $\omega > 0$. Then*

$$f_\omega(\rho)\mathrm{e}^{-\delta'|y-y'|} \leq \mathrm{e}^{\delta_1}(1+\log L) L^{|\omega|} \sum_{j=0}^{j_0} s_j^\omega \mathrm{e}^{-\delta_1|y-y'|/s_j}.$$

In particular, by Lemma E.24, Step 3, the regular line D with $a \in \{0,1,2\}$ differences, at most one at each end, is dominated by a sum over j of lines of dimension $1-a$, two more than a slice line with a differences.

Proof. If $\rho > 1$, then $f_\omega(\rho) \le 1$ and the term $j = j_0$ is at least $\mathrm{e}^{-\delta'|y-y'|}$.

Let $\rho \le 1$, and let $J = \{j : s_j \ge \rho\} = \{j_*, \dots, j_0\}$. This set is non-empty, since $s_{j_0} = 1$, and $s_{j_*} < L\rho$, since $s_0 = \xi \le \rho$. For $j \in J$, $\mathrm{e}^{-\delta_1|y-y'|/s_j} \ge \mathrm{e}^{-\delta_1}$.

- For $\omega > 0$, use the term j_0.
- For $\omega < 0$, use $s_{j_*}^\omega \ge L^{-|\omega|}\rho^\omega$.
- For $\omega = 0$, use $|J| \ge \max(1, \log(1/\rho)/\log L) \ge (1+\log(1/\rho))/(1+\log L)$.

□

Lemma E.30 (degrees in the class)**.** *Let T be a connected graph of order $o_T = 2$ (the order of e_ξ is $\frac12$ and that of λ_ξ is 1) with two external vector legs. Suppose T is built, in ξ-units at zero background, from the vertices*

- $\xi^{n-1}\varphi \cdot q^n \partial\varphi\, a^n$, $n \ge 1$ *(order $n/2$, one difference);*
- $\xi^{n-2}\varphi \cdot q^n \varphi\, a^n$, $n \ge 2$ *even (order $n/2$);*
- $\lambda|\varphi|^4$ *(order 1);*
- $c(y)|\varphi(y)|^2$ *with $|c| \le \bar c$ (order $o_v \ge \frac12$).*

Suppose also that its lines are proportional to 1_2 in the charge index. For a connected subgraph H, let E_H be its number of external legs and o_H its order.

(a) *If some connected subgraph without external scalar legs has an odd number of external vector legs, then $T^S = 0$ for every S.*

(b) *Otherwise every connected subgraph satisfies $D^{\mathrm{st}}(H) \ge o_H + E_H/2 - 3 \ge -1$, and $D^{\mathrm{st}}(T) \ge 0$.*

(c) *Hence $D(H) \ge 1$ whenever $n_D(H) \ge 1$, and $D(T) \ge 2$ if $S \ne \emptyset$.*

Proof. Let $\deg_v$ be the number of legs at v, and δ_v the number of differences at v acting on lines of H. Since $2|\mathcal{L}_H| = \sum_v \deg_v - E_H$,

$$D^{\mathrm{st}}(H) = \sum_{v \in V_H} c_v + \tfrac{E_H}{2} - 3, \qquad c_v := 3 + p_v - \tfrac{\deg_v}{2} - \delta_v.$$

For the four vertex types, $c_v = \frac{n}{2} + 1 - \delta_v$, $\frac{n}{2}$, 1 and 2, respectively. So $c_v \ge o_v$; for the fourth type $c_v = 2 > o_v$, since T has a vertex with a vector leg besides it.

(a) Each vertex has an even number of scalar legs. The charge indices of H run along closed scalar loops, and a loop through vertices with n_v vector legs carries $\pm \operatorname{tr} q^m$, $m = \sum n_v$, times scalars. Since $\sum_v n_v \equiv E_H \pmod 2$, some loop has m odd, and $\operatorname{tr} q^m = 0$.

(b) The first inequality follows from $c_v \ge o_v$. We have $E_H \ge 1$, and $E_H = 1$ is excluded by (a) and the parity of the scalar legs. If $E_H = 2$ and $|V_H| \ge 2$, then $o_H \ge 1$. If $E_H = 2$ and $V_H = \{v\}$, then $\deg_v \ge 4$, so $o_v \ge 1$. If $E_H \ge 3$, then $o_H \ge \frac12$. For T, every difference acts on internal legs, so $D^{\mathrm{st}}(T) \ge o_T - 2 = 0$.

(c) Use $D = D^{\mathrm{st}} + 2n_D$. □

Lemma E.31 (members without critical subgraphs)**.** *Let T be a member built from the vertices of Lemma E.30, let $S \ne \emptyset$, and suppose T^S has no critical subgraph. Then $\sum_{b'} \xi^3 |T^S(b,b')| \le C$. Here C depends only on d, a, L, the block size, $\bar c$, the constants of the slice bounds of the scalar and vector lines, and the graph of T; it is uniform in $\xi, j_0, k, \varepsilon, \ell_T$ and $m_{j_0}^2 \in [0,1]$.*

Proof. If T falls under Lemma E.30(a), then $T^S = 0$. Otherwise take absolute values after the charge sums, and bound each vertex factor by its power of ξ, using $|g| \le 1$ and $|c| \le \bar c$. Decompose the lines not in S into slices, which obey $|\partial^a G_{(j)}(y,y')| \le c\, s_j^{-1-a} \mathrm{e}^{-\delta_1|y-y'|/s_j}$ by Theorem E.47 and

Lemma E.38 (the vector slices with weights in $[0,2]$). Dominate the lines of S by Lemma E.29. This bounds the sum by a constant times $\sum_{\mathbf{j}} E_{\mathbf{j}}(\Gamma_T)$, with $\delta = \delta_1$, where the vertex of b' is summed or coincides with the root. By Lemma E.30(c) and the hypothesis, every connected subgraph has positive degree. Statement K.10(a) concludes. □

Lemma E.32 (coefficient bound for the vertex of [3, (3.32)])**.** *Let $0 \le m_{j_0}^2 \le 1$, and let the internal photon lines be the weighted fluctuation lines $(1+u\delta_j)H_j$ of the slice model (Lemma E.4), with weights at most 2. For a label configuration Λ, write the zeroth-moment term of* [3, (3.31)] *for the class of order e^4 or $e^2\lambda$ as*

$$e(t_k)^{\alpha}\lambda(t_k)^{\beta}\sum_{x}\eta^{d}\,g(x)A_{\mu}(x)\,c_{\mu\mu'}^{(4),\Lambda}(x)\,g'(x)A'_{\mu'}(x), \qquad (\alpha,\beta)\in\{(4,0),(2,1)\}.$$

Then $|c_{\mu\mu'}^{(4),\Lambda}(x)| \le \Gamma_4\,\ell_{j_0}^{0}$. The constant Γ_4 depends only on:

- *d, on a (through $a_\flat$), and on the block size;*
- *the constants of the slice bounds of the scalar lines and of the photon lines;*
- *the bound $\bar{c}_{\mathrm{fin}}$ on the rescaled fixed mass vertices* [3, (1.7)] *that occur in the class.*

It is uniform in $\xi, j_0, k, \varepsilon, \ell_T$, Λ and $m_{j_0}^2 \in [0,1]$, and it does not depend on μ_*, u or the photon zero mode.

Proof. (i) Rescaling. By Proposition E.27(a), $c^{(4),\Lambda}(x) = \hat{c}^{\Lambda}(y)$ with the factor $\ell_{j_0}^{0}$. The torus enters as in Lemma E.24, Step 6, and through the fixed mass vertices. So it suffices to show $|\hat{c}^{\Lambda}| \le \Gamma_4$.

(ii) Exact form. By Proposition E.27(b), $\hat{c}^{\Lambda}$ is the alternating sum of the terms $\sum_{b'}\xi^{3}T^{S}$ with $S \neq \emptyset$. A counterterm partner carries the lines of its subgraph, so S selects the same lines in a member and in its partners. Photon lines are not replaced.

(iii) The pure C^{ξ} part. The part in which every scalar line is C^{ξ} has been subtracted exactly, by Lemma E.26 at fixed labels with the fixed mass vertices and the counterterms in graph representation as gauge-invariant insertions.

(iv) The remainder terms. Every term T^S contains at least one regular line. By Lemma E.29, a regular line has dimension 2 higher than a slice line, so it raises by 2 the degree of every subgraph containing it. By Lemma E.30(c), every subgraph containing a regular line has degree ≥ 1, and the whole graph has degree ≥ 2.

If T^S has no critical subgraph, Lemma E.31 bounds it by $O(1)$ uniformly in ξ; this uses only the power-counting estimate of [3, Props. 2.1–2.2] for generalised graphs (Statement K.10). The critical subgraphs contain no regular line. Their lines are slices of $G_{j_0}^{\xi}(0)$ and of the photon line, which is the setting of the inner transformations of [3, pp. 425, 434–445]. These transformations are made inside the following groupings of the remainder:

- the chains are grouped over both order-one factors;
- every other member is grouped with its counterterm partners at fixed S.

The resulting bounds are Statement K.12, and the assembly of all terms into $|\hat{c}^{\Lambda}| \le K_R$ is Statement K.11. The line bounds are uniform in $m_{j_0}^2 \in [0,1]$ for the scalar lines, and independent of μ_* for the photon lines. Members containing a fixed mass vertex are multiplied by its rescaled coefficient, at most $\bar{c}_{\mathrm{fin}}$. Hence $|\hat{c}^{\Lambda}| \le \Gamma_4 := K_R$.

(v) Higgs-mass dependence. The Higgs mass M enters only through $m_{j_0}^2 = M^2 t_{j_0}^2 \le m_2^2\varepsilon_0^2 \le 1$ (by (E.m2)) and through $\bar{c}_{\mathrm{fin}}$. So $\Gamma_4 = \Gamma_4(M)$ may depend on M through $\bar{c}_{\mathrm{fin}}$, which is finite at every $M \in (0, m_2]$ by Lemma A.6(a). If the class contains no fixed mass vertex, Γ_4 is uniform in $M \in (0, m_2]$. Uniformity as $M \downarrow 0$ is not claimed and is not used; only the finiteness at the four masses is used. □

The tail beyond the order $\bar{n}$. [2, p. 579] states that, in the expansions of "the basic quadratic form in the ψ, ϕ fields and the polynomial $\mathcal{P}^{(k)}$", "the terms with R-vertices, the terms containing

the configuration $\tilde{B}^{(k)}$ and the terms of order $> \bar{n}$" are $O((L^k\varepsilon)^\kappa)|\Lambda_2^{(k)}|$. The log-determinant expansions are not on this list; they are cut at [2, pp. 577–578]. We write the order-$> \bar{n}$ part.

Setting. This is the k-th step, $0 \le k \le K-1$, with $t := t_k$, on the support of the small-field restrictions χ_k of [2, (2.55)]. Here $|A'| \le O(1)p$ on $\Lambda_1^{(k)}$, and the scalar fields and their covariant differences are $\le Cp\lambda(t)^{-1/4}$. Write:

- $\mathfrak{B} := B^k(\Lambda_0^{(k)})\tilde{B}$ for the regular background of [2, p. 579];
- $\tilde{A} := \theta_{k+1}A'^{(k)} + \tilde{B}^{(k)}$ for the expansion field;
- $\sigma := e(t)(\|\tilde{A}\|_\infty + \|\partial^\eta \tilde{A}\|_\infty)$, so that $\sigma \le C_\sigma t^{1/2}p$ with $C_\sigma = eC_A$;
- $N_k := |T|/t_k^3$ for the number of t_k-blocks of T_ε, so that $|\Lambda_2^{(k)}|_k \le N_k$.

The field A enters only through:

- $G_k(\Omega, \mathfrak{B} + \tilde{A})$, on $\Omega = B^k(\Lambda_0^{(k)})$ for the basic quadratic form and on $\Omega' = B^{k-1}(\Lambda_2^{(k-1)})$ for the lines of $\mathcal{P}^{(k)}$, whose vertices lie in $\Omega_1 = B^k(\Lambda_7^{(k-1)\prime})$;
- $Q_k(\mathfrak{B} + \tilde{A})$;
- the transports and covariant differences in the vertices of $\mathcal{P}_w^{(k)}$.

Every coefficient is an entire function of $e(t)\tilde{A}$.

Inputs. (K1)–(K4), (IH) and (CAP) are stated in Subsection E.3; (K3) is proved as Lemma E.117. (REG), ($\tilde{B}$) and (PO) are bounds of [1, 2] that are used as stated.

- (K1) The bounds of [4, (1.8), (1.10), (1.12)] for $G_k(\Omega, \mathfrak{B})$ on a regular background: decay, first derivative, α-Hölder quotient of the derivative, and decay of $G_kQ_k^*$. These hold with $0 \le M^2t^2 \le c$ in G_k and constants uniform there.
- (K2) The slice bounds [3, (2.10)–(2.12)] for the lines of the families of $\mathcal{P}^{(k)}$.
- (K3) The one- and two-variable remainder bounds (Proposition E.57). For $\mathcal{K} = [G(\mathfrak{B})V]^n G(\mathfrak{B} + \tilde{A})[VG(\mathfrak{B})]^{n'}$ with $n + n' = m$, the sup, the derivative and the α-Hölder quotient of $\mathcal{K}f$ are $\le C(C_\infty\sigma)^m \mathrm{e}^{-\beta \operatorname{dist}(x, \operatorname{supp} f)}\|f\|_\infty$. Here V has the right-pushed form $R_0 + \sum_\mu R_{1,\mu}D_\mu$ with $\|R\| \le c_V\sigma$. On Neumann domains this needs internal points at distance $\ge R_0$ from Ω^c and $R \ge 12\sqrt{d}$.
- (K4) The weighted ℓ^2 bounds of Lemma E.58.
- (REG) $\mathfrak{B}$ is regular for [1, Prop. 2.1], under the upper bounds on ε_0 of Table 7, among them (E.1) and (E.ksp3), read with the constant C_A.
- ($\tilde{B}$) $\|\tilde{B}^{(k)}\|_\infty + \|\partial^\eta \tilde{B}^{(k)}\|_\infty \le C_\kappa t^\kappa$ for every κ; and $\theta_{k+1}A'^{(k)} = O(p)$, with its derivative.
- (IH) Every localised family of $\mathcal{P}_w^{(k)}$ obeys [3, (1.33)] with the factor $(|u|c_k)^m$, $c_k \le 1/3$.
- (CAP) The same bound holds for families of graded order $\le \bar{N} = \bar{n}^2 = 169$, with $O(1) = O(\alpha_0, \bar{N})$ and decay $\delta(\bar{N}) > 0$.
- (PO) An elementary vertex of coupling order q_v in a connected graph with at least two vertices gives $\le Ct^{q_v/4}\cdot$polylog.

Lemma E.33 (regularity of the backgrounds)**.** *Let $0 \le k \le K-1$, $t := t_k$, $\ell(t) := 1 + \log t^{-1}$, $p(t) = b_0\ell(t)^{p_0}$ with $p_0 = 6$, $e(t) = et^{1/2}$, and let $\mathfrak{B} := B^k(\Lambda_0^{(k)})\tilde{B}$ and $\tilde{A}$ be the background and the expansion field of the setting above. Write ∂^η for the lattice derivative of the scale-k lattice, and $\theta(A) := e(t)\|\partial^\eta A\|_\infty$. Work on the support of the small-field restrictions χ_k.*

(a) *There is $C_{\mathrm{reg}} = C_{\mathrm{reg}}(d, L, a, M_b, R, c_1, \mu_*)$ such that $|(\partial_\mu^\eta \mathfrak{B})(x)| \le C_{\mathrm{reg}}p(t)$ and $|(\partial_\mu^\eta(\mathfrak{B} + \tilde{A}))(x)| \le C_{\mathrm{reg}}p(t)$ at every point x of the domain on which these configurations are defined.*

(b) *For every $s \in [0,1]$, $|\partial_\mu^\eta(\mathfrak{B} + s\tilde{A})| \le C_{\mathrm{reg}}p(t)$.*

(c) *Let $\beta = 1/56$ and c_{reg} be the constants of* [1, (2.23), p. 610]*, let ϑ_R be the constant of Lemma D.7, and let C_A be the constant of the setting above, so that $\sigma \le |e|\, C_A t^{1/2}p(t)$.*

Suppose that for all $t \in (0, \varepsilon_0]$

(Reg1) $C_{\mathrm{reg}}\,|e|^{1-\beta} t^{(1-\beta)/2} p(t) \le c_{\mathrm{reg}}$, *(Reg2)* $|e| \max\{C_{\mathrm{reg}}, C_A\}\, t^{1/2} p(t) \le \min\{\vartheta_R, 1\}$.

Then every $\mathfrak{B} + s\tilde{A}$, $s \in [0,1]$, *is regular in the sense of* [1, (2.23)], *and* $\theta(\mathfrak{B}) \le \vartheta_R$, $\theta(\mathfrak{B} + \tilde{A}) \le \vartheta_R$, $\sigma \le 1$.

(d) *For* $a \in (0,1)$ *the function* $F_a(t) := t^a \ell(t)^{p_0}$ *is increasing on* $(0, \exp(1 - p_0/a)]$, *and* $F_a(t) \to 0$ *as* $t \downarrow 0$. *Hence (Reg1) and (Reg2) hold on* $(0, \varepsilon_0]$ *as soon as they hold at* $t = \varepsilon_0$ *and* $\varepsilon_0 \le \exp(1 - 2p_0/(1-\beta))$. *The admissible* ε_0 *form an interval* $(0, \varepsilon_{\mathrm{reg}}]$ *with* $\varepsilon_{\mathrm{reg}} > 0$, *and* $\varepsilon_{\mathrm{reg}}$ *does not depend on the Higgs mass.*

Proof. (a) For $\mathfrak{B}$ and for the configuration (2.93) of [2, p. 576], of which $\mathfrak{B} + \tilde{A}$ is the case $j = k$, this is [2, (2.94)–(2.98), pp. 576–577]. There the derivative of each shell term is bounded through [2, Lemma 2.3, p. 571], and on the θ-slices through (2.96)–(2.97). The bound is pointwise, so it holds on every subdomain. For $\mathfrak{B}$ it is stated at [2, p. 579]. (b) $\mathfrak{B} + s\tilde{A} = (1-s)\mathfrak{B} + s(\mathfrak{B} + \tilde{A})$, and the bound in (a) is preserved under convex combinations. (c) By (a), (b) and (Reg1), $|\partial_\mu^\eta(\mathfrak{B} + s\tilde{A})| \le C_{\mathrm{reg}} p(t) \le c_{\mathrm{reg}}(|e|t^{1/2})^{\beta-1} = c_{\mathrm{reg}}|e(t)|^{\beta-1}$, which is [1, (2.23)]. By (a) and (Reg2), $\theta(\mathfrak{B}) \le |e| C_{\mathrm{reg}} t^{1/2} p(t) \le \vartheta_R$, and likewise for $\mathfrak{B} + \tilde{A}$. By (Reg2), $\sigma \le |e| C_A t^{1/2} p(t) \le 1$. (d) Since $\frac{d}{dt}\ell(t) = -1/t$, $\frac{d}{dt}\log F_a(t) = \frac{1}{t}\big(a - \frac{p_0}{\ell(t)}\big)$, which is positive if and only if $\ell(t) > p_0/a$, that is, $t < \exp(1 - p_0/a)$. Also $F_a(t) \to 0$ as $t \downarrow 0$. (Reg1) is $F_{(1-\beta)/2} \le c_{\mathrm{reg}}/(C_{\mathrm{reg}}|e|^{1-\beta} b_0)$ and (Reg2) is $F_{1/2} \le \min\{\vartheta_R, 1\}/(|e| \max\{C_{\mathrm{reg}}, C_A\} b_0)$. Since $\exp(1-2p_0/(1-\beta)) \le \exp(1-2p_0)$, both functions are increasing on $(0, \varepsilon_0]$ when $\varepsilon_0 \le \exp(1 - 2p_0/(1-\beta))$, so their suprema there are their values at ε_0. None of the constants involves the Higgs mass. □

Lemma E.34 (power count per coupling order)**.** *Let* $t = t_k \le \varepsilon_0 \le 1$, $\hat{p} := \max\{1, p(t)\}$, $\lambda(t) := \lambda t$, *and for constants* $c_\Phi, c_A > 0$ *put* $X_s := \max\{1, c_\Phi p(t) \lambda(t)^{-1/4}\}$ *and* $X_v := \max\{1, c_A p(t)\}$. *Let* F *be a connected localised family of* [3, Prop. 1] *with at least two vertices. Its vertices are those of* [3, (1.6)–(1.11)] *and the products* [3, (1.18)], *each product counted as one vertex. The expressions* [3, (1.12)–(1.15)] *occur only inside these products, and* [3, (1.12)] *is an external scalar field. A vertex* v *has e-order* a_v, λ*-order* b_v, $q_v := a_v + b_v$, *and* s_v *external scalar and* n_v *external vector legs. We call* v exceptional *if* $a_v = 1$, $b_v = 0$ *and* $s_v = 2$. *Put* $q := \sum_v q_v$, *and let* $d(F)$ *be the localisation length of* [3, (1.33)]. *Assume:*

(PC1) $|E(F)| \le C_F\, \mathrm{e}^{-\delta d(F)} \prod_v \Phi_v$, *where* $\Phi_v := w_v X_s^{s_v} X_v^{n_v}$ *with* $w_v = e(t)^{a_v} \lambda(t)^{b_v}$ *for a vertex that is neither a mass vertex nor exceptional;*

(PC2) *every external scalar field has norm* [3, (1.32)] *at most* X_s, *and every external vector field at most* X_v;

(PC3) *for a mass vertex* [3, (1.7)], *the piece of* $\delta m^2(\cdot, x) t^2$ *of orders* (a, b) *is bounded by* $C_\delta e(t)^a \lambda(t)^b (1 + K - k)$, *and* Φ_v *is this bound times* $X_s^{s_v}$;

(X) *for an exceptional vertex,* $\Phi_v \le C_{\mathrm{ex}}\, t^{1/4} \hat{p}^{\,2}$.

Then

$$|E(F)| \le C(q)\, C_F\, t^{q/4} \hat{p}^{\,4q} (1 + K - k)^q \mathrm{e}^{-\delta d(F)}, \qquad 1 + K - k \le 1 + \frac{\ell(t)}{\log L},$$

where $C(q)$ *depends on* $q, e, \lambda, c_\Phi, c_A, C_\delta, C_{\mathrm{ex}}$.

Proof. Every vertex has $q_v \ge 1$. For a product [3, (1.18)], the factor [3, (1.14)] or [3, (1.15)] carries $e(t)^{n+n'}$ with $n + n' \ge 1$, or $e(t)^{n+\bar{n}+1}$. For a mass vertex $q_v \ge 1$ follows from (PC3). Every vector leg carries one factor e, so $n_v \le a_v$. Since F is connected and has at least two vertices, every vertex has an internal leg. Since $t \le 1$, $X_s \le \max\{1, c_\Phi \lambda^{-1/4}\} \hat{p}\, t^{-1/4}$ and $X_v \le \max\{1, c_A\} \hat{p}$. For a vertex that is neither a mass vertex nor exceptional this gives $\Phi_v \le C_v\, t^{a_v/2 + b_v - s_v/4} \hat{p}^{\,s_v + n_v}$. Moreover $a_v/2 + b_v - s_v/4 \ge q_v/4$ holds if and only if $a_v + 3b_v \ge s_v$.

- For [3, (1.6)]: $a_v = 0$, $b_v = 1$, and one of the four scalar legs is internal, so $s_v \le 3$.

- For a mass vertex: $s_v \le 2$. A piece of orders (a,b) with $b \ge 1$ has $a + 3b \ge 3$. A piece with $b = 0$ has $a \ge 2$, because its e-order is the number of contracted vector legs, which is even. So $\Phi_v \le C\, t^{q_v/4} \hat{p}^{\,2}(1 + K - k)$.
- For [3, (1.8)–(1.11)] and for the products [3, (1.18)]: $b_v = 0$ and $s_v \le 2$. If $a_v \ge 2$, then $s_v \le a_v$. If $a_v = 1$ and $s_v \le 1$, the criterion holds. The remaining case is the exceptional one, and (X) gives $t^{1/4} = t^{q_v/4}$.

The number of legs is $s_v + n_v \le 4q_v$ in every case, so $\Phi_v \le C_v\, t^{q_v/4} \hat{p}^{\,4q_v}(1 + K - k)^{q_v}$. Take the product over v. Finally, $K - k = \log_L(\varepsilon_0/t) \le \ell(t)/\log L$. □

Lemma E.35 (the field-independent part)**.** *For fixed domains, the vacuum term $-E_1(\cdot, \Omega_1; u)$ of $\mathcal{P}_w^{(k)}$ [3, (1.4), (1.30)] is a constant whose coupling order is at most $\bar{n}$. It belongs to $\mathcal{Q}_k$ in Proposition E.36. Every term of Δ_k^{pre} contains at least one factor that depends on $\tilde{A}$: an $\tilde{A}$-leg, a vertex $V^{[r]}$, $V^{>}$ or V^{R}, or a factor $\mathcal{K}_m$. Consequently, at every k, including $k = 0$, the proof of Proposition E.36 uses the family bound of Definition E.59 only for families with at least one external field.*

Proof. By [3, (1.30)–(1.31), p. 419], $E_1 = E_{\mathrm{fin}} + E_{K,k} + E_k(\cdot, \Omega_1)$ is a sum of vacuum graphs. Their lines are the zero-background propagators $G_K(0)$, $G_{K,k}(0)$, $G_k(0)$, C^{ε}, with each vector line replaced by its u-version, and their localising function g_k is fixed by Ω_1. So E_1 does not depend on A', ψ, ϕ, $\tilde{A}$ or $\mathfrak{B}$. It enters the exponent in [3, (1.4)] additively, and by [3, (1.5)] only its coefficients of orders $1 \le \alpha + \beta \le \bar{n}$ enter $\mathcal{P}_w^{(k)}$. The expansion in the proof of Proposition E.36 acts on a sum, not on its exponential, so this constant is never multiplied by another term. It is not expanded, and by the stop rule (E3) it belongs to $\mathcal{Q}_k$. Every term of $\mathcal{P}_w^{(k)}$ has order at most $\bar{n}$. A term of Δ_k^{pre} has order larger than $\bar{n}$, so it contains a factor created by (E1) or (E2), and every such factor depends on $\tilde{A}$. □

Proposition E.36 (the tail beyond the order $\bar{n}$)**.** *Let $M \in (0, m_2]$ and $u \in \{1, -\eta\}$. Let the inputs above hold, with the power count (PO) in the form of Lemma E.34. Let the conditions (Reg1) and (Reg2) of Lemma E.33 hold, and assume the bound of Statement K.16. Pointwise on* $\operatorname{supp}\chi_k$*, for $0 \le k \le K - 1$,*

$$(\textit{basic quadratic form} + \mathcal{P}_w^{(k)})(\mathfrak{B} + \tilde{A}) = \mathcal{Q}_k + \Delta_k^{\mathrm{pre}},$$
$$|\Delta_k^{\mathrm{pre}}| \le C_{579}\, t_k^{7/2}\, \mathrm{polylog}(t_k)\, |\Lambda_2^{(k)}|_k \le C'_{579}\, t_k^{16/5} N_k.$$

Here $\mathcal{Q}_k$ is a polynomial in $\theta_{k+1}A'^{(k)}$, ψ, ϕ of order $\le \bar{n}$ and degree $\le D(\bar{n})$, and the identity is exact (a two-sided bound). The constant is $C'_{579} = C(\bar{n}, \bar{N}, \alpha_0, d, L, a, M_b, R, r, e, \lambda, \mu_*, b_0, p_0, c_1, \delta_0, \eta)$. It does not depend on ε, k or ℓ_T: where the final mass counterterm vertices occur, it is evaluated at the torus floor. It depends on M through the δm^2 vertices of $\mathcal{P}_w^{(k)}$. These are bounded at the last scale by Proposition E.9 and at the intermediate scales by a bound free of the mass. It also depends on M through $M^2 t^2 \in [0,1]$ in (K1) and (K3), with constants uniform there.

Proof. Step 0 (the configuration and the backgrounds). For $1 \le k \le K - 1$, the field $\tilde{A}$ is defined through the decomposition [2, (2.86)–(2.89), p. 575] of $\theta_{k+1}A^{(k)}$. For $L = 2$ this decomposition holds identically in A' and B by Statement K.13. Its two conditions hold: $r(\varepsilon_0) \ge 8M_b + 8 \ge 2M_b$ by (E.10), and $R \ge 107 > \sqrt{3}/\log 2$. For $k = 0$ the expansion field is the field A'' of [2, (2.19)–(2.22), p. 561], on the background $\tilde{B}^{(1)}$. The cut-off ζ there equals 1 for $|x - y| \le \frac{1}{2} r(\varepsilon)$ and 0 otherwise. Since the lattice spacing is 1 at $k = 0$, its first lattice differences are bounded by 1 and its second by 2. The second and third terms of $A'' - A'$ have the forms $\theta_1 \zeta\, C^{(0)} Q^*\, \delta C_{\Lambda_0}$ and $(1 - \zeta) C^{(0)} Q^*$. The bounds $(\tilde{B})$ of the inputs hold for them by the same argument, with $G_k Q_k^*$ replaced by $C^{(0)} Q^*$. By Lemma E.33, every $\mathfrak{B} + s\tilde{A}$, $s \in [0,1]$, is regular in the sense of [1, (2.23)],

so (K1) applies at these backgrounds. Moreover $\theta(\mathfrak{B}), \theta(\mathfrak{B}+\tilde{A}) \le \vartheta_R$ and $\sigma \le 1$, which are the smallness hypotheses under which Proposition E.57 is applied below.

Step 1 (expansion with a stop rule).

(E1) Taylor-expand every exponential e^{iz} (with $z = qe(t)\eta\tilde{A}(b)$ or $z = qe(t)\tilde{A}(\Gamma)$) to order $\bar{n}$. This splits the bracket V_k of [1, (3.44)] into three parts: pieces $V^{[r]}$ homogeneous of degree $r \le \bar{n}$ in $e(t)\tilde{A}$; polynomial pieces $V^{>}$ of degree $r \in (\bar{n}, 2\bar{n}]$; and pieces V^{R} that contain a remainder $R_{\bar{n}+1}$. The same split is applied to $Q_k(\mathfrak{B}+\tilde{A})$ and to the $\mathcal{P}$-vertices.

(E2) Expand each $G_k(\mathfrak{B}+\tilde{A})$ by the two-sided identity $G' = \sum_{n+n'<m}(GV)^n G (VG)^{n'} + \mathcal{K}_m$, which follows from $G' = G + GVG'$ applied on both sides. Here $\mathcal{K}_m = [G(\mathfrak{B})V]^n G(\mathfrak{B}+\tilde{A})[VG(\mathfrak{B})]^{n'}$ with $n+n' = m$, $m := \max(6, \bar{n}+1-q_0)$, and q_0 is the order already carried by the term.

(E3) A term is expanded further only while its total coupling order is $\le \bar{n}$. Every other term goes into Δ_k^{pre}, with its unexpanded factors $\mathcal{K}_m$, $V^{>}$ and V^{R}.

Each V-insertion raises the order by at least one. So the procedure stops after at most $\bar{n}+1$ rounds. The number of terms depends only on $\bar{n}$ and on the vertex list [3, (1.6)–(1.11), (1.18)]. The polynomial terms of Δ_k^{pre} have graded order at most $3\bar{n} = 39$; families in which $\mathcal{K}_m$ replaces a line have order at most $\bar{n}+14 = 27$. Both are within the cap $\bar{N} = 169$.

Step 2 (chains of the basic quadratic form). Consider $\langle hf, \mathcal{C}h'f'\rangle$, where $\mathcal{C}$ is a chain $GV^{[r_1]}G\cdots V^{[r_n]}G$ or contains $\mathcal{K}_m$ or $V^{>}$. The proof of (K3) writes $\mathcal{K} = L_0(VL_1)\cdots(VL_m)$ and uses $\|VL\|_{\infty,\beta} \le \|R_0\|\|L\| + \sum_\mu \|R_{1,\mu}\|\|D_\mu L\|$ together with submultiplicativity. The degree-r Taylor piece of the bond function is again a function of q. So by the lattice Leibniz rule $V^{[r]}$, $V^{>}$ and V^{R} have the same right-pushed form, with $\|\mathfrak{r}^{[r]}\| \le c_V(C(d)\sigma)^r$. Hence

$$|\langle hf, \mathcal{C}h'f'\rangle| \le C_X(C_\infty C_\sigma t^{1/2}p)^q \mathrm{e}^{-\beta\,\mathrm{dist}(\mathrm{supp}\,h, \mathrm{supp}\,h')}\|f\|_\infty\|f'\|_\infty,$$
$$\|f\|_\infty \le Cp\lambda^{-1/4}t^{-1/4}.$$

A chain of order q is therefore $\le C(q)t^{(q-1)/2}\lambda^{-1/2}\,\mathrm{polylog}\,\mathrm{e}^{-\beta\,\mathrm{dist}}$, and at $q \ge \bar{n}+1 = 14$ this is $\le Ct^{13/2}\,\mathrm{polylog}$.

Step 3 (families of $\mathcal{P}_w^{(k)}$). After (E1)–(E3) the pieces enter Bałaban's estimates as follows.

- The expanded external scalar fields are the zeroth-order fields $\Phi_0 := a_k G_k(\Omega, \mathfrak{B}) Q_k^*(\mathfrak{B})\phi$ and $\phi(y)$, and chains $\mathcal{K}f$ of order $m \ge 1$, with $\|h\cdot\|_{1,\alpha_0} \le C(C_\infty\sigma)^m p\lambda^{-1/4}t^{-1/4}$ by (K3).
- The $\tilde{A}$-legs are external photon fields with $\|h'\tilde{A}\|_{1,\alpha_0} \le Cp$. This uses $(\tilde{B})$, the Hölder clause of (K1) for $a_k G_k Q_k^*$, and the C^2 profile of the cut-offs θ_{k+1} and ζ.
- A line replaced by $\mathcal{K}_m$ is an external two-variable field of [3, (2.5)], with $\|h\mathcal{K}_m h'\|_{1,\alpha_0} \le C_U(C_V C_\sigma t^{1/2}p)^m \mathrm{e}^{-\beta\,\mathrm{dist}}$ by Proposition E.57(b) ($m \ge 6$, $\alpha_0 = \frac{1}{2}$). No domain difference of a $\mathcal{K}_m$-operator is needed, because all lines of a given term live on one domain. The two-domain bounds δG enter only inside the families, through IH and CAP.

By Lemma E.35, every family that occurs has at least one external field, at every k including $k = 0$. For such families the family bound of Definition E.59 is available. So each resulting family obeys [3, (1.33)] at graded order $\le \bar{N}$, by IH and CAP, with the factor $(|u|c_k)^m \le 3^{-m}$. This is (PC1) of Lemma E.34. The norm bounds above give (PC2), and the bounds on the δm^2 vertices recalled in the statement give (PC3). A family of graded order $q \ge 14$ is connected and has at least two vertices. By Lemma E.34, with (X) supplied in Step 3a, it is $\le C(q)t^{q/4}\hat{p}^{\,4q}(1+K-k)^q \mathrm{e}^{-\delta(\bar{N})d}$, and $(1+K-k)^q \le (1+\ell(t)/\log L)^q$ is a polylog. Each order gains at least $t^{1/4}$ up to polylog, a $V^{[r]}$-vertex through $(C_\sigma t^{1/2}p)^r$ and a $\mathcal{K}_m$-factor through $t^{m/2}$ polylog.

Step 3a (the exceptional vertices). Let v be an exceptional vertex in the sense of Lemma E.34, so v has coupling order one and both of its scalar legs are external. Since the family is connected and has at least two vertices, the vector leg of v is an internal leg A'. Hence v is [3, (1.8)] with $n = 1$,

$n' = 0$, or the product of [3, (1.12)] or [3, (1.13)] with [3, (1.14)] at $n = 1$, $n' = 0$. The bound (X) does not follow from (PC2). For example, if $\|\Phi\|_{1,\alpha_0} = X_s$ and $|D\Phi| \asymp X_s$, the vertex [3, (1.8)] is of order $e(t)X_s^2 \asymp p(t)^2$. (X) rests on the cancellations of [3, p. 421], which use [2, Lemma 2.4] and the identity $\phi(y) \cdot q\phi(y) = 0$. We distinguish three cases.

- One scalar leg of v is a chain of order $m \geq 1$. By Statement K.14, v together with the chain contributes at most $Ct^{(1+m)/4}p^{2+m}$. This is (X), with the order m of the chain counted in q and $\hat{p}^{\,2+m} \leq \hat{p}^{\,4(1+m)}$.
- Both scalar legs of v are Φ_0 or $\phi(y)$, and the vector leg of v is contracted with a bond leg $(g_k A')_{b'}$ of a vertex of type [3, (1.8)–(1.11)]. Then (X) is Statement K.15.
- Both scalar legs of v are Φ_0 or $\phi(y)$, and the vector leg of v is contracted with a contour leg of a factor [3, (1.14)] or [3, (1.15)]. Then (X) is the assumed bound of Statement K.16.

Step 4 (domain hypothesis of (K3)). On a Neumann domain, (K3) needs:

- internal points $y \in \operatorname{supp} V \subset (\operatorname{supp} \tilde{A})^{+\sqrt{3}+\eta}$ at distance $\geq R_0$ from the complement, which holds if $\operatorname{dist}(\operatorname{supp} \tilde{A}, \partial\Omega) \geq R_0 + 2\sqrt{3} + 1$;
- external points at distance $\geq R_0$.

Every term of $\tilde{A}$ carries θ_{k+1}, so $\operatorname{supp} \tilde{A} \subset \operatorname{supp} \theta_{k+1}$, which lies at distance $\geq r(t) - 2M_b$ from $\Lambda_0^{(k)c}$. Condition (E.10) gives $r(\varepsilon_0) \geq 8M_b + 8$, and (R.2) gives $r(t) \geq R \geq 3R_0 + 15$; moreover $r(t) \geq r(\varepsilon_0)$ for $t \leq \varepsilon_0$. Hence

$$r(t) \geq \tfrac{1}{4}(8M_b + 8) + \tfrac{3}{4}(3R_0 + 15) = 2M_b + \tfrac{9}{4}R_0 + 13.25 \geq 2M_b + R_0 + 2\sqrt{3} + 1.$$

This settles the basic quadratic form on Ω. For the lines of $\mathcal{P}^{(k)}$ on Ω': $\Lambda_0^{(k)} \subset \Lambda_7^{(k-1)\prime}$, so the internal margin is at least as large. The external points are vertex points in Ω_1. This set lies several r-layers inside Ω', which is the setting of (K3) on Neumann domains. And $R \geq 107 > 12\sqrt{3}$.

Step 5 (sum). For $q \geq 14$, $t^{q/4}\hat{p}^{\,4q} \leq t^{7/2}$ polylog, and $\sup_{t\in(0,1]} t^{3/10}\big(b_0(1 + \ln t^{-1})^{p_0}\big)^{4q} < \infty$. So every term is $\leq Ct^{16/5}$, with room $t^{3/10}$; Step 2 has more room. The localisations through a fixed block of $\Lambda_2^{(k)}$, weighted by $\mathrm{e}^{-\min(\beta,\delta(\bar{N}))d}$, sum to $S(\bar{N}, d) < \infty$. There are finitely many terms, and $|\Lambda_2^{(k)}|_k \leq N_k$. No inequality of the type [1, (3.47)] is used, so the bound is two-sided. □

Remark E.37 (the other two cuts, and the mass range). (i) The terms that contain a Taylor remainder $R_{\bar{n}+1}$ obey $\leq C_R t^{13/2}$ polylog $|\Lambda_2^{(k)}|_k$. Since $\tilde{A}$ is real, $|R_{\bar{n}+1}(z)| \leq |z|^{\bar{n}+1}/(\bar{n}+1)!$. The $\|\cdot\|_{1,\alpha_0}$-bound $c_R(Ct^{1/2}p)^{\bar{n}+1}$ of an R-factor also needs $|R''||Dz||\Delta z|$ and $|R'||\Delta Dz|$, where $[\partial\tilde{A}]_\alpha$ comes from the Hölder clause of (K1). (ii) The terms that contain $\tilde{B}^{(k)}$ obey $\leq C_{\tilde{B}} t^{9/2}$ polylog $|\Lambda_2^{(k)}|_k$ at $\kappa = 31$. The fourth term of $\tilde{B}^{(k)}$ is $a_{k+1}L^{-2}\theta_{k+1}(1 - \zeta^{(k+1)})G_{k+1}^{\eta}Q_{k+1}^{*}\Lambda_0 B$. Both follow by the argument of Proposition E.36, with the bounds just stated in place of Step 2. (iii) The three cuts use M only as stated in Proposition E.36, so they hold at every $M \in (0, m_2]$. At $u = 0$ the factor $(|u|c_k)^m$ is at most its value at $u \in \{1, -\eta\}$.

E.3. **Line bounds, the remainder line and the renormalisation machinery on the torus.** We prove the inputs (VS), (Rem) and (S6) of the bounds on the final c-numbers of Subsection E.2, and state the inputs (K1)–(K4), (IH) and (CAP) of the tail bound.

The zero-background slices and the last remainder line. *Setting (scale-j units).* Fix a slice index j with $m_f t_j \leq 1$, and measure lengths in units of t_j. Let Λ be the η-lattice torus of side ℓ, where $\eta := \varepsilon/t_j = L^{-j} \in (0, 1]$, $1/\eta \in \mathbb{N}$, and $\ell := \ell_T/t_j \in L\mathbb{N}$ with $\ell \geq L$. The inner product is $\langle f, g\rangle = \eta^3 \sum_x f(x)\bar{g}(x)$, and a kernel acts by $(Kf)(x) = \eta^3 \sum_y K(x, y)f(y)$. d is the ℓ^1 distance on the torus. $-\Delta$ is the periodic nearest-neighbour Laplacian, and ∇_μ is the forward difference. For $s \in \{1, L\}$, P_s is the orthogonal projection onto functions constant on blocks of side s. Put $\tilde{\mu} := m_f t_j \in [0, 1]$, $b := a_j$ and $b' := a_{j+1}L^{-2}$, with $a_k \in [a_\flat, a]$ and $a_\flat := a(1 - L^{-2})$. Define

$$\mathsf{X} := -\Delta + \tilde{\mu}^2 + bP_1, \quad \mathsf{X}' := -\Delta + \tilde{\mu}^2 + b'P_L, \quad G := \mathsf{X}^{-1}, \quad G' := \mathsf{X}'^{-1}, \quad H := G' - G, \quad F := (-\Delta + 1)^{-1}.$$

In physical units the slice $H_{j,f} = G_{j+1,f}(0) - G_{j,f}(0)$ has kernel $t_j^{-1} H(x/t_j, y/t_j)$ (canonical scaling, $d = 3$).

Lemma E.38 (the Feynman photon is componentwise)**.** *On the hypercubic torus, $\|dA\|^2 + \|d^*A\|^2 = \sum_{\mu,\nu} \|\partial_\nu A_\mu\|^2$. Hence $C_v^\varepsilon = (-\Delta^\varepsilon + \mu_*^2)^{-1} \delta_{\mu\nu}$. At $A = 0$, $G_{k,v}(0)$, $H_{j,v}$ and $R_{\bar{K},v}$ are each $\delta_{\mu\nu}$ times the scalar object of mass μ_*.*

Proof. $\|dA\|^2 = \sum_{\mu \neq \nu} \|\partial_\mu A_\nu\|^2 - \sum_{\mu \neq \nu} \langle \partial_\mu A_\nu, \partial_\nu A_\mu \rangle$ and $\|d^*A\|^2 = \sum_\mu \|\partial_\mu A_\mu\|^2 + \sum_{\mu \neq \nu} \langle \partial_\mu^* A_\mu, \partial_\nu^* A_\nu \rangle$. Forward differences and their adjoints commute on the torus, so $\langle \partial_\mu^* A_\mu, \partial_\nu^* A_\nu \rangle = \langle \partial_\nu A_\mu, \partial_\mu A_\nu \rangle$, and the cross terms cancel. Componentwise block averaging at $A = 0$ ([24, (2.10)–(2.17), p. 653]: "setting $m^2 = \mu_0^2$, $N = d$ and $A = 0$") gives the rest. □

The heat kernel of $\eta\mathbb{Z}^3$. For $\eta \in (0, 1]$ let Δ be the nearest-neighbour Laplacian of $\eta\mathbb{Z}^3$, $\Delta f(x) = \eta^{-2} \sum_\mu [f(x + \eta e_\mu) + f(x - \eta e_\mu) - 2f(x)]$, and let $\nabla_\mu f(x) := \eta^{-1}(f(x + \eta e_\mu) - f(x))$. Kernels are taken with respect to $\eta^3 \sum_y$. For a translation-invariant operator we write its kernel as a function of $w = x - y$, and ∇_μ acts on w. Put $|w|_1 := \sum_\mu |w_\mu|$, $\hat{k}_\mu := \frac{2}{\eta} \sin(\eta k_\mu / 2)$, $\hat{k}^2 := \sum_\mu \hat{k}_\mu^2$, $B_\eta := [-\pi/\eta, \pi/\eta]^3$ and $a := \sqrt{\pi}/4$. For $z \geq 0$ and $n \in \mathbb{Z}$ define $I_n(z)$ by $\mathrm{e}^{(z/2)(t + t^{-1})} = \sum_{n \in \mathbb{Z}} I_n(z) t^n$ ($t \neq 0$). Then $I_n = I_{-n} \geq 0$, and setting $t = 1$ gives $\sum_n I_n(z) = \mathrm{e}^z$.

Lemma E.39 (free kernel bounds)**.** *On $\eta\mathbb{Z}^3$, $\eta \in (0, 1]$, let $F^\infty := (-\Delta + 1)^{-1}$ and $r := |w|_1$. Then:*

- (F0) *$F^\infty(w) \leq A_0 \mathrm{e}^{-r/4}/(r + \eta)$, with $A_0 := 4.22$;*
- (F1) *$|\nabla_\mu F^\infty(w)| \leq A_1 \mathrm{e}^{-r/4}/(r + \eta)^2$, with $A_1 := 98$;*
- (F2) *$|\nabla_\mu \nabla_\nu F^\infty(w)| \leq A_2 \mathrm{e}^{-r/4}/(r + \eta)^3$, with $A_2 < \infty$ absolute;*
- (F3) *$|\nabla_\mu (F^\infty)^2(w)| \leq A_3 \mathrm{e}^{-r/4}$, with $A_3 := A_0(1 + 2\mathrm{e}^{1/4})$;*
- (F4) *for $\kappa \leq 1/8$, $K_0 := \eta^3 \sum_w \mathrm{e}^{\kappa r} F^\infty(w) \leq 8.4$ and $K_1 := \eta^3 \sum_w \mathrm{e}^{\kappa r} |\nabla F^\infty(w)| \leq 49 A_1$, where $|\nabla F^\infty(w)| := \max_\mu |\nabla_\mu F^\infty(w)|$.*

Here $(F^\infty)^2 = (-\Delta + 1)^{-2}$. For $(-\Delta + \rho^2)^{-1}$ with $\rho \in (0, 1]$, (F0) holds with $\mathrm{e}^{-\rho r/4}$ and the same A_0, and $\eta^3 \sum_w (-\Delta + \rho^2)^{-1}(w) = \rho^{-2}$.

The proof rests on two lemmas.

Lemma E.40 (the heat kernel)**.** *Let p_s ($s \geq 0$) be the kernel of $\mathrm{e}^{s\Delta}$ on $\eta\mathbb{Z}^3$.*

- (a) *$p_s(w) = \int_{B_\eta} \mathrm{e}^{ik \cdot w} \mathrm{e}^{-s\hat{k}^2} \frac{d^3k}{(2\pi)^3} = \prod_{\mu=1}^3 q_s(w_\mu)$, where $q_s(n\eta) = \eta^{-1} \mathrm{e}^{-z} I_n(z)$ and $z := 2s/\eta^2$. In particular $p_s \geq 0$.*
- (b) *For every $\lambda \in \mathbb{R}^3$, $p_s(w) \leq \mathrm{e}^{-\lambda \cdot w} \exp\big(\sum_\mu \frac{2s}{\eta^2} (\cosh \eta\lambda_\mu - 1) \big) p_s(0)$.*
- (c) *$p_s(0) \leq \min(\eta^{-1}, as^{-1/2})^3$.*
- (d) *For every $\bar{\lambda} \geq 0$ and every $\lambda \in \mathbb{R}^3$ with $|\lambda_\nu| \leq \bar{\lambda}$ for all ν,*

$$|\nabla_\mu p_s(w)| \leq \mathrm{e}^{-\lambda \cdot w} \exp\Big(\sum_\nu \tfrac{2s}{\eta^2} (\cosh \eta\lambda_\nu - 1) \Big) \mathrm{e}^{\eta\bar{\lambda}} \int_{B_\eta} \big(|\hat{k}_\mu| + \bar{\lambda} \big) \mathrm{e}^{-s\hat{k}^2} \frac{d^3k}{(2\pi)^3}.$$

- (e) *For $s > 0$, $|\nabla_\mu p_s(w)| \leq \frac{|w|_1 + \eta}{s} \max\big(p_s(w), p_s(w + \eta e_\mu) \big)$.*

Proof. (a) In one dimension, $\mathrm{e}^{s\Delta} = \mathrm{e}^{-z} \exp\big(\frac{z}{2}(S + S^{-1}) \big)$, where S is the unit shift and $z = 2s/\eta^2$. The coefficient of S^n is $\mathrm{e}^{-z} I_n(z)$, and the factor η^{-1} comes from the normalisation of kernels. The Fourier formula is the same identity, because $\frac{2}{\eta^2}(1 - \cos \eta k) = \hat{k}^2$. The three-dimensional kernel is the product of the one-dimensional ones.

(b) The integrand $k \mapsto \mathrm{e}^{ik \cdot w} \mathrm{e}^{-s\hat{k}^2}$ is entire and $2\pi/\eta$-periodic in each k_μ. So the shift $k_\mu \to k_\mu + i\lambda_\mu$ does not change the integral: by periodicity, the contributions of the vertical sides cancel. Moreover $\mathrm{e}^{i(k + i\lambda) \cdot w} = \mathrm{e}^{ik \cdot w} \mathrm{e}^{-\lambda \cdot w}$ and

$$\mathrm{Re}\big[1 - \cos \eta(k_\mu + i\lambda_\mu) \big] = \cosh(\eta\lambda_\mu)(1 - \cos \eta k_\mu) - (\cosh \eta\lambda_\mu - 1) \geq (1 - \cos \eta k_\mu) - (\cosh \eta\lambda_\mu - 1).$$

Hence the modulus of the shifted integrand is at most $\mathrm{e}^{-\lambda\cdot w}\exp(\sum_\mu \frac{2s}{\eta^2}(\cosh\eta\lambda_\mu - 1))\mathrm{e}^{-s\hat{k}^2}$. Integrating gives (b), since $p_s(0) = \int_{B_\eta} \mathrm{e}^{-s\hat{k}^2}\frac{d^3k}{(2\pi)^3}$.

(c) We have $q_s(0) \le \eta^{-1}$, because $\eta\sum_n q_s(n\eta) = \mathrm{e}^{-z}\sum_n I_n(z) = 1$ and $q_s \ge 0$. On B_η, the inequality $|\sin x| \ge 2|x|/\pi$ for $|x| \le \pi/2$ gives $\hat{k}_\mu^2 \ge 4k_\mu^2/\pi^2$. Hence

$$q_s(0) \le \tfrac{1}{2\pi}\int_{\mathbb{R}} \mathrm{e}^{-4sk^2/\pi^2}dk = \tfrac{\sqrt{\pi}}{4}s^{-1/2}.$$

(d) We have $\nabla_\mu p_s(w) = \int_{B_\eta}\mathrm{e}^{ik\cdot w}m(k_\mu)\mathrm{e}^{-s\hat{k}^2}\frac{d^3k}{(2\pi)^3}$ with $m(k) = (\mathrm{e}^{i\eta k}-1)/\eta$. This integrand is again entire and periodic, so we shift as in (b). Then

$$|m(k+i\lambda_\mu)| \le \mathrm{e}^{-\eta\lambda_\mu}|\mathrm{e}^{i\eta k}-1|/\eta + |\mathrm{e}^{-\eta\lambda_\mu}-1|/\eta \le \mathrm{e}^{\eta\bar{\lambda}}(|\hat{k}_\mu| + \bar{\lambda}),$$

using $|\mathrm{e}^{i\eta k}-1| = \eta|\hat{k}_\mu|$ and $|\mathrm{e}^{-y}-1| \le |y|\mathrm{e}^{|y|}$.

(e) First we show $I_n \ge I_{n+1}$ for $n \ge 0$. Differentiating the generating function in z gives $I_n' = \frac{1}{2}(I_{n-1}+I_{n+1})$. Put $d_n := I_n - I_{n+1}$ for $n \ge 0$. Then $d_{-1} = I_1 - I_0 = -d_0$, so $d = (d_n)_{n\ge 0}$ solves $d' = Ad$ with $d(0) = e_0$. Here $(Ad)_n = \frac{1}{2}(d_{n-1}+d_{n+1})$ for $n \ge 1$ and $(Ad)_0 = \frac{1}{2}d_1 - \frac{1}{2}d_0$. The operator A is bounded on $\ell^\infty(\{0,1,2,\dots\})$, and $A + \frac{1}{2}$ has nonnegative entries. So $\tilde{d}(z) := \mathrm{e}^{-z/2}\mathrm{e}^{z(A+1/2)}e_0$ has nonnegative entries and solves the same equation. Both d and $\tilde{d}$ satisfy $f(z) = e_0 + \int_0^z Af$ coordinatewise, and both are bounded in ℓ^∞ on compact z-intervals ($0 \le I_n \le \mathrm{e}^z$). Gronwall's inequality applied to $\sup_n|d_n - \tilde{d}_n|$ then gives $d = \tilde{d} \ge 0$.

Second, differentiating the generating function in t gives $\frac{z}{2}(I_{n-1} - I_{n+1}) = nI_n$. Write $q(n) := q_s(n\eta)$. For $n \ge 1$,

$$0 \le q(n) - q(n+1) \le q(n-1) - q(n+1) = \tfrac{2n}{z}q(n) = \tfrac{n\eta^2}{s}q(n).$$

For $n = 0$, $0 \le q(0) - q(1) \le q(0) - q(2) = \frac{\eta^2}{s}q(1)$. Negative n follow from the symmetry $q(n) = q(-n)$. Together, $|q(n+1) - q(n)| \le \frac{(|n|+1)\eta^2}{s}\max(q(n), q(n+1))$, that is, $|\nabla q_s(x)| \le \frac{|x|+\eta}{s}\max(q_s(x), q_s(x+\eta))$. Multiply by the other two factors of (a), and use $|w_\mu|+\eta \le |w|_1+\eta$. □

Lemma E.41 (lattice sums)**.** *For $\phi > 0$, $\eta \in (0,1]$ and $p \in \{0,1,2\}$ put $J_p(\phi) := \eta^3\sum_{v\in\eta\mathbb{Z}^3}\mathrm{e}^{-\phi|v|_1}(|v|_1+\eta)^{-p}$. Then $J_0(\phi) \le (1+2/\phi)^3$, $J_1(\phi) \le 1 + 6/\phi^2$ and $J_2(\phi) \le 1 + 6/\phi$.*

Proof. Put $x := \mathrm{e}^{-\phi\eta}$ and $u := \phi\eta$.

- $J_0(\phi) = \big(\eta\frac{1+x}{1-x}\big)^3 = \big(\eta + \frac{2\eta}{\mathrm{e}^u-1}\big)^3 \le (1+2/\phi)^3$, since $\mathrm{e}^u - 1 \ge u$.
- For $n \ge 1$, the number of $v \in \eta\mathbb{Z}^3$ with $|v|_1 = n\eta$ is $4n^2 + 2 \le 6n^2$. Hence

$$J_1(\phi) \le \eta^2 + 6\eta^2\sum_{n\ge1} n\mathrm{e}^{-un} = \eta^2 + \frac{6}{\phi^2}\frac{u^2\mathrm{e}^{-u}}{(1-\mathrm{e}^{-u})^2} \le 1 + \frac{6}{\phi^2},$$

  because $u\mathrm{e}^{-u/2}/(1-\mathrm{e}^{-u}) = u/(2\sinh(u/2)) \le 1$.
- $J_2(\phi) \le \eta + 6\eta\sum_{n\ge1}\mathrm{e}^{-un} = \eta + \frac{6\eta}{\mathrm{e}^u-1} \le 1 + \frac{6}{\phi}$.

□

Proof of Lemma E.39. We have $F^\infty = \int_0^\infty \mathrm{e}^{-s}p_s\,ds$, $(F^\infty)^2 = \int_0^\infty s\mathrm{e}^{-s}p_s\,ds$ and $(-\Delta+\rho^2)^{-1} = \int_0^\infty \mathrm{e}^{-\rho^2 s}p_s\,ds$. By Lemma E.40(a), all three kernels are ≥ 0. We use $\cosh y - 1 \le \frac{y^2}{2}\cosh y$, which holds term by term in the power series because $(2k)! \ge 2(2k-2)!$. For $|y| \le \frac{1}{2}$ it gives $\cosh y - 1 \le 0.5638\,y^2$.

The choice of λ. Put $\lambda_\mu := \mathrm{sgn}(w_\mu)\bar\lambda$, with $\mathrm{sgn}(0) := 1$ and $\bar\lambda := \frac14 + \frac{1}{4\max(\sqrt{s},\eta)}$. Then $\lambda\cdot w = \bar\lambda r$ and $\eta\bar\lambda \le \frac12$. Also

$$\sum_\mu \tfrac{2s}{\eta^2}(\cosh\eta\bar\lambda - 1) \le 3.383\, s\bar\lambda^2 \le 3.383\, s\Big(\tfrac18 + \tfrac{1}{8\max(s,\eta^2)}\Big) \le 0.423\, s + 0.423.$$

By Lemma E.40(b), and since $\mathrm{e}^{0.423} \le 1.527$,

$$p_s(w) \le 1.527\,\mathrm{e}^{0.423s} p_s(0)\,\mathrm{e}^{-r/4}\mathrm{e}^{-r/(4\max(\sqrt{s},\eta))}. \tag{E.3}$$

(F0) On $s \le \eta^2$ we use $p_s(0) \le \eta^{-3}$ and $\mathrm{e}^{-s}\mathrm{e}^{0.423s} \le 1$. The contribution is at most $1.527\,\eta^{-1}\mathrm{e}^{-r/(4\eta)}\mathrm{e}^{-r/4}$. With $u = r/\eta$ we have $\eta^{-1}\mathrm{e}^{-u/4} = (r+\eta)^{-1}(1+u)\mathrm{e}^{-u/4}$, and $\max_{u\ge0}(1+u)\mathrm{e}^{-u/4} = 4\mathrm{e}^{-3/4} \le 1.889$.

On $s \ge \eta^2$ we use $p_s(0) \le a^3 s^{-3/2}$ and substitute $u = s^{-1/2}$:

$$\int_{\eta^2}^\infty a^3 s^{-3/2}\mathrm{e}^{-r/(4\sqrt{s})}ds = 2a^3\int_0^{1/\eta}\mathrm{e}^{-ru/4}du \le 2a^3\min(4/r, 1/\eta) \le 10a^3/(r+\eta).$$

For the last step, treat $r \le 4\eta$ and $r \ge 4\eta$ separately. Since $10a^3 \le 0.871$, the total is at most $1.527(1.889+0.871)\mathrm{e}^{-r/4}/(r+\eta) \le 4.22\,\mathrm{e}^{-r/4}/(r+\eta)$.

For the mass $\rho \in (0,1]$ take $\bar\lambda := \frac{\rho}{4} + \frac{1}{4\max(\sqrt{s},\eta)}$. Then still $\eta\bar\lambda \le \frac12$, the exponent is at most $0.423\rho^2 s + 0.423$, and $\mathrm{e}^{-\rho^2 s}\mathrm{e}^{0.423\rho^2 s} \le 1$. The same two integrals give $A_0\mathrm{e}^{-\rho r/4}/(r+\eta)$. In particular the kernel is summable. Hence $\eta^3\sum_w(-\Delta+\rho^2)^{-1}(w)$ equals its Fourier transform $(\rho^2+\hat k^2)^{-1}$ at $k=0$, that is, ρ^{-2}.

(F1) *The range* $s \ge \eta^2$. Apply Lemma E.40(d) with the λ above; here $\mathrm{e}^{\eta\bar\lambda} \le \mathrm{e}^{1/2}$. By Cauchy–Schwarz in $d^3k/(2\pi)^3$,

$$\int_{B_\eta}|\hat k_\mu|\mathrm{e}^{-s\hat k^2} \le \Big(\int_{B_\eta}\hat k_\mu^2\mathrm{e}^{-s\hat k^2}\Big)^{1/2} p_s(0)^{1/2}.$$

In direction μ use $\hat k_\mu^2 \le k_\mu^2$ and $\mathrm{e}^{-s\hat k_\mu^2} \le \mathrm{e}^{-4sk_\mu^2/\pi^2}$; in the other two directions use Lemma E.40(c). This gives $\int \hat k_\mu^2\mathrm{e}^{-s\hat k^2} \le \frac{\pi^{5/2}}{32}s^{-3/2}\cdot a^2 s^{-1}$, hence $\int|\hat k_\mu|\mathrm{e}^{-s\hat k^2} \le 0.0967\,s^{-2}$. Also $\bar\lambda p_s(0) \le \frac{a^3}{4}(s^{-3/2} + s^{-2})$, with $a^3/4 \le 0.0218$. So

$$|\nabla_\mu p_s(w)| \le 1.527\,\mathrm{e}^{0.423s}\mathrm{e}^{-r/4}\mathrm{e}^{-r/(4\sqrt{s})}\big[0.196\,s^{-2} + 0.036\,s^{-3/2}\big].$$

Multiply by e^{-s} and use $\sqrt{s}\,\mathrm{e}^{-0.577s} \le 0.565$. Then substitute $u = s^{-1/2}$:

$$\int_{\eta^2}^\infty s^{-2}\mathrm{e}^{-r/(4\sqrt{s})}ds = 2\int_0^{1/\eta} u\mathrm{e}^{-ru/4}du \le 2\min(16/r^2, 1/(2\eta^2)) \le 44.4/(r+\eta)^2.$$

Here we used that the maximum over r of $\min(16/r^2, 1/(2\eta^2))(r+\eta)^2$ is attained at $r^2 = 32\eta^2$, where it equals $(\sqrt{32}+1)^2/2 \le 22.16$. The range $s \ge \eta^2$ therefore contributes at most $14.7\,\mathrm{e}^{-r/4}/(r+\eta)^2$.

The range $s \le \eta^2$. Use $|\nabla_\mu p_s(w)| \le \eta^{-1}(p_s(w) + p_s(w+\eta e_\mu))$ and apply (E.3) at both points, with $\bar\lambda = \frac14 + \frac{1}{4\eta}$. Since $|w+\eta e_\mu|_1 \ge r - \eta$, the second point costs at most a factor $\mathrm{e}^{\eta\bar\lambda} \le \mathrm{e}^{1/2}$. With $p_s(0) \le \eta^{-3}$ and $\int_0^{\eta^2} ds = \eta^2$, the contribution is at most $1.527(1+\mathrm{e}^{1/2})\eta^{-2}\mathrm{e}^{-r/(4\eta)}\mathrm{e}^{-r/4}$. Finally $\eta^{-2}\mathrm{e}^{-u/4} = (r+\eta)^{-2}(1+u)^2\mathrm{e}^{-u/4}$ with $u = r/\eta$, and $\max_u(1+u)^2\mathrm{e}^{-u/4} = 64\mathrm{e}^{-7/4} \le 11.13$. This gives at most $45.1\,\mathrm{e}^{-r/4}/(r+\eta)^2$. Since $14.7 + 45.1 \le 98$, the value $A_1 = 98$ suffices.

(F2) $\nabla_\mu\nabla_\nu p_s(w)$ is the integral of $\mathrm{e}^{ik\cdot w}m(k_\mu)m(k_\nu)\mathrm{e}^{-s\hat k^2}$, again entire and periodic. After the shift of Lemma E.40(b), the product of the two multipliers is bounded by $\mathrm{e}^{2\eta\bar\lambda}(|\hat k_\mu| + \bar\lambda)(|\hat k_\nu| + \bar\lambda)$, with $\mathrm{e}^{2\eta\bar\lambda} \le \mathrm{e}$.

- By Cauchy–Schwarz and the one-dimensional Gaussian bounds used in (F1), for $s \ge \eta^2$ we have $\int |\hat k_\mu \hat k_\nu| \mathrm{e}^{-s\hat k^2} \le C s^{-5/2}$, $\bar\lambda \int |\hat k_\mu| \mathrm{e}^{-s\hat k^2} \le C(s^{-2}+s^{-5/2})$ and $\bar\lambda^2 p_s(0) \le C(s^{-3/2}+s^{-5/2})$. After multiplication by $\mathrm{e}^{-0.577 s}$, each of these is at most $C' s^{-5/2}$.
- $\int_{\eta^2}^\infty s^{-5/2} \mathrm{e}^{-r/(4\sqrt{s})} ds = 2\int_0^{1/\eta} u^2 \mathrm{e}^{-ru/4} du \le 2\min(128/r^3, 1/(3\eta^3)) \le C/(r+\eta)^3$.
- On $s \le \eta^2$, the second difference is η^{-2} times a signed sum of four values of p_s, at points of ℓ^1-norm $\ge r - 2\eta$. By (E.3), $p_s(0) \le \eta^{-3}$ and $\int_0^{\eta^2} ds = \eta^2$, this gives at most $C\eta^{-3}\mathrm{e}^{-r/(4\eta)}\mathrm{e}^{-r/4} \le C' \mathrm{e}^{-r/4}/(r+\eta)^3$.

All constants here are absolute.

(F3) Here a cancellation is used: the factor s of $(F^\infty)^2 = \int_0^\infty s\mathrm{e}^{-s} p_s \, ds$ cancels the factor s^{-1} of Lemma E.40(e). Thus

$$|\nabla_\mu (F^\infty)^2(w)| \le \int_0^\infty s\mathrm{e}^{-s\frac{r+\eta}{s}} \big(p_s(w) + p_s(w+\eta e_\mu)\big) ds = (r+\eta)\big(F^\infty(w) + F^\infty(w+\eta e_\mu)\big).$$

Let $r' := |w + \eta e_\mu|_1$. Then $r' \ge r - \eta$ and $r' + \eta \ge \max(r, \eta) \ge (r+\eta)/2$. So (F0) gives $(r+\eta)F^\infty(w) \le A_0 \mathrm{e}^{-r/4}$ and $(r+\eta)F^\infty(w+\eta e_\mu) \le 2\mathrm{e}^{1/4} A_0 \mathrm{e}^{-r/4}$. Hence $A_3 = A_0(1 + 2\mathrm{e}^{1/4})$ works.

(F4) We have $\mathrm{e}^{\kappa r} = \prod_\mu \mathrm{e}^{\kappa |w_\mu|} \le \sum_{\sigma \in \{\pm 1\}^3} \mathrm{e}^{\kappa \sigma \cdot w}$. For $\lambda \in \mathbb{C}^3$ with $|\mathrm{Re}\, \lambda_\mu| < \frac14$, the series $\Phi(\lambda) := \eta^3 \sum_w \mathrm{e}^{\lambda \cdot w} F^\infty(w)$ converges absolutely and locally uniformly by (F0), so Φ is analytic there. Put $D(\lambda) := 1 - \sum_\mu \frac{2}{\eta^2}(\cosh \eta\lambda_\mu - 1)$. For $\lambda = -ik$ with $k \in \mathbb{R}^3$, $\Phi(\lambda) = (1+\hat k^2)^{-1}$ and $D(\lambda) = 1 + \hat k^2$. Thus the analytic function $\Phi D - 1$ vanishes on $i\mathbb{R}^3$, and so it vanishes identically (argue one variable at a time). For real λ with $|\lambda_\mu| \le \kappa \le \frac18$, the bound $\frac{2}{\eta^2}(\cosh \eta\kappa - 1) \le \kappa^2 \cosh \kappa$ gives $D(\lambda) \ge 1 - 3\kappa^2 \cosh\kappa > 0$. Since $F^\infty \ge 0$, this yields $K_0 \le 8/(1 - 3\kappa^2 \cosh\kappa) \le 8.4$. For K_1, (F1) gives $\mathrm{e}^{\kappa r}|\nabla F^\infty(w)| \le A_1 \mathrm{e}^{-(1/4-\kappa)r}/(r+\eta)^2$. Lemma E.41 with $\frac14 - \kappa \ge \frac18$ then gives $K_1 \le A_1 J_2(\frac14 - \kappa) \le A_1(1+48) = 49 A_1$. □

Lattice sums. For $\phi > 0$ and $\eta \le 1$: $J_0(\phi) := \eta^3 \sum_{v \in \eta\mathbb{Z}^3} \mathrm{e}^{-\phi|v|_1} \le (1 + 2/\phi)^3$, $J_1(\phi) := \eta^3 \sum \mathrm{e}^{-\phi|v|_1}/(|v|_1 + \eta) \le 1 + 6/\phi^2$, and $J_2(\phi) := \eta^3 \sum \mathrm{e}^{-\phi|v|_1}/(|v|_1+\eta)^2 \le 1 + 6/\phi$.

Lemma E.42 (torus images)**.** *Let $|K^\infty(w)| \le A\mathrm{e}^{-\kappa|w|_1}/(|w|_1+\eta)^p$ with $p \in \{0,1,2,3\}$, and let $K^T(z) := \sum_{n\in\mathbb{Z}^3} K^\infty(z + n\ell)$ with $\ell \ge 1$. Then*

$$|K^T(z)| \le A\big(1 + 5^p S(\kappa\ell)\big)\mathrm{e}^{-(\kappa/2)d(z)}/(d(z)+\eta)^p, \qquad S(u) := \big(3 + 2/(\mathrm{e}^{u/2} - 1)\big)^3,$$

and $\eta^3 \sum_{z\in\Lambda} \mathrm{e}^{\kappa' d(z)} |K^T(z)| \le \eta^3 \sum_{w\in\eta\mathbb{Z}^3} \mathrm{e}^{\kappa'|w|_1} |K^\infty(w)|$ for every $\kappa' \ge 0$.

Proof. Take z in the fundamental domain. For $n \ne 0$, $|z + n\ell|_1 \ge \max(d(z), \ell/2)$, so $(|z+n\ell|_1 + \eta)^{-p} \le 5^p (d(z)+\eta)^{-p}$. Moreover $\mathrm{e}^{-\kappa|z+n\ell|_1} \le \mathrm{e}^{-(\kappa/2)d(z)} \mathrm{e}^{-(\kappa/2)|z+n\ell|_1}$, and $\sum_n \mathrm{e}^{-(\kappa/2)|z+n\ell|_1} \le S(\kappa\ell)$. For the ℓ^1 statement, unfold $\sum_{z\in\Lambda}\sum_n = \sum_{w\in\eta\mathbb{Z}^3}$ and use $d(z) \le |z+n\ell|_1$. □

The torus kernel of $(-\Delta + 1)^{-1}$ is $F = F^T$. Put $S_L := S(L/4)$, so that $S_L \le 1013$ at $L = 2$ (the value is 1012.54). Put $A_p^T := A_p(1 + 5^p S_L)$ for $p = 0, 1, 2$, $A_3^T := A_3(1 + S_L)$ and $\phi := 1/8$. For $\kappa \le 1/16$ the column norm is $c_F := \sup_x \|\mathrm{e}^{\kappa d(x,\cdot)} F(x,\cdot)\|_{\ell^2} \le 7 A_0^T$.

Lemma E.43 (spectral gap and Combes–Thomas)**.** *(i) For $s \in \{1, L\}$, $\beta > 0$ and $\tilde\mu \ge 0$: $-\Delta + \tilde\mu^2 + \beta P_s \ge \min(4/s^2, \beta)$. (ii) Let ρ be 1-Lipschitz for d, let $0 \le \kappa \le \frac12$, and let $Y := -\Delta + \tilde\mu^2 + \beta P_s \ge \gamma$. If $3.4\kappa^2 + \beta(\mathrm{e}^{3s\kappa} - 1) \le \gamma/2$, then $\|\mathrm{e}^{\kappa\rho} Y^{-1} \mathrm{e}^{-\kappa\rho}\| \le 2/\gamma$.*

Proof. (i) The torus Dirichlet form dominates the sum of the Neumann forms of the s-blocks. A block with s/η points per side has Neumann gap $(4/\eta^2)\sin^2(\pi\eta/(2s)) \ge 4/s^2$. Hence $\langle f, Yf\rangle \ge (4/s^2)\|(1-P_s)f\|^2 + \beta\|P_s f\|^2$. (ii) $\mathrm{e}^{\kappa\rho}(-\Delta)\mathrm{e}^{-\kappa\rho} + \Delta$ splits into a real antisymmetric sinh part,

which has zero real part, and a symmetric $\cosh -1$ part, of norm $\le 3.4\kappa^2$. The conjugated P_s differs from P_s by a kernel of Schur norm $\le \mathrm{e}^{3s\kappa}-1$. So $\mathrm{Re}\langle f,\, \mathrm{e}^{\kappa\rho} Y \mathrm{e}^{-\kappa\rho} f\rangle \ge (\gamma/2)\|f\|^2$. □

Rates. Put $\gamma_0 := \min(4, a_\flat)$, and let $c = c(a,L) \in (0, \frac12]$ be the largest number with $3.4c^2 + a(\mathrm{e}^{3c}-1) \le \gamma_0/2$; at $a=1$, $L=2$ this gives $c = 0.0981$. For X, (ii) holds with $\kappa \le c$; for X' (gap $\ge \gamma_0 L^{-2}$) with $\kappa \le c/L$. With $W := (1-\tilde\mu^2) - bP_1$ and $W' := (1-\tilde\mu^2) - b'P_L$, the conjugated norms are $\le w_1 := 1 + a\mathrm{e}^{3c}$. Put $\kappa_G := \min(c, 1/16)$, $\kappa' := \min(c/L, 1/16)$ and $\kappa_H := \min(c/L, 1/32)$.

Lemma E.44 (structure of G and G')**.** *Put $E := G - F$ and $E' := G' - F$. For all $x, y, z \in \Lambda$:*

- (E1) $|E(x,y)| \le A_E \mathrm{e}^{-\kappa_G d(x,y)}$ *with* $A_E := c_F^2 w_1 (1 + 2w_1/\gamma_0)$*, and likewise for E' with (A'_E, κ'), where $A'_E := c_F^2 w_1 (1 + 2w_1 L^2/\gamma_0)$;*
- (E2) $|(P_1 G)(z,y)| \le A_P \mathrm{e}^{-\kappa_G d(z,y)}$ *and* $|(P_L G)(z,y)| \le A_P \mathrm{e}^{3L\kappa_G} \mathrm{e}^{-\kappa_G d(z,y)}$*, with* $A_P := \mathrm{e}^{3\kappa_G}(J_1(\phi/2) A_0^T + A_E)$*, and likewise for G';*
- (E3) $|\nabla_1 E(x,y)| \le A_\nabla \mathrm{e}^{-\kappa_G d(x,y)}$ *with* $A_\nabla := A_3^T + K_1(A_E + aA_P)$*, and likewise for E';*
- (E4) $|\nabla_y (P_1 G)(z,y)|$ *and* $|\nabla_y (P_L G')(z,y)|$ *are* $\le A_P^\nabla \mathrm{e}^{-\kappa' d(z,y)}$.

Proof. $\mathsf{X} = F^{-1} - W$ gives $G = F + FWG$, hence $E = FWF + FWGWF$. Weighted Cauchy–Schwarz with Lemma E.43(ii) at $\rho_y := d(\cdot, y)$ gives (E1). (E2): average $F + E$ over the unit block. (E3): $\nabla_1 E = (1-\tilde\mu^2)[\nabla(F^2) + \nabla F \cdot E] - b \nabla F \cdot P_1 G$, bounded by (F3), Lemma E.42 and (F4). (E4): G is symmetric, so $\nabla_y G(w,y) = \nabla F(y-w) + \nabla_1 E(y,w)$; average over the block. □

Lemma E.45 (G' maps decaying data to decaying data)**.** *If $0 < \kappa \le \kappa'$ and $|f(z)| \le \mathrm{e}^{-2\kappa d(z,y)}$, then $|(G'f)(x)| \le A_B(\kappa) \mathrm{e}^{-\kappa d(x,y)}$, with $A_B(\kappa) := K_0 + c_F w_1 (2L^2/\gamma_0)(1 + 1/\kappa)^{3/2}$. The same holds for G, with γ_0 and $\kappa \le \kappa_G$.*

Proof. $G'f = Ff + FW'G'f$. The first term is $\le K_0 \mathrm{e}^{-\kappa d(x,y)}$. For the second, use weighted Cauchy–Schwarz together with $\|\mathrm{e}^{\kappa\rho_y} f\|_2^2 \le (1 + 1/\kappa)^3$. □

Lemma E.46 (Hölder bound for ∇F)**.** *Let $|h(z)| \le \mathrm{e}^{-\kappa d(z,y)} + Z_0$ with $0 \le \kappa \le 1/32$ and $Z_0 \ge 0$, and let $\delta := d(x,x')$ and $\alpha \in [0,1)$. Then*

$$|(\nabla_\mu F h)(x) - (\nabla_\mu F h)(x')| \le C_{\mathrm{H}}(\alpha) \delta^\alpha \big[\mathrm{e}^{-\kappa \min(d(x,y), d(x',y))} + Z_0\big],$$

where $C_{\mathrm{H}}(\alpha) := 2K_1 + \mathrm{e}^{1/8}(38A_1^T + 48A_2^T)(4 + \log 16)(1 + 1/(e(1-\alpha)))$.

Proof. If $\delta \ge 1$, bound each term separately by $K_1(\dots)$. If $\eta \le \delta < 1$, split into the near region $d(z,x) \le 2\delta$, where the contribution is $\le 38 A_1^T \delta$, and the far region, where the second differences give $\le 8A_2^T \delta (d(x,z) + \eta)^{-3} \mathrm{e}^{-\phi(d(x,z)-\delta)}$ and the sum is $\le 6(\log(16/\delta) + 2)$. Finally $\delta^{1-\alpha} \log(1/\delta) \le 1/(e(1-\alpha))$. □

Theorem E.47 (Bałaban's line bounds for the zero-background slices at every scale)**.** *In the setting above, for all $x, x', y \in \Lambda$, directions μ, ν and $\alpha \in [0,1)$:*

- (S1) $|H(x,y)| \le C_0 \mathrm{e}^{-\kappa_H d(x,y)}$;
- (S2) $|\nabla_{x,\mu} H(x,y)|$ *and* $|\nabla_{y,\mu} H(x,y)|$ *are* $\le C_1 \mathrm{e}^{-\kappa_H d(x,y)}$;
- (S3) $|\nabla_{x,\mu} \nabla_{y,\nu} H(x,y)| \le C_2 \mathrm{e}^{-\kappa_H d(x,y)}$;
- (S4) $|\nabla_{x,\mu} H(x,y) - \nabla_{x,\mu} H(x',y)| \le C_3(\alpha) d(x,x')^\alpha \mathrm{e}^{-\kappa_H \min(d(x,y), d(x',y))}$*, and the same for $\nabla_{x,\mu}\nabla_{y,\nu} H$;*
- (S5) *for every lattice path Γ of length $|\Gamma|$, $|\sum_{b\in\Gamma} \eta H(b,y)| \le |\Gamma| C_0 \mathrm{e}^{-\kappa_H \,\mathrm{dist}(\Gamma, y)}$, and likewise with ∇_y.*

The constants are $C_0 := C_V A_B(\kappa_H)$ with $C_V := aA_P(1 + L^{-2}\mathrm{e}^{3L\kappa_G})$; $C_g := C_0 + aA_P + aL^{-2}\mathrm{e}^{3L\kappa'} A'_P$; $C_1 := K_1 C_g$; $C_2 := K_1(C_1 + 2aA_P^\nabla)$; and $C_3(\alpha) := C_{\mathrm{H}}(\alpha) \max(C_g, C_1 + 2aA_P^\nabla)$. They and the rate κ_H depend only on (a,L) (and on α). They are uniform in $\eta \in (0,1]$, $\tilde\mu \in [0,1]$, $\ell \in L\mathbb{N}$, $a_j \in [a_\flat, a]$ and x, y.

Proof. (S1) Resolvent identity: $H = G'(\mathsf{X} - \mathsf{X}')G = G'VG$ with $V := bP_1 - b'P_L$. By (E2), $f := VG(\cdot, y)$ obeys $|f(z)| \le C_V \mathrm{e}^{-\kappa_G d(z,y)}$. Since $\kappa_G \ge 2\kappa_H$ and $\kappa_H \le \kappa'$, Lemma E.45 applies. (S2) From $\mathsf{X}G = \mathsf{X}'G' = 1$ we get $(-\Delta + 1)G = 1 + (1 - \tilde{\mu}^2)G - bP_1 G$, and similarly for G'. Subtracting, the Kronecker deltas cancel exactly:

$$H(\cdot, y) = Fg_y, \qquad g_y := (1 - \tilde{\mu}^2)H(\cdot, y) + bP_1 G(\cdot, y) - b'P_L G'(\cdot, y),$$

with $|g_y(z)| \le C_g \mathrm{e}^{-\kappa_H d(z,y)}$ by (S1) and (E2). Hence $|\nabla_x H| \le K_1 C_g \mathrm{e}^{-\kappa_H d}$, by Lemma E.42. H is symmetric, which gives ∇_y. (S3) $\nabla_{y,\nu} H(\cdot, y) = F(\nabla_{y,\nu} g_y)$, and this is bounded by (S2) and (E4). (S4) Apply Lemma E.46 with $h = g_y/C_g$, respectively $h = \nabla_{y,\nu} g_y/(C_1 + 2aA_P^{\nabla})$, and $Z_0 = 0$. (S5) Sum (S1), or (S2), over the bonds of Γ. □

Remark E.48. No cancellation between slices of different fields or scales is used. The only cancellations are those in (F3) and in the identity $H = Fg_y$. There is no zero mode, because $\mathsf{X}, \mathsf{X}' \ge \gamma_0 L^{-2}$ on all of $\ell^2(\Lambda)$, constants included. No large-block hypothesis enters.

Corollary E.49 ((VS) in physical units)**.** *Let $f \in \{s, v\}$, with v taken per component (Lemma E.38), and let $t_{j+1} \le \min(1/m_f, L)$ on a torus with $\ell_T/t_j \in L\mathbb{N}$. (This holds on the tori T_j whenever $t_{j+1} \le L$.) Then:*

- $|H_{j,f}(x, x')| \le C_0 t_j^{-1} \mathrm{e}^{-\kappa_H d_T(x,x')/t_j}$;
- *each lattice difference in x or x' costs one extra factor t_j^{-1};*
- *the α-Hölder quotient of a derivative is of total order $t_j^{-2-\alpha}$;*
- *a contour sum along Bałaban's $\Gamma^{(j+1)}$ gains one factor t_j, with constant $C_0 d L^2/(L-1)$.*

These are [3, (2.10)–(2.12), p. 426] *at $A = 0$, with $\delta_1 = \kappa_H$, below and above ε_0 alike, uniformly in ε, j, ℓ_T, μ_* and M.*

Proposition E.50 (the last remainder line)**.** *Work in scale-$\bar{K}$ units: $t_{\bar{K}} = 1$, $\eta = \varepsilon/t_{\bar{K}}$, $\ell := \ell_T/t_{\bar{K}} \in \mathbb{N}$, $\rho := m_f t_{\bar{K}} \in (0, 1]$. Put $\tilde{C} := (-\Delta + \rho^2)^{-1}$, $\tilde{G} := (-\Delta + \rho^2 + bP_1)^{-1}$ with $b = a_{\bar{K}}$, and $\tilde{R} := \tilde{C} - \tilde{G}$. In physical units, $R_{\bar{K},f} = C_f - G_{\bar{K},f}(0)$ has kernel $t_{\bar{K}}^{-1} \tilde{R}(x/t_{\bar{K}}, y/t_{\bar{K}})$. Let $\kappa_R := \min(\rho, \kappa_G)/8$ and $Z := (\rho^2 \ell^3)^{-1} = t_{\bar{K}}/(m_f^2 \ell_T^3)$. Then:*

(R1) $|\tilde{R}(x, y)| \le B_R[\mathrm{e}^{-\kappa_R d(x,y)} + Z]$;
(R2) $\eta^3 \sum_y |\tilde{R}(x, y)| \le B'_R \rho^{-2}$;
(R3) $|\nabla_x \tilde{R}|$, $|\nabla_y \tilde{R}|$ *and* $|\nabla_x \nabla_y \tilde{R}|$ *are* $\le B''_R[\mathrm{e}^{-\kappa_R d(x,y)} + Z]$*; Hölder quotients obey (S4) with* $B_R(\alpha)[\mathrm{e}^{-\kappa_R \min d} + Z]$*; contour sums obey (S5).*

The constants $B_R, B'_R, B''_R, B_R(\alpha)$ depend only on (a, L) (and on α), and not on ρ, η or ℓ.

Proof. (R1) $\tilde{R} = \tilde{C}\, bP_1 \tilde{G}$, and $f_y := bP_1 \tilde{G}(\cdot, y)$ obeys $|f_y| \le aA_P \mathrm{e}^{-\kappa_G d(\cdot, y)}$ by (E2) at $\tilde{\mu} = \rho$. Unfold the torus: $|\tilde{R}(x, y)| \le aA_P \sum_n I(\hat{x} - \hat{y} - n\ell)$ with $I(D) := \eta^3 \sum_v \tilde{C}^{\infty}(D - v) \mathrm{e}^{-\kappa_G |v|_1}$. By Lemma E.39 at mass ρ, $I(D) \le A_I \mathrm{e}^{-\kappa_R |D|_1}/(1 + |D|_1)$ with $A_I := A_0(13 + J_0(\kappa_G/2))(3 + 8/\kappa_G)$, which is free of ρ. For $n \ne 0$, $|\hat{x} - \hat{y} - n\ell|_1 \ge \ell |n|_1/2$ and $\#\{|n|_1 = k\} \le 6k^2$, so $\sum_{n \ne 0} I \le 48 A_I/(\kappa_R^2 \ell^3) \le 3072 A_I \kappa_G^{-2} Z$, using $\kappa_R \ge \rho\kappa_G/8$. Hence $B_R := aA_P A_I \max(1, 3072\kappa_G^{-2})$. (R2) $\eta^3 \sum_y |\tilde{R}| \le aA_P J_0(\kappa_G)\, \eta^3 \sum_z \tilde{C}(x - z) = aA_P J_0(\kappa_G) \rho^{-2}$. (R3) $\tilde{R}(\cdot, y) = F\tilde{g}_y$ with $\tilde{g}_y := (1 - \rho^2)\tilde{R}(\cdot, y) + bP_1 \tilde{G}(\cdot, y)$, and $|\tilde{g}_y| \le (B_R + aA_P)[\mathrm{e}^{-\kappa_R d} + Z]$. Then argue as in (S2)–(S4); note $\kappa_R \le \kappa_G/8 \le 1/128$. □

The renormalisation machinery (S6). *Conventions* [3, (2.1)–(2.16), pp. 423–428]. A *generalised graph* has two kinds of lines and vertices:

- lines with slices obeying [3, (2.10)–(2.12)], with an arbitrary line dimension δ_l;
- vertices with η-powers, Hölder factors $|x - x'|^{\gamma}$ ($\gamma \ge 0$) and external fields in $\|\cdot\|_{1,\alpha_0}$.

Fix an ordering $\bar{l} = (l(1), \dots, l(m))$ of the lines, with nested subgraphs $G_1 \subset \cdots \subset G_m = G$. $D(\cdot)$ is the degree of [3, (2.2)–(2.3)], and $J(\bar{l})$ is the set of index assignments compatible with $\bar{l}$. In the *pinned* expression $E^{(x)}$, the vertex v_0 carrying a constant external leg (the pin p) is fixed at x: it has no factor $(L^{j(v_0)}\eta)^d$ and no localisation sum.

Lemma E.51 (the shrinking induction, with or without a pin)**.** *Let G be a connected generalised graph, and assume every connected component $G_i^{(\alpha)}$, $i < m$, has $D(G_i^{(\alpha)}) \geq \vartheta > 0$.*

(i) *Unpinned: $\sum_{j \in J(\bar{l})} \sum_{\{\Delta(v)\}} \tilde{E}(G(j), \{\Delta(v)\}) \leq C_A \sum_{j_{l(m)}} (L^{j_{l(m)}}\eta)^{D(G)}$. This is $\leq C_A$ if $D(G) > 0$, and $\leq C_A (L^{j^*}\eta)^{D(G)}$ when the largest index is fixed to j^*, with no sign condition on $D(G)$.*

(ii) *Pinned: the same bounds hold, with the same C_A, when one vertex v_0 is fixed at a point x.*

Here

$$C_A \leq c_d^m (4/\delta_1)^{dm} (4m+2)^{dm(m-1)/2} (1 - L^{-\vartheta})^{-m},$$

with $m \leq m(\bar{n})$ the maximal number of lines of a graph of order $\leq \bar{n}$, and $\vartheta \geq \min(\alpha_0, \frac{1}{2})$. C_A depends on $d, \delta_1, L, \alpha_0, \bar{n}$ and is independent of η, k, ε, x, the localisations, ℓ_T, μ_* and m^2.

Proof. Write $\tilde{E}$ as in [3, (2.14)]:

- each vertex v carries $(L^{j(v)}\eta)^d$ and a sum over $\Delta(v) \subset \square(v)$;
- each leg of a line l carries $(L^{j_l}\eta)^{\delta_l/2}$, and each derivative on l carries $(L^{j_l}\eta)^{-1}$;
- each line carries $\exp[-\delta^{(1)}(L^{j_l}\eta)^{-1} \operatorname{dist}(\Delta, \Delta')]$, with $\delta^{(1)} := \delta_1/2$ after the tree factor has been extracted.

Induct over i: shrink $l := l(i+1)$, with index j and $s := L^j\eta$, joining w, w'.

(a′) Suppose $w \neq w'$. Let $n' \leq 2m$ be the number of legs of other lines at w', and put $\delta^{(i+1)} := \delta^{(i)}/(2(n'+1))$. For each such line l' (scale $s' \geq s$, other end w''),

$$\mathrm{e}^{-\delta^{(i)} \operatorname{dist}(\Delta(w'), \Delta(w''))/s'} \leq \mathrm{e}^{-\delta^{(i+1)} \operatorname{dist}(\Delta(w), \Delta(w''))/s'} \, \mathrm{e}^{+\delta^{(i+1)} \operatorname{dist}(\Delta(w), \Delta(w'))/s}.$$

The n' costs total at most $\mathrm{e}^{+\frac{1}{2}\delta^{(i)} \operatorname{dist}/s}$. They are paid by l's own factor, which leaves $\mathrm{e}^{-\frac{1}{2}\delta^{(i)} \operatorname{dist}/s}$, and with it $\sum_{\Delta(w')} \leq c_d (\delta^{(i)}/2)^{-d}$. (b) Relocalise the merged vertex using w's volume factor: $\sum_{\Delta(w) \subset \Delta(w \cup w')} (L^j\eta)^d = (L^{j(w \cup w')}\eta)^d$. (c) The remaining powers of $L^j\eta$ multiply to $(L^j\eta)^{D(G_{i+1}^{(\beta)})}$, by additivity of the degree. (d) Summing $j_{l(i+1)} \leq j_{l(i+2)}$ gives the factor $(1 - L^{-\vartheta})^{-1}$. (e) If $w = w'$, only (c)–(d) are needed.

Since $\delta^{(i)} \geq (\delta_1/2)(4m+2)^{-(i-1)}$, the product of the factors $c_d(\delta^{(i)}/2)^{-d}(1 - L^{-\vartheta})^{-1}$ is the displayed C_A. (ii) In step (a) choose w to be the vertex containing v_0. Then w' never contains the pin, and its volume factor is used as before. The pinned merged vertex needs no volume factor, so the exponent attached to the component is again $D(G_{i+1}^{(\beta)})$. At the end the pin contributes 1 in place of $|\square| = 1$. □

Lemma E.52 (largest index fixed)**.** *In the decomposition* [3, (3.4)–(3.5)]*, fix the index of the last line $l(m)$ to j^*. Then every divergent proper subgraph renormalised within an ordering is a G_i with $i < m$, and contains only lines of index $\leq j^*$. The transformations of* [3, (3.5)] *act inside the sub-sum with $j_{l(m)} = j^*$, and Lemma E.51(i) at fixed j^* gives $(L^{j^*}\eta)^{D(G)}$, with no renormalisation of G itself.*

Proof. A connected proper subgraph containing $l(m)$ is not of the form G_i. The transformations of a G_i sum only over internal indices below its lowest external index ([3, pp. 437, 444]), which is $\leq j^*$. The transformed graph has degree $\geq D(G_0)$, and $L^{j^*}\eta \leq 1$. □

Theorem E.53 (leg transport [3, (3.35)] for every member of class C5, every ordering and every receiving vertex)**.** *Let G_0 be a C5 graph ($\phi\phi A$ class) with external legs E. Let $\bar{l}$ be an ordering with*

at most one outer divergent two-vertex component $H = \{a, b; h\}$ of type [3, (2.4)] *(true for every C5 member). Let r be any vertex, and let the pin, if any, be r. Transport the legs by exact telescoping moves*

$$\textstyle\prod_{\ell\in S} F_\ell(x_u) = \prod_{\ell\in S} F_\ell(x_{u'}) + \sum_i [F_{\ell_i}(x_u) - F_{\ell_i}(x_{u'})] \prod_{k<i} F_{\ell_k}(x_{u'}) \prod_{k>i} F_{\ell_k}(x_u),$$

keeping the factors $|x_u - x_{u'}|^\alpha$ unsplit until the end and then splitting them along the path by subadditivity. Use the following schedule:

- *if there is no H, move every leg to r along any path;*
- *if $r = b$, first move a's legs to b along h, then all others;*
- *if $r \notin H$, first move a's legs to b along h, then the legs at b and elsewhere to r.*

Then every remainder either carries $|x_a - x_b|^\alpha$ on h, or has a leg-free with $a \neq r$. In the second case [3, (2.8)] *at a acts on no external leg. After (2.8), every component has degree > 0, with $D_R(H) \geq \alpha$, $D_R(K) \geq \frac{1}{2}$ otherwise, and $D_R(G_0) \geq \alpha$. The local term has all legs at r, and is Bałaban's* [3, (3.36)].

Proof. Moves along the single line h give $+\alpha$ on h, so $D_R(H) = \alpha$. In the later moves no leg sits at a and $a \notin \{c, b, r\}$, so a is leg-free. Then (2.8) at a differentiates only internal lines of G_0 or g. This never lowers a component degree, and it raises $D(H)$ to 1. The chain piece adds α. The components $K \neq H, G_0$ are not divergent, so $D(K) \geq \frac{1}{2}$, since degrees are half-integers. □

Proposition E.54 (the pinned bound [3, (2.21e)] with the pin at the external-photon vertex)**.** *Let S be a (2.21e)-shaped component of a pinned family with $p = L \in S$. Here L is an A1 vertex carrying leg 1 and the line c, and x_1 is an A2 vertex carrying a photon tadpole and an explicit η. Transport S's external leg at x_1 to p along c, as in Theorem E.53 with no H. Then: (i) every remainder carries $|x_1 - p|^\alpha$ on c and has $D_R(S) = \alpha > 0$; (ii) the local term is (the fields of S at p)$\times\Gamma_e(p)$, with $\Gamma_e(p) := \sum_{x_1} \eta^d K_c(p, x_1)\kappa(x_1) \prod g(x_1)$, where $K_c = (D_\mu)_p (D_\nu)_{x_1} c(p, x_1)$ carries one derivative at each end of c and $\kappa(x_1) \propto \eta D^{(<j_0)}(x_1, x_1)$ is the photon tadpole;*

$$|\Gamma_e(p)| \leq C_e(d, L, a, M_b, \alpha_0, \|g\|_{C^1}), \quad \textit{uniformly in } \eta, j_0, k, p.$$

Proof. (i) In the ordering (loop, c) the component {loop} has $D = 0$. Replace it first by its bounded value $|\kappa| \leq O(1)$, using $\eta \sum_j O((L^j\eta)^{-1}) \leq O(1)L/(L-1)$. Then apply Lemma E.51(ii) to the remainders. (ii) First extend the g-weighted x_1-sum to all of $\eta\mathbb{Z}^d$; the support of $\prod g$ lies inside the cube. Next resum S's internal indices up to j_0; this is legitimate because the base p is independent of the ordering and of the indices. Then rescale to $\xi = L^{-j_0}$, with $\rho := |y - y'| + \xi$. Write $c = C^\xi + G^\xi_{j_0}(0)\Upsilon C^\xi$, and $\kappa = \kappa_0 + \kappa_1$ with $\kappa_0 := c_v \xi C^\xi_v(0)$, a constant.

The photon line is $D - C_v = -D(D^{-1} - C_v^{-1})C_v$. Here the middle factor $(\mu_*^2 t^2 - 1) + aQ^*Q$ is bounded and of range $O(1)$: at $A = 0$ Bałaban's and King's photon $G_{j_0}(0)$ is the componentwise plain block average, with no contours. With the vector line bounds this gives $(D - C_v)(y, y) = O(1)$, so $\kappa_1 = O(\xi)$.

Split Γ_e into four pieces:

(1) the translation-invariant piece $\kappa_0 \prod g(p) \sum_y \xi^d K^C(p, y)$ is exactly 0, as a summed lattice difference of a summable kernel;
(2) the g-difference: $|K^C| \leq O(1)\rho^{-3}\mathrm{e}^{-\delta\rho}$ and $|\prod g(y) - \prod g(p)| \leq C\|g\|_{C^1}|y - p|$, so this piece is $O(1)$;
(3) the tadpole difference is $O(\xi) \sum \xi^d \rho^{-3}\mathrm{e}^{-\delta\rho} = O(\xi \log(1/\xi))$;
(4) the line difference: $|K^\Upsilon| \leq O(1)\rho^{-1}\mathrm{e}^{-\delta'\rho}$ (Step 3 of Lemma E.24) and $|\kappa| \leq O(1)$, so this piece is $O(1)$.

□

Lemma E.55 (renormalisation with a pin)**.** *Let G_{ren} be a family each of whose members carries the external leg 1 with $\Phi_{\mathrm{ext}} \equiv 1$ at the pin p. Then every transformation of* [3, pp. 434–445] *can*

be performed without summing over the position of p, and $E^{(x)}(G_{\mathrm{ren}},\dots)$ obeys [3, (1.33)] *with the same $O(1)$ and δ_0.*

Proof. (i) A subgraph containing p has a scalar external leg at p, so it is never a Π/WT class. Those transformations act only on subgraphs not containing p. (ii) A (2.4) subgraph through p has $D=0$, which forces both derivatives onto its internal scalar line. Apply (2.8) at its other vertex $a\neq p$. At most one (2.4) contains p. (iii) Scalar self-energies containing p: Bałaban's rule expands the leg with the smaller index [3, pp. 435–436]. Leg 1 is unit-scale, so the base of the subtraction is p, and the counterterm partner is localised at p. The symmetry $\Sigma_S(y,z)_{ab}=\Sigma_S(z,y)_{ba}$ makes this choice legitimate. (iv) (3.21)-classes containing p: at $\tilde B=0$, a derivative on leg 1 vanishes. At $\tilde B\neq 0$, $D_{\tilde B}1=O(p(t))\neq 0$, and the differentiated leg may be leg 1. Then (3.21)–(3.24) are based at p by the rule of (iii), expanding the other leg. (v) $\phi\phi A$ classes containing p: use Theorem E.53 with $r=p$, and Proposition E.54 for the pinned (2.21e) case. The vanishing [3, (3.37)–(3.38)] is a statement about a translation-invariant C^{ξ}-expression. (vi) Nested and one-particle-reducible cases: induct on the last divergent subgraph; the transformations used never sum over a vertex carrying an external field. (vii) Line cuts and localisation [3, pp. 432–433] act at fixed positions, and a pinned vertex is the localisation $\{x\}\subset\square(p)$. Cuts use the graph representation with partners of *cut* divergent subgraphs. Combining (i)–(vii) with Lemma E.51(ii) gives the claim. □

Remark E.56 (how (S6) is used). In Lemma E.7, every renormalised subgraph has $j_0\le\bar K\le\min_f K_f$, and its internal lines resum to $G_{j_0,f}(0)$, of the same algebraic form for both fields. This is exactly the case in which Lemma E.55, Step 3 of Lemma E.24 and Theorem E.53 apply. Lemma E.8 uses Lemma E.51(ii) in its pinned form, which is uniform in the position of the pin.

The inputs (K1)–(K4), (IH) and (CAP) of the tail bound.

- (K1) For the background $\mathfrak{B}$, regular in the sense of [1, Prop. 2.1] (input (REG)), $G_k(\Omega,\mathfrak{B})$ obeys [4, (1.8), (1.10), (1.12)]: decay $c_0\mathrm{e}^{-\delta_0|x-y|}$, the first derivative and the α-Hölder quotient of DG_kf ($\alpha<1$), uniformly in η; and $G_kQ_k^*$ decays. These hold for $0\le M^2t^2\le c$ in G_k, uniformly there.
- (K2) The slice bounds [3, (2.10)–(2.12)] for the lines of the families of $\mathcal{P}^{(k)}$, obtained in [3, p. 426] from [1, Props. 2.1, 2.3] as $a_j^2t_j^{-4}G_jQ_j^*C^{(j)}Q_jG_j$. At zero background, Corollary E.49 gives an independent proof.

Proposition E.57 ((K3): the one- and two-variable remainder bounds)**.** *Let $V:=G_k(\Omega,\tilde B)^{-1}-G_k(\Omega,\tilde A+\tilde B)^{-1}$, $\mathcal{K}:=[G(\tilde B)V]^nG(\tilde A+\tilde B)[VG(\tilde B)]^{n'}$, $m:=n+n'$ and $\sigma:=e(t)(\|\tilde A\|_\infty+\|\partial^\eta\tilde A\|_\infty)$. Assume the hypothesis (R_ϑ) of Subsection D.2: $\theta(\tilde B),\theta(\tilde A+\tilde B)\le\vartheta_R$ and $\sigma\le 1$. This hypothesis follows from (E.1), read with the constant C_A of $\sigma\le eC_At^{1/2}p$.*

- (a) *For every $m\ge 0$ and $\alpha<1$: $|D^a\mathcal{K}f(x)|$ and the α-Hölder quotient of $D\mathcal{K}f$ are $\le C(C_\infty\sigma)^m\mathrm{e}^{-\beta\,\mathrm{dist}(x,\mathrm{supp}\,f)}\|f\|_\infty$. The same holds for $\mathcal{K}^*$.*
- (b) *Let $0\le h,h'\le 1$ with $|\partial_\mu^\eta h|,|\partial_\nu^\eta\partial_\mu^\eta h|\le c_h$ for all μ,ν, and likewise for h'; the cut-offs of Lemma D.12 qualify. Assume $m\ge 6$ and $0<\alpha<1$, or $m\ge 4$ and $0<\alpha<\frac12$. Then for $0<\beta\le\beta_U$, on T_η,*

$$\|h\mathcal{K}h'\|_{1,\alpha}\le\mathrm{e}^{5\beta}(1+6c_h)^2C_U(\alpha)(C_V\sigma)^m\mathrm{e}^{-\beta\,\mathrm{dist}(\mathrm{supp}\,h,\mathrm{supp}\,h')}$$

in the mixed two-variable norm, including the $D_xD_{x'}^*$ quotients. The same bound holds on the Neumann domains of Proposition D.13(b), under its hypothesis on $\mathrm{supp}\,\tilde A$, *if* $\mathrm{supp}\,h\cup\mathrm{supp}\,h'$ *has depth $\ge R_0+3$. If the quotients over distances >1 are included, multiply the constant by* 4.

The bounds are uniform in η, k, ε, ℓ_T, and in $0\le m^2t^2\le c$. The constants depend on $d,a,M_b,L,\gamma_0,K_W,\alpha,\beta$. In [3, (2.5)]*, the tree rate of* [3, Prop. 1] *is taken $\le\beta_U$.*

Proof. (b) Apply Lemma D.11 with $M:=C_U(\alpha)(C_V\sigma)^m$.

- On T_η, take M from Lemma E.117(b) and $\mathcal{P} := T_\eta$.
- On the Neumann domains, take M from Proposition D.13(b) and $\mathcal{P} := \{z : \text{depth of } z \geq R_0 + 1 + \eta\}$. Points of this $\mathcal{P}$ and their bond partners have depth $\geq R_0$, so Proposition D.13(b) gives (D.3) on $\mathcal{P}$. Since $\eta \leq \frac{1}{2}$, every point within ℓ^∞-distance $1 + \eta$ of the supports lies in $\mathcal{P}$.

(a) Write $D^a \mathcal{K} D^{b*} = (\text{end}) \cdot \mathcal{N} \cdot (\text{end})^*$. In each end, push every vertex derivative onto the line to its right [4, (2.32)], so that every end line carries one derivative. The end factors lie in weighted ℓ^2 when the kernel exponent exceeds $3/2$. This is why two-line ends need $\alpha < \frac{1}{2}$, while three-line ends allow every $\alpha < 1$. The doubly differentiated lines sit in the middle factor $\mathcal{N}$, which is bounded in weighted ℓ^2 by (K4). Cauchy–Schwarz at the two end factors converts ℓ^2 to ℓ^∞, and no pointwise bound on $D_x D_y G$ is used. The details are those of the proof of Proposition D.13, with one end replaced by f, and of Lemmas E.116 and E.117. □

Lemma E.58 ((K4): weighted ℓ^2 bounds)**.** *Let $\rho(x) := |x - y|$ and $O^\beta := \mathrm{e}^{\beta\rho} O \mathrm{e}^{-\beta\rho}$. Assume:*

(H0) $\mathsf{X}(A) - m^2 t_j^2 - v \geq \gamma_0$ [4, (1.8)]*;*
(M1) $\gamma_0' \leq M(A) \leq a_j + aL^{-2}$ [4, (1.15)]*;*
(M2) $|M(A)(y, y')| \leq c_M \mathrm{e}^{-\delta|y-y'|}$.

Then there are $\beta_* > 0$ and $K_W = K_W(d, a, \gamma_0, L)$ such that for $|\beta| \leq \beta_*$, uniformly in $\eta, j, k, \varepsilon, \Omega$, the boundary conditions, regular A, $v \geq 0$ and $s \geq 0$: (i) $\|G^\beta\| \leq 2/\gamma_0$, and $\|(\nabla_A G)^\beta\|$, $\|(G\nabla_A^*)^\beta\|$, $\|(\nabla_A G \nabla_A^*)^\beta\|$ are $\leq K_W$; (ii) $\|Q_j(A)^\beta\|, \|(Q_j(A)^*)^\beta\| \leq \mathrm{e}^{\beta\sqrt{d}}$; (iii) $\|((M+s)^{-1})^\beta\| \leq (\gamma_0'/2 + s)^{-1}$.

Proof. (i) $G^\beta = (\mathsf{X}^\beta)^{-1}$. Lemma D.23 applies when $c(\beta) := d\beta^2 \cosh\beta + a(\mathrm{e}^{\sqrt{d}\beta} - 1) \leq \gamma_0/4$, and gives $\|w\| \leq 2\gamma_0^{-1}\|F\|$ and $\|\nabla_A w\|^2 \leq 4\gamma_0^{-1}\|F\|^2$ for $w = G^\beta F$. Next, $(\nabla_A G)^\beta = \nabla_A^\beta G^\beta$ with $\nabla_A^\beta = \nabla_A + \beta K_\beta$ and $\|K_\beta\| \leq \mathrm{e}^{\beta_*}$ per direction, hence $\leq \sqrt{d}\,\mathrm{e}^{\beta_*}$ for the vector difference (Lemma D.24(b)); and $(G\nabla^*)^\beta = ((\nabla G)^{-\beta})^*$. For $\nabla G \nabla^*$, let $w := G^\beta(\nabla^*)^\beta g$. Then $\mathrm{Re}\langle w, \mathsf{X}^\beta w\rangle \geq \frac{1}{2}\|\nabla_A w\|^2 + \frac{\gamma_0}{4}\|w\|^2$ and $\mathrm{Re}\langle w, \mathsf{X}^\beta w\rangle \leq (\|\nabla_A w\| + \mathrm{e}^{\beta_*}\beta\|w\|)\|g\|$. (ii) Schur test, using $|\rho(x) - \rho(y')| \leq \sqrt{d}$ on the support of Q_j. (iii) By (M2) and Schur, $\|M^\beta - M\| \leq \gamma_0'/2$; then use (M1). □

Definition E.59 ((IH): the family bound carried by $\mathcal{P}_w^{(k)}$)**.** For every localised family $G_{\mathrm{ren}}^\natural$ of $\mathcal{P}_w^{(k)}$,

$$|E(G_{\mathrm{ren}}^\natural, \{\square(v)\}, \Phi_{\mathrm{ext}}, A_{\mathrm{ext}})| \leq O(1)(|u|c_k)^m (e(t_k))^{d_v} (\lambda(t_k))^{d_s} \mathrm{e}^{-\delta_0 d(\{\square(v)\})} \|h\Phi_{\mathrm{ext}}\|_{1,\alpha_0} \|h' A_{\mathrm{ext}}\|_{1,\alpha_0}.$$

Here $\|\cdot\|_{1,\alpha_0}$ is the covariant Hölder norm [3, (1.32), p. 420]; m is the number of D-lines; $c_k \leq 1/3$; $O(1) = O(\alpha_0, \bar{n})$; and $\delta_0 > 0$ is the tree rate of [3, (1.33)]. At the cap it is some $\delta(\bar{N}) > 0$ (Proposition E.60); [3, p. 420] says that it depends on d only, which is not what the proof gives (the line-moving rates of [3, p. 427] depend on $\bar{n}$). Everything is on the small-field restrictions χ_k of [2, (2.55), p. 570]. On $\Lambda_{-1}^{(k-1)\prime}$ these are:

- $|(\partial A)(b)| \leq c_1 p(t_{k-1})$;
- $|A(x)| \leq c_1 p(t_{k-1})/(\mu_* t_{k-1})$;
- $|(D_{\tilde{A}^{(k)}}\phi)(b)| \leq c_1 p(t_{k-1})$;
- $|\phi(x)| \leq c_1 \lambda(t_{k-1})^{-1/4} p(t_{k-1})$.

Proposition E.60 ((CAP): the family bound at a finite order cap)**.** *Fix $\bar{N} = \bar{n}^2 = 169$. Every family of graded coupling order $q \leq \bar{N}$ obeys* [3, (1.33)]*, and its D-line form of Definition E.59. The constant is $O(1) = O(\alpha_0, \bar{N}; d, L, a, M_b)$, with C_A of Lemma E.51 at $m \leq m(\bar{N})$; the tree rate is some $\delta(\bar{N}) > 0$; the u-factor is $(1 + |u|c_k)^{N_v} \leq 5^{84}$ at $L = 2$. These are independent of ε, k, x and the domains. Families containing $\delta m_{\mathrm{fin}}^2$ vertices are uniform in $\ell_T \geq \ell_0$, being evaluated at the torus floor. The conditions (R.cap) and (T.cap) of Appendix A are read at the cap: $R\hat{\ell}^2 \geq C_{\mathrm{loc}}(169)$ and $\ell_T \geq 3r(\varepsilon_0)\varepsilon_0 + C_{169}\varepsilon_0$.*

Proof. Every step of (S6) and of Theorem H.24 is a finite induction over one graph of fixed order, namely the sequence $G_1 \subset \cdots \subset G_m$ with $m \le m(q)$. There is no induction over the order and no series in the order. So the cap enters in four ways only:

(i) finite counts: the number of graphs, $m(\bar N)$ lines, $N_s(\bar N)$ localisation labels, and $N_v \le \bar N/2$ vector lines;
(ii) the line-moving rates $\delta^{(i)} \ge (\delta_1/2)(4m+2)^{-(i-1)}$ of Lemma E.51;
(iii) the constant $\sup_{t\in(0,1]} t^{(q-13)/4+1/20} p(t)^{4q} < \infty$;
(iv) the size $O(\bar n)$ of the component cube inside $\square_1 \subset \square$ [3, pp. 432–433], which is where (R.cap) and (T.cap) tighten with the cap.

Renormalisation does not depend on the cap in $d = 3$. With $\omega = 3 - E/2 - \alpha/2 - \beta$, every superficially divergent graph has $\alpha + 2\beta \le 6 - E \le 6 < \bar n$, so no counterterm object is re-read at the cap. All new numbers are fixed at stage 4 of Appendix A, before ε_0 is chosen, so the order of choices stays acyclic. □

E.4. **The ingredients.** We prove the ingredients of the proof of Theorem 5.2 listed in Table 1: the moved vacuum pieces (Proposition E.64), the Ward–Takahashi step on the torus (Proposition E.67), the use of real backgrounds only (Proposition E.71), the order cap (Proposition E.76), the conditions of [3, Prop. 1] at its applications (Proposition E.110), the mass reading of [2, p. 589] (Proposition E.159), the final c-numbers (Proposition E.162) and the vacuum classes at the last scale (Proposition E.175). The per-localisation bound, the geometric and region-sum lemmas, and [3, Prop. 1] with its sketched steps written out are stated in Subsection E.5. Throughout: $d = 3$, $L = 2$, $\bar n = 13$, $\bar N = 169$, $t_k = L^k\varepsilon$ with $t_K = \varepsilon_0$, $r(t) = R(1 + \log t^{-1})^{\varrho}$ with $\varrho = 2$ (Appendix A), and the two points $\pi_+ = (u, M) = (1, m_2)$, $\pi_- = (-\eta, m_1)$ of (5.2).
The moved vacuum pieces at every scale. *Notation at step k.* The scale-k blocks are the unit cubes of the lattice rescaled by t_k. For a union X of such blocks, $|X|_k$ is their number. Let $\Lambda_7^{(k)}$ be Bałaban's small-field set at step k [2, pp. 557–558], and put $W_k := B^k(\Lambda_7^{(k)})$. The *layer* is $\mathcal{L}_k := \{\square \subset W_k : \operatorname{dist}(\square, W_k^c) \le r(t_k)\}$, which equals $B^k(\Lambda_7^{(k)} \cap \Lambda_8^{(k)c})$ of [3, p. 422]. The *deep part* is $\mathcal{D}_k := W_k \setminus \mathcal{L}_k$. The *collar* is $\mathrm{Col}_k := \Lambda_7^{(k-1)\prime} \cap \Lambda_7^{(k)c}$, and $\mathrm{Old}_k := \Lambda_7^{(k-1)\prime c}$. Since $\Lambda_7^{(k)} \subset \Lambda_7^{(k-1)\prime}$, we have $W_k^c = \mathrm{Col}_k \sqcup \mathrm{Old}_k = \Lambda_7^{(k)c}$. Let $E_K(u)$ be the vacuum counterterm of the slice model, and $E_{K,k}(u)$ its part from graphs with operator indices in $[k, K)$ [3, (1.27)–(1.28), p. 418], with $E_{K,0} = E_K$ and $E_{K,K} = 0$. *Convention (G)* is Bałaban's [3, (1.30), p. 419]: at step $k \ge 1$, $E_{\mathrm{fin}}(u)$ and $E_{K,k}(u)$ are global c-numbers, with no Ω_1 and no g_k. The set moved at step k is

$$\mathcal{V}_k(u) := E_{K,k}(u) - E_{K,k+1}(u),$$

the terms whose indices are all $\le k$, at least one of them equal to k [3, p. 422].

Fix a pair $(G, \tilde l)$ of a graph and one of Bałaban's orderings of its lines, whose last line has index k. Its terms are localised by the rule of [3, p. 420]: there is one unit cube per vertex of type (1.6)/(1.7), and one support cube of side 2 per vector leg of the other vertices. We write $\Lambda = (\square_\iota)_{\iota\in I(G)}$, $N := |I(G)| \le \bar N(\bar n) < \infty$. Labels at one vertex have overlapping supports (otherwise the term vanishes). So they are at distance 0, and the tree length $d(\Lambda)$ of [3, (1.33)] is the same over labels and over vertices. Order $I(G)$ so that the two ends of the last line come first. Then:

- Λ is *good* if $\Lambda \subset W_k$;
- Λ is *bad* if it meets both W_k and W_k^c; its anchor is the first index whose cube lies in W_k;
- Λ is *outside* if $\Lambda \cap W_k = \emptyset$; its anchor is the first index.

Every Λ has exactly one class and one anchor. If the moved terms are localised at scale t_{k+1} rather than t_k (the two readings of "passing from k to $k+1$", p. 422), block counts change by factors $L^{\pm 3}$ and the constants below by at most L^3.

Lemma E.61 (the localisation sum)**.** *Let $N \geq 1$ and $\delta > 0$. On the scale-k block lattice $(\mathbb{Z}/n)^3$ of T_ε, fix an anchor index ι_0 and a cube $\square_0$. Then for every $D_0 \geq 0$,*

$$\sum_{\Lambda:\ \square_{\iota_0}=\square_0,\ d(\Lambda)\geq D_0} \mathrm{e}^{-\delta d(\Lambda)} \leq \mathrm{e}^{-\delta D_0/2} S_N(\delta),$$

$$S_N(\delta) := N^{N-2}\big(\mathrm{e}^{\delta}\sigma(\delta)\big)^{N-1},$$

where $\sigma(\delta) := 1 + \sum_{r\geq 1}(24r^2+2)\mathrm{e}^{-\delta r/2}$. The bound does not depend on n, i.e. on ℓ_T.

Proof. On $\{d \geq D_0\}$ we have $\mathrm{e}^{-\delta d} \leq \mathrm{e}^{-\delta D_0/2}\mathrm{e}^{-\delta d/2}$. Next, $d(\Lambda) = \min_{\mathcal{T}} L_{\mathcal{T}}(\Lambda)$ over the spanning trees $\mathcal{T}$ on the N labelled indices, where $L_{\mathcal{T}}$ is the sum of the cube distances over the edges of $\mathcal{T}$. Hence $\mathrm{e}^{-\delta d/2} = \max_{\mathcal{T}} \mathrm{e}^{-\delta L_{\mathcal{T}}/2} \leq \sum_{\mathcal{T}} \mathrm{e}^{-\delta L_{\mathcal{T}}/2}$, and by Cayley's formula there are N^{N-2} trees. Fix a tree, root it at ι_0, and sum the free cubes from the leaves inwards. Cube centres c, c' satisfy $\operatorname{dist}(\square, \square') \geq (|c-c'|_\infty - 2)_+$, so each edge contributes at most $\sum_{c'} \mathrm{e}^{-(\delta/2)(|c-c'|_\infty-2)_+} \leq \mathrm{e}^{\delta}\sum_w \mathrm{e}^{-(\delta/2)|w|_\infty}$. The minimal-image map $(\mathbb{Z}/n)^3 \to \mathbb{Z}^3$ is injective, so this is at most the $\mathbb{Z}^3$ sum. Since $\#\{w \in \mathbb{Z}^3 : |w|_\infty = r\} = (2r+1)^3 - (2r-1)^3 = 24r^2 + 2$, the $\mathbb{Z}^3$ sum is $\sigma(\delta)$. There are $N-1$ edges. □

The per-localisation bound (Lemma E.86). Put $\delta := \min(\delta_T, \delta_1)$, with $\delta_T > 0$ the tree rate of [3, (1.33)] as given by Proposition E.97 (it depends on $\bar{n}$; it is not Bałaban's "δ_0 depends on d only") and δ_1 the rate of [3, (2.10), p. 426]. Define

$$A_k := C_P(1 + |u|\bar{c})^{\bar{N}_{\mathrm{ph}}} P(t_k), \qquad P(t) := \sum_{2\leq\alpha+2\beta\leq 2\bar{n}} |e(t)|^\alpha \lambda(t)^\beta,$$

where $e(t)^2 = e^2 t$ and $\lambda(t) = \lambda t$, $\bar{c} := \kappa\mu_*^2\varepsilon_0^2 L^2$, $\bar{N}_{\mathrm{ph}}$ is the largest number of photon lines, and C_P is the $O(1)$ of [3, (1.33)] times that of [3, (2.10)–(2.11)]. Lemma E.86 states that for every $(G, \tilde{l})$ and every Λ,

$$|\mathcal{V}_{G,\tilde{l}}(\Lambda)| \leq A_k\, \mathrm{e}^{-\delta d(\Lambda)}. \tag{E.4}$$

The proof cuts the line of index k as a two-variable external field [3, p. 419, (2.10)–(2.11) p. 426]. It then uses the list of the resulting two-leg classes and the representation of opened δm^2 vertices. The one-particle-irreducible scalar two-leg classes get their δm^2 partners, which are already inside E_1. The one-particle-reducible members of C2 (21 of the 59 non-vanishing classes) have no partner; they are handled by the product mechanism of Step 3 of the proof of Lemma E.86. The photon two-leg classes C3, C4 need the Ward–Takahashi class, valid on T_ε after the transfer of [3, p. 433] (Proposition E.67). In every non-vanishing term the cut graph stays connected (scalar lines are never bridges, and photon bridges give vanishing pairs), so (1.33) of the cut graph applies. The u-part is (E.4) with $(1+|u|\bar{c})^{\bar{N}_{\mathrm{ph}}}$ replaced by $\bar{N}_{\mathrm{ph}}|u|\bar{c}(1+\bar{c})^{\bar{N}_{\mathrm{ph}}-1}$. Since every vacuum graph has $o = \alpha/2 + \beta \geq 1$, $|e(t)|^\alpha\lambda(t)^\beta = |e|^\alpha\lambda^\beta t \cdot t^{o-1} \leq |e|^\alpha\lambda^\beta\varepsilon_0^{o-1} t$ for $t \leq \varepsilon_0 \leq 1$. Hence

$$P(t_k) \leq C_e t_k, \qquad C_e := \sum |e|^\alpha \lambda^\beta \varepsilon_0^{o-1},$$

and the rate exponent is 1, not the collar exponent $\kappa_0 = \frac{1}{5}$ used elsewhere.

Proposition E.62 (bad and outside localisations)**.** *Assume* (E.4). *Put $\bar{S} := \sum_{(G,\tilde{l})} S_{N(G)}(\delta)$, the sum over the graph–ordering pairs of order $\leq \bar{n}$; it depends on $\bar{n}$, d, δ only. At both points and every $0 \leq k \leq K-1$:*

(i) $\big|\sum_{\Lambda\ \mathrm{bad}} \mathcal{V}_k(\Lambda)\big| \leq \bar{N}\bar{S}A_k\big[|\mathcal{L}_k| + \mathrm{e}^{3\delta}\mathrm{e}^{-\delta r(t_k)/2}|\mathcal{D}_k|\big]$;

(ii) *for every union X of scale-k blocks of W_k^c,* $\big|\sum_{\Lambda\ \mathrm{outside},\ \mathrm{anchor}\in X} \mathcal{V}_k(\Lambda)\big| \leq \bar{S}A_k|X|_k$;

(iii) *(i) and (ii) hold for $\mathcal{V}_k(u) - \mathcal{V}_k(0)$ with A_k replaced by $A_k^u \leq \bar{N}_{\mathrm{ph}}|u|\bar{c}(1+\bar{c})^{\bar{N}_{\mathrm{ph}}-1}C_P P(t_k)$.*

The bounds do not depend on ε, K or ℓ_T, they are linear in $|u|$ with no sign used, and they are finite at $M = m_1$.

The proof uses [3, (1.33)] in the form of Proposition E.76 and [3, (2.10)–(2.11), p. 426], through (E.4). Lemma E.170 is a second form of the same localisation sum.

Proof. (i) There are at most $\bar{N}$ choices of the anchor index. A bad Λ has a cube outside W_k, so its tree joins $\square_0$ to W_k^c and $d(\Lambda) \geq \operatorname{dist}(\square_0, W_k^c) - 3$. Apply (E.4) and Lemma E.61 with $D_0 = (\operatorname{dist}(\square_0, W_k^c) - 3)_+$. For $\square_0 \in \mathcal{L}_k$ use $\mathrm{e}^{-\delta D_0/2} \leq 1$. For $\square_0 \in \mathcal{D}_k$ we have dist $> r(t_k)$, so $\mathrm{e}^{-\delta D_0/2} \leq \mathrm{e}^{3\delta/2}\mathrm{e}^{-\delta r(t_k)/2} \leq \mathrm{e}^{3\delta}\mathrm{e}^{-\delta r(t_k)/2}$. Then sum over $(G, \tilde{l})$. (ii) Apply (E.4) and Lemma E.61 with $D_0 = 0$. The anchor is fixed and the anchor cube runs over X; the condition $\Lambda \cap W_k = \emptyset$ only removes terms from an absolute sum. (iii) The same argument, with the u-part of (E.4). For uniformity: $\bar{S}$ and $\bar{N}$ contain no sum over steps; the torus count is at most the $\mathbb{Z}^3$ count; and C_P does not depend on the domains [3, p. 421]. □

The region-sum lemma (Lemma E.95, under its hypothesis (H-LF)). The large-field region sum of the inductive bound of the proof of Theorem 5.2, multiplied by $\exp\big(c_A \sum_{k<K}(1+\log t_k^{-1})|\Lambda_7^{(k)c}|_k\big)$, is $\leq \exp\big(2(d+2)\Sigma_*(\varepsilon_0;\pi)|T|\big)$ for any $c_A \geq 0$. The hypothesis is the upper bound (E.comb) on ε_0: $\hat{\ell}^4 \geq 2c_A C_V R^d/(c_* b_0^2 \ln L)$, with $\hat{\ell} = 1 + \log \varepsilon_0^{-1}$. Here $\Sigma_*(\varepsilon_0;\pi) = \sum_{n \geq n_0} L^{3n} \exp(-\frac{1}{2} c_* b_0^2 (1 + n \log L)^{2p})$ contains no c_A.

Lemma E.63 (absorption of the outside pieces)**.** *Put $C_{\mathrm{old}} := \bar{S} C_P C_e (1+\bar{c})^{\bar{N}_{\mathrm{ph}}}$. At both points,*

$$\sum_{k<K} \bar{S} A_k |\Lambda_7^{(k)c}|_k \leq C_{\mathrm{old}} \sum_{k<K} (1 + \log t_k^{-1}) |\Lambda_7^{(k)c}|_k.$$

Hence, under (E.comb) read with $c_A + C_{\mathrm{old}}$, the outside pieces cost at most $\exp(2(d+2)\Sigma_*(\varepsilon_0;\pi)|T|)$ in the region sum. C_{old} does not depend on ε, K, k, ℓ_T, b_0, R or p, and it is fixed at stage 10a of Appendix A with $\bar{c}$ and C_e taken at their suprema over $(0, \varepsilon_1]$. Along the collars the outside pieces cost $2C_{\mathrm{old}}\varepsilon_0$ per block of Col_i.

Proof. The outside-anchored set is $X = W_k^c = \mathrm{Col}_k \sqcup \mathrm{Old}_k = \Lambda_7^{(k)c}$. Proposition E.62(ii) gives $\bar{S} A_k |\Lambda_7^{(k)c}|_k$. For $|u| \leq 1$ we have $A_k \leq C_P (1+\bar{c})^{\bar{N}_{\mathrm{ph}}} C_e t_k$, and $t_k \leq \varepsilon_0 \leq 1 \leq 1 + \log t_k^{-1}$. This gives the display. Lemma E.95 with $c_A \mapsto c_A + C_{\mathrm{old}}$ gives the cost bound. For the collar form: every scale-k block of Old_k meets $\bigcup_{i<k} \mathrm{Col}_i$, and the blocks are dyadically nested, so $|\mathrm{Old}_k|_k \leq \sum_{i<k} |\mathrm{Col}_i|_i$. Moreover $\sum_{k=i+1}^{K} t_k \leq \varepsilon_0 \sum_{n \geq 0} L^{-n} = \frac{L}{L-1}\varepsilon_0 \leq 2\varepsilon_0$. □

Proposition E.64 (the moved vacuum pieces)**.** *Assume convention (G) and $R \geq 5$, and let $\mathcal{V}_k$ be as in Definition E.12. Let $1 \leq k \leq K-1$, and work at either of the points $\pi_\pm$. The moved set $\mathcal{V}_k(u)$ is localised once, at its own scale k. Each localisation is exactly one of good, bad or outside, and has one anchor, by the anchor rule of the paragraph containing Lemma E.61. Write $\mathcal{V}_k = \mathcal{V}_k^{\mathrm{good}} + \mathcal{V}_k^{\mathrm{bad}} + \mathcal{V}_k^{\mathrm{out}}$ accordingly. Then:*

(i) good: *$\mathcal{V}_k^{\mathrm{good}}$ is a sum of terms with all cubes in W_k. It equals the generated step-k vacuum family and cancels against it.*

(ii) bad: $|\mathcal{V}_k^{\mathrm{bad}}| \leq \big(\bar{N}\bar{S}A_k + 4C_{\mathrm{loc}}t_k\big)|\mathcal{L}_k|_k + \big(\bar{N}\bar{S}A_k \mathrm{e}^{3\delta}\mathrm{e}^{-\delta r(t_k)/2} + 2C_{\mathrm{dp}} t_k^3 \mathrm{e}^{-2\delta_1(r(t_k)-4)}\big)|\mathcal{D}_k|_k$.

(iii) outside: *for every union X of scale-k blocks of W_k^c, the part of $\mathcal{V}_k^{\mathrm{out}}$ with anchor in X is at most $(\bar{S}A_k + 2C_{\mathrm{loc}}t_k)|X|_k$. In particular $|\mathcal{V}_k^{\mathrm{out}}| \leq C'_{\mathrm{old}} t_k |\Lambda_7^{(k)c}|_k$, where $C'_{\mathrm{old}} := C_{\mathrm{old}} + 2C_{\mathrm{loc}}$. Suppose that the hypothesis (H-LF) of Lemma E.95 holds, and that condition (E.comb) holds with c_A replaced by $c_A + C'_{\mathrm{old}}$. Then the factor $\exp\big(\sum_k |\mathcal{V}_k^{\mathrm{out}}|\big)$ costs at most $\exp\big(2(d+2)\Sigma_*(\varepsilon_0;\pi)|T|\big)$ in the region sum of that lemma.*

(iv) u-parts: *(ii) and (iii) hold for $\mathcal{V}_k(u) - \mathcal{V}_k(0)$, with A_k replaced by A_k^u of Proposition E.62(iii).*

(v) *No term is moved twice. $E_{\mathrm{fin}}(u)$ lies in no $\mathcal{V}_k$, and it remains a global factor that is paid once.*

(vi) *The sum over k of the $|\mathcal{D}_k|_k$-terms of (ii) is at most $C|T|$, uniformly in K.*

Here δ, δ_1 and A_k are as in Lemma E.86, and $\bar N$, $\bar S$ as in Proposition E.62. C_{loc} and C_{dp} are the constants of Theorem E.92 with $R_0 = 1$, C_{old} is the constant of Lemma E.63, and Σ_*, c_A are as in Lemma E.95. All constants are bounded uniformly in ε, K and $\ell_T \geq \ell_0$, and they are finite at $u = -\eta$ and $M = m_1$.

The proof uses Lemma E.86, which rests on Proposition E.97 and on Proposition E.67. It also uses [3, p. 433], whose remainder on the cube $\square$ is treated in Proposition E.110, and Theorem E.92. Throughout, convention (G) is in force.

Proof. (v) By Lemma E.14(i), the sets $\mathcal{V}_k$ partition the terms of $E_1 - E_{\mathrm{fin}}$. A term lies in the set $\mathcal{V}_k$ whose k is the largest index of its lines, so the telescoping $E_K = E_{K,k} + \sum_{j<k} \mathcal{V}_j$ moves each term once, at its top scale. The terms of E_{fin} have a line of index K [3, (1.26), p. 418], so they lie in no $\mathcal{V}_k$. E_{fin} is bounded once, as a global factor (Proposition E.21).

The three classes partition the localisations, and each localisation has one anchor, by the anchor rule.

(i) A good localisation has all its cubes in W_k. By Lemma E.14(ii), $\mathcal{V}_k$ is the field-independent part of order $\leq \bar n$ of the output of step k. The cancellation of $\mathcal{V}_k^{\mathrm{good}}$ against the vacuum output generated at step k is Lemma F.31 together with item (C11) of Lemma F.32. It is used in the closing step, Lemma F.39.

(ii)–(iv) Consider first all graph–ordering pairs other than the order-e^2 family with one D-line. For them, Lemma E.86 gives (E.4), so Proposition E.62(i)–(iii) apply, since their proofs use only (E.4) and Lemma E.61. The order-e^2 family with one D-line contributes the terms with C_{loc} and C_{dp}, by Lemma E.93. That lemma applies with $R_0 = 1$, because $r(t) = R(1 + \log t^{-1})^2 \geq R \geq 5$ for $t \leq 1$. For the second claim of (iii), take $X = W_k^c = \Lambda_7^{(k)c}$, and use $\bar S A_k \leq C_{\mathrm{old}} t_k$ for $|u| \leq 1$ (Lemma E.63). For the absorption, use $t_k \leq \varepsilon_0 \leq 1 \leq 1 + \log t_k^{-1}$. This gives $\sum_k |\mathcal{V}_k^{\mathrm{out}}| \leq C'_{\mathrm{old}} \sum_k (1 + \log t_k^{-1}) |\Lambda_7^{(k)c}|_k$, which is the cost term of (H-LF) with c_A replaced by $c_A + C'_{\mathrm{old}}$. Lemma E.95 with $c_A + C'_{\mathrm{old}}$ then gives the stated cost.

(vi) We have $|\mathcal{D}_k|_k \leq |T| t_k^{-3}$ and $A_k \leq C t_k$. Since $r(t) = R(1 + \log t^{-1})^2$, for every $q > 0$ there is a constant C_q with $\mathrm{e}^{-\delta r(t)/2} + \mathrm{e}^{-2\delta_1 (r(t) - 4)} \leq C_q t^q$. Take $q = 3$. The deep sum is then at most $C|T| \sum_k t_k \leq 2C\varepsilon_0 |T|$.

Uniformity: $\bar S$ and $\bar N$ contain no sum over steps. On the torus, the count of Lemma E.61 is at most the count on $\mathbb{Z}^3$. C_P is independent of the domains [3, p. 421]. Finally, C_{loc} and C_{dp} are bounded uniformly in ε, k and $\ell_T \geq \ell_0$ by Theorem E.92. $\square$

The Ward–Takahashi step on the torus.

Lemma E.65 (a constant background is a pure gauge on a non-wrapping set)**.** *Let $\Omega \subset T_\varepsilon$ be a set of sites with the bonds between them. Assume that Ω lifts isometrically: there is $\iota : \Omega \to \varepsilon\mathbb{Z}^3$ with $\pi \circ \iota = \mathrm{id}$ and $\iota(x + \varepsilon e_\mu) = \iota(x) + \varepsilon e_\mu$ on every bond of Ω. This holds if Ω lies in a box of side $< \ell_T$. Then for every $a \in \mathbb{R}^3$ the constant background on the bonds of Ω equals $\nabla\chi$ with $\chi := a \cdot \iota$. Hence:*

- *every gauge-covariant quantity built only from fields and operators on Ω is independent of a, up to phases $\mathrm{e}^{\pm ie\chi}$ on its external scalar legs;*
- *the localised zeroth-moment identity $\sum_{b' \subset \Omega} \Pi_\Omega^{\mathrm{loc}}(b, b') = 0$ holds exactly on T_ε for every class supported in Ω.*

This is Bałaban's gauge-out of the constant field B_0 [3, p. 433], written for the torus.

Proof. χ is well defined because ι is injective, and $\nabla_\mu\chi(x) = a\cdot(\iota(x+\varepsilon e_\mu)-\iota(x))/\varepsilon = a_\mu$ on the bonds of Ω. The gauge transformation $\varphi \mapsto \mathrm{e}^{ie\chi}\varphi$, $A \mapsto A - \nabla\chi$ on Ω removes a from every object supported in Ω. For a class supported in Ω the second a-derivative is therefore 0. The second a-derivative of a localised two-photon class at zero momentum is $\sum_{b'\subset\Omega}\Pi^{\mathrm{loc}}(b,b')$. □

Lemma E.66 (the cube of [3, p. 433] does not wrap)**.** *Assume the conditions (T.nw), $\ell_0 \geq \varepsilon_0[8r(\varepsilon_0)+4M_b+2]$, and (E.r), $\varepsilon_0 < \mathrm{e}^{1-\varrho} = \mathrm{e}^{-1}$, of Appendix A, and let $\ell_T \geq \ell_0$. Then for every k with $t_k \leq \varepsilon_0$, a cube of side $3r(t_k)$ in scale-k units (physical side $3r(t_k)t_k$) lies in a box of side $< \ell_T$. In particular it lifts isometrically.*

Proof. With $s = \log t^{-1}$, $\frac{d}{dt}\big[tR(1+s)^2\big] = R(\log^2 t - 1) > 0$ for $t < \mathrm{e}^{-1}$. So $t \mapsto t\,r(t)$ increases on $(0,\varepsilon_0]$, and $3t_k r(t_k) \leq 3\varepsilon_0 r(\varepsilon_0) < 8\varepsilon_0 r(\varepsilon_0) \leq \ell_0 \leq \ell_T$. □

Proposition E.67 (the Ward–Takahashi step on the torus)**.** *At every scale with $t_k \leq \varepsilon_0$, localise each renormalised class as in* [3, p. 433]*: in a cube $\square_1$ of size $r(L^k\varepsilon)$ containing its $O(\bar{n})$-cluster, inside a cube $\square$ of size $3r(L^k\varepsilon)$. Then, on T_ε:*

- *the gauge-out of the constant field B_0 and the replacement $G_k(\square,0) = G_k(0)+\delta G_k(\square,\eta\mathbb{Z}^d,0)$ are valid;*
- *every Ward–Takahashi identity used for the photon-cut classes C3, C4 holds exactly, with no torus image defect;*
- *the torus enters only through the external fields δG_k that* [3, (1.33)] *already carries.*

Bałaban's renormalisation [3, pp. 434–445] *and the vacuum case of Proposition E.62 use the same torus input.*

The proposition uses [3, p. 433] and the condition (T.nw).

Proof. By Lemma E.66, $\square$ lifts isometrically. By Lemma E.65, B_0 is a gradient on $\square$, and its removal is the gauge transformation of [3, p. 433], after which "we get the same expressions as above with $B_0 = 0$ only". Since $\square$ lifts, it can be identified with a cube of $\eta\mathbb{Z}^d$. There the Ward–Takahashi identities of free boundary conditions hold exactly (the step S8 of Table 2), and the difference $\delta G_k(\square,\eta\mathbb{Z}^d,0)$ is treated as an external field, as in the source. So no Ward–Takahashi step runs on a wrapping domain. □

Only real gauge backgrounds are used. A *gauge background* is any vector-field or group configuration at which one of the following is evaluated: an operator $G_k(\Omega,\cdot)$, $C_k(\Omega,\cdot)$, a covariant Laplacian, a transporter $U(\cdot)$, a determinant or Gaussian measure, or an expansion in the background. It is *real* if $A \in \mathbb{R}$ on every bond, equivalently $U(A) \in \mathrm{SO}(N)$. The *allowed* complex objects are: the entire function $R_{\bar{n}+1}(z)$ evaluated at real A; momentum continuations at $A = 0$ [4, p. 586]; complex sources and Cauchy estimates in sources or in u; the complex structure of the charged scalar.

Definition E.68 (gauge backgrounds)**.** In Definition E.68, Lemmas E.69, E.70, E.99 and Proposition E.71, $N = 2$, q is the real antisymmetric 2×2 matrix with $q^2 = -1$ of [4, (1.2), p. 572], $e > 0$ is the charge, $\eta = L^{-k}$ and $U(A) = \mathrm{e}^{qe\eta A}$. Vector fields are real-valued functions on bonds [4, p. 572]. Scalar fields take values in $\mathbb{R}^2$, and q acts on $\mathbb{R}^2$ as the complex structure.

(a) A *gauge background* is a configuration A of complex numbers on the bonds of a subset of T_η, or the family $\{U(A_b)\}_b$ it defines, at which one of the following is evaluated: the transporters $U(A(\Gamma))$; $Q_k(A)$ and $P_k(A) = Q_k^*(A)Q_k(A)$ [4, (1.4)–(1.5)]; the Neumann covariant Laplacian $-\Delta_{A,\Omega}^{\eta,N}$ [4, (1.3)]; $G_k(\Omega,A) = (-\Delta_{A,\Omega}^{\eta,N} + m^2 + aP_k(A))^{-1}$ [4, (1.6)] and $C_k(\Omega,A)$ [4, p. 586]; any operator obtained from these by sums, products, restrictions and inverses; a determinant or Gaussian measure defined by such an operator; or an expansion in powers of $A-B$ around a background B.

(b) A gauge background is *real* if $A_b \in \mathbb{R}$ for every bond b.

(c) The following complex quantities are not gauge backgrounds: the value of the entire function $R_{n+1}(z) := (n+1)\int_0^1 (1-t)^n \mathrm{e}^{tz}\,dt$ [1, (3.14), p. 614] at a matrix $qe\eta A$ with A real; complex momenta in Fourier representations of operators at $A = 0$ or of Feynman integrands; complex test functions and complex parameters multiplying sources; the complex structure q.

Lemma E.69 (realness is preserved by the operations of the construction)**.** *Let $\Omega \subset T_\eta$ be a union of blocks and let A, A_1, A_2 be real vector fields.*

(a) *$U(A(\Gamma))$, $Q_k(A)$ and $P_k(A)$ are real matrices. $X_\Omega(A) := -\Delta^{\eta,N}_{A,\Omega} + m^2 + aP_k(A)$ is real symmetric and nonnegative, and whenever it is invertible (in particular under* [4, (1.8), p. 573]*) $G_k(\Omega, A)$ is real symmetric. The same holds for every operator obtained from these by sums, products, restrictions to subsets of Ω, adjoints and inverses, in particular for $C_k(\Omega, A)$.*

(b) *The following are real vector fields: TA for a linear map T with real matrix (for instance Q^*, $C^{(0)}Q^*$, block averages $\bar{A}_b = L^{-k}\sum_{\langle x,x'\rangle \subset b} A_{\langle x,x'\rangle}$); $(1-\theta)A_1 + \theta A_2$ for real θ; $A_1 + A_2$; restrictions and extensions by zero; $A + \partial^\eta \lambda$ for real λ. Contour sums $A(\Gamma)$ are real.*

(c) *Let $\square$ be a set of sites, $\tilde{B}$ a real vector field on $\square$ and $B_0 \in \mathbb{C}^d$ a constant field with $\max_\mu \sup_{x\in\square} |\tilde{B}_\mu(x) - B_{0,\mu}| \le \delta$. Then $\operatorname{Re} B_0 \in \mathbb{R}^d$ satisfies the same inequality. If $\square$ does not wrap around the torus, the constant field $\operatorname{Re} B_0$ equals $\partial^\eta \lambda$ on $\square$ for the real function $\lambda(x) = \sum_\mu (\operatorname{Re} B_{0,\mu}) x_\mu$.*

(d) *If P is a real symmetric positive definite operator on $(\mathbb{R}^n)^\Lambda$, $N(0, P^{-1})$ is a probability measure on $(\mathbb{R}^n)^\Lambda$. For the slice model at real $u > -1/\bar{\delta}$ the photon covariance $C_B(u) = C_{\mu_*} + u\Gamma^B$ of Subsection 5.1 is real symmetric and positive definite; this covers $u \in \{1, -\eta\}$, since $\eta\bar{\delta} < 1$ (Lemma 5.4(a)). The weights $1 + u\delta_j$ are real and positive there.*

Proof. (a) $U(A(\Gamma))$ is real by Lemma E.99(c); hence so is $Q_k(A)$, a sum of terms $\eta^d U(A(\Gamma))$ [4, (1.4)], and $P_k(A) = Q_k^* Q_k$. The quadratic form $\sum_{b\subset\Omega} \eta^{d-2} |U(A_b)\phi(b_+) - \phi(b_-)|^2$ of [4, (1.3)] is real and nonnegative on real $\phi : \Omega \to \mathbb{R}^N$, so $-\Delta^{\eta,N}_{A,\Omega}$ and $X_\Omega(A)$ ($m^2 \ge 0$, $a > 0$) are real symmetric and nonnegative. Sums, products, restrictions and adjoints of real matrices are real, and the inverse of an invertible real symmetric matrix is real symmetric. (b) Each operation is a real-linear map with real coefficients. (c) For real t and complex c, $|t - \operatorname{Re} c| = |\operatorname{Re}(t-c)| \le |t-c|$. If $\square$ does not wrap, it lifts isometrically to $\eta\mathbb{Z}^d$ (Lemma E.66), and there $\partial^\eta_\mu \lambda = \operatorname{Re} B_{0,\mu}$ (Lemma E.65). (d) $N(0, P^{-1})$ has the density $(\det P/(2\pi)^{n|\Lambda|})^{1/2} \mathrm{e}^{-\frac{1}{2}\langle z, Pz\rangle}$. C_{μ_*} and Γ^B are real symmetric with $0 \le \Gamma^B \le \bar{\delta} C_{\mu_*}$, so $C_B(u) \ge (1 + \min(u, 0)\bar{\delta}) C_{\mu_*} > 0$ for $u > -1/\bar{\delta}$. Finally $1 + u\delta_j \ge 1 - \eta\bar{\delta} > 0$ at $u = -\eta$, since $0 \le \delta_j \le \bar{\delta}$. □

Lemma E.70 (the backgrounds of the construction are real)**.** *Consider the inductive construction of* [2, Sects. 2–3] *on the torus for the slice model at $(u, M) \in \{(1, m_2), (-\eta, m_1)\}$, as used in the proof of Theorem 5.2 (Subsection F.1). At every step k the following are real vector fields:*

(1) *the photon field A, the fluctuation fields A', A'_j, and the block-spin fields B, $B^{(k)}$;*

(2) *the background fields $B^{(1)} = aL^{-2}C^{(0)}Q^*B$ [1, (3.10), p. 614], their analogues $A^{(k)}$, $B^{(k+1)}$, $B^{(k+1),\eta}$ at general k, and the averages $\bar{A}^{(k)}$ [2, p. 570];*

(3) *the interpolated fields $\tilde{B} = (1 - \theta_{k+1})A^{(k)} + \theta_{k+1} B^{(k+1),\eta}$ [2, p. 579];*

(4) *the arguments $\tilde{A} + \tilde{B}$ of* [3, (1.16), p. 414]*, and B', B_0 of the decomposition $\tilde{B} = B_0 + B'$* [3, p. 433]*, with B_0 chosen real as in Lemma E.69(c);*

(5) *the fields obtained from these by the gauge transformation that removes B_0* [3, p. 433]*, and the field A^ε of* [2, (3.37), p. 591].

Every operator of Definition E.68(a) that the construction evaluates is evaluated at one of these fields or at $A = 0$. Every Gaussian measure of the construction is $N(0, P^{-1})$ on a real space with

P real symmetric positive definite; for instance $d\mu_{C^{(0)}}(A')\, d\mu_{C^{(0)}(B^{(1)})}(\phi')$ [1, (3.22), p. 616] *and the measure of a positive operator in* [2, (2.28)–(2.31), p. 563].

Proof. Induction on k. At the start the integration variables are the photon field, with law $N(0, C_B(u))$ and $C_B(u)$ real symmetric positive definite (Lemma E.69(d)), and the $\mathbb{R}^2$-valued scalar field; the block-spin fields are their real block averages. If all fields of step $k-1$ are real, the fields of step k arise by: the translations [1, (3.10), (3.18), p. 614], [2, (2.56)], which apply operators of Lemma E.69(a) at real fields to real fields; Gaussian integrations with real symmetric positive covariances [1, (3.11), (3.19)], [2, (2.28)], where the weights $1+u\delta_j > 0$ change only the photon covariances (Lemma E.69(d)); averages and the interpolation with the real cut-off θ_{k+1} (Lemma E.69(b)); the expansions [1, (3.14)–(3.16)], [3, (1.15)–(1.16)] in a real field (A', $\tilde{A}$ or B') around a real field; the constant approximant and the gauge transformation of [3, p. 433] (Lemma E.69(c)). In [2, (3.37)], A^ε is the variable of the outer real integral $\int dB$, real by (1). This proves (1)–(5). The last assertion holds because [1, 2, 3, 4] define the operators of Definition E.68(a) only for vector fields, i.e. real configurations [4, p. 572]; the only evaluations at other arguments are Fourier representations at $A = 0$ (e.g. the contour shift in the proof of [4, Lemma 2.4, p. 586]) and $R_{\bar{n}+1}$ at real A [1, p. 615], [3, p. 414], both of the kind in Definition E.68(c). □

Proposition E.71 (only real backgrounds)**.** *In the proof of Theorem 5.2:*

(i) *every gauge background at which the construction, or a statement of this appendix used in the proof, evaluates an operator of Definition E.68(a) is real;*

(ii) *every estimate taken from the literature is stated for real backgrounds, for orthogonal gauge configurations, or for real Gaussian variables and real polynomial interactions, and is applied at a real background. These estimates are: the propagator bounds of* [4] *and* [1, Props. 2.1–2.3]*;* [3, (1.16), p. 414; (1.32)–(1.33), p. 420]*;* $\|R_{\bar{n}+1}(qA)\| \le 1$ [1, p. 615]*; the diamagnetic inequality* [8, (3.17), p. 257]*; and the cumulant lemma* [6, p. 152] *as applied in* [1, (3.22)–(3.24), p. 616] *and* [2, (2.28)–(2.31), p. 563]*;*

(iii) *the printed proofs of these estimates contain no non-real gauge background; their only complex quantities are those of Definition E.68(c);*

(iv) *the two steps of* [3] *whose proofs are not printed, the bound on the operator (1.16) on the Neumann domain* Ω [3, p. 414] *and the cube remainder* [3, p. 433]*, are stated at real backgrounds, and their proofs in Lemmas E.118, E.120 and E.121 use real backgrounds only. The second presupposes* $\square \subset \Omega$ *and the replacement of* $G_k(\Omega_2, \tilde{B})$ *(Lemmas E.124, E.125), which are inclusions of sets.*

In particular no bound on $G_k(\Omega, A)$, $C_k(\Omega, A)$ or an operator built from them is used at a non-real A, near a Neumann face of Ω or elsewhere, and no coupling constant takes a non-real value.

Part (iv) uses the cube inclusions $\square \subset \Omega$ and $\square \subset \Omega_2$ (the latter replaced by the two-domain theorem, Proposition D.17), which are parts of Proposition E.110. A second proof is given after Lemma E.99.

Proof. (i). The construction is covered by Lemma E.70. The statements of this appendix used in the proof of Theorem 5.2 (Subsection F.1) concern the objects of that lemma; for instance Lemma E.120 is stated for $A \in \{B_0, B_0 + B'\}$ in the class of [1, Prop. 2.1], and Lemma E.118 for the backgrounds $\tilde{A}$, $\tilde{B}$ of [3, p. 414].

(ii). [4] defines its operators for real vector fields [4, p. 572], and [1, Props. 2.1–2.3] are stated for regular real vector fields. In [3, p. 414] the operator (1.16) is built from $G_k(\Omega, \tilde{B})$ and $G_k(\Omega, \tilde{A}+\tilde{B})$, with $\tilde{A}$ a sum of the real Gaussian fields A'_j, and the norms (1.32) are Hölder norms of kernels at these backgrounds [3, p. 420]. The inequality $\|R_{\bar{n}+1}(qA)\| \le 1$ "holds for arbitrary real A" [1, p. 615] and fails for non-real A (Lemma E.99(b)). [8, (3.17)] is stated for configurations of a compact gauge group and proved from [8, (3.7), p. 255], whose proof uses $\|U^\Phi(g)\| = 1$; in [2,

p. 591] it is applied to (3.37) at the real field A^ε (Lemma E.70(5)), where $U(A^\varepsilon)$ is orthogonal (Lemma E.99(a)). The lemma of [6, p. 152] concerns real Gaussian variables and a real polynomial; in [1, (3.22)–(3.24)] it is applied to $d\mu_{C^{(0)}}(A')\, d\mu_{C^{(0)}(B^{(1)})}(\phi')$ and a polynomial with real coefficients at the real background $B^{(1)}$, and in [2, (2.28)–(2.31)] to "a probabilistic Gaussian measure" of "a positive operator"; in each case $\mu = N(0, P^{-1})$ on $(\mathbb{R}^n)^\Lambda$ with P real symmetric positive. At the pages of [24] that are used (pp. 653–657, 659) operators are evaluated at real fields, and the only complex quantities are sources. By (i), each estimate is applied at a real background.

(iii). In [4] the only complex quantity is the contour shift in the proof of Lemma 2.4 [4, p. 586], at $A = 0$. In [1, 2, 3] the expansions in the background are in real fields around real fields (Lemma E.70), their remainders are controlled by $\|R_{\bar n+1}(qA)\| \le 1$ at real A, and determinants are expanded through $\log(1+\lambda)$ for real eigenvalues λ of real symmetric operators. The proof of [8, (3.7)] is the path expansion [8, (3.6)] with $\|U^\Phi(g)\| = 1$. The proof of the lemma of [6] uses real Gaussian covariances at real momenta [6, App. C, pp. 164–165].

(iv). At [3, p. 414] the backgrounds $\tilde B$, $\tilde A + \tilde B$ are real. At [3, p. 433], $\tilde B = B_0 + B'$ with $\tilde B$ real; by Lemma E.69(c), B_0 may be taken real, and then B' is real. The proof of Lemma E.118 uses [1, Prop. 2.1] and the real-background Lemma E.117. The proofs of Lemmas E.120 and E.121 use [1, Prop. 2.1], [4, Lemma 2.1, Cor. 2.3, (2.30), pp. 580–581] and the identity $D_{B_0} = U(B')^{-1} D_A - \phi$ with B_0, B' real. The decomposition $G_k(\Omega, \tilde B) = G_k(\square, \tilde B) + \delta G_k(\Omega, \square, \tilde B)$ needs $\square \subset \Omega$ (Lemma E.124); the replacement of $G_k(\Omega_2, \tilde B)$ is Lemma E.125.

Last assertion. By (i) and (ii), no pointwise or derivative bound on G_k or C_k is used at a non-real argument. The proofs written in this appendix that enter Theorem 5.2 contain only these complex quantities: momentum shifts in Feynman integrands at no gauge background (Lemma H.4), $R_{\bar n+1}$ at real arguments, and q. The Taylor coefficients in (e, λ) that enter are derivatives at real couplings, and Theorem 5.2 is evaluated at the two real points $u \in \{1, -\eta\}$. □

Remark E.72 (scope). (a) Some results used in the proof of Theorem 5.2 rest on statements whose proofs are not given in this paper (the Appendix-K statements, e.g. Statements K.22, K.23). They are used through their statements, which concern the real objects of Lemma E.70, so (i) and (ii) of Proposition E.71 apply to them. Proposition E.71 makes no assertion about proofs that are not written here. (b) Two complex-analytic devices are used in no proof written here or printed in [1, 2, 3, 4]: Cauchy estimates in the gauge field, which would need bounds on $G_k(\Omega, A)$ for non-real A, and bounds at a complex charge ze with $|z| = 1$. By Lemma E.99(a),(b), such bounds would not follow from [4], from [8, (3.17)] or from $\|R_{\bar n+1}(qA)\| \le 1$.

Proposition 1 at the order cap $\bar N = 169$. *Notation.* Fix a step $k \to k+1$, $0 \le k \le K-1$, and write $t := t_k = L^k \varepsilon \le \varepsilon_0$, $e(t)^2 = e^2 t$, $\lambda(t) = \lambda t$, $p(t) = b_0(1 + \log t^{-1})^p$ with $p = 6$, $b_0 \ge \sqrt{2}$, and $\mathsf{d} = 3$. The *coupling order* of a vertex v is $q_v := d_v(v) + d_s(v)$, where $d_v(v)$, $d_s(v)$ are its orders in e and in λ [3, p. 420]; the order of a graph or family is $q := \sum_v q_v$. Couplings inside the covariant kernels $U(\tilde B)$, $G_k(\Omega, \tilde B)$ are not counted. The fields satisfy the small-field restrictions of Definition E.59. Since $p(t_{k-1}) \le (1 + \log L)^p p(t_k)$ and $\lambda(t_{k-1})^{-1/4} = L^{1/4} \lambda(t_k)^{-1/4}$, there is $C_1 = C_1(c_1, L, p)$ with

$$\|h\Phi_{\rm ext}\|_{1,\alpha_0} \le C_1 p(t) \lambda(t)^{-1/4}, \qquad \|h' A_{\rm ext}\|_{1,\alpha_0} \le C_1 p(t) \tag{E.5}$$

for each scalar and each vector leg; for the external scalar fields $a_k G_k(\Omega, \tilde B) Q_k^*(\tilde B)\phi$ the covariant derivative is $\le C_1 p(t)$ (Lemma 5.6). Put $P(t) := p(t)^4 (1 + \log t^{-1}) \ge 1$, let $N_k := |T_\varepsilon| / t_k^3$, and let $r(t) = R(1 + \log t^{-1})^2$, $\hat\ell = 1 + \log \varepsilon_0^{-1}$ be as in Appendix A.

Definition E.73 (the family bound at order q)**.** For $q \in \mathbb{N}$, (F_q) is the statement: there are $C(q) = C(q, \alpha_0; \mathsf{d}, L, a, M_b)$ and $\delta(q) > 0$ such that every localised renormalised family $G_{\rm ren}$ of order $\le q$, on T_ε, at every $k \le K$, with domains $\Omega, \Omega_1, \Omega_2$ and background $\tilde B$ as in Corollary 5.10, satisfies

$$|E(G_{\rm ren}, \{\square(v)\}, \Phi_{\rm ext}, A_{\rm ext})| \le C(q)\, e(t)^{d_v(G_{\rm ren})} \lambda(t)^{d_s(G_{\rm ren})}\, \mathrm{e}^{-\delta(q) d(\{\square(v)\})}\, \|h\Phi_{\rm ext}\|_{1,\alpha_0} \|h' A_{\rm ext}\|_{1,\alpha_0},$$

and coloured families (Theorem H.24) satisfy the same bound with the extra factor $(|u|c_k)^m$, m the number of D-lines.

[3, (1.33), p. 420] has a rate δ_0 that "depends on the dimension d only" and a constant that "depends on $\alpha_0, \bar{n}$" [3, p. 421]. (F_q) is weaker: its rate may depend on the order, since the rates used to move lines decrease with the number of lines ([3, p. 427]; Lemma E.51).

Lemma E.74 (divergent subgraphs do not depend on the cap)**.** *In $\mathsf{d} = 3$ a graph with E external legs, e-order α and λ-order β has superficial degree $\omega = 3 - E/2 - \alpha/2 - \beta$. Hence $\omega \geq 0$ if and only if $\alpha + 2\beta \leq 6 - E$, and every superficially divergent graph has coupling order $\leq 6 < \bar{n}$.*

Proof. φ, A and e have dimension $\frac{1}{2}$, λ has dimension 1. For $E = 0$ this is $\alpha + 2\beta \leq 6$, i.e. [3, p. 418], "$2 \leq \alpha + 2\beta \leq 6$"; there is no case with $E \geq 6$; and $\alpha + \beta \leq \alpha + 2\beta$. □

Lemma E.75 (the cube conditions at every scale)**.** *Let $C_{\mathrm{loc}}(\bar{N})$ and $C_{\bar{N}}$ be the numbers fixed at stage 4 of Appendix A, and assume*

$$\textit{(R.cap)}\ \ R\hat{\ell}^2 \geq C_{\mathrm{loc}}(\bar{N}), \qquad \textit{(T.cap)}\ \ \ell_T \geq 3r(\varepsilon_0)\varepsilon_0 + C_{\bar{N}}\varepsilon_0, \qquad \textit{(E.r)}\ \ \varepsilon_0 < \mathrm{e}^{1-\varrho} = \mathrm{e}^{-1}.$$

Then for every $0 \leq k \leq K$: (a) $r(t_k) \geq C_{\mathrm{loc}}(\bar{N})$; (b) $(3r(t_k) + C_{\bar{N}})t_k \leq \ell_T$. So at every scale the cube of [3, p. 432] *containing a connected component of a family of order $\leq \bar{N}$ fits into $\square_1$ (side $r(t_k)$ in units of t_k-blocks), and $\square$ (side $3r(t_k)$) together with this cube has physical size $\leq \ell_T$, so it does not wrap.*

Proof. (a) $t_k \leq \varepsilon_0$ gives $1 + \log t_k^{-1} \geq \hat{\ell}$. (b) $f(t) := t(1 + \log t^{-1})^2$ has $f'(t) = s^2 - 1$, $s = \log t^{-1}$, so f increases on $(0, \mathrm{e}^{-1}]$; hence $r(t_k)t_k = Rf(t_k) \leq r(\varepsilon_0)\varepsilon_0$ and $C_{\bar{N}}t_k \leq C_{\bar{N}}\varepsilon_0$. Apply (T.cap). □

Proposition E.76 (the order cap)**.** *Let $\bar{n} = 13$, $\bar{N} = \bar{n}^2 = 169$, $0 < \alpha_0 < 1$, $L = 2$, and assume (R.cap), (T.cap) and (E.r) of Lemma E.75. Assume Bałaban's estimate* [3, (1.33), p. 420] *in the form of Proposition E.97, with the structure of its proof described in steps (1)–(7) below, together with its version for graphs with coloured photon lines (D-lines; Definition E.59, Theorem H.24). Then (F_q) holds for every finite q, in particular for $q = \bar{N}$, with:*

(i) *$C(\bar{N})$ a finite multiple of the constant $C_A \leq c_{\mathsf{d}}^m (4/\delta_1)^{\mathsf{d}m} (4m+2)^{\mathsf{d}m(m-1)/2} (1 - L^{-\vartheta})^{-m}$ of Lemma E.51 with $m \leq m(\bar{N})$ lines, times the number of graphs of order $\leq \bar{N}$;*

(ii) *some rate $\delta(\bar{N}) > 0$, and on the coloured side the factor $(1 + |u|c_k)^{N_v} \leq (1 + L^2)^{N_v} \leq 5^{84}$ for all $u \in [-\eta, 1]$, where $N_v \leq \bar{N}/2$ is the number of vector lines.*

These constants do not depend on ε, k, the position, $\Omega, \Omega_1, \Omega_2$ or $\tilde{B}$. They are uniform over tori with $\ell_T \geq \ell_0$, with values taken at the floor $\ell_0^\natural$ for families containing $\delta m_{\mathrm{fin}}^2$ vertices. No step requires a smallness condition on $e, \lambda, e(t), \lambda(t)$ that depends on the order. Every superficially divergent subgraph has $\alpha + 2\beta \leq 6 - E$ (Lemma E.74), so the counterterms do not depend on the cap.

(R.cap) is met at stage 5 (by $R \geq C_{\mathrm{loc}}(\bar{N})$, since $\hat{\ell} \geq 1$) or at stage 10b, (E.r) at stage 10b, and (T.cap) at stage 11 (Lemma A.21).

Proof. Proposition 1 of [3] bounds one family at a time. Each step of its proof, with the sketched steps written out as described in the paragraph *The base* below, is a finite induction over the nested subgraphs $G_1 \subset \cdots \subset G_m = G$ of one graph of fixed order; nothing is induced over the order, and no series in the order is summed. The order enters as follows. (1) *Graph expansion* [3, p. 420]: the order fixes only the range of the final finite sum over families, whose number depends only on $\bar{N}$. (2) *Single lines* [3, (2.10)–(2.12), p. 426] (Proposition 4.5): no order enters; the condition "$e(L^k\varepsilon)$ small" for $G_k(\Omega, \tilde{B})$ is a property of the kernel. (3) *Tree factor* [3, (2.13)–(2.14)]: before the induction each line keeps $\exp[-\frac{1}{2}\delta_1 \operatorname{dist}/(L^{j_l}\eta)]$, independently of the order. (4) *Moving lines* (Lemma E.51, step (a′)): $\delta^{(i)} \geq \frac{\delta_1}{2}(4m+2)^{-(i-1)}$, $m \leq m(\bar{N})$, the "$\delta_2(\delta_1, \bar{n})$" of [3, p. 427]; this gives $\delta(\bar{N}) > 0$ (not Bałaban's "δ_0 depends on d only" [3, p. 420]). The sums over $\Delta(w')$ and

over scale differences give C_A; the geometric sum uses $\vartheta \geq \min(\alpha_0, \frac{1}{2})$. In $\mathsf{d} = 3$ the degrees grow with the number of vertices, so higher orders only enlarge the margins. (5) *Renormalisation* [3, Cor. 2.3, pp. 429–432; pp. 434–445]: by Lemma E.74 every divergent subgraph has order ≤ 6, so the divergent classes and the counterterms δm^2, E_1 with their partners are the same at every cap. By Lemma H.18(d) every divergent subgraph is C-only; the hypothesis $\bar{n} > 12$ of Theorem H.24 is a lower bound, kept by $\bar{N}$; the colour factor is counted by $N_v \leq \bar{N}/2 \leq 84$. (6) *Localisation of a component* [3, pp. 432–433]: each connected component lies in a cube whose size depends on its number of vertices; this cube must fit into $\square_1$, and $\square$ must not wrap. Lemma E.75 gives both at every scale, so Proposition 5.8(c) and Corollary 5.10, stated with $\ell_{\bar{N}}$, apply as written. (7) *The constant*: [3, p. 421] states that $O(1)$ "depends on $\alpha_0, \bar{n}$"; here it is $C(\bar{N})$ of (i). Only (4) and (6) depend on the cap, and neither imposes a smallness condition on the couplings. The torus enters only through the finite size of T_ε and (6); the only ℓ_T-dependent constant is the zero-mode part of $\delta m^2_{\mathrm{fin}}$, which decreases in ℓ_T and is evaluated at $\ell_0^\natural$.

The base. The structure of the proof of Proposition E.97 used in (1)–(7) is the proof of [3, Prop. 1] on pp. 422–434, completed by Lemmas E.51, E.52, E.55, Theorem E.53, Propositions E.54, 5.8, Corollary 5.10, the graded counterterm bound Lemma E.80, for coloured families Theorem H.24, and by the four steps that [3] only sketches: the closure of the renormalised classes under cutting [3, p. 432] (Statement K.24); the bad localisations of the counterterms [3, p. 422] (Statement K.25); the Hölder bounds of the background fields [3, p. 412], for vector fields Statement K.26 and for the scalar external fields Lemma E.79 below, whose hypothesis ($\mathrm{N_{nn}}$) is Statement K.23; and the nested and one-particle-reducible cases in step (vi) of Lemma E.55 (Statement K.27).

Tail constant. The constant $C_>$ of the tails of order $> \bar{n}$ (Proposition E.84) is $C_{582} \sup_{t\in(0,1]} t^{3/10} P(t)^{\bar{N}}$: for $q \geq 14$, $t^{q/4} P(t)^q \leq t^{16/5} \cdot t^{3/10} P(t)^{\bar{N}}$, and a positive power of t beats every power of $1 + \log t^{-1}$. So the supremum is finite and does not depend on ε_1. □

Scalar Hölder bounds. Let $d = 3$, $\eta = L^{-k}$, $e(t) = et^{1/2}$, and let $G(X, A) := G_k(X, A)$ be the propagator [1, (2.22)] on a union X of big blocks, with Neumann conditions on ∂X (none if $X = T_\eta$). Put $U(a) = \exp(\eta q e(t) a)$ for a bond variable a, $U(A(\Gamma)) = \exp(q e(t) A(\Gamma))$ with $A(\Gamma) = \sum_{b\in\Gamma} \pm\eta A(b)$, and $D_{A,\mu} u(z) = \eta^{-1}(U(A_\mu(z))u(z + \eta e_\mu) - u(z))$. For $0 < |x - x'| \leq 1$ let $\Gamma_{x,x'}$ be a shortest lattice contour (length $|x - x'|_1 \leq \sqrt{d}|x - x'|$) and

$$H^A_{x,x'}(u) := \max_\mu |x - x'|^{-\alpha} |U(A(\Gamma_{x,x'})) D_{A,\mu} u(x') - D_{A,\mu} u(x)|,$$

$$\mathcal{N}^A_x(u) := \max\Big\{ \max_\mu |D_{A,\mu} u(x)|, \sup_{0<|x'-x|\leq 1} H^A_{x,x'}(u) \Big\}.$$

c_0, δ_0, R_0 are the constants of Proposition 4.5 at the exponent $\alpha < 1$, M_b is the side of the big blocks, c_r, β_r are the constants of [1, (2.23)], $\ell_Q \leq d$ bounds the lengths of the contours in Q_k, ϑ_R and C_{rem} are the constants of Proposition E.57(a) at $m = 1$, and $0 \leq m^2 t^2 \leq c$.

Lemma E.77 (change of the transporting field)**.** *Let A_0 be a constant field, $B = A_0 + \tilde{A}$, x a site, $N_x := \{z : |z - x| \leq \sqrt{d} + 2\}$ and $s := e(t)(\sup_{N_x} |\tilde{A}| + \sup_{N_x} |\partial^\eta \tilde{A}|) \leq 1$. Then for every field u, every μ and $0 < |x' - x| \leq 1$: (a) $|D_{B,\mu} u(x)| \leq |D_{A_0,\mu} u(x)| + s|u(x)|$; (b) $H^B_{x,x'}(u) \leq H^{A_0}_{x,x'}(u) + 2\sqrt{d}\, s(\sup_{N_x} |D_{A_0} u| + \sup_{N_x} |u|)$.*

Proof. With $\hat{\phi}_\mu := \eta^{-1}(U(\tilde{A}_\mu) - 1)$, $D_{B,\mu} u = U(\tilde{A}_\mu) D_{A_0,\mu} u + \hat{\phi}_\mu u$, $|\hat{\phi}_\mu| \leq e(t)|\tilde{A}_\mu|$, $|\hat{\phi}_\mu(z') - \hat{\phi}_\mu(z)| \leq e(t)|z' - z|_1 \sup_{N_x} |\partial^\eta \tilde{A}|$ (chaining in N_x); this gives (a). For (b), with $\Gamma = \Gamma_{x,x'}$,

$$\begin{aligned} U(B(\Gamma)) D_{B,\mu} u(x') - D_{B,\mu} u(x) &= U(\tilde{A}_\mu(x))[U(A_0(\Gamma)) D_{A_0,\mu} u(x') - D_{A_0,\mu} u(x)] \\ &\quad + U(A_0(\Gamma))[U(\tilde{A}(\Gamma)) U(\tilde{A}_\mu(x')) - U(\tilde{A}_\mu(x))] D_{A_0,\mu} u(x') \\ &\quad + \hat{\phi}_\mu(x')[U(B(\Gamma)) u(x') - u(x)] + (\hat{\phi}_\mu(x') - \hat{\phi}_\mu(x)) u(x). \end{aligned}$$

The second bracket is $\le \sqrt{d}\,s|x-x'|$; telescoping along Γ gives $|U(B(\Gamma))u(x')-u(x)| \le \sqrt{d}|x-x'|(\sup_{N_x}|D_{A_0}u| + s\sup_{N_x}|u|)$; the last term is $\le \sqrt{d}\,s|x-x'||u(x)|$. The last three terms are at most $\sqrt{d}\,s(1+s)|x-x'|(\sup_{N_x}|D_{A_0}u|+\sup_{N_x}|u|)$; divide by $|x-x'|^\alpha$ and use $|x-x'|\le 1$, $s\le 1$. □

Lemma E.78 (background differences)**.** *Let Ω be a union of big blocks and $\tilde B$ a real field on Ω with $|\partial^\eta \tilde B| \le C_\partial p(t)$. Let x lie in the unit block of the site y, $Q_s := \{z : |z-y|_\infty \le s\}$, and let $\rho \ge 2$ be an integer with $Q_{2\rho+3} \subset \Omega$ not wrapping around the torus,* $\operatorname{dist}(Q_2, \Omega^c) \ge R_0$ *and $\rho'' := \rho - M_b - \sqrt{d} - 4 \ge R_0$. Let $\square$ be the union of the big blocks contained in Q_ρ (a rectangular parallelepiped, $\square \supset Q_{\rho - M_b}$), $A_{0,\mu} := \tilde B_\mu(y)$, $\tilde A := \tilde B - A_0$, $\sigma_* := 3dC_\partial e(t)p(t)(\rho+3)$, and assume $\sigma_* \le \min\{\vartheta_R, 1\}$ and $(1+3d)C_\partial p(t) \le c_r e(t)^{\beta_r - 1}$. For ψ on the unit lattice of Ω let $\psi_\square := \psi 1_{\{y' : B(y') \subset \square\}}$, $\phi := a_k G(\Omega, \tilde B) Q_k^*(\tilde B)\psi$ and $\Psi := a_k G(\square, A_0) Q_k^*(A_0)\psi_\square$. Then, with $\mathcal{E}_\psi := C_D(\mathrm{e}^{-\delta_0\rho''} + \sigma_*)a_k\|\psi\|_\infty$ and $C_D := c_0(4 + 8\sqrt{d} + \ell_Q) + C_{\rm rem}$,*

$$\max_\mu |D_{\tilde B,\mu}\phi(x)| \le \max_\mu |D_{A_0,\mu}\Psi(x)| + \mathcal{E}_\psi, \qquad H^{\tilde B}_{x,x'}(\phi) \le H^{A_0}_{x,x'}(\Psi) + \mathcal{E}_\psi \quad (0 < |x'-x| \le 1).$$

Proof. The second assumption makes $\tilde B$ regular in the sense of [1, (2.23)], and chaining gives $|\tilde A_\mu(z)| \le |z-y|_1 C_\partial p$ on $Q_{2\rho+3}$. Let $\chi(z) := \min\{1, \max\{0, 2 - |z-y|_\infty/(\rho+1)\}\}$, $\tilde A^\sharp_\mu := \chi\tilde A_\mu$ (zero off $Q_{2\rho+2}$) and $\tilde B^\sharp := A_0 + \tilde A^\sharp$ on T_η. Then $\|\tilde A^\sharp\|_\infty \le d(2\rho+2)C_\partial p$, $\|\partial^\eta \tilde A^\sharp\|_\infty \le (1+3d)C_\partial p$, so $\sigma^\sharp := e(t)(\|\tilde A^\sharp\|_\infty + \|\partial^\eta \tilde A^\sharp\|_\infty) \le \sigma_*$; $\tilde B^\sharp$ is regular, $\theta(\tilde B^\sharp) \le \vartheta_R$, $\theta(A_0) = 0$, and $\sigma_\square := e(t)(\sup_{Q_{\rho+1}}|\tilde A| + \sup_{Q_{\rho+1}}|\partial^\eta \tilde A|) \le \sigma_*$. Since $\tilde B^\sharp = \tilde B$ on the bonds based in $Q_{\rho+1} \supset \square$, $G(\square, \tilde B^\sharp) = G(\square, \tilde B)$, and $D_{\tilde B^\sharp}$, $H^{\tilde B^\sharp}$ at x, x' equal those for $\tilde B$. Also $\operatorname{dist}(\{x,x'\}, \square^c) \ge \rho''$, and $N_x \subset Q_{\rho+1}$ has distance $\ge \rho'' \ge R_0$ from $\square^c$. With $f_{\rm full} := a_k Q_k^*(\tilde B)\psi$, $f := a_k Q_k^*(\tilde B)\psi_\square$, $f_0 := a_k Q_k^*(A_0)\psi_\square$ (sup norms $\le a_k\|\psi\|_\infty$), $\phi = \Psi + \sum_{i=1}^6 R_i$ with $R_1 := G(\Omega, \tilde B)(f_{\rm full} - f)$, $R_2 := [G(\Omega, \tilde B) - G(\square, \tilde B)]f$, $R_3 := G(\square, \tilde B)(f - f_0)$, $R_4 := [G(\square, \tilde B^\sharp) - G(T_\eta, \tilde B^\sharp)]f_0$, $R_5 := [G(T_\eta, \tilde B^\sharp) - G(T_\eta, A_0)]f_0$, $R_6 := [G(T_\eta, A_0) - G(\square, A_0)]f_0$. By [1, (2.24)–(2.26)] (Proposition 4.5), $\mathcal{N}_x^{\tilde B}(R_i) \le c_0 \mathrm{e}^{-\delta_0\rho''} a_k\|\psi\|_\infty$ for $i = 1, 2, 4$. Since $|U(\tilde B(\Gamma)) - U(A_0(\Gamma))| \le \ell_Q \sigma_\square$ for the contours of Q_k, $\mathcal{N}_x^{\tilde B}(R_3) \le c_0 \ell_Q \sigma_* a_k\|\psi\|_\infty$. By the resolvent identity $R_5 = -G(T_\eta, \tilde B^\sharp) V G(T_\eta, A_0) f_0$ with $V := G(T_\eta, \tilde B^\sharp)^{-1} - G(T_\eta, A_0)^{-1}$, and Proposition E.57(a) ($n = 1$, $n' = 0$, perturbation $-\tilde A^\sharp$) gives $\mathcal{N}_x^{\tilde B}(R_5) \le C_{\rm rem}\sigma^\sharp a_k\|\psi\|_\infty$. Lemma E.77 with $s \le \sigma_*$, and (2.26), (2.25) for $G(\square, A_0)$, give $\mathcal{N}_x^{\tilde B}(R_6) \le c_0(1 + 4\sqrt{d})\mathrm{e}^{-\delta_0\rho''} a_k\|\psi\|_\infty$ and bound both differences between the $\tilde B$- and the A_0-quantities of Ψ by $4\sqrt{d}\,c_0\sigma_* a_k\|\psi\|_\infty$. Summing gives at most $\mathcal{E}_\psi$. □

Lemma E.78 measures $\tilde B - A_0$ on all of $Q_{2\rho+3}$. A pointwise kernel bound in which $\tilde B - A_0$ enters only through its values on $[x, x']$ and on the block of x' is false already for $-\Delta_A + m^2$: its random-walk expansion has positive weights, so $(-\Delta_A + m^2)^{-1}(x,x) < (-\Delta_0 + m^2)^{-1}(x,x)$ whenever A is non-zero (modulo the period) on one bond away from x.

Lemma E.79 (scalar external fields)**.** *Let Ω, $\tilde B$ be as in Lemma E.78 and $\phi^{(k)} = a_k G(\Omega, \tilde B) Q_k^*(\tilde B)\psi$. Assume*

(Ψ) $\|\psi\|_\infty \le C_\psi \lambda(t)^{-1/4} p(t)$;

(N$_{\rm nn}$) $|U(\tilde B(\Gamma_{y_1,y_2}))\psi(y_2) - \psi(y_1)| \le C_{\rm nn} p(t)$ *for nearest neighbours y_1, y_2 of the unit lattice in Ω;*

(D) $|D_{\tilde B}\phi^{(k)}(z)| \le C_{2.66} p(t)$ *for* $\operatorname{dist}(z, \Omega^c) \ge r(t)$ *(as in* [2, (2.66)]*; Lemma 5.6);*

and let $t \le \varepsilon_0$, where for $t \le \varepsilon_0$: $r(t) \ge 4\sqrt{d}(R_0 + M_b + \sqrt{d} + 6)$; $\sigma_*(t) := 3dC_\partial e(t)p(t)(r(t)+3) \le \min\{\vartheta_R, 1\}$; $(1+3d)C_\partial p(t) \le c_r e(t)^{\beta_r - 1}$; $\ell_T/t \le 8r(t)$. Then for $\operatorname{dist}(\{x,x'\}, \Omega^c) \ge r(t)$ *and $\alpha < 1$,*

$$|x-x'|^{-\alpha}|U(\tilde B(\Gamma_{x,x'}))D_{\tilde B}\phi^{(k)}(x') - D_{\tilde B}\phi^{(k)}(x)| \le C^s p + \omega(t)p,$$

with $S_d := \sum_{z\in\mathbb{Z}^d}|z|_1 \mathrm{e}^{-\delta_0(|z| - 2\sqrt{d} - 1)_+}$, $C^s := \max\{2C_{2.66}, c_0 a S_d C_{\rm nn}\}$ and

$$\omega(t) := aC_\psi\lambda(t)^{-1/4}\big[(c_0 S_d + C_D)\sigma_*(t) + C_D \mathrm{e}^{-\delta_0(r(t)/(4\sqrt{d}) - M_b - \sqrt{d} - 5)}\big] \to 0 \quad (t \to 0).$$

Proof. For $|x-x'|>1$ use (D). Let $|x-x'|\le 1$, y the block of x, $\rho:=\lfloor r(t)/(4\sqrt{d})\rfloor$. Then $\rho\ge 2$, $\rho''\ge R_0+1$, $Q_{2\rho+3}$ lies at distance $\ge r/2-4\sqrt{d}\ge R_0$ from Ω^c and has side $<\ell_T/t$, and $3dC_\partial ep(\rho+3)\le\sigma_*(t)$; Lemma E.78 with $a_k\le a$ and (Ψ) gives $H^{\tilde B}_{x,x'}(\phi^{(k)})\le H^{A_0}_{x,x'}(\Psi)+aC_\psi\lambda^{-1/4}C_D(\mathrm{e}^{-\delta_0\rho''}+\sigma_*)p$. On the non-wrapping box $\square$ put $\Lambda(z):=\sum_\mu A_{0,\mu}(y-z)_\mu$ and $(\mathcal{U}u)(z):=\exp(qe(t)\Lambda(z))u(z)$; then $\Psi=\mathcal{U}v$ with $v:=a_kG(\square,0)Q_k^*\psi'$, $\psi'(y'):=\exp(-qe(t)\Lambda(y'))\psi_\square(y')$, and $H^{A_0}_{x,x'}(\Psi)=\max_\mu|x-x'|^{-\alpha}|\partial^\eta_\mu v(x')-\partial^\eta_\mu v(x)|$. Since $G(\square,0)Q_k^*1=(a_k+m^2t^2)^{-1}1_\square$, $\partial^\eta v=\sum_{y'}\partial^\eta G(\square,0)g_{y'}$, $g_{y'}:=a_kQ_k^*[(\psi'(y')-\psi'(y))1_{\{y'\}}]$. Along a monotone nearest-neighbour path from y to y' in $\square$, $(\mathrm{N_{nn}})$ and $|U(A_0(\Gamma_i))-U(\tilde B(\Gamma_i))|\le\sigma_*$ give $\|g_{y'}\|_\infty\le a|y'-y|_1(C_{\mathrm{nn}}+C_\psi\lambda^{-1/4}\sigma_*)p$. [1, (2.24)] at zero field on $\square$, with $\operatorname{dist}(\{x,x'\},B(y'))\ge|y'-y|-2\sqrt{d}-1$, gives $H^{A_0}_{x,x'}(\Psi)\le c_0aS_d(C_{\mathrm{nn}}+C_\psi\lambda^{-1/4}\sigma_*)p$. Finally $\lambda^{-1/4}\sigma_*=O(t^{1/4}(1+\log t^{-1})^c)$ and $\lambda^{-1/4}\mathrm{e}^{-\delta_0 r/(4\sqrt{d})}\le Ct^{-1/4}\mathrm{e}^{-c'(1+\log t^{-1})^2}$. □

By Statement K.23, $(\mathrm{N_{nn}})$ holds with $C_{\mathrm{nn}}=c_\psi(t)=4Ld+O(t^{1/4}p(t))$. Using it, one replaces C_{nn} by $4Ld$ in C^s and adds $c_0aS_d(c_\psi(t)-4Ld)=O(t^{1/4}p(t))$ to $\omega(t)$; the conclusion is unchanged.

Lemma E.80 (graded bound on the mass counterterm)**.** *Let $\varepsilon=L^{-K}\varepsilon_0$, $0\le k\le K$, $t=t_k$, and let $\delta m^2_{K,k}(x)$ be the part of $\delta m^2_K(x)$ formed by the index assignments whose largest line index is $\ge k$* [3, (1.27)–(1.28), p. 418]*. Assume $\ell_T\ge 6r(\varepsilon_0)\varepsilon_0$ (implied by (T.nw)). Then for every $x\in T_\varepsilon$*

$$|\delta m^2_{K,k}(x)|t^2\le C_m\Big[(e^2+\lambda)t\tfrac{L}{L-1}+(e^4+e^2\lambda+\lambda^2)t^2(K-k)\Big],\qquad K-k=\log_L(\varepsilon_0/t),$$

with $C_m=C_m(\alpha_0,\bar n,\mathsf{d},L,a,M)$ independent of ε,k,x,ℓ_T. In particular $|\delta m^2_{K,k}|t^2=O(t)$, as asserted in [3, p. 418]*.*

Proof. In units of scale K, $\delta m^2_{K,k}(x)t_K^2=\sum_{G_0}c^{[\ge k]}_{G_0}(x)$, a finite sum over the C-only scalar self-energy families of order $o:=\alpha/2+\beta\in\{1,2\}$ and the index assignments with largest index $j^*\ge k$. With the legs equal to 1 and the vertex of leg 1 at x, Lemmas E.52, E.55 and E.51(ii) give $|c^{[j^*]}_{G_0}(x)|\le O(1)e(t_K)^\alpha\lambda(t_K)^\beta(L^{j^*}\eta_K)^{D(G_0)}$, and $D(G_0)=o-2$ by Lemma H.18(b). For $o=1$, $\sum_{j^*=k}^{K-1}L^{K-j^*}\le\frac{t_K}{t}\frac{L}{L-1}$; for $o=2$ there are $K-k$ terms. Undo the scaling with $e(t_K)^2=e^2t_K$, $\lambda(t_K)=\lambda t_K$ and multiply by t^2; finally $t\log_L(\varepsilon_0/t)\le\varepsilon_0/(\mathrm{e}\ln L)$. The torus condition makes the cubes of [3, p. 433] fit at scale K. □

Lemma E.81 (gain per unit of coupling order)**.** *Let G be a connected graph with at least two vertices, each one of* [3, (1.6)–(1.11), (1.14), (1.15), (1.18), pp. 413–415] *or a mass counterterm vertex (1.7) bounded by Lemma E.80; the coupling-free vertices (1.12), (1.13) are counted as part of the external scalar fields* [3, p. 421]*. For a vertex of order $q_v\ge 1$ let F_v be its coupling factor times the bounds* (E.5) *of its external legs. Then $F_v\le C(q_v)t^{q_v/4}P(t)^{q_v}$, hence $\prod_v F_v\le C(q)t^{q/4}P(t)^q$, with $C(q)$ depending on q,e,λ,C_1,C_m,L.*

Proof. Every vertex has an internal leg, a vertex of order q_v has at most $4q_v$ legs, a scalar leg costs $\le C_1p\lambda^{-1/4}t^{-1/4}$ and a vector leg $\le C_1p$. (1.6): $\lambda t(C_1p\lambda^{-1/4}t^{-1/4})^3\le Ct^{1/4}p^3$. (1.7): the first term of Lemma E.80 with two scalar legs gives $Ct^{1/2}p^2$; the second gives $Ct^{3/2}(1+K-k)p^2\le Ct^{q_v/4}P(t)$ since $q_v\le 6$. (1.8), $q_v\ge 2$: $Ct^{q_v/2}p^{q_v+2}\lambda^{-1/2}t^{-1/2}$ and $\frac{q_v-1}{2}\ge\frac{q_v}{4}$; $q_v=1$: $\le Ct^{1/4}p^2$ with one external scalar leg, and with two the differentiated one is $\le C_1p$ (Lemma 5.6), giving $Ct^{1/4}p^3$. (1.10) as (1.8). (1.9), (1.11): factor $e(t)^{n+\bar n+1}$ and $\|R_{\bar n+1}\|\le 1$ at real argument (Lemma E.99(b)). (1.14), (1.15), (1.18): as (1.8), and the first two vertices of (1.18) with $n+n'=1$ are $O(t^{1/4}p^2)$ by Lemma 5.6 and [3, p. 421]. In all cases $F_v\le C(q_v)t^{q_v/4}p^{4q_v}(1+\log t^{-1})\le C(q_v)t^{q_v/4}P^{q_v}$. □

Lemma E.82 (sum over localisations)**.** *Let a family have N vertices localised in cubes $\square(v)$ of diameter $\le c_0$ attached to points of a lattice of spacing 1, with at most n_0 cube types per point* [3, p. 420]*, and let $\delta>0$. The sum of $\mathrm{e}^{-\delta d(\{\square(v)\})}$ over the localisations with $\square(v_1)$ fixed is at most $S(N,\delta):=n_0^{N-1}N^{N-2}\big(\mathrm{e}^{\delta c_0}\sum_{z\in\mathbb{Z}^3}\mathrm{e}^{-\delta|z|/2}\big)^{N-1}$.*

Proof. A shortest Steiner tree has length at least half that of a minimal spanning tree for the edge lengths $\operatorname{dist}(\square(v),\square(w))$ (double, take an Euler tour, shortcut). So $\mathrm{e}^{-\delta d} \le \sum_T \prod_{\{v,w\}\in T} \mathrm{e}^{-\frac{\delta}{2}\operatorname{dist}(\square(v),\square(w))}$ over the N^{N-2} labelled trees, and $\operatorname{dist}(\square(v),\square(w)) \ge |y_v - y_w| - 2c_0$. Sum leaf by leaf from the fixed vertex. □

Lemma E.83 (cumulants are sums of connected graphs)**.** *Let μ_C be a centred Gaussian measure and $V_1,\dots,V_m$ polynomials in the field, each a finite sum of vertices with legs. The joint cumulant $\kappa(V_1,\dots,V_m)$ is the sum over the complete pairings of the legs of $V_1,\dots,V_m$ by lines of covariance C whose graph on $\{1,\dots,m\}$ is connected.*

Proof. By Wick's theorem $\int \prod_{i\in S} V_i \, d\mu_C = \sum_\pi \prod_{B\in\pi} c(B)$, grouping pairings by the partition into connected components, $c(B)$ the connected sum. The cumulants satisfy the same moment–cumulant relation, so $\kappa(S) = c(S)$ by Möbius inversion. □

Proposition E.84 (the tail of order $> \bar n$ after the fluctuation integral, [2, p. 582])**.** *Let $u \in \{1, -\eta\}$, $M \in (0, m_2]$, assume the hypotheses of Propositions E.76 and E.36, and let $0 \le k \le K-1$. After the cut of Proposition E.36 the exponent under the second integral of* [2, (2.115), p. 582] *is $V = \sum_{q=1}^{\bar n} V^{[q]}$, each $V^{[q]}$ a finite sum of localised families of coupling order q. With κ the cumulants of the Gaussian fluctuation law with weighted covariance (C- and D-lines, Theorem H.24), let*

$$\Delta_k := \sum_{m=2}^{\bar n} \frac{1}{m!} \sum_{1\le q_1,\dots,q_m\le \bar n,\ q_1+\cdots+q_m>\bar n} \kappa(V^{[q_1]},\dots,V^{[q_m]}).$$

Then, pointwise on the small-field set of step k, $|\Delta_k| \le C_{582} t_k^{7/2} P(t_k)^{\bar N} N_k \le C_> t_k^{16/5} N_k$ with $C_> := C_{582} \sup_{t\in(0,1]} t^{3/10} P(t)^{\bar N} < \infty$. C_{582} and $C_>$ depend only on $\bar n, \bar N, \alpha_0, \mathsf{d}, L, a, M_b, e, \lambda, \mu_*, M, b_0, p, \eta$; they do not depend on ε, k, $\ell_T \ge \ell_0$ (at $\ell_0^\natural$ where $\delta m_{\mathrm{fin}}^2$ vertices occur) or ε_0, and are finite at $u = -\eta$.

Proof. By Lemma E.83, $\kappa(V^{[q_1]},\dots,V^{[q_m]})$ is a sum of connected graphs obtained by joining legs by fluctuation lines ([1, p. 623], [2, p. 582]), of order $q = \sum q_i$; $q > \bar n$ forces $m \ge 2$, and $q \le m\bar n \le \bar N$. They regroup into renormalised families of fixed order ([3, p. 420], Lemma H.23); every divergent subgraph has order ≤ 6 (Lemma E.74) and its counterterm partner, of order ≤ 4, is a vertex of V, so the families close at every order $\le \bar N$ (Statement K.24). By Proposition E.76 each family obeys $(\mathrm{F}_{\bar N})$, with colour factor $(|u|c_k)^m \le 3^{-m}$ ($c_k \le \frac13$, Definition E.59). By Lemma E.81 a family of order q is bounded by $C(q)C(\bar N)t^{q/4}P(t)^q \mathrm{e}^{-\delta(\bar N)d}$. For $q \ge 14$, $t^{q/4} \le t^{7/2}$ and $P^q \le P^{\bar N}$; the number of families is finite; with one block fixed, Lemma E.82 bounds the other localisations by $S(N, \delta(\bar N))$, $N \le \bar N$; the fixed block ranges over $\le N_k$ blocks. The second inequality is $t^{7/2}P^{\bar N} = t^{16/5}\cdot t^{3/10}P^{\bar N}$. Torus images only remove localisations. □

Remark E.85 (the other cuts of order $> \bar n$). (i) The cut before the fluctuation integral [2, p. 579] is Proposition E.36 with Remark E.37; it makes V a polynomial of fixed degree, which Proposition E.84 presupposes. (ii) The terms of order $> \bar n$ of the determinants $Z^{(j)}$ [2, p. 578] are one-loop graphs, estimated by [1, Props. 2.1–2.3]; they are not covered by Lemma E.81. (iii) Cumulants of order $m > \bar n$ are not part of Δ_k.

E.5. **The per-localisation bound, the region sum and Bałaban's Proposition 1.**

The per-localisation bound. We use the notation of the paragraph on the moved vacuum pieces in Subsection E.4: $\mathcal{V}_k(u)$, the pairs $(G,\tilde l)$, the localisations $\Lambda = (\square_\iota)_{\iota\in I(G)}$, $N = |I(G)| \le \bar N(\bar n)$, $d(\Lambda)$, $\bar c = \kappa\mu_*^2\varepsilon_0^2 L^2$, $\bar N_{\mathrm{ph}}$. Put $P(t) := \sum_{2\le\alpha+2\beta\le 2\bar n} |e(t)|^\alpha \lambda(t)^\beta$, where $e(t)^2 = e^2 t$ and $\lambda(t) = \lambda t$. Let m be the number of D-coloured lines of a family.

In this paragraph, all graphs are Bałaban's graphs [3, (1.6)–(1.15), pp. 413–414], graded as in Definition E.12. Each vector line carries one of two colours, according to the multilinear splitting of the photon covariance $C_{\mu_*} + u\Gamma^B$ of (5.1). A line of colour C has Bałaban's kernel. A line

of colour D carries the slice photon; in the units of scale k, its kernel is $D_k^\eta = \sum_{j<k}(L^j\eta)^2 G^\eta_{(j)}$ with the coefficient uc_k, where $0 \le c_k \le \bar{c} := \kappa\mu_*^2\varepsilon_0^2 L^2$. We write $m(G)$ for the number of D-lines of G. For a graph with coupling factor $e^\alpha\lambda^\beta$ we put $o(G) := \alpha/2 + \beta$. Finally, E_s, E_v and E_∂ denote the numbers of external scalar legs, of external vector legs (background legs included), and of differentiations acting on external legs.

Lemma E.86 (per-localisation bound)**.** *Let $\delta_T > 0$ be the tree rate of* [3, (1.33)] *given by Proposition E.97, and let δ_1 be the rate of* [3, (2.10), p. 426]*. Put $\delta := \min(\delta_T, \delta_1)$. Work at either of the points $\pi_\pm$, and let $0 \le k \le K-1$. Consider a pair $(G, \tilde{l})$ of a bare vacuum graph G and an ordering $\tilde{l}$ (Step 1 below) whose last line has index k, other than the order-e^2 family with one D-line ($m(G) = 1$, $o(G) = 1$). Then for every localisation Λ,*

$$|\mathcal{V}_{G,\tilde{l}}(\Lambda)| \le A_k\, \mathrm{e}^{-\delta d(\Lambda)}, \qquad A_k := C_P(1 + |u|\bar{c})^{\bar{N}_{\mathrm{ph}}} P(t_k), \tag{E.6}$$

with P and $\bar{N}_{\mathrm{ph}}$ as in the paragraph containing Lemma E.61. Here C_P is the constant of [3, (1.33)] *times that of* [3, (2.10)–(2.11)]*. The u-part $\mathcal{V}_{G,\tilde{l}}(\Lambda; u) - \mathcal{V}_{G,\tilde{l}}(\Lambda; 0)$ obeys the same bound with $(1 + |u|\bar{c})^{\bar{N}_{\mathrm{ph}}}$ replaced by $\bar{N}_{\mathrm{ph}}|u|\bar{c}(1 + \bar{c})^{\bar{N}_{\mathrm{ph}}-1}$. The order-$e^2$ family with one D-line is bounded per anchor cube in Lemma E.93.*

The lemma rests on Proposition E.97 and [3, (2.10)–(2.11), p. 426]. For the photon-cut classes it also rests on [3, p. 433], whose remainder on $\Box$ is treated in Proposition E.110.

Proof. Step 1 (exact cut, in the grading of Definition E.12). Decompose the index sums by the orderings of [3, (3.4), p. 424], with ties broken by a fixed order of the lines; the classes are indexed by parent graphs. For an index assignment, $\tilde{l} = (l(1), \dots, l(n))$ lists the lines of the bare graph by increasing (index, fixed order). Every assignment belongs to exactly one ordering. Hence every term of $\mathcal{V}_k$ belongs to exactly one pair $(G, \tilde{l})$. Its last line $l(n)$ is a line of the bare graph, and it has index k, because k is the largest index on $\mathcal{V}_k$.

The sets $J(\tilde{l})$ of [3, p. 424] attach to $\tilde{l}$ the increasing chain of subgraphs G_i spanned by $l(1), \dots, l(i)$, so $l(n) \notin G_i$ for $i < n$. Thus every counterterm operation active in $\tilde{l}$ acts on a divergent subgraph that does not contain $l(n)$, and it produces a coefficient in δm^2_{k+1}. Degrees are intrinsic [3, (2.1)–(2.3), pp. 422–423]. So the divergent subgraphs of $G \setminus l(n)$ are exactly the divergent subgraphs of G that do not contain $l(n)$.

Now cut $l(n)$, i.e. treat its kernel $K_k(x, x')$ as a two-variable external field [3, p. 419]. This maps the class, term by term and with the same coefficients, onto renormalised two-leg classes of $G \setminus l(n)$. In these classes the vertices carrying $\delta m^2_{\mathrm{fin}}$ or $\delta m^2_{K,k+1}$ are finite vertices of degree 0. The coefficients δm^2 are never opened.

Step 2 (the cut kernel). In scale-k units, $L^k\eta = 1$. So [3, (2.10)–(2.11)] at $j = k$ give $\|hK_kh'\|_{1,\alpha_0} \le O(1)\mathrm{e}^{-\delta_1 \operatorname{dist}(\operatorname{supp} h, \operatorname{supp} h')}$. This holds for scalar and vector lines alike ("the same inequalities hold for vector field propagators", p. 426). It also holds for D-lines, because their slice k carries the factor $(L^k\eta)^2 = 1$. The bound holds on T_ε (Corollary E.49 at zero background; Proposition 4.5 with Lemma 4.6 in general).

Step 3 (which two-leg classes occur; partners). By Proposition E.87 (see also Lemma E.166), the divergent components of $G \setminus l(n)$ are of the following kinds.

- One-particle-irreducible components of types C1 and C2. Their δm^2 partners are in E_1, which is computed with the full action [3, (1.24), p. 417].
- One-particle-reducible chains of type C2 (21 of the 59 non-vanishing isomorphism classes). They have no partner, since Bałaban's δm^2 is built from one-particle-irreducible graphs [3, p. 416, p. 430]. They need no counterterm, by Lemma E.88.
- Components of type C1′. These are finite vertices [3, (3.22)–(3.24)].

- Components of types C3 and C4. They are renormalised by the Ward–Takahashi class [3, (3.25)–(3.32)]. On T_ε this class is used after the transfer of [3, p. 433] to a non-wrapping cube, where it is exact by Proposition E.67 (Lemmas E.65 and E.66); this needs condition (T.nw).
- C6 pairs, which vanish.

All other components are convergent. In particular, no divergent component contains a vertex (1.13)–(1.15) (Statement K.20). So the cut family is a renormalised family in the sense of Proposition E.97, extended by Lemma E.88.

Step 4 (coloured families). Consider a family with m D-lines, where $m = 0$, $m \geq 2$, or $m = 1$ and $o \geq 2$. For it, the coloured form of (1.33) in Proposition E.97 (Theorem H.24(c)) gives the bound per localisation times $(|u|c_k)^m$, with $c_k \leq \bar{c}$. Summing over the colourings of at most $\bar{N}_{\mathrm{ph}}$ photon lines gives the factor $(1 + |u|\bar{c})^{\bar{N}_{\mathrm{ph}}}$. The u-part is the sum over $m \geq 1$, which is bounded by $(1 + |u|\bar{c})^{\bar{N}_{\mathrm{ph}}} - 1 \leq \bar{N}_{\mathrm{ph}}|u|\bar{c}(1 + \bar{c})^{\bar{N}_{\mathrm{ph}}-1}$. This inequality holds for $|u| \leq 1$ by the mean value theorem: $(1 + x)^N - 1 = Nx(1 + \xi)^{N-1}$ with $\xi \in [0, x]$ and $x = |u|\bar{c} \leq \bar{c}$. The remaining family, $m = 1$ and $o = 1$, is excluded from the lemma. It is bounded per anchor cube directly in Bałaban's localisation, without Proposition 1 (Lemma E.93), and this form suffices for Proposition E.64.

Step 5 (connectedness). In a non-vanishing term, $G \setminus l(n)$ is connected. A scalar line is never a bridge, because every vertex has an even number of scalar legs, so the scalar lines form closed charge loops. A photon bridge would leave two pieces with one external vector leg each, i.e. a C6 pair. Each such piece has an odd number of vector legs on a closed scalar strand, so it vanishes ($\operatorname{tr} q^{\mathrm{odd}} = 0$) after the gauge-out of B_0 at p. 433. So the tree of (1.33) for $G \setminus l(n)$ joins all cubes of Λ. Labels at one vertex have overlapping supports, since the term vanishes otherwise. Hence d over labels equals d over vertices, and the tree length d' of $G \setminus l(n)$ satisfies $d' \geq d(\Lambda)$.

Step 6 (assembly). Apply Proposition E.97 to the cut class, with the external field of Step 2 and the colour factor of Step 4. This gives (E.6), where $e(t_k)^{d_v}\lambda(t_k)^{d_s} \leq P(t_k)$ and $\mathrm{e}^{-\delta_T d'} \leq \mathrm{e}^{-\delta d(\Lambda)}$. The quantity $\mathcal{V}$ is a c-number, so no field restriction and no factor $p(t)$ or $\lambda^{-1/4}$ enters. □

We name the divergent two-leg types by $(E_s, E_v, E_\partial, o)$, following [3, Cor. 2.3, p. 429; (3.6), (3.18), (3.21), (3.25), (3.31)]:

- C1 $= (2, 0, 0, 1)$ and C2 $= (2, 0, 0, 2)$, the scalar self-energies;
- C1′ $= (2, 0, 1, 1)$, the self-energy with a differentiated leg;
- C3 $= (0, 2, \cdot, 1)$ and C4 $= (0, 2, \cdot, 2)$, the vacuum polarisations;
- a C6 pair: two pieces with one external vector leg each, joined by a photon line.

Proposition E.87 (degrees and the cut table)**.** *Let $d = 3$, and let D be Bałaban's degree* [3, (2.1)–(2.3)]. *A mass vertex carrying $\delta m^2_{\mathrm{fin}}$ or $\delta m^2_{K,k+1}$ (a* finite mass vertex*) has degree* 0. *A mass vertex carrying the contribution of a graph G_0 to δm^2_{k+1} has degree $D(G_0)$.*

(a) *Let G be a connected graph built from the vertices (1.6), (1.8), (1.10) and from mass vertices of divergent self-energy graphs. Then $D(G) = o(G) + \frac{1}{2}(E_s + E_v) + E_\partial - 3$. If $D(G) \leq 0$ and G has V vertices, then $V + E_s + E_v + 2E_\partial \leq 6$.*

(b) *Let G be a vacuum graph with $D(G) \leq 0$, and let ℓ be a line of G. The components of $G \setminus \ell$ with $D \leq 0$ are of the following kinds only:*
- *if ℓ is a scalar line that is not differentiated: C1, C2, or a bare vertex;*
- *if ℓ is a scalar line differentiated once: C1′;*
- *if ℓ is a scalar line differentiated twice: none;*
- *if ℓ is a photon line: C3, C4, or the two pieces of a C6 pair.*

Of the 59 *non-vanishing isomorphism classes of type C2,* 38 *are one-particle irreducible. The other* 21 *are chains H_1–ℓ'–H_2 of two order-*1 *two-scalar-leg pieces joined by one scalar line. Vacuum graphs with $o \geq 4$ give $D(G \setminus \ell) \geq 2$. Vacuum graphs with a finite mass*

vertex give only bare vertices, convergent pieces, and vanishing pieces with an odd number of vector legs on a closed scalar strand.

Proof. (a) By [3, (2.1)], a leg external to G contributes 0 in place of $-\frac{1}{2}$, and a differentiation of an external leg contributes 0 in place of -1. Summing over the vertices and using [3, (2.3)] gives $D(G) = \sum_v D(v) + \frac{1}{2}(E_s + E_v) + E_\partial - 3 + \sum \deg$. By the list on [3, p. 423], the vertex (1.6) has $D = 1 = o(v)$, and the vertices (1.8), (1.10) with n vector legs have $D = n/2 = o(v)$. For a mass vertex, $D(v) + \deg = 2 + D(G_0) = o(v)$, since $D(G_0) = o(G_0) - 2$ by induction on the order. Every such vertex has $o(v) \geq \frac{1}{2}$, so $0 \geq D \geq \frac{1}{2}V + \frac{1}{2}(E_s + E_v) + E_\partial - 3$.

(b) A vertex (1.9) or (1.11) has at least $\bar{n} + 1 > 13$ vector legs, so a graph containing one has $D > 0$ [3, p. 423, (iii)–(iv)]. A connected graph containing a vertex (1.13)–(1.15) has $D \geq 1$ by Statement K.20. A finite mass vertex has $D_G(v) \geq 2$. By (a), the vacuum graphs with $D \leq 0$ therefore lie in an explicit finite set: 129 vertex multisets, with all pairings of their legs. For each graph and each line, the components of $G \setminus \ell$ and their degrees are computed from [3, (2.1)–(2.3)], and they are cross-checked against (a). We carried out this finite enumeration by computer, with two independent programs whose tables agree. The counts of (graph, line) pairs are: C1 8, C2 1069, C1′ 5, C3 4, C4 751. All other non-vanishing components are convergent. Cutting a line adds two external legs, so for $o \geq 4$ part (a) gives $D(G \setminus \ell) \geq o - 3 + 1 \geq 2$. □

Lemma E.88 (one-particle-reducible chains need no counterterm)**.** *Work after the transfer of* [3, p. 433] *to a cube* □ *of* $\eta\mathbb{Z}^3$ *(Proposition E.67). Let* $S^{(1)}$ *be the amputated order-*1 *one-particle-irreducible two-scalar-point operator at fixed localisation labels. It consists of the order-*e^2 *and order-*λ *graphs and of the order-*1 *counterterm vertex, with derivative legs written as lattice differences. Write* $S^{(1)} = \Sigma_{nn} + \Sigma_{nd}\partial + \partial^*\Sigma_{dn} + \partial^*\Sigma_{dd}\partial$ *according to where the differences act* [3, (1.21)]*. Then:*

(a) *the sum of the one-particle-reducible members of a C2 family at fixed labels is* $\mathcal{K} := (S_1^{(1)} - \mathrm{ct}_1)\, K_{\ell'}\, (S_2^{(1)} - \mathrm{ct}_2)$*, where* $K_{\ell'}$ *is the kernel of the scalar line* ℓ'*;*

(b) $\sum_{x'} \mathcal{K}(x, x') = \sum_{y,y'} (S_1^{(1)} - \mathrm{ct}_1)(x, y) K_{\ell'}(y, y') (\partial_\nu^* c_\nu)(y')$*, where* $c_\nu(y') := \sum_{x'} \Sigma_{dn,\nu}(y', x')$ *is bounded and Lipschitz at the unit scale;*

(c) [3, (3.19)] *applies to* $\mathcal{K}$ *with zero counterterm. Both resulting terms have positive degree, and all their proper divergent subgraphs are renormalised. Hence* $\mathcal{K}$ *obeys bounds of the type* (E.6)*, by* [3, Props. 2.1–2.2, pp. 424–428].

Proof. (a) As formal power series, coefficient by coefficient, the amputated connected two-point function is $S + SK_{\ell'}S + \cdots$. At $o = 2$, a one-particle-reducible two-scalar-leg graph is cut by a scalar bridge into two connected two-leg pieces, each of order ≥ 1. A photon bridge would leave a piece with one external vector leg, which vanishes by $\mathrm{tr}\, q^{\mathrm{odd}} = 0$ after the gauge transformation of [3, p. 433]. A two-leg piece of order $\frac{1}{2}$ is a single vertex (1.8) with its vector leg external, and this is not of type C2. The subgraph counterterms are present in the family, and the labels factor because they are attached to the pieces.

(b) The counterterm is the zeroth moment of the nn part [3, p. 417, (3.9)]. So $\sum_{x'}(\Sigma_{nn} - \mathrm{ct})(y', x') = 0$, and tadpole pieces cancel their counterterms exactly [3, p. 435]. A term with a difference acting at x' has zero sum, because a sum of differences of a summable function vanishes, on $\eta\mathbb{Z}^3$ and over a period. The remaining terms give $\partial_{y'}^* c(y')$. The propagator $G_k(0)$ is only block periodic [3, p. 418]. After the replacement of [3, p. 437], however, the C^ξ part is translation invariant and the remainder is regular. So c depends on y' only through the unit-scale localisation functions and the regular part, and it is bounded and Lipschitz at the unit scale [3, (3.22)–(3.24)].

(c) [3, (3.19)] is the identity

$$\varphi(x)\mathcal{K}(x, x')\varphi'(x') = \varphi(x)\mathcal{K}(x, x')|x' - x|^\alpha \frac{\varphi'(x') - \varphi'(x)}{|x' - x|^\alpha} + \varphi(x)\varphi'(x) \sum_{x'} \mathcal{K}(x, x').$$

The first term has degree $\alpha > 0$. By (b), in the second term one difference on the line is replaced by a unit-scale difference of localisation functions, so its degree is ≥ 1. The inner pieces are of types C1, C1′ or tadpole, with their counterterms present by (a). The torus counterterm δm_k^2 is built with the torus propagator [3, (1.28)], while the members after the transfer use $G_k(0)$ on $\eta\mathbb{Z}^3$. Their difference is a finite mass vertex whose coefficient is exponentially small in ℓ_T/t_k, by [3, (2.10), p. 426]. The vacuum cuts of this vertex produce no divergent class (Proposition E.87). □

The order-e^2 family with one D-line. *Setting.* On T_ε we use $\langle f, g\rangle = \sum_x \varepsilon^3 f(x)g(x)$, and kernels are taken relative to $\sum \varepsilon^3$. We put $\nabla_\mu = \nabla_\mu^+$, $\Delta_\mu = \nabla_\mu^+\nabla_\mu^-$ and $-\Delta_\varepsilon = -\sum_\mu \Delta_\mu$. For $z \in T_\varepsilon$, $|z|_1$ and $|z|_\infty$ are the norms of the minimal representative, and $r := |z|_\infty$. Let c be the kernel of $C = (-\Delta_\varepsilon + M^2)^{-1}$. Let $G_k = G_k^M$ be King's scalar operators (Statement K.17(S4)), with slices $G_{(i)} := G_{i+1} - G_i$, $G_0 := 0$, and put $\mathcal{C}_k := C - G_k \geq 0$. The photon slices are $H_j \geq 0$ (Statement K.17), $\delta_j := \kappa\mu_*^2 L^2 t_j^2$, and $\Gamma^B = \sum_{j<K} \delta_j H_j$. For a two-variable function f, ∇f and $\nabla' f$ are the forward differences in the first and second variable.

For a real symmetric kernel $D \geq 0$ and a real symmetric scalar covariance S put

$$\mathrm{Bub}(S,S)(x,y) := 2\varepsilon^{-2}\big[S(x+\varepsilon e_\mu, y+\varepsilon e_\mu)S(x,y) - S(x+\varepsilon e_\mu, y)S(x, y+\varepsilon e_\mu)\big],$$

$$V(D;S) := \sum_\mu \Big\{ -e^2 \sum_x \varepsilon^3 D(x,x)S(x+\varepsilon e_\mu, x) + \frac{e^2}{2}\sum_{x,y} \varepsilon^6 D(x,y)\,\mathrm{Bub}(S,S)(x,y)\Big\}.$$

This is the order-e^2 vacuum coefficient produced by a photon of covariance $\delta_{\mu\nu}D$; it consists of the seagull and the bubble whose two vector legs both end on the photon line. The order-e^2 part of the u-dependence of the vacuum terms is $u\sum_{j<K}\delta_j V(H_j;C)$. A graph has top index $\geq k$ exactly when its D-slice has index $\geq k$, or when that index is $< k$ and the graph contains a line of $\mathcal{C}_k$.

Leg partition [3, p. 420]. Fix $t \in \{t_i\}$. Put $\phi(s) := \cos^2(\pi s/2)\mathbf{1}_{|s|<1}$ and $h_a(x) := \prod_{i=1}^3 \phi(x_i/t - a_i)$ for $a \in (\mathbb{Z}/(\ell_T/t))^3$. Then $\sum_a h_a = 1$ and $0 \leq h_a \leq 1$, and h_a is supported in the open cube S_a of side $2t$ centred at ta. At most $N_{\mathrm{ov}} = 8$ of the h_a are non-zero at any point, and $|h_a(x) - h_a(y)| \leq m_1(x-y) := \min(c_{h1}|x-y|_1/t, 1)$ with $c_{h1} = \pi/2$. Each vector leg is placed at the base point of its bond. Put

$$F_{D,S}(a,a') := \sum_\mu \Big\{ -e^2 \sum_x \varepsilon^3 h_a h_{a'}(x) D(x,x) S(x+\varepsilon e_\mu, x) + \frac{e^2}{2}\sum_{x,y} \varepsilon^6 h_a(x) h_{a'}(y) D(x,y)\,\mathrm{Bub}(S,S)(x,y)\Big\}.$$

With the partition at scale $t = t_k$, put

$$F^{\geq k} := u\Big[\sum_{k\leq j<K} \delta_j F_{H_j,C} + \sum_{j<k} \delta_j\big(F_{H_j,C} - F_{H_j,G_k}\big)\Big], \qquad F^{=k} := F^{\geq k} - F^{\geq k+1},$$

where $F^{\geq k+1}$ uses the same scale-t_k partition.

Lemma E.89 (the localised family and its Ward form)**.** *For real symmetric D and S, $F_{D,S}(a,a') = F_{D,S}(a',a)$ and $\sum_{a,a'} F_{D,S}(a,a') = V(D;S)$. Moreover*

$$F_{D,C}(a,a') = -\tfrac{1}{2}\sum_\mu \sum_{x,y} \varepsilon^6 h_a(x) h_{a'}(y) D(x,y) \Pi_\mu(x-y),$$

where $\Pi_\mu(z) := 2e^2 c(\varepsilon e_\mu)\varepsilon^{-3}\delta_{z,0} - B_\mu(z)$ and $B_\mu(z) := 2e^2\varepsilon^{-2}[c(z)^2 - c(z+\varepsilon e_\mu)c(z-\varepsilon e_\mu)]$. If in addition $D \geq 0$, let $A \sim N(0,D)$, and put $g := h_a \otimes h_{a'} + h_{a'} \otimes h_a$, $\varphi := gD$, $\Delta\varphi(x,y) := \varphi(x,x) + \varphi(y,y) - 2\varphi(x,y)$ and $\widehat{\Pi}_\mu(0) := \sum_z \varepsilon^3 \Pi_\mu(z)$. Then

$$F_{D,C}(a,a') = -\tfrac{1}{4}\sum_\mu \Big[\sum_x \varepsilon^3 g(x,x) D(x,x)\widehat{\Pi}_\mu(0) + \tfrac{1}{2}\sum_{x,y} \varepsilon^6 \Delta\varphi(x,y) B_\mu(x-y)\Big].$$

Here $\Delta\varphi = g(x,y)\Delta D + D(x,y)\Delta g + (g(x,x)-g(x,y))(D(x,x)-D(x,y)) + (g(y,y)-g(x,y))(D(y,y)-D(x,y))$, *with* $\Delta D(x,y) = \mathbb{E}(A(x)-A(y))^2$ *and* $\Delta g(x,y) = 2(h_a(x)-h_a(y))(h_{a'}(x)-h_{a'}(y))$.

Proof. $\mathrm{Bub}(S,S)$ is symmetric for symmetric S. Exchanging x and y and using $D(y,x)=D(x,y)$ gives the symmetry; the seagull sum is symmetric in (a,a'). The total follows from $\sum_a h_a = 1$. For $S=C$, we have $S(x+\varepsilon e_\mu,x) = c(\varepsilon e_\mu)$ and $\frac{e^2}{2}\,\mathrm{Bub}(C,C)(x,y) = \frac12 B_\mu(x-y)$.

For the Ward form, symmetrise: $F = -\frac14\sum_\mu\sum\varepsilon^6\varphi\Pi_\mu$. Then insert $\varphi(x,y) = \frac12\varphi(x,x) + \frac12\varphi(y,y) - \frac12\Delta\varphi(x,y)$. The first two terms give $\widehat{\Pi}_\mu(0)$, because Π_μ is even. In the third, $\Delta\varphi(x,x)=0$ and $\Pi_\mu = -B_\mu$ off the diagonal. The decomposition of $\Delta\varphi$ follows by writing $g(x,x) = g(x,y)+\alpha$, $g(y,y) = g(x,y)+\beta$, $D(x,x) = D(x,y)+\gamma$ and $D(y,y) = D(x,y)+\gamma'$. □

Proposition E.90 (per-anchor bound)**.** *Let $D \ge 0$ be real symmetric, put $s_0 := s_0(D)$ and $s_1 := s_1(D)$, and let the partition have any scale t. Let w_T, $\hat B$ and $I = I(\ell_T)$ be as in Statement K.18, and put*

$$\hat S := 312\hat B + 156 I, \quad C_1 := 4N_{\mathrm{ov}}c_{h1}(156\hat B + 78 I), \quad C_2 := 104(N_{\mathrm{ov}}+1)\sqrt{18c_{h1}}\,\hat B, \quad C_3 := 312(N_{\mathrm{ov}}+1)c_{h1}I.$$

Then for every label a,

$$\sum_{a'}|F_{D,C}(a,a')| \le \tfrac32 e^2(2t)^3\Big\{s_0 w_T + \sqrt{s_0 s_1}\,\hat S + (C_1+C_3)\frac{s_0}{t} + C_2 s_0^{3/4}s_1^{1/4}t^{-1/2}\Big\}.$$

Proof. Put $z = y-x$ and $q := \min(|z|_1^2 s_1, 4s_0)$. Write $A(y)-A(x)$ as a sum of $|z|_1/\varepsilon$ differences and apply Cauchy–Schwarz. This gives $|D(x,y)| \le s_0$, $\Delta D \le q$ and $|D(x,x)-D(x,y)| \le \sqrt{s_0 q}$. Put $\delta h := h(x)-h(y)$. At most $2N_{\mathrm{ov}}$ labels have $\delta h_{a'} \neq 0$, so $\sum_{a'}|\delta h_{a'}| \le 2N_{\mathrm{ov}}m_1$, and $\sum_{a'} g(x,y) = h_a(x)+h_a(y)$. By Lemma E.89,

$$\sum_{a'}|\Delta\varphi| \le (h_a(x)+h_a(y))[q + 2N_{\mathrm{ov}}m_1\sqrt{s_0 q}] + |\delta h_a|[4N_{\mathrm{ov}}s_0 m_1 + 2\sqrt{s_0 q}].$$

We have $|\delta h_a| \le m_1(z)(\mathbf{1}_{S_a}(x)+\mathbf{1}_{S_a}(y))$, $\sum_{a'} g(x,x) = 2h_a(x)$, and $\sum_x \varepsilon^3 h_a \le (2t)^3$. With Statement K.18(a), this yields

$$\sum_{a'}|F| \le \tfrac34(2t)^3[2e^2 s_0 w_T + \Sigma_A + 4N_{\mathrm{ov}}s_0\Sigma_B + 2(N_{\mathrm{ov}}+1)\Sigma_C],$$

where $\Sigma_A, \Sigma_B, \Sigma_C$ are the suprema over μ of $\sum_z \varepsilon^3|B_\mu|w$ with the weights $w = q$, m_1^2 and $\sqrt{s_0 q}\,m_1$, respectively.

On the fundamental cell, Statement K.18(b) gives $|B_\mu| \le 2e^2[(g_1+\rho_1)^2 + (g_0+\rho_0)(g_2+\rho_2)]$ with $g_a \le \hat G_a r^{-a-1}$, and $|z|_1 \le 3r$. The $\varepsilon\mathbb{Z}^3$ parts are estimated with the shell sums of Statement K.18(e) and $\int_0^\infty \min(A, B/r^2)\,dr = 2\sqrt{AB}$. The image and cross terms are estimated with $\min(9r^2 s_1, 4s_0) \le 6r\sqrt{s_0 s_1}$, $m_1^2 \le 3c_{h1}r/t$ and $\sqrt{s_0 q}\,m_1 \le 6c_{h1}s_0 r/t$. The result is

$$\Sigma_A \le 2e^2\sqrt{s_0 s_1}\hat S, \qquad \Sigma_B \le 2e^2 c_{h1}(156\hat B + 78 I)/t,$$

$$\Sigma_C \le 2e^2\big[52\sqrt{18c_{h1}}\hat B s_0^{3/4}s_1^{1/4}t^{-1/2} + 156 c_{h1}s_0 I/t\big].$$

Collecting terms with $\frac34\cdot 2e^2 = \frac32 e^2$ proves the bound. Every weight vanishes as $r\to 0$, so no logarithm of t/ε appears. □

Lemma E.91 (members with a coarse scalar line)**.** *Let $1 \le k \le K$ and $0 \le j < k$, and put $F^{\mathrm{cs}}_{j,k} := F_{H_j,C} - F_{H_j,G_k}$. Let $s_0 = s_0(H_j)$, $d_0 = s_0(\mathcal{C}_k)$ and $d_1 = s_1(\mathcal{C}_k)$. Then for every label a and every partition scale t,*

$$\sum_{a'}|F^{\mathrm{cs}}_{j,k}(a,a')| \le (2t)^3 e^2\Big[5s_0 d_0 + 3s_0 d_1 M^{-2} + 6s_0\sqrt{d_0 d_1}\Lambda_1 + 3d_1 M^{-1}\sqrt{\|H_j\| s_0 d_0} + 3\sqrt{d_0 d_1}\sqrt{\|H_j\| s_0 d_0}\Big].$$

Consequently $|u|\sum_{j<k}\delta_j\sum_{a'}|F^{\mathrm{cs}}_{j,k}(a,a')| \le (2t)^3|u|e^2\kappa\mu_*^2L^2A_<t_k^{-2}$, *where*

$$A_< := 5\sigma_0\gamma\varepsilon_0 + 3\sigma_0\theta M^{-2} + 6\sigma_0\sqrt{\gamma\theta}\Lambda_1\varepsilon_0^{1/2} + \frac{3\theta\sqrt{h\sigma_0\gamma}\,\varepsilon_0^{1/2}}{(L^{5/2}-1)M} + \frac{3\gamma\sqrt{\theta h\sigma_0}\,\varepsilon_0}{L^{5/2}-1}.$$

Proof. Write $D = H_j$ and $\mathcal{C} = \mathcal{C}_k$, so that $G_k = C - \mathcal{C}$. By bilinearity, $F^{\mathrm{cs}}_{j,k}(a,a')$ is

$$\sum_\mu\Big\{-e^2\sum_x\varepsilon^3h_ah_{a'}D(x,x)\mathcal{C}(x+\varepsilon e_\mu,x) + e^2\sum_{x,y}\varepsilon^6h_a(x)h_{a'}(y)D(x,y)J(x,y)\Big\},$$

where $J := C\nabla\nabla'\mathcal{C} + \mathcal{C}\nabla\nabla'C - \nabla C\nabla'\mathcal{C} - \nabla\mathcal{C}\nabla'C - \mathcal{C}\nabla\nabla'\mathcal{C} + \nabla\mathcal{C}\nabla'\mathcal{C}$. Call the seagull term τ_1 and the six terms of J τ_2–τ_7.

All terms except τ_3 are bounded in absolute value, after summing over a' with $\sum_{a'}h_{a'}=1$, by Statement K.18(c),(d):

- τ_1 by s_0d_0, and τ_2 by $d_1s_0M^{-2}$;
- τ_4 and τ_5 by $s_0\sqrt{d_0d_1}\Lambda_1$ each;
- τ_6 by $d_1(\|D\|s_0)^{1/2}(M^{-2}d_0)^{1/2}$, and τ_7 by $\sqrt{d_0d_1}(\|D\|s_0d_0)^{1/2}$.

Each of these carries a factor 3 from the sum over μ.

τ_3 is summed over μ first. Since $\nabla_x\nabla'_yc(x-y) = -(\Delta_\mu c)(x-y)$ and $(-\Delta_\varepsilon + M^2)c = \varepsilon^{-3}\delta$, it equals $h_{a'}(x)D(x,x)\mathcal{C}(x,x) - M^2\sum_y\varepsilon^3h_{a'}(y)D\mathcal{C}\,c(x-y)$, which is at most $2s_0d_0$ after the sum over a'. Finally $\sum_x\varepsilon^3h_a \le (2t)^3$.

For the sum over j, insert $\delta_j \le \kappa\mu_*^2L^2t_j^2$, $s_0 \le \sigma_0/t_j$, $\|H_j\| \le ht_j^2$, $d_0 \le \gamma t_k^{-2}$ and $d_1 \le \theta t_k^{-3}$. Then use $\sum_{j<k}t_j \le t_k$, $\sum_{j<k}t_j^{5/2} \le t_k^{5/2}/(L^{5/2}-1)$ and $t_k \le \varepsilon_0$. □

Theorem E.92 (order-e^2 anchor bounds)**.** *Let* $1 \le k \le K-1$, $|u| \le 1$ *and* $M \in \{m_1, m_2\}$, *and use the partition at scale* $t = t_k$. *Put*

$$C_W := 12e^2|u|\kappa\mu_*^2L^2\varepsilon_0^2\big[\sigma_0w_T\varepsilon_0 + 0.2654\sqrt{\sigma_0\theta_H}\,\hat{S} + (C_1+C_3)\sigma_0 + 2.415\,C_2\sigma_0^{3/4}\theta_H^{1/4}\big]$$

and $C_{\mathrm{loc}} := C_W + 8|u|e^2\kappa\mu_*^2L^2A_<$.

(a) $F^{=k}(a,a') = F^{=k}(a',a)$, *and*

$$\sum_{a,a'}F^{=k}(a,a') = u\Big[\delta_kV(H_k;G_{k+1}) + \sum_{j<k}\delta_j\big(V(H_j;G_{k+1}) - V(H_j;G_k)\big)\Big].$$

This is the part carried by graphs of top index exactly k; *all its lines have index* $\le k$.

(b) *For every label* a, *the sums* $\sum_{a'}|F^{\ge k}(a,a')|$ *and* $\sum_{a'}|F^{\ge k+1}(a,a')|$ *are at most* $C_{\mathrm{loc}}t_k$, *and* $\sum_{a'}|F^{=k}(a,a')| \le 2C_{\mathrm{loc}}t_k$.

(c) *Let* c_G, c_H, δ_1 *be as in Statement K.19, and fix* $R_0 > 0$. *Put* $Z_p := \sum_{n\ge0}2^{np}\mathrm{e}^{-\delta_1R_0(2^n-1)}$, $C_b := 2c_G^2(Z_1Z_3 + Z_2^2)$ *and* $d_{aa'} := \mathrm{dist}(S_a, S_{a'})$. *If* $d_{aa'} \ge R_0t_k$, *then* $|F^{=k}(a,a')| \le C_{\mathrm{pair}}t_k^3\mathrm{e}^{-3\delta_1d_{aa'}/t_k}$, *with* $C_{\mathrm{pair}} := 384c_HC_be^2|u|\kappa\mu_*^2L^2$. *For every* $R \ge R_0t_k$,

$$\sum_{a':\,d_{aa'}\ge R}|F^{=k}(a,a')| \le C_{\mathrm{dp}}t_k^3\mathrm{e}^{-2\delta_1R/t_k}, \qquad C_{\mathrm{dp}} := C_{\mathrm{pair}}\sum_{w\in\mathbb{Z}^3}\mathrm{e}^{-\delta_1(|w|_\infty-2)_+}.$$

If the scale-k block containing the centre of S_a *lies in* $\mathcal{D}_k$, *and* $S_{a'}$ *meets* W_k^c, *then* $d_{aa'} > (r(t_k)-3)t_k$.

The constants C_{loc}, C_{pair} *and* C_{dp} *do not depend on* ε, k *or* K. *They are bounded for* $\ell_T \ge \ell_0$, *linear in* $|u|$ *(no sign of* u *is used), and finite at* $M = m_1$.

Proof. (a) $F^{\ge k}$ and $F^{\ge k+1}$ are linear combinations of the $F_{H_j,S}$ with symmetric $S \in \{C, G_k, G_{k+1}\}$, so Lemma E.89 gives the symmetry and the totals. The difference of the totals is u times

$$\delta_kV(H_k;C) - \delta_k\big(V(H_k;C) - V(H_k;G_{k+1})\big) + \sum_{j<k}\delta_j\big(V(H_j;G_{k+1}) - V(H_j;G_k)\big),$$

and $G_{k+1} = \sum_{i\le k} G_{(i)}$.

(b) Let $k \le j < K$. Apply Proposition E.90 to $D = H_j$ at $t = t_k$, with $s_0 \le \sigma_0/t_j$ and $s_1 \le \theta_H/t_j^3$ (Statement K.18(c)), and multiply by δ_j. The four terms become proportional to t_j, 1, t_j/t_k and $t_j^{1/2}t_k^{-1/2}$. Then use:

- $\sum_j t_j \le \varepsilon_0$, $\#\{j\} \le \log_2(\varepsilon_0/t_k)$ and $\sum_j t_j^{1/2} \le 2.415\,\varepsilon_0^{1/2}$;
- the factor $(2t_k)^3 = 8t_k^3$, the bound $t_k \le \varepsilon_0$, and $t_k^2 \log_2(\varepsilon_0/t_k) \le \varepsilon_0^2/(2\mathrm{e}\ln 2) \le 0.2654\,\varepsilon_0^2$.

This gives at most $C_W t_k$. The members $j < k$ add at most $8t_k^3|u|e^2\kappa\mu_*^2L^2A_<t_k^{-2}$, by Lemma E.91. For $F^{\ge k+1}$ at the same scale t_k, the slices $j \ge k+1$ form a subset of those above. The coarse members carry $\mathcal{C}_{k+1}$, and Lemma E.91 at $k+1 \le K$ gives $(2t_k)^3A_<t_{k+1}^{-2} \le 8A_<t_k$, up to the common factor. The third bound follows from $F^{=k} = F^{\ge k} - F^{\ge k+1}$.

(c) If $d_{aa'} > 0$, there is no seagull term, and

$$F^{=k}(a,a') = u\frac{e^2}{2}\sum_\mu \sum_{x,y} \varepsilon^6 h_a(x)h_{a'}(y)\Big[\delta_k H_k\,\mathrm{Bub}_{k+1} + \sum_{j<k}\delta_j H_j(\mathrm{Bub}_{k+1} - \mathrm{Bub}_k)\Big](x,y),$$

where $\mathrm{Bub}_{k'} := \mathrm{Bub}(G_{k'}, G_{k'})$. For $|x-y| \ge d_{aa'} \ge R_0 t_k$ we have $\sum_{i\le k} t_i^{-p}\mathrm{e}^{-\delta_1|x-y|/t_i} \le Z_p t_k^{-p}\mathrm{e}^{-\delta_1 d_{aa'}/t_k}$. With Statement K.19, this gives $|\mathrm{Bub}_{k'}| \le C_b t_k^{-4}\mathrm{e}^{-2\delta_1 d_{aa'}/t_k}$ for $k' \in \{k, k+1\}$. Moreover $\delta_j|H_j| \le \kappa\mu_*^2L^2c_H t_j\mathrm{e}^{-\delta_1 d_{aa'}/t_k}$ for $j \le k$, and $\sum_{j\le k} t_j \le 2t_k$. With $\sum_x \varepsilon^3 h_a \le (2t_k)^3$,

$$|F^{=k}| \le \tfrac{3}{2}e^2|u|(2t_k)^6\kappa\mu_*^2L^2c_H \cdot 2t_k \cdot 2C_b t_k^{-4}\mathrm{e}^{-3\delta_1 d_{aa'}/t_k},$$

and $\frac{3}{2}\cdot 64\cdot 4 = 384$. For the partner sum, use $d_{aa'} \ge R$ and $d_{aa'} \ge (|a'-a|_\infty - 2)_+ t_k$ in the minimal-image metric, and note that the minimal-image map from the label torus to $\mathbb{Z}^3$ is injective. For the last claim: every point of S_a is within ∞-distance t_k of the block, and every point of $S_{a'}$ is within $2t_k$ of W_k^c.

Uniformity: w_T, $\hat{S}$, I, Λ_1, σ_0, θ_H, γ, θ and h are free of ε and bounded for $\ell_T \ge \ell_0$ (Statement K.18). The only count of scales is absorbed by t_k^2. The factors M^{-1} and M^{-2} are finite at m_1. □

Each localisation of the seagull alone is of order ε^{-1}. In Lemma E.89 this part cancels inside the single localisation (a,a') against the zeroth moment of the bubble, and only the image sum $\widehat{\Pi}_\mu(0)$ remains.

Lemma E.93 (the order-e^2 family with one D-line)**.** *Let $1 \le k \le K-1$, $|u| \le 1$ and $r(t_k)-4 \ge R_0$. Let C_{loc} and C_{dp} be as in Theorem E.92. For the order-e^2 vacuum family with one D-line at step k, the following hold.*

(i) *The sum over the bad localisations is at most $4C_{\mathrm{loc}}t_k|\mathcal{L}_k|_k + 2C_{\mathrm{dp}}t_k^3\mathrm{e}^{-2\delta_1(r(t_k)-4)}|\mathcal{D}_k|_k$.*

(ii) *For every union X of scale-k blocks of W_k^c, the sum over the outside localisations with anchor in X is at most $2C_{\mathrm{loc}}t_k|X|_k$.*

Both bounds are proportional to $|u|$.

Proof. Under convention (G), every moved piece of this family is localised once, at its own scale k, by the pair of labels of its two photon legs. This localisation is $F^{=k}(a,a')$ (Theorem E.92(a)). A pair (a,a') is bad if exactly one of $S_a, S_{a'}$ lies in W_k. By the symmetry in Theorem E.92(a), the bad sum is at most $2\sum_{S_a\subset W_k}\sum_{S_{a'}\not\subset W_k}|F^{=k}(a,a')|$. For a with block in $\mathcal{L}_k$, use $\sum_{a'}|F^{=k}| \le 2C_{\mathrm{loc}}t_k$. For a with block in $\mathcal{D}_k$, Theorem E.92(c) gives $d_{aa'} > (r(t_k)-4)t_k \ge R_0 t_k$, so its partner sum applies with $R = (r(t_k)-4)t_k$. For (ii), the anchor a runs over X, and $\sum_{a'}|F^{=k}| \le 2C_{\mathrm{loc}}t_k$. The family carries exactly one D-line, so it is proportional to u. □

The region sum. Notation at scale k (Bałaban's large-field sets on T_ε, all distances being torus distances):

- A *tuple* $\mathfrak{t}_k = (P_v, Q_v, R_v, P_s, Q_s, R_s)$ is a 6-tuple of subsets of $\Lambda_7^{(k-1)\prime}$ (scale-$(k+1)$ blocks, bonds and sites of scale k; $\Lambda_7^{(-1)\prime} := T_1^{(0)}$), the set of violated restrictions. We write $|\mathfrak{t}|$ for its total number of elements.
- $\Lambda_0^{(k)}$ is the union of the large blocks contained in $\Lambda_7^{(k-1)\prime}$ at distance $\geq r(t_k)$ from the elements of $\mathfrak{t}_k$, and $\Lambda_{i+1}^{(k)c}$ is the union of the large blocks within distance $r(t_k)$ of $\Lambda_i^{(k)c}$ [2, (2.7)–(2.8), p. 558; p. 570].
- Given $\Lambda_7^{(k-1)\prime}$, the set $\Lambda_0^{(k)}$ is a function of $\mathfrak{t}_k$. Hence a sequence $\mathfrak{S} = (\mathfrak{t}_0, \dots, \mathfrak{t}_{K-1})$ determines at most one admissible region sequence.
- $|\cdot|_k$ counts scale-k points, and $N_i := |T_1^{(i)}| = |T|/t_i^3$.

Lemma E.94 (geometric lemma)**.** *Put $g(t) := 8r(t) + 8\sqrt{d}\,M_b + \sqrt{d}(L+1)$, $g_0 := 8 + (8\sqrt{d}M_b + \sqrt{d}(L+1))/R$, $C_r := \sum_{n\geq 0} L^{-n}(1+n\log L)^{\varrho}$ and $C_V := (2g_0C_r+1)^d$. For every admissible region sequence generated by $(\mathfrak{t}_0, \dots, \mathfrak{t}_{K-1})$ and every $k < K$,*

$$|\Lambda_7^{(k)c}|_k \leq C_V\, r(t_k)^d \sum_{i\leq k} |\mathfrak{t}_i|.$$

Proof. (i) The set $\Lambda_0^{(k)c}$ is contained in the union of the large blocks not contained in $\Lambda_7^{(k-1)\prime}$ and the large blocks at distance $< r(t_k)$ from an element of $\mathfrak{t}_k$. Moreover $\Lambda_7^{(k)c}$ lies in the $(7r(t_k)+7\sqrt{d}M_b)$-neighbourhood of $\Lambda_0^{(k)c}$ (scale-k units). A point of the second kind lies within $r(t_k)+\sqrt{d}M_b+\sqrt{d}L$ of an element of $\mathfrak{t}_k$. A point z of the first kind lies in a large block containing a point $z' \notin \Lambda_7^{(k-1)\prime}$, with $|z-z'| \leq \sqrt{d}M_b$; the unit block of z' contains a scale-$(k-1)$ point of $\Lambda_7^{(k-1)c}$, at distance $\leq \sqrt{d}$. (ii) By induction on k, every point of $\Lambda_7^{(k)c}$ lies within physical distance $D(i,k) := \sum_{m=i}^{k} t_m g(t_m)$ of an element of some $\mathfrak{t}_i$, $i \leq k$. (iii) Since $r \geq R$, $g(t) \leq g_0 r(t)$. For $t_k \leq 1$, $1 + \log t_{k-n}^{-1} = (1+\log t_k^{-1}) + n\log L \leq (1+\log t_k^{-1})(1+n\log L)$, so $r(t_{k-n}) \leq (1+n\log L)^{\varrho} r(t_k)$. With $t_{k-n} = L^{-n}t_k$ this gives $D(i,k) \leq g_0 C_r t_k r(t_k)$. (iv) A physical ball of radius D contains at most $(2D/t_k+1)^d$ scale-k points of $\mathbb{Z}^3$, hence at most that many of $T_1^{(k)}$, since projection to the torus does not increase the count. As $r \geq R \geq 1$, this is $\leq C_V r(t_k)^d$. A union bound over the elements of the tuples concludes. □

Lemma E.95 (region-sum lemma)**.** *Hypothesis (H-LF) at a point π: the large-field region sum of the closing estimate (*[2, (2.117), p. 582; (3.40)–(3.42), pp. 591–592] *for the slice model) is bounded by $\mathrm{e}^{C_0(\pi)|T|}$ times*

$$\sum_{\mathfrak{S}=(\mathfrak{t}_0,\dots,\mathfrak{t}_{K-1}) \text{ admissible}} \prod_{k<K} \mathrm{e}^{-c_*(\pi)p(t_k)^2|\mathfrak{t}_k|} \exp\Big(c_A(\pi)\sum_{k<K}(1+\log t_k^{-1})|\Lambda_7^{(k)c}|_k\Big),$$

with $c_*(\pi) > 0$, $c_A(\pi) < \infty$ and $C_0(\pi)$ independent of ε, ℓ_T and $\mathfrak{S}$. Assume $2p > d\varrho + 2$ and

$$\textit{(E.comb)} \qquad \hat{\ell}^{\,2p-d\varrho-2} \geq \frac{2c_A(\pi)C_V R^d}{c_*(\pi)b_0^2 \log L}, \qquad \hat{\ell} := 1 + \log \varepsilon_0^{-1}.$$

Then, uniformly in $\varepsilon = L^{-K'}$ and ℓ_T, the sum is at most $\exp\big(2(d+2)\Sigma_*(\varepsilon_0;\pi)|T|\big)$, where

$$\Sigma_*(\varepsilon_0;\pi) := \sum_{n\geq n_0} L^{3n} \exp\big(-\tfrac{1}{2}c_*(\pi)b_0^2(1+n\log L)^{2p}\big), \qquad n_0 := \log_L \varepsilon_0^{-1}.$$

Σ_* is finite, contains no c_A, and decreases to 0 as $\varepsilon_0 \downarrow 0$. At $p=6$, $\varrho=2$, $d=3$ we have $2p - d\varrho - 2 = 4$.

The constants c_A, c_*, C_V are evaluated at the bound ε_1 (stage 10a of Appendix A) before ε_0 is chosen by (E.comb) at stage 10b, so there is no cycle. The hypothesis (H-LF) is the form of the region sum in the closing step of the proof of Theorem 5.2 (Lemma F.39).

Proof. (1) *Costs to elements.* Put $x_i := \log t_i^{-1}$. For $k \geq i$ we have $t_k \geq t_i$, so $1 + \log t_k^{-1} \leq 1 + x_i$ and $r(t_k) \leq R(1+x_i)^{\varrho}$. By Lemma E.94,

$$\sum_{k<K} (1 + \log t_k^{-1}) |\Lambda_7^{(k)c}|_k \leq C_V R^d \sum_i |\mathfrak{t}_i| (K-i)(1+x_i)^{1+d\varrho},$$

and $K - i = \log_L(\varepsilon_0/t_i) \leq (1+x_i)/\log L$ since $\varepsilon_0 \leq 1$. (2) *Absorption.* The summand is therefore at most $\prod_i \exp\big(-[c_* b_0^2 (1+x_i)^{2p} - c_A C_V R^d (\log L)^{-1} (1+x_i)^{d\varrho+2}] |\mathfrak{t}_i|\big)$. Since $2p > d\varrho + 2$, $x \mapsto (1+x)^{2p-d\varrho-2}$ is increasing, and $x_i \geq \log \varepsilon_0^{-1}$. So (E.comb) makes the bracket at least $\frac{1}{2} c_* b_0^2 (1+x_i)^{2p} = \frac{1}{2} c_* p(t_i)^2$. (3) *Entropy.* By determinism, each $\mathfrak{S}$ occurs at most once. The region expansion [2, (2.9)] holds in the "$\leq$" direction, which is the only one used, and dropping admissibility only adds non-negative terms. Hence, with $a_i := \frac{1}{2} c_* p(t_i)^2$,

$$\sum_{\mathfrak{S}} \prod_i \mathrm{e}^{-a_i |\mathfrak{t}_i|} \leq \prod_i (1 + \mathrm{e}^{-a_i})^{n_i} \leq \exp\Big(\sum_i n_i \mathrm{e}^{-a_i} \Big),$$

where n_i is the number of possible elements of a tuple at scale i: P_v, P_s give at most N_i/L^d each, Q_v, Q_s at most dN_i each, and R_v, R_s at most N_i each, so $n_i \leq 2(d+2)N_i$. (4) *Scales.* $t_i = L^{-n}$ with distinct integers $n \geq n_0$ for distinct i, and $N_i = L^{3n}|T|$, so $\sum_i n_i \mathrm{e}^{-a_i} \leq 2(d+2)|T|\, \Sigma_*(\varepsilon_0; \pi)$. Σ_* converges because $2p > 1$. □

Remark E.96. Lemma E.95 holds for every c_A. Lemma E.63 uses it with $c_A \mapsto c_A + C_{\mathrm{old}}$. Under convention (G), the cost of the vacuum counterterm on Col_k is contained in C_{old}.

Bałaban's Proposition 1 with its sketched steps written out.

Proposition E.97 (Bałaban's Proposition 1 on the torus)**.** *Let $\bar{n}$, α_0, k, η, the domains $\Omega, \Omega_1, \Omega_2$, $\tilde{B}$, the renormalised families and the external fields be as in Statement K.21. Then the bound of Statement K.21 holds on T_ε, and on King's torus with photon mass $\mu_* > 0$ and D-coloured lines. It holds with a tree rate $\delta_T > 0$ and a constant $O(1) = O(\alpha_0, \bar{n}; d, L, a, M_b)$, both independent of ε, k, the domains and $\tilde{B}$. The rate δ_T depends on $\bar{n}$ through the number of lines; it is not asserted to depend on d only. The coloured families with $m = 0$, $m \geq 2$, or $m = 1$ and $o \geq 2$ carry the extra factor $(|u|c_k)^m$. This proposition is Statement K.21.*

The proposition rests on the written formulas of [3] (including p. 433) and on [4]. Two items are treated separately, in Proposition E.110:

- the two-variable form of the remainder kernel of [3, (1.16), (2.5)] (Lemmas E.117 and E.118);
- the remainder on the cube □ of [3, p. 433] (Lemma E.121).

Remark E.98 (the sketched steps of [3, Prop. 1])*.* The steps of [3, pp. 418–445] that are only sketched there fall into the items below. For each item we name the closest step of Table 2 (by page), and the results of this paper that treat it. The domain conditions of all items are Proposition 5.8 and Corollary 5.10.

- *I1* (p. 432, family combinatorics, "inspecting all possible cases"; cf. S10). For the vacuum families of the moved pieces, the classes, their cut table and the one-particle-reducible C2 chains are given by Proposition E.87, Lemma E.166 and Lemma E.88. The closure of the classes under the operations of pp. 434–445 in general, including the nested cases, is part of Statement K.21.
- *I2* (p. 422, bad localisations). The vacuum part is Lemma E.86 with Proposition E.62, together with Lemma E.93 for the order-e^2 family with one D-line. The mass part is the same localisation sum for the two-leg classes (Lemma E.170).

- *I3* (p. 418, pointwise counterterm bounds). The final counterterm bounds are Propositions E.9 and E.21. The bounds at the intermediate scales are part of Statement K.21.
- *I4* ([1, Props. 2.1–2.3], behind (2.10)–(2.12)). This item is cited, as Proposition 4.5. The slices at zero background are also Theorem E.47.
- *I5* (p. 412, Hölder norms of the background). The Hölder bounds of the vector and scalar backgrounds in the norm [3, (1.32)] are part of Statement K.21.
- *(3.35)* (p. 443, leg transport; inside S21). This is Theorem E.53, with the pinned case [3, (2.21e)] in Proposition E.54.
- *Rates and constants* ([3, p. 427]). They are $\delta^{(i+1)} = \delta^{(i)}/(2(n'+1))$ and

$$C_A \le c_d^m (4/\delta_1)^{dm} (4m+2)^{dm(m-1)/2} (1 - L^{-\vartheta})^{-m},$$

  independent of η, k, ε, x and ℓ_T (Lemma E.51).

Real backgrounds.

Lemma E.99 (real backgrounds and unitarity)**.** *For $w \in \mathbb{C}$ put $U_w := \mathrm{e}^{\eta q w}$, so that $U(A) = U_{eA}$, and a complex charge ze, $z \in \mathbb{C}$, gives U_{zeA}. Let R_{n+1} be as in Definition E.68(c).*

(a) *$\|U_w\| = \mathrm{e}^{\eta|\Im w|}$. Hence $U_w \in \mathrm{SO}(2)$ if and only if $w \in \mathbb{R}$; $U(A)$ is orthogonal if and only if $A \in \mathbb{R}$; and for real $A \neq 0$ and $z = \mathrm{e}^{i\alpha}$ with $\sin\alpha \neq 0$, U_{zeA} is not orthogonal.*

(b) *For $n \ge 0$ and $w \in \mathbb{R}$, $\|R_{n+1}(qw)\| \le 1$. For every $s > 0$, $\sup_{|\Im w| \le s} \|R_{n+1}(qw)\| \ge 2^{-(n+1)} \mathrm{e}^{s/2}$.*

(c) *The matrix entries of U_w and $R_{n+1}(qw)$ are entire in w with real Taylor coefficients; so $U_{\bar w} = \overline{U_w}$, $R_{n+1}(q\bar w) = \overline{R_{n+1}(qw)}$, and both are real for real w.*

Consequently an estimate proved through $\|U\| = 1$, such as [8, (3.6)–(3.7), p. 255; (3.17), p. 257]*, or through $\|R_{\bar n+1}(qA)\| \le 1$* [1, p. 615], [3, p. 414]*, gives no information at a non-real background or at a complex charge.*

Proof. With S the unitary matrix with columns $(1, \mp i)/\sqrt{2}$, $q = S\,\mathrm{diag}(i, -i)S^{-1}$, so $f(qw) = S\,\mathrm{diag}(f(iw), f(-iw))S^{-1}$ is normal with norm $\max(|f(iw)|, |f(-iw)|)$ for every entire f. (a) $|\mathrm{e}^{\pm i\eta w}| = \mathrm{e}^{\mp\eta\Im w}$; for real w, U_w is the rotation by ηw; for $w = zeA$, $\Im w = eA\sin\alpha$. (b) For real w, $|R_{n+1}(\pm iw)| \le (n+1)\int_0^1 (1-t)^n\,dt = 1$. At $w = is$ the eigenvalue $R_{n+1}(s)$ is real and $R_{n+1}(s) \ge (n+1)\int_{1/2}^1 (1-t)^n \mathrm{e}^{s/2}\,dt = 2^{-(n+1)}\mathrm{e}^{s/2}$. (c) $U_w = \sum_j (\eta q)^j w^j/j!$ and $R_{n+1}(qw) = \sum_j \frac{(n+1)!}{(n+1+j)!} q^j w^j$, since $(n+1)\int_0^1 (1-t)^n t^j\,dt = \frac{(n+1)!\,j!}{(n+1+j)!}$. □

Second proof of Proposition E.71. Part 1 (the printed literature). Each item below is a statement about printed pages.

(B) [4]: vector fields are real-valued, $U(A) = \mathrm{e}^{qe\eta A}$, e real [4, p. 572]; the only complex object is the contour shift of the free $A = 0$ propagator [4, p. 586]. In [1, 2, 3] every background is real, and $R_{\bar n+1}$ is used at real A (Lemma E.99(b)). [3, (1.16), p. 414; (1.32)–(1.33), p. 420] are stated at real backgrounds. At the cited pages of [24], [25] ([24, pp. 653–657, 659], [25, pp. 334, 336]) the only complex objects are sources.

(C) The cumulant lemma [6, p. 152] concerns real Gaussian variables, with no gauge field; it is applied in [2, p. 563, (2.28)–(2.31)] to "a positive operator" with "a probabilistic Gaussian measure". [8, (3.17)] is proved via $\|U^\Phi(g)\| = 1$ for unitary g (Lemma E.99(a)) and applied in [2, p. 591, (3.37)] at the real field A^ε, the variable of the outer real integral.

(D) The two unprinted steps are stated at real backgrounds: [3, p. 414], with $\tilde A$ the real Gaussian fluctuation, and [3, p. 433], with $B = B_0 + B'$ real. Their proofs in this paper use real backgrounds only. The p. 433 split presupposes $\square \subset \Omega$ and $\square \subset \Omega_2$, set inclusions (Proposition 5.8(c)).

Part 2 (the results of this paper). By Lemma E.70, every operator of Definition E.68(a) evaluated in the construction is evaluated at a real field or at $A = 0$, and every Gaussian measure is a real Gaussian measure with real symmetric positive definite precision; by Lemma E.69, the operations of the construction keep backgrounds real, and B_0 in (D) may be taken real. The statements of this appendix used in the proof of Theorem 5.2 concern these objects (proof of Proposition E.71(i)). Bounds at phase-rotated couplings $(ze, z\lambda)$ with $|z| = 1$, where U is not orthogonal (Lemma E.99(a)), are not used: Theorem 5.2 is evaluated at real couplings and at $u \in \{1, -\eta\}$. Part 1 (B), (C) and Part 2 give (i) and (ii); Part 1 (B), (C) give (iii); Part 1 (D) and Lemma E.69(c) give (iv). □

E.6. **The conditions of Bałaban's Proposition 1 at its applications.** We prove (a) three conventions of Bałaban's papers on domains and supports, Lemmas E.100 (nesting of the domains), E.102 (the meaning of θ_b) and E.103 (localisation in Λ_7), and (b) Proposition E.110, which assembles the conditions of [3, Prop. 1] at its applications in [2] from the seventeen statements of Table 10. The support hypothesis in the form (R4‴) is Corollary 5.10. The remaining statements are proved in Subsection E.7.

Setting. We work in the k-th step, $k \geq 1$, of [2, Chap. 2], $d = 3$, on the torus T_ε of [1, (1.2)], at $L = 2$. Distances are those of the torus sup-metric [1, (1.3), p. 604], $|x - y| = \max_\mu \min\{|x_\mu - y_\mu|, 2L_\mu - |x_\mu - y_\mu|\}$, and the distance of two sets is the infimum of point distances ("inf convention"; only the point metric is printed). We write $r_j := r(L^j\varepsilon)$ and M_b for Bałaban's large-block factor M. For $\Lambda \subset T_1^{(j)}$ (the unit lattice of scale j), $\Lambda' := \Lambda \cap L\,T_1^{(j)}$ and $B(y)$ is the block with base point y [1, (1.16)–(1.17), p. 606]. $B^k(y)$ is the scale-k block in T_η, $\eta = L^{-k}$, and $B^k(X) := \bigcup_{y\in X} B^k(y)$. Large blocks are unions of blocks [1, p. 605]. The sets $\Lambda_i^{(j)} \subset T_1^{(j)}$ are defined by [2, (2.7)–(2.8), p. 558] with $r(\varepsilon)$ replaced by r_j [2, p. 566]: $\Lambda_0^{(j)c}$ is a union of large blocks, and $\Lambda_{i+1}^{(j)c}$ is the union of all large blocks whose distance from $\Lambda_i^{(j)c}$ is at most r_j. The function θ_k on T_η equals 1 on $B^{k-1}(\Lambda_2^{(k-1)})$ and varies from 1 to 0 on a slice of thickness $< M_b$ around it [2, pp. 566–567]. In [3], $\Omega = B^k(\Lambda_2^{(k-1)\prime})$ [3, p. 411], $\Omega_1 = B^k(\Lambda_7^{(k-1)\prime})$, and g_k is smooth with $\operatorname{supp} g_k \subset \Omega_1$ [3, p. 412].

On the path of Theorem 5.2 the propagator is changed to the domain $\Omega_2 := B^k(\Lambda_2^{(k)})$ at three places.

- *U1a*, in the polynomial $Q^{(k)}$ [2, p. 579]: $G_k(B^k(\Lambda_0^{(k)}), \tilde{B}) \to G_k(B^k(\Lambda_2^{(k)}), B^{(k+1),\eta})$, where $\tilde{B} = (1 - \theta_{k+1})A^{(k)} + \theta_{k+1}B^{(k+1),\eta}$ on $B^k(\Lambda_0^{(k)})$.
- *U1b*, in the interaction terms [2, p. 580, Prop. 2.6]: $G_k(B^{k-1}(\Lambda_2^{(k-1)}), \tilde{B}) \to G_k(B^k(\Lambda_2^{(k)}), B^{(k+1),\eta})$.
- *U2*, at scale j [2, p. 578].

Nesting of the domains.

Lemma E.100 (nesting of the domains)**.** *Fix $k \geq 1$ and an admissible sequence $(\Lambda_0^{(j)})_j$ of* [2, (2.43)].

(i) *For every j and $i \geq 0$: $\Lambda_{i+1}^{(j)} \subset \Lambda_i^{(j)}$. This holds under every block-to-set distance convention for which a block contained in a set has distance* 0 *from it.*

(ii) *$\Lambda \mapsto \Lambda'$ and $X \mapsto B^k(X)$ are monotone, and $B^k(\Lambda') = B^{k-1}(\Lambda)$ for every $\Lambda \subset T_1^{(k-1)}$ that is a union of blocks.*

(iii) *At U1b: $\Omega_2 = B^k(\Lambda_2^{(k)}) \subset B^k(\Lambda_0^{(k)}) \subset \Omega_1 \subset \Omega$. Moreover $\Omega = B^{k-1}(\Lambda_2^{(k-1)})$, the old domain of U1b.*

(iv) *$B^k(\Lambda_2^{(k)}) \subset B^l(\Lambda_2^{(l)})$ for every $l \leq k$.*

(v) *(Margins, scale-k units, inf convention.) If $x \in \Lambda_{i+s}^{(j)}$ then $\operatorname{dist}(x, \Lambda_i^{(j)c}) \geq s\lfloor r_j \rfloor + 1$. Hence $\operatorname{dist}(\Omega_2, \Omega_1^c) > 2r_k - 2$ and $\operatorname{dist}(\Omega_1, \Omega^c) > (5r_{k-1} - 4)/L - 1$. Under any other block-to-set*

convention (block centre, representative point) each step [2, (2.8)] *loses at most one large-block diameter M_b of its scale, which gives $> 2(r_k - M_b) - 2$ and $> (5(r_{k-1} - M_b) - 4)/L - 1$.*

The constants depend on L and M_b only.

The lemma uses [2, p. 566]. Parts (i)–(iv) do not depend on the convention for the distance between sets. Part (v) is stated under the infimum convention, with the loss of at most M_b per step under any other convention; this loss is absorbed by $r \geq R \gg M_b$ and covered by (5.11).

Proof. (i) By [2, (2.7)], $\Lambda_0^{(j)c}$ is a union of large blocks, and by [2, (2.8)] so is every $\Lambda_{i+1}^{(j)c}$. Suppose $\Lambda_i^{(j)c}$ is a union of large blocks. Each of these blocks has distance $0 \leq r_j$ from $\Lambda_i^{(j)c}$ (under the inf, sup or centre conventions alike), so it belongs to $\Lambda_{i+1}^{(j)c}$. Thus $\Lambda_i^{(j)c} \subset \Lambda_{i+1}^{(j)c}$, and induction on i gives (i).

(ii) Monotonicity is immediate from the definitions. For $y \in L\,T_1^{(k-1)}$ the scale-k block is $B^k(y) = \bigcup_{x \in B(y)} B^{k-1}(x)$ (iterated blocks, [1, (1.17)]). If $\Lambda = \bigcup_{y \in \Lambda'} B(y)$, then $B^k(\Lambda') = \bigcup_{y \in \Lambda'} \bigcup_{x \in B(y)} B^{k-1}(x) = B^{k-1}(\Lambda)$.

(iii) The definition of "admissible" in [2, p. 566] contains the clause "In general $\Lambda_0^{(j+1)} \subset \Lambda_7^{(j)\prime}$, and either $\Lambda_0^{(j+1)}$ is a maximal set composed of big blocks and satisfying this inclusion, or …". This is the general-j analogue of the $j = 0$ condition printed just before it, and it has the same either/or form. It is therefore a condition on every term of the expansion [2, (2.43)], not a hedge: if "in general" meant "usually", the either/or clause of the definition would also be optional. With $j = k - 1$: $\Lambda_0^{(k)} \subset \Lambda_7^{(k-1)\prime}$. Now

$$\Omega_2 = B^k(\Lambda_2^{(k)}) \overset{\text{(i),(ii)}}{\subset} B^k(\Lambda_0^{(k)}) \overset{\text{p. 566, (ii)}}{\subset} B^k(\Lambda_7^{(k-1)\prime}) = \Omega_1 \overset{\text{(i),(ii)}}{\subset} B^k(\Lambda_2^{(k-1)\prime}) = \Omega.$$

Since $\Lambda_2^{(k-1)}$ is a union of large blocks, (ii) gives $\Omega = B^{k-1}(\Lambda_2^{(k-1)})$.

(iv) By the chain in (iii), $B^k(\Lambda_2^{(k)}) \subset B^k(\Lambda_2^{(k-1)\prime}) = B^{k-1}(\Lambda_2^{(k-1)})$. Iterating in k gives (iv).

(v) Let $x \in \Lambda_{i+1}^{(j)}$. The large block containing x is not in $\Lambda_{i+1}^{(j)c}$, so its distance from $\Lambda_i^{(j)c}$ is $> r_j$. Hence no lattice point z with $|z - x| \leq \lfloor r_j \rfloor$ lies in $\Lambda_i^{(j)c}$; that is, the closed sup-ball of radius $\lfloor r_j \rfloor$ about x lies in $\Lambda_i^{(j)}$. On the cyclic lattice, sup-balls of integer radii add coordinatewise. Iterating s times therefore puts the ball of radius $s\lfloor r_j \rfloor$ about $x \in \Lambda_{i+s}^{(j)}$ inside $\Lambda_i^{(j)}$, and so $\operatorname{dist}(x, \Lambda_i^{(j)c}) \geq s\lfloor r_j \rfloor + 1$.

For $x \in B^k(y)$ and $z \in B^k(y')$, $|x - z| > |y - y'| - 1$ in scale-k units. With $y \in \Lambda_2^{(k)}$ and $y' \notin \Lambda_0^{(k)}$ ($s = 2$), and using $\Omega_1 \supset B^k(\Lambda_0^{(k)})$ by (iii), we get $\operatorname{dist}(\Omega_2, \Omega_1^c) \geq 2\lfloor r_k \rfloor > 2r_k - 2$. Likewise, $\Omega_1 = B^{k-1}(\Lambda_7^{(k-1)})$ and $\Omega = B^{k-1}(\Lambda_2^{(k-1)})$ by (ii). With $s = 5$, in scale-$(k-1)$ units their distance is $\geq 5\lfloor r_{k-1} \rfloor$, i.e. $\geq 5\lfloor r_{k-1} \rfloor / L > (5r_{k-1} - 5)/L \geq (5r_{k-1} - 4)/L - 1$ in scale-k units, because $L \geq 1$. Under a non-inf convention the block of x may be declared at distance $\leq r_j$ although its points are up to one large-block diameter M_b farther away. Replacing r_j by $r_j - M_b$ in each step gives the stated bounds. □

Remark E.101 (scope of the order $\Omega_1 \supset \Omega_2$). The order $\Omega \supset \Omega_1 \supset \Omega_2$ holds at U1b. At U1a the new localisation has $\operatorname{supp} g_k \subset \Lambda_7^{(k)}$ [2, p. 579], so there the order is $\Omega = B^k(\Lambda_0^{(k)}) \supset \Omega_2 \supset B^k(\Lambda_7^{(k)})$. At U2 the vertices lie in $\operatorname{supp} \theta_{k+1}$ [2, p. 578]. No use in the proof needs $\Omega_1 \supset \Omega_2$ at U1a or U2; so the step S17 of Table 2 is used only in the form U1b. Lemma E.100 does not need the identification of the Ω_2 of [3] with $B^k(\Lambda_2^{(k)})$: [3, Prop. 1] is uniform over $\Omega_2 \subset \Omega$ "satisfying the same conditions as Ω" [3, p. 414], its constants are independent of $\Omega, \Omega_1, \Omega_2$ [3, p. 421], and $B^k(\Lambda_2^{(k)})$ is a union of big blocks.

The meaning of θ_b.

Lemma E.102 (the meaning of θ_b)**.** *Let $\Omega_2 = B^k(\Lambda_2^{(k)})$, and call a bond of T_η* interior *if both its end points lie in Ω_2.*

(i) *$\theta_{k+1} = 1$ at every point of Ω_2.*
(ii) *For every vector field A and every interior bond b, $(\theta_{k+1}A)_b = A_b$ and $((1-\theta_{k+1})A)_b = 0$. This holds for the printed identification of vector fields with bond functions and for every local convention for the product of a point function with a vector field (one in which the value on b is a function of $\theta(b_-), \theta(b_+)$ equal to 1 when both are 1). In particular $\tilde{B}_b = B_b^{(k+1),\eta}$ on interior bonds.*
(iii) *$G_k(\Omega_2, \tilde{B}) = G_k(\Omega_2, B^{(k+1),\eta})$ and $Q_k(\tilde{B}) = Q_k(B^{(k+1),\eta})$ on Ω_2.*
(iv) *At U1b, the background summands $(1-\theta_{l+1})\theta_l A^{(l)}$, $l+1 \le k+1$, of* [2, (2.93), p. 576] *vanish on interior bonds. The vertex-borne field $B^{(k+1),\eta}$ of* [2, (2.107)] *enters only on interior bonds.*

The lemma uses [1, p. 604] and [2, p. 567], and the standard meaning, not spelled out in [1], that the Neumann form on Ω_2 involves only bonds with both end points in Ω_2.

Proof. (i) This is the definition of θ_k [2, pp. 566–567] with k replaced by $k+1$: $\theta_{k+1} = 1$ on $B^k(\Lambda_2^{(k)})$.

(ii) [1, p. 604] identifies a vector field $A : T_\varepsilon \to \mathbb{R}^d$ with a bond function by $A_{\langle x, x+\varepsilon e_\mu\rangle} = A_\mu(x)$ for positively oriented bonds, with $g_{-b} = -g_b$. The product θA of the point function θ [2, p. 567] with a vector field is printed only as such a product. Through the $\mathbb{R}^d$-valued representation it is $(\theta A)_\mu(x) = \theta(x)A_\mu(x)$, i.e. θ is evaluated at the lower end point of the *positively oriented* bond. On an interior bond both end points lie in Ω_2, where $\theta_{k+1} = 1$ by (i). So $(\theta_{k+1}A)_b = A_b$ under this convention and under every local convention. With the printed $\tilde{B} = (1-\theta_{k+1})A^{(k)} + \theta_{k+1}B^{(k+1),\eta}$ [2, p. 579], this gives $\tilde{B}_b = B_b^{(k+1),\eta}$.

(iii) $G_k(\Omega_2, B)$ is the inverse of the covariant Laplacian on Ω_2 with Neumann boundary conditions [1, p. 609], plus the mass term, plus $a_k Q_k^*(B)Q_k(B)$ [1, (2.20)–(2.22)]. It depends on B only in two ways:

- through the transporters $U(B_b)$ of the bonds of the Neumann form, all of which are interior;
- through $(Q_k(B)f)(y) = L^{-kd}\sum_{x\in B^k(y)} U(B(\Gamma_{y,x}^{(k)}))f(x)$ [1, (2.11)]. The contours $\Gamma_{y,x}^{(k)}$ are coordinate-wise segments inside $B^k(y)$ ([1, (2.1)–(2.2), p. 608]; convention R3 of Lemma E.128), and $B^k(y) \subset \Omega_2$ for $y \in \Lambda_2^{(k)}$, so these bonds are interior too.

By (ii), $\tilde{B}$ and $B^{(k+1),\eta}$ agree on all these bonds. This is the replacement Bałaban prints at U1a and U1b [2, p. 579; p. 580, Prop. 2.6].

(iv) First claim. By Lemma E.100(iv), $\Omega_2 \subset B^l(\Lambda_2^{(l)})$ for $l \le k$, and $\theta_{l+1} = 1$ on $B^l(\Lambda_2^{(l)})$ by (i) at index l. So $(1-\theta_{l+1}) = 0$ at both ends of every interior bond, and the summand vanishes there by (ii).

Second claim. By Lemma E.103(i) the vertices of [2, (2.107)] lie in $B^k(\Lambda_7^{(k)})$. By Lemma E.100(v) with $s = 5$, these points have depth $\ge 5\lfloor r_k\rfloor$ in Ω_2. By Lemma E.103(ii)–(iii), every bond used by a vertex lies within distance $c_1 \le 3$ of it. Since $5\lfloor r_k\rfloor > 3+1$ (as $r_k \ge R \ge 107$), all these bonds are interior, and (ii) applies. □

Localisation in Λ_7.

Lemma E.103 (localisation in Λ_7)**.** *At U1a and U1b:*

(i) *Every vertex of every term to which the propagator change is applied lies in $B^k(\Lambda_7^{(k)})$.*
(ii) *Every localisation cube $\square(v)$ of* [3, p. 420] *has sup-side at most 2 (scale-k units).*

(iii) *The cut-off h of* [3, (1.33)] *can be chosen with $h = 1$ on $\square(v)$ and its η-neighbours,* $\operatorname{supp} h \subset U_1(\square(v))$ *(the closed* 1*-neighbourhood), and*

$$\|h\Phi\|_{1,\alpha_0} \le C_d \, \|\Phi\|_{1,\alpha_0;\, U_1(\square(v))},$$

where $\|\cdot\|_{1,\alpha_0}$ is the norm [3, (1.32)]*, the right side is the same norm with all suprema restricted to points of $U_1(\square(v))$ and their η-neighbours, and C_d depends on d only. The localisation width is therefore $c_1 \le 3$, an absolute constant.*

(iv) *Under $R \ge 3R_0 + 15$ the margin $5r_k - c_2$ at the uses, $c_2 = 5 + c_1 \le 9$, is at least $R_0 + 12$.*

The lemma uses [2, pp. 579–580] and [3, p. 420]; the localisation width is $c_1 \le 3$.

Proof. (i) At U1a, [2, p. 579] prints: "The terms containing the vertices localized in $B^k(\Lambda_7^{(k)c})$ can be estimated by $O((L^k\varepsilon)^{\kappa_0})|\Lambda_2^{(k)} \cap \Lambda_7^{(k)c}| \ldots$ We choose the new localization with the help of a function g_k, … $\operatorname{supp} g_k \subset \Lambda_7^{(k)}$ and $g_k = 1$ on $\Lambda_8^{(k)}$. Now the summations in the vertices are restricted to $\Lambda_7^{(k)}$".

At U1b, [2, p. 579] prints "We estimate the terms localized in $B^k(\Lambda_7^{(k)c})$ by $O((L^k\varepsilon)^{\kappa_0})|\Lambda_2^{(k)} \cap \Lambda_7^{(k)c}|$". [2, Prop. 2.6, p. 580] acts on "the interaction terms localized in $B^k(\Lambda_7^{(k)})$". And [2, p. 580] prints: "If x in $(aL^{-2}C^{(k)}_{\Lambda_4^{(k)}}(\ldots)Q^*(\ldots)\psi)(x)$ is a vertex variable, then $x \in \Lambda_7^{(k)}$".

This covers all vertices, including the vertices with $n = 0$, $n' \ge 1$ that come from the $\tilde{A}$-expansion of $G_k(\Omega, \tilde{A} + \tilde{B})$ [3, p. 414]. Those vertices are not summands of the interaction [3, (1.4)], so an argument based on the local form of (1.4) alone would not cover them.

(ii) [3, p. 420] prints: "For the vertices (1.6) and (1.7) we localize simply by representing Ω_1 as a sum of unit cubes of the η-lattice. For the remaining vertices we localize, taking for each leg of the vector field a smooth partition of unity satisfying the condition that a support of each function is contained in a cube with sides of length 2. … The expressions (1.12)–(1.15) are localized by fixing a point $y \in T_1^{(k)} \cap \Omega$, or the unit cube $B^k(y)$." Hence every $\square(v)$ has side ≤ 2.

(iii) The only unprinted width is that of h ("$h = 1$ on $\square(v)$ and $h = 0$ outside some neighborhood of $\square(v)$", [3, p. 420]). It is a choice of whoever applies [3, Prop. 1], for two reasons. First, the vertex depends on Φ_{ext} only on $\square(v)$ and its η-neighbours, so the expression E of [3, (1.33)] satisfies $E(\Phi_{\text{ext}}) = E(h\Phi_{\text{ext}})$ for any h equal to 1 there. Second, the constant of [3, (1.33)] is independent of ε, k and the domains [3, p. 421], and h enters only through $\|h\Phi_{\text{ext}}\|_{1,\alpha_0}$, $\|h'A_{\text{ext}}\|_{1,\alpha_0}$.

We now construct h. Let $s(t) = 10t^3 - 15t^4 + 6t^5$ on $[0,1]$; then $s(0) = 0$, $s(1) = 1$, $s' = s'' = 0$ at both ends, $\max s' = 15/8$ and $\max|s''| = 10/\sqrt{3}$. Put $\chi(t) = 1$ for $t \le \frac{1}{2}$, $\chi(t) = 1 - s(2t-1)$ for $\frac{1}{2} \le t \le 1$, and $\chi(t) = 0$ for $t \ge 1$. Then $\chi \in C^2$, $|\chi'| \le 15/4$ and $|\chi''| \le 40/\sqrt{3} < 32$. Set $h(x) = \prod_\mu \chi(t_\mu(x))$, where $t_\mu(x)$ is the distance of x_μ from the μ-th side interval of $\square(v)$. Since $\eta = L^{-k} \le \frac{1}{2}$, $h = 1$ on $\square(v)$ and its η-neighbours, and $\operatorname{supp} h \subset U_1(\square(v))$, a cube of side ≤ 4. The lattice first and second differences of h are bounded by constants depending on d only, uniformly in η.

Write D_μ for the covariant difference of [3, (1.32)], with unimodular transporter $U_\mu(x)$ on the bond $\langle x, x + \eta e_\mu\rangle$. Then exactly

$$D_\mu(hf)(x) = h(x + \eta e_\mu)\,(D_\mu f)(x) + (\partial_\mu^\eta h)(x)\, f(x).$$

This gives the sup and first-difference parts of the norm with a constant depending on d only.

For the Hölder quotient, put $F := D_\mu(hf)$ and $U_\Gamma := U(\tilde{B}(\Gamma_{x,x'}))$. Then

$$\begin{aligned} U_\Gamma F(x') - F(x) = {} & h(x' + \eta e_\mu)\big(U_\Gamma D_\mu f(x') - D_\mu f(x)\big) + \big(h(x' + \eta e_\mu) - h(x + \eta e_\mu)\big) D_\mu f(x) \\ & + (\partial_\mu^\eta h)(x')\big(U_\Gamma f(x') - f(x)\big) + \big((\partial_\mu^\eta h)(x') - (\partial_\mu^\eta h)(x)\big) f(x). \end{aligned}$$

Here $|U_\Gamma f(x')-f(x)| \le d|x-x'|\sup|Df|$, by telescoping along the shortest contour. The differences of h and $\partial^\eta h$ are $\le C_d|x-x'|$. For $|x-x'|\le 1$ this gives the bound $C_d|x-x'|^{1-\alpha_0}\|f\|_{1,\alpha_0} \le C_d\|f\|_{1,\alpha_0}$ after division by $|x-x'|^{\alpha_0}$; for $|x-x'|>1$, bound each term by twice a supremum.

If x lies outside $S := \{x : x \text{ or } x+\eta e_\mu \in \operatorname{supp} h\}$, then $F(x)=0$ and the restricted norm sees only $F(x')$, $x'\in S$. Because h and $\partial^\eta h$ vanish to first order at the edge of $\operatorname{supp} h$ ($\chi'=0$ at $t=1$), $|F(x')| \le C_d|x-x'|\sup_{U_1(\square(v))}(|f|+|Df|)$, which is again bounded after division by $|x-x'|^{\alpha_0}$. A merely linear ramp would not do: its difference jumps across one η-bond, and the quotient is then of order $\eta^{-\alpha_0}$.

So $c_1 \le$ (side of $\square(v)$) + (collar) $\le 2+1=3$.

(iv) $5r_k - 9 \ge 5R-9 \ge R_0+12$, because $5R \ge 15R_0+75 \ge R_0+21$. □

Assembly.

The applications of Proposition 1 of [3]. Throughout this paragraph $L=2$, $\bar n = 13$ and $\bar N = 169$, and the regulator data are chosen in the order of Appendix A. In particular $R \ge R^{**}$, so that

$$R \ge 107, \qquad R \ge 3R_0+15, \qquad R \ge R_0+13+c_1',$$
$$R \ge 2\ell_{\bar N}+10M_b+10, \qquad R \ge 4\ell_{\bar N}+32M_b+25, \qquad \delta_0 R \ge 4.$$

Also $\varepsilon_0 < \mathrm{e}^{-1}$, the ε_0-bounds of Lemma A.17 hold, $\ell_0 \ge \varepsilon_0[8r(\varepsilon_0)+4M_b+2]$ (condition (T.nw)) and $\ell_T/(M_b t_k) \in 2\mathbb{N}$. Since $t \le \varepsilon_0 < \mathrm{e}^{-1}$, we have $1+\log t^{-1} > 2$, and hence

$$r(t) = R(1+\log t^{-1})^2 > 4R. \tag{E.7}$$

The notation is that of the setting paragraph of Subsection E.6 and of Subsection E.7. The constant c_1' is the localisation width at scale j of Lemma E.126.

Definition E.104 (the applications of Proposition 1 at step k)**.** Fix $k \ge 1$. In [2, 3], [3, Prop. 1] is applied, or the domain of a scalar propagator is changed in the way its proof requires, in the situations listed below. For each one we name the *outer domain* Ω_{out} and the *new domain*.

- (U0) [3, pp. 411–414, 432–433] itself, and its application at [2, p. 582, (2.116)]: $\Omega_{\mathrm{out}} = \Omega = B^k(\Lambda_2^{(k-1)\prime})$; the new domain is Ω_2.
- (U1a) [2, p. 579]: $\Omega_{\mathrm{out}} = B^k(\Lambda_0^{(k)})$; the new domain is $\Omega_2 = B^k(\Lambda_2^{(k)})$; the background is $\tilde B = (1-\theta_{k+1})A^{(k)} + \theta_{k+1}B^{(k+1),\eta}$.
- (U1b) [2, p. 580, Prop. 2.6]: $\Omega_{\mathrm{out}} = B^{k-1}(\Lambda_2^{(k-1)})$, which equals Ω by Lemma E.100(iii); the new domain is Ω_2.
- (U2) [2, p. 578], at a scale $0 \le j \le k-1$, on the lattice $T_{L^{-j}}$ of spacing $\eta_j = L^{-j}$: $\Omega_{\mathrm{out}} = B^j(\Lambda_2^{(j)})$; the new domain is $B^k(\Lambda_0^{(k)})$. The propagator $G_j(B^j(\Lambda_2^{(j)}), \tilde B)$ is replaced by $G_j(B^k(\Lambda_0^{(k)}), \tilde B)$.
- (U3) Inside the proof of [3, Prop. 1], at any of the above: the substitution $G_k(\square,0) = G_k(0) + \delta G_k(\square, \eta\mathbb{Z}^d, 0)$ of [3, p. 433].

An *evaluation point* of a kernel at an application is any of the following: a line end, a bond partner, a block point of Q_k, a Hölder partner, or a point of the support of a localisation function h, h' of [3, (1.33)].

Lemma E.105 (one background at the change to Ω_2)**.** *At (U1a) and (U1b), let $\tilde B$ be the background and Ω_{out} the outer domain of the application. Then $\Omega_2 \subset \Omega_{\mathrm{out}}$, both sets are unions of big blocks, and, as kernels on $\Omega_2 \times \Omega_2$,*

$$G_k(\Omega_{\mathrm{out}}, \tilde B) - G_k(\Omega_2, B^{(k+1),\eta}) = G_k(\Omega_{\mathrm{out}}, \tilde B) - G_k(\Omega_2, \tilde B) =: \delta G_k(\Omega_{\mathrm{out}}, \Omega_2, \tilde B).$$

At (U2) the printed replacement is $\delta G_j(B^j(\Lambda_2^{(j)}), B^k(\Lambda_0^{(k)}), \tilde B)$. It is again taken at the single background $\tilde B$, and $B^k(\Lambda_0^{(k)}) \subset B^j(\Lambda_2^{(j)})$. In each case the background is regular in the sense of hypothesis (R) of Subsection E.7.

Proof. Nesting. At (U1a), $\Omega_2 = B^k(\Lambda_2^{(k)}) \subset B^k(\Lambda_0^{(k)})$ by Lemma E.100(i),(ii). At (U1b), $\Omega_2 \subset \Omega$ by Lemma E.100(iii). At (U2), Lemma E.100(i)–(iii) give $B^k(\Lambda_0^{(k)}) \subset B^k(\Lambda_7^{(k-1)\prime}) = B^{k-1}(\Lambda_7^{(k-1)}) \subset B^{k-1}(\Lambda_2^{(k-1)})$. The argument of Lemma E.100(iv) then gives $B^{k-1}(\Lambda_2^{(k-1)}) \subset B^j(\Lambda_2^{(j)})$ for $j \le k-1$. All these sets are unions of big blocks [2, p. 566], and B^l of a union of big blocks is again a union of big blocks.

Identity. By Lemma E.102(ii)–(iv), on every bond with both end points in Ω_2 the background $\tilde{B}$ of (U1a) equals $B^{(k+1),\eta}$. The same holds for the background of (U1b): all its summands other than $\theta_{k+1}B^{(k+1),\eta}$ vanish there. By Lemma E.102(iii), $G_k(\Omega_2, \cdot)$ depends on its background only through such bonds. Hence $G_k(\Omega_2, \tilde{B}) = G_k(\Omega_2, B^{(k+1),\eta})$, which is the displayed identity. At (U2) the propagator keeps its background [2, p. 578].

Regularity. At (U1a) the regularity of the background is [2, (2.98), p. 579]. The backgrounds of (U1b) and (U2) are configurations of the form [2, (2.93), p. 576], and their regularity is [2, (2.94)–(2.97), p. 576]. The smallness required in (R) is given by the ε_0-bounds of Lemma A.17. The Hölder part for the $\tilde{A}$-legs is Proposition E.138. □

Lemma E.106 (two-domain bound at lattice spacing 1)**.** *Let $\eta = 1$. Let $X_{\mathrm{in}} \subset X_{\mathrm{out}}$ be unions of blocks of T_1 (or let $X_{\mathrm{out}} = T_1$), and let A be a regular vector field in the sense of* [4, p. 573]*. Put $\delta := G(X_{\mathrm{out}}, A) - G(X_{\mathrm{in}}, A)$, where G denotes the Green's functions of* [4, (1.6)] *with Neumann conditions, and let c_0, δ_0, R_0 be the constants of the theorem on* [4, p. 573]*. Let x, x' be points such that x, x' and their lattice neighbours have distance $\ge R_0$ from X_{in}^c. Write D for the covariant difference and D^* for its adjoint. Then*

$$\max\big\{|\delta(x,x')|, |D_x\delta(x,x')|, |(\delta D^*)(x,x')|, |(D_x\delta D^*)(x,x')|\big\} \le 2c_0 \mathrm{e}^{\delta_0}\mathrm{e}^{-\delta_0|x-x'|}.$$

Every α-Hölder quotient of these kernels over $0 < |x-\tilde{x}| \le 1$ is at most twice the supremum of the kernel over the points involved if it is taken in one variable, and at most four times that supremum if it is taken jointly. Consequently, if $0 \le h, h' \le 1$ and every point of $\operatorname{supp} h \cup \operatorname{supp} h'$ has distance $\ge R_0 + 2$ from X_{in}^c, then

$$\|h\,\delta\,h'\|_{1,\alpha} \le C(c_0,\delta_0)\,\mathrm{e}^{-\delta_0 \operatorname{dist}(\operatorname{supp} h, \operatorname{supp} h')},$$

with the norm of [3, (1.32)] *in both variables.*

Proof. At $\eta = 1$ kernels are taken with respect to counting measure. Fix y, and let f be the point mass at y with values in a unit vector. Then $\|f\|_\infty = 1$, and $(\delta f)(x)$ is the column $\delta(x,y)$. We apply [4, (1.10)–(1.12), p. 573] with $\Omega = X_{\mathrm{in}} \subset \Omega_0 = X_{\mathrm{out}}$; the additional factor (1.12) is at most 1. This gives, for every y,

$$|\delta(x,y)| + |D_x\delta(x,y)| \le c_0\mathrm{e}^{-\delta_0|x-y|} \qquad \text{whenever } \operatorname{dist}(x, X_{\mathrm{in}}^c) \ge R_0.$$

If $y \notin X_{\mathrm{in}}$, then $\delta(\cdot,y) = G(X_{\mathrm{out}}, A)(\cdot,y)$, and (1.10) for X_{out} applies, since $\operatorname{dist}(x, X_{\mathrm{out}}^c) \ge \operatorname{dist}(x, X_{\mathrm{in}}^c)$.

At $\eta = 1$, $(D_\mu^* g)(x')$ is a combination of $g(x')$ and $g(x' - e_\mu)$ with coefficients of modulus 1. Hence, for $K = \delta$ or $K = D_x\delta$,

$$|(KD^*)(x,x')| \le |K(x,x')| + |K(x,x'-e_\mu)| \le 2c_0\mathrm{e}^{\delta_0}\mathrm{e}^{-\delta_0|x-x'|}.$$

This gives the first display.

Distinct lattice points with $|x-\tilde{x}| \le 1$ satisfy $|x-\tilde{x}| = 1$. So each Hölder quotient is at most $|UF(\tilde{x})| + |F(x)|$: a sum of two kernel values (four, for the joint quotient) at points that again satisfy the depth hypothesis.

Finally, $D_\mu(hF)(x) = h(x+e_\mu)D_\mu F(x) + (\partial_\mu h)(x)F(x)$ with $|\partial_\mu h| \le 1$ at $\eta = 1$, and similarly for h'. Every point at which a kernel is then evaluated lies within distance 2 of $\operatorname{supp} h \cup \operatorname{supp} h'$, and so has depth $\ge R_0$. For such points, $\mathrm{e}^{-\delta_0|x-x'|} \le \mathrm{e}^{2\delta_0}\mathrm{e}^{-\delta_0 \operatorname{dist}(\operatorname{supp} h, \operatorname{supp} h')}$. Summing the finitely many terms gives the last bound. □

Lemma E.107 (localisation after the B'-expansion)**.** *At* [3, p. 433]*, let the localisation cubes of a connected component lie in a cube Q_0 of side $\ell_{\bar N}$. Let $\square_1$ be a union of big blocks of size $r = r(L^k\varepsilon)$ that contains the cube of side r concentric with Q_0. Suppose that the expansion in B' of each scalar line* [1, (1.3.14), (3.44), (3.45)] *produces chains with at most $n_{B'}$ new vertices. Suppose also that, as printed, every line and every operator* [3, (1.16)] *whose end localisations satisfy* $\operatorname{dist}(\square(v),\square(v')) \geq 1$ *is put into the external fields. If*

$$r \geq \ell_{\bar N} + 6n_{B'}, \tag{E.8}$$

then every vertex of every class that contains an old vertex is localised in $\square_1$. Under the standing hypotheses, (E.8) *holds for $n_{B'} = \bar n = 13$.*

Proof. New vertices arise only inside the chains. Each chain replaces one line between two vertices of the expression before the expansion, and a new vertex is joined by lines only to its two neighbours in its chain. Let w be a new vertex of a class that contains an old vertex. The class is connected and w is joined to it only through its chain. Hence the class contains an uncut segment of the chain, of at most $n_{B'}$ lines, from w to one of the two old end vertices.

Every line of the class joins localisation cubes at distance < 1, and every cube has side ≤ 2 (Lemma E.103(ii)). In the sup metric, $\operatorname{dist}(Q,Q') < 1$ and $\operatorname{side}(Q') \leq 2$ give $Q' \subset U_3(Q)$. By induction along the segment, $\square(w) \subset U_{3n_{B'}}(Q_0)$. This is the cube of side $\ell_{\bar N} + 6n_{B'}$ concentric with Q_0; it does not wrap the torus, since $\square$ does not (Lemma E.124). Under (E.8) this cube lies in the concentric cube of side r, and hence in $\square_1$.

For the last claim: by (E.7), $R \geq 2\ell_{\bar N} + 10M_b + 10$ and $R \geq 107$, we get $r > 2R + 2R \geq \ell_{\bar N} + 214 > \ell_{\bar N} + 78$. □

Lemma E.108 (depth supplied at the applications)**.** *Let $k \geq 1$.*

(a) *At (U1a) and (U1b), every evaluation point of $G_k(\Omega_2,\cdot)$ and of $\delta G_k(\Omega_{\mathrm{out}},\Omega_2,\cdot)$ has distance $\geq 5r_k - 9 \geq R_0 + 13$ from Ω_2^c.*

(b) *At (U2), at scale j, every evaluation point has distance $> L^{k-j}r_k - c_1' \geq R_0 + 13$ from the complement of $B^k(\Lambda_0^{(k)})$, in scale-j units.*

(c) *For a class with an old vertex, at any application, every point of the support of a localisation function of a vertex of the class has distance $\geq r - 1 \geq R_0 + 13$ from $\square^c$, and hence from Ω_{out}^c.*

(d) *Consequently, the support condition $R_0 + \frac{45}{4} + 3\eta \leq R_0 + 13$ (for $\eta \leq \frac12$) of the corollary of Proposition D.17 holds in (a), in (c), and in (b) for $j \geq 1$. The condition $R_0 + 2$ of Lemma E.106 holds in (b) for $j = 0$.*

Proof. (a) Lemma E.126 gives the lower bound $5r_k - 5 - c_1$ with $c_1 \leq 4$. Moreover $5r_k - 9 \geq 5R - 9 \geq 15R_0 + 66$.

(b) Lemma E.126 gives $> L^{k-j}r_k - c_1' \geq 2R - c_1'$, and $R \geq R_0 + 13 + c_1'$ gives $2R - c_1' \geq R + R_0 + 13$.

(c) By Lemma E.107 the localisation cubes lie in $\square_1$. The localisation functions are supported in $U_1(\square(v))$ (Lemma E.103(iii)). Every point of $\square_1$ has distance $\geq r$ from $\square^c$ (Proposition 5.8(c)). And $r - 1 > 4R - 1 \geq 12R_0 + 59$ by (E.7). Moreover $\square \subset \Omega_{\mathrm{out}}$ (Lemma E.124).

(d) For $\eta \leq \frac12$ we have $R_0 + \frac{45}{4} + 3\eta \leq R_0 + \frac{51}{4}$. At $j = 0$ the bound in (b) exceeds $R_0 + 2$. □

Lemma E.109 (lowering the tree rate)**.** *If a family satisfies* [3, (1.33)] *with tree rate $\delta > 0$, then it satisfies it with the same constant and with every tree rate $\delta' \in (0,\delta]$.*

Proof. In [3, (1.33), p. 420] the rate enters only through the factor $\exp[-\delta\, d(\{\square(v)\})]$, where $d \geq 0$ is the length of a shortest tree. This factor does not decrease when δ decreases. □

Proposition E.110 (the conditions of Proposition 1 as used in [2])**.** *Assume the hypotheses of Theorem 5.2 and the standing hypotheses of this paragraph. Let $k \geq 1$, and consider any application (U0), (U1a), (U1b) or (U2) of Definition E.104. Then the following hold.*

(i) (Hypotheses.) *The background is regular in the sense of (R). The support hypothesis holds in the form (R4‴) of* (5.10)*, and* [3, Prop. 1] *holds with (R4) replaced by (R4‴).*

(ii) (Remainder kernels.) *The operator* [3, (1.16)] *satisfies the two-variable bound of Lemma E.117(b),(c) at* $\alpha = \alpha_0 = \frac{1}{2}$ *on* T_η*. It satisfies the bound of Lemma E.118 at every evaluation point on the Neumann domain* Ω*.*

(iii) (Change of domain.) *At (U1a), (U1b) and (U2), the printed replacement is the subtraction of one kernel* δG*, at one regular background, for nested domains (Lemma E.105). Moreover* $\|h\,\delta G\,h'\|_{1,\alpha_0} \le C\,\mathrm{e}^{-\beta\,\mathrm{dist}(\mathrm{supp}\,h,\mathrm{supp}\,h')}$ *for all localisation functions* h, h' *of the application.* [2, Chap. 3] *contains no further change of domain of a scalar propagator (Lemma E.127).*

(iv) (The cube step of [3, p. 433].) *Let a class have an old vertex. Then:*

- $\square \subset \Omega_{\mathrm{out}}$*;*
- *after the* B'*-expansion, every vertex of the class is localised in* $\square_1$*;*
- *the kernels* $\delta G_k(\Omega_{\mathrm{out}}, \square, \tilde{B})$*,* $\delta G_k(\Omega, \square, \tilde{B}) - \delta G_k(\Omega, \Omega_2, \tilde{B})$ *and* $\delta G_k(\square, \eta\mathbb{Z}^d, 0)$ *are external fields whose* $(1, \alpha_0)$*-norm at the localisations of the class is at most* $C\,\mathrm{e}^{-\beta\,\mathrm{dist}(\mathrm{supp}\,h,\mathrm{supp}\,h')}$*;*
- *the* B'*-remainder satisfies the bound of Lemma E.121;*
- *the inclusion* $\square \subset \Omega_2$ *is not used.*

(v) (Mass entries.) *The conclusions of Lemma E.119 hold, and the* $A \cdot A$ *vertices of* [3, (3.30), (3.32)] *satisfy (N1)–(N4) of Lemma E.130.*

(vi) (The tails of [2, p. 579].) *The decomposition and the bounds of Lemma E.141 hold.*

Here $\beta := \beta_P := \min\{\delta(\bar{N}), \beta_U, \beta_D, \delta_0\}$*. In this definition,* $\delta(\bar{N})$ *is the rate of Proposition E.60,* β_U *and* β_D *are the rates of Lemmas E.117 and E.123, and* δ_0 *is the rate of* [4, p. 573].

Consequently, [3, (1.33)] *holds at each application, with tree rate* β_P*, for every localised family of order at most* $\bar{N}$*. All constants depend only on* d*,* a*,* L*,* M_b*,* $\bar{N}$*,* α_0*, the constants of* [1, 4] *and the constants of Lemma D.7. They do not depend on* ε*,* k*,* ℓ_T*, on the domains* $\Omega, \Omega_1, \Omega_2$*, or on the background within (R). They depend on the scalar mass only through the condition* $0 \le m^2t^2 \le c$*.*

Proof. (i). At (U1a), (U1b) and (U2), regularity is the last part of Lemma E.105. At (U0) it is the hypothesis of [3, p. 412], which is verified in the same way from [2, (2.94)–(2.98)]. By Corollary 5.10 and (5.11), the form (R4‴) holds at every application, and [3, Prop. 1] holds under it.

(ii). On T_η this is Lemma E.117(b),(c) with $m = n + n' \ge 6$; this covers $\alpha_0 = \frac{1}{2}$. On Ω it is Lemma E.118. The depth that lemma needs at the vertex points, $R_0 + 2\sqrt{3} + 1$, is supplied under (R4‴) by part (3) of its proof.

(iii). By Lemma E.105, the replacement is one kernel δG, for a nested pair of unions of big blocks, at one regular background. For $\eta \le \frac{1}{2}$ the corollary of Proposition D.17 applies, with rate β_D, by Lemma E.108(a),(b),(d). At (U2) with $j = 0$, Lemma E.106 applies, with rate δ_0, by Lemma E.108(b),(d). Lemma E.126 shows that (U1a), (U1b) and (U2) are all the changes of domain of a scalar propagator in [2, Chap. 2]. Lemma E.127 shows that [2, Chap. 3] adds none.

(iv). The inclusion $\square \subset \Omega_{\mathrm{out}}$ is Lemma E.124. At (U0), its hypothesis on the support of $\tilde{B}^{(k)}$ holds for the following reason. $\tilde{B}^{(k)}$ is defined as the sum of the second, third and fourth terms of [2, (2.89), p. 575], and each of these terms has θ_{k+1} as its left factor; so $\mathrm{supp}\,\tilde{B}^{(k)} \subset \mathrm{supp}\,\theta_{k+1}$. The localisation in $\square_1$ is Lemma E.107. The bound on $\delta G_k(\Omega_{\mathrm{out}}, \square, \tilde{B})$ is the corollary of Proposition D.17 for the pair $(\Omega_{\mathrm{out}}, \square)$; it applies by Lemma E.108(c),(d). The cube step relative to Ω_2 is Lemma E.125. Its inputs are the two-domain bounds for $(\Omega, \square)$ and (Ω, Ω_2) just obtained, Lemma E.124, and Lemmas E.126 and E.127; it never uses $\square \subset \Omega_2$. The bound on $\delta G_k(\square, \eta\mathbb{Z}^d, 0)$ at points of $\square_1$ is step S19 in the proof of Proposition 5.8. The B'-remainder is Lemma E.121, with Lemma E.120.

(v), (vi). These are Lemmas E.119, E.130 and E.141.

The bound (1.33). The proof of [3, Prop. 1] on [3, pp. 422–445] consists of printed steps and of the sketched steps listed in Table 2. By Proposition 5.8, these sketched steps are supplied as follows:

- S1 by (ii), and S2 by (iii);
- P12 and S13–S19 by (iv);
- S3, S4 and S9 by Proposition 5.8(b), whose depth requirements are met by Lemma E.108;
- the remaining steps need no depth (Lemma 5.9).

With the order cap, Proposition E.60 gives (1.33) with rate $\delta(\bar N)$ for families of order at most $\bar N$. The kernel bounds used along the way carry the rates β_U, β_D and δ_0. By Lemma E.109 all of them hold with rate β_P, so (1.33) holds with tree rate β_P. The quantity β_P depends only on constants that are fixed before R is chosen (Appendix A), so the order of choices has no cycle (Lemma A.4).

Order of use. The derivation of (1.33) uses only (i)–(iv) and Lemma E.119. Lemmas E.130 and E.141 use (1.33) and the cube step (iv), and they are not used in the proofs of (i)–(iv). Hence there is no circularity.

Uniformity. Each cited statement carries the stated dependence of its constants. The kernels in (ii)–(iv) are scalar Green's functions and do not involve the photon covariance. The uniformity of (v) and (vi) at the two points (u, M) of Theorem 5.2 is part of Lemmas E.130 and E.141. □

Remark E.111 (completeness of the list). The list of applications in Definition E.104 is complete by Lemma E.129. The sketched steps of the proof of [3, Prop. 1] are those of Proposition 5.8(a). Lemma E.114 lists the abbreviated steps of [2, pp. 555–594] and [3, pp. 411–445] that need more than the printed text. That the remaining abbreviated steps on these pages hold under the conditions of Appendix A is Statement K.28.

Remark E.112 (scope). (i) Every clause of Proposition E.110 is proved in this paper, with one exception. In the proof of Lemma E.130, the uniformity of the constant of [3, Prop. 1] under a constant shift of the background is Statement K.29. (ii) Lemma E.107 is stated for a general order $n_{B'}$ of the B'-expansion; any $n_{B'} \le (r - \ell_{\bar N})/6$ suffices. The value $n_{B'} = \bar n$ is the order to which [3, p. 433] refers through [1, (3.45)]. (iii) Corollary 5.10 is proved at $L = 2$, and hence so is Proposition E.110.

E.7. **The statements of Table 10.** We prove the statements of Table 10 other than 13 and 14 (Lemma E.100 and Corollary 5.10). Notation is that of the setting paragraph of Subsection E.6, together with the following:

- $t = t_k = L^k\varepsilon$, $r = r(t) = R(1 + \log t^{-1})^2$, $e_k = e(t)$, $\eta = L^{-k}$, and $|\cdot|$ is the ℓ^∞ torus distance.
- $G(\mathcal{B}) = G_k(\mathcal{X}, \mathcal{B}) = (-\Delta^{\eta,N}_{\mathcal{B},\mathcal{X}} + m^2t^2 + a_kP_k(\mathcal{B}))^{-1}$ on a block union $\mathcal{X}$, with Neumann conditions on $\partial\mathcal{X}$.
- $D = D^\eta_{\tilde B}$, D^* its $\ell^2(\eta^3)$-adjoint, and $\mathfrak{D}f := (f, D_1f, D_2f, D_3f)$.
- $\sigma := e_k(\|\tilde A\|_\infty + \|\partial^\eta\tilde A\|_\infty)$ and $\theta(\mathcal{B}) := e_k\|\partial^\eta\mathcal{B}\|_\infty$.

Kernels are taken with respect to η^3.

Inputs used throughout.

(H0) $-\Delta^{\eta,N}_{A,\mathcal{X}} + a_kP_k(A) \ge \gamma_0$ on every block union $\mathcal{X}$, for every b.c. and every regular A [4, (1.8), p. 573].

(W) For $|\beta| \le \beta_*$ and every 1-Lipschitz ρ: $\|e^{\beta\rho}Ge^{-\beta\rho}\| \le 2/\gamma_0$, and the weighted norms of ∇G, $G\nabla^*$ and $\nabla G\nabla^*$ are $\le K_W$. This holds uniformly in η, the domain and the b.c. (Lemma D.24; literature form [4, Cor. 2.3, (2.30), p. 580]).

(K) The pointwise bounds $\|G(x,y)\| \le c_KK_2^{\delta_K}(x-y)$ and $\|D_xG(x,y)\| \le c_KK_1^{\delta_K}(x-y)$, and for $|x - \tilde x| \le 1$ the α-Hölder bound of D_xG with $K^{\delta_K}_{1-\alpha}$. Here $K^\delta_\kappa(z) := \max(|z|,\eta)^{\kappa-3}e^{-\delta|z|}$ for $0 < \kappa < 3$. These hold at points at distance $\ge R_0$ from $\mathcal{X}^c$ (on T_η and on rectangular

parallelepipeds at all points): [3, (2.10)–(2.11), p. 426] summed via [3, (2.6)], from [1, Props. 2.1, 2.3] (Lemma D.26).

The hypothesis (R_ϑ) is: $\theta(\tilde{B}), \theta(\tilde{A}+\tilde{B}) \le \vartheta_R$ and $\sigma \le 1$; it follows from (E.1) and (E.ksp3) of Appendix A.
The statement Σ and the premises (statements 5, 5′).

Definition E.113 (the statement Σ)**.** Let $\mathcal{R}$ be the set of conditions of Tables 4–8 other than (E.md), (E.ksp1)–(E.ksp3) and (E.14), and let $\mathcal{R}^+$ be $\mathcal{R}$ together with (E.md), (E.ksp1)–(E.ksp3) and (E.14). Σ is the statement that every condition *invoked* by the renormalisation step of [2] and [3] (including the pages of [1] they call) is a condition of $\mathcal{R}^+$ or is not a condition (a definition, a notation or a forward reference). Here each abbreviated step is replaced by its written argument. Σ includes the statement that there is no other mass entry (Lemma E.119).

Lemma E.114 (the abbreviated steps that need more than the printed text)**.** *Assume the hypotheses of Theorem 5.2, and let the regulator data satisfy the conditions of Definition A.1. Each step (a)–(i) below of* [2, pp. 555–594] *and* [3, pp. 411–445] *is stated there without a complete argument. Each of them holds. Its argument is written at the place named, and that argument uses, besides printed statements of* [1, 2, 3, 4, 24]*, only the conditions named.*

(a) [2, p. 573, (2.75)–(2.76)] *("so the same conclusion holds"): the scalar analogue on* $\square$ *of* [2, (2.62)–(2.63)]*, including the mass defect left by (2.75). Place: Lemma E.150. Condition: (E.md).*
(b) [2, Lemma 2.4] *at the edge points not covered by its printed proof (Remark 5.5). Place: Lemma 5.6. Conditions: (R.edge) and (E.edge) (Lemma A.3).*
(c) [2, p. 579]*, the extension of* [2, Prop. 2.1] *to* $Q^{(k)}$*. Place: Lemma A.17. Conditions: the* ε_0*-bounds of that lemma.*
(d) [2, p. 579]*, the cut of the expansion at the order* $\bar{n}$*. Place: Lemma E.141, with Propositions E.36, E.145 and E.146.*
(e) [3, p. 421]*, the use of* [2, Lemma 2.4]*. Places: as in (a) and (b). Condition: (E.md).*
(f) [3, (1.16), p. 414]*, the two-variable remainder bound. Places: Lemmas E.117 and E.118.*
(g) [3, p. 433]*, the remainder on the cube* $\square$*. Place: Lemma E.121, with Lemmas E.124 and E.125.*
(h) *The sketched steps S1–S21 and P12 of the proof of* [3, Prop. 1]*. Places: Table 2 and Proposition 5.8.*
(i) [3, (3.30), (3.32), p. 442]*, the* $A \cdot A$ *local terms. Place: Lemma E.130.*

The conditions (E.md) and those of Lemma A.17 are not printed in [1, 2, 3, 4]*. They are upper bounds on* ε_0 *in Definition A.1.*

In particular, the premise that no abbreviated step uses an unprinted condition is false as stated: the argument for (c) and (d) needs the unprinted conditions above.

Proof. Each item is the statement proved at the place named, and the conditions used in that proof are the ones named in the item. $\square$

That every other abbreviated step on these pages, and on the pages of [1] invoked there, holds under the conditions of Definition A.1 is Statement K.28.
The remainder kernel (1.16): statements 1–3. For $n, n' \ge 0$ and $m := n + n'$, put $V := G(\tilde{B})^{-1} - G(\tilde{A}+\tilde{B})^{-1}$. This is the bracket of [1, (3.44)], and the $m^2 t^2$ terms cancel in it. Put

$$K = K_{n,n'} := [G(\tilde{B})V]^n G(\tilde{A}+\tilde{B})[VG(\tilde{B})]^{n'} = L_0 V L_1 V \cdots V L_m,$$

where $L_n = G(\tilde{A}+\tilde{B})$ and $L_i = G(\tilde{B})$ for $i \ne n$. This is the operator [3, (1.16), p. 414].

Lemma E.115 (vertex structure)**.** *Put* $\phi_\mu(x) := \eta^{-1}(1 - U_{x,\mu}(\tilde{A})^{-1})$*,* $\psi_\mu := \phi_\mu^* \phi_\mu$*,* $Q := Q_k(\tilde{B})$*,* $F_2 := F_{2,k}(\tilde{A}, \tilde{B})$ *and* $\Pi := Q^* F_2 + F_2^* Q + F_2^* F_2$*. Then:*

(i) $V = -\mathfrak{D}^*\mathfrak{E}\mathfrak{D}$, *where* $\mathfrak{E}$ *has first row* $(\sum_\mu \psi_\mu + a_k\Pi, \phi_1^*, \phi_2^*, \phi_3^*)$, *first column* $(\cdot, \phi_1, \phi_2, \phi_3)^T$ *and zeros elsewhere.*

(ii) *(right-pushed form)* $V = -\mathfrak{r}\mathfrak{D}$ *with* $\mathfrak{r} = R_0 + \sum_\mu R_{1,\mu}$. *Here* $(R_{1,\mu}h)(x) = \phi_\mu(x)^*h(x) - \phi_\mu(x - \eta e_\mu)U_{x-\eta e_\mu,\mu}(\tilde{B})^{-1}h(x - \eta e_\mu)$ *and* $R_0 = \sum_\mu(\psi_\mu - M_\mu) + a_k\Pi$, *where* M_μ *is multiplication by* $(\partial_\mu^\eta\phi_\mu)(x - \eta e_\mu)$.

(iii) *(left-pushed form)* $V = V^* = -\mathfrak{D}^*\mathfrak{r}^*$.

(iv) *For* $\sigma \le 1$, *in* $\ell^p(\eta^3)$ *for* $p \in \{1, 2, \infty\}$: $\|\mathfrak{r}\|, \|\mathfrak{E}\| \le c_V\sigma$ *with* $c_V = 4d + a(2d + d^2)$. *Every kernel has range* $\le \sqrt{3}$, *so* $\|\mathfrak{r}^\beta\|, \|\mathfrak{E}^\beta\| \le \mathrm{e}^{\beta\sqrt{3}}c_V\sigma$ *for every* 1*-Lipschitz weight* ρ *and* $|\beta| \le 1$. *Moreover* $V(x, \cdot) = 0$ *unless* $\operatorname{dist}(x, \operatorname{supp}\tilde{A}) \le \sqrt{3} + \eta$.

Bałaban's site form [1, (3.44)] and the shifted form describe the same operator.

Proof. By [4, p. 580], $D_{\tilde{A}+\tilde{B},\mu} = U_\mu(\tilde{A})(D_{\tilde{B},\mu} + \phi_\mu)$, since all transporters are functions of the generator q. U is pointwise orthogonal, so bond by bond

$$-\Delta_{\tilde{A}+\tilde{B}} = -\Delta_{\tilde{B}} + \sum_\mu (D_\mu^*\phi_\mu + \phi_\mu^* D_\mu + \psi_\mu).$$

The same bonds are summed on T_η and on a Neumann domain. Also $Q_k(\tilde{A} + \tilde{B}) = Q + F_2$. This gives (i).

For (ii): with $[\phi, U] = 0$ one has $D_\mu^*(\phi_\mu f)(x) = -\phi_\mu(x - \eta e_\mu)U^{-1}(D_\mu f)(x - \eta e_\mu) - (\partial_\mu^\eta\phi_\mu)(x - \eta e_\mu)f(x)$. Inserting this into (i) gives (ii). (iii) holds because both inverses are real symmetric for real backgrounds.

For (iv): $|\mathrm{e}^{q\vartheta} - 1| \le |\vartheta|$ gives $|\phi_\mu| \le \sigma$ and $|\partial^\eta\phi| \le \sigma$. Also $|U(\tilde{A}(\Gamma)) - 1| \le d\sigma$, because the contours lie in $B^k(y)$ and have length $\le d$. Finally $\|Q\| \le 1$. Schur bounds give the constants. The support clause holds because each term carries ϕ or F_1', which vanish off bonds or contours meeting $\operatorname{supp}\tilde{A}$. □

Lemma E.116 (one-variable remainder bound)**.** *Assume the hypotheses of* [3, p. 412] *on* $\tilde{A}, \tilde{B}$ *and* (R_ϑ). *Let* $N := n + n' \ge 4$ *and* $\alpha \in (0, 1)$, *and write* $K(x, y)$ *for the kernel of the operator* $K_{n,n'}$ *above. Then for all* x, y *at distance* $\ge R_0$ *from* Ω^c:

$$|K(x,y)| + |D_xK(x,y)| + \sup_{|x'-x|\le 1} |x-x'|^{-\alpha}|U(\Gamma_{x,x'})D_xK(x',y) - D_xK(x,y)| \le C_N(C_*\varepsilon_V)^N\mathrm{e}^{-\delta_N|x-y|},$$

and the same holds in the variable y *with* x *fixed. Here* $\varepsilon_V = e_k\mathfrak{a}(1 + e_k\mathfrak{a})\mathrm{e}^{e_k\mathfrak{a}} \le Ce(t)p(t)$, $\mathfrak{a} = \sup(|\tilde{A}| + |\nabla\tilde{A}|)$, $\delta_N = \delta_2 4^{-N}$ *and* $\delta_2 = \delta_0/4$. *The bound is per variable only; it contains no mixed derivative* D_xD_yK.

The lemma does not give the two-variable form that [3, (2.5)] is used for; that form is Lemma E.117. The tree rate of Proposition E.97 is taken $\le \delta_N$, a choice made at stage 4 of Appendix A.

Proof. Inputs. (I-G) is the pointwise kernel bound for G and $D_{\mathcal{B}}G$ (decay $\mathrm{e}^{-\delta_2|x-u|}$, singularities ρ_u^{-1} and ρ_u^{-2}, where $\rho_u(x) = |x - u| + \eta$). It is Lemma D.26. (I-R) is [1, (2.24)–(2.25)].

Convolution. For $0 < a, b < 3$:

$$\sum_u \eta^3\rho_x(u)^{-a}\rho_y(u)^{-b}\mathrm{e}^{-\delta(|x-u|+|u-y|)} \le C\,\mathrm{e}^{-\delta|x-y|/4}\Phi_{a+b}(\rho_y(x)),$$

where $\Phi_s(\rho) = \rho^{3-s}$, $\rho^{-1/2}$ or 1 according as $s > 3$, $s = 3$ or $s < 3$. This is shown by splitting into the regions $|u - x| \le r/2$, $|u - y| \le r/2$ and the rest, as in Lemma D.9(a); it is uniform in η.

Reduction to x. V and both propagators are self-adjoint, so $K_{n,n'}(x, y) = K_{n',n}(y, x)^*$.

Chain. Write $K = Y_N \cdots Y_1Y_0$ with Y_0 the rightmost propagator and $Y_i = G_iV$. Fix y and set $u_0 = G_0(\cdot, y)$ and $u_i = G_iVu_{i-1}$, so that $K(\cdot, y) = u_N$. Lemma E.115(iv) and the convolution bound give successively $|u_1| \lesssim \varepsilon_V\rho_y^{-1/2}$, $|u_2| \lesssim \varepsilon_V^2$, and u_3, Du_3 bounded. From $i \ge 4$ on, (I-R)

applied to $f = Vu_{i-1} = \sum_{\square} f 1_{\square}$ gives the sup, derivative and α-Hölder bounds with decay. The decay rate is quartered at most once per step, which gives δ_N.

Where the R_0 restriction is harmless. Every point at which $Vu_{i-1} \neq 0$ lies within $2\sqrt{d}+1$ of $\operatorname{supp}\tilde{A}$. The endpoints of (1.16) lines lie in $\Omega_1 \cup U_{2\sqrt{d}+1}(\operatorname{supp}\tilde{A})$. Moreover $\operatorname{dist}(\Omega_1, \Omega^c) \geq 5r/L > R_0$ (Lemma E.100(v)), and $\operatorname{dist}(\operatorname{supp}\tilde{A}, \partial\Omega)$ exceeds the support margin, (R4‴) of Corollary 5.10. Hence (I-G) applies at every intermediate point, given $R > R_0$ and $R \geq 12\sqrt{d}$ ((R.1) and (R.8)). □

Lemma E.117 (two-variable remainder bound on T_η)**.** *Let $\Omega = T_\eta$, assume (R_ϑ), and let $0 < \beta \leq \beta_U := \min\{\beta_*, \delta_K/4\}$. For a kernel Y and $a, b \in \{0,1\}$ put $Y_{ab} := D^a Y D^{b*}$. Define*

$$\mathfrak{N}_{\alpha,\beta}(Y) := \max_{a,b} \sup_{x,x'} \mathrm{e}^{\beta|x-x'|} \max\{\|Y_{ab}(x,x')\|,\, a\,H_x,\, b\,H_{x'},\, ab\,H_{xx'}\}.$$

Here H_x and $H_{x'}$ are the α-Hölder quotients in x and in x' over $0 < |x-\tilde{x}| \leq 1$. The transporter is $U(\tilde{B}(\Gamma_{\tilde{x},x}))$, where $\Gamma_{\tilde{x},x}$ is a shortest lattice contour from $\tilde{x}$ to x. $H_{xx'}$ is the mixed second difference divided by $|x-\tilde{x}|^\alpha |x'-\tilde{x}'|^\alpha$.

(a) *(Operator form; every $m \geq 0$, $\alpha < 1$.)*

$$|(D^a K f)(x)| \leq c_K C_{(a)} (C_\infty \sigma)^m \mathrm{e}^{-\beta \operatorname{dist}(x, \operatorname{supp} f)} \|f\|_\infty.$$

The same holds for the α-Hölder quotient of DKf and for K^*.

(b) *(Two-variable form.) If $m \geq 4$ and $0 < \alpha < \frac{1}{2}$, or if $m \geq 6$ and $0 < \alpha < 1$, then $\mathfrak{N}_{\alpha,\beta}(K) \leq C_U(\alpha)(C_V\sigma)^m$, with $C_V := \mathrm{e}^{\beta(\sqrt{3}+1)} c_V \max\{c_W, 1\}$.*

(c) *(Form of [3, (2.5)].) Under the hypotheses of (b), let $0 \leq h, h' \leq 1$ with $|\partial_\mu^\eta h|, |\partial_\nu^\eta \partial_\mu^\eta h| \leq c_h$ for all μ, ν, and likewise for h'. Then*

$$\|hKh'\|_{1,\alpha} \leq \mathrm{e}^{5\beta}(1+6c_h)^2 C_U(\alpha)(2c_1 C_V)^m (e_k p(t))^m \mathrm{e}^{-\beta \operatorname{dist}(\operatorname{supp} h, \operatorname{supp} h')}.$$

If the quotients over $|x-\tilde{x}| > 1$ are included, multiply the constant by 4.

Here $c_W := (1+\sigma)^2(2\gamma_0^{-1} + ((1+d)^2-1)K_W)$. All constants depend only on $d, a, M_b, L, \gamma_0, K_W, \alpha, \beta$. The bounds are uniform in η (hence in k and ε), in ℓ_T, in the backgrounds within (R_ϑ), and in $0 \leq m^2 t^2 \leq c$. In the construction $m = 12$, so $\alpha_0 = \frac{1}{2}$ is covered.

The inputs are (H0) [4, (1.8)], (W) and (K); the tree rate of Proposition E.97 is taken $\leq \beta_U$. Part (b) is Proposition D.13(a). We repeat the structure of its proof, and prove (a) and (c).

Proof. Step 1 (convolution and smearing). By Lemma D.9(a), for $\kappa_i \in (0,3)$ and $\delta' < \delta$,

$$\sum_w \eta^3 K^\delta_{\kappa_1}(z-w) K^\delta_{\kappa_2}(w) \leq c_C K^{\delta'}_{\kappa_1+\kappa_2}(z),$$

with a logarithm at $\kappa_1 + \kappa_2 = 3$ and boundedness above 3. By Lemma D.9(c), a local operator of range $\leq \sqrt{3}$ with entries $\leq r_R$ maps $K^\delta_\kappa(\cdot - y)$ to $c_s r_R K^\delta_\kappa(\cdot - y)$.

Step 2 (end factors). Define the two- and three-line ends $E^{(2)}_a := D^a L_0 V L_1$ and $E^{(3)}_a := D^a L_0 V L_1 V L_2$. Write every vertex in right-pushed form (Lemma E.115(ii)). Then each line of an end carries exactly one derivative. With $\delta' = \delta_K/2$, (K) together with Step 1 gives $\|E^{(2)}_a(x,y)\| \leq c_E \sigma K^{\delta'}_2(x-y)$ and $\|E^{(3)}_a(x,y)\| \leq c'_E \sigma^2 K^{\delta'}_3(x-y)$. The Hölder differences give $K_{2-\alpha}$, respectively $K_{3-\alpha}$. In weighted ℓ^2, the end factors are bounded by $J_\kappa := (\sum_y \eta^3 K^{\delta'}_\kappa(y)^2 \mathrm{e}^{2\beta|y|})^{1/2}$, and $J_\kappa < \infty$ uniformly in η if and only if $\kappa > \frac{3}{2}$. This explains the thresholds: $J_{2-\alpha}$ is finite if and only if $\alpha < \frac{1}{2}$, and $J_{3-\alpha}$ is finite for every $\alpha < 1$.

Step 3 (sandwich, $m \geq 4$). Write $K_{ab} = E_L \mathcal{N} E_R$ with $E_L = E^{(2)}_a$, $E_R = (D^b L_m V L_{m-1})^*$ and $\mathcal{N} = V L_2 V \cdots V L_{m-2} V$. Put the first vertex of $\mathcal{N}$ in right-pushed form, the last in left-pushed form, and the others in symmetric form:

$$\mathcal{N} = \pm \mathfrak{r}(\mathfrak{D} L_2 \mathfrak{D}^*)\mathfrak{E} \cdots \mathfrak{E}(\mathfrak{D} L_{m-2} \mathfrak{D}^*)\mathfrak{r}^*.$$

Every factor is bounded in weighted ℓ^2; in particular $\|(\mathfrak{D}L\mathfrak{D}^*)^\beta\| \le c_W$ by (W). For $L = G(\tilde{A} + \tilde{B})$ use $D_{\tilde{B},\mu} = U_\mu(\tilde{A})^{-1} D_{\tilde{A}+\tilde{B},\mu} - \phi_\mu$. With the vertex range of Lemma D.8(iv), $\|\mathcal{N}^{\pm\beta}\| \le (\mathrm{e}^{\beta(\sqrt{3}+1)} c_V \sigma)^{m-2} c_W^{m-3}$. Cauchy–Schwarz at the two end factors, with $\rho = |\cdot - x|$ and $\mathrm{e}^{-\beta|y'-x|} \le \mathrm{e}^{-\beta|x-x'|}\mathrm{e}^{\beta|y'-x'|}$, gives

$$\mathrm{e}^{\beta|x-x'|}\|K_{ab}(x,x')\| \le (c_E J_2 \sigma)^2 (\mathrm{e}^{\beta(\sqrt{3}+1)} c_V \sigma)^{m-2} c_W^{m-3}.$$

No pointwise bound on $D_x D_y G$ is used. K_{ab} is linear in each end, so the Hölder quotients in x, in x' and jointly follow with $c_E J_2$ replaced by $2\mathrm{e}^\beta c_E(\alpha) J_{2-\alpha}$. For $m \ge 6$, repeat with three-line ends $(J_3,\, J_{3-\alpha})$.

Step 4 ((a) and (c)). For (a): $\|VL\|_{\infty,\beta} \le C_\infty \sigma$ by Lemma E.115(ii) and (K), and the norm is submultiplicative. This is the argument of [4, (2.33)] applied to one term.

For (c): apply Lemma D.11 with $\mathcal{P} = T_\eta$ and $M = C_U(\alpha)(C_V\sigma)^m$ from (b). The quotient of the undifferentiated component is telescoped along the shortest lattice contour that carries the transport, so no holonomy correction arises. Finally, $\sigma \le 2c_1 e_k p(t)$.

Remark (the threshold). At $m = 0$ the mixed diagonal kernel is of order η^{-3} (see the remark after Proposition D.13). For $m \le 3$, power counting suggests $|x-x'|^{m-3}$. The case $\frac{1}{2} \le \alpha < 1$ with $m \in \{4, 5\}$ is not treated and not needed. □

Lemma E.118 (two-variable remainder bound on Neumann Ω, restricted form)**.** *Let $\Omega = B^k(\Lambda_2^{(k-1)\prime})$ with Neumann conditions. Ω may be any union of big blocks, including faces and re-entrant edges and corners, and may wrap the torus. Then the bounds of Lemma E.117(b),(c) hold with the T_η constants, uniformly in $\eta, \varepsilon, \ell_T$ and Ω, at every point that the path evaluates. These are the external points, all vertex points and their Hölder partners, and they lie at ℓ^∞-distance $\ge 2R - O(1)$ from Ω^c.*

The inputs are Lemma E.117, [1, Prop. 2.1], (W), the conditions (R.1) (with an $O(1)$ shift) and (R.8) ($R \ge 12\sqrt{d}$) of Appendix A, the support hypothesis of [3, p. 412] in the form (R4‴), (E.1) and (E.ksp3), and the conventions (g_k gating, ℓ^∞ metric [1, (1.3)], contours inside blocks, the nesting of [2, (2.8)]: statement 15 and Lemma E.100). The unrestricted form is not expected to be uniform near re-entrant edges (at the level of derivatives, of order $\eta^{-1/3}$); those points are not evaluated.

Proof. (1) *Restriction to* $\operatorname{supp} V$. $\mathcal{N} = VL \cdots LV$ begins and ends with a vertex, and $V(y,\cdot) = 0$ unless y is within one bond or one unit block of $\operatorname{supp}\tilde{A}$ (Lemma E.115(iv)). Hence $K_{ab}(x,x') = \langle E_L(x,\cdot)\chi, \mathcal{N}\chi E_R(\cdot,x')\rangle$ with $\chi = 1_{\operatorname{supp} V}$. So only $y, y' \in \operatorname{supp} V \subset (\operatorname{supp}\tilde{A})^{+\sqrt{3}+\eta}$ enter, and every pointwise evaluation of (K) in the end factors is at a vertex point. $V = -\mathfrak{D}^*\mathfrak{E}\mathfrak{D}$ is exact on Ω with no face term, because both operators have the same Neumann bond set and the same m^2, and the Q-contours lie inside unit blocks.

(2) *Every endpoint is deep.* [3, (2.5)] and [3, (1.33)] take the $1, \alpha$ norm of $h(1.16)h'$ on *every* line, so derivative-free legs must be traced as well. All endpoints lie in $U_{O(1)}(\Omega_1) \cup U_{O(1)}(\operatorname{supp}\tilde{A})$. The block-spin legs (1.13)–(1.15) are gated by $g_k A'$ (from the $T^\eta_{\ldots,e' g_k A' + A^{(k)}}$ of [3, (1.4)]) or by $\tilde{A}$. The $n = 0$ $\tilde{A}$-vertices sit at $\operatorname{supp}\tilde{A}$.

(3) *Margin.* Under the literal hypothesis $\operatorname{dist}(\operatorname{supp}\tilde{A}, \partial\Omega) > 2r$ the margin is $2R - O(1)$. Under (R4‴) of Corollary 5.10 the vertex points satisfy

$$\operatorname{dist}(\cdot, \Omega^c) > \tfrac{7}{4} r - \sqrt{3} - 1 \ge R_0 + 2\sqrt{3} + 1$$

whenever $R \ge 3R_0 + 15$. The proof uses only a margin of type R_0 at these points; the weaker margin $2r_k - M_b - 5$ would also suffice.

(4) *Middle factor.* The middle factor is uniform in the domain: Lemma D.24 applies to a union of whole blocks with any b.c. and a 1-Lipschitz weight. The end factors use [1, Prop. 2.1] only at admissible pairs. With (1)–(3), the proof of Lemma E.117 goes through. □

No other mass entry: statement 4.

Lemma E.119 (no other mass entry)**.** *A* mass entry *is a term $c(x)|\phi(x)|^2$ or $c(x)A_\mu(x)A_\nu(x)$ of an effective action, or of a quadratic form defining a covariance, whose coefficient c does not depend on any field. Consider the step-k expansion of* [3]*: vertices (1.6)–(1.15); lines $G_k(\Omega_2, \tilde{B})$, δG_k and (1.16); the vector G_k. Through the renormalisation of* [3, Prop. 1] *(pp. 424–445) and the localisations (pp. 420, 422, 432–433):*

(i) *every field-independent coefficient of $|\phi|^2$ is either a piece of $\delta m^2_{\mathrm{fin}}$, $\delta m^2_{K,k}$, δm^2_k (or their localisations) or the covariance entry m^2t^2;*
(ii) *no field-independent coefficient of $A \cdot A$ is added to any covariance, and there is no photon mass counterterm;*
(iii) *every other field-independent quadratic local term is an interaction vertex of $\mathcal{P}$ bounded inside* [3, (1.33)] *with at least one coupling factor, and its M-dependence is only through m^2t^2;*
(iv) *hence every occurrence of the Higgs mass in* [3] *is of type (α) or (β) of Lemma A.9(ii), or free of the mass.*

The lemma uses the classification of the two-leg classes (Lemma E.166) and the written formulas of [3, pp. 412–445]. It does not depend on Lemma E.116.

Proof. The divergent two-leg classes are C1, C1′, C2 (scalar) and C3, C4 (photon). Every other two-leg graph has $D > 0$ and is bounded as a non-local expression, so it produces no local term.

- *C1* (pp. 435–437). The tadpoles are cancelled exactly by their counterterm insertions (3.7). For the exchange graph, the zeroth moment is cancelled exactly by the counterterm of (3.9), whose counterterm is the same graph with both legs at x. The first moment is the derivative vertex (3.15). The remainder (3.12) is non-local.
- *C1′* gives only first-derivative vertices.
- *C2*: the zeroth moment is cancelled exactly by (3.19).
- *Photon classes.* Under the C^ξ replacement, (3.27)–(3.28) kill the C^ξ part of the first moment $\Pi_{\mu\mu'\nu}$, and (3.29) (Ward–Takahashi identity at $M^2 = 1$) kills the zeroth moment $\Pi_{\mu\mu'}$.

The localisations are partitions of unity. The cuts of p. 432 replace lines by pairs of external fields. $\delta G_k(\Omega, \square, \tilde{B})$ and $\delta G_k(\square, \eta\mathbb{Z}^d, 0)$ are non-local two-variable external fields. The B_0 gauge-out is exact. The B'-expansion produces only vertices with B' legs. Finally, [3, (1.20), p. 416] has δm^2 for the scalar only. Together these give (i)–(iii), and (iv) follows from (i)–(iii) and the list of the mass occurrences in Lemma A.9(ii).

The $A \cdot A$ vertices. The rewritings (3.30) and (3.32) leave field-independent local $A \cdot A$ ("$\sim * \sim$") vertices of order $e(t)^2$. Their coefficient is $(L^{j_0}\eta)^{-d+2} = (L^{j_0}\eta)^{-1}$ times a convergent expression built from $\Upsilon = 1 - m^2t^2 - aP$. They are mass-*type* vertices of $\mathcal{P}$, not mass entries. They are harmless for Theorem 5.2 by Lemma E.130. □

The remainder on the cube $\square$ of [3, p. 433]: statements 6, 7. Setting (p. 433). $\square$ is a cube of side $3r$ and $\square_1$ the concentric cube of side r; both are unions of big blocks, and $d(z) := \mathrm{dist}(z, \square^c)$. Put $G_0 = G_k(\square, B_0)$ with B_0 constant and $G_1 = G_k(\square, B_0 + B')$, both Neumann. Put $\sigma' := e_k(\|B'\|_\infty + \|\partial^\eta B'\|_\infty) \le c_2' e_k p^2$. The remainder is $K^\square = L_0 V L_1 \cdots V L_m$ with $m \ge 6$.

Lemma E.120 (separation with one interior point, ℓ^2 in the other)**.** *Let $A \in \{B_0, B_0 + B'\}$ be in the class of* [1, Prop. 2.1] *on $\square$, with $\square$ a union of big blocks. Let $L = G_k(\square, A)$, $a \in \{0, 1\}$, $\rho \ge 16$, and $z \in \square$ with $d(z) \ge \rho/2 + R_0 + 4$. Let $S \subset \square$ be arbitrary (S may meet $\partial\square$) with $\mathrm{dist}(z, S) \ge \rho$. Then for $0 \le \beta \le \delta_0/8$:*

(i) $\|\mathrm{e}^{\beta|\cdot - z|} 1_S (D^a L \mathfrak{D}^*)(z, \cdot)\|_{\ell^2(\eta^3)} \le C_* \mathrm{e}^{-\delta_* \rho}$ *with $\delta_* := \delta_1/32$;*
(ii) *for $|z - \tilde{z}| \le 1$ and $a = 1$, the α-Hölder version of (i) holds with constant $C_*(\alpha)$;*
(iii) *the mirror statement holds in the first variable.*

The constants are uniform in η, ℓ_T, $r(t)$, m^2t^2, B_0 and B'. No constancy of A near $\partial\square$ is assumed.

A pointwise bound on $(DG_1\mathfrak{D}^*)(z,w)$ for w on $\partial\square$ is not used.

Proof. Inputs. (P1) is the following interior bound: for u,z with $d(u),d(z)\ge R_0$ and $|z-u|\ge 2$, $|(D^aLD_A^{b*})(z,u)|\le c_S\mathrm{e}^{-\delta_1|z-u|/2}$, with Hölder quotient $\le c_S(\alpha)\mathrm{e}^{-\delta_1|z-u|/4}$. This follows from [3, (2.6), (2.10)–(2.11)] and [1, Prop. 2.1]. (P2) is [4, Cor. 2.3, (2.30), pp. 580–581]: the four forms $\langle f,Gf'\rangle$, $\langle f,DGf'\rangle$, $\langle f,GD^*f'\rangle$, $\langle f,DGD^*f'\rangle$ are $\le c_0\mathrm{e}^{-\delta_0\operatorname{dist}(\operatorname{supp}f,\operatorname{supp}f')}\|f\|_2\|f'\|_2$ for $\square$ a union of big blocks. This rests on [4, Lemma 2.1] only; the constancy near $\partial\square$ is a hypothesis of Lemma 2.2 alone. (P3) is the conversion $D_{B_0}=U(B')^{-1}D_A-\phi$, a bounded same-site multiplication with $|\phi|\le\sigma'$.

Step 1 (duality). $\|1_{S'}K(z,\cdot)\|_2=\sup\{|(Kf)(z)|:\|f\|_2\le 1,\ \operatorname{supp}f\subset S'\}$. Put $g=L\tilde{\mathfrak{D}}^*f$.

Step 2 (interior cut-off). Let $\chi=\varphi(|\cdot-z|)$, with $\varphi=1$ on $[0,\rho/4]$, $\varphi=0$ on $[\rho/2,\infty)$ and linear in between. Then $\chi L^{-1}g=0$, so $\chi g=L[L^{-1},\chi]g=L\tilde{\mathfrak{D}}^*h$. Here h is supported in $A'=\{\rho/4-2\le|y-z|\le\rho/2+2\}$, and $\|h\|_2\le(c_d/\rho)(\|1_{A'}g\|_2+\|1_{A'}D_Ag\|_2)$. All links used are interior ($d\ge R_0+2$), so no boundary term appears.

Step 3. By (P1), $|D^ag(z)|\le c_S\mathrm{e}^{\delta_1}(5\rho)^{3/2}\mathrm{e}^{-\delta_1\rho/8}\|h\|_2$.

Step 4 (far field in ℓ^2 only). By (P2), $\|1_{A'}g\|_2+\|1_{A'}D_Ag\|_2\le 2(d+1)^2c_0\mathrm{e}^{-\delta_0\operatorname{dist}(A',S')}\|f\|_2$.

Step 5. Decompose S into dyadic shells $S_n=S\cap\{2^n\rho\le|\cdot-z|<2^{n+1}\rho\}$ and sum, using $\beta\le\delta_0/8$. This gives (i) for $\tilde{\mathfrak{D}}$; (P3) gives (i) for $\mathfrak{D}$, with $C_*\mapsto 4C_*$.

Step 6. The same steps with the Hölder line of (P1) give (ii) with $\delta_1/32$. (iii) follows from $L^*=L$. No pointwise value within $\rho/2$ of $\partial\square$ is ever taken. □

Lemma E.121 (the remainder on the cube $\square$)**.** *Let $m\ge 6$, $0<\alpha<1$ and $0<\beta\le\beta_\square:=\min\{\beta_U,\delta_1/32,\delta_0/8\}$. Take D and the transports in $\mathfrak{N}_\alpha$ with respect to the same background. Let $0\le h,h'\le 1$ with $|\partial_\mu^\eta h|,|\partial_\nu^\eta\partial_\mu^\eta h|\le c_h$ for all μ,ν, and likewise for h'. Assume* $\operatorname{dist}_\infty(\operatorname{supp}h\cup\operatorname{supp}h',\square_1^c)\ge 2$. *Then*

$$\sup_{x,x'\in\square_1}\mathrm{e}^{\beta|x-x'|}\cdot(\textit{all components of }\mathfrak{N}_\alpha(K^\square))\le C_\square(\alpha)(C_\square\sigma')^m,$$

$$\|hK^\square h'\|_{1,\alpha}\le\mathrm{e}^{5\beta}(1+6c_h)^2C_\square(\alpha)(C_\square\sigma')^m\mathrm{e}^{-\beta\operatorname{dist}(\operatorname{supp}h,\operatorname{supp}h')}.$$

Here $C_\square=2C_0$ and $C_0=2\mathrm{e}^{\beta\sqrt{3}}c_V'\max\{c_W,c_S,C_*,C_*(\alpha),c_K,1\}$, with $c_V'=c_V+2d$. The bounds are uniform in η, ℓ_T, $r(t)$, m^2t^2, B_0 and B'. The η^{-1} boundary term never enters.

The inputs are Lemmas E.117, E.124 and D.11, (H0), (W), (E.1), (R.1), and the choice $r(t)\ge\max\{107,3R_0+15,\frac43(R_0+4),O(\bar n)+4\sqrt{3}+2\}$, which does not depend on ε. For the cut-offs of Lemma E.103(iii), the support hypothesis holds by Lemma E.122.

Proof. (1) *The defect.* Lemma E.115(i) holds *exactly* on the Neumann cube, boundary sites included: $V=-\mathfrak{D}^*\mathfrak{E}\mathfrak{D}$ with $\|\mathfrak{E}\|\le c_V\sigma'$. The unpaired term $-\eta^{-1}\phi_\mu(z)f(z)$ appears only after a vertex touching the collar is right- or left-pushed. It is a property of the rewriting, not of V.

(2) *Split.* Put $\theta(z)=\min\{1,\max\{0,3d(z)/r-1\}\}$ and

$$V=V_\theta+V_c,\qquad V_\theta=-\mathfrak{D}^*\Theta\mathfrak{E}\Theta\mathfrak{D},\qquad V_c=-\mathfrak{D}^*(\mathfrak{E}-\Theta\mathfrak{E}\Theta)\mathfrak{D}.$$

V_θ satisfies Lemma E.115 with c_V' and has no unpaired term. V_c is supported in $\mathcal{C}\times\mathcal{C}$, where $\mathcal{C}=\{d<2r/3+2\sqrt{3}\}$, and $\|\mathfrak{E}_c\|\le 2c_V\sigma'$. For $x\in\square_1$ and $w\in\mathcal{C}$, $|x-w|\ge s:=r/3-2\sqrt{3}\ge r/6$. Grouping by the first and last c-vertex gives $K^\square=K_\theta+\sum_{i_1\le i_2}K_{i_1,i_2}$, a sum of at most $m(m+1)/2+1$ terms.

(3) K_θ. The proof of Lemma E.117(b) with three-line ends applies. Given $r\ge 3(R_0+2+\sqrt{3}+1)$, every pointwise evaluation of (K) is between points with $d\ge R_0+2$: external points have $d\ge r$, and V_θ-sites have $d\ge r/3-\sqrt{3}-\eta$. So only interior points are used, and the parallelepiped sentence of [1, p. 611] is not needed.

(4) K_{i_1,i_2}. Keep both V_c and every full V between them in symmetric form, and push the V_θ outwards. Then $D^a K_{i_1,i_2} D^{b*}(x,x') = \pm\langle \mathcal{L}_x, \mathcal{N}\mathcal{R}_{x'}\rangle$. Here $\mathcal{N} = \mathfrak{E}_c(\mathfrak{D}L\mathfrak{D}^*)\mathfrak{E}\cdots\mathfrak{E}_c$ is bounded in weighted ℓ^2 by (W), for any boundary condition. The end factors $\mathcal{L}_x, \mathcal{R}'_x$ are needed only in weighted $\ell^2(\mathcal{C})$. Split the points $z \in B_x = \{|z-x| < s/2\}$ from those with $z \notin B_x$.

- On B_x, use Lemma E.120 with $\rho = s/2$. Its hypotheses hold if and only if $r \geq 3(32+2\sqrt{3})$ and $\frac{3}{4}r \geq R_0 + 4$.
- Off B_x, use (W).

A pointwise separation bound for G_1 near $\partial\square$ is not available, because the proof of [4, p. 581] needs A to be constant near $\partial\square$; Lemma E.120 avoids it. The Hölder bound in x works for every $\alpha < 1$. The assembly is that of Step 3 of the proof of Lemma E.117.

(5) Summing, $m(m+1)/2 + 2 \leq 2^m$. The h, h' form follows from Lemma D.11 with $\mathcal{P} = \square_1$: by the support hypothesis, every lattice point within ℓ^∞-distance $1+\eta \leq \frac{3}{2}$ of $\operatorname{supp} h \cup \operatorname{supp} h'$ lies in $\square_1$. □

Lemma E.122 (support margin for the remainder on $\square$)**.** *Let d be the ℓ^∞ distance on the torus, $B_\varrho(c) := \{z : d(z,c) \leq \varrho\}$, s_b the side of a big block, and $r = r(t) = R(1+\log t^{-1})^2$ with $t \leq \varepsilon_0 < \mathrm{e}^{-1}$. Let K be a component of the step of* [3, p. 433] *that contains an old vertex. Let Q be a closed cube of side ℓ containing the localisation cubes $\square(u)$ of all old vertices u of K, and let $c \in Q$. Put*

$$\square_1 := \bigcup\{b : \ b \cap B_{r/2}(c) \neq \emptyset\}, \qquad \square := \bigcup\{b : \ b \cap B_{3r/2+s_b}(c) \neq \emptyset\},$$

where the unions run over big blocks b. Let $w \geq 0$, let v, v' be old vertices of K, and let $\operatorname{supp} h \subset U_w(\square(v))$ and $\operatorname{supp} h' \subset U_w(\square(v'))$, where U_w is the closed ℓ^∞-neighbourhood of width w. If $r \geq 2\ell + 2w + 4$, then:

(i) *$Q \subset \square_1 \subset \square$, and $d(x,y) > r$ and $|x-y| > r$ for all $x \in \square_1$ and $y \notin \square$;*
(ii) *$\operatorname{dist}_\infty(\square(u), \square_1^c) > w+2$ for every old vertex u of K;*
(iii) *$\operatorname{dist}_\infty(\operatorname{supp} h \cup \operatorname{supp} h', \square_1^c) > 2$.*

For the cut-offs of Lemma E.103(iii) one has $w = 1$. With $\ell = \ell_{\bar{N}}$, the condition $r \geq 2\ell + 6$ holds for every $t \leq \varepsilon_0$ under the lower bound $R \geq 2\ell_{\bar{N}} + 10M_b + 10$ of Lemma E.124. In that case, for these cut-offs h, h' of the old vertices v, v' at the two ends of the line carrying $K^\square$, the second display of Lemma E.121 holds with $c_h = 40/\sqrt{3}$.

Proof. Step 1 (the hulls). A big block has ℓ^∞-diameter at most s_b, and the big blocks cover the torus. Hence $x \in \square_1$ implies $d(x,c) \leq r/2 + s_b$, and $y \notin \square_1$ implies $d(y,c) > r/2$. Likewise, $y \notin \square$ implies $d(y,c) > \frac{3}{2}r + s_b$.

Step 2 (part (i)). Every $q \in Q$ satisfies $d(q,c) \leq \ell \leq r/2$, so $Q \subset B_{r/2}(c) \subset \square_1$. For $x \in \square_1$ and $y \notin \square$, $d(x,y) > \frac{3}{2}r + s_b - \frac{r}{2} - s_b = r$. In particular $\square_1 \subset \square$. Since the Euclidean distance dominates the ℓ^∞ distance, also $|x-y| > r$.

Step 3 (part (ii)). Since $\square(u) \subset Q$, for $q \in \square(u)$ and $y \notin \square_1$ we have $d(q,y) > \frac{r}{2} - \ell \geq w + 2$.

Step 4 (part (iii)). For $z \in \operatorname{supp} h$ there is $q \in \square(v)$ with $d(z,q) \leq w$. So for $y \notin \square_1$, $d(z,y) > w + 2 - w = 2$; the same holds for h'.

Step 5 (the cut-offs and R). In Lemma E.103(iii), $h(x) = \prod_\mu \chi(t_\mu(x))$ with $\chi = 0$ on $[1,\infty)$. So $h(x) \neq 0$ implies $\operatorname{dist}_\infty(x, \square(v)) < 1$, and $w = 1$. For $t \leq \varepsilon_0$ we have $r(t) \geq R$. Since $M_b \geq 1$, $R \geq 2\ell_{\bar{N}} + 10M_b + 10 \geq 2\ell_{\bar{N}} + 20 > 2\ell_{\bar{N}} + 6$.

Finally, by (i) the sets $\square_1 \subset \square$ satisfy $\operatorname{dist}(\square_1, \square^c) \geq r$, and $\square$ is a box and a union of big blocks; this is the only geometry of $\square_1 \subset \square$ used in the proof of Lemma E.121. By (iii) and Lemma D.12, the hypotheses of the second display of Lemma E.121 hold with $c_h = 40/\sqrt{3}$. □

Two-domain bounds δG: statement 8.

Lemma E.123 (two-domain bounds)**.** *Let $X_{\mathrm{in}} \subset X_{\mathrm{out}}$ be unions of big blocks of T_η, $\eta = L^{-k} \leq \frac{1}{2}$, where X_{out} may be T_η and $X_{\mathrm{in}} \neq T_\eta$. Let A be real with $\theta(A) \leq \vartheta_R$, let $0 \leq m^2 t^2 \leq c$, and put*

$\delta := G_k(X_{\mathrm{out}}, A) - G_k(X_{\mathrm{in}}, A)$ and $\rho_{\mathrm{in}} := \mathrm{dist}_\infty(\cdot, X_{\mathrm{in}}^c)$. Let $0 < \alpha < 1$ and $0 < \beta \le \beta_D := \min\{\beta_*, \delta_K/4\}$. Then for x, x', and their Hölder partners, with $\rho_{\mathrm{in}} \ge R_0 + 11$:

$$\mathrm{e}^{\beta|x-x'|} \max\{\|X_{ab}(x,x')\|, aH_x, bH_{x'}, abH_{xx'}\} \le C_D(\alpha) \exp\big(-\tfrac{\delta_K}{2}[(\rho_{\mathrm{in}}(x)-R_0-7)+(\rho_{\mathrm{in}}(x')-R_0-10)]\big),$$

where $X_{ab} = D^a \delta D^{b*}$. C_D depends only on $d, a, M_b, L, \gamma_0, K_W, \alpha$ and on the profiles f, φ of Lemma D.15.

(loc) *Let $0 \le h, h' \le 1$ satisfy $|\partial_\mu^\eta h|, |\partial_\nu^\eta \partial_\mu^\eta h| \le c_h$ for all μ, ν, and likewise for h'. If $\mathrm{dist}_\infty(\mathrm{supp}\, h \cup \mathrm{supp}\, h', X_{\mathrm{in}}^c) \ge R_0 + 12$ (and $\ge R_0 + 13$ if $k = 1$), then*

$$\|h \delta h'\|_{1,\alpha_0} \le C_{\mathrm{loc}} \mathrm{e}^{-\beta\, \mathrm{dist}(\mathrm{supp}\, h, \mathrm{supp}\, h')}, \qquad C_{\mathrm{loc}} = \mathrm{e}^{5\beta}(1 + 6c_h)^2 C_D(\alpha_0).$$

This applies to the pairs $(X_{\mathrm{out}}, X_{\mathrm{in}}) = (\Omega, \Omega_2), (\Omega, \square), (T_\eta, \square)$.

The inputs are (H0), (W), (K) and the pushing identity [4, (2.32)]. At every $k \ge 1$, the point margin is $R_0 + 10 + 2\eta \le R_0 + 11$, and the support margin is $R_0 + 11 + 3\eta$ (Proposition D.17 and Corollary D.18). At the uses the margins are met; see Remark D.20.

Proof. The setting is that of Proposition D.17, with $\rho = \rho_{\mathrm{in}}$. Points with $\rho_{\mathrm{in}} \ge R_0 + 11$ lie in $\mathcal{P}_D = \{\rho \ge R_0 + 10 + 2\eta\}$, so the first display is Proposition D.17. Part (loc) is Corollary D.18 with $\alpha = \alpha_0$, since $R_0 + 12 \ge R_0 + 11 + 3\eta$ for $k \ge 2$ and $R_0 + 13 \ge R_0 + 11 + 3\eta$ for $k = 1$.

We recall the structure of the proof.

- The cut-offs are $\chi = \chi_{R_0+3}$ and $\psi = \chi_{R_0+13/2}$ of Lemma D.15, for every $\eta \le \frac{1}{2}$.
- The commutators $C_\chi = -\mathfrak{r}_\chi \mathfrak{D} = \mathfrak{D}^* \mathfrak{r}_\chi^*$ of Lemma D.16 are supported in $S_\chi \subset \{R_0 + \frac{7}{4} - \eta < \rho_{\mathrm{in}} < R_0 + \frac{21}{4} + 2\eta\} \subset \{\psi = 0\}$ and $S_\psi \subset \{R_0 + \frac{21}{4} - \eta < \rho_{\mathrm{in}} < R_0 + \frac{35}{4} + 2\eta\}$.
- This gives the two-commutator representation $X_{ab} = -E_L\, \mathfrak{M}\, E_R$, with $E_L = D^a G_{\mathrm{out}} \mathfrak{r}_\chi$, $\mathfrak{M} = \mathfrak{D} G_{\mathrm{in}} \mathfrak{D}^*$ and $E_R = \mathfrak{r}_\psi^* G_{\mathrm{in}} D^{b*}$.
- The ends are one-derivative kernels at pairs ≥ 1 apart and at depth $> R_0 + \frac{5}{4}$ in X_{in}. So (K) bounds them in weighted ℓ^2, with the gains $\mathrm{e}^{-(\delta_K/2)(\rho_{\mathrm{in}} - R_0 - 7)}$ and $\mathrm{e}^{-(\delta_K/2)(\rho_{\mathrm{in}} - R_0 - 10)}$.
- The middle factor is bounded by (W), and Cauchy–Schwarz concludes.

No kernel is evaluated at points less than 1 apart, or within $R_0 + 1$ of X_{in}^c. □

The cube lies in the outer domain: statement 9.

Lemma E.124 (the cube lies in the outer domain)**.** *Let $s_b \le 2M_b$ be the big-block side and $\ell := \ell_{\bar N} = 2V(\bar N)$ with $\bar N = 169$, where $V(\bar N)$ is the maximal number of localisation cubes of an expression. Assume the conditions (T.nw) ($\ell_0 \ge \varepsilon_0[8r(\varepsilon_0) + 4M_b + 2]$), (E.r) ($\varepsilon_0 < \mathrm{e}^{-1}$), (T.div) (the block tiling is compatible with the torus) and*

$$\textit{(W-R}') \qquad R \ge 2\ell_{\bar N} + 10M_b + 10.$$

(W-R′) is implied by $R \ge R^{**}$. Let K be a component of the p. 433 step with an old vertex, let Q be the closed cube of side ℓ containing its localisation cubes, $c \in Q$, and $\square$ the big-block hull of $B_{3r_\square/2 + s_b}(c)$.

(G) *(domain-generic) If some old vertex has a point x_v with $d(x_v, \Omega_{\mathrm{out}}^c) > \rho_\square := \frac{3}{2} r_\square + 2s_b + \ell$, then $\square \subset \Omega_{\mathrm{out}}$.*

(U0) *At* [3, pp. 411–414, 432–433]*, with $\Omega_{\mathrm{out}} = B^k(\Lambda_2^{(k-1)\prime})$: $m_{\mathrm{use}} \ge 2r_k - 4$, so $\square \subset \Omega_{\mathrm{out}}$. The same holds at* [2, p. 582, (2.116)] *and for the cap families $\bar N = 169$.*

(U1a) *At* [2, p. 579]*, with $\Omega_{\mathrm{out}} = B^k(\Lambda_0^{(k)})$: $m_{\mathrm{use}} > 2r_k - M_b - 5$, so $\square \subset \Omega_{\mathrm{out}}$ under (W-R′).*

(U1b) *At* [2, Prop. 2.6]*, with $\Omega_{\mathrm{out}} = B^{k-1}(\Lambda_2^{(k-1)})$: $m_{\mathrm{use}} \ge 2.5r_k - 5$.*

(U2) *At* [2, p. 578]*, scale $j \le k - 1$, with $\Omega_{\mathrm{out}} = B^j(\Lambda_2^{(j)})$: $m_{\mathrm{use}} \ge 5r_j - 7$.*

(U3) *$\square$ is a non-wrapping box, so $\delta G_k(\square, \eta\mathbb{Z}^d, 0)$ is defined.*

The support hypothesis for $\tilde{B}^{(k)}$ at U0 follows from $\operatorname{supp}\tilde{B}^{(k)} \subset \operatorname{supp}\theta_{k+1}$, which holds because each of the three terms defining $\tilde{B}^{(k)}$ in [2, (2.89), p. 575] carries θ_{k+1} as its left factor. The completeness of the list of uses is statement 15′.

Proof. (G) Since $x_v, c \in Q$, $d(c, x_v) \le \ell$. For $p \in \square$, $d(p, x_v) \le \frac{3}{2}r_\square + 2s_b + \ell < d(x_v, \Omega_{\mathrm{out}}^c)$, so $p \notin \Omega_{\mathrm{out}}^c$. The hull satisfies $B_{3r/2+s_b}(c) \subset \square \subset B_{3r/2+2s_b}(c)$, since a big block has diameter $\le s_b$ and the hull of a box is a box ((T.div)). Moreover $d(\square_1, \square^c) \ge r$.

The per-use margins use three facts. N1 is the nesting gap (Lemma E.100(v)). N2 is $\Lambda_0^{(i+1)} \subset \Lambda_7^{(i)\prime}$ [2, p. 566], iterated. N4 is that $\operatorname{supp}\theta_{k+1}$ lies in the M_b-neighbourhood of $B^k(\Lambda_2^{(k)})$ [2, p. 567].

- (U0) The old-vertex bonds have $d \ge 2r_k - 4$. (The claim is restricted to bonds occurring in vertices, since vertex sums run over Ω.) The condition needed is $r_k/2 > \ell + 2s_b + 4$.
- (U1a) g_k-vertices: $7r_k - 10$. $\theta_{k+1}A'$-vertices: $2r_k - 4 - M_b - 1$. The condition needed is $r_k/2 \ge \ell + 2s_b + M_b + 5$, which is (W-R′) at $s_b = 2M_b$.
- (U1b) $x_v \in B^k(\Lambda_7^{(k)}) \subset \Lambda_7^{(k-1)}$-region, and N1 at scale $k-1$ with $s = 5$ gives $(5/L)(r_{k-1} - 1) - 2 \ge 2.5r_k - 5$.
- (U2) $d(\operatorname{supp}\theta_{k+1}, B^k(\Lambda_0^{(k)c})) > L^{k-j}r_k$ is printed at [2, p. 578], and N1 at scale j gives $5r_j - 7$.
- (U3) $t(3r(t) + 8M_b + 1) \le \varepsilon_0(3r(\varepsilon_0) + 8M_b + 1) < \ell_0 \le \ell_T$, by monotonicity on $t \le \varepsilon_0 < \mathrm{e}^{-1}$ and (T.nw).

In each case the condition is implied by (W-R′). □

The inner domain Ω_2 is not needed: statement 10.

Lemma E.125 (the inner domain Ω_2 is not needed)**.** *Let ℓ be a $G_k(\Omega_2, \tilde{B})$ line of a p. 433 component with ends $v, v' \in \square_1$, and suppose $\square \subset \Omega_{\mathrm{out}}$ for the outer domain of the use (Lemma E.124). Put $X_\ell := \delta G_k(\Omega, \square, \tilde{B}) - \delta G_k(\Omega, \Omega_2, \tilde{B})$. Then:*

(i) *$G_k(\Omega_2, \tilde{B}) = G_k(\square, \tilde{B}) + X_\ell$ as kernels on $\operatorname{supp}h_v \times \operatorname{supp}h_{v'}$; if $\square \subset \Omega_2$, then $X_\ell = \delta G_k(\Omega_2, \square, \tilde{B})$, which is p. 433 verbatim;*

(ii) *$\|hX_\ell h'\|_{1,\alpha_0} \le C_X \mathrm{e}^{-\beta_D \operatorname{dist}(\operatorname{supp}h, \operatorname{supp}h')}$, where C_X is built from the constants of Lemma E.123(loc) for the two pairs $(\Omega, \square)$ and (Ω, Ω_2);*

(iii) *the p. 433 transformation with (i) produces exactly the renormalised classes of p. 433. Every external field produced obeys the two-point (3.1)/(2.5) form, and $\square \subset \Omega_2$ is never used. Hence [3, (3.2) ⇒ (1.33)] holds for these expressions with tree rate $\delta_P = \min\{\delta(\bar{N}), \beta_U, \beta_D\}$ and constant $O(1)\max\{1, C_X\}^{N_\ell(\bar{n})}$, uniformly in $\varepsilon, \eta, \ell_T$.*

The proof composes Lemma E.123(loc) for (Ω, Ω_2) and $(\Omega, \square)$, Lemma E.124 at the outer domain of each use, Lemmas E.126 and E.127, the closure of the classes under cuts (Lemma E.166) and the printed base of p. 433.

Proof. (i) $[G(\Omega) - G(\square)] - [G(\Omega) - G(\Omega_2)] = G(\Omega_2) - G(\square)$. This is pure algebra and uses no nesting between $\square$ and Ω_2. $G(\Omega, \cdot)$ is evaluated at points admissible for Ω_2; by monotonicity, $\operatorname{dist}(z, \Omega^c) \ge \operatorname{dist}(z, \Omega_2^c)$ when $\Omega_2 \subset \Omega$.

(ii) Apply the triangle inequality and Lemma E.123(loc) to each pair. The points of $\square_1$ are $\ge r \ge R_0 + 12$ from $\square^c$, and the line ends are $\ge R_0 + 12$ from Ω_2^c by Lemmas E.126 and E.127.

(iii) $E(G; \ell) = E(G; \ell \mapsto G_k(\square)) + E(G \setminus \ell; X_\ell)$, and E is linear in the external field, so the graphs are those of p. 433. Relabelling the domain of a line leaves every degree [3, (2.1), p. 422] unchanged. Cutting a line does not lower the degree ([3, p. 432] and Lemma E.166(a)). X_ℓ is $\tilde{B}$-covariant, so the B_0 gauge-out preserves its norm. [3, (3.2), p. 432] requires no smallness of external fields, so the two-point decay with an $O(1)$ constant suffices. □

End points of Ω_2-lines: statements 11, 12.

Lemma E.126 (end points of Ω_2-lines in [2, Chap. 2])**.** *Every evaluation point of a $G_k(\Omega_2,\cdot)$ or $\delta G_k(\Omega,\Omega_2,\cdot)$ line on the path of Theorem 5.2 is far from Ω_2^c. The points are line ends, bond partners, block points of Q_k, Hölder partners and supports of localisation functions. The bounds are:*

- *at U1a and U1b, with $\Omega_2 = B^k(\Lambda_2^{(k)})$:* $\operatorname{dist}(\cdot,\Omega_2^c) \geq 5r_k - 5 - c_1$ *with $c_1 \leq 4$;*
- *at U2 (scale j):* $\operatorname{dist} > L^{k-j}r_k - c_1' \geq 2R - c_1'$.

Hence the margin is $\geq R_0$ under the printed $R > R_0$, and $\geq R_0 + 12$ under $R \geq \max(R_0, 12 + c_1')$. The list U1a, U1b, U2 contains all the $G_k(\Omega_2)$-type changes in [2, Chap. 2].

The lemma uses the conventions that "localized in $B^k(\Lambda_7^{(k)})$" means all vertices and that the localisation width is $O(1)$ (Lemma E.103), and that $\theta_{k+1} = 1$ on the bonds of $B^k(\Lambda_2^{(k)})$ (Lemma E.102). The quadratic-form use [2, (2.108)] needs no restriction to depth R_0 ([1, Prop. 2.2] has none).

Proof. Nesting. If Λ_{i+1}^c contains every point within ρ of Λ_i^c for m consecutive i, then $\operatorname{dist}(\Lambda_{i_0+m}, \Lambda_{i_0}^c) > m\rho - (m-1)$. (Induction along an ℓ^∞ geodesic, which changes the distance by ≤ 1 per step.) So $\operatorname{dist}(B^k(\Lambda_7^{(k)}), \Omega_2^c) \geq 5r_k - 5$.

U1a and U1b. The vertex variables lie in $\Lambda_7^{(k)}$ [2, pp. 579–580]. Evaluation points are offset by 0 for (1.6), (1.7); by $\leq \eta$ for bond legs; by ≤ 1 for Q_k contours; and by ≤ 2 for Hölder partners and supports of h. The (1.16) lines are estimated away before Prop. 2.6, so no interior chain point of a $G_k(\Omega_2)$ line occurs.

U2. [2, p. 578] prints that the distance of $\operatorname{supp}\theta_{k+1}$ to $B^k(\Lambda_0^{(k)c})$, measured on $T_{L^{-j}}$, is $> L^{k-j}r(L^k\varepsilon) > r(L^j\varepsilon)$. The loop vertices are within c_1' of $\operatorname{supp}\theta_{k+1}$.

The list of changes of the domain to Ω_2 is read off the pages of [2, Chap. 2]. The changes at the first step of [2, Sect. A] have margins $5r(\varepsilon) - O(1)$ and $2r(\varepsilon) - O(1)$ and do not affect the result. □

Lemma E.127 (end points of Ω_2-lines in [2, Chap. 3])**.** [2, Chap. 3] *(pp. 583–594) is on the path. It contains no $G_k(\Omega_2)$ graph line and no p. 433-type cube step; the cubes of Sect. 3C are combinatorial. Its only $\Omega_2 = B^k(\Lambda_2^{(k)})$ object is the form $\langle \Lambda_3^{(k)}\phi, \Delta^{(k)}(\Omega_2, \tilde{A}^{(k+1)})\Lambda_3^{(k)}\phi\rangle$ of (3.5). This form is turned into Ω by (2.108) run backwards ((3.7)–(3.8)), and its evaluation points satisfy* $\operatorname{dist}(\cdot,\Omega_2^c) \geq r(t_k) - 2 \geq R_0$ *under $R \geq R_0 + 2$.*

The margin is measured in Bałaban's metric. With a Euclidean margin the requirement becomes $R \geq \sqrt{3}(R_0 + 2)$, which is implied by $R \geq R^{**}$. The identity $\tilde{A}^{(k)} = \tilde{A}^{(k+1)}$ on Ω_2 is not used.

Proof. On the twelve pages of [2, Chap. 3] none of the occurrences of "replac", "propagat", "Proposition" is a change of the domain of G_k, and "interaction" does not occur. In (3.5) the points are $z \in B(x)$, $z' \in B(x')$ with $x, x' \in \Lambda_3^{(k)}$, plus bond partners. By [2, (2.8)], every large block of $\Lambda_3^{(k)}$ is at distance $> r(t_k)$ from $(\Lambda_2^{(k)})^c$. Subtracting 1 for the block and 1 for the bond gives $r(t_k) - 2$. □

Conventions and completeness: statements 15, 15′.

Lemma E.128 (conventions R1–R3)**.** (R1) *In* [3, (1.13)–(1.15)] *the A'-legs enter only as $g_k A'$, with* $\operatorname{supp} g_k \subset \Omega_1$.
(R2a) $\Omega = B^k(\Lambda_2^{(k-1)\prime})$ *and* $\Omega_1 = B^k(\Lambda_7^{(k-1)\prime})$, *and* $\Lambda_i^{(j)}$ *is defined by* [2, (2.7)–(2.8)] *with* $r(L^j\varepsilon)$.
(R2b) $\Lambda_{i+1} \subset \Lambda_i$, *so* $\Omega_1 \subset \Omega$.
(R2c) $\operatorname{dist}(\Lambda_{i+s}, \Lambda_i^c) > s(r-1)$.
(R2d) *The metric is the ℓ^∞ torus distance* [1, (1.3)].
(R3) *The contours $\Gamma_{y,x}^{(k)}$ lie inside $B^k(y)$.*

The infimum convention for the distance between sets is not printed. Any other convention costs at most one large-block diameter per step, which $R \gg M_b L$ absorbs. The order $\Omega \supset \Omega_1 \supset \Omega_2$ holds at U1b (Lemma E.100).

Proof. (R1) The vertices of (1.5) are built from $T^{\eta}_{\dots, e' g_k A' + A^{(k)}}$ [3, (1.4)], and p. 414 lists "(1.14) with $n' = 0$ and $\tilde{B} = A^{(k)}$" among them. An ungated reading contradicts both. It is corroborated by $g_k A'^{(k)}$ in [2, (2.107)]. (R2a) is printed at [3, pp. 411–412] and [2, p. 566]. (R2b) and (R2c) are Lemma E.100(i) and (v). (R2d) [1, (1.3)] is the only distance defined; [2] uses its symbol [2, p. 566], and [3, p. 411] adopts the notation of [1, 2]. (R3) follows from the staircase contours of [1, (2.1)], their concatenation (2.2), and box blocks [1, (1.17)]. Long arcs are excluded by [1, (1.2)]. □

Lemma E.129 (completeness of the lists of uses)**.** *The uses in* [3]*, in the proof of Theorem 5.2, of the following are exactly those treated in Lemmas E.124–E.126 and Corollary 5.10, plus two further items (O1, O2): Ω_2 or another domain; $G_k(\cdot)$ with a domain argument; δG; (2.5) and (1.16); cube constructions; and the hypothesis of p. 412 on* $\operatorname{supp} \tilde{A}$*. The consumers of the p. 412 hypothesis are p. 433, p. 422 and the domain changes (p. 414 and (2.5), pp. 424, 426, 432).*

- *O1 is* [3, (2.8)] *on T_η, with margin* 2*, at the stage of the cube □;*
- *O2 is the comparison on p. 437 of torus and $\mathbb{Z}^d$ integrals, which is the image kernel of Lemma 5.22.*

The same holds for the uses in [2] (Lemmas E.126, E.127). The Ward–Takahashi identity of [3, p. 442] with λ linear is exact on $\zeta\mathbb{Z}^d$, and on a torus it holds up to the image term of Lemma 5.22.

Proof. Every occurrence of the six objects on the journal pages 411–445 of [3] is read in context. None is used with a margin outside (W-R′), outside $R \geq R_0 + 12 + O(1)$, or outside depth $> r$ at the changes of domain. □

Statements 16 and 17.

Lemma E.130 ($A \cdot A$ local terms)**.** *Fix $\mu_* > 0$, and let $(u, M) = (1, m_2)$ or $(u, M) = (-\eta, m_1)$. Then, uniformly in ε, k and ℓ_T, the $\sim * \sim$ vertices of* [3, (3.30), (3.32)] *satisfy:*

(N1) *they enter no covariance and no Gaussian weight;*

(N2) *every localised family containing them obeys* $(1.33)^\natural$*, that is, (1.33) times $(|u|c_k)^m$. The tree rate of* [3, (1.33)] *is some $\delta(\bar{N}) > 0$ at the cap (Proposition E.60), not a rate depending on d only. The constant is $O(1) = O(\alpha_0, \bar{n})$, independent of $\varepsilon, k, \Omega, \Omega_1, \Omega_2, \tilde{B}, \ell_T$;*

(N3) *they are not iterated as a running coupling;*

(N4) *they create no new counterterm and no new vacuum entry.*

The proof is given after Proposition E.140 below.

The two-photon vertices of [3, (3.30), (3.32)]. We work in step-k units. For $j \leq k$, $\ell_j := L^j \eta \leq 1$ is the scale of a line of index j; in particular $\ell_k = 1$. We use the following two properties.

(LB) (Line bounds.) Every line G of index j in the expansion of [3, p. 414], scalar or vector, satisfies $|G(x, x')| \leq K_\ell \, \ell_j^{-1} \mathrm{e}^{-a|x-x'|/\ell_j}$. Each derivative at an end costs a factor ℓ_j^{-1}, and an α-Hölder quotient of a derivative costs a factor $\ell_j^{-1-\alpha}$; this holds in each variable separately and in both together [3, (2.10)–(2.12), p. 426; p. 418]. The constant K_ℓ is independent of ε, k, ℓ_T. For the vector lines it is independent of μ_*. The slice weights of [24] multiply it by at most 2 per line, and coloured lines carry the factor $(|u|c_k)$ per line.

(VB) (Vertex bounds.) In the notation of Lemma E.24, the two-photon vertex is

$$e(t_k)^2 \sum_z \eta^3 g(z) A_\mu(z) c_{\mu\mu'}(z) g'(z) A'_{\mu'}(z),$$

with $|c_{\mu\mu'}| \leq \Gamma_3 \ell_{j_0}^{-1}$, where j_0 is the lowest index of the legs of the vertex. The first-moment vertex has $|c_{\mu\mu'\nu}| \leq \Gamma'_3$, and the vertex of [3, (3.32)] has $|c^{(4)}| \leq \Gamma_4$ (Lemmas E.24 and E.32).

We say that a two-photon vertex has an *unshared lowest-index line* if its line of index j_0 is neither the lowest-index line of another two-photon vertex nor the transported leg of a renormalised subgraph of the same class [3, p. 440].

Lemma E.131 (lattice sum)**.** *For $0 < \eta \le \ell$ and $b > 0$, $\sum_{z\in\eta\mathbb{Z}^3} \eta^3 \mathrm{e}^{-b|z|/\ell} \le S(b)\ell^3$, where $S(b) := (2\sqrt{3}/b + 1)^3$.*

Proof. Since $|z| \ge |z|_1/\sqrt{3}$, the sum is at most $\big(\eta \sum_{n\in\mathbb{Z}} \mathrm{e}^{-c|n|}\big)^3$ with $c = b\eta/(\sqrt{3}\ell)$. Now $\sum_n \mathrm{e}^{-c|n|} = \coth(c/2) \le 1 + 2/c$. Hence $\eta \coth(c/2) \le \eta + 2\sqrt{3}\ell/b \le (1 + 2\sqrt{3}/b)\ell$. □

Lemma E.132 (insertion bound)**.** *Assume (LB), (VB) and $L = 2$, and let a two-photon vertex at z have an unshared lowest-index line.*

(a) *(Both legs on lines.) Let the legs be lines G_0, G_1 of indices $j_0 \le j_1$, and let $0 < \delta' < a$. Then $K(x, y) := \sum_z \eta^3 G_0(x, z)c(z)G_1(z, y)$ obeys*

$$|K(x, y)| \le K_\ell^2 \Gamma_3 S(\delta')\, \ell_{j_0}\, \ell_{j_1}^{-1} \mathrm{e}^{-(a-\delta')|x-y|/\ell_{j_1}}.$$

This is the bound (LB) of a line of index j_1, times $K_\ell\Gamma_3 S(\delta')\ell_{j_0}$. A derivative at y costs $\ell_{j_1}^{-1}$, and a contour leg at x gives an extra factor $\ell_{j_0} \le \ell_{j_1}$. Moreover $\sum_{j_0\le j_1} \ell_{j_0} < 2\ell_{j_1}$.

(b) *(One external leg.) For an external field $\tilde{A}$ at z and a line of index j_0 to x,*

$$\Big|\sum_z \eta^3 G_0(x, z)c(z)g(z)\tilde{A}(z)\Big| \le K_\ell\Gamma_3 S(a/2)\ell_{j_0}\|\tilde{A}\|_\infty \mathrm{e}^{-\frac{a}{2}\operatorname{dist}(x,\operatorname{supp} g)},$$

and $\sum_{j_0\le k} \ell_{j_0} < 2$.

(c) *(Both legs external; then $j_0 = k$.) Per unit cube $\square$, $\big|\sum_{x\in\square} \eta^3 g\tilde{A}_\mu c_{\mu\mu'} g'\tilde{A}'_{\mu'}\big| \le \Gamma_3\|h\tilde{A}\|_{1,\alpha_0}\|h'\tilde{A}'\|_{1,\alpha_0}$.*

(d) *(First-moment vertex.) With a derivative on the j_1-leg, the relative factor is $K_\ell\Gamma'_3 S\,\ell_{j_0}^2/\ell_{j_1} \le K_\ell\Gamma'_3 S\,\ell_{j_0}$. With a derivative on the j_0-leg, it is $K_\ell\Gamma'_3 S\,\ell_{j_0}$.*

(e) *(Chains.) If N_v insertions occur in series on one path, the choice $\delta' = a/(2N_v)$ at each insertion keeps the decay rate $\ge a/2$, at the price of a factor $S(a/(2N_v))^{N_v}$.*

(f) *(Transported legs.) Let a renormalised subgraph H of internal scale ℓ_i, $i < j_0$, transport its leg G_0 to order $1 + \alpha$* [3, (3.26), p. 440]*. Then the composite of (a), with H contracted to its local vertex at scale j_0, is bounded by the bound of (a) times $C(\ell_i/\ell_{j_0})^\alpha$, and $\sum_{i<j_0}(\ell_i/\ell_{j_0})^\alpha \le (1 - L^{-\alpha})^{-1}$.*

Proof. (a) Since $\ell_{j_0} \le \ell_{j_1}$, we have $\mathrm{e}^{-a|x-z|/\ell_{j_0}} \le \mathrm{e}^{-(a-\delta')|x-z|/\ell_{j_1}} \mathrm{e}^{-\delta'|x-z|/\ell_{j_0}}$ and $\mathrm{e}^{-a|z-y|/\ell_{j_1}} \le \mathrm{e}^{-(a-\delta')|z-y|/\ell_{j_1}}$. By the triangle inequality, the dependence on y is bounded by $\mathrm{e}^{-(a-\delta')|x-y|/\ell_{j_1}}$. By Lemma E.131, the z-sum is at most $\ell_{j_0}^{-1}\Gamma_3\ell_{j_0}^{-1} S(\delta')\ell_{j_0}^3$. Finally, $\sum_{j_0\le j_1} \ell_{j_0} = \ell_{j_1} \sum_{n=0}^{j_1} 2^{-n} < 2\ell_{j_1}$.

(b) Since $\ell_{j_0} \le 1$, $\mathrm{e}^{-a|x-z|/\ell_{j_0}} \le \mathrm{e}^{-\frac{a}{2}|x-z|/\ell_{j_0}} \mathrm{e}^{-\frac{a}{2}|x-z|}$. Sum over z as in (a).

(c) Here $\ell_k = 1$, the cube has volume 1, and $\|\cdot\|_\infty \le \|\cdot\|_{1,\alpha_0}$.

(d) As in (a), with one more factor ℓ^{-1} from the differentiated line.

(e) Apply (a) N_v times.

(f) The remainder factor $|x' - x|^{1+\alpha}$ is absorbed by the lines of H, which give $\le C\ell_i^{1+\alpha}$ [3, p. 438]. By (LB), the quotient of order $1 + \alpha$ of G_0 costs $\ell_{j_0}^{-1-\alpha}$ and keeps the decay $\mathrm{e}^{-a|x-z|/\ell_{j_0}}$. So in the transported term the factor of G_0 relative to (LB) is $(C\ell_i)^{1+\alpha}\ell_{j_0}^{-1-\alpha}$. The subgraph H, of degree -1, contributes ℓ_i^{-1} in place of the local vertex factor $\ell_{j_0}^{-1}$. The product is $C(\ell_i/\ell_{j_0})^\alpha$ times the unmoved expression. The z-sum of (a) is unchanged, because the quotient keeps the decay of G_0, and the sum over i is geometric. □

Remark E.133 (two configurations not covered by Lemma E.132). (i) *Shared lines.* Let two two-photon vertices V_1, V_2 be joined by a line of index j_0 that lies below their other legs. A

bound by absolute values gives $e(t_k)^4\Gamma_3^2\ell_{j_0}^{-3}S(a)\ell_{j_0}^3 = e(t_k)^4\Gamma_3^2 S(a)$ per unit cube for each j_0, and the sum over $j_0 < k$ is of order k. Such families must therefore be bounded as whole classes (Lemma E.134 and Proposition E.140). (ii) *Transport charged to the composite.* Suppose the transport of Lemma E.132(f) is charged to the composite line of index j_1 instead of to G_0. Then the ratio to the required bound is $L^{n\alpha}\ell_{j_1}/(1-L^{-\alpha})$ with $n = j_1 - i$, and this ratio is unbounded at $j_1 = k$. This is why Lemma E.132(f) charges the gain to G_0.

Lemma E.134 (zeroth moment of a one-particle-reducible chain)**.** *Let Π^1, Π^2 be kernels with two vector legs and C_γ a vector line on $\xi\mathbb{Z}^3$, all with absolutely summable products. Then*

$$\sum_{x'}(\Pi^1 C_\gamma \Pi^2)(x,x') = \sum_{y,y'}\Pi^1(x,y)C_\gamma(y,y')\,m^{(2)}(y'), \qquad m^{(2)}(y') := \sum_{x'}\Pi^2(y',x').$$

Suppose in particular that every scalar line of Π^2 is the propagator C^ξ, and that Π^2 is the sum of the order-e^2 class with two vector legs at fixed labels. Then $m^{(2)} \equiv 0$ by [3, (3.29), p. 441]*, that is, by the Ward–Takahashi identity* [3, (2.26)] *with $M^2 = 1$. Hence the one-particle-reducible members of the order-e^4 class with two vector legs contribute nothing to the part of its zeroth moment in which all scalar lines are C^ξ.*

Proof. Use Fubini's theorem, then [3, (3.29)] at fixed labels. □

The external vector fields of the $A \cdot A$ vertices. Fix a step k with $1 \le k \le K-1$, put $t := t_k$, and work pointwise on the support of the small-field restrictions of [2, (2.55)] at scales k and $k+1$. As in [2, (2.80)–(2.89), pp. 574–575], A is the block field of scale k on $T_1^{(k)}$, B is the block field of scale $k+1$, and

$$A = A' + aL^{-2}C^{(k)}_{\Lambda_0^{(k)}}Q^*B, \qquad \theta_{k+1}A'^{(k)} = a_k\theta_{k+1}\zeta^{(k)}G_kQ_k^*A', \qquad \tilde{A} := \theta_{k+1}A'^{(k)} + \tilde{B}^{(k)}.$$

Here $(\zeta^{(k)}G_kQ_k^*)(x,y) = \zeta^{(k)}(x,y)(G_kQ_k^*)(x,y)$, and $\tilde{B}^{(k)}$ is the sum of the second, third and fourth terms of [2, (2.89)]. Everything else in $A^{(k)}$ is the background $\mathfrak{B}$. For a vector field f on the η-lattice put

$$\|f\|_{1,\alpha} := \sup_x|f(x)| + \sup_{x,\mu}|\partial_\mu f(x)| + [\partial f]_\alpha, \qquad [\partial f]_\alpha := \sup_{x\neq x',\mu}\frac{|\partial_\mu f(x') - \partial_\mu f(x)|}{|x-x'|^\alpha}.$$

This is the norm [3, (1.32)] for external vector fields (there with $\tilde{B} = 0$). We use the following properties of the construction of [1, 2].

(F1) On $\Lambda_0^{(k)\prime}$, $|\partial B| \le c_1 p(t)$ and $|B| \le c_1 p(t)/(\mu_0 t)$. For $x \in B(y)$ with $y \in \Lambda_1^{(k)\prime}$, $|A(x) - (Q^*B)(x)| \le c_2 p(t)$, and the three remainder terms of [2, (2.82)] are at most $c_3 p(t)$. These are the restrictions and the estimates of [2, (2.81)–(2.86)]. They involve the values of A and B only through differences, or multiplied by exponentially small kernels.

(F2) Let $W \in \{a_kG_kQ_k^*,\ a_{k+1}L^{-2}G_{k+1}^\eta Q_{k+1}^*\}$. There are $c_0, \delta_0 > 0$, independent of ε, k, ℓ_T, such that $|W(x,y)| + |\partial_x W(x,y)| \le c_0 \mathrm{e}^{-\delta_0|x-y|}$, and, for $|x-x'| \le 1$,

$$|\partial_x W(x',y) - \partial_x W(x,y)| \le c_0|x-x'|^{\alpha_0}\big(\mathrm{e}^{-\delta_0|x-y|} + \mathrm{e}^{-\delta_0|x'-y|}\big)$$

[1, Props. 2.1–2.3], [2, Prop. 2.2].

(F3) θ_{k+1}, $\zeta^{(k)}(\cdot,y)$ and $\zeta^{(k+1)}(\cdot,y)$ take values in $[0,1]$, and their first and second differences in x are bounded by c_θ.

(F4) If $\operatorname{dist}(x, \operatorname{supp}\theta_{k+1}) \le 2\eta$ and $\zeta^{(k)}(x,y) \neq 0$, then $y \in \Lambda_1^{(k)}$.

(F5) Each term of $\tilde{B}^{(k)}$ has the form $\theta_{k+1}\sum_y \chi(\cdot,y)W(\cdot,y)F(y)$, with $\chi \in \{\zeta^{(k)}, 1-\zeta^{(k)}, 1-\zeta^{(k+1)}\}$, W as in (F2), and $\mathrm{e}^{-\delta_0 d_F/2}M_F \le C_\kappa t^\kappa$ for every κ. Here M_F and d_F are defined in Lemma E.136. That (F5) holds is shown in Lemma E.144.

Lemma E.135 (product rule)**.** *Let $d = 3$ and $0 < \alpha < 1$, and let h be real with $|h| \le 1$ and with first and second differences bounded by c_h. Then $\|hf\|_{1,\alpha} \le (3+5c_h)\|f\|_{1,\alpha}$ for every vector field f.*

Proof. Write $\|\cdot\|$ for sup norms. We have $\partial_\mu(hf) = u\,\partial_\mu f + v\,f$ with $u := h(\cdot + \eta e_\mu)$ and $v := \partial_\mu h$. Hence $\|hf\| \le \|f\|$ and $\|\partial(hf)\| \le \|\partial f\| + c_h\|f\|$.

Let w be a lattice function with $\|\partial w\| \le c$. Joining x to x' by a monotone lattice path gives $|w(x') - w(x)| \le \min(2\|w\|, \sqrt{3}\,c|x - x'|)$. Treating the cases $|x - x'| \le 1$ and $|x - x'| > 1$ separately, this is $\le \max(2\|w\|, \sqrt{3}\,c)|x - x'|^\alpha$.

Put $g := \partial_\mu(hf)$ and write

$$g(x') - g(x) = u(x')\big(\partial f(x') - \partial f(x)\big) + \partial f(x)\big(u(x') - u(x)\big) + v(x')\big(f(x') - f(x)\big) + f(x)\big(v(x') - v(x)\big).$$

This gives $[g]_\alpha \le [\partial f]_\alpha + (2 + \sqrt{3}c_h)\|\partial f\| + c_h(2\|f\| + \sqrt{3}\|\partial f\|) + 2c_h\|f\|$. Adding the three parts of the norm gives $\|hf\|_{1,\alpha} \le (1 + 5c_h)\|f\| + (3 + 2\sqrt{3}c_h)\|\partial f\| + [\partial f]_\alpha$, which implies the claim. □

Lemma E.136 (cut-off kernels)**.** *Let θ, χ satisfy (F3), let W satisfy (F2), and let F be a function on the unit lattice. Put $f(x) := \theta(x)\sum_y \chi(x,y)W(x,y)F(y)$. Let Y be the set of y such that $\chi(\hat{x}, y) \ne 0$ for some $\hat{x}$ with $\operatorname{dist}(\hat{x}, \operatorname{supp}\theta) \le 2\eta$. Put $M_F := \sup_Y |F|$ and*

$$d_F := \inf\{|\hat{x} - y| :\ \operatorname{dist}(\hat{x}, \operatorname{supp}\theta) \le 2\eta,\ \chi(\hat{x}, y) \ne 0\}.$$

Then $\|f\|_{1,\alpha_0} \le C_K\, \mathrm{e}^{-\delta_0 d_F/2} M_F$, where C_K depends only on c_0, δ_0, c_θ.

Proof. Let $K(x,y) := \theta(x)\chi(x,y)W(x,y)$. By the Leibniz rule,

$$\partial_x K = \theta(\cdot + \eta e)\chi(\cdot + \eta e, y)\partial W + \theta(\cdot + \eta e)(\partial\chi)W + (\partial\theta)\chi W,$$

so $|\partial_x K(x,y)| \le (1 + 2c_\theta)c_0 \mathrm{e}^{-\delta_0|x-y|}$. In each term, every factor is of one of two kinds. Either it is bounded and Lipschitz in x, with constants depending on c_θ, c_0, δ_0 (for W this follows from (F2), along a path of length $\le \sqrt{3}|x - x'|$). Or it is α_0-Hölder in x for $|x - x'| \le 1$; this is the factor ∂W. Hence, for $|x - x'| \le 1$,

$$|\partial_x K(x', y) - \partial_x K(x, y)| \le C|x - x'|^{\alpha_0}\big(\mathrm{e}^{-\delta_0|x-y|} + \mathrm{e}^{-\delta_0|x'-y|}\big).$$

For $|x - x'| > 1$ we bound the difference by twice the supremum of ∂f.

Only $y \in Y$ contribute. For these, $\min(|x - y|, |x' - y|) \ge d_F - 2\eta$ whenever the summand is non-zero. So $\mathrm{e}^{-\delta_0|x-y|} \le \mathrm{e}^{3\delta_0}\mathrm{e}^{-\delta_0 d_F/2}\mathrm{e}^{-\delta_0|x-y|/2}$, and similarly at x'. Summing over y, with $S_0 := \sup_x \sum_y \mathrm{e}^{-\delta_0|x-y|/2} < \infty$, gives the claim. □

Lemma E.137 (the translation removes the constant mode)**.** *Assume (F1). Then, for $x \in B(y)$ and $y \in \Lambda_1^{(k)\prime}$,*

$$|A'(x)| \le C_1 p(t), \qquad C_1 := c_2 + c_3 + a^{-1}L^2 c_1\,\mu_0\varepsilon_0.$$

The constant C_1 does not depend on the values of A and B; in particular it does not depend on their constant parts.

Proof. By [2, (2.82)], $aL^{-2}(C^{(k)}_{\Lambda_0^{(k)}}Q^*B)(x) = aL^{-2}(C^{(k)}Q^*1)(x)\,B(y) + \mathcal{R}(x)$, where $|\mathcal{R}(x)| \le c_3 p(t)$ by (F1). By [2, (2.83)–(2.85)], $aL^{-2}C^{(k)}Q^*1 = (1 + X)^{-1}$ with

$$0 \le X = a^{-1}L^2 a_k \mu_0^2 t^2/(a_k + \mu_0^2 t^2) \le a^{-1}L^2\mu_0^2 t^2,$$

so $|aL^{-2}C^{(k)}Q^*1 - 1| \le a^{-1}L^2\mu_0^2 t^2$. Since $(Q^*B)(x) = B(y)$,

$$A'(x) = \big(A(x) - (Q^*B)(x)\big) - \big(aL^{-2}C^{(k)}Q^*1 - 1\big)B(y) - \mathcal{R}(x).$$

The bound $|B| \le c_1 p(t)/(\mu_0 t)$ bounds the middle term by $a^{-1}L^2 c_1\mu_0 t\, p(t) \le a^{-1}L^2 c_1\mu_0\varepsilon_0\, p(t)$. □

Proposition E.138 (size of the external vector fields)**.** *Assume (F1)–(F5), and let ε_0 be so small that $C_\kappa t^\kappa \le p(t)$ for $t \le \varepsilon_0$, at one fixed κ. Then, for every localisation function h' of* [3, p. 420] *whose first and second differences are bounded by c_h,*

$$\|h'\tilde{A}\|_{1,\alpha_0} \le (3 + 5c_h)\big(C_K C_1 + 3C_K\big)\, p(t).$$

The constant depends only on $d, a, L, c_0, \delta_0, c_\theta, c_1, c_2, c_3, \mu_0\varepsilon_0, c_h$. It does not depend on ε, k, ℓ_T, the domains $\Omega, \Omega_1, \Omega_2$, or the constant parts of A and B. In particular, the hypothesis of [3, p. 412] *on the external vector fields holds uniformly in the photon zero mode and in the twist on the torus.*

Proof. For $\theta_{k+1}A'^{(k)}$, apply Lemma E.136 with $\chi = \zeta^{(k)}$, $W = a_k G_k Q_k^*$ and $F = A'$. By (F4), $Y \subset \Lambda_1^{(k)}$, so $M_F \le C_1 p(t)$ by Lemma E.137; also $d_F \ge 0$. Hence $\|\theta_{k+1}A'^{(k)}\|_{1,\alpha_0} \le C_K C_1 p(t)$. For each of the three terms of $\tilde{B}^{(k)}$, Lemma E.136 and (F5) give $\|\cdot\|_{1,\alpha_0} \le C_K C_\kappa t^\kappa \le C_K p(t)$. Lemma E.135 with $h = h'$ concludes.

The bound involves only three kinds of input: the vector operators $G_k, Q_k, C^{(k)}_{\Lambda_0^{(k)}}, G_{k+1}, Q_{k+1}$, which do not depend on any background field; the fixed cut-offs; and the bounds (F1), which see the values of A and B only through $|B| \le c_1 p/(\mu_0 t)$. So the bound is the same for every constant part. □

Remark E.139 (where the constant mode goes). (i) By Lemma E.137, the constant part of the block fields, including the constant photon mode of the torus, is carried by the background $\mathfrak{B}$; only $O(\mu_0^2 t^2)|B| = O(\mu_0 t\, p)$ of it reaches A'. Hence no vertex of [3, (3.30), (3.32)] carries the torus constant on an external leg. For the background, [3, p. 412] asks only for bounds on $\partial^\eta \mathfrak{B}$ and on its Hölder quotient, and these do not change under $\mathfrak{B} \mapsto \mathfrak{B} + c$ with c constant. A constant c' with $|c'| \le \|\tilde{A}\|_\infty$ can be moved from $\tilde{A}$ to the background by passing to $(\mathfrak{B} + \theta_{k+1}c',\ \tilde{A} - \theta_{k+1}c')$. This keeps $\operatorname{supp}\tilde{A} \subset \operatorname{supp}\theta_{k+1}$, and by (F3) it changes all the bounds above by at most a factor depending on c_θ. (ii) Proposition E.138 does not use the statement that the constant $O(1)$ of [3, Prop. 1] is uniform under $\mathfrak{B} \mapsto \mathfrak{B} + c$ on T_η. That statement is Statement K.29. (iii) At $L = 3$, $\tilde{B}^{(k)}$ has a fourth term of the same form, (F5) holds for it, and the proof is unchanged. At $k = 0$ the same argument applies to [2, (2.19)–(2.22)].

Proposition E.140 (the two-photon vertices)**.** *Fix $\mu_* > 0$ and $(u, M) \in \{(1, m_2), (-\eta, m_1)\}$. Assume (LB) and (VB). Assume that the family bounds of Definition E.59 and Proposition E.60 hold for families of graded order $\le \bar{N}$. Assume also the following two hypotheses on the frame of the expansion on T_ε.*

- (ZM) *The constant mode c of the photon field is carried as a constant background in $\tilde{B}$, equivalently as a twist $\vartheta = ec\ell_T$ of the scalar boundary condition. Every bound on scalar operators used in* [3, Prop. 1] *is uniform in ϑ, provided the localisation cubes do not wrap the torus.*
- (EXT) *The external vector fields of the families satisfy $|\tilde{A}|, \|h'\tilde{A}\|_{1,\alpha_0} \le O(1)p(t_k)$, with the constant mode removed as in (ZM). This is the hypothesis of* [3, p. 412].

Then the vertices of [3, (3.30), (3.32)] *satisfy (N1)–(N4) of Lemma E.130, uniformly in ε, k, ℓ_T.*

Proof. (N1). The covariances of the expansion are fixed before any graph is formed: "A'_j are independent Gaussian fields with covariances $C^{(j),L^j\eta}$ … the field A' has the covariance G_k. The basic propagator for the scalar field ϕ' is $G_k(\Omega, \tilde{B})$" [3, p. 414]. Moreover, "all the A'-legs are contracted, i.e. they are divided into pairs and each pair is replaced by the corresponding propagator" (ibid.). The vertex list [3, (1.6)–(1.15), pp. 413–414] contains no two-photon term with a coefficient proportional to $\ell_{j_0}^{-1}$. The vertices of (3.30) and (3.32) arise only on p. 442, where the values of subgraph classes are rewritten inside the estimate. So they enter no covariance and no Gaussian weight.

(N2). Every renormalised family is bounded by absolute values [3, p. 426], using the counting (2.13)–(2.16). By [3, Prop. 2.2, p. 428], this counting needs only lines with representations (2.6) and bounds (2.10)–(2.12) ("lines with the same exponential factors but with arbitrary dimensions"),

and vertices whose coefficients are bounded times powers of the scale. The two-photon vertex is the only vertex whose coefficient carries a negative power $\ell_{j_0}^{-1}$ of a line scale. There are two cases.

- *The lowest-index line is unshared.* Lemma E.132(a),(d),(e),(f) replaces the vertex, together with that line, by a generalised line with an extra positive power ℓ_{j_0}, which is summable over j_0. If one or both legs are external, Lemma E.132(b),(c) replaces the vertex by a bounded smeared external leg, respectively by a bounded local vertex. These steps use $\|\tilde{A}\|_\infty \le \|h'\tilde{A}\|_{1,\alpha_0} \le O(1)p$, which is (EXT). By (ZM), the constant mode is not a vector leg; it enters only through scalar operators, uniformly in ϑ. So the family obeys $(1.33)^\natural$ with $O(1)$ replaced by $O(1)(1+C_{\mathrm{INS}})^{N_v}$, where
$$C_{\mathrm{INS}} := 2K_\ell^2 \max(\Gamma_3, \Gamma_3', \Gamma_4)(1-2^{-\alpha})^{-1} S(a/(2N_v))^{N_v}$$
and $N_v \le N(\bar{N})$.
- *The lowest-index line is shared* (Remark E.133(i)). Then the family is a one-particle-reducible or nested member of the class of order e^4 (or $e^2\lambda$) with two external vector legs, and its lowest external index m exceeds the index of the shared line. Such members are treated in [3] as whole graphs: "we resum over proper orderings and j-indices, we replace the propagators $G_{j_0}(0)$, G_{j_0} by C^ξ ... we apply a proper Ward–Takahashi identity" [3, p. 442]. Only the last divergent subgraph of an ordering is decomposed, and the remaining divergent graphs are the whole graphs, to which (3.31) applies [3, p. 445]. So the index of the shared line is summed inside the resummed propagators at index m, and no sum over j_0 of terms of size $O(1)$ arises. By (3.31), the family is the sum of a generalised graph of degree α and a zeroth-moment term. The coefficient of the zeroth-moment term is bounded by Lemma E.32, and its pure-C^ξ part vanishes; for the one-particle-reducible part this is already Lemma E.134. The inner transformations are exact identities on the set of orderings in which the inner subgraph is the last divergent one, so the whole-class sums are unchanged.

In both cases the bound is uniform in ε, k, ℓ_T and finite at both points (u, M). The only dependence on u is the factor $(|u|c_k)^m$ of the coloured lines. The only dependence on M is through $m_{j_0}^2 = M^2 t_{j_0}^2 \le 1$ in Lemmas E.24 and E.32.

(N3). The interaction after k steps is given by formula (1.5) of [3, p. 412], built from the original vertices (1.6)–(1.15). By (N1) the two-photon vertices are not among them. So they are created anew inside the estimate of each step and are never carried to step $k+1$ as a coupling. Hence no factor L^2 per step can accumulate, and their weight per step is bounded through (N2).

(N4). (3.30)–(3.32) are identities between values of classes [3, p. 442], so the total of a family is unchanged. The mass counterterm is defined from complete scalar classes (Lemma C.1), and inserting these vertices into a scalar two-leg family changes neither the family nor its zeroth moment. The photon has no mass counterterm. Contracting the two legs of such a vertex with each other gives a vacuum graph, which belongs to the vacuum classes of Proposition E.175. The torus counterterm is built with the torus propagator [3, (1.28)]. Its difference from the members on $\eta\mathbb{Z}^3$ after [3, p. 433] is an image term of order $\mathrm{e}^{-\delta\ell_T/t}$. This term enters as a finite vertex of type [3, (1.7)], not as a new counterterm. □

Proof of Lemma E.130. We apply Proposition E.140 and verify its hypotheses.

- (LB) is [3, (2.10)–(2.12), p. 426], together with Theorem E.47 for the vector lines.
- (VB) is Lemmas E.24 and E.32.
- The family bounds hold at step k by Definition E.59 and Proposition E.60, through the bound (1.33) of Proposition E.110; that bound does not use the present lemma.
- (EXT) is Proposition E.138. Its hypothesis (F5) holds by Lemma E.144, and its smallness condition on ε_0 holds since $C_\kappa t^\kappa / p(t) \to 0$ as $t \to 0$.

- In (ZM), the first part (the constant mode is carried by the background) is Lemma E.137 and Remark E.139(i). The localisation cubes do not wrap the torus, by (T.nw) and Lemmas E.66 and E.65. The uniformity in ϑ is Statement K.29.

□

Lemma E.141 (the tails of [2, p. 579])**.** *Pointwise on the small-field support, at step k,*

$$(\textit{basic quadratic form} + \mathcal{P}_w^{(k)}) = (\textit{polynomial in } \theta_{k+1}A'^{(k)}, \psi, \phi \textit{ of order} \le \bar{n}) + \Delta_k^{\mathrm{pre}} + \Delta_k^R + \Delta_k^{\tilde{B}},$$

with the following two-sided bounds:

- $|\Delta_k^{\mathrm{pre}}| \le C t_k^{7/2} \operatorname{polylog} |\Lambda_2^{(k)}|_k$ *(the terms of order $> \bar{n}$, cut before integration);*
- $|\Delta_k^{R}| \le C t_k^{13/2} \operatorname{polylog} |\Lambda_2^{(k)}|_k$ *(the terms with Taylor remainders $R_{\bar{n}+1}$);*
- $|\Delta_k^{\tilde{B}}| \le C t_k^{9/2} \operatorname{polylog} |\Lambda_2^{(k)}|_k$ *at $\kappa = 31$ (the terms with $\tilde{B}^{(k)}$, [2, (2.90)]).*

Each of these is $\le C' t_k^{16/5} N_k$. The constants do not depend on ε, k or ℓ_T. They are finite at $u = -\eta$ and at $M = m_1$. Where $\delta m^2_{\mathrm{fin}}$ vertices occur, they are evaluated at the torus floor ℓ_0. The expansion of the logarithm of the determinant [1, (3.46)–(3.47)] *is not part of this cut; it is cut at* [2, pp. 577–578].

The inputs are those of Proposition E.36:

- the kernels of [4] and Lemma D.26;
- Lemma E.117(a), and Lemma E.118 on Neumann domains under (R.1) and (R.8);
- Lemma D.24;
- the regularity of the background under the conditions of Definition A.1;
- the cap $\bar{N} = 169$, with (R.cap) and (T.cap);
- the per-order bound of Lemma E.142 below;
- (E.10), and $R \ge R^{**}$.

Proposition E.76 uses Proposition E.110 only through item 7 of Table 10, and item 7 does not use item 17, so there is no cycle.

The remainder and $\tilde{B}$ parts. The setting, the fields $\mathfrak{B}$ and $\tilde{A}$, the bound $\sigma \le C_\sigma t^{1/2} p$, the number N_k and the inputs (K1)–(K4), (REG), $(\tilde{B})$, (IH), (CAP) are those of Proposition E.36. The expansion (E1)–(E3) is that of its proof; here $\bar{n} = 13$ and $\bar{N} = 169$. We write $t = t_k$ and $p = p(t) = b_0(1 + \ln t^{-1})^{p_0}$. "Polylog" means a factor bounded by $C(1 + \ln t^{-1})^c$, with constants independent of ε, k, ℓ_T. The order of a term is $\alpha + \beta$ for a factor $e^\alpha \lambda^\beta$, and each factor $e_k \tilde{A}$ counts 1. On $\operatorname{supp} \chi_k$, scalar external fields obey $\|h\Phi_{\mathrm{ext}}\|_{1,\alpha_0} \le Cp\lambda(t)^{-1/4}$, and vector ones obey $\|h' A_{\mathrm{ext}}\|_{1,\alpha_0} \le Cp$ (Proposition E.138). The mass vertices satisfy

$$(\delta m) \qquad |\delta m^2_{K,k}| t^2 \le O(1)\big[e(t)^2 + \lambda(t) + (e(t)^4 + e(t)^2\lambda(t) + \lambda(t)^2)(1 + K - k)\big],$$

with $K - k \le \log_2(\varepsilon_0/t)$, $e(t) = et^{1/2}$ and $\lambda(t) = \lambda t$.

Lemma E.142 (per-order power count)**.** *Consider a connected family with at least two vertices. After its external legs are bounded as above, every elementary vertex of* [3, (1.6)–(1.15), (1.7), (1.18)] *of order q_v contributes at most $Ct^{q_v/4}$ times a polylog. Hence a family of total order q is $\le C(q) t^{q/4} p^{4q} \mathrm{e}^{-\delta(\bar{N})d}$.*

Proof. Every vertex has at least one internal leg. We go through the vertices.

- $\lambda|\phi|^4$ (order 1) has at most three external scalar legs, which gives $\lambda t \cdot t^{-3/4} = t^{1/4}$.
- $e^2 A^2 |\phi|^2$ (order 2) with two external scalar legs gives $e^2 t \cdot t^{-1/2} = t^{1/2}$.
- $e\bar{\phi} A \cdot D\phi$ (order 1) gives $O(t^{(4-d)/4} p^2) = O(t^{1/4} p^2)$ by [3, p. 421], via [2, Lemma 2.4].
- For the mass vertex with two external scalar legs, (δm) gives $t^{1/2}$ for the λ-part (order 1), $t^{1/2}$ for the e^2-part (order 2), and $t^{3/2}(1 + K - k)$, that is $t^{3/2}$ times a polylog, for the parts of order 4.

- The vertices (1.18) with $n+n' \geq 2$ give $t^{(n+n')/2-1/2} \leq t^{(n+n')/4}$.
- The remaining vertices are better [3, p. 421].

Internal lines and integrated fluctuations contribute to the $O(1)$ of $(1.33)^\natural$, by (IH) and (CAP). Multiplying over the vertices gives the claim. □

Lemma E.143 (Taylor-remainder factors)**.** *Let $R_{n+1}(z) := \mathrm{e}^{iz} - \sum_{r\leq n}(iz)^r/r!$, and let $\tilde{A}$ be real. Then $|R_{n+1}(z)| \leq |z|^{n+1}/(n+1)!$, $|R'_{n+1}(z)| \leq |z|^n/n!$ and $|R''_{n+1}(z)| \leq |z|^{n-1}/(n-1)!$. Assume that the cut-offs θ_{k+1}, ζ have bounded second differences and that $[\partial^\eta \tilde{A}]_\alpha \leq Cp$ (the Hölder clause of (K1); see Proposition E.138). Consider each factor $R_{\bar{n}+1}(z)$ with $z = qe_k\tilde{A}(\Gamma)$ for a contour of length $O(1)$, and each bond factor $\eta^{-1}R_{\bar{n}+1}(\eta q e_k \tilde{A}(b))$. As a coefficient or as an external field, each such factor satisfies*

$$\|\cdot\|_{1,\alpha_0} \leq c_R(Ct^{1/2}p)^{\bar{n}+1}, \qquad c_R = c_R(\bar{n}, d, L).$$

Proof. We have $R'_{n+1} = iR_n$ and $R''_{n+1} = -R_{n-1}$, and $|R_m(z)| \leq |z|^m/m!$ for real z by Taylor's formula with integral remainder. On $\operatorname{supp}\chi_k$, $|z| \leq Ce_k\|\tilde{A}\|_\infty \leq C'_\sigma t^{1/2}p$ and $|\partial^\eta z| \leq Ce_k\|\partial^\eta\tilde{A}\|_\infty \leq C'_\sigma t^{1/2}p$. The lattice difference of $R(z(\cdot))$ is $\int_0^1 R'(z_s)\,ds\cdot\partial^\eta z$, where z_s interpolates between the two values. Its Hölder difference over $|x-\tilde{x}|\leq 1$ is bounded by

$$\sup|R''|\,|z(\tilde{x})-z(x)|\,|\partial^\eta z| + \sup|R'|\,|\partial^\eta z(\tilde{x}) - \partial^\eta z(x)| \leq \big[|z|^{\bar{n}-1}(Ct^{1/2}p)^2 + |z|^{\bar{n}}e_k[\partial^\eta\tilde{A}]_\alpha\big]|x-\tilde{x}|^\alpha,$$

where $|x-\tilde{x}| \leq |x-\tilde{x}|^\alpha$ was used. This is $\leq (Ct^{1/2}p)^{\bar{n}+1}|x-\tilde{x}|^\alpha$. For the bond factor, each estimate carries an extra factor $\eta^{\bar{n}} \leq 1$. □

Lemma E.144 ($\tilde{B}^{(k)}$ in the norm (1.32))**.** *By* [2, (2.89), p. 575], $\tilde{B}^{(k)} = \sum_{i=2}^4 \theta_{k+1}\zeta_i u_i$, *with*

$$\zeta_2 u_2 = a_k a L^{-2}\zeta^{(k)}G_k Q_k^* \delta C^{(k)}_{\Lambda_0^{(k)}} Q^* B, \qquad \zeta_3 u_3 = -a_k a L^{-2}(1-\zeta^{(k)})G_k Q_k^* C^{(k)} Q^* \Lambda_0^{(k)\prime} B,$$

$$\zeta_4 u_4 = a_{k+1}L^{-2}(1-\zeta^{(k+1)})G^\eta_{k+1}Q^*_{k+1}\Lambda_0^{(k)\prime}B.$$

Assume that $\theta_{k+1}, \zeta^{(k)}, \zeta^{(k+1)}$ take values in $[0,1]$ and have first and second lattice differences $\leq c_\theta$. Assume further that, on the support of $\theta_{k+1}\zeta_i$, for every κ,

$$(\tilde{B}_0) \qquad |u_i| + |\partial^\eta u_i| + [\partial^\eta u_i]_{\alpha_0} \leq C_\kappa t^\kappa.$$

Then $\|h\tilde{B}^{(k)}\|_{1,\alpha_0} \leq C(c_\theta)C_\kappa t^\kappa$ for every κ.

Hypothesis $(\tilde{B}_0)$ holds, and so does (F5) of Subsection E.7. The sup and derivative parts of $(\tilde{B}_0)$ are [2, (2.90)] *("$\tilde{B}^{(k)}(x) = O((L^k\varepsilon)^\kappa)$ for every κ, and similar estimates for the derivative"). For the remaining part: each u_i is a kernel of $G_kQ_k^*$ (respectively $G^\eta_{k+1}Q^*_{k+1}$) applied to a source of size $\leq c_1 p/(\mu_0 t)$ supported at distance $\geq r(t)/2 - 2M_b$ from $\operatorname{supp}(\theta_{k+1}\zeta_i)$. The decay, derivative and Hölder clauses of (K1) carry the same factor $\mathrm{e}^{-\delta_0(r(t)/2-2M_b)}$. Since $r(t) = R(1+\ln t^{-1})^2$, this factor is $\leq C_\kappa t^{\kappa+1}$ for every κ.*

Proof. $\partial^\eta(\theta\zeta u)$ is the sum of the three terms $(\partial^\eta\theta)\zeta u$, $\theta(\partial^\eta\zeta)u$ and $\theta\zeta\,\partial^\eta u$, with shifted arguments. Their Hölder quotients over $|x-\tilde{x}|\leq 1$ are bounded using $(\tilde{B}_0)$ and the Lipschitz bounds on $\partial^\eta\theta$ and $\partial^\eta\zeta$, which follow from the bound c_θ on the second differences. □

Proposition E.145 (the Taylor-remainder cut)**.** *In the expansion (E1)–(E3), let Δ_k^R be the sum of the terms that contain at least one factor $R_{\bar{n}+1}$: in the bracket V_k, in $Q_k(\mathfrak{B}+\tilde{A})$, or in a transport of a vertex of $\mathcal{P}_w^{(k)}$. Then, pointwise on $\operatorname{supp}\chi_k$ and for $0\leq k\leq K-1$,*

$$|\Delta_k^R| \leq C_R t_k^{13/2}\operatorname{polylog}|\Lambda_2^{(k)}|_k \leq C'_R t_k^{16/5}N_k.$$

The constant C_R depends on $\bar{n}, \bar{N}, d, L, a, M_b, e, \lambda, \mu_*, b_0, p_0, c_1, R, \delta_0, \eta$, and not on ε, k or ℓ_T. It is finite at $u=-\eta$ and at $M=m_1$, with the same dependence on u and M as in Proposition E.36.

Proof. (1) Each R-factor obeys Lemma E.143.

(2) *An R-factor in V_k.* By the lattice Leibniz rule, the piece V^R has the right-pushed form $R_0 + \sum_\mu R_{1,\mu} D_\mu$ of Lemma E.115(ii), with coefficients $\le c_V (Ct^{1/2}p)^{\bar n+1}$. Consider a chain with one V^R-vertex and further vertices of orders r_i. As in Step 2 of the proof of Proposition E.36, it is bounded by a product of per-vertex factors, namely by $C(Ct^{1/2}p)^{\bar n+1+\sum r_i}\|f\|_\infty \|f'\|_\infty \mathrm{e}^{-\beta\,\mathrm{dist}}$. With end fields $\le (Cp\lambda^{-1/4}t^{-1/4})^2$, this is $\le Ct^{(\bar n+1)/2-1/2} = Ct^{13/2}$ times a polylog. Inside a line of a family of $\mathcal{P}_w^{(k)}$, the same chain is an external two-variable field of [3, (2.5)]. It is bounded by (K3) (Lemmas E.117 and E.118) with one coefficient replaced, and the family is then bounded by (IH) and (CAP).

(3) *An R-factor in a vertex transport.* $(1.33)^\natural$ is linear in the norm of each external field. So the family receives the factor $c_R(Ct^{1/2}p)^{\bar n+1}$, that is t^7 times a polylog. At most two scalar legs sit at that vertex, and they give $t^{-1/2}$. By Lemma E.142, each of the other vertices gives $\ge t^{q_v/4}$ times a polylog. The total is $\le Ct^{13/2}$ times a polylog times $\mathrm{e}^{-\delta(\bar N)d}$.

(4) There are finitely many terms, each with at most $D(\bar n)$ factors. The localisations through a fixed block of $\Lambda_2^{(k)}$ sum to $\le S(\bar N, d)$, and $|\Lambda_2^{(k)}|_k \le N_k$. Finally, $\sup_{t\in(0,1]} t^{13/2-16/5}\,\mathrm{polylog} < \infty$. The bound is two-sided, because the expansion is an exact identity and every term is bounded in absolute value. □

Proposition E.146 (the $\tilde B$ cut)**.** *In every polynomial term of order $\le \bar n$ of the expansion (E1)–(E3), split $\tilde A = \theta_{k+1}A'^{(k)} + \tilde B^{(k)}$ multilinearly. Terms of higher order are part of Δ_k^{pre} in Proposition E.36, with $\tilde B^{(k)}$ included in $\tilde A$. Let $\Delta_k^{\tilde B}$ be the sum of the terms with at least one factor $\tilde B^{(k)}$. Let $s(\bar n)$ be the largest number of scalar legs of a term of order $\le \bar n$. Every elementary vertex of* [3, (1.6)–(1.15)] *has order ≥ 1 and at most four scalar legs, so $s(\bar n) \le 8\bar n$. Put $\kappa := \lceil s(\bar n)/4\rceil + 5$; this is 31 when $s(\bar n) = 8\bar n$. Then, pointwise on* $\operatorname{supp}\chi_k$*,*

$$|\Delta_k^{\tilde B}| \le C_{\tilde B} t_k^{9/2}\,\mathrm{polylog}\,|\Lambda_2^{(k)}|_k \le C'_{\tilde B} t_k^{16/5} N_k.$$

The constant $C_{\tilde B}$ depends on $\bar n, \bar N, \kappa, \mu_*, c_1, b_0, p_0, R, \delta_0, M_b, \alpha_0, e, \lambda, \eta, d, L, a$, and not on ε, k or ℓ_T.

Proof. (1) By Lemma E.144, $\|h\tilde B^{(k)}\|_{1,\alpha_0} \le C'_\kappa t^\kappa$.

(2) In a term with a factor $\tilde B^{(k)}$, replace one vector-leg bound Cp by $C'_\kappa t^\kappa$, and bound every other factor without using any gain. The factors $e_k\tilde A$ are $\le Ct^{1/2}p$, which is at most a polylog. The end fields are $\le (Cp\lambda^{-1/4}t^{-1/4})^2$. By (IH) and (CAP), the families obey $(1.33)^\natural$, since their graded order is $\le 3\bar n \le \bar N$. They have at most $s(\bar n)$ scalar legs, which contribute $\le t^{-s(\bar n)/4}$ times a polylog. Hence every such term is $\le C(\bar n)C'_\kappa t^{\kappa - s(\bar n)/4 - 1/2}\mathrm{e}^{-\delta d}$ times a polylog, and $\kappa - s(\bar n)/4 - \frac12 \ge \frac92$. A factor $\tilde B$ inside a V of a line is covered by the same multilinear count, through the product structure of Step 2 of the proof of Proposition E.36.

(3) There are at most $2^{D(\bar n)}$ splittings per term. The localisation sum and the volume are treated as in Proposition E.145. Finally, $\sup_t t^{9/2-16/5}\,\mathrm{polylog} < \infty$. □

Remark E.147 (scope of the three cuts). (i) Propositions E.36, E.145 and E.146 do not cover two operations: the propagator change $G_k(B^k(\Lambda_0^{(k)}), \tilde B) \to G_k(B^k(\Lambda_2^{(k)}), B^{(k+1),\eta})$ at the end of [2, p. 579] ([2, Prop. 2.6]), and the cuts of the determinant expansions [2, (2.103)–(2.104), p. 578]. (ii) Through (IH) and (CAP), the three cuts depend on the two-domain bounds (Proposition D.17, Lemma E.123) and on the □-remainder of [3, p. 433], but only inside the families of [3, Prop. 1]. No domain difference of an operator (1.16) occurs, because all lines of a given term live on one domain. So this dependence enters Proposition E.110 once, through items 2–10 of Table 10.

Proof of Lemma E.141. The decomposition is the exact expansion (E1)–(E3) of the proof of Proposition E.36, in which $\tilde A = \theta_{k+1}A'^{(k)} + \tilde B^{(k)}$ is split multilinearly in the terms of order $\le \bar n$ (Proposition E.146).

- The bound on Δ_k^{pre} is Proposition E.36; its per-order input is Lemma E.142.
- The bound on Δ_k^{R} is Proposition E.145.
- The bound on $\Delta_k^{\tilde B}$ is Proposition E.146, at $\kappa = 31$.

The hypotheses on $\tilde A$ and on $\tilde B^{(k)}$ used in Lemmas E.143 and E.144 are supplied by Proposition E.138 and by Lemma E.144 itself. The two-domain bounds enter only inside the families (Remark E.147(ii)). They hold at every $k \geq 1$, including $k = 1$, where $\eta = \frac{1}{2}$, by Lemma E.108(d). Each of the three bounds is two-sided, is $\leq C' t_k^{16/5} N_k$, and has the stated dependence of its constant. □

E.8. **The mass reading, the final c-numbers and the vacuum classes at the last scale.** We prove Propositions E.159, E.162 and E.175 of Table 1; the region entropy is Proposition 5.21. Throughout: $d = 3$; Bałaban's periodic lattice torus T of side ℓ_T, $|T| = \ell_T^d$; $L = 2$; $\varepsilon = 2^{-K}\varepsilon_0$, $t_k := L^k\varepsilon$ ($t_K = \varepsilon_0$); $\ell_k := 1 + \ln t_k^{-1}$; $r_k := r(t_k) = R\ell_k^{\varrho}$, $\varrho = 2$; $p(t) = b_0(1 + \ln t^{-1})^p$, $p = 6$; M_b is Bałaban's block parameter.
The mass bound "$m^2 \leq O(1)$" of [2, p. 589]. [2, Prop. 3.1, p. 589] ends: "We assume $m^2 \leq O(1)$ also." (read at the page). Write $\nu_k := M^2 t_k^2$ for the rescaled Higgs mass at scale k, $\lambda(t) := \lambda t^{4-d}$, $a_\flat := a(1 - L^{-2}) < a_k \leq a$ ($a_\flat = 3/4$), and c_1 for the constant of [2, (2.55)]. Since $t_k \leq \varepsilon_0$, $\sup_{k \leq K} \nu_k \leq M^2\varepsilon_0^2$.

Lemma E.148 (where no mass bound is needed)**.** *(i)* [4, Theorem (1.9)–(1.12), p. 573] *(=*[1, Prop. 2.1]*) holds for every $\nu \geq 0$, with constants depending on d, M_b (c_0 also on α). (ii) For $0 \leq \nu \leq \nu'$: $\Delta_0 \leq \Delta_\nu \leq \Delta_{\nu'} \leq a_k I$, where $\Delta_\nu := a_k - a_k^2 Q_k G_\nu Q_k^*$ and $G_\nu := (-\Delta_{A,\Omega}^{\eta,N} + \nu + a_k P_k(A))^{-1}$. Hence* [1, (2.33), p. 611] *holds for every $\nu \geq 0$, with its $\nu = 0$ lower constant and upper constant $a(1 + L^{-2}) = 5/4$.*

Proof of (ii). $X_\nu \leq X_{\nu'}$ as forms, and $X_0 > 0$ by [4, (1.8)]. Operator monotonicity of the inverse gives $G_{\nu'} \leq G_\nu$, and $G_\nu \geq 0$ gives $\Delta_\nu \leq a_k \leq a$. Finally $\|P\| \leq 1$. □

Lemma E.149 (Prop. 3.1 of [2] with explicit constant)**.** *Let $0 \leq \nu \leq c_m$ and $\gamma_0(c_m) := \frac{1}{2}\min\{\gamma_0^{(0)}, a_\flat/(a_\flat + c_m)\}$, where $\gamma_0^{(0)}$ is the constant of* [4, (1.22), p. 574] *at $m^2 = 0$. Under the hypotheses of that proposition, $\langle\varphi, \Delta_\nu\varphi\rangle \geq \gamma_0(c_m)\big(\sum_{\langle x,x'\rangle}|U(A)\varphi(x') - \varphi(x)|^2 + \nu\sum_x|\varphi(x)|^2\big) - \frac{1}{2}O(1)\mathrm{e}^{2-\alpha}\sum_x|\varphi(x)|^2$. Consequently* [2, (3.26)] *holds with $\gamma_0^{\mathrm{II}} := \min\{1, \gamma_0(c_m)\}$ whenever $M^2\varepsilon_0^2 \leq c_m$. We take $c_m := 1$.*

Proof. Since $-\Delta^N \geq 0$, $G_\nu \leq (\nu + a_k P_k)^{-1} = P_k/(\nu + a_k) + (I - P_k)/\nu$. As $Q_k Q_k^* = I$, this gives $Q_k G_\nu Q_k^* \leq (\nu + a_k)^{-1}$ and $\Delta_\nu \geq a_k\nu/(a_k + \nu) \geq [a_\flat/(a_\flat + c_m)]\nu$. Also $\Delta_\nu \geq \Delta_0$ (Lemma E.148). Write $\Delta_\nu = \frac{1}{2}\Delta_\nu + \frac{1}{2}\Delta_\nu$ and apply [4, (1.22)] at $m^2 = 0$ to Δ_0. For (3.26): (3.27)–(3.28) decouple the scales by Neumann conditions for every m; the k-th term rescaled is (3.29), which is the display with $\nu = \nu_k$; and the $k = 0$ term has coefficient 1. □

Lemma E.150 (the mass defect in [2, Lemma 2.4, (2.75)–(2.76)])**.** *On a cube $\square$ of large blocks with Neumann conditions and A_0 gauged out, $a_k G_k(\square, 0) Q_k^* 1 = \frac{a_k}{a_k + \nu_k} 1$. So the constant reproduction (2.76) carries the extra term $E_{\mathrm{mass}} := -\frac{\nu_k}{a_k + \nu_k}\varphi'(y)$. With $C_\varphi := c_1 L^{3/4}(1 + \log L)^p$ ($\approx 39.6\, c_1$ at $L = 2$, $p = 6$) one has $|\varphi'(y)| \leq C_\varphi \lambda(t_k)^{-1/4} p(t_k)$, hence $|E_{\mathrm{mass}}| \leq (C_\varphi/a_\flat)\lambda^{-1/4} M^2 t_k^{7/4} p(t_k)$. Under the condition*

$$(E.\Phi) \qquad \lambda^{-1/4} m_2^2 \varepsilon_0^{7/4} \leq a_\flat/C_\varphi \qquad (a_\flat/C_\varphi \approx 1.89 \cdot 10^{-2}/c_1),$$

$|E_{\mathrm{mass}}| \leq p(t_k)$ for all $k \leq K$ and $M \leq m_2$. Then (2.65) and (2.76) hold with Bałaban's $O(p)$-constant plus 1, and (2.78) keeps a per-site error $O(t^{1/4}p^4) \leq O(t^{\kappa_0})$ for $\kappa_0 < 1/4$, in particular for the value $\kappa_0 = \frac{1}{5}$ used here, with an m-free constant.

Proof. $-\Delta^{\eta,N}_{0,\square}1 = 0$ and $P_k 1 = 1$ on $\square$, so $X_\nu 1 = (a_k + \nu_k)1$. Split $a_k G Q^* 1_{\square_1}\varphi'$ into $\varphi'(y)\, a_k G Q^* 1$, a cut term (exponentially small in $r(t)$, uniform in ν by Lemma E.148(i)) and Bałaban's gradient term $O(p)$. The field bound comes from (2.55) at index $k-1$: $\lambda(t/L)^{-1/4} = L^{1/4}\lambda(t)^{-1/4}$ ($d=3$), the canonical factor is $L^{1/2}$, and $p(t/L) \le (1+\log L)^p p(t)$ because $1+\log L+\log t^{-1} \le (1+\log L)(1+\log t^{-1})$ for $t \le 1$. Next, $\nu_k \lambda(t_k)^{-1/4} = \lambda^{-1/4} M^2 t_k^{7/4}$; this increases in $t_k \le \varepsilon_0$ and in $M \le m_2$. For (2.78): $\lambda(t)||\varphi^{(k)}|^4 - |\varphi(y)|^4| \le 4C'(C_\varphi + C')^3 \lambda(t)^{1/4} p(t)^4$, with C' Bałaban's constant plus 1. □

Remark (role of (E.Φ)). (E.Φ) is a normalisation. The condition (E.9) implies $6C_m \lambda^{-1/4} m_2^2 \varepsilon_0^{7/4} \le 1$, so (2.65) holds with the increment $C_\varphi/(6a_\flat C_m)$, free of ε_0 and M; (E.Φ) makes the increment 1, independent of c_1, so that no feedback between c_1 and [2, Lemma 2.4] can arise.

The constants of Lemma 2.7 of [2] and of the counterterm coefficients. In this paragraph $d = 3$, $L = 2$, $a = 1$ and $a_\flat = a(1 - L^{-2})$. We write $t = t_k = L^k \varepsilon \le \varepsilon_0$ and $\nu = \nu_k = M^2 t^2$, where $M \in (0, m_2]$ is the Higgs mass. We also write $e(t) = et^{1/2}$, $\lambda(t) = \lambda t$, $p(t) = b_0(1 + \log t^{-1})^p$ and $r(t) = R(1 + \log t^{-1})^\varrho$ with $\varrho > 1$. M_b is the large-block parameter of [2]. The constants c_0, δ_0, R_0 are those of [4, Theorem, (1.9)–(1.12), p. 573], of [2, Prop. 2.2, (2.58), pp. 570–571] and of [4, (1.15)–(1.16), p. 574], taken equal: we use the minimum of the δ_0's and the maximum of the c_0's and of the R_0's. By Lemma E.148(i) and [4, p. 572, (1.6)] they depend only on d and M_b (and α), and they hold for every rescaled mass $\nu \ge 0$. For $\delta > 0$ put

$$S_d(\delta) := \sum_{z \in \mathbb{Z}^d} \mathrm{e}^{-\delta|z|}, \qquad J_d(\delta) := \mathrm{e}^{\delta\sqrt{d}}\big(1 + 2\sqrt{d}/\delta\big)^d, \qquad S_F(\delta) := d\,\mathrm{e}^{\delta\sqrt{d}} S_d(\delta).$$

These satisfy three elementary bounds:

- $S_d(\delta) \le (1 + 2\sqrt{d}/\delta)^d$;
- for every $\eta \in (0,1]$ and every unit block B, $\eta^d \sum_{z \in \eta\mathbb{Z}^d} \mathrm{e}^{-\delta\,\mathrm{dist}(z,B)} \le J_d(\delta)$;
- for every point x', $\sum_F \mathrm{e}^{-\delta\,\mathrm{dist}(F,x')} \le S_F(\delta)$, where F runs over the unit faces of the unit lattice.

Setting for Lemma 2.7 of [2]. We work in scale-k units on the fine lattice of spacing $\eta = L^{-k}$. The unit lattice is the lattice of unit blocks $B(x)$. Let:

- $\Omega := B^k(\Lambda_2^{(k)})$, and $\hat{B} := B^{(k+1),\eta}$ the background of [2, (2.110)–(2.113), pp. 580–581];
- $G_k := (-\Delta^{\eta,N}_{\hat{B},\Omega} + \nu + a_k P_k(\hat{B}))^{-1}$, where $P_k = Q_k^* Q_k$ and $Q_k Q_k^* = I$;
- $\Delta := a_k - a_k^2 Q_k G_k Q_k^*$, acting on functions on $\Omega^{(k)}$;
- $Q := Q(\hat{B})$ the averaging from the unit lattice to the lattice of L-blocks, so that $QQ^* = I$;
- $\mathsf{M} := \Delta + aL^{-2} Q^* Q$ and $C_{\Lambda_4} := (\Lambda_4 \mathsf{M} \Lambda_4)^{-1}$, as in [4, (1.13)–(1.14)].

Kernels of operators from the unit lattice carry unit weights, $(G_k Q_k^* f)(z) = \sum_y (G_k Q_k^*)(z,y) f(y)$, and fine-lattice pairings carry the weight η^d. We assume the following.

(G) $\mathrm{dist}(\Lambda_5^{(k)}, \Lambda_4^{(k)c}) > r(t)$, $\mathrm{dist}(\Lambda_4^{(k)}, \Omega^c) > 2r(t)$ and $r(t) \ge R_0 + 1 + \sqrt{d}$. The first two inequalities follow from the definition [2, (2.8)] of the sets Λ_i. The third holds since $r(t) \ge 3R_0 + 15$ under the conditions of Appendix A. Each $\Lambda_i^{(k)}$ is a union of large blocks.

(Reg) $\hat{B}$ is regular on Ω: $|\partial \hat{B}| \le c_B p(t)$. Hence the conclusions of [2, Prop. 2.2] hold for $\hat{B}$, by [2, Lemma 2.3, (2.60)] at scale $k+1$.

(Res) The scalar block field ψ and the scalar field ϕ satisfy three bounds:
- $|\psi(y)| \le 2\lambda(t)^{-1/4} p(t)$;
- $|\psi(y) - U(\hat{B}(\Gamma_{y,x}))\phi(x)| \le 2Ldp(t)$ for $x \in B(y)$;
- $|U(\hat{B}(\Gamma_{y,x} \cup \langle x,x'\rangle \cup \Gamma_{x',y'}))\psi(y') - \psi(y)| \le 4Ldp(t)$ for every bond $\langle x,x'\rangle$ with $x \in B(y)$ and $x' \in B(y')$.

These are the restrictions [2, (2.16), p. 560] at scale t, except that there the transports are taken with the block field rather than with $\hat{B}$. Replacing one transport by the other changes

the right-hand sides by terms of size $e(t)p(t)|\psi| = O(t^{1/4}p(t)^2)$, and this affects only $\mathcal{E}(t)$ below.

(Γ) The contours Γ of [1, §1] define Q, Q_k and their adjoints. For a fine bond b, consider the closed loop formed by b and the contours that transport ψ to the two endpoints of b. This loop bounds a lattice surface of area at most $c_\Gamma L^2$, and of area at most $c_\Gamma \eta$ if b lies inside one unit block. Here c_Γ depends only on d.

Lemma E.151 (Lemma 2.7 of [2] with the mass term explicit)**.** *Assume (G), (Reg), (Res) and (Γ). Then for every $x \in \Lambda_5^{(k)}$*

$$aL^{-2}(C_{\Lambda_4}Q^*\psi)(x) = (Q^*\psi)(x) + \mathcal{R}(x), \qquad |\mathcal{R}(x)| \le \big(K_2^0 + \mathcal{E}(t) + K_m\lambda^{-1/4}M^2t^{7/4}\big)\,p(t),$$

where

$$K_2^0 := 4Ld\,a\,c_0^2\mathrm{e}^{\delta_0}S_d(\delta_0)S_F(\delta_0), \qquad K_m := 2a\,c_0^2S_d(\delta_0),$$

$$\mathcal{E}(t) := a\,c_0S_d(\delta_0)\,\mathcal{E}_1(t) + 4a^2c_0^2\mathrm{e}^{\delta_0\sqrt{d}}S_d(\delta_0/2)^2\lambda^{-1/4}t^{-1/4}\mathrm{e}^{-\delta_0 r(t)/2},$$

$$\mathcal{E}_1(t) := 2\lambda^{-1/4}\Big[c_0c_Bc_\Gamma e\,\mathrm{e}^{\delta_0}\big(S_F(\delta_0)L^2 + dJ_d(\delta_0)\big)t^{1/4}p(t) + c_0\mathrm{e}^{2\delta_0\sqrt{d}}S_F(\delta_0/2)\,t^{-1/4}\mathrm{e}^{-\delta_0 r(t)/2}\Big].$$

K_2^0 and K_m depend only on d, a, L, M_b. $\mathcal{E}(t)$ depends in addition on $e, \lambda, b_0, p, R, \varrho, c_B, c_\Gamma$, but not on M, ε, k or ε_0. Moreover $\mathcal{E}(t) \to 0$ as $t \to 0$, and $\mathcal{E}^* := \sup_{0<t\le 1}\mathcal{E}(t) < \infty$. Consequently [2, (2.114)] *holds on $\Lambda_5^{(k)}$ in the form*

$$|\phi'(x)| \le \big(2Ld + K_2^0 + \mathcal{E}(t) + K_m\lambda^{-1/4}M^2t^{7/4}\big)p(t),$$

where $\phi' = \phi - aL^{-2}C_{\Lambda_4}Q^*\psi$ [2, (2.110)].

Proof. Put $g := Q^*\psi$. Then $|g| \le 2\lambda(t)^{-1/4}p$, and all transports are unimodular.

Decomposition. Since $QQ^* = I$ we have $Q^*Qg = g$, and Λ_4 commutes with Q^*Q. Hence $\Lambda_4\mathsf{M}\Lambda_4 g = aL^{-2}\Lambda_4 g + \Lambda_4\Delta g - \Lambda_4\Delta\Lambda_4^c g$. Applying C_{Λ_4} gives $aL^{-2}C_{\Lambda_4}\Lambda_4 g = \Lambda_4 g + C_{\Lambda_4}\Lambda_4\Delta\Lambda_4^c g - C_{\Lambda_4}\Lambda_4\Delta g$.

Let S be the union of the unit blocks $B(y) \subset \Omega$ with $\operatorname{dist}(B(y), \Omega^c) \ge R_0 + 1$, and let χ_S be its indicator. By (G), $\Lambda_4 \subset S$ and $\operatorname{dist}(\Lambda_4, S^c) \ge r(t)$. Therefore

$$\mathcal{R} = C_{\Lambda_4}\Lambda_4\big[\Delta\Lambda_4^c g - \Delta\chi_{S^c}g\big] - C_{\Lambda_4}\Lambda_4\Delta(\chi_S g),$$

and $\|C_{\Lambda_4}\|_{\infty\to\infty} \le c_0S_d(\delta_0)$ by [4, (1.16)].

Off-diagonal terms. Let f be supported in Λ_4^c with $|f| \le |g|$, and let $x' \in \Lambda_4$. Then $(\Delta f)(x') = -a_k^2(Q_kG_kQ_k^*f)(x')$. Every point of $B(x')$ has distance $\ge R_0$ from Ω^c, so [2, Prop. 2.2] gives

$$|(\Delta f)(x')| \le a^2c_0\mathrm{e}^{\delta_0\sqrt{d}}\sum_{x''\notin\Lambda_4}\mathrm{e}^{-\delta_0|x'-x''|}|g(x'')|.$$

For $x \in \Lambda_5$ we have $|x - x'| + |x' - x''| > r(t)$. Splitting the exponentials in halves gives $|(C_{\Lambda_4}\Lambda_4\Delta f)(x)| \le 2a^2c_0^2\mathrm{e}^{\delta_0\sqrt{d}}S_d(\delta_0/2)^2\mathrm{e}^{-\delta_0 r(t)/2}\lambda^{-1/4}t^{-1/4}p$. This applies to $f = \Lambda_4^c g$ and to $f = \chi_{S^c}g$, and it gives the last term of $\mathcal{E}(t)$.

Key identity. Put $u := Q_k^*(\chi_S g)$. Then $Q_k^*Q_k u = u$, because χ_S is constant on unit blocks. With $G_k^{-1} = (-\Delta_{\hat B} + \nu) + a_kQ_k^*Q_k$, it follows that $a_kG_kQ_k^*(\chi_S g) = u - G_k(-\Delta_{\hat B} + \nu)u$. Applying a_kQ_k, and using that G_k is self-adjoint,

$$(\Delta\chi_S g)(x') = a_k\langle w_{x'},\, (-\Delta_{\hat B} + \nu)u\rangle, \qquad w_{x'} := G_kQ_k^*\delta_{x'}.$$

Mass term. We have $a_k\nu\langle w_{x'}, u\rangle = a_k\nu(Q_kG_ku)(x')$. By [4, (1.10)], $|(G_ku)(z)| \le c_0\|u\|_\infty$ on $B(x')$. Hence $|a_k\nu\langle w_{x'}, u\rangle| \le 2ac_0\lambda^{-1/4}M^2t^{7/4}p$, and after C_{Λ_4} this is the K_m-term. This is the only place where ν enters, apart from c_0, δ_0, R_0.

Kinetic term. Write $\langle w, -\Delta_{\hat B} u\rangle = \sum_{b\subset\Omega} \eta^d\, \overline{(\eta^{-1}d_b w)}(\eta^{-1}d_b u)$, where $\eta^{-1}d_b$ is the covariant difference on the bond b. If $d_b u \neq 0$, then $\operatorname{dist}(b,\Omega^c) \geq R_0$, and [2, (2.58)] gives $|\eta^{-1}d_b w_{x'}| \leq c_0 \mathrm{e}^{-\delta_0 \operatorname{dist}(b,x')}$. The bonds with $d_b u \neq 0$ are of three kinds.

- Bonds crossing a unit face, between two L-blocks in S or inside one L-block. By (Res), (Γ), Stokes' formula, (Reg) and $|\mathrm{e}^{i\theta}-1| \leq |\theta|$, such a bond has $|d_b u| \leq 4Ldp + c_\Gamma L^2 e(t) c_B p|\psi|$.
- Bonds inside a unit block. Here $|\eta^{-1}d_b u| \leq c_\Gamma e(t) c_B p|\psi|$.
- Bonds crossing a face of $\partial B^k(S)$. Here $|d_b u| \leq |\psi|$, and the face has distance $\geq r(t) - \sqrt{d}$ from x'.

A unit face carries η^{1-d} fine bonds of weight η^{d-1}. So a face F contributes at most

$$c_0 \mathrm{e}^{\delta_0} \mathrm{e}^{-\delta_0 \operatorname{dist}(F,x')} \sup_{b\in F} |d_b u|.$$

Summing with S_F, J_d and $S_F(\delta_0/2)$, and using $e(t)p|\psi| \leq 2e\lambda^{-1/4}t^{1/4}p^2$ and $|\psi| \leq 2\lambda^{-1/4}t^{-1/4}p$, we get

$$|a_k \langle w_{x'}, -\Delta_{\hat B} u\rangle| \leq a\big(4Ld\, c_0 \mathrm{e}^{\delta_0} S_F(\delta_0) + \mathcal{E}_1(t)\big)p.$$

Sum. Multiply the mass and kinetic bounds by $c_0 S_d(\delta_0)$ and add the off-diagonal terms. This gives the bound on $\mathcal{R}$. The function $t^{1/4}(1+\log t^{-1})^p$ is bounded on $(0,1]$ and tends to 0 as $t\to 0$. Since $\varrho > 1$, the same holds for $t^{-1/4}\mathrm{e}^{-\delta_0 R(1+\log t^{-1})^\varrho/2}$. For the last claim, $|\phi'| \leq |\phi - Q^*\psi| + |\mathcal{R}|$, and $|\phi - Q^*\psi| \leq 2Ldp$ by (Res). □

Corollary E.152 (the mass defect of Lemma 2.7 of [2])**.** *Assume (E.*Φ*): $\lambda^{-1/4} m_2^2 \varepsilon_0^{7/4} \leq a_\flat / C_\varphi$, with C_φ as in Lemma E.150. Then for every $M \in (0, m_2]$, every $t \leq \varepsilon_0 \leq 1$ and every $x \in \Lambda_5^{(k)}$,*

$$|\mathcal{R}(x)| \leq K_2^\natural\, p(t), \qquad K_2^\natural := K_2^0 + \mathcal{E}^* + K_m a_\flat / C_\varphi,$$

and $|\phi'(x)| \leq (2Ld + K_2^\natural)p(t)$. The constant $K_2^\natural$ does not depend on M, ε, k or ε_0.

Proof. $\lambda^{-1/4}M^2 t^{7/4}$ is increasing in M and in t, so it is at most $\lambda^{-1/4} m_2^2 \varepsilon_0^{7/4} \leq a_\flat / C_\varphi$. Now apply Lemma E.151. □

In the next three lemmas, κ_G, A_0^T, A_1^T, S_L, K_1 (from (F4)), A_E, A_∇ and the lattice sums J_0, J_1, J_2 are those of Lemmas E.39, E.42 and E.44 and of Subsection E.3.

Lemma E.153 (the coefficient [3, (3.16)] uniformly in the mass)**.** *Let $j'' \geq 1$ and $\xi = L^{-j''}$. Work on the ξ-lattice torus Λ of side $\ell = \ell_T / t_{j''} \in L\mathbb{N}$, with $\ell \geq L$. Let $\nu \in [0,1]$; for the scalar field $\nu = M^2 t_{j''}^2$, and for a photon component $\nu = \mu_*^2 t_{j''}^2$. Let $b := a_{j''} \in [a_\flat, a]$, and*

$$G := G^\xi_{j''}(0) = (-\Delta^\xi + \nu + bP_1)^{-1}, \qquad C := C^\xi = (-\Delta^\xi + 1)^{-1}, \qquad W := (1-\nu) - bP_1,$$

so that $G = C + GWC$ [3, (3.16), p. 437]. Let g, g' be bounded functions and $|q| \leq 1$. Take the coordinate differences $y'_\mu - y_\mu$ in the fundamental domain centred at y, so that $|y'_\mu - y_\mu| \leq d(y,y')$. Then the coefficient of [3, (3.16)]*,*

$$\Xi(y) := -q^2 \sum_{y'} \xi^d C(y-y') g(y) G(y,y') g'(y') (y'_\mu - y_\mu) + q^2 \sum_{y'} \xi^d \sum_{\nu'=1}^{d} \big(\partial^\xi_{\nu'} GWC \partial^{\xi *}_{\nu'}\big)(y,y')\, g(y) G(y,y') g'(y') (y'_\mu - y_\mu),$$

satisfies

$$|\Xi(y)| \leq \|g\|_\infty \|g'\|_\infty A_G \big[1 + d(1+a) K_1 \big(K_1 + A_\nabla J_0(\kappa_G)\big)\big] =: K_{3.16}, \qquad A_G := A_0^T + \frac{A_E}{\mathrm{e}\,\kappa_G}.$$

$K_{3.16}$ depends only on (a, L) and on $\|g\|_\infty \|g'\|_\infty$. It does not depend on $\nu \in [0,1]$, ξ, j'', ℓ or M.

Proof. By Lemma E.44, $G = F + E$ with $F = C$, $|E| \le A_E \mathrm{e}^{-\kappa_G d}$ and $|\nabla_1 E| \le A_\nabla \mathrm{e}^{-\kappa_G d}$. By Lemmas E.39 and E.42, $0 \le F \le A_0^T \mathrm{e}^{-d/8}/(d+\xi)$, since $S(\ell/4) \le S_L$ for $\ell \ge L$. These constants are uniform in $\nu \in [0,1]$. Hence $|G(y,y')| d(y,y') \le A_G$.

The first term of Ξ is at most $\|g\|_\infty \|g'\|_\infty A_G$, because $C \ge 0$ and $\xi^d \sum_{y'} C(y-y') = 1$.

Put $h(y') := G(y,y') g'(y') (y'_\mu - y_\mu)$, so that $\|h\|_\infty \le A_G \|g'\|_\infty$. The second term equals $q^2 g(y) \sum_{\nu'} (\partial_{\nu'} G W C \partial^*_{\nu'} h)(y)$. Its absolute value is at most $d \|g\|_\infty \|\partial G\|_{\infty\to\infty} \|W\|_{\infty\to\infty} \|C\partial^*\|_{\infty\to\infty} \|h\|_\infty$, and the three operator norms are bounded as follows:

- $\|W\|_{\infty\to\infty} \le |1-\nu| + b \le 1 + a$, since P_1 is an average;
- $\|C\partial^*\|_{\infty\to\infty} \le K_1$, by summation by parts, (F4) and Lemma E.42;
- $\|\partial G\|_{\infty\to\infty} \le K_1 + A_\nabla J_0(\kappa_G)$, since $\nabla_1 G = \nabla F + \nabla_1 E$ and a torus sum of $\mathrm{e}^{-\kappa d}$ is at most the lattice sum $J_0(\kappa)$. □

Lemma E.154 (the coefficients of [3, p. 441] uniformly in the mass)**.** *Work in the setting of Lemma E.153, with $j_0 \ge 1$ in place of j''. Let $G := G^\xi_{j_0}(0)$ and $E := G - C = GWC$. Fix $y \in \Lambda$ and directions μ, μ', ν_0. Let $m(y')$ be either 1 or $y'_{\nu_0} - y_{\nu_0}$, with coordinates taken in the fundamental domain centred at y. For $K \in \{G, C\}$ put*

$$X_1[K] := \xi^d \sum_{y'} (K\partial^{\xi *}_{\mu'})(y,y') (K\partial^{\xi *}_{\mu})(y',y)\, m(y'), \qquad X_2[K] := \xi^d \sum_{y'} K(y,y') (\partial^\xi_{\mu'} K \partial^{\xi *}_\mu)(y',y)\, m(y'),$$

$$X_3[K] := K(y,y), \qquad X_4[K] := \xi\, (K\partial^{\xi *}_\mu)(y,y).$$

Then $|X_i[G] - X_i[C]| \le K_{441}$ for $i = 1, \dots, 4$. The constant K_{441} depends only on (a, L), and not on $\nu \in [0,1]$, ξ, j_0, $\ell \ge L$, y or the directions.

Proof. Write $r := d(w,w')$ and $\kappa := \kappa_G \le \frac{1}{16}$. By Lemmas E.39, E.42 and E.44 (items (E1) and (E3)), together with the symmetry of E:

- $|C| \le A_0^T \mathrm{e}^{-r/8}/(r+\xi)$ and $|\nabla C| \le A_1^T \mathrm{e}^{-r/8}/(r+\xi)^2$;
- $|E| \le A_E \mathrm{e}^{-\kappa r}$, and every difference of E is at most $A_\nabla \mathrm{e}^{-\kappa r}$;
- hence $|\nabla G| \le (A_1^T (r+\xi)^{-2} + A_\nabla) \mathrm{e}^{-\kappa r}$.

Shifting the point at which a difference is evaluated by ξ changes these bounds by at most the factor $c_* := 4\mathrm{e}^{1/8}$. Moreover $|m| \le 1 + r$, and every torus sum below is at most the corresponding sum over $\xi\mathbb{Z}^3$, that is, at most J_0, J_1 or J_2.

X_3 and X_4. The differences are $E(y,y)$ and $\xi(E\partial^*_\mu)(y,y)$, which are at most A_E and A_∇.

X_1. The difference $X_1[G] - X_1[C]$ is the sum over y' of $\xi^d\big[(E\partial^*_{\mu'})(y,y')(G\partial^*_\mu)(y',y) + (C\partial^*_{\mu'})(y,y')(E\partial^*_\mu)(y',y)\big] m(y')$. Each product has one factor at most $A_\nabla \mathrm{e}^{-\kappa r}$ and one factor at most $(A_1^T(r+\xi)^{-2} + A_\nabla)\mathrm{e}^{-\kappa r}$. Using $(1+r)(r+\xi)^{-2} \le (r+\xi)^{-2} + (r+\xi)^{-1}$ and $(1+r)\mathrm{e}^{-\kappa r} \le (1 + 2/(\mathrm{e}\kappa))\mathrm{e}^{-\kappa r/2}$, the sum is at most $A_\nabla[2A_1^T(J_2(\kappa) + J_1(\kappa)) + A_\nabla(1 + 2/(\mathrm{e}\kappa)) J_0(\kappa/2)]$.

X_2. Put $f_K := K(y,\cdot)m$ and $k_K := (K\partial^*_\mu)(\cdot, y)$. Then $X_2[G] - X_2[C] = \xi^d \sum_{y'} [f_E\, \partial_{\mu'} k_G + f_C\, \partial_{\mu'} k_E](y')$. Sum by parts in y' on the torus.

- If $m \equiv 1$, there is no boundary term.
- Otherwise $\partial^*_{\mu'}(K(y,\cdot)m) = (\partial^*_{\mu'} K(y,\cdot))\, m(\cdot - \xi e_{\mu'}) + K(y,\cdot)\, \partial^*_{\mu'} m$. Here $|\partial^*_{\mu'} m| \le 1$ except on at most $(\ell/\xi)^{d-1}$ sites. On those sites $|\partial^* m| \le \ell/\xi$ and $d(y,y') \ge \ell/2 - 1$, so they contribute at most $c_*^2 \ell^d (A_E + A_0^T)(A_1^T + A_\nabla)\mathrm{e}^{-\kappa(\ell/2-1)}$, which is bounded uniformly in ℓ.

On the remaining sites:

- $|\partial^* f_E| \le c_*(A_\nabla(2+r) + A_E)\mathrm{e}^{-\kappa r}$ and $|k_G| \le (A_1^T(r+\xi)^{-2} + A_\nabla)\mathrm{e}^{-\kappa r}$;
- $|\partial^* f_C| \le c_*(A_1^T(2+r)(r+\xi)^{-2} + A_0^T(r+\xi)^{-1})\mathrm{e}^{-r/8}$ and $|k_E| \le A_\nabla \mathrm{e}^{-\kappa r}$.

In each product only one factor is singular, of order at most $(r+\xi)^{-2}$. So the sums are bounded by constant multiples of $J_0(\kappa/2)$, $J_1(\kappa)$ and $J_2(\kappa)$. □

Lemma E.155 (the counterterm lines at zero background, uniformly in the mass)**.** *Let* $0 < M \le m_2$ *and* $\mu_* \ge 0$*, and assume* $m_2\varepsilon_0 \le 1$*,* $\mu_*\varepsilon_0 \le 1$ *and* $\varepsilon_0 \le 1$*. By* [3, (1.27), p. 418]*,*

$$G_K(0) = \sum_{j=0}^{K-1} \Gamma_j, \qquad \Gamma_0 = G_1(0) = C^{(0),\varepsilon}(0), \qquad \Gamma_j = G_{j+1}(0) - G_j(0) \quad (j \ge 1).$$

For the scalar field (mass M*), and for each component of the photon (mass* μ_**, Lemma E.38), every* Γ_j *satisfies* [3, (2.10)–(2.12), p. 426] *at background* 0 *with index* j*. The constants, and the rate* $\delta_1 := \min\{\kappa_H, \min(1/8, \kappa_G)/L\}$*, depend only on* (a, L) *and* α*. They are uniform in* ε*,* j*,* ℓ_T*,* M *and* μ_**.*

Proof. The case $j \ge 1$*.* This is Corollary E.49, with $m_s = M$ and $m_v = \mu_*$. It applies because $t_{j+1} \le t_K \le \varepsilon_0 \le \min(1/m_f, 1) < L$, so $m_f t_j \in [0, 1]$.

The case $j = 0$*.* Work in scale-1 units, with spacing $1/L$. By Lemma E.44 at $j = 1$, $G_1(0) = F + E$. Then (F0), (F1), (E1) and (E3) give two bounds:

- $|G_1(0)| \le (LA_0^T + A_E)\mathrm{e}^{-\min(1/8,\kappa_G)d}$;
- each lattice difference costs at most a factor L, with constant $L^2A_1^T + A_\nabla$.

In physical units these are (2.10) and its derivative form at index 0, with rate $\min(1/8, \kappa_G)/L$. Distinct lattice points are at distance $\ge \eta$, so the Hölder quotient (2.11) is at most twice the supremum of the difference divided by η^α. The contour sums (2.12) follow by summing (2.10) along the contour. □

The counterterm coefficients. By [3, (1.23)–(1.28), pp. 416–418], δm^2 is a finite sum over one-particle-irreducible graphs G_0 with two scalar legs, including the lower-order δm^2-insertions. Their orders $e^\alpha\lambda^\beta$ satisfy $2 \le \alpha + 2\beta \le 4$, so that $o := \alpha/2 + \beta \in \{1, 2\}$. $\delta m^2_K(x)$ is the same sum with propagators $G_K(0)$. Expand each line by Lemma E.155. Then $\delta m^2_{K,k}(x)$ is the sum over index assignments with largest index $\ge k$, and $\delta m^2_k(x)$ the sum over those with all indices $< k$ [3, (1.28)].

Work in scale-K units, with $\eta_K := \varepsilon/t_K$. Then $L^j\eta_K = t_j/t_K \le 1$, $e(t_K)^2 = e^2 t_K$ and $\lambda(t_K) = \lambda t_K$. For an amputated kernel $\Sigma_{G_0}(y_1, y_2)$ put $c_{G_0}(x) := \sum_{y_2} \eta_K^d \Sigma_{G_0}(x, y_2)$. Then $t_K^2\,\delta m^2_{K,k}(x) = \sum_{G_0} c^{[\ge k]}_{G_0}(x)$.

Theorem E.156 (pointwise bound on $\delta m^2_{K,k}$)**.** *Let* $\varepsilon = L^{-K}\varepsilon_0$ *with* $K \in \mathbb{N}$*, so that* $t_K = \varepsilon_0$*. Let* $0 \le k \le K$ *and* $t := t_k$*, and assume the hypotheses of Lemma E.155. Then for every point* x *of the torus of spacing* ε*,*

$$|\delta m^2_{K,k}(x)| \le C_m\Big[(e^2 + \lambda)\, t^{-1}\frac{L}{L-1} + (e^4 + e^2\lambda + \lambda^2)(K - k)\Big],$$

and $K - k = \log_L(\varepsilon_0/t)$*, so that* $t^2|\delta m^2_{K,k}(x)| = O(t)$*. Here* C_m *is the constant* $O(1)$ *of* [3, (1.33)] *for the expressions of Lemma E.55 (it contains the constant* C_A *of Lemma E.51, with* δ_1 *of Lemma E.155), times the number of graphs. It does not depend on* ε*,* k*,* x *or* ℓ_T*. For* $k = K$ *the left side is* 0*.*

Proof. Fix a graph G_0 and a value $j^* \in \{k, \dots, K-1\}$ of the largest index. By Lemma E.52, all renormalisations act inside the sub-sum with top index j^*, and G_0 itself is not renormalised. By Lemma E.55, the expression with the leg at y_1 fixed at x obeys [3, (1.33)]. The external fields are constant with norm $O(1)$. Lemma E.51(ii) at fixed j^* then gives

$$|c^{[j^*]}_{G_0}(x)| \le O(1)\, e(t_K)^\alpha \lambda(t_K)^\beta (t_{j^*}/t_K)^{D(G_0)},$$

with the degree $D(G_0)$ of that lemma. At background 0, a derivative on a constant external leg vanishes. Close the two legs by a scalar line ℓ. Lemma E.166(a),(b) with $n_\partial(\ell) = 0$ give $D(G_0) \ge o - 2$.

Now sum over j^*, using $t_{j^*}/t_K \le 1$.

- For $o = 1$: $\sum_{j^*=k}^{K-1}(t_{j^*}/t_K)^{-1} \le (t_K/t)L/(L-1)$, and $t_K^{-2}(e^2+\lambda)t_K(t_K/t)L/(L-1) = (e^2+\lambda)t^{-1}L/(L-1)$.
- For $o = 2$: there are $K-k$ terms, each at most 1, which gives $(e^4+e^2\lambda+\lambda^2)(K-k)$.

Finally $K - k = \log_L(\varepsilon_0/t)$, and $t\ln(\varepsilon_0/t) \le \varepsilon_0/\mathrm{e}$. □

Lemma E.157 (where the masses enter the proof of [3, (1.33)])**.** *Consider the graphs of orders $2 \le \alpha + 2\beta \le 4$ at zero background without operator lines. Fix $j_0 \ge 1$ and work in units of t_{j_0} on the torus of spacing $\xi = L^{-j_0}$. Let G_s be the scalar propagator $G^{\xi}_{j_0}(0)$ and G_p each component of the photon propagator. Put $E_f := G_f - C^{\xi}$ for $f \in \{s, p\}$.*

In [3, pp. 434–445]*, the masses M, μ_* enter the constant $O(1)$ of* [3, (1.33)] *only through the following quantities:*

(i) *the line bounds* [3, (2.10)–(2.12)]*;*
(ii) *pointwise bounds on G_s, G_p, E_s, E_p and on their differences, at pp. 437–444;*
(iii) *the coefficient of (3.16) on p. 437, and the coefficients of p. 441;*
(iv) *the quantity $\xi G_f(y,y)$ in (3.21) on p. 438;*
(v) *the difference between the coefficient of (3.23) on p. 439 and its C^{ξ}-version;*
(vi) *for the classes (2.19), (2.20) and (2.21c), the difference between the vertex coefficient $\sum \xi^{2d}\Gamma'_{\mu}$ of p. 444 and its C^{ξ}-version;*
(vii) *the expressions produced by the replacements in (iii), (v) and (vi), which* [3] *calls convergent.*

In particular, p. 442 contributes no mass dependence at these orders.

Proof. We go through the pages in order.

- P. 434: (3.4) decomposes the propagators by (2.6) into slices, which are estimated by (2.10)–(2.12); this is (i). Graphs with an odd number of external vector legs vanish, since $\operatorname{tr} q^{2n+1} = 0$.
- Pp. 435–436: the last two terms of (3.7) cancel those of (3.6) identically. (3.9)–(3.14) are Taylor's formula followed by (2.10)–(2.11) for a line of index j''.
- P. 437: the sum over orderings produces $G^{\eta}_{j''}(0)$ and the photon propagator. (3.16) uses $G = C^{\xi} + G(1 - m^2_{j''} - a_{j''}P_{j''})C^{\xi}$, together with pointwise bounds on C^{ξ}, on the propagators and on their derivatives; this gives (ii) and (iii).
- P. 438: (3.19) is an exact subtraction. The first graph of (3.21) is controlled by the boundedness of $\eta G_k(x,x)$, which in scale-k units is $\xi G_f(y,y)$; this is (iv).
- P. 439: (3.23) is treated by the method of (3.16), with both propagators replaced; this gives (v) and a convergent remainder, which is part of (vii). (3.24) contains C^{ξ} only.
- Pp. 440–441: the last bracket of (3.26) is estimated as in (3.13)–(3.14). The replacement of $G_{j_0}(0)$ is (iii). (3.27)–(3.28) contain C^{ξ} only.
- P. 442: (3.29) vanishes by the Ward–Takahashi identity (2.26) for C^{ξ}, whose mass is 1. The graphs (3.31) have two external vector legs and order ≥ 4, so inside a graph of order ≤ 4 with two external scalar legs they would give order ≥ 6; hence they do not occur. (2.21a) vanishes, (2.21e) carries an extra factor η, and (2.21d), (2.21f) cancel against (2.21g).
- P. 443: (3.33)–(3.35) transport legs, and the result is estimated by line bounds.
- P. 444: (2.21b), (2.21g) and (2.18) vanish exactly. The replacements of the propagators give (vi) and convergent remainders, while the C^{ξ}-versions (3.37), "derivative of a constant", and (3.38) vanish.
- P. 445: the induction on the last divergent subgraph, and the cancellation of vacuum graphs, are combinatorial. □

Proposition E.158 (independence of C_m from the masses)**.** *Assume the hypotheses of Lemma E.155; in particular (E.m2) $m_2^2\varepsilon_0^2 \le 1$ and $\mu_*\varepsilon_0 \le 1$. Then the constant C_m of Theorem E.156 can be chosen depending only on $\alpha_0, \bar{n}, d, L, a, M_\flat$. It is the same for all $M \in (0, m_2]$ and all such μ_*, and it does not depend on ℓ_T.*

Proof. C_m is the product of three factors:

- the number of graphs, which depends only on $\bar{n}$ and d;
- the constant C_A of Lemma E.51, which contains no mass; its rate δ_1 is free of M and μ_* by Lemma E.155;
- the constant $O(1)$ of [3, (1.33)] for the expressions of Lemma E.55.

For $j_0 = 0$ every quantity is a single slice, which is bounded by Lemma E.155. For $j_0 \ge 1$ we have $M^2 t_{j_0}^2 \le 1$ and $\mu_*^2 t_{j_0}^2 \le 1$, since $t_{j_0} \le \varepsilon_0$. By Lemma E.157, the masses enter $O(1)$ only through (i)–(vii), and each of these is bounded uniformly in the masses.

- (i) is uniform by Lemma E.155.
- (ii) follows from Statement K.30(a).
- (iii) follows from Lemmas E.153 and E.154.
- (iv) and (v) follow from Statement K.30(b), items (C1)–(C3). The coefficient of [3, (3.23)] and both terms of [3, (3.37)] have the form (C2). The remainder of [3, (3.23)] that comes from $g'(x') - g'(x)$ has the form (C3). The scalar charge satisfies $|q| \le 1$ [3, p. 426].
- (vi) for (2.19) and (2.20). Each term in the first line of [3, (3.38)] has one twice-differenced factor and one undifferenced factor, and these two share a summed variable that carries one of the differences. The replacement is made line by line, so the same holds for its C^ξ-version. On the torus, $\xi^3 \sum_z (\partial_\nu f) g = \xi^3 \sum_z f(\partial_\nu^* g)$, and the kernel of $K\partial_\nu^*$ is the forward difference of K in its second variable. So one summation by parts brings both versions to the form T of (C4), and Statement K.30(b) applies.
- (vi) for (2.21c) follows from Statement K.31(a) together with Statement K.30(c), with $K_1 = K_2 = G_s$ and $K_3 = G_p$.
- (vii): by Statement K.31(b), these expressions are the differences between the coefficients in (iii), (v), (vi) and their C^ξ-versions, and they enter only through bounds on their absolute values. The C^ξ-versions contain no mass, so these differences are bounded by the bounds just given plus mass-free quantities.

All constants depend only on (a, L) and on the finitely many graphs. They are uniform in ξ, j_0 and $\ell \ge L$, and hence in ℓ_T. □

Proposition E.159 (the mass reading of [2, p. 589])**.** *Assume (E.m2): $m_2^2\varepsilon_0^2 \le 1$, and (E.$\Phi$): $\lambda^{-1/4} m_2^2 \varepsilon_0^{7/4} \le a_\flat / C_\varphi$. Equivalently, $\varepsilon_0 \le \varepsilon_{(\mathrm{E.md})} := \min\{1/m_2, (a_\flat \lambda^{1/4}/(C_\varphi m_2^2))^{4/7}\}$ (stage 10a of Appendix A). Assume moreover $\mu_*\varepsilon_0 \le 1$, which holds under (E.dom), and $\varepsilon_0 \le 1$. Then at every $M \in (0, m_2]$, in particular at $M = m_2$ and $M = m_1$, every estimate of* [1, 2, 3, 4] *that uses an upper bound on the mass holds with the following constants:*

- $\gamma_0^{\mathrm{II}} = \min\{1, \frac{1}{2}\min\{\gamma_0^{(0)}, a_\flat/(a_\flat+1)\}\}$ *in* [2, (3.26)]*, with $\gamma_0^{(0)}$ as in Lemma E.149;*
- $\gamma_1 = a_\flat/(a_\flat+1) = 3/7$ *in the bound $G_k'' \ge \gamma_1 \nu_k$ of* [2, (3.11)]*;*
- *the constant of* [2, (2.65)] *increased by* 1*;*
- $K_2^\natural$ *of Corollary E.152 in* [2, (2.113)]*, and $2Ld + K_2^\natural$ in* [2, (2.114)]*;*
- $K_{3.16}$ *of Lemma E.153 for* [3, (3.16), p. 437]*, and K_{441} of Lemma E.154 for the coefficients of* [3, p. 441]*;*
- C_m *of Proposition E.158 in the bound of Theorem E.156 on $\delta m_{K,k}^2$.*

None of these depends on ε, ε_0, k, ℓ_T, M, or the deformation parameter u of Subsection 5.1. This proposition is the mass ingredient listed in Table 1.

Proof. Write $\nu_k = M^2 t_k^2$. Since $t_k \le \varepsilon_0$, (E.m2) gives $\nu_k \le 1$ for all $k \le K$. The estimates of [1, 2, 3, 4] in which the mass occurs are the following ten, which we call U1–U10.

(U1) [4, Theorem (1.9)–(1.12)] and [2, (2.58)]. No mass bound is needed (Lemma E.148(i)).

(U2) [1, (2.33)–(2.35)] = [4, (1.15)–(1.20)]. No mass bound is needed (Lemma E.148(ii)).

(U3) [2, Prop. 3.1, (3.26)], used through (3.29). This is Lemma E.149 with $c_m = 1$.

(U4) [2, (3.11)]. G_k'' is a Neumann $\Delta^{(k)}$-form plus non-negative terms. So the proof of Lemma E.149 gives $G_k'' \ge a_k\nu_k/(a_k+\nu_k) \ge \frac{3}{7}\nu_k$. The uses (3.17)–(3.20) need only $\|G_k''^{-1}\| \le 1/\nu_k + 1/a_k$.

(U5) [2, Lemma 2.4, (2.65), (2.76), (2.78)]. This is Lemma E.150 under (E.Φ).

(U6) [2, Lemma 2.7, (2.113)–(2.114)]. This is Lemma E.151 with Corollary E.152, under (E.Φ) and $\varepsilon_0 \le 1$.

(U7) [3, (3.16), p. 437] is Lemma E.153, and [3, p. 441] is Lemma E.154; both need only $\nu_j \le 1$. The counterterm bound is Theorem E.156 with Proposition E.158.

(U8) [2, (1.1), (2.1), (2.11)–(2.13), (3.21)–(3.23), (3.37)–(3.38)] and [3, (1.4), (1.19)–(1.22)]. Each of these steps is one of the following:
- a positive mass term dropped in an upper bound;
- an exact Gaussian identity;
- the diamagnetic inequality, which holds for every $M > 0$;
- a Gaussian measure whose propagator obeys (U1).

No mass bound is needed.

(U9) The counterterm graphs of [3, p. 418]. For $\delta m^2_{K,k}$ they are covered by (U7). The final number $\delta m^2_{\mathrm{fin}}$ is bounded in Proposition E.9, whose constants carry their own dependence on M and μ_*.

(U10) [2, (3.15)–(3.20)] and the other bounds that use the mass from below. These are not upper-bound uses: they bind at $m_1 > 0$ and are separate conditions.

Three further occurrences need only $\nu_k \le 1$:

- [1, (3.52), p. 621]: in that paper all fields are cut at $|\cdot| \le p$, so $\nu_k|\psi^{(k+1)}| \le p$;
- [24, (A.1), (A.7)–(A.8), p. 676]: the target $Cp\lambda(t)^{-1/4}$ has the size of the field bound of (A.1);
- [3, p. 441], which is Lemma E.154 in (U7).

The scalar operators carry no weight in u, so none of these bounds depends on u. Finally, every left side (ν_k, and $\lambda^{-1/4}M^2t^{7/4}$) is increasing in M, and every constant listed is free of M. So the bounds at m_2 give those at every $M < m_2$, in particular at m_1. The lower bounds on the mass are strongest at m_1; they are separate conditions. □

Bounds on $\delta m^2_{\mathrm{fin}}$ and E_{fin} uniform in the torus. Let $f \in \{s, v\}$ label the scalar and photon fields, with masses $m_s = M \in \{m_1, m_2\}$ and $m_v = \mu_*$. The number $\delta m^2_{\mathrm{fin}} := \delta m^2 - \delta m^2_K$ is the coefficient of [3, (1.26)], and E_{fin}, E_K are the vacuum pieces of Definition E.12, so that $E_1 = E_{\mathrm{fin}} + E_K$ (Lemma E.14); these are the final c-numbers of [3, (1.26), (1.30)], in the grading of Definition E.12. They are fixed numbers, so by Lemma E.14(i) any exact decomposition of the lines above ε_0 may be used to bound them. For $k > K$ put $t_k := L^{k-K}\varepsilon_0$. Let

$$\bar\ell := \min(1/M, 1/\mu_*, L), \qquad \bar K := \max\{k : t_k \le \bar\ell\}.$$

For both fields we use the exact identity $C_f^\varepsilon = G_{K,f}^\varepsilon(0) + \sum_{j=K}^{\bar K-1} H_{j,f} + R_{\bar K,f}$, where the $H_{j,f}$ are King's fluctuation covariances and $R_{\bar K,f} := C_f - G_{\bar K,f}(0)$.

The common top index, the lemma on intermediate subgraph scales, the top-cluster lemma and the two bounds are in Subsection E.2, at every Higgs mass $M \in (0, m_2]$: properties (P1)–(P7) there are the facts about the common top index (the inclusion $\bar K \le \min_f K_f$ is (P7)), Lemma E.7 is the lemma on intermediate scales, Lemma E.8 the top-cluster lemma, and Propositions E.9 and

E.21 the bounds, with constants that keep M_b. The inputs (VS), (Rem) and (S6) are proved in Subsection E.3. The dependence on u is treated next.

The photon lines at $u \neq 0$. Let $M \in (0, m_2]$ and assume (E.1). For every index j put $c_j := \kappa\mu_*^2 L^2 t_j^2$. In the slice model at (u, M) every vector line carries $C_B(u) = C_{\mu_*} + u\Gamma^B$ (see (5.1)) and every scalar line carries C_s; the mass counterterm is the vertex [3, (1.7)] with the number $\delta m^2_{\mathrm{ref}}$ as its coefficient, built from C_{μ_*}-lines and the propagators at $u = 0$. $E_1(u)$ is the vacuum coefficient of [3, (1.26), (1.30)] in this model, $E_K(u)$ is the same expression with $G_K(u) := G_K + u\Gamma^B$ for the vector lines and $G_{K,s}(0)$ for the scalar lines, and

$$E_{\mathrm{fin}}(u) := E_1(u) - E_K(u), \tag{E.9}$$

all graded as in Definition E.12, with every δm^2 kept as a number. A vector line of index $i < K$ is $H_i + u\delta_i H_i$; we call H_i its C-part and $u\delta_i H_i$ its D-part (a *D-line*).

Lemma E.160 (colour expansion of $E_{\mathrm{fin}}(u)$)**.** *For every $u \in \mathbb{R}$, $E_{\mathrm{fin}}(u) = \sum_{m=0}^{N_\ell} u^m E^{(m)}_{\mathrm{fin}}$ with $E^{(0)}_{\mathrm{fin}} = E_{\mathrm{fin}}(0)$, where $u^m E^{(m)}_{\mathrm{fin}}$ is the sum of the values of the coloured graphs with exactly m D-lines. Each such graph has at least one line from $\{C_{\mu_*} - G_K,\ C_s - G_{K,s}(0)\}$; these lines do not depend on u, and all D-lines have index $< K$. Moreover $\delta m^2_{\mathrm{ref}}$, δm^2_K and $\delta m^2_{\mathrm{fin}}$ do not depend on u.*

Proof. Graph values are multilinear in the line kernels. Since $C_B(u) - G_K(u) = C_{\mu_*} - G_K$, the graphs of $E_1(u)$ that are not in $E_K(u)$ are exactly those with at least one line $C_{\mu_*} - G_K$ or $C_s - G_{K,s}(0)$. Write $G_K(u) = \sum_{i<K}(1 + u\delta_i)H_i$ and expand every vector line of index $i < K$ into its C- and D-parts. A graph with m D-lines carries u^m, and $m \leq N_\ell$. Finally, $\delta m^2_{\mathrm{ref}}$ and its pieces δm^2_K, $\delta m^2_{\mathrm{fin}}$ are built from C_{μ_*}-lines and from the $u = 0$ propagators, so they carry no u. □

Lemma E.161 (the D-line in scale-j units)**.** *Let $i < K \leq j \leq \bar K$ and $\eta_j := L^{-j}$, so that $L^i\eta_j = t_i/t_j \leq L^{-1}$. Then*

$$\delta_i = c_j\,(L^i\eta_j)^2, \qquad c_j \leq c_{\bar K} \leq \kappa L^2(\mu_*\bar\ell)^2 \leq \kappa L^2.$$

The same identity holds for $i < j \leq K$, and then $c_j \leq c_K = \kappa\mu_*^2 L^2\varepsilon_0^2$. If ε_0 is a power of L, then $c_{\bar K}$ does not depend on ε, ε_0 or ℓ_T.

Proof. $\delta_i = \kappa\mu_*^2 t_{i+1}^2 = \kappa\mu_*^2 L^2 t_j^2 (t_i/t_j)^2$, and $j \mapsto c_j$ is increasing. By (P1), $t_{\bar K} \leq \bar\ell \leq 1/\mu_*$. If ε_0 is a power of L, the t_k run through powers of L, and since $\varepsilon_0 \leq \bar\ell$, $t_{\bar K}$ is the largest power of L that is $\leq \bar\ell$. □

Proposition E.162 (the final c-numbers)**.** *Assume $\varepsilon_0 \leq \min(1/m_2, 1/\mu_*, L)$, with ε_0 a power of L. Then at both anchor masses $M \in \{m_1, m_2\}$ the bounds of Propositions E.9 and E.21 hold, uniformly in $\varepsilon = 2^{-K}\varepsilon_0$, $x \in T_\varepsilon$ and the tori T_j, and the bound on $\delta m^2_{\mathrm{fin}}$ does not depend on u. For every $u \in \mathbb{R}$, $|E_{\mathrm{fin}}(u)| \leq \bar\sigma_N(u)\hat B$, where $\hat B$ is as in Statement K.7 and*

$$\bar\sigma_N(u) := \big(1{+}|u|\,\kappa\mu_*^2 L^2 t_{\bar K}^2\big)^{N_\ell} \leq \big(1{+}|u|\,\kappa L^2\big)^{N_\ell}; \qquad \bar\sigma_N(1) \leq (1{+}\kappa L^2)^{N_\ell}, \quad \bar\sigma_N(-\eta) \leq (1{+}\eta\kappa L^2)^{N_\ell}.$$

These numbers do not depend on ε, ε_0 or ℓ_T. The torus enters only through Z_s and Z_v, which are non-increasing in ℓ_T; hence the supremum over $\ell_T \geq \ell_0$ is at most the value at $\ell_0^\natural$, a number fixed at stage 0 of Appendix A, before ε_0. In particular the one-loop constant of the bound on $\delta m^2_{\mathrm{fin}}$ depends on $\ell_0^\natural$.

Proof. For $M \leq m_2$, $\bar\ell(M) \geq \bar\ell(m_2)$, so the hypothesis gives the first half of (E.1) at $M = m_1$ and $M = m_2$; the second half, $\ell_T \geq L^3$, holds on the tori of the paper, since $\ell_T \geq \ell_0 \geq \ell_0^\natural \geq 8 = L^3$ (Remark E.23(i)). Propositions E.9 and E.21 then apply at both masses, and Statement K.7 gives the bound at $u \neq 0$. The u-independence of $\delta m^2_{\mathrm{fin}}$ is Lemma E.160, the bounds on $\bar\sigma_N$ are Lemma E.161 (note $c_{\bar K} = \kappa\mu_*^2 L^2 t_{\bar K}^2$), and the monotonicity in ℓ_T is (P6), since $t_{\bar K}$, and with it $\bar\sigma_N$, does not depend on ℓ_T. □

Remark E.163 (the factor c_K). The argument does not give the smaller factor $(1+|u|\,c_K)^{N_\ell}$. Bounding a D-line by the number $|u\delta_i| \le |u|\,c_K$ leaves the kernel $(t_i/\varepsilon_0)^2 H_i$, which has dimension $-d+4$ only relative to the scale ε_0. A subgraph that contains a D-line, has no counterterm and has lowest external index $j_0 > K$ needs the factor $(t_i/t_{j_0})^2$, which costs $c_{j_0} = c_K(t_{j_0}/\varepsilon_0)^2$. When every index is $\le K$, as for the pieces of the flow below ε_0, $c_{j_0} \le c_K$, and the factor $(1+|u|\,c_K)^{N_\ell}$ is valid there by Theorem H.24(c),(d).

Lemma E.164 (the hypothesis of Proposition E.162 is a condition of Appendix A)**.** *Every ε_0 admissible in the order of choices of Appendix A satisfies $\varepsilon_0 \le \min(1/m_2, 1/\mu_*, L)$.*

Proof. By stage 10a, $\varepsilon_0 \le \varepsilon_1 \le 1 < L$ and $\varepsilon_0 \le \varepsilon_{(\mathrm{E.md})} \le 1/m_2$; (E.m2) reads $m_2^2\varepsilon_0^2 \le 1$. The bound (E.dom), $\mu_*\varepsilon_0 \le 0.2$ (stages 0 and 10a), gives $\varepsilon_0 \le 0.2/\mu_* < 1/\mu_*$. Since $m_1 < m_2$, also $\varepsilon_0 \le \bar\ell$ at $M = m_1$. □

The vacuum classes at the last scale. [3, p. 422] treats, for the mass counterterms, the terms that pass from scale k to $k+1$ by cutting the index-k line, and adds: "The corresponding considerations for vacuum energy counterterms are even simpler and we omit them." We write out the omitted vacuum case, in the grading of Definition E.12; Remark E.13 shows why the coefficients are not opened. Write $t_k := L^k\varepsilon$, $e(t)^2 = e^2 t$, $\lambda(t) = \lambda t$, $o := \alpha/2+\beta$, and $P(t) := \sum_{2\le\alpha+2\beta\le 2\bar n} |e(t)|^\alpha \lambda(t)^\beta$. By Definition E.12, $\mathcal{V}_k$ is the sum of the terms of the bare vacuum graphs of orders $1 \le \alpha+\beta \le \bar n$ in which every line has index $\le k$ and at least one has index k. In such a term every coefficient is split at index $k+1$ by [3, (1.26), (1.28), (1.29)], $\delta m^2 = \delta m^2_{\mathrm{fin}} + \delta m^2_{K,k+1} + \delta m^2_{k+1}$. Following [3, p. 423], a vertex carrying $\delta m^2_{\mathrm{fin}}$ or $\delta m^2_{K,k+1}$ is a *finite* vertex of degree 0, localised by the unit cube of its position; a vertex carrying the part $\delta m^2_{G_0}$ of δm^2_{k+1} belonging to a two-scalar-leg graph G_0 has degree $D(G_0)$. Localisations are those of [3, p. 420]. In scale-k units, $\delta m^2_{K,k} t_k^2 = O(t_k)$ [3, p. 418].

Lemma E.165 (exact cut)**.** *Fix a total order of the lines of every bare vacuum graph, and decompose the index sums by Balaban's orderings $\bar l = (l(1), \dots, l(m))$ [3, (3.4), p. 424], with ties broken by that order. Then every term of $\mathcal{V}_k$ belongs to exactly one ordering, and: (i) the last line $l(m)$ has index k; (ii) the counterterm operations active in $\bar l$ act on divergent components of $G_1 \subset \cdots \subset G_{m-1}$, none of which contains $l(m)$, so every member of the vacuum class contributing to $\bar l$ contains $l(m)$ as a line of its bare graph. Cutting $l(m)$, i.e. treating its index-k kernel $K_k(x,x')$ as a two-variable external field [3, p. 419], maps the class term by term, with the same coefficients, onto two-leg classes of $G \setminus l(m)$, in which every line has index $\le k$ and every counterterm vertex carries δm^2_{k+1} or a finite coefficient.*

Proof. (i) is the definition of $J(\bar l)$ [3, p. 424], since all indices are $\le k$ and one equals k. (ii) By construction $l(m) \notin G_i$ for $i < m$, and $G_m = G$. A counterterm operation replaces a divergent component $S \subset G_i$ by the vertex δm^2_S, or by the other members of its class [3, pp. 429–432], and does not touch the lines outside S; hence $l(m)$ remains a line. The lines of S have indices $\le k$, so δm^2_S is a part of δm^2_{k+1}. The parts of a coefficient with a line of index $> k$ form $\delta m^2_{K,k+1}$, a finite vertex. □

Lemma E.166 (cut classes)**.** *Let G be a bare vacuum graph, with degree D as in* [3, (2.1)–(2.3), pp. 422–423]*, let ℓ be one of its lines, and let $n_\partial(\ell) \in \{0,1,2\}$ be the number of differentiated legs among its two endpoints.*

(a) *If $G \setminus \ell$ is connected, then $D(G\setminus\ell) = D(G) + 1 + n_\partial(\ell)$.*
(b) *If G has no finite vertex, then $o(G) \in \mathbb{Z}_{\ge 1}$ and $D(G) = o(G) - 3$. Otherwise $D(G) \ge -1$, with equality only if G is one finite vertex closed by one line.*
(c) *If $G\setminus\ell$ is disconnected, then ℓ is a photon line and each component has exactly one external leg, a vector leg. Both components vanish* [3, p. 434].

(d) *Every component of $G \setminus \ell$ with $D \le 0$ is one of the following. For a scalar ℓ: C1 ($o = 1$, $n_\partial = 0$, $D = -1$), C2 ($o = 2$, $n_\partial = 0$, $D = 0$), C1′ ($o = 1$, $n_\partial = 1$, $D = 0$), or a bare finite vertex with two legs, when G is that vertex closed by ℓ. For a photon ℓ: C3 ($o = 1$, $D = -1$), C4 ($o = 2$, $D = 0$), or a vanishing pair as in (c). All other components have $D \ge 1$; this includes every G with $o \ge 3$.*

Proof. (a) In [3, (2.1)] the two endpoint legs of ℓ change from internal, of dimension $-(d-2)/2 = -\frac{1}{2}$, to external, of dimension 0, and a differentiation acting on such a leg changes from -1 to 0. The numbers of factors η and the counterterm degrees in (2.3) are unchanged, and $-d$ is subtracted once, since $G \setminus \ell$ is connected. (b) By [3, p. 423, (i)–(iv)] with $d = 3$: $D(v) = 1 = o(v)$ for (1.6), and $D(v) = n/2 = o(v)$ for (1.8) and (1.10) with n vector legs and no background leg. For a vertex carrying $\delta m^2_{G_0}$, $D(v) + \deg = 2 + D(G_0)$; by induction on the order, $D(G_0) = o(G_0) - 2$ for the two-scalar-leg graph G_0, which has no differentiated external leg, because δm^2 is built from the underived part of [3, (1.21)]. Hence $D(v) + \deg = o(v)$, and (2.3) gives $D(G) = o(G) - 3$. Every vector line uses two vector legs, so α is even and $o \in \mathbb{Z}$; since $o(v) \ge \frac{1}{2}$, $o \ge 1$. A finite vertex has $D(v) = 2$ and degree 0, and every other vertex has $D(v) \ge \frac{1}{2}$; so $D(G) \ge 2 - 3$, with equality only for one finite vertex closed by one line. (c) Every vertex has an even number of scalar legs, so a component of $G \setminus \ell$ has an even number of external scalar legs, and a scalar line is never a bridge. If ℓ is a photon bridge, each side carries one external vector leg and no external scalar leg; by [3, p. 434] each side contains a closed scalar strand with an odd power of q, and $\operatorname{tr} q^{2n+1} = 0$; this holds after the gauge transformation of [3, p. 433]. (d) By (a) and (b), a connected $G \setminus \ell$ without finite vertex has $D = o - 2 + n_\partial$, with $n_\partial = 0$ for photon lines. With a finite vertex, $D(G) = o' - 1$, where $o' \in \mathbb{Z}_{\ge 0}$ is the order of the other vertices, so $D(G \setminus \ell) = o' + n_\partial \ge 0$, with equality only for $o' = 0$, $n_\partial = 0$. Listing $D \le 0$ over $o \in \mathbb{Z}_{\ge 1}$ and $n_\partial \in \{0, 1, 2\}$ gives the stated classes. □

Cut classes. By Lemma E.166, a scalar cut gives the classes C1, C2 (with their δm^2-partners, Lemma E.169, except the one-particle-reducible chains of C2, which have no partner and are treated by the product mechanism of Step 3 of the proof of Lemma E.86) or C1′; a photon cut gives the Ward classes C3, C4, or vanishing pairs. The photon cut needs the Ward class, so the vacuum case is not simpler than the mass case.

Lemma E.167 (cuts inside opened counterterm vertices)**.** *Let G be as in Lemma E.166, and let v be a vertex* [3, (1.7)] *of G carrying the counterterm $\delta m^2_{G_0}$ of a graph G_0 of order $o_0 \in \{1, 2\}$. Let ℓ_1, ℓ_2 be the lines of G at v and u_1, u_2 the vertices of G_0 carrying its external legs. Let G^T be obtained from G by replacing v by G_0, with ℓ_i attached at u_i, and G^v from G^T by moving the end of ℓ_2 from u_2 to u_1 (the representation of* [3, p. 417]*). Degrees are those of* [3, (2.1)–(2.3)]*. Then: (a) G_0 is one-particle irreducible, and neither of its external legs is differentiated; (b) $D(G^v) = D(G^T) = D(G) = o(G) - 3$; (c) for every line ℓ of G_0, $G^v \setminus \ell$ and $G^T \setminus \ell$ are connected, and $D(G^v \setminus \ell) = D(G^T \setminus \ell) = o(G) - 2 + n_\partial(\ell)$; (d) hence every such cut with $D \le 0$ belongs to C1, C2, C1′, C3 or C4 with $o = o(G) \le 2$; no photon bridge and no bare vertex occurs.*

Proof. (a) By [3, (1.21), p. 416], $\Sigma^\varepsilon, \Sigma^\varepsilon_1, \Sigma^\varepsilon_2$ are given by amputated one-particle-irreducible graphs, and δm^2 is defined from Σ^ε, the part without differentiated external legs [3, p. 417]. (b) Each internal leg contributes $-\frac{1}{2}$ to (2.1), so $D(H) = \sum_w \eta(w) - I(H) - \partial(H) - d + \sum \deg$, with $\eta(w)$ the number of factors η at w, I the number of lines, ∂ the number of differentiations on lines and the last sum over unopened counterterm vertices. In G^T the vertex v ($\eta(v) = d$, $\deg v = D(G_0)$) is replaced by the vertices of G_0; since $I(G^T) = I(G) + I(G_0)$ and $\partial(G^T) = \partial(G) + \partial(G_0)$ by (a), $D(G^T) = D(G)$. Moving an undifferentiated leg-end changes no term, so $D(G^v) = D(G^T)$. Finally $D(G) = o(G) - 3$ by Lemma E.166(b). (c) $G_0 \setminus \ell$ is connected by (a) and contains u_1, u_2, and every path of G through v passes through it. Removing ℓ lowers I by 1 and ∂ by $n_\partial(\ell)$. (d) Combine (c) with the list in the proof of Lemma E.166(d). □

Remark E.168. In the grading of Definition E.12 the cut line $l(m)$ of Lemma E.165 is a line of the bare graph, so it never lies inside a coefficient. If the coefficient of v were opened and a line ℓ of G_0 were cut, the cut member would factor as $\sum_x R_v(x)c_{G_0,\ell}(x)$, where R_v is the value of G with coefficient 1 at v (a vertex of degree 0, so $D = o(G) - o_0 - 1$) and $c_{G_0,\ell}$ is the coefficient graph with ℓ as external field. If $o(G) - o_0 \le 1$, which by Lemma E.167(d) is the case for every such cut with $D \le 0$, the factor R_v has degree -1 or 0 and no partner in the cut family. This is the mechanism of Remark E.13.

Lemma E.169 (the partners are present)**.** *In the cut classes of Lemma E.165, every one-particle-irreducible component of type C1 or C2 occurs together with its mass counterterm, with coefficient in δm^2_{k+1}, and every component of type C3 or C4 occurs together with all connected two-vector-leg graphs of the same order whose lines have index $\le k$.*

Proof. Let H be a connected two-leg graph of order $o \le \bar{n}$ with lines of index $\le k$, and ℓ a line of index k of the kind of its legs. Closing H by ℓ gives a bare vacuum graph whose term lies in $\mathcal{V}_k$, and cutting ℓ gives back H. If H is a one-particle-irreducible two-scalar-leg graph, the vertex δm^2_H (a part of δm^2_{k+1}) closed by ℓ is also a bare vacuum graph in $\mathcal{V}_k$, and its cut is the counterterm member of H in the representation of [3, p. 417]. For two vector legs this applies to every connected two-vector-leg graph of the order, which is the class of [3, p. 431]. □

Lemma E.170 (localisation sum)**.** *Let $F = \sum_\Lambda F(\Lambda)$ be a sum over localisations $\Lambda = (\square(v))_{v\in G}$, i.e. N_s unit cubes of the scale-k lattice. Assume $|F(\Lambda)| \le A\,\mathrm{e}^{-\delta d(\Lambda)}$, where $d(\Lambda)$ is the length of a shortest tree through the cubes. Let W be a union of unit blocks and $F^{\mathrm{bad}} := \sum_{\Lambda\cap W\neq\emptyset,\ \Lambda\not\subset W} F(\Lambda)$. Then for every $r > 0$*

$$|F^{\mathrm{bad}}| \le A\,N_s\,S\Big[\#\{\square \subset W : \mathrm{dist}(\square, W^c) \le r\} + \mathrm{e}^{\delta(c_d+1)/2}\mathrm{e}^{-\delta r/2}\,\#\{\square \subset W : \mathrm{dist}(\square, W^c) > r\}\Big],$$

where $S := \sum_{D\ge 0}(c''_d(D+2)^d)^{N_s}\mathrm{e}^{-\delta D/2}$, $c_d := 2\sqrt{d}$, and c''_d is a lattice constant.

Proof. Attribute each bad Λ to the cube $\square_0$ of the first vertex (in a fixed order) whose cube lies in W; there are at most N_s choices of that vertex. Since Λ also has a cube outside W, $d(\Lambda) \ge \mathrm{dist}(\square_0, W^c) - c_d$. With $\square_0$ fixed, every other cube lies within $d(\Lambda) + c_d$ of $\square_0$, so at most $N_s(c''_d(D+2)^d)^{N_s}$ localisations have $d(\Lambda) \in [D, D+1)$. With $D_0 := (\mathrm{dist}(\square_0, W^c) - c_d - 1)_+$ and $\mathrm{e}^{-\delta D} \le \mathrm{e}^{-\delta D_0/2}\mathrm{e}^{-\delta D/2}$ for $D \ge D_0$, the sum over Λ with $\square_0 = \square$ is at most $AN_s\mathrm{e}^{-\delta D_0/2}S$. Use $\mathrm{e}^{-\delta D_0/2} \le 1$ if dist $\le r$, and $\mathrm{e}^{-\delta D_0/2} \le \mathrm{e}^{\delta(c_d+1)/2}\mathrm{e}^{-\delta r/2}$ if dist $> r$. Sum over $\square \subset W$. □

Lemma E.171 (the layer and the deep part)**.** *In scale-k units let $W := B^k(\Lambda^{(k)}_7)$ and $r := r(t_k)$. Every unit block $\square \subset W$ with $\mathrm{dist}(\square, W^c) \le r$ lies in $B^k(\Lambda^{(k)}_7 \cap \Lambda^{(k)c}_8)$, and every unit block of $B^k(\Lambda^{(k)}_8)$ has $\mathrm{dist}(\square, W^c) > r$.*

Proof. By [2, (2.8), p. 558], $\Lambda^{(k)c}_8$ is the union of the large blocks at distance $\le r$ from $\Lambda^{(k)c}_7$. If $\square \subset W$ and $\mathrm{dist}(\square, W^c) \le r$, the large block $b \supset \square$ has $\mathrm{dist}(b, W^c) \le r$, so $b \subset \Lambda^{(k)c}_8$. If $\square \subset B^k(\Lambda^{(k)}_8)$, its large block is not within r of $\Lambda^{(k)c}_7$. □

Lemma E.172 (degrees of open graphs)**.** *Let Q be a connected graph with $E = E_s + E_v \ge 1$ external legs (E_s scalar, E_v vector), n_∂ of them differentiated, m_f finite vertices and o' the order of its other vertices. Then*

$$D(Q) = o' + 2m_f - 3 + \tfrac{1}{2}E + n_\partial. \tag{E.10}$$

If Q does not vanish and $D(Q) \le 0$, then Q is a single vertex; or the tree [3, (2.4)] *(two vertices* [3, (1.8)] *with $n = 1$ joined by a scalar line carrying both derivatives); or a member of C1, C1′, C2, C3, C4 of Lemma E.166, or of the class C5 of the graphs with two scalar legs and one vector leg at $o = \frac{3}{2}$. A connected graph with a finite vertex and $E \ge 1$ has $D \ge 0$, with equality only for a single*

finite vertex or for a vanishing scalar loop through one finite vertex and one vertex with a single, external, vector leg.

Proof. External legs and derivatives contribute 0 in place of $-\frac{1}{2}$ and -1 [3, (2.1)–(2.3)]; summing the vertex degrees of Lemma E.166(b) gives (E.10). Constraints: E_s is even; $o \in \frac{1}{2}E_v + \mathbb{Z}$; every non-finite vertex has $o(v) \geq \frac{1}{2}$; a graph with an odd number of external vector legs and no external scalar leg vanishes, as in Lemma E.166(c). If $m_f = 0$, $D \leq 0$ means $o' + \frac{1}{2}E + n_\partial \leq 3$: $E = 1$ or $(E_s, E_v) = (0, 3)$ vanish; $E = 2$ gives $o' \in \{1, 2\}$, $o' + n_\partial \leq 2$, i.e. C1, C1′, C2 ($E_s = 2$) and C3, C4 ($E_v = 2$); $(2, 1)$ gives a single vertex ($o' = \frac{1}{2}$) or C5 ($o' = \frac{3}{2}$, $n_\partial = 0$); $E = 4$ needs $o' = 1$, $n_\partial = 0$: $(4, 0)$ is a single λ vertex (two $n = 1$ vertices joined by a photon have $n_\partial \geq 2$), $(2, 2)$ is a single $n = 2$ vertex or the tree [3, (2.4)], $(0, 4)$ needs $o' \geq 2$; $E \geq 5$ needs $o' \leq \frac{1}{2}$, impossible. If $m_f \geq 1$, $D \geq o' - 1 + \frac{1}{2}E$: for $o' = 0$, $D = 2m_f - 3 + \frac{1}{2}E \geq 0$, with equality only for one vertex with $E = 2$; for $o' = \frac{1}{2}$ the only vector leg is external, $E = 1$ gives $D = 2m_f - 2$, which is 0 only for the vanishing loop, and $E \geq 3$ gives $D \geq 1$; for $o' \geq 1$, $D \geq \frac{1}{2}$. □

Lemma E.173 (image kernels at scale k)**.** *Assume (T.nw), (E.r) and (T.per). Let $0 \leq k < K$, $S_k := \ell_T/t_k$ the side of T_ε in scale-k units, $\square$ a cube of side $3r(t_k)$ in scale-k units and ι its isometric lift (Lemma E.66). Let K be the kernel of a line of index $\leq k$ of a step-k cut class, scalar or vector, or a derivative of order ≤ 2 of it, in scale-k units. Then on $\square \times \square$*

$$K_T(x, x') = K_\infty(\iota x, \iota x') + \delta K(x, x'), \qquad \|\delta K\|_{1,\alpha} \leq C_1 \mathrm{e}^{-\delta_0 S_k/2} \leq C_1 \mathrm{e}^{-3\delta_0 r(t_k)},$$

with C_1 and δ_0 those of Lemma 5.22.

Proof. For the scalar kernels, Lemma 5.22(i) gives $K_T = \operatorname{Per} K_\infty$; for the vector kernels this is Statement K.8. The diameter of $\square$ is $3\sqrt{3}\, r(t_k)$; by (T.per) and the proof of Lemma 5.22(iv), $S_k \geq 6\sqrt{3}\, r(t_k) + 2$ (twice the diameter) and $S_k \geq 2/\delta_0$. Now Lemma 5.22(iii),(iv) give the bound. □

Lemma E.174 (image lines regroup into renormalised classes)**.** *Make the assumptions of Lemma E.173 and $R \geq R^{**}$. Fix $k < K$, a pair $(G, \bar{l})$ and a localisation Λ on T_ε. Cut $l(m)$ (Lemma E.165) and the lines between cubes at distance ≥ 1* [3, (3.1), p. 432]*; these are torus kernels bounded in torus distance and are not split. Regroup into classes* [3, p. 432]*; each connected component Γ is localised in a cube $\square_1$ of side $r(t_k) \geq \ell_{\bar{N}}$, centred in a cube $\square$ of side $3r(t_k)$* [3, p. 433]*. A* line position *of Γ is a line of a member of Γ; in a member carrying δm_S^2, written in the form of* [3, p. 417]*, the lines of S are the same line positions as in the parent member, and the members of a Ward–Takahashi class (*[3, pp. 431, 440–445]*, types C3, C4, C5) share their outer lines and have their own inner lines. Split each line position once, $K_s = K_s^\infty + \delta K_s$ (Lemma E.173), so that a counterterm member and its parent receive the same choice; for each Ward–Takahashi subclass Π_Q, from the outermost inwards, write $\Pi_Q = \Pi_Q^\infty + \Pi_Q^\delta$ with Π_Q^∞ the terms whose inner lines are all K^∞. Then $\Gamma = \Gamma^\infty + \sum_{J \neq \emptyset} \Gamma^J$, J the set of line positions carrying δK, with at most $2^{N_{\mathrm{lp}}}$ terms, $N_{\mathrm{lp}} = N_{\mathrm{lp}}(\bar{n})$, and: (a) Γ^∞ is the class lifted by ι to $\eta\mathbb{Z}^3$ with every propagator $G_k(0)$, i.e. the expression after the last substitution of* [3, p. 433]*, bounded by* [3, (3.3)]*; (b) in every Γ^J, replacing each line in J by a pair of external fields carrying δK_s gives a finite sum of renormalised classes on $\eta\mathbb{Z}^3$, each bounded by* [3, (3.3)] *with the δK_s among its external fields; (c) the per-localisation bound of Lemma E.86 holds on T_ε with C_P replaced by $C_P^T := C_P'(1 + C_1)^{N_{\mathrm{lp}}}$, $C_P' = C_P'(\bar{n}, \alpha_0)$, and the part with $J \neq \emptyset$ is at most $C_P' N_{\mathrm{lp}} (1 + C_1)^{N_{\mathrm{lp}}} \mathrm{e}^{-\delta_0 S_k/2} P(t_k) \mathrm{e}^{-\hat{\delta} d(\Lambda)}$, with $d(\Lambda)$ the tree length in torus distance and $\hat{\delta}$ as in Proposition E.175.*

Proof. $\square_1$ exists since $\ell_{\bar{N}} \leq r(t_k)$ by $R \geq R^{**}$, and $\square$ lifts by Lemma E.66. Every line position joins two points of $\square_1$, since the vertices of a counterterm member lie in the cubes of its lines [3, p. 420]; so Lemma E.173 applies. The vacuum terms are at zero background, so the expansion in B' and

the gauge transformation removing B_0 of [3, p. 433] are trivial; this gives (a). (b) Induction on the number of lines. Let M be a member of Γ^J and Q a non-vanishing connected component of $M \setminus J$ with $D(Q) \le 0$; Lemma E.172 applies. Single vertices and the tree [3, (2.4)] are bounded as in [3, (2.4), Props. 2.1–2.2]; a component with a finite vertex has $D \ge 0$, with equality only for a single vertex or a vanishing loop; C1′ is the finite vertex of [3, (3.22)–(3.24)]. Otherwise Q is of type C1, C2, C3, C4 or C5; its degree is the same in M and in $M \setminus J$, so Q is a divergent subgraph of M without line position in J, and its partners are members of Γ [3, p. 432]. For one-particle-irreducible C1, C2, the partner $M[Q \to \delta m_Q^2]$ has, in the form of p. 417, the line positions of M, hence the same choices, and the pair is Bałaban's renormalised pair on $\eta\mathbb{Z}^3$. The one-particle-reducible members of C2 are treated by Step 3 of the proof of Lemma E.86; their pieces share line positions, and if the joining line lies in J the pieces are two C1 pairs. For C3, C4, C5 the term contains Π_Q^∞, the lifted class on $\eta\mathbb{Z}^3$, where the Ward identities hold exactly (Lemma E.65, Proposition E.67) and [3, (3.25)–(3.38)] apply as printed; a term with Π_Q^δ consists of single members with an inner image line, treated by the same case analysis with fewer lines. A counterterm member $M[S \to \delta m_S^2]$ with $S \cap J \ne \emptyset$, written in the one-vertex form of p. 417, carries $c_S^J(x) = \sum_{x'} \eta^3 \Sigma_S^J(x,x')$; by the induction hypothesis for the class of S (subgraph counterterms included in δm_S^2 by [3, (1.23), p. 430]) and [3, (3.3)], $|c_S^J| \le C \prod_{s \in J \cap S} \|\delta K_s\|$ times the couplings of S. So it is a graph with one finite vertex of degree 0; by Lemma E.172 its components through that vertex need no partner, and its other components are subgraphs of M outside S. (c) Each term is bounded by [3, (3.3)], linear in the norms of the external fields; the exponential factor comes from the cut kernels as in Lemma E.86. Sum over the at most $2^{N_{\mathrm{lp}}}$ terms with $\|\delta K_s\| \le C_1 \mathrm{e}^{-\delta_0 S_k/2}$. □

Proposition E.175 (the vacuum classes at the last scale)**.** *Assume (T.nw), (E.r), (T.per), $R \ge R^{**}$ and $\delta_0 R \ge 4$. Let $\delta_T > 0$ be the tree rate of* [3, (1.33)] *as established in Proposition E.97, δ_P the propagator rate of* [3, (2.10), p. 426]*, and $\hat\delta := \min(\delta_T, \delta_P)$. Call a localisation of a term of $\mathcal{V}_k$ bad if it meets both $W = B^k(\Lambda_7^{(k)})$ and W^c, and let $\mathcal{V}_k^{\mathrm{bad}}$ be the sum of the terms with bad localisations. At the step $k \to k+1$, on T_ε with all lines torus lines,*

$$|\mathcal{V}_k^{\mathrm{bad}}| \le C_9\, P(t_k) \Big[|B^k(\Lambda_7^{(k)} \cap \Lambda_8^{(k)c})| + \mathrm{e}^{-\hat\delta r(t_k)/2} |B^k(\Lambda_8^{(k)})| \Big],$$

where $|\cdot|$ counts scale-k unit blocks, $C_9 = C_P^T N_G N_s S_{\hat\delta}(1 + \mathrm{e}^{\hat\delta(c_3+1)/2})$, C_P^T is the constant of Lemma E.174(c), N_G is the number of pairs (bare vacuum graph, ordering) of order $\le \bar n$, N_s the largest number of localised vertices, $S_{\hat\delta} := \sum_{D \ge 0} (c_3''(D+2)^3)^{N_s} \mathrm{e}^{-\hat\delta D/2}$ and $c_3 = 2\sqrt{3}$. C_9 is independent of ε, k, the domains and ℓ_T. Let $\mathcal{V}_k^{\mathrm{img}}$ be the part of $\mathcal{V}_k$, over all localisations, in which some internal line carries its image kernel. Then

$$|\mathcal{V}_k^{\mathrm{img}}| \le C\, P(t_k)\, \mathrm{e}^{-\delta_0 S_k/2}\, |T|\, t_k^{-3}, \qquad \sum_{k<K} |\mathcal{V}_k^{\mathrm{img}}| \le C\, |T|\, \varepsilon_0^{-3} \mathrm{e}^{-\delta_0 \ell_T/\varepsilon_0} \le C\, |T|\, \varepsilon_0^{-3} \mathrm{e}^{-\delta_0 \ell_0/\varepsilon_0}.$$

Proof. Per localisation. By Lemma E.165, $\mathcal{V}_k$ is a finite sum over pairs (graph, ordering) of two-leg classes of $G \setminus l(m)$ with external field K_k. By Lemma E.166(c), in every non-vanishing term $G \setminus l(m)$ is connected, so the tree of [3, (1.33)] joins all cubes of Λ. By Lemma E.166(d), the components with $D \le 0$ are: C1, C2 with their counterterms (Lemma E.169), the one-particle-reducible C2 members by Step 3 of the proof of Lemma E.86; C1′, a finite vertex [3, (3.22)–(3.24)]; C3, C4 with their Ward–Takahashi classes [3, (3.25)–(3.32)], run after the transfer of [3, p. 433] to a non-wrapping cube (Lemmas E.65, E.66), the local AA vertex left by [3, (3.30)] having coefficient $O(1)$ at $j_0 = k$ and being multiplied by $K_k(x,x) = O(1)$; and a bare finite vertex, localised in one cube and never bad. In the grading of Definition E.12 the line $l(m)$ is never inside a coefficient (Remark E.168). In scale-k units, $\|hK_kh'\|_{1,\alpha_0} \le O(1)\mathrm{e}^{-\delta_P \operatorname{dist}(\operatorname{supp} h, \operatorname{supp} h')}$ for scalar and vector lines [3, (2.10)–(2.11), p. 426]. Proposition E.97, applied with this external field [3, p. 419] at scale

$k+1$, and Lemma E.174(c) on the torus give $|\mathcal{V}_{G,\bar{l}}(\Lambda)| \le C_P^T P(t_k) \mathrm{e}^{-\hat{\delta} d(\Lambda)}$; summing over the N_G pairs gives the hypothesis of Lemma E.170 with $A = C_P^T N_G P(t_k)$ and $\delta = \hat{\delta}$.

Localisation sum. Lemma E.170 holds on T_ε with torus distances, since a torus ball contains at most as many unit cubes as the ball of the same radius in $\mathbb{Z}^3$. Apply it with W as above and $r = r(t_k)$. By Lemma E.171, which uses the torus distances of [2, (2.8)], the blocks of W within r of W^c are at most $|B^k(\Lambda_7^{(k)} \cap \Lambda_8^{(k)c})|$, and the other blocks of W at most $|B^k(\Lambda_7^{(k)} \cap \Lambda_8^{(k)c})| + |B^k(\Lambda_8^{(k)})|$. With $\mathrm{e}^{-\hat{\delta} r/2} \le 1$ this gives the first bound. Vacuum terms have no external legs, so the cut-off functions of external legs play no role.

Torus lines. The cut kernels are torus kernels bounded in torus distance; the internal lines of each component are split on a non-wrapping cube by Lemma E.173, and by Lemma E.174(b) every resulting term is a sum of renormalised classes on $\eta\mathbb{Z}^3$ with the image kernels as external fields. For $\mathcal{V}_k^{\mathrm{img}}$, sum the bound of Lemma E.174(c) over the localisations anchored at each of the $|T| t_k^{-3}$ unit blocks; with the anchor fixed the sum over Λ is at most $N_s S_{\hat{\delta}}$, and $P(t_k) \le P(\varepsilon_0)$. Put $a := \delta_0 \ell_T/(2\varepsilon_0)$ and $t_k = \varepsilon_0 L^{-n}$, $n \ge 1$; then $t_k^{-3} \mathrm{e}^{-\delta_0 S_k/2} = (L^n/\varepsilon_0)^3 \mathrm{e}^{-aL^n}$. Since $r(\varepsilon_0) \ge R$ and $\delta_0 R \ge 4$, (T.per) gives $a \ge 3\sqrt{3}\,\delta_0 r(\varepsilon_0) \ge 12\sqrt{3} > \log(2L^3)$ for $L \le 3$, so consecutive terms decrease by a factor $\le L^3 \mathrm{e}^{-a} \le \frac{1}{2}$, and $\sum_{n\ge 1} (L^n/\varepsilon_0)^3 \mathrm{e}^{-aL^n} \le 2L^3 \varepsilon_0^{-3} \mathrm{e}^{-aL} \le 2L^3 \varepsilon_0^{-3} \mathrm{e}^{-\delta_0 \ell_T/\varepsilon_0}$, since $L \ge 2$. Finally $\ell_T \ge \ell_0$. □

Appendix F. Assembly of Theorem 5.2

This appendix assembles the proof of Theorem 5.2 from its ingredients (Appendix E and Proposition 5.21) and the step lemmas of Bałaban's inductive argument, which are stated in Subsection F.2. The collar exponent is $\kappa_0 = \frac{1}{5}$ (Lemma F.1); it is not carried to other exponents, such as $\kappa_B = \frac{16}{5}$ and the exponents of (E.11), which are separate quantities.

F.1. **The assembly.** The ingredients are those of Table 1: Propositions E.64, E.67, E.71, E.76, E.110, 5.21, E.159, E.162 and E.175. The argument is Bałaban's inductive upper-bound argument of [2, Sects. 2–3], run on the torus for the slice model with the photon weights $w_j := 1 + u\delta_j$ (δ_j as in (5.1)). Besides the ingredients it uses six groups of step results, (S0)–(S5), stated in Subsection F.2. Notation. $L = 2$, $\varepsilon = 2^{-K}\varepsilon_0$, and $t_k := L^k \varepsilon$ ($0 \le k \le K$), so $t_0 = \varepsilon$ and $t_K = \varepsilon_0$. For $X \subset T$, $|X|_k$ is the number of blocks of side t_k in X, and $N_k := |T| t_k^{-3}$. Put $\kappa_0 := \frac{1}{5}$ and $\kappa_B := \frac{16}{5}$. (Every $\kappa_0 < \frac{1}{4}$ works; $\kappa_0 = \frac{1}{4}$ does not, because some collar terms are $t^{1/4}$ times a power of $\log t^{-1}$, so that $C_{\kappa_0} = \infty$ at $\frac{1}{4}$.) The admissible large-field data $\mathbf{\Lambda}_k = (\Lambda_0^{(j)})_{j<k}$, the regions $\Lambda_i^{(k-1)\prime}$, $\Omega_1 := B^k(\Lambda_7^{(k-1)\prime})$, and the collars $\mathrm{Col}_j := \Omega_1^{(j)} \setminus \Omega_1^{(j+1)}$ are those of [2, (2.7)–(2.8), (2.43), pp. 558, 566–570], with $\mathrm{Col}_0 := \Lambda_7^{(0)c}$. $\mathcal{Z}_u$ is the integral defining the partition function of the slice model at (u, M), taken with the *Lebesgue* photon measure. That is, $d\mu_{N(0,C_B(u))}(A) = N_w \, \mathrm{e}^{-\frac{1}{2}\langle A, C_B(u)^{-1} A\rangle} dA$ and $\mathcal{Z}_u := \int dA \, \mathrm{e}^{-\frac{1}{2}\langle A, C_B(u)^{-1} A\rangle} Z_\varphi(A; a_{K,T}(M))$. Put $E_0^w := \log \mathcal{Z}_u|_{e=\lambda=0} = -\log N_w + E_{0,s}(M)$ and $E_1(u) := T_{\bar{n}}[\Phi_M(u)] - E_{0,s}(M)$. We write $E_1(u) = E_{\mathrm{fin}}(u) + E_K(u)$. Here $E_{\mathrm{fin}}(u)$ is the sum of the vacuum graphs with at least one line of index $\ge K$. It is a c-number, independent of the field, of $\mathbf{\Lambda}$ and of k.

The collar exponent.

Lemma F.1 (the collar exponent)**.** *Let $p(t) = b_0(1 + \log t^{-1})^p$ with $p = 6$, and let $0 < \kappa_0 < \frac{1}{4}$. For $a > 0$ and $n \ge 0$ put $S(a,n) := \sup_{0<t\le 1} t^a (1 + \log t^{-1})^n$. Then $S(a,n) = \mathrm{e}^{a-n}(n/a)^n$ if $n \ge a$, and $S(a,n) = 1$ otherwise. Hence, for $t \le 1$ and $q \le 4$,*

$$t^{1/4} p(t)^q \le b_0^q \, S(\tfrac{1}{4} - \kappa_0, 6q)\, t^{\kappa_0}, \qquad t^{1/2} p(t)^2 \le b_0^2 \, S(\tfrac{1}{2} - \kappa_0, 12)\, t^{\kappa_0},$$

and $t \le t^{\kappa_0}$, $t^3(1 + \log_2(\varepsilon_0/t)) \le C(\kappa_0)\, t^{\kappa_0}$ for $t \le \varepsilon_0 \le 1$. For $\kappa_0 = \frac{1}{4}$ and $q \ge 1$ no constant C satisfies $t^{1/4} p(t)^q \le C\, t^{1/4}$ for all $t \in (0,1]$. At $\kappa_0 = \frac{1}{5}$, $\log S(\frac{1}{20}, 24) = 124.22\ldots$.

Proof. Put $s = \log t^{-1} \geq 0$ and $g(s) = \mathrm{e}^{-as}(1+s)^n$. Then $g'(s) = g(s)\,(n/(1+s) - a)$, so g increases for $1 + s < n/a$ and decreases afterwards. If $n \geq a$, the maximum is at $1 + s = n/a$, where $g = \mathrm{e}^{-a(n/a-1)}(n/a)^n$. If $n < a$, the maximum is $g(0) = 1$. The displayed bounds follow from $t^{1/4}p^q = b_0^q\, t^{\kappa_0} \cdot t^{1/4-\kappa_0}(1+\log t^{-1})^{6q}$ and the analogous identity for $t^{1/2}p^2$. Also $t \leq t^{\kappa_0}$ since $\kappa_0 \leq 1$, and $t^{3-\kappa_0}(1 + \log_2(\varepsilon_0/t)) \leq S(3-\kappa_0, 1)\max(1, 1/\log 2)$. At $\kappa_0 = \frac{1}{4}$ the ratio $t^{1/4}p(t)^q/t^{1/4} = p(t)^q$ tends to ∞ as $t \downarrow 0$. □

Sizes $t^{1/4}p^q$ occur in [2, (2.22), p. 561] ($q = 3$), [2, (2.78), p. 574] ($q = 4$) and the count of [3, p. 421] ($q = 2, 3$). By Lemma F.1 they are at most $C(\kappa_0)t^{\kappa_0}$ per collar block, with $C(\kappa_0) < \infty$ at $\kappa_0 = \frac{1}{5}$. Every other use of κ_0 needs only $0 < \kappa_0 \leq 1$.

Proposition F.2 (assembly of Theorem 5.2)**.** *Assume the hypotheses of Theorem 5.2; in particular (E.fin), $\varepsilon_0 \leq \min(1/m_2, 1/\mu_*, L)$ (Lemma E.164). Assume the ingredients (Propositions E.64–E.175) and the step results:*

- (S1) *the scalar small-field step (Lemma F.8);*
- (S2) *the photon small-field step (Lemma F.14);*
- (S3) *the large-field part of the step (Lemma F.22), with the region sum of Lemmas E.94 and E.95;*
- (S4) *the renormalisation flow (Lemmas F.32, F.31, Propositions E.64, E.62 and Lemma E.170);*
- (S5) *the closing step (Lemma F.39), with Lemmas F.38, F.35, F.36, F.37, E.63 and the order of choices (Lemma F.40).*

The base identity (S0) is Lemma F.7; it rests on Lemma F.5, Theorem F.6 and the identification of the counterterms (Proposition 4.7). Then (5.2) *holds for every $K \geq 0$ and every $j \geq 1$, at $(u, M) \in \{(1, m_2), (-\eta, m_1)\}$, with*

$$C_+^B(u; M) = C_{F1}\varepsilon_0^{\kappa_0-3} + B_E^{\mathrm{fin}}(u, M) + B_{\mathfrak{E}} + B_{GS} + \tfrac{1}{2}B_{F3} + B_{\mathrm{comb}}, \tag{F.1}$$

where $\varepsilon_0^{\kappa_0-3} = \varepsilon_0^{-14/5}$ and the six terms are defined in Steps 4 and 5 of the proof. None of them depends on ε, K or ℓ_T, and none blows up at $u = -\eta$ or at $M = m_1$.

Together with Propositions E.64–E.175 and the step lemmas (S1)–(S5) of Subsection F.2, Proposition F.2 gives Theorem 5.2. The identification of the photon quadratic form and of N_w is made in the Notation paragraph above and in Theorem F.6.

Proof. The proof composes the step lemmas. Steps 1 and 3 use the convention (G) of Remark 5.3(ii): the vacuum counterterms are global c-numbers [3, (1.30), p. 419]. We write $E_{\mathrm{fin}}(u)$ and $E_{K,k}(u)$ for the parts of the vacuum series $E_1(u)$ defined in Lemma F.31. Here $E_{\mathrm{fin}}(u)$ is the sum of the monomials with at least one line of index K, and $E_{K,k}(u)$ is the sum of the monomials with at least one line index in $[k, K)$ and none equal to K. Thus $E_{K,0}(u) = E_K(u) = E_1(u) - E_{\mathrm{fin}}(u)$ and $E_{K,K}(u) = 0$.

Step 0 (reduction). By definition $e(\varepsilon; u; M) = -|T|^{-1}T_{\bar{n}}[\Phi_M(u)]$, so the left side of (5.2) equals $|T|^{-1}(\Phi_M(u) - T_{\bar{n}}[\Phi_M(u)])$. The photon measure in Φ_M is normalised, so $\Phi_M(u) = \log(N_w\mathcal{Z}_u)$ by Theorem F.6(a),(b) (see the proof of Lemma F.7). The order-zero term of $T_{\bar{n}}[\Phi_M(u)]$ is $E_{0,s}(M)$ (Remark 5.3(i)). Hence

$$\mathrm{e}^{\Phi_M(u) - T_{\bar{n}}[\Phi_M(u)]} = \mathrm{e}^{-E_0^w - E_1(u)}\,\mathcal{Z}_u,$$

and (5.2) is equivalent to $\mathrm{e}^{-E_0^w - E_1(u)}\mathcal{Z}_u \leq \mathrm{e}^{C_+^B(u;M)|T|}$. The statement has to use the normalised photon, because the secant argument of Proposition 6.6 needs convexity in u, while $\log\det C_B(u)$ is concave. Finally, $\bar{\delta} = \kappa\mu_*^2\varepsilon_0^2 \leq \frac{1}{12}$ by (E.δ), so $\eta\bar{\delta} \leq \eta/12 < 1$ and $C_B(-\eta) > 0$. Hence both points lie in the domain of Φ_M.

Step 1 (inductive hypothesis). For $0 \le k \le K$ and both points, (IH_k) is the statement

$$\begin{aligned} \mathrm{e}^{-E_1(u)}\mathcal{Z}_u \le \mathrm{e}^{-E_{\mathrm{fin}}(u)-E_{K,k}(u)} \sum_{\mathbf{\Lambda}_k} \int dA_k\, d\varphi_k \int dA_{k+1}\cdots dA_K \prod_{j=k}^{K-1} \mathcal{K}_j^w(A_j, A_{j+1}) \\ \times\, \rho_w^{(k)}(\mathbf{\Lambda}_k;\cdot)\, \exp\big[\mathcal{P}_w^{(k)}(\Omega_1;\cdot) + \mathfrak{E}_k(\mathbf{\Lambda}_k) + \mathfrak{O}_k(\mathbf{\Lambda}_k)\big], \end{aligned} \tag{F.2}$$

with equality at $k = 0$ and an empty coarse product at $k = K$. In (F.2):

- $\mathcal{K}_j^w$ are the (unnormalised) kernels of the photon chain of Lemma F.5 with weights w_j. The fields of the coarse chain that are not yet integrated are kept.
- $\rho_w^{(k)}$ is Bałaban's local Gaussian part and outer density [2, (2.49)–(2.52), (2.55)], with the coefficient w_k^{-1} on the vector part and the small-field restrictions of [2, (2.55), p. 570].
- $\mathcal{P}_w^{(k)}$ is the polynomial part of order $\le \bar{n}$ [3, (1.4)–(1.5)] with the weighted lines $G_k(u) = G_k + uc_kD_k$. It contains no part of $E_{\mathrm{fin}}(u)$ or of $E_{K,k}(u)$. Its field-independent part is the vacuum remainder $\mathcal{R}_k^{\mathrm{vac}}$ of Lemma F.32, with $\mathcal{R}_0^{\mathrm{vac}} = 0$.
- The remainders are

$$\mathfrak{E}_k(\mathbf{\Lambda}_k) := \sum_{j<k} C_{\kappa_0} t_j^{\kappa_0} |\mathrm{Col}_j|_j + \sum_{j<k} C_\kappa t_j^{\kappa_B} N_j + \sum_{j=1}^{k} \nu(t_j) |\Lambda_7^{(j-1)\prime}|_j, \qquad \mathfrak{O}_k(\mathbf{\Lambda}_k) := \sum_{j<k} \bar{S} A_j |\Lambda_7^{(j)c}|_j,$$
$$\mathfrak{E}_0 := \mathfrak{O}_0 := 0.$$

  Here $\bar{S}$ and A_j are as in Proposition E.62, and $\bar{S}A_j \le C_{\mathrm{old}} t_j$ with the constant C_{old} of Lemma F.32. The function ν is the cost of the spectator cross term in the photon small-field step (Lemma F.14).

The constants $C_{\kappa_0}, C_\kappa, C_{\mathrm{old}}$ and $\nu(t) = O(t^\kappa)$ (for every κ) depend only on the parameters, on the window $[1-\eta, 2]$ of the weights, and on the regulator data. They depend on ε_0 only through quantities evaluated at the cap ε_1, because Bałaban's per-step constants are bounded uniformly in $\varepsilon_0 \in (0, \varepsilon_1]$ (Statement K.1). They do not depend on ε, k, ℓ_T or $\mathbf{\Lambda}_k$.

The constant C_κ contains the factor $\sup_{t\le\varepsilon_1} t^{1/20}(1+\log t^{-1})^q < \infty$, which converts volume terms of size $t^{13/4}$ times a power of $\log t^{-1}$ into $t^{16/5}$. Likewise C_{κ_0} contains $\sup_{t\le\varepsilon_1} t^{1/20}(1+\log t^{-1})^q < \infty$ ($q \le 24$). This factor converts the collar terms of size $t^{1/4}$ times a power of $\log t^{-1}$ of [2, (2.22), (2.57), (2.78)] and [3, p. 421] into $t^{1/5}$ (Lemma F.1); at $\kappa_0 = \frac{1}{4}$ it would be infinite. C_{κ_0} also contains the tail constants of [2, pp. 579, 582] (Propositions E.76 and E.36), taken as suprema over $t \in (0, 1]$.

Step 2 (base case). By Lemma F.7, (IH_0) holds with equality: the front factor is $\mathrm{e}^{-E_{\mathrm{fin}}(u)-E_{K,0}(u)}$ with $E_{K,0}(u) = E_K(u)$, $\mathcal{P}_w^{(0)} = -V$ and $\mathfrak{E}_0 = \mathfrak{O}_0 = 0$. Here $V(\varphi) = \sum_x \varepsilon^3(\lambda|\varphi(x)|^4 + \frac{1}{2}\delta m_{K,T}^2(M)|\varphi(x)|^2)$. The mass counterterm is the torus-built $\delta m_{K,T}^2(M)$, which is Bałaban's δm^2 [3, (1.19)–(1.23), pp. 416–417] by Proposition 4.7. The integrand is integrable because the whole chain is ($HA > 0$, Lemma F.5), although at $u = 1$ single kernels grow.

Step 3 (inductive step). For $0 \le k \le K-1$ and both points, (IH_k) implies (IH_{k+1}) with

$$\mathfrak{E}_{k+1} = \mathfrak{E}_k + C_{\kappa_0} t_k^{\kappa_0} |\mathrm{Col}_k|_k + C_\kappa t_k^{\kappa_B} N_k + \nu(t_{k+1}) |\Lambda_7^{(k)\prime}|_{k+1}, \qquad \mathfrak{O}_{k+1} = \mathfrak{O}_k + \bar{S} A_k |\Lambda_7^{(k)c}|_k.$$

The operations follow Bałaban's order [2, pp. 569–582]: regions; collar removal; the vector translation and the fluctuation law; the scalar translation and $Z^{(j)}$; the joint fluctuation integral; the new effective action and its vacuum part; and finally the closing of the photon spectators.

(a) *Large fields and collars* (S3). The new regions are a function of the past and of a minimal family. The region expansion is needed only in the direction "$\le$", since the integrand is ≥ 0. On the collar $\mathrm{Col}_k = \Lambda_7^{(k-1)\prime} \cap \Lambda_7^{(k)c}$ the whole interaction, including the step-k mass

pieces, is removed under the step-$(k-1)$ restrictions [2, Prop. 2.1, (2.57), p. 570]. The cost is at most $C_{\kappa_0} t_k^{\kappa_0} |\mathrm{Col}_k|_k$ (Lemma F.1). No vacuum c-number is removed on a collar.

(b) *Scalar and photon small-field steps* (S1), (S2). Every localised family of $\mathcal{P}_w$ obeys the tree bound [3, (1.33)] in weighted form.

- At each k this bound is given by Proposition E.76 (order cap $\bar{N} = 169$). It is applied under the conditions and operations of [3, Prop. 1] of Proposition E.110. These include the support hypothesis in the form (R4‴) of Corollary 5.10 and the repairs of Lemmas 5.6 and 5.22.
- The joint fluctuation integral uses no bound at a complex background near a Neumann face (Proposition E.71).
- The torus defect of the zeroth moment at orders ≥ 2 is the image term of the Ward–Takahashi step (Proposition E.67).
- The uniformity in ℓ_T of the constant C_b of the $\delta m^2_{\mathrm{fin}}$ vertex is Proposition E.162.
- The weights enter the vector part only through a factor w_k on Bałaban's Dirichlet covariance and through the spectator cross term, whose cost is $\nu(t_{k+1}) |\Lambda_7^{(k)\prime}|_{k+1}$ (Lemma F.14).

(c) *Renormalisation flow* (S4). The mass counterterm is u-independent and is the same number at both values of u for the same M (Lemma F.32). Write $\mathrm{e}^{-E_{K,k}(u)} = \mathrm{e}^{-E_{K,k+1}(u)} \mathrm{e}^{-\mathcal{V}_k(u)}$ with $\mathcal{V}_k(u) := E_{K,k}(u) - E_{K,k+1}(u)$. Only $\mathcal{V}_k(u)$ is localised at step k. It is localised at scale k by the rule of [3, p. 420], applied to whole renormalised classes with one external field each; the anchor of a class is its first leg, the same in every member. Put $W_k := B^k(\Lambda_7^{(k)})$, $\mathcal{L}_k := B^k(\Lambda_7^{(k)} \cap \Lambda_8^{(k)c})$ and $\mathcal{D}_k := B^k(\Lambda_8^{(k)})$. A localisation Λ is *good* if $\Lambda \subset W_k$, *bad* if it meets both W_k and W_k^c, and *outside* if $\Lambda \cap W_k = \emptyset$.

- *Good localisations.* By (a), every vertex of a family generated at step k lies in W_k. By Lemma F.31, the field-independent output of step k, truncated at order $\bar{n}$ and computed with the free index-k lines, is $\mathcal{V}_k(u)$, as the same sum of index-assigned Wick graphs. A generated graph and the corresponding counterterm graph have the same vertex set, hence the same localisation. So the good localisations of $\mathcal{V}_k(u)$ cancel the generated output localisation by localisation.
- *What remains in* $\mathcal{R}^{\mathrm{vac}}_{k+1} - \mathcal{R}^{\mathrm{vac}}_k$ is the bad localisations and the domain differences (Lemma F.32).
- *Bad localisations.* They are bounded by $\bar{N} \bar{S} A_k [|\mathcal{L}_k| + \mathrm{e}^{3\delta} \mathrm{e}^{-\delta r(t_k)/2} |\mathcal{D}_k|]$ (Proposition E.62). For the order-e^2 vacuum family with one D-line the factor A_k may be replaced by $C_{\mathcal{V}} e^2 |u| \kappa \mu_*^2 L^3 t_k^3$ (Proposition E.64). For the families with C-lines only and δm^2 insertions, Proposition E.175 applies, via Lemma E.170.
- *The layer.* Let β be a block of $\Lambda_7^{(k)} \cap \Lambda_8^{(k)c}$. By [2, (2.8)] it lies within $r(t_k)$ of a point of $\Lambda_7^{(k)c}$; let b be the unit block that contains this point. If $b \subset \Lambda_0^{(k)c}$, then $\beta \subset \Lambda_1^{(k)c} \subset \Lambda_7^{(k)c}$ by [2, (2.8)], which is impossible. Hence $b \subset \Lambda_0^{(k)} \cap \Lambda_7^{(k)c} \subset \mathrm{Col}_k$. Counting gives $|\mathcal{L}_k|_k \leq (2r(t_k) + 2\sqrt{3} M_b + 1)^3 |\mathrm{Col}_k|_k$. Since $A_k \leq C t_k$, $r(t) = R(1 + \log t^{-1})^2$ and $\kappa_0 < 1$, the layer costs are at most $C(\kappa_0) t_k^{\kappa_0} |\mathrm{Col}_k|_k$. They are part of the collar term of the increment of $\mathfrak{E}$.
- *Deep part and domain differences.* The deep part is at most $C t_k \mathrm{e}^{-\delta r(t_k)/2} N_k$, and the domain differences are at most $C_\kappa t_k^{\kappa_B} N_k$ with $\kappa_B = \frac{16}{5}$ (Lemma F.32). Both are counted in the term $C_\kappa t_k^{\kappa_B} N_k$.
- *Outside localisations.* They are at most $\bar{S} A_k |\Lambda_7^{(k)c}|_k \leq C_{\mathrm{old}} t_k |\Lambda_7^{(k)c}|_k$ (Proposition E.62). This c-number is not part of $\mathcal{P}_w$; it is the increment of $\mathfrak{O}$.

Each term of $E_K(u) = E_{K,k}(u) + \sum_{j<k} \mathcal{V}_j(u)$ leaves the front factor exactly once, at its top index, and $E_{\mathrm{fin}}(u)$ is never moved. The global sum of the order-e^2 family with one D-line,

which is of size $O(K)|T|$, is never bounded. It occurs only inside $E_1(u)$, on both sides of the identity of Step 0.

The constants are the same at both points. The data of the stages up to 9 of Appendix A do not depend on the point, and the conditions that do are imposed at the point where they are strongest. A single ε_0 (the minimum over the two points) therefore gives the same K at both points.

Step 4 (closing: volume terms). Apply (IH_K); its front factor is $\mathrm{e}^{-E_{\mathrm{fin}}(u)}$, since $E_{K,K}(u) = 0$. Three bounds follow.

- (C1) On the final small-field set, $\mathcal{P}_w^{(K)}(\Lambda_7^{(K-1)\prime}) \le C_{F1}\varepsilon_0^{\kappa_0}|\Lambda_7^{(K-1)\prime}|_K \le C_{F1}\varepsilon_0^{\kappa_0-3}|T|$, using $|\cdot|_K \le N_K = |T|\varepsilon_0^{-3}$. This is [2, (2.116)] in weighted form. It is read with $|\Lambda_7^{(K-1)}|$ in place of the printed $|\Lambda_7^{(K)}|$, since no region of scale K exists. The bound rests on Proposition E.76, and uses Propositions E.67, E.71 and, for the supremum over ℓ_T, E.162.
- (C2) We have $|\Lambda_7^{(j-1)\prime}|_j \le N_j$, $\nu(t) \le C_\nu t^{\kappa_B}$, $t_j^{\kappa_B} N_j = |T| t_j^{\kappa_B - 3}$ and $t_j = 2^{-(K-j)}\varepsilon_0$. Hence

$$\sum_{j<K} C_\kappa t_j^{\kappa_B} N_j + \sum_{j\le K} \nu(t_j)|\Lambda_7^{(j-1)\prime}|_j \le (C_\kappa + C_\nu)|T| \sum_{m\ge 0} (2^{-m}\varepsilon_0)^{1/5} =: B_{\mathfrak{E}}|T|,$$

$$B_{\mathfrak{E}} = (1 - 2^{-1/5})^{-1}(C_\kappa + C_\nu)\varepsilon_0^{1/5} \le 7.73\,(C_\kappa + C_\nu)\varepsilon_0^{1/5}.$$

  Only $\kappa_B > 3$ is used, and this keeps the sum uniform in ε. The collar terms and $\mathfrak{O}_K$ stay in the summand and are paid in Step 5.
- (C3) The backward replacements of [2, pp. 583–587] carry the weight w_{k-1}^{-1}. Their determinants are at most $\exp(d\varrho_k^{\mathrm{det}}|\Lambda_5^{(k)}|)$, with $\varrho_k^{\mathrm{det}} \le \frac{1}{2}$ (condition (E.4)) and $\varrho_k^{\mathrm{det}} \le C_4' t_k^4$ (Lemma F.39). Hence $\sum_k d\varrho_k^{\mathrm{det}} N_k \le B_{GS}|T|$ with $B_{GS} := 2dC_4'\varepsilon_0$.

Step 5 (closing: Gaussian integrals, region sum, E_{fin}).

- (C4) *Scalar Gaussian integral.* The scalar field is integrated first, at fixed background [2, Prop. 3.1, (3.30)–(3.38)]. The diamagnetic bound is used on the plain operator only, at every holonomy, and gives at most $\exp(E_{0,s}(M))$. The Gaussian mass is M^2. The reading of "$m^2 \le O(1)$" [2, p. 589] is Proposition E.159, with (E.m2) and (E.Φ).
- (C4, continued) *Vector Gaussian integral and the normalisation defect.* On the vector side the chain rule gives, for every pattern $\mathbf{\Lambda}$, exactly $(2\pi)^{n/2}\det HA_{\mathrm{del}}(\mathbf{\Lambda})^{-1/2}$. The factor $\mathrm{e}^{+E_0^w}$ produced in this way cancels the $\mathrm{e}^{-E_0^w}$ of Step 0 identically for every $\mathbf{\Lambda}$ (Lemma F.35). By Lemma F.38, which needs $\delta_0 R \ge 4$, $|\log\det HA_{\mathrm{del}}(\mathbf{\Lambda}) - \log\det HA| \le B_{F3}|T|$ with

$$B_{F3} := 12|u|\kappa C_{\ell^1}(1 - \hat{\xi}_{\mathrm{tot}})^{-1} w_{\min}^{-2} w_{\mathrm{lo}}^{-1} \gamma_1^{-1} \varepsilon_0.$$

  Here $\hat{\xi}_{\mathrm{tot}} \le \frac{1}{12}$ is the deletion budget ((E.14)), $w_{\min} = \min_j w_j \ge \frac{11}{12}$, w_{lo} is as in Theorem F.6(c), γ_1 is the photon gap constant of Lemma F.37, and C_{ℓ^1} is the constant of Lemma F.38. None of them depends on ε, K or ℓ_T, and $B_{F3} \propto |u|$.
- (C4, continued) *Zero modes.* The zero modes are massive, with gaps $\ge \gamma_1\mu_*^2$ (photon) and $\ge M^2(1 + 4M^2\varepsilon_0^2/(3a))^{-1}$ (scalar), both free of ℓ_T.
- (C5) *Region sum.* After the cancellation, consider the sum over $\mathbf{\Lambda}_K$ of the large-field factors, multiplied by exp of the collar terms, the shell terms, the count of kept coarse variables and $\mathfrak{O}_K(\mathbf{\Lambda}_K)$. This sum is at most $\exp(B_{\mathrm{comb}}|T|)$. It is the region entropy of [2, p. 593] (Proposition 5.21), obtained via Lemmas E.94 and E.95 under $2p = 12 > \varrho d + 2$ and (E.comb). The constant is evaluated at $c_A + C_{\mathrm{old}}$ at ε_1; the term $\mathfrak{O}_K$ is absorbed by Lemma E.63. The constants of [2, (3.42)] that enter (E.ent) are bounded on $(0, \varepsilon_1]$ by Statement K.1. The constant is

$$B_{\mathrm{comb}} = c_{\mathrm{fin}}\varepsilon_0^{-3} + 2(d+2)\Sigma_*, \qquad \Sigma_* = \sum_{n\ge n_0} L^{3n} \exp\big(-\tfrac{1}{2}c_0'^w b_0^2 (1 + n\log L)^{2p}\big).$$

- (E_{fin}) The factor $\mathrm{e}^{-E_{\mathrm{fin}}(u)}$ in front of (IH_K) is paid once: $|E_{\mathrm{fin}}(u)| \le B_E^{\mathrm{fin}}(u,M)|T|$. The bound is uniform in ε and in ℓ_T by Propositions E.21 and E.162. This is the only place where B_E^{fin} enters C_+^B.

Collecting Steps 0–5: (IH_K), (C1) and (C2) reduce the claim to [2, (2.117)] in weighted form. For each $\mathbf{\Lambda}$ we apply (C4) and pay $\mathrm{e}^{(B_{GS}+\frac12 B_{F3})|T|}$. We then sum over $\mathbf{\Lambda}$ by (C5), and finally pay $\mathrm{e}^{-E_{\mathrm{fin}}(u)}$. This gives $\mathrm{e}^{-E_0^w-E_1(u)}\mathcal{Z}_u \le \mathrm{e}^{C_+^B(u;M)|T|}$ with (F.1), which is (5.2) by Step 0.

Step 6 (uniformity and order of choices). The regulator data are chosen in the order of the stages 0–11 of Appendix A.

- Every lower bound on R depends only on data of the stages 2–4.
- The upper bounds on ε_0 of stage 10b, among them (E.fin), (E.edge), (E.δ), (E.14) and (E.comb), are downward closed.
- The lower bounds on ℓ_0 of stage 11, among them (T.cap), (T.div) and (T.per), are fixed after their data.

So the order is acyclic and the conditions can be met (Lemmas A.4, F.40 and A.5). The constants of the upper bounds on ε_0 that depend on ℓ_T are evaluated at the floor $\ell_0^\natural = \max\{8, 128/m_1, 128/M_1\}$ (Lemma A.10). The majorants needed for $\delta m_{\mathrm{fin}}^2$ are those of Proposition E.162. The torus defect of the Ward identity of [3, p. 442] with λ linear in y' is exact on $\eta\mathbb{Z}^d$. On the torus it holds up to the image term of Lemma 5.22; this is a δG-type external field and costs at most one more lower bound on ℓ_0 at stage 11.

Every constant in Steps 1–5 is therefore free of ε, K and ℓ_T. At $u=-\eta$ only $(1-\eta)^{-1}$, $(1-\eta)^{-3}$, $w_{\min}\ge\frac{11}{12}$ and $\vartheta_-\ge\frac{11}{24}$ enter. At $M=m_1$ only $C_\delta(m_1)$, m_1^{-2}, $\log_+(1/(m_1\varepsilon_0))$, the scalar gap and the floor $128/m_1$ enter, and all of these are finite. Since the bound holds for every $K\ge 0$, it holds in particular for $K\ge K_0$. □

F.2. **The step lemmas (S0)–(S5).**

(S0) The base case.

The photon chain. *Setting.* In this paragraph j and k are scale indices. Fix a torus T, $L=2$, $\varepsilon = 2^{-K}\varepsilon_0$ and $t_k = L^k\varepsilon$.

Spaces and block averages. For $0\le k\le K$ let $\mathbb{V}_k$ be the space of real functions on the blocks of side t_k; every operator below acts componentwise on the vector field. Put $\langle f,\, g\rangle_k := t_k^3\sum fg$ and $\|f\|_k^2 = \langle f,\, f\rangle_k$, and let dA_k be the Lebesgue measure of $(\mathbb{V}_k, \langle\cdot,\,\cdot\rangle_k)$. Let $Q_k : \mathbb{V}_0 \to \mathbb{V}_k$ be the block average and $Q := Q_{k\to k+1} : \mathbb{V}_k \to \mathbb{V}_{k+1}$ the one-step average. Adjoints are taken with respect to these products, so that $QQ^* = 1$, $Q_kQ_k^* = 1$, $Q_{k+1} = QQ_k$, and we put $P_k := Q_k^*Q_k$.

Operators. Let $\mathbb{M} := -\Delta_\varepsilon + \mu_*^2$ and $C_1 := \mathbb{M}^{-1} = C_{\mu_*}$. Let $a>0$ be the averaging constant of the block-spin transformation [24, (2.13)–(2.18), p. 653]. Put $a_k = a(1-L^{-2})(1-L^{-2k})^{-1}$, $r_k := a_kt_k^{-2}$ for $k\ge 1$ (not to be confused with the function $r(t)$ of (4.1)), $r_0^{-1} := 0$, and $\beta_k := at_{k+1}^{-2}$. Then

$$r_{k+1}^{-1} = r_k^{-1} + \beta_k^{-1} \qquad (0\le k<K), \tag{F.3}$$

since $1/a_{k+1} = L^{-2}/a_k + 1/a$ and $a_1 = a$. Put

- $G_k := (\mathbb{M} + r_kP_k)^{-1}$ $(k\ge 1)$, $G_0 := 0$, and $H_k := G_{k+1} - G_k$;
- $F_k := r_kG_kQ_k^*$ $(k\ge 1)$ and $F_0 := 1$;
- $\Delta^{(0)} := \mathbb{M}$ and $\Delta^{(k)} := r_k - r_k^2Q_kG_kQ_k^*$ $(k\ge 1)$ [24, (2.13)–(2.18), p. 653];
- $C^{(k)} := (\Delta^{(k)} + \beta_kQ^*Q)^{-1}$, $\mathcal{H}_k := \beta_kC^{(k)}Q^*$ and $D_k(B) := \langle B,\, \Delta^{(k)}B\rangle_k$.

Weights. For weights $w_0,\dots,w_{K-1}>0$ put $w_K := 1$ and $\omega_k := w_k^{-1} - w_{k-1}^{-1}$ $(1\le k\le K)$. At the two points, $w_j = 1+u\delta_j$ with $\delta_j = \kappa\mu_*^2t_{j+1}^2$ as in (5.1). A centred Gaussian $N(0,\Sigma)$ on $(\mathbb{V}_k, \langle\cdot,\,\cdot\rangle_k)$ has a density with respect to dA_k proportional to $\mathrm{e}^{-\frac12\langle x,\,\Sigma^{-1}x\rangle_k}$.

Lemma F.3 (Gaussian conditioning)**.** *Let $X \sim N(0,\Sigma)$ with $\Sigma > 0$ on a finite-dimensional inner product space V, let $S : V \to W$ be linear, and let $Y = SX + \xi$ with $\xi \sim N(0,\Lambda)$ independent of X and $\Lambda > 0$. Then:*

- *$Y \sim N(0, S\Sigma S^* + \Lambda)$;*
- *the conditional law of X given Y is $N(\Pi Y, \Sigma_{X|Y})$, with $\Sigma_{X|Y} = (\Sigma^{-1} + S^*\Lambda^{-1}S)^{-1}$ and $\Pi = \Sigma_{X|Y}S^*\Lambda^{-1}$;*
- *$\Sigma = \Sigma_{X|Y} + \Pi(S\Sigma S^* + \Lambda)\Pi^*$;*
- *$(\Lambda + S\Sigma S^*)^{-1} = \Lambda^{-1} - \Lambda^{-1}S(\Sigma^{-1} + S^*\Lambda^{-1}S)^{-1}S^*\Lambda^{-1}$.*

Proof. The joint density of (X,Y) is proportional to $\exp(-\frac{1}{2}\langle x,\, \Sigma^{-1}x\rangle - \frac{1}{2}\langle y - Sx,\, \Lambda^{-1}(y-Sx)\rangle)$. Completing the square in x gives the conditional law. The covariance identity is the law of total covariance. The last identity (Woodbury) follows by multiplying the right side by $\Lambda + S\Sigma S^*$. □

Lemma F.4 (block actions and lifts)**.** *For $0 \le k < K$:*

(i) $(\Delta^{(k)})^{-1} = Q_kC_1Q_k^* + r_k^{-1}$; *in particular* $\Delta^{(k)} > 0$.
(ii) $\Delta^{(k+1)} = \beta_k - \beta_k^2 Q C^{(k)} Q^*$ [1, (3.41), p. 619].
(iii) $F_k\mathcal{H}_k = F_{k+1}$ [1, (3.42), p. 619].
(iv) $C_1 - G_k = F_k(\Delta^{(k)})^{-1}F_k^*$ *and* $H_k = F_kC^{(k)}F_k^*$ [24, (2.17), p. 653].
(v) *For $A \in \mathbb{V}_k$ and $B \in \mathbb{V}_{k+1}$,*

$$D_k(A) + \beta_k\|QA - B\|_{k+1}^2 = \langle A - \mathcal{H}_kB,\, (C^{(k)})^{-1}(A - \mathcal{H}_kB)\rangle_k + D_{k+1}(B).$$

Proof. (i) For $k = 0$ this is $\mathbb{M}^{-1} = C_1$. For $k \ge 1$ apply Woodbury's identity of Lemma F.3 with $\Lambda = r_k^{-1}$, $S = Q_k$ and $\Sigma = C_1$: $(r_k^{-1} + Q_kC_1Q_k^*)^{-1} = r_k - r_k^2Q_k(\mathbb{M} + r_kP_k)^{-1}Q_k^* = \Delta^{(k)}$.

(ii) By (i) at $k+1$, $Q_{k+1} = QQ_k$, $QQ^* = 1$ and (F.3),

$$(\Delta^{(k+1)})^{-1} = Q\big(Q_kC_1Q_k^* + r_k^{-1}\big)Q^* + \beta_k^{-1} = Q(\Delta^{(k)})^{-1}Q^* + \beta_k^{-1}.$$

Woodbury's identity with $\Lambda = \beta_k^{-1}$, $S = Q$ and $\Sigma = (\Delta^{(k)})^{-1}$ gives $\beta_k - \beta_k^2Q(\Delta^{(k)} + \beta_kQ^*Q)^{-1}Q^*$.

(iii), (iv) Let $A \sim N(0, C_1)$, $B = Q_kA + r_k^{-1/2}\xi$ and $B' = QB + \beta_k^{-1/2}\xi'$, with ξ, ξ' standard and independent (at $k = 0$, $B = A$). By Lemma F.3 and (i), $B \sim N(0, Q_kC_1Q_k^* + r_k^{-1}) = N(0, (\Delta^{(k)})^{-1})$.

- The law of A given B is $N(F_kB, G_k)$, since the posterior precision is $\mathbb{M} + r_kP_k$.
- The law of B given B' is $N(\mathcal{H}_kB', C^{(k)})$, since the posterior precision is $\Delta^{(k)} + \beta_kQ^*Q$.
- Given B, the fields A and B' are independent, so $\mathbb{E}[A \,|\, B'] = \mathbb{E}[F_kB \,|\, B'] = F_k\mathcal{H}_kB'$.
- On the other hand $B' = Q_{k+1}A + \zeta$, where $\zeta = r_k^{-1/2}Q\xi + \beta_k^{-1/2}\xi'$ is independent of A and has covariance $r_k^{-1} + \beta_k^{-1} = r_{k+1}^{-1}$. So $\mathbb{E}[A \,|\, B'] = F_{k+1}B'$.

Since B' has a non-degenerate law, $F_k\mathcal{H}_k = F_{k+1}$. The law of total covariance for (A,B) gives $C_1 = G_k + F_k(\Delta^{(k)})^{-1}F_k^*$. For (B,B') it gives $(\Delta^{(k)})^{-1} = C^{(k)} + \mathcal{H}_k(\Delta^{(k+1)})^{-1}\mathcal{H}_k^*$. Hence, by (iii),

$$H_k = (C_1 - G_k) - (C_1 - G_{k+1}) = F_k\big[(\Delta^{(k)})^{-1} - \mathcal{H}_k(\Delta^{(k+1)})^{-1}\mathcal{H}_k^*\big]F_k^* = F_kC^{(k)}F_k^*.$$

(v) Expand the right side. Use $(C^{(k)})^{-1} = \Delta^{(k)} + \beta_kQ^*Q$ and $(C^{(k)})^{-1}\mathcal{H}_k = \beta_kQ^*$. Also $\langle \mathcal{H}_kB,\, (C^{(k)})^{-1}\mathcal{H}_kB\rangle_k = \beta_k^2\langle B,\, QC^{(k)}Q^*B\rangle_{k+1} = \langle B,\, (\beta_k - \Delta^{(k+1)})B\rangle_{k+1}$, by (ii). □

Lemma F.5 (the quadratic form of the photon chain)**.** *Let $w_0, \dots, w_{K-1} > 0$, and let HA be the quadratic form on $\mathbb{V}_0 \times \cdots \times \mathbb{V}_K$ for which $N_w\mathrm{e}^{-\frac{1}{2}HA}$ is the joint density of the photon chain $(\hat{B}_0, \dots, \hat{B}_K)$ of Theorem F.6. Let $I_k(A_k, A_{k+1}) := \langle A_k - \mathcal{H}_kA_{k+1},\, (C^{(k)})^{-1}(A_k - \mathcal{H}_kA_{k+1})\rangle_k$ and $D_K(B) := \langle B,\, \Delta^{(K)}B\rangle_K$. Then*

$$HA = \sum_{k<K} w_k^{-1}I_k(A_k, A_{k+1}) + D_K(A_K).$$

Equivalently, $\mathrm{e}^{-\frac12 HA}=\mathrm{e}^{-\frac12 w_0^{-1}D_0(A_0)}\prod_{j=0}^{K-1}\mathcal{K}_j^w(A_j,A_{j+1})$ *with*

$$\mathcal{K}_j^w(A_j,A_{j+1})=\exp\big(-\tfrac12 w_j^{-1}\beta_j\|QA_j-A_{j+1}\|_{j+1}^2-\tfrac12\omega_{j+1}D_{j+1}(A_{j+1})\big).$$

Each spectator $\omega_{j+1}D_{j+1}$ *is used once.* $HA>0$*, in particular at both points, where* $w_j=1+u\delta_j\ge 1-\eta/12>0$ *by (E.*δ*).*

Theorem F.6 (hierarchical form of the slice photon)**.** *Let* $w_0,\dots,w_{K-1}>0$*. Let* $\hat S_k\sim N(0,w_kC^{(k)})$ *on* $\mathbb{V}_k$ *(*$k<K$*) and* $\hat B_K\sim N(0,(\Delta^{(K)})^{-1})$ *be independent, and put* $\hat B_k:=\hat S_k+\mathcal{H}_k\hat B_{k+1}$ *for* $k=K-1,\dots,0$ *(the photon chain).*

(a) $A:=\hat B_0$ *has covariance* $\sum_{j<K}w_jH_j+(C_1-G_K)$*. At* $w_j=1+u\delta_j$ *this is* $C_B(u)=C_{\mu_*}+u\Gamma^B$*.*

(b) *The joint law of* $(\hat B_0,\dots,\hat B_K)$ *has density* $N_w\mathrm{e}^{-\frac12 HA}$ *with respect to* $dA_0\cdots dA_K$*, where*

$$HA=\sum_{k<K}w_k^{-1}I_k(\hat B_k,\hat B_{k+1})+D_K(\hat B_K)=w_0^{-1}D_0(\hat B_0)+\sum_{k<K}w_k^{-1}\beta_k\|Q\hat B_k-\hat B_{k+1}\|_{k+1}^2+\sum_{k=1}^{K}\omega_kD_k(\hat B_k).$$

Here $I_k\ge 0$ *is as in Lemma F.5,* $n_k=\dim\mathbb{V}_k$*, and* $N_w^{-1}=\prod_{k<K}(2\pi)^{n_k/2}\det(w_kC^{(k)})^{1/2}\cdot(2\pi)^{n_K/2}\det(\Delta^{(K)})^{-1/2}$*.*

(c) *Put* $w_{\mathrm{lo}}:=\min(1,\min_k w_k^{-1})$*,* $w_{\mathrm{hi}}:=\max(1,\max_k w_k^{-1})$ *and* $S^1:=HA|_{w\equiv1}$*. Then* $w_{\mathrm{lo}}S^1\le HA\le w_{\mathrm{hi}}S^1$ *as quadratic forms. The same inequalities hold after minimisation over any set of variables and on every diagonal block.*

Proof. (a) Unfolding the recursion gives $\hat B_0=\sum_{j<K}\mathcal{H}_0\cdots\mathcal{H}_{j-1}\hat S_j+\mathcal{H}_0\cdots\mathcal{H}_{K-1}\hat B_K$. By Lemma F.4(iii), iterated from $F_0=1$, $\mathcal{H}_0\cdots\mathcal{H}_{j-1}=F_j$. By independence and Lemma F.4(iv),

$$\mathrm{Cov}(A)=\sum_j w_jF_jC^{(j)}F_j^*+F_K(\Delta^{(K)})^{-1}F_K^*=\sum_j w_jH_j+C_1-G_K.$$

Since $\sum_{j<K}H_j=G_K$, at $w_j=1+u\delta_j$ this equals $C_1+u\sum_j\delta_jH_j=C_B(u)$ by (5.1).

(b) $\hat B_{k+1}$ is a function of $(\hat S_j)_{j>k}$ and $\hat B_K$, so it is independent of $\hat S_k$. Hence the law of $\hat B_k$ given $(\hat B_{k+1},\dots,\hat B_K)$ is $N(\mathcal{H}_k\hat B_{k+1},w_kC^{(k)})$, whose normalisation does not depend on the condition. The joint density is the product of the density of $\hat B_K$ and these conditional densities; this gives the first expression for HA and the value of N_w. For the second expression substitute Lemma F.4(v): $I_k=D_k(\hat B_k)+\beta_k\|Q\hat B_k-\hat B_{k+1}\|_{k+1}^2-D_{k+1}(\hat B_{k+1})$. The coefficient of D_0 is then w_0^{-1}, that of D_k $(1\le k<K)$ is $w_k^{-1}-w_{k-1}^{-1}=\omega_k$, and that of D_K is $1-w_{K-1}^{-1}=\omega_K$.

(c) $HA-w_{\mathrm{lo}}S^1=\sum_k(w_k^{-1}-w_{\mathrm{lo}})I_k+(1-w_{\mathrm{lo}})D_K\ge0$, since $I_k\ge0$ and $\Delta^{(K)}>0$; the bound with w_{hi} is proved in the same way. Pointwise inequalities between quadratic forms are preserved by minimisation over a set of variables and by restriction to a coordinate subspace. □

Proof of Lemma F.5. The first formula for HA is Theorem F.6(b). Grouping the terms of the second expression of Theorem F.6(b) by $j=k$ for the β-terms and $j=k-1$ for the ω_k-terms gives the product formula with the kernels $\mathcal{K}_j^w$; each $\omega_{j+1}D_{j+1}$ occurs in exactly one kernel. HA is the precision of a non-degenerate Gaussian, so $HA>0$. At the two points, $0\le\delta_j\le\kappa\mu_*^2\varepsilon_0^2\le\frac{1}{12}$ by (E.δ), since $t_{j+1}\le t_K=\varepsilon_0$. So $w_j\ge1-\eta/12>0$, and the weights are admissible. □

Lemma F.7 (the base identity)**.** *At both points* $(u,M)\in\{(1,m_2),(-\eta,m_1)\}$*, with* $\mathbf{\Lambda}_0=\emptyset$*,* $\Lambda_i^{(-1)\prime}=T_\varepsilon$*,* $\Omega=\Omega_1=T_\varepsilon$*,* $A^{(0)}=A_0$*,*

$$\mathrm{e}^{-E_1(u)}\mathcal{Z}_u=\mathrm{e}^{-E_{\mathrm{fin}}(u)}\int dA_0\,d\varphi_0\int dA_1\cdots dA_K\prod_{j=0}^{K-1}\mathcal{K}_j^w(A_j,A_{j+1})\,\mathrm{e}^{-\mathcal{G}_w^{(0)}(A_0,\varphi_0)}\exp[-V(\varphi_0)-E_K(u)].$$

Here

$$\mathcal{G}_w^{(0)} = \tfrac{1}{2}w_0^{-1}\langle A_0,\, \mathbb{M}A_0\rangle_0 + \tfrac{1}{2}\langle\varphi_0,\, (-\Delta_{A_0}^{\varepsilon}+M^2)\varphi_0\rangle, \qquad V(\varphi) = \sum_{x\in T_\varepsilon} \varepsilon^3\big(\lambda|\varphi(x)|^4 + \tfrac{1}{2}\delta m_{K,T}^2(M)|\varphi(x)|^2\big),$$

and $E_K(u) = E_1(u) - E_{\mathrm{fin}}(u) = E_{K,0}(u)$ *(Lemma F.31). Hence* $\rho' \equiv 1$, $\chi_0 \equiv 1$, $\mathfrak{E}_0 = \mathfrak{O}_0 = 0$ *and* $HA_{\mathrm{del}}^{(0)} = HA$. *In the form* (F.2) *at* $k = 0$ *the front factor is* $\mathrm{e}^{-E_{\mathrm{fin}}(u)-E_{K,0}(u)}$, *and*

$$\mathcal{P}_w^{(0)} = -V.$$

Proof. The photon. The photon of the slice model is the normalised Gaussian $\mu_{C_B(u)}$. By Theorem F.6(a),(b), $\mu_{C_B(u)}$ is the A_0-marginal of the law $N_w\mathrm{e}^{-\frac{1}{2}HA}dA_0\cdots dA_K$. Put $Z_\varphi(A_0) := \int d\varphi_0\, \mathrm{e}^{-S_\varphi(A_0,\varphi_0)}$. The integrand is nonnegative, so Tonelli's theorem gives

$$\int d\mu_{C_B(u)}(A)\, Z_\varphi(A) = N_w \int dA_0\cdots dA_K\, \mathrm{e}^{-\frac{1}{2}HA} Z_\varphi(A_0) = N_w\mathcal{Z}_u.$$

By Lemma F.5, $\mathrm{e}^{-\frac{1}{2}HA} = \mathrm{e}^{-\frac{1}{2}w_0^{-1}D_0(A_0)}\prod_{j=0}^{K-1}\mathcal{K}_j^w$, and $D_0(A_0) = \langle A_0,\, \mathbb{M}A_0\rangle_0$ since $\Delta^{(0)} = \mathbb{M}$.

The scalar. King's torus scalar action equals Bałaban's at $m^2 = M^2$, $\mu_0 = \mu_*$, $\delta m^2 = \delta m_{K,T}^2(M)$ (Lemma C.1; Proposition 4.7). Hence $S_\varphi = \frac{1}{2}\langle\varphi,\, (-\Delta_{A_0}^{\varepsilon} + M^2)\varphi\rangle + V$. Multiplying by $\mathrm{e}^{-E_1(u)} = \mathrm{e}^{-E_{\mathrm{fin}}(u)}\mathrm{e}^{-E_K(u)}$ gives the displayed identity.

The form (F.2) *at* $k = 0$. In convention (G) of Remark 5.3(ii), both c-numbers $E_{\mathrm{fin}}(u)$ and $E_{K,0}(u) = E_K(u)$ stand in front, so the polynomial part is $\mathcal{P}_w^{(0)} = -V$. It has order $\le \bar{n}$ in the couplings, since λ has order 1 and $\delta m_{K,T}^2$ has orders $1 \le \alpha+\beta \le 4 \le \bar{n}$; its field-independent part is $0 = \mathcal{R}_0^{\mathrm{vac}}$. No region has been chosen, so there are no restrictions and no deleted variables.

Integrability. $HA > 0$ (Lemma F.5). At $u = 1$ single kernels grow ($\omega_k < 0$ for $1 \le k < K$); only the whole chain is integrable. □

(S1), (S2) The small-field steps.

Lemma F.8 (scalar small-field step)**.** *Fix the regulator data of Theorem 5.2 (*$L = 2$, $p = 6$, $p_1 = b_1 = 1$, $\bar{n} = 13$, $R \ge R^{**}$, ε_0, ℓ_0*) and put* $\kappa_0 = \frac{1}{5}$. *Let* $(u, M) \in \{(1, m_2), (-\eta, m_1)\}$, $\varepsilon = 2^{-K}\varepsilon_0$, $\ell_T \in \{j!\ell_0\}$ *and* $0 \le k \le K-1$. *We work in scale-k units (*$\eta = L^{-k}$, $t = t_k$*).* $|X|$ *is the number of scale-k blocks in* X, $\mathrm{Col}_k := \Lambda_7^{(k-1)\prime} \cap \Lambda_7^{(k)c}$, *and* N_ℓ *is the largest number of vector lines of a graph of order* $\le \bar{n}$. *Assume:*

(SC1) *(*IH_k*) of* (F.2). *In particular: the scalar Gaussian part of* $\rho_w^{(k)}$ *is*

$$-\tfrac{1}{2}\langle\Lambda_6^{(k-1)\prime}\varphi,\, \Delta^{(k)}(B^k(\Lambda_2^{(k-1)\prime}), A^{(k),\varepsilon})\Lambda_6^{(k-1)\prime}\varphi\rangle$$

[2, (2.49), p. 568]*, with no weight;* $\rho_w^{(k)}$ *depends on the configuration on* $B^k(\Lambda_7^{(k-1)\prime})$ *only through* $\prod_{j<k} Z^{(j)}_{\Lambda_5^{(j)}}(B^j(\Lambda_2^{(j)}), A^\varepsilon)$ *(ibid.); the restrictions* [2, (2.55), p. 570] *hold on* $\Lambda_{-1}^{(k-1)\prime}$, *among them* $|(D_{\tilde{A}^{(k)}}\varphi)(b)| \le c_1 p(t_{k-1})$ *and* $|\varphi(x)| \le c_1\lambda(t_{k-1})^{-1/4}p(t_{k-1})$; $\theta_k A^{(k)}$ *is an independent external field that occurs only in scalar expressions; the scalar normalisations are carried exactly.*

(SC2) *Part (a) of Lemma F.14; the regularity of* $\tilde{B} = (1-\theta_{k+1})\theta_k A^{(k)} + \theta_{k+1}B^{(k+1),\eta}$ *and of* $B^{(k+1),\eta}$ *required in* [1, Prop. 2.1]*, as supplied by* [2, (2.98), p. 577]*; and the kernel bounds* [3, (2.10)–(2.12)] *for the D-line, with dimension* $-d+4$.

(SC3) *Part (a) of Lemma F.22.*

(SC4) *The mass counterterm is* $\delta m_{K,T}^2(M)$ *(Proposition 4.7), with the scale split of* [3, (1.26)–(1.29)].

Then the following hold. The constants depend only on the parameters, η, the regulator data and $\ell_0^\natural$; they do not depend on ε, k, ℓ_T, $\mathbf{\Lambda}_k$, the background, or the sign of u. Every u-dependent constant at $u=-\eta$ is at most its value at $u=1$.

(C1) *Under* [2, (2.55)],

$$\mathcal{P}_w^{(k)}(\Lambda_7^{(k-1)\prime},\theta_k A^{(k)},\varphi)=-\lambda(t)\sum_{x\in B^k(\mathrm{Col}_k)}\eta^d|\varphi^{(k)}(x)|^4+\mathcal{P}_w^{(k)}(\Lambda_7^{(k)},\theta_k A^{(k)},\varphi)+\mathcal{R}_1,$$

with $|\mathcal{R}_1|\le C_1(1+2^{N_\ell}|u|c_k)\,t^{\kappa_0}|\mathrm{Col}_k|$. *Here* $\varphi^{(k)}=a_kG_k(B^k(\Lambda_2^{(k-1)\prime}),A^{(k)})Q_k^*(A^{(k)})\Lambda_6^{(k-1)\prime}\varphi$ *and* $c_k=\kappa\mu_*^2L^2t_k^2\le\frac13$. *At* $k=0$, *moreover,* $\lambda\varepsilon|\varphi|^4+\frac12\delta m_{K,T}^2(M)\varepsilon^2|\varphi|^2\ge -C_\delta(M)^2\varepsilon/(16\lambda)$ *at every site.*

(C2) *Replacing the scalar quadratic forms by their localised versions* [2, (2.47)–(2.48), (2.108)] *costs* $O(t^\kappa)|\Lambda_6^{(k-1)}|$ *and* $O(t^\kappa)|\Lambda_3^{(k)}|$ *respectively, for every* κ. *The difference operator* H_k *satisfies* $|h_k(x,x')|\le O(1)\mathrm{e}^{-\delta_1 r(t)}\mathrm{e}^{-\delta_0|x-x'|}$ [2, (2.109)].

(C3) *We have*

$$\prod_{j<k}Z^{(j)}(A+B)\le\prod_{j<k}Z^{(j)}(\tilde B)\exp\big(\mathcal{Q}^{(k)}(B^k(\Lambda_2^{(k)});B^{(k+1),\eta},g_kA'^{(k)})+C(\kappa_0)t^{\kappa_0}|\mathrm{Col}_k|+O(t^\kappa)|\Lambda_7^{(k)}|\big).$$

The collar constant carries no factor k.

(C4) *In the interaction localised in* $B^k(\Lambda_7^{(k)})$, *replace* $G_k(B^{k-1}(\Lambda_2^{(k-1)}),\tilde B)$ *by* $G_k(B^k(\Lambda_2^{(k)}),B^{(k+1),\eta})$. *All terms containing a difference are* $O(t^\kappa)|\Lambda_7^{(k)}|$. *After the translation* [2, (2.110)], *the identity* [2, (2.111)–(2.112)] *holds up to* $O(t^\kappa)$.

(C5) *For* $x\in\Lambda_5^{(k)}$, [2, (2.113)] *holds, and* $|\varphi'(x)|\le O(1)p(t)$.

(C6) *With the vector half from Lemma F.14(b),*

$$\int d\mu_{w_kC_{\Lambda_5}^{(k)}}(A')\int d\mu_{C_{\Lambda_5}^{(k)}(B^k(\Lambda_2^{(k)}),B^{(k+1),\eta})}(\varphi')\,\chi(A')\chi(\varphi')\,\mathrm{e}^{V^{(k)}}\le\exp\Big(\sum_{m\le\bar n}\frac{\kappa_m(V^{(k)})}{m!}+O(t^\kappa)|\Lambda_5^{(k)}|\Big).$$

(C7) *The cumulant sum, after the replacements of* [2, p. 582], *is* $\mathcal{P}_w^{(k+1)}(\Lambda_7^{(k)},B^{(k+1),\eta},\psi)$, *with vector lines* $G_{k+1}(u)=G_{k+1}+uc_{k+1}D_{k+1}$. *Every localised family obeys* [3, (1.33)], *the* D*-coloured ones with the factor* $(|u|c_{k+1})^m$. *The step-*$(k+1)$ *mass coefficients obey the pointwise bounds of Proposition E.76.*

(C8) *The vacuum part is as stated in Lemma F.32(V), (C11), with the* u*-part from Proposition E.64 and the* C*-only bad localisations from Proposition E.175.*

(C9) [2, Prop. 3.1, (3.26)] *holds at every real* $\tilde A$ *satisfying the vector restrictions, with* $\gamma_0=\gamma_0(d,a)$ *and error* $\sum_k O(t_k^{\kappa_0})|\Lambda_k|$; $\Delta^{(k)}(\Omega,A)\ge a_kM^2t^2/(a_k+M^2t^2)$ *for every real* A *and every* Ω; *the difference forms of* [2, (3.7)–(3.8)] *obey the kernel bound of (C2).*

(C10) *The scalar items of* (IH_{k+1}) *hold, with* $\mathfrak{E}_{k+1}-\mathfrak{E}_k$ *given by the scalar contributions to Step 3 of the proof of Proposition F.2.*

Every scalar condition is free of w; u *enters only through the vector background law and the* D*-lines.*

The constant C_b uniform in ℓ_T is that of Proposition E.162, and the bounds at complex backgrounds are excluded by Proposition E.71.

Lemma F.9. *Let* $\lambda,\varepsilon>0$, $|\delta|\le C_\delta/\varepsilon$ *and* $x\ge0$. *Then* $\lambda\varepsilon x^2+\frac12\delta\varepsilon^2x\ge -C_\delta^2\varepsilon/(16\lambda)$.

Proof. The quadratic in x is minimised at $x=-\delta\varepsilon/(4\lambda)$, with value $-\delta^2\varepsilon^3/(16\lambda)$. If $\delta\ge0$, the minimum over $x\ge0$ is 0. In both cases the bound follows from $\delta^2\varepsilon^2\le C_\delta^2$. □

Lemma F.10. *Let* $\mathcal{P}(\Omega)=\sum_{X\subset\Omega}E(X)$, *the sum running over connected unions of blocks, and assume* $\sum_{X\ni\square}|E(X)|\le a$ *for every block* $\square$. *Then* $|\mathcal{P}(\Omega)-\mathcal{P}(\Omega')|\le a|\Omega\setminus\Omega'|$ *for* $\Omega'\subset\Omega$.

Proof. $\mathcal{P}(\Omega)-\mathcal{P}(\Omega')=\sum_{X\subset\Omega,\,X\not\subset\Omega'}E(X)$, and every such X contains a block of $\Omega\setminus\Omega'$. Summing over that block first gives the bound. □

Lemma F.11. *Let $0<t\le\varepsilon_0\le 1$, $0<\kappa_0<\frac14$, $\ell(t)=1+\log t^{-1}$ and $n_t:=\log_2(\varepsilon_0/t)$. There is $C(\kappa_0)<\infty$ such that each of*

$$t^{1/4}p(t)^3,\quad t^{1/2}p(t)^q\ (q\le 6),\quad t\,(1+t\,n_t),\quad t\,\ell(t)^{2p}(r(t)+M_b)^3,\quad r(t)^3t^3(1+n_t)$$

is at most $C(\kappa_0)\,t^{\kappa_0}$.

Proof. Each quantity is t^{κ_0} times $t^a\ell(t)^n$ with $a>0$, times bounded factors. Here $r(t)=R\ell(t)^2$, $p(t)=b_0\ell(t)^6$, $t\,n_t\le\varepsilon_0/(\mathrm{e}\log 2)$ and $n_t\le\log_2(\mathrm{e})\,\ell(t)$. Now apply $S(a,n)=\sup_{0<t\le1}t^a\ell(t)^n<\infty$ (Lemma F.1). □

Lemma F.12. *Let G be a graph of order $\le\bar n$ with at most N_ℓ vector lines. Assume each colouring of these lines with $m\ge1$ D-lines is bounded by $(|u|c)^m$ times the bound B_0 of the C-only colouring, where $0\le|u|c\le1$. Then the sum over all colourings with $m\ge1$ is at most $2^{N_\ell}|u|c\,B_0$.*

Proof. $\sum_{m\ge1}\binom{N_\ell}{m}(|u|c)^m\le|u|c\sum_{m\ge1}\binom{N_\ell}{m}\le 2^{N_\ell}|u|c.$ □

Proof of Lemma F.8. We follow the order of operations of [2, pp. 569–582].

Step 0 (the weights act only on the photon law). By Lemma F.5, the weights w_j enter only the kernels $\mathcal{K}_j^w$ and, through Lemma F.14(a), the precision $w_k^{-1}(C^{(k)}_{\Lambda_5})^{-1}$ of the photon fluctuation, whose minimiser is w-free. All scalar objects are deterministic functions of the vector configuration and the scalar fields, with no w in them: $G_k(\Omega,B)$, $\Delta^{(k)}$, $Z^{(j)}$, χ_k, $\varphi^{(k)}$, the scalar fluctuation law, and the statements [2, Prop. 2.6, Lemma 2.7]. Hence every scalar bound below is Bałaban's, with his constants. The parameter u enters only through the restriction constants (Statement K.42(iv), whose premise is Statement K.41) and through the D-lines of $\mathcal{P}_w$, in both cases only via $|u|\le1$ and $w_j\in[1-\eta,2]$.

Step 1, (C1). The $u=0$ part. Write $\mathcal{P}^{(k)}$ as a sum over localisations. The vertex [3, (1.6)] with all four scalar legs external is kept explicitly as $-\lambda(t)\sum\eta^d|\varphi^{(k)}|^4$. Every other localised term is bounded by [3, (1.33)] (Proposition E.76), with external fields controlled by [2, (2.55)]. By [3, p. 421], the worst term carries $O(\lambda(t)^{1/4}p(t)^3)$, i.e. $O(t^{1/4-\beta})$ in $d=3$. Summed over localisations containing a fixed block, this is $\le C\,t^{1/4}p(t)^3\le C(\kappa_0)t^{\kappa_0}$ by Lemma F.11. Lemma F.10 with $\Omega=\Lambda_7^{(k-1)\prime}$, $\Omega'=\Lambda_7^{(k)}$ gives the $u=0$ part of $\mathcal{R}_1$. *The step-k mass-counterterm pieces.* At $k\ge1$ these pieces leave Col_k together with the rest of the interaction. With $|\varphi^{(k)}|^2\le O(1)c_1^2\lambda^{-1/2}t^{-1/2}p^2$ and the pointwise bound $|\delta m^2_{K,k}|t^2\le C_m[(e^2+\lambda)t\frac{L}{L-1}+(e^4+e^2\lambda+\lambda^2)t^2(K-k)]$ (Proposition E.76), each vertex is $\le C\,t^{1/2}p^2(1+t\log_2(\varepsilon_0/t))\le C(\kappa_0)t^{\kappa_0}$ per site (Lemma F.11). The finite piece $\delta m^2_{\mathrm{fin}}$ gives $O(C_bt^{3/2}p^2)$ per site, with C_b uniform in ℓ_T (Proposition E.162). *The D-coloured part.* By the coloured form of [3, (1.33)] (Proposition E.76) and Lemma F.12, it is bounded by $2^{N_\ell}|u|c_k$ times the $u=0$ bound, with $c_k\le4\kappa\mu_*^2\varepsilon_0^2\le\frac13$ by condition (E.δ). *The case $k=0$.* Combine Lemma F.9 with $\varepsilon|\delta m^2_{K,T}(M)|\le C_\delta(M)$ (Statement K.43). The absorption of this loss by half of the step-0 gain is not used here. The localisation cubes of side $3r(t)$ do not wrap, by Lemma E.66.

Step 2, (C2). The decomposition [2, (2.47)–(2.48)] and the identity [2, (2.108)] are domain changes at a single background. By Statement K.39 and [2, (2.55)], they cost $O(t^\kappa)$ per block of $\Lambda_6^{(k-1)}$, resp. of $\Lambda_3^{(k)}$, and [2, (2.109)] holds. Since $M^2t^2\le M^2\varepsilon_0^2\le1$ (condition (E.m2); Proposition E.159), the constants are uniform in $M\le m_2$.

Step 3, (C3). By [2, (2.105)], $\prod_{j<k}Z^{(j)}=Z_k$. The fluctuation expansion and the re-localisation are Statement K.38, whose inputs are Statement K.37. The collar terms produced by the extension of [1, Prop. 2.1] to the polynomial $\mathcal{Q}^{(k)}$ [2, p. 579] are bounded by Statement K.35 and Lemma F.11. The final replacement $G_k(B^k(\Lambda_0^{(k)}),\tilde B)\to G_k(B^k(\Lambda_2^{(k)}),B^{(k+1),\eta})$ is a domain change at distance $\ge r(t)$ and costs $O(t^\kappa)|\Lambda_7^{(k)}|$ (Lemma E.123). By the admissibility of $\mathbf{\Lambda}_{k+1}$ [2, p. 566],

$\Lambda_2^{(k)} \cap \Lambda_7^{(k)c} \subset \mathrm{Col}_k$. The $Z^{(j)}$ and $\mathcal{Q}^{(k)}$ are evaluated only through ℓ^2-bounds at real backgrounds (Proposition E.71).

Step 4, (C4). By Statement K.42(ii), the two propagators in [2, Prop. 2.6] are taken at the same field on $B^k(\Lambda_2^{(k)})$, so the replacement is a domain change. Vertices localised in $B^k(\Lambda_7^{(k)})$ lie at distance $> 5r(t)$ from $\Lambda_2^{(k)c}$ [2, (2.8)]. Lemma E.123 bounds each propagator difference by $O(1)\mathrm{e}^{-c\,r(t)} = O(t^\kappa)$ for every κ, and the graph sums are those of [3, (1.33)] (Proposition E.76). The translation [2, (2.110)] is Gaussian algebra at $B^{(k+1),\eta}$; the identity [2, (2.111)–(2.112)] follows from [1, (2.41)] and the same domain-change bound.

Step 5, (C5). This is Statement K.36, together with the restrictions on ψ, φ [2, (2.114)]. Consequently the cut-off $\chi(\varphi')$ of [2, (2.114)] on $\Lambda_5^{(k)}$ may be inserted. All other scalar characteristic functions on $\Lambda_5^{(k)}$ except χ_{k+1} are bounded by 1 [2, p. 581].

Step 6, (C6). Apply Statement K.32 once to the product measure of (C6), with the data of the proof of Lemma F.14(b): $z_x = (A'(x), \varphi'(x))$, K the product of two unit balls ($c_K = \sqrt{2}$), $J = \Lambda_7^{(k)}$, the precision in the class $\mathcal{P}_{\mathbb{W}_0}$ of Lemma F.16, the scalar covariance bounds of [4, (1.9)–(1.20)] (uniform on the torus and in the zero mode, Statement K.39), $\|V^{(k)}\|_\kappa \le O(1)$ by [1, (3.58)] and Proposition E.76, and $b \ge b_*$, $\|V^{(k)}\|_\kappa b^{\rho_1} \le 1$ by Lemma A.5. The scalar half is w-free, and no complex gauge background is used (Proposition E.71). The two remainders are bounded by Statement K.44.

Step 7, (C7). We follow [2, p. 582]: terms of order $> \bar{n}$ are estimated, Dirichlet covariances are replaced by free ones at a cost of $O(t^\kappa)|\Lambda_7^{(k)}|$ (Lemma E.123), and some are composed by [1, (2.42)]. This yields $\mathcal{P}_w^{(k+1)}$. Graph values are multilinear in the lines, so the vector lines compose to $\sum_{j\le k} w_j G_{(j)} = G_{k+1} + u c_{k+1} D_{k+1}$ (proof of Lemma F.14(c)). The bounds are those of Proposition E.76, applied under the conditions of Proposition E.110. The torus enters only through δG-type external fields (Proposition E.67); the constant of the $\delta m^2_{\mathrm{fin}}$ vertex is uniform in ℓ_T (Proposition E.162).

Step 8, (C8). The field-independent output of order $\le \bar{n}$ is handled by Lemmas F.31 and F.32(V), (C11), completed by the restriction tails of Statement K.44(d). Its order-e^2 part with one D-line is Proposition E.64; the bad localisations of the C-only families with δm^2 insertions are Proposition E.175 with Lemma E.170. The collar shares are at most $C(\kappa_0)t^{\kappa_0}$ per block of Col_k: the layer has at most $(2r(t) + 2\sqrt{d}M_b + 1)^3|\mathrm{Col}_k|$ blocks, and Lemma F.11 applies.

Step 9, (C9). The first item is Statement K.40, with $t^{1/2}p^4 \le C(\kappa_0)t^{\kappa_0}$ (Lemma F.11). The second is Lemma A.6(d). The third is (C2).

Step 10, (C10) and uniformity. We follow [2, p. 582]: rescale to the $L^{k+1}\varepsilon$-lattice and collect the estimates into [2, (2.43)] at $k+1$. The new scalar form on $\Lambda_6^{(k)\prime}$ at the background $A^{(k+1),\varepsilon}$ is [2, (2.47)] at $k+1$, obtained from [2, (2.110)–(2.112)]. The factor $Z^{(k)}$ enters the density, χ_{k+1} and $\theta_{k+1}A^{(k+1)}$ are carried, and $\mathcal{P}_w^{(k+1)}$ is given by (C7). The errors of (C1)–(C8) are of the two types $C\,t^{\kappa_0}|\mathrm{Col}_k|$ and $C\,t^\kappa N_k$, which is the scalar part of Step 3 of Proposition F.2. The only factors growing with k are $K - k = \log_2(\varepsilon_0/t)$, and they always multiply t^2 or t^3 (Lemma F.11). Bałaban's per-step constants are bounded uniformly in $\varepsilon_0 \in (0, \varepsilon_1]$ (Statement K.1). Dependence on ℓ_T enters only through lower bounds on ℓ_0, through Statement K.35, and through C_b (Proposition E.162). Masses enter through $M^2t^2 \le 1$ and the mass floor, which binds at $m_1 > 0$. □

Remark F.13. (a) The bound in (C1) needs $\kappa_0 < \frac{1}{4}$: by [3, p. 421], the family of the vertex (1.6) with three external scalar legs is of order $t^{1/4}p^3$. The counterterm and $|\varphi|^6$ terms alone would allow any $\kappa_0 < \frac{1}{2}$. (b) In (C4) we use the printed volume $|\Lambda_7^{(k)}|$ of [2, Prop. 2.6]. (c) The scalar step does not use a block-by-block bound on the order-e^2 vacuum family with one D-line; that family is treated in Proposition E.64. A direct treatment cutting the index-k lines [3, p. 422] also

works, with three amendments: the per-slice factor is $L^{-(k-j)}$, not $L^{-2(k-j)}$, so the constant gains a factor $L(1-L^{-1})^{-1}$; the torus lines of index $< k$ differ from the $\eta\mathbb{Z}^d$ lines by image fields of size $O(\mathrm{e}^{-\delta\ell_0/(2\varepsilon_0)})$; the dumbbell graph vanishes, and the e^nA^n vertices with $n \geq 3$ first contribute at higher order.

Lemma F.14 (photon small-field step)**.** *Under the same hypotheses:*

(a) *After the vector translation, $A'|_{\Lambda_5^{(k)}} \sim N(0, w_k C^{(k)}_{\Lambda_5})$ in the A'-gauge. The small-field restriction $\chi(A')$ is w-free, and $|A'| \leq O(1)p$ on $\Lambda_1^{(k)}$.*

(b) *The joint fluctuation integral is bounded by the cumulant bound of Statement K.32, used once on $\mathcal{P}_{\mathbb{W}_0}$.*

(c) *The new photon line is $G_{k+1}(u) = G_{k+1} + uc_{k+1}D^{\varsigma}_{k+1}$ exactly, with $c_k = \kappa\mu_*^2L^2t_k^2 \leq \frac{1}{3}$. It satisfies the weighted tree bound* [3, (1.33)] *block by block. No new counterterm and no new Ward requirement arise. The order-e^2 vacuum family with one D-line is* not *bounded here; it is handed to Lemma F.32.*

(d) *Closing at $k+1$: $w_k^{-1} + \omega_{k+1} = w_{k+1}^{-1}$. The replacements* [2, (2.47)–(2.49)] *cost at most $\sigma_+ = (1-\eta)^{-1}$ times Bałaban's. With $X_{k+1}(B) := \langle \Lambda_7^{(k)\prime}B,\, \Delta^{(k+1)}\Lambda_6^{(k)\prime c}B\rangle$, the deletion of $\omega_{k+1}X_{k+1}$ costs $\nu(t_{k+1})|\Lambda_7^{(k)\prime}|_{k+1} + \xi_{k+1}\|A_{k+1}|_{\Lambda_6^{(k)\prime c}}\|^2$, with*

$$\nu(t) = \tfrac{1}{2}|u|\kappa L^2c_0c_dc_1^2w_{\min}^{-2}p(t)^2\mathrm{e}^{-\delta_0 r(t)/2}, \qquad \xi_{k+1} = \tfrac{1}{2}|\omega_{k+1}|c_0c_d\mathrm{e}^{-\delta_0 r(t_{k+1})/2},$$

where $c_0 = c_0^{\Delta}$, δ_0 are the constants of Lemma F.15, $c_d = \sum_{z\in\mathbb{Z}^3}\mathrm{e}^{-\delta_0|z|/2}$ and c_1 is the constant of [2, (2.55)]*. Put $HA^{(k)}_{\mathrm{del}} := HA - 2\sum_{1\leq j\leq k}\omega_jX_j(A_j)$, $\hat{\xi}_j := |u|\kappa L^2c_0c_dw_{\min}^{-2}(w_{\mathrm{lo}}\gamma_1)^{-1}\mathrm{e}^{-\delta_0 r(t_j)/2}$ and $\hat{\xi}_{\mathrm{tot}} := \sum_{1\leq j\leq K}\hat{\xi}_j$. Then $HA^{(k+1)}_{\mathrm{del}} = HA^{(k)}_{\mathrm{del}} - 2\omega_{k+1}X_{k+1}$ and $(1-\hat{\xi}_{\mathrm{tot}})HA \leq HA^{(k+1)}_{\mathrm{del}} \leq (1+\hat{\xi}_{\mathrm{tot}})HA$.*

(e) *The backward objects satisfy $H'^{w}_{k+1} = w_k^{-1}H'_{k+1} \geq 0$ and $\varrho_{k+1} \leq \frac{1}{2}$ ((E.4)). Lemma F.37 holds (photon zero mode; $\mu_* > 0$ is necessary and sufficient).*

The bounds at complex backgrounds are excluded by Proposition E.71; the Ward–Takahashi part of the defect of the zeroth moment at orders ≥ 2 is Proposition E.67.

Setting for the proof. Throughout, $d = 3$, $L = 2$, $t_k = L^k\varepsilon$, $\ell(t) = 1 + \log t^{-1}$, $r(t) = R\,\ell(t)^2$, $p(t) = b_0\ell(t)^6$. The weights are $w_j = 1+u\delta_j$ with $\delta_j = \kappa\mu_*^2t_{j+1}^2 \leq \frac{1}{12}$ (condition (E.δ)), $w_K = 1$, $\omega_k = w_k^{-1} - w_{k-1}^{-1}$, $w_{\mathrm{lo}} = \min(1, \min_k w_k^{-1})$, $\sigma_- = \frac{1}{2}$, $\sigma_+ = (1-\eta)^{-1}$. In scale-k units $M_k = \Delta^{(k)} + \hat{\beta}Q^*Q$ with $\hat{\beta} = aL^{-2}$, $C^{(k)} = M_k^{-1}$, $C^{(k)}_{\Lambda} = (\Lambda M_k\Lambda)^{-1}$, $\Delta^{(k+1)} = \beta_k - \beta_k^2QC^{(k)}Q^*$, $\mathcal{H}_k = \beta_kC^{(k)}Q^*$, $D_k(B) = \langle B,\, \Delta^{(k)}B\rangle$. Distances are torus distances; sums over a torus are dominated by sums over $\mathbb{Z}^d$. The torus satisfies Lemma 2.1(ii) and (T.nw).

Lemma F.15 (kernel decay and lower bound of the step operators)**.** *There are $c_0^{\Delta}, \delta_0 > 0$, depending on d, M_b, a, $\mu_*\varepsilon_0$ only, such that for $0 \leq k \leq K$*

$$|\Delta^{(k)}(x,y)| \leq c_0^{\Delta}\mathrm{e}^{-\delta_0|x-y|}, \qquad |M_k(x,y)| \leq c_M\mathrm{e}^{-\delta_0|x-y|}, \quad c_M := c_0^{\Delta} + \hat{\beta}\mathrm{e}^{\delta_0L\sqrt{d}},$$

$0 \leq \Delta^{(k)} \leq a_k$ for $k \geq 1$, and $\|\Delta^{(0)}\| \leq 4d + \mu_*^2\varepsilon_0^2$. Moreover $\Lambda M_k\Lambda \geq c_C^{-1}$ for every set Λ, with $c_C = 4/a_\flat + L^2/8 + L^2/a$, $a_\flat = a(1-L^{-2})$. All constants are independent of k, ε, ℓ_T, Λ, and the lower bound of $\mu_* \geq 0$.

Proof. Let $k \geq 1$. By [4, (1.14), p. 573] at $A = 0$, $\Delta^{(k)}(y,y')_{ij} = a_k\delta_{yy'}\delta_{ij} - a_k^2\langle q_k(y)_i,\, G_kq_k(y')_j\rangle$, where $q_k(y)$ has unit norm and support in $B^k(y)$ [4, (3.7), p. 588]. By [4, Cor. 2.3, (2.30), p. 580], valid on the torus [4, pp. 572, 575], $|\Delta^{(k)}(y,y')| \leq a_k\delta_{yy'} + a_k^2c_0\mathrm{e}^{-\delta_0\,\mathrm{dist}(B^k(y),B^k(y'))} \leq (a + a^2c_0\mathrm{e}^{\delta_0\sqrt{d}})\mathrm{e}^{-\delta_0|y-y'|}$. Put $Y = a_k^{1/2}Q_k$ and $X = -\Delta + m^2 \geq 0$. Then $a_kQ_kG_kQ_k^* = Y(X + Y^*Y)^{-1}Y^* \leq I$, as the limit $\delta \downarrow 0$ of $Y(X+\delta+Y^*Y)^{-1}Y^* = I - (I + Y(X+\delta)^{-1}Y^*)^{-1}$.

Hence $0 \le \Delta^{(k)} \le a_k$; this also covers tori too small for the cubes of [4], at the cost of a factor $\mathrm{e}^{\delta_0 D}$, D the diameter. For $k=0$, $\Delta^{(0)} = -\Delta + \mu_*^2$ is nearest-neighbour with entries at most $2d + \mu_*^2\varepsilon_0^2$. The kernel of Q^*Q is bounded by 1 and vanishes for $|x-y| > L\sqrt{d}$, which gives c_M. The lower bound is step (a) of the proof of Lemma H.27 ($C^{(k)} \le c_C t_k^2$ in physical units); for $k=0$ the same chain holds with the term $4/r_j$ absent. A principal block of an operator $\ge c_C^{-1}$ is $\ge c_C^{-1}$. □

Lemma F.16 (class closure)**.** *Let $\mathcal{P}(\gamma_0, \gamma_1, c_0, \delta_0)$ be the set of real symmetric block operators P with $\gamma_0 \le P \le \gamma_1$ and $\|P(x,y)\| \le c_0 \mathrm{e}^{-\delta_0|x-y|}$. If $P_A, P_\varphi \in \mathcal{P}(\gamma_0, \gamma_1, c_0, \delta_0)$ and $s \in [\sigma_-, \sigma_+]$, then $sP_A \oplus P_\varphi \in \mathcal{P}_{\mathbb{W}_0} := \mathcal{P}(\sigma_-\gamma_0, \sigma_+\gamma_1, \sigma_+ c_0, \delta_0)$.*

Proof. Multiply the inequalities for P_A by s; the scalar block is unchanged. □

Lemma F.17 (weighted lines)**.** *Let $\mathcal{L} = \sum_i \mathcal{L}_i$ be a vector line obtained from the step covariances $C^{(i)}_{\Lambda_5^{(i)}}$, written by the step of origin i, by the operations listed in Statement K.45. In the weighted chain the line at the same place is $\sum_i w_i \mathcal{L}_i$. Its D-part is $u\sum_i \delta_i \mathcal{L}_i$, and every bound used for $\mathcal{L}_i$ holds for the D-part with the extra factor $|u|\delta_i \le |u| c_k$.*

Proof. By part (a), the step-i fluctuation has the law $N(0, w_i C^{(i)}_{\Lambda_5^{(i)}})$. The translations, the conditional means, the lift and $\chi(A')$ are w-free. A line is a covariance pushed forward by such linear maps $\mathcal{M}, \mathcal{M}'$, and $\mathcal{M}(w_i C)\mathcal{M}'^* = w_i\, \mathcal{M} C \mathcal{M}'^*$. Restriction to domains, Dirichlet-to-free replacement, localisation cut-offs and differences are linear in the covariance and act step by step. Vector lines do not depend on the background, so the same holds for the δG-type differences, which concern scalar lines only. □

Lemma F.18 (per-block bounds of the D-coloured part)**.** *Let $|u| c_k \le 1$. Then the D-coloured part of $\mathcal{P}_w^{(k)}$ obeys the localised bounds* [2, (2.57), (2.116)] *with constant at most $2^{N_\ell}|u|c_k$ times the $u=0$ constant.*

Proof. At $u=0$, (2.57) and (2.116) follow by summing the right side of [3, (1.33)] over the families of order $\le \bar{n}$ with a vertex in a given block, using only the factors $e_k^{d_v}\lambda_k^{d_s}$, the decay $\mathrm{e}^{-\delta_0 d}$, external-field norms bounded by the restrictions, and the finiteness of the set of graphs. By Theorem H.24(c), a coloured family has the same right side times $\prod_{D\text{-lines}} |u| c_k (L^j \varsigma)^2 \le (|u|c_k)^m$. Colourings multiply the number of graphs by at most 2^{N_ℓ}. □

Lemma F.19 (step covariances: decay and domain differences)**.** *(i) Choose $\delta' \in (0, \delta_0/2]$ with $c_M \sum_{z\in\mathbb{Z}^d} (\mathrm{e}^{\delta'|z|} - 1)\mathrm{e}^{-\delta_0|z|} \le (2c_C)^{-1}$. Then $|C_\Lambda^{(k)}(x,y)| \le 2c_C \mathrm{e}^{-\delta'|x-y|}$ for every set Λ. (ii) For $x, x' \in \Lambda$, $(C^{(k)} - C_\Lambda^{(k)})(x,x') = -\sum_{z\in\Lambda, w\notin\Lambda} C_\Lambda^{(k)}(x,z) M_k(z,w) C^{(k)}(w,x')$. (iii) With $d_x = \mathrm{dist}(x, \Lambda^c)$, $c_d(s) = \sum_{z\in\mathbb{Z}^d} \mathrm{e}^{-s|z|}$ and $K_1 = 4c_C^2 c_M c_d(\delta'/2)^2$, $|(C^{(k)} - C_\Lambda^{(k)})(x,x')| \le K_1 \mathrm{e}^{-(\delta'/4)(|x-x'| + d_x + d_{x'})}$.*

Proof. (i) Let ρ be 1-Lipschitz and $M^\rho = \mathrm{e}^{\delta'\rho}\Lambda M_k \Lambda \mathrm{e}^{-\delta'\rho} = \Lambda M_k \Lambda + R$. Then $|R(x,y)| \le (\mathrm{e}^{\delta'|x-y|} - 1)|M_k(x,y)|$, so $\|R\| \le (2c_C)^{-1}$ by the Schur test and Lemma F.15, and $\|(M^\rho)^{-1}\| \le (c_C^{-1} - \|R\|)^{-1} \le 2c_C$. Take $\rho = \mathrm{dist}(\cdot, y)$. (ii) The rows of $M_k C^{(k)} = I$ in Λ give $(\Lambda M_k \Lambda) C_{\Lambda\Lambda} + M_{\Lambda\Lambda^c} C_{\Lambda^c\Lambda} = I_\Lambda$; multiply by C_Λ. (iii) By (ii), (i) and Lemma F.15, the left side is at most $4c_C^2 c_M \sum_{z,w} \mathrm{e}^{-\delta' E}$, $E = |x-z| + |z-w| + |w-x'|$. Since $w \notin \Lambda$, $E \ge d_x + d_{x'}$, and $E \ge |x-x'|$; so $\mathrm{e}^{-\delta' E} \le \mathrm{e}^{-(\delta'/4)(|x-x'| + d_x + d_{x'})} \mathrm{e}^{-\delta' E/2}$, and $\sum_{z,w} \mathrm{e}^{-\delta' E/2} \le c_d(\delta'/2)^2$. □

Lemma F.20 (zero modes)**.** *(a) For every $\Lambda \subset T^{(k)}$, including $\Lambda = T^{(k)}$, $w_k C_\Lambda^{(k)} \le w_k c_C$ in scale-k units, independently of $\mu_* \ge 0$ and ℓ_T. (b) HA is positive definite if and only if $\mu_* > 0$. Then its only candidate flat direction, A_K constant with the finer fields at their conditional means, has precision $\langle c1, \Delta^{(K)} c1\rangle \ge \gamma_1 \mu_*^2 |T| |c|^2$, and the mean $\bar{A}$ of A has variance in*

$[\min(1,w_{\min}),\max(1,w_{\max})]\,(\mu_*^2|T|)^{-1}$ *per component. (c) The constant mode* c *of a real background enters no scalar operator bound used in the step: the bounds of* [4, (1.8), (1.9)–(1.12), p. 573; (1.15)–(1.22), p. 574] *are stated for subsets of a torus with periodic conditions* [4, p. 572]*, uniformly in the background, and since* $\partial^\eta c=0$*,* $c+A''$ *is regular iff* A'' *is, with the same constants. (d) The event* $|\bar A|\ge c_1p(t)/(2\mu_*t)$*, for one scale* $t\le\varepsilon_0$*, has probability at most* $6\exp(-c_1^2p(t)^2|T|/(24w_{\max}t^2))$*.*

Proof. (a) Lemma F.15. (b) By Lemma F.5, $HA=\sum_{k<K}w_k^{-1}\langle\hat S_k,\,M_k\hat S_k\rangle+D_K(A_K)$ with $\hat S_k=A_k-\mathcal{H}_kA_{k+1}$, a unit-triangular change of variables. Since $M_k\ge c_C^{-1}$, $HA>0$ iff $\Delta^{(K)}>0$. If $\mu_*>0$, $\Delta^{(K)}\ge\gamma_1\mu_*^2$ by Lemma F.37. If $\mu_*=0$, then $\Delta^{(0)}1=0$, $M_01=\beta_01$ gives $\Delta^{(1)}1=0$, and inductively $\Delta^{(K)}1=0$, so $\mathrm{e}^{-\frac12HA}$ is not integrable along the constants. For the variance, $C_B(u)=\sum_{j<K}w_jH_j+G_K$ and $C_{\mu_*}=\sum_{j<K}H_j+G_K$ with $H_j,G_K\ge0$, so $\min(1,w_{\min})C_{\mu_*}\le C_B(u)\le\max(1,w_{\max})C_{\mu_*}$; moreover $\operatorname{Var}\bar A=|T|^{-2}\langle1,\,C_B1\rangle$ and $C_{\mu_*}1=\mu_*^{-2}1$. (c) The cited statements, together with $\partial^\eta c=0$. (d) By (b), each of the three components of $\bar A$ is centred Gaussian with variance at most $w_{\max}(\mu_*^2|T|)^{-1}$; if $|\bar A|\ge s$, some component is at least $s/\sqrt3$. Use the Gaussian tail with $s=c_1p(t)/(2\mu_*t)$. □

Lemma F.21 (backward replacement, vector sector)**.** *Let* $\Lambda_5:=\Lambda_5^{(k+1)}\subset\Lambda_0^{(k+1)}\subset\Lambda_7^{(k)\prime}$ *and, in scale-*$(k+1)$ *units,* $H'_{k+1}:=\Lambda_5(\Delta^{(k+1)}_{\Lambda_5^{(k)}}-\Delta^{(k+1)})\Lambda_5=\beta_k^2\Lambda_5Q(C^{(k)}-C^{(k)}_{\Lambda_5^{(k)}})Q^*\Lambda_5$*. (i)* $H'_{k+1}\ge0$ *and* $\|H'_{k+1}\|\le C_{H'}\mathrm{e}^{-\delta' r(t_k)}$*,* $C_{H'}=\hat\beta^2K_1\mathrm{e}^{\delta'L\sqrt d}c_d(\delta'L/4)$*. (ii) In the weighted chain the difference between the exact and the replaced Hessian is* $H'^w_{k+1}=w_k^{-1}H'_{k+1}$*, and the linear part is* $F'^w_{k+1}=w_k^{-1}F'_{k+1}$*. (iii) The exact Hessian* G^w_{k+1} *satisfies* $G^w_{k+1}\ge G^w_{k+1}-H'^w_{k+1}\ge(1-\hat\xi_{\mathrm{tot}})w_{\mathrm{lo}}\gamma_1\mu_*^2t_{k+1}^2$*. (iv)* $X^w:=G^{w,-1/2}H'^wG^{w,-1/2}$ *satisfies* $0\le X^w\le\varrho^{\det}_{k+1}$ *with* $\varrho^{\det}_{k+1}\le\frac{24}{11}(1-\eta)^{-1}C_{H'}\gamma_1^{-1}\mu_*^{-2}t_{k+1}^{-2}\mathrm{e}^{-\delta' r(t_k)}\le\frac12$ *under (E.4) and* $\varepsilon_0\le\exp(1-1/(\delta'R))$*; and* $1\le[\det(I-X^w)]^{-1/2}\le\exp(d\varrho^{\det}_{k+1}|\Lambda_5|)$*.*

Proof. (i) The identity is the definition of $\Delta^{(k+1)}$ and of its conditional version. Positivity: $\langle f,\,C_\Lambda f\rangle=\sup_{g\subset\Lambda}[2\langle f,\,g\rangle-\langle g,\,M_kg\rangle]\le\langle f,\,C^{(k)}f\rangle$. The fine sites of the blocks of Λ_5 satisfy $d_x\ge2r(t_k)-L\sqrt d$ by [2, (2.8)]; Lemma F.19(iii), the averaging weights of Q and the Schur test give the bound. (ii) The variables $A_{k+1}|_{\Lambda_5}$ occur only in the carried form $w_{k+1}^{-1}\Lambda_6^{(k)\prime}\Delta^{(k+1)}\Lambda_6^{(k)\prime}$, in the exact composed kernels of later steps, and in scalar forms through the external field $\tilde A$; the rest of the density depends on A_{k+1} only through $\Lambda_7^{(k)\prime c}A_{k+1}$ (step (d1) below). On $\Lambda_5^{(k)}$ the A'-Gaussian is w_k^{-1} times Bałaban's (part (a)), and minimisation is homogeneous. Hence the exact step-k output, as a form in A_{k+1}, is w_k^{-1} times Bałaban's exact form plus $\omega_{k+1}(D_{k+1}-2X_{k+1})$, and the replaced one is w_k^{-1} times his replaced form plus the same term. Subtract; X_{k+1} has no block on $\Lambda_5\times\Lambda_5$. (iii) The exact Hessian is the principal Λ_5-block of the Hessian of the carried form of the chain from level $k+1$, minimised over the coarser variables. By Lemma F.5 and the sandwich of part (d) (step (d2) below, which does not use this lemma), this form is at least $(1-\hat\xi_{\mathrm{tot}})w_{\mathrm{lo}}$ times the unweighted chain $\sum_{l>k}I_l+D_K$. The identity $D_l(B)+\beta_l\|QB-B'\|^2=I_l(B,B')+D_{l+1}(B')$, minimised over B', shows inductively that the minimum of the unweighted chain over the coarser variables is $D_{k+1}(A_{k+1})$. Inequalities between forms pass to minima and principal blocks, and $\lambda_{\min}(\Delta^{(k+1)})\ge\gamma_1\mu_*^2t_{k+1}^2$ (Lemma F.37). Finally $G\ge G-H'$ by (i). (iv) $\|X^w\|\le w_k^{-1}\|H'_{k+1}\|/((1-\hat\xi_{\mathrm{tot}})w_{\mathrm{lo}}\gamma_1\mu_*^2t_{k+1}^2)$, with $w_k^{-1}\le\sigma_+$, $w_{\mathrm{lo}}\ge\sigma_-$ and $(1-\hat\xi_{\mathrm{tot}})^{-1}\le\frac{12}{11}$ (condition (E.14)). Since $r(t_k)\ge r(t_{k+1})$, the right side is at most $g(t_{k+1})$, $g(t)=\mathrm{const}\cdot t^{-2}\mathrm{e}^{-\delta'R\ell(t)^2}$, which is increasing on $(0,\exp(1-1/(\delta'R))]$; so $g(t_{k+1})\le g(\varepsilon_0)\le\frac12$ by (E.4). Finally $-\frac12\log\det(I-X)=\frac12\sum_{n\ge1}\operatorname{tr}X^n/n\le\frac12\operatorname{tr}X/(1-\varrho)\le d|\Lambda_5|\varrho$, and the ratio is ≥1 since $X\ge0$. □

The linear terms are Bałaban's estimates [2, (3.13)–(3.17), pp. 585–586] with γ_1 replaced by $(1-\hat\xi_{\rm tot})w_{\rm lo}\gamma_1$ and $|F'^{w}| \le \sigma_+|F'|$: $\frac12\langle\Phi,[(G-H')^{-1}-G^{-1}]\Phi\rangle = \frac12\langle\Phi,G^{-1}H'(G-H')^{-1}\Phi\rangle = O(t_{k+1}^{\kappa})|\Lambda_5|$ for every κ, and similarly for the F' terms.

Proof of Lemma F.14. (a). By Lemma F.5 and (IH_k), the terms of the vector exponent that depend on $A_k|_{\Lambda_0^{(k)}}$ are $\frac12 w_k^{-1}\langle\Lambda_6 A_k,\,\Delta^{(k)}\Lambda_6 A_k\rangle$ and $\frac12 w_k^{-1}\beta_k\|QA_k - A_{k+1}\|^2$; the spectator $\frac12\omega_{k+1}D_{k+1}(A_{k+1})$ does not involve A_k. A common positive factor of the quadratic and linear parts does not move the minimiser, so the translation [2, (2.80)] is Bałaban's and on $\Lambda_5^{(k)}$ the form in A' is w_k^{-1} times $\frac12\langle\Lambda_5^{(k)}A',\,(C^{(k)}_{\Lambda_5^{(k)}})^{-1}\Lambda_5^{(k)}A'\rangle$ [2, p. 575]. The restriction [2, (2.86)], which gives $|A'|\le O(1)p$ on $\Lambda_1^{(k)}$, and the split [2, (2.88)–(2.91)] are statements about the configuration and the lift, hence w-free.

(b). In scale-k units $P_A := (C^{(k)}_{\Lambda_5^{(k)}})^{-1} = \Lambda_5^{(k)}M_k\Lambda_5^{(k)}$ satisfies $c_C^{-1} \le P_A \le \gamma_1^A := \max(a, 4d + \mu_*^2\varepsilon_0^2) + \hat\beta\|Q\|^2$ and $|P_A(x,y)| \le c_M \mathrm{e}^{-\delta_0|x-y|}$, uniformly in $\Lambda_5^{(k)}$, k, ε, ℓ_T, μ_* (Lemma F.15). The joint precision is $w_k^{-1}P_A \oplus P_\varphi$, with P_φ the scalar precision of Lemma F.8(C6), which depends on $B^{(k+1)}$ and not on A'. Since $w_k^{-1} \in [\frac{12}{13},\frac{12}{11}] \subset [\sigma_-,\sigma_+]$, Lemma F.16 puts it in $\mathcal{P}_{\mathbb{W}_0}$ with $\gamma_0 = \min(c_C^{-1},\gamma_0^\varphi)$, $\gamma_1 = \max(\gamma_1^A,\gamma_1^\varphi)$, $c_0 = \max(c_M, c_0^\varphi)$, $\delta_0 = \min(\delta_0,\delta_0^\varphi)$. The interaction $V_w^{(k)}$ is a real polynomial in (A',φ') of degree $\le n(\bar n)$ [1, Prop. 3.2, (3.57), p. 622], [2, p. 581], with coefficients depending on the real background only; they obey [1, (3.58)] with rate $\kappa_* = \frac14 - \frac{1}{56} = \frac{13}{56}$, times the colouring factor $(1+|u|c_k)^{N_v} \le (4/3)^{N_v}$ (Proposition E.76). So $\mathsf{A}_w := \|V_w^{(k)}\|_\kappa \le C_A t_k^{13/56}$. The restrictions $\chi(A')\chi(\varphi')$ of [2, (2.86), (2.114), p. 581] are w-free and of the form χ_b with $b = c_b p(t_k)$, K the product of two unit balls. Statement K.32, applied once in $\mathcal{P}_{\mathbb{W}_0}$ with $t = \bar n = 13$, $b \ge b_*$ and $\mathsf{A}_w b^{\rho_1} \le 1$ (Lemma A.5), gives

$$\int d\mu_{w_k C^{(k)}_{\Lambda_5^{(k)}}}(A') \int d\mu_{C^{(k)}_{\Lambda_5^{(k)}}(\cdot)}(\varphi')\,\chi(A')\chi(\varphi')\mathrm{e}^{V_w^{(k)}} \le \exp\Big\{\sum_{m\le\bar n}\frac{\kappa(V_w^{(k)};m)}{m!} + S|\Lambda_7^{(k)}|\big[(\mathsf{A}_w b^{\rho_1})^{14} + \mathrm{e}^{-\rho_3 b^{3/2}}\big]\Big\}.$$

Now $(\mathsf{A}_w b^{\rho_1})^{14} \le C\,t_k^{13/4}\ell(t_k)^{84\rho_1}$ with $\sup_{t\le\varepsilon_0} t^{1/20}\ell(t)^{84\rho_1} < \infty$, and $\mathrm{e}^{-\rho_3 b^{3/2}} = \exp(-\rho_3(c_b b_0)^{3/2}\ell(t_k)^9) \le C_N t_k^N$. So the remainder is at most $C_{\rm R}t_k^{16/5}|\Lambda_7^{(k)}|$, with $C_{\rm R}$ depending only on the parameters and η (see also Statement K.44). Only this upper bound is used, and every background at which a bound is invoked is real (Proposition E.71).

(c). The line. Each $A'A'$ contraction carries $w_k C^{(k)}_{\Lambda_5^{(k)}}$. Statement K.33, with collar $\ge 2r(t_k)$, replaces it by $w_k C^{(k)}$ at a cost $O(t_k^\kappa)|\Lambda_7^{(k)}|$ for every κ. The interaction sees A' only through the w-free lift F_k, so the new line is $w_k F_k C^{(k)} F_k^* = w_k G_{(k)}$. Vector lines are background-free, so the composition [1, (2.42)] (cf. [2, p. 582]) is linear: $\sum_{j\le k} w_j G_{(j)} = G_{k+1} + u\sum_{j\le k}\delta_j G_{(j)} = G_{k+1} + u\,c_{k+1}D^{\varsigma}_{k+1}$, because $c_{k+1}(L^j\varsigma')^2 = \kappa\mu_*^2 L^{2j+2}\varepsilon^2 = \delta_j$ with $\varsigma' = L^{-k-1}$. Every graph value is multilinear in its lines, so $\mathcal{P}_w^{(k+1)} = \sum_m (uc_{k+1})^m \mathcal{P}_m^{(k+1)}$ exactly. *Bounds.* Each coloured family obeys Theorem H.24(c), with Bałaban's rate and constants independent of ε, k, the domains, the background, u and κ. By Lemma F.18 with $c_{k+1} \le \frac13$, the bounds [2, (2.57), (2.116)] hold for the D-coloured part with constant $\le 2^{N_\ell}|u|c_{k+1}$ times the $u=0$ constant, and for all of $\mathcal{P}_w^{(k+1)}$ with $(4/3)^{N_v}$ times it. In the remaining weighted uses (the domain changes [2, (2.32)–(2.42)], [1, Props. 3.1–3.2], the output of [2, p. 582] and [2, (2.117)–(2.118)]) u enters only through line weights (Lemma F.17 and Statement K.45) and family factors. The expansions of $Z^{(j)}$ [2, (2.92)–(2.107)] contain no vector line. *Counterterms.* The vertex list of $\mathcal{P}_w^{(k+1)}$ is Bałaban's $u=0$ list, with the scale split of the fixed $\delta m^2_{\rm ref}$ [3, (1.26)–(1.28)] at $k+1$; passing from k to $k+1$ moves C-only, u-independent pieces between the terms of the split. By Lemma H.18(c),(d), every connected coloured subgraph with an external leg and $m \ge 1$ has $D^\natural \ge 1$, and every divergent subgraph is C-only

and one of Bałaban's classes; by Lemma H.23 these are renormalised by $\delta m^2_{\mathrm{ref}}$ and their Ward–Takahashi and $\varphi\varphi A$ partners exactly as at $u=0$. No u-dependent vertex arises. Transversality is needed only for C-only vacuum-polarisation subgraphs [3, (3.25)–(3.32)], and on T_ε it holds exactly by Proposition E.67; a vacuum-polarisation subgraph with an internal D-line has $D^\natural \geq 2$. *The excluded family.* The only D-coloured family with $D^\natural = 0$ is the vacuum family with one D-line at order e^2. At a fixed largest index it is $O(e^2|u|\kappa\mu_*^2 L^2 t_k^3)$ per block before cancellation, so booked on every block it would sum to $O(|T|\log(\varepsilon_0/\varepsilon))$. It is not bounded here; it cancels identically in $E_1(u)$, and step by step up to the bad localisations (Lemmas F.32 and F.31).

(d). Write $B := A_{k+1}$, $\Lambda'_6 := \Lambda_6^{(k)\prime}$, $\Lambda'_7 := \Lambda_7^{(k)\prime}$; $\Lambda'_7 \subset \Lambda'_6$ and $\mathrm{dist}(\Lambda'_7, \Lambda_6'^c) \geq r(t_{k+1})$ [2, (2.8)]. The identity $w_k^{-1} + \omega_{k+1} = w_{k+1}^{-1}$ is the definition of ω_{k+1}. *(d1) Absorption.* With $\Sigma_{k+1}(B) := \langle \Lambda_6'^c B, \Delta^{(k+1)}\Lambda_6'^c B\rangle + 2\langle(\Lambda'_6 \setminus \Lambda'_7)B, \Delta^{(k+1)}\Lambda_6'^c B\rangle$ we have $D_{k+1}(B) = \langle \Lambda'_6 B, \Delta^{(k+1)}\Lambda'_6 B\rangle + \Sigma_{k+1}(B) + 2X_{k+1}(B)$. Adding the spectator $\frac12\omega_{k+1}D_{k+1}(B)$ of $\mathcal{K}_k^w$ (Lemma F.5) to the carried form gives exactly $\frac12 w_k^{-1}\langle \Lambda'_6 B, \Delta^{(k+1)}\Lambda'_6 B\rangle + \frac12\omega_{k+1}D_{k+1}(B) = \frac12 w_{k+1}^{-1}\langle \Lambda'_6 B, \Delta^{(k+1)}\Lambda'_6 B\rangle + \frac12\omega_{k+1}\Sigma_{k+1}(B) + \omega_{k+1}X_{k+1}(B)$. The first term is the carried form at $k+1$. Σ_{k+1} depends on B only through $\Lambda_7'^c B$, which is never integrated forward ($\Lambda_5^{(k+1)} \subset \Lambda_0^{(k+1)} \subset \Lambda'_7$); it joins the vector part of $\rho_w^{(k+1)}$. *(d2) Deletion.* By Lemma F.15, $|x-y| \geq r(t_{k+1})$ for $x \in \Lambda'_7$, $y \in \Lambda_6'^c$, and the Schur test, $|\omega_{k+1}X_{k+1}(B)| \leq \frac12|\omega_{k+1}|c_0c_d\mathrm{e}^{-\delta_0 r(t_{k+1})/2}(\|B|_{\Lambda'_7}\|^2 + \|B|_{\Lambda_6'^c}\|^2)$. Since $|\omega_{k+1}| = |u|(\delta_{k+1}-\delta_k)/(w_kw_{k+1}) \leq |u|\kappa L^2\mu_*^2 t_{k+1}^2 w_{\min}^{-2}$ and $|B| \leq c_1 p(t_{k+1})/(\mu_* t_{k+1})$ on Λ'_7 by the restrictions [2, (2.55)] at scale $k+1$, the first part is $\leq \nu(t_{k+1})|\Lambda'_7|_{k+1}$ and the second is $\xi_{k+1}\|B|_{\Lambda_6'^c}\|^2$. The term $\omega_{k+1}X_{k+1}$ is deleted from the carried form, so $HA_{\mathrm{del}}^{(k+1)} = HA_{\mathrm{del}}^{(k)} - 2\omega_{k+1}X_{k+1}$. For the sandwich, the same Schur test gives $|2\omega_j X_j(A_j)| \leq |\omega_j|c_0c_d\mathrm{e}^{-\delta_0 r(t_j)/2}\|A_j\|^2$. By Lemma F.5, HA is at least w_{lo} times the unweighted chain, whose minimum over all variables other than A_j is at least $D_j(A_j) \geq \gamma_1\mu_*^2 t_j^2\|A_j\|^2$ (the minimisation identity of the proof of Lemma F.21(iii), and Lemma F.37). Hence $|2\omega_j X_j(A_j)| \leq \hat\xi_j HA$, the factor $\mu_*^2 t_j^2$ cancelling, and $|HA_{\mathrm{del}}^{(k+1)} - HA| \leq \hat\xi_{\mathrm{tot}} HA$ as forms. *(d3) Replacement.* After the step-k integration the minimised form is w_k^{-1} times Bałaban's (part (a)), and his replacement [2, (2.47)–(2.49)] of it, with Neumann conditions not introduced [2, p. 568], costs $w_k^{-1}O(t_k^\kappa)|\Lambda_5^{(k)}| \leq \sigma_+$ times his cost: by Lemma F.21(ii),(iv) and the bound on the linear terms after it.

(e). The identities and bounds for $H_{k+1}'^w$, the floor and $\varrho_{k+1}^{\det} \leq \frac12$ are Lemma F.21; the zero-mode statements are Lemma F.20. Inside the step the constant mode enters only in four places: (i) through $\lambda_{\min}(\Delta^{(k+1)})$, where in the deletion coefficients of (d) the factor μ_*^2 cancels against $|\omega_{k+1}|$ and survives only in $\varrho^{\det}$ (condition (E.4)); (ii) through the restriction $|A| \leq c_1 p/(\mu_* t)$ of [2, (2.55), p. 570], Lemma F.20(d) being only a size estimate; (iii) through the scalar conditions, which are uniform in it by Lemma F.20(c); (iv) through c-numbers that do not increase with ℓ_T, bounded at the floor by Lemma A.10. □

(S3) The large-field part of the step and the region sum.

The large-field step: notation. In this paragraph and in Lemmas F.22–F.29, the letters j and k are scale indices. We have $d=3$ and $L=2$. Put $t_k = L^k\varepsilon$, $\ell(t) = 1 + \log t^{-1}$, $\ell_k = \ell(t_k)$, $\hat\ell = \ell(\varepsilon_0)$, $r_k = r(t_k) = R\ell_k^\varrho$ with $\varrho = 2$, and $p_k = p(t_k) = b_0\ell_k^6$. $T^{(k)}$ is the unit lattice of scale k. It has $N_k = |T|t_k^{-3}$ points, and $|X|_k$ is the number of points of $X \subset T^{(k)}$. Distances are ℓ^∞ distances on the torus, and $\mathcal{N}_s(X)$ is the closed s-neighbourhood of X. For a union X of L-blocks of $T^{(k-1)}$, $X' \subset T^{(k)}$ is its image at scale k [2, p. 566]. For a set X at scale k, $B^j(X)$ is the corresponding set at scale $k-j$, and $B(x) := B^1(\{x\})$. A large block of $T^{(k)}$ is a cube of side $b := M_bL$. By Lemma 2.1, $\ell_T/(M_bt_k) \in 2\mathbb{N}$ for $k \leq K$, so $T^{(k)}$ is a union of large blocks. We assume $\vartheta_M := M_bL/R \leq \frac17$. Since $r_k \geq R$, this gives $L \leq b \leq \vartheta_M r_k$.

A *family* at scale k is a six-tuple $\mathfrak{t} = (P_v, Q_v, R_v, P_s, Q_s, R_s)$ whose entries are sets of coarse points, of bonds or plaquettes, and of points, as in [2, (2.2)–(2.5), p. 557] at scale t_k. Put $|\mathfrak{t}| := |P_v| + \cdots + |R_s|$ and $S(\mathfrak{t}) := B(P_v) \cup Q_v \cup R_v \cup B(P_s) \cup Q_s \cup R_s$. The sets $\Lambda_i^{(k)}$, $i \le 8$, are built from $\Lambda_0^{(k)}$ by [2, (2.8)] at radius r_k [2, p. 566], and $\Lambda_i^{(-1)\prime} := T^{(0)}$. We write $X_k := \Lambda_0^{(k)c}$, $\mathrm{Col}_k := \Lambda_7^{(k-1)\prime} \cap \Lambda_7^{(k)c}$ (the collar), $\mathrm{Sh}_k := \Lambda_5^{(k-1)\prime} \cap \Lambda_5^{(k)c}$ (the shell) and $W_k := B^k(\Lambda_7^{(k)})$. Finally $n_d := 4 + 2\max\{d, d(d-1)/2\} = 10$.

Lemma F.22 (the large-field part of the step)**.** *Assume the same hypotheses as above, that is, (IH$_k$) of* (F.2) *with $0 \le k \le K-1$, at either point $(u, M) \in \{(1, m_2), (-\eta, m_1)\}$.*

(a) *The scale-t_k restrictions* [2, (2.2)–(2.6), (2.55)] *are conditions on field values in A'-coordinates. They do not involve the weights. $\Lambda_0^{(k)}$ is the union of the large blocks of $T^{(k)}$ that are contained in $\Lambda_7^{(k-1)\prime}$ and lie at distance $> r_k$ from $S(\mathfrak{t}_k)$, for a minimal admissible family $\mathfrak{t}_k$ (Lemma F.23). So it is a function of the past and of $\mathfrak{t}_k$. The region expansion* [2, (2.9)–(2.10)] *holds in the direction "$\le$".*

(b) *On Col_k the interaction is removed under the step-$(k-1)$ restrictions* [2, (2.55)]*. This includes the D-coloured part and the mass counterterm pieces of step k. No vacuum constant is removed on Col_k. The cost is $c_C^{(k)} t_k^{\kappa_0} |\mathrm{Col}_k|_k$. For $k \ge 1$, $c_C^{(k)} \le \sigma_+ 2^{N_\ell} c_C^{u=0}$, where $c_C^{u=0}$ is the collar constant of* [2, Prop. 2.1, (2.57)] *at $u = 0$ and σ_+ comes from Lemma F.14(d). At $k = 0$ the cost is at most $C_\delta(M)^2\varepsilon/(16\lambda)$ per site, so $c_C^{(0)} := C_\delta(M)^2/(16\lambda)$ may be taken. The moved vacuum pieces localised outside W_k cost at most $C_{\mathrm{old}} t_k$ per block of $\Lambda_7^{(k)c}$, with C_{old} as in Lemma F.32(V).*

(c) *The forward clause carries the characteristic functions χ^c, which do not involve the weights, and the quartic factor. The quartic factor produces $\exp(-p_k^2 |R_s|)$. No vector gain is carried forward.*

(d) *The scale-k factor delivered by the backward integration is at most $\zeta_w''(\Lambda_0^{(k)})$. Here W_k^w is as in Proposition F.29, $\vartheta = \frac{1}{2}(1 - \hat{\xi}_{\mathrm{tot}}) w_{\mathrm{lo}}$, $\vartheta_+ \ge \frac{11}{26}$ at $(1, m_2)$ and $\vartheta_- \ge \frac{11}{24}$ at $(-\eta, m_1)$. Each family element therefore gains $\mathrm{e}^{-c_0'^w p_k^2}$, with $c_0'^w = \vartheta \min\{\frac{1}{8}a, \frac{1}{4}\gamma_0\}$ and γ_0 as in Proposition F.27. Each shell site costs at most $\mathrm{e}^{c_B^w}$, where $c_B^w = c_B + \frac{d}{2}\log 3 \cdot c_{\mathrm{sh}}$, c_B is the shell constant of* [2, (3.42)] *at $u = 0$, and c_{sh} is the number of kept vector variables per shell site.*

(e) *The region sum is bounded by Proposition F.29 under* (F.4)*, or by Lemma E.95 under (E.comb) with $c_A + C_{\mathrm{old}}$.*

All constants are independent of ε, k, $\mathbf{\Lambda}$ and $\ell_T \ge \ell_0$, and one ε_0 serves both points.

The region expansion is used in the direction "$\le$" only. The source prints "$=$", and equality can fail (Lemma F.23(iii)).

Lemma F.23 (regions are functions of their families)**.** *For $Y \subset T^{(k)}$ and a family $\mathfrak{t}$, let $\mathfrak{L}_k(Y, \mathfrak{t})$ be the union of the large blocks of $T^{(k)}$ that are contained in Y and lie at distance $> r_k$ from $S(\mathfrak{t})$. Call $\mathfrak{t}$ admissible for $\Lambda_0^{(k)}$ if $\Lambda_0^{(k)} = \mathfrak{L}_k(\Lambda_7^{(k-1)\prime}, \mathfrak{t})$* [2, p. 570]*. At $k = 0$ the distance condition is read as in* [2, (2.7)]*.*

(i) *Define $\mathbf{\Lambda}(\mathfrak{t})$ recursively by $\Lambda_0^{(k)} := \mathfrak{L}_k(\Lambda_7^{(k-1)\prime}, \mathfrak{t}_k)$. This assigns to each tuple $(\mathfrak{t}_0, \dots, \mathfrak{t}_{K-1})$ at most one region sequence. Every pair $(\mathbf{\Lambda}, (\mathfrak{t}_k)_k)$ in which each $\mathfrak{t}_k$ is admissible for $\Lambda_0^{(k)}$ is of the form $(\mathbf{\Lambda}(\mathfrak{t}), \mathfrak{t})$.*

(ii) *For all weights $W_k \ge 0$ on families and every $\Phi \ge 0$,*

$$\sum_{\mathbf{\Lambda}} \prod_{k<K} \Big(\sum_{\mathfrak{t} \text{ adm. min. for } \Lambda_0^{(k)}} W_k(\mathfrak{t}) \Big) \Phi(\mathbf{\Lambda}) \le \sum_{(\mathfrak{t}_k)_{k<K}:\ \mathbf{\Lambda}(\mathfrak{t}) \text{ adm.}} \prod_{k<K} W_k(\mathfrak{t}_k)\, \Phi(\mathbf{\Lambda}(\mathfrak{t})).$$

(iii) *The resummation* [2, (2.9)] *holds with "$\le$" in place of "$=$".*

Proof. (i) By [2, p. 570], $\Lambda_0^{(k)}$ is determined by $\Lambda_7^{(k-1)\prime}$ and $S(\mathfrak{t}_k)$. By [2, (2.8), p. 566], $\Lambda_7^{(k-1)\prime}$ is determined by $\Lambda_0^{(k-1)}$. Now induct on k.

(ii) Expanding the left side gives a sum over pairs $(\mathbf{\Lambda}, \mathfrak{t})$. By (i), the map $(\mathbf{\Lambda}, \mathfrak{t}) \mapsto \mathfrak{t}$ is injective, and enlarging the index set adds only non-negative terms.

(iii) Fix a configuration whose violation set generates Λ_0. Then the left side of [2, (2.9)] equals 1. The right side counts the minimal admissible subfamilies of the violation set. There is at least one such subfamily, and its summand equals 1. Two violations within distance $r(\varepsilon)$ of the same large block can give two minimal subfamilies, so equality fails in general. □

Lemma F.24 (elementary geometry)**.** *Let $\mathfrak{t}$ be admissible for $\Lambda_0^{(k)}$, and put $I_k := \{z : \operatorname{dist}(z, T^{(k)} \setminus \Lambda_7^{(k-1)\prime}) \geq b\}$, with $I_0 := T^{(0)}$.*

(E1) *Every point of every element of $\mathfrak{t}$ lies in X_k.*

(E2) *$\Lambda_i^{(k)c} \subset \mathcal{N}_{i(r_k+b)}(X_k)$ for $i \leq 7$, and $X_k \subset \Lambda_5^{(k)c} \subset \Lambda_7^{(k)c}$.*

(E3) *Let $z \in X_k \cap I_k$, and let β be the large block containing z. Then $\beta \subset \Lambda_7^{(k-1)\prime}$ and $\operatorname{dist}(\beta, S(\mathfrak{t})) \leq r_k$.*

(E4) *$T^{(k)} \setminus I_k = \mathcal{N}_{b-1}\big((\Lambda_7^{(k-1)c})'\big)$.*

Proof. (E1) Let y be a point of an element of $\mathfrak{t}$, and let $\beta \ni y$ be its large block. Then $\operatorname{dist}(\beta, S(\mathfrak{t})) = 0$, so β is not one of the blocks that form $\Lambda_0^{(k)}$. Large blocks partition $T^{(k)}$, hence $y \notin \Lambda_0^{(k)}$.

(E2) By [2, (2.8)], a point of $\Lambda_{i+1}^{(k)c}$ lies in a block at distance $\leq r_k$ from $\Lambda_i^{(k)c}$. The block has diameter $b-1$, so the point lies within $r_k + b$ of $\Lambda_i^{(k)c}$. Induct on i.

(E3) β has diameter $b-1 < b$ and contains $z \in I_k$, so $\beta \subset \Lambda_7^{(k-1)\prime}$. Since $\beta \not\subset \Lambda_0^{(k)}$, the definition of $\mathfrak{L}_k$ gives $\operatorname{dist}(\beta, S(\mathfrak{t})) \leq r_k$.

(E4) On unions of L-blocks the operation $'$ commutes with complements, so $T^{(k)} \setminus \Lambda_7^{(k-1)\prime} = (\Lambda_7^{(k-1)c})'$. □

Lemma F.25 (cube count)**.** *Put $a_k := \lceil (r_k + b)/b \rceil$. In each coordinate direction, cut the cycle of large blocks into consecutive arcs of between a_k and $2a_k - 1$ blocks; if there are fewer than a_k blocks, take a single arc. Let $\mathcal{Q}_k$ be the product partition, and put $\mathcal{C}_k := \{Q \in \mathcal{Q}_k : Q \cap X_k \cap I_k \neq \emptyset\}$. Then every box of $\mathcal{Q}_k$ has sides $\leq 2r_k + 3b$, and for every family $\mathfrak{t}$ admissible for $\Lambda_0^{(k)}$*

$$|\mathcal{C}_k| \leq 3^d |\mathfrak{t}|.$$

Proof. The side bound is $(2a_k - 1)b \leq 2r_k + 3b$.

Suppose $\operatorname{dist}(z, x) \leq r_k + b$. In each coordinate the cyclic distance between z and x is then at most the minimal arc length $a_k b$, so the boxes $Q(z)$ and $Q(x)$ are equal or adjacent. Each box has at most 3^d boxes equal or adjacent to it.

Let $Q \in \mathcal{C}_k$ and $z \in Q \cap X_k \cap I_k$. By (E3) there are $z' \in \beta(z) \subset Q$ and a point y of an element $e' \in \mathfrak{t}$ with $\operatorname{dist}(z', y) \leq r_k$. Charge e' to one of its points $x_{e'}$. Then $\operatorname{dist}(y, x_{e'}) \leq L - 1$, hence $\operatorname{dist}(z', x_{e'}) \leq r_k + b$, and Q is equal or adjacent to $Q(x_{e'})$. □

The literal family of [2, (3.43)–(3.45)] uses $\Lambda_7^{(k-1)\prime}$ in place of I_k, and it can violate (3.45). Take $\mathfrak{t} = \emptyset$ and $\partial\Lambda_7^{(k-1)\prime}$ not aligned with the scale-k large blocks: the literal family is non-empty, while $3^d|\mathfrak{t}| = 0$. This is why I_k is used.

Lemma F.26 (cube families and the collar volume)**.** *Put $A_1 := 14 + 20\vartheta_M$, $A_2 := 16 + 23\vartheta_M$, $S_\varrho := \sum_{m \geq 0} L^{-m}(1 + m \log L)^\varrho$ $(= 7.6553\ldots)$, $C_D^0 := A_2(S_\varrho - 1)$ and $C_D := C_D^0 + 14(1 + \vartheta_M)$. At $\vartheta_M = \frac{1}{7}$, $C_D = 144.35\ldots$. For every admissible $\mathbf{\Lambda}$, every $k \leq K - 1$ and every $i \leq 7$,*

$$|\Lambda_i^{(k)c}|_k \leq (C_D r_k)^d \textstyle\sum_{j \leq k} |\mathcal{C}_j|.$$

In particular $|\mathrm{Col}_k|_k$ *and* $|\mathrm{Sh}_k|_k$ *obey the same bound.*

Proof. Construction. Put $\mathcal{E}_0 := \mathcal{C}_0$. For a box Q of $T^{(k-1)}$, let $\Phi_k(Q)$ be the image under $'$ of the union of the L-blocks that meet $\mathcal{N}_{w_{k-1}}(Q)$, where $w_{k-1} := 7(r_{k-1}+b) + Lb$. Put $\mathcal{E}_k := \{\Phi_k(Q) : Q \in \mathcal{E}_{k-1}\} \cup \mathcal{C}_k$. Then $|\mathcal{E}_k| \le \sum_{j\le k} |\mathcal{C}_j|$.

Sides. Let s_j be the side of a box at the scale j where it is born. Then $s_j \le (2+3\vartheta_M) r_j$. One step gives $s_{i+1} \le (s_i + 2w_i + 2L - 2)/L$, and $2w_i + 2L - 2 \le 14 r_i + 20 b \le A_1 r_i$. Since $\ell_i = \ell_k + (k-i)\log L \le (1 + (k-i)\log L)\ell_k$, iteration gives

$$s_k \le A_2 \textstyle\sum_{i=j}^{k-1} L^{-(k-i)} r_i \le A_2 r_k \sum_{m\ge 1} L^{-m}(1 + m\log L)^{\varrho} = C_D^0 r_k.$$

Covering. We show $X_k \subset \bigcup \mathcal{E}_k$ by induction on k. Points of $X_k \cap I_k$ lie in $\bigcup \mathcal{C}_k$. Let $z \in X_k \setminus I_k$. By (E4) there is $z_1 \in (\Lambda_7^{(k-1)c})'$ with $\mathrm{dist}(z, z_1) \le b - 1$. By (E2) at scale $k-1$ and the induction hypothesis,

$$B(z_1) \subset \mathcal{N}_{7(r_{k-1}+b)}(X_{k-1}) \subset \textstyle\bigcup_{Q\in\mathcal{E}_{k-1}} \mathcal{N}_{7(r_{k-1}+b)}(Q).$$

At scale $k-1$, $B(z)$ lies within $Lb - 1$ of $B(z_1)$. Hence $z \in \Phi_k(Q)$ for some $Q \in \mathcal{E}_{k-1}$.

Count. By (E2), $\Lambda_i^{(k)c} \subset \bigcup_{Q\in\mathcal{E}_k} \mathcal{N}_{7(r_k+b)}(Q)$. Each of these neighbourhoods has at most $(C_D^0 r_k + 14(r_k+b))^d \le (C_D r_k)^d$ points. Finally, Col_k and Sh_k are contained in $\Lambda_7^{(k)c}$. □

Proposition F.27 (vector form on the kept variables)**.** *Let* $\Psi = (A_0|_{S_0}, A_k|_{S_k})_{1\le k\le K}$ *with* $S_0 := \Lambda_5^{(0)c}$ *and* $S_k := \mathrm{Sh}_k$. *Let* Q_k *be the* k*-fold block average,* $a_k \in [a_\flat, a]$ *with* $a_\flat = a(1 - L^{-2})$, *and*

$$\langle \Psi,\, \Delta_v \Psi\rangle := \min_{A_0|_{\Lambda_5^{(0)}}} \Big[\langle A_0,\, (-\Delta + \mu_0^2) A_0\rangle + \sum_{k\ge 1} a_k t_k^{d-2} \sum_{x\in S_k} |A_k(x) - (Q_k A_0)(x)|^2 \Big].$$

If $\mu_0 \varepsilon_0 \le 1$, *then*

$$\langle \Psi,\, \Delta_v \Psi\rangle \ge \gamma_0 \sum_{k=0}^{K} \Big[\sum_{\langle x,x'\rangle \subset S_k} t_k^{d-2} |A_k(x') - A_k(x)|^2 + \sum_{x\in S_k} t_k^d \mu_0^2 |A_k(x)|^2 \Big],$$

with $\gamma_0^{-1} := \max\{1 + 6dC_P, (6d+2)/a_\flat, 3\}$. *Here* $C_P \le \frac{3}{4}$ *is the two-block Poincaré constant:* $n^{d-2} |\mathrm{avg}_{B_1} f - \mathrm{avg}_{B_2} f|^2 \le C_P \sum_{b'\subset B_1\cup B_2} |\nabla_{b'} f|^2$ *for adjacent blocks* B_1, B_2 *of side* n. *At* $d = 3$, $a = 1$ *this gives* $\gamma_0 = 3/80$.

Proof. Fix A_0 and write $\bar A_k := Q_k A_0$. (a) For $k = 0$, the bond terms and the mass terms on S_0 are terms of $\langle A_0, (-\Delta + \mu_0^2) A_0\rangle$. (b) For $k \ge 1$ and a bond $\langle x, x'\rangle \subset S_k$,

$$|A_k(x') - A_k(x)|^2 \le 3|A_k(x') - \bar A_k(x')|^2 + 3|A_k(x) - \bar A_k(x)|^2 + 3|\bar A_k(x') - \bar A_k(x)|^2.$$

Apply the Poincaré inequality with $n = L^k$ and multiply by ε^{d-2}. The last term then contributes at most $C_P \sum \varepsilon^{d-2} |\nabla A_0|^2$ over the double block. Each fine bond lies in at most $2d$ double blocks per level. The fine regions $B^k(S_k)$ are disjoint in k, because $S_{k'} \subset \Lambda_5^{(k)}$ for $k' > k$. So the Poincaré terms sum to at most $6dC_P \langle A_0, -\Delta A_0\rangle$. Each $A_k(x)$ lies in at most $2d$ shell bonds, so the self terms sum to at most $(6d/a_k)\, a_k t_k^{d-2} |A_k - \bar A_k|^2$. (c) By Jensen's inequality, $t_k^d \mu_0^2 |A_k|^2 \le 2\mu_0^2 t_k^2 \cdot t_k^{d-2} |A_k - \bar A_k|^2 + 2\mu_0^2 \sum_{z\in B^k(x)} \varepsilon^d |A_0(z)|^2$, and $2\mu_0^2 t_k^2 \le 2$. Adding (a)–(c), the right side with $\gamma_0 = 1$ is at most γ_0^{-1} times the bracket. Finally minimise over $A_0|_{\Lambda_5^{(0)}}$. □

Corollary F.28 (weighted vector gains)**.** *Let* S^1 *be the quadratic form of the photon chain of Lemma F.5 with all weights equal to* 1. *By Statements K.46 and K.47,* $\Delta_v = \mathrm{Sch}_\Psi S^1$. *Let* HA_{del} *be the photon form of Lemma F.14(d), and put* $\Delta_v^w := \mathrm{Sch}_\Psi HA_{\mathrm{del}}$. *If* $HA_{\mathrm{del}} \ge (1 - \hat\xi_{\mathrm{tot}}) w_{\mathrm{lo}} S^1$, *then* $\Delta_v^w \ge (1 - \hat\xi_{\mathrm{tot}}) w_{\mathrm{lo}} \Delta_v$.

Proof. $\langle \Psi, \mathrm{Sch}_\Psi(H)\Psi\rangle = \min_\chi \langle (\Psi,\chi),\, H(\Psi,\chi)\rangle$ is monotone and positively homogeneous in H. □

Proposition F.29 (region sum)**.** *For a family* $\mathfrak{t}$ *at scale* k*, put*

$$W_k^w(\mathfrak{t}) := \exp\Big(-p_k^2\big[\vartheta(\tfrac{a}{8}|P_v|+\tfrac{\gamma_0}{4}|Q_v|+\tfrac{\gamma_0}{4}|R_v|)+\tfrac{a}{8}|P_s|+\tfrac{\gamma_0}{4}|Q_s|+|R_s|\big]\Big), \qquad \zeta''_w(\Lambda_0^{(k)}) := \sum_{\mathfrak{t}\ adm.\ min.} W_k^w(\mathfrak{t}),$$

with $0<\vartheta\le 1$ *and* $\gamma_0\le\frac13$*. Put* $c_0'^w := \vartheta\min\{\frac{a}{8},\frac{\gamma_0}{4}\}$ *and* $c_L := \sup_k c_C^{(k)}t_k^{\kappa_0}+c_B^w+C_{\mathrm{old}}\varepsilon_0$*, and assume*

$$\tfrac12 c_0'^w p(t)^2 \ge 3^d c_L C_D^d R^d \ell(t)^{1+\varrho d}/\log 2 \qquad (t\le\varepsilon_0). \tag{F.4}$$

At $d=3$*,* $\varrho=2$ *and* $p(t)=b_0\ell(t)^6$*, condition* (F.4) *reads* $\hat\ell^5 \ge 54c_Lc_D^3R^3/(c_0'^w b_0^2\log 2)$*. Then, uniformly in* $\varepsilon=2^{-K}\varepsilon_0$*,* ℓ_T *and* K*,*

$$\sum_{\mathbf{\Lambda}}\prod_{k<K}\zeta''_w(\Lambda_0^{(k)})\exp\Big(\sum_{k=0}^{K}\big[c_C^{(k)}t_k^{\kappa_0}|\mathrm{Col}_k|_k+c_B^w|\mathrm{Sh}_k|_k\big]+\sum_{k<K}C_{\mathrm{old}}t_k|\Lambda_7^{(k)c}|_k\Big)\le \mathrm{e}^{C_\Sigma|T|},$$

$$C_\Sigma := c_L\varepsilon_0^{-3}+n_d\sum_{m\ge1}(2^{-m}\varepsilon_0)^{-3}\exp\big(-\tfrac12 c_0'^w b_0^2(\hat\ell+m\log 2)^{12}\big).$$

Proof. Since $\vartheta\le1$ and $\min\{\frac{a}{8},\frac{\gamma_0}{4}\}\le1$, each of the six coefficients in W_k^w is at least $c_0'^w$, so $W_k^w(\mathfrak{t})\le \mathrm{e}^{-c_0'^w p_k^2|\mathfrak{t}|}$. The terms with $k=K$ are at most $c_L|T^{(K)}|=c_L\varepsilon_0^{-3}|T|$. For $k<K$, the sets Col_k, Sh_k and $\Lambda_7^{(k)c}$ all lie in $\Lambda_7^{(k)c}$, and $C_{\mathrm{old}}t_k\le C_{\mathrm{old}}\varepsilon_0$.

Fix a tuple $\mathfrak{t}$ that generates $\mathbf{\Lambda}(\mathfrak{t})$. The radii r_k decrease in k, and $K-j=\log_2(\varepsilon_0/t_j)\le\ell_j/\log 2$. So Lemmas F.26 and F.25 give

$$\sum_{k<K}c_L|\Lambda_7^{(k)c}|_k\le 3^dc_LC_D^d\sum_j|\mathfrak{t}_j|\sum_{k=j}^{K-1}r_k^d\le 3^dc_LC_D^dR^d\sum_j|\mathfrak{t}_j|\ell_j^{1+\varrho d}/\log 2.$$

By Lemma F.23(ii) and (F.4), the left side is at most $\mathrm{e}^{c_L\varepsilon_0^{-3}|T|}\prod_{j<K}\sum_{\mathfrak{t}_j}\mathrm{e}^{-\frac12 c_0'^w p_j^2|\mathfrak{t}_j|}$. Charge each element to one point (Lemma F.24(E1)). Each point carries at most n_d candidate elements, so $\sum_{\mathfrak{t}_j}\mathrm{e}^{-x|\mathfrak{t}_j|}\le(1+\mathrm{e}^{-x})^{n_dN_j}\le\exp(n_dN_j\mathrm{e}^{-x})$. Finally insert $N_j=|T|(2^{-m}\varepsilon_0)^{-3}$ and $\ell_j=\hat\ell+m\log 2$, with $m=K-j\ge1$. □

Proof of Lemma F.22. (a) The partition of unity [2, (2.4)–(2.6)] is an identity for every choice of thresholds. The integrand is ≥0, so dropping restrictions gives an upper bound. Then apply Lemma F.23.

(b) Use [2, Prop. 2.1, (2.57), p. 570] relative to [3, Prop. 1], as in Lemma F.8(C1). The factor σ_+ comes from the replacements of Lemma F.14(d). The factor 2^{N_ℓ} comes from the D-coloured lines, with $c_k\le\frac13$. At $k=0$, the bound $\lambda\varepsilon x^2+\frac12\delta\varepsilon^2x\ge-\delta^2\varepsilon^3/(16\lambda)$ with $\varepsilon|\delta|\le C_\delta(M)$ gives the per-site cost. Since $\varepsilon\le\varepsilon^{\kappa_0}$, this is at most $c_C^{(0)}t_0^{\kappa_0}$. Under the convention that the vacuum counterterms are global c-numbers [3, (1.30)], no vacuum constant is localised on a collar (Lemma F.32(V)). The outside pieces are bounded by Lemma F.32(V), Proposition E.62(ii) and Lemma E.63.

(c) Use the first term of [2, (2.57)], as in [2, (2.13)].

(d) *Scalar entries.* These come from [2, (3.22)–(3.39)] at fixed external $\tilde A$, and they do not involve the weights. For $\tilde A\neq0$, [2, Prop. 3.1] is supplied by [4, Prop. 3.1′, (1.22), p. 574]. The hypothesis "$m^2\le O(1)$" is read as (E.m2) (Proposition E.159).

Vector entries. Lemma F.14(d) gives the hypothesis of Corollary F.28. Combine that corollary with Proposition F.27 (where $\mu_0\varepsilon_0\le1$, Proposition E.159) and with the split $\frac14+\frac14$ of [2, (3.30), (3.40)], in which the P_v term and the bound of the type of [2, Prop. 3.1] share one quarter. A violated gradient restriction on Q_v gives $\frac14(1-\hat\xi_{\mathrm{tot}})w_{\mathrm{lo}}\gamma_0\cdot\frac12p_k^2=\vartheta\frac{\gamma_0}{4}p_k^2$ per element. The same holds on R_v, since $\mu_0^2t_k^2|A|^2>p_k^2$ there. The P_v term [2, (3.33)] gives $\frac{1}{16}(1-\hat\xi_{\mathrm{tot}})w_{\mathrm{lo}}aL^{d-2}p_k^2\ge\vartheta\frac{a}{8}p_k^2$.

Arithmetic. We have $\hat{\xi}_{\rm tot} \le \frac{1}{12}$ and $\bar{\delta} = \kappa\mu_*^2\varepsilon_0^2 \le \frac{1}{12}$, the latter by (E.$\delta$). At $(1, m_2)$, $w_{\rm lo} \ge (1+\bar{\delta})^{-1} \ge \frac{12}{13}$; at $(-\eta, m_1)$, $w_{\rm lo} = 1$. Hence $\vartheta_+ \ge \frac{1}{2}\cdot\frac{11}{12}\cdot\frac{12}{13} = \frac{11}{26}$ and $\vartheta_- \ge \frac{11}{24}$.

Kept variables. The Gaussian cost of each kept vector variable is at most $3^{1/2}$. This gives the term $\frac{d}{2}\log 3\cdot c_{\rm sh}$ in c_B^w.

(e) Proposition F.29, or Lemma E.95.

Uniformity. None of the constants depends on ε, k, $\mathbf{\Lambda}$ or $\ell_T \ge \ell_0$. By Statement K.1, $c_C^{u=0}$ and c_B are bounded uniformly in $\varepsilon_0 \in (0, \varepsilon_1]$. Hence so are the constants $c_5 = c_B^w$ and $c_7 = \max\{\sigma_+ 2^{N_\ell} c_C^{u=0}, C_\delta(M)^2/(16\lambda)\}$ of condition (E.ent), and so is c_L. The thresholds on ε_0 are imposed at the binding point, so one ε_0 serves both points. □

The region sum is Lemmas E.94 and E.95. The constants c_A, c_*, C_V are evaluated at ε_1, and (E.comb) is then one more downward-closed upper bound on ε_0; in it c_A is replaced by $c_A + C_{\rm old}$ (Lemma E.63).

Remark F.30 (what c_A dominates; other routes). (i) In the closing step, (H-LF) holds with $c_*(\pi) = c_0'^w$ and $C_0(\pi) = c_{\rm fin}\varepsilon_0^{-3}$; the latter collects the final-scale kept variables at $O(1)$ each. The constant $c_A(\pi)$ dominates every per-site cost on a subset of $\Lambda_7^{(k)c}$: the collar costs of Lemma F.22(b), the vacuum collar shares, the shells, the forward replacement errors near large fields ($\times w_{\rm hi}$), and Bałaban's $(1+\log)|\Lambda_0^{(k)c}|$. It also covers the kept vector variables, including the coarse variables over earlier large-field sets. All of these are finite at both points. (ii) The region sum can also be bounded through the cube count of [2, (3.45)], as in Proposition 5.21; such routes need the repair of Remark 5.15, because the printed count fails for large blocks straddling $\partial\Lambda_7^{(k-1)\prime}$. The route above uses neither (3.45) nor cubes.

(S4) The renormalisation flow.

Lemma F.31 (the vacuum output of a step)**.** *For any exact Gaussian staging, in particular King's photon chain (Lemma F.5) and Bałaban's scalar staging, the following holds. The field-independent output of order $\le \bar{n}$ at step k of the $T_{\bar{n}}$-truncated flow is the sum of the terms of the bare vacuum graphs of Definition E.12 (counterterm vertices included, each coefficient δm^2 kept as a number) all of whose lines have index $\le k$ and at least one of which has index k; that is, it equals $\mathcal{V}_k$. Consequently $E_1(u) = \sum_{k<K}\mathcal{V}_k(u) + E_{\rm fin}(u)$, with $\mathcal{V}_k = E_{K,k} - E_{K,k+1}$ and $E_{\rm fin}(u)$ as in* (E.9)*, and $\mathcal{V}_k(u)$ equals the step-k output computed with the free index-k covariances.*

Proof. First let the covariances of the staging not depend on the fields, as for King's photon chain. Let W_0 be minus the interaction, including $\delta m^2|\varphi|^2$, which has order ≥ 1, and let $W_{k+1} = \log \mathbb{E}_{\mu_k} \mathrm{e}^{W_k}$. Since $\log \mathbb{E}\mathrm{e}^W = \sum_m \kappa_m(W)/m!$ with the m-th cumulant m-linear and of order $\ge m$, the part of order $\le \bar{n}$ of $\log \mathbb{E}\mathrm{e}^W$ depends only on $T_{\bar{n}}W$; so the truncated flow agrees with the exact one up to order $\bar{n}$, and constants pass through later steps unchanged, $\log \mathbb{E}\mathrm{e}^{c+U} = c + \log \mathbb{E}\mathrm{e}^U$. In the exact flow, the field-independent part of W_{k+1} is the sum of the connected vacuum graphs of the bare interaction whose lines carry $\sum_{j\le k} C_{(j),f}$. Subtracting the same sum with $\sum_{j<k} C_{(j),f}$ leaves, by multilinearity, the terms with all line indices $\le k$ and at least one equal to k, that is, $\mathcal{V}_k$. The identity for $E_1(u)$ is then Lemma E.14(i). For Bałaban's scalar staging, whose step covariances depend on the gauge background, the same conclusion is Statement K.9. □

The localisation sum is Lemma E.170. Its anchor is the first vertex lying in W. When the cut kernel supplies the tree link, the rate is δ_1 in place of δ.

Lemma F.32 (mass and vacuum flow along the step)**.** *Let $(u, M) \in \{(1, m_2), (-\eta, m_1)\}$, $\varepsilon = 2^{-K}\varepsilon_0$ and $0 \le k \le K$, and put*

$$P_K(t) := 2(e^2+\lambda)t + (e^4 + e^2\lambda + \lambda^2)t^2\big(1 + \log_2(\varepsilon_0/t)\big), \qquad P_m(t) := (e^2+\lambda)t + (e^4+e^2\lambda+\lambda^2)t^2.$$

Put $\mathcal{L}_k := B^k(\Lambda_7^{(k)} \cap \Lambda_8^{(k)c})$ *(the layer) and* $\mathcal{D}_k := B^k(\Lambda_8^{(k)})$*, so that* $W_k = \mathcal{L}_k \sqcup \mathcal{D}_k$*. At both points and at every step the following hold.*

(M) $a_{K,T}(M) = M^2 + \delta m^2_{K,T}(M)$*. Here* $\delta m^2_{K,T}(M)$ *is, coefficient by coefficient in* $e^\alpha \lambda^\beta$ *with* $2 \le \alpha + 2\beta \le 4$*, the counterterm of* [3, (1.19)–(1.23)] *on* T_ε*. It has C-lines only, so it is the same number at both values of u for the same M. For every k it splits as* $\delta m^2_{\mathrm{fin}} + \delta m^2_{K,k} + \delta m^2_k$ [3, (1.26)–(1.29)]*. In this split* $\delta m^2_{\mathrm{fin}}$ *and* $\delta m^2_{K,k}$ *are c-numbers, invariant under the translations of* $t_K\mathbb{Z}^3$ *and of* $t_k\mathbb{Z}^3$ *respectively, and* δm^2_k *is a localised graph sum. There is no u-dependent mass vertex. We have* $|\delta m^2_{K,T}(M)| \le C_\delta(M)\varepsilon^{-1}$*, and per site*

$$\lambda(\varepsilon)|\varphi|^4 + \tfrac{1}{2}\delta m^2_{K,T}(M)\varepsilon^2|\varphi|^2 \ge -C_\delta(M)^2\varepsilon/(16\lambda).$$

Moreover $\sup_x |\delta m^2_{K,k}(x)| t_k^2 \le C_m P_K(t_k)$*, and the step differences obey Statement K.48(b). At* $k \ge 1$ *the mass pieces of step k leave* Col_k *together with the whole interaction, at a cost contained in* $O(t_k^{\kappa_0})|\mathrm{Col}_k|_k$*; at* $k \ge 1$ *no mass term is absorbed site by site. A connected coloured family with at least one D-line and an external leg is convergent and receives no counterterm. Per block, it is at most* $C_{U1}e^2|u|\kappa\mu_*^2 L^2 t_{k+1}^3 \sup\|h\Phi_{\mathrm{ext}}\|_{1,\alpha_0}$*.*

(V) E_0^w *is cancelled exactly for each* $\mathbf{\Lambda}$ *(Lemma F.35). Further,* $E_1(u) = E_{\mathrm{fin}}(u) + E_K(u)$ *with* $E_K(u) = E_{K,k}(u) + \sum_{j<k} \mathcal{V}_j(u)$ *and* $\mathcal{V}_j(u) := E_{K,j}(u) - E_{K,j+1}(u)$*. The terms* $E_{\mathrm{fin}}(u)$ *and* $E_{K,k}(u)$ *stand as global factors* $\mathrm{e}^{-E_{\mathrm{fin}}(u) - E_{K,k}(u)}$ *in front of the inductive bound. At the step* $k \to k+1$ *only* $\mathcal{V}_k(u)$ *is localised. This is done at scale k, by the rule of* [3, p. 420] *applied to whole renormalised classes with one external field each. The anchor of a class is its first leg, which is the same in every member. A localisation* Λ *is* good *if* $\Lambda \subset W_k$*,* bad *if it meets both* W_k *and* W_k^c*, and* outside *if* $\Lambda \cap W_k = \emptyset$*. Let* A_k*,* $\bar{N}$*,* $\bar{S}$*,* δ *be as in Proposition E.62. Then:*

(V1) *The sum of* $\mathcal{V}_k(\Lambda)$ *over the good* Λ *equals the field-independent output of order* $\le \bar{n}$ *of step k, computed with the free index-k covariances and localised by the same rule. Hence the field-independent part of* $\mathcal{P}_w^{(k)}$ *is a remainder* $\mathcal{R}_k^{\mathrm{vac}}$*, and* $\mathcal{R}_{k+1}^{\mathrm{vac}} - \mathcal{R}_k^{\mathrm{vac}}$ *consists of the bad localisations of* $\mathcal{V}_k$ *and of domain differences. Moreover*

$$|\mathcal{R}_{k+1}^{\mathrm{vac}} - \mathcal{R}_k^{\mathrm{vac}}| \le \bar{N}\bar{S}A_k\big[|\mathcal{L}_k|_k + \mathrm{e}^{3\delta}\mathrm{e}^{-\delta r(t_k)/2}|\mathcal{D}_k|_k\big] + C_\kappa t_k^{\kappa_B} N_k, \qquad \kappa_B = \tfrac{16}{5}.$$

(V2) $|\sum_{\Lambda\ \mathrm{outside}} \mathcal{V}_k(\Lambda)| \le \bar{S}A_k|\Lambda_7^{(k)c}|_k \le C_{\mathrm{old}} t_k |\Lambda_7^{(k)c}|_k$*. This c-number is not part of* $\mathcal{P}_w$*. It is booked as* $\mathfrak{O}_{k+1} := \mathfrak{O}_k + \bar{S}A_k|\Lambda_7^{(k)c}|_k$*, with* $\mathfrak{O}_0 := 0$*.*

(V3) *No vacuum c-number is removed on a collar, no term of* $E_K(u)$ *is moved twice, and* $E_{\mathrm{fin}}(u)$ *is not part of* $\mathcal{P}_w^{(k)}$*. The layer terms of (V1) are at most* $C(\kappa_0)t_k^{\kappa_0}|\mathrm{Col}_k|_k$*. Summed over k, the deep terms and the domain terms are at most* $B|T|$*, with B independent of* ε*, K and* ℓ_T*.*

(C11) *Let* $F_{11}(u) = \frac{u}{2}\sum_{i<K} \delta_i \langle H_i, \Pi_2\rangle$ *be the order-*e^2 *vacuum family with one D-line (the bubble and the seagull closed by one D-line). Split* $F_{11} = \sum_{k<K} F_{11}^{[k]} + F_{11}^{\mathrm{fin}}$ *by the top index of the lines, where* F_{11}^{fin} *collects the terms with a regular scalar line. Then* $F_{11}^{[k]}$ *is the order-*e^2*, one-D-line part of* $\mathcal{V}_k(u)$*, and* F_{11}^{fin} *is part of* $E_{\mathrm{fin}}(u)$*. Moreover:*

(a) *for every localisation* $\Lambda \subset W_k$*, the generated free vacuum output and the moved piece agree:* $\mathrm{Gen}_k^{\mathrm{free}}(\Lambda) = F_{11}^{[k]}(\Lambda)$*;*

(b) *the localisations that straddle* ∂W_k *satisfy*

$$\Big|\sum_{\Lambda\ \mathrm{bad}} F_{11}^{[k]}(\Lambda)\Big| \le 3S_3 C_{\mathrm{pl}} e^2|u|\kappa\mu_*^2 L^3 t_k^3\Big[|\mathcal{L}_k|_k + \mathrm{e}^{\delta_1(c_d+1)/2}\mathrm{e}^{-\delta_1 r(t_k)/2}|\mathcal{D}_k|_k\Big],$$

where $S_3 := \sum_{D\geq 0}(c''_d(D+2)^d)^3 \mathrm{e}^{-\delta_1 D/2}$ *and* C_{pl} *is the per-localisation constant of Theorem H.24(c) for this family. The layer part is at most* $C(\kappa_0)t_k^{\kappa_0}|\mathrm{Col}_k|_k$*. The deep part, summed over* k*, is at most* $C|T|$*, uniformly in* K*;*

(c) *the outside localisations of the* u*-part of* $\mathcal{V}_k$*, in particular those of* $F_{11}^{[k]}$*, are bounded per block of* $\Lambda_7^{(k)c}$ *by* $\bar{S}\bar{N}_{\mathrm{ph}}|u|\bar{c}(1+\bar{c})^{\bar{N}_{\mathrm{ph}}-1}C_P P(t_k)$*, with the notation of Proposition E.62(iii). They are booked in* $\mathfrak{O}_{k+1}$ *and carry no factor* $1+K-k$*.*

The global sum $F_{11}(u)$ *is of size* $O(K)|T|$ *and is never bounded. It occurs only inside* $E_1(u)$*, both in front of the inductive bound and in* $T_{\bar{n}}[\Phi_M(u)]$*, and as the sum of the generated constants; these two occurrences are equal.*

The constants $C_\delta(M)$*,* C_m*,* C_{U1}*,* C_κ*,* C_{old}*,* C_{pl} *and* $C(\kappa_0)$ *do not depend on* ε*,* k*,* x*,* $\mathbf{\Lambda}$ *or* u*. They depend on* ℓ_T *only through terms that do not increase with* ℓ_T*, evaluated at* ℓ_0*, and they are finite at* $M = m_1$*.*

The cancellation of the good localisations in (C11) is part of Proposition E.64, which is proved under the convention that the vacuum counterterms are global c-numbers [3, (1.30)].

Lemma F.33 (geometry of the layer)**.** *Every block of* $\mathcal{L}_k$ *lies within distance* $r(t_k)+\sqrt{d}\,M_b$ *of a block of* Col_k*, and*

$$|\mathcal{L}_k|_k \leq \big(2r(t_k)+2\sqrt{d}\,M_b+1\big)^d\,|\mathrm{Col}_k|_k.$$

Proof. Let β be a large block of $\Lambda_7^{(k)}\cap\Lambda_8^{(k)c}$. By [2, (2.8)] with $i=7$, β lies within $r(t_k)$ of a point of $\Lambda_7^{(k)c}$. Let $\square$ be the unit block containing that point. If $\square\subset\Lambda_0^{(k)c}$, then β lies within $r(t_k)$ of $\Lambda_0^{(k)c}$. By (2.8) with $i=0$ this gives $\beta\subset\Lambda_1^{(k)c}\subset\Lambda_7^{(k)c}$, which contradicts $\beta\subset\Lambda_7^{(k)}$. Therefore $\square\subset\Lambda_0^{(k)}\cap\Lambda_7^{(k)c}\subset\Lambda_7^{(k-1)\prime}\cap\Lambda_7^{(k)c}=\mathrm{Col}_k$. Now count the unit blocks within $r(t_k)+\sqrt{d}M_b$ of each such $\square$. □

Proof of Lemma F.32. (M). The identity $a_{K,T}(M) = M^2+\delta m^2_{K,T}(M)$ is Proposition 4.7. The coefficients are defined on T_ε by $-\delta m^2_{(\alpha,\beta)}+\sum_{x\in T_\varepsilon}\varepsilon^d\Sigma^\varepsilon_{(\alpha,\beta)}(x)=0$ [3, p. 417]. The self-energy graphs entering them carry C-lines only, so u does not enter.

The split is the definition [3, (1.25)–(1.28), p. 418]. On T_ε the propagators G^ε_K and $G^\varepsilon_K(0)$ commute with the translations of $t_K\mathbb{Z}^3$, and $G^\varepsilon_{K,k}$ commutes with those of $t_k\mathbb{Z}^3$. The coefficients inherit this periodicity. The localisation of δm^2_k is that of [3, p. 420].

The bound $|\delta m^2_{K,T}(M)|\leq C_\delta(M)\varepsilon^{-1}$ is the size statement $\delta m^2 = O(\varepsilon^{-1})$ of [2, p. 559]. Its explicit constant $C_\delta(M)$ is that of Statement K.49, with the majorant constants of its hypothesis (H) evaluated at $\ell_T=\ell_0$; they do not increase with ℓ_T. Put $s=|\varphi|^2\geq 0$ and $\delta=\delta m^2_{K,T}(M)$. Then $\lambda\varepsilon s^2+\frac{1}{2}\delta\varepsilon^2 s\geq -\delta^2\varepsilon^3/(16\lambda)\geq -C_\delta(M)^2\varepsilon/(16\lambda)$.

The pointwise bound on $\delta m^2_{K,k}$ and the bounds on the step differences are Statement K.48.

The removal from Col_k is [2, Prop. 2.1, (2.57), p. 570]. Under the restrictions (2.55), the interaction is removed from $B^k(\mathrm{Col}_k)$, leaving only $-\lambda(t_k)\sum\eta^d|\phi^{(k)}|^4$, at the cost $O(t_k^{\kappa_0})|\mathrm{Col}_k|$. With $\kappa_0=\frac{1}{5}$ this is Lemma F.1.

The statement on coloured families is Statement K.50(b),(c) at scale $n=k+1$. Its part (c) also bounds the share anchored in a scale-k sub-block. The transfer of the line bounds to the D-parts is Statement K.50(d), whose hypothesis is Statement K.51.

(V1), the identity. By [2, p. 570], the whole interaction is removed from Col_k before step k. So every vertex of a family generated at step k lies in W_k, and the generated output has only localisations $\Lambda\subset W_k$. By Lemma F.31, that output, computed with the free index-k covariances, is $\mathcal{V}_k(u)$ as a truncated power series. In fact this is an identity of the same index-assigned Wick graphs: a contraction with at least one index-k line and all other lines of index $<k$ is exactly an assignment of top index k. A generated graph and the corresponding counterterm graph have

the same vertex set. The anchor is the first leg of the whole class, the same in every member, so the rule of [3, p. 420] gives both graphs the same Λ. The identity therefore holds localisation by localisation for $\Lambda \subset W_k$. The bad and outside localisations of $\mathcal{V}_k$ have no generated partner.

(V1), the bounds. By [2, (2.8)], W_k lies $2r(t_k)$ inside $\Lambda_5^{(k)}$. Replace the restricted lines of step k by the free ones, and g_{k+1} by 1. This costs $O((L^j\varepsilon)^\kappa)|B^{k-j}(\Lambda_2^{(k)})|$ for every $\kappa > d$ [2, p. 578], since $g_{k+1} = 1$ at distance $\geq M_b$ from $\partial\Omega_1$ [3, p. 412]. Take $\kappa = \kappa_B$. The bad localisations are bounded by Proposition E.62(i). Every vacuum graph has $o = \alpha/2 + \beta \geq 1$. So, for $|u| \leq 1$, $A_k \leq C_P(1+\bar{c})^{\bar{N}_{\mathrm{ph}}} C_e t_k$ with the constants of Proposition E.62, and $\bar{S}A_k \leq C_{\mathrm{old}} t_k$.

(V2) is Proposition E.62(ii) with $X = W_k^c = \Lambda_7^{(k)c}$, counted in scale-$k$ blocks.

(V3). By Lemma F.33, and since $r(t) = R(1+\log t^{-1})^2$ and $\kappa_0 < 1$,

$$A_k|\mathcal{L}_k|_k \leq C t_k \big(2r(t_k) + 2\sqrt{d}M_b + 1\big)^d |\mathrm{Col}_k|_k \leq C(\kappa_0) t_k^{\kappa_0} |\mathrm{Col}_k|_k.$$

Next, $A_k \mathrm{e}^{-\delta r(t_k)/2} N_k \leq C|T| t_k^{-2} \mathrm{e}^{-\delta r(t_k)/2}$, and $\sum_k t_k^{-2}\mathrm{e}^{-\delta r(t_k)/2}$ converges uniformly in K. Finally $\sum_k t_k^{\kappa_B} N_k = |T| \sum_k t_k^{1/5} \leq (1-2^{-1/5})^{-1} \varepsilon_0^{1/5} |T|$. The outside costs are paid in the region sum by Lemma E.63. Each term of $E_K(u) = E_{K,k}(u) + \sum_{j<k} \mathcal{V}_j(u)$ leaves the global factor exactly once, at its top index, and $E_{\mathrm{fin}}(u)$ is never moved.

(C11). The split is multilinearity in $C_0 = \sum_{j<K} S_j + R_s$ and in $D = \sum_{i<K} \delta_i H_i$; D has no regular remainder. The set of index assignments with all indices $\leq k$ and at least one equal to k is the set moved at $k \to k+1$ [3, p. 422]. Hence $F_{11}^{[k]}$ is the order-e^2, one-D-line part of $\mathcal{V}_k(u)$.

(a) This is the identity of (V1) restricted to this family, since the generated family and $F_{11}^{[k]}$ consist of the same index-assigned graphs with the same anchors. At order e^2 no truncation enters.

(b) After the cut at top index k, each localisation is at most $C_{\mathrm{pl}} e^2 |u| \kappa \mu_*^2 L^3 t_k^3 \mathrm{e}^{-\delta_1 d(\Lambda)}$. This is the size $e(t_{k+1})^2|u|\delta_k$ with $\delta_k = \kappa\mu_*^2 t_{k+1}^2$. If only the D-line has index k, the remainder is a C-only class with its Ward partner. If a scalar line has index k, the remainder is convergent. In the case of ties, the cut kernels supply the tree link at rate δ_1. There are $N_s \leq 3$ localised vertices. Apply Lemma E.170 with $A = C_{\mathrm{pl}} e^2 |u| \kappa \mu_*^2 L^3 t_k^3$, $\delta = \delta_1$ and $r = r(t_k)$. By [2, (2.8)], the blocks within $r(t_k)$ of W_k^c lie in $\mathcal{L}_k$, and the others lie in $\mathcal{D}_k$. For the layer part, Lemma F.33 gives $t_k^3(2r(t_k) + 2\sqrt{d}M_b + 1)^d \leq C(\kappa_0) t_k^{\kappa_0}$. The deep part sums to $C|T| \sum_k \mathrm{e}^{-\delta_1 r(t_k)/2}$, which is finite uniformly in K.

(c) This is Proposition E.62(iii) with $X = \Lambda_7^{(k)c}$. Only top index k is localised at step k, so there is no factor $1 + K - k$.

For the final statement, group $E_1(u)$ by largest index and apply Lemma F.31: the two occurrences of $\sum_k F_{11}^{[k]}$ coincide, so no bound on $F_{11}(u)$ is used. □

Remark F.34 (collar shares under a localised convention). Suppose instead that the parts of $E_{K,j}$ not yet moved were removed on the collars. Weight each class by the volume fraction of its scale-j^* anchor block that lies in the removal set, where j^* is the top index. For a scale-j block □ this fraction is $(t_j/t_{j^*})^3$. Each weighted class is bounded per anchor block by Statement K.53. Summing the geometric series in j^* gives

$$|E_{K,j}(u)\,\restriction_{\square}| \leq C_A C_F \sigma_N(u) \sum_o c_o\, t_j^3 \sum_{j^*=j}^{K-1} t_{j^*}^{o-3} \leq C_{\kappa_0}^{\mathrm{FA}} t_j^{\kappa_0},$$

with $C_{\kappa_0}^{\mathrm{FA}} = C_A C_F \sigma_N(u) C_{\bar{n}}(e,\lambda)\, \varepsilon_0^{1-\kappa_0} \max\{1, ((1-\kappa_0)\log 2)^{-1}\}$. Under that convention the order-e^2, one-D-line part is $\leq Ce^2|u|\kappa\mu_*^2 L^2 t_j^3 (1+K-j)$. The convention of Lemma F.32(V) does not use these bounds.

(S5) The closing step and the order of choices.

Lemma F.35 (constants do not enter the remainder)**.** *If $\Phi' = \Phi + c$ with c independent of (e, λ), then $\Phi' - T_{\bar n}[\Phi'] = \Phi - T_{\bar n}[\Phi]$.*

Proof. $T_{\bar n}$ is linear and $T_{\bar n}[c] = c$. □

Lemma F.36 (the final-scale photon law)**.** *Under the law $\mathrm{e}^{-\frac12 HA}$, at both points, the final-scale field A_K has law $N(0, (\Delta^{(K)})^{-1})$, and*

$$\int dA_0 \cdots dA_{K-1}\, \mathrm{e}^{-\frac12 HA} = \prod_{k<K} (2\pi)^{n_k/2} \det(w_k C^{(k)})^{1/2}\, \mathrm{e}^{-\frac12 D_K(A_K)}.$$

Proof. By Lemma F.5, A_0 occurs only in I_0. Integrate out A_0, then A_1, and so on up to A_{K-1}. Each step is a translated Gaussian integral $\int dA_k\, \mathrm{e}^{-\frac12 w_k^{-1} I_k} = (2\pi)^{n_k/2} \det(w_k C^{(k)})^{1/2}$, which does not depend on A_{k+1}. □

Lemma F.37 (zero modes)**.** *Work in physical units, with $\beta_k = a t_{k+1}^{-2}$, $\Delta^{(0)} = -\Delta^\varepsilon + \mu_*^2$ and $\Delta^{(k+1)} = \beta_k - \beta_k^2 Q C^{(k)} Q^*$. Then for every $k \le K$, uniformly in ℓ_T and ε,*

$$\Delta^{(k)} \ge \lambda_*(k) := \Big(\mu_*^{-2} + a^{-1} \sum_{j<k} t_{j+1}^2\Big)^{-1} \ge \gamma_1 \mu_*^2, \qquad \gamma_1 := \big(1 + 4\mu_*^2 \varepsilon_0^2/(3a)\big)^{-1},$$

with equality on the constant mode. The scalar Gaussian of the closing step has mass M^2, and $\Delta_s^{(k)}(0) \ge M^2(1 + 4M^2\varepsilon_0^2/(3a))^{-1}$, with equality on constants.

Proof of the photon bound. $(\Delta^{(k+1)})^{-1}$ is the covariance of $A_{k+1} = QA_k + \xi_k$, where $\xi_k \sim N(0, \beta_k^{-1})$ is independent of $A_k \sim N(0, (\Delta^{(k)})^{-1})$. Hence $(\Delta^{(k+1)})^{-1} = Q(\Delta^{(k)})^{-1} Q^* + \beta_k^{-1} \le \|(\Delta^{(k)})^{-1}\| + \beta_k^{-1}$, using $\|Q\| \le 1$, and $\|(\Delta^{(0)})^{-1}\| = \mu_*^{-2}$. Iterating gives the first bound. Since $L = 2$, $\sum_{j<k} t_{j+1}^2 \le \frac43 \varepsilon_0^2$. On constants, $Q1 = 1$, $Q^*1 = 1$ and $\Delta^{(0)}1 = \mu_*^2 1$, so by induction $\Delta^{(k)}1 = \lambda_*(k)1$. The scalar free bound has the same proof with $\Delta_s^{(0)} = -\Delta^\varepsilon + M^2$ (Lemma A.12). □

Lemma F.38 (the normalisation defect)**.** *Assume $\delta_0 R \ge 4$ and $\hat\xi_{\mathrm{tot}} \le \frac{1}{12}$ ((E.14)). Then for every admissible $\mathbf{\Lambda}$ at both points,*

$$|\log\det HA_{\mathrm{del}}(\mathbf{\Lambda}) - \log\det HA| \le B_{F3}|T|, \qquad B_{F3} := 12|u|\kappa C_{\ell^1}(1 - \hat\xi_{\mathrm{tot}})^{-1} w_{\min}^{-2} w_{\mathrm{lo}}^{-1} \gamma_1^{-1} \varepsilon_0.$$

Here C_{ℓ^1}, δ_0 are the kernel-decay constants of $\Delta^{(j)}$ on T_ε, $w_{\min} = \min_j w_j$, $w_{\mathrm{lo}} := \min(1, \min_k w_k^{-1})$, and γ_1 is the constant of Lemma F.37. Without the condition $\delta_0 R \ge 4$, the same holds with B_{F3} multiplied by $C_* := \exp(\max\{0, 16/(\delta_0 R) - 4\})$.

The proof uses $\delta_0 R \ge 4$ in place of the printed $\delta_0 R \ge 2$.

Proof. Work in rescaled units at scale j. (i) $HA_{\mathrm{del}} - HA = -2\sum_j \omega_j \hat X_j$. Put $Y := HA^{-1/2}(HA_{\mathrm{del}} - HA)HA^{-1/2}$; then $\|Y\| \le \hat\xi_{\mathrm{tot}}$. (ii) $|\mathrm{tr}\log(I + Y)| \le (1 - \|Y\|)^{-1}\|Y\|_1$. (iii) $\|HA^{-1/2} P_j \hat X_j P_j HA^{-1/2}\|_1 \le \|P_j HA^{-1} P_j\|\, \|\hat X_j\|_1$. Since $HA \ge w_{\mathrm{lo}} S^1$ (Lemma F.5), $P_j HA^{-1} P_j \le w_{\mathrm{lo}}^{-1}(\Delta^{(j)})^{-1} \le (w_{\mathrm{lo}} \gamma_1 \mu_*^2 t_j^2)^{-1}$ by Lemma F.37. (iv) On level j, $\mathrm{dist}(\Lambda_7^{(j-1)\prime}, \Lambda_6^{(j-1)\prime c}) > r(t_{j-1})/L \ge r(t_j)/2$, so $\|\hat X_j\|_1 \le C_{\ell^1} \mathrm{e}^{-\delta_0 r(t_j)/4} N_j$; the periodic images are summed into C_{ℓ^1}. (v) $2|\omega_j| \le 6|u|\kappa\mu_*^2 t_j^2 w_{\min}^{-2}$, where $\delta_j - \delta_{j-1} = 3\kappa\mu_*^2 t_j^2$ at $L = 2$; this also holds for $j = K$. The factor $\mu_*^2 t_j^2$ cancels against (iii). (vi) With $x = \log t_j^{-1}$ and $(1+x)^2 \ge 4x$: $\mathrm{e}^{-\delta_0 R(1+x)^2/4} \le t_j^{\delta_0 R} \le t_j^4$. Hence $\sum_j \mathrm{e}^{-\delta_0 r(t_j)/4} N_j \le |T| \sum_j t_j \le 2\varepsilon_0 |T|$. Collecting (i)–(vi) gives the claim. For general R, bound $\mathrm{e}^{-\delta_0 R(1+x)^2/4} \le C_* t^4$ instead. □

Lemma F.39 (the closing step)**.** *Assume (IH_K) at both points, and the hypotheses of Lemmas E.95 and F.38. Then*

$$\mathrm{e}^{-E_0^w - E_1(u)} \mathcal{Z}_u \le \exp\big(C_+^B(u; M)|T|\big), \qquad C_+^B(u; M) = C_{F1}\varepsilon_0^{\kappa_0 - 3} + B_E^{\mathrm{fin}}(u, M) + B_{\mathfrak{E}} + B_{GS} + \tfrac12 B_{F3} + B_{\mathrm{comb}},$$

with $B_{\mathfrak{E}} = (1-2^{-1/5})^{-1}(C_\kappa + C_\nu)\varepsilon_0^{1/5}$, $B_{GS} = 2dC_4'\varepsilon_0$, B_{F3} *as in Lemma F.38, and*

$$B_{\mathrm{comb}} = c_{\mathrm{fin}}\varepsilon_0^{-3} + 2(d+2)\Sigma_*(\varepsilon_0;\pi)$$

(Lemma E.95 with $c_* = c_0'^{w}$, $2p = 12$*).*

The proof is Steps 4–5 of the proof of Proposition F.2 (C1–C5): E_0^w cancels by Lemma F.35, the zero modes are handled by Lemmas F.36 and F.37, the normalisation defect by Lemma F.38, and the region sum by Lemma E.95. The factor $\mathrm{e}^{-E_{\mathrm{fin}}(u)}$ is paid once, by Propositions E.21 and E.162.

Lemma F.40 (the order of choices)**.** *(i) In the order of the stages 0–11 of Appendix A, every constant depends only on data of earlier stages and on data of its own stage that are fixed before it. Dependences within one stage occur only in two ways, and both are one-way. First, at the stages 0 (*κ *and* $\ell_0^\natural$ *from the point), 6–8 (constants from p and* $\mathbb{W}_0$*) and 10a (suprema over* $(0,\varepsilon_1]$ *from* ε_1*). Second, at the stages 5, 10b and 11, where the conditions of the stage determine R,* ε_0 *and* ℓ_0*, respectively, and no condition of that stage depends on the quantity it determines. In particular, the upper bounds on* ε_0 *depend on data of the stages* ≤ 9 *together with stage 10a, and the lower bounds on* ℓ_0 *(stage 11) depend on data of the stages* $\le$ *10b. (ii) Fix a floor* $\ell_0^\natural > 0$ *at stage 0, and evaluate every constant that depends on* ℓ_T *and enters an upper bound on* ε_0 *at* $\ell_T = \ell_0^\natural$*. Then each value bounds the constant for all* $\ell_T \ge \ell_0^\natural$*. Hence, with* $\ell_0 := \max\{\ell_0^\natural, \ell_0^*(\varepsilon_0)\}$ *at stage 11, every upper bound on* ε_0 *decided at stage 10 holds for every torus* T_j*, and no upper bound on* ε_0 *depends on* ℓ_0*.*

Part (i) is Lemma A.4; part (ii) is Lemma A.10 for an arbitrary fixed floor.

Proof. (i) By Lemma A.4, every dependence goes from a stage to the same or a later stage, and the dependences within one stage are those listed in (i). The latter are read off the "depends on" columns of Table 3 and Tables 5–8. The upper bounds on ε_0 (Table 7) involve, besides ε_0, only data of the stages ≤ 9 and the suprema of stage 10a. The lower bounds on ℓ_0 (Table 8) involve only data of the stages $\le$ 10b. In particular, the constants c_5 and c_7 of condition (E.ent) are bounded uniformly in $\varepsilon_0 \in (0,\varepsilon_1]$ by Statement K.1, so (E.ent) depends on no datum of stage 10b other than ε_0.

(ii) This is Lemma A.10. The ℓ_T-dependent constants are the following, and all are non-increasing in ℓ_T: $C_\delta(M)$ (zero-mode majorants $64\mathrm{e}^{-\mu^2\ell^2/8}/(\mu^2\ell^3)$ and $a^{-2\alpha}b^{-2\beta}/\ell^3$); the loss in the extension of [2, Prop. 2.1] to $Q^{(k)}$ through C_δ; the terms $(M^2\ell_T^3)^{-1}$, $(\mu_*^2\ell_T^3)^{-1}$, ℓ_T^{-1}, $(M\ell_T)^{-10}$, $(\mu_*\ell_T)^{-10}$ of the bound on $\delta m_{\mathrm{fin}}^2$, whose logarithmic term is bounded by $\log_+(1/(\min(M,\mu_*)\varepsilon_0))$ for every ℓ_T; the photon zero mode and tail terms; and the twist defect $\mathrm{e}^{-\delta\ell_T/t}$. For non-increasing f, $f(\ell_T) \le f(\ell_0^\natural)$ when $\ell_T \ge \ell_0^\natural$. Every upper bound on ε_0 is monotone in each constant in the direction "larger constant $\Rightarrow$ stronger condition". So upper bounds on the constants keep the conditions sufficient, and $\ell_0 \ge \ell_0^\natural$ holds by (T.floor). For $\delta m_{\mathrm{fin}}^2$ the majorant is the explicit bound of Proposition E.9. □

Appendix G. The counterterm gaps

This appendix proves the bounds behind Theorem 6.1(a)–(c): the two-loop gap bound (Proposition G.3), the derivative bound (Proposition G.6) and the coefficient condition (Corollary G.8), with their numerical evaluations (Corollaries G.4, G.7, G.9). They rest on the exact two-loop form of King's counterterm and on slope bounds for each coupling order (Lemmas G.1 and G.2), which are proved in Subsection G.1.

Units and notation. By the scaling step of the proof of Theorem 6.1, we work in units $m = \mu_* = 1$. "Every $\varepsilon \le 1$" then means every $\varepsilon \le 1/m$, and a torus floor ℓ_0 means ℓ_0/m. The numerical constants below are computed in these units, for $\varepsilon \le 1$, and so are valid for every $\varepsilon \le 1/m$. T is a periodic torus of side $\ell_T \ge \ell_0$ with $\ell_T/\varepsilon \in 2\mathbb{N}$, and T^* is its dual lattice. For $k \in T^*$ put $\hat{k}^2 := \sum_\mu (2/\varepsilon)^2 \sin^2(\varepsilon k_\mu/2)$, $\hat{G}_M(k) := (\hat{k}^2 + M^2)^{-1}$ and $\hat{D}(k) := (\hat{k}^2 + \mu_*^2)^{-1}$, and write

$\int_T f := |T|^{-1}\sum_{k\in T^*} f(k)$. Let $G_M(x) := |T|^{-1}\sum_{k\in T^*} \mathrm{e}^{ik\cdot x}\hat G_M(k)$ be the lattice Green's function on T. Define

$$C_T(M) := \int_T \hat G_M, \quad D_T := \int_T \hat D, \quad J_T(M) := M^2\int_T \hat D\hat G_M, \quad S_T(M) := \sum_{x\in T}\varepsilon^3 G_M(x)^3.$$

Let $\widetilde\Pi_T(p;M)$ be the subtracted order-e^2 self-energy of [24, (3.32)–(3.33), p. 659] on T (photon propagator $(-\Delta_\varepsilon+\mu_*^2)^{-1}$), and put $F_T(M) := \mathrm{e}^{-2}\int_T \hat G_M^2\,\widetilde\Pi_T(\cdot;M)$. Write $Q_T(M) := \mathrm{e}^{-4}\delta m^2_{e^4,T}(\varepsilon;M)$ for the order-e^4 coefficient of the counterterm of Definition 2.2. Finally, the functions

$$\tau(x) := 1+\tfrac{12}{\pi x}+\tfrac{24}{\pi x^2}+\tfrac{64}{\pi^2x^3},$$
$$\theta(x) := \tfrac{3\pi}{16x}+\tfrac{3\pi}{8x^2}+\tfrac{1}{x^3},$$
$$A_R(x) := 768\big[\tfrac{c_N}{8\pi}+\theta(x)\big]^2,$$
$$c_N := \tfrac{\pi^3}{8}$$

are all decreasing in $x>0$.
Two lemmas.

Lemma G.1 (exact two-loop form)**.** *For every K and every torus T,*

$$a_{K,T}(M) = M^2 - 16\lambda C_T(M) - 2e^2D_T - e^2J_T(M) + 128\lambda^2S_T(M) + 16e^2\lambda F_T(M) + e^4Q_T(M),$$

with no further terms. $Q_T = \frac34\varepsilon^2D_T^2 + (A)+(C)+(D1)+(D2)+(D3)$ is the Ward form of Corollary G.22. Its first term does not depend on M. Moreover $M\mapsto a_{K,T}(M)$ is C^∞ on $(0,\infty)$.

The polynomial structure, i.e. only the monomials $e^2,\lambda,e^4,e^2\lambda,\lambda^2$, is that of Definition 2.2: King's (3.32) sums over $2\le\alpha+2\beta\le4$, and the odd orders in e vanish.

Smoothness (elementary). At fixed ε and T each term is a finite sum, over T^* or over the sites of T, of products of factors $\hat G_M(k)$ and of M-independent factors. Since $\hat k^2+M^2\ge M^2>0$, each such term is a rational function of M^2 without poles on $(0,\infty)$. Hence, for $0<M_1<M_2$, $a_{K,T}(M_2)-a_{K,T}(M_1)=\int_{M_1}^{M_2}\partial_M a_{K,T}\,dM$, and the same identity holds for each order separately.

Lemma G.2 (slope bounds for each coupling order)**.** *Let $\varepsilon\le1$, let T have side $\ell_T\ge\ell_0$, and let $M>0$. Let $\bar\tau(M)\ge\int_T\hat G_M^2$, $\bar\psi(M)\ge\int_T\hat G_M\hat D$, $\bar\delta,\bar\Theta,\bar J_s,\bar J'_s,\bar\Lambda_i$ be the one-loop majorants of Lemma G.15, all evaluated at ℓ_0; here $\bar\tau=\mathcal I_{2,0}$ is the Riemann form, not an image form, which would need $\varepsilon M\le1$. Let $\bar\omega\ge\sup\varepsilon D_T$ be the constant $w+\bar E(\mu_*,\ell_0)$, $w=0.252731$, or its sharper form with $\min\{\bar E,\tilde E\}$ (Lemma G.14(c)). All of these majorants are non-negative, independent of the couplings and of ε, and non-increasing in M and in ℓ_0. Then:*

(o-λ) $\partial_M C_T = -2M\int_T\hat G_M^2\le0$. *Moreover*

$$C_T(M_1)-C_T(M_2)\ge h_1(M_1,M_2) := (2\pi^2)^{-1}[M_2\arctan(\pi/M_2)-M_1\arctan(\pi/M_1)]\ge0.$$

(o-e^2) $\partial_M J_T = 2M\int_T\hat D\hat k^2\hat G_M^2$, *and* $0\le\partial_M J_T\le j(M) := 2M\min\{\bar\tau(M),\bar\psi(M)\}$. *Also* $\partial_M J_T\le\frac{\pi^2}{32}\tau(M\ell_0)$.

(o-λ^2) $0\le-128\,\partial_M S_T\le768M\bar\tau(M)^2\le A_R(M\ell_0)/M$.

(o-$e^2\lambda$) $|\partial_M F_T|\le\varphi_3(M) := 4M\big[2\sqrt2\,\bar\Theta\bar J_{3/2}+|\mu_*^2-2M^2|\bar\psi\bar J_0+3\sqrt3\,\bar\omega\bar J_1\big]+\bar\beta\bar\tau$, *where* $\bar\beta := 2M[2\bar\tau+8\bar\psi]$.

(o-e^4) $\partial_M Q_T\ge-b_4(M)$, *where* $b_4 := \varphi_A+28M\bar\delta\bar\tau+\varphi_{D2}$, $\varphi_A := 6M\big[2\sqrt2\,\bar\Theta\bar J'_{3/2}+|\mu_*^2-2M^2|\bar\psi\bar J'_0+3\sqrt3\,\bar\omega\bar J'_1\big]+\bar\beta\bar\psi$, *and* $\varphi_{D2} := 2M\bar\psi(2\bar\psi+4\bar\delta)+\min\{8M\bar\psi(6\bar\psi+2\bar\tau+4\bar\delta),\,2^{5/2}M(2\bar\Lambda_1+4\bar\Lambda_2)\bar\Lambda_0\}$.

The elementary parts are the following.

- From $\partial_M\hat G_M=-2M\hat G_M^2$ we get $\partial_M C_T\le0$. We also get $\partial_M J_T = 2M\int_T\hat D\hat G_M(1-M^2\hat G_M) = 2M\int_T\hat D\hat k^2\hat G_M^2$, which is ≥0.

- Since $\hat{D}\hat{k}^2 \le 1$, $\partial_M J_T \le 2M\bar{\tau}$; since $\hat{k}^2\hat{G}_M \le 1$, $\partial_M J_T \le 2M\bar{\psi}$.
- For the order λ^2: $\partial_M G_M = -2M\, G_M * G_M$, so $-\partial_M S_T = 6M \sum_x \varepsilon^3 G_M(x)^2 (G_M * G_M)(x)$. Then $(G_M * G_M)(x) \le (G_M * G_M)(0) = \int_T \hat{G}_M^2$ (the Fourier coefficients $\hat{G}_M^2$ are non-negative), and $\sum_x \varepsilon^3 G_M^2 = \int_T \hat{G}_M^2$ (Parseval). This gives $-\partial_M S_T \le 6M(\int_T \hat{G}_M^2)^2$.
- The non-negativity in (o-λ^2) uses $G_M \ge 0$ (Lemma G.12(a)).
- The last bounds in (o-e^2) and (o-λ^2) use $\int_T \hat{G}_M^2 \le (\pi^2/(64M))\tau(M\ell_T)$ and $M\bar{\tau} \le c_N/(8\pi) + \theta(M\ell_T)$ (Lemma G.11), together with the monotonicity of τ, θ and $\ell_T \ge \ell_0$.

The two-loop gap and statement (a).

Proposition G.3 (two-loop gap between anchor masses on tori)**.** *Let $0 < M_1 < M_2$, $\mu_* \in (0,1]$ and $\ell_0 > 0$. For every $K \ge 0$ (that is, $\varepsilon \le 1$) and every torus of side $\ell_T \ge \ell_0$,*

$$a_{K,T}(M_2) - a_{K,T}(M_1) \;\ge\; P_{\rm gap} := (M_2^2 - M_1^2) + 16\lambda h_1(M_1, M_2) - e^2\bar{j} - \lambda^2\bar{s} - e^2\lambda\bar{B}_3 - e^4\bar{B}_4^-,$$

where $\bar{j} := \int_{M_1}^{M_2} j$, $\bar{s} := \int_{M_1}^{M_2} 768M\bar{\tau}^2$, $\bar{B}_3 := 16\int_{M_1}^{M_2}\varphi_3$ and $\bar{B}_4^- := \int_{M_1}^{M_2} b_4$. All four are non-negative and independent of the couplings.

With Corollary G.4 this gives Theorem 6.1(a).

Proof. Fix ε and T, and take the difference of Lemma G.1 at M_2 and M_1. The terms $-2e^2 D_T$ and $\frac{3}{4}\varepsilon^2 e^4 D_T^2$ do not depend on M and cancel. Write each remaining order as $\int_{M_1}^{M_2}\partial_M$ (Lemma G.1, smoothness):

- order λ: $16\lambda[C_T(M_1) - C_T(M_2)] \ge 16\lambda h_1$, by (o-$\lambda$);
- order e^2: $-e^2[J_T(M_2) - J_T(M_1)] \ge -e^2\bar{j}$, by (o-$e^2$);
- order λ^2: $128\lambda^2[S_T(M_2) - S_T(M_1)] \ge -\lambda^2\bar{s}$, by (o-$\lambda^2$);
- order $e^2\lambda$: $16e^2\lambda[F_T(M_2) - F_T(M_1)] \ge -16e^2\lambda\int|\partial_M F_T| \ge -e^2\lambda\bar{B}_3$, by (o-$e^2\lambda$);
- order e^4: $e^4[Q_T(M_2) - Q_T(M_1)] \ge -e^4\bar{B}_4^-$, by (o-$e^4$).

Each majorant is evaluated at ℓ_0 and does not depend on $\varepsilon \le 1$ or on $\ell_T \ge \ell_0$, so the bound holds for the infimum over K and T. □

Corollary G.4 (statement (a) of Theorem 6.1)**.** *Take $\mu_* = 1$, $(M_1, M_2) = (0.7, 1)$ and $\ell_0 = 96$. On the closed box $0 \le \lambda \le 0.1$, $0 \le e^2 \le 0.1$, for every $\varepsilon \le 1$ and every torus of side ≥ 96, we have $a_{K,T}(1) - a_{K,T}(0.7) \ge 0.3190043 > 0.319$.*

Proof. Cell upper sums. Fix μ_* and ℓ_0, and let ϕ be one of the functions j, $768M\bar{\tau}^2$, $16\varphi_3$, b_4 of Lemma G.2. Each of them is built from the majorants $\bar{\tau}, \bar{\psi}, \bar{\delta}, \bar{\Theta}, \bar{J}_s, \bar{J}'_s, \bar{\Lambda}_i$ and the constant $\bar{\omega}$ by sums, products and minima with non-negative numerical coefficients, multiplied by the prefactors M and $|\mu_*^2 - 2M^2|$. The majorants are non-negative and non-increasing in M (Lemma G.2), and $\bar{\omega}$ does not depend on M. The prefactor M is increasing. The prefactor $|\mu_*^2 - 2M^2|$ is monotone on each side of its zero $M = \mu_*/\sqrt{2}$, so its maximum on an interval is attained at an endpoint. For an interval $[a,b] \subset (0,\infty)$ let $\phi^{\rm cell}[a,b]$ be the number obtained from the formula for ϕ by evaluating every majorant at a and every prefactor at its maximum on $[a,b]$. Sums, products and minima of non-negative numbers are non-decreasing in each argument, so $\phi(M) \le \phi^{\rm cell}[a,b]$ for every $M \in [a,b]$. Hence, for every partition $a = M^{(0)} < M^{(1)} < \cdots < M^{(n)} = b$,

$$\int_a^b \phi\, dM \le \sum_{i=0}^{n-1} \big(M^{(i+1)} - M^{(i)}\big)\, \phi^{\rm cell}[M^{(i)}, M^{(i+1)}], \qquad \sup_{[a,b]}\phi \le \max_{0\le i<n} \phi^{\rm cell}[M^{(i)}, M^{(i+1)}]. \tag{G.1}$$

Take $[a,b] = [0.7, 1]$, the uniform partition of $[0.7, 1]$ into 400 cells, and all majorants evaluated at $\ell_0 = 96$. Then the upper sums (G.1) give

$$\bar{j} \le 0.0886293, \quad \bar{s} \le 7.179064, \quad \bar{B}_3 \le 9.330427, \quad \bar{B}_4^- \le 1.703784.$$

These four upper sums are evaluated in floating-point arithmetic with 40 significant digits. In this evaluation the constant w in $\bar{\omega}$ is replaced by the upper bound 0.25273101, every term that is subtracted in P_{gap} is multiplied by $1+10^{-25}$, and each result is rounded upward in the last digit shown.

Next, $h_1 \geq 0$, because $x \mapsto x\arctan(\pi/x)$ is increasing: its derivative is $\arctan y - y/(1+y^2) > 0$ with $y = \pi/x > 0$. Drop the term $16\lambda h_1$. Each remaining negative term is non-decreasing in $\lambda \geq 0$ and in $e^2 \geq 0$. So on the box P_{gap} is at least its value at the corner $(\lambda, e^2) = (0.1, 0.1)$:

$$P_{\mathrm{gap}} \;\geq\; 0.51 - 0.1 \cdot 0.0886293 - 0.01\,(7.179064 + 9.330427 + 1.703784) = 0.31900432.$$

The majorants are non-increasing in ℓ_0, so the bound at $\ell_0 = 96$ holds on every torus of side ≥ 96. Undoing the scaling gives (a): $2\sigma_+ \geq 0.319\, m^2$ for $\varepsilon \leq 1/m$ and $\ell_T \geq 96/m$. □

Remark G.5. The proof of Theorem 6.1(a) uses the rounded values $\bar{j} \leq 0.088629$ and $\bar{B}_3 \leq 9.3304$; they are slightly below the computed upper sums $0.08862928\ldots$ and $9.330426\ldots$. With the outward-rounded values above the corner value is 0.3190043 instead of 0.3190041, so the conclusion is unchanged.

The derivative bound and statement (b).

Proposition G.6 (derivative bound and gap between anchor masses above the target mass)**.** *Let $\lambda \geq 0$, $e \in \mathbb{R}$, $\mu_* > 0$, $\ell_0 > 0$ and $0 < m_1 < m_2$. Put*

$$p(M) := e^2 \tfrac{\pi^2}{32}\tau(M\ell_0) + \lambda^2 \frac{A_R(M\ell_0)}{M} + 16e^2\lambda\,\varphi_3(M) + e^4 b_4(M).$$

Then for every $K \geq 0$ ($\varepsilon \leq 1$) and every torus of side $\ell_T \geq \ell_0$:

(i) *$\partial_M a_{K,T}(M) \geq 2M - p(M)$ for every $M > 0$;*

(ii) *$a_{K,T}(m_2) - a_{K,T}(m_1) \geq (m_2 - m_1)(m_2 + m_1 - S)$ for every $S \geq \sup_{[m_1,m_2]} p$;*

(iii) *$\sup_{M \geq m_1} p(M) \leq e^2 s_1 + \lambda^2 s_2 + e^2\lambda s_3 + e^4 s_4$, where $s_1 = \frac{\pi^2}{32}\tau(m_1\ell_0)$, $s_2 = A_R(m_1\ell_0)/m_1$, $s_3 = 16\sup_{M\geq m_1}\varphi_3$ and $s_4 = \sup_{M\geq m_1} b_4$.*

With Corollary G.7 this gives Theorem 6.1(b).

Proof. (i): Differentiate Lemma G.1. The M-free terms drop out. The order-λ term contributes $-16\lambda\,\partial_M C_T \geq 0$ by (o-λ), and we drop it. The order-e^2 term contributes $-e^2\partial_M J_T \geq -e^2\frac{\pi^2}{32}\tau(M\ell_0)$. The order-$\lambda^2$ term contributes $128\lambda^2\partial_M S_T \geq -\lambda^2 A_R(M\ell_0)/M$. The order-$e^2\lambda$ term is at least $-16e^2\lambda\varphi_3$, and the order-e^4 term is at least $-e^4 b_4$, by Lemma G.2. (ii): By smoothness (Lemma G.1), $a(m_2) - a(m_1) = \int_{m_1}^{m_2} \partial_M a\, dM \geq \int_{m_1}^{m_2}(2M - S)\, dM = (m_2^2 - m_1^2) - (m_2 - m_1)S$. The bound does not depend on $\varepsilon \leq 1$ or on $\ell_T \geq \ell_0$. (iii): The supremum of a sum is at most the sum of the suprema. Since τ and $x \mapsto A_R(x\ell_0)/x$ are decreasing, the first two suprema are attained at $M = m_1$. □

Corollary G.7 (statement (b) of Theorem 6.1)**.** *Take $\mu_* = 1$ and $\ell_0 = 64$. On the box $0 \leq \lambda, e^2 \leq 0.1$, for $m_2 > m_1 > 2$, every $\varepsilon \leq 1$ and every torus of side ≥ 64 (in particular ≥ 96), we have $a_{K,T}(m_2) - a_{K,T}(m_1) \geq (m_2 - m_1)(m_2 + m_1 - 0.3622) \geq (m_2 - m_1)(m_2 + m_1 - 0.3631) > 0$.*

Here s_1 and s_2 are closed forms. The numbers s_3 and s_4 are bounded by upper sums over the cells of a partition of a bounded interval of masses, together with a bound on the tail of large masses.

Proof. At $m_1 = 2$, $\ell_0 = 64$: $s_1 = \frac{\pi^2}{32}\tau(128) = 0.317774\ldots \leq 0.3178$ and $s_2 = A_R(128)/2 = 9.694097\ldots \leq 9.6941$.

For $s_3 = 16\sup_{M\geq 2}\varphi_3$ and $s_4 = \sup_{M\geq 2} b_4$ choose $M_c \geq 2$ and a partition of $[2, M_c]$. On $[2, M_c]$ the supremum is bounded by the maximum of the cell values, by the second inequality in (G.1). On $[M_c, \infty)$ it is bounded by the monomial bound of Statement K.54 at M_c, because $M_c \geq 2 > \mu_*$. The resulting upper bounds are evaluated numerically, in floating-point arithmetic,

from the explicit majorants of Lemma G.2 at $\ell_0 = 64$ and from the monomials of Statement K.54. Rounded to nearest at four decimals, the values are 17.3775 and 5.9678. Hence $s_3 \le 17.37755$ and $s_4 \le 5.96785$.

Each term of p has a non-negative coefficient times a non-negative monomial in (e^2, λ), so p is non-decreasing in $\lambda, e^2 \ge 0$. Hence, on the box,

$$\sup_{M\ge 2} p \le 0.1 \cdot 0.3178 + 0.01\,(9.6941 + 17.37755 + 5.96785) = 0.362175 < 0.3622.$$

For $m_1 > 2$ we have $[m_1, m_2] \subset [2, \infty)$, so $S := 0.3622$ is admissible in (ii). Finally, $m_2 + m_1 - 0.3631 > 4 - 0.3631 > 0$. □

The coefficient condition and statement (c).

Corollary G.8 (the coefficient condition)**.** *Let $m > 0$, $c := 2H \ge 0$, and let $S(m) \ge \sup_{M\ge m} p(M)$. For every $\varepsilon \le 1$ and every torus of side $\ge \ell_0$, and every $m_1 \ge m$,*

$$a_{K,T}(m_1) - a_{K,T}(m) \ge (m_1 - m)(m_1 + m - S(m)).$$

Hence $a_{K,T}(m_1) \ge a_{K,T}(m) + 2H$ whenever $m_1 \ge m_1^c := \frac{1}{2}\big[S(m) + \big((S(m) - 2m)^2 + 4c\big)^{1/2}\big]$. Moreover $m_1^c \ge m$, and m_1^c is non-decreasing in $S(m)$.

This is Proposition G.6 with the anchors (m, m_1), written as an inequality between King's c-numbers; with Corollary G.9 it gives Theorem 6.1(c). In Section 6 the bare coefficient is $b_T = a_{K,T}(m)$ (Definition 2.2), which may be used on every torus by Lemma B.19(b).

Proof. The inequality is (ii) of Proposition G.6 with (m_1, m_2) replaced by (m, m_1) and $S = S(m)$ (it is trivial if $m_1 = m$). Put $q(x) := (x - m)(x + m - S) - c = x^2 - Sx + (Sm - m^2 - c)$. Its roots are $\frac{1}{2}[S \pm (S^2 - 4Sm + 4m^2 + 4c)^{1/2}] = \frac{1}{2}[S \pm ((S - 2m)^2 + 4c)^{1/2}]$. Since the leading coefficient is positive, $q(x) \ge 0$ for $x \ge m_1^c$. Also $((S - 2m)^2 + 4c)^{1/2} \ge |S - 2m| \ge 2m - S$, so $m_1^c \ge m$. Finally,

$$\partial_S m_1^c = \tfrac{1}{2}\big[1 + (S - 2m)\big((S - 2m)^2 + 4c\big)^{-1/2}\big] \ge 0.$$

□

Corollary G.9 (statement (c) of Theorem 6.1)**.** *Take $\mu_* = m = 1$, $H = 1$ and $\ell_0 = 64$. On the box $0 \le \lambda, e^2 \le 0.1$ we have $m_1^c \le 1.882 < 2$. Hence $a_{K,T}(m_1) \ge a_{K,T}(1) + 2H$ for every $m_1 > 2$, every $\varepsilon \le 1$ and every torus of side ≥ 64.*

Proof. Bound $\sup_{M\ge 1} p(M)$ at the corner $(0.1, 0.1)$ as follows. The terms $e^2 \frac{\pi^2}{32} \tau(M\ell_0)$ and $\lambda^2 A_R(M\ell_0)/M$ are decreasing in M, so their suprema are their values at $M = 1$. For $16e^2\lambda\varphi_3 + e^4 b_4$, choose $M_c \ge 1$ and use the cell values of (G.1) on a partition of $[1, M_c]$, and the monomial bound of Statement K.54 on $[M_c, \infty)$ (here $M_c \ge 1 = \mu_*$). This upper bound is evaluated numerically, in floating-point arithmetic, from the explicit majorants of Lemma G.2 at $\ell_0 = 64$. Rounded to nearest at four decimals it is 0.6119, so $S(1) \le 0.61195 \le 0.612$. Since S is non-decreasing in $\lambda, e^2 \ge 0$, $S(1) \le 0.612$ on the box. By the monotonicity in Corollary G.8,

$$m_1^c \le \tfrac{1}{2}\big[0.612 + (1.388^2 + 8)^{1/2}\big] = 1.88132\ldots \le 1.882.$$

□

G.1. **Proofs of Lemmas G.1 and G.2.** Notation is that of the beginning of this appendix, together with the following. $\mathsf{X} \in \{T, f\}$, where f stands for $\varepsilon\mathbb{Z}^3$; $\mathrm{BZ}_\varepsilon := [-\pi/\varepsilon, \pi/\varepsilon)^3$ and $\int_f := \int_{\mathrm{BZ}_\varepsilon} d^3k/(2\pi)^3$. $G_{M,\mathsf{X}}(x)$ and $D_\mathsf{X}(x)$ are the position kernels of $\hat{G}_M$ and $\hat{D}$, so $G_{M,T} = G_M$, $C_\mathsf{X}(M) = G_{M,\mathsf{X}}(0)$ and $D_\mathsf{X} := D_\mathsf{X}(0) = \int_\mathsf{X} \hat{D}$. $\mu := \mu_*$. Products of kernels are convolutions: $(K * K')(x) := \sum_y \varepsilon^3 K(x - y)K'(y)$. Finally $E_\mathsf{X}(M)$ (for $\mathsf{X} = T$) is $E_T(M) := C_T(M) - C_f(M)$, and $w := \varepsilon \int_f \hat{k}^{-2} = 0.252731$, which does not depend on ε.

Lattice tools.

Lemma G.10 (density of states)**.** *For every $\varepsilon > 0$ and every non-increasing $F : [0,\infty) \to [0,\infty)$,*

$$\int_{\mathrm{BZ}_\varepsilon} F(\hat{k}^2)\, d^3k \le c_N \int_{\mathbb{R}^3} F(|k|^2)\, d^3k, \qquad c_N = \tfrac{\pi^3}{8}.$$

Proof. On $[-\pi,\pi]$ we have $|\sin(y/2)| \ge |y|/\pi$, so $\hat{k}^2 \ge (4/\pi^2)|k|^2$ on BZ_ε. Hence $\{\hat{k}^2 \le E\} \subset \{|k| \le \frac{\pi}{2}E^{1/2}\}$, and this set has volume at most $\frac{4\pi}{3}(\frac{\pi^2 E}{4})^{3/2} = c_N \cdot \frac{4\pi}{3}E^{3/2}$.

For the integral, use the layer-cake formula: $\int_{\mathrm{BZ}} F(\hat{k}^2) = \int_0^\infty |\{F(\hat{k}^2) > t\}|\, dt$. Since F is non-increasing, $\{F(\hat{k}^2) > t\}$ is $\{\hat{k}^2 < E_t\}$ or $\{\hat{k}^2 \le E_t\}$. The level sets of $\hat{k}^2$ are null, so its measure is at most $c_N|\{k \in \mathbb{R}^3 : F(|k|^2) > t\}|$. If $E_t = \infty$ the right side is infinite. Integrate in t. □

Lemma G.11 (torus Riemann sums)**.** *Let F be as in Lemma G.10, and let T have side $\ell = N\varepsilon$ with N even. Then*

$$\int_T F(\hat{k}^2) \le c_N \int_{\mathbb{R}^3} F(|k|^2)\tfrac{d^3k}{(2\pi)^3} + \tfrac{3\pi}{16\ell}\int_0^\infty F(u)\, du + \tfrac{3}{4\ell^2}\int_{\mathbb{R}} F(q^2)\, dq + \tfrac{F(0)}{\ell^3}.$$

In particular $\int_T \hat{G}_M^2 \le M^{-1}[c_N/(8\pi) + \theta(M\ell)] = \frac{\pi^2}{64M}\tau(M\ell)$.

Proof. One coordinate. Let g be even and non-increasing in $|k|$ on $[-\pi/\varepsilon, \pi/\varepsilon]$, and put $h = 2\pi/\ell$. For $1 \le j \le N/2$ we have $h\, g(jh) \le \int_{(j-1)h}^{jh} g$, and symmetrically for $j < 0$. Summing, $\ell^{-1}\sum_j g(jh) \le \int g\, dk/(2\pi) + g(0)/\ell$; the edge point $j = N/2$ is included.

Three coordinates. $\hat{k}^2$ is increasing in each $|k_\mu|$ on the Brillouin zone. So $F(\hat{k}^2)$, and every partial sum or integral of it, stays even and non-increasing in each remaining coordinate. Iterating over the three coordinates gives $\sum_{S\subset\{1,2,3\}} \ell^{-|S|} \int F(\hat{k}^2)|_{k_S=0}$.

Bounding the pieces. The term $S = \emptyset$ is bounded by Lemma G.10. In the lower-dimensional integrals use $\hat{k}_\mu^2 \ge (4/\pi^2)k_\mu^2$ and extend to $\mathbb{R}^d$. This gives $\int_{\mathbb{R}^2} F(\frac{4}{\pi^2}|k|^2)\frac{d^2k}{(2\pi)^2} = \frac{\pi}{16}\int_0^\infty F$ and $\int_{\mathbb{R}} F(\frac{4}{\pi^2}k^2)\frac{dk}{2\pi} = \frac{1}{4}\int F(q^2)dq$.

The instance. For $F = (u + M^2)^{-2}$ the four terms are $\frac{c_N}{8\pi M}$, $\frac{3\pi}{16\ell M^2}$, $\frac{3\pi}{8\ell^2 M^3}$ and $\frac{1}{\ell^3 M^4}$. Their sum is $M^{-1}[c_N/(8\pi) + \theta(M\ell)]$. With $c_N = \pi^3/8$ this equals $\frac{\pi^2}{64M}\tau(M\ell)$, since $\frac{\pi^2}{64}\tau(x) = \frac{\pi^2}{64} + \theta(x)$. □

Lemma G.12 (positivity and images)**.** *Let $M, \mu > 0$ and $\mathsf{X} \in \{T, f\}$.*

(a) *$G_{M,\mathsf{X}}(x) > 0$ for every x, and $-\partial_{M^2} G_{M,\mathsf{X}} = G_{M,\mathsf{X}} * G_{M,\mathsf{X}} \ge 0$ pointwise.*

(b) *$G_{M,T}(x) = \sum_{n\in\mathbb{Z}^3} G_{M,f}(x + n\ell)$, and likewise for D and for $G * G$. In particular $E_T(M) = \sum_{n\ne 0} G_{M,f}(n\ell) \ge 0$ and $D_T - D_f = E_T(\mu)$.*

(c) *For $s, t > 0$ the position kernels of $\hat{G}^s$ and $\hat{D}^t$ are pointwise positive. Hence, for every q (with $q \in T^*$ on a torus),*

$$\int_{\mathsf{X},P} \hat{G}(P)^s \hat{D}(P \pm q)^t \le \int_{\mathsf{X}} \hat{G}^s \hat{D}^t, \quad \int_{\mathsf{X},P} \hat{G}(P)\hat{G}(P+q) \le \int_{\mathsf{X}} \hat{G}^2, \quad \int_{\mathsf{X},P} \hat{D}(P-q)\hat{D}(P+q) \le \int_{\mathsf{X}} \hat{D}^2.$$

Also $|(K * K')(x)| \le (K * K')(0)$ whenever $\hat{K}, \hat{K}' \ge 0$.

Proof. (a) Write $-\Delta_\varepsilon + M^2 = (6/\varepsilon^2 + M^2)\big(1 - P/(6 + \varepsilon^2 M^2)\big)$, where P is the nearest-neighbour adjacency matrix: its entries are ≥ 0 and its norm is 6. The Neumann series therefore has non-negative terms, its zeroth term is positive on the diagonal, and its n-th term reaches distance n. Moreover $-\partial_{M^2}(-\Delta + M^2)^{-1} = (-\Delta + M^2)^{-2}$.

(b) Both sides solve $(-\Delta + M^2)G = \varepsilon^{-3}\delta$ on T, and that solution is unique. The image sum converges absolutely by Lemma G.14(b) below.

(c) Positivity follows from subordination, $(-\Delta_\varepsilon + M^2)^{-s} = \Gamma(s)^{-1}\int_0^\infty \tau^{s-1} \mathrm{e}^{-\tau(-\Delta_\varepsilon + M^2)} d\tau$: the lattice heat kernel is positive, and on T one uses images. Then Parseval gives $\int_P \hat{K}(P)\hat{K}'(P-q) =$

$\sum_x \varepsilon^3 K(x)K'(x)\mathrm{e}^{-iqx} \le \sum_x \varepsilon^3 K(x)K'(x)$. For the last claim, $K * K'$ has non-negative Fourier transform $\hat K \hat K'$. □

Lemma G.13 (splitting and two lattice identities)**.** *Write* $\hat x := (2/\varepsilon)\sin(\varepsilon x/2)$ *componentwise.*

(a) *For* $q \neq 0$ *and* $0 < \theta \le \min(s,t)$*:* $\int_{\mathsf{X},P} \hat G(P)^s \hat D(P-q)^t \le (2/\hat q)^{2\theta}\big[\int_{\mathsf{X}} \hat G^{s-\theta}\hat D^t + \int_{\mathsf{X}} \hat G^s \hat D^{t-\theta}\big]$.

(b) $|\widehat{(a+b)}_\nu| \le |\hat a_\nu| + |\hat b_\nu|$*, hence* $\widehat{(a+b)}^2 \le 2\hat a^2 + 2\hat b^2$*. Also* $\widehat{(p+p')}^2 \le 2\widehat{(p-p')}^2 + 8\hat p'^2$.

(c) $|\hat D(P-q) - \hat D(P+q)| \le 4\hat P \hat q\, \hat D(P-q)\hat D(P+q)$.

(d) *Componentwise,* $\widehat{(a+b)}_\mu^2 + \widehat{(a-b)}_\mu^2 = 2\hat a_\mu^2 + 2\hat b_\mu^2 - \varepsilon^2 \hat a_\mu^2 \hat b_\mu^2$.

Proof. (a) By $|\sin(x+y)| \le |\sin x| + |\sin y|$ componentwise and Minkowski, $\hat q \le \hat P + \widehat{(q-P)}$. So either $\hat P \ge \hat q/2$, and then $\hat G(P)^\theta \le (2/\hat q)^{2\theta}$; or $\widehat{(P-q)} \ge \hat q/2$, and then $\hat D(P-q)^\theta \le (2/\hat q)^{2\theta}$. Bound the integrand by the sum of the two cases and apply Lemma G.12(c). When $s = \theta$, the factor $\hat G^0$ is the δ-kernel.

(b) The first claim is the same subadditivity. The second uses $|\sin 2x| \le 2|\sin x|$, so $\widehat{2v} \le 2\hat v$, followed by $(a+b)^2 \le 2a^2 + 2b^2$.

(c) Use $\sin^2 A - \sin^2 B = \sin(A+B)\sin(A-B)$, then $|(2/\varepsilon)\sin\varepsilon x| \le 2|\hat x|$ and Cauchy–Schwarz.

(d) This is $4 - 4\cos a\cos b = 2(2-2\cos a) + 2(2-2\cos b) - (2-2\cos a)(2-2\cos b)$, divided by ε^2. □

Lemma G.14 (torus images; the constant $\bar\omega$)**.** (a) *For every* ε *with* $\ell/\varepsilon \in \mathbb{N}$ *and every* $M > 0$*,*

$$E_T(M) \le \bar E(M,\ell) := \tfrac{7.37}{\ell} + \tfrac{3\pi}{4\ell} E_1\big(\tfrac{M^2\ell^2}{8}\big) + \tfrac{12\pi\,\mathrm{erfc}(M\ell/(2\sqrt2))}{M\ell^2} + \tfrac{64\mathrm{e}^{-M^2\ell^2/8}}{M^2\ell^3},$$

and $\bar E$ *is decreasing in* ℓ*.*

(b) *Let* $z \in \varepsilon\mathbb{Z}^3 \setminus \{0\}$*,* $r := |z|_\infty$*, and* $\kappa(x) := \ln(1 + x/(2\sqrt2))/(x/(2\sqrt2))$*. Then* $G_{M,f}(z) \le \frac{c_N}{\pi(2r-\varepsilon)}(1 + \frac{\varepsilon M}{2\sqrt2})^{1-2r/\varepsilon}$*. Hence, for* $\varepsilon \le 1$*,*

$$E_T(M) \le \tilde E(M,\ell) := \sum_{j\ge1} (24j^2+2)\frac{c_N(1+M/(2\sqrt2))}{\pi(2j\ell-1)}\mathrm{e}^{-\kappa(M)Mj\ell/\sqrt2},$$

which is decreasing in ℓ*.*

(c) *Put* $\hat E := \min\{\bar E, \tilde E\}$*. For* $\varepsilon \le 1$ *and* $\ell_T \ge \ell_0$*:* $\varepsilon D_T \le \bar\omega := w + \hat E(\mu,\ell_0)$.

In (c), "$\varepsilon \le 1$" is in the units in which $\bar E$ and $\tilde E$ are evaluated; see the units paragraph of Appendix G.

Proof. (a) This is Lemma H.31, with $7.364 \le 7.37$. Its hypothesis $\ell/\varepsilon \ge 2$ holds on the tori of this paper, where $\ell_T/\varepsilon \in 2\mathbb{N}$.

(b) First, $G_M = \int_{M^2}^\infty (G_s * G_s)\,ds$ by Lemma G.12(a). Let $z_1 = r > 0$. Shift $k_1 \to k_1 + i\eta$ in $(G_s * G_s)(z) = \int_f \mathrm{e}^{ikz}\hat G_s(k)^2$; the integrand is periodic, so the vertical sides cancel. Since $\mathrm{Re}\,\frac{4}{\varepsilon^2}\sin^2\frac{\varepsilon(k_1+i\eta)}{2} \ge \hat k_1^2 - \frac{4}{\varepsilon^2}\sinh^2\frac{\varepsilon\eta}{2}$, the choice $\frac{4}{\varepsilon^2}\sinh^2\frac{\varepsilon\eta_s}{2} = \frac{s}{2}$ gives $|\hat G_s| \le (\hat k^2 + s/2)^{-1}$. Lemma G.10 then gives $|(G_s * G_s)(z)| \le \mathrm{e}^{-\eta_s r} c_N/(8\pi(s/2)^{1/2})$, with $\eta_s \ge (2/\varepsilon)\ln(1 + \varepsilon\sqrt{s}/(2\sqrt2))$. Integrating over s gives the first bound.

For $\tilde E$: use the images of Lemma G.12(b), $\#\{n : |n|_\infty = j\} = 24j^2 + 2$, and $r = j\ell$. Each factor of the first bound is non-decreasing in ε (κ is decreasing), so $\varepsilon \le 1$ gives the value at $\varepsilon = 1$.

(c) $D_T = D_f + E_T(\mu)$ by Lemma G.12(b). Also $\varepsilon D_f \le \varepsilon \int_f \hat k^{-2} = w$. Finally $\varepsilon E_T(\mu) \le \hat E(\mu,\ell_T) \le \hat E(\mu,\ell_0)$, using $\varepsilon \le 1$ and monotonicity in ℓ. □

Lemma G.15 (one-loop majorants)**.** *Let* $\alpha,\beta \ge 0$ *with* $g := \alpha + \beta > 3/2$*, let* $a,b > 0$*, let* $\nu := \min(a,b)$*, and put* $F(u) := (u+a^2)^{-\alpha}(u+b^2)^{-\beta}$*. Define*

$$\mathcal{I}_{\alpha,\beta}(a,b;\ell) := c_N C_3 + \tfrac{3\pi}{16\ell} C_2 + \tfrac{3}{4\ell^2} C_1 + \tfrac{a^{-2\alpha}b^{-2\beta}}{\ell^3},$$

where each C_d is the minimum of the following admissible closed forms.

- C_3*:*
  - $\Gamma(g-\frac{3}{2})\nu^{3-2g}/((4\pi)^{3/2}\Gamma(g))$*;*
  - *if* $\beta<\frac{3}{2}<2g$*:* $\frac{1}{4\pi^2}B(c,\alpha-c)a^{2c-2\alpha}$ *with* $c=\frac{3-2\beta}{2}$*, and the same with* $(a,\alpha)\leftrightarrow(b,\beta)$*;*
  - *the exact values* $\frac{1}{4\pi(a+b)}$ *at* $(1,1)$ *and* $\frac{1}{8\pi a(a+b)^2}$ *at* $(2,1)$*.*
- C_2*:* $\nu^{2-2g}/(g-1)$*; or, if* $0<\beta<1$*,* $a^{2(1-g)}B(1-\beta,g-1)$*, and symmetrically.*
- C_1*:* $\sqrt{\pi}\Gamma(g-\frac{1}{2})\nu^{1-2g}/\Gamma(g)$*; or, if* $0<\beta<\frac{1}{2}$*,* $a^{1-2g}B(\frac{1}{2}-\beta,g-\frac{1}{2})$*, and symmetrically.*
- *For every* d*, the Hölder form* $C_d(g,a;0)^{\alpha/g}C_d(g,b;0)^{\beta/g}$*.*

Then:

(a) *For every* ε *and every torus of side* ℓ*:* $\int_T F(\hat{k}^2)\le\mathcal{I}_{\alpha,\beta}(a,b;\ell)$*. On* $\varepsilon\mathbb{Z}^3$ *the same holds with* $\ell=\infty$*.*

(b) $\mathcal{I}_{\alpha,\beta}$ *is decreasing in* ℓ*.*

(c) *At* $a=M$*,* $b=\mu$*, every admissible form is non-increasing in* M*, and hence so is* $\mathcal{I}_{\alpha,\beta}(M,\mu;\ell)$*.*

A factor $\hat{p}^s$ *with* $0\le s<2\alpha$ *is absorbed by* $\hat{p}^s\le(\hat{p}^2+a^2)^{s/2}$*, i.e.* $\alpha\to\alpha-s/2$*. The majorants used in Lemma G.2 are the following, all at* $(M,\mu;\ell_0)$*:*

symbol	*bounds*	*majorant*
$\bar\tau(M)$	$\tau_\Lambda:=\int_\Lambda\hat{G}^2$	$\mathcal{I}_{2,0}$
$\bar\psi(M)$	$\Xi_\Lambda(0):=\int_\Lambda\hat{D}\hat{G}$	$\mathcal{I}_{1,1}$
$\bar\delta$	$\delta_\Lambda:=\int_\Lambda\hat{D}^2$	$\mathcal{I}_{0,2}$
$\bar\Theta(M)$	$\Theta_\Lambda:=\int_\Lambda\hat{D}^{3/4}\hat{G}+\int_\Lambda\hat{D}\hat{G}^{3/4}$	$\mathcal{I}_{1,3/4}+\mathcal{I}_{3/4,1}$
$\bar J_s(M)$	$J_s:=\int_\Lambda\hat{G}^3\hat{p}^s$*,* $s\in\{0,1,\frac{3}{2}\}$	$\mathcal{I}_{3-s/2,0}$
$\bar J'_s(M)$	$J'_s:=\int_\Lambda\hat{D}\hat{G}^2\hat{p}^s$	$\mathcal{I}_{2-s/2,1}$
$\bar\Lambda_0$	$\int\hat{k}^{1/2}\hat{G}\hat{D}$	$\mathcal{I}_{3/4,1}$
$\bar\Lambda_1$	$\int\hat{G}^{5/4}\hat{D}+\int\hat{G}^2\hat{D}^{1/4}$	$\mathcal{I}_{5/4,1}+\mathcal{I}_{2,1/4}$
$\bar\Lambda_2$	$\int\hat{G}^{1/4}\hat{D}^2+\int\hat{G}\hat{D}^{5/4}$	$\mathcal{I}_{1/4,2}+\mathcal{I}_{1,5/4}$

In particular $\bar\tau(M)=\mathcal{I}_{2,0}(M;\ell_0)$ *is the Riemann form* $M^{-1}[c_N/(8\pi)+\theta(M\ell_0)]$ *of Lemma G.11. It is* not *an image form, which would need* $\varepsilon M\le1$*.*

Proof. (a) Apply Lemma G.11. Each closed form is the integral of a pointwise larger function: $F\le(u+\nu^2)^{-g}$, or $(k^2+b^2)^{-\beta}\le k^{-2\beta}$, or Hölder with exponents g/α and g/β. Each is evaluated by $\int_0^\infty t^{c-1}(t+a^2)^{-x}dt=a^{2(c-x)}B(c,x-c)$.

(b) The ℓ-terms are positive multiples of $\ell^{-1},\ell^{-2},\ell^{-3}$.

(c) Check each form in turn:

- the ν-forms: $\nu=\min(M,\mu)$ is non-decreasing and the exponent $d-2g$ is negative;
- the Beta forms: $M^{2c-2\alpha}$ with $c<\alpha$, or independent of M;
- the exact forms are decreasing;
- the Hölder forms are $M^{(d-2g)\alpha/g}$;
- the zero-mode term is $M^{-2\alpha}\mu^{-2\beta}$.

A minimum of non-increasing functions is non-increasing. □

The low orders and the order $e^2\lambda$. Write $\psi:=(\varphi_1+i\varphi_2)/\sqrt{2}$, $U_\mu(x)=\mathrm{e}^{i\varepsilon eA_\mu(x)}$, $s_\mu(q):=(2/\varepsilon)\sin(\varepsilon q_\mu/2)$. King's hopping term $\frac{1}{2}\sum_{x,\mu}\varepsilon|U_\mu(x)\varphi(x+\varepsilon e_\mu)-\varphi(x)|^2$ expands into:

- the free part $\sum_x\varepsilon^3\bar\psi(-\Delta_\varepsilon+M^2)\psi$, so $\langle\psi\bar\psi\rangle=\hat{G}_M$;
- the vertices $V_1=\sum_x\varepsilon^3(2e/\varepsilon)\sum_\mu A_\mu\,\mathrm{Im}(\bar\psi\psi')$ and $V_2=\sum_x\varepsilon^3e^2\sum_\mu A_\mu^2\,\mathrm{Re}(\bar\psi\psi')$;
- $V_3=-(\varepsilon e^3/3)\sum_x\varepsilon^3\sum_\mu A_\mu^3\,\mathrm{Im}(\bar\psi\psi')$ and $V_4=-(\varepsilon^2e^4/12)\sum_x\varepsilon^3\sum_\mu A_\mu^4\,\mathrm{Re}(\bar\psi\psi')$,

where $\psi' = \psi(x + \varepsilon e_\mu)$. The quartic term is $\lambda|\varphi|^4 = 4\lambda(\bar\psi\psi)^2$. A self-energy Σ is a shift of the coefficient of $|\psi|^2$, and $\delta m^2 = -\Sigma(0)$ order by order. Every V_n carries e^n and n photon legs, and all photon legs are contracted, so only even powers of e occur.

Lemma G.16 (orders λ, e^2, λ^2, $e^2\lambda$)**.** *On* $\mathsf{X} \in \{T, f\}$*:*

(i) $\delta m^2_{\lambda,\mathsf{X}} = -16\lambda C_{\mathsf{X}}(M)$.

(ii) *The order-*e^2 *self-energy is*

$$\Sigma_{e^2,\mathsf{X}}(p; M) = e^2 D_{\mathsf{X}} \sum_\mu \cos(\varepsilon p_\mu) - e^2 \int_{\mathsf{X},p'} \hat D(p - p')\widehat{(p + p')}^2 \hat G_M(p'),$$

and $\delta m^2_{e^2,\mathsf{X}} = -2e^2 D_{\mathsf{X}} - e^2 J_{\mathsf{X}}(M)$*: the seagull gives* $-3e^2 D_{\mathsf{X}}$ *and the exchange gives* $e^2 D_{\mathsf{X}} - e^2 J_{\mathsf{X}}$.

(iii) $\delta m^2_{\lambda^2,\mathsf{X}} = +128\lambda^2 S_{\mathsf{X}}(M)$.

(iv) $\delta m^2_{e^2\lambda,\mathsf{X}} = 16\lambda \int_{\mathsf{X}} \hat G_M^2 \tilde\Pi_{\mathsf{X}}(\cdot; M) = 16e^2\lambda F_{\mathsf{X}}(M)$*, where* $\tilde\Pi_{\mathsf{X}}(p) := \Sigma_{e^2,\mathsf{X}}(p) - \Sigma_{e^2,\mathsf{X}}(0)$*. Explicitly,*

$$\tilde\Pi_{\mathsf{X}}(p; M) = -e^2 D_{\mathsf{X}} \sum_\mu (1 - \cos\varepsilon p_\mu) - e^2 \int_{\mathsf{X},p'} \big[\hat D(p - p')\widehat{(p + p')}^2 - \hat D(p')\hat p'^2\big]\hat G_M(p').$$

The proof of (iv) uses Lemma G.19 below, which lists all one-particle-irreducible graphs of order $e^2\lambda$.

Proof. (i) Contracting one pair in $4\lambda(\bar\psi\psi)^2$ gives $16\lambda C\bar\psi\psi$.

(ii) The seagull with its photon pair contracted gives $e^2 D_{\mathsf{X}} \sum_\mu \cos(\varepsilon p_\mu)$, which is $3e^2 D_{\mathsf{X}}$ at $p = 0$. The second-order term $-\frac12\langle V_1 V_1\rangle$, with one photon and one scalar contraction, has vertex factors whose product is $e^2\widehat{(p + p')}^2_\mu$. At $p = 0$, $\int \hat D \hat p'^2 \hat G = D_{\mathsf{X}} - J_{\mathsf{X}}$, because $\hat p'^2 \hat G = 1 - M^2\hat G$.

(iii) Only the sunset survives: the cactus is cancelled exactly by the δm^2_λ insertion, since both contain the same C_{X}. The Wick count is $2 \cdot 2 \cdot 2$ per orientation, times 2 orientations, times $\frac12(4\lambda)^2$, i.e. $32(N + 2) = 128$ for $N = 2$.

(iv) By definition, the counterterm $\delta m^2_{e^2\lambda,\mathsf{X}}$ is minus the zero-momentum value of the sum of the 1PI graphs of order $e^2\lambda$, the vertex $c_{2,1}$ itself excluded (Corollary G.18(c) below). By Lemma G.19 these graphs are (a_1)–(a_3), (b_1), (b_2) and (c). The argument goes graph by graph, so it holds for $\mathsf{X} \in \{T, f\}$.

- The graphs (a_1), (a_2), (a_3) are the order-λ tadpole $16\lambda \int_{\mathsf{X}} \hat G$ with one of three insertions on its loop line: the seagull, the exchange, or the counterterm $\delta m^2_{e^2,\mathsf{X}}$. Their sum is the tadpole with $\hat G$ replaced by $-\hat G(\Sigma_{e^2,\mathsf{X}} + \delta m^2_{e^2,\mathsf{X}})\hat G = -\hat G^2\tilde\Pi_{\mathsf{X}}$, by (ii) and $\delta m^2_{e^2,\mathsf{X}} = -\Sigma_{e^2,\mathsf{X}}(0)$. So their contribution to the self-energy is $-16\lambda \int_{\mathsf{X}} \hat G^2\tilde\Pi_{\mathsf{X}}$.
- The graph (b_1) is the exchange graph with the insertion $16\lambda C_{\mathsf{X}}$ on its internal scalar line, and (b_2) is the same graph with the insertion $\delta m^2_{\lambda,\mathsf{X}} = -16\lambda C_{\mathsf{X}}$. Their sum is 0.
- Each graph (c) vanishes before the sum over the position of its λ-vertex. Let w be that position, and let V_1' sit on the bond $(z, z + \varepsilon e_\mu)$; both of its scalar lines end at w. By (G.2), the sum over the two orientations of V_1' is

$$c_1\big[G_M(z + \varepsilon e_\mu - w)\, G_M(z - w) - G_M(z - w)\, G_M(z + \varepsilon e_\mu - w)\big] = 0,$$

because G_M is even.

Hence $\delta m^2_{e^2\lambda,\mathsf{X}} = 16\lambda \int_{\mathsf{X}} \hat G_M^2 \tilde\Pi_{\mathsf{X}}$. □

Graphs, irreducibility and one-photon parts. In this paragraph T is a torus of side ℓ_T with $\ell_T/\varepsilon \in 2\mathbb{N}$, $\psi = \varphi/\sqrt{2}$, and

$$(-\Delta_A\psi)(x) = \varepsilon^{-2}\sum_\mu \big[2\psi(x) - \mathrm{e}^{i\varepsilon e A_\mu(x)}\psi(x+\varepsilon e_\mu) - \mathrm{e}^{-i\varepsilon e A_\mu(x-\varepsilon e_\mu)}\psi(x-\varepsilon e_\mu)\big],$$

so that the hopping term of the action of [24] is $\langle\psi, -\Delta_A\psi\rangle$. Let V_n be the part of $-\Delta_A - (-\Delta_\varepsilon)$ of degree n in A:

(G.2)

$$(V_n\psi)(x) = c_n\sum_\mu \Big[A_\mu(x)^n\psi(x+\varepsilon e_\mu) + (-1)^n A_\mu(x-\varepsilon e_\mu)^n\psi(x-\varepsilon e_\mu)\Big], \qquad c_n := -\varepsilon^{-2}\frac{(i\varepsilon e)^n}{n!}.$$

The quadratic forms $\langle\psi, V_n\psi\rangle$, $n = 1,\dots,4$, are the vertices $V_1,\dots,V_4$ written before Lemma G.16. We read (G.2) as follows. A vertex of type V_n sits on a bond $b = (x, x+\varepsilon e_\mu)$ and carries the field $A_\mu(x)^n$ of that bond. Its factor is c_n if its $\bar\psi$-leg is at the lower end x and its ψ-leg at the upper end $x+\varepsilon e_\mu$, and $(-1)^n c_n$ in the opposite case.

Let $\mathbb{E}_0$ be the Gaussian expectation under which ψ is a complex field with $\mathbb{E}_0[\psi(x)\bar\psi(y)] = G_M(x-y)$ and $\mathbb{E}_0[\psi\psi] = 0$, and A is an independent real field with $\mathbb{E}_0[A_\mu(x)A_\nu(y)] = \delta_{\mu\nu}D_T(x-y)$. The interaction is

$$\mathcal{V} := \langle\psi, (-\Delta_A + \Delta_\varepsilon)\psi\rangle + 4\lambda\sum_x \varepsilon^3|\psi(x)|^4 + \delta m^2\sum_x \varepsilon^3|\psi(x)|^2, \qquad \delta m^2 = \sum e^\alpha\lambda^\beta\,\delta m^2(\alpha,\beta).$$

The full two-point function $G_{\mathrm{full}}(x-y) := \mathbb{E}_0[\psi(x)\bar\psi(y)\mathrm{e}^{-\mathcal{V}}]/\mathbb{E}_0[\mathrm{e}^{-\mathcal{V}}]$ is a formal power series in (e,λ), and each coefficient is a finite sum of Gaussian moments.

By Wick's theorem each coefficient is a finite sum of values of *graphs*. A graph has vertices of three kinds. A vertex of type V_n has one ψ-leg, one $\bar\psi$-leg and n photon legs. A vertex λ sits at a site and has two ψ-legs and two $\bar\psi$-legs. A vertex $c_{\alpha,\beta}$ sits at a site, has one ψ-leg and one $\bar\psi$-leg, and has the factor $\delta m^2(\alpha,\beta)$. A graph has two external scalar legs. All its other legs are paired into lines. A scalar line joins a ψ-leg at u with a $\bar\psi$-leg at v and has the value $G_M(u-v)$. A photon line joins legs on the bonds (x,μ) and (y,ν) and has the value $\delta_{\mu\nu}D_T(x-y)$. The value of a graph is the sum, over the positions of its vertices, of the product of its line values and vertex factors. Only connected graphs contribute to G_{full}.

A connected two-point graph is *one-particle irreducible* (1PI) if it stays connected after the removal of any one line. It is *scalar-irreducible* if it stays connected after the removal of any one scalar line. A *one-photon part* of a graph is a connected set γ of vertices that carries no external leg and is joined to the other vertices by exactly one line, this line being a photon line. By [24, p. 659], the function Γ^ε in (3.33) there is the sum of the 1PI graphs.

Lemma G.17 (scalar parity and lattice inversion)**.** (a) *For every set S of vertices of a graph, the number of scalar legs at vertices of S that are not paired with a scalar leg at a vertex of S is even.*

(b) *Let γ be a connected configuration of vertices and lines in which all scalar legs are paired and exactly one photon leg is unpaired. This leg belongs to a vertex v_0 of type V_n. For a bond (z,μ) let $w_\mu(z)$ be the sum, over all positions of the vertices of γ with v_0 on (z,μ), of the product of the line values and vertex factors of γ. Then $w_\mu(z) = 0$.*

(c) *Every graph that contains a one-photon part has value* 0.

Proof. (a) Every vertex has two or four scalar legs, and every scalar line with both ends in S uses two of them. So the number of unpaired legs is an even number minus an even number.

(b) The value $w_\mu(z)$ does not depend on z, because all line values are translation invariant. Let ι be the inversion of T: $\iota(x) = -x$ on sites, and ι maps the bond $(x, x+\varepsilon e_\mu)$ to the bond $(-x-\varepsilon e_\mu, -x)$. Then ι maps the lower end of a bond to the upper end of its image and vice versa, and it preserves the direction μ. Apply ι to every position in a position assignment of γ, carrying

each leg to the image of its site. This is a bijection from the assignments with v_0 on (z,μ) onto those with v_0 on $\iota(z,\mu)$. Under it:

- a scalar line value $G_M(u-v)$ becomes $G_M(v-u)=G_M(u-v)$;
- a photon line value $\delta_{\mu\nu}D_T(x-y)$ becomes $\delta_{\mu\nu}D_T(y-x)$, which is the same;
- the factors of the vertices λ and $c_{\alpha,\beta}$ do not change;
- by (G.2), the factor of each vertex of type V_{n_v} is multiplied by $(-1)^{n_v}$, because its $\bar\psi$- and ψ-legs exchange the roles of lower and upper end.

Every photon line uses two photon legs and one photon leg is unpaired, so $\sum_v n_v$ is odd. Hence $w_\mu(z)=-w_\mu(\iota(z,\mu))=-w_\mu(z)$.

(c) Let γ be a one-photon part, joined to the rest by the photon line from the leg of $v_0\in\gamma$ on (z,μ) to a leg on (y,μ). By definition, every other line at a vertex of γ has both ends in γ. Fix z and the positions outside γ, and sum first over the positions of the vertices of γ other than v_0. This inner sum is $w_\mu(z)$ times factors that do not involve γ, and $w_\mu(z)=0$ by (b). Here γ, with the joining line removed, satisfies the hypotheses of (b): its scalar legs are all paired, by (a) applied to γ and because γ carries no external leg. □

Corollary G.18 (1PI and scalar-irreducible self-energies)**.** (a) *Let Γ be a connected two-point graph. Every scalar line whose removal disconnects Γ separates the two external legs.*

(b) *A scalar-irreducible two-point graph that is not 1PI contains a one-photon part, and its value is 0.*

(c) *Let $\Sigma_s(p)$ be the sum of the amputated scalar-irreducible two-point graphs other than the single counterterm vertices, with the sign convention of Lemma G.16. Then, as formal power series,*

$$\hat G_{\text{full}}(p)^{-1}=\hat p^2+M^2+\delta m^2+\Sigma_s(p),$$

and at every order $\Sigma_s(0)$ is the zero-momentum value $\Sigma_{1\text{PI}}(0)$ of the sum of the amputated 1PI graphs other than the counterterm vertices, i.e. of the function of [24, (3.33)]*. Hence the renormalisation condition of* [24, (3.33)]*, $\delta m^2(\alpha,\beta)+\Sigma_{1\text{PI}}(0;\alpha,\beta)=0$ for $2\le\alpha+2\beta\le4$, where $\Sigma_{1\text{PI}}(0;\alpha,\beta)$ is the coefficient of $e^\alpha\lambda^\beta$, holds if and only if the coefficients of $e^\alpha\lambda^\beta$, $2\le\alpha+2\beta\le4$, of $\hat G_{\text{full}}(0)-M^{-2}$ vanish.*

Proof. (a) Removing such a line leaves two components H_1,H_2, and each contains one end of the line. By Lemma G.17(a), the number of external legs in H_i plus one is even. So each H_i contains exactly one external leg.

(b) Let a photon line disconnect Γ into H_1,H_2. By Lemma G.17(a) the number of external (scalar) legs in each H_i is even, so one of them, say H_1, contains none. Then H_1 is a one-photon part, and Lemma G.17(c) applies.

(c) By (a), the scalar bridges of a connected two-point graph lie on every path between its external legs. So the amputated graph is a chain of scalar-irreducible two-point parts joined by single scalar lines. For a complex field the Wick weight of such a chain is the product of the weights of its parts, and summing the chains gives the geometric series $\hat G\sum_{n\ge0}\big(-(\delta m^2+\Sigma_s)\hat G\big)^n$. By (b), Σ_s and the sum of the 1PI graphs have the same value at every order. The last statement follows by induction on the order, because $\hat G(0)=M^{-2}$. □

Lemma G.19 (the graphs of order $e^2\lambda$)**.** *The vertex sets of order $e^2\lambda$ are*

$$S_1=\{\lambda,V_2\},\ S_2=\{\lambda,V_1,V_1'\},\ S_3=\{\lambda,c_{2,0}\},\ S_4=\{V_2,c_{0,1}\},\ S_5=\{V_1,V_1',c_{0,1}\},\ S_6=\{c_{2,0},c_{0,1}\}.$$

The 1PI two-point graphs on these sets are exactly the following six:

(a_1) *on S_1: both external legs at λ, two scalar lines λ–V_2, and the photon line from V_2 to itself;*

(a_2) *on S_2: both external legs at λ, scalar lines λ–V_1, V_1–V_1', V_1'–λ, and the photon line V_1–V_1';*

(a_3) *on S_3: both external legs at λ, and two scalar lines λ–$c_{2,0}$;*

(b_1) *on S_2: external legs at V_1 and V_1', scalar lines V_1–λ, λ–V_1' and a scalar line from λ to itself, and the photon line V_1–V_1';*
(b_2) *on S_5: external legs at V_1 and V_1', scalar lines V_1–$c_{0,1}$, $c_{0,1}$–V_1', and the photon line V_1–V_1';*
(c) *on S_2: external legs at λ and at V_1 (or at V_1'), one scalar line λ–V_1, two scalar lines λ–V_1', and the photon line V_1–V_1' (and the same with V_1 and V_1' exchanged).*

Every other graph on $S_1, \dots, S_6$ is disconnected or has a bridge.

Proof. The power λ^1 comes from exactly one vertex λ or $c_{0,1}$. The power e^2 comes from V_2, from V_1V_1, or from $c_{2,0}$, because V_n carries e^n. This gives $S_1, \dots, S_6$.

On each set the photon legs can be paired in only one way: the two legs of V_2 with each other, or the legs of V_1, V_1' with each other. Once the external legs are placed, the scalar lines form a multigraph whose degrees are the numbers of free scalar legs. We list all such multigraphs, writing uu for a scalar line from u to itself.

- S_1. External legs both at λ, degrees $(2,2)$: either $\{\lambda V_2, \lambda V_2\}$, which is ($\mathrm{a}_1$), or $\{\lambda\lambda, V_2V_2\}$, which is disconnected. Both at V_2, degrees $(4,0)$: λ is isolated. One at each, degrees $(3,1)$: $\{\lambda\lambda, \lambda V_2\}$, and λV_2 is a bridge.
- S_3. As for S_1, with $c_{2,0}$ in place of V_2 and no photon line. This gives (a_3), a disconnected graph, or a graph with a bridge.
- S_4 and S_6. There is one scalar line. If the external legs are at different vertices this line is a bridge; otherwise the other vertex is isolated.
- S_2, with degrees listed in the order (λ, V_1, V_1').
  - External legs (λ, λ), degrees $(2,2,2)$. The triangle is (a_2). In $\{\lambda V_1, \lambda V_1, V_1'V_1'\}$ and $\{\lambda V_1', \lambda V_1', V_1V_1\}$ the photon line is a bridge. The graphs $\{\lambda\lambda, V_1V_1', V_1V_1'\}$ and $\{\lambda\lambda, V_1V_1, V_1'V_1'\}$ are disconnected.
  - External legs (V_1, V_1'), degrees $(4,1,1)$. Either $\{\lambda\lambda, \lambda V_1, \lambda V_1'\}$, which is ($\mathrm{b}_1$), or $\{\lambda\lambda, \lambda\lambda, V_1V_1'\}$, which is disconnected.
  - External legs (V_1, V_1), degrees $(4,0,2)$. Either $\{\lambda\lambda, \lambda V_1', \lambda V_1'\}$, where the photon line is a bridge, or $\{\lambda\lambda, \lambda\lambda, V_1'V_1'\}$, which is disconnected. The case (V_1', V_1') is the same.
  - External legs (λ, V_1), degrees $(3,1,2)$. The graph $\{\lambda V_1, \lambda V_1', \lambda V_1'\}$ is (c). In $\{\lambda\lambda, \lambda V_1, V_1'V_1'\}$ the line λV_1 and the photon line are bridges. In $\{\lambda\lambda, \lambda V_1', V_1'V_1\}$ the line $\lambda V_1'$ is a bridge. There are no other cases: λ has odd degree, so it has one or three lines to the other vertices, at most one of them to V_1 and at most two to V_1'. The case (λ, V_1') is the same.
- S_5, with degrees listed in the order $(c_{0,1}, V_1, V_1')$ and $c := c_{0,1}$.
  - External legs (V_1, V_1'): either $\{V_1c, cV_1'\}$, which is (b_2), or $\{V_1V_1', cc\}$, which is disconnected.
  - External legs (V_1, V_1): either $\{cV_1', cV_1'\}$, where the photon line is a bridge, or $\{cc, V_1'V_1'\}$, which is disconnected.
  - External legs (c, c): c is isolated.
  - External legs (c, V_1): either $\{cV_1', V_1'V_1\}$ or $\{cV_1, V_1'V_1'\}$, and in both the line at c is a bridge.

□

Lemma G.20 (structure of $\tilde\Pi$)**.** *Put $\Xi_{\mathsf X}(p) := \int_{\mathsf{X},p'} \hat D(p-p')\hat G_M(p') \in [0, \Xi_{\mathsf X}(0)]$ and $Y_\mu(p) := \int_{\mathsf{X},p'} \hat D(p-p')\hat p_\mu'^2 \hat G_M(p') \in [0, D_{\mathsf X}]$.*

(a) $\tilde\Pi_{\mathsf X}(p;M) = -e^2\big\{2\hat p^2\Xi_{\mathsf X}(p) + (\mu^2 - 2M^2)[\Xi_{\mathsf X}(p) - \Xi_{\mathsf X}(0)] - \varepsilon^2 \sum_\mu \hat p_\mu^2 Y_\mu(p) + D_{\mathsf X}\tfrac{\varepsilon^2}{2}\hat p^2\big\}$.
(b) $|\tilde\Pi_{\mathsf X}(p;M)|/e^2 \le 2\sqrt{2}\,\Theta_{\mathsf X}(M)\hat p^{3/2} + |\mu^2 - 2M^2|\,\Xi_{\mathsf X}(0) + 3\sqrt{3}\,(\varepsilon D_{\mathsf X})\,\hat p$.
(c) $0 \le \partial_M \Sigma_{e^2,\mathsf X}(p;M) \le e^2\beta_{\mathsf X}(M)$ *with* $\beta_{\mathsf X} := 2M[2\tau_{\mathsf X} + 8\Xi_{\mathsf X}(0)]$. *Hence* $|\partial_M\tilde\Pi_{\mathsf X}(p;M)| \le e^2\beta_{\mathsf X}(M)$ *for all p.*

In (b) the seagull and Y terms could be sharpened from $\frac{3}{2}\varepsilon^2 D\hat{p}^2$ to $\frac{1}{2}\varepsilon^2 D\hat{p}^2$; this is not used.

Proof. (a) Apply Lemma G.13(d) to $\widehat{(p+p')}^2$ in Lemma G.16(iv), and use $\hat{k}^2\hat{D} = 1-\mu^2\hat{D}$ and $\hat{p}'^2\hat{G} = 1 - M^2\hat{G}$. The linearly divergent constants $2D_{\mathsf{X}} - C_{\mathsf{X}}$ cancel in $\Sigma(p) - \Sigma(0)$. The identity is algebraic, so it holds on T and on $\varepsilon\mathbb{Z}^3$. The ranges of Ξ_{X} and Y_μ follow from Lemma G.12(c) and $\hat{p}'^2_\mu\hat{G} \le 1$.

(b) Lemma G.13(a) with $s = t = 1$ and $\theta = \frac{1}{4}$ gives $\Xi_{\mathsf{X}}(p) \le \sqrt{2}\hat{p}^{-1/2}\Theta_{\mathsf{X}}$, which bounds $2\hat{p}^2\Xi_{\mathsf{X}}(p)$. Next, $|\Xi_{\mathsf{X}}(p) - \Xi_{\mathsf{X}}(0)| \le \Xi_{\mathsf{X}}(0)$. The seagull and Y terms together are at most $\frac{3}{2}\varepsilon^2 D_{\mathsf{X}}\hat{p}^2$, and $\varepsilon\hat{p} \le 2\sqrt{3}$ turns this into $3\sqrt{3}(\varepsilon D_{\mathsf{X}})\hat{p}$.

(c) We have $\partial_M\Sigma_{\mathrm{ex}}(p) = 2Me^2\int \hat{D}(p-p')\widehat{(p+p')}^2\hat{G}(p')^2 \ge 0$. Bound it with Lemma G.13(b), $\hat{D}\hat{k}^2 \le 1$, $\hat{p}'^2\hat{G} \le 1$ and Lemma G.12(c). The seagull and the counterterm do not depend on M, or do not depend on p, and so drop out of $\partial_M\tilde{\Pi}$. Then $\partial_M\Sigma_{\mathrm{ex}}(p)$ and $\partial_M\Sigma_{\mathrm{ex}}(0)$ both lie in $[0, e^2\beta_{\mathsf{X}}]$, so their difference is at most $e^2\beta_{\mathsf{X}}$ in absolute value. □

The order e^4: Ward form.

Proposition G.21 (Ward form of King's counterterm)**.** *Take $\lambda = 0$, a torus T and any ε. Let $a = M^2 + \delta m^2$ and $\mathcal{G}_{A,a} := (-\Delta_A + a)^{-1}$. Let $\langle\cdot\rangle$ be the photon expectation of the model at $\lambda = 0$: the Feynman Gaussian with mass μ, times $\det(-\Delta_A + a)^{-1}$. Put $\hat{G}_{\mathrm{full}} := (\langle\mathcal{G}_{A,a}\rangle)^\wedge$, $W_A := -\Delta_A\mathbf{1}$ and $R_A := W_A + ie\,\mathrm{div}A$. Then, as formal power series in e,*

$$\delta m^2 = e^2 D_T - e^2M^2\int_T \hat{D}\,\hat{G}_{\mathrm{full}} - M^2\langle(\mathcal{G}_{A,a}R_A)(x)\rangle.$$

King's δm^2, truncated at $\alpha + 2\beta \le 4$, satisfies this identity through order e^4, which is all that is used. Moreover $R_A = R^{(2)} + R^{(3)} + R^{(4)} + O(e^5)$ with

- $R^{(2)} = \frac{e^2}{2}\sum_\mu [A_\mu(x)^2 + A_\mu(x-\varepsilon e_\mu)^2]$,
- $R^{(3)} = \frac{ie^2\varepsilon^3}{6}\sum_\mu \nabla^-_\mu(A^3_\mu)(x)$,
- $R^{(4)} = -\frac{\varepsilon^2 e^4}{24}\sum_\mu [A_\mu(x)^4 + A_\mu(x-\varepsilon e_\mu)^4]$.

Proof. Step 1. King's condition [24, (3.33), p. 659] reads $\delta m^2(\alpha,\beta) + \Sigma_{1\mathrm{PI}}(0;\alpha,\beta) = 0$ at each order. By Dyson at $p = 0$, $\hat{G}_{\mathrm{full}}(0)^{-1} = M^2 + \delta m^2 + \Sigma_{1\mathrm{PI}}(0)$. So King's condition is equivalent to $\hat{G}_{\mathrm{full}}(0) = 1/M^2$, order by order.

Step 2. $(-\Delta_A + a)\mathbf{1} = a + W_A$, so $a\,\mathcal{G}_{A,a}\mathbf{1} = 1 - \mathcal{G}_{A,a}W_A$. Averaging gives $\delta m^2 = -M^2\langle(\mathcal{G}_{A,a}W_A)(x)\rangle$.

Step 3. Gauge covariance: $\mathcal{G}_{A+\nabla^+\chi,a}(x,y) = \mathrm{e}^{-ie(\chi(x)-\chi(y))}\mathcal{G}_{A,a}(x,y)$, and the determinant is invariant.

Step 4. Gaussian integration by parts with the Feynman covariance, together with Step 3 (the determinant contributes 0), gives $\langle \mathrm{div}A(y)F\rangle = \langle \frac{d}{dt}F(A + t\nabla^+\chi_y)|_0\rangle$ with $\chi_y(z) = -D_T(y-z)$.

Step 5. Applied to $F = \mathcal{G}_{A,a}(x,y)$: $\langle\mathcal{G}(x,y)\mathrm{div}A(y)\rangle = -ie(D_T(0) - D_T(x-y))G_{\mathrm{full}}(x-y)$. Multiplying by $-ie\varepsilon^3$, summing over y, and using Parseval and Step 1 gives $-M^2\langle\mathcal{G}\cdot(-ie\,\mathrm{div}A)\rangle = e^2D_T - e^2M^2\int\hat{D}\hat{G}_{\mathrm{full}}$.

Step 6. Add Steps 2 and 5. The expansion of R_A is that of $1-\cos\vartheta$ and $\sin\vartheta/\varepsilon - eA$, with $\vartheta = \varepsilon eA$. □

Corollary G.22 (the e^4 coefficient)**.** *On $\mathsf{X} \in \{T, \varepsilon\mathbb{Z}^3\}$, $\delta m^2_{e^4,\mathsf{X}}(M) = e^4 Q_{\mathsf{X}}(M)$ with*

$$Q_{\mathsf{X}} = \tfrac{3}{4}\varepsilon^2 D^2_{\mathsf{X}} + (A) + (C) + (D1) + (D2) + (D3),$$

where, with $s_\mu(k) := (2/\varepsilon)\sin(\varepsilon k_\mu/2) = \hat{k}_\mu$:

- $(A) = M^2\int_{\mathsf{X}}\hat{D}\hat{G}^2\tilde{\Pi}_{\mathsf{X}}/e^2$;

- $(C) = -\frac{\varepsilon^2 D_{\mathsf{X}}}{2}(D_{\mathsf{X}} - J_{\mathsf{X}}(M))$;
- $(D1) = \frac{1}{2}\sum_\mu \sum_z \varepsilon^3 [D_{\mathsf{X}}(z)^2 + D_{\mathsf{X}}(z+\varepsilon e_\mu)^2][G_{M,\mathsf{X}}(z) + G_{M,\mathsf{X}}(z+\varepsilon e_\mu)]$;
- $(D2) = -\sum_\mu \int_k \int_P s_\mu(k)[s_\mu(k) + s_\mu(2P+k)]\hat G(k)\hat G(P)\hat D(k)\hat D(P-k)$;
- $(D3) = \int_{\mathsf{X}} \hat D(k)^2 \sum_\mu \Pi_{\mu\mu}(k;M)/e^2$, *where* $\Pi_{\mu\mu}(k;M)/e^2 = 2G_{M,\mathsf{X}}(\varepsilon e_\mu) - \int_{\mathsf{X},q} \widehat{(2q+k)}_\mu^2 \hat G(q)\hat G(q+k)$.

The first term does not depend on M.

The proof expands Proposition G.21 to order e^4 on T; the case $\varepsilon\mathbb{Z}^3$ follows by a limit.

Proof. We work on $\mathsf{X} = T$, with the notation of Proposition G.21 ($G_{A,a}$, R_A and its parts $R^{(n)}$ of order e^n) and of the paragraph before Lemma G.17. Fourier transforms are $\hat f(k) = \sum_x \varepsilon^3 \mathrm{e}^{-ik\cdot x} f(x)$ and $f(x) = \int_T \mathrm{e}^{ik\cdot x}\hat f(k)$. The photon field is expanded at the lower ends of the bonds, $A_\mu(x) = \int_T \mathrm{e}^{ik\cdot x}\hat A_\mu(k)$, so all momentum-space factors below are periodic. By (G.2), the kernel of V_1 between an outgoing $\bar\psi$-momentum p and an incoming ψ-momentum p', with photon momentum $p-p'$ entering, is

$$v_1(p,p') = -\tfrac{ie}{\varepsilon}\big(\mathrm{e}^{i\varepsilon p'_\mu} - \mathrm{e}^{-i\varepsilon p_\mu}\big), \qquad v_1(p',p) = \overline{v_1(p,p')}, \qquad |v_1(p,p')|^2 = e^2 s_\mu(p+p')^2. \tag{G.3}$$

Write $G := (-\Delta_\varepsilon + M^2)^{-1}$, so that $G\mathbf{1} = M^{-2}\mathbf{1}$ and $\hat G(0) = M^{-2}$, and write $(KR)(x) := \sum_y \varepsilon^3 K(x,y)R(y)$.

Step 0 (the order-e^4 terms). Expand Proposition G.21 as a formal series in e, at $\lambda = 0$, with $\delta m^2 = e^2 d_2 + e^4 d_4$. The Neumann series gives $G_{A,a} = G^{(0)} + G^{(1)} + G^{(2)} + O(e^3)$ with

$$G^{(0)} = G, \qquad G^{(1)} = -GV_1G, \qquad G^{(2)} = -G(V_2 + e^2 d_2)G + GV_1GV_1G.$$

The weight is $\det(-\Delta_A+a)^{-1} = \det(-\Delta_\varepsilon+a)^{-1}\mathrm{e}^{-U}$ with $U = \operatorname{Tr}\log(1+(-\Delta_\varepsilon+a)^{-1}(-\Delta_A+\Delta_\varepsilon)) = U_1 + U_2 + U_3 + O(e^4)$, where U_m has order e^m and

$$U_1 = \operatorname{Tr}(GV_1) = 0, \qquad U_2 = \operatorname{Tr}(GV_2) - \tfrac{1}{2}\operatorname{Tr}(GV_1GV_1).$$

Here $U_1 = 0$ because, by (G.2), $\operatorname{Tr}(GV_1) = c_1 \sum_{x,\mu} \varepsilon^3 A_\mu(x)\big[G_M(\varepsilon e_\mu) - G_M(-\varepsilon e_\mu)\big]$ and G_M is even. Hence $\langle F\rangle = \mathbb{E}_0 F - \operatorname{Cov}_0(F,U_2) - \operatorname{Cov}_0(F,U_3) + \dots$, and $\operatorname{Cov}_0(F,U_m)$ has order at least (order of F)$+m$. The function $R_A = R^{(2)} + R^{(3)} + R^{(4)} + O(e^5)$ starts at order e^2. So the order-e^4 part of $\langle (G_{A,a}R_A)(x)\rangle$ is

$$\mathbb{E}_0\big[(GR^{(4)})(x)\big] + \mathbb{E}_0\big[(G^{(1)}R^{(3)})(x)\big] + \mathbb{E}_0\big[(G^{(2)}R^{(2)})(x)\big] - \operatorname{Cov}_0\big((GR^{(2)})(x), U_2\big). \tag{G.4}$$

Similarly $[\hat G_{\mathrm{full}}]_{e^2}$ is the Fourier transform of $\mathbb{E}_0 G^{(2)}$, because $\operatorname{Cov}_0(G,U_2) = 0$ for the A-independent G. By Lemma G.16(ii) and $d_2 = -\Sigma_{e^2}(0)$, this transform is $-\hat G^2\tilde\Pi_T$. All expectations are invariant under the translations of T. Since $\sum_x \varepsilon^3 G(x-z) = M^{-2}$, every term of the form $-M^2\,\mathbb{E}[(GY)(x)]$ equals $-\mathbb{E}[Y(0)]$; we use this repeatedly.

Step 1 (the term $-e^2M^2\int_T \hat D\hat G_{\mathrm{full}}$). Its order-$e^4$ part is $e^2M^2\int_T \hat D\hat G^2\tilde\Pi_T$, which is $e^4(A)$.

Step 2 (the constant). $-M^2\mathbb{E}_0[(GR^{(4)})(x)] = -\mathbb{E}_0 R^{(4)}(0) = \frac{\varepsilon^2 e^4}{24}\sum_\mu 2\cdot 3D_T^2 = \frac{3}{4}\varepsilon^2 e^4 D_T^2$, because $\mathbb{E}_0 A_\mu^4 = 3D_T^2$.

Step 3 (the term (C)). For L linear in A, $\mathbb{E}_0[L\,A_\mu(z)^3] = 3D_T\,\mathbb{E}_0[L\,A_\mu(z)]$. So in $\mathbb{E}_0[G^{(1)}R^{(3)}]$ the factor $R^{(3)} = \frac{i\varepsilon^2 e^3}{6}\sum_\mu \nabla^-_\mu(A_\mu^3)$ may be replaced by $\frac{i\varepsilon^2 e^3 D_T}{2}\operatorname{div}A = -\frac{\varepsilon^2 e^2 D_T}{2}(-ie\operatorname{div}A)$. Steps 3 and 4 of the proof of Proposition G.21, applied to the Gaussian $\mathbb{E}_0$ and to $G^{(1)}(x,y)$, which is linear in A, give

$$\mathbb{E}_0\big[G^{(1)}(x,y)\operatorname{div}A(y)\big] = -ie\big(D_T(0) - D_T(x-y)\big)G_M(x-y).$$

Hence $-M^2\mathbb{E}_0[(G^{(1)}(-ie\operatorname{div}A))(x)] = e^2M^2\big(D_T\hat G(0) - \int_T \hat D\hat G\big) = e^2(D_T - J_T)$, and the term is $-\frac{\varepsilon^2 e^4 D_T}{2}(D_T - J_T) = e^4(C)$.

Step 4 (the term $\mathbb{E}_0[G^{(2)}R^{(2)}]$). The two photon fields of $R^{(2)}$ are paired either with each other or with photon fields of the vertices in $G^{(2)}$.

- *Pairing with each other.* $\mathbb{E}_0 R^{(2)} = 3e^2 D_T$ is constant, so this part is $3e^2 D_T \sum_y \varepsilon^3 \mathbb{E}_0 G^{(2)}(x,y) = 3e^2 D_T [\hat G_{\mathrm{full}}]_{e^2}(0)$. It vanishes, because $\hat G_{\mathrm{full}}(0) = M^{-2}$ at order e^2 (Corollary G.18(c)).
- *Both fields paired with V_2.* This gives $e^4(D1)$. Indeed, by (G.2), $(V_2 f)(0) = \frac{e^2}{2}\sum_\mu [A_\mu(0)^2 f(\varepsilon e_\mu) + A_\mu(-\varepsilon e_\mu)^2 f(-\varepsilon e_\mu)]$, and the connected moment is $\mathbb{E}_0[A_\mu(u)^2 R^{(2)}(y)]_{\mathrm c} = e^2[D_T(u-y)^2 + D_T(u-y+\varepsilon e_\mu)^2]$. Therefore, after the shift $z = y - \varepsilon e_\mu$ in the first term and with $G_M(-z) = G_M(z)$,

$$-M^2\,\mathbb{E}_0\big[(-GV_2GR^{(2)})(x)\big]_{\mathrm c} = \mathbb{E}_0\big[(V_2GR^{(2)})(0)\big]_{\mathrm c} = \frac{e^4}{2}\sum_\mu\sum_z \varepsilon^3\big[D_T(z)^2 + D_T(z+\varepsilon e_\mu)^2\big]\big[G_M(z)+G_M(z+\varepsilon e_\mu)\big].$$

- *One field paired with each V_1.* This gives $e^4(D2)$. In $\mathbb{E}_0[(V_1GV_1GR^{(2)})(0)]$, let P be the momentum of the right propagator and k that of the left one. The left vertex has factor $v_1(0,k)$ and the right vertex $v_1(k,P)$. The photon lines carry k and $P-k$, with values $\hat D(k)$ and $\hat D(P-k)$. The field $R^{(2)} = \frac{e^2}{2}\sum_\mu[A_\mu(y)^2 + A_\mu(y-\varepsilon e_\mu)^2]$, with total photon momentum P and both fields paired, contributes $e^2(1+\mathrm e^{-i\varepsilon P_\mu})$. From $\mathrm e^{i\varepsilon k_\mu} - 1 = i\varepsilon\,\mathrm e^{i\varepsilon k_\mu/2} s_\mu(k)$ and $\mathrm e^{i\varepsilon P_\mu} + 1 - \mathrm e^{-i\varepsilon k_\mu} - \mathrm e^{-i\varepsilon(k+P)_\mu} = i\varepsilon\,\mathrm e^{-i\varepsilon k_\mu/2}[s_\mu(2P+k) + s_\mu(k)]$ we get

$$(\mathrm e^{i\varepsilon k_\mu}-1)\big(\mathrm e^{i\varepsilon P_\mu} - \mathrm e^{-i\varepsilon k_\mu}\big)\big(1+\mathrm e^{-i\varepsilon P_\mu}\big) = (i\varepsilon)^2\, s_\mu(k)\big[s_\mu(k) + s_\mu(2P+k)\big],$$

so the product of the three factors is $e^4 s_\mu(k)[s_\mu(k) + s_\mu(2P+k)]$. Hence

$$-M^2\,\mathbb{E}_0\big[(GV_1GV_1GR^{(2)})(x)\big]_{\mathrm c} = -e^4\sum_\mu \int_{T,k}\int_{T,P} s_\mu(k)\big[s_\mu(k)+s_\mu(2P+k)\big]\hat G(k)\hat G(P)\hat D(k)\hat D(P-k).$$

Step 5 (the measure term, $(D3)$). U_2 is a quadratic form in A, $U_2 = \frac12\langle A, \Pi A\rangle$. By (G.2), $\mathrm{Tr}(GV_2) = \sum_{x,\mu}\varepsilon^3 e^2 G_M(\varepsilon e_\mu) A_\mu(x)^2$. By (G.3), the diagonal symbol of $-\frac12\,\mathrm{Tr}(GV_1GV_1)$ is $-\frac{e^2}{2}\int_{T,q} s_\mu(2q+k)^2 \hat G(q)\hat G(q+k)$ per component. So

$$\Pi_{\mu\mu}(k) = 2e^2 G_M(\varepsilon e_\mu) - e^2\int_{T,q} \widehat{(2q+k)}_\mu^{\,2}\,\hat G(q)\hat G(q+k).$$

For a Gaussian, $\mathrm{Cov}_0(A_\mu(y)^2, \frac12\langle A,\Pi A\rangle) = (D\Pi D)_{\mu\mu}(y,y) = \int_T \hat D^2 \Pi_{\mu\mu}$, and the same holds for $A_\mu(y-\varepsilon e_\mu)^2$. Only the diagonal of Π enters, because the covariance is $\delta_{\mu\nu} D_T$. Therefore

$$M^2\,\mathrm{Cov}_0\left((GR^{(2)})(x), U_2\right) = \mathrm{Cov}_0\left(R^{(2)}(0), U_2\right) = e^2\sum_\mu\int_T \hat D(k)^2\Pi_{\mu\mu}(k),$$

and the term $-M^2\big(-\mathrm{Cov}_0((GR^{(2)})(x), U_2)\big)$ is $e^4(D3)$.

Step 6 (conclusion). The term $e^2 D_T$ of Proposition G.21 has order e^2. By (G.4), the order-e^4 part of $-M^2\langle (G_{A,a}R_A)(x)\rangle$ is the sum of Steps 2–5. Adding Step 1 gives $\delta m^2_{e^4,T} = e^4\big[\frac34\varepsilon^2 D_T^2 + (A) + (C) + (D1) + (D2) + (D3)\big]$. The first term does not depend on M. The identity of Proposition G.21 holds through order e^4 for the counterterm of Definition 2.2: by Corollary G.18(c), the renormalisation condition fixes d_2, d_4 through $\hat G_{\mathrm{full}}(0) = M^{-2}$ through that order.

The lattice $\varepsilon\mathbb{Z}^3$. Lemma G.1, and through it Propositions G.3 and G.6 and Corollary G.8, use the present corollary only for $\mathsf X = T$. For $\mathsf X = \varepsilon\mathbb{Z}^3$ the coefficient is defined graph by graph, with $\int_f$ in place of $\int_T$. Then both sides of the identity are finite sums of torus Riemann sums $|T|^{-L}\sum_{(T^*)^L}$ of continuous periodic functions on $\mathrm{BZ}^L_\varepsilon$, and all denominators are at least $\min(M^2, \mu_*^2)$. As $\ell_T \to \infty$

at fixed ε, these Riemann sums converge to the corresponding integrals, and the identity passes to the limit. □

Proof of Lemma G.1.

Proof. By [24, (3.32), p. 659] the counterterm has the orders $2 \le \alpha + 2\beta \le 4$, and the odd powers of e vanish (above, or Definition 2.2). So $\delta m^2_{K,T}$ is exactly the sum of its five orders λ, e^2, λ^2, $e^2\lambda$, e^4, with no remainder.

Lemma G.16 gives the first four orders. Corollary G.22 gives the order e^4: the coefficient of $e^4\lambda^0$ comes from graphs and insertions without a λ-vertex, so it is the e^4 coefficient of the $\lambda = 0$ model. Hence

$$a_{K,T} = M^2 - 16\lambda C_T - 2e^2 D_T - e^2 J_T + 128\lambda^2 S_T + 16e^2\lambda F_T + e^4 Q_T.$$

Smoothness. At fixed ε and T each term is a finite sum of products of factors $\hat G_M(k)$ and M-independent factors; $G_M(x)$ is itself such a finite sum. Since $\hat k^2 + M^2 \ge M^2$, each term is a rational function of M^2 with no pole on $(0,\infty)$. □

Proof of Lemma G.2.

Proof. Throughout, the barred majorants are those of Lemma G.15 at ℓ_0, and $\bar\omega$ is that of Lemma G.14(c). Every actual one-loop sum on T ($\ell_T \ge \ell_0$) is at most its majorant, by Lemma G.15(a),(b). The bound $\varepsilon D_T \le \bar\omega$ is the only place where $\varepsilon \le 1$ enters, apart from $h_\varepsilon \ge h_1$. No step needs $\varepsilon M \le 1$.

(o-λ). $\partial_M \hat G_M = -2M\hat G_M^2$. By Lemma G.12(b),(c), $C_T(M_1) - C_T(M_2) = (M_2^2 - M_1^2)(G_{M_1} * G_{M_2})_T(0)$, which is at least its value on $\varepsilon\mathbb{Z}^3$. On BZ_ε we have $\hat k^2 \le |k|^2$, and BZ_ε contains the ball of radius $R = \pi/\varepsilon$. So

$$C_f(M_1) - C_f(M_2) \ge \tfrac{1}{2\pi^2}\int_0^R \Big[\tfrac{M_2^2}{k^2+M_2^2} - \tfrac{M_1^2}{k^2+M_1^2}\Big]dk =: h_\varepsilon(M_1, M_2).$$

The integrand is ≥ 0, so h_ε is non-decreasing in R, and $h_\varepsilon \ge h_1$ for $\varepsilon \le 1$.

(o-e^2). $\partial_M J_T = 2M\int \hat D\hat G(1 - M^2\hat G) = 2M\int \hat D\hat k^2\hat G^2 \ge 0$. With $\hat D\hat k^2 \le 1$ this is at most $2M\tau_T \le 2M\bar\tau$; with $\hat k^2\hat G \le 1$ it is at most $2M\Xi_T(0) \le 2M\bar\psi$. Lemma G.11 gives $2M\tau_T \le \frac{\pi^2}{32}\tau(M\ell_T) \le \frac{\pi^2}{32}\tau(M\ell_0)$.

(o-λ^2). By Lemma G.12(a), $-\partial_M S_T = 3\sum_x \varepsilon^3 G_M^2(-\partial_M G_M) = 6M\sum_x \varepsilon^3 G_M(x)^2 (G_M * G_M)(x) \ge 0$. Next, $(G*G)(x) \le (G*G)(0) = \tau_T$ by Lemma G.12(c), and $\sum_x \varepsilon^3 G_M^2 = \tau_T$ by Parseval. Hence $-128\partial_M S_T \le 768M\tau_T^2 \le 768M\bar\tau^2 = A_R(M\ell_0)/M$.

(o-$e^2\lambda$). We have $\partial_M F_T = -4M\int \hat G^3\tilde\Pi/e^2 + \int \hat G^2\partial_M\tilde\Pi/e^2$. Lemma G.20(b) bounds the first term by $4M[2\sqrt{2}\Theta_T J_{3/2} + |\mu^2 - 2M^2|\Xi_T(0)J_0 + 3\sqrt{3}(\varepsilon D_T)J_1]$, and Lemma G.20(c) bounds the second by $\beta_T\tau_T$. Replace each factor by its majorant and εD_T by $\bar\omega$; this gives φ_3.

(o-e^4). By Corollary G.22, $\partial_M Q_T$ is the sum of the following five derivatives.

(C) $\partial_M(C) = \frac{\varepsilon^2 D_T}{2}\partial_M J_T \ge 0$ by (o-e^2). It is dropped.

(A) $(A) = M^2 H$ with $H := \int \hat D\hat G^2\tilde\Pi/e^2$, so $\partial_M(M^2H) = 2MH - 4M^3\int \hat D\hat G^3\tilde\Pi/e^2 + M^2\int \hat D\hat G^2\partial_M\tilde\Pi/e^2$. Use $M^2\hat G \le 1$ twice and Lemma G.20(b),(c): $|\partial_M(A)| \le 6M\int \hat D\hat G^2|\tilde\Pi|/e^2 + \beta_T\Xi_T(0) \le \varphi_A$, with $J'_s \le \bar J'_s$.

(D1) $\partial_M G(z) = -2M(G*G)(z) \in [-2M\tau_T, 0]$ by Lemma G.12(a),(c), and $\sum_z \varepsilon^3 D(z)^2 = \sum_z \varepsilon^3 D(z+\varepsilon e_\mu)^2 = \delta_T$. Over three directions this gives $-12M\bar\delta\bar\tau \le \partial_M(D1) \le 0$.

(D3) $\sum_\mu \partial_M[2G(\varepsilon e_\mu)] \in [-12M\tau, 0]$. The bubble part has derivative $2M\int \widehat{(2q+k)}_\mu^2(\hat G^2\hat G' + \hat G\hat G'^2) \ge 0$, where $\hat G' := \hat G(q+k)$. Summed over μ, by Lemma G.13(b) (with $a = q$, $b = q+k$), $\hat q^2\hat G \le 1$ and Lemma G.12(c), it is at most $2M[4\int \hat G\hat G' + 4\tau] \le 16M\tau$. Since $\int \hat D^2 = \delta$, $|\partial_M(D3)| \le 16M\bar\delta\bar\tau$.

(D2) Write $(D2) = -\sum_\mu \int_k s_\mu(k)\hat G(k)\hat D(k) S_\mu(k)$, where $S_\mu(k) := \int_P [s_\mu(k) + s_\mu(2P+k)]\hat G(P)\hat D(P-k)$. Decompose $s_\mu(2P+k) = (2/\varepsilon)[\sin(\varepsilon P_\mu)\cos(\varepsilon k_\mu/2) + \cos(\varepsilon P_\mu)\sin(\varepsilon k_\mu/2)]$.
- The first piece is odd in P except through $\hat D(P-k)$. Symmetrising $P \to -P$ and applying Lemma G.13(c) bounds it by $\int 2|\hat P_\mu| \cdot 2\hat P\hat k\, \hat G\hat D(P-k)\hat D(P+k)$.
- The other two pieces are each at most $|s_\mu(k)| \int \hat G\hat D(\cdot - k)$.

With $\sum_\mu |\hat k_\mu||\hat P_\mu| \le \hat k \hat P$, $\hat P^2 \hat G \le 1$ and Lemma G.12(c):

$$\sum_\mu |s_\mu||S_\mu| \le \hat k^2 (2\Xi(0) + 4\delta).$$

Likewise, using also $\hat D\hat D' \le \frac12(\hat D^2 + \hat D'^2)$ and Lemma G.13(a) with $\theta \in (\frac12, 1]$:

$$\sum_\mu |s_\mu||\partial_M S_\mu| \le 2M\hat k^2 (2/\hat k)^{2\theta} \big[2\big(\textstyle\int \hat G^{2-\theta}\hat D + \int \hat G^2 \hat D^{1-\theta}\big) + 4\big(\int \hat G^{1-\theta}\hat D^2 + \int \hat G \hat D^{2-\theta}\big)\big].$$

When ∂_M hits $\hat G(k)$, use $\hat k^2 \hat G^2 \hat D \le \hat G\hat D$: the contribution is at most $2M\Xi(0)(2\Xi(0) + 4\delta)$. When ∂_M hits $\hat G(P)$: at $\theta = 1$ the contribution is at most $8M\Xi(0)(6\Xi(0) + 2\tau + 4\delta)$, and at $\theta = \frac34$ it is at most $2^{5/2} M(2\Lambda_1 + 4\Lambda_2)\Lambda_0$. Together these give $|\partial_M(D2)| \le \varphi_{D2}$.

Summing, $\partial_M Q_T \ge -[\varphi_A + (12+16)M\bar\delta\bar\tau + \varphi_{D2}] = -b_4$.

Properties of the majorants. All of them are non-negative and coupling-free. The $\mathcal{I}$'s and $\bar\omega$ do not depend on ε, and they are non-increasing in M and in ℓ_0 (Lemmas G.15(b),(c) and G.14). The densities φ_3 and b_4 contain the increasing prefactors M and $|\mu^2 - 2M^2|$; these are handled by the upper sums over cells in Corollaries G.4 and G.7. □

Appendix H. The c-number bounds

This appendix proves the c-number bounds used in Section 6: the normalisation identities behind Propositions 6.3 and 6.6 (Lemma H.1), the bound on c-number differences of Proposition 6.3 (Proposition H.8), and the two bounds (6.1) and (6.2) of Proposition 6.6 (Lemma H.9 and Proposition H.11). The inputs from power counting, the weighted-line form of [3, Prop. 1] and the kernel bounds are in Subsection H.1.

Notation. Every symbol of Section 6 keeps its meaning. In addition: $\hat p^2 := \sum_\mu (4/\varepsilon^2)\sin^2(\varepsilon p_\mu/2)$; $\hat G_M(p) := (\hat p^2 + M^2)^{-1}$ is the free scalar propagator per real component; $T^* := (2\pi/\ell_T)\mathbb{Z}^3 \bmod (2\pi/\varepsilon)\mathbb{Z}^3$ is the set of torus momenta, and $\int_p$ means $|T|^{-1}\sum_{p\in T^*}$ on T and $\int_{[-\pi/\varepsilon,\pi/\varepsilon)^3} d^3p/(2\pi)^3$ on $\varepsilon\mathbb{Z}^3$. Write $C^T(s) := |T|^{-1}\sum_{p\in T^*}\hat G_{\sqrt s}(p)$ and $C^\infty(s) := \int_{\mathrm{BZ}} \hat G_{\sqrt s}$. $X_\varepsilon^T(M)$ is the order-e^2 coefficient of $\langle r_0^2\rangle$ on T at Higgs mass M with King's torus counterterm, and $X_\varepsilon(M)$ is the same on $\varepsilon\mathbb{Z}^3$, so $z_\varepsilon = e^2 X_\varepsilon(m)$. $[F]_{\alpha\beta}$ is the coefficient of $e^\alpha\lambda^\beta$ in a polynomial F in (e,λ), and $T_{\bar n}$ is the Taylor polynomial of total order $\le \bar n$ in (e,λ) at 0, with $\delta m^2_{K,T}(M)$ expanded as well. $Z_T^F(a) := \int \mathrm{e}^{-S_{\mu_*,a}} \prod_b dA_b \prod_x d^2\varphi(x)$ is the Lebesgue integral of the model (2.1) at photon mass μ_*. Note that the photon measure is *not* normalised here.

Normalisation identities.

Lemma H.1 (vacuum normalisations; identity (3.3))**.** *For every torus $T = T_j$, every spacing of the family and every $M > 0$, put $\gamma_T^\circ(M) := |T|^{-1}[E_0(M) + E_1(M)] + \frac12 a_{K,T}(M)(W_\varepsilon + z_\varepsilon)$. Then:*

(a) *$f_T(a_{K,T}(M)) = |T|^{-1}\log Z_K(T;M) + \gamma_T^\circ(M)$; in particular the $\gamma_T(M)$ of Proposition 6.3 equals $\gamma_T^\circ(M)$;*

(b) *$|T|^{-1}\log Z_K(T;M) = |T|^{-1}\Phi_M(0) + e(\varepsilon;0;M)$, with Φ_M and e as in Subsection 5.1;*

(c) *$e(\varepsilon;0;M') - e(\varepsilon;0;M) + \frac12\big(a_{K,T}(M) - a_{K,T}(M')\big)(W_\varepsilon + z_\varepsilon) = \gamma_T(M) - \gamma_T(M')$ for all $M, M' > 0$.*

Consequently, the hypothesis "Theorem 4.1 at m_1 in the form $|T|^{-1}\Phi_{m_1}(0) + e(\varepsilon;0;m_1) \geq -E_-(m_1)$" of Proposition 6.6 is exactly Theorem 4.1 at $M = m_1$.

The lemma uses Proposition 4.7. It gives the identity used in Proposition 6.3 and the form of the lower vacuum bound used in Proposition 6.6.

Proof. Write $a := a_{K,T}(M)$ and $N_{\mathrm{ph}} := \int \exp(-\frac{1}{2}\langle A,\, (-\Delta_\varepsilon + \mu_*^2)A\rangle) \prod_b dA_b$. This normalisation does not depend on M, e or λ. (i) By the definition of $Z(\Gamma;a)$ in Subsection 5.1 with $\Gamma = C_B(0) = C_{\mu_*}$, $Z_T^F(a) = N_{\mathrm{ph}}\, Z(C_{\mu_*};a)$, so $\log Z_T^F(a) = \log N_{\mathrm{ph}} + \Phi_M(0)$. (ii) King's vacuum constant is the Taylor polynomial $E_0(M) + E_1(M) = T_{\bar n}[\log Z_T^F(a)]$ [24, (2.6), (3.35)]. By Proposition 4.7, i.e. Lemma C.2(a),(b), it is Bałaban's $E_0 + E_1$ of [1, (1.12)–(1.13), p. 606]; the pairing with the normalised photon is Lemma C.2(c). There $\exp(E_0) = \exp(E_{0,v})\exp(E_{0,s})$ is the free Lebesgue normalisation, with $E_{0,v} = \log N_{\mathrm{ph}}$, and E_1 is the sum of the Taylor coefficients of orders $1 \leq \alpha+\beta \leq \bar n$ of $\log \int\int \exp(-S+E)$. Since $\log N_{\mathrm{ph}}$ is independent of (e,λ), $T_{\bar n}[\log N_{\mathrm{ph}} + \Phi_M(0)] = \log N_{\mathrm{ph}} + T_{\bar n}[\Phi_M(0)]$, and the order-zero term of $T_{\bar n}[\Phi_M(0)]$ is $E_{0,s}(M)$. (iii) By definition $Z_K(T;M) = Z_T^F(a)\,\mathrm{e}^{-E_0(M)-E_1(M)}$ (Subsection 4.2), so by (i)–(ii)

$$\log Z_K(T;M) = \Phi_M(0) - T_{\bar n}[\Phi_M(0)] = \Phi_M(0) + |T|\, e(\varepsilon;0;M),$$

which is (b). (iv) $f_T(a) = |T|^{-1}\log Z_T^F(a) + \frac{1}{2}a(W_\varepsilon + z_\varepsilon) = |T|^{-1}[\log Z_K(T;M) + E_0(M) + E_1(M)] + \frac{1}{2}a(W_\varepsilon + z_\varepsilon)$, which is (a). (v) By (ii), $e(\varepsilon;0;M) = -|T|^{-1}[E_0(M) + E_1(M)] + |T|^{-1}E_{0,v}(\mu_*)$. The last term is free of M and cancels in the difference in (c); what remains is $\gamma_T^\circ(M) - \gamma_T^\circ(M')$. The final sentence of the lemma follows from (b). □

Lattice power counting in three dimensions. *Reisz's theorems in $d = 3$* are Proposition C.3, item 3 of Subsection 4.6.

Three inputs. The proof of Proposition H.8 uses three inputs, which we name. (q1) The transfer from $d = 4$ to $d = 3$ of Proposition C.3(a)–(b) (Lemma H.12). (q2) The identification of King's counterterm expansion with a Reisz forest sum (Lemmas H.3 and H.13), by the standard counterterm ⇔ forest combinatorics (Bogoliubov–Zimmermann). (q3) The constant z_ε is the graphical one, $z_\varepsilon = e^2 X_\varepsilon(m)$ (Subsection 2.4); this is the scheme of Subsection 6.1.

The c-number differences. Fix anchor masses $0 < \underline{M} < \overline{M}$ with $\varepsilon\overline{M} \leq 1$; for the spacings of the construction $\varepsilon \leq 1/m_2$ by Lemma A.5. Put $\hat\Delta := \hat G_{\underline{M}} - \hat G_{\overline{M}} = (\overline{M}^2 - \underline{M}^2)\hat G_{\underline{M}}\hat G_{\overline{M}}$ and $C_t := \hat G_{\overline{M}} + t\hat\Delta$ for $t \in [0,1]$. Let δm_t^2 be King's recursion [24, (3.33)] with scalar lines C_t and photon lines C_{μ_*}, and write $\delta\dot m_t^2 := \partial_t \delta m_t^2$. Let $\langle\cdot\rangle_t$ be the model with scalar Gaussian $C_t \otimes \mathbf{1}_2$, photon Gaussian C_{μ_*} and interaction V_t (the e-dependent hopping terms, $\lambda\sum\varepsilon^3 r^4$ and $\frac{1}{2}\delta m_t^2 \sum \varepsilon^3 r^2$). Write $\mathcal{G}_t := \langle\varphi\varphi^{\mathrm{T}}\rangle_t$, $\mathcal{A}_t := C_t^{-1}(\mathcal{G}_t - C_t)C_t^{-1}$, $\mathcal{P}_t := \int_p \mathrm{tr}[\hat G_{\underline{M}}\hat G_{\overline{M}}\mathcal{A}_t] - z_\varepsilon$ and $\rho_t := \langle r_0^2\rangle_t - W_\varepsilon - z_\varepsilon$.

Lemma H.2 (covariance path)**.** *For every $(\alpha,\beta) \neq (0,0)$,*

$$[\gamma_T(\underline{M}) - \gamma_T(\overline{M})]_{\alpha\beta} = \int_0^1 \Big\{\tfrac{1}{2}(\overline{M}^2 - \underline{M}^2)[\mathcal{P}_t]_{\alpha\beta} - \tfrac{1}{2}[\delta\dot m_t^2\, \rho_t]_{\alpha\beta}\Big\} dt.$$

Every coefficient of $\mathcal{P}_t, \rho_t, \delta\dot m_t^2$ is a polynomial in t. The $(0,0)$ coefficient equals $\int_{\underline{M}^2}^{\overline{M}^2}[C^T(s) - C^\infty(s)]\,ds + \int_{\underline{M}^2}^{\overline{M}^2}[C^\infty(s) - C^\infty(m^2)]\,ds$.

Proof. Put $s(t) := \overline{M}^2 + t(\underline{M}^2 - \overline{M}^2)$ and $\Gamma(t) := |T|^{-1}T^{\geq 1}\log\langle \mathrm{e}^{-V_t}\rangle_{C_t} + \frac{1}{2}s(t)z_\varepsilon + \frac{1}{2}\delta m_t^2(W_\varepsilon + z_\varepsilon)$, where $T^{\geq 1}$ keeps the orders $1 \leq \alpha+\beta \leq \bar n$. At $t = 0,1$ the Gaussian is the free measure at $\overline{M}$, resp. $\underline{M}$, so $|T|^{-1}T^{\geq 1}\log\langle \mathrm{e}^{-V}\rangle = |T|^{-1}E_1$, and the c-numbers equal $\frac{1}{2}a_{K,T}(M)(W_\varepsilon + z_\varepsilon) - \frac{1}{2}M^2 W_\varepsilon$. Hence $\Gamma(1) - \Gamma(0)$ is the order-≥ 1 part of $\gamma_T(\underline{M}) - \gamma_T(\overline{M})$. Each coefficient of Γ is a polynomial in t (a finite Wick sum), so $\Gamma(1) - \Gamma(0) = \int_0^1 \Gamma'$ coefficientwise. Γ' has three parts. (i) The covariance: $\frac{d}{dt}\log\int d\mu_{C_t} F = \frac{1}{2}\,\mathrm{tr}[\dot C_t C_t^{-1}(\mathcal{G} - C_t)C_t^{-1}]$ (differentiate the density $\propto \det C^{-1/2}\mathrm{e}^{-\frac{1}{2}\langle\varphi, C^{-1}\varphi\rangle}$ in finite

dimension). With $\dot C_t = \hat\Delta$ this gives, per volume, $\frac12(\overline{M}^2 - \underline{M}^2)\int_p \mathrm{tr}[\hat G_{\underline{M}}\hat G_{\overline{M}}\mathcal{A}_t]$. (ii) The vertex: $-\frac12\delta\dot m_t^2\langle r_0^2\rangle_t$. (iii) The c-numbers: $\frac12(\underline{M}^2 - \overline{M}^2)z_\varepsilon + \frac12\delta\dot m_t^2(W_\varepsilon + z_\varepsilon)$. Summing (i)–(iii) gives the identity. For the $(0,0)$ coefficient: $|T|^{-1}E_0(M) = -\int_p \log(\hat p^2 + M^2)$ + (an M-free term), so $\partial_{M^2}|T|^{-1}E_0 = -C^T(M^2)$. Also $\frac12(\underline{M}^2 - \overline{M}^2)W_\varepsilon = -\int_{\underline{M}^2}^{\overline{M}^2} C^\infty(m^2)\,ds$. □

Lemma H.3 (graph bookkeeping on the torus)**.** *Each $t^j e^\alpha \lambda^\beta$-coefficient of $\delta m_t^2 - \delta m_0^2$, $\delta\dot m_t^2$, $\mathcal{P}_t$ and ρ_t (minus its W_ε, z_ε terms) is a finite rational combination of terms. Each term is a product of three kinds of factor: lower-order coefficients of $\delta m_t^2 - \delta m_0^2$; Ward defects $\Pi_c^T(0)$; and a torus sum $S_T(\mathcal{S}) := |T|^{-L}\sum_{k\in(T^*)^L} R_{\mathcal{S}}(k)$ of a Reisz forest integrand of a reduced graph $\mathcal{S}$. Here:*

- *a reduced graph is a graph with lines $\hat G_{\overline{M}}$ or $\hat\Delta$ and C_{μ_*} in which pairwise disjoint subgraphs of class R3 of Lemma H.14 (two-photon parts of order $o \in \{1,2\}$) are contracted to constant vertices;*
- *$\Pi_c^T(0)$ is the zero-momentum value of the* full *order-o Taylor coefficient of the $\hat\Delta$-free, δm^2-difference-free 1PI two-photon function of the uniform-mass-$\overline{M}$ model, with King's counterterm δm_0^2 inside;*
- *the forests subtract with $\delta(\gamma) = \omega(\gamma)$ every 1PI proper part γ that contains no $\hat\Delta$-line, no δm^2-difference vertex and not the insertion, where $\omega = 3 - \frac12(E_\varphi + E_A) - o_V - 2v_2$. By Lemma H.14, only the parts of the classes R2–R6 are actually subtracted.*

Every forest of a reduced graph is Reisz-admissible. Every reduced graph has $\omega \le -1$, apart from the overall non-negative pieces listed in step (iv) of Proposition H.8.

The lemma uses Lemma H.15, Lemma H.17 and Lemma H.14, i.e. step (a) of the proof of Lemma H.13 in its torus form. It also uses the torus Ward identity, Statement K.55.

Proof. By Lemma H.15, which holds on T with sums over T^*, the insertion of King's counterterm δm_0^2 is exactly the sum over the t^0-forests of the blocks. By the remark after that lemma and by Lemma H.17, the blocks are exactly the scalar self-energy parts that contain no $\hat\Delta$-line, no f-vertex and not the insertion and have order at most 2, i.e. the parts of class R2 of Lemma H.14. Also, the forests of blocks are Reisz's forests restricted to these parts. So King's counterterm expansion on T is the forest sum of [33] with t^0 on the R2 parts. We now add the subtractions of the remaining divergent parts, together with the t^1 subtractions of the R2 parts. On T the $t^1 - t^0$ terms of R2 parts, and the t^0 terms of R4, R5, R6 parts, are odd under $k \mapsto -k$ of the internal momenta. Since $T^* = -T^*$, their torus sums vanish exactly. Adding them does not change the sum. For R3 this cancellation fails: a constant photon field on T is a twist, not a gauge transformation, so the class sum is $\Pi_c^T(0) \neq 0$. Group the forests by their set P of R3 elements. These are pairwise disjoint, since at $o \le 2$ no R3 part contains another. This gives $\mathrm{Full}(G) = \sum_P (-\Pi)^P \mathrm{Phys}(G/P)$. The positions form a Boolean lattice, so Möbius inversion gives $\mathrm{Phys}(G) = \sum_P \Pi^P \mathrm{Full}(G/P)$. Admissibility: $\omega(\tau/P) = \omega(\tau) - \sum_{\gamma\in P,\gamma\subset\tau}\omega(\gamma) \le \delta(\tau)$. This is an over-subtraction, which [33, (4-3)] allows. Condition (4-4) holds because the dropped elements have $\delta \ge 0$. The Ward identity of Lemma H.5, i.e. Statement K.55 with $a = \overline{M}^2 + \delta m_0^2$, holds only for the full class coefficient of the gauge-invariant model of uniform mass. The sum above runs over exactly that class. The assertions on ω follow from Lemma H.14(b). □

Lemma H.4 (torus sums)**.** *Let R be a forest integrand as in Proposition C.3(b)–(c) with L loops, $3h + \deg_H R \le -1$ for all H, and bounded external momenta. Put $\mathfrak{s}(x) := \sum_{n\in\mathbb{Z}^3\setminus 0} \mathrm{e}^{-x|n|}$. For $\ell_T \ge \ell_0$ and $\varepsilon < a_0$, $|S_T - I_\infty| \le C'[(1+\mathfrak{s}(\eta_0\ell_T))^L - 1]$, where $I_\infty := \int_{\mathrm{BZ}^L} R$. Hence $\sup_{\varepsilon,\ell_T\ge\ell_0}|S_T| < \infty$.*

The lemma uses Proposition C.3.

Proof. Poisson summation on the lattice torus gives $|T|^{-L}\sum_{(T^*)^L} f = \sum_{n\in\mathbb{Z}^{3L}}\int_{\mathrm{BZ}^L} f(k)\mathrm{e}^{i\ell_T n\cdot k}$; check it on the characters $\mathrm{e}^{-ik\cdot x}$. For $n\neq 0$, shift $k_j \mapsto k_j + i\eta_0 n_j/|n_j|$. This is allowed because R is analytic in the strip and periodic, so the vertical sides of the contour cancel. The shift gives $|\int R\mathrm{e}^{i\ell_T n\cdot k}| \le \mathrm{e}^{-\eta_0\ell_T\sum_j|n_j|}C'$ by Proposition C.3(c). Finally $\sum_{n\neq 0}\mathrm{e}^{-x\sum_j|n_j|} = (1+\mathfrak{s}(x))^L - 1$, and $|I_\infty| \le C$ by Proposition C.3(a). □

Lemma H.5 (torus Ward identity and its defect)**.** *(a) For $o\in\{1,2\}$, the order-o 1PI two-photon coefficient $\Pi^{(o)}$ of the uniform-mass model on T satisfies $\sum_\nu \Pi^{(o)}_{\mu\nu}(q)\tilde q_\nu = 0$ for all $q\in T^*$, with $\tilde q_\nu := (\mathrm{e}^{i\varepsilon q_\nu}-1)/\varepsilon$. In particular $\Pi^{(o)}_{\mu\nu}(q_1 e_\nu) = 0$ for $q_1 := 2\pi/\ell_T$. The value $\Pi^{(o)}(0)$ is not constrained. (b) For $\overline{M}\in\{m,m_1,m_2\}$, $\sup_{\varepsilon<a_0,\ell_T\ge\ell_0}|\Pi^T_c(0)| < \infty$.*

Part (a) is unconditional; part (b) uses Proposition C.3.

Proof. (a) Every coefficient is a Gaussian moment on the finite torus. The change of variables $\delta\varphi = e\chi q\varphi$, $\delta A = -d\chi$ (χ periodic) leaves the matter action invariant (Subsection 2.2) and has Jacobian 1. It changes $\frac12\langle A, (-\Delta_\varepsilon+\mu_*^2)A\rangle$ by a term *linear* in A. Gaussian integration by parts then shows that Γ_{int} (the Legendre transform minus the free quadratic form) is invariant to first order under $A\mapsto A - d\chi$, so $\Pi\, d\chi = 0$. The gradients $d\chi$ span exactly the non-constant modes. (b) By (a), $\Pi^T(0) = -[\Pi^T(q_1e_\nu) - \Pi^T(0)]$. The bracket is the torus sum of the $(1-t^\omega)$-subtracted class integrand at the fixed external momentum q_1e_ν; for $o=1$ the extra t^1-t^0 part is odd and sums to 0. This integrand is Reisz-admissible with $3h+\deg_H \le -1$, so Lemma H.4 applies. □

Lemma H.6 (one propagator)**.** *For $\varepsilon M \le 1$ and $x\in\varepsilon\mathbb{Z}^3\setminus 0$, $|G^\varepsilon_M(x)| \le 77.1\, M^{-3}|x|^{-4}$, where $G^\varepsilon_M(x) := \int_{\mathrm{BZ}}\hat G_M \mathrm{e}^{ik\cdot x}$. Hence $|C^T(M^2) - C^\infty(M^2)| \le 77.1\,\zeta_3(4) M^{-3}\ell_T^{-4}$, where $\zeta_3(4) := \sum_{n\in\mathbb{Z}^3\setminus 0}|n|^{-4} \le 16.6$.*

Proof. $|x|^4|G(x)| \le \int_{\mathrm{BZ}}|\Delta_k^2\hat G_M|$, by two periodic integrations by parts. Put $h=\hat k^2$ and $s = h+M^2$. Then $|\partial_\mu h| \le 2h_\mu^{1/2}$, $|\partial_\mu^2 h| \le 2$, $|\partial_\mu^3 h| \le 2\varepsilon$, $|\partial_\mu^4 h| \le 2\varepsilon^2$, and $\varepsilon^2 \le 24/s$. Faà di Bruno gives $|\Delta^2\hat G| \le 2000 s^{-3}$. Since $h \ge (4/\pi^2)|k|^2$, $\int_{\mathrm{BZ}} s^{-3} \le \pi^2/(256M^3)$, and $2000\pi^2/256 \le 77.1$. The torus bound follows from $C^T - C^\infty = \sum_{n\neq 0} G^\varepsilon_M(n\ell_T)$. □

Lemma H.7 (the order e^2 of $\langle|\varphi(0)|^2\rangle$ on the torus)**.** *For $\varepsilon M\le 1$ and $\ell_T\ge\ell_0$, $|X^T_\varepsilon(M) - X_\varepsilon(M)| \le C_z(M,\mu_*)\,\zeta_6(8)\,\ell_T^{-8}$, where $\zeta_6(8) := \sum_{N\in\mathbb{Z}^6\setminus 0}|N|^{-8} \le 22.1$.*

The lemma uses Proposition C.3(a)–(b); the constant is Reisz's and is not explicit.

Proof. Let $\sigma(k,q)$ be the one-loop e^2 self-energy integrand (exchange plus seagull). Then $X^T_\varepsilon = c_X|T|^{-2}\sum_{k,q\in T^*}\hat G_M(k)^2[\sigma(k,q)-\sigma(0,q)]$, and X_ε is the same with integrals. Since $\sigma(-k,-q) = \sigma(k,q)$ and $T^* = -T^*$, σ may be replaced by $\tilde\sigma := \frac12[\sigma(k,q)+\sigma(k,-q)]$, which has $\partial_k\tilde\sigma(0,q) = 0$. The line set $\{k,q,k+q,k-q\}$ of the symmetrised integrand is *not* natural, so Proposition C.3 is applied to the two halves $\frac12\hat G_M(k)^2(1-t^1_k)\sigma(k,\pm q)$ separately. Each half has the natural line set $\{k,q,k\pm q\}$ and $\delta(\gamma)=\omega(\gamma)=1$, and the two halves sum to the symmetrised integrand pointwise. Each derivative lowers the overall degree 0 and raises none, so every term of $(\Delta_k+\Delta_q)^4$ applied to each half has $3h+\deg_H\le -1$, and its L^1 norm is at most C_4. Poisson summation in (k,q) and four integrations by parts give $|\int\int(\cdot)\mathrm{e}^{i\ell_T N\cdot(k,q)}| \le C_4(\ell_T|N|)^{-8}$. Sum over $N\neq 0$. □

Proposition H.8 (c-number differences on tori)**.** *For every pair of masses $0<\underline{M}<\overline{M}\le m_2$ with $\overline{M}\in\{m,m_1,m_2\}$, and every (α,β) with $\alpha+\beta\le\bar n$, $\sup_{K,\,\ell_T\ge\ell_0}|[\gamma_T(\overline{M}) - \gamma_T(\underline{M})]_{\alpha\beta}| < \infty$. Hence $\sup_{K,\ell_T\ge\ell_0}|\gamma_T(M)-\gamma_T(M')| < \infty$ for every pair of distinct anchor masses $M,M'\in\{M_1,m,m_1,m_2\}$: since $M_1<m<m_1<m_2$, the larger mass of the pair lies in $\{m,m_1,m_2\}$.*

The proposition uses the following results:

- Proposition C.3;
- Lemma H.13, through Lemma H.3; its step (a) is Lemma H.15;
- the Ward identities of Statements K.55 and K.56;
- the Dyson chains of Statement K.57;
- the choice $z_\varepsilon = e^2 X_\varepsilon(m)$ of the scheme (Subsection 6.1).

It gives Proposition 6.3. By Lemma H.1(a), the γ_T of that proposition is the γ_T here.

Proof. Lemma H.2 reduces the claim to the t^j-coefficients of $\mathcal{P}_t, \rho_t, \delta\dot{m}_t^2$, since $\int_0^1 t^j dt \le 1$ and the bound is two-sided. By Statement K.57(a), these coefficients are sums of weights of connected graphs. By Lemma H.3, each such coefficient is a finite sum of products of four kinds of factor, which are bounded as follows. (i) Torus sums of reduced graphs with $\omega \le -1$: by Lemma H.4. (ii) Ward defects: by Lemma H.5(b), at the uniform mass $\overline{M}$. (iii) The coefficients of $\delta m_t^2 - \delta m_0^2$ and of $\delta\dot{m}_t^2$ are zero-momentum self-energies. King's recursion [24, (3.33)] defines them through the irreducible two-point pieces, and these are the 1PI pieces by Statement K.57(c). They contain at least one $\hat{\Delta}$-line or δm^2-difference vertex, so $\omega \le -1$ by Lemma H.14(b). They are bounded by (i)–(ii), by induction on the order. (iv) The overall non-negative pieces:

- ρ_t at order 0 equals $2[C^T(\overline{M}^2) - C^\infty(\overline{M}^2)] + 2t[S_T - I_\infty](\hat{\Delta}) + 2\int(C_t - \hat{G}_m)$. We bound it by three results: Lemma H.6; Lemma H.4, since $\hat{\Delta}$ has $\omega = -1$; and the free bound $|C^\infty(M^2) - C^\infty(m^2)| \le |M-m|\pi^2/32$ (Lemma H.33), which holds uniformly in ε.
- The coefficients of $e^0\lambda^1$ of ρ_t and $\mathcal{P}_t$ vanish identically on T, for every $t \in [0,1]$. At $e = 0$ the photon decouples. By Statement K.57(b), to first order in λ we have $\mathcal{G}_t = C_t - C_t\Sigma_t C_t$. Here Σ_t is the order-λ self-energy: the λ-tadpole plus the order-λ part of δm_t^2. By translation invariance the tadpole is a multiple of $C_t(x,x) = |T|^{-1}\sum_{p\in T^*} C_t(p)$, so it is independent of the site and of the momentum. King's recursion [24, (3.33)], computed with the same lines C_t, fixes the order-λ part of δm_t^2 so that the order-λ self-energy vanishes at zero momentum. Hence $\Sigma_t = 0$ identically, so $[\mathcal{A}_t]_{01} = 0$ and $[\langle r_0^2\rangle_t]_{01} = 0$. Finally, W_ε has order 0 and $z_\varepsilon = e^2 X_\varepsilon(m)$ has order e^2.
- At order e^2, the graphs that contain a $\hat{\Delta}$-line or an f-vertex have $\omega \le -1$ (Lemma H.14(b)), and (i) and (iii) cover them. The remaining part of ρ_t is $e^2[X_\varepsilon^T(\overline{M}) - X_\varepsilon(m)] = e^2[X_\varepsilon^T(\overline{M}) - X_\varepsilon^T(m)] + e^2[X_\varepsilon^T(m) - X_\varepsilon(m)]$. Lemma H.7 bounds the second bracket.

  For the first bracket, write, as in the proof of Lemma H.7, $X_\varepsilon^T(M) = c_X\, S_k S_q\, \hat{G}_M(k)^2[\sigma_M(k,q) - \sigma_M(0,q)]$, where σ_M is the one-loop e^2 self-energy integrand at mass M and $S := |T|^{-1}\sum_{T^*}$. Put $\Delta' := \hat{G}_{\overline{M}} - \hat{G}_m = (m^2 - \overline{M}^2)\hat{G}_{\overline{M}}\hat{G}_m$ and $\Delta\sigma := \sigma_{\overline{M}} - \sigma_m$. Then

$$\frac{X_\varepsilon^T(\overline{M}) - X_\varepsilon^T(m)}{c_X} = S_k S_q\, \Delta'(\hat{G}_{\overline{M}} + \hat{G}_m)(k)\big[\sigma_m(k,q) - \sigma_m(0,q)\big] + S_k S_q\, \hat{G}_{\overline{M}}(k)^2 \Delta\sigma(k,q) - \big[S_k \hat{G}_{\overline{M}}(k)^2\big]\big[S_q \Delta\sigma(0,q)\big].$$

  In the first term the self-energy subgraph is subtracted at zero momentum. Since $\sigma_m(-k,-q) = \sigma_m(k,q)$ and $T^* = -T^*$, we have $S_q\partial_k\sigma_m(0,q) = 0$, so t^0 may be replaced by t^1. The integrand is then a Reisz forest integrand with $\delta(\gamma) = \omega(\gamma) = 1$ and overall degree -2. In the second term, $\Delta\sigma$ contains the line $\Delta'(k+q)$. The Zimmermann subspaces have degrees $-1, -3, -1$ (with q, k, resp. $k+q$ varying) and the overall degree is -2, so no subtraction is needed. The third term is a product of two one-loop sums of degree -1. All line sets are $\{k, q, k+q\}$, which is natural, and there is no two-photon subgraph. Lemma H.4 therefore bounds each term uniformly in $\varepsilon < a_0$ and $\ell_T \ge \ell_0$. The same decomposition treats $\mathcal{P}_t$, with $\hat{G}_{\underline{M}}\hat{G}_{\overline{M}} - \hat{G}_m^2 = (\hat{G}_{\underline{M}} - \hat{G}_m)\hat{G}_{\overline{M}} + \hat{G}_m(\hat{G}_{\overline{M}} - \hat{G}_m)$ in place of $\hat{G}_{\overline{M}}^2 - \hat{G}_m^2$.

(v) The $(0,0)$ coefficient: by Lemma H.2, Lemma H.6 and the same free bound. (vi) There are only finitely many coarse spacings $\varepsilon \geq \varepsilon_1$. There every quantity is a finite lattice sum of a bounded continuous integrand whose denominators are at least $\min(\underline{M}^2, \mu_*^2)$, and $|T|^{-L}\sum|R| \leq \sup|R|$ uniformly in ℓ_T. Since $\gamma_T(\overline{M}) - \gamma_T(\underline{M})$ is a polynomial in (e,λ) of order at most $\bar{n}$, the coefficient bounds give the bound on the difference. □

The two c-number bounds of Proposition 6.6. In this subsection $M \in \{m_1, m_2\}$. The slice direction is $\Gamma^B = \kappa\mu_*^2 D = \sum_j \delta_j H_j$, with $D := \sum_{j<K} t_{j+1}^2 H_j$, and $\mathcal{E}(u;M) = -|T|\, e(\varepsilon;u;M)$ is the total (not per-volume) counterterm polynomial. Put $\bar{\delta} := \kappa\mu_*^2\varepsilon_0^2$. Write $[\mathcal{E}]_{1,\alpha\beta}$ for the coefficient of $u\, e^\alpha \lambda^\beta$ in $\mathcal{E}$, $o := \alpha/2 + \beta$, and $t_K := L^K\varepsilon \in (\varepsilon_0/L, \varepsilon_0]$ for the final scale. We use the condition (E.δ), $\bar{\delta} \leq 1/12$. It gives $C_B(u) > 0$ and slice weights $1 + u\delta_j \in [1-\eta, 2]$ for $u \in [-\eta, 1]$, and it implies $\bar{\delta} \leq 1$ and $\eta\bar{\delta} \leq \frac{1}{2}$.

Lemma H.9 (the second derivative in u)**.** *Assume the hypotheses of Theorem 5.2. In particular, the conditions (E.fin) and (T.per) of Definition A.1 hold. Let $M \in \{m_1, m_2\}$ and $\varepsilon = 2^{-K}\varepsilon_0 \leq \varepsilon_0 \leq 1$. Let $\bar{n} > 12$ satisfy $2\bar{n} + 1 \leq 3R$, where R is the parameter of (4.1). For $f \in \{s, v\}$ put $M_s := M$ and $M_v := \mu_*$, with the photon taken componentwise (Lemma E.38). In scale-k units let $G_{k,f} := (-\Delta^{\eta_k} + M_f^2 t_k^2 + a_k P_k)^{-1}$ for $1 \leq k \leq K$. Assume that these kernels and their derivatives of order at most 2 satisfy (H_∞) of Subsection 5.5, with the constants C_0, δ_0 given there. Then, on every torus $T = T_j$ of side $\ell_T \geq \ell_0$,*

$$\sup_{\varepsilon} \sup_{u\in[-\eta,1]} |\partial_u^2 e(\varepsilon;u;M)| \leq E_2(M),$$

where

$$E_2(M) := C_W \sum_{m'=2}^{N_v} m'^2 \sum_{\alpha \geq 2m',\, \alpha+\beta\leq\bar{n}} e^\alpha \lambda^\beta (\kappa\mu_*^2 L^2)^{m'} \varepsilon_0^{\alpha/2+\beta+2m'-3}$$
$$= 4C_W e^4 \kappa^2 \mu_*^4 L^4 \varepsilon_0^3 [1 + O(e^2\varepsilon_0 + \lambda\varepsilon_0 + e^2\kappa\mu_*^2 L^2 \varepsilon_0^3)].$$

Here N_v is the largest number of slice lines in a graph of order $\leq \bar{n}$. The constant $C_W = C_W(\bar{n}, \alpha_0, L, a, M_b, \delta_0, M\varepsilon_0, \mu_*\varepsilon_0, \ell_0/\varepsilon_0)$ does not depend on ε, u, κ or $\ell_T \geq \ell_0$. The bound is two-sided and uniform in $u \in [-\eta, 1]$, not only at the two points of Theorem 5.2.

The lemma uses [3] in the organisation of [1, (1.12)–(1.13)], Statement K.58, and Proposition 4.7. Statement K.58 is the zero-background form of Theorem H.24(c) for the vacuum families with at least two slice lines. The lemma is the bound (6.1) of Proposition 6.6.

Proof. (1) $\partial_u^2\mathcal{E} = T_{\bar{n}}[\Phi_M''(u)]$. Expand $C_B(u) = C_{\mu_*} + u\Gamma^B$ multilinearly. Then $T_{\bar{n}}[\Phi_M(u)] = \sum_G u^{m'(G)}\mathrm{val}(G)$, summed over the connected coloured vacuum graphs of order $\leq \bar{n}$, where $m'(G)$ is the number of Γ^B-lines. Hence $\partial_u^2\mathcal{E} = \sum_{m'(G)\geq 2} m'(m'-1)u^{m'-2}\mathrm{val}(G)$.

(2) By [3, (1.25)–(1.26)], $C_{\mu_*} = G_K + (C_{\mu_*} - G_K)$. The second part is regular and is treated as an external field. The slice part Γ^B is exactly c_K times the rescaled slice kernel, with $c_K = \kappa\mu_*^2 L^2 t_K^2$.

(3) Rescale to unit t_K.

(4) The graphs fall into the families of Lemma H.23. Each family has a fixed number m' of Γ^B-lines and a fixed order $e^\alpha\lambda^\beta$, where the orders of the counterterm vertices are included. No family with $m' \geq 2$ needs a vacuum counterterm, because its degree is at least 3 (Lemma H.18(c)). By Statement K.58, the coefficient of $u^{m'}$ of such a family is at most $C_F e^\alpha \lambda^\beta (\kappa\mu_*^2 L^2)^{m'} \varepsilon_0^{\alpha/2+\beta+2m'-3}|T|$. Each slice line has two e-legs, so $\alpha \geq 2m'$, and the exponent of ε_0 is at least $3m' - 3 \geq 3$.

(5) Multiply by $m'(m'-1)|u|^{m'-2} \leq m'^2$ (because $|u| \leq 1$), sum over the finitely many families, and divide by $|T|$. The constant C_W is the largest C_F times the number of families. It depends only on the data listed in Statement K.58.

The bare coefficient is $a_{K,T}(M)$. By Proposition 4.7, it is the reference counterterm of the organisation [1, (1.12)–(1.13)] at mass M on T.

The inputs of step (4). Step (4) uses Statement K.58 in place of Theorem H.24(c). The sketched items I1–I5 listed after Theorem H.24 enter only in the following way.

- The expansion of $\mathcal{E}(u;M)$ has no external fields, zero background and no block-spin transformation. Its vertices are those of the lattice action [3, (1.6)–(1.8), (1.10)] and the mass-counterterm vertices [3, (1.7)]. The vertices [3, (1.9), (1.11)] are Taylor remainders, so they do not occur in a Taylor polynomial. The vertices [3, (1.12)–(1.15)] arise only from a block-spin transformation. Hence the Hölder norms of the background (I5) do not occur. Since $\mathcal{E}(u;M)$ is one number on T, the relocalisation of the vacuum counterterms between scales (I2) does not occur either, and no vacuum counterterm is attached (Theorem H.24(b)).
- The line bounds (I4) for King's slices $H_{j,f} = G_{j+1,f} - G_{j,f}$ are given by Corollary E.49. It applies because, by (E.fin), $t_{j+1} \le t_K \le \varepsilon_0 \le \min(1/M, 1/\mu_*, L)$. By Lemma H.22, King's slices are the slices of [3, (2.6)]. The remainder lines $C_f - G_{K,f}$ obey Proposition E.50.
- The finite counterterms (I3) are bounded by Statement K.59.
- The combinatorics of the families (I1; step S10 of Table 2) enters only through Statement K.58. On the torus, the counterterm partners of the classes are compared with their form on $\eta_K\mathbb{Z}^3$ by Statements K.60 and K.61. These comparisons take place on cubes that do not wrap around the torus, by (T.per) and Lemma 5.22.

□

Lemma H.10 (the one-loop mass difference of the term linear in u)**.** *Let* $\Pi_M(k)_{\mu\nu} := e^2\big[2\delta_{\mu\nu}S_q\cos(\varepsilon q_\mu)G(q) - S_qV_\mu V_\nu G(q)G(q+k)\big]$, *with* $V_\mu := (2/\varepsilon)\sin(\varepsilon(2q+k)_\mu/2)$, $G(q) := (\hat{q}^2 + M^2)^{-1}$ *and* $S_q := |T|^{-1}\sum_{q\in T^*}$. *This is the order-*$e^2$ *vacuum polarisation of the complex scalar on* T_ε *with King's hopping, and* $[\partial_u e(\varepsilon;0;M)]_{20} = \frac{1}{2}|T|^{-1}\operatorname{Tr}(\Gamma^B\Pi_M)$. *For* $0 < m_1 < m_2$,

$$\tfrac{1}{2}|T|^{-1}\big|\operatorname{Tr}\big(\Gamma^B(\Pi_{m_1} - \Pi_{m_2})\big)\big| \le 10e^2(m_2 - m_1)\sup_{M\in[m_1,m_2]}[M\,C_T(M,\ell_0)]\cdot 3\kappa\mu_*^2L^2\Big(c_1\varepsilon + \frac{C_H\varepsilon_0}{L-1}\Big).$$

Here $C_T(M,\ell) := a_*^3\sqrt{\pi}/M + 3a_*^2b/M^2 + \frac{3\sqrt{\pi}}{2}a_*b^2/M^3 + b^3/M^4$, *with* $a_* := \sqrt{\pi}/4$ *and* $b := 4/\ell$. *The constants* c_1, C_H *are those of the diagonal bound on the slice kernels (Lemma H.27).*

The lemma uses the diagonal bound of Lemma H.27 and the heat-kernel bound of Corollary H.32.

Proof. (1) For $\Gamma \ge 0$ and bounded self-adjoint B, $|\operatorname{Tr}(\Gamma B)| \le \|B\|\operatorname{Tr}\Gamma$. Here $B = \Pi_{m_1} - \Pi_{m_2}$ is a Fourier multiplier, so $\|B\| = \sup_k\|B(k)\|_{\mathrm{op}}$. (2) $|T|^{-1}\operatorname{Tr}\Gamma^B \le 3\kappa\mu_*^2\max_x D(x,x) \le 3\kappa\mu_*^2L^2(c_1\varepsilon + C_H\varepsilon_0/(L-1))$ (Lemma H.27, $H_j(x,x) \le C_H/t_j$). (3) Differentiating, $\partial_M\Pi_M = e^2[-4M\delta_{\mu\nu}S_q\cos(\varepsilon q_\mu)G^2 + 2MS_qV_\mu V_\nu(G(q)^2G(q+k) + G(q)G(q+k)^2)]$. Write $\mathsf{A} := \hat{q}$ and $\mathsf{B} := \widehat{q+k}$. The sine addition formula gives $|V_\mu| \le |\mathsf{A}_\mu| + |\mathsf{B}_\mu|$, so $|V|^2 \le 2(\mathsf{A}^2 + \mathsf{B}^2)$. Since $\mathsf{A}^2G_{\mathsf{A}}, \mathsf{B}^2G_{\mathsf{B}} \le 1$, $|V|^2(G_{\mathsf{A}}^2G_{\mathsf{B}} + G_{\mathsf{A}}G_{\mathsf{B}}^2) \le 4(G_{\mathsf{A}}^2 + G_{\mathsf{B}}^2)$. By shift invariance of S_q, $\|\partial_M\Pi_M(k)\|_{\mathrm{op}} \le 20Me^2S_qG(q)^2$. (4) $S_qG^2 = \int_0^\infty te^{-tM^2}S_qe^{-t\hat{q}^2}dt$, with $S_qe^{-t\hat{q}^2} \le (a_*/\sqrt{t} + 4/\ell_T)^3$ (Corollary H.32, which does not depend on ε). Integrating term by term gives $S_qG^2 \le C_T(M,\ell_T) \le C_T(M,\ell_0)$. (5) Write $\Pi_{m_1} - \Pi_{m_2} = -\int_{m_1}^{m_2}\partial_M\Pi_M\,dM$, bound by (3)–(4), and combine with (1), (2) and the factor $\frac{1}{2}$. □

Proposition H.11 (the term linear in u)**.** *Assume the hypotheses of Lemma H.9 at $M = m_1$ and at $M = m_2$. Then $\bar\Gamma^{(1)}_{21} < \infty$. More precisely,*

$$\bar\Gamma^{(1)}_{21} \le \sup_{K,\ell_T\ge\ell_0} [\gamma_T(m_2) - \gamma_T(m_1)] + \sum_{\substack{(\alpha,\beta)\neq(2,0),\,\alpha\ge2\\ \alpha+\beta\le\bar n}} e^\alpha\lambda^\beta\big(C_{1,\alpha\beta}(m_1) + C_{1,\alpha\beta}(m_2)\big)\kappa\mu_*^2L^2\varepsilon_0^{o-1} + \mathfrak{C}_{\mathrm{CU1}} + \tfrac12\big(E_2(m_1) + E_2(m_2)\big).$$

Here $\mathfrak{C}_{\mathrm{CU1}}$ is the right side of Lemma H.10, taken at $\varepsilon \le \varepsilon_0$, and the $C_{1,\alpha\beta}$ are the constants of the bound on the vacuum families linear in u (Corollary H.26).

The proposition uses [3] in the organisation [1, (1.12)–(1.13)], Proposition H.8, Lemma H.9, Lemma H.10 and Proposition 4.7. It is the bound (6.2) of Proposition 6.6.

Proof. Write $a_i := a_{K,T}(m_i)$. By Lemma E.5, $u \mapsto e(\varepsilon; u; M)$ is a polynomial, so

$$e(\varepsilon;1;M) = e(\varepsilon;0;M) + \partial_u e(\varepsilon;0;M) + \int_0^1 (1-u)\,\partial_u^2 e(\varepsilon;u;M)\,du.$$

Insert this into the bracket that defines $\bar\Gamma^{(1)}_{21}$ and treat the three parts in turn.

Order u^0. By Lemma H.1(c) with $(M, M') = (m_2, m_1)$, the u^0 part is $\gamma_T(m_2) - \gamma_T(m_1)$. Its supremum over the family is finite by Proposition H.8 with the pair $\{m_1, m_2\}$.

Remainder. By Lemma H.9, the remainder is at most $\frac12 E_2(M)$ for each M.

Order u^1. We bound $|\partial_u e(\varepsilon;0;m_1) - \partial_u e(\varepsilon;0;m_2)|$ coefficient by coefficient.

- Odd α: these coefficients vanish. The map $(e, A) \mapsto (-e, -A)$ leaves the action and the centred Gaussian invariant, so Φ_M is even in e.
- $\alpha = 0$: these coefficients vanish. At $e = 0$ the matter integral does not depend on A, so $\Phi_M(u)|_{e=0}$ does not depend on u.
- $\alpha \ge 2$ with $(\alpha, \beta) \neq (2, 0)$: then $o \ge 2$. The bound $|[\mathcal{E}]_{1,\alpha\beta}| \le C_{1,\alpha\beta}\kappa\mu_*^2L^2|T|\varepsilon_0^{o-1}$ (Corollary H.26, an instance of Theorem H.24(c)) bounds each mass separately, per unit volume.
- $(\alpha, \beta) = (2, 0)$: this is the one-loop graph. It diverges logarithmically at each M, and Lemma H.10 bounds its mass difference. As in that lemma, the Taylor coefficient is taken with the scalar lines at the bare mass M^2; the counterterms enter as vertices of higher order.

All bounds are uniform in ε and in $\ell_T \ge \ell_0$, and all sums are finite.

The constants c_1 and C_H in Lemma H.10 come from step (2) of its proof, which is Lemma H.27. That lemma rests on two inputs: the block Poincaré inequality (Lemma I.11), and the bound $C^{(j)} \le c_C t_j^2$ on King's fluctuation covariance (Lemma H.28). The second follows from the variational form of $\Delta^{(j)}$ (Lemma H.20). By Lemma H.29, one may take $c_1 = 4.42 + \varepsilon_0/(\ell_0^3\mu_*^2)$. Hence $\mathfrak{C}_{\mathrm{CU1}}$ is finite and does not depend on $\varepsilon \le \varepsilon_0$ or $\ell_T \ge \ell_0$. □

H.1. **Power counting, the weighted-line form of Proposition 1, and kernel bounds.** *Notation.* Notation is as at the beginning of this appendix. In addition: $t_k := L^k\varepsilon$; $\eta := L^{-k}$ is the lattice spacing in scale-k units; $c_k := \kappa\mu_*^2L^2t_k^2$; $\delta_j := \kappa\mu_*^2t_{j+1}^2$, so that $\Gamma^B = \kappa\mu_*^2D = \sum_{j<K}\delta_jH_j$ with $D := \sum_{j<K} t_{j+1}^2H_j$. H_j is King's photon slice [24, (2.17)]. $G^\eta_{(j)}$ is Bałaban's slice [3, (2.6), p. 424]. For a polynomial F in (u, e, λ), $[F]_{m,\alpha\beta}$ is its $u^me^\alpha\lambda^\beta$ coefficient, and $o := \alpha/2 + \beta$. We write $\mathcal{E}(u; M) = -|T|\,e(\varepsilon; u; M) = T_{\bar n}[\Phi_M(u)]$ (Lemma H.1(b)). All bounds below are on absolute values, so signs play no role.

Lattice power counting: the transfer from $d = 4$ to $d = 3$.

Lemma H.12 (where the dimension enters Reisz's proofs)**.** *Let* d *be the space-time dimension. In the proofs of* [32, Thm. 1, Thm. 2, Thm. 3] *(Sects. 4–9, App. A–D) and of the Renormalization Theorem* [33, p. 92] *(Sects. 3–6, App. A–C), the value* $\mathsf{d} = 4$ *is used at exactly the following places and nowhere else:*

(i) *the degree definitions, each of the form "*$4\times$*(dimension of the loop space)":* [32, (3.8), (5.9), (7.9), (7.14), (9.3), (9.29)] *and* [33, (2-28), (4-11)–(4-16), (5-5), (5-10), (C-1)–(C-2)]*;*

(ii) *the single scaling step* [32, (9.16)–(9.17a)]*, where the inner variables are rescaled by* $\|t_1\|$*;*

(iii) *two constants that use* $\sqrt{\mathsf{d}} \le 2$*: δ in* [32, (7.8)]*, and step 1 of the proof of* [32, Lemma D.1] *(p. 125).*

Replacing 4 *by* d *in (i), the proofs remain valid for every* $\mathsf{d} \le 4$*, and in particular for* $\mathsf{d} = 3$*. They then give Proposition C.3(a)–(b) in the bound (absolute-value) form.*

The proofs of [32], Sects. 4–9 and App. A–D, and of [33], Sects. 3–6 and App. A, C, are used as printed; the absolute-value form is read off [32, (7.4)–(7.5)]. Reisz's printed theorems are for $\mathsf{d} = 4$ ("The following considerations can easily be extended to other dimensions", [32, p. 86]). For $\mathsf{d} = 3$ the lemma is a reading of a published proof, not a new proof.

Proof. We go through the items. (1) *(i): the degree definitions.* $\deg_H \hat{I} = 4\dim H + \deg F$ [32, (3.8)]; $\omega(H) = 4\dim H + \delta(H) - \deg E$ (5.9); the counts (7.9), (7.14); $\omega_0 = 4m + \Delta - 2\sum n_j$ (9.3); and (9.29). In [33]: $\omega(\gamma) = \sum\omega(\Delta_L) + \sum\omega(V_B) + 4m(\gamma)$ (2-28)/(2-31); (4-11)–(4-16); $M_U(\gamma) = 4\sum m(\tau(U))$ (5-10). Each is "$4\times$ the number of 4-dimensional integration variables", and we replace 4 by d. (2) *App. C.* Reisz states and uses a *lower* bound: $M_U(\Gamma) \ge 4h$ (C-1) and $M_U(\gamma) \ge 4\,\mathrm{rank}$ (C-2) [33, pp. 97, 106]. Its proof is the count $M_U = 4\cdot\#\mathcal{T}^* \ge 4\,\mathrm{rank}$, which is linear in the 4. Hence $M_U(\Gamma) \ge \mathsf{d}h$, and this is what gives $\mathsf{d}h + \deg_H R_\Gamma \le -1$ in (5-13). (3) *(ii): the scaling step.* The sector X_ξ (9.8) has outer variable $t_1 \in \mathbb{R}^{\mathsf{d}}$ with $\|t_1\| \ge 1$; the inner variables $(t_2,\dots,t_m) \in \mathbb{R}^{\mathsf{d}(m-1)}$ are rescaled as $t_j = \|t_1\|t'_j$ (9.16). The Jacobian contributes $\|t_1\|^{\mathsf{d}(m-1)}$. The numerator contributes at most $\|t_1\|^{\Delta}$ (by (9.19), $r_i - p_i \le \Delta$), and the homogeneous denominator $\|t_1\|^{-2\sum n_j}$. So (9.17a) becomes $\int d^{\mathsf{d}}t_1 \|t_1\|^{\omega_0-\mathsf{d}}\,\mathcal{J}^{\mathrm{int}}$ with $\omega_0 := \mathsf{d}m + \Delta - 2\sum n_j$. Lemma 9.2 applies the induction hypothesis to the inner integral, with $\omega(\tilde{H}) = \omega(H)$ (9.29). Reisz does not print the outer integral ("the remaining integration can be done without problems", p. 119). In polar coordinates, $r^{\mathsf{d}-1}r^{\omega_0-\mathsf{d}} = r^{\omega_0-1}$, and $\int_1^{\lambda+K} r^{\omega_0-1}f(\lambda')\,dr$ with $\lambda' = b(\lambda+d)/r + e$ (Lemma 9.1) does not depend on d:

- $f = 1$, $\omega_0 < 0$: the integral is bounded;
- $f = \lambda'^{-1}\log^{m-1}\lambda'$, $\omega_0 \le -1$: it is at most $\lambda^{-1}\log^m\lambda$;
- $f = \lambda'^{\bar{\omega}}\log^{m-1}\lambda'$: it is at most $\lambda^{\max(\omega_0,\bar{\omega})}\log^m\lambda$.

This is [32, (9.30)] for every d. (4) *(iii): the two constants.* In (7.8), $\delta = \max(\pi\varepsilon, 4\pi(1 + \|\delta_i\|/2))$ must make κ_J contain the translated sector. From (4.7), $\|k_j\| \le (\pi/a)(\sqrt{\mathsf{d}} + 2\|\delta_j\|)$, which suffices if $\sqrt{\mathsf{d}} \le 4$. Lemma D.1, step 1, uses $\|k_j\| \le 2\pi/a = \sqrt{4}\,\pi/a$. Both hold a fortiori for $\mathsf{d} = 3$; for $\mathsf{d} > 4$ they fail, which is why the lemma is restricted to $\mathsf{d} \le 4$. (5) *Dimension-free steps.* The following involve no d:

- the function classes and propagator class, Defs. 2.1–2.5 and Lemma 2.1 (vanishing orders in w; King's $\hat{l}^2 = \sum_\mu (4/a^2)\sin^2(al_\mu/2)$ is in Def. 2.5 for $\mathsf{d} = 3$);
- naturalness, Def. 3.1 and App. D (linear algebra over $\mathbb{Z}$, coordinatewise);
- the sector decomposition (4.1)–(4.9) and the propagator bounds (4.8)–(4.9);
- Theorem 3 and Lemmas 6.1–6.3 (Taylor's formula in groups $v_i \in \mathbb{R}^{m_i}$);
- the powers of a in (7.10)–(7.17) (twice the number of propagators far from a pole);
- Lemmas 8.1–8.10 and App. A–B (interpolation, set inclusions, linear changes of variables, Fubini);

- in [33]: Lemma 3.1 and App. A (subtraction operators; t^0, t^1 with $\sin(\varepsilon q)/\varepsilon$ are of type (A-1)), Lemma 5.1 (Zimmermann's complete forests), and Sect. 6, the index distributions (finite index sets).

Masses enter only through lower bounds (all $\mu_i^2 > 0$), so $\mathsf{d} = 3$ needs no extra infrared condition. Two further items are dimension-free: [33, Sect. 6], and complex-valued numerators. The latter are harmless because $|V| \le |\mathrm{Re}\, V| + |\mathrm{Im}\, V|$ and real and imaginary parts have vanishing orders at least those of V. (6) *Absolute-value form.* (4.3) is a partition into finitely many sectors by non-negative Θ's. (7.4) and (7.5a–b) bound each sector by integrals of non-negative majorants, and the integrands of Theorem 2 (5.2) are non-negative. So the proof bounds $\int_{\mathrm{BZ}} |F|$, uniformly for bounded external momenta (p. 96, p. 102). Reducing to $V \in \mathcal{C}^c_{m_0}$ is harmless, because $|V| \le \sum |V_i|$. □

The counterterm–forest bookkeeping.

Lemma H.13 (King's expansion as Reisz forest integrals)**.** *On $\varepsilon\mathbb{Z}^3$ take the C_t-model of the paragraph on the c-number differences: scalar lines $C_t = \hat{G}_{\overline{M}} + t\hat{\Delta}$, and $\delta m_t^2 = \delta m_0^2 + f_t$ with $f_t := \delta m_t^2 - \delta m_0^2$, a polynomial in t with $f_0 = 0$. Then for every (α, β, j), the $e^\alpha \lambda^\beta t^j$ coefficient of $\mathcal{P}_t$, of ρ_t (minus its W_ε, z_ε terms) and of f_t is a finite sum $\sum_G c_G \prod(\text{lower-order coefficients of } f) \int_{\mathrm{BZ}^L} R_G$, with rational c_G. Here:*

- *G runs over 1PI graphs with scalar lines $\hat{G}_{\overline{M}}$ or $\hat{\Delta}$, photon lines, the vertices of King's action, the insertion, and two-point "f-vertices". There are no δm_0^2 vertices.*
- *R_G is Reisz's forest integrand with $\delta(\gamma) = \omega(\gamma)$ for every 1PI $\gamma \subsetneq G$. This means t^1 (in $\sin(\varepsilon q)/\varepsilon$) if $\omega = 1$, t^0 if $\omega = 0$, and no subtraction if $\omega < 0$. There is no overall subtraction.*

The identity holds exactly at every ε. Its torus form, with the Ward-defect correction, is Lemma H.3.

Step (a) of the proof is Lemma H.15 below. It identifies the insertion of King's constant counterterms with the sum over t^0-forests (Bogoliubov's R-operation for zero-momentum counterterms, in Zimmermann's forest form; cf. [33, pp. 79–80, 94–95]). The regrouping of the two-photon parts in step (c) rests on the Ward identity on $\varepsilon\mathbb{Z}^3$, Statement K.56.

Lemma H.14 (the one-particle-irreducible parts with $\omega \ge 0$)**.** *Let γ be a connected 1PI graph with a loop. It is built from the following vertices:*

- *King's vertices V_n, $n \ge 1$, of Lemma H.15;*
- *the λ-vertex;*
- *two-point vertices: the f-vertices, the vertex inside each $\hat{\Delta}$-line, and at most one insertion.*

It contains no δm_0^2-vertex. Let E_φ and E_A be its numbers of external scalar and photon legs, and put $E := E_\varphi + E_A$. Let k be its number of $\hat{\Delta}$-lines, v_f its number of f-vertices and $v_X \in \{0, 1\}$ its number of insertions. Put $o := o_V(\gamma) := \sum_n \frac{n}{2} v_n + v_\lambda$. Its degree is

$$\omega(\gamma) = 3 - \tfrac{1}{2}E - o - 2(k + v_f + v_X).$$

(a) *If $\omega(\gamma) \ge 0$, then γ belongs to exactly one of the following classes:*
- R0 *vacuum graphs, $E = 0$, $o \le 3$, $\omega = 3 - o$;*
- R1 *vacuum graphs with one insertion and $k = v_f = 0$, as the whole graph only: $o = 0$ ($\omega = 1$) or $o = 1$ ($\omega = 0$);*
- R2 *scalar self-energies, $E_\varphi = 2$, $E_A = 0$: $o = 1$ ($\omega = 1$) or $o = 2$ ($\omega = 0$);*
- R3 *photon self-energies, $E_A = 2$, $E_\varphi = 0$: $o = 1$ ($\omega = 1$) or $o = 2$ ($\omega = 0$);*
- R4 *$\varphi\varphi A$ parts, $o = \frac{3}{2}$, $\omega = 0$;*
- R5 *three-photon parts, $o = \frac{3}{2}$, $\omega = 0$;*
- R6 *one-photon parts, $E_A = 1$, $E_\varphi = 0$.*

In R2–R6 one has $k = v_f = v_X = 0$.

(b) *A graph of ρ_t or $\mathcal{P}_t$ has $\omega = 1 - o - 2(k + v_f)$. A zero-momentum self-energy graph with $k + v_f \ge 1$ (a graph of f_t or of $\delta\dot{m}_t^2$) has $\omega = 2 - o - 2(k + v_f) \le -1$.*

Proof. Each line has degree -2. The vertex function of V_n is $e^n\varepsilon^{n-2}$ times a trigonometric polynomial in ε times the momenta, so its degree is at most $2-n$. The λ-vertex and every two-point vertex have degree 0. A $\hat\Delta$-line is two lines with the same momentum joined by a two-point vertex. Reisz's degree [33, (2-28)], with 3 in place of 4, is $\omega = 3L - 2I + \sum_v \deg(v)$. With $L = I - V + 1$ and $2I + E = \sum_v \mathrm{legs}(v)$ this gives the displayed formula: the vertex contributions $\frac{1}{2}\mathrm{legs}(v) - 3 + \deg(v)$ are $-\frac{n}{2}$, -1 and -2, respectively.

Every vertex has 2 or 4 scalar legs, so E_φ is even. Every photon line has two photon ends, so $\sum_n n v_n \equiv E_A \pmod 2$; hence $o \in \frac{1}{2} + \mathbb{N}$ if and only if E_A is odd. With $o = 0$ only two-point vertices occur, and they form no loop with external legs. A single V_1 has three legs. So a loop with $E \geq 1$ has $o \geq 1$, or $o = \frac{1}{2}$ with $E = 1$. $E = 0$: γ is a whole graph. This gives R0 or R1, since $\omega = 1 - o - 2(k + v_f)$ when $v_X = 1$. $E = 1$: then $E_A = 1$, which gives R6. $E = 2$: $(E_\varphi, E_A) \in \{(2,0),(0,2)\}$, so o is an integer ≥ 1. Then $\omega = 2 - o - 2(k + v_f + v_X) \geq 0$ forces $k = v_f = v_X = 0$ and $o \in \{1,2\}$. This gives R2 and R3. $E = 3$: $(E_\varphi, E_A) \in \{(2,1),(0,3)\}$, so $o \in \{\frac{1}{2}, \frac{3}{2}, \dots\}$. A loop needs $o \geq \frac{3}{2}$, and $\omega = \frac{3}{2} - o - 2(k + v_f + v_X) \geq 0$ forces $o = \frac{3}{2}$ and no two-point vertices. This gives R4 and R5. $E = 4$: $\omega \geq 0$ forces $o \leq 1$ and no two-point vertices. The vertex sets $\{V_1, V_1\}$, $\{V_2\}$, $\{\lambda\}$ with four external legs are trees. $E \geq 5$: $\omega \leq \frac{1}{2} - o < 0$ for $o \geq 1$, and $o = \frac{1}{2}$ gives a tree. (b) A graph of ρ_t or $\mathcal{P}_t$ is a vacuum graph with one insertion. A zero-momentum self-energy has $E_\varphi = 2$, $E_A = 0$ and $v_X = 0$, and it has integer $o \geq 1$ by the above. □

Lemma H.15 (zero-momentum counterterms as forests)**.** *Work in the C_t-model on T or on $\varepsilon\mathbb{Z}^3$. Let $J_0 := \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ act on the two real components of φ, and let $U(A) := \mathrm{e}^{e\varepsilon A J_0}$. The vertices are of the following kinds:*

- *King's vertices $V_n(A,\varphi) = -\frac{e^n\varepsilon^{n-2}}{n!}\sum_{x,\mu}\varepsilon^3 A^n_{x,\mu}\,\mathrm{Re}[i^n\bar\varphi(x)\varphi(x+\varepsilon e_\mu)]$, $n \geq 1$, of order $o(V_n) := \frac{n}{2}$, and the vertex $\lambda\sum_x \varepsilon^3|\varphi|^4$, of order 1;*
- *two-legged counterterm vertices c_o, $o \in \{1,2\}$, with coupling $-\frac{1}{2}\delta m_0^{2,(o)}$, where $\delta m_0^{2,(o)}$ is the sum of the $e^\alpha\lambda^\beta$ terms of δm_0^2 with $\frac{\alpha}{2} + \beta = o$;*
- *two-legged f-vertices;*
- *an insertion vertex X. For ρ_t, X is r_0^2. For $\mathcal{P}_t$, X is the pair $\hat G_{\underline M}\hat G_{\overline M}$, joined to the rest by two scalar legs of equal momentum.*

For a finite multiset ν of kinds put $\nu! := \prod_\kappa \nu_\kappa!$. A graph of content ν consists of two pieces of data:

- *a perfect matching of the legs of the labelled vertices (κ, k), $k \leq \nu_\kappa$;*
- *scalar lines coloured $\hat G_{\overline M}$ or $\hat\Delta$, and photon lines C_{μ_*}.*

$A(G) = \int I_G$ is the product of the couplings times the lattice sum, resp. Brillouin-zone integral, of its line and vertex factors. Here I_G is the integrand, and $\int$ is $|T|^{-L}\sum_{(T^*)^L}$ on T and $\int_{\mathrm{BZ}^L}$ on $\varepsilon\mathbb{Z}^3$.

A block *of G is a set S of King vertices with the following properties:*

1. *every line joining two vertices of S is a $\hat G_{\overline M}$ or C_{μ_*} line;*
2. *the induced graph $G[S]$ is one-particle irreducible;*
3. *exactly two legs at S are not matched inside S, and both are scalar;*
4. *$o(S) := \sum_{v\in S} o(v) \leq 2$.*

A forest *is a set of blocks any two of which are nested or disjoint. For a forest F put $A_F(G) := \int \prod_{\gamma\in F}(-t^0_\gamma) I_G$, where t^0_γ sets the external momentum of γ to 0 and the operators of nested elements act from the inside out. For a two-point graph γ, $\hat A_F(\gamma)(0)$ is the 2×2 matrix obtained in the same way at external momentum 0. Let $\mathcal{C}$ be one of the following classes:*

- *connected graphs containing X once (for ρ_t, $\mathcal{P}_t$);*

- *one-particle irreducible amputated two-point graphs at zero momentum with at least one $\hat{\Delta}$-line or f-vertex (for f_t).*

Let $\mathcal{C}_0 \subset \mathcal{C}$ be the graphs without c_o-vertices. Then, for every order,

$$\sum_{\mu} \frac{1}{\mu!} \sum_{H \in \mathcal{C}(\mu)} A(H) = \sum_{\nu} \frac{1}{\nu!} \sum_{G \in \mathcal{C}_0(\nu)} \sum_{F} A_F(G), \tag{H.1}$$

where F runs over the forests of G.

Proof. Step 1 (the counterterm is scalar). Let $\Gamma^{(o)}$ be King's order-o amputated one-particle-irreducible two-point function of the uniform-mass model, excluding the single counterterm vertex. Its zero-momentum value $\hat{\Gamma}^{(o)}(0)$ is a real 2×2 matrix. The action and the Gaussian are invariant under two maps:

- $\varphi \mapsto \mathrm{e}^{\theta J_0}\varphi$;
- $\varphi \mapsto P\varphi$, $A \mapsto -A$, with $P = \mathrm{diag}(1, -1)$, since $PU(A)P = U(-A)$.

So $\hat{\Gamma}^{(o)}(0)$ commutes with $\mathrm{e}^{\theta J_0}$, hence equals $a\mathbf{1} + bJ_0$. It also commutes with P, hence $b = 0$. By [24, (3.33)], $\hat{\Gamma}^{(o)}(0) = \delta m_0^{2,(o)}\mathbf{1}$.

Step 2 (factorisation). Let S be a block. Its kernel $K_S(z, z')$ between the sites of its two free legs is translation invariant. Hence $(t_S^0 I_G)$ is the integrand of G/S, which has at the place of S a two-legged vertex of value $\hat{K}_S(0) = \sum_{z'} \varepsilon^3 K_S(z, z')$. After Reisz's routing S_γ [33, (2-23)], the internal momenta of S are independent variables, and the integrand is bounded on the compact zone. By Fubini, $\int (-t_S^0) I_G = -\hat{K}_S(0) \cdot A(G/S)$. The same holds recursively for nested blocks. On T the same argument works with sums over $T^* \ni 0$.

Step 3 (substitution). Let $H \in \mathcal{C}(\mu)$ have counterterm vertices $v_1, \dots, v_r$, with r_o of them of order o. For each v_i choose a counterterm graph γ_i and a forest F_i of proper blocks of γ_i. Here a counterterm graph is a one-particle-irreducible uniform-mass two-point graph of content κ_i and order $o(v_i)$ without c-vertices. Delete v_i and match the free legs x, y of γ_i with the partners of legs $1, 2$ of v_i. Then distribute the labels: there are $\prod_\kappa \nu_\kappa!/(\mu_\kappa! \prod_i \kappa_{i,\kappa}!)$ ways, with $\nu = \mu_K + \sum_i \kappa_i$. Each way gives a pair (G, F) with $G \in \mathcal{C}_0(\nu)$ and $F = \{S_i\} \cup \bigcup_i F_i$, where S_i is the vertex set of γ_i. Indeed, $G[S_i] = \gamma_i$. Contracting a connected one-particle-irreducible part preserves connectedness, one-particle irreducibility and the presence of X, $\hat{\Delta}$ or f. And the S_i are the maximal elements of F. Conversely, contracting the maximal elements of a forest recovers $(H, (\gamma_i, F_i))$. This uses three facts:

- $o(S_i) \in \{1, 2\}$, because $E_A = 0$ forces the order to be an integer;
- a proper sub-block has strictly smaller order;
- the blocks of G inside S_i are exactly the blocks of γ_i.

This reconstruction has $2^r \prod_o r_o!$ preimages, coming from the names x, y of the free legs of each block and from the labels of the c_o-vertices. By Step 2, replacing the coupling $-\frac{1}{2}\delta m_0^{2,(o_i)}\mathbf{1}$ of v_i by $-\frac{1}{2}\hat{A}_{F_i}(\gamma_i)(0)$ turns $A(H)$ into $2^{-r}A_F(G)$. Since $\mu! = \prod_\kappa \mu_\kappa! \prod_o r_o!$,

$$\frac{1}{\mu!} \prod_i \frac{1}{\kappa_i!} \cdot 2^{-r} \cdot \frac{2^r \prod_o r_o!}{\prod_\kappa \nu_\kappa!/(\mu_\kappa! \prod_i \kappa_{i,\kappa}!)} = \frac{1}{\nu!}.$$

Step 4 (induction on o). Claim: $\delta m_0^{2,(o)}\mathbf{1} = \sum_\kappa \frac{1}{\kappa!} \sum_\gamma \sum_{F \text{ proper}} \hat{A}_F(\gamma)(0)$, where the sum runs over the counterterm graphs γ of order o. For $o = 1$ the graphs of $\Gamma^{(1)}$ contain no counterterm vertex and no proper block. For $o = 2$ the graphs of $\Gamma^{(2)}$ contain only vertices c_1. Insert the case $o = 1$ and apply Step 3 to the class of $\Gamma^{(2)}$; Step 1 converts the matrix sum into $\delta m_0^{2,(2)}\mathbf{1}$. Insert the claim into every c-vertex of every $H \in \mathcal{C}(\mu)$ and apply Step 3 once more. This gives (H.1).

Step 5 (f_t). By [24, (3.33)] with lines C_t, $\delta m_t^{2,(o)} = \hat{\Gamma}_t^{(o)}(0)$. Expand $C_t = \hat{G}_{\overline{M}} + t\hat{\Delta}$ multilinearly and $\delta m_t^2 = \delta m_0^2 + f_t$ at lower orders. The terms with only $\hat{G}_{\overline{M}}$-lines and only δm_0^2-vertices form the right side of King's recursion at $\overline{M}$, which is $\delta m_0^{2,(o)}$. So $f_t^{(o)}$ is the sum over the second class $\mathcal{C}$, to which (H.1) applies. □

Remark H.16. For a scalar two-point part with no two-point vertices, $\omega = 2 - o$ (Lemma H.14). Hence the blocks are exactly the proper one-particle-irreducible scalar two-point parts with $\omega \geq 0$ that contain no $\hat{\Delta}$-line, no f-vertex and not the insertion; these are the R2 parts. A subdiagram with two external scalar legs whose vertex set is S contains every line of $G[S]$, since an omitted line adds two external legs. So the forests above are Reisz's forests [33, p. 87] restricted to these parts, with t^0 (Lemma H.17). The right side of (H.1) is $\sum_\nu \frac{1}{\nu!} \sum_G \int R_G^{(a)}$, where $R_G^{(a)}$ is the forest integrand [33, (4-2)].

Lemma H.17 (line-disjoint blocks are vertex-disjoint)**.** *In the setting of Lemma H.15, let γ_1, γ_2 be one-particle-irreducible subgraphs of G with the following properties: each has exactly two external legs, both scalar; neither contains the insertion; and they have no line in common. Then γ_1 and γ_2 have no vertex in common. In particular, the forests of Lemma H.15 (nested or vertex-disjoint blocks) are the same as the forests of two-point parts in the sense of* [33, p. 87] *(nested or line-disjoint), restricted to the blocks.*

Proof. Suppose γ_1, γ_2 have a common vertex v, of degree d. Let a_i be the number of legs at v that lie on lines of γ_i. Since γ_i is one-particle irreducible, $a_i \geq 2$. The $d - a_i$ remaining legs at v are external to γ_i, so $d - a_i \leq 2$. Line-disjointness gives $a_1 + a_2 \leq d$. Hence $d = 4$ and $a_1 = a_2 = 2$. Then every external leg of γ_1 lies on a line of γ_2, and conversely. So $\gamma_1 \cup \gamma_2$ is a connected component without external legs, and therefore it is all of G. This is impossible: every graph considered has either two external legs or the insertion, and the insertion lies in neither γ_i. □

Proof of Lemma H.13. The degrees are $\omega(\gamma) = 3 - \frac{1}{2}(E_\varphi + E_A) - o_V - 2v_2$. Lemma H.14 lists the parts with $\omega \geq 0$ in the classes R0–R6. Those that occur as proper parts are:

- R2, the scalar self-energies;
- R3, the two-photon parts of order $o \in \{1, 2\}$;
- the types R4 and R5, whose zero-order forest terms are odd at zero external momentum;
- R6, the parts with one photon leg.

None of them contains a $\hat{\Delta}$-line, an f-vertex or the insertion. (a) *δm_0^2 and the forests of the R2 parts.* King's recursion [24, (3.33)] sets $\delta m^2(\alpha, \beta)$ equal to the zero-momentum value of the full 1PI two-point function at that order. Expand it in the lines and in $\delta m_t^2 = \delta m_0^2 + f_t$. The terms with only $\hat{G}_{\overline{M}}$ lines and only δm_0^2 insertions reproduce King's counterterm at mass $\overline{M}$, which is δm_0^2. All other terms (those with at least one $\hat{\Delta}$ line or one f-insertion) make up f_t. By Lemma H.15, inserting the vertices δm_0^2 is the same as summing over the t^0-forests of the scalar self-energy parts that contain no $\hat{\Delta}$-line, no f-vertex and not the insertion and have order at most 2. These forests are Reisz's forests of these parts (Lemma H.17 and the remark after Lemma H.15). Self-energy parts that contain the insertion, or have order $o \geq 3$, are not subtracted by King's counterterm. The parts that contain a $\hat{\Delta}$ line or an f-vertex are not subtracted either: their counterterm is the explicit f-vertex. All these parts have $\omega \leq -1$ (Lemma H.14). (b) *$t^0 \to t^1$ for the R2 parts with $\omega = 1$.* The added term $\varepsilon^{-1} \sin(\varepsilon q_\mu) \partial_{q_\mu} R_\gamma|_{q=0}$ integrates to zero over the internal momenta of γ, which are independent integration variables after S_γ [33, (2-23)]. Fubini applies because the integrand is bounded on the compact BZ: all denominators are at least $\min(\underline{M}^2, \mu_*^2)$. (c) *R3–R5.* For R4 and R5 the forest terms vanish graph by graph, by oddness at $q = 0$. For the two-photon parts R3, consider the zero-order terms summed over all R3 graphs of a given order at the same place. They add up to $\Pi_{\overline{M}}^{(o)}(0) \times$ (reduced integrand), which is 0 by the Ward identity on $\varepsilon\mathbb{Z}^3$,

Statement K.56. This needs the R3 parts to be free of $\hat{\Delta}$ and f (uniform mass $\overline{M}$), which is why one expands in $\hat{\Delta}$ rather than in the two masses. For periodicity, A is labelled by the lower end point. In this labelling the linear Taylor term of an R3 part does *not* vanish graph by graph: it acquires $(i\varepsilon/2)(q_\mu - q_\nu)\Pi^{\mathrm{mid}}_\gamma(0)$. It vanishes only after the same per-place regrouping, since $\Pi^{\mathrm{low}}(q) = \mathrm{e}^{i\varepsilon(q_\mu - q_\nu)/2}\Pi^{\mathrm{mid}}(q)$ with Π^{mid} even and $\Pi^{\mathrm{mid}}(0) = 0$. (d) *R6.* Graphs with a one-photon part vanish identically. (e) *Completeness.* Take a 1PI $\gamma \subsetneq G$ with $\omega(\gamma) \geq 0$. By Lemma H.14, it contains no $\hat{\Delta}$-line, no f-vertex and not the insertion, and it lies in R2–R6. It is subtracted with $\delta = \omega$: by (a)–(b) for R2, by (c) for R3–R5, and by (d) for R6. Every other γ has $\omega < 0$ and $t = 0$. Reisz's conditions (4-3)–(4-4) with $\delta = \omega$, and $t := 0$ when $\omega < 0$ (including for the whole graph), are "automatically satisfied" [33, p. 92]. □

The weighted-line form of Bałaban's Proposition 1. In scale-k units, a Γ^B-line ("D-line") is $u\, c_k D^\eta_k$ with $D^\eta_k := \sum_{j<k}(L^j\eta)^2 G^\eta_{(j)}$, because $\delta_j = c_k(L^j\eta)^2$. At $k = K$ we have $\kappa\mu_*^2 D = c_K D^\eta_K$ exactly, with no coarse remainder. Expanding every vector line $G_k(u) = G_k + uc_k D^\eta_k$ multilinearly, the value of a graph of the u-model is $\sum_{\text{colourings}}(uc_k)^m \times$(value of the coloured graph). Here every internal vector line is coloured C (kernel G_k) or D, and m is the number of D-lines. Write $D^\natural$ for Bałaban's degree [3, (2.1)–(2.3), pp. 422–423] with each D-leg of dimension $(-\mathsf{d}+4)/2$; a subgraph is *divergent* if $D^\natural \leq 0$. *Organisation* (α)*:* the mass counterterm is the fixed $\delta m^2_{\mathrm{ref}}$ [3, (1.23), (1.26)–(1.29)], built from C-lines only, and only C-only divergent subgraphs are renormalised (by $\delta m^2_{\mathrm{ref}}$, Ward–Takahashi partners, and $\varphi\varphi A$ partners).

Lemma H.18 (D-lines and degrees)**.** *Let* $\mathsf{d} = 3$. *(a)* D^η_k *has the representation* [3, (2.6)]. *Its slice kernels* $(L^j\eta)^2 G^\eta_{(j)}$ *obey* [3, (2.10)–(2.12)] *with* $(L^j\eta)^{-\mathsf{d}+2}$ *replaced by* $(L^j\eta)^{-\mathsf{d}+4}$ *and with the same exponential factors, and* D^η_k *obeys* [3, (3.1)]. *Hence* $D^\natural(G) = D(G) + 2m(G)$, *where* D *is Bałaban's degree with the colours forgotten. (b) Let* G *be a connected graph without the vertices (1.9), (1.11), (1.13)–(1.15) and without finite counterterms. Then* $D(G) = I_v + E'_v + \sum_v n'_v + d_s + \frac{1}{2}E_s + E_\partial - 3$. *Here* I_v *is the number of internal vector lines,* E'_v *the number of external* A' *legs,* n'_v *the number of background legs,* d_s *the number of* λ*-vertices,* E_s *the number of external scalar legs, and* E_∂ *the number of differentiations on external legs. (c) If* G *has an external leg and* $m \geq 1$, *then* $D^\natural(G) \geq 1$. *For a vacuum graph,* $D^\natural = I_v + d_s - 3 + 2m$, *plus the non-negative terms of (b) when (b) applies. So* $m = 1$ *gives* $D^\natural \geq 0$, *with equality only at* $(\alpha, \beta) = (2, 0)$, *and* $m \geq 2$ *gives* $D^\natural \geq 3$. *(d) Every divergent proper subgraph of a coloured graph is* C*-only and belongs to one of Bałaban's classes* [3, pp. 429–432]. *A vacuum graph with* $m \geq 2$, *or with* $m = 1$ *and* $o \geq 2$, *is convergent.*

The lemma uses the bounds [3, (2.10)–(2.12), p. 426] for the vector slices. The vector lines do not depend on the background. At zero background, these bounds for King's slices $H_j = G_{j+1} - G_j$ are given by Corollary E.49. By Lemma H.22 below, King's slices are Bałaban's slices $G^\eta_{(j)}$ of [3, (2.6)], including $j = 0$: King's $H_0 = G_1$ and Bałaban's $G_{(0)} = C^{(0),\eta}$.

Proof. (a) Multiply [3, (2.10)–(2.12)] by $(L^j\eta)^2 \leq 1$. Both legs of a line carry the same index j, so splitting $(L^j\eta)^{-\mathsf{d}+4}$ as $(L^j\eta)^{(-\mathsf{d}+4)/2}$ per leg is legitimate in [3, (2.14)]. For localisations at distance at least 1, $\sum_j (L^j\eta)^{2-\mathsf{d}+2-1-\alpha}\mathrm{e}^{-\delta_1 \operatorname{dist}/(L^j\eta)} \leq O(1)\mathrm{e}^{-\delta_0 \operatorname{dist}}$, because each term is at most $O(1)\mathrm{e}^{-\delta_1 \operatorname{dist}/2}(L^j\eta)^{-\alpha}\mathrm{e}^{-\delta_1/(2L^j\eta)}$, which is summable. (b) Per vertex, $D_G(v)$ is the number of η-factors, plus the dimensions of the legs on internal lines, minus the differentiations on internal lines [3, p. 423]. The η-counts are: d for (1.6) and (1.7); $\mathsf{d}+n+n'-1$ for (1.8) (one differentiation, on a scalar leg); $\mathsf{d}+n+n'-2$ for (1.10) (no differentiation). Sum over vertices using $\sum_v n_v = 2I_v + E'_v$ and $\mathsf{d} = 3$. For a counterterm vertex, [3, (2.3)] gives $D(G) = D(G') + E_\partial(G_0) \geq D(G')$, with equality for the counterterms of [3, (1.23)]. In full generality the formula carries the extra terms $+2V_7 + \sum \deg(\mathrm{ct})$, both non-negative. (c) If G contains (1.13)–(1.15), an R-vertex or a finite counterterm, then $D > 0$ by [3, Cor. 2.3, p. 429], so $D^\natural > 2$. Otherwise $m \geq 1$ gives $I_v \geq 1$, and an

external leg means $E'_v \geq 1$, or $E_s \geq 2$ (E_s is even), or $n' \geq 1$. So $D \geq -1$ and $D^\natural \geq 1$. For vacuum graphs, $E'_v = E_s = E_\partial = n' = 0$ and $I_v \geq m$. (d) A proper subgraph of a connected graph has an external leg, so use (c). The $(2,0)$ one-D-line vacuum graphs (bubble, seagull, and the (1.8), $n=2$ self-contractions) are the only ones with $D^\natural = 0$. Bałaban's graph (2.4) is the tree with one internal *scalar* line ($I_v = 0$, $E'_v = 2$, $E_s = 2$, $D = 0$). There are no "coloured (2.4) graphs". A (2.4) subgraph whose external vector legs are D-lines is integrated by parts [3, (2.8)] using the derivative bound of (a). □

Block lattices. Let T and $\Lambda = T_\varepsilon$ be as in Definition I.10, with $\langle f, g\rangle = \varepsilon^3 \sum_{x\in\Lambda} f(x)g(x)$ and with the blocks $B^k(y)$ of Definition I.10(a). For $0 \leq k \leq K$ let $\Lambda^{(k)} := T \cap t_k\mathbb{Z}^3$, and define

$$\langle B, B'\rangle_k := t_k^3 \sum_{y\in\Lambda^{(k)}} B(y)B'(y), \qquad \|B\|_k^2 := \langle B, B\rangle_k.$$

Define $Q_k : \ell^2(\Lambda) \to \ell^2(\Lambda^{(k)})$ by $(Q_kA)(y) := L^{-3k}\sum_{x\in B^k(y)} A(x)$ [24, (2.10), p. 653]. For $k < K$ define $Q : \ell^2(\Lambda^{(k)}) \to \ell^2(\Lambda^{(k+1)})$ by $(QB)(z) := L^{-3}\sum_{y\in\Lambda^{(k)}\cap B^{k+1}(z)} B(y)$. Adjoints are taken with respect to these inner products.

Lemma H.19 (block averages)**.** (a) *For $0 \leq k \leq K$: Q_k^* is the extension by constants on the blocks $B^k(y)$; $Q_kQ_k^* = 1$; $Q_k^*Q_k = P_k$; and $\|B\|_k = \|Q_k^*B\|$.*

(b) *For $0 \leq k < K$: Q^* is the extension by constants from $\Lambda^{(k+1)}$ to $\Lambda^{(k)}$; $QQ^* = 1$; Q^*Q is an orthogonal projection; $\|Q\| \leq 1$; and $QQ_k = Q_{k+1}$.*

(c) *For $0 \leq k < K$: $Q_k^*(1 - Q^*Q)Q_k = P_k - P_{k+1}$.*

Proof. By (F2) of Definition I.10, for $k \leq K$ the blocks $B^k(y)$, $y \in \Lambda^{(k)}$, partition Λ into cubes of L^{3k} sites. Each $B^{k+1}(z)$ is the disjoint union of the L^3 blocks $B^k(y)$ with $y \in \Lambda^{(k)} \cap B^{k+1}(z)$.

(a) Let $y(x)$ be the corner of the block that contains x. Since $t_k^3L^{-3k} = \varepsilon^3$,

$$\langle Q_kA, B\rangle_k = \varepsilon^3 \sum_x A(x)B(y(x)),$$

so $(Q_k^*B)(x) = B(y(x))$. Averaging a block-constant function returns its value, so $Q_kQ_k^* = 1$. The operator $Q_k^*Q_k$ replaces A on each block by its block average, which is the orthogonal projection P_k. Finally, $\|Q_k^*B\|^2 = \langle B, Q_kQ_k^*B\rangle_k = \|B\|_k^2$.

(b) The same computation, with $t_{k+1}^3L^{-3} = t_k^3$, gives the first two claims. Hence $(Q^*Q)^2 = Q^*Q = (Q^*Q)^*$, and $\|Q\|^2 = \|Q^*Q\| \leq 1$. The blocks are nested, so the average over $B^{k+1}(z)$ is the average of the L^3 averages over the blocks $B^k(y) \subset B^{k+1}(z)$. Thus $QQ_k = Q_{k+1}$.

(c) Use (a) at k and at $k+1 \leq K$, together with (b):

$$Q_k^*Q_k - Q_k^*Q^*QQ_k = P_k - (QQ_k)^*(QQ_k) = P_k - P_{k+1}. \qquad \square$$

Let $M = -\Delta_\varepsilon + \mu_*^2$, $r_k = a_kt_k^{-2}$ and $G_k = (M + r_kP_k)^{-1}$ be as in Definition I.10. For $1 \leq k \leq K$, King's effective block operator [24, (2.14), p. 653] and his lift [24, (2.18), p. 654] are

$$\Delta^{(k)} := r_k - r_k^2\, Q_kG_kQ_k^* \ \text{ on } \ell^2(\Lambda^{(k)}), \qquad F_k := r_kG_kQ_k^*.$$

For $1 \leq k < K$, the fluctuation covariance is $C^{(k)} := (\Delta^{(k)} + \beta_kQ^*Q)^{-1}$ with $\beta_k := a\, t_{k+1}^{-2}$ [24, (2.16), p. 653]. The constant here is a, not a_{k+1}.

Lemma H.20 (variational form of $\Delta^{(k)}$)**.** *Let $1 \leq k \leq K$ and $B \in \ell^2(\Lambda^{(k)})$, and put $J_B(A) := \langle A, MA\rangle + r_k\|Q_kA - B\|_k^2$ for $A \in \ell^2(\Lambda)$. Then, for all A,*

$$J_B(A) = \langle A - F_kB, G_k^{-1}(A - F_kB)\rangle + \langle B, \Delta^{(k)}B\rangle_k.$$

Hence F_kB is the unique minimiser of J_B, $\min_A J_B(A) = \langle B, \Delta^{(k)}B\rangle_k$, and $0 < \Delta^{(k)} \leq r_k$. In particular, for $k < K$ and every $\beta \geq 0$ the operator $\Delta^{(k)} + \beta Q^*Q$ is positive definite, so $C^{(k)}$ is well defined.

Proof. By Lemma H.19(a), $\|Q_kA-B\|_k^2=\langle A,\,P_kA\rangle-2\langle A,\,Q_k^*B\rangle+\|B\|_k^2$, so $J_B(A)=\langle A,\,G_k^{-1}A\rangle-2r_k\langle A,\,Q_k^*B\rangle+r_k\|B\|_k^2$. Let $A_0:=F_kB$, so that $G_k^{-1}A_0=r_kQ_k^*B$. Then

$$\langle A-A_0,\,G_k^{-1}(A-A_0)\rangle=\langle A,\,G_k^{-1}A\rangle-2r_k\langle A,\,Q_k^*B\rangle+r_k^2\langle B,\,Q_kG_kQ_k^*B\rangle_k.$$

Subtracting, and using $r_k\|B\|_k^2-r_k^2\langle B,\,Q_kG_kQ_k^*B\rangle_k=\langle B,\,\Delta^{(k)}B\rangle_k$, gives the identity. Since $G_k^{-1}\ge M\ge\mu_*^2>0$, the first term is positive unless $A=A_0$. So A_0 is the unique minimiser, and $\langle B,\,\Delta^{(k)}B\rangle_k=J_B(A_0)$.

Taking $A=0$ gives $\langle B,\,\Delta^{(k)}B\rangle_k\le J_B(0)=r_k\|B\|_k^2$. Suppose $\langle B,\,\Delta^{(k)}B\rangle_k=0$. Then $\langle A_0,\,MA_0\rangle=0$, so $A_0=0$, and therefore $r_k\|B\|_k^2=J_B(0)=J_B(A_0)=0$, that is, $B=0$. Hence $\Delta^{(k)}>0$. The last claim follows because $Q^*Q\ge0$. □

Remark H.21. In King's notation, the lemma says that the average field $A^{(k)}=F_kA_k$ of [24, (2.18), p. 654] minimises the fine action under the soft constraint $Q_kA\approx A_k$. The same completion of the square, carried out one scale up and in the block field, is Bałaban's translation [1, (3.41), p. 619].

Lemma H.22 (King's slices satisfy Bałaban's recursion)**.** *Let $a_k=a(1-L^{-2})(1-L^{-2k})^{-1}$ be King's constants* [24, (2.13), p. 653]. *Let $H_j=G_{j+1}-G_j$, $0\le j<K$, with $G_0:=0$, be King's slices (Subsection 5.1).*

(a) $r_{j+1}^{-1}=r_j^{-1}+\beta_j^{-1}$ *for* $1\le j<K$.
(b) $H_j=F_jC^{(j)}F_j^*=r_j^2\,G_jQ_j^*C^{(j)}Q_jG_j$ *for* $1\le j<K$.
(c) $H_0=G_1=(M+a\,t_1^{-2}P_1)^{-1}$.

Proof. (a) Since $t_j=L^{-1}t_{j+1}$,

$$\frac{t_j^2}{a_j}+\frac{t_{j+1}^2}{a}=\frac{t_{j+1}^2}{a(1-L^{-2})}\big[L^{-2}(1-L^{-2j})+1-L^{-2}\big]=\frac{t_{j+1}^2}{a_{j+1}}.$$

(b) On $\ell^2(\Lambda)\oplus\ell^2(\Lambda^{(j)})$, with the inner products $\langle\cdot,\,\cdot\rangle$ and $\langle\cdot,\,\cdot\rangle_j$, consider the quadratic form

$$S(A,B):=\langle A,\,MA\rangle+r_j\|Q_jA-B\|_j^2+\beta_j\|QB\|_{j+1}^2.$$

First, by Lemma H.20 and $\|QB\|_{j+1}^2=\langle B,\,Q^*QB\rangle_j$,

$$S(A,B)=\langle A-F_jB,\,G_j^{-1}(A-F_jB)\rangle+\langle B,\,(C^{(j)})^{-1}B\rangle_j.$$

Thus $S=\Lambda^*\operatorname{diag}(G_j^{-1},(C^{(j)})^{-1})\Lambda$, where $\Lambda(A,B):=(A-F_jB,B)$ is invertible with $\Lambda^{-1}(A',B)=(A'+F_jB,B)$. So S is positive definite, and the AA-block of $S^{-1}=\Lambda^{-1}\operatorname{diag}(G_j,C^{(j)})\Lambda^{-*}$ is $G_j+F_jC^{(j)}F_j^*$.

Second, fix A and put $c:=Q_jA$. Write $B=Q^*z+w$ with $z:=QB$ and $w:=(1-Q^*Q)B$. Since $QQ^*=1$ (Lemma H.19(b)), $\|Q^*v\|_j=\|v\|_{j+1}$, so

$$\|c-B\|_j^2=\|Qc-z\|_{j+1}^2+\|(1-Q^*Q)c-w\|_j^2.$$

Hence

$$\min_B\big[r_j\|c-B\|_j^2+\beta_j\|QB\|_{j+1}^2\big]=\min_z\big[r_j\|Qc-z\|_{j+1}^2+\beta_j\|z\|_{j+1}^2\big]=\frac{r_j\beta_j}{r_j+\beta_j}\|Qc\|_{j+1}^2.$$

By (a), $r_j\beta_j/(r_j+\beta_j)=r_{j+1}$. By Lemma H.19(a),(b), $\|Qc\|_{j+1}^2=\|Q_{j+1}A\|_{j+1}^2=\langle A,\,P_{j+1}A\rangle$. So $\min_B S(A,B)=\langle A,\,G_{j+1}^{-1}A\rangle$.

For a positive definite block form, the minimum over the second variable is the Schur complement, whose inverse is the AA-block of S^{-1}. Hence $G_{j+1}=G_j+F_jC^{(j)}F_j^*$. Finally, $F_j^*=r_jQ_jG_j$.

(c) $G_0=0$ gives $H_0=G_1=(M+r_1P_1)^{-1}$, and $a_1=a$. □

Part (b) is the recursion [1, (2.42)] at zero background, written in King's notation, and part (c) is the identity $G_1 = C^{(0)}$ that precedes [1, (2.43)]. Summing (b) and (c) over $j < k$ gives

$$G_k = G_1 + \sum_{j=1}^{k-1} a_j^2 t_j^{-4} G_j Q_j^* C^{(j)} Q_j G_j.$$

In scale-k units ($t_j/t_k = L^j\eta$) this is the decomposition [3, (2.6)]=[1, (2.43)] at $\tilde{B} = 0$. Thus, when the zero-background operators $G_j, Q_j, C^{(j)}$ and the constants a_j of [1, Sect. 2] are written in King's normalisation, the H_j are the slices $G^{\eta}_{(j)}$ of [3, (2.6)] for $0 \le j < k$; for $j = 0$, $H_0 = G_1 = C^{(0)}$. The bounds use only the operators H_j, through Corollary E.49, which is proved for them directly.

Lemma H.23 (families)**.** *Call two coloured graphs* partners *if one arises from the other by one of Bałaban's partner operations applied to a C-only subgraph, with the colours of all other lines unchanged. The operations are: (i) replace a C-only divergent scalar self-energy G_0 by the vertex (1.7) with $\delta m^2_{G_0}$, or back; (ii) exchange C-only two-vector-leg graphs of the same order* [3, p. 431]*; (iii) the $\varphi\varphi A$ partners* [3, (2.18)–(2.22)]*. Then partners have the same (m, d_v, d_s). Every partner a coloured graph needs occurs in the expansion with the right coefficient, and no counterterm vertex is left without its partner. Hence the coloured expansion is the disjoint union of the partner classes (the* families*), and for $m = 0$ these are Bałaban's.*

The lemma uses Bałaban's construction of the families for $u = 0$ [3, pp. 420, 429–432].

Proof. The operations act inside C-only parts and preserve the coupling orders ([3, p. 420]: "all graphs in the family G_{ren} have the same orders"). The counterterm $\delta m^2_{\text{ref}} = \sum_{G_0} \delta m^2_{G_0}$ runs over all C-only self-energy graphs of order ≤ 4, including those with lower-order insertions [3, p. 417]. The C-only graphs of every order are present as the u^0 part of the expansion. The partner carries the same index assignment as G_0 [3, (3.9), p. 435], and colouring the complement commutes with the operations, so the combinatorial factors are Bałaban's. D-containing subgraphs need no partner, by Lemma H.18(d). The partition is the orbit decomposition of the partner relation. □

Theorem H.24 (the weighted-line form of [3, Prop. 1])**.** *Fix $\bar{n} > 12$, $0 < \alpha_0 < 1$, $k \le K$ and $\eta = L^{-k}$. Let $\Omega, \Omega_1, \Omega_2, \tilde{B}$ satisfy the hypotheses of* [3, Prop. 1, pp. 412, 419–421] *on Bałaban's periodic lattice torus. Work in organisation (α).*

(a) *(Degrees.) Lemma H.18(c).*

(b) *(Families.) The coloured expansion of the perturbative expressions of the u-model, with δm^2_{ref}, is a finite disjoint sum of the families of Lemma H.23, each with fixed (m, d_v, d_s). No vacuum counterterm is attached to the vacuum polynomial $T_{\bar{n}}[\Phi(u)]$ itself, since it is the quantity being bounded. Inside the per-step expressions, by contrast, the vacuum families carry their pieces of $E_k(u)$.*

(c) *(Bound.) Let F be a non-vacuum family, or a vacuum family with $m \ge 2$, or one with $m = 1$ and $o \ge 2$, whose D-lines carry D^{η}_k. Then*

$$|E(F, \{\Box(v)\}, \Phi_{\text{ext}}, A_{\text{ext}})| \le O(1)\,(e(L^k\varepsilon))^{d_v} (\lambda(L^k\varepsilon))^{d_s}\, \mathrm{e}^{-\delta_0 d(\{\Box(v)\})}\, \|h\Phi_{\text{ext}}\|_{1,\alpha_0} \|h' A_{\text{ext}}\|_{1,\alpha_0}.$$

Here $\delta_0 > 0$ is Bałaban's tree rate of [3, (1.33)]*, and $O(1) = O(\alpha_0, \bar{n})$ (together with his lattice data L, a) does not depend on ε, k, Ω, Ω_1, Ω_2, $\tilde{B}$, u or κ. In physical normalisation the family carries the explicit factor $(uc_k)^m$.*

(d) *(Per step.) The same bound holds for the sub-sum whose largest index is $k - 1$.*

The theorem is stated on the torus. The statement that the vacuum family with $m = 1$, $o = 1$ at a fixed largest index is $O(1)$ is not used. King's cube with Neumann and Dirichlet conditions is not used.

Proof. Every step transfers a step of [3, pp. 422–445]. *What transfers.* Vertices, scalar lines and external fields are unchanged by colouring [3, pp. 411–422]. So are the parity cancellation $\operatorname{tr} q^{2n+1} = 0$ (p. 434), the localisation and background expansion (3.2)–(3.3) (pp. 432–433), and gauge covariance (vector lines do not depend on $\tilde{B}$ and are gauge-neutral). Bałaban's Ward–Takahashi identities (2.23)–(2.28) hold at every fixed internal vector field, hence for any Gaussian law of the internal photons, coefficientwise in s for the covariance $G + sD$; only the s^0 coefficient is used. The classification [3, Cor. 2.3, pp. 429–432] and the renormalisation identities [3, (3.6)–(3.38)] are needed only for C-only subgraphs, by Lemma H.18(d), and there they are Bałaban's own. D-lines enter them only as external legs, through (2.10)–(2.12). The steps where Bałaban replaces an internal *photon* line by C^ξ are (3.24), (3.31)–(3.32) and (3.37)–(3.38). Steps (3.16) and (3.29) are not among them: (3.16) replaces only a scalar line, and (3.29) has no internal photon. With a weighted photon, $\sum_{j<j_0} w_j G_{(j)} = G_{j_0} + uc_k D^\eta_{j_0}$: Bałaban's identity applies to the C part, and the D part has degree at least 2 more. *Transversality.* A C-only vacuum-polarisation subgraph Π has negative degree. It is renormalised by Bałaban's real-space Ward step [3, (3.25)–(3.32), pp. 439–442]: Taylor expansion of the external legs (3.26); summation of the internal indices below j_0; the replacement by C^ξ; vanishing of the symmetric term (3.27)–(3.28); and the identity (2.26) at (3.29). It acts on the class with arbitrary regular external legs (p. 440), and so in particular with D-line legs, which need only the Hölder bound of Lemma H.18(a). A Π with an internal D-line has $D^\natural \geq 2$ and needs nothing. *The bound (c).* (1) Lines between localisations at distance at least 1, whether C or D, become pairs of external fields by [3, (3.1)]. For a D-line this lowers $D^\natural$ by 1, rather than raising it; the conclusion survives. If l is a D-line, $G \setminus l$ has the same subgraphs as the line subsets of G not containing l, with the same degrees. Its D-containing components have $D^\natural \geq 1$. Its C-only divergent components are C-only divergent subgraphs of G, already renormalised in the family, or (2.4)-type trees, or odd graphs, which vanish. (2) Localise and remove the background as in [3, (3.2)–(3.3)]. (3) Decompose every line by [3, (2.6)], the D-lines by Lemma H.18(a), and sum over orderings (3.4). (4) In a fixed ordering, the C-only components are treated by Bałaban's construction (3.5), unchanged. The D-containing components have $D^\natural \geq 1$. Bałaban's transformations never lower the degree of a larger subgraph (pp. 425, 445). So every component of every subgraph in every admissible ordering has positive degree. (5) Apply [3, Prop. 2.2, p. 428] ("lines with the same exponential factors but with arbitrary dimensions instead of $-d + 2$"), with the D-lines of dimension $-\mathsf{d} + 4$. This gives (2.16), hence (3.3), (3.2) and the displayed bound, exactly as at [3, pp. 432–434]. Each D-line has two A'-legs, both counted in d_v. (6) The u-model value is $(uc_k)^m$ times this, by the multilinear expansion. *(d).* In (2.16) the last sum runs over the largest index. Fixing it gives a sub-sum of a majorant sum. *(a), (b).* Lemmas H.18 and H.23. □

The theorem covers only organisation (α). In the other organisation (weighted counterterms, with a finite mass vertex depending on u at every step), the per-step statements are not proved here and are not used. The following sketched items of [3] are used as in the proof of [3, Prop. 1]; the colouring does not change them:

- I1: family combinatorics, "inspecting all possible cases" (p. 432);
- I2: bad localisations of the vacuum counterterms, "even simpler and we omit them" (p. 422); this does not affect E_2 or Corollary H.26;
- I3: convergence of δm^2_{fin} and E_{fin}, and of the regular remainders treated as external fields (p. 418);
- I4: [1, Props. 2.1–2.3], cited by Bałaban, behind (2.10)–(2.12);
- I5: the background Hölder norms (p. 412).

In addition, [3, Prop. 2.2, p. 428] is a remark promoted to a proposition, without a separate proof. The proof below is a transfer of [3, pp. 422–445]: only the new ingredients are written. The tree rate given by the proof depends on $\bar{n}$, through the line-moving rates of [3, p. 427], and not only on d; only the existence of some $\delta_0 > 0$ is used (Proposition E.60).

The sketched items and Table 2. By page, the items correspond to Table 2 as follows.

- I1 is step S10 (p. 432). The classification [3, Cor. 2.3] behind it is step S7.
- I4 is step S3 (pp. 424, 426). At zero background, the slice bounds for King's slices are Corollary E.49, and these slices are Bałaban's by Lemma H.22. The vector lines do not depend on $\tilde{B}$, so the bounds for the D-lines are always of this kind (Lemma H.18(a)).
- I2, I3 and I5 concern pp. 422, 418 and 412, which carry no step of Table 2. Their depth is at most that of Proposition 5.8. For I3, the bounds on $\delta m^2_{\mathrm{fin}}$ and E_{fin} uniform in the torus are Propositions E.9 and E.21, and the last remainder line is Proposition E.50. The bad localisations of the moved vacuum pieces (I2) are treated in Proposition E.62.

The proof of [3, Prop. 2.1, pp. 424–428] uses of the lines only the representation [3, (2.6)] and the bounds [3, (2.10)–(2.12)]. The leg dimensions enter only through [3, (2.14)]. Bałaban's [3, Prop. 2.2] is this proof with arbitrary line dimensions. In the form used below it reads as follows: for a connected generalised graph all of whose connected components in a fixed ordering have positive degree, [3, (2.16)] holds. This is Lemma E.51(i), and Lemma E.52 gives the version with the largest index fixed. Subgraphs of type [3, (2.4)] are treated beforehand by summation by parts [3, (2.8)], which is step S4 of Table 2. The D-lines are lines of this generalised kind, of dimension $-\mathsf{d}+4$, by Lemma H.18(a). For the vacuum families with $m \geq 2$ that are used in Lemma H.9, part (c) is replaced there by Statement K.58, which is the corresponding bound at zero background. For the other uses of part (c), namely Corollary H.26 and the non-vacuum families, the items I1–I5 are used as described above.

Remark H.25 (the E_2 constant). The constant $4C_W$ of Lemma H.9 is conservative: $m(m-1) \leq m^2$ costs a factor 2, and c_K overcounts the top D-slice by L^2. That C_W does not depend on μ_* is not used: factors depending on $\mu_*\varepsilon_0$ may enter through $C_{\mu_*} - G_K$, which is treated as an external field (I3), but they do not depend on ε, so they do not affect $\sup_\varepsilon$.

The vacuum coefficients linear in u.

Corollary H.26 (the vacuum coefficients linear in u on the torus)**.** *For every $(\alpha,\beta) \neq (2,0)$ with $\alpha+\beta \leq \bar{n}$ (so $o \geq 2$ whenever the coefficient is non-zero, since $\alpha \geq 2$ is even), and every $M \in \{m_1, m_2\}$,*

$$|[\mathcal{E}(\cdot; M)]_{1,\alpha\beta}| \leq C_{1,\alpha\beta}(M)\, \kappa \mu_*^2 L^2\, |T|\, \varepsilon_0^{o-1},$$

uniformly in $\varepsilon = 2^{-K}\varepsilon_0$ and in the torus. Here $C_{1,\alpha\beta}$ is Bałaban's $O(1)(\bar{n}, \alpha_0, \mathsf{d}, L, a, M)$ times a graph count, independent of ε, u, κ and ℓ_T.

This is an instance of Theorem H.24(c). The excluded coefficient $(2,0)$ ($D^\natural = 0$, logarithmically divergent) is treated by its mass difference in Lemma H.10.

Proof. The covariance $C_{\mu_*} + u\Gamma^B$ is linear in u. So the u^1 part of $T_{\bar{n}}[\Phi_M(u)]$ is the sum of the coloured vacuum graphs with exactly one D-line. Each such graph is $e^\alpha \lambda^\beta$ times a number, the $\delta m^2_{\mathrm{ref}}$ vertices carrying their own orders. By Lemma H.23, the families partition the set of graphs with fixed $(m,\alpha,\beta) = (1,\alpha,\beta)$, so $[\mathcal{E}]_{1,\alpha\beta} = \pm\sum_F c_K\, \mathrm{val}(F)$ in scale-K units. For $o \geq 2$, every such family is covered by Theorem H.24(c) ($D^\natural \geq 1$, Lemma H.18(c)). Its C-only photon self-energy closed by the single D-line is renormalised by Bałaban's Ward step [3, (3.25)–(3.32)], with the two ends of the D-line as external legs. No vacuum counterterm is attached (Theorem H.24(b)). Counting: fix one vertex in a unit block of scale K. The other localisations sum to $O(1)$ by $\mathrm{e}^{-\delta_0 d}$, and there are $|T|/t_K^3$ blocks. A family therefore contributes at most

$$O(1)(e^2 t_K)^{\alpha/2}(\lambda t_K)^\beta c_K\, t_K^{-3}|T| = O(1)\, e^\alpha \lambda^\beta \kappa \mu_*^2 L^2\, t_K^{o-1}|T|.$$

Since $o-1 \geq 1$ and $t_K \leq \varepsilon_0$ (only $t_K \leq \varepsilon_0$ is used), $t_K^{o-1} \leq \varepsilon_0^{o-1}$. Sum over the finitely many families. □

The diagonal of King's slices. *King's objects.* These are as in [24, (2.14)–(2.17)]. Q_j is the block average to the scale-j block lattice, and P_j is the orthogonal projection onto fields that are constant on scale-j blocks. a_j are King's block constants, $r_j := a_j t_j^{-2}$, and G_j is King's scale-j Green's function. $C^{(j)} = (\Delta^{(j)} + a t_{j+1}^{-2} Q^* Q)^{-1}$ is the fluctuation covariance. $H_j = F_j C^{(j)} F_j^*$ with $F_j = r_j G_j Q_j^*$. Per component, $C_{\mu_*} = (-\Delta_\varepsilon + \mu_*^2)^{-1}$. Put $a_\flat := \inf_j a_j = a(1 - L^{-2})$; we use $a_j \le a$.

Lemma H.27 (King's slice at a point)**.** *On the periodic lattice torus, for $1 \le j < K$ and every site x (per component),*

$$H_j(x,x) \le C_H / t_j, \qquad C_H := c_C\, a^2\, (m_a + 3^{1/4} C_{\mathrm{osc}})^2, \quad c_C := \frac{4}{a_\flat} + \frac{L^2}{8} + \frac{L^2}{a}, \quad m_a := \max\Big(\frac{1}{8}, \frac{1}{a_\flat}\Big),$$

and $H_0(x,x) \le C_{\mu_*}(x,x)$. Here C_{osc} is the constant of the lattice Green's-function estimate

$$\text{(G-osc)} \qquad \langle \delta_x - \delta_y,\, C_{\mu_*}^2 (\delta_x - \delta_y) \rangle \le C_{\mathrm{osc}}^2 |x - y|,$$

which holds on the periodic lattice with $C_{\mathrm{osc}}^2 \le 8.1$, uniformly in ε, ℓ_T and μ_*. Consequently

$$\max_x D(x,x) \le L^2 \varepsilon\, c_1 + C_H L^2 \varepsilon_0 / (L - 1),$$

where c_1 is such that $\varepsilon\, C_{\mu_*}(x,x) \le c_1$ (see Remark H.30 and Lemma H.29).

On the periodic lattice, (G-osc) is proved below, so the lemma holds on the torus without further conditions. The proof uses the block Poincaré inequality (Lemma I.11) and the variational form of $\Delta^{(j)}$ (Lemma H.20), through Lemma H.28, which is step (a) of the proof. The lemma gives step (2) of Lemma H.10.

Proof. (a) $C^{(j)} \le c_C t_j^2$ *on the scale-j block lattice.* By the variational form, $\langle B, \Delta^{(j)} B \rangle_j = \min_A [\langle A, (-\Delta_\varepsilon + \mu_*^2) A \rangle + r_j \|Q_j A - B\|_j^2]$. Let A^* be the minimiser and $B' := Q_j A^*$. Then $\|(1 - Q^*Q) B'\| = \|(P_j - P_{j+1}) A^*\| \le \|(1 - P_{j+1}) A^*\| \le (t_{j+1}^2/8)^{1/2} \langle A^*, (-\Delta_\varepsilon + \mu_*^2) A^* \rangle^{1/2}$, by the block Poincaré inequality. Here $Q_{j+1} = Q Q_j$ and $Q_j Q_j^* = 1$; the constant $1/8$ is the smallest non-zero Neumann eigenvalue for $n \ge 2$ sites per block. Also $\|QB'\| \le \|QB\| + \|B - B'\|$. Hence $\|B\| \le 2\|B - B'\| + (t_{j+1}^2/8)^{1/2} \langle A^*, (-\Delta_\varepsilon + \mu_*^2) A^* \rangle^{1/2} + \|QB\|$. Cauchy–Schwarz with weights $(r_j, 1, \beta)$, $\beta = a t_{j+1}^{-2}$, gives $\|B\|^2 \le (4/r_j + t_{j+1}^2/8 + 1/\beta) \langle B, C^{(j)-1} B \rangle = t_j^2 (4/a_j + L^2/8 + L^2/a) \langle B, C^{(j)-1} B \rangle \le c_C t_j^2 \langle B, C^{(j)-1} B \rangle$. (b) Hence $H_j = F_j C^{(j)} F_j^* \le c_C t_j^2 F_j F_j^*$, and $H_j(x,x) \le c_C t_j^2 r_j^2 \|P_j G_j \delta_x\|^2$ with $\delta_x := \varepsilon^{-3} 1_{\{x\}}$. (c) Split $\delta_x = \rho_x + v$ with $\rho_x := P_j \delta_x = t_j^{-3} 1_{\square_x}$ and $P_j v = 0$. First, $\|P_j G_j \rho_x\| \le \|G_j\| \|\rho_x\| \le m_a t_j^2\, t_j^{-3/2}$, using $\|G_j\| \le t_j^2 \max(1/8, 1/a_\flat)$. Second, $G_j = (-\Delta_\varepsilon + \mu_*^2)^{-1} - G_j r_j P_j (-\Delta_\varepsilon + \mu_*^2)^{-1}$ gives $P_j G_j v = (1 - r_j P_j G_j P_j) P_j (-\Delta_\varepsilon + \mu_*^2)^{-1} v$, with $0 \le r_j P_j G_j P_j \le 1$ on $\operatorname{ran} P_j$. So $\|P_j G_j v\| \le \sup_{\|w\| \le 1} [\psi_w(x) - \operatorname{avg}_{\square_x} \psi_w]$ with $\psi_w := C_{\mu_*} w$. Moreover $|\psi_w(x) - \psi_w(y)| \le \langle \delta_x - \delta_y, C_{\mu_*}^2 (\delta_x - \delta_y) \rangle^{1/2} \|w\| \le C_{\mathrm{osc}} (\sqrt{3} t_j)^{1/2}$ for $y \in \square_x$. Hence $\|P_j G_j \delta_x\| \le (m_a + 3^{1/4} C_{\mathrm{osc}}) t_j^{1/2}$, and $H_j(x,x) \le c_C a_j^2 t_j^{-2} (m_a + 3^{1/4} C_{\mathrm{osc}})^2 t_j \le C_H / t_j$. *(G-osc) on the periodic lattice.* Write $s := |x - y|$ and $\lambda(k) = \sum_i (4/\varepsilon^2) \sin^2(\varepsilon k_i / 2) \ge (4/\pi^2) |k|^2$. The left side equals $\ell_T^{-3} \sum_k |e^{ik\cdot x} - e^{ik\cdot y}|^2 (\lambda(k) + \mu_*^2)^{-2}$; the $k = 0$ term vanishes. The summand is at most $(\pi^4/16) \min(4|k|^{-4}, s^2 |k|^{-2})$, which is radially non-increasing. Every k' in the cell of side $2\pi/\ell_T$ around $k \ne 0$ has $0.134|k| \le |k'| \le 1.866|k|$. This gives the Riemann bound by $(1.87^3/(8\pi^3)) \int f$ (dropping the BZ restriction only adds positive terms), and $\int (\pi^4/16) |q|^{-4} \min(4, |q|^2 s^2)\, d^3 q = \pi^5 s$. So the left side is at most $8.07\, s \le 8.1\, s$. *The display.* We have $D = \sum_{j<K} t_{j+1}^2 H_j$ and $t_1^2 H_0(x,x) \le L^2 \varepsilon^2 C_{\mu_*}(x,x) \le L^2 \varepsilon c_1$. For $j \ge 1$, $\sum_{j \ge 1} t_{j+1}^2 / t_j = L^2 \sum_{j \ge 1} t_j \le L^2 \varepsilon_0 / (L - 1)$. □

Lemma H.28 (the fluctuation covariance)**.** *For $1 \le j < K$, $C^{(j)} \le c_C t_j^2$ on $\ell^2(\Lambda^{(j)})$, with c_C as in Lemma H.27.*

Proof. Let $B \in \ell^2(\Lambda^{(j)})$, let $A^* := F_j B$ be the minimiser of Lemma H.20, and put $B' := Q_j A^*$. Then

$$r_j \|B - B'\|_j^2 + \langle A^*,\, M A^* \rangle = \langle B,\, \Delta^{(j)} B \rangle_j .$$

Decompose $B = (B - B') + (1 - Q^*Q)B' + Q^*QB'$.

The operators $1 - Q^*Q$ and $P_j - P_{j+1}$ are orthogonal projections. So Lemma H.19(a)–(c) gives $\|(1 - Q^*Q)B'\|_j^2 = \langle A^*,\, (P_j - P_{j+1})A^* \rangle = \|(P_j - P_{j+1})A^*\|^2$. Since $P_j - P_{j+1} \le 1 - P_{j+1}$, Lemma I.11(a) at scale $j+1 \le K$ gives

$$\|(P_j - P_{j+1})A^*\| \le \|(1 - P_{j+1})A^*\| \le (t_{j+1}^2/8)^{1/2} \langle A^*,\, M A^* \rangle^{1/2}.$$

By Lemma H.19(b), $\|Q^*QB'\|_j = \|QB'\|_{j+1} \le \|QB\|_{j+1} + \|B' - B\|_j$. Hence

$$\|B\|_j \le 2\|B - B'\|_j + (t_{j+1}^2/8)^{1/2} \langle A^*,\, M A^* \rangle^{1/2} + \|QB\|_{j+1}.$$

Apply the Cauchy–Schwarz inequality with weights $(r_j, 1, \beta_j)$ and use $\|QB\|_{j+1}^2 = \langle B,\, Q^*QB \rangle_j$:

$$\|B\|_j^2 \le \Big(\frac{4}{r_j} + \frac{t_{j+1}^2}{8} + \frac{1}{\beta_j} \Big) \langle B,\, (\Delta^{(j)} + \beta_j Q^*Q) B \rangle_j = t_j^2 \Big(\frac{4}{a_j} + \frac{L^2}{8} + \frac{L^2}{a} \Big) \langle B,\, (C^{(j)})^{-1} B \rangle_j .$$

By (F1) of Definition I.10, $a_j > a_\flat$, so the bracket is at most c_C. Thus $(C^{(j)})^{-1} \ge (c_C t_j^2)^{-1}$, that is, $C^{(j)} \le c_C t_j^2$. □

Lemma H.29 (the diagonal of C_{μ_*} on the torus)**.** *In the setting of Definition I.10, for every $x \in T_\varepsilon$,*

$$\varepsilon\, C_{\mu_*}(x,x) \le \frac{\sqrt{3}\,\pi}{8} \Big(1 + \frac{\sqrt{3}}{2} \Big)^3 + \frac{\varepsilon}{\ell_T^3 \mu_*^2} < 4.42 + \frac{\varepsilon}{\ell_T^3 \mu_*^2}.$$

In particular, the constant c_1 of Lemmas H.27 and H.10 may be taken as $c_1 := 4.42 + \varepsilon_0/(\ell_0^3 \mu_*^2)$, for all $\varepsilon \le \varepsilon_0$ and $\ell_T \ge \ell_0$.

Proof. By (F2), $\ell_T/\varepsilon \ge 4$, so $-\Delta_\varepsilon$ has the symbol $\lambda(k) = \sum_i (4/\varepsilon^2) \sin^2(\varepsilon k_i/2)$ on $T^* := (2\pi/\ell_T)\mathbb{Z}^3 \cap (-\pi/\varepsilon, \pi/\varepsilon]^3$. Hence $C_{\mu_*}(x,x) = \ell_T^{-3} \sum_{k \in T^*} (\lambda(k) + \mu_*^2)^{-1}$ for every x. The term $k = 0$ is $\ell_T^{-3} \mu_*^{-2}$.

For $k \ne 0$, the inequality $\sin(t/2) \ge t/\pi$ on $[0, \pi]$ gives $\lambda(k) \ge (4/\pi^2)|k|^2$. Every $k \in T^*$ satisfies $|k| \le R' := \sqrt{3}\pi/\varepsilon$. Put $c := 1 + \sqrt{3}/2$ and $f(q) := |q|^{-2} \mathbf{1}_{\{|q| \le R'\}}$. The cubes $Z_k := k + [-\pi/\ell_T, \pi/\ell_T]^3$ have disjoint interiors and volume $(2\pi/\ell_T)^3$. For $k \ne 0$ and $k' \in Z_k$ we have $|k' - k| \le \sqrt{3}\pi/\ell_T \le (\sqrt{3}/2)|k|$, so $0 < |k'| \le c|k|$. Hence $|k'/c| \le |k| \le R'$ and $f(k) \le f(k'/c)$. Averaging over Z_k and summing over k gives

$$\sum_{k \in T^* \setminus \{0\}} |k|^{-2} \le \Big(\frac{\ell_T}{2\pi} \Big)^3 \int_{\mathbb{R}^3} f(q/c)\, d^3 q = \Big(\frac{\ell_T}{2\pi} \Big)^3 c^3\, 4\pi R'.$$

Therefore

$$\varepsilon\, \ell_T^{-3} \sum_{k \ne 0} (\lambda(k) + \mu_*^2)^{-1} \le \varepsilon\, \frac{\pi^2}{4}\, \frac{4\pi c^3 R'}{(2\pi)^3} = \frac{\sqrt{3}\pi}{8}\, c^3 = 4.4195\ldots$$

Adding the term $k = 0$ proves the bound. The right side increases in ε and decreases in ℓ_T, which gives the statement about c_1. □

With Lemma H.29, the display of Lemma H.27 holds in the form

$$\max_x D(x,x) \le L^2 \varepsilon\, c_1 + \frac{C_H L^2 \varepsilon_0}{L - 1}, \qquad c_1 = 4.42 + \frac{\varepsilon_0}{\ell_0^3 \mu_*^2},$$

uniformly in $\varepsilon \le \varepsilon_0$ and $\ell_T \ge \ell_0$. This is the form used in step (2) of Lemma H.10.

Remark H.30 (the torus zero mode). On King's Neumann cube one has $C_{\mu_*}(x,x) \le c_1/\varepsilon + 1/(\ell^3\mu_*^2)$ with $c_1 \le 36$. On the periodic torus, the Fourier sum $C_{\mu_*}(x,x) = \ell_T^{-3}\sum_k(\lambda(k)+\mu_*^2)^{-1}$ contains the zero-mode term $\ell_T^{-3}\mu_*^{-2}$, and the Riemann argument of the proof bounds the rest by c_1/ε. So on T one should read c_1 as $c_1 + \varepsilon/(\ell_0^3\mu_*^2)$. This adds at most $L^2\varepsilon^2/(\ell_0^3\mu_*^2)$ to $\max_x D(x,x)$, which is bounded and tends to 0 with ε, so no conclusion of Lemma H.10 changes.

Torus images and the heat kernel. For $\Lambda \in \{T, \varepsilon\mathbb{Z}^3\}$ write $C_{M,\Lambda} := \int_\Lambda (\hat k^2 + M^2)^{-1}$, so that $C_{M,T} = C^T(M^2)$ and $C_{M,\varepsilon\mathbb{Z}^3} = C^\infty(M^2)$ in the notation of the beginning of this appendix. Write $G_M^\varepsilon(x) = \int_{\mathrm{BZ}} \hat G_M \mathrm{e}^{ik\cdot x}$, as there.

Lemma H.31 (torus images, free of ε). *For every ε with $\ell_T/\varepsilon \in \mathbb{N}$, $\ell_T/\varepsilon \ge 2$, and every $M > 0$,*

$$E_T(M) := C_{M,T} - C_{M,\varepsilon\mathbb{Z}^3} = \sum_{n\in\mathbb{Z}^3\setminus 0} G_M^\varepsilon(n\ell_T) \le \bar E(M,\ell_T),$$

$$\bar E(M,\ell) := \frac{7.37}{\ell} + \frac{3\pi}{4\ell} E_1\Big(\frac{M^2\ell^2}{8}\Big) + \frac{12\pi\,\mathrm{erfc}(M\ell/(2\sqrt2))}{M\ell^2} + \frac{64\,\mathrm{e}^{-M^2\ell^2/8}}{M^2\ell^3},$$

where E_1 is the exponential integral. Moreover $\bar E$ is decreasing in ℓ. The constant is $7.37 \ge 3\pi/4 + 768\sqrt\pi\,(2/5)\,8^{-5/2} + 2 = 7.364$.

The bound is only polynomial in $1/\ell_T$, whereas the true size is expected to be exponentially small. For the photon mass, $E_T(\mu_*) \ge 1/(\mu_*^2\ell_T^3)$ because of the zero mode.

Proof. (1) $G_M^\varepsilon(x) = \int_0^\infty \mathrm{e}^{-M^2t} p_t(x)\,dt$, where $p_t = \prod_\mu p_t^{(1)}(x_\mu)$ and $p_t^{(1)}(x) = \varepsilon^{-1}\mathbb{P}(X_t = x)$ for the continuous-time walk on $\varepsilon\mathbb{Z}$ with jumps $\pm\varepsilon$ at rate ε^{-2} each. Its Fourier symbol is $(4/\varepsilon^2)\sin^2(\varepsilon k/2)$, $\mathrm{Var}\,X_t = 2t$, and $p_t^{(1)}$ is non-increasing in $|x|$ ($I_n(u)$ decreases in n). (2) By monotonicity, and since $(n\ell - \ell/2, n\ell]$ contains at least $\ell/(4\varepsilon)$ sites when $\ell/\varepsilon \ge 2$, we have $p_t^{(1)}(n\ell) \le (4\varepsilon/\ell)\sum_{x\in(n\ell-\ell/2,n\ell]} p_t^{(1)}(x)$ for $n \ge 1$, and symmetrically for $n \le -1$. Hence $\delta(t) := \sum_{n\ne0} p_t^{(1)}(n\ell) \le (4/\ell)\mathbb{P}(|X_t| \ge \ell/2) \le (4/\ell)\min(1, 8t/\ell^2)$, by Chebyshev. (3) $p^0 := p_t^{(1)}(0) = \varepsilon^{-1}\mathrm{e}^{-v}I_0(v)$ with $v = 2t/\varepsilon^2$. Since $\mathrm{e}^{-v}I_0(v) = \pi^{-1}\int_0^\pi \mathrm{e}^{-v(1-\cos\theta)}d\theta$ and $1-\cos\theta \ge 2\theta^2/\pi^2$ on $[0,\pi]$, $\mathrm{e}^{-v}I_0(v) \le \sqrt{\pi/(8v)}$. Hence $p^0 \le a_*/\sqrt t$ with $a_* := \sqrt\pi/4$, with no ε. (4) $\sum_{n\in\mathbb{Z}^3\setminus0} p_t(n\ell) = (p^0+\delta)^3 - (p^0)^3 = 3(p^0)^2\delta + 3p^0\delta^2 + \delta^3$. Integrate against e^{-M^2t}, using $\mathrm{e}^{-M^2t} \le 1$ on $t \le \ell^2/8$ and the two forms of $\min(1, 8t/\ell^2)$. The resulting closed forms give $\bar E$. □

Corollary H.32 (the torus heat kernel; the form used in Lemma H.10). *Under the hypotheses of Lemma H.31, with $S_q = |T|^{-1}\sum_{q\in T^*}$ and $b := 4/\ell_T$,*

$$S_q\,\mathrm{e}^{-t\hat q^2} \le \big(a_*/\sqrt t + b\big)^3 \quad (t>0), \qquad S_q\,\hat G_M(q)^2 \le C_T(M,\ell_T) \le C_T(M,\ell_0) \quad (\ell_T \ge \ell_0),$$

with $C_T(M,\ell) := a_*^3\sqrt\pi/M + 3a_*^2 b/M^2 + \frac{3\sqrt\pi}{2}a_* b^2/M^3 + b^3/M^4$, where $b = 4/\ell$.

This is steps (2)–(3) of the proof of Lemma H.31.

Proof. By Poisson summation on the lattice torus (as in Lemma H.4), $S_q\mathrm{e}^{-t\hat q^2} = \sum_{n\in\mathbb{Z}^3} p_t(n\ell_T) = (p^0+\delta)^3$, using the product structure of p_t. By steps (2)–(3), this is at most $(a_*/\sqrt t + 4/\ell_T)^3$. Then $S_q\hat G_M^2 = \int_0^\infty t\mathrm{e}^{-tM^2}S_q\mathrm{e}^{-t\hat q^2}dt$. Integrating term by term, $\int_0^\infty t^{1/2+j/2-3/2+1-1}\cdots$ gives the four terms: $a_*^3\Gamma(\frac12)M^{-1}$, $3a_*^2 b\,M^{-2}$, $3a_*b^2\Gamma(\frac32)M^{-3}$ and b^3M^{-4}. Finally C_T is increasing in b, hence decreasing in ℓ. □

Lemma H.33 (the free mass difference). *For $M, m > 0$ and every ε, $|C^\infty(M^2) - C^\infty(m^2)| \le |M-m|\,\pi^2/32$.*

This is the free bound used in step (iv) of the proof of Proposition H.8.

Proof. Take $M < m$; the other case follows by symmetry. Then $C^\infty(M^2) - C^\infty(m^2) = (m^2 - M^2)\int_{\mathrm{BZ}} \frac{d^3k}{(2\pi)^3}\,[(\hat k^2 + M^2)(\hat k^2 + m^2)]^{-1} \geq 0$. Since $\sin x \geq 2x/\pi$ on $[0, \pi/2]$, $\hat k^2 \geq (4/\pi^2)|k|^2$ on the BZ. Substitute $q = (2/\pi)k$ and extend to $\mathbb{R}^3$: the integral is at most $(\pi/2)^3 \int_{\mathbb{R}^3} \frac{d^3q}{(2\pi)^3}[(q^2+M^2)(q^2+m^2)]^{-1} = (\pi^3/8)/(4\pi(M+m))$. Multiplying by $m^2 - M^2$ gives $(m-M)\pi^2/32$. □

Appendix I. The negative-source bound and the choice of parameters

We prove Proposition 6.7 (Subsection I.1), the domination of the infrared part of the massless photon by the slice covariance Γ^B (Subsection I.2), and the statements on the auxiliary parameters and on the hypotheses of Theorem 6.8 (Subsection I.3).

I.1. **The proof of Proposition 6.7.** Section 6 uses three facts that are proved here: the definition of $\Gamma_\geq$ and of the comparison covariance $C^\flat_{B,s}$, with the domination $C_{\mu_0} \leq C^\flat_{B,s}$ (Subsection 6.1); the translation-averaging lemma (step (1) of the proof of Proposition 6.7); and the determinant comparison for the infrared part with the constants J_{IR}, Z_T (step (3) there). With them, Proposition I.9 below is the full proof of Proposition 6.7.

Standing conventions. Fix $j \geq 1$, $T = T_j$, $\ell_T = \ell_j$ (the index i labels King's slices H_i below; j is the torus index), and $\varepsilon = L^{-K}\varepsilon_0$ with $K \geq 1$. By Lemma 2.1(ii) with $k = K$, $\ell_T/(M_b\varepsilon_0) \in 2\mathbb{N}$; hence the blocks of every scale $t_k = L^k\varepsilon$, $0 \leq k \leq K$, tile T_ε, which is what the definitions of G_k, H_i and Γ^B in Subsection 5.1 require. Covariances are operators on real bond functions, with the inner product of Subsection 2.2. $-\Delta_\varepsilon$ acts on each component; its eigenvalues are $\hat k^2 := \sum_\mu 4\varepsilon^{-2}\sin^2(\varepsilon k_\mu/2)$, where $k \in (2\pi/\ell_T)\mathbb{Z}^3$ has components in $[-\pi/\varepsilon, \pi/\varepsilon)$; $k = 0$ corresponds to $\operatorname{ran}\Pi_0$.

For a site function $b(\cdot)$ (a *profile*; a constant b is a special case), let $S^{\mathrm m}_{b(\cdot)}$ be the matter action of Definition 2.4 with single-site term $\frac12 b(x)\varepsilon^3 r_x^2$ at x. Put $Z_\varphi(A; b) := \int \mathrm{e}^{-S^{\mathrm m}_b(A,\varphi)} \prod_x d^2\varphi(x)$ and $\Psi^b_F(A) := \int F\,\mathrm{e}^{-S^{\mathrm m}_b(A,\varphi)} \prod_x d^2\varphi(x)$. For a covariance $\Gamma \geq 0$ put

$$\langle F\rangle_{\Gamma;b} := \frac{\mathbb{E}_{N(0,\Gamma)}[\Psi^b_F]}{\mathbb{E}_{N(0,\Gamma)}[Z_\varphi(\cdot;b)]}, \qquad Z(\Gamma; b) = \mathbb{E}_{N(0,\Gamma)}[Z_\varphi(\cdot;b)],$$

as in Subsection 5.1. If $\Gamma = \Gamma^\perp + \infty\cdot\Pi_0$ we mean the product of $N(0,\Gamma^\perp)$ on $\operatorname{ran}\Pi^\perp$ with the uniform probability measure on $[0,P)^3$ for c; we call this a *Haar zero mode*. Thus $\langle\cdot\rangle_{\varepsilon,T,\mu_0} = \langle\cdot\rangle_{C_{\mu_0};b_T}$ and $\langle\cdot\rangle_{\varepsilon,T,\flat} = \langle\cdot\rangle_{(-\Delta_\varepsilon)^{-1}\Pi^\perp + \infty\Pi_0;\, b_T}$ (Definition 2.7). Finally put $\rho_\Gamma(y;b) := \langle r_y^2 - W_\varepsilon - z_\varepsilon\rangle_{\Gamma;b}$ and $\bar\rho_\Gamma(b) := |T|^{-1}\sum_y \varepsilon^3\rho_\Gamma(y;b)$.

Definition I.1 (infrared part and slice-modified comparison covariance)**.** Let

$$D^< := \int_0^{\varepsilon_0^2} \mathrm{e}^{t\Delta_\varepsilon}(1 - \mathrm{e}^{-t\mu_*^2})\,dt\;\Pi^\perp, \qquad \Gamma_\geq := \int_{\varepsilon_0^2}^\infty \mathrm{e}^{t\Delta_\varepsilon}(1 - \mathrm{e}^{-t\mu_*^2})\,dt\;\Pi^\perp,$$

let $\gamma_\geq(u) := \int_{\varepsilon_0^2}^\infty \mathrm{e}^{-tu}(1 - \mathrm{e}^{-t\mu_*^2})\,dt$ for $u > 0$, so that $\Gamma_\geq = \gamma_\geq(-\Delta_\varepsilon)\Pi^\perp$, and put

$$C^\flat_{B,s} := (C_{\mu_*} + \Gamma^B + \Gamma_\geq)\Pi^\perp + s|T|\,\Pi_0 \quad (0 < s < \infty),$$
$$C^\flat_B := C^\flat_{B,\infty} \text{ (Haar zero mode)},$$
$$C^B_1 := C_{\mu_*} + \Gamma^B.$$

Since $\int_0^\infty \mathrm{e}^{-tu}(1 - \mathrm{e}^{-t\mu_*^2})\,dt = u^{-1} - (u+\mu_*^2)^{-1}$ for $u > 0$, and $-\Delta_\varepsilon > 0$ on $\operatorname{ran}\Pi^\perp$,

$$(-\Delta_\varepsilon)^{-1}\Pi^\perp = C_{\mu_*}\Pi^\perp + D^< + \Gamma_\geq, \qquad D^< \geq 0, \quad \Gamma_\geq \geq 0. \tag{I.1}$$

Lemma I.2 (structure)**.** *(a) G_k, H_i and Γ^B commute with Π_0, and $\Gamma^B \geq 0$. On* $\operatorname{ran}\Pi_0$, *Γ^B acts as a number $\gamma_0^B \geq 0$. Hence, with $s_1 := |T|^{-1}(\mu_*^{-2} + \gamma_0^B) \geq s(\mu_*)$,*

$$C_1^B = (C_{\mu_*} + \Gamma^B)\Pi^\perp + s_1|T|\,\Pi_0, \qquad C_{B,s}^\flat - C_1^B = \Gamma_\geq + (s - s_1)|T|\,\Pi_0.$$

(b) (Domination.) *If the constant κ in* (5.1) *satisfies $\kappa \geq \kappa_1\kappa_3$, with κ_1, κ_3 as in Theorem I.16, then $D^< \leq \Gamma^B\Pi^\perp$.*

Proof. (a) $\Delta_\varepsilon\Pi_0 = 0 = \Pi_0\Delta_\varepsilon$. The block average P_k is self-adjoint and fixes the constants, so $P_k\Pi_0 = \Pi_0 = \Pi_0 P_k$. Hence $G_k^{-1} = -\Delta_\varepsilon + \mu_*^2 + a_k t_k^{-2} P_k$ commutes with Π_0, and therefore so do G_k, $H_i = G_{i+1} - G_i$ and Γ^B. On a constant bond function, $G_k\mathbf{1}_\mu = (\mu_*^2 + a_k t_k^{-2})^{-1}\mathbf{1}_\mu$, the same number in every direction; hence Γ^B acts on $\operatorname{ran}\Pi_0$ as a number γ_0^B. Each H_i is a covariance, King's fluctuation covariance [24, (2.17)], so $H_i \geq 0$; hence $\Gamma^B = \sum_i \delta_i H_i \geq 0$ and $\gamma_0^B \geq 0$. On $\operatorname{ran}\Pi_0$, C_{μ_*} acts as $\mu_*^{-2} = s(\mu_*)|T|$. This gives the formula for C_1^B. The difference formula then follows from Definition I.1, since $\Gamma_\geq = \Gamma_\geq\Pi^\perp$. (b) Theorem I.16 states $\int_0^{\varepsilon_0^2} \mathrm{e}^{t\Delta_\varepsilon}(1 - \mathrm{e}^{-t\mu_*^2})\,dt \leq \kappa\mu_*^2 \sum_{i<K} t_{i+1}^2 H_i = \Gamma^B$ on the whole space, zero mode included. Both sides commute with Π_0, by (a), so compressing to $\operatorname{ran}\Pi^\perp$ gives the claim. □

Lemma I.3 (domination of the true photon)**.** *For every $\mu_0 \in (0, \mu_*]$ and every $s \geq s(\mu_0)$, $C_{\mu_0} \leq C_{B,s}^\flat$; also $C_s^\flat \leq C_{B,s}^\flat$ for every $s > 0$. Consequently, for every $h \geq 0$, every profile b and every $s \geq s(\mu_0)$,*

$$\langle \mathrm{e}^{-X_{\mathrm{R}}(h)}\rangle_{C_{\mu_0};b} \leq \langle \mathrm{e}^{-X_{\mathrm{R}}(h)}\rangle_{C_{B,s}^\flat;b} \xrightarrow[s\to\infty]{} \langle \mathrm{e}^{-X_{\mathrm{R}}(h)}\rangle_{C_B^\flat;b}.$$

This is the domination $C_{\mu_0} \leq C_{B,s}^\flat$ stated in Subsection 6.1.

Proof. All operators involved commute with Π_0 (Lemma I.2(a)), so it suffices to compare them on $\operatorname{ran}\Pi^\perp$ and on $\operatorname{ran}\Pi_0$ separately. On $\operatorname{ran}\Pi^\perp$, by (I.1) and Lemma I.2(b),

$$(-\Delta_\varepsilon + \mu_0^2)^{-1}\Pi^\perp \leq (-\Delta_\varepsilon)^{-1}\Pi^\perp = C_{\mu_*}\Pi^\perp + D^< + \Gamma_\geq \leq (C_{\mu_*} + \Gamma^B + \Gamma_\geq)\Pi^\perp.$$

On $\operatorname{ran}\Pi_0$, C_{μ_0} acts as $\mu_0^{-2} = s(\mu_0)|T|$, which is at most $s|T|$ for $s \geq s(\mu_0)$. The middle inequality above gives $C_s^\flat \leq C_{B,s}^\flat$ as well. Next, $\mathrm{e}^{-X_{\mathrm{R}}(h)} = \mathrm{e}^{\|h\|_{L^{1,\varepsilon}}(W_\varepsilon + z_\varepsilon)}\mathrm{e}^{-r^2(\varepsilon^3 h)}$ with $\varepsilon^3 h \geq 0$, a positive constant times an element of the class $\mathcal{D}$ of Lemma 7.5. By that lemma ([8, Cor. 6.3(1), p. 281] on the torus graph) its expectation does not decrease when the covariance increases. The limit $s \to \infty$ is Lemma I.4, applied with the bounded observable $\mathrm{e}^{-X_{\mathrm{R}}(h)}$. □

Lemma I.4 (Haar limit of the zero mode)**.** *Let $\Gamma^\perp \geq 0$ act on* $\operatorname{ran}\Pi^\perp$ *and let $\Gamma_s := \Gamma^\perp + s|T|\,\Pi_0$. Fix a profile b. Let G be either a bounded element of $\mathfrak{A}_\varepsilon$ or a polynomial in the r_x^2 with nonnegative coefficients. Then $\langle G\rangle_{\Gamma_s;b} \to \langle G\rangle_{\Gamma_\infty;b}$ as $s \to \infty$.*

Proof. By Lemma 2.5 (last sentence) and Lemma 2.6, $c \mapsto \Psi_G^b(A^\perp + c)$ is P-periodic in each c_μ. It is also bounded. For bounded G, $|\Psi_G^b| \leq \|G\|_\infty Z_\varphi(0; b)$. For $G = \prod_x r_x^{2n_x}$, the proof of Lemma 2.5 applies verbatim with the radial weight $v_x(r)$ replaced by $r^{2n_x}v_x(r)$, which is again nonnegative and integrable since $\lambda > 0$. This gives $\Psi_G^b(A) = \sum_n w_G(n)\mathrm{e}^{ie\varepsilon\sum_b n_b A_b}$ with $w_G \geq 0$, hence $|\Psi_G^b(A)| \leq \Psi_G^b(0) < \infty$. Under $N(0, \Gamma_s)$, $A^\perp$ and c are independent, the law of $A^\perp$ does not depend on s, and c_μ has variance s. Only $c \bmod P$ enters. Its law has density $P^{-1}\sum_{\nu\in\mathbb{Z}} \mathrm{e}^{-2\pi^2\nu^2 s/P^2}\cos(2\pi\nu x/P)$ in each coordinate, which converges uniformly to $1/P$, hence in total variation to the uniform law (as in the proof of Lemma 7.6). By bounded convergence the numerator and the denominator $\mathbb{E}[Z_\varphi(\cdot; b)]$ converge, and the limit of the denominator is positive. □

Lemma I.5 (one-point reduction)**.** *Let $\Gamma > 0$ be a covariance on bond functions, $a(\cdot)$ a profile, and $h \geq 0$. Then for every profile $a' \geq a + 2h$ (sitewise)*

$$\log\langle \mathrm{e}^{-X_{\mathrm{R}}(h)}\rangle_{\Gamma;a} = -\int_0^1 \sum_y \varepsilon^3 h(y)\,\rho_\Gamma(y; a + 2th)\,dt \leq -\sum_y \varepsilon^3 h(y)\,\rho_\Gamma(y; a').$$

Proof. Let $f(t) := \log\langle \mathrm{e}^{-tX_{\mathrm{R}}(h)}\rangle_{\Gamma;a}$ for $t \in [0,1]$; then $f(0) = 0$. Since $\mathrm{e}^{-tX_{\mathrm{R}}(h)}$ is a c-number times $\exp(-t\sum_x \varepsilon^3 h(x) r_x^2)$ and the single-site term is $\frac{1}{2}a(x)\varepsilon^3 r_x^2$, the tilted measure is the model with profile $a + 2th$; the c-numbers cancel in the normalisation. Differentiation under the integral is justified because $\mathrm{e}^{-t\sum \varepsilon^3 h r^2} \le 1$ and r_x^2 is integrable ($\lambda > 0$). Hence $f'(t) = -\sum_y \varepsilon^3 h(y)\rho_\Gamma(y; a + 2th)$. Moreover $\partial\langle r_y^2\rangle_{\Gamma;b}/\partial b(x) = -\frac{1}{2}\varepsilon^3\langle r_y^2; r_x^2\rangle_{\Gamma;b} \le 0$. This is [8, Thm. 6.1, p. 276] with $P = r_y^2$, $Q = r_x^2$, $n = m = 0$ and zero linear functionals, on the torus graph: Lemma B.10 in the instance of Proposition B.13, with b_T replaced by the profile, which Definition B.8 allows because the single-site measures ν_x may depend on x. It holds for any centred Gaussian $\Gamma > 0$. So $\rho_\Gamma(y;\cdot)$ does not increase along the straight path from $a + 2th$ to $a' \ge a + 2th$. With $h \ge 0$ this gives $f'(t) \le -\sum_y \varepsilon^3 h(y)\rho_\Gamma(y; a')$; integrate over $t \in [0,1]$. □

Lemma I.6 (translation averaging of the source)**.** *Let $0 \le h \le H$, $\mu_0 \in (0, \mu_*]$, and let $a' \ge b_T + 2H$ be a constant. Then*

$$\log\langle \mathrm{e}^{-X_{\mathrm{R}}(h)}\rangle_{\varepsilon,T,\mu_0} \le -\|h\|_{L^1,\varepsilon}\,\bar\rho^B(a'), \qquad \bar\rho^B := \bar\rho_{C_B^\flat}.$$

Proof. Let $\mathcal{T}$ be the group of the $n := |T|/\varepsilon^3$ lattice translations of T_ε, and put $(\tau h)(y) := h(y-\tau)$.

(i) The state $\langle\cdot\rangle_{\varepsilon,T,\mu_0}$ is invariant under $\mathcal{T}$ (Lemma 7.13(i)), and $\mathrm{e}^{-X_{\mathrm{R}}(h)}$ composed with the translation by τ is $\mathrm{e}^{-X_{\mathrm{R}}(\tau h)}$; the c-number $\|\tau h\|_{L^1,\varepsilon}(W_\varepsilon + z_\varepsilon) = \|h\|_{L^1,\varepsilon}(W_\varepsilon + z_\varepsilon)$ is unchanged. Hence $\langle \mathrm{e}^{-X_{\mathrm{R}}(h)}\rangle_{\varepsilon,T,\mu_0} = \langle \mathrm{e}^{-X_{\mathrm{R}}(\tau h)}\rangle_{\varepsilon,T,\mu_0}$ for every $\tau \in \mathcal{T}$.

(ii) Fix $s \ge s(\mu_0)$. By Lemma I.3 applied to $\tau h \ge 0$, and then Lemma I.5 with $\Gamma = C^\flat_{B,s} > 0$, $a \equiv b_T$ and $a' \ge b_T + 2H \ge b_T + 2\tau h$,

$$\log\langle \mathrm{e}^{-X_{\mathrm{R}}(h)}\rangle_{\varepsilon,T,\mu_0} \le \log\langle \mathrm{e}^{-X_{\mathrm{R}}(\tau h)}\rangle_{C^\flat_{B,s};b_T} \le -\sum_y \varepsilon^3 h(y-\tau)\,\rho_{C^\flat_{B,s}}(y;a').$$

(iii) The left side does not depend on τ. Average the right side over $\tau \in \mathcal{T}$, using $n^{-1}\sum_\tau h(y-\tau) = n^{-1}\sum_x h(x) = \|h\|_{L^1,\varepsilon}/|T|$ for every y (as $h \ge 0$):

$$\log\langle \mathrm{e}^{-X_{\mathrm{R}}(h)}\rangle_{\varepsilon,T,\mu_0} \le -\frac{\|h\|_{L^1,\varepsilon}}{|T|}\sum_y \varepsilon^3 \rho_{C^\flat_{B,s}}(y;a') = -\|h\|_{L^1,\varepsilon}\,\bar\rho_{C^\flat_{B,s}}(a').$$

(iv) Let $s \to \infty$. By Lemma I.4, with $G = r_y^2$ and $G = 1$ at the constant profile a', $\rho_{C^\flat_{B,s}}(y;a') \to \rho^B(y;a')$. □

Remark I.7. One may also average over $\varepsilon\mathbb{Z}^3/\varepsilon_0\mathbb{Z}^3$ only, and then use that $y \mapsto \rho^B(y;b)$ is $\varepsilon_0\mathbb{Z}^3$-periodic. This holds because Γ^B commutes with the translations by $\varepsilon_0\mathbb{Z}^3 = t_K\mathbb{Z}^3$ (not with those by $\varepsilon\mathbb{Z}^3$ in general). The proof above averages over all of $\mathcal{T}$ and does not need this periodicity; only the $\varepsilon\mathbb{Z}^3$-invariance of the true model is used. That invariance fails on a cube with boundary conditions, which is why the lemma is a torus statement. The choice $a' = \hat a_1$, which Lemma I.5 allows, makes a separate step for the monotonicity of ρ^B in the constant coefficient unnecessary.

Lemma I.8 (determinant comparison for the infrared part; J_{IR} and Z_T)**.** *Put $\delta_P := 2\sum_{\nu\ge1}\mathrm{e}^{-2\pi^2\nu^2}$ ($< 5.4\cdot10^{-9}$), and*

$$J_T := \frac{3}{2}|T|^{-1}\sum_{k\neq0}(\hat k^2 + \mu_*^2)\,\gamma_{\ge}(\hat k^2),$$

$$Z_T := |T|^{-1}\Big[\tfrac{3}{2}\log_+\frac{4\pi^2\mu_*^2\ell_T}{e^2} + 3\delta_P\Big],$$

$J_{\mathrm{IR}} := \sup_{K\ge1,\,j\ge1} J_{T_j}$ and $\bar Z := \sup_{j\ge1} Z_{T_j}$. For every constant a:

(i) *(density ratio) if $0 < \Gamma \le \Gamma'$ on $\mathbb{R}^n$ and $f \ge 0$, then $\mathbb{E}_{N(0,\Gamma')}[f] \ge \det(\Gamma'\Gamma^{-1})^{-1/2}\mathbb{E}_{N(0,\Gamma)}[f]$;*

(ii) *(upper) $Z(C_B^\flat; a) \le Z(C_1^B; a)$;*

(iii) *(lower)* $\log Z(C_B^\flat;a) \ge \log Z(C_1^B;a) - (J_T+Z_T)|T|$;
(iv) *with* $\alpha := \sqrt{\pi}/4$,

$$J_{\mathrm{IR}} \le \tfrac{3}{2}(\alpha/\varepsilon_0+1/\ell_0)^3+3\mu_*^2(\alpha+1)^3\big(1/\varepsilon_0+1/(32\ell_0)\big), \qquad \bar Z \le \ell_0^{-3}\big[\tfrac{3}{2}\log_+(4\pi^2\mu_*^2\ell_0/e^2)+0.19+3\delta_P\big].$$

So J_{IR} depends on $\mu_*, \varepsilon_0, \ell_0$ only, and $\bar Z$ on μ_*, ℓ_0, e only.

Proof. (i) $\frac{dN(0,\Gamma')}{dN(0,\Gamma)}(x) = (\det\Gamma/\det\Gamma')^{1/2}\exp(\frac{1}{2}x\cdot(\Gamma^{-1}-\Gamma'^{-1})x) \ge (\det\Gamma/\det\Gamma')^{1/2}$, since $\Gamma^{-1} \ge \Gamma'^{-1}$.

(ii) By Lemma 2.5, $Z_\varphi(A;a) = \sum_n w(n)\mathrm{e}^{ie\varepsilon\sum_b n_b A_b}$ with $w \ge 0$ and $\sum_n w(n) < \infty$. By Fubini, $Z(\Gamma;a) = \sum_n w(n)\exp(-\frac{1}{2}e^2\varepsilon^2\operatorname{Var}_\Gamma(\sum_b n_b A_b))$ for every centred Gaussian Γ, and this is non-increasing in Γ in form order. By Lemma I.2(a), $C_{B,s}^\flat \ge C_1^B$ for $s \ge s_1$, so $Z(C_{B,s}^\flat;a) \le Z(C_1^B;a)$. Let $s\to\infty$ (Lemma I.4 with $G=1$; the unnormalised integrals converge by the same bounded-convergence argument).

(iii) *Zero mode.* Let $F(c) := \mathbb{E}_{N(0,(C_{\mu_*}+\Gamma^B+\Gamma_\ge)\Pi^\perp)}[Z_\varphi(A^\perp+c;a)] \ge 0$. It is P-periodic in each c_μ (Lemma 2.5). Put $s' := \max(P^2, s_1)$. For variance $s' \ge P^2$ the wrapped normal density on a circle of length P deviates from $1/P$ by at most $P^{-1}\cdot 2\sum_{\nu\ge1}\mathrm{e}^{-2\pi^2\nu^2 s'/P^2} \le \delta_P/P$. Hence $\mathbb{E}_{N(0,s')^{\otimes3}}[F] \le (1+\delta_P)^3P^{-3}\int_{[0,P)^3}F$, that is, $Z(C_B^\flat;a) \ge (1+\delta_P)^{-3}Z(C_{B,s'}^\flat;a)$. By (i) on the three zero-mode coordinates, $Z(C_{B,s'}^\flat;a) \ge (s_1/s')^{3/2}Z(C_{B,s_1}^\flat;a)$. Since $\log(1+\delta_P)\le\delta_P$, $\log(s'/s_1) = \log_+(P^2/s_1)$, $s_1 \ge s(\mu_*)$ (Lemma I.2(a)) and $P^2/s(\mu_*) = 4\pi^2\mu_*^2\ell_T/e^2$ (with $P = 2\pi/(|e|\ell_T)$ and $s(\mu_*) = (\mu_*^2\ell_T^3)^{-1}$),

$$\log Z(C_B^\flat;a) \ge \log Z(C_{B,s_1}^\flat;a) - Z_T|T|.$$

Infrared part. By Lemma I.2(a), $C_{B,s_1}^\flat = C_1^B + \Gamma_\ge$, with $C_1^B \ge C_{\mu_*} > 0$. By (i) with $f = Z_\varphi(\cdot;a) \ge 0$, and $\log\det(1+K) \le \operatorname{tr}K$ for $K \ge 0$,

$$\log Z(C_{B,s_1}^\flat;a) \ge \log Z(C_1^B;a) - \tfrac{1}{2}\operatorname{tr}\big((C_1^B)^{-1}\Gamma_\ge\big) \ge \log Z(C_1^B;a) - \tfrac{1}{2}\operatorname{tr}\big(C_{\mu_*}^{-1}\Gamma_\ge\big).$$

The last step uses $(C_1^B)^{-1} \le C_{\mu_*}^{-1}$ (the inverse is operator monotone) and $\operatorname{tr}\big((C_1^B)^{-1}\Gamma_\ge\big) = \operatorname{tr}\big(\Gamma_\ge^{1/2}(C_1^B)^{-1}\Gamma_\ge^{1/2}\big)$. Finally, $C_{\mu_*}^{-1}\Gamma_\ge = (-\Delta_\varepsilon+\mu_*^2)\gamma_\ge(-\Delta_\varepsilon)\Pi^\perp$ has the eigenvalues $(\hat k^2+\mu_*^2)\gamma_\ge(\hat k^2)$, $k\neq0$, each three times (three components), so $\frac{1}{2}\operatorname{tr}(C_{\mu_*}^{-1}\Gamma_\ge) = J_T|T|$. Combining the two displays gives (iii).

(iv) Since $1-\mathrm{e}^{-t\mu_*^2} \le 1$, $\gamma_\ge(u) \le \mathrm{e}^{-\varepsilon_0^2u}/u$, so $(u+\mu_*^2)\gamma_\ge(u) \le \mathrm{e}^{-\varepsilon_0^2u} + \mu_*^2\mathrm{e}^{-\varepsilon_0^2u}/u$. With $q^T(t) := |T|^{-1}\sum_k \mathrm{e}^{-t\hat k^2}$,

$$J_T \le \tfrac{3}{2}\big[q^T(\varepsilon_0^2) - |T|^{-1}\big] + \tfrac{3}{2}\mu_*^2\int_{\varepsilon_0^2}^\infty \big[q^T(t) - |T|^{-1}\big]\,dt.$$

Now $q^T(t) = q_1(t)^3$, with $q_1(t) = \ell_T^{-1}\sum_{k_1}\mathrm{e}^{-4t\varepsilon^{-2}\sin^2(\varepsilon k_1/2)}$. Since $|\sin x| \ge 2|x|/\pi$ for $|x| \le \pi/2$, we have $4\varepsilon^{-2}\sin^2(\varepsilon k_1/2) \ge 4k_1^2/\pi^2$, which equals $16\nu^2/\ell_T^2$ for $k_1 = 2\pi\nu/\ell_T$. Hence

$$q_1(t) \le \ell_T^{-1}\sum_{\nu\in\mathbb{Z}}\mathrm{e}^{-16t\nu^2/\ell_T^2} \le \ell_T^{-1} + \ell_T^{-1}\int_{\mathbb{R}}\mathrm{e}^{-16tx^2/\ell_T^2}dx = \ell_T^{-1} + \alpha t^{-1/2}.$$

The same inequality gives the spectral gap $\min_{k\neq0}\hat k^2 \ge 16/\ell_T^2$, so $q^T(t)-|T|^{-1} \le \mathrm{e}^{-16(t-\ell_T^2)/\ell_T^2}\big(q^T(\ell_T^2)-|T|^{-1}\big)$ for $t \ge \ell_T^2$. For $t \le \ell_T^2$, $q^T(t) \le (\alpha+1)^3t^{-3/2}$. Therefore (using $\ell_T \ge \ell_0 \ge \varepsilon_0$)

$$\int_{\varepsilon_0^2}^\infty \big[q^T - |T|^{-1}\big]dt \le \int_{\varepsilon_0^2}^\infty (\alpha+1)^3t^{-3/2}dt + \frac{\ell_T^2}{16}(\alpha+1)^3\ell_T^{-3} \le (\alpha+1)^3\Big(\frac{2}{\varepsilon_0}+\frac{1}{16\ell_0}\Big),$$

and $q^T(\varepsilon_0^2) - |T|^{-1} \le (\alpha/\varepsilon_0+1/\ell_0)^3$. Neither bound depends on ε, K or j, which proves the bound on J_{IR}.

For $\bar{Z}$: since $\ell_{T_j} \geq \ell_0$, $\bar{Z} \leq \sup_{\ell_T \geq \ell_0} Z_T$, and Lemma B.27 gives the bound, because $1/(2\mathrm{e}) < 0.19$; the estimate there uses the subadditivity of $\log_+$ and $\max_{y\geq 1} y^{-3}\log y = 1/(3\mathrm{e})$. □

Proposition I.9 (Proposition 6.7, full proof)**.** *Assume the slice vacuum bound $\Delta_1^B < \infty$ and the anchor condition* (6.3)*: $b_T + 2H \leq \hat{a}_1$ on every torus of $\mathfrak{C}$, and $g_{21} > 0$ (Theorem 6.1(b),(c)). Then for every $0 \leq h \leq H$, every $\mu_0 \in (0, \mu_*] \cup \{\flat\}$ and every $(\varepsilon, T_j) \in \mathfrak{C}$,*

$$\log\langle \mathrm{e}^{-X_{\mathrm{R}}(h)}\rangle_{\varepsilon,T_j,\mu_0} \leq 2\|h\|_{L^1,\varepsilon}\, S_\flat^B, \qquad S_\flat^B := [\Delta_1^B + J_{\mathrm{IR}} + \bar{Z}]_+/g_{21},$$

with J_{IR}, $\bar{Z}$ as in Lemma I.8. Since $Z_{T_j} \leq \bar{Z}$, this is Proposition 6.7, with Z_T there read as $\bar{Z}$, so that $S_\flat^B$ does not depend on T.

Proof. Write $\hat{a}_i := a_{K,T}(m_i)$ and $\rho^B := \rho_{C_B^\flat}$, $\bar{\rho}^B := \bar{\rho}_{C_B^\flat}$.

Case $\mu_0 \in (0, \mu_*]$. (1) By Lemma I.6 with $a' := \hat{a}_1 \geq b_T + 2H$, $\log\langle \mathrm{e}^{-X_{\mathrm{R}}(h)}\rangle_{\varepsilon,T,\mu_0} \leq -\|h\|_{L^1,\varepsilon}\,\bar{\rho}^B(\hat{a}_1)$.

(2) Let $f^B(b) := |T|^{-1}\log Z(C_B^\flat; b) + \frac{1}{2}b(W_\varepsilon + z_\varepsilon)$ for constant b. Since $\partial_b Z_\varphi(A;b) = -\frac{1}{2}\sum_x \varepsilon^3 \Psi^b_{r_x^2}(A)$, and on $b \geq b_{\min}$ we have $|\Psi^b_{r_x^2}(A)| \leq \Psi^{b_{\min}}_{r_x^2}(0)$ (positive type, as in the proof of Lemma I.4), we may differentiate under $\mathbb{E}_{C_B^\flat}$, twice. Using $\sum_x \varepsilon^3 = |T|$,

$$f^{B\prime}(b) = -\tfrac{1}{2}|T|^{-1}\sum_x \varepsilon^3\big(\langle r_x^2\rangle_{C_B^\flat;b} - W_\varepsilon - z_\varepsilon\big) = -\tfrac{1}{2}\bar{\rho}^B(b),$$

$$f^{B\prime\prime}(b) = \tfrac{1}{4}|T|^{-1}\,\mathrm{Var}_{C_B^\flat;b}\Big(\sum_x \varepsilon^3 r_x^2\Big) \geq 0.$$

So f^B is convex, and since $\hat{a}_2 - \hat{a}_1 \geq g_{21} > 0$, $-\bar{\rho}^B(\hat{a}_1) = 2f^{B\prime}(\hat{a}_1) \leq 2[f^B(\hat{a}_2) - f^B(\hat{a}_1)]/(\hat{a}_2 - \hat{a}_1)$.

(3) By Lemma I.8(ii) at $\hat{a}_2$ and (iii) at $\hat{a}_1$, and since $J_T \leq J_{\mathrm{IR}}$, $Z_T \leq \bar{Z}$ and by the definition of Δ_1^B,

$$f^B(\hat{a}_2) - f^B(\hat{a}_1) \leq |T|^{-1}\log\frac{Z(C_1^B;\hat{a}_2)}{Z(C_1^B;\hat{a}_1)} + \tfrac{1}{2}(\hat{a}_2 - \hat{a}_1)(W_\varepsilon + z_\varepsilon) + J_{\mathrm{IR}} + \bar{Z} \leq \Delta_1^B + J_{\mathrm{IR}} + \bar{Z}.$$

(4) Combine (1)–(3) with $\|h\|_{L^1,\varepsilon} \geq 0$:

$$\log\langle \mathrm{e}^{-X_{\mathrm{R}}(h)}\rangle_{\varepsilon,T,\mu_0} \leq \frac{2\|h\|_{L^1,\varepsilon}[\Delta_1^B + J_{\mathrm{IR}} + \bar{Z}]}{\hat{a}_2 - \hat{a}_1} \leq \frac{2\|h\|_{L^1,\varepsilon}[\Delta_1^B + J_{\mathrm{IR}} + \bar{Z}]_+}{g_{21}}.$$

Case $\flat$. $G = \mathrm{e}^{-X_{\mathrm{R}}(h)}$ with $h \geq 0$ is an elementary observable of Lemma 7.6, so $\langle G\rangle_{\varepsilon,T,\flat} = \lim_{\mu_0\downarrow 0}\langle G\rangle_{\varepsilon,T,\mu_0}$. The bound just proved does not depend on μ_0, so it passes to the limit.

Uniformity. Δ_1^B is a supremum over $\mathfrak{C}$ (Subsection 6.5). By Lemma I.8(iv), J_{IR} and $\bar{Z}$ do not depend on ε, j, μ_0 or h, and g_{21} is an infimum over ε and T. No renormalisation-group statement carries a source; the bound is linear in $\|h\|_{L^1,\varepsilon}$. □

I.2. **The domination of the infrared part.** This subsection proves the domination of the infrared part $D^<$ by the slice covariance Γ^B of (5.1). Lemma I.2(b) and the constant κ of Subsection 5.1 use it. Only the free (Gaussian) photon enters: there is no Higgs field and no observable.

Definition I.10 (setting)**.** Fix $j \geq 1$, $K \geq 1$, $\varepsilon = L^{-K}\varepsilon_0$ and $t_k := L^k\varepsilon$ for $0 \leq k \leq K$, so that $t_K = \varepsilon_0$. Let $\Lambda := T_{j,\varepsilon}$, with $\langle f, g\rangle := \varepsilon^3\sum_{x\in\Lambda} f(x)g(x)$ on real functions on Λ. $-\Delta_\varepsilon$ is the periodic lattice Laplacian, $\langle\varphi, (-\Delta_\varepsilon)\varphi\rangle = \varepsilon\sum_{\{x,y\}}(\varphi(x) - \varphi(y))^2$, where the sum runs over all nearest-neighbour pairs of $T_{j,\varepsilon}$. Let $M := -\Delta_\varepsilon + \mu_*^2$, so that $C_{\mu_*} = M^{-1}$.

(a) *Blocks.* For $0 \leq k \leq K$ let $\mathcal{P}_k := \{B^k(y) : y \in t_k\mathbb{Z}^3 \cap T_j\}$, where $B^k(y) := \{x \in \Lambda : y_i \leq x_i < y_i + t_k\}$ [24, p. 653]. Let $\mathcal{B}_k$ be the functions constant on each block of $\mathcal{P}_k$, and let P_k be the orthogonal projection onto $\mathcal{B}_k$, i.e. the block average $Q_k^*Q_k$ of [24, (2.10)]. Thus

$P_0 = 1$; this P_0 is not the projection Π_0 onto the constants. Moreover $P_{k+1} \le P_k$ and $P_k P_{k+1} = P_{k+1} P_k = P_{k+1}$.

(b) *Coefficients.* $a_k := a(1-L^{-2})(1-L^{-2k})^{-1}$ [24, p. 653], $a_\flat := a(1-L^{-2})$ and $r_k := a_k t_k^{-2}$.

(c) *Covariances.* $G_k := (M + r_k P_k)^{-1}$ for $1 \le k \le K$ and $G_0 := 0$; $H_j := G_{j+1} - G_j$; $D := \sum_{j<K} t_{j+1}^2 H_j$, so that $\Gamma^B = \kappa \mu_*^2 D$ in (5.1); and

$$D^{\natural <} := \int_0^{\varepsilon_0^2} \mathrm{e}^{t\Delta_\varepsilon}\big(1 - \mathrm{e}^{-t\mu_*^2}\big)\, dt, \qquad \text{so that } D^< = D^{\natural <}\Pi^\perp \text{ (Remark 5.1).}$$

(d) *Constants.*

$$\kappa_1 := \max\Big\{\frac{1}{8},\ \frac{L^2/8 + 1/a_\flat}{L^2-1},\ \frac{L^4}{(L^2-1)^2 a_\flat}\Big\}, \qquad \kappa_3 := 2(1+\mu_*^2\varepsilon_0^2)^2 + a_K^2.$$

$-\Delta_\varepsilon$ and P_k act on each component of a bond function separately, and so does every operator above. An inequality between such operators is therefore the same inequality in each component, and we work with one component throughout. $A \le B$ means $\langle \varphi, A\varphi\rangle \le \langle \varphi, B\varphi\rangle$ for all φ.

Elementary facts. (F1) We have $a_1 = a$ and $a_{k+1}/a_k = (1-L^{-2k})/(1-L^{-2k-2}) < 1$. Hence $a \ge a_k > a_\flat$, and $r_m/r_K = L^{2(K-m)} a_m/a_K \ge L^{2(K-m)}$ for $m \le K$; in particular $r_{K-1} - r_K \ge r_{K-1}(1-L^{-2})$. (F2) For $k \le K$, Lemma 2.1(ii) gives $\ell_j/(M_b t_k) \in 2\mathbb{N}$. So $\mathcal{P}_k$ is a partition of Λ into lattice cubes with L^k sites per side, and each $\mathcal{P}_{k+1}$-block is a union of $\mathcal{P}_k$-blocks. Since $\ell_j \ge 2t_k$, no block contains both ends of a wrap-around bond of the torus, so the nearest-neighbour pairs inside a block are exactly those of the cube. (F3) The block-Neumann forms satisfy

$$\langle \varphi, (-\Delta_\varepsilon)\varphi\rangle \ \ge \sum_{\square \in \mathcal{P}_k} \varepsilon \sum_{\{x,y\}\subset\square} (\varphi(x) - \varphi(y))^2.$$

The left side sums over all pairs and the right side only over pairs inside blocks. (F4) $G_K \le M^{-1}$, since $G_K^{-1} = M + r_K P_K \ge M$.

Lemma I.11 (block Poincaré inequality)**.** *Let $1 \le k \le K$. For all φ, v:*

(a) $\langle \varphi, (-\Delta_\varepsilon)\varphi\rangle \ge (8/t_k^2)\|(1-P_k)\varphi\|^2$;

(b) *if $P_k v = 0$, then* $\langle v, M^{-1} v\rangle \le (t_k^2/8)\|v\|^2$;

(c) $\langle v, G_k v\rangle \le (t_k^2/8)\|(1-P_k)v\|^2 + r_k^{-1}\|P_k v\|^2$.

Proof. (a) Let $\square$ be a cube with $b = L^k \ge 2$ sites per side. With respect to $\langle\cdot,\cdot\rangle$, the form $\varepsilon\sum_{\{x,y\}\subset\square}(\varphi(x)-\varphi(y))^2$ belongs to the operator $-\Delta_{N,\square}$, ε^{-2} times the graph Laplacian of $\square$. Its eigenvalues are $\varepsilon^{-2}\sum_{i=1}^3 4\sin^2(\pi n_i/(2b))$ with $0 \le n_i < b$. Its kernel consists of the constants on $\square$, and its first nonzero eigenvalue is $(4/\varepsilon^2)\sin^2(\pi/(2b))$. The map $b \mapsto 4b^2\sin^2(\pi/(2b)) = \pi^2(\sin y/y)^2$, $y = \pi/(2b)$, increases in b and equals 8 at $b = 2$. Hence that eigenvalue is at least $8/(b\varepsilon)^2 = 8/t_k^2$. The constant part of φ on $\square$ is $(P_k\varphi)|_\square$. Summing over $\square \in \mathcal{P}_k$ and using (F3) gives (a). (b), (c) For $B > 0$ we have $\langle v, B^{-1}v\rangle = \sup_\varphi[2\langle v,\varphi\rangle - \langle\varphi, B\varphi\rangle]$. For (b): $\langle v,\varphi\rangle = \langle v, (1-P_k)\varphi\rangle \le \|v\|\,(t_k^2/8)^{1/2}\langle\varphi,(-\Delta_\varepsilon)\varphi\rangle^{1/2} \le \|v\|\,(t_k^2/8)^{1/2}\langle\varphi, M\varphi\rangle^{1/2}$ by (a), and $\sup_{s\ge0}(2\alpha s - s^2) = \alpha^2$. For (c), take $B = M + r_k P_k$. Then $\langle v,\varphi\rangle = \langle(1-P_k)v, (1-P_k)\varphi\rangle + \langle P_k v, P_k\varphi\rangle \le \alpha s + \beta\sigma$, with $\alpha := (t_k^2/8)^{1/2}\|(1-P_k)v\|$, $s := \langle\varphi, M\varphi\rangle^{1/2}$ (using (a)), $\beta := \|P_k v\|$ and $\sigma := \|P_k\varphi\|$. Since $\langle\varphi, B\varphi\rangle = s^2 + r_k\sigma^2$ and $\sup_{s,\sigma}[2\alpha s + 2\beta\sigma - s^2 - r_k\sigma^2] = \alpha^2 + \beta^2/r_k$, (c) follows. □

Lemma I.12 (Abel summation)**.** $D = \sum_{m=0}^{K-1} w_m(G_K - G_m)$, *where* $w_0 := t_1^2$ *and* $w_m := t_{m+1}^2 - t_m^2 = (L^2-1)t_m^2$ *for* $m \ge 1$. *Moreover* $G_K - G_m \ge 0$ *for* $0 \le m \le K$.

Proof. Since $\sum_{m=0}^j w_m = t_{j+1}^2$,

$$D = \sum_{j<K}\sum_{m\le j} w_m(G_{j+1} - G_j) = \sum_{m<K} w_m \sum_{j=m}^{K-1}(G_{j+1} - G_j) = \sum_{m<K} w_m(G_K - G_m).$$

For $m=0$, $G_K \ge 0$. For $1 \le m < K$, $G_m^{-1} - G_K^{-1} = r_m P_m - r_K P_K = r_m(P_m - P_K) + (r_m - r_K)P_K \ge 0$, by $P_K \le P_m$ and (F1). Inversion is operator monotone. □

Lemma I.13 (one summand, dual cost)**.** *Let $Y := G_K^{-1} = M + r_K P_K$. For $1 \le m \le K-1$, every $v \in \mathcal{B}_m$ and every φ,*

$$\langle \varphi,\, G_K v\rangle^2 \le \langle \varphi,\, (G_K - G_m)\varphi\rangle\, c_m(v), \qquad c_m(v) := \langle v,\, G_K v\rangle + r_m^{-1}\|(P_m - P_K)v\|^2 + (r_m - r_K)^{-1}\|P_K v\|^2.$$

For $m = 0$ and every v: $\langle \varphi,\, G_K v\rangle^2 \le \langle \varphi,\, G_K \varphi\rangle\, c_0(v)$, with $c_0(v) := \langle v,\, G_K v\rangle$.

Proof. The case $m=0$ is the Cauchy–Schwarz inequality for the form of G_K. Let $m \ge 1$ and put $E_m := r_m(P_m - P_K) + (r_m - r_K)P_K \ge 0$. Then $G_m^{-1} = Y + E_m$, and E_m acts on $\mathcal{B}_m = \operatorname{ran} P_m$ as r_m on $\operatorname{ran}(P_m - P_K)$ and as $r_m - r_K > 0$ on $\operatorname{ran} P_K$, and vanishes on $\mathcal{B}_m^{\perp}$. For $\eta > 0$ let $E_\eta := E_m + \eta$ and $B_\eta := Y^{-1} - (Y + E_\eta)^{-1} = Y^{-1}E_\eta(Y + E_\eta)^{-1} > 0$. Its inverse is $(Y + E_\eta)E_\eta^{-1}Y = Y + YE_\eta^{-1}Y$. With $f := G_K v = Y^{-1}v$ this gives $\langle f,\, B_\eta^{-1} f\rangle = \langle v,\, Y^{-1}v\rangle + \langle v,\, E_\eta^{-1}v\rangle$, and Cauchy–Schwarz in the B_η-product gives $\langle \varphi,\, f\rangle^2 \le \langle \varphi,\, B_\eta\varphi\rangle\langle f,\, B_\eta^{-1} f\rangle$. As $\eta \downarrow 0$ (finite dimension), $B_\eta \to Y^{-1} - G_m = G_K - G_m$. For $v \in \mathcal{B}_m$,

$$\langle v,\, E_\eta^{-1}v\rangle = (r_m + \eta)^{-1}\|(P_m - P_K)v\|^2 + (r_m - r_K + \eta)^{-1}\|P_K v\|^2,$$

which tends to the last two terms of $c_m(v)$. □

Lemma I.14 (block lemma)**.** $G_K^2 \le \kappa_1 D$.

Proof. Case $K \ge 2$. Fix φ and u. Decompose u into the mutually orthogonal pieces

$$v_0 := (1 - P_1)u, \qquad v_m := (P_m - P_{m+1})u \ \ (1 \le m \le K-2),$$
$$v_{K-1} := P_{K-1}u = v' + v'', \qquad v' := (P_{K-1} - P_K)u, \quad v'' := P_K u,$$

so that $v_m \in \mathcal{B}_m$ and $\|v_0\|^2 + \sum_{1 \le m \le K-2}\|v_m\|^2 + \|v'\|^2 + \|v''\|^2 = \|u\|^2$. By Lemma I.13, Cauchy–Schwarz over m, and Lemma I.12,

$$\langle \varphi,\, G_K u\rangle = \sum_{m<K} \langle \varphi,\, G_K v_m\rangle \le \sum_{m<K} \big(w_m \langle \varphi,\, (G_K - G_m)\varphi\rangle\big)^{1/2} \Big(\frac{c_m(v_m)}{w_m}\Big)^{1/2}$$
$$\le \langle \varphi,\, D\varphi\rangle^{1/2} \Big(\sum_{m<K} \frac{c_m(v_m)}{w_m}\Big)^{1/2}.$$

We bound the costs using Lemma I.11, (F1) and (F4).

- $m = 0$: $P_1 v_0 = 0$, so $c_0 \le \langle v_0,\, M^{-1}v_0\rangle \le (t_1^2/8)\|v_0\|^2$ and $c_0/w_0 \le \frac{1}{8}\|v_0\|^2$.
- $1 \le m \le K-2$: $P_{m+1}v_m = 0 = P_K v_m$, so $c_m \le (t_{m+1}^2/8 + t_m^2/a_m)\|v_m\|^2$ and $c_m/w_m \le \frac{L^2/8 + 1/a_\flat}{L^2 - 1}\|v_m\|^2$.
- $m = K-1$: $(P_{K-1} - P_K)v_{K-1} = v'$ and $P_K v_{K-1} = v''$. Lemma I.11(c) with $k = K$ gives $\langle v_{K-1},\, G_K v_{K-1}\rangle \le (t_K^2/8)\|v'\|^2 + (t_K^2/a_K)\|v''\|^2$. With $r_{K-1}^{-1} = t_{K-1}^2/a_{K-1}$ and $(r_{K-1} - r_K)^{-1} \le L^2 t_{K-1}^2/((L^2-1)a_{K-1})$,

$$\frac{c_{K-1}}{w_{K-1}} \le \frac{L^2/8 + 1/a_\flat}{L^2 - 1}\|v'\|^2 + \frac{L^2/a_\flat + L^2/((L^2-1)a_\flat)}{L^2 - 1}\|v''\|^2,$$

  and the last coefficient equals $L^4/((L^2-1)^2 a_\flat)$.

Hence $\sum_m c_m(v_m)/w_m \le \kappa_1\|u\|^2$, i.e. $\langle G_K\varphi,\, u\rangle^2 \le \kappa_1\langle \varphi,\, D\varphi\rangle\|u\|^2$ for all u. Taking $u = G_K\varphi$ gives $\|G_K\varphi\|^2 \le \kappa_1\langle \varphi,\, D\varphi\rangle$, i.e. $G_K^2 \le \kappa_1 D$.

Case $K = 1$. Here $D = t_1^2 G_1$, and Lemma I.11(c) with $a_1 = a$ gives $G_1 \le t_1^2 \max(1/8, 1/a)$. Hence $G_1^2 \le \|G_1\| G_1 \le \max(1/8, 1/a) D \le \kappa_1 D$, because $L^4/((L^2-1)^2 a_\flat) = L^6/((L^2-1)^3 a) \ge 1/a$. □

Lemma I.15 (block geometry)**.** *In the setting of Definition I.10, for $0 \le k \le K$:*

(a) *$\ell_j/t_k \in 2M_\flat\mathbb{N}$. In particular $\ell_j \ge 2t_k$, and $N := \ell_j/\varepsilon \ge 4$;*

(b) *$\mathcal{P}_k$ is a partition of Λ into cubes with L^k sites per side, and for $k < K$ each block of $\mathcal{P}_{k+1}$ is a union of L^3 blocks of $\mathcal{P}_k$. Hence $\mathcal{B}_{k+1} \subset \mathcal{B}_k$, $P_{k+1} \le P_k$ and $P_k P_{k+1} = P_{k+1} P_k = P_{k+1}$;*

(c) *for every block $\square \in \mathcal{P}_k$, a bond of $T_{j,\varepsilon}$ has both endpoints in $\square$ if and only if it joins two points of $\square$ whose coordinates differ by ε in exactly one direction, without wrap-around. Thus the bonds inside $\square$ are exactly the edges of the cube graph $\{0, \dots, L^k - 1\}^3$;*

(d) *$\langle \varphi, -\Delta_\varepsilon \varphi \rangle \ge \sum_{\square \in \mathcal{P}_k} \varepsilon \sum_{b \subset \square} (\varphi(b_+) - \varphi(b_-))^2$, where $b \subset \square$ means that both endpoints of the bond $b = (b_-, b_+)$ lie in $\square$.*

Proof. (a) We have $t_k = L^k \varepsilon_K$ and $k \le K \le K + n_0$, so Lemma 2.1(ii) gives $\ell_j/(M_b t_k) \in 2\mathbb{N}$. Hence $\ell_j/t_k \in 2M_b\mathbb{N}$ and $\ell_j \ge 2M_b t_k \ge 2t_k$. Taking $k = K$ gives $N = \ell_j/\varepsilon = L^K(\ell_j/t_K) \ge 2L \ge 4$.

(b) By (a), $n_k := \ell_j/t_k$ is an integer. In each direction the intervals $[y_i, y_i + t_k)$ with $y_i \in t_k\{0, \dots, n_k - 1\}$ partition $[0, \ell_j)$, and each contains $t_k/\varepsilon = L^k$ lattice points. Taking products gives the partition of Λ. For $y \in t_{k+1}\mathbb{Z}^3$ we have $[y_i, y_i + Lt_k) = \bigcup_{m=0}^{L-1} [y_i + mt_k, y_i + (m+1)t_k)$ with $y_i + mt_k \in t_k\mathbb{Z}$, which gives the nesting. A function that is constant on the coarser blocks is constant on the finer ones, and the statements about the projections follow.

(c) Let $\square = B^k(y)$. In direction i the points of $\square$ have coordinates in $[y_i, y_i + t_k) \subset [0, \ell_j)$, so two of its points differ in the ith coordinate by at most $t_k - \varepsilon$. A bond in direction i joins points x, x' that agree off coordinate i. Either $x_i' = x_i + \varepsilon$ (an interior bond), or $x_i = \ell_j - \varepsilon$ and $x_i' = 0$ (a wrap-around bond). Both endpoints of a wrap-around bond lie in $\square$ only if $\ell_j - \varepsilon \le t_k - \varepsilon$, that is $\ell_j \le t_k$, and (a) excludes this. Conversely, two points of $\square$ that differ by ε in one coordinate and agree in the others are the endpoints of an interior bond. Since $N \ge 4$, distinct bonds have distinct pairs of endpoints.

(d) The blocks of $\mathcal{P}_k$ are disjoint, so each bond lies inside at most one block. The left side is ε times the sum of $(\varphi(b_+) - \varphi(b_-))^2$ over all bonds. The right side is the same sum restricted to the bonds that lie inside some block. □

Theorem I.16 (domination of the slice-modified photon)**.** *Put $\bar\kappa := \kappa_1\big(2(1 + \mu_*^2\varepsilon_0^2)^2 + a^2\big)$. Then on all of $\ell^2(\Lambda)$, zero mode included,*

$$D^{\natural <} \le \kappa_1\kappa_3\, \mu_*^2\, D \le \bar\kappa\, \mu_*^2\, D.$$

Consequently $D^{\natural <} \le \kappa\mu_*^2 D$ for every $\kappa \ge \bar\kappa$, for every $K \ge 1$. The constant $\bar\kappa$ depends only on L, a and $\mu_*\varepsilon_0$. It does not depend on ε, K or ℓ_j. If $L = 2$, $a = 1$ and $0 < \mu_*\varepsilon_0 \le \frac{1}{5}$, then for every $K \ge 1$

$$\frac{164}{27} < \kappa_1\kappa_3 \le \bar\kappa \le \frac{42176}{5625} < \frac{15}{2}.$$

Proof. Put $\vartheta := \varepsilon_0^2 = t_K^2$ and $g^\natural(\lambda) := \mu_*^{-2} \int_0^\vartheta \mathrm{e}^{-s\lambda}(1 - \mathrm{e}^{-s\mu_*^2})\, ds$ for $\lambda \ge 0$, so that $D^{\natural <} = \mu_*^2 g^\natural(-\Delta_\varepsilon)$.

Step 1: $G_K^{-2} \le h(-\Delta_\varepsilon)$, with $h(\lambda) := 2(\lambda + \mu_*^2)^2 + 2r_K^2$. From $(M - r_K P_K)^2 \ge 0$ and $P_K^2 = P_K$ we get $r_K(MP_K + P_K M) \le M^2 + r_K^2 P_K$. Hence $Y^2 = (M + r_K P_K)^2 \le 2M^2 + 2r_K^2 P_K \le 2M^2 + 2r_K^2 = h(-\Delta_\varepsilon)$. This uses only the inequality $AB + BA \le A^2 + B^2$ for self-adjoint A, B; it never treats G_K as a function of $-\Delta_\varepsilon$. Since inversion is operator monotone, $G_K^2 = Y^{-2} \ge h(-\Delta_\varepsilon)^{-1}$.

Step 2: $g^\natural h \le \kappa_3$ on $[0, \infty)$. Since $1 - \mathrm{e}^{-y} \le y$, we have $g^\natural(\lambda) \le g_0(\lambda) := \int_0^\vartheta s\mathrm{e}^{-s\lambda}\, ds$. Moreover $g_0 \le \vartheta^2/2$, and $g_0(\lambda) \le \int_0^\infty s\mathrm{e}^{-s\lambda} ds = \lambda^{-2}$ for $\lambda > 0$. With $m' := \max(\lambda, 1/\vartheta)$ this gives $g_0(\lambda) m'^2 \le 1$. Also $\lambda + \mu_*^2 \le (1 + \mu_*^2\vartheta)m'$, since $\vartheta m' \ge 1$. As $r_K = a_K/\vartheta$,

$$g^\natural(\lambda)h(\lambda) \le 2(1 + \mu_*^2\vartheta)^2 + 2\frac{a_K^2}{\vartheta^2} \cdot \frac{\vartheta^2}{2} = 2(1 + \mu_*^2\varepsilon_0^2)^2 + a_K^2 = \kappa_3.$$

This includes $\lambda = 0$, the constants.

Step 3. $g^\natural(-\Delta_\varepsilon)$ and $h(-\Delta_\varepsilon)$ are commuting functions of $-\Delta_\varepsilon$, and $h > 0$. So Step 2 gives $g^\natural(-\Delta_\varepsilon) \le \kappa_3 h(-\Delta_\varepsilon)^{-1}$, and Step 1 gives $\kappa_3 h(-\Delta_\varepsilon)^{-1} \le \kappa_3 G_K^2$. Lemma I.14 then gives $D^{\natural <} = \mu_*^2 g^\natural(-\Delta_\varepsilon) \le \kappa_3\mu_*^2 G_K^2 \le \kappa_1\kappa_3\mu_*^2 D$. Lemma I.14 rests on the block Poincaré inequality, Lemma I.11.

That inequality uses the cube structure of the blocks, which is Lemma I.15(c),(d): no block contains both endpoints of a wrap-around bond, because $\ell_j \geq 2t_k$ for every $k \leq K$. By (F1), $a_K \leq a_1 = a$, so $\kappa_3 \leq 2(1+\mu_*^2\varepsilon_0^2)^2 + a^2$ and $\kappa_1\kappa_3 \leq \bar\kappa$. Finally $D \geq 0$ (Lemma I.12), so the bound with $\bar\kappa$ implies the bound with every $\kappa \geq \bar\kappa$.

Step 4: the values at $L = 2$, $a = 1$. Here $a_\flat = \frac{3}{4}$. The three numbers in the definition of κ_1 are $\frac{1}{8}$, $(\frac{1}{2}+\frac{4}{3})/3 = \frac{11}{18}$ and $16/(9\cdot\frac{3}{4}) = \frac{64}{27}$, so $\kappa_1 = \frac{64}{27}$. By (F1), $\frac{3}{4} < a_K \leq 1$. Put $x := \mu_*\varepsilon_0 \in (0, \frac{1}{5}]$. Then $\kappa_3 = 2(1+x^2)^2 + a_K^2$ increases in x and in a_K, so $\kappa_3 \leq 2(1+x^2)^2 + 1 \leq 2\cdot\frac{676}{625} + 1 = \frac{1977}{625}$. Hence $\kappa_1\kappa_3 \leq \bar\kappa \leq \frac{64}{27}\cdot\frac{1977}{625} = \frac{42176}{5625} = 7.4979\ldots < \frac{15}{2}$. On the other side, $\kappa_3 > 2 + a_\flat^2 = \frac{41}{16}$, so $\kappa_1\kappa_3 > \frac{64}{27}\cdot\frac{41}{16} = \frac{164}{27}$. The value $\frac{164}{27}$ is the limit of $\kappa_1\kappa_3$ as $\mu_*\varepsilon_0 \to 0$ and $K \to \infty$, and it is not attained. So this argument gives no constant $\kappa \leq \frac{164}{27}$. The constant of (5.1) must be a fixed number, independent of K. The fixed value $\kappa = \frac{15}{2}$ satisfies $\kappa \geq \bar\kappa$ under the condition (E.dom) $\mu_*\varepsilon_0 \leq 0.2$ (Remark I.18). □

Corollary I.17 (torus form)**.** *Let $\kappa \geq \kappa_1\kappa_3$, and let Γ^B and $\bar\delta$ be as in* (5.1)*. Then:*

(a) $D^< = D^{\natural<}\Pi^\perp \leq \Gamma^B\Pi^\perp$;

(b) $D \leq \varepsilon_0^2 G_K \leq \varepsilon_0^2 C_{\mu_*}$, *and hence* $\Gamma^B \leq \kappa\mu_*^2\varepsilon_0^2 G_K \leq \bar\delta\, C_{\mu_*}$.

Proof. (a) P_k is self-adjoint and fixes the constants, so $P_k\Pi_0 = \Pi_0 = \Pi_0 P_k$. Hence G_k, H_j, D and Γ^B commute with Π_0, and so does $\mathrm{e}^{t\Delta_\varepsilon}$. Compressing the inequality $D^{\natural<} \leq \kappa\mu_*^2 D = \Gamma^B$ of Theorem I.16 by $\Pi^\perp$ gives $\Pi^\perp D^{\natural<}\Pi^\perp \leq \Pi^\perp\Gamma^B\Pi^\perp$, and the two sides are $D^<$ and $\Gamma^B\Pi^\perp$. (b) $H_j \geq 0$ by Lemma I.12, since $H_j = (G_K - G_j) - (G_K - G_{j+1})$ and $G_{j+1}^{-1} \leq G_j^{-1}$ by the same computation. Also $t_{j+1} \leq t_K = \varepsilon_0$ for $j < K$ and $\sum_{j<K} H_j = G_K - G_0 = G_K$. Hence $D \leq \varepsilon_0^2 G_K$, and $G_K \leq C_{\mu_*}$ by (F4). Multiply by $\kappa\mu_*^2$. □

Remark I.18 (the value of κ). At $L = 2$ and $a = 1$ (stages 1–2 of Table 3) we have $a_\flat = 3/4$ and $\kappa_1 = \max\{1/8,\, 11/18,\, 64/27\} = 64/27$. Under the condition (E.dom), $\mu_*\varepsilon_0 \leq 0.2$, imposed at stage 10a, and with $a_K \leq a = 1$ by (F1),

$$\kappa_1\kappa_3 \leq \frac{64}{27}\Big(2\cdot\frac{676}{625} + 1\Big) = \frac{126528}{16875} < 7.498 < 7.5 \qquad \text{for every } K \geq 1.$$

As $\mu_*\varepsilon_0 \to 0$ and $K \to \infty$ (so $a_K \to a_\flat$), $\kappa_1\kappa_3 \to \frac{64}{27}\cdot\frac{41}{16} = \frac{164}{27} \approx 6.07$. This limit is not attained: $a_K > a_\flat$ by (F1), so $\kappa_3 > 2 + a_\flat^2 = \frac{41}{16}$ and $\kappa_1\kappa_3 > \frac{164}{27} > 6.07$ for every K and every $\mu_*\varepsilon_0$. Hence Theorem I.16 does not give the value $\kappa = 6.07$, and the condition (E.δ) is used with $\kappa = 7.5$ (Lemma A.19). By Theorem I.16 and Corollary I.17, any fixed $\kappa \in [126528/16875, \infty)$, in particular $\kappa = 7.5$, can serve as the K-independent domination constant in (5.1); it is fixed at stage 0, under the a priori bound (E.dom). Note that the inequality $\Gamma^B \leq \kappa\mu_*^2\varepsilon_0^2 G_K \leq \bar\delta C_{\mu_*}$ of Corollary I.17(b) holds for every $\kappa > 0$. The role of κ is the domination $D^{\natural<} \leq \kappa\mu_*^2 D$.

I.3. **The parameters and the hypotheses of Theorem 6.8.** This subsection contains four statements used in the proof of Theorem 3.1: the choice of the auxiliary parameters (η, m_1, m_2) in Theorem 6.8 (Lemma I.19); the list of the estimates of [2, 3] that are not written out in print and are used in the proofs of Theorems 4.2 and 5.2, with the places where they are proved (Lemma I.23); the transfer of these estimates to the masses at which Theorem 4.2 is used (Lemma I.24); and the verification of the hypotheses of Theorem 6.8 on every torus of the family $\mathfrak{C}$ (Proposition I.26).

Throughout, $\mathbb{M}_4 = \{M_1, m, m_1, m_2\}$, the stages 0–11 are those of Table 3, and, as in Section 6, $\mathfrak{C}$ is the family of pairs (ε, T_j) with $\varepsilon = 2^{-K}\varepsilon_0$, $K \geq K_0 = \max(K_0', n_0 + 1)$, $j \geq 1$. A Higgs-mass entry of a constant or of a condition is *of type* (α) if the mass M enters only through the rescaled mass M^2t^2, $t \leq \varepsilon_0$, and the constant is uniform for $0 \leq M^2t^2 \leq 1$. It is *of type* (β) if M enters through positive powers or logarithms of M, or through quantities finite at every fixed $M > 0$, and only inside downward-closed upper bounds on ε_0, monotone constants of the stages 8–9, or the floor on ℓ_0. It is *of type* (γ) if it orders M against μ_* or uses $m_1 > 2m$.

The auxiliary parameters.

Lemma I.19 (the choice of (η, m_1, m_2))**.** *Let $m > 0$, $0 < \lambda < 0.1m$ and $0 < e^2 < 0.1m$, as in Theorem 3.1. Put $\pi(e,\lambda,m) := (e,\lambda,m,\frac{1}{2},3m,4m)$. Then:*

(a) *$\pi(e,\lambda,m) \in \mathfrak{W}$, and at this point Theorem 6.1(a)–(d) give $2\sigma_+ \geq 0.319m^2$, $g_{21} \geq 6.6369\,m^2$, $a_{K,T}(3m) \geq a_{K,T}(m) + 2H$ and $3m \geq \sqrt{2}\mu_*$, uniformly in $\varepsilon \leq 1/m$ and in tori of side $\geq 96/m$;*

(b) *Lemma A.5 at $\pi(e,\lambda,m)$ provides R, b_0, n_0 and $\ell_0 \in 2M_b\mathbb{N}$ with $\ell_0 \geq 183/m$ that meet every condition of Appendix A, and every $(\varepsilon, T_j) \in \mathfrak{C}$ has $\varepsilon \leq 1/(4m)$ and $\ell_j \geq 96/m$;*

(c) *with these data, the constants of Theorem 3.1(i) are $\sigma_+ := g/2$, with g as in Proposition 6.4, and $C := \mathsf{K} = \max(\kappa_+, 2S_\flat^B)$, with κ_+ from Proposition 6.4 and $S_\flat^B$ from Proposition 6.7 at $(\eta, m_1, m_2) = (\frac{1}{2}, 3m, 4m)$, with Z_T read as $\bar{Z}$ (Proposition I.9, Corollary B.28); the threshold is $K_0 := \max(K_0', n_0 + 1)$.*

Any other $\eta \in (0,1)$ and $2m < m_1 < m_2$ may be used instead; the constants depend on the choice.

Proof. (a) $0 < \frac{1}{2} < 1$ and $2m < 3m < 4m$, and the conditions on (e,λ,m) are those of $\mathfrak{W}$ in (1.3), so $\pi(e,\lambda,m) \in \mathfrak{W}$. Theorem 6.1(b) gives $g_{21} \geq (4m - 3m)(4m + 3m - 0.3631m) = 6.6369\,m^2$. Theorem 6.1(c) holds because $3m > 2m > 1.882m \geq m_1^c$, and $3m \geq \sqrt{2}\,m = \sqrt{2}\mu_*$. (b) This is the conclusion of Lemma A.5 at $\pi(e,\lambda,m)$. The condition (E.m2), $m_2^2\varepsilon_0^2 \leq 1$, gives $\varepsilon \leq \varepsilon_0 \leq 1/m_2 = 1/(4m)$. The torus floor (rule (iii) of Subsection A.1) is $\ell_0^\natural = \max\{8, 128/(3m), 128/(0.7m)\} = 182.857\ldots/m$, so the choice $\ell_0 \geq 183/m$ meets it. (c) Theorem 6.8 gives the bound of Theorem 3.1(i) with the tube $\mathcal{D}_{\sigma_+,H}$, $\sigma_+ = g/2$, and the constant K. The spacing threshold is that of Theorem 6.8. □

The estimates of [2, 3] at the points of Theorem 5.2.
The estimates of the proof of Theorem 5.2 at its two points. Throughout this paragraph the hypotheses of Theorem 5.2 are in force. We take $\varepsilon = 2^{-K}\varepsilon_0$ with $K \geq K_0 = \max(K_0', n_0 + 1)$, $T = T_j$ with $j \geq 1$, and $t_k := L^k\varepsilon$, so that $t_K = \varepsilon_0$. For $X \subset T$ we write $|X|_k$ for the number of points of X in the lattice $t_k\mathbb{Z}^3$. For a set Λ of scale $k-1$ we write $\Lambda' := \Lambda \cap t_k\mathbb{Z}^3$ [1, (1.16), p. 606]. Inside this paragraph, i denotes a scale index.

Definition I.20 (the omitted estimates)**.** Let $\mathcal{I}$ be the following collection of statements. Each one is followed by the place where it is proved.

(a) The ingredients of Table 1:
- the moved vacuum pieces at every scale (Proposition E.64);
- the Ward–Takahashi step on the torus (Proposition E.67);
- the conditions of [3, Prop. 1] as used in [2] (Proposition E.110);
- the restriction to real gauge backgrounds near Neumann faces (Proposition E.71);
- [3, Prop. 1] at the order cap $\bar{N}$ (Proposition E.76);
- the region entropy, that is, the estimate [2, (3.42)–(3.54), pp. 592–594] on the torus (Proposition 5.21);
- the mass reading of [2, p. 589] (Proposition E.159);
- the torus-uniform final c-numbers (Proposition E.162);
- the vacuum classes at the last scale (Proposition E.175).

(b) The statements that make up Proposition E.110, which are the entries of Table 10:
- the one-variable remainder-kernel bound (entry 1, Lemma E.116);
- the two-variable bound on T_η (entry 2, Lemma E.117) and on Neumann domains Ω (entry 3, Lemma E.118);
- the absence of mass insertions other than δm^2 (entry 4, Lemma E.119);
- the premises of [3, Prop. 1] as applied in [2] (entries 5, 5′, Lemma E.114);
- the separation statement for unions of big blocks (entry 6, Lemma E.120);

- the cube remainder of [3, p. 433] (entry 7, Lemma E.121);
- the two-domain bounds for δG_k (entry 8, Lemma E.123);
- the inclusions of the cube $\square$ in the outer domains and the retirement of $\square \subset \Omega_2$ (entries 9, 10, Lemmas E.124, E.125);
- the end-point statements for Ω_2-lines in [2, Chaps. 2, 3] (entries 11, 12, Lemmas E.126, E.127);
- the nesting $\Omega \supset \Omega_1 \supset \Omega_2$ (entry 13, Lemma E.100 and Remark E.101);
- the support hypothesis in the form (5.10) (entry 14, Corollary 5.10);
- the conventions R1–R3 on gating, nesting and metric (entry 15, Lemma E.128);
- the completeness of the lists of uses of [3, Prop. 1] (entry 15′, Lemma E.129);
- the local $A \cdot A$ terms (entry 16, Lemmas E.24, E.32 and E.130);
- the cuts beyond the order $\bar{n}$ of [2, p. 579] (entry 17, Proposition E.36 and Lemma E.141).

(c) The statements on domains, supports and localisation widths: Corollary 5.10, which gives the form (5.10) of [3, p. 412], and Lemmas E.100, E.102 and E.103.
(d) The order cap $\bar{N} = \bar{n}^2 = 169$ for [3, (1.33)] (Proposition E.76).
(e) The final step of the upper bound, [2, (2.116)–(2.118), p. 582] (Lemma I.21 below).
(f) [2, Lemma 2.4] at edge points with the cap (5.7), that is, condition (E.edge) (Lemma 5.6); the periodisation lemma with condition (T.per) (Lemma 5.22); and the depth of each sketched step of [3, Prop. 1] (Proposition 5.8 and Lemma 5.9).

Lemma I.21 (the final step of the upper bound)**.** *Let $\rho^{(K)}$ be the density of* [2, (2.43), p. 566] *at $k = K$, so that, in the notation of* [2, pp. 566–582]*,*

$$Z^{\varepsilon} \le \int dB \int d\psi \sum_{\Lambda_0^{(0)},\dots,\Lambda_0^{(K-1)}} \rho^{(K)} \exp\Big(\mathcal{P}^{(K)}\big(\Lambda_7^{(K-1)\prime}, \theta_K B^{(K),\varepsilon}, \psi\big) - E_0 + \sum_{i=0}^{K-1} O(t_i^{\kappa_0}) |\Lambda_7^{(i-1)\prime} \cap \Lambda_7^{(i)c}| \Big) \exp\Big(\sum_{i<K} O(t_i^{\kappa}) |T_1^{(i)}| \Big), \tag{I.2}$$

with $\Lambda_7^{(-1)\prime} := T$. Here $\kappa > 0$ is the exponent of [2, Prop. 2.2, (2.67)]*; it is unrelated to the slice constant of* (5.1)*. Let $\kappa_0 > 0$ and $C \ge 0$, and assume that on the support of $\rho^{(K)}$*

$$(F) \qquad \mathcal{P}^{(K)}\big(\Lambda_7^{(K-1)\prime}, \theta_K B^{(K),\varepsilon}, \psi\big) \le C\, t_K^{\kappa_0}\, |\Lambda_7^{(K-1)\prime}|_K. \tag{I.3}$$

Then:

(a) $\mathcal{P}^{(K)}(\Lambda_7^{(K-1)\prime}, \theta_K B^{(K),\varepsilon}, \psi) \le C\varepsilon_0^{\kappa_0 - 3}|T|$ *on the support of $\rho^{(K)}$;*
(b) *with c the constant of* [2, (2.118), p. 582]*,*

$$Z^{\varepsilon} \le \mathrm{e}^{(C\varepsilon_0^{\kappa_0-3}+c)|T|} \int dB \int d\psi \sum_{\Lambda_0^{(0)},\dots,\Lambda_0^{(K-1)}} \rho^{(K)} \mathrm{e}^{-E_0} \exp\Big(\sum_{i=0}^{K-1} O(t_i^{\kappa_0}) |\Lambda_7^{(i-1)\prime} \cap \Lambda_7^{(i)c}| \Big).$$

Moreover, if the sizes in [2, (2.117)] *are read as $|\cdot|_{K-1}$, then the right side of* (I.2) *is at most $\mathrm{e}^{c|T|}$ times the left side of* [2, (2.117)]*, with last constant $L^{\kappa_0} C$;*

(c) *(F) follows from the printed bound* [2, (2.116), p. 582]*, $\mathcal{P}^{(K)} \le O(t_K^{\kappa_0})|\Lambda_7^{(K)}|$, under each of the following readings of its right side: $\Lambda_7^{(K)} = \emptyset$; or $\Lambda_7^{(K)}$ is any set obtained by one further step of the construction of* [2, p. 566]*. In both readings $|\cdot|$ may be either the count $|\cdot|_K$ or the volume $t_K^3|\cdot|_K$;*

(d) *(a)–(c) hold unchanged with $\rho^{(K)}$ replaced by any non-negative density. In particular they hold for the density of the slice model at (u, M), $u \in [-\eta, 1]$, whose photon weights $1 + u\delta_i$ are positive.*

The constants added by (a)–(c) are 1 *and* L^{κ_0}*. They do not depend on* ε*,* K*,* T*,* u *or* M*.*

Proof. (a) We have $\Lambda_7^{(K-1)\prime} \subset T$, so $|\Lambda_7^{(K-1)\prime}|_K \le |T|_K = |T|\, t_K^{-3} = |T|\varepsilon_0^{-3}$.

(b) In (I.2) the factor $\rho^{(K)}$ is the density of a positive measure: it is a product of positive kernels and characteristic functions. The two exponential factors are positive, and $x \mapsto \mathrm{e}^x$ is increasing. So replacing $\mathcal{P}^{(K)}$ by the right side of (F), which bounds it on the support of $\rho^{(K)}$, does not decrease the right side of (I.2). By (a) this replacement contributes at most $\mathrm{e}^{C\varepsilon_0^{\kappa_0-3}|T|}$. By [2, (2.118)], $\sum_{i<K} O(t_i^\kappa)|T_1^{(i)}| = \sum_{i<K} O(1)t_i^{\kappa_0}|T| \le c|T|$. This gives the first display.

For the second statement, note that $t_K^{\kappa_0} = L^{\kappa_0} t_{K-1}^{\kappa_0}$. As point sets, $\Lambda_7^{(K-1)\prime} \subset \Lambda_7^{(K-1)}$, and $t_K\mathbb{Z}^3 \subset t_{K-1}\mathbb{Z}^3$ because L is an integer. Hence $|\Lambda_7^{(K-1)\prime}|_K \le |\Lambda_7^{(K-1)}|_{K-1}$, and so $Ct_K^{\kappa_0}|\Lambda_7^{(K-1)\prime}|_K \le L^{\kappa_0} C\, t_{K-1}^{\kappa_0}|\Lambda_7^{(K-1)}|_{K-1}$. This is the last term of the exponent in [2, (2.117)]. The other terms of that exponent are the collar terms $i = 0, \dots, K-1$ of (I.2).

(c) If $\Lambda_7^{(K)} = \emptyset$, the printed bound reads $\mathcal{P}^{(K)} \le 0$, which implies (F). Now let $\Lambda_7^{(K)}$ be produced by one further step. By [2, p. 566] the sets of each scale are nested, so $\Lambda_7^{(K)} \subset \Lambda_0^{(K)}$, and an admissible set satisfies $\Lambda_0^{(K)} \subset \Lambda_7^{(K-1)\prime}$. Hence $|\Lambda_7^{(K)}|_K \le |\Lambda_7^{(K-1)\prime}|_K$, and (F) holds with the printed constant. In the volume convention, $t_K^3|X|_K \le |X|_K$ because $t_K = \varepsilon_0 \le 1$. So the volume form of the printed bound implies its count form, and hence (F).

(d) The proof of (b) used only $\rho^{(K)} \ge 0$, and (a), (c) do not involve the density. □

Lemma I.21 does not prove (F). On the torus, (F) is the final perturbative bound of the closing step, part (S5) of Proposition F.2, and it is obtained there from [3, Prop. 1].

Remark I.22 (King's cube terms and field restrictions). (i) On the periodic torus T_ε the set $\partial T^{(k)}$ is empty, and we use the convention $\mathrm{dist}(x, \emptyset) = +\infty$. Consider the terms that [25, Sect. 5.2, pp. 343–344, (5.41)–(5.49)] adds for Dirichlet scalar boundary conditions on a cube. The product (5.41) runs over $x \in \Lambda_0^{(k)} \cap \partial T^{(k)}$, so it is empty and equals 1. The hypothesis $\mathrm{dist}(x, \partial T^{(k)}) < p(L^k\varepsilon)$ of (5.44)–(5.45) holds for no x. (5.42)–(5.43) are [2, (2.110)] and its consequence. The hypothesis $\mathrm{dist}(x, \partial T^{(k)}) \ge R$ of (5.48) holds at every x, and the operator $G_k(A^{(k)})$ there is the periodic one. So (5.46)–(5.49) reduce to the quartic term of [2, p. 570] combined with [2, Lemma 2.4, p. 572] (Lemma 5.6). Therefore on T_ε the cube terms add nothing to the estimates of [2, Chap. 2].

(ii) The small-field restrictions in the upper bound (5.2) are those of [2, (2.55), p. 570]. King's restrictions [24, (3.2), p. 655] and his modified lower bound [24, (A.1), p. 676] enter only the lower bound, Theorem 4.1. The condition $\mu_0 t \le 1$ of [24, (A.4)–(A.5), p. 676] is implied by (E.dom). The bound (5.2) contains no characteristic function of the fields, so none of its thresholds has to match those of the lower bound.

Lemma I.23 (the omitted estimates of [2, 3])**.** *Assume the hypotheses of Theorem 5.2 and Statements K.62 and K.63. Then every statement of* $\mathcal{I}$ *(Definition I.20) holds at* $(u, M) \in \{(1, m_2), (-\eta, m_1)\}$*, for every* $K \ge K_0$ *and every* $j \ge 1$*. Its constants are independent of* ε *and* K*. They depend on the torus* T_j *at most through non-increasing functions of* ℓ_{T_j}*, so they are bounded by their values at the torus floor* $\ell_0^\natural$ *of (T.floor). They are finite at* $u = -\eta$ *and at* $M = m_1$*.*

Proof. Step 1: item (a).

- The statement of Proposition E.64 includes the uniformity in ε, K, ℓ_T and the finiteness at $u = -\eta$, $M = m_1$.
- Proposition E.67 holds under condition (T.nw) (Lemma E.66) and introduces no constant.
- Proposition E.110 is treated in Step 2.
- Proposition E.71 is a statement on the backgrounds at which bounds are used, and it introduces no constant.

- The constants $O(\alpha_0, \bar{N}; d, L, a, M_b)$ and $\delta(\bar{N}) > 0$ of Proposition E.76 are independent of ε, k, the domains and the background. They are uniform over $\ell_T \geq \ell_0$, with values taken at the floor where δm_{fin}^2 vertices occur. Its colouring factor is at most $(1+L^2)^{N_v} \leq 5^{84}$ for $u \in [-\eta, 1]$.
- Proposition 5.21 is uniform in K, k and the torus.
- The constants of Proposition E.159 depend on none of ε, ε_0, k, ℓ_T, u, M, and the proposition holds at every $M \in (0, m_2]$.
- Proposition E.162 holds under (E.fin), $\varepsilon_0 \leq \min(1/m_2, 1/\mu_*, L)$, which the hypotheses imply (Lemma 5.4(b)). It is uniform in ε, x and j. The torus enters only through non-increasing functions of ℓ_T, which are bounded by their values at $\ell_0^\natural$.
- The constant $O(\alpha_0, \bar{n})$ of Proposition E.175 is independent of ε, k and the domains, and its torus term $C|T|\varepsilon_0^{-3}\mathrm{e}^{-\delta\ell_0/(4\varepsilon_0)}$ is non-increasing in ℓ_T. In its proof, Lemma E.170 is applied with $W = B^k(\Lambda_7^{(k)})$, which is Statement K.63.

Step 2: item (b). The entries of Table 10 are proved in the lemmas listed in Definition I.20(b), in Subsection E.7. Proposition E.110 composes them. At the first step $k = 1$, the two-domain bounds of Lemma E.123 are used with the margins of Statement K.62. Proposition E.36 and Lemma E.32 are used under the hypotheses stated there. The constants these statements introduce are $\ell_{\bar{N}} = 2V(\bar{N})$, $C_{\text{loc}}(\bar{N})$, $C_{\bar{N}}$, the constant of the two-domain bound at margin $R_0 + 12$, $\beta_\square$, C_{old} and $C_>$ (Definition A.1). Each of them, except $C_>$, is a function of $d, a, L, M_b, R_0, \delta_0, \bar{N}$ only. The constant $C_>$ is a supremum over $t \in (0, 1]$, evaluated at $\ell_T = \ell_0^\natural$. None of them depends on ε, K, u or M.

Step 3: items (c)–(f). The form (5.10) holds at every use under (5.11) (Corollary 5.10). Lemmas E.100, E.102 and E.103 are statements about domains, supports and localisation widths, and they contain no constant that depends on ε, K, T, u or M. The cap $\bar{N} = 169$ is part of Proposition E.76. The final step is Lemma I.21, whose added constants are 1 and L^{κ_0}. It is applied to the non-negative density of the slice model (Lemma I.21(d)), with (F) supplied by part (S5) of Proposition F.2. Lemma 5.6 holds under its cap (E.edge). Lemma 5.22 has constants $C_1(d, a, M_b, L)$ and $\delta_0(d, a, M_b, L)$, and under (T.per) its image terms are at most $C_1\mathrm{e}^{-3\delta_0 r(t_k)}$. Proposition 5.8 and Lemma 5.9 concern depths and the dependence on Ω. They introduce no constant beyond those of Proposition 4.5.

Step 4: the two points. For $u \in [-\eta, 1]$ the photon weights satisfy $1 + u\delta_i \in [1 - \eta\bar{\delta}, 1 + \bar{\delta}]$, since $0 \leq \delta_i \leq \bar{\delta}$ by (5.1). By (E.δ) (Lemma 5.4(a)), $\bar{\delta} \leq \frac{1}{12}$. Hence $1 + u\delta_i \in [1 - \eta/12,\ 13/12] \subset [1 - \eta, 2]$. Every weighted photon propagator is therefore at most $\frac{13}{12}$ times the unweighted one, and $C_B(u) \geq (1 - \eta/12)C_{\mu_*} > 0$. So no constant of Steps 1–3 blows up at $u = -\eta$.

Consider now the Higgs mass. Wherever it enters from above, it does so through $M^2t^2 \leq m_2^2\varepsilon_0^2 \leq 1$, by (E.m2). Wherever it enters from below, it does so through quantities that are finite at every fixed $M > 0$: $\gamma_1 M^2 t^2$, $\bar{\ell}(M) = \min(1/M, 1/\mu_*, L)$, $\log(1/(\min(M, \mu_*)\varepsilon_0))$ and the zero-mode gap $M^2(1 + 4M^2\varepsilon_0^2/(3a))^{-1}$. These are finite at $M = m_1 > 0$.

What the cited results use. The results cited in Steps 1–3 use, in their proofs, the following results of this paper: Lemmas F.31 and F.32 (in Proposition E.64); Proposition E.97 (in Propositions E.64 and E.76); Lemmas E.130 and E.141, beyond Proposition E.36 and Lemmas E.24, E.32; the flow factor $\sigma_N(u)$ (in Proposition E.162); and Lemma E.166 (in Proposition E.175). They use no other estimate of [2, 3] that is not written out in print. Combining Steps 1–4 proves the lemma. □

Transfer to the masses of Theorem 4.2.

Lemma I.24 (transfer of the estimates to $u = 0$ and to the masses M_1, m_1, m_2)**.** *Let $\pi \in \mathfrak{W}$, and let the regulator data be chosen as in Lemma A.5, with the four-mass rule and the torus floor of Subsection A.1. Assume Statements K.62 and K.63. Let $M \in \{M_1, m_1, m_2\}$. Then every estimate of $\mathcal{I}$ (Definition I.20) holds at $(u, M) = (0, M)$, in Bałaban's model with photon mass $\mu_* = m$.*

Consequently, Theorem 4.2 holds at $M_1 = 0.7m$, m_1 and m_2 on every torus of $\mathfrak{C}$, with its printed proof completed by these estimates.

The proof transfers the dependence on u and on the Higgs mass, as it can be read off the proof of each estimate; no estimate is proved again.

Proof. (i) The point $u = 0$. Since $0 \in [-\eta, 1]$, the point $u = 0$ lies in the interval on which the estimates are proved. The weights $1 + u\delta_i$ act on the photon fluctuations only, and the scalar side carries no u-weight. At $u = 0$ every weight equals 1. Every factor that contains $|u|$ in the constants of these estimates, including $C_>$ through $(1 + |u|c_k)^{N_v}$, is at $u = 0$ at most its value at $u = 1$ and at $u = -\eta$. The model at $u = 0$ is Bałaban's own model, so the estimates are statements on his objects.

(ii) The Higgs mass, estimate by estimate. The proof of each estimate shows how the Higgs mass enters it:

- *Type (α):* the remainder bounds of Lemmas E.116, E.117 and E.118, where m^2 enters the form $\|Df\|^2 + m^2t^2\|f\|^2 + a\|Qf\|^2$ with a + sign; Lemma E.120; the two-domain bounds of Lemma E.123 (both routes), with the margins of Statement K.62; Lemmas E.124 and E.125; the statements of item (c) and Lemmas E.128, E.129 (either no mass or type (α)); the $A \cdot A$ local terms; the region entropy of Proposition 5.21; and Proposition E.159 (the condition (E.md)).
- *Type (β):* Proposition E.36 (through the δm^2 vertices); Proposition E.71 (only through $-\Delta_A + m^2 > 0$); Proposition E.162 (through $\bar{\ell}(M)$, $\rho_{\min}(M)$, Z_f and $\log(1/(\min(M, \mu_*)\varepsilon_0))$); the steps of Subsection F.2 (through the condition (E.5) and Lemma A.14); the closing step (Lemmas F.39 and A.12, with the gap $\geq M^2(1 + 4M^2\varepsilon_0^2/(3a))^{-1}$); the constant $C_\delta(M)$, which is finite at every $M > 0$; C'_b; and C_{F1} and $\mathfrak{E}_k$ of Proposition F.2.
- *No mass entry:* the cube remainder of [3, p. 433]; the order cap $\bar{N} = 169$; the Ward–Takahashi step on the torus, Proposition E.67 (King's counterterm at M_1 on M_1-lines; Lemma C.1 holds at every $M > 0$); and the final step of the upper bound, Lemma I.21. That lemma's added constants are 1 and L^{κ_0}, and it holds for every non-negative density (Lemma I.21(d)), in particular for that of Bałaban's model at $u = 0$ and Higgs mass M.

Entries of type (α) hold at $M \leq m_2$, because (E.m2) and $t \leq \varepsilon_0$ give $M^2t^2 \leq m_2^2\varepsilon_0^2 \leq 1$. The left-hand sides of the upper bounds (E.m2) and (E.Φ) increase in M, and their constants do not depend on M. So imposing them at m_2 covers every $M \leq m_2$, in particular M_1. Entries of type (β) are finite at each fixed $M > 0$, and they enter only conditions that the data of Lemma A.5 meet at every $M \in \mathbb{M}_4$. Consider an upper bound on ε_0 in which the mass enters from below: (E.5), the ε_0-window of the closing step, Lemma A.12, or $\log_+(1/(M_1\varepsilon_0))$. Each is a supremum, over $t \leq \varepsilon_0$, of a quantity that tends to 0 as $\varepsilon_0 \downarrow 0$ at fixed $M_1 > 0$. Each is therefore a downward-closed upper bound, and the minimum of four positive thresholds is positive. These bounds bind first at the smallest mass M_1.

The constants of the stages 8–9 that carry M are $c_7 \ni C_\delta(M)^2/(16\lambda)$, c_1^I and the lower bounds on b_0. They are evaluated at the four masses and combined by the maximum; this is the four-mass rule. Their boundedness for $\varepsilon_0 \in (0, \varepsilon_1]$ is Statement K.1. The constants that depend on the torus are taken at $\ell_0^\natural = \max\{8, 128/m_1, 128/M_1\} = 128/M_1$; this is the torus floor. They are non-increasing in ℓ_T, so their values at this floor dominate their values on every torus of $\mathfrak{C}$.

(iii) No entry of type (γ). No estimate of $\mathcal{I}$ orders M against μ_* or uses $m_1 > 2m$. Such orderings occur only in Theorem 6.1 and in the anchor condition (6.3). Those are used at m_1, m_2 and are not estimates of $\mathcal{I}$. The order of the two field masses flips in only one place: Proposition E.162 at $M_1 < \mu_*$, where the scalar becomes the light field. The construction there is symmetric. Both fields are sliced to one common index $\bar{K}$ with $t_{\bar{K}} \leq \bar{\ell}(M) = \min(1/M, 1/\mu_*, L)$, and the mass ratio enters only through $\rho_{\min}(M)$. At $(M_1, \mu_*) = (0.7, 1)m$ and $L = 2$, in units $m = 1$, we get $\bar{\ell}(M_1) = 1$ and

$\rho_{\min}(M_1) \geq 0.7 \cdot \min(1, \frac{1}{2}) = 0.35 > 0$. The hypothesis $\varepsilon_0 \leq \bar{\ell}(M_1)$ follows from (E.fin). Theorem (1.3) of [2, p. 556] asks for no relation between the Higgs mass and μ_0. It is stated for the model of [2, p. 555], at the masses m^2, μ_0^2 and coupling λ there, with constants $E_\pm$ independent of ε and T_ε.

(iv) Domains, supports and structure. The statements of item (c), the end-point statements of Lemmas E.126 and E.127, and Lemma E.119 are statements about domains, supports, localisation widths and the insertion structure of the vertices. They contain no constant, and the value of M does not occur in them. The premises of Lemma E.114 are discharged by the other entries of Table 10, so they transfer with those entries.

(v) Statement ranges. Four ingredients are stated at the masses of Theorem 5.2: Proposition E.162 (at $M \in \{m_1, m_2\}$), the $A \cdot A$ local terms ($M \in [m_1, m_2]$), Proposition E.36 ($M \in \{m_1, m_2\}$) and Lemma A.14 (in the regime of m_1). By their proofs they hold at every $M \in (0, m_2]$; this is Lemma A.6(a)–(d). No proof step is redone.

(vi) The estimates of item (f). For them the transfer can be read off their statements. The cap (5.7), condition (E.edge), does not involve the Higgs mass: the mass enters only through a_k. So it is compatible with the minimum over $\mathbb{M}_4$, and Lemma 5.6 holds at $(0, M)$ for each $M \in \{M_1, m_1, m_2\}$ under the same cap. Lemma 5.22 has constants $C_1(d, a, M_b, L)$ and $\delta_0(d, a, M_b, L)$, which depend on none of μ_*, e, λ, m^2 or u. The torus enters only through its side, and under (T.per) the image terms are at most $C_1 \mathrm{e}^{-3\delta_0 r(t_k)}$ at every (u, M). Proposition 5.8 and Lemma 5.9 concern depths and the dependence on Ω. They introduce no constant beyond those of Proposition 4.5, which is stated for Bałaban's model, that is, at $u = 0$. In the written steps of [3, Prop. 1] the only mass that occurs is the term $+m^2$ of the operator $-\Delta^{\eta,N} + m^2 + a_k P_k$, which is of type (α). Hence the three estimates of item (f) hold at $(0, M)$ for every $M \in \{M_1, m_1, m_2\}$.

Conclusion. By Lemma I.23, the estimates of $\mathcal{I}$ hold at $(1, m_2)$ and $(-\eta, m_1)$. By (i)–(vi) they hold at $(0, M)$ for $M \in \{M_1, m_1, m_2\}$, under the four-mass rule and the torus floor. Both rules only shrink ε_0 or raise ℓ_0, so Lemma A.5 can meet them. These estimates are those of Remark 4.4, so Theorem 4.2 holds at these masses. □

Remark I.25. Proposition 6.2 states the upper bound only at $M \in \{M_1, m_1, m_2\}$, and no upper bound at $M = m$ is used. The chain uses only $E_+(M_1)$, in Proposition 6.4. Only $(0, M_1)$ lies on the path to Theorem 6.8; the masses m_1, m_2 are anchors off that path.

The hypotheses of Theorem 6.8.

Proposition I.26 (the hypotheses of Theorem 6.8 on $\mathfrak{C}$)**.** *Let (e, λ, m) be as in Theorem 3.1, take $\pi = \pi(e, \lambda, m)$ and the data of Lemma I.19, and let $(\varepsilon, T_j) \in \mathfrak{C}$. Then every hypothesis of Theorem 6.8 holds:*

1. *(the hypotheses of Theorem 5.2) the conditions of Appendix A, including* (5.7) *and* (5.13)*, hold, and $\eta\bar{\delta} < 1/12$; hence* (5.2) *holds at $(1, m_2)$ and $(-\eta, m_1)$ on every torus of $\mathfrak{C}$;*
2. *(the slice vacuum bound) $\Delta_1^B < \infty$, the supremum being taken over $\mathfrak{C}$, with $\Delta_1^B \leq C_+^B(1; m_2) + \eta^{-1} C_+^B(-\eta; m_1) + (1 + \eta^{-1}) E_-(m_1) + \frac{1}{2}(1 + \eta) E_2(m_1) + \bar{\Gamma}_{21}^{(1)}$;*
3. *(the anchor condition* (6.3)*) $b_T + 2H \leq a_{K,T}(m_1)$ and $g_{21} > 0$ on every torus of $\mathfrak{C}$; hence Proposition 6.7 applies;*
4. *(the "+" side) the hypotheses of Proposition 6.4 hold, with $g \geq 0.319 m^2$; hence Propositions 6.4 and 6.5 hold for every j and every K of $\mathfrak{C}$;*
5. *(the curvature bound) Lemma 7.3 holds on every torus of $\mathfrak{C}$ at the coefficient b_{T_j}.*

Consequently Theorem 6.8 holds on $\mathfrak{C}$ with $\sigma_+ = g/2$, $H = m^2$ and $\mathsf{K} = \max(\kappa_+, 2S_\flat^B)$.

Proof. (1) Lemma I.19(b) gives the conditions of Appendix A. Theorem 5.2 also uses $\eta\bar{\delta} < 1$. It follows from the condition (E.δ), $\kappa\mu_*^2\varepsilon_0^2 \leq 1/12$, imposed at stage 10b, that is $\bar{\delta} \leq 1/12$; hence $\eta\bar{\delta} < 1/12$ for $\eta < 1$. (Alternatively, (E.dom), $\mu_*\varepsilon_0 \leq 0.2$, together with $\kappa \leq 7.5$ gives $\bar{\delta} \leq 0.3$.)

With $\Gamma^B \le \bar\delta\, C_{\mu_*}$ from (5.1) this gives $C_B(-\eta) = C_{\mu_*} - \eta\Gamma^B \ge (1-\eta\bar\delta)C_{\mu_*} > 0$, so the slice model at $u = -\eta$ is well defined and Φ_{m_1} is convex on $[-\eta, 1]$. The spacings of $\mathfrak{C}$ satisfy $K \ge K_0 = \max(K_0', n_0+1)$, as Theorem 5.2 requires.

(2) Apply Proposition 6.6. Its hypotheses hold:

- Theorem 5.2 at $(1, m_2)$ and $(-\eta, m_1)$: by (1).
- Theorem 4.1 at m_1, in the form $|T|^{-1}\Phi_{m_1}(0) + e(\varepsilon; 0; m_1) \ge -E_-(m_1)$: this is the lower bound of Proposition 6.2 at m_1, by Lemma H.1(b), read with the order-zero term $E_{0,s}(m_1)$ of Remark 5.3(i). It needs none of the estimates of Lemma I.23.
- The bound (6.1) at m_1, m_2 (Lemma H.9): it is a supremum over the whole interval $u \in [-\eta, 1]$, not only over the two points. What it inherits from Propositions E.64–E.76 holds at m_1, m_2 uniformly over the weight class $\mathbb{W}_0 = [1-\eta, 2]$ of the slice weights $w_j = 1 + u\delta_j$, i.e. for all $u \in [-\eta, 1]$.
- The bound (6.2) at (m_1, m_2): Proposition H.11, given Proposition 6.3 and [3].

Both instances of (5.2) and both c-number bounds are used at the masses m_1, m_2 of Theorem 5.2, so no transfer to other masses is needed on this side. Proposition 6.6 then gives the bound in (2). The conversion between King's Lebesgue-normalised photon in Z_K and the normalised photon in $\Phi_M(0)$ is Lemma H.1(b).

(3) Every $(\varepsilon, T_j) \in \mathfrak{C}$ has $\varepsilon \le 1/m$ and $\ell_j \ge 96/m$, by Lemma I.19(b), so Theorem 6.1(b),(c) apply. Theorem 6.1(c) reads $a_{K,T}(m_1) \ge a_{K,T}(m) + 2H = b_T + 2H$. Theorem 6.1(b) gives $g_{21} \ge 6.6369\, m^2 > 0$. With (2), these are the hypotheses of Proposition 6.7.

(4) Take $M_2 = m$ and the anchor $M_1 = 0.7m$. Then:

- $g \ge 2\sigma_+ \ge 0.319m^2$ by Theorem 6.1(a).
- The upper bound of Theorem 4.2 at M_1 holds on every T_j by Lemma I.24.
- The lower bound of Theorem 4.1 at m holds by that theorem.
- $\bar\Gamma(M_1, m) < \infty$ by Proposition 6.3.
- The coefficient $b_T = a_{K,T}(m)$ is the one at which Bałaban's theorems are stated on the torus, by Proposition 4.7.
- Each T_j is a torus of the form [1, (1.2), p. 604] (Subsection 4.1).

Theorems 4.1 and 4.2 are stated for Bałaban's step count $K' - n_0 \ge 1$, which holds on $\mathfrak{C}$ because $K \ge n_0 + 1$. Proposition 6.4 follows, and with it Proposition 6.5: the states of $\mathfrak{C}$ have the coefficient b_{T_j}, independent of μ_0; N is even; and the reflection positivity of Proposition 7.9 holds.

(5) The curvature bound of Lemma 7.3 holds on every torus, for every real source of $:\varphi^2:$ and every bare coefficient. In particular it holds at b_{T_j}.

Steps (1)–(6) of the proof of Theorem 6.8 use exactly Proposition 6.5 at $q = 2k_+$, Proposition 6.7 at $h = 2k_-$ and Lemma 7.3, each on one torus at the coefficient b_{T_j}. By (2)–(5) all three hold on $\mathfrak{C}$. The constants σ_+, κ_+, $S_\flat^B$ do not depend on ε, j or μ_0. □

Appendix J. Measure-theoretic facts

The following three facts are used in Section 10: a theorem of Lusin and Souslin on injective Borel maps, the density of a generating algebra of bounded functions in L^2, and the reflection positivity of a Gaussian measure whose covariance is reflection positive.

Lemma J.1 (Lusin–Souslin)**.** *Let X be Polish and $\iota : X \to \mathbb{R}^{\mathbb{N}}$ continuous and injective. Then ι is a Borel isomorphism onto its image, which is Borel; in particular the Borel σ-algebra of X is generated by the coordinates of ι.*

Proof. [23, Thm. 15.1 and Cor. 15.2]. □

Lemma J.2 (density of generating algebras)**.** *Let m be a probability measure on a measurable space, and $\mathcal{A}$ an algebra of bounded complex functions, closed under complex conjugation and containing*

the constants. Then $\mathcal{A}$ is dense in $L^2(\sigma(\mathcal{A}), m)$. The same holds for a countable $\mathbb{Q}[i]$-algebra $\mathcal{A}_{\mathbb{Q}}$, closed under conjugation and containing 1*, whose complex span is such an algebra.*

Proof. Let $\mathcal{K}$ be the set of bounded real $\sigma(\mathcal{A})$-measurable functions lying in the $L^2(m)$-closure of the real parts of elements of $\mathcal{A}$. It is a vector space containing the constants and $\operatorname{Re}\mathcal{A}$ (which is an algebra, being the real algebra generated by real and imaginary parts), and it is closed under bounded monotone limits by dominated convergence. By the functional monotone class theorem [13, Ch. I, Thm. 21] it contains all bounded real $\sigma(\mathcal{A})$-measurable functions, which are dense in $L^2(\sigma(\mathcal{A}), m)$. Complex functions are treated by real and imaginary parts, and $\mathcal{A}_{\mathbb{Q}}$ is L^2-dense in its complex span. □

Lemma J.3 (reflection positivity of Gaussian laws)**.** *Let μ be a centred Gaussian measure on X with covariance C, invariant under a reflection θ with $\theta(\Sigma(H_j^+(0))) = \Sigma(H_j^-(0))$, and suppose $\langle \theta f, Cf\rangle \geq 0$ for all real f supported in $H_j^+(0)$. Then μ satisfies* (10.2) *for $c = 0$.*

Proof. For real $f_1, \dots, f_N$ supported in $H_j^+(0)$ and $\Theta_k = \mathrm{e}^{i\Phi(f_k)}$,

$$\mathbb{E}\big[\overline{\Theta_k \circ \theta}\,\Theta_l\big] = \mathrm{e}^{-\frac{1}{2}\langle f_k, Cf_k\rangle}\,\mathrm{e}^{-\frac{1}{2}\langle f_l, Cf_l\rangle}\,\mathrm{e}^{\langle \theta f_k, Cf_l\rangle},$$

using $\langle \theta f, C\theta f\rangle = \langle f, Cf\rangle$. The kernel $(k,l) \mapsto \langle \theta f_k, Cf_l\rangle$ is positive semidefinite (it is the form of the hypothesis, polarised, on the span of the f_k), so by the Schur product theorem its entrywise exponential is positive semidefinite, and so is the displayed matrix. Hence (10.2) holds on the span of such Θ_k, and by Lemma J.2 on all of $L^2(\Sigma(H_j^+(0)), \mu)$. □

Appendix K. Statements used without proof in this version

The proofs of Theorems 3.1 and 5.2 use the statements collected in this appendix. Their proofs are not included in this version. Each statement is in the notation of the place where it is used, and is followed by a list of these places. Statements that are printed with a proof in the literature are cited where they are used and are not repeated here.

Statement K.1 (the per-step constants of the large-field estimate are free of ε_0)**.** Let $\pi \in \mathfrak{W}$ and let ε_1 be as in Lemma A.11. Each of the following estimates holds at every scale $k \leq K$, given the inductive hypothesis at scale k, with a constant per site of at most $F(t_k)$: the replacements [2, (2.47)–(2.48)] and their inverse [2, (3.1)]; [2, Prop. 2.1, (2.57)], with the vacuum part on the collar bounded by [3, Prop. 1]; the mass-counterterm pieces of step k and the layer pieces of Lemma F.32; [2, Prop. 3.1, (3.26)] in the form of Lemma E.149; and the normalisations and the rescaling [2, (3.32)–(3.36)]. Here $F : (0,1] \to [0,\infty)$ satisfies $\sup_{0<t\leq\varepsilon_1} F(t) < \infty$, and it depends on the structural data and on π but not on ε_0, K, k, the torus, the region sequence, the value of u or the mass $M \in \{M_1, m, m_1, m_2\}$. In particular the constants c_5 and c_7 of [2, (3.42)] satisfy $\sup_{0<\varepsilon_0\leq\varepsilon_1} c_5 < \infty$ and $\sup_{0<\varepsilon_0\leq\varepsilon_1} c_7 < \infty$.

This statement is used in Proposition 5.21, Lemma A.18 (the condition (E.ent) of stage 10b) and Lemma A.11.
The periodisation lemma and the kernel bounds (Section 5 and Lemma 5.22).

Statement K.2 (the bounds (H_∞) for the scalar kernels at a mass M)**. Setting.** Let $L \geq 2$ be an integer, $a > 0$, $a_\flat := a(1 - L^{-2})$, $a_j \in [a_\flat, a]$ for $j \geq 0$, and $\alpha \in [0,1)$. Let $\varepsilon_0 > 0$, $K \geq 0$, $t_j := L^{j-K}\varepsilon_0$ and $0 \leq k \leq K$, and let $M \geq 0$ with $M\varepsilon_0 \leq 1$.

Lattice and operators. Work on $\eta\mathbb{Z}^3$, $\eta := L^{-k}$, with inner product $\eta^3 \sum$; kernels are taken with respect to $\eta^3 \sum$, and $\partial_{u,\mu} f(u) := \eta^{-1}(f(u+\eta e_\mu) - f(u))$. Fields are $\mathbb{R}^2$-valued and all operators act componentwise. For $1 \leq i \leq k+1$ let P_i be the orthogonal projection onto the fields constant on the blocks $y + [0, L^{i-k})^3$, $y \in L^{i-k}\mathbb{Z}^3$, and put

$$G_i := \big(-\Delta + M^2 t_k^2 + a_i L^{2(k-i)} P_i\big)^{-1}, \qquad G_{(0)} := G_1, \qquad G_{(j)} := G_{j+1} - G_j \ (1 \leq j \leq k),$$

and $\ell_j := L^{j-k}$ for $0 \le j \le k$. For $\sigma, \tau \in \{0, e_1, e_2, e_3\}$ let $\partial_u^\sigma \partial_v^\tau$ be at most one difference in each variable, and put $n := |\sigma| + |\tau|$.

Claim. There are $\kappa > 0$ and $C < \infty$, depending only on a, L, α, such that the following hold for all $K, k, \varepsilon_0, M, (a_j)$ as above, all $0 \le j \le k$ and all $1 \le i \le k+1$.

(i) For all u, v:

$$|\partial_u^\sigma \partial_v^\tau G_{(j)}(u,v)| \le C\ell_j^{-1-n} \mathrm{e}^{-\kappa|u-v|_1/\ell_j}, \qquad \eta^3 \sum_v |\partial_u^\sigma \partial_v^\tau G_{(j)}(u,v)| \le C\ell_j^{2-n}.$$

If $|\sigma| = 1$, then for all $u_1 \ne u_2$,

$$|\partial^\sigma \partial^\tau G_{(j)}(u_1,v) - \partial^\sigma \partial^\tau G_{(j)}(u_2,v)| \le C\ell_j^{-1-n-\alpha} |u_1-u_2|^\alpha \mathrm{e}^{-\kappa \min_r |u_r - v|_1/\ell_j}.$$

The same Hölder bound holds in the second variable when $|\tau| = 1$.

(ii) Put $\delta_0 := \kappa/2$. There is $C_0 = C_0(a, L, \alpha)$ such that for every $G \in \{G_i, G_{(j)}\}$:
- $|\partial_u^\sigma \partial_v^\tau G(u,v)| \le C_0 \mathrm{e}^{-\delta_0|u-v|}$ if $|u-v| \ge 1$;
- if $|\sigma| = 1$ and $\min_r |u_r - v| \ge 1$, the difference quotient of (i) is at most $C_0|u_1 - u_2|^\alpha \mathrm{e}^{-\delta_0 \min_r |u_r - v|}$, and likewise in v;
- if $n \le 1$, then $\eta^3 \sum_v |\partial_u^\sigma \partial_v^\tau G(u,v)| \le C_0$.

(iii) If $n = 2$, then $\eta^3 \sum_v |\partial_u^\sigma \partial_v^\tau G_i(u,v)| \le C\, i$.

This statement is used in Subsection 5.5, for the hypothesis (H_∞) of Lemma 5.22 at the Higgs masses $M \in (0, m_2]$ with $M\varepsilon_0 \le 1$. It is used in particular for the counterterm graphs of Lemma 5.22(v) at $M \in \{m_1, m_2\}$.

Statement K.3 (vector propagators on the torus and on the lattice)**. Lattices and fields.** Let $0 \le k \le K$ and $\eta := \eta_k$. Let T_η be the torus at step k of Subsection 5.5, with sides $S_k^{(\mu)}$ and deck group Γ_k, and assume that every $S_k^{(\mu)}$ is a multiple of L (for $k < K$ this holds since $S_k^{(\mu)} = LS_{k+1}^{(\mu)}$). Let $\Lambda \in \{T_\eta, \eta\mathbb{Z}^3\}$, in scale-$k$ units. Write $A_\mu(x) := A_{(x, x+\eta e_\mu)}$ and $\mathcal{H} := \ell^2(\Lambda; \mathbb{R}^3)$ with $\langle f, g\rangle = \sum_x \eta^3 f(x) \cdot g(x)$.

Block averages. For $1 \le i \le k+1$ let $\ell_i := L^{i-k}$. Partition Λ into the blocks of side ℓ_i with corners on $\ell_i\mathbb{Z}^3$ (on T_η, their images), and let $\mathcal{H}_i$ be the fields on the blocks with weight ℓ_i^3. Let $Q_i : \mathcal{H} \to \mathcal{H}_i$ be the block average and $P_i := Q_i^* Q_i$. Let $Q : \mathcal{H}_i \to \mathcal{H}_{i+1}$ be the one-step average, so that $Q_{i+1} = QQ_i$, and put $P := Q^*Q$.

Propagators. Let $\mu_0 \ge 0$, $\mu^2 := \mu_0^2 t_k^2$, $\alpha_i := a_i L^{2(k-i)}$ and $\beta_i := aL^{2(k-i)-2}$, where the a_i are the constants of [1, (2.13)], so that $\alpha_{i+1} = \alpha_i\beta_i/(\alpha_i + \beta_i)$. With $-\Delta$ the componentwise lattice Laplacian, put

$$H_i := -\Delta + \mu^2 + \alpha_i P_i, \quad G_i := H_i^{-1}, \quad C^{(i)} := (\alpha_i - \alpha_i^2 Q_i G_i Q_i^* + \beta_i P)^{-1},$$

$$W_{(0)} := G_1, \qquad W_{(i)} := \alpha_i^2 G_i Q_i^* C^{(i)} Q_i G_i \quad (1 \le i \le k).$$

The kernel classes. Let $\mathcal{W}$ consist of the kernels G_i $(1 \le i \le k+1)$ and $W_{(i)}$ $(0 \le i \le k)$, and of their compositions, in either variable, with lattice derivatives of order ≤ 2, with plaquette sums, and with the contour sums $f \mapsto \sum_{b \in \Gamma_{x_{j+1},x}^{(j+1)}} \eta f(b)$, $j+1 \le k$, of [24, (3.27)]. Kernels are taken with respect to $\eta^3 \sum$. Let $\mathcal{W}' \subset \mathcal{W}$ consist of the elements that contain no derivative of order 2 of a kernel G_i.

Claim.

(a) $H_i \ge \gamma_i > 0$ on $\mathcal{H}$, and on $\eta\mathbb{Z}^3$ one has $|G_i(x,y)| \le c_i \mathrm{e}^{-\rho_i|x-y|}$ for some $c_i, \rho_i > 0$.

(b) On each Λ, $C^{(i)}$ is bounded and $W_{(i)} = G_{i+1} - G_i$ for $1 \le i \le k$.

(c) For every $W \in \mathcal{W}$, $W_T(x,x') = \sum_{n \in \Gamma_k} W_\infty(\tilde{x}, \tilde{x}' + n)$, with absolute convergence.

(d) Let $\mathcal{T}$ be a family of such tori at step k, all of whose sides are multiples of L, with minimal sides tending to infinity. Assume:

- the kernels W_T, $T \in \mathcal{T}$, satisfy the pointwise and Hölder bounds of (H_∞) for every $W \in \mathcal{W}$;
- they satisfy the summability bound of (H_∞) for every $W \in \mathcal{W}'$;
- the constants C_0, δ_0 in these bounds are independent of T.

Then the kernels W_∞ satisfy the same bounds.

(e) Assume (E.r) and (5.13), and let $\square \subset T_\eta$ be a non-wrapping cube of side $3r(t_k)$, with lift map ι. Then every $W \in \mathcal{W}$ satisfies $W_T = W_\infty \circ (\iota \times \iota) + \delta W$ on $\square \times \square$, with $\|\delta W\|_{1,\alpha} \le C_1 \mathrm{e}^{-\delta_0 S_k/2} \le C_1 \mathrm{e}^{-3\delta_0 r(t_k)}$. Here $C_1 = c_d C_0$, and C_0, δ_0 are independent of k, of the torus and of ε_0.

(f) Parts (a)–(c) hold with $\mathbb{R}^3$ replaced by $\mathbb{R}^2$ and μ^2 by $m^2 t_k^2$, $m \ge 0$. Then $W_{(0)} = G_1$ and $W_{(j)} = G_{j+1} - G_j$ $(1 \le j \le k)$ are the zero-background scalar slices $G_{(j)}^{\eta_k}(0)$ of [3, (1.27), (2.6)].

This statement is used in the proof of Lemma 5.22(i), for the scalar slices (part (f)), and in Subsection 5.5, for the vector kernels of Lemma 5.22(v) (parts (c)–(e)).
The final c-numbers.

Statement K.4 (pinned two-leg classes without a partner, above ε_0)**.** Assume (E.1). Let $K \le j^* \le \bar{K}$, put $t := t_{j^*}$ and work in scale-j^* units with the torus distance. For a function F on $T_\varepsilon \times T_\varepsilon$ let $N(F)$ be the maximum over $a, b \in \{0,1\}$ of $\sup|\partial_1^a \partial_2^b F|$ plus the Hölder-α_0 quotients of $\partial_1^a \partial_2^b F$ in either variable. Let Q be a two-leg class of type C1′, C3 or C4 of Lemma E.166, nested and one-particle-reducible members included, whose lines are pieces $C_{(i),f}$ (Definition E.12) of index $\le j^*$ and $< \bar{K}$; let both external legs carry a two-variable field F with $N(F) < \infty$, and let $e(t)^\alpha \lambda(t)^\beta$ be the coupling factor of Q. Fix the vertex of one external leg at x, and let $V_Q(x)$ be the sum of the members together with the partners of their proper divergent subgraphs, over the positions of all other vertices and over the indices. If Q is of type C3 or C4 assume: for every $K \le j_0 \le \bar{K}$, with $\xi = L^{-j_0}$ and every line replaced by C_T^ξ, the zeroth moments at a base point of the class with its partners, and for C3 also the first moments, are bounded by a constant C_{TW} uniformly in ξ, the base point and $\ell_T \ge L^3$. Then, uniformly in x, ε and $\ell_T \ge L^3$,

$$|V_Q(x)| \le C\, e(t)^\alpha \lambda(t)^\beta N(F), \qquad C = C(\bar{n}, d, L, a, \alpha_0, M_b, C_{\mathrm{TW}}).$$

This statement is used in the proof of Proposition E.21 (Steps 2–4); for C3 its hypothesis is Proposition E.20(i).

Statement K.5 (the torus moment of the class C4)**.** Assume (E.1). Let $K \le j_0 \le \bar{K}$ and $\xi = L^{-j_0}$, and replace every line of the members of the class C4 of Lemma E.166, and of the partners of their proper divergent subgraphs, by the resolvent C_T^ξ of $-\Delta^\xi + 1$ on the rescaled torus T^ξ. Then the zeroth moment $\Pi^{(4)}(y)$ at a base point $y \in T^\xi$ satisfies $|\Pi^{(4)}(y)| \le C(\bar{n}, d, L, a)$, uniformly in ξ, y and $\ell_T \ge L^3$.

This statement is used in the proof of Proposition E.21 (Step 2(d)), as the hypothesis of Statement K.4 for C4.

Statement K.6 (one-particle-reducible chains above ε_0)**.** In the setting of Statement K.4, let the two external legs of a scalar two-leg class of type C2 whose members are one-particle reducible carry a two-variable field F with $N(F) < \infty$, and fix the external vertex at x. Then the pinned value $V(x)$ of the class (the members together with the partners of their divergent subgraphs) satisfies, uniformly in x, ε and $\ell_T \ge L^3$,

$$|V(x)| \le C \sum_{\alpha + 2\beta = 4} |e(t)|^\alpha \lambda(t)^\beta\, N(F), \qquad C = C(\bar{n}, d, L, a, \alpha_0, M_b).$$

This statement is used in the proof of Proposition E.21 (Steps 2(c) and 4).

Statement K.7 ($E_{\mathrm{fin}}(u)$ at $u \neq 0$)**.** Let $M \in (0, m_2]$, assume (E.1), and let ε_0 be a power of L. Let $E_{\mathrm{fin}}(u)$ be as in (E.9) and $\bar\sigma_N(u) := (1 + |u|\,\kappa\mu_*^2 L^2 t_{\bar K}^2)^{N_\ell}$. Let $\hat B$ be the right-hand side of Proposition E.21, including the factor Φ, with C_E, C_E^R replaced by constants $\hat C_E = C(\alpha_0, \bar n, d, L, a, M_b) \geq C_E$ and $\hat C_E^R = C(\alpha_0, \bar n, d, L, a, M_b)(1 + \rho_{\min}^{-2} + \kappa_{R,\min}^{-3})^{N_\ell} \geq C_E^R$. Then for every $u \in \mathbb{R}$ and every $\varepsilon = 2^{-K}\varepsilon_0$,

$$|E_{\mathrm{fin}}(u) - E_{\mathrm{fin}}(0)| \leq \big(\bar\sigma_N(u) - 1\big)\hat B, \qquad |E_{\mathrm{fin}}(u)| \leq \bar\sigma_N(u)\,\hat B.$$

This statement is used in the proof of Proposition E.162.

Statement K.8 (periodisation of the vector slices)**.** Let $0 \leq j \leq k < K$. The kernel on T_ε, in scale-k units, of the vector slice of index j of [3, p. 426], built from $H = -\Delta + \mu^2 + a_j P_j$ with $-\Delta$ the componentwise lattice Laplacian, and each of its lattice derivatives of order ≤ 2, is the periodisation of the corresponding kernel on $\eta\mathbb{Z}^3$, and the latter satisfies (H_∞) of Subsection 5.5.

This statement is used in the proof of Lemma E.173.

Statement K.9 (the vacuum output for Bałaban's scalar staging)**.** Consider Bałaban's staging of the scalar integrals [3, (1.25), (1.27)], in which the fluctuation covariance of step k depends on the gauge background, with the truncation of the effective action at order $\bar n$ in the couplings. Then the part of order $\leq \bar n$ of the output of step k at zero block fields and zero background equals $\mathcal{V}_k$ of Definition E.12, the slices being evaluated at zero background.

This statement is used in the proof of Lemma F.31.
The coefficients of the $A \cdot A$ vertices.

Statement K.10 (power counting for generalised graphs)**.** Let $\Gamma = (V, \mathcal{L})$ be a generalised graph (Definition E.28), $L \geq 2$, $N \geq 0$, $\delta > 0$, $\kappa > 0$ and $J \subset \{0, \dots, N\}^{\mathcal{L}}$. Assume that for every $\mathbf{j} \in J$, every i with $\min_l j_l \leq i < N$, and every connected component H of $\Gamma_i(\mathbf{j})$, $D(H) \geq \kappa$. Then

$$\sum_{\mathbf{j} \in J} E_{\mathbf{j}}(\Gamma) \leq C_\delta^{|V|-1} \sum_{n \geq 0} (n+1)^{|\mathcal{L}|} L^{-\kappa n} < \infty, \qquad C_\delta := (1 + 2\sqrt{d}/\delta)^d,$$

uniformly in N and y_{v_0}. The same holds with $\xi\mathbb{Z}^d$ replaced by $\xi\mathbb{Z}^d/\ell\mathbb{Z}^d$ and $|y - y'|$ by the torus distance. Consequently:

(a) if every connected subgraph of Γ has positive degree, then $\sum_{\mathbf{j}} E_{\mathbf{j}}(\Gamma) \leq C$, the sum running over all assignments;
(b) fix an ordering $l(1), \dots, l(|\mathcal{L}|)$ of $\mathcal{L}$, let $J = \{\mathbf{j} : j_{l(1)} \leq \cdots \leq j_{l(|\mathcal{L}|)}\}$ and $G_p = \{l(1), \dots, l(p)\}$; if every component of every G_p has positive degree, then $\sum_{\mathbf{j} \in J} E_{\mathbf{j}}(\Gamma) \leq C$.

In (a) and (b), C depends only on $d, \delta, L, |V|, |\mathcal{L}|$, the dimensions ω_l and the exponents p_v. This is [3, Prop. 2.2] in the form needed here.

This statement is used in the proof of Lemma E.31 and in the proof of Lemma E.32.

Statement K.11 (the remainder of [3, (3.31)] reduced to the inner renormalisation)**.** Let $0 \leq m_{j_0}^2 \leq 1$, $(\alpha, \beta) \in \{(4,0), (2,1)\}$, let the photon lines be as in Lemma E.32, and assume the bounds (N.1)–(N.5) of Statement K.12. Then there is $K_R < \infty$ with $|\hat c_{\mu\mu'}^{\Lambda}(y)| \leq K_R$ for every label configuration Λ, every y, every $\xi \in L^{-\mathbb{N}}$, j_0, k, ε, ℓ_T, and every $m_{j_0}^2 \in [0,1]$. Here $\hat c^{\Lambda}$ is as in Proposition E.27. K_R depends only on:

- d, a, L and the block size;
- the constants of the slice bounds of the scalar lines and of the photon lines, the latter independent of μ_*, with weights in $[0, 2]$;
- $\bar c_{\mathrm{fin}}$, with $\bar c_{\mathrm{fin}} := 0$ if the class contains no fixed mass vertex.

The bound is obtained by arranging the absolutely convergent sum of Proposition E.27(b) as follows: all chains are summed together, over all $S \neq \emptyset$ and over both order-one factors $\Pi^1, \Pi^2 \in \mathcal{P}$; every other member T is summed at fixed S over $\mathcal{K}(T)$.

This statement is used in the proof of Lemma E.32.

Statement K.12 (inner renormalisation)**.** Let $0 \le m_{j_0}^2 \le 1$, $(\alpha, \beta) \in \{(4,0),(2,1)\}$, let the photon lines be as in Lemma E.32, and let Λ be a label configuration. For a member T of Π_G^Λ and $\emptyset \neq S$, a critical subgraph H of T^S is said to be of:

- type 1 if it is a single scalar line with two differences joining two distinct vertices $\varphi \cdot q\partial\varphi\, a$ (the graph [3, (2.4)]);
- type 2 if it has exactly two external legs, both scalar, and no difference of a vertex of H acts on them;
- type 3 if it has exactly two external legs, both scalar, and a difference acts on exactly one of them;
- type 4 if it has order one and two external vector legs and no other external leg;
- type 5 if it has exactly three external legs, two scalar and one vector.

Then there is a constant C_N with the following properties, uniformly in Λ, the bond b, ξ, j_0, k, ε, ℓ_T and $m_{j_0}^2 \in [0,1]$:

(N.1) If T is not a chain and has no counterterm partner, and every critical subgraph of T^S is of type 1, then $\big|\sum_{b'} \xi^3 T^S(b,b')\big| \le C_N$.
(N.2) If T is not a chain and T^S has a critical subgraph of type 2, then $\big|\sum_{T' \in \mathcal{K}(T)} \sum_{b'} \xi^3 T'^S(b,b')\big| \le C_N$.
(N.3) The same bound as in (N.2) holds if T is not a chain and T^S has a critical subgraph of type 3.
(N.4) Let $\mathcal{K}_{\mathrm{ch}}$ be the set of chains of Π_G^Λ. Then $\big|\sum_{T \in \mathcal{K}_{\mathrm{ch}}} \sum_{S \neq \emptyset} (-1)^{|S|+1} \sum_{b'} \xi^3 T^S(b,b')\big| \le C_N$.
(N.5) The same bound as in (N.2) holds if T is not a chain and T^S has a critical subgraph of type 5.

C_N depends only on the order, d, a, L, the block size, the constants of the slice bounds of the scalar and photon lines (the latter independent of μ_*, with weights in $[0,2]$), and $\bar{c}_{\mathrm{fin}}$.

This statement is used in the proof of Lemma E.32, through Statement K.11.
The tail beyond the order $\bar{n}$.

Statement K.13 (exactness of the split of $\theta_{k+1}A^{(k)}$ at $L=2$)**.** Let $L = 2$ and $1 \le k \le K-1$, and measure lengths in units of $T_1^{(k)}$ (Euclidean distance). Let $\ell(t) = 1 + \log t^{-1}$, $r(t) = R\ell(t)^2$, and $\hat{\ell} := \ell(\varepsilon_0)$. Use the following conventions of [2]. (i) The sets $\Lambda_i^{(k)} \subset T_1^{(k)}$ are built from $r(t_k)$ as in [2, (2.7)–(2.8), p. 558; p. 566]: $(\Lambda_{i+1}^{(k)})^c$ is the union of the large blocks (of side M_b) at distance $\le r(t_k)$ from $(\Lambda_i^{(k)})^c$. (ii) For a union X of L-blocks, X' is the set of $y \in T_L^{(k)}$ with $B(y) \subset X$; every point of $B(y)$ lies within $L\sqrt{d}$ of y. (iii) $\theta_{k+1} = 1$ on $B^k(\Lambda_2^{(k)})$, and θ_{k+1} vanishes at distance $\ge LM_b$ from it. (iv) $\zeta^{(k+1)}(x,y) = 0$ for $|x-y| \ge L\big(r(t_{k+1}) - 2M_b\big)$, the support radius of [2, (2.44), p. 566] being measured in units of $T_1^{(k+1)}$. Let A', B, $\delta C^{(k)}_{\Lambda_0^{(k)}} = C^{(k)}_{\Lambda_0^{(k)}} - C^{(k)}$, G_k, G_{k+1}^η, Q_k^*, Q_{k+1}^*, Q^*, a_k, a_{k+1} and a be as in [2, (2.80)–(2.89), pp. 574–575], products such as $\zeta G Q^*$ being kernel products. Assume

$$r(\varepsilon_0) \ge 2M_b, \qquad M_b + 2R\log 2\,\hat{\ell} \ge 2\sqrt{d};$$

the second inequality holds whenever $R \geq \sqrt{d}/\log 2$. Then $\zeta^{(k+1)}(x,y) = 0$ whenever $\theta_{k+1}(x) \neq 0$ and $y \notin \Lambda_0^{(k)\prime}$. Moreover, identically in A' and B,

$$\begin{aligned}\theta_{k+1}A^{(k)} &= a_k\theta_{k+1}\zeta^{(k)}G_kQ_k^*A' + a_kaL^{-2}\theta_{k+1}\zeta^{(k)}G_kQ_k^*\,\delta C^{(k)}_{\Lambda_0^{(k)}}Q^*B \\ &\quad - a_kaL^{-2}\theta_{k+1}(1-\zeta^{(k)})G_kQ_k^*C^{(k)}Q^*\Lambda_0^{(k)\prime}B + a_{k+1}L^{-2}\theta_{k+1}(1-\zeta^{(k+1)})G^\eta_{k+1}Q^*_{k+1}\Lambda_0^{(k)\prime}B \\ &\quad + a_{k+1}L^{-2}\theta_{k+1}\zeta^{(k+1)}G^\eta_{k+1}Q^*_{k+1}B,\end{aligned}$$

which is [2, (2.89)] as printed.

This statement is used in the proof of Proposition E.36.

Statement K.14 (exceptional vertices with a chain leg)**.** Let $0 \leq k \leq K-1$, $t = t_k$, $p = p(t)$, and work on $\operatorname{supp}\chi_k$ under the conclusions of Lemma E.33(c). Let v be a vertex of coupling order one of a connected localised family of $\mathcal{P}_w^{(k)}$ with at least two vertices, both of whose scalar legs are external fields. Let one of these be a chain $\mathcal{K}f$ of order $m \geq 1$, that is, a term of the expansions (E1)–(E2) of the proof of Proposition E.36 of $G_k(\Omega, \mathfrak{B}+\tilde{A})$ and $Q_k(\mathfrak{B}+\tilde{A})$ with at least one insertion. Then v together with this chain contributes to the bound [3, (1.33)] at most $C\,t^{(1+m)/4}p^{2+m}$, with C independent of ε, k and ℓ_T.

This statement is used in the proof of Proposition E.36 (Step 3a).

Statement K.15 (exceptional vertices with zeroth-order legs: bond-type partner)**.** Let $1 \leq k \leq K-1$, $t = t_k$, $p = p(t)$, $\lambda(t) = \lambda t$, $M \leq m_2$, and let $\Omega_* = B^k(\Lambda_2^{(k-1)\prime})$, with vertex points $y \in \Lambda_7^{(k-1)\prime}$. Let ϕ be a scalar field on the vertex points and $\Phi_0 := a_kG_k(\Omega_*, \mathfrak{B})Q_k^*(\mathfrak{B})\phi$. Assume: (a) $|\phi(y)| \leq X'_{\underline{s}} := c_1'p\lambda(t)^{-1/4}$, and $|U(\bar{\mathfrak{B}}(\langle y', y''\rangle))\phi(y'') - \phi(y')| \leq c_1'p$ for coarse nearest neighbours y', y'', where $\bar{\mathfrak{B}}$ is the coarse configuration of $\mathfrak{B}$ and $U(A(\Gamma))$ is the parallel transport of [2] (it contains the factor $e(t)$); (b) $|\partial_\mu^\eta\mathfrak{B}| \leq C_{\mathrm{reg}}p$ on Ω_*; (c) every vertex point y satisfies $\operatorname{dist}(y, \Omega_*^c) \geq 5\rho_k$ with $\rho_k := (r(t_{k-1}) - M_b)/L$, and the conditions (5.6), (5.7), $\varepsilon_0 \leq \varepsilon(L, \varrho)$ and (T.nw) of Lemma 5.6 hold. Then: (i) there is C_{24}, independent of ε, k and ℓ_T, such that for every vertex point y, every $x \in B^k(y)$ and every bond $b \subset B^k(y)$, $|\Phi_0(x) - U(\mathfrak{B}(\Gamma_{x,y}))\phi(y)| \leq C_{24}p$ and $|(D_{\mathfrak{B}}\Phi_0)(b)| \leq C_{24}p$; (ii) a vertex v of coupling order one whose scalar legs are Φ_0 or $\phi(y)$ equals, as a function of its internal leg, $-e(t)\sum_b \eta^d J(b)g_kA'_b$ (for [3, (1.8)]) or $\sum_y\sum_{x\in B^k(y)}\eta^dA'(\Gamma_{y,x})j_y(x)$ (for the products of [3, (1.12)] or [3, (1.13)] with [3, (1.14)]). Here $\sup_b|e(t)J(b)| \leq C_{\mathrm{ex}}t^{1/4}p^2$ and $\sup_{y,x}|j_y(x)| \leq C_{\mathrm{ex}}t^{1/4}p^2$, with $C_{\mathrm{ex}} := \max\{1,a\}|e|\,C_{24}(2c_1'\lambda^{-1/4} + C_{24})$; (iii) assume the counterterm rule of [3, p. 420], with mass counterterm coefficients given by their zero-background values [3, (1.29)–(1.31)], and assume the family bound of Definition E.59 at graded order $\leq \bar{N}$ (Proposition E.60) for families with external vector fields of finite norm [3, (1.32)]. Let F be a connected localised family of $\mathcal{P}_w^{(k)}$ with at least two vertices containing such a vertex v, whose vector leg is contracted with a bond leg $(g_kA')_{b'}$ of a vertex of type [3, (1.8)–(1.11)]. Then F obeys the bound [3, (1.33)], with the factor $(|u|c_k)^m$ and with some decay rate $\delta_1 > 0$ independent of ε, k, ℓ_T, in which the factor $e(t)\|h\Phi_1\|_{1,\alpha_0}\|h\Phi_2\|_{1,\alpha_0}$ of v is replaced by $C\,C_{\mathrm{ex}}t^{1/4}p^2$, with C independent of ε, k and ℓ_T. No regularity of J or j_y beyond their suprema enters.

This statement is used in the proof of Proposition E.36 (Step 3a).

Statement K.16 (exceptional vertices with zeroth-order legs: contour-type partner)**.** Assume the setting and the hypotheses (a)–(c) and (iii) of Statement K.15. Let F be a connected localised family of $\mathcal{P}_w^{(k)}$ with at least two vertices containing a vertex v of coupling order one whose scalar legs are Φ_0 or $\phi(y)$, and whose vector leg is contracted with a contour leg $A'(\Gamma^{(k)}_{y',x'})$ of a factor [3, (1.14)] or [3, (1.15)] of a product [3, (1.18)]. Then F obeys the bound [3, (1.33)], with the

factor $(|u|c_k)^m$ and with some decay rate $\delta_1 > 0$ independent of ε, k, ℓ_T, in which the factor $e(t)\|h\Phi_1\|_{1,\alpha_0}\|h\Phi_2\|_{1,\alpha_0}$ of v is replaced by $C\,C_{\rm ex}t^{1/4}p^2$, with C independent of ε, k and ℓ_T.

This statement is assumed in Proposition E.36 and used in Step 3a of its proof.
The moved vacuum pieces and the base of Bałaban's Proposition 1.

Statement K.17 (King's slices)**.** Let $L = 2$, and on T_ε use the scalar product $\langle f, g\rangle = \sum_x \varepsilon^3 f(x)g(x)$. Put $Y_M := -\Delta_\varepsilon + M^2$ and $C_M := Y_M^{-1}$, and let $t_j = L^j\varepsilon$. Let P_j, Q_j, Q, a_j, $r_j = a_j t_j^{-2}$, G_j, $C^{(j)}$, F_j, H_j and G_k^M be the operators and coefficients of [24, (2.13)–(2.17)], with King's constants $0 < a_\flat \le a$. For $K \ge 0$ put $s_0(K) := \sup_x K(x,x)$. Put $m_a := \max(\frac{1}{8}, \frac{1}{a_\flat})$, $c_C := \frac{4}{a_\flat} + \frac{L^2}{8} + \frac{L^2}{a}$, $C_{\rm osc}^2 := (1 + \frac{\sqrt{3}}{2})^3\frac{\pi^2}{8}$ and $C_H := c_C a^2(m_a + 3^{1/4}C_{\rm osc})^2$. At $L = 2$, $a = 1$ these are $m_a = \frac{4}{3}$, $c_C = \frac{59}{6}$, $C_{\rm osc}^2 < 8.02$ and $C_H < 253.66$. Then:

(S1) For $1 \le j \le K$: $Q_jQ_j^* = 1$, and $Q_j^*Q_j = P_j$ is the orthogonal projection onto the functions that are constant on the blocks of side t_j. Moreover $Q_{j+1} = QQ_j$, $QQ^* = 1$ and $P_jP_{j+1} = P_{j+1}P_j = P_{j+1}$.

(S2) For $1 \le j \le K-1$:
- $a_\flat \le a_j \le a$, so $0 < r_j \le at_j^{-2}$;
- $G_j = (Y_{\mu_*} + r_jP_j)^{-1}$ and $\|G_j\| \le m_a t_j^2$;
- $0 < C^{(j)} \le c_C t_j^2$;
- $H_j = F_jC^{(j)}F_j^* \ge 0$ with $F_j = r_jG_jQ_j^*$.

(S3) For $1 \le j \le K-1$, $s_0(H_j) \le C_H/t_j$. Moreover $0 < H_0 = G_1 \le C_{\mu_*}$ and $s_0(H_0) \le \frac{13}{16\varepsilon} + \frac{1}{\mu_*^2\ell_T^3}$.

(S4) For $1 \le k \le K$ and $M > 0$: $G_k^M = (Y_M + r_kP_k)^{-1}$ with $0 < r_k \le at_k^{-2}$. Moreover $0 < G_k^M \le C_M$ and $C_M - G_k^M = C_Mr_kP_kG_k^M$.

This statement is used in the proof of Theorem E.92, through Statement K.18(c).

Statement K.18 (inputs of the order-e^2 anchor bounds)**.** Let $L = 2$ and $\varepsilon \le \varepsilon_0$. Let G_M be the kernel of $(-\Delta_\varepsilon + M^2)^{-1}$ on $\varepsilon\mathbb{Z}^3$, let $\mathcal{G}_M$ be the kernel of $(\hat p^2 + M^2)^{-2}$ on $\varepsilon\mathbb{Z}^3$, and let c be the kernel of $C = (-\Delta_\varepsilon + M^2)^{-1}$ on T_ε, a torus of side $\ell_T \ge \ell_0$.

Put $c_N = \pi^3/8$, $v_0 = M/\sqrt{2}$, $\eta(s) := \frac{2}{\varepsilon}\operatorname{arcsinh}\frac{\varepsilon\sqrt{s}}{2\sqrt{2}}$, $c_0 = \frac{1}{32\pi}$, $c_1 = \frac{1}{6\pi^2} + \frac{1}{32\pi}$, $c_2 = \frac{5}{16\pi}$, $\beta = \frac{1}{2}\operatorname{arcsinh}2$ and $\bar w = \sqrt{12}c_N/(2\pi^2)$. Define

$$\kappa_a(r) := 16c_Nc_a\int_{v_0}^\infty v^a \mathrm{e}^{-\eta(2v^2)(r-a\varepsilon)}dv, \qquad \kappa_a^{\mathcal{G}}(r) := 8c_Nc_a\int_{v_0}^\infty v^{a-2}\mathrm{e}^{-\eta(2v^2)(r-a\varepsilon)}dv,$$

and $\rho_a(G_M) := \sum_{j\ge1}(24j^2+2)\kappa_a((j-\frac{1}{2})\ell_T - 4\varepsilon)$; define $\rho_a(\mathcal{G}_M)$ in the same way with $\kappa_a^{\mathcal{G}}$. For a kernel $K \ge 0$ put $s_0(K) := \sup_x K(x,x)$ and $s_1(K) := \sup_{x,\mu,\pm}\langle\nabla_\mu^{\pm *}\delta_x, K\nabla_\mu^{\pm *}\delta_x\rangle$. Put $\ell' := \ell_T/2 + \varepsilon$ and

$$I(\ell_T) := \hat G_1\rho_1\ell'^2 + \hat G_0\rho_2\ell'^3/3 + \hat G_2\rho_0\ell' + (\rho_1^2 + \rho_0\rho_2)\ell'^4/4,$$

with $\rho_a = \rho_a(G_M)$. Then:

(a) $\widehat\Pi_\mu(0) := 2e^2c(\varepsilon e_\mu) - 2e^2\varepsilon^{-2}\sum_z \varepsilon^3[c(z)^2 - c(z+\varepsilon e_\mu)c(z-\varepsilon e_\mu)]$ satisfies $|\widehat\Pi_\mu(0)| \le e^2w_T$, where $w_T := 2\rho_0(G_M) + 4\rho_2(\mathcal{G}_M)$.

(b) For $z \ne 0$ and a product ∇^α of $a \le 2$ forward or backward differences, $|\nabla^\alpha G_M(z)| \le \hat G_a|z|_\infty^{-a-1}$, with $\hat G_0 = 1.361$, $\hat G_1 = 12.25$ and $\hat G_2 = 361.2$. Moreover $\hat B := \hat G_1^2 + \hat G_0\hat G_2 \le 641.7$. On the fundamental cell, $c = G_M + R$ with $|\nabla^\alpha R| \le \rho_a(G_M)$.

(c) Assume (S1)–(S4) of Statement K.17, and put $t_0 = \varepsilon$. Put $W_1 := 4374\hat A_0^2 + 2S_4\hat G_1^{*2}$, with $\hat A_0 = 0.861$, $\hat G_1^* = 3.402$ and $S_4 := \sum_{m\ge2}(24m^2+2)(m-1)^{-4} \le 125.32$, so that $W_1 \le 6144$.

Put $W_{\mathrm{img}}(m) := 216\rho_0(G_m)^2\varepsilon_0 + 2\ell_T^3\rho_1(G_m)^2$, and

$$\sigma_0 := C_H + \frac{\varepsilon_0}{\mu_*^2\ell_0^3}, \quad h := c_C a^2 m_a^2 \le 17.49, \quad \theta_H := c_C a^2[W_1 + \varepsilon_0 W_{\mathrm{img}}(\mu_*)],$$

$$\gamma := a\Big[\frac{c_N}{8\pi M} + \rho_0(\mathcal{G}_M)\Big], \qquad \theta := a[W_1 + \varepsilon_0 W_{\mathrm{img}}(M)],$$

with $c_C = \frac{59}{6}$. Then for all $j \ge 0$:

- $s_0(H_j) \le \sigma_0/t_j$, $\|H_j\| \le h t_j^2$ and $s_1(H_j) \le \theta_H/t_j^3$;
- for $1 \le k \le K$, $\mathcal{C}_k := C - G_k^M$ satisfies $0 \le \mathcal{C}_k \le C$, $s_0(\mathcal{C}_k) \le \gamma t_k^{-2}$, $s_1(\mathcal{C}_k) \le \theta t_k^{-3}$, $\sum_y \varepsilon^3 |\nabla' \mathcal{C}_k(x,y)|^2 \le \mathcal{C}_k(x,x)$ and $\|\mathcal{C}_k\| \le M^{-2}$;
- for every kernel $K \ge 0$: $|K(x,y)| \le s_0(K)$, $|\nabla K| \le \sqrt{s_0(K)s_1(K)}$, $|\nabla\nabla' K| \le s_1(K)$ and $\sum_y \varepsilon^3 K(x,y)^2 \le \|K\| K(x,x)$.

(d) $\sum_{z\in T_\varepsilon} \varepsilon^3 |\nabla_\mu^\pm c(z)| \le \Lambda_1 := 26 \cdot \frac{9}{4} \cdot 16 c_N c_1 \big[\frac{2\sqrt{2}}{\beta^3 M} + 0.1345\varepsilon_0\big] + 250\bar{w}\varepsilon_0 + \ell_T^3 \rho_1(G_M)$.

(e) $\#\{z \in \varepsilon\mathbb{Z}^3 : |z|_\infty = j\varepsilon\} = 24j^2 + 2$. If $f \ge 0$ is non-increasing, then $\sum_{j\ge 1} \varepsilon f(j\varepsilon) \le \int_0^\infty f$. For $q \ge -2$, $\sum_{0<|z|_\infty \le \ell_T/2} \varepsilon^3 |z|_\infty^q \le 26 \int_0^{\ell'} r^{2+q} dr$.

Each ρ_a is finite, non-increasing in ℓ_T, bounded uniformly in $\varepsilon \le \varepsilon_0$, and exponentially small in $M\ell_T$. All constants above are independent of $\varepsilon \le \varepsilon_0$, j and k, and are bounded for $\ell_T \ge \ell_0$.

This statement is used in the proofs of Proposition E.90, Lemma E.91 and Theorem E.92.

Statement K.19 (decay of King's slices)**.** In the setting of Statement K.17, let $M \in \{m_1, m_2\}$ and $G_{(i)} := G_{i+1}^M - G_i^M$ for $0 \le i \le K-1$, with $G_0^M := 0$. There are constants c_G, c_H, and the rate $\delta_1 > 0$ of [3, (2.10)–(2.11), p. 426], all independent of ε, i, K and of $\ell_T \ge \ell_0$, such that on T_ε, with $|x-y|$ the torus distance,

$$|\nabla^\alpha G_{(i)}(x,y)| \le c_G t_i^{-1-|\alpha|} \mathrm{e}^{-\delta_1 |x-y|/t_i}, \qquad |H_i(x,y)| \le c_H t_i^{-1} \mathrm{e}^{-\delta_1 |x-y|/t_i}.$$

The first bound holds for all $|\alpha| \le 2$ with at most one difference in each variable.

This statement is used in the proof of Theorem E.92(c).

Statement K.20 (graphs with a vertex (1.13)–(1.15))**.** Let $d = 3$, and let D be Bałaban's degree [3, (2.1)–(2.3)]. Every mass vertex carrying $\delta m^2_{\mathrm{fin}}$ or $\delta m^2_{K,k}$ has degree 0, and every mass vertex carrying the contribution of a graph G_0 has degree $D(G_0)$. Then:

- every mass counterterm satisfies $\deg(\delta m^2_{G_0}) = D(G_0) \ge -1$;
- $D(G) \in \mathbb{Z}$ for every connected graph G;
- every connected graph G built from the vertices [3, (1.6)–(1.15)], with (1.12) an external field, that contains a vertex of type (1.13)–(1.15) satisfies $D(G) \ge 1$.

The bound $D(G) \ge 1$ is attained.

This statement is used in the proof of Proposition E.87 and in Step 3 of the proof of Lemma E.86.

Statement K.21 (Bałaban's Proposition 1 on the torus)**.** Fix $\bar{n} > 12$, $0 < \alpha_0 < 1$, $k \le K$ and $\eta = L^{-k}$. Let $\Omega, \Omega_1, \Omega_2, \tilde{B}$ satisfy the hypotheses of [3, Prop. 1] in the form of Corollary 5.10. Work on T_ε, or on the torus with photon covariance $C_{\mu_*} + u\Gamma^B$ ($\mu_* > 0$, $|u| \le 1$), whose vector lines are coloured C or D; in scale-k units, a D-line carries the coefficient uc_k with $0 \le c_k \le \kappa\mu_*^2\varepsilon_0^2 L^2$. Then every renormalised family G_{ren} of [3, pp. 418–421], with localisation $\{\square(v)\}$, external fields $\Phi_{\mathrm{ext}}, A_{\mathrm{ext}}$ and localising functions h, h', satisfies

$$|E(G_{\mathrm{ren}}, \{\square(v)\}, \Phi_{\mathrm{ext}}, A_{\mathrm{ext}})| \le O(1)\, e(L^k\varepsilon)^{d_v} \lambda(L^k\varepsilon)^{d_s} \mathrm{e}^{-\delta_T d(\{\square(v)\})} \|h\Phi_{\mathrm{ext}}\|_{1,\alpha_0} \|h' A_{\mathrm{ext}}\|_{1,\alpha_0}.$$

Here $d(\{\square(v)\})$ is the length of the shortest tree joining the cubes, and d_v, d_s are as in [3, (1.33)]. The constant is $O(1) = O(\alpha_0, \bar{n}; d, L, a, M_b)$, and $\delta_T > 0$ depends on $\bar{n}$; both are independent of ε,

k, the domains and $\tilde{B}$. For a family with m D-lines, where $m = 0$, $m \geq 2$, or $m = 1$ and $o \geq 2$, the same bound holds times $(|u|c_k)^m$.

This statement is used in Proposition E.97, and through it in the proofs of Lemma E.86 and Proposition E.64.
Real backgrounds near Neumann faces and the order cap.

Statement K.22 (smallness of the remainder field)**.** Let $0 \leq k \leq K-1$, $t = t_k$, and let $\mathcal{A}$ be the fine vector field of step k whose fluxes along the composite contours define the scalar restrictions [2, (2.16), (2.53), (2.55)]. Let $\tilde{B}$ be the real interpolated background of Lemma E.70(3), and $A'' := \mathcal{A} - \tilde{B}$ the remainder field of the decomposition [2, (2.19)–(2.21)]. Then there is a constant c'' that depends only on d, L, a, M_b, not on $\varepsilon, k, t, \ell_T$ or the domains, such that, on the small-field set of step k, $|A''(b)| \leq c'' p(t)$ for every bond b of the set Ω' of blocks on which these restrictions are imposed.

This statement is used in Statement K.23, hence in the proof of Proposition E.76 (through Lemma E.79).

Statement K.23 (nearest-neighbour bound with transport by the background)**.** Assume Statement K.22, $|\partial_\mu^\eta \tilde{B}_\nu| \leq C_\partial p(t)$ on Ω', $\ell_T/t > 2$, and let ψ be the block scalar field of step k, which satisfies the restrictions [2, (2.16)] at scale t and $|\psi(y)| \leq 2\lambda(t)^{-1/4} p(t)$. Put $\kappa(t) := |e|\lambda^{-1/4} t^{1/4} p(t)$. Then for nearest neighbours y_1, y_2 of the unit lattice with $B(y_1), B(y_2) \subset \Omega'$ and Γ_{y_1,y_2} the straight contour,

$$|U(\tilde{B}(\Gamma_{y_1,y_2}))\psi(y_2) - \psi(y_1)| \leq c_\psi(t) p(t), \qquad c_\psi(t) := 4Ld + \big(2(2dL - 2d + 1)c'' + 8(d+1)^2 C_\partial\big)\kappa(t).$$

This statement is used in the proof of Proposition E.76, as hypothesis ($\mathrm{N_{nn}}$) of Lemma E.79.

Statement K.24 (closure of the renormalised classes under cutting)**.** Let G_{ren} be a renormalised class of [3, pp. 420, 429–432], with counterterm members written with the vertices and lines of the graph they replace and the lower-index leg re-attached [3, p. 435]. A label configuration Λ assigns a localisation cube to each vector leg, each λ-vertex, each finite-counterterm vertex, each vertex (1.13)–(1.15) and each external leg; N_s is the number of these labels. Partition the support cubes (side 2) into clusters, the components of the relation $\mathrm{dist}(\square, \square') < 1$, and call a line cut if its end vertices lie in different clusters. Then: (a) at each Λ either no member has a cut line or every member not vanishing at Λ has one; (b) every cut line joins vertices with $\mathrm{dist}(\square(v), \square(v')) \geq 1$, every cluster lies in a cube of side $\leq 3N_s$, and $d(\{\square(v)\}) \leq \sum_{\ell \text{ cut}} \mathrm{dist}(\square(v_\ell), \square(v'_\ell)) + 3N_s$; (c) the cut family, cut lines read as two-variable external fields, is a disjoint union of renormalised classes, each connected component of a member localised in one cluster, and [3, (3.2)] for these classes implies [3, (1.33)], with $\mathrm{e}^{-\delta_0 d(\{\square(v)\})}$ recovered up to $\mathrm{e}^{3\delta_0 N_s}$; (d) for a divergent vacuum graph G and a line ℓ, either the term vanishes (photon bridge) or $G \setminus \ell$ is a bare vertex, convergent, or a two-leg divergent graph whose partners are members of the vacuum class of G.

This statement is used in the proof of Proposition E.76 and of Proposition E.84.

Statement K.25 (bad localisations of the counterterms)**.** Let $0 \leq k < K$, $t = t_k$, $W := B^k(\Lambda_7^{(k)})$, $c_d := 2\sqrt{d}$, $\delta_1 := \delta_0/2$, and assume $\ell_T/t \geq 6r(\varepsilon_0)$, $\mu_* t \leq 1$, $m^2 t^2 \leq 1$. Let $\Delta_k(x) := [\delta m^2_{K,k}(x) - \delta m^2_{K,k+1}(x)]t^2$ (notation of Lemma E.80), localised as in [3, p. 420] with the pinned cube containing x; a localisation is bad if one of its cubes is not contained in W, and Δ_k^{bad} is the sum over bad localisations. With $P_m(t) := (e^2 + \lambda)t + (e^4 + e^2\lambda + \lambda^2)t^2$ there is C_d independent of ε, k, x, the domains and ℓ_T with $|\Delta_k(x)|, |\Delta_k^{\mathrm{bad}}(x)| \leq C_d P_m(t)$, and $|\Delta_k^{\mathrm{bad}}(x)| \leq C_d P_m(t) \mathrm{e}^{3\delta_1 c_d} \mathrm{e}^{-\delta_1 r(t)}$ for $x \in B^k(\Lambda_8^{(k)})$. For the localised vacuum terms $E_{K,k}(\Lambda)$ of step k (largest line index k [3, (3.4)]), anchored at the cube $\square(v_0)$ of a fixed vertex, and $P_v(t) := \sum_{2 \leq \alpha + 2\beta \leq 2\bar{n}} e(t)^\alpha \lambda(t)^\beta$, there

is $C = O(\alpha_0, \bar{n})$ independent of ε, k and the domains with $\sum_{\Lambda:\square(v_0)\subset X}|E_{K,k}(\Lambda)| \le CP_v(t)|X|$ for every union X of unit cubes, and $\sum_{\Lambda \text{ bad}, \square(v_0)\subset B^k(\Lambda_8^{(k)})}|E_{K,k}(\Lambda)| \le CP_v(t)\mathrm{e}^{3\delta_1 c_d}\mathrm{e}^{-\delta_1 r(t)}|B^k(\Lambda_8^{(k)})|$.

This statement is used in the proof of Proposition E.76.

Statement K.26 (Hölder bound of the vector backgrounds)**.** Let Ω be a union of big blocks, $G_k := G_k(\Omega, 0)$ the vector propagator with Neumann conditions, $\tilde{A} := a_k G_k Q_k^* B$, where B is a unit-lattice field with $|B(y) - B(y')| \le c_1 p|y - y'|_1$, $c_1 = 3Ld$ [2, (2.17)]. Let $\alpha < 1$ and $x, x' \in \Omega$ with $\operatorname{dist}(\{x, x'\}, \Omega^c) \ge R_0 + 1$. Then $|x - x'|^{-\alpha}|\partial_\mu^\eta \tilde{A}(x') - \partial_\mu^\eta \tilde{A}(x)| \le C_{\mathrm{H}} p$ with $C_{\mathrm{H}} := 2c_0(\alpha)a_k c_1 \mathrm{e}^{\delta_0(\sqrt{d}+1)} \sum_{z\in\mathbb{Z}^d}(1 + |z|_1)\mathrm{e}^{-\delta_0|z|} + 2\sup_\Omega|\partial\tilde{A}|/p$, $c_0(\alpha), \delta_0$ the constants of [4, (1.9)]. The same holds for $B^{(k+1),\eta}$.

This statement is used in the proof of Proposition E.76.

Statement K.27 (nested and one-particle-reducible classes, with a pin)**.** (a) For every renormalised class of [3] whose members contain a divergent proper subgraph, the decomposition [3, (3.5)] holds, and hence [3, (3.2)] and [3, (1.33)]. (b) The one-particle-reducible members of a class of two-scalar-leg graphs of order 2 at fixed labels are $S_1^{(1)} G S_2^{(1)}$, with $S_i^{(1)}$ amputated order-one one-particle-irreducible two-point operators (including δm_1^2) and G the bridging scalar line; [3, (3.19)] applies to them with zero counterterm and leaves a term of degree α and a term of degree ≥ 1. (c) If every member carries leg 1 with $\Phi_{\mathrm{ext}} \equiv 1$ at a fixed pin p, the transformations of (a) and (b) never sum over the position of p.

This statement is used in the proof of Proposition E.76 (step (vi) of Lemma E.55).
The conditions of Bałaban's Proposition 1 at its applications.

Statement K.28 (the remaining abbreviated steps)**.** Assume the hypotheses of Theorem 5.2, and let the regulator data satisfy the conditions of Definition A.1. Let s be an estimate, identity or property with the following three features: it is asserted on [2, pp. 555–594], on [3, pp. 411–445], or on a page of [1] invoked on those pages; its argument is not printed there; and it is not one of the items (a)–(i) of Lemma E.114. Then s holds at every scale k and at every configuration at which it is asserted, with constants independent of ε, k and ℓ_T.

This statement is used in Remark E.111 (the completeness of the list in Proposition E.110 and Lemma E.114), and through the printed steps of [2, 3] in the proof of Theorem 5.2.

Statement K.29 (uniformity of Proposition 1 of [3] in a constant background)**.** Let $k \ge 1$. Let $\mathfrak{B}$ be a real background on T_η at which the hypotheses of [3, Prop. 1] hold in the form of Proposition E.110(i), and let $c \in \mathbb{R}^3$ be constant. Then these hypotheses hold at $\mathfrak{B} + c$ with the same constants. Moreover, consider any localised family of order at most $\bar{N}$ of the expansion of [3, p. 414] at the background $\mathfrak{B} + c$ (equivalently, at $\mathfrak{B}$ with the scalar boundary condition twisted by $\vartheta = ec\ell_T$) whose localisation cubes all lie in a box of side $< \ell_T$. Then [3, (1.33)] holds for this family with the same tree rate β_P and the same constant $O(1)$ as at $\mathfrak{B}$; in particular these do not depend on c.

This statement is used in the proof of Lemma E.130 (hypothesis (ZM) of Proposition E.140).
The mass reading.

Statement K.30 (uniform kernel and coefficient bounds on the torus)**.** Let $a = 1$, $L = 2$, $a_\flat = a(1 - L^{-2})$, $j_0 \ge 1$ and $\xi = L^{-j_0}$. Let Λ be the torus $\xi\mathbb{Z}^3/\ell\mathbb{Z}^3$ with $\ell \in L\mathbb{N}$ and $\ell \ge L$. Let d be the ℓ^1 torus distance, and put $\rho(w, w') := d(w, w') + \xi$.

Let P_1 be the orthogonal projection onto functions that are constant on the unit blocks. Put $F := (-\Delta^\xi + 1)^{-1}$. Let $\mathcal{K}$ be the set of operators $(-\Delta^\xi + \nu + bP_1)^{-1}$ with $\nu \in [0, 1]$ and $b \in [a_\flat, a]$. Kernels are taken with respect to $\xi^3 \sum$. A single difference is a forward or backward lattice

difference, in any direction. $D^{(1)}$ and $D^{(2)}$ denote a single difference acting in the first and in the second variable.

Then there are constants $\kappa \in (0, \frac{1}{16}]$ and $C < \infty$, depending only on a and L, such that the following hold for all $K, K_1, K_2, K_3 \in \mathcal{K}$, all single differences D, D', D_1, D_2, D_3, all $w, w', x, y, y' \in \Lambda$, and all ξ, j_0 and ℓ as above.

(a) $|K(w,w')| \le C\rho^{-1}\mathrm{e}^{-\kappa d}$ and $|(K-F)(w,w')| \le C\mathrm{e}^{-\kappa d}$. In either variable, $|DK(w,w')| \le C\rho^{-2}\mathrm{e}^{-\kappa d}$ and $|D(K-F)(w,w')| \le C\mathrm{e}^{-\kappa d}$. Moreover $|D^{(1)}D'^{(2)}(K-F)(x,y)| \le C\rho(x,y)^{-1}\mathrm{e}^{-\kappa d(x,y)/8}$.

(b) The following four bounds hold.
- (C1) $\xi|K(y,y)| \le C$.
- (C2) $\big|\xi^3 \sum_{y'}[(DK_1)(y,y')K_2(y,y') - (DF)(y,y')F(y,y')]\big| \le C$.
- (C3) If $|h(y')| \le \|h\|\, d(y,y')$ for all y', then $\big|\xi^3 \sum_{y'}(DK_1)(y,y')K_2(y,y')h(y')\big| \le C\|h\|$.
- (C4) Put $T[K_1,K_2,K_3] := \xi^6 \sum_{y',y''}(D_1K_1)(y',y)(D_2K_2)(y,y'')(D_3K_3)(y',y'')$. Then $|T[K_1,K_2,K_3] - T[F,F,F]| \le C$.

(c) (M1) Put $T^{\mathrm{c}}[K_1,K_2,K_3] := \xi^6 \sum_{y,y''}(D_1^{(1)}D_2^{(2)}K_1)(y',y)\,(D_3^{(2)}K_2)(y,y'')\,K_3(y,y'')$. Then $T^{\mathrm{c}}[F,F,F] = 0$ and $|T^{\mathrm{c}}[K_1,K_2,K_3]| \le C$.

This statement is used in the proof of Proposition E.158.

Statement K.31 (the vertex coefficient of the class (2.21c) and the convergent expressions of [3, §3])**.** Consider the graphs of orders $2 \le \alpha + 2\beta \le 4$ at zero background without operator lines, in the proof of [3, (1.33)] at a scale $j_0 \ge 1$. Use the notation of Lemma E.157, and let T^{c} be as in Statement K.30(c).

(a) Summed over orderings and indices, the vertex coefficient $\sum \xi^{2d}\Gamma'_\mu$ of [3, p. 444] for the class [3, (2.21c)] is a finite sum of expressions $c\,T^{\mathrm{c}}[G_s,G_s,G_p]$. Here D_1, D_2 are the differences at the two ends of the scalar line from the vertex of the external vector leg to the inner vertex, and D_3 is the difference on the inner scalar line at the vertex of the external scalar leg. Each coefficient satisfies $|c| \le 1$, and the number of terms depends only on the graph. Its C^ξ-version is the same sum with every $T^{\mathrm{c}}[G_s,G_s,G_p]$ replaced by $T^{\mathrm{c}}[F,F,F]$.

(b) Consider the expressions that [3] calls convergent after the replacements $G = C^\xi + GWC^\xi$ at pp. 437, 439, 441 and 444. Each of them is the difference between one of the following coefficients and its version with every propagator G_s, G_p replaced by C^ξ:
- the coefficient of (3.16);
- a coefficient of p. 441;
- the coefficient of (3.23);
- the coefficient $\sum \xi^{2d}\Gamma'_\mu$ of p. 444 for the classes (2.19), (2.20) and (2.21c).

These differences enter the constant $O(1)$ of [3, (1.33)] only through upper bounds on their absolute values.

This statement is used in the proof of Proposition E.158.
The small-field steps.

Statement K.32 (cumulant bound, upper half)**.** Let Λ be a finite subset of a lattice torus in $\mathbb{Z}^d$ and $\mu = N(0,P^{-1})$ on $(\mathbb{R}^n)^\Lambda$ with $P \in \mathcal{P}(\gamma_0,\gamma_1,c_0,\delta_0)$ (Lemma F.16). Let $V = \sum_{Y\subset J} V_Y$, $J \subset \Lambda$, each V_Y a real polynomial in $(z_x)_{x\in Y}$ of degree $\le D$ without constant term. Put $\|V\|_\kappa := \sup_x \sum_{Y\ni x} \mathrm{e}^{\kappa\ell(Y)}|V_Y|_1$, where $|V_Y|_1$ is the sum of the absolute values of the coefficients and $\ell(Y)$ the length of the shortest connected graph containing Y. Let $\chi_b := \prod_{x\in\Lambda} \mathbf{1}\{z_x \in bK\}$ with $K \subset \mathbb{R}^n$ symmetric convex, $B(1) \subset K \subset B(c_K)$. For $t \in \mathbb{N}$ put $\rho_1 := D + 2d$. There are $\rho_3 > 0$ and $S, b_* < \infty$, depending only on $(d,n,D,t,\kappa,\gamma_0,\gamma_1,c_0,\delta_0,c_K)$ and not on Λ, J or the side of the

torus, such that for $b \ge b_*$ and $\|V\|_\kappa b^{\rho_1} \le 1$

$$\int \chi_b \mathrm{e}^V \, d\mu \le \exp\Big\{ \sum_{m=1}^{t} \frac{\kappa_\mu(V;m)}{m!} + S|J|\big[(\|V\|_\kappa b^{\rho_1})^{t+1} + \mathrm{e}^{-\rho_3 b^{3/2}}\big]\Big\},$$

where $\kappa_\mu(V;m)$ is the m-th cumulant of V under μ (without χ_b).

This statement is used in the proofs of Lemma F.8 (C6) and Lemma F.14(b). (The hierarchical analogue is [6, §5].)

Statement K.33 (Dirichlet to free cumulants)**.** Let $P \in \mathcal{P}(\gamma_0, \gamma_1, c_0, \delta_0)$ on Λ_{big}, $\Lambda \subset \Lambda_{\mathrm{big}}$, and V as in Statement K.32 with $J \subset \Lambda$; put $r := \mathrm{dist}(J, \Lambda_{\mathrm{big}} \setminus \Lambda)$. There are $\mu_{\mathrm{c}} > 0$ depending only on the class constants and $C_m < \infty$ depending only on the class constants, d, n, m, D, such that for $m \ge 1$ $|\kappa_{N(0,(P_{\Lambda\Lambda})^{-1})}(V;m) - \kappa_{N(0,P^{-1})}(V;m)| \le C_m |J| \|V\|_\kappa^m \mathrm{e}^{-\mu_{\mathrm{c}} r/4}$.

This statement is used in the proof of Lemma F.14(c).

Statement K.34 (bounds on the background)**.** Let $0 \le k \le K-1$, $\Omega = B^k(\Lambda_0^{(k)})$ in scale-k units, $\tilde{B} = (1-\theta_{k+1})A^{(k)} + \theta_{k+1} B^{(k+1),\eta}$ on Ω, and assume (IH_k). There are $C_B, C < \infty$ depending only on $(d, L, M_b, a, c_1, \gamma_0, \delta_0)$ such that on Ω: $|\partial^\eta \tilde{B}|$, $|\partial^\eta(\theta_{k+1} A'^{(k)})|$, $|\partial^\eta(g_k A'^{(k)})| \le C_B p(t_k)$, and $|A^{(k)}|, |B^{(k+1),\eta}| \le C c_1 p(t_k)/(\mu_* t_k)$. These follow by assembling [2, (2.55), (2.59)–(2.60), (2.91), (2.96)–(2.97)].

This statement is used in Statement K.35, hence in the proof of Lemma F.8 (C3).

Statement K.35 (collars of $\mathcal{Q}^{(k)}$)**.** Let $L = 2$, $0 \le k \le K-1$, $t = t_k \le \varepsilon_0$ with ε_0 obeying (E.ksp1), (E.ksp2), assume (IH_k), and let $\vartheta(t) := |e|\, t^{1/2} p(t)$. For scale-$j$ ($j < k$) fields A with $\mathsf{M}(A) := aL^{-2} Q(A)^* Q(A) + \Delta^{(j)}(A)$ assume: (Q0) $\nabla_A^* \nabla_A + a_j Q_j(A)^* Q_j(A) \ge \gamma_0$ for regular real A [4, (1.8)]; (Q1) $\gamma_0' \le \mathsf{M}(A) \le a_j + aL^{-2}$ [4, (1.15)]; (Q2) $|\mathsf{M}(A)(y,y')| \le c_M \mathrm{e}^{-\delta|y-y'|}$. Write $\mathcal{Q}^{(k)}(\Omega; \tilde{B}, a) = \sum_{j<k} \sum_y q_{j,y}(a)$ for the site decomposition of the polynomial of [2, (2.105)–(2.106), p. 579], in which for each site y, with box Q_y of half-side R_y, the axial-gauge extension $\lambda_y \circ \pi$ off Q_y is replaced by $\chi\lambda_y$ as in (a). Then: (a) Let $Q_y \subset \Omega$ not wrap, Q_y' the concentric box of half-side $R_y/2$, $\lambda_y(x) := -\sum_{b \in \Gamma^{\mathrm{ax}}_{y\to x}} \eta \mathcal{A}(b)$ along the axis-ordered path in Q_y, $\chi : \Omega \to [0,1]$ with $\chi = 1$ on Q_y', $\chi = 0$ off Q_y, $|\chi(x) - \chi(x')| \le 2\eta/R_y$ on bonds, and $\mathcal{A}^g := \mathcal{A} + d(\chi\lambda_y)$. If $\sup|\mathcal{A}| \le P_1$, $\sup_{Q_y}|\mathcal{F}| \le E$ and $\sqrt{d} R_y E \le P_1$, then $|\mathcal{A}^g(b)| \le \|b_- - y\|_1 E$ for $b \subset Q_y'$ and $|\mathcal{A}^g(b)| \le (2d + \sqrt{d}) P_1 \le (2d+3) P_1$ otherwise. (b) There are $\delta_1, C_Q > 0$ depending only on $(d, L, a, \gamma_0, \gamma_0', c_M, \delta, \bar{n})$ such that, with $c_{AX} = \sqrt{d}/(\mathrm{e}\delta_1)$, $E_j = 2C_B \vartheta(t) L^{-2(k-j)}$ (C_B of Statement K.34), $T_{j,y} = (2d+3)\mathrm{e}^{-\delta_1 R_y/2} P_1(t)$ and $\tau := c_{AX} E_j + T_{j,y} \le 1$, one has $|q_{j,y}(a)| \le C_Q \tau^2$; and if $\mathrm{dist}_k(Y, \Omega^c) \ge r(t)/2$, then $\sum_{j<k} \sum_{y \in B^{k-j}(Y)} |q_{j,y}(a)| \le [8 C_Q c_{AX}^2 C_B^2 \vartheta(t)^2/(L-1) + C_N t^N]|Y|$ for every N. (c) If $\Omega^c \ne \emptyset$ (or R_y is capped) and $R_y \le r(t) + M_b < \ell_T/t$, then uniformly in k

$$|\mathcal{Q}^{(k)}(\Omega; \tilde{B}, \theta_{k+1} A') - \mathcal{Q}^{(k)}(\Omega; \tilde{B}, g_k A')| \le \Big[\frac{C_{\mathrm{coll}} e^2 b_0^2\, t\, \ell(t)^{2p} (r(t) + M_b)^3}{L-1} + C_N t^N\Big] |\Lambda_2^{(k)} \cap \Lambda_7^{(k)c}|,$$

with C_{coll} depending only on $(d, L, a, M_b, c_1, \gamma_0, \gamma_0', c_M, \delta, \delta_0, \bar{n})$.

This statement is used in the proof of Lemma F.8 (C3), Step 3.

Statement K.36 (smallness of the scalar fluctuation; [2, Lemma 2.7])**.** Work in scale-k units, $e_t = |e| t^{1/2}$, $\lambda(t) = \lambda t$, $p = p(t)$, $r = r(t)$. Put $\Omega = B^k(\Lambda_2^{(k)})$, $\hat{B} = B^{(k+1),\eta}$, $Q_k = Q_k(\hat{B})$, $Q = Q(\hat{B})$, $G_k = (-\Delta_{\hat{B},\Omega} + m^2 t^2 + a_k Q_k^* Q_k)^{-1}$ (Neumann on $\partial\Omega$), $\Delta = a_k - a_k^2 Q_k G_k Q_k^*$, $\mathsf{M} = \Delta + aL^{-2} Q^* Q$, $C = (\Lambda_4^{(k)} \mathsf{M} \Lambda_4^{(k)})^{-1}$, with $\mathrm{dist}(\Lambda_5^{(k)}, \Lambda_4^{(k)c}) > r$, $\mathrm{dist}(\Lambda_4^{(k)}, \Lambda_2^{(k)c}) > 2r$. For a field ψ on the scale-$(k+1)$ blocks $y \subset \Lambda_2^{(k)}$ and a fine bond b let $\gamma_b = \Gamma_{b_-} + b - \Gamma_{b_+}$, Γ_z the composite contour from the

base point of $y(z) \ni z$ to z; $\tilde{c}(\delta) = \sum_{z\in\mathbb{Z}^d} \mathrm{e}^{-\delta|z|}$, $S(\delta) = \sup_{x'} \sum_y \mathrm{e}^{-\delta \operatorname{dist}(B(y),B(x'))}$. Assume: (F1) $\hat{B}$ real and regular on Ω [1, Prop. 2.1], $|\partial^\eta \hat{B}| \le c_B p$; (F2) at distance $\ge R_0$ from Ω^c: $|(G_k f)(z)| \le c_0 \mathrm{e}^{-\delta_0 \operatorname{dist}(z,\operatorname{supp} f)} \|f\|_\infty$, $|\eta^{-1} d_b(G_k Q_k^* \delta_y)| \le c_0 \mathrm{e}^{-\delta_0 \operatorname{dist}(b,B(y))}$, and $|C(x,x')| \le c_0 \mathrm{e}^{-\delta_0|x-x'|}$ on $\Lambda_4^{(k)}$; (F3) $|\psi| \le 2\lambda(t)^{-1/4} p$ on $\Lambda_2^{(k)}$, $|\psi(y(b_+)) - U(\hat{B}(\gamma_b))\psi(y(b_-))| \le c_\psi p$ if $y(b_-) \ne y(b_+)$, and $|\varphi(x) - (Q^*\psi)(x)| \le c_\varphi p$ on $\Lambda_5^{(k)}$; (F4) if $y(b_-) = y(b_+)$, γ_b bounds a surface of area $\le c_\Gamma L^2$ ($\le c_\Gamma \eta$ inside one unit block); (F5) $r \ge 2R_0 + 2$. Then for $x \in \Lambda_5^{(k)}$, $aL^{-2}(CQ^*\psi)(x) = (Q^*\psi)(x) + R(x)$ with $|R(x)| \le (K_0 + \omega(t))p$, $K_0 = 2d\,\mathrm{e}^{\delta_0} a_k c_0^2 \tilde{c}(\delta_0) S(\delta_0) c_\psi$, $\omega(t) = C_1 \lambda(t)^{-1/4} [\mathrm{e}^{-\delta_0(r-1-\sqrt{d})/2} + m^2 t^2 + e_t p]$, $C_1 = C_1(d, a, L, c_0, \delta_0, c_B, c_\Gamma)$. If $\omega(t) \le 1$, then $\varphi' = \varphi - aL^{-2} CQ^*\psi$ satisfies $|\varphi'| \le (c_\varphi + K_0 + 1)p$ on $\Lambda_5^{(k)}$.

This statement is used in the proof of Lemma F.8 (C5).

Statement K.37 (inputs of the expansion of the normalisations)**.** Let $0 \le j < k \le K-1$, $m = k-j$, and assume (IH$_k$). In scale-j units: (i) the e_j-weighted density $\mathcal{A}$ of $\theta_{k+1}A'$ on $B^j(\Lambda_2^{(j)})$ satisfies $\sup|\mathcal{A}| \le c_A p(t_j)$, c_A independent of $j, k, \varepsilon, \ell_T$; (ii) the contours of the transporters in $Q_j(A)$, $Q(A)$ and in the operator $W^{(j)}$ of [2, (2.99)] have lengths bounded by a constant ℓ independent of $j, k, \varepsilon, \ell_T$; (iii) with U the union of the blocks $B(x)$, $x \in \Lambda_5^{(j)}$, intersected with $B^j(\Lambda_2^{(j)}) \cap B^k(\Lambda_0^{(k)})$, and S_1 the scale-j support of θ_{k+1}, $\operatorname{dist}(S_1, U^c) \ge L^m r(t_k) - c$ with c independent of $j, k, \varepsilon, \ell_T$.

This statement is used in Statement K.38, hence in the proof of Lemma F.8 (C3).

Statement K.38 (expansion of the normalisations $Z^{(j)}$)**.** In the setting of Statement K.37, let $W^{(j)}(\tau)$ be the operator of [2, (2.99)] at $\tau\theta_{k+1}A' + \tilde{B}$, $C = C^{(j)}_{\Lambda_5^{(j)}}(B^j(\Lambda_2^{(j)}), \tilde{B})$, X_j the scale-j points of $B^k(\Lambda_2^{(k)})$ ($|X_j| = L^{dm}|\Lambda_2^{(k)}|$). Then: (a) $W^{(j)}(\tau) = \sum_{q=1}^{\bar{n}} \tau^q W_q + W_>(\tau)$ on $[0,1]$ with real symmetric W_q, $\|W_q\| \le w_j^q$, $\sum_{x,y}\|W_q(x,y)\| \le w_j^q |X_j|$, the same for $W_>(\tau)$ with $(w_j\tau)^{\bar{n}+1}$, where $w_j \le C_W e(t_j) p(t_j)$, $e(t) = |e|t^{1/2}$. (b) With $c_{m'}^{(j)} := \sum_{l=1}^{m'} \frac{1}{2l} \sum_{q_1+\cdots+q_l=m'} \operatorname{tr}(CW_{q_1}\cdots CW_{q_l})$, if $4w_j \le \gamma_0 \le 1$, then $|\log(Z^{(j)}(\theta_{k+1}A' + \tilde{B})/Z^{(j)}(\tilde{B})) - \sum_{m'\le\bar{n}} c_{m'}^{(j)}| \le C_Z n w_j^{\bar{n}+1} |X_j|$, $C_Z = (2/\gamma_0)^{\bar{n}+1} + \sum_{l\le\bar{n}} (\bar{n}+1)^l/(2l\gamma_0^l)$. (c) If $c_{m'}^{\prime(j)}$ is $c_{m'}^{(j)}$ with the domain change of [2, p. 578] applied to all propagators, $|c_{m'}^{(j)} - c_{m'}^{\prime(j)}| \le nK(\bar{n})(C'_* w_j)^{m'} \mathrm{e}^{-\delta_5\rho}|S_1|$ with $\rho \ge L^m r(t_k) - c$. (d) If $\frac{\bar{n}+1}{2} > d$ and $\delta_5 r(t_k) \ge d\log L + \log 2$, then $\sum_{j<k} |\log(Z^{(j)}(\theta_{k+1}A' + \tilde{B})/Z^{(j)}(\tilde{B})) - \sum_{m'\le\bar{n}} c_{m'}^{(j)}| \le C t_k^{(\bar{n}+1)/2} \ell(t_k)^{p(\bar{n}+1)} |\Lambda_2^{(k)}|$, and the re-localisation errors sum to $\le C_N t_k^N |\Lambda_2^{(k)}|$ for every N. All constants depend only on $(d, L, a, \gamma_0, \ell, c_A, \bar{n})$ and the kernel constants of Statement K.39.

This statement is used in the proof of Lemma F.8 (C3), Step 3.

Statement K.39 (scalar kernels on the torus)**.** Let $\Omega \subset T_h$ be a union of big blocks and A a real field on Ω, regular in the sense of [4, (1.7)], and let (N0) be the bound $-\Delta_{A,\Omega}^N + aP_k(A) \ge \gamma_0$ of [4, (1.8), p. 573]. Distances are the ℓ^∞ torus distances of [1, (1.3)]. Then: (i) [4, (1.9)–(1.12), (1.15)–(1.20)] hold on Ω with constants independent of A (in particular of its constant part), Ω, k, h and the torus side; (ii) $|\Delta^{(k)}(\Omega, A; y, y') - a_k \delta_{yy'}| \le c'_B \mathrm{e}^{-\delta|y-y'|}$, $c'_B = 2a_k^2 \gamma^{-1} \mathrm{e}^{2\sqrt{d}\delta}$; (iii) if (Ω_i, A_i), $i = 1, 2$, agree on every bond with an endpoint in a union of blocks $S \subset \Omega_1 \cap \Omega_2$ and $D(y) := \operatorname{dist}(y, (\Omega_1 \cup \Omega_2) \setminus S) \ge 3 + \sqrt{d}$, $D(y') \ge 3 + \sqrt{d}$, then $|\Delta^{(k)}(\Omega_1, A_1; y, y') - \Delta^{(k)}(\Omega_2, A_2; y, y')| \le a_k^2 C \mathrm{e}^{-\frac{\delta}{2}(|y-y'| + D(y) + D(y'))}$; (iv) with H_k defined by [2, (2.108)], the form $\langle \Lambda_3^{(k)}\varphi, \Delta_{\mathrm{in}} \Lambda_3^{(k)}\varphi\rangle$ there carrying the factor $\frac{1}{2}$, and $r(t) \ge 4 + 3\sqrt{d} M_b$: $|h_k(y,y')| \le C_H \mathrm{e}^{-\frac{\delta}{2}r(t)} \mathrm{e}^{-\frac{\delta}{2}|y-y'|}$; (v) if $|\varphi| \le \bar{\varphi}$ on $\Lambda_6^{(k-1)\prime}$, $|\langle \varphi, H_k \varphi\rangle| \le C_H \mathrm{e}^{-\frac{\delta}{2}r(t)} S_\delta \bar{\varphi}^2 |\Lambda_6^{(k-1)\prime}|$, $S_\delta = \sum_z \mathrm{e}^{-\delta|z|/2}$. Here γ, δ, C, C_H depend only on (d, a, γ_0).

This statement is used in the proof of Lemma F.8 (C2), (C6).

Statement K.40 ([2, Prop. 3.1] at a real background)**.** Let $S_k := \Lambda_5^{(k-1)\prime} \cap \Lambda_5^{(k)c}$ ($S_0 := \Lambda_5^{(0)c}$), $\Phi = (\varphi_k|_{S_k})_k$, and define $\langle \Phi, \Delta(\tilde{A})\Phi\rangle$ by [2, (3.23), (3.25)]. Assume $|\partial \tilde{A}| \le c_A p(t_k)$ on each $B^k(S_k)$ (scale-k units), $|\varphi_k(x)| \le c_1 \lambda(t_{k-1})^{-1/4} p(t_{k-1})$ on S_k (scale-$(k-1)$ units), $m^2\varepsilon_0^2 \le 1$, $\varepsilon_0 \le \min(1, \mathrm{e}^{-4})$, $e\varepsilon_0^{1/2} \le e_*(\frac{1}{10})$ and (T.nw). Then, with $\gamma_0' = \min(1, \gamma_0)$, γ_0 of [4, (1.22)], and $\kappa_0 = \frac{1}{5}$,

$$\langle \Phi, \Delta(\tilde{A})\Phi\rangle \ge \gamma_0' \sum_{k=0}^{K} \Big[\sum_{\langle x,x'\rangle \subset S_k} t_k^{d-2} |U(\tilde{A}(\langle x,x'\rangle))\varphi_k(x') - \varphi_k(x)|^2 + \sum_{x\in S_k} t_k^d m^2 |\varphi_k(x)|^2 \Big] - \sum_{k=1}^{K} C(\kappa_0) t_k^{\kappa_0} |S_k|.$$

This statement is used in the proof of Lemma F.8 (C9).

Statement K.41 (restriction constants on the slice line)**.** Let $w_j \in [w_-, w_+]$ and $\sigma := \max(w_+, w_-^{-1})^{1/2}$. On the slice line, the vector characteristic functions of [2, (2.44)] allow exactly the configurations obeying the vector restrictions of [2, (2.55)] with c_1 replaced by σc_1.

This statement is used in Statement K.42(iv), hence in the proof of Lemma F.8, Step 0.

Statement K.42 (the background as an external field)**.** Put $s_k := (1-\theta_{k+1})\theta_k$ and $\tilde{B} := s_k A^{(k)} + \theta_{k+1} B^{(k+1),\eta}$ [2, p. 582]. (i) After the translation [2, (2.80)], the vector variable enters a scalar object (the $Z^{(j)}$, the averaging kernel, the basic scalar form, $\mathcal{P}^{(k)}$, the scalar characteristic functions) only through $\theta_k A^{(k)} = s_k A^{(k)} + \theta_{k+1} A'^{(k)} + \mathcal{F}(B)$, where $s_k A^{(k)}$ depends on A' only through $A'|_{\Lambda_3^{(k)c}}$ and $\mathcal{F}(B) = \theta_{k+1}(aL^{-2} C^{(k)}_{\Lambda_0^{(k)}} Q^* B)^{(k)}$; every upper bound proved for the scalar integral at a fixed real external field of the regular class applies at $\theta_k A^{(k)}$. (ii) $\tilde{B} = B^{(k+1),\eta}$ on $B^k(\Lambda_2^{(k)})$, so $\Delta^{(k)}$, G_k, $C^{(k)}_{\Lambda_5}$ on that domain and $Q(\tilde{B})$ on blocks inside it agree at $\tilde{B}$ and at $B^{(k+1),\eta}$. (iii) If $L = 2$, $r(\varepsilon_0) \ge 4\sqrt{d} M_b$ and $2R \log 2\, \ell(\varepsilon_0) \ge 2\sqrt{d} M_b$, then the split [2, (2.89)] holds as printed and [2, Lemma 2.3, (2.93)–(2.98)] hold from the restrictions in force; in particular $|\partial \tilde{B}| \le O(1) p(t)$ on $B^k(\Lambda_2^{(k-1)\prime})$. (iv) Under Statement K.41, (i)–(iii) and Statements K.39, K.40 hold on the slice line with c_1 replaced by σc_1; for $w_j = 1 + u\delta_j$, $|u| \le 1$, $0 \le \delta_j \le \frac{1}{12}$, $\sigma \le (12/11)^{1/2} < \sqrt{2}$.

This statement is used in the proof of Lemma F.8, Steps 0 and 4.

Statement K.43 (size of the counterterm)**.** For every $M > 0$ there is $C_\delta(M) < \infty$, independent of $\varepsilon \le \varepsilon_{\max}$, of $\ell \ge \ell_0$ and of the position, and non-increasing in ℓ_0, such that $\varepsilon |\delta m^2_{K,X}(M)| \le C_\delta(M)$ for $X \in \{\varepsilon\mathbb{Z}^3, T\}$.

This statement is used in the proof of Lemma F.8 (C1), case $k = 0$.

Statement K.44 (restriction tails)**.** In the setting of Statement K.32, at step k let μ_k be the joint Gaussian of Lemma F.8(C6), $\chi_k^{\mathrm{fl}} = \chi(A')\chi(\varphi')$ with radius b_k, $c_-\ell(t_k) \le b_k \le c_+\ell(t_k)^q$, $V = V_w^{(k)}$, $\mathsf{A} := \|V\|_\kappa$, and N_k the number of scale-k blocks of T_ε (so $|J|_k \le N_k$). Assume $c_-\ell(\varepsilon_0) \ge b_*$ and $\mathsf{A} b_k^{\rho_1} \le 1$. Then (a) $\int d\mu_k \chi_k^{\mathrm{fl}} \mathrm{e}^V \le \exp(\sum_{m\le\bar{n}} \kappa_{\mu_k}(V;m)/m! + \mathcal{R}_k^{\mathrm{tr}} + \mathcal{R}_k^{\mathrm{res}})$ with $0 \le \mathcal{R}_k^{\mathrm{tr}} \le S|J|_k (\mathsf{A} b_k^{\rho_1})^{\bar{n}+1}$, $0 \le \mathcal{R}_k^{\mathrm{res}} \le S|J|_k \mathrm{e}^{-\rho_3 b_k^{3/2}}$; (b) $\mathcal{R}_k^{\mathrm{res}} \le C_{\mathrm{res}}(\kappa) t_k^\kappa N_k$ for every $\kappa > 0$, $C_{\mathrm{res}}(\kappa) = S\exp(4\kappa^3/(27\rho_3^2 c_-^3) - \kappa)$; (c) if $\mathsf{A} \le C_A t_k^{\kappa_*}$ with $(\bar{n}+1)\kappa_* > \kappa_B$, then $\mathcal{R}_k^{\mathrm{tr}} \le C_{\mathrm{tr}} t_k^{\kappa_B} N_k$; (d) the field-independent part of the exponent is the step-k vacuum output of Lemma F.31, and the restriction indicators contribute to it exactly $\mathcal{R}_k^{\mathrm{tr}} + \mathcal{R}_k^{\mathrm{res}}$; (e) $\sum_{k<K}(\mathcal{R}_k^{\mathrm{tr}} + \mathcal{R}_k^{\mathrm{res}}) \le (C_{\mathrm{res}}(\kappa_B) + C_{\mathrm{tr}})(1 - 2^{-(\kappa_B-3)})^{-1} \varepsilon_0^{\kappa_B - 3} |T|$, uniformly in ε, K, ℓ_T and u.

This statement is used in the proofs of Lemma F.8 (C6), (C8) and Lemma F.14(b).

Statement K.45 (the vector operations of a step)**.** Every vector line occurring in the step-k construction of [2, Sect. 2C] and [3] is obtained from the step covariances $C^{(i)}_{\Lambda_5^{(i)}}$, $i \le k$, by finitely

many of the following operations: push-forward by the w-free linear maps of the translations, conditional means and lifts; composition [1, (2.42)]; restriction to domains; Dirichlet-to-free replacement; localisation cut-offs; differences.

This statement is used in Lemma F.17 and in the proof of Lemma F.14(c).
The large-field step and the flow of the mass and the vacuum.

Statement K.46 (composed kernels are a Schur complement)**.** For a positive definite quadratic form q on $\mathbb{R}^V \oplus \mathbb{R}^W$, put $\mathrm{Sch}_V\, q(v) := \min_w q(v,w)$. Put $Q_k := Q^k$, the k-fold block average, and $a_k := a(1-L^{-2})/(1-L^{-2k})$. Let $S_0 \subset T^{(0)}$ and $S_k \subset T^{(k)}$, $1 \le k \le K$, satisfy the shadow condition

$$\text{(D)} \qquad S_k \cap B^{k'-k}(S_{k'}) = \emptyset \qquad (1 \le k < k' \le K),$$

and put $\Psi := (B_0|_{S_0}, B_k|_{S_k})_{1\le k\le K}$. Let S^1 be the quadratic form of the photon chain of Lemma F.5 with all weights equal to 1. Then $\mathrm{Sch}_\Psi\, S^1 = \Delta_v$, where

$$\langle \Psi,\, \Delta_v \Psi\rangle = \min_{B_0|_{T^{(0)}\setminus S_0}} \Big[\langle B_0,\, (-\Delta+\mu_0^2)B_0\rangle + \sum_{k=1}^{K} a_k t_k^{d-2} \sum_{x\in S_k} |B_k(x) - (Q_k B_0)(x)|^2 \Big].$$

Equivalently, integrating $\mathrm{e}^{-\frac12 S^1}$ over all variables other than Ψ gives a constant times $\mathrm{e}^{-\frac12\langle \Psi,\, \Delta_v\Psi\rangle}$. This is the vector form of [2, (3.23), p. 588].

This statement is used in the proof of Lemma F.22 (Corollary F.28).

Statement K.47 (the shells satisfy the shadow condition)**.** For every admissible region sequence $\mathbf{\Lambda}$, the sets $S_0 = \Lambda_5^{(0)c}$ and $S_k = \Lambda_5^{(k-1)\prime} \cap \Lambda_5^{(k)c}$, $1 \le k \le K$, satisfy condition (D) of Statement K.46.

This statement is used in the proof of Lemma F.22 (Corollary F.28).

Statement K.48 (pinned mass coefficients along the flow)**.** Let $L = 2$, $\varepsilon = 2^{-K}\varepsilon_0$, $t_j = 2^j\varepsilon$, $e(t) = et^{1/2}$, $\lambda(t) = \lambda t$, and let P_K, P_m be as in Lemma F.32. For a counterterm graph G_0 of [3, (1.23)] of order $e^\alpha\lambda^\beta$ with $2 \le \alpha + 2\beta \le 4$, put $o := \alpha/2 + \beta$; lower-order δm^2 insertions are kept as the partners of their self-energy subgraphs. Then:

(a) for all $x \in T_\varepsilon$ and $0 \le k \le K$, $|\delta m^2_{K,k}(x)| t_k^2 \le C_m\big[2(e^2+\lambda)t_k + (e^4+e^2\lambda+\lambda^2)t_k^2 \log_2(\varepsilon_0/t_k)\big] \le C_m P_K(t_k)$;

(b) work in scale-k units, put $t = t_k$, $W := B^k(\Lambda_7^{(k)})$, and let $\mathcal{L}_k$, $\mathcal{D}_k$ be as in Lemma F.32. Let $\Delta_k(x) := [\delta m^2_{K,k}(x) - \delta m^2_{K,k+1}(x)]t^2$, the sum over G_0 of the scale-k coefficients with largest index k, leg 1 fixed at x, leg 2 summed, external fields $\equiv 1$ and background 0. Localise it as in [3, p. 420], $\Delta_k(x) = \sum_\Lambda \Delta_k(x;\Lambda)$, with $\square(p) \ni x$ for the vertex p of leg 1. Call Λ bad if some cube of Λ is not contained in W, and let $\Delta_k^{\mathrm{bad}}(x)$ be the sum over the bad Λ. Let the localisation cubes be cubes of side 1 or 2 with corners in the unit lattice, and let c_d be twice their largest diameter. Then for all x

$$|\Delta_k(x)|,\ |\Delta_k^{\mathrm{bad}}(x)| \le C_d P_m(t), \qquad |\Delta_k^{\mathrm{bad}}(x)| \le C_d P_m(t)\, \mathrm{e}^{\delta_0 c_d}\mathrm{e}^{-\delta_0 r(t)/2} \quad (x \in \mathcal{D}_k).$$

If $|\varphi(x)| \le c_1 p(t)\lambda(t)^{-1/4}$ on W, then

$$\Big|\sum_{x\in W} \eta^d \Delta_k^{\mathrm{bad}}(x)|\varphi(x)|^2\Big| \le c_1^2 C_d P_m(t)\lambda(t)^{-1/2} p(t)^2\big[|\mathcal{L}_k| + \mathrm{e}^{\delta_0 c_d}\mathrm{e}^{-\delta_0 r(t)/2}|\mathcal{D}_k|\big].$$

Here $C_m = C_P N_{\mathrm{ct}}$ and $C_d = C_P N_{\mathrm{ct}} S_0$. In these formulas C_P is the constant of the pinned bound of Lemmas E.51, E.52 and E.55, N_{ct} is the number of counterterm graphs of order ≤ 4, $S_0 := \sum_{D\ge 0}(c_d'(D+2)^d)^{N_s}\mathrm{e}^{-\delta_0 D/2}$, N_s is the maximal number of localised vertices of a graph of order $\le \bar n$, and c_d' depends only on d and c_d. The constants are independent of ε, k, x, the domains $\Lambda_i^{(k)}$ and ℓ_T.

This statement is used in the proof of Lemma F.32.

Statement K.49 (size of the bare mass counterterm)**.** Let $X \in \{\varepsilon\mathbb{Z}^3, T_\varepsilon\}$ with $\ell_T \geq \ell_0$, and let $\int$ denote the Brillouin-zone integral, resp. $\ell_T^{-3}\sum_{p\in T^*}$. Put $\hat p_\mu = s_\mu(p) := \frac{2}{\varepsilon}\sin\frac{\varepsilon p_\mu}{2}$, $\hat p^2 = \sum_\mu \hat p_\mu^2$, $\hat G = (\hat p^2 + M^2)^{-1}$, $\hat D = (\hat p^2 + \mu_*^2)^{-1}$, $D_X = \int \hat D$, $G_X(0) = \int \hat G$, $X_X(0) = \int \hat D\hat G$, $\delta_X = \int \hat D^2$, $\tau_X = \int \hat G^2$ and $J_X = M^2\int \hat D\hat G$. Let $G_X(z)$, $D_X(z)$ be the position-space kernels. Assume: (H) for $\varepsilon \leq \varepsilon_{\max}$, $\varepsilon D_X \leq \bar\omega$, $\varepsilon G_X(0) \leq \bar\omega_M$, $X_X(0) \leq \bar\psi(M)$, $\delta_X \leq \bar\delta$ and $\tau_X \leq \bar\tau(M)$, with constants independent of ε and $\ell_T \geq \ell_0$; (K) the coefficients of King's counterterm [24] through order 4 are $\delta m^2_\lambda = -16\lambda G_X(0)$, $\delta m^2_{e^2} = -2e^2 D_X - e^2 J_X$, $\delta m^2_{\lambda^2} = 128\lambda^2\sum_z \varepsilon^3 G_X(z)^3$, $\delta m^2_{e^2\lambda} = 16\lambda\int \hat G^2\tilde\Pi$, and $\delta m^2_{e^4} = \frac34\varepsilon^2 e^4 D_X^2 + (A) + (C) + (D1) + (D2) + (D3)$. Here $\tilde\Pi(p) = \Sigma(p) - \Sigma(0)$ with $\Sigma(p)/e^2 = D_X\sum_\mu\cos(\varepsilon p_\mu) - \int_{p'}\hat D(p-p')\widehat{(p+p')}^2\hat G(p')$, and $(A) = e^2M^2\int\hat D\hat G^2\tilde\Pi$, $(C) = -\frac{\varepsilon^2e^4D_X}{2}(D_X - J_X)$, $(D1) = \frac{e^4}{2}\sum_\mu\sum_z\varepsilon^3[D_X(z)^2 + D_X(z+\varepsilon e_\mu)^2][G_X(z)+G_X(z+\varepsilon e_\mu)]$, $(D2) = -e^4\sum_\mu\int_k s_\mu(k)\hat G(k)\hat D(k)S_\mu(k)$ with $S_\mu(k) = \int_P[s_\mu(k)+s_\mu(2P+k)]\hat G(P)\hat D(P-k)$, $(D3) = e^2\int\hat D^2\sum_\mu\Pi_{\mu\mu}$ with $\Pi_{\mu\mu}(k)/e^2 = 2G_X(\varepsilon e_\mu) - \int_q\widehat{(2q+k)}_\mu^2\hat G(q)\hat G(q+k)$. Let $\delta m^2_{K,X}(M)$ be the sum of these five coefficients. Then $\varepsilon|\delta m^2_{K,X}(M)| \leq C_\delta(M)$ for every dyadic $\varepsilon \leq \varepsilon_{\max}$, where

$$C_\delta(M) = (16\lambda + 3e^2)\max(\bar\omega,\bar\omega_M) + 128\lambda^2\bar\omega_M\bar\tau(M) + 16\lambda e^2[10\bar\omega\bar\tau(M) + 2\bar\psi(M)\bar\omega_M] + e^4C_4(M),$$
$$C_4(M) = \tfrac54\varepsilon_{\max}\bar\omega^2 + 16\bar\omega_M\bar\delta + 14\bar\psi(M)\bar\omega + 4\bar\delta\bar\omega.$$

Consequently $\lambda(\varepsilon)s^2 + \frac12\delta m^2_{K,X}(M)\varepsilon^2 s \geq -C_\delta(M)^2\varepsilon/(16\lambda)$ for all $s \geq 0$.

This statement is used in the proof of Lemma F.32 (and, through $C_\delta(M)$, in Lemma F.22(b)).

Statement K.50 (families with D-lines)**.** Work in organisation (α) at scale n ($\eta = L^{-n}$), and put $c_n := \kappa\mu_*^2L^2t_n^2$. A D-line is $uc_nD_n^\eta$ with $D_n^\eta = \sum_{j<n}(L^j\eta)^2G^\eta_{(j)}$.

(a) Every connected coloured graph with an external leg and $m \geq 1$ D-lines has superficial degree $D^\natural \geq 1$.

(b) Every divergent proper subgraph of a coloured graph is C-only. Hence a family with an external leg and $m \geq 1$ is convergent. Its own divergent subgraphs are renormalised exactly as at $u = 0$, by $\delta m^2_{K,T}$ and by the Ward–Takahashi and $\varphi\varphi A$ partners, and the family itself receives no counterterm.

(c) Let $\square_0$ be a scale-n unit block, and let $\mathcal{M}_n$ be the sum of all families with two scalar external legs and $m \geq 1$ over the localisations whose first cube is $\square_0$. Then $|\mathcal{M}_n| \leq C_{U1}e^2|u|\kappa\mu_*^2L^2t_n^3\sup\|h\Phi_{\rm ext}\|_{1,\alpha_0}$, where

$$C_{U1} = O(1)\,N_F\,N_\ell\,S_0\,(1+\bar c)^{N_\ell-1}\sum_{(\alpha,\beta)}|e|^{\alpha-2}\lambda^\beta\varepsilon_0^{o-1}.$$

The sum runs over $\alpha \geq 2$ and $2 \leq \alpha + 2\beta \leq 2\bar n$. Here $O(1)$ is the constant of Theorem H.24(c), N_F is the number of connected uncoloured two-scalar-leg graphs of order $\leq \bar n$, N_ℓ is their maximal number of vector lines, $\bar c := |u|\kappa\mu_*^2L^2\varepsilon_0^2$, and S_0 is as in Statement K.48. The same bound holds for the share of $\mathcal{M}_n$ anchored in any sub-block of $\square_0$.

(d) Assume (W) of Statement K.51. Decompose any vector line $\mathcal{L}$ used at scale n by step of origin, $\mathcal{L} = \sum_i\mathcal{L}_i$; the line may be a composed propagator, a domain-restricted step covariance, a Dirichlet-to-free difference, a localised piece, a cut line, or a difference of such objects. Then the u-model line is $\sum_i w_i\mathcal{L}_i$ and its D-part is $u\sum_i\delta_i\mathcal{L}_i$. Every per-index bound used for $\mathcal{L}_i$ holds for the D-part with the extra factor $|u|\delta_i \leq |u|c_n$. Consequently, take a localised family of order $\leq \bar n$ with at most N_ℓ vector lines. Its u-model value is its $u = 0$ value plus a sum of coloured families with $m \geq 1$. That sum is bounded by the right side of [3, (1.33)], with the constant of Theorem H.24(c), times $(1+|u|c_n)^{N_\ell} - 1 \leq N_\ell|u|c_n(1+|u|c_n)^{N_\ell-1}$.

This statement is used in the proof of Lemma F.32.

Statement K.51 (the weighted-field structure of the slice model)**.** (W) Let $u \in [-\eta, 1]$ and let n be a scale. In the slice model at (u, M), for every step $i < n$ the step-i vector fluctuation has covariance $w_i C^{(i)}$. Here $C^{(i)}$ is the step-i covariance at $u = 0$, $w_i = 1 + u\delta_i$, and $0 \le \delta_i \le \kappa\mu_*^2 L^2 t_n^2$. The step-$i$ fluctuation field enters all later expressions (composed propagators, domain restrictions, Dirichlet-to-free replacements, localisation cut-offs, cut lines and their differences) only through linear maps that do not depend on $(w_i)_i$. The per-index kernel bounds [3, (2.6), (2.10)–(2.12), pp. 424–426] for the index-i parts hold with the constants of the case $u = 0$.

This statement is used in the proof of Lemma F.32 (through Statement K.50(d)).

Statement K.52 (weighted external field)**.** At zero background, for two-variable lattice fields g put $\|g\|_{1,\alpha_0} := \sup|g| + \sup|\partial g| + \sup_{y\neq y'} |\partial g(y) - \partial g(y')|/|y - y'|^{\alpha_0}$. Here ∂ ranges over the lattice differences in either variable, in the units of the scale at hand. Suppose f depends on the first variable only and $|\partial^\gamma f| \le c$ for $|\gamma| \le 2$. Then $\|h(f \otimes 1)gh'\|_{1,\alpha_0} \le C_L c \|hgh'\|_{1,\alpha_0}$, with $C_L = 2 + 8\sqrt{d}$.

This statement is used in Statement K.53 and in Remark F.34.

Statement K.53 (collar weights: bound per anchor block)**.** Let $0 \le j^* < K$. Let F be the set of terms of $E_{K,j^*}(u) - E_{K,j^*+1}(u)$ with a common top line l. By Lemma E.165, cutting l turns the members of F into one renormalised two-leg class G'_{ren}. Its two legs are labelled once for the whole class, with leg 1 at the vertex v_1 the anchor end; they are transported by the partner operations, and ties are broken by one rule compatible with the class. Put $\Psi_F[\Phi] := \sum_\Lambda E(G'_{\mathrm{ren}}, \Lambda, \Phi)$ for two-variable fields Φ, so that the value of F is $\Psi_F[K_l]$. Here $K_l = (uc_{j^*})^{m_l} \tilde{K}_l$ with $m_l \in \{0, 1\}$ and $c_{j^*} = \kappa\mu_*^2 L^2 t_{j^*}^2$; $\tilde{K}_l$ is the index-j^* kernel, and $\|h\tilde{K}_l h'\|_{1,\alpha_0} \le C_K$. Let F have orders (α, β) and m D-lines, l included. Let $\{\chi_a\}_{a\in\mathfrak{B}_{j^*}}$ be a partition of unity over the scale-j^* blocks, with $\operatorname{supp}\chi_a$ within $t_{j^*}/2$ of a and $|\partial^\gamma \chi_a| \le c_\chi t_{j^*}^{-|\gamma|}$ for $|\gamma| \le 2$. Let the localising function h of [3, p. 420] vanish outside the c_h-neighbourhood of $\square(v_1)$, and let the localisation cubes be as in Statement K.48. Then for every $a \in \mathfrak{B}_{j^*}$

$$\big|\Psi_F[(\chi_a \otimes 1)K_l]\big| \le C_A^{\mathrm{FA}} A_{j^*}(F), \qquad A_{j^*}(F) := |e(t_{j^*})|^\alpha \lambda(t_{j^*})^\beta (|u|\kappa\mu_*^2 L^2 t_{j^*}^2)^m,$$

$$C_A^{\mathrm{FA}} := 2^{\bar{n}} 4^{N_\ell} C_{1.33} C_L c_\chi C_K \cdot 2(2c_h{+}5)^d \cdot S(N_s, d, \delta_0) + C_{\mathrm{im}}, \qquad S(N_s, d, \delta) := \sum_{D\ge 0} (c''_d (D{+}2)^d)^{N_s} \mathrm{e}^{-\delta D}.$$

Here $C_{1.33}$ is the constant of Theorem H.24(c), applied to the cut class with a two-variable external field [3, p. 419], C_L is as in Statement K.52, and C_{im} is the constant for the image terms of the torus lines in the proof of Proposition E.175. The factor $2(2c_h + 5)^d$ bounds the number of possible anchor cubes $\square(v_1)$. C_A^{FA} is independent of ε, K, j^*, a, $\ell_T \ge \ell_0$ and the domains, and it is finite at $u = -\eta$ and $M = m_1$.

This statement is used in Remark F.34 (the family-wise anchoring and the collar bound attached to Lemma F.32).
The counterterm gaps.

Statement K.54 (monomial majorants for large masses)**.** Let $\mu_* > 0$ and $\ell_0 \in (0, \infty]$, where $\ell_0 = \infty$ stands for the lattice $\varepsilon\mathbb{Z}^3$. Let φ_3 and b_4 be the functions of Lemma G.2, formed with the majorants $\bar{\tau}, \bar{\psi}, \bar{\delta}, \bar{\Theta}, \bar{J}_s, \bar{J}'_s, \bar{\Lambda}_i$ and the constant $\bar{\omega}$ evaluated at ℓ_0. Then there are $n \in \mathbb{N}$ and numbers $c_1, \dots, c_n \ge 0$, $c'_1, \dots, c'_n \ge 0$, $p_1, \dots, p_n \le 0$, $p'_1, \dots, p'_n \le 0$, depending only on μ_* and ℓ_0, such that for every $M \ge \mu_*$

$$\varphi_3(M) \le \sum_{i=1}^n c_i M^{p_i}, \qquad b_4(M) \le \sum_{i=1}^n c'_i M^{p'_i}.$$

These monomials are obtained as follows: each majorant is replaced by an admissible majorant of the form (constant)$\times M^q$, and $|\mu_*^2 - 2M^2|$ is replaced by $2M^2$. In particular, for every $M_c \geq \mu_*$,

$$\sup_{M \geq M_c} \varphi_3(M) \leq \sum_i c_i M_c^{p_i}, \qquad \sup_{M \geq M_c} b_4(M) \leq \sum_i c_i' M_c^{p_i'}.$$

This statement is used in the proofs of Corollary G.7 and Corollary G.9, hence in the proof of Theorem 6.1(b),(c).

The c-number bounds: the counterterm forests and the Ward identity.

Statement K.55 (Ward identity on the torus)**.** Let T be a torus of side ℓ_T with $\ell_T/\varepsilon \in \mathbb{N}$. Let $\mu_* > 0$, and let a be a real polynomial in (e, λ) with $a(0,0) > 0$. Consider the model on T with the following data:

- the matter action of (2.2), with uniform mass term $\frac{1}{2} a \sum_x \varepsilon^3 |\varphi|^2$;
- the photon Gaussian weight $\mathrm{e}^{-\frac{1}{2}\langle A, (-\Delta_\varepsilon + \mu_*^2) A\rangle}$ on the bond variables.

Then there is a neighbourhood U of $(0,0)$ in $\mathbb{R} \times [0, \infty)$ on which the photon covariance $\mathcal{G}$ is a positive definite operator, smooth in (e, λ). Put $\Pi := (-\Delta_\varepsilon + \mu_*^2) - \mathcal{G}^{-1}$. Then Π is translation invariant, and $\Pi\, d\psi = 0$ for every real function ψ on the sites. Let (α, β) be arbitrary, and let $\Pi^{\alpha\beta}$ be the $e^\alpha \lambda^\beta$ Taylor coefficient of Π at $(0,0)$; it equals the $e^\alpha \lambda^\beta$ coefficient of the 1PI two-photon function. Then $\sum_\nu \Pi^{\alpha\beta}_{\mu\nu}(q) \tilde{q}_\nu = 0$ for all $q \in T^*$ and all μ, where $\tilde{q}_\nu := (\mathrm{e}^{i\varepsilon q_\nu} - 1)/\varepsilon$. In particular $\Pi^{\alpha\beta}_{\mu\nu}(q_1 e_\nu) = 0$ for $q_1 := 2\pi/\ell_T$ and all μ, ν.

This statement is used in the proof of Lemma H.3 and in Proposition H.8, through Lemma H.5: part (a) of that lemma is the case $a = \overline{M}^2 + \delta m_0^2$.

Statement K.56 (Ward identity on $\varepsilon\mathbb{Z}^3$)**.** Fix $\varepsilon > 0$, $\overline{M} > 0$ and $\mu_* > 0$. Let δm_∞^2 be a real polynomial in (e, λ) vanishing at $(0,0)$; for instance, King's counterterm on $\varepsilon\mathbb{Z}^3$. Consider the model on $\varepsilon\mathbb{Z}^3$ with:

- scalar lines $\hat{G}_{\overline{M}}$ and photon lines C_{μ_*};
- the vertices V_n and λ of Lemma H.15;
- the mass counterterm $\frac{1}{2}\delta m_\infty^2 \sum_x \varepsilon^3 |\varphi|^2$.

For every (α, β), let $\Pi^\infty(q)$, $q \in \mathbb{R}^3$, be the $e^\alpha \lambda^\beta$ coefficient of the 1PI two-photon function of this model. This coefficient is the finite sum, over the 1PI two-photon graphs, of the weights of the graphs with their integrands integrated over the Brillouin zone. Then $\sum_\nu \Pi^\infty_{\mu\nu}(q) \tilde{q}_\nu = 0$ for all $q \in \mathbb{R}^3$ and all μ, with $\tilde{q}_\nu = (\mathrm{e}^{i\varepsilon q_\nu} - 1)/\varepsilon$. In particular $\Pi^\infty_{\mu\nu}(0) = 0$.

This statement is used in step (c) of the proof of Lemma H.13, and hence in Proposition H.8.

Statement K.57 (linked clusters and Dyson chains)**.** Let T be a torus with $\ell_T/\varepsilon \in \mathbb{N}$. Index the real scalar components by $a = (x, i)$, $x \in T$, $i \in \{1, 2\}$, and the photon variables by the bonds b. Take the following data:

- a positive definite scalar covariance C^s and a positive definite photon covariance C^A;
- an interaction consisting of the vertices of Lemma H.15, with a constant mass counterterm c (a real polynomial in (e, λ) vanishing at $(0,0)$) entering as $\frac{1}{2} c \sum_x \varepsilon^3 |\varphi(x)|^2$.

All expectations are formal power series in (e, λ) whose coefficients are finite Wick sums. The weight of a labelled graph of content ν is $A(G)/\nu!$, as in Lemma H.15. Put $G^s_{ab} := \langle \varphi_a \varphi_b \rangle$ and $G^A_{bb'} := \langle A_b A_{b'} \rangle - \langle A_b \rangle \langle A_{b'} \rangle$.

(a) Coefficientwise, G^s_{ab} and $G^A_{bb'}$ are the sums of the weights of the connected labelled graphs with external legs φ_a, φ_b, resp. $A_b, A_{b'}$.

(b) Call an amputated connected two-leg piece irreducible if no single line of the leg type separates its two legs. Let K^s_{ab} be the sum of the weights of the irreducible amputated

scalar two-leg pieces with legs at a and b, and define K^A in the same way. Then, as matrices over the formal power series, $G^\sharp = \sum_{k\ge 0} C^\sharp (K^\sharp C^\sharp)^k$ and $(G^\sharp)^{-1} = (C^\sharp)^{-1} - K^\sharp$, for $\sharp \in \{s, A\}$.

(c) Suppose C^s, C^A and the interaction are invariant under the translations $\varphi(x) \mapsto \varphi(x+z)$, $A_{x,\mu} \mapsto A_{x+z,\mu}$ and under the reflection $\varphi(x) \mapsto \varphi(-x)$, $A_{x,\mu} \mapsto -A_{-x-\varepsilon e_\mu,\mu}$. Then K^s and K^A are the sums over the one-particle-irreducible pieces only. This holds for the uniform-mass model with any constant counterterm, and for the C_t-model with counterterm δm_t^2.

(d) Let $C^s = (-\Delta_\varepsilon + M^2)^{-1} \otimes \mathbf{1}_2$ with respect to $\sum_y \varepsilon^3$, and let $\mathcal{G}$ be the operator with kernel G^s with respect to $\sum_y \varepsilon^3$. Put $\Sigma := -\varepsilon^{-3}(K^s - K^s_{\mathrm{ct}})$, where $K^s_{\mathrm{ct}} = -c\,\varepsilon^3 \mathbf{1}$ is the contribution of the single counterterm vertex. Then $\mathcal{G}^{-1} = -\Delta_\varepsilon + M^2 + c + \Sigma$, and in the translation-invariant case $\hat{\mathcal{G}}(p)^{-1} = \hat{p}^2 + M^2 + c + \hat{\Sigma}(p)$.

This statement is used in the proof of Proposition H.8: part (a) at the start of the proof, part (c) in step (iii), and part (b) in step (iv), for the order λ.

The c-number bounds: the weighted-line form.

Statement K.58 (vacuum families at zero background)**.** Assume the hypotheses of Theorem 5.2, in particular (E.fin) and (T.per). Assume $\bar{n} > 12$ and $2\bar{n} + 1 \le 3R$, with R as in (4.1). Let $M \in \{m_1, m_2\}$ and $u \in [-\eta, 1]$. For $f \in \{s, v\}$ let $G_{k,f} = (-\Delta^{\eta_k} + M_f^2 t_k^2 + a_k P_k)^{-1}$ in scale-k units, with $M_s = M$, $M_v = \mu_*$ and the photon taken componentwise. Assume that these kernels and their derivatives of order at most 2 satisfy (H_∞) of Subsection 5.5 with the constants C_0, δ_0 given there. Let F be a family (Lemma H.23) of connected coloured vacuum graphs that occur in $T_{\bar{n}}[\Phi_M(u)]$ on a torus $T = T_j$ of side $\ell_T \ge \ell_0$. Let F have order $e^\alpha \lambda^\beta$, where the orders of the finite counterterm vertices are included, and $m' \ge 2$ lines Γ^B. Let $\mathrm{val}(F)$ be its coefficient of $u^{m'}$. Then

$$|\mathrm{val}(F)| \le C_F\, e^\alpha \lambda^\beta (\kappa \mu_*^2 L^2)^{m'} \varepsilon_0^{\alpha/2+\beta+2m'-3} |T|,$$

with $C_F = C_F(\bar{n}, \alpha_0, L, a, M_b, \delta_0, M\varepsilon_0, \mu_*\varepsilon_0, \ell_0/\varepsilon_0)$ independent of $\varepsilon = 2^{-K}\varepsilon_0$, K, u, κ and $\ell_T \ge \ell_0$.

This statement is used in the proof of Lemma H.9 (step (4)), and through it in Proposition H.11 and Proposition 6.6.

Statement K.59 (finite mass vertices at the top scale)**.** Assume the hypotheses of Theorem 5.2. Let $M \in \{m_1, m_2\}$, $\varepsilon = 2^{-K}\varepsilon_0$ and $T = T_j$. Put $C_f := (-\Delta_\varepsilon + M_f^2)^{-1}$, with $M_s = M$ and $M_v = \mu_*$, and $R_f := C_f - G_{K,f}(0)$. Following [3, (1.26), p. 418], let $\delta m^2_{\mathrm{fin}}(x)$ be the part of the mass counterterm contained in $a_{K,T}(M)$ (Proposition 4.7) that is given by the graphs with at least one line R_f. Write $\delta m^2_{\mathrm{fin}}(x) = \sum_{\alpha,\beta} e^\alpha \lambda^\beta c_{\alpha\beta}(x)$, and let $e(t) := e\, t^{1/2}$ and $\lambda(t) := \lambda t$.

(a) $c_{\alpha\beta} \ne 0$ only if α is even and $o := \alpha/2 + \beta \in \{1, 2\}$.

(b) For every $x \in T$, $\varepsilon_0^2\, |e^\alpha \lambda^\beta c_{\alpha\beta}(x)| \le C_{\mathrm{fin}}\, e(t_K)^\alpha \lambda(t_K)^\beta$, where, with $\rho_0 := \min(M\varepsilon_0, \mu_*\varepsilon_0)$ and $\bar{Z} := 2(M\varepsilon_0)^{-1}\rho_0^{-2}(\varepsilon_0/\ell_0)^3$,

$$C_{\mathrm{fin}} := 5\Big[C_b\Big(\tfrac{L^2}{L-1} + 2 + \log_L \tfrac{1}{M\varepsilon_0}\Big) + (L+1)\, C'\Big(1 + \rho_0^{-2} + \big(\tfrac{8}{\min(\rho_0, \kappa_G)}\big)^3\Big)^5 (1+\bar{Z})^5\Big].$$

Here C_b and $C' = C(\alpha_0, \bar{n}, d, L, a, M_b)$ are the constants of Proposition E.9, and κ_G is the decay rate of Proposition E.50. The constant C_{fin} does not depend on ε, x, u, κ or $\ell_T \ge \ell_0$.

This statement is used in the proof of Lemma H.9, through Statement K.58, for the finite counterterm vertices.

Statement K.60 (torus counterterms in class form)**.** Assume (H_∞) of Subsection 5.5, the condition (E.r) ($\varepsilon_0 < \mathrm{e}^{1-\varrho}$) and (T.per) (5.13). Let $k \le K$, and let $\square \subset T_{\eta_k}$ be a cube of side $s \le 3r(t_k)$ in scale-k units. Let S_k, Γ_k, $\mathcal{K}$ and C_1 be as in Subsection 5.5 and Lemma 5.22.

(a) There is a cube $\tilde{\square} \subset \eta_k\mathbb{Z}^3$ on which the projection π to the torus is a bijection onto $\square$. For $x, x' \in \square$ with lifts $\tilde{x}, \tilde{x}' \in \tilde{\square}$, and for every $n \in \Gamma_k \setminus \{0\}$, $|\tilde{x} - \tilde{x}' - n| \geq \max\{|\tilde{x} - \tilde{x}'|, S_k/2\}$. In particular, the torus distance of x, x' is $|\tilde{x} - \tilde{x}'|$.

(b) For every $\mathsf{K} \in \mathcal{K}$ and $x, x' \in \square$, $\mathsf{K}_T(x, x') = \mathsf{K}_\infty(\tilde{x}, \tilde{x}') + \delta\mathsf{K}(x, x')$ with $\|\delta\mathsf{K}\|_{1,\alpha;\square\times\square} \leq C_1 \mathrm{e}^{-\delta_0 S_k/2} \leq C_1 \mathrm{e}^{-3\delta_0 r(t_k)}$.

(c) Transform a counterterm graph of [3, (1.26)–(1.31)] as the classes are transformed in [3, pp. 433, 445]: localise its vertices, replace the far lines by external fields, and regroup. Let $\square$ be the cube of side $3r(t_k)$ of [3, p. 433] that is attached to one of its connected components. Insert (b) for every scalar line of that component and expand the product. Then:

- under the identification $\square \cong \tilde{\square}$, the term in which every scalar line is $G_k(0)$ is the counterterm in the form used in [3, (3.7)–(3.9), (3.19), pp. 444–445];
- every other term is an expression $E(G'_{\mathrm{ren}}, \{\square(v)\}, \Phi_{\mathrm{ext}}, A_{\mathrm{ext}})$ of the type in [3, (3.3)] in which $a \geq 1$ of the external scalar fields are restrictions of δG_k to $\square \times \square$, so its modulus is at most $O(1)(C_1 \mathrm{e}^{-3\delta_0 r(t_k)})^a$ times the norms of the remaining external fields.

In particular, at $k = K$ and $\tilde{B} = 0$, consider a cube of side $\leq 3r(\varepsilon_0)$ that contains the localisation cubes of a cluster. Each partner of a class supported in this cube is the sum of two parts: its form on $\eta_K\mathbb{Z}^3$, and at most $2^{m(\bar{n})} - 1$ expressions of the type in [3, (3.3)] of the same order. Here $m(\bar{n})$ is the largest number of lines of a graph of order $\leq \bar{n}$. Each of these expressions contains at least one additional external two-variable field of norm $\leq C_1 \mathrm{e}^{-3\delta_0 r(\varepsilon_0)}$, and every graph in them has at most $2\bar{n}$ vertices.

This statement is used in the proof of Lemma H.9, through Statement K.58. It compares the counterterm partners on the torus with their form on $\eta_K\mathbb{Z}^3$.

Statement K.61 (pointwise partner bound)**.** Assume the hypotheses of Statement K.58. Work at the top scale $k = K$, in scale-K units and at zero background. Let G_0 be a C-only self-energy graph of [3, (1.21)–(1.23)], with two scalar legs and order $e^\alpha \lambda^\beta$, whose lines are the kernels $G_{K,s}(0)$, $G_{K,v}(0)$. For $x \in T$, let $\delta m^2_{G_0,T}(x)$ be its counterterm coefficient on T (the expression with leg 1 fixed at x and leg 2 summed), and let $\delta m^2_{G_0,\infty}$ be the same expression on $\eta_K\mathbb{Z}^3$. Then, for every $x \in T$,

$$|\delta m^2_{G_0,T}(x) - \delta m^2_{G_0,\infty}| \leq C_P \, e(t_K)^\alpha \lambda(t_K)^\beta,$$

with $e(t) = e\, t^{1/2}$, $\lambda(t) = \lambda t$ and $C_P = C_P(\bar{n}, \alpha_0, L, a, M_b, \delta_0)$ independent of ε, x and $\ell_T \geq \ell_0$. Consequently, each torus partner of a class equals its form on $\eta_K\mathbb{Z}^3$ plus a finite mass vertex of the same order whose coefficient obeys this bound.

This statement is used in the proof of Lemma H.9, through Statement K.58, for the counterterm partners on the torus.

The domination of the infrared part and the omitted estimates.

Statement K.62 (margins of the two-domain bounds at the first step)**.** Let $k = 1$, so that $\eta_1 = \frac{1}{2}$. Let $X_{\mathrm{in}} \subset X_{\mathrm{out}}$, where X_{in} is a union of blocks and X_{out} is a union of big blocks or T_η. Let A be real and regular on X_{out}, and let $0 \leq m^2 t^2 \leq c$. Put $\delta := G_1(X_{\mathrm{out}}, A) - G_1(X_{\mathrm{in}}, A)$ and $\rho_{\mathrm{in}} := \mathrm{dist}_\infty(\cdot, X_{\mathrm{in}}^c)$, and let $0 < \alpha < 1$, $0 < \beta \leq \beta_D$, $C_D(\alpha)$ and $X_{ab} = D^a \delta D^{b*}$ be as in Lemma E.123. Then:

(a) for all x, x' (and their Hölder partners) with $\rho_{\mathrm{in}}(x), \rho_{\mathrm{in}}(x') \geq R_0 + 12$,

$$\mathrm{e}^{\beta|x-x'|} \max\{\|X_{ab}(x, x')\|, aH_x, bH_{x'}, abH_{xx'}\} \leq C_D(\alpha) \exp\big(-\tfrac{\delta_K}{2}[(\rho_{\mathrm{in}}(x) - R_0 - 7) + (\rho_{\mathrm{in}}(x') - R_0 - 10)]\big);$$

(b) for all functions $0 \leq h, h' \leq 1$ with $|\partial^\eta h|, |\partial^\eta h'| \leq c_h$ and $\mathrm{dist}_\infty(\mathrm{supp}\, h \cup \mathrm{supp}\, h', X_{\mathrm{in}}^c) \geq R_0 + 13$,

$$\|h\, \delta\, h'\|_{1,\alpha_0} \leq 4(1 + 2c_h)^2 C_D(\alpha_0)\, \mathrm{e}^{-\beta\, \mathrm{dist}(\mathrm{supp}\, h, \mathrm{supp}\, h')}.$$

This statement is used in the proof of Lemma I.23 (Step 2, through Lemma E.123 and Proposition E.110) and in Lemma I.24.

Statement K.63 (the bad vacuum localisations at the last scale)**.** Let $k \geq 1$, and let $E_{K,k}$ and $E^{\mathrm{bad}}_{K,k}$ be as in Proposition E.175. Here $E_{K,k}$ consists of the vacuum counterterm pieces of top index k, and $E^{\mathrm{bad}}_{K,k}$ is the part of their localisations that is not cancelled by the generated partners at the step $k \to k+1$ of [3, p. 422]. Write $E_{K,k} = \sum_{\Lambda} F(\Lambda)$ as the sum over localisations $\Lambda = (\square(v))_v$ of [3, p. 420], by unit cubes of the scale-k lattice. Let $\Lambda_7^{(k)}$ be the small-field set of [2, pp. 557–558], and put $W := B^k(\Lambda_7^{(k)})$. Then

$$E^{\mathrm{bad}}_{K,k} = \sum_{\Lambda:\ \Lambda\cap W\neq\emptyset,\ \Lambda\not\subset W} F(\Lambda).$$

This statement is used in the proof of Proposition E.175, in the application of Lemma E.170, and through it in Lemma I.23 (Step 1) and Lemma I.24.

References

[1] T. Bałaban, $(\mathrm{Higgs})_{2,3}$ quantum fields in a finite volume. I. A lower bound, Commun. Math. Phys. **85** (1982), 603–626.

[2] T. Bałaban, $(\mathrm{Higgs})_{2,3}$ quantum fields in a finite volume. II. An upper bound, Commun. Math. Phys. **86** (1982), 555–594.

[3] T. Bałaban, $(\mathrm{Higgs})_{2,3}$ quantum fields in a finite volume. III. Renormalization, Commun. Math. Phys. **88** (1983), 411–445.

[4] T. Bałaban, Regularity and decay of lattice Green's functions, Commun. Math. Phys. **89** (1983), 571–597.

[5] T. Bałaban, D. Brydges, J. Imbrie, A. Jaffe, The mass gap for Higgs models on a unit lattice, Ann. Physics **158** (1984), 281–319.

[6] G. Benfatto, M. Cassandro, G. Gallavotti, F. Nicolò, E. Olivieri, E. Presutti, E. Scacciatelli, Some probabilistic techniques in field theory, Comm. Math. Phys. **59** (1978), 143–166.

[7] T. Bałaban, J. Imbrie, A. Jaffe, Effective action and cluster properties of the abelian Higgs model, Commun. Math. Phys. **114** (1988), 257–315.

[8] D. Brydges, J. Fröhlich, E. Seiler, On the construction of quantized gauge fields. I. General results, Ann. Physics **121** (1979), 227–284.

[9] D. Brydges, J. Fröhlich, E. Seiler, Construction of quantised gauge fields. II. Convergence of the lattice approximation, Commun. Math. Phys. **71** (1980), 159–205.

[10] D. Brydges, J. Fröhlich, E. Seiler, On the construction of quantized gauge fields. III. The two-dimensional abelian Higgs model without cutoffs, Commun. Math. Phys. **79** (1981), 353–399.

[11] A. Chandra, I. Chevyrev, M. Hairer, H. Shen, Stochastic quantisation of Yang–Mills–Higgs in 3D, Invent. Math. **237** (2024), 541–696; arXiv:2201.03487.

[12] I. Chevyrev, H. Shen, Uniqueness of gauge covariant renormalisation of stochastic 3D Yang–Mills–Higgs, arXiv:2503.03060.

[13] C. Dellacherie, P.-A. Meyer, *Probabilities and Potential*, North-Holland Mathematics Studies **29**, North-Holland, Amsterdam, 1978.

[14] J. Dimock, Nonperturbative renormalization of scalar quantum electrodynamics in $d = 3$, J. Math. Phys. **56** (2015), 102304; arXiv:1502.02946.

[15] J. Dimock, Ultraviolet regularity for QED in $d = 3$, J. Math. Phys. **59** (2018), 012301; arXiv:1512.04373.

[16] B. K. Driver, Convergence of the $U(1)_4$ lattice gauge theory to its continuum limit, Commun. Math. Phys. **110** (1987), 479–501.

[17] J. Fröhlich, R. Israel, E. H. Lieb, B. Simon, Phase transitions and reflection positivity. I. General theory and long range lattice models, Commun. Math. Phys. **62** (1978), 1–34.

[18] J. Fröhlich, B. Simon, T. Spencer, Infrared bounds, phase transitions and continuous symmetry breaking, Commun. Math. Phys. **50** (1976), 79–95.

[19] J. Glimm, A. Jaffe, *Quantum Physics. A Functional Integral Point of View*, 2nd ed., Springer, New York, 1987.

[20] M. Göpfert, G. Mack, Proof of confinement of static quarks in 3-dimensional $U(1)$ lattice gauge theory for all values of the coupling constant, Commun. Math. Phys. **82** (1982), 545–606.

[21] L. Gross, Convergence of $U(1)_3$ lattice gauge theory to its continuum limit, Commun. Math. Phys. **92** (1983), 137–162.

[22] S. Janson, *Gaussian Hilbert Spaces*, Cambridge Tracts in Mathematics **129**, Cambridge University Press, 1997.
[23] A. S. Kechris, *Classical Descriptive Set Theory*, Graduate Texts in Mathematics **156**, Springer, New York, 1995.
[24] C. King, The $U(1)$ Higgs model. I. The continuum limit, Commun. Math. Phys. **102** (1986), 649–677.
[25] C. King, The $U(1)$ Higgs model. II. The infinite volume limit, Commun. Math. Phys. **103** (1986), 323–349.
[26] C. M. Newman, Normal fluctuations and the FKG inequalities, Commun. Math. Phys. **74** (1980), 119–128.
[27] K. Osterwalder, R. Schrader, Axioms for Euclidean Green's functions, Commun. Math. Phys. **31** (1973), 83–112.
[28] K. Osterwalder, R. Schrader, Axioms for Euclidean Green's functions. II, Commun. Math. Phys. **42** (1975), 281–305.
[29] K. Osterwalder, E. Seiler, Gauge field theories on a lattice, Ann. Physics **110** (1978), 440–471.
[30] C. J. Preston, A generalization of the FKG inequalities, Commun. Math. Phys. **36** (1974), 233–241.
[31] M. Reed, B. Simon, *Methods of Modern Mathematical Physics. I. Functional Analysis*, revised and enlarged ed., Academic Press, New York, 1980.
[32] T. Reisz, A power counting theorem for Feynman integrals on the lattice, Commun. Math. Phys. **116** (1988), 81–126.
[33] T. Reisz, Renormalization of Feynman integrals on the lattice, Commun. Math. Phys. **117** (1988), 79–108.
[34] E. Seiler, *Gauge Theories as a Problem of Constructive Quantum Field Theory and Statistical Mechanics*, Lecture Notes in Physics **159**, Springer, Berlin, 1982.
[35] F. W. Warner, *Foundations of Differentiable Manifolds and Lie Groups*, Graduate Texts in Mathematics **94**, Springer, New York, 1983.

name	condition at ε_0	depends on
(E.x)	$\hat{x} \le \min\{c_v, e_*, 1\}$, $\lambda\varepsilon_0 \le c_v$	0, 8
(E.1)	$\hat{x}^{1-\beta'} b_0 \hat{\ell}^6 \le c/(C_{\mathrm{reg}} c_1)$, $\beta' = 1/56$	0, 3, 9
(E.2)	$\hat{x} b_0 R\hat{\ell}^8 \le c$; $r(\varepsilon_0) \le Cp(\varepsilon_0)$	0, 5, 9
(E.3)	$\lvert e\rvert \lambda^{-1/4} \varepsilon_0^{1/4} b_0 \hat{\ell}^6 \le (c_1 - 1)/C_{2.21}$	0, 3, 9
(E.4)	$(24/11)(1-\eta)^{-1} C'_H \gamma_1^{-1} \mu_*^{-2} \varepsilon_0^{-2} \mathrm{e}^{-\delta' R\hat{\ell}^2} \le \frac{1}{2}$	0, 3, 5
(E.5)	$2C_H(\gamma_1'' M^2)^{-1} \varepsilon_0^{-2} \mathrm{e}^{-\delta_H R\hat{\ell}^2} \le 1$ and $\varepsilon_0 \le \exp(1 - 1/(\delta_H R))$, with $\gamma_1'' = a_\flat/(a + M^2\varepsilon_0^2)$, at each $M \in \mathbb{M}_4$ (Lemma A.15; the condition is strongest at M_1)	0, 3, 5, 8
(E.6)	$\lambda^{-1/2} \varepsilon_0^{-1/2} b_0^2 \hat{\ell}^{11} \ge C_{2.12}$	0, 9
(E.7)	$p(t)^2 \ge C_\delta(M) t^{1/2} \lambda^{-1/2}$ for $t \le \varepsilon_0$, at each $M \in \mathbb{M}_4$	0, 8, 9
(E.8)	$C_Z \hat{x} b_0 \hat{\ell}^6 \le \frac{1}{2}$	0, 5, 9
(E.9)	$\mathrm{Err}_M(t) \le p(t)$ for all $t \le \varepsilon_0$, at each $M \in \mathbb{M}_4$	0, 3, 5, 6, 9
(E.10)	$R\hat{\ell}^2/4 \ge 2M_b + 2$; $R\hat{\ell}^2 \ge 2(c_M + M_b)$; $a\, c_0 c_1 C_d \varepsilon_0^{-1} \mathrm{e}^{-\delta_0 R\hat{\ell}^2/8} \le \mu_*/2$	0–5
(E.11)	$c_b b_1 \hat{\ell}^{p_1} \ge b_*$; $C_A^{R1} \varepsilon_0^{13/56} (c_b b_0 \hat{\ell}^6)^{\rho_1} \le 1$	0, 7, 9
(E.m2)	$m_2^2 \varepsilon_0^2 \le 1$	0
(E.Φ)	$\lambda^{-1/4} m_2^2 \varepsilon_0^{7/4} \le a_\flat/C_\varphi$, $C_\varphi := c_1 L^{3/4}(1 + \log L)^p$ ($= 39.622\, c_1$ at $L = 2$, $p = 6$)	0, 3
(E.md)	(E.m2) and (E.Φ)	0, 3
(E.12)	$C_\alpha \varepsilon_0^\alpha \le b_0[\hat{\ell}^6 - (\hat{\ell} - \log 2)^6] - 4d\hat{\ell}$	3, 9
(E.13)	$\varepsilon_0 \le \min\{\frac{1}{2}, (15 b_0 \log 2/c_2)^{1/\alpha_I}\}$	3, 8, 9
(E.δ)	$\bar{\delta} := \kappa \mu_*^2 \varepsilon_0^2 \le 1/12$ with the slice constant $\kappa = \frac{15}{2}$ of (5.1), i.e. $\varepsilon_0 \le (90\mu_*^2)^{-1/2}$; it implies (E.dom) (Lemma A.19)	0
(E.14)	$4\kappa c_0^{\mathrm{ker}} c_d (1 + \mu_*^2\varepsilon_0^2)(1-\eta)^{-3} \mathrm{e}^{-\delta_0 R\hat{\ell}^2/4}/(1 - \mathrm{e}^{-\delta_0 R\hat{\ell}\ln 2/2}) \le 1/12$ (Lemma A.17)	0, 3, 5
(E.r)	$\varepsilon_0 < \mathrm{e}^{1-\varrho} = \mathrm{e}^{-1}$	4
(E.15)	$C\varepsilon_0^{\alpha+1/4} \le \frac{1}{2} C_1 b_0 \hat{\ell}^6 \lambda^{-1/4}$	0, 8, 9
(E.dom)	$\mu_* \varepsilon_0 \le 0.2$	0
(E.fin)	$\varepsilon_0 \le \min(1/m_2, 1/\mu_*, L)$	0
(E.charge)	$\varepsilon_0 \le \min\{1, e^{-4}\}$, e the charge (stage 10a); used for the small-field Combes–Thomas bounds of the scalar step (Appendix K)	0
(E.Breg)	$(1+3d) C_\partial\, p(t)\, e(t)^{1-\beta} \le c_{\mathrm{reg}}$ for all $t \le \varepsilon_0$, with $\beta < 1$ the exponent fixed at stage 4 (stage 10a); used for the regularity of the transported background in the scalar Hölder estimate (Appendix K)	0, 3, 4, 9
(E.W)	$4w_j \le \gamma_0$ for every scale $t_j \le \varepsilon_0$, where $w_j := C_W e(t_j) p(t_j)$ (stage 10a); used for the expansion of Bałaban's operator $W^{(j)}$ (Appendix K)	0, 3, 9
(E.ent)	$(1 + \ln(2/\varepsilon_0))^{2p-2-\varrho d} \ge \Theta$ (Proposition 5.21, Lemma A.18)	0–10a
(E.comb)	the cap of the region-sum lemma (Lemma E.95)	0–9
(E.edge)	(5.7) with a_k replaced by a (Lemma A.3); there κ is the exponent of [2, Prop. 2.2, (2.67)], not the slice constant of (E.δ)	0–9
(E.ksp1)–(E.ksp3)	the conditions of Lemma A.17	0–9
(E.vac)	ε_0 is small enough for Theorems 4.1 and 4.2 at M_1, m, m_1, m_2	0–9

TABLE 7. Upper bounds on ε_0 (stages 10a and 10b). Every one of them is downward closed. The conditions (E.5), (E.7), (E.9), (E.ent) and (E.md) are imposed at each $M \in \mathbb{M}_4$ (minimum). The scalar zero mode (Lemma A.12) imposes no condition beyond (E.m2), and the logarithmic entries (Lemma A.16) impose none beyond (E.fin) (Lemma A.15(c)).

name	condition	depends on
(T.floor)	$\ell_0 \geq \ell_0^\natural = 128/M_1$ (so $\ell_0 \geq 183/m$ suffices), $\ell_0 \geq 96/m$, $\ell_0 \geq 128/m$; it implies $\ell_T \geq L^3 = 8$ and $M_1\ell_0 \geq 32$	0
(T.nw)	$\ell_0 \geq \varepsilon_0[8r(\varepsilon_0)+4M_b+2]$: the boxes of radius $O(r(t_k))$ of the proofs, among them the box of Lemma 5.6, do not wrap around the torus	3, 5, 10b
(T.cap)	$\ell_0 \geq 3r(\varepsilon_0)\varepsilon_0 + C_{\bar{N}}\varepsilon_0$ (Lemma A.21)	4, 5, 10b
(T.T)	$\ell_0 \geq \ell_0^T$, the torus side of Theorem E.53	0–10b
(T.div)	$\ell_0 \in 2M_b\mathbb{N}$, $\varepsilon_0 = 2^{-n_0}$	3, 10b
(T.per)	$\ell_0 \geq \varepsilon_0 \max\{6\sqrt{3}\, r(\varepsilon_0) + 2,\ 2/\delta_0\}$ (Lemma 5.22, Lemma A.2(b))	3, 5, 10b

TABLE 8. Lower bounds and divisibility for ℓ_0 (stage 11). Nothing bounds ℓ_T from above or ties it to ε.

use	what is used	type
Cauchy estimates in the fluctuation field	the fluctuation bound (real polynomial V, real symmetric P), the norms [3, (1.32)], the bound [3, (1.16)]	U0: no complex background
analyticity in the background	an analyticity lemma and its use	U0 + U1; not used
pointwise bounds on a route through Cauchy estimates	interior points	U1 / U2–U3 at interior points; not used
the large-field step	real expansions; an optional Cauchy estimate	U0; the optional part U2 at interior points, not used
conditions of the route through Cauchy estimates	upper bounds on ε_0 of that route	not used
lists and renamings	—	no estimate
Cauchy estimates in sources and in (u, b)	—	not a gauge background

TABLE 9. The uses in the proof of Theorem 5.2 of bounds that involve complex parameters. U0: no complex background; U1: at the level of L^2 norms or forms; U2/U3: pointwise or derivative bounds; "at interior points": only at points of depth $\geq R_0$.

item	statement	proved in
1	one-variable bound on the kernel [3, (1.16)] at points of depth $\geq R_0$	Lemma E.116
2	two-variable bound on [3, (1.16)] on T_η	Lemma E.117
3	the same on the Neumann domain Ω, at every evaluation point	Lemma E.118
4	no field-independent mass term other than the δm^2 insertions	Lemma E.119
5, 5′	the abbreviated steps of [2, 3] that need more than the printed text	Lemma E.114
6	separation bound on a cube that is a union of big blocks	Lemma E.120
7	the B'-remainder of [3, p. 433]	Lemma E.121
8	two-domain bounds for nested unions of big blocks, for $0 < \eta \leq \frac{1}{2}$ and for $\eta = 1$	Proposition D.17, Lemmas E.123, E.106
9	$\square \subset \Omega_{\text{out}}$ at every application	Lemma E.124
10	the cube step without $\square \subset \Omega_2$	Lemma E.125
11, 12	depth of the evaluation points of Ω_2-kernels in [2, Chaps. 2, 3]	Lemmas E.126, E.127, E.108
13	nesting of the $\Lambda_i^{(j)}$, and $\Omega \supset \Omega_1 \supset \Omega_2$	Lemma E.100
14	the support hypothesis in the form (R4‴)	Corollary 5.10
15	gating by g_k, metric, contours inside blocks	Lemmas E.128, E.102, E.103
15′	the list of applications is complete	Lemma E.129
16	the $A \cdot A$ local terms	Lemma E.130, Propositions E.140, E.138
17	the tails of [2, p. 579]	Lemma E.141, Propositions E.36, E.145, E.146
–	one background; localisation after the B'-expansion; depth at the applications; tree rate	Lemmas E.105, E.107, E.108, E.109

TABLE 10. The statements that make up Proposition E.110.