

# Nonparametric comparison of epidemic time trends: the case of COVID-19

---

Marina Khismatullina

Erasmus University Rotterdam

Michael Vogt

University of Ulm

IASC-ARS Interim Conference 2022

12 December, 2022

# Table of contents

1. Introduction
2. Model
3. Testing procedure
4. Application

# Introduction

---


## Aim of the paper

To develop new inference methods that allow to *identify* and *locate* differences between epidemic time trends.

# Motivation

## Aim of the paper


To develop new inference methods that allow to *identify* and *locate* differences between epidemic time trends.



# Motivation

## Aim of the paper


To develop new inference methods that allow to *identify* and *locate* differences between epidemic time trends.



# Motivation

## Aim of the paper


To develop new inference methods that allow to *identify* and *locate* differences between epidemic time trends.



# Motivation

## Aim of the paper

To develop new inference methods that allow to *identify* and *locate* differences between epidemic time trends.



**Research question:** Out of many given intervals, how to find those where the trends are significantly different?



## Why is it relevant?

Finding systematic differences between trends = basis for further research

⇒ understanding which government policies are more effective.

## **Why is it relevant?**

Finding systematic differences between trends = basis for further research

⇒ understanding which government policies are more effective.

## **Why is it difficult?**

Testing many hypotheses at the same time = multiple testing problem

⇒ large probability of one true null hypothesis being rejected.

## **Why is it relevant?**

Finding systematic differences between trends = basis for further research

⇒ understanding which government policies are more effective.

## **Why is it difficult?**

Testing many hypotheses at the same time = multiple testing problem

⇒ large probability of one true null hypothesis being rejected.

## **Is it limited to COVID-19?**

No! Our method = general method for comparing nonparametric trends

⇒ new statistical test for equality of nonparametric trend curves.

Comparison of deterministic trends:

- Park et al. (2009), Degras et al. (2012), Zhang et al. (2012), Hidalgo and Lee (2014), Chen and Wu (2019).

Comparison of deterministic trends:

- Park et al. (2009), Degras et al. (2012), Zhang et al. (2012), Hidalgo and Lee (2014), Chen and Wu (2019).

Comparison of deterministic trends:

- Park et al. (2009), Degras et al. (2012), Zhang et al. (2012), Hidalgo and Lee (2014), Chen and Wu (2019).

Multiscale tests:

- Chaudhuri and Marron (1999, 2000), Hall and Heckman (2000), Dümbgen and Spokoiny (2001), Park et al. (2009).

Comparison of deterministic trends:

- Park et al. (2009), Degras et al. (2012), Zhang et al. (2012), Hidalgo and Lee (2014), Chen and Wu (2019).

Multiscale tests:

- Chaudhuri and Marron (1999, 2000), Hall and Heckman (2000), Dümbgen and Spokoiny (2001), Park et al. (2009).

Comparison of deterministic trends:

- Park et al. (2009), Degras et al. (2012), Zhang et al. (2012), Hidalgo and Lee (2014), Chen and Wu (2019).

Multiscale tests:

- Chaudhuri and Marron (1999, 2000), Hall and Heckman (2000), Dümbgen and Spokoiny (2001), Park et al. (2009).

Studies of COVID-19:

- Dong et al. (2020), Gu et al. (2020), Li and Linton (2020), Jiang et al. (2020) and many others.



# Model

---

# Motivation for the model

We observe  $n$  time series  $\mathcal{X}_i = \{X_{it} : 1 \leq t \leq T\}$  of length  $T$ .

# Motivation for the model

We observe  $n$  time series  $\mathcal{X}_i = \{X_{it} : 1 \leq t \leq T\}$  of length  $T$ .

$X_{it}$  are non-negative integers  $\Rightarrow$  can be modelled by a Poisson distribution with time-varying parameter  $\lambda_i(t/T)$ :  $X_{it} \sim P_{\lambda_i(t/T)}$ .

# Motivation for the model

We observe  $n$  time series  $\mathcal{X}_i = \{X_{it} : 1 \leq t \leq T\}$  of length  $T$ .

$X_{it}$  are non-negative integers  $\Rightarrow$  can be modelled by a Poisson distribution with time-varying parameter  $\lambda_i(t/T)$ :  $X_{it} \sim P_{\lambda_i(t/T)}$ .

Since  $\lambda_i(t/T) = \mathbb{E}[X_{it}] = \text{Var}(X_{it})$ , we can rewrite  $X_{it}$  as

$$X_{it} = \lambda_i\left(\frac{t}{T}\right) + u_{it}$$

# Motivation for the model

We observe  $n$  time series  $\mathcal{X}_i = \{X_{it} : 1 \leq t \leq T\}$  of length  $T$ .

$X_{it}$  are non-negative integers  $\Rightarrow$  can be modelled by a Poisson distribution with time-varying parameter  $\lambda_i(t/T)$ :  $X_{it} \sim P_{\lambda_i(t/T)}$ .

Since  $\lambda_i(t/T) = \mathbb{E}[X_{it}] = \text{Var}(X_{it})$ , we can rewrite  $X_{it}$  as

$$X_{it} = \lambda_i\left(\frac{t}{T}\right) + \sqrt{\lambda_i\left(\frac{t}{T}\right)}\eta_{it},$$

where  $\eta_{it}$  has zero mean and unit variance.

# Motivation for the model

We observe  $n$  time series  $\mathcal{X}_i = \{X_{it} : 1 \leq t \leq T\}$  of length  $T$ .

$X_{it}$  are non-negative integers  $\Rightarrow$  can be modelled by a Poisson distribution with time-varying parameter  $\lambda_i(t/T)$ :  $X_{it} \sim P_{\lambda_i(t/T)}$ .

Since  $\lambda_i(t/T) = \mathbb{E}[X_{it}] = \text{Var}(X_{it})$ , we can rewrite  $X_{it}$  as

$$X_{it} = \lambda_i\left(\frac{t}{T}\right) + \sqrt{\lambda_i\left(\frac{t}{T}\right)}\eta_{it},$$

where  $\eta_{it}$  has zero mean and unit variance.

In applications the variance can be larger than the mean  $\Rightarrow$  quasi-Poisson models.

Quasi-Poisson model:

$$X_{it} = \lambda_i\left(\frac{t}{T}\right) + \sigma\sqrt{\lambda_i\left(\frac{t}{T}\right)}\eta_{it},$$

Quasi-Poisson model:

$$X_{it} = \lambda_i \left( \frac{t}{T} \right) + \sigma \sqrt{\lambda_i \left( \frac{t}{T} \right)} \eta_{it},$$

where

- $\sigma$  is the overdispersion parameter;



Quasi-Poisson model:

$$X_{it} = \lambda_i\left(\frac{t}{T}\right) + \sigma \sqrt{\lambda_i\left(\frac{t}{T}\right)} \eta_{it},$$

where

- $\sigma$  is the overdispersion parameter;
- $\lambda_i$  are unknown trend functions on  $[0, 1]$ ;

Quasi-Poisson model:

$$X_{it} = \lambda_i\left(\frac{t}{T}\right) + \sigma \sqrt{\lambda_i\left(\frac{t}{T}\right)} \eta_{it},$$

where

- $\sigma$  is the overdispersion parameter;
- $\lambda_i$  are unknown trend functions on  $[0, 1]$ ;
- $\eta_{it}$  are error terms that are independent across  $i$  and  $t$  and have zero mean and unit variance.

# Testing procedure

---

$$H_0 : \lambda_1 = \lambda_2 = \dots = \lambda_n$$

$$H_0 : \lambda_1 = \lambda_2 = \dots = \lambda_n$$

**Question:** if we reject the global null, how to locate the differences between the trends?

$$H_0 : \lambda_1 = \lambda_2 = \dots = \lambda_n$$

**Question:** if we reject the global null, how to locate the differences between the trends?

Let  $\mathcal{F} := \{\mathcal{I}_k \subseteq [0, 1] : 1 \leq k \leq K\}$  be a family of rescaled time intervals on  $[0, 1]$ , and for each triplet  $(i, j, k)$  consider the null hypothesis that the functions  $\lambda_i$  and  $\lambda_j$  are equal on an interval  $\mathcal{I}_k$

$$H_0 : \lambda_1 = \lambda_2 = \dots = \lambda_n$$

**Question:** if we reject the global null, how to locate the differences between the trends?

Let  $\mathcal{F} := \{\mathcal{I}_k \subseteq [0, 1] : 1 \leq k \leq K\}$  be a family of rescaled time intervals on  $[0, 1]$ , and for each triplet  $(i, j, k)$  consider the null hypothesis that the functions  $\lambda_i$  and  $\lambda_j$  are equal on an interval  $\mathcal{I}_k$ , i.e.

$$H_0^{(ijk)} : \lambda_i(w) = \lambda_j(w) \text{ for all } w \in \mathcal{I}_k$$

# Testing problem

$$H_0 : \lambda_1 = \lambda_2 = \dots = \lambda_n$$

**Question:** if we reject the global null, how to locate the differences between the trends?

Let  $\mathcal{F} := \{\mathcal{I}_k \subseteq [0, 1] : 1 \leq k \leq K\}$  be a family of rescaled time intervals on  $[0, 1]$ , and for each triplet  $(i, j, k)$  consider the null hypothesis that the functions  $\lambda_i$  and  $\lambda_j$  are equal on an interval  $\mathcal{I}_k$ , i.e.

$$H_0^{(ijk)} : \lambda_i(w) = \lambda_j(w) \text{ for all } w \in \mathcal{I}_k$$

We want to test  $H_0^{(ijk)}$  simultaneously for all pairs of countries  $i$  and  $j$  and all intervals  $\mathcal{I}_k$  in the family  $\mathcal{F}$  and we want to control the familywise error rate (FWER) at level  $\alpha$ :

$$\text{FWER}(\alpha) = \mathbb{P}\left(\exists(i, j, k) : \text{we wrongly reject } H_0^{(ijk)}\right).$$



## Test statistic

For a given interval  $\mathcal{I}_k$  and a pair of time series  $i$  and  $j$  we calculate

$$\hat{s}_{ijk} = \frac{1}{Th_k} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right)(X_{it} - X_{jt}),$$

where  $h_k$  is the length of  $\mathcal{I}_k$ .

## Test statistic

For a given interval  $\mathcal{I}_k$  and a pair of time series  $i$  and  $j$  we calculate

$$\hat{s}_{ijk} = \frac{1}{Th_k} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right)(X_{it} - X_{jt}),$$

where  $h_k$  is the length of  $\mathcal{I}_k$ .  $\hat{s}_{ijk}$  estimates the average distance between  $\lambda_i$  and  $\lambda_j$  on  $\mathcal{I}_k$ .

# Test statistic

For a given interval  $\mathcal{I}_k$  and a pair of time series  $i$  and  $j$  we calculate

$$\hat{s}_{ijk} = \frac{1}{Th_k} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right) (X_{it} - X_{jt}),$$

where  $h_k$  is the length of  $\mathcal{I}_k$ .  $\hat{s}_{ijk}$  estimates the average distance between  $\lambda_i$  and  $\lambda_j$  on  $\mathcal{I}_k$ . Under certain assumptions,

$$\text{Var}(\hat{s}_{ijk}) = \frac{\sigma^2}{T^2 h_k^2} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right) \left\{ \lambda_i\left(\frac{t}{T}\right) + \lambda_j\left(\frac{t}{T}\right) \right\},$$

# Test statistic

For a given interval  $\mathcal{I}_k$  and a pair of time series  $i$  and  $j$  we calculate

$$\hat{s}_{ijk} = \frac{1}{Th_k} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right) (X_{it} - X_{jt}),$$

where  $h_k$  is the length of  $\mathcal{I}_k$ .  $\hat{s}_{ijk}$  estimates the average distance between  $\lambda_i$  and  $\lambda_j$  on  $\mathcal{I}_k$ . Under certain assumptions,

$$\text{Var}(\hat{s}_{ijk}) = \frac{\sigma^2}{T^2 h_k^2} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right) \left\{ \lambda_i\left(\frac{t}{T}\right) + \lambda_j\left(\frac{t}{T}\right) \right\},$$

which can be estimated by

$$\widehat{\text{Var}}(\hat{s}_{ijk}) = \frac{\hat{\sigma}^2}{T^2 h_k^2} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right) (X_{it} + X_{jt}),$$

with  $\hat{\sigma}^2$  being an appropriate estimator of  $\sigma^2$ . [Details](#)

Test statistic for the hypothesis  $H_0^{(ijk)}$  is then defined as

$$\hat{\psi}_{ijk} := \frac{\hat{S}_{ijk}}{\sqrt{\widehat{\text{Var}}(\hat{S}_{ijk})}} = \frac{\sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right)(X_{it} - X_{jt})}{\hat{\sigma}\left\{\sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right)(X_{it} + X_{jt})\right\}^{1/2}}$$

How to construct critical values  $c_{ijk}(\alpha)$ ?

How to construct critical values  $c_{ijk}(\alpha)$ ?

- Traditional approach:  $c_{ijk}(\alpha) = c(\alpha)$  for all  $(i, j, k)$ .

How to construct critical values  $c_{ijk}(\alpha)$ ?

- Traditional approach:  $c_{ijk}(\alpha) = c(\alpha)$  for all  $(i, j, k)$ .
- More modern approach:  $c_{ijk}(\alpha)$  depend on the length  $h_k$  of the time interval (Dümbgen and Spokoiny (2001)):

$$c_{ijk}(\alpha) = c(\alpha, h_k) := b_k + q(\alpha)/a_k,$$

where  $a_k$  and  $b_k$  are scale-dependent constants and  $q(\alpha)$  is chosen such that we control FWER. [Details](#)



## Critical values, part 2

We want to control FWER.

## Critical values, part 2

We want to control FWER. Let  $\mathcal{M}_0 := \{(i, j, k) | H_0^{(ijk)} \text{ is true}\}$ , then

$$\text{FWER}(\alpha) = \mathbb{P}\left(\exists(i, j, k) \in \mathcal{M}_0 : |\hat{\psi}_{ijk}| > c_{ijk}(\alpha)\right)$$

## Critical values, part 2

We want to control FWER. Let  $\mathcal{M}_0 := \{(i, j, k) | H_0^{(ijk)} \text{ is true}\}$ , then

$$\begin{aligned}\text{FWER}(\alpha) &= \mathbb{P}\left(\exists(i, j, k) \in \mathcal{M}_0 : |\hat{\psi}_{ijk}| > c_{ijk}(\alpha)\right) \\&= 1 - \mathbb{P}\left(\forall(i, j, k) \in \mathcal{M}_0 : |\hat{\psi}_{ijk}| \leq c_{ijk}(\alpha)\right) \\&= 1 - \mathbb{P}\left(\forall(i, j, k) \in \mathcal{M}_0 : a_k(|\hat{\psi}_{ijk}| - b_k) \leq q(\alpha)\right) \\&= 1 - \mathbb{P}\left(\max_{(i, j, k) \in \mathcal{M}_0} a_k(|\hat{\psi}_{ijk}| - b_k) \leq q(\alpha)\right) \\&\leq 1 - \mathbb{P}\left(\max_{(i, j, k)} a_k(|\hat{\psi}_{ijk}^0| - b_k) \leq q(\alpha)\right)\end{aligned}$$

## Critical values, part 2

We want to control FWER. Let  $\mathcal{M}_0 := \{(i, j, k) | H_0^{(ijk)} \text{ is true}\}$ , then

$$\begin{aligned}\text{FWER}(\alpha) &= \mathbb{P}\left(\exists(i, j, k) \in \mathcal{M}_0 : |\hat{\psi}_{ijk}| > c_{ijk}(\alpha)\right) \\&= 1 - \mathbb{P}\left(\forall(i, j, k) \in \mathcal{M}_0 : |\hat{\psi}_{ijk}| \leq c_{ijk}(\alpha)\right) \\&= 1 - \mathbb{P}\left(\forall(i, j, k) \in \mathcal{M}_0 : a_k(|\hat{\psi}_{ijk}| - b_k) \leq q(\alpha)\right) \\&= 1 - \mathbb{P}\left(\max_{(i, j, k) \in \mathcal{M}_0} a_k(|\hat{\psi}_{ijk}| - b_k) \leq q(\alpha)\right) \\&\leq 1 - \mathbb{P}\left(\max_{(i, j, k)} a_k(|\hat{\psi}_{ijk}^0| - b_k) \leq q(\alpha)\right)\end{aligned}$$

## Critical values, part 2

We want to control FWER. Let  $\mathcal{M}_0 := \{(i, j, k) | H_0^{(ijk)} \text{ is true}\}$ , then

$$\begin{aligned}\text{FWER}(\alpha) &= P\left(\exists(i, j, k) \in \mathcal{M}_0 : |\hat{\psi}_{ijk}| > c_{ijk}(\alpha)\right) \\&= 1 - P\left(\forall(i, j, k) \in \mathcal{M}_0 : |\hat{\psi}_{ijk}| \leq c_{ijk}(\alpha)\right) \\&= 1 - P\left(\forall(i, j, k) \in \mathcal{M}_0 : a_k(|\hat{\psi}_{ijk}| - b_k) \leq q(\alpha)\right) \\&= 1 - P\left(\max_{(i, j, k) \in \mathcal{M}_0} a_k(|\hat{\psi}_{ijk}| - b_k) \leq q(\alpha)\right) \\&\leq 1 - P\left(\max_{(i, j, k)} a_k(|\hat{\psi}_{ijk}^0| - b_k) \leq q(\alpha)\right)\end{aligned}$$

Hence, we choose  $q(\alpha)$  as the  $(1 - \alpha)$ -quantile of the statistic

$$\hat{\Psi}_T = \max_{(i, j, k)} a_k(|\hat{\psi}_{ijk}^0| - b_k),$$

where  $\hat{\psi}_{ijk}^0$  is equal to  $\hat{\psi}_{ijk}$  under the null.

But we do not know the distribution of  $\hat{\Psi}_T$  in practice!

But we do not know the distribution of  $\hat{\Psi}_T$  in practice!

$\Rightarrow$  the quantiles  $q(\alpha)$  are also not known. How to approximate them?

## Critical values, part 3

But we do not know the distribution of  $\hat{\Psi}_T$  in practice!

$\Rightarrow$  the quantiles  $q(\alpha)$  are also not known. How to approximate them?

Under our assumptions,

$$\hat{\psi}_{ijk}^0 \approx \frac{1}{\sqrt{2Th_k}} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right)(\eta_{it} - \eta_{jt}),$$

can be approximated by a Gaussian version of the test statistic:

$$\phi_{ijk} = \frac{1}{\sqrt{2Th_k}} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right)(Z_{it} - Z_{jt}),$$

where  $Z_{it}$  are independent standard normal random variables.



# Test procedure

1. Consider the Gaussian test statistic

$$\Phi_T = \max_{(i,j,k)} a_k (|\phi_{ijk}| - b_k),$$

where  $a_k$  and  $b_k$  are scale-dependent constants and  $\phi_{ijk}$  are weighted averages of the differences of standard normal random variables.

# Test procedure

1. Consider the Gaussian test statistic

$$\Phi_T = \max_{(i,j,k)} a_k (|\phi_{ijk}| - b_k),$$

where  $a_k$  and  $b_k$  are scale-dependent constants and  $\phi_{ijk}$  are weighted averages of the differences of standard normal random variables.

2. Compute a  $(1 - \alpha)$ -quantile  $q_{\text{Gauss}}(\alpha)$  of  $\Phi_T$  by Monte Carlo simulations.

# Test procedure

1. Consider the Gaussian test statistic

$$\Phi_T = \max_{(i,j,k)} a_k (|\phi_{ijk}| - b_k),$$

where  $a_k$  and  $b_k$  are scale-dependent constants and  $\phi_{ijk}$  are weighted averages of the differences of standard normal random variables.

2. Compute a  $(1 - \alpha)$ -quantile  $q_{\text{Gauss}}(\alpha)$  of  $\Phi_T$  by Monte Carlo simulations.
3. Adjust  $q_{\text{Gauss}}(\alpha)$  by the scale-dependent constants

$$c_{\text{Gauss}}(\alpha, h_k) = b_k + q_{\text{Gauss}}(\alpha)/a_k$$

# Test procedure

1. Consider the Gaussian test statistic

$$\Phi_T = \max_{(i,j,k)} a_k (|\phi_{ijk}| - b_k),$$

where  $a_k$  and  $b_k$  are scale-dependent constants and  $\phi_{ijk}$  are weighted averages of the differences of standard normal random variables.

2. Compute a  $(1 - \alpha)$ -quantile  $q_{\text{Gauss}}(\alpha)$  of  $\Phi_T$  by Monte Carlo simulations.
3. Adjust  $q_{\text{Gauss}}(\alpha)$  by the scale-dependent constants

$$c_{\text{Gauss}}(\alpha, h_k) = b_k + q_{\text{Gauss}}(\alpha)/a_k$$

## Test procedure

For the given significance level  $\alpha \in (0, 1)$  and for each  $(i, j, k)$ , reject  $H_0^{(ijk)}$  if  $|\hat{\psi}_{ijk}| > c_{\text{Gauss}}(\alpha, h_k)$ .

## Proposition

*Let  $\mathcal{M}_0$  be the set of triplets  $(i, j, k)$  for which  $H_0^{(ijk)}$  holds true. Then under certain assumptions, it holds that*

$$P\left(\forall (i, j, k) \in \mathcal{M}_0 : |\hat{\psi}_{ijk}| \leq c_{\text{Gauss}}(\alpha, h_k)\right) \geq 1 - \alpha + o(1)$$

## Proposition

*Let  $\mathcal{M}_0$  be the set of triplets  $(i, j, k)$  for which  $H_0^{(ijk)}$  holds true. Then under certain assumptions, it holds that*

$$P\left(\forall (i, j, k) \in \mathcal{M}_0 : |\hat{\psi}_{ijk}| \leq c_{\text{Gauss}}(\alpha, h_k)\right) \geq 1 - \alpha + o(1)$$

## Corollary

$$FWER(\alpha) \leq \alpha.$$

### Proposition

Consider a sequence of functions  $\lambda_i = \lambda_{i,T}$ ,  $\lambda_j = \lambda_{j,T}$  such that

$$\exists \mathcal{I}_k : \lambda_i(w) - \lambda_j(w) \geq c_T \sqrt{\log T / (Th_k)} \quad \forall w \in \mathcal{I}_k, \quad (1)$$

and  $c_T \rightarrow \infty$  faster than  $\frac{\sqrt{\log T} \sqrt{\log \log T}}{\log \log \log T}$ . Let  $\mathcal{M}_1$  be the set of triplets  $(i, j, k)$  for which (1) holds true. Then under certain assumptions, it holds that

$$\mathbb{P}\left(\forall (i, j, k) \in \mathcal{M}_1 : |\hat{\psi}_{ijk}| > c_{\text{Gauss}}(\alpha, h_k)\right) = 1 - o(1)$$

# Application


---





# Application setting

- Five countries: Germany, Italy, Spain, France and the UK.
- $T = 150$  days.
- The data is aligned by weekdays: first Monday after reaching 100 cases as  $t = 1$ .
- Lengths of time intervals 7, 14, 21, 28 days. The intervals start at days 1, 8, 15, ... and 4, 11, 19, ...
- $\alpha = 0.05$ .
- 5000 Monte Carlo simulation runs to produce critical values.


# Application results







(b) (minimal) intervals produced by our test



(b) (minimal) intervals produced by our test



# Application results, part 2



We can claim, with confidence of about 95%, that the null hypothesis is violated for all intervals (and all pairs of countries) for which our test rejects the null.

We can claim, with confidence of about 95%, that the null hypothesis is violated for all intervals (and all pairs of countries) for which our test rejects the null.

However, we can not say anything about the causes of such differences. This question requires further (probably not purely statistical) analysis.

We can claim, with confidence of about 95%, that the null hypothesis is violated for all intervals (and all pairs of countries) for which our test rejects the null.

However, we can not say anything about the causes of such differences. This question requires further (probably not purely statistical) analysis.

Further possible extensions:

- introduce scaling factor in the trend function, that will allow to adjust for the size of the country (population, density, testing regimes, etc.);
- include dependence in the error terms;
- cluster the countries based on the trends they exhibit.

# Where to find more?

## Contact information:

- <https://marina-khi.github.io>
- <https://github.com/marina-khi/multiscale>
- [khismatullina@ese.eur.nl](mailto:khismatullina@ese.eur.nl)

## Reference:

- Khismatullina, M. and Vogt, M. (2021). Nonparametric comparison of epidemic time trends: the case of COVID-19. *Journal of Econometrics*.

**Thank you!**




# Assumptions


- $\mathcal{C1}$  The functions  $\lambda_i$  are uniformly Lipschitz continuous:  
 $|\lambda_i(u) - \lambda_i(v)| \leq L|u - v|$  for all  $u, v \in [0, 1]$ .
- $\mathcal{C2}$   $0 < \lambda_{\min} \leq \lambda_i(w) \leq \lambda_{\max} < \infty$  for all  $w \in [0, 1]$  and all  $i$ .
- $\mathcal{C3}$   $\eta_{it}$  are independent both across  $i$  and  $t$ .
- $\mathcal{C4}$   $\mathbb{E}[\eta_{it}] = 0$ ,  $\mathbb{E}[\eta_{it}^2] = 1$  and  $\mathbb{E}[|\eta_{it}|^\theta] \leq C_\theta < \infty$  for some  $\theta > 4$ .
- $\mathcal{C5}$   $h_{\max} = o(1/\log T)$  and  $h_{\min} \geq CT^{-b}$  for some  $b \in (0, 1)$ .
- $\mathcal{C6}$   $p := \{\#(i, j, k)\} = O(T^{(\theta/2)(1-b)-(1+\delta)})$  for some small  $\delta > 0$ .

# Family of time intervals

The family of intervals  $F$




# Simulation results for the size of the test



**Table 1:** Size of the multiscale test

	$n = 5$			$n = 10$			$n = 50$		
	significance level $\alpha$			significance level $\alpha$			significance level $\alpha$		
	0.01	0.05	0.1	0.01	0.05	0.1	0.01	0.05	0.1
$T = 100$	0.011	0.047	0.093	0.010	0.044	0.087	0.008	0.037	0.075
$T = 250$	0.009	0.047	0.091	0.009	0.046	0.087	0.008	0.035	0.069
$T = 500$	0.010	0.044	0.083	0.008	0.048	0.093	0.007	0.035	0.077



# Simulation results for the power of the test



**Table 2:** Power of the multiscale test for scenario A

	$n = 5$			$n = 10$			$n = 50$		
	significance level $\alpha$			significance level $\alpha$			significance level $\alpha$		
	0.01	0.05	0.1	0.01	0.05	0.1	0.01	0.05	0.1
$T = 100$	0.335	0.518	0.597	0.306	0.474	0.545	0.212	0.352	0.418
$T = 250$	0.615	0.790	0.836	0.580	0.764	0.800	0.470	0.648	0.705
$T = 500$	0.736	0.905	0.917	0.738	0.884	0.890	0.636	0.799	0.830

# Simulation results for the power of the test



**Table 3:** Power of the multiscale test for scenario B

	$n = 5$			$n = 10$			$n = 50$		
	significance level $\alpha$			significance level $\alpha$			significance level $\alpha$		
	0.01	0.05	0.1	0.01	0.05	0.1	0.01	0.05	0.1
$T = 100$	0.824	0.910	0.903	0.812	0.893	0.890	0.738	0.847	0.857
$T = 250$	0.991	0.972	0.941	0.991	0.960	0.920	0.991	0.965	0.933
$T = 500$	0.997	0.973	0.949	0.995	0.961	0.923	0.996	0.969	0.932

## Estimator of $\sigma^2$

We estimate the overdispersion parameter  $\sigma^2$  by

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n \hat{\sigma}_i^2 \text{ and } \hat{\sigma}_i^2 = \frac{\sum_{t=2}^T (X_{it} - X_{it-1})^2}{2 \sum_{t=1}^T X_{it}}$$

## Estimator of $\sigma^2$

We estimate the overdispersion parameter  $\sigma^2$  by

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n \hat{\sigma}_i^2 \text{ and } \hat{\sigma}_i^2 = \frac{\sum_{t=2}^T (X_{it} - X_{it-1})^2}{2 \sum_{t=1}^T X_{it}}$$

We assume that  $\lambda_i$  is Lipschitz continuous. Then

$$X_{it} - X_{it-1} = \sigma \sqrt{\lambda_i \left( \frac{t}{T} \right)} (\eta_{it} - \eta_{it-1}) + r_{it},$$

where  $|r_{it}| \leq C(1 + |\eta_{it-1}|)/T$  with a sufficiently large  $C$ .

## Estimator of $\sigma^2$

We estimate the overdispersion parameter  $\sigma^2$  by

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n \hat{\sigma}_i^2 \text{ and } \hat{\sigma}_i^2 = \frac{\sum_{t=2}^T (X_{it} - X_{it-1})^2}{2 \sum_{t=1}^T X_{it}}$$

We assume that  $\lambda_i$  is Lipschitz continuous. Then

$$X_{it} - X_{it-1} = \sigma \sqrt{\lambda_i \left( \frac{t}{T} \right)} (\eta_{it} - \eta_{it-1}) + r_{it},$$

where  $|r_{it}| \leq C(1 + |\eta_{it-1}|)/T$  with a sufficiently large  $C$ . Hence,

$$\frac{1}{T} \sum_{t=2}^T (X_{it} - X_{it-1})^2 = 2\sigma^2 \left\{ \frac{1}{T} \sum_{t=2}^T \lambda_i(t/T) \right\} + o_p(1)$$



## Estimator of $\sigma^2$

We estimate the overdispersion parameter  $\sigma^2$  by

$$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n \hat{\sigma}_i^2 \text{ and } \hat{\sigma}_i^2 = \frac{\sum_{t=2}^T (X_{it} - X_{it-1})^2}{2 \sum_{t=1}^T X_{it}}$$

We assume that  $\lambda_i$  is Lipschitz continuous. Then

$$X_{it} - X_{it-1} = \sigma \sqrt{\lambda_i \left( \frac{t}{T} \right)} (\eta_{it} - \eta_{it-1}) + r_{it},$$

where  $|r_{it}| \leq C(1 + |\eta_{it-1}|)/T$  with a sufficiently large  $C$ . Hence,

$$\frac{1}{T} \sum_{t=2}^T (X_{it} - X_{it-1})^2 = 2\sigma^2 \left\{ \frac{1}{T} \sum_{t=2}^T \lambda_i(t/T) \right\} + o_p(1)$$

Together with

$$\frac{1}{T} \sum_{t=1}^T X_{it} = \frac{1}{T} \sum_{t=1}^T \lambda_i(t/T) + o_p(1),$$

we get that  $\hat{\sigma}_i^2 = \sigma^2 + o_p(1)$  for any  $i$  and thus  $\hat{\sigma}^2 = \sigma^2 + o_p(1)$ .

[Go back](#)

# Notation

In order to proceed with the proof, we will need the following notation:

$$\begin{aligned}\hat{\psi}_{ijk,T} &= \frac{\sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right)(X_{it} - X_{jt})}{\hat{\sigma}\left\{\sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right)(X_{it} + X_{jt})\right\}^{1/2}} \\ \hat{\psi}_{ijk,T}^0 &= \frac{\sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right) \sigma \bar{\lambda}_{ij}^{-1/2}\left(\frac{t}{T}\right)(\eta_{it} - \eta_{jt})}{\hat{\sigma}\left\{\sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right)(X_{it} + X_{jt})\right\}^{1/2}} & \hat{\Psi}_T &= \max_{(i,j,k)} a_k(|\hat{\psi}_{ijk,T}^0| - b_k) \\ \psi_{ijk,T}^0 &= \frac{1}{\sqrt{2Th_k}} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right)(\eta_{it} - \eta_{jt}) & \Psi_T &= \max_{(i,j,k)} a_k(|\psi_{ijk,T}^0| - b_k) \\ \phi_{ijk,T} &= \frac{1}{\sqrt{2Th_k}} \sum_{t=1}^T 1\left(\frac{t}{T} \in \mathcal{I}_k\right)(Z_{it} - Z_{jt}) & \Phi_T &= \max_{(i,j,k)} a_k(|\phi_{ijk,T}| - b_k)\end{aligned}$$

## Strategy of the proof

1. We prove that  $|\hat{\psi}_T^0 - \psi_T| = o_p(r_T)$ , where  $\{r_T\}$  is some null sequence.

## Strategy of the proof

1. We prove that  $|\hat{\Psi}_T^0 - \Psi_T| = o_p(r_T)$ , where  $\{r_T\}$  is some null sequence.
2. With the help of results from Chernozhukov et al. (2017), we prove

$$\sup_{q \in \mathbb{R}} \left| P(\Psi_T \leq q) - P(\Phi_T \leq q) \right| = o(1)$$

# Strategy of the proof

1. We prove that  $|\hat{\Psi}_T^0 - \Psi_T| = o_p(r_T)$ , where  $\{r_T\}$  is some null sequence.
2. With the help of results from Chernozhukov et al. (2017), we prove

$$\sup_{q \in \mathbb{R}} \left| P(\Psi_T \leq q) - P(\Phi_T \leq q) \right| = o(1)$$

3. By using these two results, we now show that

$$\sup_{q \in \mathbb{R}} \left| P(\hat{\Psi}_T^0 \leq q) - P(\Phi_T \leq q) \right| = o(1) \quad (2)$$

# Strategy of the proof

1. We prove that  $|\hat{\Psi}_T^0 - \Psi_T| = o_p(r_T)$ , where  $\{r_T\}$  is some null sequence.
2. With the help of results from Chernozhukov et al. (2017), we prove

$$\sup_{q \in \mathbb{R}} \left| P(\Psi_T \leq q) - P(\Phi_T \leq q) \right| = o(1)$$

3. By using these two results, we now show that

$$\sup_{q \in \mathbb{R}} \left| P(\hat{\Psi}_T^0 \leq q) - P(\Phi_T \leq q) \right| = o(1) \quad (2)$$

4. It can be shown that  $P(\Phi_T \leq q_{\text{Gauss}}(\alpha)) = 1 - \alpha$ . From this and (2), it immediately follows that

$$P(\hat{\Psi}_T^0 \leq q_{\text{Gauss}}(\alpha)) = 1 - \alpha + o(1),$$

which in turn implies the desired statement.

## Idea behind $a_k$ and $b_k$

Dümbgen and Spokoiny (2001): the critical values  $c_{ijk}(\alpha)$  depend on the scale of the testing problem, i.e. the length  $h_k$  of the time interval.

## Idea behind $a_k$ and $b_k$

Dümbgen and Spokoiny (2001): the critical values  $c_{ijk}(\alpha)$  depend on the scale of the testing problem, i.e. the length  $h_k$  of the time interval.

Specifically,


$$c_{ijk}(\alpha) = c(\alpha, h_k) := b_k + q(\alpha)/a_k,$$

where  $a_k = \{\log(e/h_k)\}^{1/2} / \log \log(e^e/h_k)$  and  $b_k = \sqrt{2 \log(1/h_k)}$  are scale-dependent constants and  $q(\alpha)$  is chosen such that we control FWER.



## Idea behind $a_k$ and $b_k$ , part 2

This choice of scale-dependent constants helps us balance the significance of hypotheses between the time intervals of different lengths  $h_k$ :



[Go back](#)

# Idea behind the additive correction

Consider the uncorrected Gaussian statistic

$$\Phi^{\text{uncor}} = \max_{(i,j,k)} |\phi_{ijk}|$$

# Idea behind the additive correction

Consider the uncorrected Gaussian statistic

$$\Phi^{\text{uncor}} = \max_{(i,j,k)} |\phi_{ijk}|$$

and let the family of intervals be

$$\mathcal{F} = \{[(m-1)h_l, mh_l] \text{ for } 1 \leq m \leq 1/h_l, 1 \leq l \leq L\}$$

# Idea behind the additive correction

Consider the uncorrected Gaussian statistic

$$\Phi^{\text{uncor}} = \max_{(i,j,k)} |\phi_{ijk}|$$

and let the family of intervals be

$$\mathcal{F} = \{[(m-1)h_l, mh_l] \text{ for } 1 \leq m \leq 1/h_l, 1 \leq l \leq L\}$$

Then we can rewrite the uncorrected test statistic as

$$\Phi^{\text{uncor}} = \max_{i,j} \max_{\substack{1 \leq l \leq L, \\ 1 \leq m \leq 1/h_l}} \left| \frac{1}{\sqrt{2Th_l}} \sum_{t=1}^T 1\left(\frac{t}{T} \in [(m-1)h_l, mh_l]\right) (Z_{it} - Z_{jt}) \right|$$

# Idea behind the additive correction

Consider the uncorrected Gaussian statistic

$$\Phi^{\text{uncor}} = \max_{(i,j,k)} |\phi_{ijk}|$$

and let the family of intervals be

$$\mathcal{F} = \{[(m-1)h_l, mh_l] \text{ for } 1 \leq m \leq 1/h_l, 1 \leq l \leq L\}$$

Then we can rewrite the uncorrected test statistic as

$$\Phi^{\text{uncor}} = \max_{i,j} \max_{\substack{1 \leq l \leq L, \\ 1 \leq m \leq 1/h_l}} \left| \frac{1}{\sqrt{2Th_l}} \sum_{t=1}^T 1\left(\frac{t}{T} \in [(m-1)h_l, mh_l]\right) (Z_{it} - Z_{jt}) \right|$$

$\Rightarrow \max_m \dots = \sqrt{2 \log(1/h_l)} + o_P(1) \rightarrow \infty$  as  $h \rightarrow 0$  and the stochastic behavior of  $\Phi^{\text{uncor}}$  is dominated by the elements with small bandwidths  $h_l$ . [Go back](#)